PSD Representations for Effective Probability Models

Abstract

Finding a good way to model probability densities is key to probabilistic inference. An ideal model should be able to concisely approximate any probability, while being also compatible with two main operations: multiplications of two models (product rule) and marginalization with respect to a subset of the random variables (sum rule). In this work, we show that a recently proposed class of positive semi-definite (PSD) models for non-negative functions is particularly suited to this end. In particular, we characterize both approximation and generalization capabilities of PSD models, showing that they enjoy strong theoretical guarantees. Moreover, we show that we can perform efficiently both sum and product rule in closed form via matrix operations, enjoying the same versatility of mixture models. Our results open the way to applications of PSD models to density estimation, decision theory and inference. Preliminary empirical evaluation supports our findings.

1   Introduction

Modeling probability distributions is a key task for many applications in machine learning [17, 5]. To this end, several strategies have been proposed in the literature, such as adopting mixture models (e.g. Gaussian mixtures) [5], exponential models [31], implicit generative models [12, 14] or kernel (conditional) mean embeddings [16]. An ideal probabilistic model should have two main features: i) efficiently perform key operations for probabilistic inference, such as sum rule (i.e. marginalization) and product rule [5] and, ii) concisely approximate a large class of probabilities. Finding models that satisfy these two conditions is challenging and current methods tend to tackle only one of the two. For instance, mixture models are designed to efficient integrate and multiply probabilities, but tend to require many kernels to approximate complicated distributions (as discussed in more detail in Sec. 3). On the opposite, exponential and implicit generative models have typically strong approximation properties (see e.g. [31, 29]) but cannot easily perform operations such as marginalization.

In this work we propose to adopt the postive semidefinite (PSD) models recently proposed in [15] to model probabilities. PSD models generalize mixture models by allowing also for negative weights, but guarantee by construction to still obtain a non-negative function. We show that this generalization is crucial to overcome the limitations of mixture models and to satisfy the two key features of an ideal model: first, we show that PSD models enjoy the same computational advantages of mixture models in terms of operations between probabilities. Then, we focus on their theoretical properties, showing that they can approximate and learn (very concisely) a wide class of target densities. In particular, we recover the optimal rates achieved by previous estimators that do not guarantee non-negativity [34].

Contributions. The main contributions of this paper are: i) showing that PSD models can perform exact sum and product rules in terms of efficient matrix operations (Sec. 2.1); ii) Providing a "compression" method to control the number of support point of a PSD model (Sec. 2.2); iii) Characterizing the approximation and learning properties of PSD models with respect to a large family of probability densities (Sec. 3); iv) Discussing a number of possible applications of PSD models (Sec. 4) and provide a preliminary empirical investigation (Sec. 5).

Submitted to 35th Conference on Neural Information Processing Systems (NeurIPS 2021). Do not distribute.

Notation. We denote by $\mathbb{R}^d_{++}$ the space vectors in $\mathbb{R}^d$ with positive entries, $\mathbb{R}^{n \times d}$ the space of $n \times d$ matrices, $\mathbb{S}^n_+ = \mathbb{S}_+(\mathbb{R}^n)$ the space of positive semidefinite $n \times n$ matrices. Given a vector $\eta \in \mathbb{R}^d$, we denote $\mathrm{diag}(\eta) \in \mathbb{R}^{d \times d}$ the diagonal matrix associated to η. We denote by $A \circ B$ and $A \otimes B$ respectively the entry-wise and Kronecker product between two matrices A and B. We denote by $\|A\|_F, \|A\|, \det(A), \mathrm{vec}(A)$ and $A^\top$ respectively the Frobenius norm, the operator norm (i.e. maximum singular value), the determinant, the (column-wise) vectorization of a matrix and the (conjugate) transpose of A. With some abuse of notation, where clear from context we write element-wise products and division of vectors $u, v \in \mathbb{R}^d$ as $uv, u/v$. Given two matrices $X \in \mathbb{R}^{n \times d_1}, Y \in \mathbb{R}^{n \times d_2}$ with same number of rows, we denote by $[X, Y] \in \mathbb{R}^{n \times (d_1 + d_2)}$ their concatenation row-wise. The term $\mathbf{1}_n \in \mathbb{R}^n$ denotes the vector with all entries equal to 1.

2 PSD Models

Following [15], in this work we consider the family of positive semi-definite (PSD) models, namely non-negative functions parametrized by a feature map $\phi : \mathcal{X} \to \mathcal{H}$ from an input space $\mathcal{X}$ to a suitable feature space $\mathcal{H}$ (a separable Hilbert space e.g. $\mathbb{R}^q$) and a linear operator $\mathsf{M} \in \mathbb{S}_+(\mathcal{H})$, of the form

$$f(x \, ; \, \mathsf{M}, \phi) = \phi(x)^\top \mathsf{M} \, \phi(x). \tag{1}$$

PSD models offer a general way to parametrize non-negative functions (since M is positive semidefinite, $f(x \, ; \, \mathsf{M}, \phi) \geq 0$ for any $x \in \mathcal{X}$) and enjoy several additional appealing properties discussed in the following. In this work, we will focus on a special family of models of the form (1) to parametrize probability densities over $\mathcal{X} = \mathbb{R}^d$. In particular, we will consider the case where: i) $\phi = \phi_\eta : \mathbb{R}^d \to \mathcal{H}_\eta$ is a feature map associated to the Gaussian kernel [26] $k_\eta(x, x') = \phi_\eta(x)^\top \phi_\eta(x) = e^{-(x-x')^\top \mathrm{diag}(\eta)(x-x')}$, with $\eta \in \mathbb{R}^d_{++}$ and, ii) the operator M lives in the span of $\phi(x_1), \ldots, \phi(x_n)$ for a given set of points $(x_i)_{i=1}^n$, namely there exists $A \in \mathbb{S}^n_+$ such that $\mathsf{M} = \sum_{ij} A_{ij} \phi(x_i) \phi(x_j)^\top$. We define a *Gaussian PSD model* by specializing the definition in (1) as

$$f(x \, ; \, A, X, \eta) = \sum_{i,j=1}^n A_{i,j} k_\eta(x_i, x) k_\eta(x_j, x), \qquad \forall \, x \in \mathbb{R}^d \tag{2}$$

in terms of the coefficient matrix $A \in \mathbb{S}^n_+$, the base points matrix $X \in \mathbb{R}^{n \times d}$, whose i-th row corresponds to the point x_i for each $i = 1, \ldots, n$ and kernel parameter η. In the following, given two base point matrices $X \in \mathbb{R}^{n \times d}$ and $X' \in \mathbb{R}^{m \times d}$, we denote by $K_{X, X', \eta} \in \mathbb{R}^{n \times m}$ the kernel matrix with entries $(K_{X, X', \eta})_{ij} = k_\eta(x_i, x'_j)$ where x_i, x'_j are the i-th and j-th rows of X, X' respectively. When clear from context, in the following we will refer to Gaussian PSD models as PSD models.

Remark 1 (PSD models generalize Mixture models)**.** *Mixture models (a mixture of Gaussian distributions) are a special case of PSD models. Let $A = \mathrm{diag}(a)$ be a diagonal matrix of n positive weights $a \in \mathbb{R}^n_{++}$. We have $f(x \, ; \, A, X, \eta/2) = \sum_{i=1}^n A_{ii} k_{\eta/2}(x_i, x)^2 = \sum_{i=1}^n a_i k_\eta(x_i, x)$.*

Remark 2 (PSD models allow negative weights)**.** *From (2), we immediately see that PSD models generalize mixture models by allowing also for negative weights: e.g., $f(\cdot \, ; \, A, X, \eta)$ with $A = (1, -\frac{1}{2}; -\frac{1}{2}, \frac{1}{4}) \in \mathbb{S}^2_+, \eta = 1, X = (x_1; x_2)$ and $x_1 = 0, x_2 = 2$, corresponding to $f(x \, ; \, A, X, \eta) = e^{-2x^2} + \frac{1}{4} e^{-2(x-2)^2} - \frac{1}{e} e^{-2(x-1)^2}$, i.e. a mixture of Gaussians with also negative weights.*

2.1 Operations with PSD models

In Sec. 3 we will show that PSD models can approximate a wide class of probability densities, significantly outperforming mixture models. Here we show that this improvement does not come at the expenses of computations. In particular, we show that PSD models enjoy the same flexibility of mixture models: i) they are closed with respect to key operations such as marginalization and multiplication and ii) these operations can be performed efficiently in terms of matrix sums/products. The derivation of the results reported in the following is provided in Appendix F. They follow from well-known properties of the Gaussian function.

Evaluation. Evaluating a PSD model in a point $x_0 \in \mathcal{X}$ corresponds to $f(x = x_0 \, ; \, A, X, \eta) = K_{X, x_0, \eta}^\top A K_{X, x_0, \eta}$. Moreover, partially evaluating a PSD in one variable yields

$$f(x, y = y_0 \, ; \, A, [X, Y], (\eta_1, \eta_2)) \, = \, f(x \, ; \, B, X, \eta_1) \quad \text{with} \quad B = A \circ (K_{Y, y_0, \eta_2} K_{Y, y_0, \eta_2}^\top). \tag{3}$$

83   Note that $f(x\,;\,B, X, \eta_1)$ is still a PSD model since B is positive semidefinite.

84   **Sum Rule (Marginalization and Integration).** The integral of a PSD model can be computed as

$$\int f(x\,;\,A, X, \eta)\,dx = c_{2\eta}\,\mathrm{Tr}(A\,K_{X,X,\frac{\eta}{2}}) \qquad \text{with} \qquad c_\eta = \int k_\eta(0, x)\,dx, \qquad (4)$$

85 where $c_\eta = \pi^{d/2}\det(\mathrm{diag}(\eta))^{-1/2}$. This is particularly useful to model probabiliy densities with
86 PSD models. Let $Z = \int f(x\,;\,A, X, \eta)dx$, then the function $f(x\,;\,A/Z, X, \eta) = \frac{1}{Z}f(x\,;\,A, X, \eta)$
87 is a probability density. Integrating only one variable of a PSD model we obtain the sum rule.

88   **Proposition 1** (Sum Rule – Marginalization). *Let $X \in \mathbb{R}^{n \times d}$, $Y \in \mathbb{R}^{n \times d'}$, $A \in \mathbb{S}_+(\mathbb{R}^n)$ and*
89 $\eta \in \mathbb{R}_{++}^d, \eta' \in \mathbb{R}_{++}^{d'}$. *Then, the following integral is a PSD model*

$$\int f(x, y\,;\,A, [X, Y], (\eta, \eta'))\,dx = f(y\,;\,B, Y, \eta'), \qquad \text{with} \qquad B = c_{2\eta}\,A \circ K_{X,X,\frac{\eta}{2}}, \qquad (5)$$

90 The result above shows that we can efficiently marginalize a PSD model with respect to one variable
91 by means of an entry-wise multiplication between two matrices.

92   **Remark 3** (Integration and marginalization on the hypercube). *The integrals in (4) and (5) can be*
93 *performed also on when $\mathcal{X}$ is a hypercube $H = \prod_{t=1}^d [a_t, b_t]$ rather than the entire space $\mathbb{R}^d$. This*
94 *leads to a closed form, where the matrix $K_{X,X,\frac{\eta}{2}}$ is replaced by a suitable $K_{X,X,\frac{\eta}{2},H}$ that can be*
95 *computed with same number of operations (the full form of such matrix is reported in Appendix F).*

96   **Product Rule (Multiplication).** Multiplying two probabilities is key to several applications in
97 probabilistic inference [5]. The family of PSD models is closed with respect to this operation.

98   **Proposition 2** (Multiplication). *Let $X \in \mathbb{R}^{n \times d_1}$, $Y \in \mathbb{R}^{n \times d_2}$, $Y' \in \mathbb{R}^{m \times d_2}$, $Z \in \mathbb{R}^{m \times d_3}$, $A \in \mathbb{S}_+^n$,*
99 $B \in \mathbb{S}_+^m$ *and* $\eta_1 \in \mathbb{R}_{++}^{d_1}$, $\eta_2, \eta_2' \in \mathbb{R}_{++}^{d_2}$, $\eta_3 \in \mathbb{R}_{++}^{d_3}$. *Then*

$$f(x, y\,;\,A, [X, Y], (\eta_1, \eta_2))f(y, z\,;\,B, [Y', Z], (\eta_2', \eta_3)) = f(x, y, z\,;\,C, W, \eta), \qquad (6)$$

100 *is a PSD model, where $C = (A \otimes B) \circ \big(\mathrm{vec}(K_{Y,Y',\tilde{\eta}_2})\mathrm{vec}(K_{Y,Y',\tilde{\eta}_2})^\top\big)$, with $\tilde{\eta}_2 = \frac{\eta_2 \eta_2'}{\eta_2 + \eta_2'}$, base*
101 *matrix $W = [X \otimes \mathbf{1}_m,\ \frac{\eta_2}{\eta_2 + \eta_2'}Y \otimes \mathbf{1}_m + \frac{\eta_2'}{\eta_2 + \eta_2'}\mathbf{1}_n \otimes Y',\ \mathbf{1}_n \otimes Z]$ and $\eta = \big(\eta_1, \eta_2 + \eta_2', \eta_3\big)$.*

102   We note that, despite the heavy notation, multiplying two PSD is performed via simple operations such
103 as tensor and entry-wise product between matrices. In particular, we note that $X \otimes \mathbf{1}_m \in \mathbb{R}^{nm \times d_1}$
104 and $\mathbf{1}_n \otimes Z \in \mathbb{R}^{nm \times d_3}$ correspond respectively to the $nm \times d_1$-matrix containing m copies of each
105 row of X and the $nm \times d_3$-matrix containing n copies of Z. Finally, $\eta Y \otimes \mathbf{1}_m + \eta' \mathbf{1}_n \otimes Y$ is the
106 $nm \times d_2$ base point matrix containing all possible sums $(\eta y_i + \eta' y_j')_{i,j=1}^{n,m}$ of points from Y and Y'.

107   **Reduction.** As observed above, when performing operations with PSD models, the resulting base
108 point matrix might be of the form $X \otimes \mathbf{1}_m$ (e.g. if we couple the multiplication in (6) with
109 marginalization as in a Markov transition, see the corollary below). In these cases we can reduce the
110 PSD model to have only n base points (rather than nm), as follows

$$f(x\,;\,A, X \otimes \mathbf{1}_m, \eta) = f(x\,;\,B, X, \eta) \qquad \text{with} \qquad B = (I_n \otimes \mathbf{1}_m^\top)A(I_n \otimes \mathbf{1}_m), \qquad (7)$$

111 where $A \in \mathbb{S}_+^{nm}$ and $I_n \in \mathbb{R}^{n \times n}$ is the $n \times n$ identity matrix. The reduction operation is useful to
112 avoid the dimensionality of the PSD model grow unnecessarily. This is for instance the case of a
113 Markov transition.

114   **Corollary 3** (Markov Transition). *Let $X \in \mathbb{R}^{n \times d_1}$, $Y \in \mathbb{R}^{n \times d_2}$, $Y' \in \mathbb{R}^{m \times d_2}$, $A \in \mathbb{S}_+^n$, $B \in \mathbb{S}_+^m$*
115 *and $\eta_1 \in \mathbb{R}_{++}^{d_1}$, $\eta_2, \eta_2' \in \mathbb{R}_{++}^{d_2}$. Then*

$$\int f(x, y\,;\,A, [X, Y], (\eta_1, \eta_2))f(y\,;\,B, Y', \eta_2')\,dy = f(x\,;\,C, X, \eta_1), \qquad (8)$$

116 *with $C \in \mathbb{S}_+^n$ obtained by applying in order, Prop. 2, Prop. 1 and reduction (7).*

117   We remark that the result of a Markov transition retains the same base point matrix X and parameters
118 η of the transition kernel. This is thanks to the reduction operation in (7), which avoids the resulting
119 matrix C to be $nm \times nm$. This fact is particularly useful in applications that require multiple Markov
120 transitions, such as in hidden Markov models (see also Sec. 4).

2.2 Compression of a PSD model

From Prop. 2 we note that the multiplication operation can rapidly yield a large number of base points and thus high computational complexity in some settings. Here we propose to adopt a dimensionality reduction strategy to mitigate this issue. In particular, given a set of points $\tilde{x}_1, \ldots, \tilde{x}_m \in \mathbb{R}^d$, we leverage the representation of the PSD model in terms of reproducing kernel Hilbert spaces [3] to use powerful sketching techniques as *Nyström projection* (see, e.g., [39]), to project the PSD model on a new PSD model now based only on the new points. Given $A \in \mathbb{S}_+^n$, $X \in \mathbb{R}^{n \times d}$, $\eta \in \mathbb{R}_{++}^d$. Let $\tilde{X} \in \mathbb{R}^{m \times d}$ be the base point matrix whose j-rows corresponds to the point $\tilde{x}_j$. The *compression* of $f(\cdot\,;A, X, \eta)$ corresponds to

$$f(x\,;\tilde{A}, \tilde{X}, \eta) \qquad \text{with} \qquad \tilde{A} = BAB^\top \in \mathbb{S}_+^m, \quad B = K_{\tilde{X}, \tilde{X}, \eta}^{-1} K_{\tilde{X}, X, \eta} \in \mathbb{R}^{m \times n}, \qquad (9)$$

which it is still a PSD model since the matrix $BAB^\top \in \mathbb{S}_+^m$. We note that it is possible to show that even with a rather simple strategy to choose the new base points $\tilde{x}_1, \ldots, \tilde{x}_m$ – such as uniform sampling – compression is an effective tool to reduce the computational complexity of costly operations. In particular, it can be shown that a compressed model with only $O(t \operatorname{polylog}(1/\varepsilon))$ centers (instead of t^2) can ε-approximate the product of two models with t points. We refer the interested reader to Appendix C for a formal statement and proof of this result.

3 Representation power of PSD models

In this section we deepen our discussion in the introduction to further motivate our choice towards PSD models over the more well-established mixture models. In principle, mixture models offer an appealing strategy to model probability densities since they allow for efficient computations when performing key operations such as sum and product rule, as well as many others often useful in Bayesian inference (the same operations in Sec. 2.1 apply). However, these advantages in terms of computations, come as a disadvantage in terms of flexibility: while mixture models are universal approximators (namely they can approximate arbitrarily well any probability density), they require a significant number of observations and centers to do that. Indeed it is known that models that are non-negative mixtures of non-negative probabilities lead to learning rates that suffer from the curse of dimensionality. For example kernel density estimation (KDE) [21] achieves a rate that is only $n^{-2/(4+d)}$ (see, e.g. [37, page 100]), when using non-negative probabilities, and cannot improve with the regularity (smoothness) of the target probability (see e.g. [9, Thm. 16.1] for impossibility results in the one dimensional case).

In the past decades, this limitation has been overcome by removing either the non-negativity of the weights of the mixture (leading to RBF networks [27]), or the non-negativity of the kernel functions used in the mixture (leading to KDE with oscillating kernels [34]), or both. On the positive side this allows to achieve a learning rate of $n^{-\beta/(2\beta+d)}$, for β-times differentiable densities. Note that such rate is *minimax optimal* [11] and overcomes the curse of dimensionality when $\beta \geq d$. Additionally the resulting model is very *concise*, since only $m = O(n^{d/\beta})$ centers are necessary to achieve the optimal rate [35]. However, on the negative side, *the resulting model is not a probability*, since it may attain negative values, and so it cannot be used where a proper probability is required.

PSD models, offer a way out from this dichotomy, since they allow negative weights (see Remark 2) while still guaranteeing a model that is non-negative everywhere. In particular, in this section we prove that PSD models take the best of both worlds: i) they lead to models that are non-negative everywhere ii) they achieve the learning rate of $O(n^{-\beta/(2\beta+d)})$ with concise models of $m = O(n^{d/\beta})$ centers, for a wide family of distributions.

3.1 Approximation properties of Gaussian PSD models

We start the section recalling that Gaussian PSD models are universal approximators for probability densities. In particular, the following result restates [15, Thm. 2] for the case of probabilities.

Proposition 4 (Universal consistency – Thm. 2 in [15])**.** *The Gaussian PSD family is a universal approximator for probabilities that admit a density.*

The result above is not surprising since Gaussian PSD models generalize classical Gaussian mixtures (see Remark 1), which are known to be universal [5]. We now introduce a mild assumption that will

enable us to complement Prop. 4 with approximation and learning results. In the rest of the section we assume that $\mathcal{X} = (-1, 1)^d$ (or more generally an open bounded subset of $\mathbb{R}^d$ with Lipschitz boundary). Here $L^\infty(\mathcal{X})$ and $L^2(\mathcal{X})$ denote respectively the space of essentially bounded and square-integrable functions over $\mathcal{X}$, while $W_2^\beta(\mathcal{X})$ denotes the *Sobolev space* of functions whose weak derivatives up to order β are square-integrable on $\mathcal{X}$ (see [1] or Appendix A for more details).

Assumption 1. *Let $\beta > 0, q \in \mathbb{N}$. There exists $f_1, \ldots, f_q \in W_2^\beta(\mathcal{X}) \cap L^\infty(\mathcal{X})$, such that the density $p : \mathcal{X} \to \mathbb{R}$ satisfies*

$$p(x) = \sum_{j=1}^q f_j(x)^2, \qquad \forall x \in \mathcal{X}. \tag{10}$$

The assumption above is quite general and satisfied by a wide family of probabilities, as discussed in the following proposition. The proof is reported in Appendix D.1

Proposition 5 (Generality of Assumption 1). *The assumption above is satisfied by*

   (a) any probability density $p \in W_2^\beta(\mathcal{X}) \cap L^\infty(\mathcal{X})$ and strictly positive on $[-1, 1]^d$,

   (b) any exponential model $p(x) = e^{-v(x)}$ with $v \in W_2^\beta(\mathcal{X}) \cap L^\infty(\mathcal{X})$,

   (c) any mixture model of Gaussians or, more generally, of exponential models from (b),

   (d) any p that is $\beta + 2$-times differentiable on $[-1, 1]^d$, with a finite set of zeros, all in $(-1, 1)^d$, and with positive definite Hessian in each zero. E.g. $p(x) \propto x^2 e^{-x^2}$.

Moreover if p is β-times differentiable over $[-1, 1]^d$, then it belongs to $W_2^\beta(\mathcal{X}) \cap L^\infty(\mathcal{X})$.

We note that in principle the class $\mathcal{C}_{\beta,d} = W_2^\beta(\mathcal{X}) \cap L^\infty(\mathcal{X})$ is larger than the Nikolskii class usually considered in density estimation [11] when $\beta \in \mathbb{N}$, but Assumption 1 imposes a restriction on it. However, Prop. 5 shows that the restriction imposed by Assumption 1 is quite mild, since it the resulting space includes all probabilities that are β-times differentiable and strictly positive (it also contains probabilities that are not strictly positive but have some zeros, see [25]). We can now proceed to the main result of this work, which characterizes the approximation capabilities of PSD models.

Theorem 6 (Conciseness of PSD Approximation). *Let p satisfy Assumption 1. Let $\varepsilon > 0$. There exists a Gaussian PSD model of dimension $m \in \mathbb{N}$, i.e., $\hat{p}_m(x) = f(x\,;\,A_m, X_m, \eta_m)$, with $A_m \in \mathbb{S}_+^m$ and $X_m \in \mathbb{R}^{m \times d}$ and $\eta_m \in \mathbb{R}_{++}^d$, such that*

$$\|p - \hat{p}_m\|_{L^2(\mathcal{X})} \le \varepsilon, \quad \text{with} \quad m = O(\varepsilon^{-d/\beta}(\log \tfrac{1}{\varepsilon})^{d/2}). \tag{11}$$

The proof of Thm. 6 is reported in Appendix D.3. According to the result, the number of base points needed for a PSD model to approximate a density up to precision ε depends on its smoothness (the smoother the better) and matches the bound $m = \varepsilon^{-d/\beta}$ that is optimal for function interpolation [18], corresponding to models allowing for negative weights, and is also optimal for convex combinations of oscillating kernels [35].

3.2 Learning a density with PSD models

In this section we study the capabilities of PSD models to estimate a density from n samples. Let $\mathcal{X} = (-1, 1)^d$ and let p be a probability on $\mathcal{X}$. Denote by $x_1, \ldots, x_n$ the samples independently and identically distributed according to p, with $n \in \mathbb{N}$. We consider a Gaussian PSD estimator $\hat{p}_{n,m} = f(x\,;\,\hat{A}, \tilde{X}, \eta)$ that is built on top of m additional points $\tilde{x}_1, \ldots, \tilde{x}_m$, sampled independenly and uniformly at random in $\mathcal{X}$. In particular, $\eta \in \mathbb{R}_{++}^d$, $\tilde{X} \in \mathbb{R}^{m \times d}$ is the base point matrix whose j-th row corresponds to the point $\tilde{x}_j$ and $\hat{A} \in \mathbb{S}_+^m$ is trained as follows

$$\hat{A} = \underset{A \in \mathbb{S}_+^m}{\operatorname{argmin}} \int_{\mathcal{X}} f(x\,;\,A, \tilde{X}, \eta)^2 dx - \frac{2}{n} \sum_{i=1}^n f(x_i\,;\,A, \tilde{X}, \eta) \, + \, \lambda \|K^{1/2} A K^{1/2}\|_F^2, \tag{12}$$

where $K = K_{\tilde{X}, \tilde{X}, \eta}$. Note that the functional is constituted by two parts. The first two elements are an empirical version of $\|f - p\|_{L^2(\mathcal{X})}^2$ modulo a constant independent of f (and so not affecting the

optimization problem), since x_i are identically distributed according to p and so $\frac{1}{n}\sum_{i=1}^{n} f(x_i) \approx \int f(x)p(x)dx$. The last term is a regularizer and corresponds to $\|K^{1/2}AK^{1/2}\|_F^2 = \text{Tr}(AKAK)$, i.e. the Frobenius norm of $\mathsf{M} = \sum_{ij=1}^{m} A_{ij}\phi_\eta(\tilde{x}_i)\phi_\eta(\tilde{x}_j)$. The problem in (12) corresponds to a quadratic problem with a semidefinite constraint and can be solved using techniques such as Newton method [25] or first order dual methods [15]. We are now ready to state our result.

Theorem 7. *Let* $n, m \in \mathbb{N}, \lambda > 0, \eta \in \mathbb{R}_{++}^d$ *and* p *be a density satisfying* Assumption 1. *With the definitions above, let* $\hat{p}_{n,m}$ *be the model* $\hat{p}_{n,m}(x) = f(x\,;\,\hat{A}, \tilde{X}, \eta)$, *with* $\hat{A}$ *the minimizer of* (12). *Let* $\eta = n^{\frac{2}{2\beta+d}}\,\mathbf{1}_d$ *and* $\lambda = n^{-\frac{2\beta+2d}{2\beta+d}}$. *When* $m \geq C'n^{\frac{d}{2\beta+d}}(\log n)^d \log(C''n(\log n))$, *the following holds with probability at least* $1 - \delta$,

$$\|p - \hat{p}_{n,m}\|_{L^2(\mathcal{X})} \leq Cn^{-\frac{\beta}{2\beta+d}}(\log n)^{d/2}, \tag{13}$$

where constant C *depends only on* β, d *and* p *and the constants* C', C'' *depend only on* β, d.

The proof of Thm. 7 is reported in Appendix E.2. The theorem guarantees that under Assumption 1, Gaussian PSD models can achieve the rate $O(n^{-\beta/(2\beta+d)})$ – that is optimal for the β-times differentiable densities – while admitting a concise representation. Indeed, it needs a number $m = O(n^{d/(2\beta+d)})$ of base points, matching the optimal rate in [35]. When $\beta \geq d$, a model with $m = O(n^{1/3})$ centers achieves optimal learning rates.

4 Applications

PSD models are a strong candidate in a variety of probabilistic settings. On the one hand, they are computationally amenable to key operations such as sum and product rules, similarly to mixture models (Sec. 2.1). On the other hand, they are remarkably flexible and can approximate/learn (coincisely) a wide family of target probability densities (Sec. 3). Building on these properties, in this section we consider different possible applications of PSD models in practice.

4.1 PSD Models for Decision Theory

Decision theory problems (see e.g. [5] and references therein) can be formulated as a minimization

$$\theta_* = \underset{\theta \in \Theta}{\text{argmin}}\; L(\theta) = \mathbb{E}_{x \sim p}\,\ell(\theta, x), \tag{14}$$

where ℓ is a loss function, Θ is the space of target parameters (decisions) and p is the underlying data distribution. When we can sample directly from p – e.g. in supervised or unsupervised learning settings – we can apply methods such as stochastic gradient descent to efficently solve (14). However, in many applications, sampling from p is challenging or computationally unfeasible. This is for instance the case when p has been obtained via inference (e.g. it is the t-th estimate in a hidden Markov model, see Sec. 4.3) or it is fully known but has a highly complex form (e.g. the dynamics of a physical system). In contexts where sampling cannot be performed efficiently, it is advisable to consider alternative approaches. Here we propose a strategy to tackle (14) when p can be modeled (or well-approximated) by a PSD model. Our method hinges on the following result.

Proposition 8. *Let* $p(x) = f(x\,;\,A, X, \eta)$ *with* $X \in \mathbb{R}^{n \times d}, A \in \mathbb{S}_+^n, \eta \in \mathbb{R}_{++}^d$. *Let* $g : \mathbb{R}^d \to \mathbb{R}$ *and define* $c_{g,\eta}(z) = \int g(x)e^{-\eta\|x-z\|^2}\,dx$ *for any* $z \in \mathbb{R}^d$. *Then*

$$\mathbb{E}_{x \sim p}\,g(x) \;=\; \text{Tr}\big((A \circ K_{X,X,\eta/2})\,G\big) \qquad \text{with} \qquad G_{ij} = c_{g,2\eta}\big(\tfrac{x_i+x_j}{2}\big). \tag{15}$$

Thanks to Prop. 8 we can readily compute several quantities related to a PSD model such as its mean $\mathbb{E}_p[x] = X^\top b$ with $b = (A \circ K_{X,\eta/2})\mathbf{1}_n$, its covariance or its characteristic function (see Appendix F for the explicit formulas and derivations). However, the result above is particularly useful to tackle the minimization in (14). In particular, since $\nabla L(\theta) = \mathbb{E}_{x \sim p}\nabla_\theta \ell(\theta, x)$, we can use Prop. 8 to directly compute the gradient of the objective function: it is sufficient to know how to evaluate (or approximate) the integral $c_{\nabla\ell_\theta,\eta}(z) = \int \nabla_\theta \ell(\theta, x)e^{-\eta\|x-z\|^2}\,dx$ for any $\theta \in \Theta$ and $z \in \mathbb{R}^d$. Then, we can use first order optimization methods, such as gradient descent, to efficiently solve (14). Remarkably, this approach works well also when we approximate p with a PSD model $\hat{p}$. If ℓ is convex, since $\hat{p}$ is non-negative, the resulting $\hat{L}(\theta) = \mathbb{E}_{x \sim \hat{p}}\,\ell(\theta, x)$ is still a convex functional (see also the discussion on structured prediction in Sec. 4.2). This is not the case if we use more general estimators of p that do not preserve non-negativity.

4.2 PSD Models for Estimating Conditional Probabilities

In supervised learning settings, one is typically interested in solving decision problems of the form $\min_{\theta \in \Theta} \mathbb{E}_{(x,y) \sim p} \ell(h_\theta(x), y)$ where p is a probability over the joint input-output space $\mathcal{X} \times \mathcal{Y}$ and $h_\theta : \mathcal{X} \to \mathcal{Y}$ is a function parameterized by θ. It is well-known [see e.g. 33] that the ideal solution of this problem is the θ_* such that for any $x \in \mathcal{X}$ the function $h_{\theta_*}(x) = \mathrm{argmin}_{z \in \mathcal{Y}} \mathbb{E}_{y \sim p(\cdot|x)} \ell(z, y)$ is the minimizer with respect to $z \in \mathcal{Y}$ of the conditional expectation of $\ell(z, y)$ given x. This leads to target functions that capture specific properties of p, such as moments. For instance, when ℓ is the squared loss, $h_{\theta_*}(x) = \mathbb{E}_{y \sim p(\cdot|x)} y$ corresponds to the conditional expectation of y given x, while for ℓ the absolute value loss, h_{θ_*} recovers the conditional median of p.

In several applications, associating an input x to a single quantity $h_\theta(x)$ in output is not necessarily ideal. For instance, when $p(y|x)$ is multi-modal, estimating the mean or median might not yield useful predictions for the given task. Moreover, estimators of the form h_θ require access to the full input x to return a prediction, and therefore cannot be used when some features are missing (e.g. due to data corruption). In these contexts, an alternative viable strategy is to directly model the conditional probability. When using PSD models, conditional estimation can be performed in two steps, by first modeling the joint distribution $p(y, x) = f(y, x\,;\, A, [Y, X], (\eta, \eta'))$ (e.g. by learning it as suggested Sec. 3) and then use the operations in Sec. 2.1 to condition it with respect to $x_0 \in \mathcal{X}$ as

$$p(y|x_0) = \frac{p(y, x_0)}{p(x_0)} = f(y\,;\, B, Y, \eta) \quad \text{with} \quad B = \frac{A \circ (K_{X, x_0, \eta'} K_{X, x_0, \eta'}^\top)}{c_{2\eta} \mathrm{Tr}\big(A \circ (K_{X, x_0, \eta'} K_{X, x_0, \eta'}^\top) K_{Y, Y, \eta}\big)}. \quad (16)$$

In case of data corruption, it is sufficient to first marginalize $p(y, x)$ on the missing variables and then apply (16). Below we discuss a few applications of the conditional estimator.

Conditional Expectation. Conditional mean embeddings [30] are a well-established tool to efficiently compute the conditional expectation $\mathbb{E}_{y \sim p(\cdot|x_0)} g(y)$ of a function $g : \mathcal{Y} \to \mathbb{R}$. However, although they enjoy good approximation properties [16], they to not guarntee the resulting estimator to take only non-negative values. In contrast, when p is a PSD model (or an approximation), we can apply Prop. 8 to $p(\cdot|x_0)$ in (16) and evaluate the conditional expectation of any g for which we know how to compute (or approximate) the integral $c_{g,\eta}(z) = \int g(y) e^{-\eta\|y-z\|^2} \, dy$. In particular we have $\mathbb{E}_{y \sim p(\cdot|x)} g(y) = \mathrm{Tr}((B \circ K_{Y,Y,\eta/2}) G)$ with B as in (16) and G the matrix with entries $G_{ij} = c_{g,2\eta}(\frac{y_i + y_j}{2})$. Remarkably, differently from conditional mean embeddings estimators, this strategy allows us to compute the conditional expectations also of functions g not in $\mathcal{H}_\eta$.

Structured Prediction. Structured prediction identifies supervised learning problems where the output space $\mathcal{Y}$ has complex structures so that it is challenging to find good parametrizations for $h_\theta : \mathcal{X} \to \mathcal{Y}$ [4, 19]. In [7], a strategy was proposed to tackle these settings by first learning an approximation $\psi_\theta(z, x) \approx \mathbb{E}_{y \sim p(\cdot|x)} \ell(z, y)$ and then model $h_\theta(x) = \mathrm{argmin}_{z \in \mathcal{Y}} \psi_\theta(z, x)$. However, the resulting function $\psi_\theta(\cdot, x)$ is not guaranteed to be convex, even when ℓ is convex. In contrast, by combining the conditional PSD estimator in (16) with the reasoning in Sec. 4.1, we have a strategy that overcomes this issue: when $p(\cdot|x)$ is a PSD model approximating p, we can compute its gradient $\mathbb{E}_{y \sim p(\cdot|x)} \nabla_z \ell(z, y)$ as mentioned in Sec. 4.1 using Prop. 8. Moreover, if $\ell(\cdot, y)$ is convex, the term $\mathbb{E}_{y \sim p(\cdot|x)} \ell(\cdot, y)$ is also convex, and we can use methods such as gradient descent to find $h(x)$ exactly.

Mode Estimation. When the output distribution $p(y|x)$ is multimodal, having access to an explicit form for the conditional density can be useful to estimate its modes. This problem is typically non-convex, yet, when the output y belongs to a small dimensional space (e.g. in classification or scalar-valued regression settings), efficient approximations exist (e.g. bisection).

4.3 Inference on Hidden Markov Models

We consider the problem of performing inference on hidden Markov models (HMM) using PSD models. Let $(x_t)_{t \in \mathbb{N}}$ and $(y_t)_{t \in \mathbb{N}}$ denote two sequences of states and observations respectively. For each $t \geq 1$, we denote by $x_{0:t} = x_0, \ldots, x_t$ and $y_{1:t} = y_1, \ldots, y_t$ and we assume that $p(x_t|x_{0:t-1}, y_{1:t-1}) = p(x_t|x_{t-1}) = \tau(x_t, x_{t-1})$ and $p(y_t|x_{0:t}, y_{1:t-1}) = p(y_t|x_t) = \omega(y_t, x_t)$ with $\tau : \mathcal{X} \times \mathcal{X} \to \mathbb{R}_+$ and $\omega : \mathcal{Y} \times \mathcal{X} \to \mathbb{R}_+$ respectively the transition and observation functions.

Our goal is to infer the distribution of possible states x_t at time t, given all the observations $y_{1:t}$ and a probability $p(x_0)$ on the possible initial states. We focus on this goal for simplicity, but other forms

Algorithm 1 PSD Hidden Markov Model

Input: Transition $\hat{\tau}(x_+, x) = f(x_+, x \,;\, B, [X_+, X], (\eta_+, \eta))$, initial $\hat{p}(x_0) = f(x_0 \,;\, A_0, X_0, \eta_0)$ and observation $\hat{\omega}(y, x) = f(y, x \,;\, C, [Y, X'], (\eta_{obs}, \eta'))$ distributions. $\tilde{X}$ as in Prop. 9. $\tilde{\eta} = \frac{\eta(\eta' + \eta_+)}{\eta + \eta' + \eta_+}$.

$\tilde{X}' = \frac{\eta}{\eta + \eta' + \eta_+} X \otimes \mathbf{1}_{nm} + \frac{\eta' + \eta_+}{\eta + \eta' + \eta_+} \mathbf{1}_n \otimes \tilde{X}, \qquad$ and $\quad \tilde{\eta}' = \frac{\eta' \eta_+}{\eta' + \eta_+}$

For any new observation y_t:

$C_t = C \circ (K_{Y, y_t, \eta_{obs}}^\top K_{Y, y_t, \eta_{obs}})$ // Partial evaluation $\hat{\omega}_t(x_+) = \hat{\omega}(y = y_t, x_+)$

$B_t = (B \otimes A_{t-1}) \circ \big(\mathrm{vec}(K_{X, \tilde{X}, \tilde{\eta}}) \mathrm{vec}(K_{X, \tilde{X}, \tilde{\eta}})^\top\big)$ // Product $\hat{\beta}_t(x_+, x) = \hat{\tau}(x_+, x)\hat{p}(x|y_{1:t-1})$

$D_t = (I_n \otimes \mathbf{1}_{nm}^\top)(B_t \circ K_{\tilde{X}', \tilde{X}', \frac{\tilde{\eta}}{2}})(I_n \otimes \mathbf{1}_{nm})$ // Marginalization $\hat{\beta}_t(x_+) = \int \beta_t(x_+, x)\, dx$

$E_t = (C_t \otimes D_t) \circ \big(\mathrm{vec}(K_{X_+, X', \frac{\tilde{\eta}'}{2}}) \mathrm{vec}(K_{X_+, X', \frac{\tilde{\eta}'}{2}})^\top\big)$ // Product $\hat{\pi}_t(x_+) = \hat{\omega}_t(x_+)\hat{\beta}_t(x_+)$

$A_t = E_t / c_t$, with $c_t = c_{2(\eta' + \eta_+)} \mathrm{Tr}(E_t K_{\tilde{X}, \tilde{X}, \frac{\eta' + \eta_+}{2}})$ // Normalization $\hat{p}(x_+|y_{1:t}) = \frac{\hat{\pi}_t(x_+)}{\int \hat{\pi}(x_+)\, dx}$

Return $f(x_t \,;\, A_t, \tilde{X}, \eta' + \eta_+)$

---

of inferences are possible (e.g. estimating x_{m+t}, namely m steps into the future or the past). We assume that the transition and observation functions can be well approximated by PSD models $\hat{\tau}$ and $\hat{\omega}$ (e.g. by learning them or known a-priori for the problem's dynamics). Then, given a PSD model estimate $\hat{p}(x_0)$ of the initial state $p(x_0)$, we can recursively define the sequence of estimates

$$\hat{p}(x_t|y_{t:1}) = \frac{\hat{\tau}(y_t, x_t)\int \omega(x_t, x_{t-1})\hat{p}(x_t|y_{1:t-1})\, dx_{t-1}}{\int \hat{\tau}(y_t, x_t)\omega(x_t, x_{t-1})\hat{p}(x_t|y_{1:t-1})\, dx_t dx_{t-1}}. \tag{17}$$

Note that when $\hat{\tau}, \hat{\omega}, \hat{p}(x_0)$ correspond to the real transition, observation and initial state probability, the formula above yields the exact distribution $p(x_t|y_{1:t})$ over the states at time t (this follows directly by subsequent applications of Bayes' rule. See also e.g. [5]). If $\hat{\tau}, \hat{\omega}, \hat{p}(x_0)$ are PSD models, then each of the $\hat{p}(x_t|y_{t:1})$ is a PSD model recursively defined only in terms of the previous estimate and the operations introduced in Sec. 2.1. In particular, we have the following result.

Proposition 9 (PSD Hidden Markov Models (HMM)). *Let* $X_0 \in \mathbb{R}^{n_0 \times d}$, $X_+, X \in \mathbb{R}^{n \times d}$, $X' \in \mathbb{R}^{m \times d}$, $Y \in \mathbb{R}^{m \times d'}$, $A_0 \in \mathbb{S}_+^{n_0}$, $A \in \mathbb{S}_+^n$, $B \in \mathbb{S}_+^m$ *and* $\eta_0, \eta, \eta', \eta_+ \in \mathbb{R}_{++}^d$, $\eta_{obs} \in \mathbb{R}_{++}^{d'}$. *Let*

$$\hat{\tau}(x_+, x) = f(x_+, x \,;\, B, [X_+, X], (\eta_+, \eta)), \qquad \hat{\omega}(y, x) = f(y, x \,;\, C, [Y, X'], (\eta_{obs}, \eta')), \quad (18)$$

be approximate transition and observation functions. Then, given the initial state probability $\hat{p}(x_0) = f(x_0 \,;\, A_0, X_0, \eta_0)$, *for any* $t \geq 1$, *the estimate* $\hat{p}$ *in* (17) *is a PSD model of the form*

$$\hat{p}(x_t|y_{t:1}) = f(x_t \,;\, A_t, \tilde{X}, \eta' + \eta_+), \tag{19}$$

where $\tilde{X} = \frac{\eta'}{\eta' + \eta_+} X' \otimes \mathbf{1}_n + \frac{\eta_+}{\eta' + \eta_+} \mathbf{1}_m \otimes X_+$ *and* A_t *is recursively obtained from* A_{t-1} *as in Alg. 1.*

Remark 4 (Sum-product Algorithm). *Eq.* (17) *is an instance of the so-called sum-product algorithm, a standard inference method for graphical models [5] (of which HMMs are a special case). The application of the sum-product algorithm relies mainly on sum and product rules for probabilities (as is the case for HMMs in* (17)*). Hence, according to Sec. 2.1, it is highly compatible with PSD models.*

5 Numerical Simulations

We discuss two simulation reporting on the behavior of PSD models in some of the setting described in Sec. 4. For brevity, we report all details in Appendix G.

Inference. We report here a qualitative experiment showing the effect of approximation in inference settings. Our goal is to show the effect of the Markov transition $p_1(y) = \int p(y|x)p_0(x)\, dx$ (see Cor. 3) for different estimators used to approximate the conditional probability $p(y|x)$. In particular, we consider $p(y|x) = \rho\, e^{-\eta_1 \|y - \mu_1 - x\|^2} + (1 - \rho)\, e^{-\eta_2 \|y - \mu_2 - x\|^2}$, with $\mu_1 = 0.0, \mu_2 = 0.5$, $\eta_1 = 7, \eta_2 = 3.5$ and $\rho = 0.3$ (essentially "splitting" the input signal in two components) and $p_0(x) = \frac{1}{2}e^{-\eta_1 \|x - \mu_1\|^2} + \frac{1}{2}e^{-\eta_2 \|x - \mu_2\|^2}$ (see Fig. 1 left). We estimate the conditional density using $n = 2000$ samples with: the strategy in (12) (PSD model), a KDE mixture model using gaussian kernels [21], a conditional mean embedding (CME) estimator [30] using the kernel bayes rule [10].

Figure 1: Inference step from a prior (Left) using different methods to estimate the conditional density.

Figure 2: (Left) Loss $L(\theta) = \mathbb{E}_{x \sim p} \ell(\theta, x)$ and (Right) distance from the true minimizer for increasing number of samples with PSD or KDE to estimate p (see Sec. 4.1), or SGD. Average and standard deviation over 5 runs.

Since CME does not allow to directly recover the target density, we report $g(y) = \mathbb{E}_{y' \sim \hat{p}_1} e^{-\eta(y'-y)^2}$ for η large, which can be considered an approximation of the $\hat{p}_1$ estimated by CME. Since the setting allows exact inference, we report also the ground truth $p_1(y)$. Fig. 1 offers a qualitative comparison between these methods. While all methods are able to capture the multi-modality of p_1, we note that KDE is overly smooth, an effect of its higher sample complexity (as discussed in Sec. 3). CME shows the issue of using estimators that do not guarantee non-negativity: it can take negative values and moreover the effect of positive and negative weights in the linear kernel combination yields modes that are more pronounced than the ground truth. In contrast, the PSD model recovers the structure of the ground truth better than the two competitors. In Appendix G we provide additional details on this setting in particular related to the effect of subsequent steps of inference on the three distributions.

Decision Theory. We perform experiments related to the discussion in Sec. 4.1. We minimize $L(\theta) = \mathbb{E}_{x \sim p} \ell(\theta, x)$ for $\ell(\theta, x) = (x - \theta)^4$ and $p = \rho\, e^{-\eta_1 \|x - \mu_1\|^2} + (1 - \rho)\, e^{-\eta_2 \|x - \mu_2\|^2}$, with $\mu_1 = -0.5, \mu_2 = 0.5, \eta_1 = 7, \eta_2 = 2.5$ and $\rho = 0.3$. We follow the strategy in Sec. 4.1: we approximate $\hat{p}$ using either a PSD model minimizing (12) or Kernel Density Estimation (KDE) with gaussian kernel and then relying on Prop. 8 to estimate the gradients of the approximate loss $\hat{L}(\theta) = \mathbb{E}_{x \sim \hat{p}} \ell(\theta, x)$ to perform gradient descent. Fig. 2 reports the comparison between these two methods and a plugin stochastic gradient descent (SGD) strategy as the number of training samples increases. Since we know p explicitly, we can also compute the minimizer θ_* of $L(\theta)$ in closed form and therefore report the distance of the estimated θ from it. We see that while the PSD model achieves good performance with very few samples, both KDE and SGD have worse sample complexity. This is in line with the approximation results reported in Sec. 3.

6   Discussion

In this work we have shown that PSD models are a strong candidate in practical application related to probabilistic inference. They satisfy both requirements for an ideal probabilistic model: i) they perform exact sum and product rule in terms of efficient matrix operations; ii) we proved that they can concisely approximate a wide range of probabilities.

Limitations and Future Directions. We identify three main limitations of the current work: i) when performing inference on large graphical models (see Remark 4) the multiplication of PSD models might lead to an inflation in the number of base points. Building on our compression strategy, we plan to further investigate low-rank approximations to mitigate this issue. ii) The current form of PSD models does not allow for direct sampling. This is an interesting question left open by our analysis that we plan to investigate in the future. iii) Due to the theoretical focus of the paper, the empirical investigation has been limited. We plan to further characterize the empirical behavior of PSD models in future work. Related to this, we plan to develop a library for operations with PSD models and make it available to the community.

References

[1] Robert A Adams and John JF Fournier. *Sobolev spaces*. Elsevier, 2003.

[2] Jason Altschuler, Francis Bach, Alessandro Rudi, and Jonathan Niles-Weed. Massively scalable sinkhorn distances via the nyström method. *arXiv preprint arXiv:1812.05189*, 2018.

[3] Nachman Aronszajn. Theory of reproducing kernels. *Transactions of the American Mathematical Society*, 68(3):337–404, 1950.

[4] G. H. Bakir, T. Hofmann, B. Schölkopf, A. J. Smola, B. Taskar, and S. V. N. Vishwanathan. *Predicting Structured Data*. MIT Press, 2007.

[5] Christopher M Bishop. Machine learning and pattern recognition. *Information Science and Statistics. Springer, Heidelberg*, 2006.

[6] René Erlín Castillo and Humberto Rafeiro. *An introductory course in Lebesgue spaces*. Springer, 2016.

[7] Carlo Ciliberto, Lorenzo Rosasco, and Alessandro Rudi. A general framework for consistent structured prediction with implicit loss embeddings. *Journal of Machine Learning Research*, 21(98):1–67, 2020.

[8] Felipe Cucker and Ding Xuan Zhou. *Learning theory: an approximation theory viewpoint*, volume 24. Cambridge University Press, 2007.

[9] Luc Devroye and Gábor Lugosi. *Combinatorial methods in density estimation*. Springer Science & Business Media, 2012.

[10] Kenji Fukumizu, Le Song, and Arthur Gretton. Kernel bayes' rule: Bayesian inference with positive definite kernels. *The Journal of Machine Learning Research*, 14(1):3753–3783, 2013.

[11] Alexander Goldenshluger and Oleg Lepski. On adaptive minimax density estimation on r^d. *Probability Theory and Related Fields*, 159(3-4):479–543, 2014.

[12] Ian Goodfellow, Jean Pouget-Abadie, Mehdi Mirza, Bing Xu, David Warde-Farley, Sherjil Ozair, Aaron Courville, and Yoshua Bengio. Generative adversarial nets. In *Advances in neural information processing systems*, pages 2672–2680, 2014.

[13] Lars Hörmander. *The analysis of linear partial differential operators II: Differential operators with constant coefficients*. Springer, 1990.

[14] Diederik P Kingma and Max Welling. Auto-encoding variational bayes. *International Conference on Learning Representation (ICLR*, 2014.

[15] Ulysse Marteau-Ferey, Francis Bach, and Alessandro Rudi. Non-parametric models for non-negative functions. 2020.

[16] Krikamol Muandet, Kenji Fukumizu, Bharath Sriperumbudur, and Bernhard Schölkopf. Kernel mean embedding of distributions: A review and beyond. *Foundations and Trends(r) in Machine Learning Series*, 2017.

[17] Kevin P Murphy. *Machine learning: a probabilistic perspective*. MIT press, 2012.

[18] Erich Novak. Deterministic and stochastic error bounds in numerical analysis. 2006.

[19] Sebastian Nowozin, Christoph H Lampert, et al. Structured learning and prediction in computer vision. *Foundations and Trends in Computer Graphics and Vision*, 2011.

[20] Nicolò Pagliana, Alessandro Rudi, Ernesto De Vito, and Lorenzo Rosasco. Interpolation and learning with scale dependent kernels. *arXiv preprint arXiv:2006.09984*, 2020.

[21] Emanuel Parzen. On estimation of a probability density function and mode. *The annals of mathematical statistics*, 33(3):1065–1076, 1962.

[22] Christian Rieger and Barbara Zwicknagl. Sampling inequalities for infinitely smooth functions, with applications to interpolation and machine learning. *Advances in Computational Mathematics*, 32(1):103, 2010.

[23] Luke Gervase Rogers. *A degree-independent Sobolev extension operator*. Yale University, 2004.

[24] A. Rudi, R. Camoriano, and L. Rosasco. Generalization Properties of Learning with Random Features. *ArXiv e-prints*, February 2016.

[25] Alessandro Rudi, Ulysse Marteau-Ferey, and Francis Bach. Finding global minima via kernel approximations. *arXiv preprint arXiv:2012.11978*, 2020.

[26] Bernhard Schölkopf and Alexander J Smola. *Learning with kernels: support vector machines, regularization, optimization, and beyond*. MIT press, 2002.

[27] John Shawe-Taylor and N Cristianini. *An introduction to support vector machines and other kernel-based learning methods*, volume 204. Volume, 2000.

[28] Winfried Sickel. Composition operators acting on sobolev spaces of fractional order—a survey on sufficient and necessary conditions. *Function spaces, differential operators and nonlinear analysis (Paseky nad Jizerou, 1995)*, pages 159–182, 1996.

[29] Shashank Singh, Ananya Uppal, Boyue Li, Chun-Liang Li, Manzil Zaheer, and Barnabás Póczos. Nonparametric density estimation with adversarial losses. In *Proceedings of the 32nd International Conference on Neural Information Processing Systems*, pages 10246–10257. Curran Associates Inc., 2018.

[30] Le Song, Kenji Fukumizu, and Arthur Gretton. Kernel embeddings of conditional distributions: A unified kernel framework for nonparametric inference in graphical models. *IEEE Signal Processing Magazine*, 30(4):98–111, 2013.

[31] Bharath Sriperumbudur, Kenji Fukumizu, Arthur Gretton, Aapo Hyvärinen, and Revant Kumar. Density estimation in infinite dimensional exponential families. *Journal of Machine Learning Research*, 18, 2017.

[32] Elias M Stein and Guido Weiss. *Introduction to Fourier Analysis on Euclidean Spaces (PMS-32), Volume 32*. Princeton university press, 1971.

[33] Ingo Steinwart and Andreas Christmann. *Support Vector Machines*. Information Science and Statistics. Springer New York, 2008.

[34] Alexandre B Tsybakov. *Introduction to nonparametric estimation*. Springer Science & Business Media, 2008.

[35] Paxton Turner, Jingbo Liu, and Philippe Rigollet. A statistical perspective on coreset density estimation. In *International Conference on Artificial Intelligence and Statistics*, pages 2512–2520. PMLR, 2021.

[36] Adrien Vacher, Boris Muzellec, Alessandro Rudi, Francis Bach, and Francois-Xavier Vialard. A dimension-free computational upper-bound for smooth optimal transport estimation. *arXiv preprint arXiv:2101.05380*, 2021.

[37] Matt P Wand and M Chris Jones. *Kernel smoothing*. CRC press, 1994.

[38] Holger Wendland. *Scattered Data Approximation*, volume 17. Cambridge University Press, 2004.

[39] Christopher Williams and Matthias Seeger. Using the nyström method to speed up kernel machines. In *Proceedings of the 14th annual conference on neural information processing systems*, number CONF, pages 682–688, 2001.

[40] Vadim Yurinsky. *Sums and Gaussian vectors*. Springer, 1995.