Polylogarithmic width suffices for gradient descent to achieve arbitrarily small test error with shallow ReLU networks

Abstract

Recent work has revealed that overparameterized networks trained by gradient descent achieve arbitrarily low training error, and sometimes even low test error. The required width, however, is always polynomial in at least one of the sample size n, the (inverse) target error $1/\epsilon$, and the (inverse) failure probability $1/\delta$. This work shows that $\widetilde{O}(1/\epsilon)$ iterations of gradient descent on two-layer networks of any width exceeding $\mathrm{polylog}(n, 1/\epsilon, 1/\delta)$ and $\widetilde{\Omega}(1/\epsilon^2)$ training examples suffices to achieve a test error of ϵ. The analysis further relies upon a margin property of the limiting kernel, which is guaranteed positive, and can distinguish between true labels and random labels.

1 Introduction

Despite the extensive empirical success of deep networks, their optimization and generalization properties are still not well understood. Recently, the neural tangent kernel (NTK) has provided some insight into the problem. In the infinite-width limit, the NTK converges to a limiting kernel which stays constant during training; on the other hand, when the width is large enough, the function learned by gradient descent follows the NTK (Jacot et al., 2018). This motivates the study of overparameterized networks trained by gradient descent, using properties of the NTK. In fact, parameters related to NTK, such as the minimum eigenvalue of the limiting kernel, appear to affect optimization and generalization (Arora et al., 2019).

However, in addition to such NTK-dependent parameters, prior work also requires the width to depend polynomially on n, $1/\delta$ or $1/\epsilon$, where n denotes the size of the training set, δ denotes the failure probability, and ϵ denotes the target error. These large widths far exceed what is used empirically, constituting a significant gap between theory and practice.

Our contributions. In this paper, we narrow this gap by showing that a two-layer ReLU network with $\Omega(\ln(n/\delta) + \ln(1/\epsilon)^2)$ hidden units trained by gradient descent achieves classification error ϵ *on test data*, meaning both optimization and generalization occur. Unlike prior work, the width is fully polylogarithmic in n, $1/\delta$, and $1/\epsilon$; the width will additionally depend on the *separation margin* of the limiting kernel, a quantity which is guaranteed positive (assuming no inputs are duplicated with noisy labels), and can distinguish between true labels and random labels. The paper organization together with some details are described below.

Section 2 studies gradient descent on the training set. Using the ℓ_1 geometry inherent in classification tasks, we prove that with any width at least polylogarithmic and any constant step size no larger than 1, gradient descent achieves training error ϵ in $\widetilde{O}(1/\epsilon)$ iterations (cf. Theorem 2.2). As is common in the NTK literature (Chizat & Bach, 2019), we also show the parameters hardly change, which will be essential to our generalization analysis.

Section 3 gives a test error bound. Concretely, using the preceding gradient descent analysis, and standard Rademacher tools and exploiting how little the weights moved, we show that with $\widetilde{\Omega}(1/\epsilon^2)$ samples and $\widetilde{O}(1/\epsilon)$ iterations, gradient descent finds a solution with ϵ test error (cf. Theorem 3.2 and Corollary 3.3). (As discussed in Remark 3.4, $\widetilde{\Omega}(1/\epsilon)$ samples

also suffice via a smoothness-based generalization bound, at the expense of large constant factors.)

Section 4 considers stochastic gradient descent (SGD) with access to a standard stochastic online oracle. We prove that with width at least polylogarithmic and sample complexity $\widetilde{O}(1/\epsilon)$, SGD achieves an arbitrarily small test error (cf. Theorem 4.1).

Section 5 discusses the separability condition, which is in general a positive number, but reflects the difficulty of the classification problem. Regarding random labels, we show that starting from a distribution with a good positive margin, but replacing the labels with random noise, the margin can degrade all the way down to $O(1/\sqrt{n})$, which (correctly) removes the possibility of generalization. In this way, our analysis can distinguish between true labels and random labels.

Section 6 concludes with some open problems.

1.1 RELATED WORK

There has been a large literature studying gradient descent on overparameterized networks via the NTK. The most closely related work is (Nitanda & Suzuki, 2019), which shows that a two-layer network trained by gradient descent with the logistic loss can achieve a small test error, under the same assumption that the neural tangent model with respect to the first layer can separate the data distribution. However, they analyze smooth activations, while we handle the ReLU. They require $\Omega(1/\epsilon^2)$ hidden units, $\widetilde{\Omega}(1/\epsilon^4)$ data samples, and $O(1/\epsilon^2)$ steps, while our result only needs polylogarithmic hidden units, $\widetilde{\Omega}(1/\epsilon^2)$ data samples, and $\widetilde{O}(1/\epsilon)$ steps.

Additionally on shallow networks, Du et al. (2018b) prove that on an overparameterized two-layer network, gradient descent can globally minimize the empirical risk with the squared loss. Their result requires $\Omega(n^6/\delta^3)$ hidden units. Oymak & Soltanolkotabi (2019); Song & Yang (2019) further reduce the required overparameterization, but it still has a $\mathrm{poly}(n)$ dependency. Using the same amount of overparameterization as (Du et al., 2018b), Arora et al. (2019) further show that the two-layer network learned by gradient descent can achieve a small test error, assuming that on the data distribution the smallest eigenvalue of the limiting kernel is at least some positive constant. They also give a fine-grained characterization of the predictions made by gradient descent iterates; such a characterization makes use of a special property of the squared loss and cannot be applied to the logistic regression setting. Li & Liang (2018) show that stochastic gradient descent (SGD) with the cross entropy loss can learn a two-layer network with small test error, using $\mathrm{poly}(\ell, 1/\epsilon)$ hidden units, where ℓ is at least the covering number of the support of the feature distribution using balls whose radii are no larger than the smallest distance between two data points with different labels. Allen-Zhu et al. (2018a) consider SGD on a two-layer network, and a variant of SGD on a three-layer network. The three-layer analysis further exhibits some properties not captured by the NTK. They assume a ground truth network with infinite-order smooth activations, and they require the width to depend polynomially on $1/\epsilon$ and some constants related to the smoothness of the activations of the ground truth network.

On deep networks, a variety of works have established low training error Allen-Zhu et al. (2018b); Du et al. (2018a); Zou et al. (2018); Zou & Gu (2019). Allen-Zhu et al. (2018c) show that SGD can minimize the regression loss for recurrent neural networks, and Allen-Zhu & Li (2019b) further prove a low generalization error. Allen-Zhu & Li (2019a) show that using the same number of training examples, a three-layer ResNet can learn a function class with a much lower test error than any kernel method. Cao & Gu (2019a) assume that the neural tangent model with respect to the second layer of a two-layer network can separate the data distribution, and prove that gradient descent on a deep network can achieve ϵ test error with $\Omega(1/\epsilon^4)$ samples and $\Omega(1/\epsilon^{14})$ hidden units. Cao & Gu (2019b) consider SGD with an online oracle and give a general result. Under the same assumption as in (Cao & Gu, 2019a), their result requires $\Omega(1/\epsilon^{14})$ hidden units and sample complexity $\widetilde{O}(1/\epsilon^2)$. By contrast, with the same online oracle, our result only needs polylogarithmic hidden units and sample complexity $\widetilde{O}(1/\epsilon)$.

1.2 NOTATION

The dataset is denoted by $\{(x_i, y_i)\}_{i=1}^n$ where $x_i \in \mathbb{R}^d$ and $y_i \in \{-1, +1\}$. For simplicity, we assume that $\|x_i\|_2 = 1$ for any $1 \le i \le n$, which is standard in the NTK literature.

The two-layer network has weight matrices $W \in \mathbb{R}^{m \times d}$ and $a \in \mathbb{R}^m$. We use the following parameterization, which is also used in (Du et al., 2018b; Arora et al., 2019).

$$f(x; W, a) := \frac{1}{\sqrt{m}} \sum_{s=1}^m a_s \sigma\left(\langle w_s, x\rangle\right),$$

with initialization

$$w_{s,0} \sim \mathcal{N}(0, I_d), \quad \text{and} \quad a_s \sim \text{unif}\left(\{-1, +1\}\right).$$

Note that in this paper, $w_{s,t}$ denotes the s-th row of W at step t. We fix a and only train W, as in (Li & Liang, 2018; Du et al., 2018b; Arora et al., 2019; Nitanda & Suzuki, 2019). We consider the ReLU activation $\sigma(z) := \max\{0, z\}$, though our analysis can be extended easily to Lipschitz continuous, positively homogeneous activations such as leaky ReLU.

We use the logistic (binary cross entropy) loss $\ell(z) := \ln\left(1 + \exp(-z)\right)$ and gradient descent. For any $1 \le i \le n$ and any W, let $f_i(W) := f(x_i; W, a)$. The empirical risk and its gradient are given by

$$\widehat{\mathcal{R}}(W) := \frac{1}{n} \sum_{i=1}^n \ell\left(y_i f_i(W)\right), \quad \text{and} \quad \nabla\widehat{\mathcal{R}}(W) = \frac{1}{n} \sum_{i=1}^n \ell'\left(y_i f_i(W)\right) y_i \nabla f_i(W).$$

For any $t \ge 0$, the gradient descent step is given by $W_{t+1} := W_t - \eta_t \nabla\widehat{\mathcal{R}}(W_t)$. Also define

$$f_i^{(t)}(W) := \left\langle \nabla f_i(W_t), W\right\rangle, \quad \text{and} \quad \widehat{\mathcal{R}}^{(t)}(W) := \frac{1}{n} \sum_{i=1}^n \ell\left(y_i f_i^{(t)}(W)\right).$$

Note that $f_i^{(t)}(W_t) = f_i(W_t)$. This property generally holds due to homogeneity: for any W and any $1 \le s \le m$,

$$\frac{\partial f_i}{\partial w_s} = \frac{1}{\sqrt{m}} a_s \mathbb{1}\left[\langle w_s, x_i\rangle > 0\right] x_i, \quad \text{and} \quad \left\langle \frac{\partial f_i}{\partial w_s}, w_s\right\rangle = \frac{1}{\sqrt{m}} a_s \sigma\left(\langle w_s, x_i\rangle\right),$$

and thus $\left\langle \nabla f_i(W), W\right\rangle = f_i(W)$.

2 EMPIRICAL RISK MINIMIZATION

In this section, we consider a fixed training set and empirical risk minimization. We first state our assumption on the separability of the neural tangent model, and then give our main result and a proof sketch.

Here is some additional notation. Let $\mu_{\mathcal{N}}$ denote the Gaussian measure on $\mathbb{R}^d$, given by the Gaussian density with respect to the Lebesgue measure on $\mathbb{R}^d$. We consider the following Hilbert space

$$\mathcal{H} := \left\{ w : \mathbb{R}^d \to \mathbb{R}^d \,\middle|\, \int \|w(z)\|_2^2 \, \mathrm{d}\mu_{\mathcal{N}}(z) < \infty \right\}.$$

For any $x \in \mathbb{R}^d$, define $\phi_x \in \mathcal{H}$ by

$$\phi_x(z) := x \mathbb{1}\left[\langle z, x\rangle > 0\right],$$

and particularly define $\phi_i := \phi_{x_i}$ for the training input x_i.

Assumption 2.1. There exists $\bar{v} \in \mathcal{H}$ and $\gamma > 0$, s.t. $\left\|\bar{v}(z)\right\|_2 \le 1$ for any $z \in \mathbb{R}^d$, and for any $1 \le i \le n$,

$$y_i \langle \bar{v}, \phi_i\rangle_{\mathcal{H}} := y_i \int \langle \bar{v}(z), \phi_i(z)\rangle \, \mathrm{d}\mu_{\mathcal{N}}(z) \ge \gamma.$$

$\diamond$

As shown in Section 5, the space $\mathcal{H}$ is the reproducing kernel Hilbert space (RKHS) induced by the infinite-width NTK with respect to W, and ϕ_x maps x into $\mathcal{H}$. Assumption 2.1 supposes that the induced training set $\{(\phi_i, y_i)\}_{i=1}^n$ can be separated by some $\bar{v}$ with bounded $\left\| \bar{v}(z) \right\|_2$ for any z. Such a condition on $\left\| \bar{v}(z) \right\|_2$ is important in our analysis. A more natural assumption is that the infinite-width NTK itself can separate the training set with a positive margin, but it actually implies Assumption 2.1 (cf. Proposition 5.1). More discussion on separability is also given in Section 5.

With Assumption 2.1, we state our main empirical risk result.

Theorem 2.2. *Under Assumption 2.1, given any risk target $\epsilon \in (0,1)$ and any $\delta \in (0, 1/3)$, let*

$$\lambda := \frac{\sqrt{2\ln(4n/\delta)} + \ln(4/\epsilon)}{\gamma/4}, \quad and \quad M := \frac{4096\lambda^2}{\gamma^6}.$$

Then for any $m \geq M$ and any constant step size $\eta \leq 1$, with probability $1 - 3\delta$ over the random initialization,

$$\frac{1}{T} \sum_{t < T} \widehat{\mathcal{R}}(W_t) \leq \epsilon, \quad where \quad T := \lceil 2\lambda^2/\eta\epsilon \rceil.$$

Moreover for any $0 \leq t < T$ and any $1 \leq s \leq m$,

$$\left\| w_{s,t} - w_{s,0} \right\|_2 \leq \frac{4\lambda}{\gamma\sqrt{m}}.$$

While the number of hidden units required by prior work all have a polynomial dependency on n, $1/\delta$ or $1/\epsilon$, Theorem 2.2 only requires $m = \Omega\left(\ln(n/\delta) + \ln(1/\epsilon)^2\right)$. In the rest of Section 2, we give a proof sketch of Theorem 2.2.

2.1 PROPERTIES AT INITIALIZATION

In this subsection, we give some nice properties of random initialization. The proofs are given in Appendix A.

Given an initialization (W_0, a), for any $1 \leq s \leq m$, define

$$\bar{u}_s := \frac{1}{\sqrt{m}} a_s \bar{v}(w_{s,0}), \tag{2.1}$$

where $\bar{v}$ is given by Assumption 2.1. Collect $\bar{u}_s$ into a matrix $\overline{U} \in \mathbb{R}^{m \times d}$. It holds that $\|\bar{u}_s\|_2 \leq 1/\sqrt{m}$, and $\left\| \overline{U} \right\|_F \leq 1$. Lemma 2.3 ensures that with high probability $\overline{U}$ has a positive margin at initialization.

Lemma 2.3. *Under Assumption 2.1, given any $\delta \in (0,1)$ and any $\epsilon_1 \in (0, \gamma)$, if $m \geq \left(2\ln(n/\delta)\right)/\epsilon_1^2$, then with probability $1 - \delta$, it holds simultaneously for all $1 \leq i \leq n$ that*

$$y_i f_i^{(0)}\left(\overline{U}\right) = y_i \left\langle \nabla f_i(W_0), \overline{U} \right\rangle \geq \gamma - \sqrt{\frac{2\ln(n/\delta)}{m}} \geq \gamma - \epsilon_1.$$

For any W, any $\epsilon_2 > 0$, and any $1 \leq i \leq n$, define

$$\alpha_i(W, \epsilon_2) = \frac{1}{m} \sum_{s=1}^m \mathbb{1}\left[\left|\langle w_s, x_i \rangle\right| \leq \epsilon_2\right].$$

Lemma 2.4 controls $\alpha_i(W_0, \epsilon_2)$. It will help us show that $\overline{U}$ has a good margin during the training process.

Lemma 2.4. *Under the condition of Lemma 2.3, for any $\epsilon_2 > 0$, with probability $1 - \delta$, it holds simultaneously for all $1 \leq i \leq n$ that*

$$\alpha_i\left(W_0, \epsilon_2\right) \leq \sqrt{\frac{2}{\pi}}\epsilon_2 + \sqrt{\frac{\ln(n/\delta)}{2m}} \leq \epsilon_2 + \frac{\epsilon_1}{2}.$$

Finally, Lemma 2.5 controls the output of the network at initialization.

Lemma 2.5. *Given any $\delta \in (0,1)$, if $m \geq 25\ln(2n/\delta)$, then with probability $1 - \delta$, it holds simultaneously for all $1 \leq i \leq n$ that*

$$\left|f(x_i; W_0, a)\right| \leq \sqrt{2\ln\left(4n/\delta\right)}.$$

2.2 CONVERGENCE ANALYSIS OF GRADIENT DESCENT

We analyze gradient descent in this subsection. First, define

$$\widehat{\mathcal{Q}}(W) := \frac{1}{n} \sum_{i=1}^{n} -\ell'\left(y_i f_i(W)\right).$$

We have the following observations.

- For any W and any $1 \leq s \leq m$, $\left\|\partial f_i / \partial w_s\right\|_2 \leq 1/\sqrt{m}$, and thus $\left\|\nabla f_i(W)\right\|_F \leq 1$. Therefore by the triangle inequality, $\left\|\nabla \widehat{\mathcal{R}}(W)\right\|_F \leq \widehat{\mathcal{Q}}(W)$.

- The logistic loss satisfies $0 \leq -\ell' \leq 1$, and thus $0 \leq \widehat{\mathcal{Q}}(W) \leq 1$.

- The logistic loss satisfies $-\ell' \leq \ell$, and thus $\widehat{\mathcal{Q}}(W) \leq \widehat{\mathcal{R}}(W)$.

In prior work (Cao & Gu, 2019a; Nitanda & Suzuki, 2019), the quantity $\widehat{\mathcal{Q}}$ is also analyzed. In our work, $\widehat{\mathcal{Q}}$ specifically helps us prove the following result, which plays an important role in obtaining a width which only depends on $\operatorname{polylog}(1/\epsilon)$.

Lemma 2.6. *For any $t \geq 0$ and any $\overline{W}$, if $\eta_t \leq 1$, then*

$$\eta_t \widehat{\mathcal{R}}(W_t) \leq \left\|W_t - \overline{W}\right\|_F^2 - \left\|W_{t+1} - \overline{W}\right\|_F^2 + 2\eta_t \widehat{\mathcal{R}}^{(t)}\left(\overline{W}\right).$$

Consequently, if we use a constant step size $\eta \leq 1$ for $0 \leq \tau < t$, then

$$\eta \left(\sum_{\tau < t} \widehat{\mathcal{R}}(W_\tau)\right) + \left\|W_t - \overline{W}\right\|_F^2 \leq \left\|W_0 - \overline{W}\right\|_F^2 + 2\eta \left(\sum_{\tau < t} \widehat{\mathcal{R}}^{(\tau)}\left(\overline{W}\right)\right).$$

The proof of Lemma 2.6 starts from the standard iteration guarantee:

$$\left\|W_{t+1} - \overline{W}\right\|_F^2 = \left\|W_t - \overline{W}\right\|_F^2 - 2\eta_t \left\langle \nabla \widehat{\mathcal{R}}(W_t), W_t - \overline{W} \right\rangle + \eta_t^2 \left\|\nabla \widehat{\mathcal{R}}(W_t)\right\|_F^2.$$

We can then handle the inner product term using the convexity of ℓ and homogeneity of ReLU, and control $\|\nabla \widehat{\mathcal{R}}(W_t)\|_F^2$ by $\widehat{\mathcal{R}}(W_t)$ using the above properties of $\widehat{\mathcal{Q}}(W_t)$. Lemma 2.6 is similar to (Allen-Zhu & Li, 2019a, Fact D.4 and Claim D.5), where the squared loss is considered.

Using Lemmas 2.3 to 2.6, we can prove Theorem 2.2. Below is a proof sketch; the full proof is given in Appendix A.

1. We first show that as long as $\|w_{s,t} - w_{s,0}\|_2 \leq 4\lambda/(\gamma\sqrt{m})$ for all $1 \leq s \leq m$, it holds that $\widehat{\mathcal{R}}^{(t)}\left(W_0 + \lambda\overline{U}\right) \leq \epsilon/4$. To see this, let us consider $\widehat{\mathcal{R}}^{(0)}$ first. For any $1 \leq i \leq n$, Lemma 2.5 ensures that $|\langle \nabla f_i(W_0), W_0 \rangle|$ is bounded, while Lemma 2.3 ensures that $\langle \nabla f_i(W_0), \overline{U} \rangle$ is concentrated around γ with a large width. As a result, with the chosen λ in Theorem 2.2, we can show that $\langle \nabla f_i(W_0), W_0 + \lambda\overline{U} \rangle$ is large, and $\widehat{\mathcal{R}}^{(0)}(W_0 + \lambda\overline{U})$ is small due to the exponential tail of the logistic loss. To further handle $\widehat{\mathcal{R}}^{(t)}$, we use a standard NTK argument to control $\langle \nabla f_i(W_t) - \nabla f_i(W_0), W_0 + \lambda\overline{U} \rangle$ under the condition that $\|w_{s,t} - w_{s,0}\|_2 \leq 4\lambda/(\gamma\sqrt{m})$.

2. We then prove by contradiction that the above bound on $\|w_{s,t} - w_{s,0}\|_2$ holds for at least the first T iterations. The key observation is that as long as $\widehat{\mathcal{R}}^{(t)}(W_0 + \lambda\overline{U}) \leq \epsilon/4$, we can use it and Lemma 2.6 to control $\sum_{\tau < t} \widehat{\mathcal{Q}}(W_\tau)$, and then just invoke $\|w_{s,t} - w_{s,0}\|_2 \leq \eta \sum_{\tau < t} \widehat{\mathcal{Q}}(W_\tau)/\sqrt{m}$.

   The quantity $\sum_{\tau < t} \widehat{\mathcal{Q}}(W_\tau)$ is also considered in prior work (Cao & Gu, 2019a; Nitanda & Suzuki, 2019), where it is bounded by $\sqrt{t}\sqrt{\sum_{\tau < t} \widehat{\mathcal{Q}}(W_\tau)^2}$ using the Cauchy-Schwarz inequality, which introduces a $\sqrt{t}$ factor. To make the required width depend only on

polylog$(1/\epsilon)$, we also need an upper bound on $\sum_{\tau < t} \widehat{\mathcal{Q}}(W_\tau)$ which depends only on polylog$(1/\epsilon)$. Since the above analysis results in a $\sqrt{t}$ factor, and in our case $\Omega(1/\epsilon)$ steps are needed, it is unclear how to get a polylog$(1/\epsilon)$ width using the analysis in (Cao & Gu, 2019a; Nitanda & Suzuki, 2019). By contrast, using Lemma 2.6, we can show that $\sum_{\tau < t} \widehat{\mathcal{Q}}(W_\tau) \leq 4\lambda/\gamma$, which only depends on $\ln(1/\epsilon)$.

3. The claims of Theorem 2.2 then follow directly from the above two steps and Lemma 2.6.

3 GENERALIZATION

To get a generalization bound, we naturally extend Assumption 2.1 to the following assumption.

Assumption 3.1. There exists $\bar{v} \in \mathcal{H}$ and $\gamma > 0$, s.t. $\left\| \bar{v}(z) \right\|_2 \leq 1$ for any $z \in \mathbb{R}^d$, and

$$ y \int \left\langle \bar{v}(z), x \right\rangle \mathbb{1}\left[\langle z, x \rangle > 0 \right] \mathrm{d}\mu_{\mathcal{N}}(z) \geq \gamma $$

for almost all (x, y) sampled from the data distribution $\mathcal{D}$.   ◇

The above assumption is also made in (Nitanda & Suzuki, 2019) for smooth activations. (Cao & Gu, 2019a) make a similar separability assumption, but in the RKHS induced by the second layer a; by contrast, Assumption 3.1 is on separability in the RKHS induced by the first layer W.

Here is our test error bound with Assumption 3.1.

Theorem 3.2. *Under Assumption 3.1, given any $\epsilon \in (0, 1)$ and any $\delta \in (0, 1/4)$, let λ and M be given as in Theorem 2.2:*

$$ \lambda := \frac{\sqrt{2\ln(4n/\delta)} + \ln(4/\epsilon)}{\gamma/4}, \quad \text{and} \quad M := \frac{4096\lambda^2}{\gamma^6}. $$

Then for any $m \geq M$ and any constant step size $\eta \leq 1$, with probability $1 - 4\delta$ over the random initialization and data sampling,

$$ P_{(x,y)\sim\mathcal{D}}\left(yf(x; W_k, a) \leq 0 \right) \leq 2\epsilon + \frac{16\left(\sqrt{2\ln(4n/\delta)} + \ln(4/\epsilon) \right)}{\gamma^2\sqrt{n}} + 6\sqrt{\frac{\ln(2/\delta)}{2n}}, $$

where k denotes the step with the minimum empirical risk before $\lceil 2\lambda^2/\eta\epsilon \rceil$.

Below is a direct corollary of Theorem 3.2.

Corollary 3.3. *Under Assumption 3.1, given any $\epsilon, \delta \in (0, 1)$, using a constant step size no larger than 1 and let*

$$ n = \widetilde{\Omega}\left(\frac{1}{\gamma^4 \epsilon^2} \right), \quad \text{and} \quad m = \Omega\left(\frac{\ln(n/\delta) + \ln(1/\epsilon)^2}{\gamma^8} \right), $$

it holds with probability $1 - \delta$ that $P_{(x,y)\sim\mathcal{D}}\left(yf(x; W_k, a) \leq 0 \right) \leq \epsilon$, where k denotes the step with the minimum empirical risk in the first $\widetilde{O}(1/\gamma^2\epsilon)$ steps.

To prove Theorem 3.2, we consider the sigmoid mapping $-\ell'(z) = e^{-z}/(1 + e^{-z})$, the empirical average $\widehat{\mathcal{Q}}(W_k)$, and the corresponding population average $\mathcal{Q}(W_k) := \mathbb{E}_{(x,y)\sim\mathcal{D}}\left[-\ell'\left(yf(x; W_k, a) \right) \right]$. First of all, since $P_{(x,y)\sim\mathcal{D}}\left(yf(x; W_k, a) \leq 0 \right) \leq 2\mathcal{Q}(W_k)$, it is enough to control $\mathcal{Q}(W_k)$. Next, as $\widehat{\mathcal{Q}}(W_k)$ is controlled by Theorem 2.2, it is enough to control the generalization error $\mathcal{Q}(W_k) - \widehat{\mathcal{Q}}(W_k)$. Moreover, since $-\ell'$ is supported on $[0, 1]$ and 1-Lipschitz, it is enough to bound the Rademacher complexity of the function space explored by gradient descent. Invoking the bound on $\left\| W_k^\top - W_0^\top \right\|_{2,\infty}$ finishes the proof. The proof details are given in Appendix B.

Remark 3.4. To get Theorem 3.2, we use the Lipschitz-based Rademacher complexity bound. One can also use the smoothness-based Rademacher complexity bound (Srebro et al., 2010, Theorem 1) and get a sample complexity $\widetilde{\Omega}(1/\gamma^4\epsilon)$. However, the bound will become complicated and some large constant will be introduced. It is an interesting open question to give a clean analysis based on smoothness.   ◇

4 STOCHASTIC GRADIENT DESCENT

There are some different formulations of SGD. In this section, we consider SGD with an online oracle. We randomly sample W_0 and a, and fix a during training. At step i, a data example (x_i, y_i) is sampled from the data distribution. We still let $f_i(W) := f(x_i; W, a)$, and perform the following update

$$W_{i+1} := W_i - \eta_i \ell' \left(y_i f_i(W_i) \right) y_i \nabla f_i(W_i).$$

Note that here i starts from 0.

Still with Assumption 3.1, we show the following result.

Theorem 4.1. *Under Assumption 3.1, given any $\epsilon, \delta \in (0, 1)$, using a constant step size and $m = \Omega\left(\left(\ln(1/\delta) + \ln(1/\epsilon)^2 \right)/\gamma^8 \right)$, it holds with probability $1 - \delta$ that*

$$\frac{1}{n} \sum_{i=1}^{n} P_{(x,y) \sim \mathcal{D}} \left(y f(x; W_i, a) \leq 0 \right) \leq \epsilon, \quad \text{for} \quad n = \widetilde{O}(1/\gamma^2 \epsilon).$$

Below is a proof sketch; the details are given in Appendix C. For any W, define

$$\mathcal{R}_i(W) := \ell\left(y_i \left\langle \nabla f_i(W_i), W \right\rangle \right), \quad \text{and} \quad \mathcal{Q}_i(W) := -\ell'\left(y_i \left\langle \nabla f_i(W_i), W \right\rangle \right).$$

Due to homogeneity, $\mathcal{R}_i(W_i) = \ell\left(y_i f_i(W_i) \right)$ and $\mathcal{Q}_i(W_i) = -\ell'\left(y_i f_i(W_i) \right)$.

The first step is an extension of Lemma 2.6 to the SGD setting. The proofs are similar.

Lemma 4.2. *With a constant step size $\eta \leq 1$, for any $\overline{W}$ and any $i \geq 0$,*

$$\eta \left(\sum_{t<i} \mathcal{R}_t(W_t) \right) + \left\| W_i - \overline{W} \right\|_F^2 \leq \left\| W_0 - \overline{W} \right\|_F^2 + 2\eta \left(\sum_{t<i} \mathcal{R}_t\left(\overline{W} \right) \right).$$

With Lemma 4.2, we can also extend Theorem 2.2 to the SGD setting and get a bound on $\sum_{i<n} \mathcal{Q}_i(W_i)$, using a similar proof. To further get a bound on the cumulative population risk $\sum_{i<n} \mathcal{Q}(W_i)$, the key observation is that $\sum_{i<n} \left(\mathcal{Q}(W_i) - \mathcal{Q}_i(W_i) \right)$ is a martingale. Using a martingale Bernstein bound, we prove the following lemma; applying it finishes the proof of Theorem 4.1.

Lemma 4.3. *Given any $\delta \in (0, 1)$, with probability $1 - \delta$,*

$$\sum_{t<i} \mathcal{Q}(W_t) \leq 4 \sum_{t<i} \mathcal{Q}_t(W_t) + 4 \ln\left(\frac{1}{\delta} \right).$$

5 ON SEPARABILITY

Given a training set $\left\{ (x_i, y_i) \right\}_{i=1}^{n}$, the linear kernel is defined as $K_0(x_i, x_j) := \left\langle x_i, x_j \right\rangle$. The maximum margin by a linear classifier is given by

$$\gamma_0 := \min_{q \in \Delta_n} \sqrt{(q \odot y)^\top K_0 (q \odot y)}. \tag{5.1}$$

where Δ_n denotes the probability simplex and $\odot$ denotes the Hadamard product. In addition to the dual definition eq. (5.1), when $\gamma_0 > 0$ there also exists a maximum margin classifier $\bar{u}$ which gives a primal characterization of γ_0: it holds that $\|\bar{u}\|_2 = 1$ and $y_i \left\langle \bar{u}, x_i \right\rangle \geq \gamma_0$ for all i.

In this paper we consider the infinite-width NTK with respect to the first layer:

$$K_1\left(x_i, x_j \right) := \mathbb{E} \left[\frac{\partial f(x_i; W_0, a)}{\partial W_0}, \frac{\partial f(x_j; W_0, a)}{\partial W_0} \right]$$

$$= \mathbb{E}_{w \sim \mathcal{N}(0, I_d)} \left[\left\langle x_i \mathbb{1}\left[\left\langle x_i, w \right\rangle > 0 \right], x_j \mathbb{1}\left[\left\langle x_j, w \right\rangle > 0 \right] \right\rangle \right] = \left\langle \phi_i, \phi_j \right\rangle_{\mathcal{H}}.$$

Here ϕ and $\mathcal{H}$ are defined at the beginning of Section 2. We can see that $\mathcal{H}$ is indeed the RKHS induced by the infinite-width NTK with respect to the first layer.

Similar to the dual definition of γ_0, the margin given by K_1 is defined as

$$\gamma_1 := \min_{q \in \Delta_n} \sqrt{(q \odot y)^\top K_1 (q \odot y)}. \tag{5.2}$$

To clarify, below we only use γ_1 with the definition eq. (5.2), and γ as given by Assumption 2.1.

We can also give a primal characterization of γ_1 when it is positive.

Proposition 5.1. *If $\gamma_1 > 0$, then there exists $\hat{v} \in \mathcal{H}$ such that $\|\hat{v}\|_{\mathcal{H}} = 1$, and $y_i \langle \hat{v}, \phi_i \rangle_{\mathcal{H}} \geq \gamma_1$ for any $1 \leq i \leq n$. Additionally $\|\hat{v}(z)\|_2 \leq 1/\gamma_1$ for any $z \in \mathbb{R}^d$.*

The proof is given in Appendix D, which uses the Fenchel duality theory. Using the upper bound $\|\hat{v}(z)\|_2 \leq 1/\gamma_1$, we can see that $\gamma_1 \hat{v}$ satisfies Assumption 2.1 with $\gamma \geq \gamma_1^2$. However, such an upper bound $\|\hat{v}(z)\|_2 \leq 1/\gamma_1$ might be too loose, which leads to a bad rate. In fact, as we show in Appendix E, in some cases we can construct $\bar{v}$ directly which satisfies Assumption 2.1 with a large γ. For this reason, we choose to assume Assumption 2.1 instead of a positive γ_1.

Below is some further analysis of γ_1 using the dual characterization eq. (5.2). These bounds mostly hold for arbitrary labels and random labels, and thus are worst case; see Appendix E for some examples of large γ.

(Oymak & Soltanolkotabi, 2019, Corollary I.2) show that if for any two feature vectors x_i and x_j, we have $\|x_i - x_j\|_2 \geq \theta$ and $\|x_i + x_j\|_2 \geq \theta$ for some $\theta > 0$, then the minimum eigenvalue of K_1 is at least $\theta/(100n^2)$. For arbitrary labels $y \in \{-1, +1\}^n$, since $\|q \odot y\|_2 \geq 1/\sqrt{n}$, we have the worst case bound $\gamma_1^2 \geq \theta/100n^3$. A direct improvement of this bound is $\theta/100n_S^3$, where n_S denotes the number of support vectors, which could be much smaller than n with real world data.

Another observation is that given any training set $\{(x_i, y_i)\}_{i=1}^n$ which may have a large margin, replacing y with random labels would destroy the margin, which is what should be expected.

Proposition 5.2. *Given any training set $\{(x_i, y_i)\}_{i=1}^n$, if we replace y with random labels $\epsilon \sim$ unif $(\{-1, +1\}^n)$, then with probability 0.9 over the random labels, it holds that $\gamma_1 \leq 1/\sqrt{20n}$.*

Proof. Let $\hat{q}$ denote the uniform probability vector $(1/n, \ldots, 1/n)$. Note that

$$\mathbb{E}_{\epsilon \sim \text{unif}(\{-1,+1\}^n)} \left[(\hat{q} \odot \epsilon)^\top K_1 (\hat{q} \odot \epsilon) \right] = \mathbb{E}_{\epsilon \sim \text{unif}(\{-1,+1\}^n)} \left[\sum_{i,j=1}^n \frac{1}{n^2} \epsilon_i \epsilon_j K_1(x_i, x_j) \right]$$

$$= \frac{1}{n^2} \sum_{i,j=1}^n \mathbb{E}_{\epsilon \sim \text{unif}(\{-1,+1\}^n)} \left[\epsilon_i \epsilon_j K_1(x_i, x_j) \right]$$

$$= \frac{1}{n^2} \sum_{i=1}^n K_1(x_i, x_i) = \frac{1}{2n}.$$

Since $0 \leq (\hat{q} \odot \epsilon)^\top K_1 (\hat{q} \odot \epsilon) \leq 1$ for any ϵ, by Markov's inequality with probability 0.9, it holds that $(\hat{q} \odot \epsilon)^\top K_1 (\hat{q} \odot \epsilon) \leq 1/(20n)$, and thus $\gamma_1 \leq 1/\sqrt{20n}$. $\qquad\qquad\square$

6 OPEN PROBLEMS

In this paper, we analyze gradient descent on a two-layer network in the NTK regime, where the weights stay close to the initialization. It is an interesting open question if gradient descent learns something beyond the NTK, after the iterates move far enough from the initial weights. It is also interesting to extend our analysis to other architectures, such as multi-layer networks, convolutional networks, and residual networks. Finally, in this paper we only discuss binary classification; it is interesting to see if it is possible to get similar results for other tasks, such as regression.

REFERENCES

Zeyuan Allen-Zhu and Yuanzhi Li. What can resnet learn efficiently, going beyond kernels? *arXiv preprint arXiv:1905.10337*, 2019a.

Zeyuan Allen-Zhu and Yuanzhi Li. Can sgd learn recurrent neural networks with provable generalization? *arXiv preprint arXiv:1902.01028*, 2019b.

Zeyuan Allen-Zhu, Yuanzhi Li, and Yingyu Liang. Learning and generalization in overparameterized neural networks, going beyond two layers. *arXiv preprint arXiv:1811.04918*, 2018a.

Zeyuan Allen-Zhu, Yuanzhi Li, and Zhao Song. A convergence theory for deep learning via over-parameterization. *arXiv preprint arXiv:1811.03962*, 2018b.

Zeyuan Allen-Zhu, Yuanzhi Li, and Zhao Song. On the convergence rate of training recurrent neural networks. *arXiv preprint arXiv:1810.12065*, 2018c.

Sanjeev Arora, Simon S Du, Wei Hu, Zhiyuan Li, and Ruosong Wang. Fine-grained analysis of optimization and generalization for overparameterized two-layer neural networks. *arXiv preprint arXiv:1901.08584*, 2019.

Peter L. Bartlett and Shahar Mendelson. Rademacher and gaussian complexities: Risk bounds and structural results. *JMLR*, 3:463–482, Nov 2002.

Alina Beygelzimer, John Langford, Lihong Li, Lev Reyzin, and Robert Schapire. Contextual bandit algorithms with supervised learning guarantees. In *Proceedings of the Fourteenth International Conference on Artificial Intelligence and Statistics*, pp. 19–26, 2011.

Jonathan M. Borwein and Qiji J. Zhu. *Techniques of Variational Analysis, volume 20 of*. CMS Books in Mathematics, 2005.

Yuan Cao and Quanquan Gu. Generalization error bounds of gradient descent for learning over-parameterized deep relu networks. *arXiv preprint arXiv:1902.01384*, 2019a.

Yuan Cao and Quanquan Gu. Generalization bounds of stochastic gradient descent for wide and deep neural networks. *arXiv preprint arXiv:1905.13210*, 2019b.

Lenaic Chizat and Francis Bach. A Note on Lazy Training in Supervised Differentiable Programming. `arXiv:1812.07956v2 [math.OC]`, 2019.

Simon S Du, Jason D Lee, Haochuan Li, Liwei Wang, and Xiyu Zhai. Gradient descent finds global minima of deep neural networks. *arXiv preprint arXiv:1811.03804*, 2018a.

Simon S Du, Xiyu Zhai, Barnabas Poczos, and Aarti Singh. Gradient descent provably optimizes over-parameterized neural networks. *arXiv preprint arXiv:1810.02054*, 2018b.

Arthur Jacot, Franck Gabriel, and Clément Hongler. Neural tangent kernel: Convergence and generalization in neural networks. In *Advances in neural information processing systems*, pp. 8571–8580, 2018.

Ziwei Ji and Matus Telgarsky. Risk and parameter convergence of logistic regression. *arXiv preprint arXiv:1803.07300v2*, 2018.

Yuanzhi Li and Yingyu Liang. Learning overparameterized neural networks via stochastic gradient descent on structured data. In *Advances in Neural Information Processing Systems*, pp. 8157–8166, 2018.

Percy Liang. Stanford CS229T/STAT231: Statistical Learning Theory, Apr 2016. URL `https://web.stanford.edu/class/cs229t/notes.pdf`.

Atsushi Nitanda and Taiji Suzuki. Refined generalization analysis of gradient descent for over-parameterized two-layer neural networks with smooth activations on classification problems. *arXiv preprint arXiv:1905.09870*, 2019.

Samet Oymak and Mahdi Soltanolkotabi. Towards moderate overparameterization: global convergence guarantees for training shallow neural networks. *arXiv preprint arXiv:1902.04674*, 2019.

Shai Shalev-Shwartz and Shai Ben-David. *Understanding Machine Learning: From Theory to Algorithms*. Cambridge University Press, 2014.

Zhao Song and Xin Yang. Quadratic suffices for over-parametrization via matrix chernoff bound. *arXiv preprint arXiv:1906.03593*, 2019.

Nathan Srebro, Karthik Sridharan, and Ambuj Tewari. Smoothness, low noise and fast rates. In *Advances in neural information processing systems*, pp. 2199–2207, 2010.

Martin J. Wainwright. UC Berkeley Statistics 210B, Lecture Notes: Basic tail and concentration bounds, Jan 2015. URL `https://www.stat.berkeley.edu/~mjwain/stat210b/Chap2_TailBounds_Jan22_2015.pdf`.

Colin Wei, Jason D Lee, Qiang Liu, and Tengyu Ma. Regularization matters: Generalization and optimization of neural nets vs their induced kernel. *arXiv preprint arXiv:1810.05369*, 2018.

Difan Zou and Quanquan Gu. An improved analysis of training over-parameterized deep neural networks. *arXiv preprint arXiv:1906.04688*, 2019.

Difan Zou, Yuan Cao, Dongruo Zhou, and Quanquan Gu. Stochastic gradient descent optimizes over-parameterized deep relu networks. *arXiv preprint arXiv:1811.08888*, 2018.

A OMITTED PROOFS FROM SECTION 2

Proof of Lemma 2.3. By Assumption 2.1, given any $1 \leq i \leq n$,

$$\mu := \mathbb{E}_{w \sim \mathcal{N}(0, I_d)} \left[y_i \left\langle \bar{v}(w), x_i \right\rangle \mathbb{1} \left[\left\langle w, x_i \right\rangle > 0 \right] \right] \geq \gamma.$$

On the other hand,

$$y_i f_i^{(0)} \left(\overline{U} \right) = \frac{1}{m} \sum_{s=1}^m y_i \left\langle \bar{v}(w_{s,0}), x_i \right\rangle \mathbb{1} \left[\left\langle w_{s,0}, x_i \right\rangle > 0 \right]$$

is the empirical mean of i.i.d. r.v.'s supported on $[-1, +1]$ with mean μ. Therefore by Hoeffding's inequality, with probability $1 - \delta/n$,

$$y_i f_i^{(0)} \left(\overline{U} \right) - \gamma \geq y_i f_i^{(0)} \left(\overline{U} \right) - \mu \geq -\sqrt{\frac{2 \ln(n/\delta)}{m}}.$$

Applying the union bound finishes the proof. $\qquad \square$

Proof of Lemma 2.4. Given any fixed ϵ_2 and $1 \leq i \leq n$,

$$\mathbb{E} \left[\alpha_i(W_0, \epsilon_2) \right] = \mathbb{P} \left(\left| \left\langle w, x_i \right\rangle \right| \leq \epsilon_2 \right) \leq \frac{2\epsilon_2}{\sqrt{2\pi}} = \sqrt{\frac{2}{\pi}} \epsilon_2,$$

because $\left\langle w, x_i \right\rangle$ is a standard Gaussian r.v. and the density of standard Gaussian has maximum $1/\sqrt{2\pi}$. Since $\alpha_i(W_0, \epsilon_2)$ is the empirical mean of Bernoulli r.v.'s, by Hoeffding's inequality, with probability $1 - \delta/n$,

$$\alpha_i(W_0, \epsilon_2) \leq \mathbb{E} \left[\alpha_i(W_0, \epsilon_2) \right] + \sqrt{\frac{\ln(n/\delta)}{2m}} \leq \sqrt{\frac{2}{\pi}} \epsilon_2 + \sqrt{\frac{\ln(n/\delta)}{2m}}.$$

Applying the union bound finishes the proof. $\qquad \square$

To prove Lemma 2.5, we need the following technical result.

Lemma A.1. *Consider the random vector $X = (X_1, \ldots, X_m)$, where $X_i = \sigma(Z_i)$ for some $\sigma : \mathbb{R} \to \mathbb{R}$ that is 1-Lipschitz, and Z_i are i.i.d. standard Gaussian r.v.'s. Then the r.v. $\|X\|_2$ is 1-sub-Gaussian, and thus with probability $1 - \delta$,*

$$\|X\|_2 - \mathbb{E}\left[\|X\|_2\right] \leq \sqrt{2\ln(1/\delta)}.$$

Proof. Given $a \in \mathbb{R}^m$, define

$$f(a) = \sqrt{\sum_{i=1}^{m} \sigma(a_i)^2} = \left\|\sigma(a)\right\|_2,$$

where $\sigma(a)$ is obtained by applying σ coordinate-wisely to a. For any $a, b \in \mathbb{R}^m$, by the triangle inequality, we have

$$\left|f(a) - f(b)\right| = \left|\left\|\sigma(a)\right\|_2 - \left\|\sigma(b)\right\|_2\right| \leq \left\|\sigma(a) - \sigma(b)\right\|_2 = \sqrt{\sum_{i=1}^{m} \left(\sigma(a_i) - \sigma(b_i)\right)^2},$$

and by further using the 1-Lipschitz continuity of σ, we have

$$\left|f(a) - f(b)\right| \leq \sqrt{\sum_{i=1}^{m} \left(\sigma(a_i) - \sigma(b_i)\right)^2} \leq \sqrt{\sum_{i=1}^{m} (a_i - b_i)^2} = \|a - b\|_2.$$

As a result, f is a 1-Lipschitz continuous function w.r.t. the ℓ_2 norm, and by Theorem 2.4 of (Wainwright, 2015), $f(X)$ is 1-sub-Gaussian. The remaining bound comes directly from Hoeffding's inequality. $\square$

Proof of Lemma 2.5. Given $1 \leq i \leq n$, let $h_i = \sigma(W_0 x_i)/\sqrt{m}$. By Lemma A.1, $\|h_i\|_2$ is sub-Gaussian with variance proxy $1/m$. By Hoeffding's inequality, with probability $1 - \delta/2n$ over W_0,

$$\|h_i\|_2 - \mathbb{E}\left[\|h_i\|_2\right] \leq \sqrt{\frac{2\ln(2n/\delta)}{m}} \leq \sqrt{\frac{2\ln(2n/\delta)}{25\ln(2n/\delta)}} \leq 1 - \frac{\sqrt{2}}{2}.$$

On the other hand, by Jensen's inequality,

$$\mathbb{E}\left[\|h_i\|_2\right] \leq \sqrt{\mathbb{E}\left[\|h_i\|_2^2\right]} = \frac{\sqrt{2}}{2}.$$

As a result, with probability $1 - \delta/2n$, it holds that $\|h_i\|_2 \leq 1$. By the union bound, with probability $1 - \delta/2$ over W_0, for all $1 \leq i \leq n$, we have $\|h_i\|_2 \leq 1$.

For any W_0 such that the above event holds, and for any $1 \leq i \leq n$, the r.v. $\langle h_i, a \rangle$ is sub-Gaussian with variance proxy $\|h_i\|_2^2 \leq 1$. By Hoeffding's inequality, with probability $1 - \delta/2n$ over a,

$$\left|\langle h_i, a \rangle\right| = \left|f(x_i; W_0, a)\right| \leq \sqrt{2\ln\left(4n/\delta\right)}.$$

By the union bound, with probability $1 - \delta/2$ over a, for all $1 \leq i \leq n$, we have $\left|f(x_i; W_0, a)\right| \leq \sqrt{2\ln\left(4n/\delta\right)}$.

The probability that the above events all happen is at least $(1 - \delta/2)(1 - \delta/2) \geq 1 - \delta$, over W_0 and a. $\square$

Proof of Lemma 2.6. We have

$$\left\|W_{t+1} - \overline{W}\right\|_F^2 = \left\|W_t - \overline{W}\right\|_F^2 - 2\eta_t \left\langle \nabla\widehat{\mathcal{R}}(W_t), W_t - \overline{W} \right\rangle + \eta_t^2 \left\|\nabla\widehat{\mathcal{R}}(W_t)\right\|_F^2. \tag{A.1}$$

The first order term of eq. (A.1) can be handled using the convexity of ℓ and homogeneity of ReLU:

$$\left\langle \nabla \widehat{\mathcal{R}}(W_t), W_t - \overline{W} \right\rangle = \frac{1}{n} \sum_{i=1}^{n} \ell' \left(y_i f_i(W_t) \right) y_i \left\langle \nabla f_i(W_t), W_t - \overline{W} \right\rangle$$

$$= \frac{1}{n} \sum_{i=1}^{n} \ell' \left(y_i f_i(W_t) \right) \left(y_i f_i(W_t) - y_i f_i^{(t)} \left(\overline{W} \right) \right)$$

$$\geq \frac{1}{n} \sum_{i=1}^{n} \left(\ell \left(y_i f_i(W_t) \right) - \ell \left(y_i f_i^{(t)} \left(\overline{W} \right) \right) \right) = \widehat{\mathcal{R}}(W_t) - \widehat{\mathcal{R}}^{(t)} \left(\overline{W} \right).$$

$$\text{(A.2)}$$

The second-order term of eq. (A.1) can be bounded as follows

$$\eta_t^2 \left\| \nabla \widehat{\mathcal{R}}(W_t) \right\|_F^2 \leq \eta_t^2 \widehat{\mathcal{Q}}(W_t)^2 \leq \eta_t \widehat{\mathcal{Q}}(W_t) \leq \eta_t \widehat{\mathcal{R}}(W_t), \tag{A.3}$$

because $\left\| \nabla \widehat{\mathcal{R}}(W_t) \right\|_F \leq \widehat{\mathcal{Q}}(W_t)$, and $\eta_t, \widehat{\mathcal{Q}}(W_t) \leq 1$, and $\widehat{\mathcal{Q}}(W_t) \leq \widehat{\mathcal{R}}(W_t)$. Combining eqs. (A.1) to (A.3) gives

$$\eta_t \widehat{\mathcal{R}}(W_t) \leq \left\| W_t - \overline{W} \right\|_F^2 - \left\| W_{t+1} - \overline{W} \right\|_F^2 + 2\eta_t \widehat{\mathcal{R}}^{(t)} \left(\overline{W} \right).$$

Telescoping gives the other claim. $\qquad \square$

Proof of Theorem 2.2. The required width ensures that with probability $1 - 3\delta$, Lemmas 2.3 to 2.5 hold with $\epsilon_1 = \gamma^2/8$ and $\epsilon_2 = 4\lambda/(\gamma\sqrt{m})$.

Let t_1 denote the first step such that there exists $1 \leq s \leq m$ with $\left\| w_{s,t_1} - w_{s,0} \right\|_2 > 4\lambda/(\gamma\sqrt{m})$. Therefore for any $0 \leq t < t_1$ and any $1 \leq s \leq m$, it holds that $\left\| w_{s,t} - w_{s,0} \right\|_2 \leq 4\lambda/(\gamma\sqrt{m})$. In addition, we let $\overline{W} := W_0 + \lambda\overline{U}$.

We first prove that for any $0 \leq t < t_1$, it holds that $\widehat{\mathcal{R}}^{(t)} \left(\overline{W} \right) \leq \epsilon/4$. Since $\ln(1+r) \leq r$ for any r, the logistic satisfies $\ell(z) = \ln(1 + \exp(-z)) \leq \exp(-z)$, and it is enough to prove that for any $1 \leq i \leq n$,

$$y_i \left\langle \nabla f_i(W_t), \overline{W} \right\rangle \geq \ln \left(\frac{4}{\epsilon} \right).$$

We will split the left hand side into three terms and control them individually:

$$y_i \left\langle \nabla f_i(W_t), \overline{W} \right\rangle = y_i \left\langle \nabla f_i(W_0), W_0 \right\rangle + y_i \left\langle \nabla f_i(W_t) - \nabla f_i(W_0), W_0 \right\rangle + \lambda y_i \left\langle \nabla f_i(W_t), \overline{U} \right\rangle.$$

$$\text{(A.4)}$$

- The first term of eq. (A.4) can be controlled using Lemma 2.5:

$$\left| y_i \left\langle \nabla f_i(W_0), W_0 \right\rangle \right| \leq \sqrt{2 \ln(4n/\delta)}. \tag{A.5}$$

- The second term of eq. (A.4) can be written as

$$y_i \left\langle \nabla f_i(W_t) - \nabla f_i(W_0), W_0 \right\rangle = y_i \frac{1}{\sqrt{m}} \sum_{s=1}^{m} a_s \left(\mathbb{1} \left[\langle w_{s,t}, x_i \rangle > 0 \right] - \mathbb{1} \left[\langle w_{s,0}, x_i \rangle > 0 \right] \right) \langle w_{s,0}, x_i \rangle.$$

Let $S_c := \left\{ s \Big| \mathbb{1} \left[\langle w_{s,t}, x_i \rangle > 0 \right] - \mathbb{1} \left[\langle w_{s,0}, x_i \rangle > 0 \right] \neq 0, 1 \leq s \leq m \right\}$. Note that $s \in S_c$ implies

$$\left| \langle w_{s,0}, x_i \rangle \right| \leq \left| \langle w_{s,t} - w_{s,0}, x_i \rangle \right| \leq \left\| w_{s,t} - w_{s,0} \right\|_2 \left\| x_i \right\|_2 = \left\| w_{s,t} - w_{s,0} \right\|_2 \leq 4\lambda/(\gamma\sqrt{m}) = \epsilon_2.$$

Therefore Lemma 2.4 ensures that

$$|S_c| \le \left| \left\{ s \,\middle|\, |\langle w_{s,0}, x_i \rangle| \le \epsilon_2 \right\} \right| \le m \left(\frac{4\lambda}{\gamma\sqrt{m}} + \frac{\epsilon_1}{2} \right) = m \left(\frac{4\lambda}{\gamma\sqrt{m}} + \frac{\gamma^2}{16} \right).$$

and thus

$$\left| y_i \left\langle \nabla f_i(W_t) - \nabla f_i(W_0), W_0 \right\rangle \right| \le \frac{1}{\sqrt{m}} \cdot |S_c| \cdot \frac{4\lambda}{\gamma\sqrt{m}} \le \frac{16\lambda^2}{\gamma^2\sqrt{m}} + \frac{\lambda\gamma}{4} \le \frac{\lambda\gamma}{2}, \quad (A.6)$$

where in the last step we use the condition that $m \ge 4096\lambda^2/\gamma^6$.

- The third term of eq. (A.4) can be bounded as follows: by Lemma 2.3,

$$y_i \left\langle \nabla f_i(W_t), \overline{U} \right\rangle = y_i \left\langle \nabla f_i(W_0), \overline{U} \right\rangle + y_i \left\langle \nabla f_i(W_t) - \nabla f_i(W_0), \overline{U} \right\rangle$$

$$\ge \gamma - \epsilon_1 + y_i \left\langle \nabla f_i(W_t) - \nabla f_i(W_0), \overline{U} \right\rangle.$$

In addition,

$$y_i \left\langle \nabla f_i(W_t) - \nabla f_i(W_0), \overline{U} \right\rangle = y_i \frac{1}{m} \sum_{i=1}^{m} \left(\mathbb{1} \left[\langle w_{s,t}, x_i \rangle > 0 \right] - \mathbb{1} \left[\langle w_{s,0}, x_i \rangle > 0 \right] \right) \left\langle \overline{v}(w_{s,0}), x_i \right\rangle$$

$$\ge -\frac{1}{m} \cdot |S_c| \ge -\frac{4\lambda}{\gamma\sqrt{m}} - \frac{\epsilon_1}{2} \ge -\frac{\gamma^2}{16} - \frac{\epsilon_1}{2},$$

where we use $m \ge 4096\lambda^2/\gamma^6$. Therefore,

$$y_i \left\langle \nabla f_i(W_t), \overline{U} \right\rangle \ge \gamma - \epsilon_1 - \frac{\gamma^2}{16} - \frac{\epsilon_1}{2} = \gamma - \frac{\gamma^2}{4} \ge \frac{3\gamma}{4}. \quad (A.7)$$

Putting eqs. (A.5) to (A.7) into eq. (A.4), we have

$$y_i \left\langle \nabla f_i(W_t), \overline{W} \right\rangle \ge -\sqrt{2\ln\left(\frac{4n}{\delta}\right)} - \frac{\lambda\gamma}{2} + \frac{3\lambda\gamma}{4} = \frac{\lambda\gamma}{4} - \sqrt{2\ln\left(\frac{4n}{\delta}\right)} = \ln\left(\frac{4}{\epsilon}\right),$$

for the λ given in the statement of Theorem 2.2. Consequently, for any $0 \le t < t_1$, it holds that $\widehat{\mathcal{R}}^{(t)}\left(\overline{W}\right) \le \epsilon/4$.

Let $T := \lceil 2\lambda^2/\eta\epsilon \rceil$. The next claim is that $t_1 \ge T$. To see this, note that Lemma 2.6 ensures

$$\left\| W_{t_1} - \overline{W} \right\|_F^2 \le \left\| W_0 - \overline{W} \right\|_F^2 + 2\eta \left(\sum_{t<t_1} \widehat{\mathcal{R}}^{(t)}\left(\overline{W}\right) \right) \le \lambda^2 + \frac{\epsilon}{2}\eta t_1.$$

Suppose $t_1 < T$, then we have $t_1 \le 2\lambda^2/\eta\epsilon$, and thus $\left\| W_{t_1} - \overline{W} \right\|_F^2 \le 2\lambda^2$. As a result, using $\|\overline{U}\|_F \le 1$ and the definition of $\overline{W}$,

$$\sqrt{2}\lambda \ge \left\| W_{t_1} - \overline{W} \right\|_F \ge \left\langle W_{t_1} - \overline{W}, \overline{U} \right\rangle = \left\langle W_{t_1} - W_0, \overline{U} \right\rangle - \left\langle \overline{W} - W_0, \overline{U} \right\rangle$$

$$\ge \left\langle W_{t_1} - W_0, \overline{U} \right\rangle - \lambda.$$

Moreover, due to eq. (A.7),

$$\left\langle W_{t_1} - W_0, \overline{U} \right\rangle = -\eta \sum_{\tau<t_1} \left\langle \nabla\widehat{\mathcal{R}}(W_\tau), \overline{U} \right\rangle = \eta \sum_{\tau<t_1} \frac{1}{n} \sum_{i=1}^{n} -\ell'\left(y_i f_i(W_\tau)\right) y_i \left\langle \nabla f_i(W_\tau), \overline{U} \right\rangle$$

$$\ge \eta \sum_{\tau<t_1} \widehat{\mathcal{Q}}(W_\tau) \frac{3\gamma}{4}.$$

As a result,

$$\eta \sum_{\tau < t_1} \widehat{\mathcal{Q}}(W_\tau) \leq \frac{4(\sqrt{2}+1)\lambda}{3\gamma} \leq \frac{4\lambda}{\gamma}.$$

Furthermore, by the triangle inequality, for any $1 \leq s \leq m$

$$\begin{aligned}
\left\| w_{s,t} - w_{s,0} \right\|_2 &\leq \eta \sum_{\tau < t} \left\| \frac{1}{n} \sum_{i=1}^n \ell' \left(y_i f_i(W_\tau) \right) y_i \frac{\partial f_i}{\partial w_{s,\tau}} \right\|_2 \\
&\leq \eta \sum_{\tau < t} \frac{1}{n} \sum_{i=1}^n \left| \ell' \left(y_i f_i(W_\tau) \right) \right| \cdot \left\| \frac{\partial f_i}{\partial w_{s,\tau}} \right\|_2 \\
&\leq \eta \sum_{\tau < t} \widehat{\mathcal{Q}}(W_\tau) \frac{1}{\sqrt{m}} \\
&\leq \eta \sum_{\tau < t_1} \widehat{\mathcal{Q}}(W_\tau) \frac{1}{\sqrt{m}} \leq \frac{4\lambda}{\gamma\sqrt{m}},
\end{aligned} \tag{A.8}$$

which contradicts the definition of t_1. Therefore $t_1 \geq T$.

Now we are ready to prove the claims of Theorem 2.2. The bound on $\left\| w_{s,t} - w_{s,0} \right\|_2$ follow by repeating the steps in eq. (A.8). The risk guarantee follows from Lemma 2.6:

$$\frac{1}{T} \sum_{t < T} \widehat{\mathcal{R}}(W_t) \leq \frac{\left\| W_0 - \overline{W} \right\|_F^2}{\eta T} + \frac{2}{T} \sum_{t < T} \widehat{\mathcal{R}}^{(t)} \left(\overline{W} \right) \leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

$\square$

B OMITTED PROOFS FROM SECTION 3

The proof of Theorem 3.2 is based on Rademacher complexity. Given a sample $S = (z_1, \ldots, z_n)$ (where $z_i = (x_i, y_i)$) and a function class $\mathcal{H}$, the Rademacher complexity of $\mathcal{H}$ on S is defined as

$$\text{Rad} \left(\mathcal{H} \circ S \right) := \frac{1}{n} \mathbb{E}_{\epsilon \sim \{-1,+1\}^n} \left[\sup_{h \in \mathcal{H}} \sum_{i=1}^n \epsilon_i h(z_i) \right].$$

We will use the following general result.

Lemma B.1. *(Shalev-Shwartz & Ben-David, 2014, Theorem 26.5) If $h(z) \in [a, b]$, then with probability $1 - \delta$,*

$$\sup_{h \in \mathcal{H}} \left(\mathbb{E}_{z \sim \mathcal{D}} \left[h(z) \right] - \frac{1}{n} \sum_{i=1}^n h(z_i) \right) \leq 2\text{Rad} \left(\mathcal{H} \circ S \right) + 3(b - a) \sqrt{\frac{\ln(2/\delta)}{2n}}.$$

We also need the following contraction lemma. Consider a feature sample $X = (x_1, \ldots, x_n)$ and a function class $\mathcal{F}$ on X. For each $1 \leq i \leq n$, let $g_i : \mathbb{R} \to \mathbb{R}$ denote a K-Lipschitz function. Let $g \circ \mathcal{F}$ denote the class of functions which map x_i to $g_i(f(x_i))$ for some $f \in \mathcal{F}$.

Lemma B.2. *(Shalev-Shwartz & Ben-David, 2014, Lemma 26.9)* $\text{Rad} \left(g \circ \mathcal{F} \circ X \right) \leq K\text{Rad} \left(\mathcal{F} \circ X \right).$

To prove Theorem 3.2, we need one more Rademacher complexity bound. Given a fixed initialization (W_0, a), consider the following classes:

$$\mathcal{W}_\rho := \left\{ W \in \mathbb{R}^{m \times d} \middle| \left\| w_s - w_{s,0} \right\|_2 \leq \rho \text{ for any } 1 \leq s \leq m \right\},$$

and

$$\mathcal{F}_\rho := \big\{ x \mapsto f(x; W, a) \big| W \in \mathcal{W}_\rho \big\} .$$

Given a feature sample X, the following Lemma B.3 controls the Rademacher complexity of $\mathcal{F}_\rho \circ X$. A similar version was given in (Liang, 2016, Theorem 43), and the proof is similar to the proof of (Bartlett & Mendelson, 2002, Theorem 18) which also pushes the supreme through and handles each hidden units separately.

Lemma B.3. $\mathrm{Rad}\left(\mathcal{F}_\rho \circ X\right) \leq \rho\sqrt{m/n}.$

Proof of Lemma B.3. We have

$$
\mathbb{E}_\epsilon \left[\sup_{W \in \mathcal{W}_\rho} \sum_{i=1}^n \epsilon_i f(x_i; W, a) \right] = \mathbb{E}_\epsilon \left[\sup_{W \in \mathcal{W}_\rho} \sum_{i=1}^n \epsilon_i \sum_{s=1}^m \frac{1}{\sqrt{m}} a_s \sigma\left(\langle w_s, x_i \rangle\right) \right]
$$

$$
= \mathbb{E}_\epsilon \left[\frac{1}{\sqrt{m}} \sup_{W \in \mathcal{W}_\rho} \sum_{s=1}^m \sum_{i=1}^n \epsilon_i a_s \sigma\left(\langle w_s, x_i \rangle\right) \right]
$$

$$
= \mathbb{E}_\epsilon \left[\frac{1}{\sqrt{m}} \sum_{s=1}^m \left(\sup_{\|w_s - w_{s,0}\|_2 \leq \rho} \sum_{i=1}^n \epsilon_i a_s \sigma\left(\langle w_s, x_i \rangle\right) \right) \right]
$$

$$
= \frac{1}{\sqrt{m}} \sum_{i=1}^m \mathbb{E}_\epsilon \left[\sup_{\|w_s - w_{s,0}\|_2 \leq \rho} \sum_{i=1}^n \epsilon_i a_s \sigma\left(\langle w_s, x_i \rangle\right) \right] .
$$

Note that for any $1 \leq s \leq m$, the mapping $z \mapsto a_s \sigma(z)$ is 1-Lipschitz, and thus Lemma B.2 gives

$$
\mathbb{E}_\epsilon \left[\sup_{W \in \mathcal{W}_\rho} \sum_{i=1}^n \epsilon_i f(x_i; W, a) \right] \leq \frac{1}{\sqrt{m}} \sum_{i=1}^m \mathbb{E}_\epsilon \left[\sup_{\|w_s - w_{s,0}\|_2 \leq \rho} \sum_{i=1}^n \epsilon_i a_s \sigma\left(\langle w_s, x_i \rangle\right) \right]
$$

$$
\leq \frac{1}{\sqrt{m}} \sum_{i=1}^m \mathbb{E}_\epsilon \left[\sup_{\|w_s - w_{s,0}\|_2 \leq \rho} \sum_{i=1}^n \epsilon_i \langle w_s, x_i \rangle \right] .
$$

Invoking the Rademacher complexity bound of linear classifiers (Shalev-Shwartz & Ben-David, 2014, Lemma 26.10) then gives

$$
\mathrm{Rad}\left(\mathcal{F}_\rho \circ X\right) = \frac{1}{n} \mathbb{E}_\epsilon \left[\sup_{W \in \mathcal{W}_\rho} \sum_{i=1}^n \epsilon_i f(x_i; W, a) \right] \leq \frac{\rho\sqrt{m}}{\sqrt{n}} .
$$

$\square$

Now we are ready to prove the main generalization result Theorem 3.2.

Proof. First fix an initialization (W_0, a), and consider $\mathcal{F}_\rho$. Let $\mathcal{H} := \big\{ (x, y) \mapsto -\ell'\left(y f(x)\right) \big| f \in \mathcal{F}_\rho \big\}$. Since for any $h \in \mathcal{H}$ and any z, $h(z) \in [0, 1]$, Lemma B.1 ensures that with probability $1 - \delta$ over the data sampling,

$$
\sup_{h \in \mathcal{H}} \left(\mathbb{E}_{z \sim \mathcal{D}}\left[h(z) \right] - \frac{1}{n} \sum_{i=1}^n h(z_i) \right) = \sup_{W \in \mathcal{W}_\rho} \left(\mathcal{Q}(W) - \widehat{\mathcal{Q}}(W) \right) \leq 2\mathrm{Rad}\left(\mathcal{H} \circ S\right) + 3\sqrt{\frac{\ln(2/\delta)}{2n}} .
$$

Since for each $1 \leq i \leq n$, the mapping $z \mapsto -\ell'(y_i z)$ is $(1/4)$-Lipschitz, Lemma B.2 further ensures that $\mathrm{Rad}\left(\mathcal{H} \circ S\right) \leq \mathrm{Rad}\left(\mathcal{F}_\rho \circ X\right)/4$, and thus

$$
\sup_{W \in \mathcal{W}_\rho} \left(\mathcal{Q}(W) - \widehat{\mathcal{Q}}(W) \right) \leq \frac{\rho\sqrt{m}}{2\sqrt{n}} + 3\sqrt{\frac{\ln(2/\delta)}{2n}} . \tag{B.1}
$$

On the other hand, Theorem 2.2 ensures that under the condition of Theorem 3.2, for any fixed dataset, with probability $1 - 3\delta$ over the random initialization, we have

$$\widehat{\mathcal{Q}}(W_k) \leq \widehat{\mathcal{R}}(W_k) \leq \epsilon, \quad \text{and} \quad \left\| w_{s,k} - w_{s,0} \right\|_2 \leq \frac{4\lambda}{\gamma\sqrt{m}}.$$

As a result, invoking eq. (B.1) with $\rho = 4\lambda/(\gamma\sqrt{m})$, with probability $1 - 4\delta$ over the random initialization and data sampling,

$$\mathcal{Q}(W_k) \leq \widehat{\mathcal{Q}}(W_k) + \frac{2\lambda}{\gamma\sqrt{n}} + 3\sqrt{\frac{\ln(2/\delta)}{2n}} \leq \epsilon + \frac{8\left(\sqrt{2\ln(4n/\delta)} + \ln(4/\epsilon)\right)}{\gamma^2\sqrt{n}} + 3\sqrt{\frac{\ln(2/\delta)}{2n}}.$$

Invoking $P_{(x,y)\sim\mathcal{D}}\left(yf(x;W,a) \leq 0\right) \leq 2\mathcal{Q}(W)$ finishes the proof. $\qquad\square$

C   OMITTED PROOFS FROM SECTION 4

Proof of Lemma 4.2. Recall that $\left\|\nabla f_t(W_t)\right\|_F \leq 1$, we have

$$\left\|W_{t+1} - \overline{W}\right\|_F^2 \leq \left\|W_t - \overline{W}\right\|_F^2 - 2\eta\ell'\left(y_t f_t(W_t)\right) y_t \left\langle \nabla f_t(W_t), W_t - \overline{W}\right\rangle + \eta^2\left(\ell'\left(y_t f_t(W_t)\right)\right)^2.$$
$$(C.1)$$

Similar to the proof of Lemma 2.6, the first order term of eq. (C.1) can be handled using the convexity of ℓ and homogeneity of ReLU as follows

$$\ell'\left(y_t f_t(W_t)\right) y_t \left\langle\nabla f_t(W_t), W_t - \overline{W}\right\rangle \geq \mathcal{R}_t(W_t) - \mathcal{R}_t\left(\overline{W}\right), \qquad (C.2)$$

and the second-order term of eq. (C.1) can be bounded as follows

$$\eta^2\left(\ell'\left(y_t f_t(W_t)\right)\right)^2 \leq -\eta\ell'\left(y_t f_t(W_t)\right) \leq \eta\ell\left(y_t f_t(W_t)\right) = \eta\mathcal{R}_t(W_t), \qquad (C.3)$$

since $\eta, -\ell' \leq 1$ and $-\ell' \leq \ell$. Combining eqs. (C.1) to (C.3) gives

$$\eta\mathcal{R}_t(W_t) \leq \left\|W_t - \overline{W}\right\|_F^2 - \left\|W_{t+1} - \overline{W}\right\|_F^2 + 2\eta\mathcal{R}_t\left(\overline{W}\right).$$

Telescoping gives the claim. $\qquad\square$

With Lemma 4.2, we give the following result, which is an extension of Theorem 2.2 to the SGD setting.

Lemma C.1. *Under Assumption 3.1, given any $\epsilon \in (0,1)$, any $\delta \in (0, 1/3)$, and any positive integer n_0, let*

$$\lambda := \frac{\sqrt{2\ln(4n_0/\delta)} + \ln(4/\epsilon)}{\gamma/4}, \quad \text{and} \quad M := \frac{4096\lambda^2}{\gamma^6}.$$

For any $m \geq M$ and any constant step size $\eta \leq 1$, if $n_0 \geq n := \lceil 2\lambda^2/\eta\epsilon \rceil$, then with probability $1 - 3\delta$,

$$\frac{1}{n}\sum_{i<n}\mathcal{Q}_i(W_i) \leq \epsilon.$$

Proof. We first sample n_0 data examples $(x_0, y_0), \ldots, (x_{n_0-1}, y_{n_0-1})$, and then feed (x_i, y_i) to SGD at step i. We only consider the first n_0 steps.

The proof is similar to the proof of Theorem 2.2. Let n_1 denote the first step before n_0 such that there exists some $1 \leq s \leq m$ with $\left\|w_{s,n_1} - w_{s,0}\right\|_2 > 4\lambda/(\gamma\sqrt{m})$. If such a step does not exist, let $n_1 = n_0$.

Let $\overline{W} := W_0 + \lambda\overline{U}$, in exactly the same way as in Theorem 2.2, we can show that with probability $1 - 3\delta$, for any $0 \leq i < n_1$,

$$y_i\left\langle\nabla f_i(W_i), \overline{W}\right\rangle \geq \ln\left(\frac{4}{\epsilon}\right), \quad \text{and thus} \quad \mathcal{R}_i\left(\overline{W}\right) \leq \epsilon/4.$$

Now consider $n := \lceil 2\lambda^2/\eta\epsilon \rceil$. Using Lemma 4.2, in the same way as the proof of Theorem 2.2 (replacing $\widehat{\mathcal{Q}}(W_\tau)$ with $\mathcal{Q}_i(W_i)$, etc.), we can show that $n \leq n_1$. Then invoking Lemma 4.2 again, we get

$$\frac{1}{n}\sum_{i<n}\mathcal{Q}_i(W_i) \leq \frac{1}{n}\sum_{i<n}\mathcal{R}_i(W_i) \leq \frac{\left\|W_0 - \overline{W}\right\|_F^2}{\eta n} + \frac{2}{n}\sum_{i<n}\mathcal{R}_i\left(\overline{W}\right) \leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

$\square$

Next we prove Lemma 4.3. We need the following martingale Bernstein bound.

Lemma C.2. *(Beygelzimer et al., 2011, Theorem 1) Let $(M_t, \mathcal{F}_t)_{t \geq 0}$ denote a martingale with $M_0 = 0$ and $\mathcal{F}_0$ be the trivial σ-algebra. Let $(\Delta_t)_{t \geq 1}$ denote the corresponding martingale difference sequence, and let*

$$V_t := \sum_{j=1}^t \mathbb{E}\left[\Delta_j^2 \Big| \mathcal{F}_{j-1}\right]$$

denote the sequence of conditional variance. If $\Delta_t \leq R$ a.s., then for any $\delta \in (0, 1)$, with probability at least $1 - \delta$,

$$M_t \leq \frac{V_t}{R}(e - 2) + R\ln\left(\frac{1}{\delta}\right).$$

Proof of Lemma 4.3. For any $i \geq 0$, let z_i denote (x_i, y_i), and $z_{0,i}$ denote $(z_0, \ldots, z_i)$. Note that $\sum_{t<i}\left(\mathcal{Q}(W_t) - \mathcal{Q}_t(W_t)\right)$ is a martingale w.r.t. the filtration $\sigma(z_{0,i-1})$. The martingale difference sequence is given by $\mathcal{Q}(W_t) - \mathcal{Q}_t(W_t)$, which satisfies

$$\mathcal{Q}(W_t) - \mathcal{Q}_t(W_t) = \mathbb{E}_{(x,y)\sim\mathcal{D}}\left[-\ell'\left(yf(x; W_t, a)\right)\right] + \ell'\left(y_t f(x_t; W_t, a)\right) \leq 1, \tag{C.4}$$

since $-1 \leq \ell' \leq 0$. Moreover, we have

$$\begin{aligned}
&\mathbb{E}\left[\left(\mathcal{Q}(W_t) - \mathcal{Q}_t(W_t)\right)^2 \Big| \sigma(z_{0,t-1})\right] \\
&= \mathcal{Q}(W_t)^2 - 2\mathcal{Q}(W_t)\mathbb{E}\left[\mathcal{Q}_t(W_t)\big|\sigma(z_{0,t-1})\right] + \mathbb{E}\left[\mathcal{Q}_t(W_t)^2\big|\sigma(z_{0,t-1})\right] \\
&= -\mathcal{Q}(W_t)^2 + \mathbb{E}\left[\mathcal{Q}_t(W_t)^2\big|\sigma(z_{0,t-1})\right] \\
&\leq \mathbb{E}\left[\mathcal{Q}_t(W_t)^2\big|\sigma(z_{0,t-1})\right] \\
&\leq \mathbb{E}\left[\mathcal{Q}_t(W_t)\big|\sigma(z_{0,t-1})\right] \\
&= \mathcal{Q}(W_t).
\end{aligned} \tag{C.5}$$

Invoking Lemma C.2 with eqs. (C.4) and (C.5) gives that with probability $1 - \delta$,

$$\sum_{t<i}\left(\mathcal{Q}(W_t) - \mathcal{Q}_t(W_t)\right) \leq (e - 2)\sum_{t<i}\mathcal{Q}(W_t) + \ln\left(\frac{1}{\delta}\right).$$

Consequently,

$$\sum_{t<i}\mathcal{Q}(W_t) \leq 4\sum_{t<i}\mathcal{Q}_t(W_t) + 4\ln\left(\frac{1}{\delta}\right).$$

$\square$

Finally, we prove Theorem 4.1.

Proof of Theorem 4.1. Suppose the condition of Lemma C.1 holds. Then we have for $n = \lceil 2\lambda^2/\eta\epsilon \rceil$, with probability $1 - 3\delta$,

$$\frac{1}{n}\sum_{i<n}\mathcal{Q}_i(W_i) \leq \epsilon.$$

Further invoking Lemma 4.3 gives that with probability $1 - 4\delta$,

$$\frac{1}{n}\sum_{i<n}\mathcal{Q}(W_i) \leq \frac{4}{n}\sum_{i<n}\mathcal{Q}_i(W_i) + \frac{4}{n}\ln\left(\frac{1}{\delta}\right) \leq 5\epsilon.$$

Since $P_{(x,y)\sim\mathcal{D}}\left(yf(x;W,a) \leq 0\right) \leq 2\mathcal{Q}(W)$, we get

$$\frac{1}{n}\sum_{i=1}^{n}P_{(x,y)\sim\mathcal{D}}\left(yf(x;W_i,a) \leq 0\right) \leq 10\epsilon.$$

For the condition of Lemma C.1 to hold, it is enough to let

$$n_0 = \Theta\left(\frac{\ln(1/\delta)}{\eta\gamma^2\epsilon^2}\right),$$

which gives

$$M = \Theta\left(\frac{\ln(1/\delta) + \ln(1/\epsilon)^2}{\gamma^8}\right) \quad \text{and} \quad n = \Theta\left(\frac{\ln(1/\delta) + \ln(1/\epsilon)^2}{\gamma^2\epsilon}\right).$$

$\square$

D   OMITTED PROOFS FROM SECTION 5

Proof of Proposition 5.1. Define $f : \mathcal{H} \to \mathbb{R}$ by

$$f(w) := \frac{1}{2}\int\|w(z)\|_2^2\,\mathrm{d}\mu_{\mathcal{N}}(z) = \frac{1}{2}\|w\|_{\mathcal{H}}^2.$$

It holds that f is continuous, and f^* has the same form. Define $g : \mathbb{R}^n \to \mathbb{R}$ by

$$g(p) := \max_{1 \leq i \leq n} p_i,$$

with conjugate

$$g^*(q) = \begin{cases} 0, & \text{if } q \in \Delta_n, \\ +\infty, & \text{o.w.} \end{cases}$$

Finally, define the linear mapping $A : \mathcal{H} \to \mathbb{R}^n$ by $(Aw)_i = y_i\langle w, \phi_i\rangle_{\mathcal{H}}$.

Since f, f^*, g and g^* are lower semi-continuous, and $\mathbf{dom}\,g - A\mathbf{dom}\,f = \mathbb{R}^n$, and $\mathbf{dom}\,f^* - A^*\mathbf{dom}\,g^* = \mathcal{H}$, Theorem 4.4.3 of (Borwein & Zhu, 2005) ensures that

$$\inf_{w\in\mathcal{H}}\left(f(w) + g(Aw)\right) = \sup_{q\in\mathbb{R}^n}\left(-f^*(A^*q) - g^*(-q)\right).$$

with optimal primal-dual solutions $(\bar{w}, \bar{q})$. Moreover

$$\inf_{w\in\mathcal{H}}\left(f(w) + g(Aw)\right) = \inf_{w\in\mathcal{H},u\in\mathbb{R}^n}\sup_{q\in\mathbb{R}^n}\left(f(w) + g(Aw+u) + \langle q, u\rangle\right)$$

$$\geq \sup_{q\in\mathbb{R}^n}\inf_{w\in\mathcal{H},u\in\mathbb{R}^n}\left(f(w) + g(Aw+u) + \langle q, u\rangle\right)$$

$$= \sup_{q\in\mathbb{R}^n}\inf_{w\in\mathcal{H},u\in\mathbb{R}^n}\left(\left(f(w) - \langle A^*q, w\rangle\right)_{\mathcal{H}} + \left(g(Aw+u) - \langle -q, Aw+u\rangle\right)\right)$$

$$= \sup_{q\in\mathbb{R}^n}\left(-f^*(A^*q) - g^*(-q)\right).$$

By strong duality, the inequality holds with equality. It follows that

$$\bar{w} = A^*\bar{q}, \quad \text{and} \quad \mathbf{supp}(-\bar{q}) \subset \arg\max_{1\leq i\leq n}(A\bar{w})_i.$$

Now let us look at the dual optimization problem. It is clear that

$$\sup_{q\in\mathbb{R}^n}\left(-f^*(A^*q) - g^*(-q)\right) = -\inf_{q\in\Delta_n}f^*(A^*q).$$

In addition, we have

$$
\begin{aligned}
f^*(A^*q) &= \frac{1}{2} \int \left\| \sum_{i=1}^n q_i y_i \phi_i(z) \right\|_2^2 \mathrm{d}\mu_{\mathcal{N}}(z) \\
&= \frac{1}{2} \int \sum_{i,j=1}^n q_i q_j y_i y_j \left\langle \phi_i(z), \phi_j(z) \right\rangle \mathrm{d}\mu_{\mathcal{N}}(z) \\
&= \frac{1}{2} \sum_{i,j=1}^n q_i q_j y_i y_j \int \left\langle \phi_i(z), \phi_j(z) \right\rangle \mathrm{d}\mu_{\mathcal{N}}(z) \\
&= \frac{1}{2} \sum_{i,j=1}^n q_i q_j y_i y_j K_1(i,j) = \frac{1}{2}(q \odot y)^\top K_1 (q \odot y),
\end{aligned}
$$

and thus $f^*(A^*\bar{q}) = \gamma_1^2/2$. Since $\bar{w} = A^*\bar{q}$, we have that $\|\bar{w}\|_{\mathcal{H}} = \gamma_1$. In addition,

$$
g(A\bar{w}) = -f^*(A^*\bar{q}) - f(\bar{w}) = -\gamma_1^2,
$$

and thus $-\bar{w}$ has margin γ_1^2. Moreover, we have

$$
\bar{w}(z) = \sum_{i=1}^n \bar{q}_i y_i \phi_i(z) = \sum_{i=1}^n \bar{q}_i y_i x_i \mathbb{1}\left[\langle z, x_i \rangle > 0 \right],
$$

and thus $\|\bar{w}(z)\|_2 \le 1$. Therefore, $\hat{v} = -\bar{w}/\gamma_1$ satisfies all requirements of Proposition 5.1. $\qquad\square$

E   EXAMPLES WHERE γ IS A LARGE, NATURAL QUANTITY

This section calculates γ for two different data distributions, the goal being to highlight that (a) γ can be large, or rather $1/\gamma$ can be polylogarithmic in the other problem parameters, and (b) γ arises naturally in these problems, and indeed can appear in lower bounds.

In more detail, a summary of the analyses here is as follows.

- Appendix E.1 provides a quick verification that, in the linearly separable case, γ (indeed 2γ) is the usual notion of margin.

- Appendix E.2 describes the *noisy 2-XOR* problem introduced in prior work (Wei et al., 2018), and shows that $\gamma = \Omega(1/d)$. For this problem we could have $n = 2^d$, in which case $1/\gamma = \mathrm{polylog}(n)$.

- Appendix E.2.1 then leverages this bound to show that γ is naturally tied to the width needed to separate data with the NTK; specifically, it is shown that if the width is less than $1/\sqrt{\gamma}$, then there exists a subset of the noisy 2-XOR problem with four points which can not be separated by the finite-width NTK with constant probability over the initialization.

- Appendix E.2.2 lastly shows that the $\gamma = \Omega(1/d)$ lower bound can be used to prove a $\widetilde{O}(d^2)$ sample complexity upper bound for the infinite-width NTK. This matches the d^2 sample complexity lower bound for the infinite-width NTK given in (Wei et al., 2018).

E.1   THE LINEARLY SEPARABLE CASE

Suppose there exists a unit vector $\bar{u}$ such that $y\langle \bar{u}, x \rangle \ge \gamma$ almost surely. Then we can define $\bar{v}(z) := \bar{u}$ for any $z \in \mathbb{R}^d$. For almost all (x, y), we have

$$
\begin{aligned}
y \int \left\langle \bar{v}(z), x \right\rangle \mathbb{1}\left[\langle z, x \rangle > 0 \right] \mathrm{d}\mu_{\mathcal{N}}(z) &= \int y \left\langle \bar{u}, x \right\rangle \mathbb{1}\left[\langle z, x \rangle > 0 \right] \mathrm{d}\mu_{\mathcal{N}}(z) \\
&\ge \gamma \int \mathbb{1}\left[\langle z, x \rangle > 0 \right] \mathrm{d}\mu_{\mathcal{N}}(z) \\
&= \frac{\gamma}{2}.
\end{aligned}
$$

E.2 THE NOISY 2-XOR DISTRIBUTION

We consider the noisy 2-XOR distribution introduced in (Wei et al., 2018). It is the uniform distribution over the following 2^d points:

$$(x_1, x_2, y, x_3, \ldots, x_d) \in \left\{ \left(\frac{1}{\sqrt{d-1}}, 0, 1 \right), \left(0, \frac{1}{\sqrt{d-1}}, -1 \right), \left(\frac{-1}{\sqrt{d-1}}, 0, 1 \right), \left(0, \frac{-1}{\sqrt{d-1}}, -1 \right) \right\}$$

$$\times \left\{ \frac{-1}{\sqrt{d-1}}, \frac{1}{\sqrt{d-1}} \right\}^{d-2}.$$

The factor $1/\sqrt{d-1}$ ensures that $\|x\|_2 = 1$, and $\times$ above denotes the Cartesian product. Here the label y only depends on the first two coordinates of the input x.

To construct $\bar{v}$, we first decompose $\mathbb{R}^2$ into four regions:

$$A_1 := \left\{ (z_1, z_2) \big| z_1 \geq 0, |z_1| \geq |z_2| \right\},$$
$$A_2 := \left\{ (z_1, z_2) \big| z_2 > 0, |z_1| < |z_2| \right\},$$
$$A_3 := \left\{ (z_1, z_2) \big| z_1 \leq 0, |z_1| \geq |z_2| \right\} \setminus \{(0,0)\},$$
$$A_4 := \left\{ (z_1, z_2) \big| z_2 < 0, |z_1| < |z_2| \right\}.$$

Then $\bar{v}$ can de defined as follows. It only depends on the first two coordinates of z.

$$\bar{v}(z) := \begin{cases} (1, 0, 0, \ldots, 0) & \text{if } (z_1, z_2) \in A_1, \\ (0, -1, 0, \ldots, 0) & \text{if } (z_1, z_2) \in A_2, \\ (-1, 0, 0, \ldots, 0) & \text{if } (z_1, z_2) \in A_3, \\ (0, 1, 0, \ldots, 0) & \text{if } (z_1, z_2) \in A_4. \end{cases}$$

The following result shows that $\gamma = \Omega(1/d)$. Note that n could be as large as 2^d, in which case γ is basically $\mathrm{polylog}(n)$.

Proposition E.1. *For any (x, y) sampled from the noisy 2-XOR distribution and any $d \geq 3$, it holds that*

$$y \int \langle \bar{v}(z), x \rangle \, \mathbb{1}\left[\langle z, x \rangle > 0 \right] \, \mathrm{d}\mu_{\mathcal{N}}(z) \geq \frac{1}{60d}.$$

Proof. By symmetry, we only need to consider an (x, y) where $(x_1, x_2, y) = (1/\sqrt{d-1}, 0, 1)$. Let $z_{p,q}$ denote $(z_p, z_{p+1}, \ldots, z_q)$, and similarly define $x_{p,q}$. We have

$$y \int \langle \bar{v}(z), x \rangle \, \mathbb{1}\left[\langle z, x \rangle > 0 \right] \, \mathrm{d}\mu_{\mathcal{N}}(z)$$

$$= y \int \left(\int \langle \bar{v}(z), x \rangle \, \mathbb{1}\left[\langle z, x \rangle > 0 \right] \, \mathrm{d}\mu_{\mathcal{N}}(z_{3,d}) \right) \mathrm{d}\mu_{\mathcal{N}}(z_{1,2}) \tag{E.1}$$

$$= y \int \langle \bar{v}(z)_{1,2}, x_{1,2} \rangle \left(\int \mathbb{1}\left[\langle z_{1,2}, x_{1,2} \rangle + \langle z_{3,d}, x_{3,d} \rangle > 0 \right] \, \mathrm{d}\mu_{\mathcal{N}}(z_{3,d}) \right) \mathrm{d}\mu_{\mathcal{N}}(z_{1,2}) \tag{E.2}$$

$$= \sum_{i=1}^{4} y \int \langle \bar{v}(z)_{1,2}, x_{1,2} \rangle \left(\int \mathbb{1}\left[\langle z_{1,2}, x_{1,2} \rangle + \langle z_{3,d}, x_{3,d} \rangle > 0 \right] \, \mathrm{d}\mu_{\mathcal{N}}(z_{3,d}) \right) \mathbb{1}\left[z_{1,2} \in A_i \right] \mathrm{d}\mu_{\mathcal{N}}(z_{1,2}), \tag{E.3}$$

where eq. (E.1) is due to the independence between $z_{1,2}$ and $z_{3,d}$, and in eq. (E.2) we use the fact that $\bar{v}(z)_{1,2}$ only depends on $z_{1,2}$ and $\bar{v}(z)_{3,d}$ are all zero. Since $\langle \bar{v}(z)_{1,2}, x_{1,2} \rangle = 0$ for $z_{1,2} \in A_2 \cup A_4$, we only need to consider A_1 and A_3 in eq. (E.3). For simplicity, we will denote $z_{1,2}$ by $p \in \mathbb{R}^2$, and $\bar{v}(z)_{1,2}$ by $\bar{v}(p)$, and $z_{3,d}$ by $q \in \mathbb{R}^{d-2}$.

For any nonzero $p \in A_1$, we have $-p \in A_3$, and $\langle \bar{v}(p), x_{1,2} \rangle = 1/\sqrt{d-1}$. Therefore

$$
y \langle \bar{v}(p), x_{1,2} \rangle \left(\int \mathbb{1} \left[\langle p, x_{1,2} \rangle + \langle q, x_{3,d} \rangle > 0 \right] \mathrm{d}\mu_{\mathcal{N}}(q) \right)
$$

$$
+ y \langle \bar{v}(-p), x_{1,2} \rangle \left(\int \mathbb{1} \left[\langle -p, x_{1,2} \rangle + \langle q, x_{3,d} \rangle > 0 \right] \mathrm{d}\mu_{\mathcal{N}}(q) \right)
$$

$$
= \frac{1}{\sqrt{d-1}} \int \left(\mathbb{1} \left[\frac{p_1}{\sqrt{d-1}} + \langle q, x_{3,d} \rangle > 0 \right] - \mathbb{1} \left[\frac{-p_1}{\sqrt{d-1}} + \langle q, x_{3,d} \rangle > 0 \right] \right) \mathrm{d}\mu_{\mathcal{N}}(q)
$$

$$
= \frac{1}{\sqrt{d-1}} \mathbb{P} \left(\frac{-p_1}{\sqrt{d-1}} \leq \langle q, x_{3,d} \rangle \leq \frac{p_1}{\sqrt{d-1}} \right). \tag{E.4}
$$

Let φ denote the density function of the standard Gaussian distribution, and for $c > 0$, let $U(c)$ denote the probability that a standard Gaussian random variable lies in the interval $[-c, c]$:

$$
U(c) := \int_{-c}^{c} \varphi(t) \, \mathrm{d}t.
$$

Since $\langle q, x_{3,d} \rangle$ is a Gaussian variable with standard deviation $\sqrt{(d-2)/(d-1)}$, we have

$$
\mathbb{P} \left(\frac{-p_1}{\sqrt{d-1}} \leq \langle q, x_{3,d} \rangle \leq \frac{p_1}{\sqrt{d-1}} \right) = U \left(\frac{p_1}{\sqrt{d-2}} \right). \tag{E.5}
$$

Plugging eqs. (E.4) and (E.5) into eq. (E.3) gives:

$$
y \int \langle \bar{v}(z), x \rangle \mathbb{1} \left[\langle z, x \rangle > 0 \right] \mathrm{d}\mu_{\mathcal{N}}(z) = \frac{1}{\sqrt{d-1}} \int U \left(\frac{p_1}{\sqrt{d-2}} \right) \mathbb{1} \left[p \in A_1 \right] \mathrm{d}\mu_{\mathcal{N}}(p)
$$

$$
= \frac{1}{\sqrt{d-1}} \int_0^{\infty} U \left(\frac{p_1}{\sqrt{d-2}} \right) \left(\int_{-p_1}^{p_1} \varphi(p_2) \, \mathrm{d}p_2 \right) \varphi(p_1) \, \mathrm{d}p_1
$$

$$
= \frac{1}{\sqrt{d-1}} \int_0^{\infty} U \left(\frac{p_1}{\sqrt{d-2}} \right) U(p_1) \varphi(p_1) \, \mathrm{d}p_1
$$

$$
\geq \frac{1}{\sqrt{d-1}} \int_0^1 U \left(\frac{p_1}{\sqrt{d-2}} \right) U(p_1) \varphi(p_1) \, \mathrm{d}p_1.
$$

For $t \in [-1, +1]$, it holds that $\varphi(t) \geq 1\sqrt{2\pi e}$, and thus

$$
U(a) = \int_{-a}^{a} \varphi(t) \, \mathrm{d}t \geq \frac{2a}{\sqrt{2\pi e}}.
$$

Therefore eq. (E.3) is lower bounded by

$$
\frac{1}{\sqrt{d-1}} \int_0^1 U \left(\frac{p_1}{\sqrt{d-2}} \right) U(p_1) \varphi(p_1) \, \mathrm{d}p_1 \geq \frac{1}{\sqrt{d-1}} \int_0^1 \frac{2}{\sqrt{2\pi e}} \cdot \frac{p_1}{\sqrt{d-2}} \cdot \frac{2p_1}{\sqrt{2\pi e}} \cdot \frac{1}{\sqrt{2\pi e}} \, \mathrm{d}p_1
$$

$$
\geq \frac{1}{20\sqrt{(d-1)(d-2)}} \int_0^1 p_1^2 \, \mathrm{d}p_1
$$

$$
= \frac{1}{60\sqrt{(d-1)(d-2)}}
$$

$$
\geq \frac{1}{60d}.
$$

$\square$

E.2.1 A LOWER BOUND ON THE WIDTH REQUIRED BY THE NTK

For the noisy 2-XOR distribution, the separator considered in Proposition E.1 has margin $\gamma = \Omega(1/d)$, or equivalently $1/\gamma = O(d)$. Below we further show that for any four points which have the same last $d - 2$ coordinates, if the width is no larger than $\sqrt{d-2}/4$, then with probability $1/2$

the finite-width NTK cannot separate them. In other words, in the NTK regime, the required width must at least depend on $1/\sqrt{\gamma}$.

The NTK classifier is given by

$$f_{\mathrm{NTK}}(x; V, W) := \sum_{s=1}^{m} \langle v_s, x_i \rangle \mathbb{1}\left[\langle w_s, x_i \rangle > 0\right].$$

Here $v_1, \ldots, v_m \in \mathbb{R}^d$ are the weights, and $w_1, \ldots, w_m \in \mathbb{R}^d$ are standard Gaussian random vectors which give the NTK features. They are collected into $V \in \mathbb{R}^{m \times d}$ and $W \in \mathbb{R}^{m \times d}$ respectively. We discuss the $0/1$ loss here and thus omit the $1/\sqrt{m}$ factor.

We consider the training set D consisting of the following four points:

$$\begin{aligned}
x_1 &= (1, 0, 1, \ldots, 1), & y_1 &= 1, \\
x_2 &= (0, 1, 1, \ldots, 1), & y_2 &= -1, \\
x_3 &= (-1, 0, 1, \ldots, 1), & y_3 &= 1, \\
x_4 &= (0, -1, 1, \ldots, 1), & y_4 &= -1.
\end{aligned}$$

The $1/\sqrt{d-1}$ factor is omitted also because we only discuss the $0/1$ loss.

Proposition E.2. *Given training set D, for any $d \geq 20$, if $m \leq \sqrt{d-2}/4$, then with probability $1/2$ over the random initialization of W, for any weights V, it holds that $y_i f_{\mathrm{NTK}}(x_i) < 0$ for at least one $i \in \{1, 2, 3, 4\}$.*

To prove Proposition E.2, we need the following technical lemma.

Lemma E.3. *Given $z_1 \sim \mathcal{N}(0, 1)$ and $z_2 \sim \mathcal{N}(0, b^2)$ that are independent where $b > 1$, we have*

$$\mathbb{P}\left(|z_1| < |z_2|\right) > 1 - \frac{1}{b}.$$

Proof. First note that for $z_3 \sim \mathcal{N}(0, 1)$ which is independent of z_1,

$$\mathbb{P}\left(|z_1| < |z_2|\right) = \mathbb{P}\left(|z_1| < b|z_3|\right) = 1 - \mathbb{P}\left(|z_3| < \frac{1}{b}|z_1|\right).$$

Still let φ denote the density of $\mathcal{N}(0, 1)$, and let $U(c)$ denote the probability that $z_3 \in [-c, c]$. We have

$$\begin{aligned}
\mathbb{P}\left(|z_3| < \frac{1}{b}|z_1|\right) &= \int \int \mathbb{1}\left[|z_3| < \frac{1}{b}|z_1|\right] \varphi(z_3) \varphi(z_1) \, \mathrm{d}z_3 \, \mathrm{d}z_1 \\
&= \int U\left(\frac{1}{b}|z_1|\right) \varphi(z_1) \, \mathrm{d}z_1 \\
&\leq \frac{2}{\sqrt{2\pi}b} \int |z_1| \varphi(z_1) \, \mathrm{d}z_1 = \frac{2}{\pi b} < \frac{1}{b},
\end{aligned}$$

where we use the facts that $U(c) \leq 2c/\sqrt{2\pi}$ and $\mathbb{E}[|z_1|] = \sqrt{2/\pi}$. $\qquad \square$

We now give the proof of Proposition E.2 using Lemma E.3.

Proof of Proposition E.2. For any s, let A_s denote the event that

$$\mathbb{1}\left[\langle w_s, x_1 \rangle > 0\right] = \mathbb{1}\left[\langle w_s, x_2 \rangle > 0\right] = \mathbb{1}\left[\langle w_s, x_3 \rangle > 0\right] = \mathbb{1}\left[\langle w_s, x_4 \rangle > 0\right].$$

We will show that if $m \leq \sqrt{d-2}/4$, then A_s is true for all $1 \leq s \leq m$ with probability $1/2$, and Proposition E.2 follows from the fact that the XOR data is not linearly separable.

For any s and i,

$$\langle w_s, x_i \rangle = w_{s,1} x_{i,1} + w_{s,2} x_{i,2} + \sum_{j=3}^{d} w_{s,j}.$$

Since $(x_{i,1}, x_{i,2})$ is $(1,0)$ or $(0,1)$ or $(-1,0)$ or $(0,-1)$, event A_s will happen as long as

$$|w_{s,1}| < \left| \sum_{j=3}^{d} w_{s,j} \right|, \quad \text{and} \quad |w_{s,2}| < \left| \sum_{s=3}^{d} w_{s,j} \right|.$$

Note that $w_{s,1}, w_{s,2} \sim \mathcal{N}(0,1)$ while $\sum_{j=3}^{d} w_{s,j} \sim \mathcal{N}(0, d-2)$. As a result, due to Lemma E.3,

$$\mathbb{P}\left(|w_{s,1}| < \left| \sum_{j=3}^{d} w_{s,j} \right| \right) = \mathbb{P}\left(|w_{s,2}| < \left| \sum_{s=3}^{d} w_{s,j} \right| \right) > 1 - \frac{1}{\sqrt{d-2}}.$$

Using a union bound, $\mathbb{P}(A_s) > 1 - 2/\sqrt{d-2}$. If $m \leq \sqrt{d-2}/4$, then by a union bound again,

$$\mathbb{P}\left(\bigcup_{1 \leq s \leq m} A_s \right) > 1 - \frac{2}{\sqrt{d-2}} m \geq 1 - \frac{2}{\sqrt{d-2}} \frac{\sqrt{d-2}}{4} = \frac{1}{2}.$$

$\square$

E.2.2 A TIGHT SAMPLE COMPLEXITY FOR THE INFINITE-WIDTH NTK

(Wei et al., 2018) give a d^2 sample complexity lower bound for any NTK classifier on the noisy 2-XOR distribution. It turns out that γ could give a *matching* sample complexity upper bound for the NTK and SGD.

(Wei et al., 2018) consider the infinite-width NTK with respect to both layers. For the first layer, the infinite-width NTK K_1 is defined in Section 5, and the corresponding RKHS $\mathcal{H}$ and RKHS mapping ϕ is defined in Section 2. For the second layer, the infinite width NTK is defined by

$$K_2(x_i, x_j) := \mathbb{E}\left[\frac{\partial f(x_i; W_0, a)}{\partial a}, \frac{\partial f(x_j; W_0, a)}{\partial a} \right]$$
$$= \mathbb{E}_{w \sim \mathcal{N}(0, I_d)}\left[\sigma\left(\langle w, x_i \rangle \right) \sigma\left(\langle w, x_j \rangle \right) \right].$$

The corresponding RKHS $\mathcal{K}$ is defined by

$$\mathcal{K} := \left\{ w : \mathbb{R}^d \to \mathbb{R} \,\middle|\, \int w(z)^2 \, \mathrm{d}\mu_{\mathcal{N}}(z) < \infty \right\},$$

and the inner product $\langle w_1, w_2 \rangle_{\mathcal{K}}$ is given by

$$\langle w_1, w_2 \rangle_{\mathcal{K}} = \int w_1(z) w_2(z) \, \mathrm{d}\mu_{\mathcal{N}}(z).$$

Given any $x \in \mathbb{R}^d$, it is mapped into $\psi_x \in \mathcal{K}$:

$$\psi_x(z) := \sigma\left(\langle z, x \rangle \right).$$

It holds that $K_2(x_i, x_j) = \langle \psi_{x_i}, \psi_{x_j} \rangle_{\mathcal{K}}$.

The infinite-width NTK with respect to both layers is just $K_1 + K_2$. The corresponding RHKS is just $\mathcal{H} \times \mathcal{K}$ with the inner product

$$\langle (v_1, w_1), (v_2, w_2) \rangle_{\mathcal{H} \times \mathcal{K}} = \langle v_1, v_2 \rangle_{\mathcal{H}} + \langle w_1, w_2 \rangle_{\mathcal{K}}.$$

The classifier $\bar{v}$ considered in Proposition E.1 has a unit norm (i.e., $\|\bar{v}\|_{\mathcal{H}} = 1$) and margin γ on the space $\mathcal{H}$. On $\mathcal{H} \times \mathcal{K}$, it is enough to consider $(\bar{v}, 0)$, which also has a unit norm and margin γ. Since the infinite-width NTK model is a linear model in $\mathcal{H} \times \mathcal{K}$, and there exists $(\bar{v}, 0)$ with margin γ, (Ji & Telgarsky, 2018, Lemma 2.5) can be used to show that SGD on the RKHS $\mathcal{H} \times \mathcal{K}$ could obtain a test error of ϵ with a sample complexity of $\widetilde{O}(1/\gamma^2 \epsilon)$. The analysis in (Ji & Telgarsky, 2018) is done in $\mathbb{R}^d$, but it still works with a well-defined inner product, and thus can be extended to $\mathcal{H} \times \mathcal{K}$ directly.

Since the margin γ given by $(\bar{v}, 0)$ is $\Omega(1/d)$, to achieve a constant test accuracy, the sample complexity upper bound becomes $\widetilde{O}(d^2)$. This upper bound is actually *tight* (up to logarithmic factors), since Wei et al. (2018) prove a sample complexity lower bound of d^2.