Out-of-sample extension of spectral embeddings: An optimization perspective

Chester Holtz[*]
University of California, San Diego
chholtz@ucsd.edu

Taanish Reja[*]
University of California, San Diego
treja@ucsd.edu

Gal Mishne[*]
University of California, San Diego
gmishne@ucsd.edu

Abstract

Graph-based manifold learning constructs / reveals low dimensional embeddings of high-dimensional data, however requires out-of-sample-extension methods to embed new data points. We propose a new framework, *ROSE* (Riemannian Out-of-Sample Extension), for out-of-sample extensions for spectral graph-based embedding algorithms. *ROSE* is motivated from an optimization perspective of the underlying eigenvector problem associated with classic manifold learning problems. Similar to graph-based semi-supervised learning, our approach exploits the geometry of the unlabeled samples in addition to the labeled samples, by treating the in-sample embedddings as *labeled* data. *ROSE* Despite its non-convexity, *ROSE* is solvable by first-order methods, which converge to global minimizers under certain assumptions.

1 Introduction

Given a graph $G = (V, E)$ on n vertices with adjacency matrix $W \in \mathbb{R}^{n \times n}$, the goal of graph embedding is to map the vertices of G to some d-dimensional vector space in such a way that geometry of the embedding preserves the geometry of G. For example, we may ask that vertices with high connectivity in G be assigned to nearby vectors in the embedding space.

Various approaches have been proposed to compute effective graph embeddings. A classic method that is particularly relevant is the manifold learning approach Laplacian Eigenmaps [1, 2], which utilizes the graph Laplacian matrix to produce low-dimensional representations of graph vertices.

One challenge of these approaches is that they do not provide an explicit mapping between the vertices and the low-dimensional embedding. Therefore, when new data is introduced or when large-scale datasets prevent an embedding of the full dataset due to computational limitations, out-of-sample extension (OOSE) methods are used to extend the embeddings from the training samples to the rest of the data or to new unseen data [3]. For instance, in the case of Laplacian Eigenmaps, out-of-sample extension can be performed using a Nyström approximation. The Nyström method [4] allows for the efficient approximation of the eigenfunctions of the graph Laplacian on the new data points by expressing them as a linear combination of the eigenfunctions computed from the original dataset [4–7]. Beyond these classical methods, more recent approaches the family of spectral and diffusion networks [3, 8, 9] have been proposed to handle out-of-sample extension in graph embeddings.

In this paper, we propose a new approach to graph-based out-of-sample extensions that builds upon these traditional methods but introduces a novel optimization perspective. Our method focuses on directly learning the mapping between the data and the embedding space through an optimization framework. This approach allows for a more principled and systematic extension of embeddings

---

[*]Equal contribution.

Y. Zhu et al., Out-of-sample extension of spectral embeddings: An optimization perspective (Extended Abstract). Presented at the Second Learning on Graphs Conference (LoG 2023), Virtual Event, November 27–30, 2023.

to new data points while preserving the topological and geometric properties of the original graph embedding. We provide one interpretation of out-of-sample extensions of spectral algorithms as a quadratic optimization problem over a smooth manifold with special quadratic constraints, for which one can implement gradient or conjugate gradient methods and Newton methods over geodesic paths on the manifold [10, 11].

2 Quadratic Minimization for Out-of-Sample-Extension

Given a training dataset $D = \{x_1, \ldots, x_n\}$ with $x_i \in \mathbb{R}^d$, D gives rise to a symmetric matrix $W \in \mathbb{R}^{n \times n}$ derived from a kernel $K : \mathcal{X} \times \mathcal{X} \to \mathbb{R}$, i.e. $w_{ij} = K(x_i, x_j) \geq 0$, that measures the similarity between pairs of datapoints x_i and x_j such that $K(x_i, x_j)$ is large if x_i and x_j are similar and small otherwise. This matrix defines a graph where nodes are data points, and edges represent pairwise similarities, with its combinatorial Laplacian $L = \text{diag}(1^\top W) - W$ capturing the local geometric structure of the data. The eigenvectors of this matrix are used to embed the data in a lower-dimensional space, preserving local neighborhood relationships.

For out-of-sample extensions, where new points need to be embedded without recomputing the entire embedding, we introduce an efficient framework derived from an optimization perspective of the underlying spectral decomposition.

Let $St(n, r)$ denote Stiefel manifold defined as

$$St(n, r) = \{X \in \mathbb{R}^{n \times r} : X^\top X = I_r\}$$

where I_r denotes the $r \times r$ identity matrix. In short, $St(n, r)$ is the set of matrices in $\mathbb{R}^{n \times r}$ whose columns are orthonormal in $\mathbb{R}^n$ with respect to the inner product $\langle x, y \rangle = tr(x^\top y)$.

In this paper, we propose one efficient algorithm to solve out-of-sample-ex tension for manifold learning problems. Our algorithm is based on a reduction to a quadratic problem of the form

$$\min_{X \in St(n,r)} \{F(X) = \langle X, AXC \rangle - \langle B, X \rangle\}, \tag{1}$$

where $A \in \mathbb{R}^{n \times n}$ is symmetric and $C \in \mathbb{R}^{r \times r}$ is positive definite and the the inner product $\langle R, T \rangle$ is the trace of the matrix $R^\top T$ for R, T of the same size.

Introduce the graph $G = (V, E, W)$ with m' vertices, where $V = \{v_1, v_2, \ldots, v_{m'}\}$ is the vertex set, E is the edge set and W is the weight matrix whose entries $w_{ij} \geq 0$ are the edge weights between v_i and v_j. Assume that the graph is symmetric, i.e., $w_{ij} = w_{ji}$. Define the degree matrix $D = \text{diag}([d_1, \ldots, d_{m'}])$ $d_i = \sum_{j=1}^{m'} w_{ij}$ for each $v_i \in V$. Let L denote the combinatorial graph Laplacian, $L = D - W$. An embedding of the vertices is given by the eigenvectors X corresponding to the smallest nontrivial eigenvalues,

$$\min_{X \in \mathbb{R}^{m' \times r}} \langle X, LX \rangle, \ 1_n^\top X = 0, \ X^\top X = I_r. \tag{2}$$

Consider the semi-supervised learning task, where a set of pre-specified labeled data is available: The first m vertices $V_l := \{v_1, v_2, \ldots, v_m\}$ are assigned labels $\{y_1, y_2, \ldots, y_m\}$, where $0 < m \ll m'$. Let n denote the number of unlabeled vertices, $n = m' - m$. Our task corresponds to smoothly propagating the observed values over the unlabeled vertices $V_u := \{v_{m+1}, v_{m+2}, \ldots, v_{m'}\}$. Let

$$L = \begin{bmatrix} L_{l,l} & L_{l,u} \\ L_{u,l} & L_{u,u} \end{bmatrix}, \ Y = \begin{bmatrix} Y_l \\ Y_u \end{bmatrix}, Y_l = [y_1, \ldots, y_m]^\top, \ X = \begin{bmatrix} X_l \\ X_u \end{bmatrix}. \tag{3}$$

where subscripts l and u correspond to labeled and unlabeled indices, respectively. Let X_l represent the **labeled vertices** with cardinality m, i.e., $X_l = Y_l$. Introduce the associated constraint set

$$\mathcal{X} := \{X \in \mathbb{R}^{n \times r} : 1_n^\top X = 0, X^\top X = X_l^\top X_l + X_u^\top X_u = pI\}, \tag{4}$$

for some scalar $p = n/r$. The following proposition indicates that the unknown matrix X_u can be computed from one quadratic minimization problem over a Stiefel manifold, i.e., (8).

Proposition 2.1. *Let p be a positive scalar. Given X, L in (3). Given the labeled matrix $X_l \in \mathbb{R}^{m \times r}$ and $c \in \mathbb{R}^r$, Consider the minimization*

$$\min_{X_u \in \mathbb{R}^{n \times r}} \{\langle X, LX \rangle : X \in \mathcal{X}\}. \tag{5}$$

Let c_l be the column sum of X_l, i.e. $= X_l^\top 1_m$ and $P = I_n - \frac{1}{n}\mathbf{1}\mathbf{1}^\top$ and

$$A = PL_{u,u}P, \ \ B = -P(n^{-1}L_{u,u}1_n c_l^\top + L_{u,l}X_l), \ \ C = pI - X_l^\top X_l - \frac{1}{n}c_l c_l^\top. \tag{6}$$

Then, $X_u = XC^{1/2} + \frac{1}{n}1_n c_l^\top$, where X is the minimizer of

$$\min_{X\in\mathbb{R}^{n\times k}} \left\{ \langle X, AXC \rangle - 2\langle X, BC^{1/2}\rangle : X \in St(n,k) \right\} \tag{7}$$

For brevity, we will recast $B = BC^{1/2}$. To reiterate, given a solution to (7), X^*, one recovers a solution to (5) via the transformation

$$X^* C^{1/2} + \frac{1}{n}\mathbf{1}r^\top \tag{8}$$

2.1 Optimality conditions

Optimization over the Stiefel manifold is a nonconvex problem. Generally, it is not possible to recover the global minimum. However, we show that in certain special cases recover of high-quality critical points is likely for first-order methods under an appropriate initialization. First, we define critical points to be those points that satisfy the following *first-order condition*

$$AXC = B + X\Lambda \tag{9}$$

for some $\Lambda \in \mathbb{R}^{n\times r}$. Points satisfying this condition can be local maximizers, minimizers, or saddle points. In general, there can be many critical points satisfying this condition. The following illustrates that a critical point X satisfying a certain second-order condition is a global minimizer of (1).

Proposition 2.2. *Let d_1 be the smallest eigenvalue of A and X' be a critical point of*

$$\min_X F(X) \quad s.t. \ X^\top X = I \tag{10}$$

and let Λ' be the associated multipliers matrix. Suppose

$$d_1 C \succcurlyeq \Lambda'. \tag{11}$$

Then X' is a global minimizer. Suppose $d_1 C \succ \Lambda'$. Then, X' is the unique global minimizer.

Note that in general, the condition in (11) could be too strict to be fulfilled for any critical points. We introduce the following proposition to describe a less restrictive spectral condition that implies global optimality for certain special quadratics. The following non-degeneracy condition on B ensures that any critical point X satisfying a certain condition is a global minimizer. What is needed is that the projection of B on V is sufficiently large, compared with the spectral gap $d_r - d_1$ for some d_r such that $\Lambda \succcurlyeq d_r C$.

Proposition 2.3. *Let $V = [v_1, v_2, \ldots, v_r] \in \mathbb{R}^{n\times r}$ be the eigenvectors of L corresponding to the smallest r nonzero eigenvalues $d_1 \le d_2 \le \ldots \le d_r$. Let X be a local solution satisfying the first order condition*

$$AXC = B + X\Lambda$$

and second order condition $\lambda_1, \ldots, \lambda_r \le d_r$. Let s_1 be the smallest singular value of $V^\top BC^{-1}$. Suppose

$$d_r - \gamma_j \ge \sigma \text{ for all } j = 1, \ldots, r \tag{12}$$

and

$$\sigma > d_r - d_1 \tag{13}$$

Then, all eigenvalues $\gamma_1, \ldots, \gamma_r$ of the matrix ΛC^{-1} are less than d_1 and X is a global minimizer.

3 Riemannian Out-of-Sample Extension (ROSE)

In this section, we describe our algorithm, termed ROSE, for semi-supervised out-Of-Sample extension. A standard algorithm for minimizing a smooth objective over the Stiefel Manifold, e.g. (1), is given in "An introduction to optimization on smooth manifolds" by Nicolas Boumal. For each $U \in \mathbb{R}^{n\times k}$, define the projection $\mathcal{P}$ onto the tangent space at $X \in St(n,k)$,

$$\mathcal{P}_X(U) = X\,skew(X^\top U) = X - X\,sym(X^\top U) \tag{14}$$

Where $sym(Z) = \frac{1}{2}(Z + Z^\top)$. Let $F(X; A, B, C) = \langle X, AX \rangle / 2 - \langle X, B \rangle$ be the objective of (1). Then, the Euclidean gradient is given by $\nabla F = AXC - B$ and the projected gradient is given by

$$grad F(X) = \mathcal{P}_X \nabla F(X) = \mathcal{P}(AXC - B) = (AXC - B) - X sym(\Lambda). \tag{15}$$

Where $\Lambda X^\top (AXC - B)$. The Riemannian Gradient method computes a sequence of iterates $X_0, X_1, \ldots, X_k$ where

$$X_{k+1} = \mathcal{R}_{X_k}(-\alpha_k grad F(X_k)) \tag{16}$$

Where $\mathcal{R}_{X_k}(g_k) = UV^\top$ for $U\Sigma V^\top = SVD(X_k - g_k)$.

3.1 Initialization of Riemannian Gradient

Convergence of first-order methods typically relies on initialization. We justify a computationally friendly initialization that relies on the approximation of an "ideal" initialization given below.

Proposition 3.1. *Let V be an isometric matrix, $V \in St(l, r), n > l \geq r$. Let $\mathcal{S}$ be an induced subspace, $X = \{V\widetilde{X} : \widetilde{X} \in St(l, r)\}$. Consider the subspace-restricted regularized problem,*

$$\min_X \{F(X; A, B, C) : X = V\widetilde{X} \in \mathcal{S}, X \in St(n, r)\}$$
$$= \min_{\widetilde{X}} \{F(\widetilde{X}; V^\top AV, V^\top B, C) : \widetilde{X} \in St(n, r)\} \tag{17}$$

Consider $V = V_g$, i.e., the subspace-restricted problem

$$\min_{X \in St(n,r)} \{F(X; A, V_g V_g^\top B, C)\} \tag{18}$$

Then the associated multiplier $\Lambda = X^\top (AXC - V_g V_g^\top B)$ satisfies $\Lambda \preccurlyeq d_r C$

Proof. Let $X = V_g Q$, where Q is one orthogonal matrix, which maximizes $\langle Q, V_g^\top B \rangle$. Note that

$$\Lambda = X^\top (AXC - B) = d_r C - Q^\top V_g^\top B \tag{19}$$

Is symmetric. Hence, Λ can be expressed as the difference between two Hermitian matrices, and by Weyl's inequality, the proof is complete. $\square$

Since computing V_g is prohibitively expensive, in practice we instead consider an estimate of $V_g Q$, $X_0 = PD_u^{-1} W_{ul} Y$.

4 Preliminary Experiments

The experiment in Figures 1 and 2 show the average-percentage of neighborhood overlap between the $k$nn graphs derived from data in the ambient space and the $k$nn graphs derived in the embedding spaces. Each line graph is generated from averages over 10 trials, where in each trial different in-sample sets were chosen uniformly. This experiment demonstrates ROSE's ability to preserve local neighborhoods in relative to a full eigenvector decomposition.

Figure 1: Neighborhood overlap at 10% in-sampled data for the noisy swiss roll dataset, MNIST, Fashion MNIST, and CIFAR-10.

# in-sample	Eigenvectors	ROSE[1]	ROSE[2]	Nyström
1%	2.24 ± 19.36	2.06 ± 3.46	3.69 ± 5.49	9.86 ± 19.13
5%	2.18 ± 16.16	2.47 ± 9.33	5.47 ± 12.66	8.37 ± 26.65
10%	2.28 ± 18.15	2.66 ± 13.64	5.01 ± 12.73	6.76 ± 25.29
25%	2.29 ± 15.68	2.90 ± 15.44	3.94 ± 9.10	4.45 ± 12.25

Table 1: Comparison of Distortion produced by various embedding methods, where $\text{Distortion}(\Phi_k, \mathcal{U}_k) = ||\Phi_k||_{Lip} \, ||\Phi_k^{-1}||_{Lip}$

References

[1] Mikhail Belkin and Partha Niyogi. Laplacian eigenmaps for dimensionality reduction and data representation. *Neural Computation*, 15(6):1373–1396, 2003. doi: 10.1162/089976603321780317. 1

[2] Mikhail Belkin, Partha Niyogi, and Vikas Sindhwani. Manifold regularization: A geometric framework for learning from labeled and unlabeled examples. *Journal of Machine Learning Research*, 7(85):2399–2434, 2006. URL http://jmlr.org/papers/v7/belkin06a.html. 1

[3] Gal Mishne, Uri Shaham, Alexander Cloninger, and Israel Cohen. Diffusion nets. *Applied and Computational Harmonic Analysis*, 47(2):259–285, 2019. ISSN 1063-5203. doi: https://doi.org/10.1016/j.acha.2017.08.007. URL https://www.sciencedirect.com/science/article/pii/S1063520317300957. 1

[4] Christopher Williams and Matthias Seeger. Using the nyström method to speed up kernel machines. In T. Leen, T. Dietterich, and V. Tresp, editors, *Advances in Neural Information Processing Systems*, volume 13. MIT Press, 2000. URL https://proceedings.neurips.cc/paper_files/paper/2000/file/19de10adbaa1b2ee13f77f679fa1483a-Paper.pdf. 1

[5] Yoshua Bengio, Jean-françcois Paiement, Pascal Vincent, Olivier Delalleau, Nicolas Roux, and Marie Ouimet. Out-of-sample extensions for lle, isomap, mds, eigenmaps, and spectral clustering. In S. Thrun, L. Saul, and B. Schölkopf, editors, *Advances in Neural Information Processing Systems*, volume 16. MIT Press, 2003. URL https://proceedings.neurips.cc/paper_files/paper/2003/file/cf05968255451bdefe3c5bc64d550517-Paper.pdf.

[6] Petros Drineas and Michael W. Mahoney. On the nystrom method for approximating a gram matrix for improved kernel-based learning. *Journal of Machine Learning Research*, 6(72):2153–2175, 2005. URL http://jmlr.org/papers/v6/drineas05a.html.

[7] C. Fowlkes, S. Belongie, F. Chung, and J. Malik. Spectral grouping using the nystrom method. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 26(2):214–225, 2004. doi: 10.1109/TPAMI.2004.1262185. 1

[8] Uri Shaham, Kelly Stanton, Henry Li, Ronen Basri, Boaz Nadler, and Yuval Kluger. Spectralnet: Spectral clustering using deep neural networks. In *International Conference on Learning Representations*, 2018. URL https://openreview.net/forum?id=HJ_aoCyRZ. 1

[9] Ziyu Chen, Yingzhou Li, and Xiuyuan Cheng. Specnet2: Orthogonalization-free spectral embedding by neural networks. In Bin Dong, Qianxiao Li, Lei Wang, and Zhi-Qin John Xu, editors, *Proceedings of Mathematical and Scientific Machine Learning*, volume 190 of *Proceedings of Machine Learning Research*, pages 33–48. PMLR, 15–17 Aug 2022. URL https://proceedings.mlr.press/v190/chen22a.html. 1

[10] Nicolas Boumal. *An Introduction to Optimization on Smooth Manifolds*. Cambridge University Press, 2023. 2

[11] P.-A. Absil, R. Mahony, and R. Sepulchre. *Optimization Algorithms on Matrix Manifolds*. Princeton University Press, Princeton, NJ, 2008. ISBN 978-0-691-13298-3. 2