Localization, Uniform Convexity, and Star Aggregation

Abstract

Offset Rademacher complexities have been shown to imply sharp, data-dependent upper bounds for the square loss in a broad class of problems including improper statistical learning and online learning. We show that the offset complexity can be generalized to any loss satisfying a certain uniform convexity condition. Surprisingly, this condition is shown to capture exponential concavity and self-concordance, uniting several apparently disparate results. By a unified argument, these bounds translate to improper learning in a non-convex class. Thus, we provide a sharp analytic tool that covers both convex empirical risk minimization and improper learning under general entropy conditions. As applications, we recover the optimal rates for proper and improper learning with the p-loss for $1 < p < \infty$, and show that improper variants of empirical risk minimization can attain fast rates for logistic regression and other generalized linear models.

1 Introduction

A central question in statistical learning theory asks which properties of the empirical risk minimizer can be inferred without regard to the underlying data distribution. In particular, the convergence rate of empirical risk minimization has been shown to depend both on distribution-specific properties, such as Tsybakov's margin condition [Tsybakov, 2004], as well as distribution-independent or "global" properties, such as curvature of the loss function [Mendelson, 2002].

The *local Rademacher complexities* of Bartlett et al. [2005] address this issue by giving bounds that are data dependent and also adapt to the global properties of the learning problem. While extremely powerful, significant work is required to apply the method on a case-by-case basis. More recently, the *offset Rademacher complexities* of Rakhlin and Sridharan [2014] and Liang et al. [2015] have been shown to provide similar bounds for the square loss in several contexts, including improper statistical learning and online learning. The method also comes with a simplified analysis that transparently relates the *geometry* of the square loss to *statistical* properties of the estimator.

We extend offset Rademacher complexities to systematically study how curvature of the loss translates to fast rates of convergence in proper and improper statistical learning. We show that classical results on *uniform convexity* of the loss [Bartlett et al., 2006, Mendelson, 2002], as well as newer results on *exponential concavity* [van Erven et al., 2015, Mehta, 2017] and *self-concordance* [Bilodeau et al., 2020, Marteau-Ferey et al., 2019], can be studied in a common geometric framework.

Remarkably, our analysis applies simultaneously to convex empirical risk minimization and to improper learning using the star algorithm of Audibert [2008]. In each case, the excess risk is controlled by a rate-adaptive offset Rademacher complexity. Thus, we give a clean and unified treatment of fast rates in proper and improper learning with curved loss. Our contributions may be summarized as follows.

1. We generalize the offset Rademacher complexities of Liang et al. [2015], showing that when the loss is (μ, d)-convex (defined below), excess risk is bounded by a corresponding offset Rademacher process.

2. We show in this general setting that the "star algorithm" of Audibert [2008] satisfies an upper bound matching the convex case, allowing us to simultaneously treat convex risk minimization and improper learning under general entropy conditions.

3. We show that (μ, d)-convexity captures the behavior of losses that are exponentially concave, uniformly convex, and self concordant, leading to fast upper bounds for learning under the p-loss, the relative entropy, and the logistic loss. Specializing to logistic regression, we have the following bound.

Proposition 1. *Let $\mathcal{F}$ be the class of k-tuples of linear predictors $\mathbb{R}^d \to \mathbb{R}$, where each predictor has norm bounded by B. Then, the regularized star estimator $\hat{g}$ satisfies the excess logistic risk bound*

$$\mathbb{P}\left(\mathcal{E}(\hat{g}) > \frac{kd \cdot \mathrm{polylog}(n, k, B, 1/\rho)}{n} \right) \leq \rho.$$

Thus, in the i.i.d. setting, fast rates that depend only logarithmically on B are achievable by simple, improper variants of empirical risk minimization, particularly the star algorithm.

1.1 Relation to Prior Work

The primary concern of this paper is the *localization* or *fast-rate* phenomenon, where the excess loss vanishes at a rate exceeding the model's uniform law of large numbers [Koltchinskii and Panchenko, 2000, Tsybakov, 2004]. This phenomenon has led to the search for global properties of the learning problem which imply localization—such as strong or uniform convexity of the loss (see Mendelson [2002], Bartlett et al. [2006])—as well as data-dependent upper bounds which could be shown to adapt to localization, notably the local Rademacher complexities of Bartlett et al. [2005].

Recently, Liang et al. [2015] showed that localization for the square loss is captured by another object, the *offset Rademacher complexity*. In addition to being analytically simpler, the offset complexity appears in a minimax analysis of *online* regression [Rakhlin and Sridharan, 2014], suggesting a unified treatment of online and statistical learning. In parallel work, van Erven et al. [2015] has showed that *exp-concavity* of the loss, often studied in online optimization, implies fast rates in the statistical setting via the probabilistic *central condition* (see also Juditsky et al. [2008], Mehta [2017]).

We attempt to unify these results. In particular, we show that the offset complexity upper bound can be generalized to arbitrary losses satisfying a certain curvature condition that includes exp-concave losses. These results considerably extend the scope of offset Rademacher complexities in the statistical setting, as we illustrate; we also believe it can refine the minimax analysis of online regression with exp-concave loss, which we leave to subsequent work.

A noteworthy aspect of our work is its treatment of *improper learning*—that is, minimizing loss compared to the class $\mathcal{F}$ with an estimator that need not belong to $\mathcal{F}$. Unlike prior work, which typically involves a reduction to finite aggregation over a discrete subset of $\mathcal{F}$ [Rakhlin et al., 2017, Audibert, 2009], we provide a completely unified analysis that simultaneously leads to sharp rates for convex empirical risk minimization *and* improper learning under general entropy conditions. In particular, we circumvent the issue that passing to the convex hull of certain classes can "blow up" their complexity and lead to slower rates [Mendelson, 2002].

Finally, we apply our analysis to generalized linear models (glms), such as logistic regression, where standard procedures surprisingly do not attain fast rates [Hazan et al., 2014]. These are notable both for their widespread use as well as for their equivalence to classical maximum likelihood under misspecification. Inspired by Foster et al. [2018], we note that glms can be recast as learning in a *non-convex* class subject to the exp-concave relative entropy. The star algorithm therefore attains fast rates for a wide class of generalized linear models, improving on prior results which exploit strong convexity (see e.g. Rigollet [2012]). Unlike Foster et al. [2018], however, our results are specific to the batch setting; whether a similar analysis extends to online learning remains open.

2 Margin inequalities

We begin our discussion with the following definition which extends the classical *uniform convexity* (see e.g. Chidume [2009]), giving a quantitative bound on the extent to which the plane tangent to f at $(y, f(y))$ lies below the graph of f.

Definition 2. *Let $\mu : \mathbb{R}_+ \to \mathbb{R}_+$ be increasing and convex, and let $d(x, y)$ be a metric on the vector space B. A convex function $f : B \to \mathbb{R}$ with sub-gradient ∇f is (μ, d)-convex if for all $x, y \in B$,*

$$f(x) - f(y) - \langle \nabla f|_y,\, x - y \rangle \geq \mu(d(x, y)) \tag{1}$$

In particular, this condition implies that the lower sets of f,

$$f_x = \left\{ z \,\middle|\, f(z) \leq f(x) \right\},$$

are strictly convex, and that for any closed and convex K there is a unique $\hat{x} \in \arg\min_{x \in K} f(x)$. We adopt the notation that for a subset $S \subset B$, ∂S denotes the boundary of S, $S^\circ = S \setminus \partial S$ denotes the interior, and $\mathrm{co}\, S$ denotes the convex hull.

Remark 3. At this point, we are intentionally vague about the role of the metric, d. We shall subsequently see that the choice of d is crucial in producing good upper bounds. The conventional choice is $d(x, y) = |x - y|$ and $\mu(z) = \alpha z^p$, corresponding to *p-uniform convexity (with modulus α)*. However, our results on self-concordant and exp-concave losses use non-standard choices of d.

By a standard argument, given in Lemma 4 below, the condition (1) on the loss implies that the empirical risk minimizer in a convex class satisfies a *geometric margin inequality*. Intuitively, if another hypothesis performs nearly as well as the empirical minimizer in the sample, it must also be d-close to the empirical minimizer.

Lemma 4. *Let K be a closed, convex subset of B. If f is (μ, d)-convex and $\hat{x}$ minimizes f over K then for all $x \in K$*

$$f(x) - f(\hat{x}) \geq \mu(d(x, \hat{x})). \tag{2}$$

Proof. Let $z_\lambda = \lambda x + (1 - \lambda)\hat{x}$ be the line segment interpolating x and $\hat{x}$, and put $\varphi(\lambda) = f(z_\lambda)$. By convexity of K, we must have $z_\lambda \in K$, hence φ is minimized at $\lambda = 0$ by optimality of $\hat{x}$. Hence $\varphi(\lambda) - \varphi(0) \geq 0$ for all λ, and $\nabla \varphi$ must satisfy

$$\nabla \varphi|_{\lambda=0} = \langle \nabla f|_{\hat{x}},\, x - \hat{x} \rangle \geq 0.$$

Thus,

$$f(x) - f(\hat{x}) \geq f(x) - f(\hat{x}) - \langle \nabla f|_{\hat{x}},\, x - \hat{x} \rangle \geq \mu(d(x, \hat{x})),$$

proving (2). $\qquad\square$

2.1 The star algorithm

We now introduce our first main result, which extends the above analysis to the problem of learning in a non-convex class. Surprisingly, in the general setting of (μ, d)-convexity, an inequality matching (2) is *always* attained by a simple, two-step variant of empirical risk minimization.

Let us now introduce the procedure. Given $x \in S$, the "star hull" of S with respect to x is defined by

$$\mathrm{star}(x, S) = \{\lambda x + (1 - \lambda)s \,|\, s \in S, \lambda \in [0, 1]\}.$$

The procedure may then be summarized as follows.

1: **procedure** STAR(S, f)
2:      Find $\hat{x}$ that minimizes f over S
3:      Find $\tilde{x}$ that minimizes f over $\mathrm{star}(\hat{x}, S)$
4:      **Output** $\tilde{x}$
5: **end procedure**

Our next goal is to show that if S is an arbitrary (not necessarily convex) set, and $\tilde{x}$ is the output of the star algorithm, then

$$f(x) - f(\tilde{x}) \geq \mu\left(\tfrac{1}{3}d(x, \tilde{x})\right) \tag{3}$$

Figure 1: Illustration of Lemma 4. Convexity implies the existence of a direct path z_λ from x to $\hat{x}$.

Figure 2: Illustration of Proposition 5. Depicted is the plane containing $(x, \hat{x}, \tilde{x})$; C is the minimal cone containing $f_{\tilde{x}}$ with vertex at $\hat{x}$. By optimality of $\tilde{x}$, S cannot intersect the interior of $C \cup f_{\hat{x}}$, implying existence of one of the indicated paths.

whenever f is (μ, d)-convex. In particular, if f is the empirical risk and S is the model class, then $\tilde{x}$ is the "star estimator" introduced by Audibert [2008]. Thus, whenever (μ, d)-convexity leads to an upper bound for empirical risk minimization in a convex class via the geometric margin inequality (2), the same bound is enjoyed (up to constants) by the star algorithm in an arbitrary class.

To prove (3), recall how we proved Lemma 4: we showed that if the line-segment $\lambda \mapsto z_\lambda$ which linearly interpolates between any two points x and y lies entirely in the set $f_x \setminus f_y^\circ$, then we have that

$$f(x) - f(y) \geq \mu(d(x, y)).$$

Unfortunately, the existence of such a line-segment z_λ connecting $\tilde{x}$ and x relies on the convexity of S. In order to extend the argument as we want to, we make two observations. Firstly, the line segment $\lambda \mapsto z_\lambda$ can be chosen to be *any* line segment connecting ∂f_x to ∂f_y and lying entirely in $f_x \setminus f_y^\circ$. Then, since ∂f_x and ∂f_y are level sets, the same argument yields

$$f(x) - f(y) = f(z_1) - f(z_0) \geq \mu(d(z_1, z_0)). \tag{4}$$

Secondly, we can show that there exist *at most three* such line-segments which together comprise a path from $\tilde{x}$ to x. By the triangle inequality, at least one of these segments has length greater than $\frac{1}{3} d(\tilde{x}, x)$. Combining this with (4) yields the desired inequality (3). We state this as a proposition and prove it formally in Appendix A. A straightforward illustration of the proof, following the preceding discussion, is provided in Figure 2.

Proposition 5. *Let f be (μ, d)-convex. Suppose $\hat{x}$ minimizes f over S and $\tilde{x}$ minimizes f over* $\mathrm{star}(\hat{x}, S)$*. Then, for any $x \in S$,*

$$f(x) - f(\tilde{x}) \geq \mu\left(\tfrac{1}{3} d(x, \tilde{x})\right).$$

3 From margins to upper bounds

We will now show how the previous section's margin inequalities lead to upper bounds on the excess risk. Our first lemma extends (μ, d)-convexity of the loss $x \mapsto \psi(x, y)$ to the empirical risk.

Lemma 6. *Suppose that for each y, $x \mapsto \psi(x, y)$ is (μ, d)-convex. Then the empirical ψ-risk, given by $f \mapsto \mathbb{E}_n[\psi(f(X_i), Y_i)]$, is (μ, d_n)-convex, where $d_n = \mu^{-1} \circ \mathbb{E}_n(\mu \circ d)$.*

Now, letting $\mathcal{F}$ be a convex class of functions, with empirical risk minimizer $\hat{f}$, applying Lemma 4 with x chosen to be the population risk minimizer f^* gives

$$\mathbb{E}_n \psi(f_i^*) - \mathbb{E}_n \psi(\hat{f}_i) \geq \mathbb{E}_n \mu(d(\hat{f}_i, f_i^*)). \tag{5}$$

Similarly, if $\mathcal{F}$ is not convex and $\tilde{f}$ is produced by the star algorithm applied to $\mathbb{E}_n \psi$, we have by Proposition 5

$$\mathbb{E}_n \psi(f_i^*) - \mathbb{E}_n \psi(\tilde{f}_i) \geq \mathbb{E}_n \mu(d(\tilde{f}_i, f_i^*)/3). \tag{6}$$

147 We note that in our applications, μ will typically have the form $x \mapsto cx^2$, so (5) and (6) are comparable
148 up to a constant factor of 9. To avoid complication, we will only work with the more general inequality
149 (6). The reader may note that in a convex class the star estimator coincides with the empirical risk
150 minimizer, so we are treating both cases simultaneously at the expense of slightly larger constants.

151 **Example 1** (Square loss). Before proceeding, it is helpful to consider the square loss $\psi(x, a) =$
152 $(x - a)^2$ as an example. A convenient aspect of (μ, d)-convexity is that the optimal $\mu \circ d$ can be
153 computed as

$$\mu(d(x, y)) = \psi(x) - \psi(y) - \langle \nabla \psi|_y, \, x - y \rangle$$
$$= \lim_{\lambda \downarrow 0} \frac{d}{d\lambda} \lambda \psi(x) + (1 - \lambda)\psi(y) - \psi(\lambda x + (1 - \lambda)y),$$

154 which in this case is simply $(x - y)^2$. Taking $d(x, y) = |x - y|$ and $\mu(z) = z^2$, we recover the
155 familiar inequality

$$\mathbb{E}_n(Y - f^*)^2 - \mathbb{E}_n(Y - \hat{f})^2 \geq \mathbb{E}_n(f^* - \hat{f})^2.$$

156 Moving to the the non-convex case with the star algorithm introduces a factor of $1/9$ on the right-hand
157 side, improving on Liang et al. [2015] by a constant. This analysis extends, with little complication
158 to the p-loss for all $1 < p < \infty$. We will discuss these results in Section 5.1.

3.1 The offset empirical process

160 We have now derived a family of margin inequalities for the star estimator,

$$\mathbb{E}_n \psi(f^*) - \mathbb{E}_n \psi(\tilde{f}) - \mathbb{E}_n \mu(d(f^*, \tilde{f})/3) \geq 0, \tag{7}$$

161 where we introduce the abbreviated notation $\psi(h) \equiv \psi(h(Z)) \equiv \psi(h(X), Y)$. Adding this to the
162 excess risk $\mathcal{E}(\tilde{f}) \equiv \mathbb{E}\psi(\tilde{f}) - \mathbb{E}\psi(f^*)$ gives us the upper bound

$$\mathcal{E}(\tilde{f}) = \mathbb{E}\psi(\tilde{f}) - \mathbb{E}\psi(f^*) \leq (\mathbb{E}_n - \mathbb{E})(\psi(f^*) - \psi(\tilde{f})) - \mathbb{E}_n \mu(d(\tilde{f}, f^*)/3)$$
$$\leq \sup_{f \in \mathcal{F}'} \left\{ (\mathbb{E}_n - \mathbb{E})(\psi(f^*) - \psi(f)) - \mathbb{E}_n \mu(d(f, f^*)/3) \right\} \tag{8}$$

163 for $\mathcal{F}' = \cup_{\lambda \in [0,1]} \lambda \mathcal{F} + (1 - \lambda)\mathcal{F}$ (the Minkowski sum). After symmetrization and contraction
164 arguments, which follow along the lines of Liang et al. [2015], we have the following result. The
165 proof is given in Appendix B.

166 **Proposition 7.** *Let $\mathcal{F}$ be a model class, ψ a (μ, d)-convex loss, and f^* the population minimizer of*
167 *the ψ-risk. Then, the star estimator $\tilde{f}$ satisfies the excess risk bound*

$$\mathbb{E}\Psi(\mathcal{E}(\tilde{f})) \leq \mathbb{E}\Psi \left(\sup_{f \in \mathcal{F}'} \left\{ \frac{1}{n} \sum_{i=1}^n 4\varepsilon_i'(\psi(f_i) - \psi(f_i^*)) - \mu(d(f_i, f_i^*)/3) \right\} \right) \tag{9}$$

168 *where the ε_i' are i.i.d. symmetric Rademacher random variables, $\mathcal{F}' = \cup_{\lambda \in [0,1]} \lambda \mathcal{F} + (1 - \lambda)\mathcal{F}$, and*
169 $\Psi : \mathbb{R} \to \mathbb{R}$ *is any increasing, convex function.*

170 **Remark 8.** Proposition 7 quantifies localization for the loss ψ. The excess risk is upper bounded
171 by a *non-centered* Rademacher complexity with the asymmetric term $-\mu(d(f_i, f_i^*)/3)$. This term
172 captures the fact that hypotheses f which are far from f^* perform worse in expectation, and are thus
173 unlikely to be selected. As a result, these hypotheses do not contribute much to the excess risk.

4 Exp-concave localization

175 We will now investigate how our upper bound (9) applies to η-exp-concave losses, namely those for
176 which $e^{-\eta\psi}$ is concave, or equivalently

$$\psi''/(\psi')^2 \geq \eta.$$

177 It turns out that this condition leads to very natural bounds on the excess risk. To provide intuition as
178 to why, note that exploiting the bound (9) requires comparing the risk increments $\psi(f) - \psi(f^*)$ to the
179 offset $\mu(d(f, f^*))$. Under strong convexity, we have an offset of the form $\alpha|f - f^*|^2$ and must resort
180 to a Lipschitz bound of the form $\psi(f) - \psi(f^*) \leq \|\psi\|_{\text{lip}} |f - f^*|$ to make the necessary comparison.

Thus, we have upper bounded the first derivative uniformly by $\|\psi\|_{\mathrm{lip}}$ and lower bounded the second derivative uniformly by α.

It turns out that exp-concave losses possess an offset term of the form $|\psi(f) - \psi(f^*)|^2$. Thus, *localization naturally occurs at the same scale as the risk increments*. Exp-concavity only requires the ratio $\psi''/(\psi')^2$ to stay bounded, which is crucial for the log loss, where $\|\psi\|_{\mathrm{lip}} = \infty$ but $\psi''/(\psi')^2 = 1$. First, we'll establish a particular version (μ, d)-convexity for the logarithmic loss. We will subsequently extend it to all exp-concave losses.

Lemma 9. *For all $x, y \in [\delta, 1]$*

$$(-\ln x) - (-\ln y) - \nabla(-\ln)|_y (x - y) \geq \frac{|\ln x - \ln y|^2}{2 \ln(1/\delta) \vee 4}. \tag{10}$$

Proof. The left hand side of (10) is equivalent to $\ln(y/x) + (x - y)/y = e^{-z} - 1 + z$ with the substitution $z \equiv \ln(y) - \ln(x)$. Finally, we verify in Lemma G3 that

$$e^{-z} - 1 + z \geq \frac{z^2}{2 \ln(1/\delta) \vee 4}, \tag{11}$$

since our assumption $x, y \in [\delta, 1]$ implies $|z| \leq \ln(1/\delta)$. $\qquad \square$

The inequality (10) establishes that the logarithmic loss, when restricted to $[\delta, 1]$, is (μ, d)-convex for $\mu : x \mapsto x^2/(2 \ln(1/\delta) \vee 4)$ and $d(x, y) = |\ln x - \ln y|$. This choice of d captures an important fact: the logarithm is *more convex* precisely when it is *faster varying*.

Remarkably, a simple reduction shows that the same property is enjoyed by bounded, η-exp-concave losses. Thus, we neatly recover past results on exp-concave learning (cf. Mehta [2017]), and extend them to the improper setting.

Proposition 10. *Suppose ψ is an η-exp-concave loss function which satisfies the uniform bound $0 \leq \psi \leq m$. Then, for each x and y,*

$$\psi(x) - \psi(y) - \langle \nabla \psi|_y, x - y \rangle \geq \frac{|\psi(x) - \psi(y)|^2}{2m \vee (4/\eta)}. \tag{12}$$

Proof. First note that since $e^{-\eta \psi}$ is concave, we have

$$\exp(-\eta \psi(\lambda x + (1 - \lambda)y)) \geq \lambda e^{-\eta \psi(x)} + (1 - \lambda)e^{-\eta \psi(y)}.$$

Note that equality holds at $\lambda = 0$. Taking logarithms of both sides and passing to the upper sub-gradient at $\lambda = 0$ therefore yields

$$-\eta \langle \nabla \psi|_y, x - y \rangle \geq \left\langle \nabla \ln |_{e^{-\eta \psi(y)}}, e^{-\eta \psi(x)} - e^{-\eta \psi(y)} \right\rangle. \tag{13}$$

Applying Lemma 9 with x and y replaced by $e^{-\eta \psi(x)}$ and $e^{-\eta \psi(y)}$ and $\delta = e^{-\eta m}$, then using (13) to bound the gradient term yields Proposition 10. $\qquad \square$

Plugging (12) into Proposition 7 and making a contraction argument yields the following upper bound for the excess ψ-risk. The full argument is in Appendix C.

Theorem 11. *Let ψ be an η-exp-concave loss function taking values in $[0, m]$. Then the star estimator in $\mathcal{F}$ satisfies the excess risk bound*

$$\mathbb{E}\Psi(\mathcal{E}(\tilde{f})) \leq \mathbb{E}\Psi\left(\sup_{f, g \in \mathcal{F}'} \left\{ \frac{1}{n} \sum_{i=1}^{n} 4\varepsilon_i'(\psi(f_i) - \psi(g_i)) - \frac{\eta(\psi(f_i) - \psi(g_i))^2}{18m\eta \vee 36} \right\} \right). \tag{14}$$

where Ψ is any increasing, convex function and $\mathcal{F}' = \cup_{\lambda \in [0,1]} \lambda \mathcal{F} + (1 - \lambda)\mathcal{F}$. Alternatively, when ψ is p-uniformly convex with modulus α and $\|\psi\|_{\mathrm{lip}}$-Lipschitz, we have

$$\mathbb{E}\Psi(\mathcal{E}(\tilde{f})) \leq \mathbb{E}\Psi\left(\sup_{f, g \in \mathcal{F}'} \left\{ \frac{1}{n} \sum_{i=1}^{n} 4 \|\psi\|_{\mathrm{lip}} (f_i - g_i)\varepsilon_i' - \frac{\alpha|f_i - g_i|^p}{3^p} \right\} \right). \tag{15}$$

Self-concordance

We close the section by considering *self-concordant* losses ψ, namely those for which

$$\psi''' \le 2(\psi'')^{\frac{3}{2}}.$$

In order to see how localization for self-concordant losses can be fit into the above framework, we rely on the following inequality which was also used by Bilodeau et al. [2020].

Lemma 12 (Nesterov [2014, Theorem 4.1.7]). *Suppose ψ is self-concordant, and define the* local norm $\|z\|_{\psi,w} \equiv \sqrt{z^2\psi''(w)}$. *Then*

$$\psi(x) - \psi(y) - \langle \nabla f|_y,\, x - y\rangle \ge \|x-y\|_{\psi,y} - \ln\left(1 + \|x-y\|_{\psi,y}\right). \tag{16}$$

The first thing that one should note is that, when one takes ψ to be the negative logarithm, the above is exactly equivalent to the modulus (11) appearing in the proof of Lemma 9. Thus, *our upper bound for exp-concave losses is a consequence of self-concordance of the logarithm.*

The path to general upper bounds via self-concordance is significantly more involved, so we will only sketch it. Applying (16) to the excess risk and performing standard manipulations, one arrives at the following upper bound, proved in Appendix D.

Proposition 13. *If ψ is a self-concordant loss and $\hat{f}$ is the empirical risk minimizer in a convex class $\mathcal{F}$, then*

$$\mathbb{E}\Psi\left(\mathcal{E}(\tilde{f})\right) \le \mathbb{E}\Psi\left(\sup_{f\in\mathcal{F}'}\left\{\frac{1}{n}\sum_{i=1}^{n}4(\psi(f_i) - \psi(f_i^*))\varepsilon_i' - \omega\left(\|f_i - f_i^*\|_{\psi,f_i^*}\right)\right\}\right), \tag{17}$$

for $\omega(z) = z - \log(1+z)$, $\|z\|_{\psi,w} \equiv \sqrt{z^2\psi''(w)}$, and $(\varepsilon_i')_{i=1}^{n}$ are independent, symmetric Rademacher random variables and Ψ is any increasing, convex function.

Comparing this to (15), the scale of the offset term has been improved from the modulus of strong convexity, which is the *minimum* second derivative, to the second derivative at the risk minimizer, f^*. Expressing the increments $\psi(f_i) - \psi(f_i^*)$ at a scale comparable to $\|f_i - f_i^*\|_{\psi,f_i^*}$ is known to be achievable when f belongs to the so-called *Dikin ellipsoid* of $(\mathbb{E}_n\psi, f^*)$, requiring a more careful argument (see e.g. Marteau-Ferey et al. [2019]). Notably, we do not know how to extend this bound to the improper setting due to the local nature of the offset term.

5 Applications

We will now discuss several applications of Theorem 11. First, we'll sketch how the previous section's results straightforwardly imply fast rates. We'll then discuss implications, first for the p-loss and then for generalized linear models. Proofs in both cases are to be found in Appendix F.

While most general bound uses chaining and is provided by Theorem 17 below, the essence of the argument is simple and we will sketch it now. Let $S_\epsilon(\mathcal{A},\mu)$ be a minimal covering of $\mathcal{A}$ in $L^2(\mu)$ at resolution ϵ, and define the entropy numbers $H_2(\epsilon,\mathcal{G}) = \sup_\mu \ln(\#S_\epsilon(\mathcal{G},\mu))$ (where the supremum is over probability measures μ). Putting $S = S_\epsilon(\psi \circ \mathcal{F}', \mathbb{P}_n)$ and applying (14) with $\Psi(z) = \exp(\lambda nz)$ gives

$$\mathbb{E}\exp(\lambda n(\mathcal{E}(\tilde{f}) - \epsilon)) \le \mathbb{E}\exp \lambda n\left(\sup_{s,t\in S_\epsilon}\left\{\frac{1}{n}\sum_{i=1}^{n}4\varepsilon_i'(s_i - t_i) - \frac{\eta(s_i - t_i)^2}{18m\eta \vee 36}\right\}\right)$$

$$\le \mathbb{E}\sum_{s,t\in S}\mathbb{E}_\varepsilon\exp\left(\sum_{i=1}^{n}4\lambda\varepsilon_i'(s_i - t_i) - \frac{\lambda\eta(s_i - t_i)^2}{18m\eta \vee 36}\right),$$

where for the second inequality we moved the supremum outside $\exp$ (by monotonicity of $\exp$) and bounded it by a sum (by positivity of the arguments). Here, $\mathbb{E}_\varepsilon$ denotes an expectation w.r.t. the multipliers ε_i', conditional upon the data. Applying Hoeffding's lemma to the inner expectations with respect to ε_i' gives

$$\le \mathbb{E}\sum_{s,t\in S}\exp\left(\sum_{i=1}^{n}2\lambda^2(s_i - t_i)^2 - \frac{\lambda\eta(s_i - t_i)^2}{18m\eta \vee 36}\right).$$

Choosing $\lambda^* = 36m \vee 72/\eta$ gives $\mathbb{E}\exp(\lambda^* n(\mathcal{E}(\tilde{f}) - \epsilon)) \leq \mathbb{E}(\#S)^2 \leq 2H_2(\epsilon, \psi \circ \mathcal{F}')$, so by a Chernoff bound

$$\mathbb{P}\left(\mathcal{E}(\tilde{f}) \geq \epsilon + \left(36m \vee \frac{72}{\eta}\right)\left(\frac{2H_2(\epsilon, \psi \circ \mathcal{F}') + \ln(1/\rho)}{n}\right)\right) \leq \rho. \tag{18}$$

The result (18) extends Mehta [2017, Theorem 1] to the non-convex and improper setting for the price of a factor 9, and differs from the standard bound in numerous ways. Firstly, the Lipschitz constant $\|\psi\|_{\mathrm{lip}}$ is replaced by the range, m. Here, $\|\psi\|_{\mathrm{lip}}$ appears only implicitly in the covering numbers, entering logarithmically in small classes. For the log loss, this is an exponential improvement.

Next, the modulus of strong convexity is replaced by the exp-concavity modulus η. This implies that the optimal $1/n$ rate can be obtained, in both proper and improper settings, for losses which are not strongly convex but are exp-concave, notably regression with p-loss for $p > 2$.

Lastly, we show in Appendix Lemma G4 that the entropy numbers of $\psi \circ \mathcal{F}'$ are equivalent to those of $\psi \circ \mathcal{F}$ in all but finite classes, where the gap is at most $\ln(1/\epsilon)$. On the other hand, when the entropy numbers of $\psi \circ \mathcal{F}$ are polynomial, particularly ϵ^{-q} with $q < 2$, then they can be *polynomially smaller* than those of $\psi \circ (\mathrm{co}\,\mathcal{F})$ [Carl et al., 1999].

5.1 Regression with p-loss

A surprising consequence of (18) concerns regression with the p-loss for $p > 2$, which is p-uniformly convex but *not* strongly convex. When the inputs to the loss are bounded, the standard argument pursued by Mendelson [2002] and Bartlett et al. [2006] (and also Rakhlin and Sridharan [2015] in the online setting) exploits uniform convexity, producing the *sub-optimal* rate exponent $p/(2p-2) < 1$. However, the p-loss for inputs in $[-B, B]$ is B^{-p}-exp-concave, so we optimal $(1/n)$-type rates matching the square loss. Indeed, the sub-optimal rates occur exactly when one plugs (15) into the above argument instead of (14). For $1 < p \leq 2$, the loss is both strongly convex and exp-concave, so the two bounds coincide up to constants.

Corollary 14 (cf. Mendelson [2002, Theorem 5.1]). *Let $\psi(f, y) = |f - y|^p$ for $p > 1$ and let the class $\mathcal{F}$ and response Y take values in $[-B, B]$. Then there exists a universal $C_{p,B} = O(p2^p B^p)$ such that the star estimator $\tilde{f}$ has excess ψ-risk bounded as*

$$\mathbb{P}\left(\mathcal{E}(\tilde{f}, \mathcal{F}) \geq \epsilon + \frac{C_{p,B}(H_2(\epsilon/C_{p,B}, \mathcal{F}) + \ln(1/\epsilon) + \ln(1/\rho))}{n}\right) \leq \rho, \tag{19}$$

By a minor extension to Tsybakov [2003, Theorem 1], this result is sharp for each p.

5.2 Improper maximum likelihood

We next derive upper bounds for learning under the log loss, $\psi : f \mapsto -\ln f$ in a class $\mathcal{L}$. If we interpret $f(Z_i)$ as the likelihood of an observation, empirical risk minimization concides with maximum likelihood. In this setting, generalized linear models for multi-class prediction correspond to cases where $\mathcal{L} = \{(x, y) \mapsto \langle \varphi(f), y \rangle \mid f \in \mathcal{F}\}$ for some base class $\mathcal{F}$. Here the *link function* φ outputs probability distributions over $[k]$ and $y \in \mathbb{R}^k$ is a basis vector encoding the label. These are often formulated as minimizing the loss $\psi^{[\varphi]} : f \mapsto -\ln \langle \varphi(f), y \rangle$ over $f \in \mathcal{F}$, which is equivalent for proper learning. Since the transformation from $\mathcal{F}$ to $\mathcal{L}$ may not preserve convexity, however, one must consider improper procedures to exploit localization of the log loss, even for convex $\mathcal{F}$.

To illustrate, we will prove Proposition 1 on logistic regression, where Hazan et al. [2014] showed that improper learning is in fact necessary for fast rates. This setting was previously considered by Foster et al. [2018], whose results fall neatly into the aforementioned recipe: the mixed prediction corresponding to an exponentially weighted average is simply another estimator taking values in the convex hull of $\mathcal{L}$. When φ is onto, these can be turned into improper predictors under $\psi^{[\varphi]}$.

For classes not bounded away from 0, where the improvement is most dramatic, we need to replace $\mathcal{L}$ by the δ-regularization $\mathcal{L}_\delta = (1 - \delta)\mathcal{L} + \delta$. The approximation error is controlled as follows.

Lemma 15 (Foster et al. [2018]). *For all f and $\delta \in (0, 1/2]$, the excess risk relative to $\mathcal{L}_\delta$ satisfies*

$$\mathcal{E}(f; \mathcal{L}_\delta) \leq \mathcal{E}(f; \mathcal{L}) + 2\delta$$

We now state our result.

Corollary 16. *With probability at least $1 - \rho$, the star estimator $\tilde{f}_\delta$ in $\mathcal{L}_\delta$ satisfies*

$$\mathcal{E}(\tilde{f}_\delta; \mathcal{L}) \leq \epsilon + 2\delta + C \ln(1/\delta) \left(\frac{H_2(\delta\epsilon, \mathcal{L}) + \ln(1/\epsilon\delta) + \ln(1/\rho)}{n} \right) \qquad (20)$$

If $\mathcal{L}$ is the generalized linear model corresponding to

$$\mathcal{F}_B = \left\{ x \mapsto Wx \,\middle|\, W \in \mathbb{R}^{k \times d}, \|W\|_{2 \to \infty} \leq B \right\}$$

with A-Lipschitz and surjective link φ (with right-inverse $\varphi^\dagger$), then with probability at least $1 - \rho$,

$$\mathbb{E}\left[-\ln \left\langle \varphi^\dagger(\tilde{f}_\delta(X)), Y \right\rangle \right] - \inf_{f \in \mathcal{F}} \mathbb{E}\left[-\ln \langle f(X), Y \rangle \right] \leq \frac{Ckd\ln(ABn)^2 \ln(1/\rho)}{n}. \qquad (21)$$

Proposition 1 follows after noting that the soft-max function is 1-Lipschitz and surjective. Before investigating how these bounds translate to larger classes—namely those with $H_2(\epsilon, \mathcal{F})$ of order ϵ^{-q}, we need to state the chaining version of our upper bound, proved in Appendix E.

Theorem 17. *Let ψ be an η-exp-concave loss taking values in $[0, m]$. Then, with probability at least $1 - (3e)e^{-z}$, the star estimator $\tilde{f}$ applied to $(\psi, \mathcal{F})$ satisfies*

$$\mathcal{E}(\tilde{f}) \leq \inf_{0 \leq \alpha \leq \gamma} \left\{ 4\alpha + \frac{10}{\sqrt{n}} \int_\alpha^\gamma \sqrt{H_2(s)} \, ds + \frac{2H_2(\gamma)}{cn} + \frac{\gamma\sqrt{8\pi}}{\sqrt{n}} + \left(\frac{2}{cn} + \frac{\gamma\sqrt{8}}{\sqrt{n}} \right) z \right\}, \qquad (22)$$

where $H_2(s) \equiv H_2(s, \psi \circ \mathcal{F}')$ and $c = 36^{-1}(1/m \wedge \eta/2)$.

In classes with entropy ϵ^{-q}, our results fall into two regimes. The first covers losses such as the logistic loss, for which $-\ln \circ \varphi$ is Lipschitz. Here, the covering numbers of $-\ln \circ \mathcal{L}_\delta$ differ from those of $\mathcal{F}$ by a constant, and we match the optimal rates for the square loss [Rakhlin et al., 2017].

For general $\mathcal{L}$, the covering numbers of $-\ln \circ \mathcal{L}_\delta$ may contain an extra factor δ^{-q}, leading to sub-optimal rates. Sub-optimality follows from combining the minimax regret analysis of Bilodeau et al. [2020], which produces the average regret $n^{-1/(1+q)}$, with a standard online-to-batch reduction. We note that the upper bound of that paper is non-constructive, whereas these upper rates come with an explicit algorithm. Tightening our analysis to close this gap, or proposing an algorithm that attains the correct rates, remains a significant open problem.

Corollary 18. *Consider a generalized linear model with A-Lipschitz loss $f \mapsto -\ln \langle \varphi(f), y \rangle$. Suppose the $L^2(\mathbb{P}_n)$ entropy numbers $H_2(\epsilon; \mathcal{F})$ are almost surely of order ϵ^{-q}. Then, the regularized star estimator $\varphi^\dagger \circ \tilde{f}_\delta$ with $\delta = 1/n$ satisfies the rates appearing on the left.*

On the other hand, for an arbitrary class $\mathcal{L}$ taking values in $[0, 1]$ subject to the log loss, the regularized star estimator $\tilde{f}'_{\delta'}$—for appropriately chosen δ'—attains the rates appearing on the right.

Here, the symbol $\lesssim_\rho$ denotes an upper bound that holds with probability $1 - \rho$, hiding universal constants and a multiplicative factor $\ln(1/\rho)$.

$$\mathcal{E}(\varphi^\dagger \circ \tilde{f}_\delta) \lesssim_\rho \begin{cases} A^q n^{-2/(2+q)} \ln(n) & q < 2 \\ A^q n^{-1/2} \ln(n) & q = 2 \\ A^q n^{-1/q} & q > 2 \end{cases} \qquad \mathcal{E}(\tilde{f}'_{\delta'}) \lesssim_\rho \begin{cases} n^{-1/(1+3q/2)} & q < 2 \\ n^{-1/4} \ln(n) & q = 2 \\ n^{-1/2q} & q > 2 \end{cases} \qquad (23)$$

In closing, we would like to highlight the additional, unresolved problem of designing a computationally efficient implementation of the regularized star estimator—in particular for log-concave link functions φ where the first step is a convex program, such as logistic regression. This problem is left to future work.

References

Jean-yves Audibert. Progressive mixture rules are deviation suboptimal. In J. C. Platt, D. Koller, Y. Singer, and S. T. Roweis, editors, *Advances in Neural Information Processing Systems 20*, pages 41–48. Curran Associates, Inc., 2008.

Jean-Yves Audibert. Fast learning rates in statistical inference through aggregation. *The Annals of Statistics*, 37 (4):1591 – 1646, 2009. doi: 10.1214/08-AOS623. URL https://doi.org/10.1214/08-AOS623.

Peter L. Bartlett, Olivier Bousquet, and Shahar Mendelson. Local Rademacher complexities. *Annals of Statistics*, 33(4):1497–1537, August 2005. ISSN 0090-5364, 2168-8966. doi: {10.1214/009053605000000282}.

Peter L. Bartlett, Michael I. Jordan, and Jon D. Mcauliffe. Convexity, Classification, and Risk Bounds. *Journal of the American Statistical Association*, 101(473):138–156, 2006. ISSN 0162-1459.

Blair Bilodeau, Dylan Foster, and Daniel Roy. Tight Bounds on Minimax Regret under Logarithmic Loss via Self-Concordance. In *International Conference on Machine Learning*, pages 919–929. PMLR, November 2020.

Bernd Carl, Ioanna Kyrezi, and Alain Pajor. Metric Entropy of Convex Hulls in Banach Spaces. *Journal of the London Mathematical Society*, 60(3):871–896, December 1999. ISSN 0024-6107. doi: {10.1112/S0024610799008005}.

Charles Chidume. Inequalities in Uniformly Convex Spaces. In Charles Chidume, editor, *Geometric Properties of Banach Spaces and Nonlinear Iterations*, Lecture Notes in Mathematics, pages 29–44. Springer, London, 2009. ISBN 978-1-84882-190-3. doi: {10.1007/978-1-84882-190-3_4}.

Dylan J. Foster, Satyen Kale, Haipeng Luo, Mehryar Mohri, and Karthik Sridharan. Logistic Regression: The Importance of Being Improper. In *Conference On Learning Theory*, pages 167–208. PMLR, July 2018.

Elad Hazan, Tomer Koren, and Kfir Y. Levy. Logistic regression: Tight bounds for stochastic and online optimization. In Maria Florina Balcan, Vitaly Feldman, and Csaba Szepesvári, editors, *Proceedings of the 27th Conference on Learning Theory*, volume 35 of *Proceedings of Machine Learning Research*, pages 197–209, Barcelona, Spain, June 2014. PMLR.

A. Juditsky, P. Rigollet, and A. B. Tsybakov. Learning by mirror averaging. *Annals of Statistics*, 36(5): 2183–2206, October 2008. ISSN 0090-5364, 2168-8966. doi: {10.1214/07-AOS546}.

Vladimir Koltchinskii and Dmitriy Panchenko. Rademacher Processes and Bounding the Risk of Function Learning. In Evarist Giné, David M. Mason, and Jon A. Wellner, editors, *High Dimensional Probability II*, Progress in Probability, pages 443–457, Boston, MA, 2000. Birkhäuser. ISBN 978-1-4612-1358-1. doi: {10.1007/978-1-4612-1358-1_29}.

Michel Ledoux and Michel Talagrand. Rademacher Averages. In Michel Ledoux and Michel Talagrand, editors, *Probability in Banach Spaces: Isoperimetry and Processes*, Ergebnisse Der Mathematik Und Ihrer Grenzgebiete, pages 89–121. Springer, Berlin, Heidelberg, 1991. ISBN 978-3-642-20212-4. doi: {10.1007/978-3-642-20212-4_6}.

Tengyuan Liang, Alexander Rakhlin, and Karthik Sridharan. Learning with Square Loss: Localization through Offset Rademacher Complexity. In *Conference on Learning Theory*, pages 1260–1285, June 2015.

Ulysse Marteau-Ferey, Dmitrii Ostrovskii, Francis Bach, and Alessandro Rudi. Beyond Least-Squares: Fast Rates for Regularized Empirical Risk Minimization through Self-Concordance. In *Conference on Learning Theory*, pages 2294–2340. PMLR, June 2019.

Nishant Mehta. Fast rates with high probability in exp-concave statistical learning. In *Artificial Intelligence and Statistics*, pages 1085–1093. PMLR, April 2017.

S. Mendelson. Improving the sample complexity using global data. *IEEE Transactions on Information Theory*, 48(7):1977–1991, July 2002. ISSN 0018-9448. doi: {10.1109/TIT.2002.1013137}.

Yurii Nesterov. *Introductory Lectures on Convex Optimization: A Basic Course*. Springer Publishing Company, Incorporated, first edition, 2014. ISBN 1-4613-4691-6.

Dmitry Panchenko. Symmetrization approach to concentration inequalities for empirical processes. *Annals of Probability*, 31(4):2068–2081, October 2003. ISSN 0091-1798, 2168-894X. doi: {10.1214/aop/1068646378}.

Alexander Rakhlin and Karthik Sridharan. Online Non-Parametric Regression. In *Conference on Learning Theory*, pages 1232–1264, May 2014.

Alexander Rakhlin and Karthik Sridharan. Online Nonparametric Regression with General Loss Functions. *arXiv:1501.06598 [cs, math, stat]*, January 2015.

Alexander Rakhlin, Karthik Sridharan, and Alexandre B. Tsybakov. Empirical entropy, minimax regret and minimax risk. *Bernoulli*, 23(2):789–824, May 2017. ISSN 1350-7265. doi: {10.3150/14-BEJ679}.

Philippe Rigollet. Kullback–Leibler aggregation and misspecified generalized linear models. *Annals of Statistics*, 40(2):639–665, April 2012. ISSN 0090-5364, 2168-8966. doi: {10.1214/11-AOS961}.

Nathan Srebro, Karthik Sridharan, and Ambuj Tewari. Smoothness, low noise and fast rates. In J. Lafferty, C. Williams, J. Shawe-Taylor, R. Zemel, and A. Culotta, editors, *Advances in Neural Information Processing Systems*, volume 23, pages 2199–2207. Curran Associates, Inc., 2010.

Alexander B. Tsybakov. Optimal aggregation of classifiers in statistical learning. *Annals of Statistics*, 32(1): 135–166, February 2004. ISSN 0090-5364, 2168-8966. doi: {10.1214/aos/1079120131}.

Alexandre B. Tsybakov. Optimal Rates of Aggregation. In Bernhard Schölkopf and Manfred K. Warmuth, editors, *Learning Theory and Kernel Machines*, Lecture Notes in Computer Science, pages 303–313, Berlin, Heidelberg, 2003. Springer. ISBN 978-3-540-45167-9. doi: {10.1007/978-3-540-45167-9_23}.

Tim van Erven, Peter D. Grünwald, Nishant A. Mehta, Mark D. Reid, and Robert C. Williamson. Fast Rates in Statistical and Online Learning. *Journal of Machine Learning Research*, 16(54):1793–1861, 2015. ISSN 1533-7928.