Last-Iterate Convergence of General Parameterized Policies in Constrained MDPs

Abstract

This paper focuses on learning a Constrained Markov Decision Process (CMDP) via general parameterized policies. We propose a Primal-Dual based Regularized Accelerated Natural Policy Gradient (PDR-ANPG) algorithm that uses entropy and quadratic regularizers to reach this goal. For parameterized policy classes with a transferred compatibility approximation error, ϵ_{bias}, PDR-ANPG achieves a last-iterate ϵ optimality gap and ϵ constraint violation with a sample complexity of $\tilde{\mathcal{O}}(\epsilon^{-2}\min\{\epsilon^{-2}, \epsilon_{\text{bias}}^{-\frac{1}{3}}\})$. If the class is incomplete ($\epsilon_{\text{bias}} > 0$), then the sample complexity reduces to $\tilde{\mathcal{O}}(\epsilon^{-2})$ for $\epsilon < (\epsilon_{\text{bias}})^{\frac{1}{6}}$. Moreover, for complete policies with $\epsilon_{\text{bias}} = 0$, our algorithm achieves a last-iterate ϵ optimality gap and ϵ constraint violation with $\tilde{\mathcal{O}}(\epsilon^{-4})$ sample complexity. It is a significant improvement over the state-of-the-art last-iterate guarantees of general parameterized CMDPs.

1 Introduction

Constrained Markov Decision Process (CMDP) is a classical framework where an agent repeatedly interacts with an unknown environment to maximize the cumulative discounted rewards while simultaneously ensuring that the cumulative observed costs are within a pre-defined boundary. It finds its application in a multitude of practical scenarios. For example, consider an autonomous vehicle that attempts to reach its destination via the shortest-time route without violating traffic rules or a corporate leader who aims to maximize revenue without crossing a monetary budget. In these cases, any departure from the boundary set by the predefined rules can be signaled by a cost while the progress towards the desired objective can be indicated by a reward.

Finding an optimal policy to navigate an unknown CMDP is a difficult task. Nevertheless, several recent articles have proposed algorithms to solve this challenging problem with optimality guarantees. For example, Mondal & Aggarwal (2024a) exhibited that their primal dual-based algorithm can achieve ϵ optimality gap and ϵ constrained violation with a $\tilde{O}(\epsilon^{-2})$ sample complexity. Unfortunately, the majority of these works define constraint violation in an average sense. In other words, if their algorithms yield $\{\pi_1, \cdots, \pi_K\}$ policies in K iterations, then the violation is defined as that experienced by a uniformly chosen policy. Since the very nature of this definition allows a large violation at one iteration to get balanced by a smaller violation later, such algorithms are not suitable for safety-critical applications.

To address this challenge, some recent articles have proposed algorithms with last-iterate guarantees. For example, Gladin et al. (2023); Ying et al. (2022) prove last-iterate guarantees for softmax policies, whereas Ding et al. (2024) establishes the same for log-linear policies. Montenegro et al. (2024) recently achieved $\tilde{\mathcal{O}}(\epsilon^{-7})$ sample complexity via general parameterized policies. It is to be emphasized that general parameterization subsumes the tabular softmax and log-linear cases and allows the policies to be represented by neural networks. Moreover, since the general parameterization uses a fixed d number of parameters where d is independent of the size of the state space, it can also be utilized for large or infinite state space. Notably, the current-state-of-the-art sample complexity $\tilde{\mathcal{O}}(\epsilon^{-7})$ is far from the lower bound $\Omega(\epsilon^{-2})$. This raises the following question.

> Is it possible to design an algorithm for general parameterized CMDPs whose last-iterate guarantee is better than the current state-of-the-art?

Algorithm	Sample Complexity	Parameterization
Dual Descent (Ying et al., 2022)	$\tilde{\mathcal{O}}(\epsilon^{-2})$	Softmax
Cutting-Plane (Gladin et al., 2023)	$\tilde{\mathcal{O}}(\epsilon^{-4})$	Softmax
RPG-PD (Ding et al., 2024)	$\tilde{\mathcal{O}}(\epsilon^{-6})$	Log-linear with $\epsilon_{\text{bias}} = \mathcal{O}(\epsilon^8)$
C-PG (Montenegro et al., 2024)	$\tilde{\mathcal{O}}(\epsilon^{-7})$	General
PDR-ANPG (**This Work**)	$\tilde{\mathcal{O}}(\epsilon^{-4})$	General with $\epsilon_{\text{bias}} = 0$
PDR-ANPG (**This Work**)	$\tilde{\mathcal{O}}(\epsilon^{-2}\min\{\epsilon^{-2}, \epsilon_{\text{bias}}^{-\frac{1}{3}}\})$	General with $\epsilon_{\text{bias}} > 0$
Lower Bound (Vaswani et al., 2022)	$\Omega(\epsilon^{-2})$	–

Table 1: List of recent papers on CMDPs with last-iterate guarantees. The term ϵ_{bias} is the expressivity error of the underlying parameterized policy class.

1.1 Contribution and Challenges

In this paper, we affirmatively answer the above question. In particular, we propose primal dual-based regularized accelerated natural policy gradient (PDR-ANPG) algorithm that uses entropy and quadratic regularizer in the Lagrangian function and momentum-based accelerated stochastic gradient descent (ASGD) process of Jain et al. (2018) as the NPG finding subroutine. We prove that, for general parameterization, our algorithm achieves $\mathcal{O}(\epsilon + (\epsilon_{\text{bias}})^{\frac{1}{6}})$ last-iterate optimality gap and the same constraint violation with a sample complexity of $\tilde{\mathcal{O}}(\epsilon^{-2}\min\{\epsilon^{-2}, \epsilon_{\text{bias}}^{-\frac{1}{3}}\})$ where ϵ_{bias} indicates the transferred compatibility approximation error of the policy class. If $\epsilon_{\text{bias}} > 0$ is independent of ϵ, the sample complexity reduces to $\tilde{\mathcal{O}}(\epsilon^{-2})$ for small ϵ. However, if $\epsilon_{\text{bias}} = 0$, i.e., the policy class is complete, then the optimality gap and constraint violation become $\mathcal{O}(\epsilon)$ and the sample complexity turns out to be $\tilde{\mathcal{O}}(\epsilon^{-4})$.

One of the main challenges in handling an entropy regularizer is that the advantage estimate can, in general, become unbounded. This is a problem since many intermediate lemmas crucial in establishing global convergence utilize its boundedness. In the tabular setup, Ding et al. (2024) circumvented this problem by allowing the policy optimization to run over a carefully chosen simplex. Such provisions are not available for general parameterized policies. Interestingly, we notice that the advantage estimates can be bounded in an average sense and that would be sufficient to arrive at our desired result, provided that the gradient[1] sampling is done in an unconventional way. In particular, on top of the standard routines, our sampling procedure (Algorithm 1) comprise an additional expectation that reduces the variance of the gradient estimate while preserving its unbiasedness.

Our improved sample complexity originates from another important observation. We noticed that the bias of the NPG estimator can be interpreted as the convergence error of an ASGD program with an exact gradient oracle. Without this reduction, the bias would be exponentially large, leading to a significant deterioration in sample complexity.

1.2 Related Works

Unconstrained MDP: Many algorithms are available in the literature that solve the unconstrained MDP via an exact gradient oracle e.g., see (Zhan et al., 2023; Lan, 2023; Cen et al., 2022; Agarwal et al., 2021; Bhandari & Russo, 2021). Among the sample-based methods, some papers demonstrate first-order convergence (Shen et al., 2019; Huang et al., 2020; Gargiani et al., 2022; Xu et al., 2020) while other works focus on global convergence (Masiha et al., 2022; Liu et al., 2020; Khodadadian et al., 2022; Chen & Maguluri, 2022). It is to be observed that Fatkhullin et al. (2023); Mondal & Aggarwal (2024b) prove the state-of-the-art $\tilde{\mathcal{O}}(\epsilon^{-2})$ sample complexity for last-iterate and average global convergence respectively. However, to the best of our understanding, the approach of Fatkhullin et al. (2023) is not extendable to CMDPs.

---

[1]Here we specifically refer to the gradients that are used in the NPG finding subroutine.

Constrained MDP: The majority of the CMDP works show average or regret-type constraint violation guarantees. Many among them design tabular model-based (He et al., 2021; Ding et al., 2021; Liu et al., 2021a; Efroni et al., 2020) and model-free (Wei et al., 2022; Bai et al., 2022; Ding et al., 2021) algorithms. Some works focus on parameterized policies as well. For example, Liu et al. (2021b); Zeng et al. (2022) deal with softmax policies while Xu et al. (2021); Bai et al. (2023); Ding et al. (2020) handle general parameterization. The optimal $\tilde{\mathcal{O}}(\epsilon^{-2})$ sample complexity for general parameterized CMDPs is proven by Mondal & Aggarwal (2024a). In comparison, the literature on last-iterate guarantees is relatively nascent. As exhibited in Table 1, Ying et al. (2022); Gladin et al. (2023) respectively establish $\tilde{\mathcal{O}}(\epsilon^{-2})$ and $\tilde{\mathcal{O}}(\epsilon^{-4})$ sample complexity for softmax policies whereas Ding et al. (2024) achieve $\tilde{\mathcal{O}}(\epsilon^{-6})$ sample complexity for the log-linear policy class assuming its expressivity error is $\epsilon_{\mathrm{bias}} = \mathcal{O}(\epsilon^8)$. Recently, Montenegro et al. (2024) showed $\tilde{\mathcal{O}}(\epsilon^{-7})$ sample complexity for general parameterization. We show an improvement over this state-of-the-art.

2 Formulation

Consider a Constrained Markov Decision Process (CMDP) characterized as $\mathcal{M} = (\mathcal{S}, \mathcal{A}, r, c, P, \gamma, \rho)$ where $\mathcal{S}$ is a (possibly infinite) state space, $\mathcal{A}$ is a finite action space with cardinality A, $r : \mathcal{S} \times \mathcal{A} \to [0, 1]$ indicates the reward function, $c : \mathcal{S} \times \mathcal{A} \to [-1, 1]$ is the cost function, $P : \mathcal{S} \times \mathcal{A} \to \Delta(\mathcal{S})$ is the transition function (where $\Delta(\cdot)$ denotes the probability simplex on its argument set), $\gamma \in [0, 1)$ defines the discount factor and $\rho \in \Delta(\mathcal{S})$ is the initial state distribution. This paper assumes $\mathcal{S}$ to be countable, though, in general, it can be taken to be compact. A (stationary) policy is defined to be a function of the form $\pi : \mathcal{S} \to \Delta(\mathcal{A})$. For a given policy, π, and a state-action pair (s, a), the Q value corresponding to a utility function $g : \mathcal{S} \times \mathcal{A} \to \mathbb{R}$ is defined as follows.

$$Q_g^\pi(s, a) = \mathbf{E}_\pi \left[\sum_{t=0}^\infty \gamma^t g(s_t, a_t) \middle| s_0 = s, a_0 = a \right] \tag{1}$$

where $\mathbf{E}_\pi$ is the expectation over all π-induced trajectories $\{(s_t, a_t)\}_{t=0}^\infty$ where $a_t \sim \pi(s_t)$, $s_{t+1} \sim P(s_t, a_t)$, $\forall t \in \{0, 1, \cdots\}$. Similarly, given a policy π and utility function g, the associated state value function is given as: $V_g^\pi(s) = \sum_a \pi(a|s) Q^\pi(s, a)$, $\forall s \in \mathcal{S}$.

Moreover, for each policy π, and utility function g, the the advantage function is computed as, $A_g^\pi(s, a) = Q_g^\pi(s, a) - V_g^\pi(s)$, $\forall (s, a)$. The state occupancy measure corresponding to π is given as

$$d^\pi(s) = (1 - \gamma) \mathrm{E}_\pi \left[\sum_{t=0}^\infty \gamma^t \mathbf{1}(s_t = s) \middle| s_0 \sim \rho \right], \ \forall s \tag{2}$$

where $\mathbf{1}(\cdot)$ is the indicator function. In this paper, we ignore the dependence on ρ for simplifying the notations whenever there are no confusions. The state-action occupancy measure induced by π is defined as $\nu^\pi(s, a) = d^\pi(s) \pi(a|s)$, $\forall (s, a)$. The expected value of $V_g^\pi(s)$ obtained over $s \sim \rho$ is denoted as J_g^π. The goal of learning CMDP is to solve the following optimization problem.

$$\max_{\pi \in \Pi} J_r^\pi \ \text{subject to:} \ J_c^\pi \geq 0 \tag{3}$$

where Π is the set of all policies. We assume that at least one interior solution exists for the above optimization, which is known as Slater's condition.

Assumption 1. *There exists $\bar{\pi} \in \Pi$ such that $J_c(\bar{\pi}) \geq c_{\mathrm{slat}}$ where $c_{\mathrm{slat}} \in (0, 1/(1 - \gamma)]$.*

Note that the policy function cannot be expressed in a tabular format for infinite states. General parameterization can be used in such cases. It indexes policies by a d-dimensional parameter θ. Let $J_g^{\pi_\theta} \triangleq J_g(\theta)$ for any $g \in \mathbb{R}^{\mathcal{S} \times \mathcal{A}}$. Problem equation 3 can now be written as:

$$\max_{\theta \in \mathbb{R}^d} J_r(\theta) \ \text{subject to:} \ J_c(\theta) \geq 0 \tag{4}$$

3 Algorithm Design

The standard approach to solve the constrained optimization equation 3 is utilizing a saddle point optimization on the Lagrangian function $J^\pi_{r+\lambda c}$ where λ is a Lagrange multiplier. We utilize π^* to denote an optimal solution to equation 3. Moreover, λ^* is its corresponding dual solution.

$$\lambda^* \in \arg\min_{\lambda \geq 0} \max_{\pi \in \Pi} J^\pi_{r+\lambda c} \tag{5}$$

The following result is well known (Ding et al., 2024).

Lemma 1. *An optimal primal-dual pair (π^*, λ^*) is guaranteed to exist if Assumption 1 holds. Moreover, it satisfies the following strong duality condition.*

$$\max_{\pi \in \Pi} J^\pi_{r+\lambda^* c} = J^{\pi^*}_{r+\lambda^* c} = \min_{\lambda \geq 0} J^{\pi^*}_{r+\lambda c} \tag{6}$$

Additionally, $0 \leq \lambda^* \leq 1/[(1-\gamma)c_{\mathrm{slat}}]$.

This paper, however, considers a regularized Lagrangian function, defined below $\forall \pi \in \Pi, \forall \lambda \geq 0$.

$$\mathcal{L}_\tau(\pi, \lambda) = J^\pi_{r+\lambda c} + \tau \left\{ \mathcal{H}(\pi) + \frac{1}{2}\lambda^2 \right\} \tag{7}$$

where τ is a tunable parameter and $\mathcal{H}(\pi)$ is the entropy corresponding to the policy π (defined below).

$$\mathcal{H}(\pi) \triangleq -\frac{1}{1-\gamma} \sum_{s,a} d^\pi(s)\pi(a|s)\log\pi(a|s)$$
$$= \mathbf{E}_\pi \left[\sum_{t=0}^\infty -\gamma^t \log\pi(a_t|s_t) \Big| s_0 \sim \rho \right] \tag{8}$$

Let $(\pi^*_\tau, \lambda^*_\tau)$ denote the primal-dual solutions corresponding to the regularized Lagrangian $\mathcal{L}_\tau$, i.e.,

$$\pi^*_\tau \triangleq \arg\max_{\pi \in \Pi} \min_{\lambda \in \Lambda} \mathcal{L}_\tau(\pi, \lambda),$$
$$\lambda^*_\tau \triangleq \arg\min_{\lambda \in \Lambda} \max_{\pi \in \Pi} \mathcal{L}_\tau(\pi, \lambda) \tag{9}$$

where $\Lambda = [0, \lambda_{\max}]$ is a carefully chosen set of positive reals and $\lambda_{\max}$ is stated later. It can be proven that for any $\tau, \lambda_{\max} > 0$, the tuple $(\pi^*_\tau, \lambda^*_\tau)$ uniquely exists and obeys a strong duality similar to Lemma 1[2]. However, such a result cannot be directly applied to the class of parameterized policies where we aim to solve equation 10 where $\mathcal{L}_\tau(\pi_\theta, \lambda)$ is denoted as $\mathcal{L}_\tau(\theta, \lambda)$.

$$\min_{\lambda \in \Lambda} \max_{\theta \in \mathbb{R}^d} \mathcal{L}_\tau(\theta, \lambda) \tag{10}$$

We aim to apply the Natural Policy Gradient (NPG)-based primal-dual updates (expressed below) to solve equation 10, starting with arbitrary θ_0 and λ_0.

$$\theta_{k+1} = \theta_k + \eta F(\theta_k)^\dagger \nabla_\theta \mathcal{L}_\tau(\theta_k, \lambda_k),$$
$$\lambda_{k+1} = \mathcal{P}_\Lambda \left[(1-\eta\tau)\lambda_k - \eta J_c(\theta_k) \right] \tag{11}$$

where $\mathcal{P}_\Lambda$ denotes the projection operation onto Λ and η is the learning rate. Note that, unlike the vanilla policy gradient iterations, the update direction of θ is not along the gradient $\nabla_\theta \mathcal{L}_\tau(\theta, \lambda)$ but rather, it is modulated by the Moore-Penrose pseudoinverse (symbolized as $\dagger$) of the Fisher matrix, $F(\theta)$ defined below.

$$F(\theta) = \mathbf{E}_{(s,a)\sim\nu^{\pi_\theta}} \left[\nabla_\theta \log\pi_\theta(a|s) \otimes \nabla_\theta \log\pi_\theta(a|s) \right] \tag{12}$$

where $\otimes$ denotes the outer product. The lemma stated below outlines a procedure to compute the gradient $\nabla_\theta \mathcal{L}_\tau(\theta, \lambda)$.

---

[2]More details are available in the appendix.

Lemma 2. *The following holds $\forall \theta \in \mathbb{R}^d$, $\forall \lambda \in \Lambda$.*

$$\nabla_\theta \mathcal{L}_\tau(\theta, \lambda) = \frac{1}{1-\gamma} H_\tau(\theta, \lambda), \tag{13}$$

$$H_\tau(\theta, \lambda) \triangleq \mathbf{E}_{(s,a)\sim\nu^{\pi_\theta}}\left[A_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}(s,a)\nabla_\theta \log \pi_\theta(a|s)\right] \tag{14}$$

where $\psi_\theta(s,a) = -\log \pi_\theta(a|s)$, $\forall(s,a)$.

The above result is similar to the well-known policy gradient theorem (Sutton et al., 1999). It is worthwhile to mention here that, since ψ_θ is unbounded, the advantage function $A_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}$ can, in general, be unbounded. This is quite disadvantageous since many intermediate results in the convergence analysis require the advantage function to be bounded. Fortunately, Lemma 2 (stated below) proves that the advantage values can be bounded in an average sense, which turns out to be sufficient for our analysis.

Lemma 3. *The following holds $\forall \theta, \lambda, \tau, s$.*

$$\mathbf{E}_{a\sim\pi_\theta(s)}\left[\left|A_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}(s,a)\right|^2\right] \leq L_{\tau,\lambda}^2 \ where$$

$$L_{\tau,\lambda}^2 \triangleq \frac{8(1+\lambda)^2}{(1-\gamma)^2} + \tau^2\left[\frac{32}{e^2}A + \frac{12(\log A)^2}{(1-\gamma)^2}\right] \tag{15}$$

Let $\omega_\tau^*(\theta, \lambda)$ denote the natural policy gradient (NPG), i.e., $\omega_\tau^*(\theta, \lambda) \triangleq F(\theta)^\dagger \nabla_\theta \mathcal{L}_\tau(\theta, \lambda)$. Define the following.

$$\mathcal{E}_\nu^\tau(\omega, \theta, \lambda) = \frac{1}{2}\mathbf{E}_{(s,a)\sim\nu}\left[\left(\omega^\mathrm{T}\nabla_\theta \log \pi_\theta(a|s)\right. \right.$$
$$\left.\left. -\frac{1}{1-\gamma}A_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}(s,a)\right)^2\right] \tag{16}$$

where $\nu \in \Delta(\mathcal{S} \times \mathcal{A})$ and $\omega \in \mathbb{R}^d$. Lemma 2 shows that $\omega_\tau^*(\theta, \lambda) = \arg\min_{\omega\in\mathbb{R}^d} \mathcal{E}_{\nu^{\pi_\theta}}^\tau(\omega, \theta, \lambda)$. This allows us to compute the NPG via a gradient-descent-based iterative algorithm. Observe that,

$$\nabla_\omega \mathcal{E}_{\nu^{\pi_\theta}}^\tau(\omega, \theta, \lambda) = F(\theta)\omega - \frac{1}{1-\gamma}H_\tau(\theta, \lambda) \tag{17}$$

Algorithm 1 Sampling Procedure

1: **Input:** $\theta, \omega, \lambda, \gamma, \tau, r, c, \rho$
2: **Define:** $g \triangleq r + \lambda c + \tau \psi_\theta$

3: $T \sim \text{Geo}(1 - \gamma), s_0 \sim \rho, a_0 \sim \pi_\theta(s_0)$
4: **for** $j \in \{0, \cdots, T-1\}$ **do**
5: $\quad s_{j+1} \sim P(s_j, a_j)$ and $a_{j+1} \sim \pi_\theta(s_{j+1})$
6: $\hat{J}_c(\theta) \leftarrow \sum_{j=0}^{T} c(s_j, a_j), \quad \hat{s} \leftarrow s_T$
7: /* Value Function Estimation /*
8: $T \sim \text{Geo}(1 - \gamma), s_0 \leftarrow \hat{s}, a_0 \sim \pi_\theta(s_0)$
9: **for** $j \in \{0, \cdots, T-1\}$ **do**
10: $\quad s_{j+1} \sim P(s_j, a_j)$, and $a_{j+1} \sim \pi_\theta(s_{j+1})$
11: $\hat{V}_g^{\pi_\theta}(\hat{s}) \leftarrow \sum_{j=0}^{T} g(s_j, a_j)$
12: /*Q and Advantage Estimation /*
13: **for** $a \in \mathcal{A}$ **do**
14: $\quad T \sim \text{Geo}(1 - \gamma), (s_0, a_0) \leftarrow (\hat{s}, a)$
15: $\quad$ **for** $j \in \{0, \cdots, T-1\}$ **do**
16: $\quad\quad s_{j+1} \sim P(s_j, a_j)$, and $a_{j+1} \sim \pi_\theta(s_{j+1})$
17: $\quad \hat{Q}_g^{\pi_\theta}(\hat{s}, a) \leftarrow \sum_{j=0}^{T} g(s_j, a_j)$
18: $\quad \hat{A}_g^{\pi_\theta}(\hat{s}, a) \leftarrow \hat{Q}_g^{\pi_\theta}(\hat{s}, a) - \hat{V}_g^{\pi_\theta}(\hat{s})$
19: /* Gradient Estimation /*

$$\hat{F}(\theta) \leftarrow \mathbf{E}_{a \sim \pi_\theta(\hat{s})} \left[\nabla_\theta \log \pi_\theta(a|\hat{s}) \otimes \nabla_\theta \log \pi_\theta(a|\hat{s}) \right]$$

$$\hat{H}_\tau(\theta, \lambda) \leftarrow \mathbf{E}_{a \sim \pi_\theta(\hat{s})} \left[\hat{A}_g^{\pi_\theta}(\hat{s}, a) \nabla_\theta \log \pi_\theta(a|\hat{s}) \right]$$

$$\hat{\nabla}_\omega \mathcal{E}_{\nu^{\pi_\theta}}^\tau (\omega, \theta, \lambda) \leftarrow \hat{F}(\theta)\omega - \frac{1}{1-\gamma} \hat{H}_\tau(\theta, \lambda) \tag{18}$$

20: **Output:** $\hat{J}_c(\theta), \hat{\nabla}_\omega \mathcal{E}_{\nu^{\pi_\theta}}^\tau (\omega, \theta, \lambda)$

Unfortunately, obtaining the exact gradient given by equation 17 might not be viable due to the lack of knowledge of the transition function P (which implies a lack of knowledge of the advantage function $A_{r+\lambda c+\tau \psi_\theta}^{\pi_\theta}$ and hence, that of $H_\tau(\theta, \lambda)$). It is also clear from equation 11 that the update of the dual variables requires an exact knowledge of $J_c(\theta_k)$, which is impossible to obtain without knowing P. Algorithm 1 provides sample-based estimates of the above quantities.

For a given set of parameters (θ, λ, τ), Algorithm 1 starts with $s_0 \sim \rho$ and rolls out a π_θ induced trajectory of length[3] $T \sim \text{Geo}(1 - \gamma)$. The total cost of this trajectory is assigned as $\hat{J}_c(\theta)$. The terminal state s_T can be shown to behave as a sample $\hat{s}$ taken from d^{π_θ}. Define $g \triangleq r + \lambda c + \tau \psi_\theta$ as a utility function. Another π_θ-induced trajectory of length $T \sim \text{Geo}(1 - \gamma)$ is rolled out next, taking $\hat{s} = s_T$ as the starting point. The total utility observed in this trajectory is assigned as the value estimate $\hat{V}_g^{\pi_\theta}(\hat{s})$. To obtain the Q estimates, for each $a \in \mathcal{A}$, a π_θ induced trajectory of length $T \sim \text{Geo}(1 - \gamma)$ is rolled out with $(\hat{s}, a)$ as the initial state-action. The total utilities observed in these trajectories are used as the estimates $\{\hat{Q}_g^{\pi_\theta}(\hat{s}, a)\}_{a \in \mathcal{A}}$. Its corresponding advantage values are estimated as $\hat{A}_g^{\pi_\theta}(\hat{s}, a) = \hat{Q}_g^{\pi_\theta}(\hat{s}, a) - \hat{V}_g^{\pi_\theta}(\hat{s}), \forall a$. Finally, the Fisher matrix estimate is obtained as the expected value of $\nabla_\theta \log \pi_\theta(a|\hat{s}) \otimes \nabla_\theta \log \pi_\theta(a|\hat{s})$ over $a \sim \pi_\theta(\hat{s})$, whereas the estimate of $H_\tau(\theta, \lambda)$ is computed as the expectation of $\hat{A}_g^{\pi_\theta}(\hat{s}, a)\nabla_\theta \log \pi_\theta(a|\hat{s})$, over $a \sim \pi_\theta(\hat{s})$. Using these, the gradient estimate is obtained via equation 18. Note that $\mathcal{O}((1 - \gamma)^{-1}A)$ samples are required on average to run one iteration of Algorithm 1.

Although our sampling procedure is similar to those used in earlier works (Liu et al., 2020; Mondal & Aggarwal, 2024b), there are some differences. In particular, the incorporation of the expectation over

---

[3]The symbol $\text{Geo}(\cdot)$ denotes the geometric distribution.

$a \sim \pi_\theta(\hat{s})$ to calculate $\hat{F}(\theta)$ and $\hat{H}_\tau(\theta, \lambda)$ is a new addition that was not present in the previous papers. Such a seemingly insignificant change has major ramifications for the global convergence result, as explained later in the paper. Lemma 4 (stated below) demonstrates that the estimates obtained from Algorithm 1 are unbiased.

Lemma 4. *The estimates obtained from Algorithm 1 are unbiased. Formally, for arbitrary $\omega, \theta, \lambda, \tau$, we get*

$$\mathbf{E}\left[\hat{J}_c(\theta)\big|\theta\right] = J_c(\theta), \ and \tag{19}$$

$$\mathbf{E}\left[\hat{\nabla}_\omega \mathcal{E}_{\nu^{\pi_\theta}}^\tau(\omega, \theta, \lambda)\big|\omega, \theta, \lambda\right] = \nabla_\omega \mathcal{E}_{\nu^{\pi_\theta}}^\tau(\omega, \theta, \lambda) \tag{20}$$

The estimates obtained from Algorithm 1 are used in Algorithm 2, which runs in nested loops. The *outer loop* executes K number of times and, at the kth instant, updates the current primal and dual parameters θ_k, λ_k via equation 26 and equation 27 respectively. The estimate $\hat{J}_c(\theta_k)$ is obtained via Algorithm 1. On the other hand, the direction ω_k, (which is used in the primal update), is computed in the inner loop by iteratively minimizing the function $\mathcal{E}_{\nu^{\pi_{\theta_k}}}^\tau(\cdot, \theta_k, \lambda_k)$ in H-steps using an Accelerated Stochastic Gradient Descent (ASGD) routine as stated in Jain et al. (2018). Each ASGD iteration comprises the updates equation 21−equation 24, followed by a tail averaging step equation 25 where the gradient estimates $\hat{G}_h$'s are yielded by Algorithm 1 and $(\alpha, \beta, \xi, \delta)$ are appropriate learning parameters.

Algorithm 2 Primal-Dual Regularized Accelerated Natural Policy Gradient (PDR-ANPG)

1: **Input:** Parameters $(\theta_0, \lambda_0, \tau, K, H)$, Distribution ρ, Learning Rates $\eta, \alpha, \beta, \xi, \delta$

2: /* Outer Loop /*

3: **for** $k \in \{0, \cdots, K-1\}$ **do**

4:     $\mathbf{x}_0, \mathbf{v}_0 \leftarrow \mathbf{0}$

5: /* Inner Loop /*

6:     **for** $h \in \{0, \cdots, H-1\}$ **do**

7:         /* Accelerated SGD /*

$$\mathbf{y}_h \leftarrow \alpha\mathbf{x}_h + (1-\alpha)\mathbf{v}_h \tag{21}$$

8:         $\hat{G}_h \leftarrow \hat{\nabla}_\omega \mathcal{E}_{\nu_\rho^{\pi_{\theta_k}}}^\tau(\omega, \theta_k, \lambda_k)\big|_{\omega=\mathbf{y}_h}$ (Algo. 1)

$$\mathbf{x}_{h+1} \leftarrow \mathbf{y}_h - \delta\hat{G}_h \tag{22}$$

$$\mathbf{z}_h \leftarrow \beta\mathbf{y}_h + (1-\beta)\mathbf{v}_h \tag{23}$$

$$\mathbf{v}_{h+1} \leftarrow \mathbf{z}_h - \xi\hat{G}_h \tag{24}$$

9:     Tail Averaging:

$$\omega_k \leftarrow \frac{2}{H} \sum_{\frac{H}{2} < h \leq H} \mathbf{x}_h \tag{25}$$

10:     Obtain $\hat{J}_{c,\rho}(\theta_k)$ via Algorithm 1.

11:     Parameter Updates:

$$\theta_{k+1} \leftarrow \theta_k + \eta\omega_k \tag{26}$$

$$\lambda_{k+1} \leftarrow \mathcal{P}_\Lambda[\lambda_k(1-\eta\tau) - \eta\hat{J}_c(\theta_k)] \tag{27}$$

12: **Output:** $\{(\theta_k, \lambda_k)\}_{k=1}^K$

---

4 Last-Iterate Convergence Analysis

This section shows the last-iterate convergence properties of Algorithm 2. Before presenting the technical results, we will outline a few key assumptions required for the analysis.

Assumption 2. *The policy is differentiable, and the score function is bounded and Lipschitz continuous, i.e., $\forall \theta, \theta_1, \theta_2 \in \mathbb{R}^d$ and $\forall (s, a)$,*

$$\|\nabla_\theta \log \pi_\theta(a|s)\| \leq G,$$
$$\|\nabla_\theta \log \pi_{\theta_1}(a|s) - \nabla_\theta \log \pi_{\theta_2}(a|s)\| \leq B\|\theta_1 - \theta_2\|$$

where $B, G > 0$ are constants.

Assumption 2 is common in the convergence analysis of policy gradient-based algorithms, e.g., see (Fatkhullin et al., 2023; Zhang et al., 2021b; Liu et al., 2020). This assumption is obeyed by many policy classes, such as neural networks with bounded parameters. The result stated below can be obtained by combining the above assumption with Lemma 2 and 3.

Lemma 5. *The following inequality holds $\forall \theta, \lambda, \tau$ where $L^2_{\tau,\lambda}$ is defined in equation 15.*

$$\|\nabla_\theta \mathcal{L}_\tau(\theta, \lambda)\|^2 \leq (1 - \gamma)^{-2} G^2 L^2_{\tau,\lambda} \tag{28}$$

Lemma 5 dictates that the norm of the gradient of the Lagrange function can be bounded above by some function of problem-dependent parameters. This result will be useful in the convergence analysis in the forthcoming section.

Assumption 3. *The approximation error function defined in equation 16 satisfies the following*

$$\mathcal{E}^\tau_{\nu^{\pi^*_\tau}}(\omega^*_\tau(\theta, \lambda), \theta, \lambda) \leq \frac{1}{2}\epsilon_{\text{bias}} \tag{29}$$

where $\theta \in \mathrm{R}^d$, $\lambda \in \Lambda$, $\tau \in [0, 1]$, and $\omega^*_\tau(\theta, \lambda)$ defines the NPG. The LHS of equation 29 is called the transferred compatibility approximation error.

The term ϵ_{bias} utilized in Assumption 3 can be interpreted as the expressivity error of the parameterized policy class. This term becomes zero for complete policy classes, e.g., softmax tabular policies Agarwal et al. (2021). One can also exhibit $\epsilon_{\text{bias}} = 0$ for linear MDPs (Yang & Wang, 2019; Jin et al., 2020). Moreover, for certain rich policy classes (e.g., wide neural networks), ϵ_{bias} turns out to be small (Wang et al., 2020).

Assumption 4. *There exists $\mu_F > 0$ such that $F(\theta) - \mu_F I_d$ is positive semidefinite, i.e., $F(\theta) \succeq \mu_F I_d$, where $F(\theta)$ is the Fisher matrix given in equation 12, I_d is a $d \times d$ the identity matrix, and $\theta \in \mathbb{R}^d$.*

Assumption 4 dictates that the Fisher matrix $F(\theta)$ is positive definite, and its eigenvalues are lower bounded by some positive constant μ_F. It is also easy to check that $F(\theta)$ is the Hessian of the function $\mathcal{E}^\tau_{\nu^{\pi_\theta}}(\cdot, \theta, \lambda)$. Assumption 4, hence, implies that the said function is μ_F-strongly convex, and its minimizer $\omega^*_\tau(\theta, \lambda)$ (the NPG) is uniquely defined. It is well-known in the optimization literature that we can approach the global minimum of a strongly convex function exponentially fast, given the availability of the exact gradient. Later, we shall see that such an insight is crucial in achieving our desired convergence rate. A policy class that obeys Assumption 4 is called a Fisher Non-Degenerate class. Such a restriction on the structure of the policy class is common in the literature (Fatkhullin et al., 2023; Zhang et al., 2021a; Bai et al., 2024).

4.1 Analysis of the Outer Loop

Let (θ_k, λ_k) denotes the parameters generated by Algorithm 2 after k iterations of the outer loop. Let

$$\Phi^\tau_k \triangleq \mathbf{E}\left[\mathrm{KL}^\tau_k\right] + \frac{1}{2}\mathbf{E}\left[(\lambda^*_\tau - \lambda_k)^2\right], \text{ where}$$
$$\mathrm{KL}^\tau_k \triangleq \sum_{s \in \mathcal{S}} d^{\pi^*_\tau}(s)\mathrm{KL}\left(\pi^*_\tau(\cdot|s)||\pi_{\theta_k}(\cdot|s)\right) \tag{30}$$

where $\mathrm{KL}(\cdot||\cdot)$ is the KL-divergence, and $\pi_\tau^*, \lambda_\tau^*$ are defined in equation 9. The expectations are computed over the distributions of (θ_k, λ_k). Intuitively, the term Φ_k^τ captures the distance between the solution provided by Algorithm 2 and the optimal regularized solution. Lemma 6 (stated below) establishes a recursive inequality obeyed by Φ_k^τ.

Lemma 6. *Let the parameters $\{\theta_k, \lambda_k\}_{k=0}^K$ be updated via equation 26, equation 27, and $\tau \in [0, 1]$. The following relation holds under assumptions 2 and 3, $\forall k \in \{0, 1, \cdots, K-1\}$.*

$$
\begin{aligned}
\Phi_{k+1}^\tau \leq{} & (1 - \eta\tau)\Phi_k^\tau + \eta\sqrt{\epsilon_{\mathrm{bias}}} \\
& + \eta G \mathbf{E} \left\| \mathbf{E}\left[\omega_k | \theta_k, \lambda_k\right] - \omega_{k,\tau}^* \right\| + \frac{B\eta^2}{2} \mathbf{E}\|\omega_k\|^2 \\
& + \eta^2 \left[\frac{2}{(1-\gamma)^2} + \tau^2 \lambda_{\max}^2 \right]
\end{aligned}
\tag{31}
$$

where $\omega_{k,\tau}^* \triangleq \omega_\tau^*(\theta_k, \lambda_k)$ is the exact NPG at (θ_k, λ_k).

Note the term ϵ_{bias} in equation 31. We later exhibit that its presence indicates that one cannot drive the optimality gap to zero due to the incompleteness of parameterized policies. Note that the term $\mathbf{E}\|\mathbf{E}[\omega_k|\theta_k, \lambda_k] - \omega_{k,\tau}^*\|$ is the expected bias of the NPG estimator. On the other hand,

$$
\begin{aligned}
\mathbf{E}\|\omega_k\|^2 &\leq 2\mathbf{E}\|\omega_k - \omega_{k,\tau}^*\|^2 + 2\|F(\theta_k)^\dagger \nabla_\theta \mathcal{L}_\tau(\theta_k, \lambda_k)\|^2 \\
&\stackrel{(a)}{\leq} 2\mathbf{E}\|\omega_k - \omega_{k,\tau}^*\|^2 + 2\mu_F^{-2}\|\nabla_\theta \mathcal{L}_\tau(\theta_k, \lambda_k)\|^2
\end{aligned}
\tag{32}
$$

where (a) follows from Assumption 4. Note that the second term in equation 32 can be bounded by Lemma 5. Next section provides bounds for both the first-order term $\mathbf{E}\|\mathbf{E}[\omega_k|\theta_k, \lambda_k] - \omega_{k,\tau}^*\|$ and the second-order term $\mathbf{E}\|\omega_k - \omega_{k,\tau}^*\|^2$ used in equation 31 and equation 32 respectively.

4.2 Analysis of the Inner Loop

We start with a statistical characterization of the noise of the gradient estimator given by Algorithm 1.

Lemma 7. *The estimate $\hat{\nabla}_\omega \mathcal{E}_{\nu^{\pi_\theta}}^\tau(\omega, \theta, \lambda)$ given by Algorithm 1 obeys the following semidefinite inequality if assumptions 2 and 4 holds.*

$$
\begin{aligned}
\mathbf{E}\left[\hat{\nabla}_\omega \mathcal{E}_{\nu^{\pi_\theta}}^\tau\left(\omega_\tau^*(\theta, \lambda), \theta, \lambda\right) \otimes \hat{\nabla}_\omega \mathcal{E}_{\nu^{\pi_\theta}}^\tau\left(\omega_\tau^*(\theta, \lambda), \theta, \lambda\right) \right] & \\
\preceq \sigma_{\tau,\lambda}^2 F(\theta)
\end{aligned}
\tag{33}
$$

where θ, λ, τ are arbitrary, $F(\theta)$ is the Fisher matrix defined in equation 12, and $\sigma_{\tau,\lambda}^2$ is given as follows.

$$
\sigma_{\tau,\lambda}^2 = \frac{48}{(1-\gamma)^4} \left[1 + \lambda^2 + 4Ae^{-2}\tau^2 \right] + \frac{2G^4 L_{\tau,\lambda}^2}{\mu_F^2(1-\gamma)^2}
\tag{34}
$$

where $L_{\tau,\lambda}^2$ is defined in equation 15.

The term $\sigma_{\tau,\lambda}^2$ can be interpreted as the scaled noise variance of the gradient estimator. Hence, if we get access to the exact gradient oracle, $\sigma_{\tau,\lambda}^2$ will be zero. This observation has been used in the subsequent lemma. Before describing that result, we would like to present some insights about the connection between Lemma 7 and the gradient estimation procedure (Algorithm 1). Define the following term.

$$
\begin{aligned}
\hat{\zeta}_{\theta,\lambda}^\tau(s, a) \triangleq{} & \nabla_\theta \log \pi_\theta(a|s) \cdot \omega_\tau^*(\theta, \lambda) \\
& - \frac{1}{1-\gamma} \hat{A}_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}(s, a), \ \forall(s, a)
\end{aligned}
\tag{35}
$$

where $\hat{A}_g^{\pi_\theta}(s,a)$, $g = r + \lambda c + \tau \psi_\theta$ is the advantage estimate yielded by Algorithm 1 for a given (s,a). Following equation 18, we can conclude that

$$\hat{\nabla}_\omega \mathcal{E}_{\nu^{\pi_\theta}}^\tau (\omega_\tau^*(\theta,\lambda),\theta,\lambda)$$
$$= \mathbf{E}_{a\sim\pi_\theta(\hat{s})} \left[\hat{\zeta}_{\theta,\lambda}^\tau(\hat{s},a) \nabla_\theta \log \pi_\theta(a|\hat{s}) \right] \tag{36}$$

where one can demonstrate that $\hat{s} \sim d^{\pi_\theta}$. Let $\hat{\mathbf{E}}_{s,a}^\theta$ be the expectation over the distribution of π_θ-induced trajectories that are used in evaluating the advantage term $\hat{A}_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}(s,a)$. Also, let $\hat{\mathbf{E}}_s^\theta$ be the expectation over the joint distribution of π_θ-induced trajectories that are used in estimating $\{\hat{A}_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}(s,a)\}_{a\in\mathcal{A}}$. Following the sampling process (Algorithm 1), one gets

$$\mathbf{E}_{a\sim\pi_\theta(s)} \hat{\mathbf{E}}_{s,a}^\theta \left[\left(\hat{\zeta}_{\theta,\lambda}^\tau(s,a) \right)^2 \right] \le \sigma_{\tau,\lambda}^2, \ \forall s \tag{37}$$

Note that the LHS of equation 33 obeys the following inequality.

$$\text{LHS} = \mathbf{E}_{s\sim d^{\pi_\theta}} \hat{\mathbf{E}}_s^\theta \left[\mathbf{E}_{a\sim\pi_\theta(s)} \left[\hat{\zeta}_{\theta,\lambda}^\tau(s,a) \nabla_\theta \log \pi_\theta(a|s) \right] \right.$$
$$\left. \otimes \mathbf{E}_{a\sim\pi_\theta(s)} \left[\hat{\zeta}_{\theta,\lambda}^\tau(s,a) \nabla_\theta \log \pi_\theta(a|s) \right] \right]$$
$$\overset{(a)}{\preceq} \mathbf{E}_{s\sim d^{\pi_\theta}} \left[\mathbf{E}_{a\sim\pi_\theta(s)} \left[\hat{\mathbf{E}}_{s,a}^\theta \left[\left(\hat{\zeta}_{\theta,\lambda}^\tau(s,a) \right)^2 \right] \right] \right.$$
$$\left. \times \mathbf{E}_{a\sim\pi_\theta(s)} \left[\nabla_\theta \log \pi_\theta(a|s) \otimes \nabla_\theta \log \pi_\theta(a|s) \right] \right]$$

where (a) is a result of the Cauchy-Schwarz inequality and interchange of expectation operations. Now, equation 33 can be proven by using equation 37 in the above relation. Notice the importance of the expectation over $a \sim \pi_\theta(\hat{s})$ in the sampling procedure. Had the gradient been estimated without this step (which is typically done in the literature, see our earlier discussion), the LHS of equation 33 would have been,

$$\text{LHS} = \mathbf{E}_{s\sim d^{\pi_\theta}} \mathbf{E}_{a\sim\pi_\theta(s)} \left[\underbrace{\hat{\mathbf{E}}_{s,a}^\theta \left[\left(\hat{\zeta}_{\theta,\lambda}^\tau(s,a) \right)^2 \right]}_{T_0(s,a)} \right.$$
$$\left. \times \nabla_\theta \log \pi_\theta(a|s) \otimes \nabla_\theta \log \pi_\theta(a|s) \right]$$

which is impossible to bound since $T_0(s,a)$ is not uniformly bounded $\forall (s,a)$ due to the presence of the term ψ_θ originating from the entropy regularizer.

Lemma 8. *If assumptions 2, 4 are true, the following holds $\forall k \in \{0,\cdots,K-1\}$ with $\alpha = \frac{3\sqrt{5}G^2}{\mu_F+3\sqrt{5}G^2}$, $\beta = \frac{\mu_F}{9G^2}$, $\xi = \frac{1}{3\sqrt{5}G^2}$, and $\delta = \frac{1}{5G^2}$ if the inner loop length of Algorithm 2 obeys $H > \max\left\{1, \bar{C}\frac{G^2}{\mu_F} \log\left(\sqrt{d}\frac{G^2}{\mu_F}\right)\right\}$ for some universal constant, $\bar{C}$.*

$$\mathbf{E}\|\omega_k - \omega_{k,\tau}^*\|^2 \le 22\frac{\sigma_{\tau,\lambda_{\max}}^2 d}{\mu_F} + \frac{CL_{\tau,\lambda_{\max}}^2}{\mu_F(1-\gamma)^2}, \tag{38}$$

$$\mathbf{E}\|\mathbf{E}\left[\omega_k|\theta_k,\lambda_k\right] - \omega_{k,\tau}^*\|$$
$$\le \sqrt{C} \exp\left(-\frac{\mu_F}{40G^2}H\right)\left[\frac{L_{\tau,\lambda_{\max}}}{\sqrt{\mu_F}(1-\gamma)}\right] \tag{39}$$

where C is a constant, ω_k is defined in equation 25, $\omega_{k,\tau}^*$ is the exact NPG, and $\sigma_{\tau,\lambda_{\max}}^2$ is given by equation 34.

Inequality equation 38 can be established utilizing Corollary 2 of Jain et al. (2018) in our framework and simplifying the upper bound. However, equation 39 follows from the same result applied to the scenario when

an exact gradient oracle is available. To understand the reasoning, note that one can write the following using conditional expectation on both sides of the update equations equation 21−equation 24, and Lemma 4.

$$
\begin{aligned}
\bar{\mathbf{y}}_h &= \alpha \bar{\mathbf{x}}_h + (1-\alpha)\bar{\mathbf{v}}_h, \\
\bar{\mathbf{x}}_{h+1} &= \bar{\mathbf{y}}_h - \delta \nabla_\omega \mathcal{E}_{\nu^{\pi_{\theta_k}}}^\tau (\omega, \theta_k, \lambda_k)\big|_{\omega = \bar{\mathbf{y}}_h} \\
\bar{\mathbf{z}}_h &= \beta \bar{\mathbf{y}}_h + (1-\beta)\bar{\mathbf{v}}_h, \\
\bar{\mathbf{v}}_{h+1} &= \bar{\mathbf{z}}_h - \xi \nabla_\omega \mathcal{E}_{\nu_\rho}^{\tau_{\pi_{\theta_k}}} (\omega, \theta_k, \lambda_k)\big|_{\omega = \bar{\mathbf{y}}_h}
\end{aligned}
\tag{40}
$$

where $\bar{l}_h = \mathbf{E}[l_h | \theta_k, \lambda_k], \forall h \in \{0, \cdots, H-1\}, l \in \{\mathbf{v}, \mathbf{x}, \mathbf{y}, \mathbf{z}\}$. Invoking equation 25, we get,

$$
\bar{\omega}_k \triangleq \mathbf{E}\left[\omega_k \big| \theta_k, \lambda_k\right] = \frac{2}{H} \sum_{\frac{H}{2} < h \leq H} \bar{\mathbf{x}}_h
\tag{41}
$$

Note that equation 40 and equation 41 emulate the steps of an ASGD process with exact (deterministic) gradients. We can now combine Lemma 6, equation 32, and Lemma 8 to arrive at the following corollary.

Corollary 1. *Let the parameters $\{\theta_k, \lambda_k\}_{k=0}^K$ be updated via equation 26, equation 27, and $\tau \in [0,1]$. The following relation holds $\forall k \in \{1, \cdots, K\}$ if assumptions 2−4 are true, the learning rates $(\alpha, \beta, \xi, \delta)$ are the same as in Lemma 8, $\eta\tau < 1$, and H is sufficiently large.*

$$
\begin{aligned}
\Phi_k^\tau \leq{}& \exp(-\eta\tau k)\Phi_0^\tau + \frac{1}{\tau}\sqrt{\epsilon_{\text{bias}}} \\
&+ \frac{1}{\tau}R_0 \exp\left(-\frac{\mu_F}{40G^2}H\right) + \frac{\eta}{\tau}(R_1 + R_2)
\end{aligned}
\tag{42}
$$

where R_0, R_1, R_2 are as follows.

$$
\begin{aligned}
R_0 &\triangleq \frac{G\sqrt{C}L_{1,\lambda_{\max}}}{\sqrt{\mu_F}(1-\gamma)}, \quad R_1 \triangleq \frac{2}{(1-\gamma)^2} + \lambda_{\max}^2, \\
R_2 &\triangleq \frac{B}{\mu_F}\left[22\sigma_{1,\lambda_{\max}}^2 \mathrm{d} + \frac{C + \mu_F^{-1}G^2}{(1-\gamma)^2}L_{1,\lambda_{\max}}^2\right]
\end{aligned}
$$

4.3 Optimality Gap and Constraint Violation

This section discusses how the optimality gap and constraint violation rates can be extracted from the convergence result of Φ_k^τ stated in Corollary 1.

Theorem 1. *Let $\{(\theta_k, \lambda_k)\}_{k=1}^K$ be the parameters generated by Algorithm 2 starting from (θ_0, λ_0). Assume that assumptions 1−4 hold, the parameters $(\alpha, \beta, \xi, \delta)$ are given by Lemma 8, $\lambda_{\max} = 4/[(1-\gamma)c_{\text{slat}}], \epsilon_{\text{bias}} < 1$, and $\epsilon < 1$ is sufficiently small. The relations stated below hold if $\tau = \max\{\epsilon, (\epsilon_{\text{bias}})^{\frac{1}{6}}\}, K = 2\epsilon^{-2}\tau^{-2}, \eta = \epsilon^2\tau$, and $H = 40G^2\mu_F^{-1}\log\left(\tau^{-1}\epsilon^{-2}\right)$.*

$$
\begin{aligned}
J_r^{\pi^*} - \mathbf{E}\left[J_r^{\pi_{\theta_K}}\right] &= \mathcal{O}(\epsilon + (\epsilon_{\text{bias}})^{\frac{1}{6}}), \quad and \\
\mathbf{E}\left[-J_c^{\pi_{\theta_K}}\right] &= \mathcal{O}(\epsilon + (\epsilon_{\text{bias}})^{\frac{1}{6}})
\end{aligned}
\tag{43}
$$

Theorem 1 essentially states that for any parameterized policy class with a transferred compatibility approximation error ϵ_{bias}, we can substitute $H = \tilde{O}(1), K = \mathcal{O}(\epsilon^{-2}\min\{\epsilon^{-2}, (\epsilon_{\text{bias}})^{-\frac{1}{3}}\})$ to ensure the (last-iterate) optimality gap and the constrained violation to be $\mathcal{O}(\epsilon + (\epsilon_{\text{bias}})^{\frac{1}{6}})$. This makes the overall sample complexity $\tilde{\mathcal{O}}(\epsilon^{-2}\min\{\epsilon^{-2}, (\epsilon_{\text{bias}})^{-\frac{1}{3}}\})$. The following remarks are worth highlighting.

Firstly, if $\epsilon_{\text{bias}} > 0$ and independent of ϵ, then the generated policies of Algorithm 2 cannot be guaranteed to be arbitrarily close to the optimal policy. This agrees with the existing literature on general parameterization (Liu et al., 2020; Fatkhullin et al., 2023; Mondal & Aggarwal, 2024b). In this case, we obtain the optimal

$\tilde{\mathcal{O}}(\epsilon^{-2})$ sample complexity for $\epsilon < (\epsilon_{\text{bias}})^{\frac{1}{6}}$. On the other hand, if $\epsilon_{\text{bias}} = 0$, i.e., the policy class is complete, the optimality gap and constraint violation can be made $\mathcal{O}(\epsilon)$ at the cost of $\tilde{\mathcal{O}}(\epsilon^{-4})$ sample complexity.

Secondly, some works have presumed ϵ_{bias} to depend on ϵ. For example, Ding et al. (2024) assumes $\epsilon_{\text{bias}} = \mathcal{O}(\epsilon^8)$ for log-linear classes to get a sample complexity of $\tilde{\mathcal{O}}(\epsilon^{-6})$ while enjoying $\mathcal{O}(\epsilon)$ optimality gap and constraint violation. If, following a similar path, one uses $\epsilon_{\text{bias}} = \Theta(\epsilon^6)$ in Theorem 1, the sample complexity becomes $\tilde{\mathcal{O}}(\epsilon^{-4})$ while the optimality gap and the constraint violation change to $\mathcal{O}(\epsilon)$.

Thirdly, although our result does not show zero constraint violation, it can be easily proven following the "conservative constraint" trick. The main idea is to use a conservative constraint $c' = c - (1 - \gamma)\delta'$ where δ' is a tunable parameter. Observe that $J_{c'}^\pi = J_c^\pi - \delta'$ for any arbitrary policy π. Therefore, once a $\mathcal{O}(\epsilon)$ constraint violation is proven for c' (utilizing the same algorithm described in our paper), one can judiciously choose δ' to prove zero constraint violation for c (see details in Ding et al. (2024); Bai et al. (2023)).

Fourthly, the choice of some of our parameters is dependent on ϵ_{bias}. We acknowledge that the knowledge of ϵ_{bias} may not be available for some parameterized policy classes. However, it might be possible to obtain an upper bound, $\bar{\epsilon}_{\text{bias}}$, of the desired quantity. For example, Wang et al. (2020) shows that for a two-layer neural parameterization of width m, the function approximation error is $\mathcal{O}(m^{-1/8})$ (see Theorem A.4 and the subsequent discussion). Our algorithm still works if this bound is used as a proxy of ϵ_{bias}. In this case, the convergence error and the sampling complexity will be functions of $\bar{\epsilon}_{\text{bias}}$, instead of ϵ_{bias}.

Finally, notice the importance of the observation that the NPG bias $\mathbf{E}[\omega_k | \theta_k, \lambda_k] - \omega_{k,\tau}^*$ is exactly the convergence error of an ASGD routine with exact gradients (see Lemma 8 and its subsequent discussion). Without this insight, the expected bias would be $\mathcal{O}(H^{-\frac{1}{2}})$, instead of its current bound $\exp(-\Theta(H))$ which would degrade the overall sample complexity by many folds.

5   Conclusions

We present an algorithm for the general parametrized CMDP that ensures $\mathcal{O}(\epsilon + \epsilon_{\text{bias}}^{\frac{1}{6}})$ last-iterate optimality gap and the same constraint violation with $\tilde{\mathcal{O}}(\epsilon^{-2} \min\{\epsilon^{-2}, (\epsilon_{\text{bias}})^{-\frac{1}{3}}\})$ sample complexity. Here ϵ_{bias} denotes the transferred compatibility approximation error of the underlying policy class. Future work includes improvement of the sample complexity to $\tilde{\mathcal{O}}(\epsilon^{-2})$ across all parameterized policy classes, extension of our ideas to general utility CMDPs, etc.

References

Alekh Agarwal, Sham M Kakade, Jason D Lee, and Gaurav Mahajan. On the theory of policy gradient methods: Optimality, approximation, and distribution shift. *The Journal of Machine Learning Research*, 22(1):4431–4506, 2021.

Qinbo Bai, Amrit Singh Bedi, Mridul Agarwal, Alec Koppel, and Vaneet Aggarwal. Achieving zero constraint violation for constrained reinforcement learning via primal-dual approach. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 36, pp. 3682–3689, 2022.

Qinbo Bai, Amrit Singh Bedi, and Vaneet Aggarwal. Achieving zero constraint violation for constrained reinforcement learning via conservative natural policy gradient primal-dual algorithm. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 37, pp. 6737–6744, 2023.

Qinbo Bai, Washim Uddin Mondal, and Vaneet Aggarwal. Regret analysis of policy gradient algorithm for infinite horizon average reward markov decision processes. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 38, pp. 10980–10988, 2024.

Jalaj Bhandari and Daniel Russo. On the linear convergence of policy gradient methods for finite mdps. In *International Conference on Artificial Intelligence and Statistics*, pp. 2386–2394. PMLR, 2021.

Shicong Cen, Chen Cheng, Yuxin Chen, Yuting Wei, and Yuejie Chi. Fast global convergence of natural policy gradient methods with entropy regularization. *Operations Research*, 70(4):2563–2578, 2022.

Zaiwei Chen and Siva Theja Maguluri. Sample complexity of policy-based methods under off-policy sampling and linear function approximation. In *International Conference on Artificial Intelligence and Statistics*, pp. 11195–11214. PMLR, 2022.

Dongsheng Ding, Kaiqing Zhang, Tamer Basar, and Mihailo Jovanovic. Natural policy gradient primal-dual method for constrained markov decision processes. *Advances in Neural Information Processing Systems*, 33:8378–8390, 2020.

Dongsheng Ding, Xiaohan Wei, Zhuoran Yang, Zhaoran Wang, and Mihailo Jovanovic. Provably efficient safe exploration via primal-dual policy optimization. In *International conference on artificial intelligence and statistics*, pp. 3304–3312. PMLR, 2021.

Dongsheng Ding, Chen-Yu Wei, Kaiqing Zhang, and Alejandro Ribeiro. Last-iterate convergent policy gradient primal-dual methods for constrained mdps. *Advances in Neural Information Processing Systems*, 36, 2024.

Yonathan Efroni, Shie Mannor, and Matteo Pirotta. Exploration-exploitation in constrained mdps. *arXiv preprint arXiv:2003.02189*, 2020.

Ilyas Fatkhullin, Anas Barakat, Anastasia Kireeva, and Niao He. Stochastic policy gradient methods: Improved sample complexity for fisher-non-degenerate policies. In *International Conference on Machine Learning*, pp. 9827–9869. PMLR, 2023.

Matilde Gargiani, Andrea Zanelli, Andrea Martinelli, Tyler Summers, and John Lygeros. Page-pg: A simple and loopless variance-reduced policy gradient method with probabilistic gradient estimation. In *International Conference on Machine Learning*, pp. 7223–7240. PMLR, 2022.

Egor Gladin, Maksim Lavrik-Karmazin, Karina Zainullina, Varvara Rudenko, Alexander Gasnikov, and Martin Takac. Algorithm for constrained markov decision process with linear convergence. In *International Conference on Artificial Intelligence and Statistics*, pp. 11506–11533. PMLR, 2023.

Jiafan He, Dongruo Zhou, and Quanquan Gu. Nearly minimax optimal reinforcement learning for discounted mdps. *Advances in Neural Information Processing Systems*, 34:22288–22300, 2021.

Feihu Huang, Shangqian Gao, Jian Pei, and Heng Huang. Momentum-based policy gradient methods. In *International conference on machine learning*, pp. 4422–4433. PMLR, 2020.

Prateek Jain, Sham M Kakade, Rahul Kidambi, Praneeth Netrapalli, and Aaron Sidford. Accelerating stochastic gradient descent for least squares regression. In *Conference On Learning Theory*, pp. 545–604. PMLR, 2018.

Chi Jin, Zhuoran Yang, Zhaoran Wang, and Michael I Jordan. Provably efficient reinforcement learning with linear function approximation. In *Proceedings of Thirty Third Conference on Learning Theory*, volume 125 of *Proceedings of Machine Learning Research*, pp. 2137–2143. PMLR, 09–12 Jul 2020.

Sajad Khodadadian, Thinh T Doan, Justin Romberg, and Siva Theja Maguluri. Finite-sample analysis of two-time-scale natural actor–critic algorithm. *IEEE Transactions on Automatic Control*, 68(6):3273–3284, 2022.

Guanghui Lan. Policy mirror descent for reinforcement learning: Linear convergence, new sampling complexity, and generalized problem classes. *Mathematical programming*, 198(1):1059–1106, 2023.

Tao Liu, Ruida Zhou, Dileep Kalathil, Panganamala Kumar, and Chao Tian. Learning policies with zero or bounded constraint violation for constrained mdps. *Advances in Neural Information Processing Systems*, 34:17183–17193, 2021a.

Tao Liu, Ruida Zhou, Dileep Kalathil, PR Kumar, and Chao Tian. Policy optimization for constrained mdps with provable fast global convergence. *arXiv preprint arXiv:2111.00552*, 2021b.

Yanli Liu, Kaiqing Zhang, Tamer Basar, and Wotao Yin. An improved analysis of (variance-reduced) policy gradient and natural policy gradient methods. *Advances in Neural Information Processing Systems*, 33: 7624–7636, 2020.

Saeed Masiha, Saber Salehkaleybar, Niao He, Negar Kiyavash, and Patrick Thiran. Stochastic second-order methods improve best-known sample complexity of sgd for gradient-dominated functions. *Advances in Neural Information Processing Systems*, 35:10862–10875, 2022.

Washim Mondal and Vaneet Aggarwal. Sample-efficient constrained reinforcement learning with general parameterization. *Advances in Neural Information Processing Systems*, 37:68380–68405, 2024a.

Washim U Mondal and Vaneet Aggarwal. Improved sample complexity analysis of natural policy gradient algorithm with general parameterization for infinite horizon discounted reward Markov decision processes. In *International Conference on Artificial Intelligence and Statistics*, pp. 3097–3105. PMLR, 2024b.

Alessandro Montenegro, Marco Mussi, Matteo Papini, and Alberto Maria Metelli. Last-iterate global convergence of policy gradients for constrained reinforcement learning. *arXiv preprint arXiv:2407.10775*, 2024.

Zebang Shen, Alejandro Ribeiro, Hamed Hassani, Hui Qian, and Chao Mi. Hessian aided policy gradient. In *International conference on machine learning*, pp. 5729–5738. PMLR, 2019.

Richard S Sutton, David McAllester, Satinder Singh, and Yishay Mansour. Policy gradient methods for reinforcement learning with function approximation. *Advances in neural information processing systems*, 12, 1999.

Sharan Vaswani, Lin Yang, and Csaba Szepesvári. Near-optimal sample complexity bounds for constrained mdps. *Advances in Neural Information Processing Systems*, 35:3110–3122, 2022.

Lingxiao Wang, Qi Cai, Zhuoran Yang, and Zhaoran Wang. Neural policy gradient methods: Global optimality and rates of convergence. In *International Conference on Learning Representations*, 2020. URL https://openreview.net/forum?id=BJgQfkSYDS.

Honghao Wei, Xin Liu, and Lei Ying. Triple-q: A model-free algorithm for constrained reinforcement learning with sublinear regret and zero constraint violation. In *International Conference on Artificial Intelligence and Statistics*, pp. 3274–3307. PMLR, 2022.

Pan Xu, Felicia Gao, and Quanquan Gu. An improved convergence analysis of stochastic variance-reduced policy gradient. In *Uncertainty in Artificial Intelligence*, pp. 541–551. PMLR, 2020.

Tengyu Xu, Yingbin Liang, and Guanghui Lan. Crpo: A new approach for safe reinforcement learning with convergence guarantee. In *International Conference on Machine Learning*, pp. 11480–11491, 2021.

Lin Yang and Mengdi Wang. Sample-optimal parametric q-learning using linearly additive features. In *International conference on machine learning*, pp. 6995–7004. PMLR, 2019.

Donghao Ying, Yuhao Ding, and Javad Lavaei. A dual approach to constrained markov decision processes with entropy regularization. In *International Conference on Artificial Intelligence and Statistics*, pp. 1887–1909. PMLR, 2022.

Sihan Zeng, Thinh T Doan, and Justin Romberg. Finite-time complexity of online primal-dual natural actor-critic algorithm for constrained markov decision processes. In *2022 IEEE 61st Conference on Decision and Control (CDC)*, pp. 4028–4033. IEEE, 2022.

Wenhao Zhan, Shicong Cen, Baihe Huang, Yuxin Chen, Jason D Lee, and Yuejie Chi. Policy mirror descent for regularized reinforcement learning: A generalized framework with linear convergence. *SIAM Journal on Optimization*, 33(2):1061–1091, 2023.

Junyu Zhang, Chengzhuo Ni, Csaba Szepesvari, Mengdi Wang, et al. On the convergence and sample efficiency of variance-reduced policy gradient method. *Advances in Neural Information Processing Systems*, 34:2228–2240, 2021a.

Junyu Zhang, Chengzhuo Ni, Zheng Yu, Csaba Szepesvari, and Mengdi Wang. On the convergence and sample efficiency of variance-reduced policy gradient method. *Advances in Neural Information Processing Systems*, 34:2228–2240, 2021b.

A Proof of Lemma 2

Fix a tuple (θ, λ, τ). The following proof assumes that the policy and all relevant value functions are differentiable w. r. t. θ. We also assume the state space, $\mathcal{S}$, to be finite to avoid technicalities associated with infinite space. Note that, $\forall s \in \mathcal{S}$, we have,

$$
\begin{aligned}
\nabla_\theta V^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s) &= \nabla_\theta \sum_{a \in \mathcal{A}} \pi_\theta(a|s) Q^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s,a) \\
&= \sum_{a \in \mathcal{A}} Q^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s,a) \nabla_\theta \pi_\theta(a|s) + \underbrace{\sum_{a \in \mathcal{A}} \pi_\theta(a|s) \nabla_\theta Q^{\pi_\theta}(s,a)}_{T_0}
\end{aligned}
\tag{44}
$$

Applying the Bellman equation, one obtains,

$$
\begin{aligned}
T_0 &= \sum_{a \in \mathcal{A}} \pi_\theta(a|s) \nabla_\theta \left\{ r(s,a) + \lambda c(s,a) - \tau \log \pi_\theta(a|s) + \gamma \sum_{s' \in \mathcal{S}} P(s'|s,a) V^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s') \right\} \\
&= -\tau \underbrace{\sum_{a \in \mathcal{A}} \pi_\theta(a|s) \nabla_\theta \log \pi_\theta(a|s)}_{=0} + \gamma \sum_{s' \in \mathcal{S}} \sum_{a \in \mathcal{A}} \pi_\theta(a|s) P(s'|s,a) \nabla_\theta V^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s') \\
&= \gamma \sum_{s' \in \mathcal{S}} \Pr\left(s_1 = s' \big| s_0 = s, \pi_\theta\right) \nabla_\theta V^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s')
\end{aligned}
\tag{45}
$$

Combining the above results, we obtain,

$$
\begin{aligned}
&\nabla_\theta V^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s) \\
&= \sum_{a \in \mathcal{A}} Q^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s,a) \nabla_\theta \pi_\theta(a|s) + \gamma \sum_{s' \in \mathcal{S}} \Pr\left(s_1 = s' \big| s_0 = s, \pi_\theta\right) \nabla_\theta V^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s') \\
&\overset{(a)}{=} \sum_{a \in \mathcal{A}} A^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s,a) \nabla_\theta \pi_\theta(a|s) + \gamma \sum_{s' \in \mathcal{S}} \Pr\left(s_1 = s' \big| s_0 = s, \pi_\theta\right) \nabla_\theta V^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s') \\
&\overset{(b)}{=} \sum_{s' \in \mathcal{S}} \sum_{t=0}^{T-1} \gamma^t \Pr(s_t = s'|s_0 = s, \pi_\theta) \sum_{a \in \mathcal{A}} A^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s,a) \nabla_\theta \pi_\theta(a|s) \\
&\qquad\qquad\qquad\qquad + \gamma^T \sum_{s' \in \mathcal{S}} \Pr(s_T = s'|s_0 = s, \pi_\theta) \nabla_\theta V^{\pi_\theta}(s')
\end{aligned}
\tag{46}
$$

where (a) uses the fact that $\sum_a \nabla_\theta \pi_\theta(a|s) = 0$, $\forall s \in \mathcal{S}$, and (b) is obtained by repeatedly applying the recursion. Notice that, for finite $\mathcal{S}$, we have the following limit.

$$
\gamma^T \sum_{s' \in \mathcal{S}} \Pr(s_T = s'|s_0 = s, \pi_\theta) \nabla_\theta V^{\pi_\theta}(s') \to \mathbf{0} \text{ as } T \to \infty
\tag{47}
$$

Finally, applying $T \to \infty$ to equation 46, we obtain the following,

$$
\begin{aligned}
\nabla_\theta \mathcal{L}_\tau(\theta, \lambda) &= \mathbf{E}_{s \sim \rho} \left[\nabla_\theta V^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s) \right] \\
&= \sum_{s' \in \mathcal{S}} \sum_{t=0}^{\infty} \gamma^t \Pr(s_t = s' | s_0 \sim \rho, \pi_\theta) \sum_{a \in \mathcal{A}} Q^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s, a) \nabla_\theta \pi_\theta(a|s) \\
&= \frac{1}{1-\gamma} \mathbf{E}_{(s,a) \sim \nu^{\pi_\theta}} \left[A^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s, a) \nabla_\theta \log \pi_\theta(a|s) \right]
\end{aligned}
\tag{48}
$$

This establishes the lemma. It is worth pointing out that, although our proof is similar to that of the standard policy gradient theorem Sutton et al. (1999), there are some differences. In particular, the standard proof assumes the reward/utility function to be independent of θ. This does not hold in our case due to the presence of ψ_θ.

B   Proof of Lemma 3

Fix a tuple $(\theta, \lambda, \tau, s)$. Since r and c are absolutely bounded in $[0, 1]$, we get,

$$
\left| A^{\pi_\theta}_{r+\lambda c}(s, a) \right| \le \left| Q^{\pi_\theta}_{r+\lambda c}(s, a) \right| + \left| V^{\pi_\theta}_{r+\lambda c}(s) \right| \le \frac{2(1+\lambda)}{1-\gamma}, \ \forall a
\tag{49}
$$

Furthermore, observe that,

$$
\begin{aligned}
0 \le V^{\pi_\theta}_{\psi_\theta}(s) &= \mathbf{E}_{\pi_\theta} \left[\sum_{t=0}^{\infty} -\gamma^t \log \pi_\theta(a_t|s_t) \bigg| s_0 = s \right] \\
&= -\frac{1}{1-\gamma} \sum_{s' \in \mathcal{S}} d^{\pi_\theta}_s(s') \sum_{a \in \mathcal{A}} \pi_\theta(a|s') \log \pi_\theta(a|s') \overset{(a)}{\le} \frac{1}{1-\gamma} \log A
\end{aligned}
\tag{50}
$$

where $d^{\pi_\theta}_s$ is defined similarly as d^{π_θ} given in equation 2 but with the conditional event $s_0 \sim \rho$ being replaced by $s_0 = s$. Inequality (a) follows from the concavity of the log function and the fact that $d^{\pi_\theta}_s \in \Delta(\mathcal{S})$. We deduce the following.

$$
\begin{aligned}
&\mathbf{E}_{a \sim \pi_\theta(s)} \left[\left| A^{\pi_\theta}_{\psi_\theta}(s, a) \right|^2 \right] \\
&\le 2\mathbf{E}_{a \sim \pi_\theta(s)} \left[\left| Q^{\pi_\theta}_{\psi_\theta}(s, a) \right|^2 \right] + 2 \left[V^{\pi_\theta}_{\psi_\theta}(s) \right]^2 \\
&\overset{(a)}{\le} 2\mathbf{E}_{a \sim \pi_\theta(s)} \left[\left| -\log \pi_\theta(a|s) + \gamma \mathbf{E}_{s' \sim P(s,a)} \left[V^{\pi_\theta}_{\psi_\theta}(s') \right] \right|^2 \right] + \frac{2(\log A)^2}{(1-\gamma)^2} \\
&\overset{(b)}{\le} 4 \sum_{a \in \mathcal{A}} \pi_\theta(a|s) \left[-\log \pi_\theta(a|s) \right]^2 + \frac{2(\log A)^2}{(1-\gamma)^2} \left[1 + 2\gamma^2 \right] \overset{(c)}{\le} \frac{16}{e^2} A + \frac{6(\log A)^2}{(1-\gamma)^2}
\end{aligned}
\tag{51}
$$

where (a) uses the Bellman equation and equation 50. Inequality (b) follows from the realization that equation 50 is satisfied by every $s \in \mathcal{S}$. Finally, (c) uses the fact that $x^2 \exp(-x) \le 4e^{-2}, \ \forall x \ge 0$. Combining equation 49 and equation 51, we arrive at,

$$
\begin{aligned}
\mathbf{E}_{a \sim \pi_\theta(s)} \left[\left| A^{\pi_\theta}_{r+\lambda c+\tau\psi_\theta}(s, a) \right|^2 \right] &\le 2\mathbf{E}_{a \sim \pi_\theta(s)} \left[\left| A^{\pi_\theta}_{r+\lambda c}(s, a) \right|^2 \right] + 2\tau^2 \mathbf{E}_{a \sim \pi_\theta(s)} \left[\left| A^{\pi_\theta}_{\psi_\theta}(s, a) \right|^2 \right] \\
&\le \frac{8(1+\lambda)^2}{(1-\gamma)^2} + \tau^2 \left[\frac{32}{e^2} A + \frac{12(\log A)^2}{(1-\gamma)^2} \right]
\end{aligned}
\tag{52}
$$

This concludes the lemma.

C Proof of Lemma 4

Fix a tuple $(\theta, \lambda, \tau, \omega)$. The first statement can be proven by observing that,

$$
\begin{aligned}
\mathbf{E}\left[\hat{J}_{c,\rho}(\theta)\big|\theta\right] &= (1-\gamma)\mathbf{E}\left[\sum_{t=0}^{\infty}\gamma^t\sum_{j=0}^{t}c(s_j, a_j)\bigg|s_0 \sim \rho, \pi_\theta\right] \\
&= (1-\gamma)\mathbf{E}\left[\sum_{j=0}^{\infty}c(s_j, a_j)\sum_{t=j}^{\infty}\gamma^t\bigg|s_0 \sim \rho, \pi_\theta\right] = \mathbf{E}\left[\sum_{j=0}^{\infty}\gamma^j c(s_j, a_j)\bigg|s_0 \sim \rho, \pi_\theta\right] = J_{c,\rho}(\theta)
\end{aligned}
\tag{53}
$$

To prove the second statement, we will first show that the state $\hat{s}$ generated by Algorithm 1 is indeed distributed according to the state-occupancy measure d^{π_θ}. Note that,

$$
\Pr(\hat{s} = s|\rho, \pi_\theta) = (1-\gamma)\sum_{t=0}^{\infty}\gamma^t\Pr(s_t = s|s_0 \sim \rho, \pi_\theta) = d^{\pi_\theta}(s)
\tag{54}
$$

Next, we will show that the Q and V function estimates obtained by Algorithm 1 for arbitrary utility function, g, are unbiased. Note the following chain of equalities for a given state-action pair (s, a).

$$
\begin{aligned}
\mathbf{E}\left[\hat{Q}_g^{\pi_\theta}(s, a)\big|\theta, s, a\right] &= (1-\gamma)\mathbf{E}\left[\sum_{t=0}^{\infty}\gamma^t\sum_{j=0}^{t}g(s_i, a_i)\bigg|s_0 = s, a_0 = a, \pi_\theta\right] \\
&= (1-\gamma)\mathbf{E}\left[\sum_{j=0}^{\infty}g(s_i, a_i)\sum_{t=j}^{t}\gamma^t\bigg|s_0 = s, a_0 = a, \pi_\theta\right] \\
&= \mathbf{E}\left[\sum_{j=0}^{\infty}\gamma^i g(s_i, a_i)\bigg|s_0 = s, a_0 = a, \pi_\theta\right] = Q_g^{\pi_\theta}(s, a)
\end{aligned}
\tag{55}
$$

Similarly, we can also demonstrate that, $\mathbf{E}[\hat{V}_g^{\pi_\theta}(s)\big|\theta, s] = V_g^{\pi_\theta}(s)$. It, hence, follows that $\mathbf{E}[\hat{A}_g^{\pi_\theta}(s, a)\big|\theta, s, a] = A_g^{\pi_\theta}(s, a)$ for each (s, a). Combining these results, we deduce the following.

$$
\mathbf{E}\left[\hat{F}(\theta)\big|\theta\right] = \mathbf{E}_{\hat{s}\sim d^{\pi_\theta}}\mathbf{E}_{a\sim\pi_\theta(\hat{s})}\left[\nabla_\theta\log\pi_\theta(a|\hat{s})\otimes\nabla_\theta\log\pi_\theta(a|\hat{s})\big|\theta\right] = F(\theta),
\tag{56}
$$

$$
\begin{aligned}
\mathbf{E}\left[\hat{H}_\tau(\theta, \lambda)\big|\theta, \lambda\right] &= \mathbf{E}_{\hat{s}\sim d^{\pi_\theta}}\mathbf{E}_{a\sim\pi_\theta(\hat{s})}\left[\mathbf{E}\left[\hat{A}_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}(\hat{s}, a)\bigg|\theta, \lambda, \hat{s}, a\right]\nabla_\theta\log\pi_\theta(a|\hat{s})\bigg|\theta, \lambda\right] \\
&= \mathbf{E}_{(\hat{s},a)\sim\nu_\rho^{\pi_\theta}}\left[A_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}(\hat{s}, a)\nabla_\theta\log\pi_\theta(a|\hat{s})\big|\theta, \lambda\right] = H_\tau(\theta, \lambda)
\end{aligned}
\tag{57}
$$

The unbiasedness of the gradient estimator can be proved by combining the above two results.

D Proof of Lemma 5

Recall the definition of $H_\tau(\theta, \lambda)$ in Lemma 2. Using the Cauchy-Schwarz inequality, we get,

$$
\begin{aligned}
\|H_\tau(\theta, \lambda)\|^2 &\leq \mathbf{E}_{(s,a)\sim\nu^{\pi_\theta}}\left[\left|A_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}(s, a)\right|^2\|\nabla_\theta\log\pi_\theta(a|s)\|^2\right] \\
&\overset{(a)}{\leq} G^2\mathbf{E}_{s\sim d^{\pi_\theta}}\mathbf{E}_{a\sim\pi_\theta(s)}\left[\left|A_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}(s, a)\right|^2\right]
\end{aligned}
$$

where (a) follows from Assumption 2. The lemma can now be concluded using Lemma 2 and 3.

E   Proof of Lemma 6

Step 1: We use the following notations: $\omega_{k,\tau}^* \triangleq \omega_\tau^*(\theta_k, \lambda_k)$, $\psi_k(s,a) \triangleq -\log \pi_{\theta_k}(a|s)$, and $\psi_\tau^*(s,a) \triangleq -\log \pi_\tau^*(a|s)$, $\forall(s,a)$. Observe the following chain of inequalities.

$$
\begin{aligned}
\mathrm{KL}_k^\tau - \mathrm{KL}_{k+1}^\tau &= \mathbf{E}_{(s,a)\sim\nu^{\pi_\tau^*}}\left[\log\frac{\pi_{\theta_{k+1}}(a|s)}{\pi_{\theta_k}(a|s)}\right] \\
&\overset{(a)}{\geq} \mathbf{E}_{(s,a)\sim\nu^{\pi_\tau^*}}[\nabla_\theta\log\pi_{\theta_k}(a|s)\cdot(\theta_{k+1}-\theta_k)] - \frac{B}{2}\|\theta_{k+1}-\theta_k\|^2 \\
&\overset{(b)}{=} \eta\mathbf{E}_{(s,a)\sim\nu^{\pi_\tau^*}}[\nabla_\theta\log\pi_{\theta_k}(a|s)\cdot\omega_k] - \frac{B\eta^2}{2}\|\omega_k\|^2 \\
&= \eta\mathbf{E}_{(s,a)\sim\nu^{\pi_\tau^*}}\left[\nabla_\theta\log\pi_{\theta_k}(a|s)\cdot\omega_{k,\tau}^*\right] + \eta\mathbf{E}_{(s,a)\sim\nu^{\pi_\tau^*}}[\nabla_\theta\log\pi_{\theta_k}(a|s)\cdot(\omega_k-\omega_{k,\tau}^*)] - \frac{B\eta^2}{2}\|\omega_k\|^2 \\
&\overset{(b)}{=} \eta\underbrace{\left[J^{\pi_\tau^*}_{r+\lambda_k c+\tau\psi_k} - J^{\pi_{\theta_k}}_{r+\lambda_k c+\tau\psi_k}\right]}_{T_0} + \eta\underbrace{\mathbf{E}_{(s,a)\sim\nu^{\pi_\tau^*}}\left[\nabla_\theta\log\pi_{\theta_k}(a|s)\cdot\omega_{k,\tau}^* - \frac{1}{1-\gamma}A^{\pi_{\theta_k}}_{r+\lambda_k c+\tau\psi_k}\right]}_{T_1} \\
&\quad + \eta\mathbf{E}_{(s,a)\sim\nu^{\pi_\tau^*}}[\nabla_\theta\log\pi_{\theta_k}(a|s)\cdot(\omega_k-\omega_{k,\tau}^*)] - \frac{B\eta^2}{2}\|\omega_k\|^2
\end{aligned}
\tag{58}
$$

where (a) follows from Assumption 2 and (b), (c) utilize equation 26 and Lemma 9 respectively. Using Assumption 3 and the Cauchy-Schwarz inequality, we derive the following.

$$
T_1 \geq -\left\{\mathbf{E}_{(s,a)\sim\nu^{\pi_\tau^*}}\left[\nabla_\theta\log\pi_{\theta_k}(a|s)\cdot\omega_{k,\tau}^* - \frac{1}{1-\gamma}A^{\pi_{\theta_k}}_{r+\lambda_k c+\tau\psi_k}\right]^2\right\}^{\frac{1}{2}} \geq -\sqrt{\epsilon_{\mathrm{bias}}}
\tag{59}
$$

Moreover, the term T_0 can be decomposed as follows,

$$
\begin{aligned}
T_0 &= \left[J^{\pi_\tau^*}_{r+\lambda_k c+\tau\psi_\tau^*} - J^{\pi_{\theta_k}}_{r+\lambda_k c+\tau\psi_k}\right] + \left[J^{\pi_\tau^*}_{r+\lambda_k c+\tau\psi_k} - J^{\pi_\tau^*}_{r+\lambda_k c+\tau\psi_\tau^*}\right] \\
&= J^{\pi_\tau^*}_{r+\lambda_k c+\tau\psi_\tau^*} - J^{\pi_{\theta_k}}_{r+\lambda_k c+\tau\psi_k} + \frac{\tau}{1-\gamma}\mathrm{KL}_k^\tau \\
&\geq J^{\pi_\tau^*}_{r+\lambda_k c+\tau\psi_\tau^*} - J^{\pi_{\theta_k}}_{r+\lambda_k c+\tau\psi_k} + \tau\mathrm{KL}_k^\tau
\end{aligned}
\tag{60}
$$

Combining the above results and taking expectations, we obtain,

$$
\begin{aligned}
\mathbf{E}\left[\mathrm{KL}_{k+1}^\tau\right] &\leq (1-\eta\tau)\mathbf{E}\left[\mathrm{KL}_k^\tau\right] + \eta\mathbf{E}\left[J^{\pi_{\theta_k}}_{r+\lambda_k c+\tau\psi_k} - J^{\pi_\tau^*}_{r+\lambda_k c+\tau\psi_\tau^*}\right] \\
&\quad + \eta\sqrt{\epsilon_{\mathrm{bias}}} + \eta\mathbf{E}_{(s,a)\sim\nu^{\pi_\tau^*}}\mathbf{E}\left[\nabla_\theta\log\pi_{\theta_k}(a|s)\cdot(\omega_{k,\tau}^* - \mathbf{E}\left[\omega_k|\theta_k,\lambda_k\right])\right] + \frac{B\eta^2}{2}\mathbf{E}\|\omega_k\|^2 \\
&\overset{(a)}{\leq} (1-\eta\tau)\mathbf{E}\left[\mathrm{KL}_k^\tau\right] + \eta\mathbf{E}\left[J^{\pi_{\theta_k}}_{r+\lambda_k c+\tau\psi_k} - J^{\pi_\tau^*}_{r+\lambda_k c+\tau\psi_\tau^*}\right] \\
&\quad + \eta\sqrt{\epsilon_{\mathrm{bias}}} + \eta G\mathbf{E}\left\|\omega_{k,\tau}^* - \mathbf{E}\left[\omega_k|\theta_k,\lambda_k\right]\right\| + \frac{B\eta^2}{2}\mathbf{E}\|\omega_k\|^2
\end{aligned}
\tag{61}
$$

where the inequality (a) is due to Assumption 2.

Step 2: Here, the dual update equation will be utilized. Observe the following.

$$
\begin{aligned}
\eta\mathbf{E}\left[\mathcal{L}_\tau(\theta_k,\lambda_k) - \mathcal{L}_\tau(\theta_k,\lambda_\tau^*)\right] &= \eta\mathbf{E}\left[(\lambda_k-\lambda_\tau^*)J_c(\theta_k) + \frac{\tau}{2}(\lambda_k)^2 - \frac{\tau}{2}(\lambda_\tau^*)^2\right] \\
&= \eta\mathbf{E}\left[(\lambda_k-\lambda_\tau^*)\left(J_c(\theta_k)+\tau\lambda_k\right)\right] - \frac{\eta\tau}{2}\mathbf{E}\left[(\lambda_k-\lambda_\tau^*)^2\right]
\end{aligned}
\tag{62}
$$

Using the dual update equation 27 and the non-expansiveness of the projection operator $\mathcal{P}_\Lambda$, we can write the following.

$$\frac{1}{2}\mathbf{E}\left[(\lambda_{k+1}-\lambda_\tau^*)^2\right] \leq \frac{1}{2}\mathbf{E}\left[\left\{\lambda_k-\lambda_\tau^*-\eta\left(\hat{J}_c(\theta_k)+\tau\lambda_k\right)\right\}^2\right]$$

$$= \frac{1}{2}\mathbf{E}\left[(\lambda_k-\lambda_\tau^*)^2\right] - \eta\mathbf{E}\left[(\lambda_k-\lambda_\tau^*)\left(\hat{J}_c(\theta_k)+\tau\lambda_k\right)\right] + \frac{\eta^2}{2}\mathbf{E}\left[\left(\hat{J}_c(\theta_k)+\tau\lambda_k\right)^2\right] \quad (63)$$

$$\overset{(a)}{\leq} \frac{1}{2}\mathbf{E}\left[(\lambda_k-\lambda_\tau^*)^2\right] - \eta\mathbf{E}\left[(\lambda_k-\lambda_\tau^*)(J_c(\theta_k)+\tau\lambda_k)\right] + \eta^2\mathbf{E}\left[\hat{J}_c^2(\theta_k)\right] + \eta^2\tau^2\lambda_{\max}^2$$

where (a) uses Lemma 4, the fact that λ_k and $\hat{J}_c(\theta_k)$ are conditionally independent given θ_k and $(a+b)^2 \leq 2(a^2+b^2)$ for any a, b. Furthermore, Algorithm 1 states that $\hat{J}_c^2(\theta_k)$ can take a value of at most $(j+1)^2$ with probability $(1-\gamma)\gamma^j$, $j \in \{0, 1, \cdots\}$. This leads to the following bound.

$$\mathbf{E}\left[\hat{J}_c^2(\theta_k)\right] \leq \sum_{j=0}^{\infty}(1-\gamma)(j+1)^2\gamma^j = \frac{1+\gamma}{(1-\gamma)^2} < \frac{2}{(1-\gamma)^2} \quad (64)$$

Combining the above results, we finally obtain,

$$\frac{1}{2}\mathbf{E}\left[(\lambda_{k+1}-\lambda_\tau^*)^2\right] \leq \frac{1}{2}(1-\eta\tau)\mathbf{E}\left[(\lambda_k-\lambda_\tau^*)^2\right] + \eta\mathbf{E}\left[\mathcal{L}_\tau(\theta_k,\lambda_\tau^*)-\mathcal{L}_\tau(\theta_k,\lambda_k)\right]$$

$$+ \eta^2\left[\frac{2}{(1-\gamma)^2}+\tau^2\lambda_{\max}^2\right] \quad (65)$$

Step 3: It is easy to verify that

$$\left[J_{r+\lambda_kc+\tau\psi_k}^{\pi_{\theta_k}} - J_{r+\lambda_kc+\tau\psi_\tau^*}^{\pi_\tau^*}\right] + \left[\mathcal{L}_\tau(\theta_k,\lambda_\tau^*)-\mathcal{L}_\tau(\theta_k,\lambda_k)\right] = \mathcal{L}_\tau(\theta_k,\lambda_\tau^*)-\mathcal{L}_\tau(\pi_\tau^*,\lambda_k) \quad (66)$$

Adding equation 61 and equation 65, and using the definition of Φ_k^τ, we obtain the following inequality.

$$\Phi_{k+1}^\tau \leq (1-\eta\tau)\Phi_k^\tau + \eta\sqrt{\epsilon_{\text{bias}}} + \eta G\mathbf{E}\left\|\omega_{k,\tau}^* - \mathbf{E}\left[\omega_k|\theta_k,\lambda_k\right]\right\| + \frac{B\eta^2}{2}\mathbf{E}\|\omega_k\|^2$$

$$+ \eta^2\left[\frac{2}{(1-\gamma)^2}+\tau^2\lambda_{\max}^2\right] + \eta\mathbf{E}\underbrace{\left[\mathcal{L}_\tau(\theta_k,\lambda_\tau^*)-\mathcal{L}_\tau(\pi_\tau^*,\lambda_k)\right]}_{T_2} \quad (67)$$

The lemma can be proven by showing that $T_2 \leq 0$ which is a direct consequence of Lemma 10. In particular,

$$\mathcal{L}_\tau(\pi_{\theta_k},\lambda_\tau^*) \leq \max_{\pi\in\Pi}\mathcal{L}_\tau(\pi,\lambda_\tau^*) = \min_{\lambda\geq0}\mathcal{L}_\tau(\pi_\tau^*,\lambda) \leq \mathcal{L}_\tau(\pi_\tau^*,\lambda_k) \quad (68)$$

F   Proof of Lemma 7

Define a function $\hat{\zeta}_{\theta,\lambda}^\tau : \mathcal{S} \times \mathcal{A} \to \mathbb{R}$ as follows.

$$\hat{\zeta}_{\theta,\lambda}^\tau(s,a) \triangleq \nabla_\theta\log\pi_\theta(a|s)\cdot\omega_\tau^*(\theta,\lambda) - \frac{1}{1-\gamma}\hat{A}_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}(s,a), \ \forall(s,a) \quad (69)$$

From equation 18, the gradient estimation can be written as follows.

$$\hat{\nabla}_\omega\mathcal{E}_{\nu^{\pi_\theta}}^\tau(\omega_\tau^*(\theta,\lambda),\theta,\lambda) = \mathbf{E}_{a\sim\pi_\theta(\hat{s})}\left[\hat{\zeta}_{\theta,\lambda}^\tau(\hat{s},a)\nabla_\theta\log\pi_\theta(a|\hat{s})\right] \quad (70)$$

where the distribution of $\hat{s}$ can be shown to be d^{π_θ} (see the proof of Lemma 4). Therefore,

$$
\begin{aligned}
&\mathbf{E}\left[\hat{\nabla}_\omega \mathcal{E}_\nu^{\tau_{\pi_\theta}}(\omega_\tau^*(\theta,\lambda),\theta,\lambda) \otimes \hat{\nabla}_\omega \mathcal{E}_\nu^{\tau_{\pi_\theta}}(\omega_\tau^*(\theta,\lambda),\theta,\lambda)\right] \\
&= \mathbf{E}_{s\sim d^{\pi_\theta}}\hat{\mathbf{E}}_s^\theta\left[\mathbf{E}_{a\sim\pi_\theta(s)}\left[\hat{\zeta}_{\theta,\lambda}^\tau(s,a)\nabla_\theta \log \pi_\theta(a|s)\right] \otimes \mathbf{E}_{a\sim\pi_\theta(s)}\left[\hat{\zeta}_{\theta,\lambda}^\tau(s,a)\nabla_\theta \log \pi_\theta(a|s)\right]\right] \\
&\preceq \mathbf{E}_{s\sim d^{\pi_\theta}}\left[\hat{\mathbf{E}}_s^\theta\left[\mathbf{E}_{a\sim\pi_\theta(s)}\left[\left(\hat{\zeta}_{\theta,\lambda}^\tau(s,a)\right)^2\right]\right] \mathbf{E}_{a\sim\pi_\theta(s)}\left[\nabla_\theta \log \pi_\theta(a|s) \otimes \log \pi_\theta(a|s)\right]\right] \\
&= \mathbf{E}_{s\sim d^{\pi_\theta}}\left[\mathbf{E}_{a\sim\pi_\theta(s)}\left[\hat{\mathbf{E}}_{s,a}^\theta\left[\left(\hat{\zeta}_{\theta,\lambda}^\tau(s,a)\right)^2\right]\right] \mathbf{E}_{a\sim\pi_\theta(s)}\left[\nabla_\theta \log \pi_\theta(a|s) \otimes \log \pi_\theta(a|s)\right]\right]
\end{aligned}
\tag{71}
$$

where the semidefinite inequality is a consequence of the Cauchy-Schwarz inequality and the expectation $\hat{\mathbf{E}}_s^\theta$ is computed over the distribution of all sample paths originating from s that are used to estimate $\{\hat{A}_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}(s,a)\}_{a\in\mathcal{A}}$. In a similar way, $\hat{\mathbf{E}}_{s,a}^\theta$ is defined as the expectation over the distribution of all sample paths that are used in estimating $\hat{A}_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}(s,a)$. Note that,

$$
\begin{aligned}
\left(\hat{\zeta}_{\theta,\lambda}^\tau(s,a)\right)^2 &\leq 2\|\nabla_\theta \log \pi_\theta(a|s)\|^2 \|\omega_\tau^*(\theta,\lambda)\|^2 + \frac{2}{(1-\gamma)^2}\left|\hat{A}_{r+\lambda c+\tau\psi_\theta}^{\pi_\theta}(s,a)\right|^2 \\
&\overset{(a)}{\leq} 2G^2\left\|F(\theta)^\dagger \nabla_\theta \mathcal{L}_\tau(\theta,\lambda)\right\|^2 + \frac{6}{(1-\gamma)^2}\left\{\left[\hat{A}_r^{\pi_\theta}(s,a)\right]^2 + \lambda^2\left[\hat{A}_c^{\pi_\theta}(s,a)\right]^2 + \tau^2\left[\hat{A}_{\psi_\theta}^{\pi_\theta}(s,a)\right]^2\right\} \\
&\overset{(b)}{\leq} \frac{2G^4}{\mu_F^2(1-\gamma)^2}L_{\tau,\lambda}^2 + \frac{6}{(1-\gamma)^2}\left\{\left[\hat{A}_r^{\pi_\theta}(s,a)\right]^2 + \lambda^2\left[\hat{A}_c^{\pi_\theta}(s,a)\right]^2 + \tau^2\left[\hat{A}_{\psi_\theta}^{\pi_\theta}(s,a)\right]^2\right\}
\end{aligned}
\tag{72}
$$

where (a) follows from Assumption 2 and the definition of $\omega_\tau^*(\theta,\lambda)$ whereas (b) results from Assumption 4 and Lemma 5. Note that, $\forall g \in \{r,c\}$, and arbitrary (s,a), both $[\hat{Q}_g^{\pi_\theta}(s,a)]^2$ and $[\hat{V}_g^{\pi_\theta}(s)]^2$ can assume a value of at most $(j+1)^2$ with probability $(1-\gamma)\gamma^j$, $j \in \{0,1,\cdots\}$ (see Algorithm 1). Therefore, the following holds $\forall (s,a)$.

$$
\begin{aligned}
\max_{g\in\{r,c\}}\left\{\hat{\mathbf{E}}_{s,a}^\theta\left[\left(\hat{A}_g^{\pi_\theta}(s,a)\right)^2\right]\right\} &\leq 2\max_{g\in\{r,c\}}\left\{\hat{\mathbf{E}}_{s,a}^\theta\left[\left(\hat{Q}_g^{\pi_\theta}(s,a)\right)^2\right] + \hat{\mathbf{E}}_{s,a}^\theta\left[\left(\hat{V}_g^{\pi_\theta}(s)\right)^2\right]\right\} \\
&\leq 4\max_{g\in\{r,c\}}\left\{\max\left\{\hat{\mathbf{E}}_{s,a}^\theta\left[\left(\hat{Q}_g^{\pi_\theta}(s,a)\right)^2\right], \hat{\mathbf{E}}_{s,a}^\theta\left[\left(\hat{V}_g^{\pi_\theta}(s)\right)^2\right]\right\}\right\} \\
&\leq 4(1-\gamma)\sum_{j=0}^\infty (j+1)^2\gamma^j = \frac{4(1+\gamma)}{(1-\gamma)^2} < \frac{8}{(1-\gamma)^2}
\end{aligned}
\tag{73}
$$

Moreover, observe that the following inequalities hold for arbitrary $s \in \mathcal{S}$.

$$
\begin{aligned}
\mathbf{E}_{a\sim\pi_\theta(s)}\hat{\mathbf{E}}_{s,a}^\theta\left[\left(\hat{Q}_{\psi_\theta}^{\pi_\theta}(s,a)\right)^2\right] &= (1-\gamma)\sum_{j=0}^\infty \gamma^j \sum_{a\in\mathcal{A}}\pi_\theta(a|s)\mathbf{E}_{j,s,a}^\theta\left[\left\{\sum_{t=0}^j -\log \pi_\theta(a_t|s_t)\right\}^2\right] \\
&\overset{(a)}{\leq} (1-\gamma)\sum_{j=0}^\infty (j+1)\gamma^j \underbrace{\sum_{a\in\mathcal{A}}\pi_\theta(a|s)\mathbf{E}_{j,s,a}^\theta\left[\sum_{t=0}^j (-\log \pi_\theta(a_t|s_t))^2\right]}_{T}
\end{aligned}
\tag{74}
$$

where $\mathbf{E}_{j,s,a}^\theta$ denotes the expectation over all $j+1$ length π_θ-induced trajectories $\{(s_t,a_t)\}_{t=0}^j$ that start with the initial condition $s_0 = s, a_0 = a$. The inequality is a consequence of Cauchy-Schwarz inequality. The term T can be bounded as follows.

$$
\begin{aligned}
T &= \sum_{a\in\mathcal{A}}\pi_\theta(a|s)\left[\left(-\log \pi_\theta(a|s)\right)^2 + \sum_{t=1}^j \sum_{s'\in\mathcal{S}}\Pr_t^{\pi_\theta}(s'|s,a)\sum_{a'\in\mathcal{A}}\pi_\theta(a'|s')\left(-\log \pi_\theta(a'|s')\right)^2\right] \\
&\overset{(a)}{\leq} \frac{4}{e^2}(j+1)A
\end{aligned}
\tag{75}
$$

where $\Pr_t^{\pi_\theta}(s'|s,a) = \Pr(s_t = s'|s_0 = s, a_0 = a, \pi_\theta)$. Inequality (a) follows from the fact that $x^2 \exp(-x) \leq 4e^{-2}, \forall x \geq 0$. Applying the above bound in equation 74, we deduce,

$$\mathbf{E}_{a\sim\pi_\theta(s)}\hat{\mathbf{E}}_{s,a}^\theta \left[\left(\hat{Q}_{\psi_\theta}^{\pi_\theta}(s,a)\right)^2\right] \leq \frac{4}{e^2}A(1-\gamma)\sum_{j=0}^\infty (j+1)^2\gamma^j < \frac{8Ae^{-2}}{(1-\gamma)^2} \tag{76}$$

In a similar manner, we can also show that,

$$\mathbf{E}_{a\sim\pi_\theta(s)}\hat{\mathbf{E}}_{s,a}^\theta \left[\left(\hat{V}_{\psi_\theta}^{\pi_\theta}(s)\right)^2\right] \leq \frac{8Ae^{-2}}{(1-\gamma)^2} \tag{77}$$

Combining the above results, one arrives at,

$$\mathbf{E}_{a\sim\pi_\theta(s)}\hat{\mathbf{E}}_{s,a}^\theta \left[\left(\hat{A}_{\psi_\theta}^{\pi_\theta}(s,a)\right)^2\right] \leq 2\mathbf{E}_{a\sim\pi_\theta(s)}\hat{\mathbf{E}}_{s,a}^\theta \left[\left(\hat{Q}_{\psi_\theta}^{\pi_\theta}(s,a)\right)^2\right] + 2\mathbf{E}_{a\sim\pi_\theta(s)}\hat{\mathbf{E}}_{s,a}^\theta \left[\left(\hat{V}_{\psi_\theta}^{\pi_\theta}(s)\right)^2\right]$$
$$\leq \frac{32Ae^{-2}}{(1-\gamma)^2} \tag{78}$$

Combining equation 72, equation 73, and equation 78, we can write,

$$\mathbf{E}_{a\sim\pi_\theta(s)}\left[\hat{\mathbf{E}}_{s,a}^\theta\left[\left(\hat{\zeta}_{\theta,\lambda}^\tau(s,a)\right)^2\right]\right] \leq \frac{2G^4}{\mu_F^2(1-\gamma)^2}L_{\tau,\lambda}^2 + \frac{48}{(1-\gamma)^4}\left[1+\lambda^2+4Ae^{-2}\tau^2\right] = \sigma_{\tau,\lambda}^2 \tag{79}$$

Substituting the above bound into equation 71, we conclude the lemma.

G Proof of Lemma 8

We prove Lemma 8 using Corollary 2 of Jain et al. (2018). Observe the following three statements for a given (λ, τ).

S1 : The following quantities exist and are finite $\forall\theta \in \mathbb{R}^{\mathrm{d}}$.

$$F(\theta) \triangleq \mathbf{E}_{(s,a)\sim\nu^{\pi_\theta}}\left[\nabla_\theta \log \pi_\theta(a|s) \otimes \nabla_\theta \log \pi_\theta(a|s)\right], \tag{80}$$

$$G(\theta) \triangleq \mathbf{E}_{(s,a)\sim\nu^{\pi_\theta}}\left[\nabla_\theta \log \pi_\theta(a|s) \otimes \nabla_\theta \log \pi_\theta(a|s) \otimes \nabla_\theta \log \pi_\theta(a|s) \otimes \nabla_\theta \log \pi_\theta(a|s)\right] \tag{81}$$

S2 : There exists $\sigma_{\tau,\lambda}^2$ such that the following is obeyed $\forall\theta \in \mathbb{R}^{\mathrm{d}}$ where $\omega_\tau^*(\theta,\lambda)$ is the minimizer of $\mathcal{E}_{\nu^{\pi_\theta}}^\tau(\cdot,\theta,\lambda)$.

$$\mathbf{E}\left[\hat{\nabla}_\omega \mathcal{E}_{\nu^{\pi_\theta}}^\tau(\omega_\tau^*(\theta,\lambda),\theta,\lambda) \otimes \hat{\nabla}_\omega \mathcal{E}_{\nu^{\pi_\theta}}^\tau(\omega_\tau^*(\theta,\lambda),\theta,\lambda)\right] \preccurlyeq \sigma_{\tau,\lambda}^2 F(\theta) \tag{82}$$

S3 : There exists $\mu_F, G > 0$ such that the following statements hold $\forall\theta \in \mathbb{R}^{\mathrm{d}}$.

$$(a)\ F(\theta) \succcurlyeq \mu_F I_{\mathrm{d}}, \tag{83}$$

$$(b)\ \mathbf{E}_{(s,a)\sim\nu^{\pi_\theta}}\left[\|\nabla_\theta \log \pi_\theta(a|s)\|^2 \nabla_\theta \log \pi_\theta(a|s) \otimes \nabla_\theta \log \pi_\theta(a|s)\right] \preccurlyeq G^2 F(\theta), \tag{84}$$

$$(c)\ \mathbf{E}_{(s,a)\sim\nu^{\pi_\theta}}\left[\|\nabla_\theta \log \pi_\theta(a|s)\|_{F(\theta)^\dagger}^2 \nabla_\theta \log \pi_\theta(a|s) \otimes \nabla_\theta \log \pi_\theta(a|s)\right] \preccurlyeq \frac{G^2}{\mu_F}F(\theta) \tag{85}$$

Statement **S1** is a consequence of Assumption 2 whereas **S2** results from Lemma 7. Statement **S3**(a) is identical to Assumption 4, **S3**(b) follows from Assumption 2, and finally, **S3**(c) can be deduced via Assumption 2 and 4. We can, hence, apply Corollary 2 of Jain et al. (2018) with $\kappa = \tilde{\kappa} = G^2/\mu_F$ and derive the following result if $H > \bar{C}\sqrt{\kappa\tilde{\kappa}}\log(\sqrt{\mathrm{d}}\sqrt{\kappa\tilde{\kappa}})$, the learning rates are set as $\alpha = \frac{3\sqrt{5}\sqrt{\kappa\tilde{\kappa}}}{1+3\sqrt{5\kappa\tilde{\kappa}}}$, $\beta = \frac{1}{9\sqrt{\kappa\tilde{\kappa}}}$, $\xi = \frac{1}{3\sqrt{5}\mu_F\sqrt{\kappa\tilde{\kappa}}}$, and $\delta = \frac{1}{5G^2}$, and $\lambda_k \in \Lambda$.

$$\mathbf{E}\left[l_k^\tau(\omega_k)\right] - l_k^\tau(\omega_{k,\tau}^*) \leq \frac{C}{2}\exp\left(-\frac{H}{20\sqrt{\kappa\tilde{\kappa}}}\right)\left[l_k^\tau(\mathbf{0}) - l_k^\tau(\omega_{k,\tau}^*)\right] + 11\frac{\sigma_{\tau,\lambda_{\max}}^2 \mathrm{d}}{H},$$
$$\text{where } l_k^\tau(\omega) \triangleq \mathcal{E}_{\nu^{\pi_{\theta_k}}}^\tau(\omega,\theta_k,\lambda_k), \ \forall\omega \in \mathbb{R}^{\mathrm{d}}, \ \text{and } \omega_{k,\tau}^* \triangleq \omega_\tau^*(\theta_k,\lambda_k) \tag{86}$$

In the above inequality, the expectation is over the distribution of ω_k for a given (θ_k, λ_k). The term C is a universal constant. Note that $l_k^\tau(\omega_{k,\tau}^*) \geq 0$ and $l_k^\tau(\mathbf{0})$ is bounded above as follows.

$$l_k^\tau(\mathbf{0}) = \frac{1}{2}\mathbf{E}_{(s,a)\sim\nu^{\pi_{\theta_k}}}\left[\frac{1}{1-\gamma}A_{r+\lambda_k c+\tau\psi_k}^{\pi_{\theta_k}}(s,a)\right]^2 \overset{(a)}{\leq} \frac{L_{\tau,\lambda_{\max}}^2}{2(1-\gamma)^2} \tag{87}$$

where (a) follows from Lemma 3. Combining *equation 86*, *equation 87*, and the fact that $l_k(\cdot)$ is μ_F-strongly convex, we establish,

$$\mathbf{E}\|\omega_k - \omega_k^*\|^2 \leq \frac{2}{\mu_F}\left[\mathbf{E}\left[l_k(\omega_k)\right] - l_k(\omega_k^*)\right] \leq 22\frac{\sigma_{\tau,\lambda_{\max}}^2\mathrm{d}}{\mu_F H} + C\exp\left(-\frac{\mu_F}{20G^2}H\right)\left[\frac{L_{\tau,\lambda_{\max}}^2}{\mu_F(1-\gamma)^2}\right] \tag{88}$$

The first statement can be proven by applying $H \geq 1$ and $\exp(-x) \leq 1$, $\forall x \geq 0$. We get the following for noiseless $(\sigma_{\tau,\lambda_{\max}}^2 = 0)$ gradient updates.

$$\mathbf{E}\|(\mathbf{E}[\omega_k|\theta_k] - \omega_k^*)\|^2 \leq C\exp\left(-\frac{\mu_F}{20G^2}H\right)\left[\frac{L_{\tau,\lambda_{\max}}^2}{\mu_F(1-\gamma)^2}\right] \tag{89}$$

The second statement can be established from *equation 89* by applying Jensen's inequality on the function $f(x) = x^2$.

H   Proof of Corollary 1

Lemma 6 establishes the following result $\forall k \in \{0, 1, \cdots, K-1\}$.

$$\Phi_{k+1}^\tau \leq (1-\eta\tau)\Phi_k^\tau + \eta\sqrt{\epsilon_{\mathrm{bias}}}$$
$$+ \underbrace{\eta G\mathbf{E}\left\|\mathbf{E}\left[\omega_k|\theta_k,\lambda_k\right] - \omega_{k,\tau}^*\right\| + \frac{B\eta^2}{2}\mathbf{E}\|\omega_k\|^2 + \eta^2\left[\frac{2}{(1-\gamma)^2} + \tau^2\lambda_{\max}^2\right]}_{T_k} \tag{90}$$

Using equation 32, and Lemma 8, we can bound the term T_k as follows.

$$T_k \leq \eta\exp\left(-\frac{\mu_F}{40G^2}H\right)\underbrace{\left[\frac{G\sqrt{C}L_{\tau,\lambda_{\max}}}{\sqrt{\mu_F}(1-\gamma)}\right]}_{\leq R_0}$$
$$+ \eta^2\left\{\underbrace{\frac{B}{\mu_F}\left[22\sigma_{\tau,\lambda_{\max}}^2\mathrm{d} + \frac{(C + \mu_F^{-1}G^2)L_{\tau,\lambda_{\max}}^2}{(1-\gamma)^2}\right]}_{\leq R_2} + \underbrace{\left[\frac{2}{(1-\gamma)^2} + \tau^2\lambda_{\max}^2\right]}_{\leq R_1}\right\} \tag{91}$$

where the upper bounds $\{R_l\}_{l\in\{0,1,2\}}$ follow from the fact that $\tau \in [0,1]$ and $L_{\tau,\lambda_{\max}}$, $\sigma_{\tau,\lambda_{\max}}$ are strictly increasing functions of τ. Therefore, we obtain the following simplified recursion $\forall k \in \{1, \cdots, K\}$.

$$\Phi_k^\tau \leq (1-\eta\tau)\Phi_{k-1}^\tau + \eta\sqrt{\epsilon_{\mathrm{bias}}} + \eta R_0\exp\left(-\frac{\mu_F}{40G^2}H\right) + \eta^2(R_1 + R_2)$$
$$\overset{(a)}{\leq} (1-\eta\tau)^k\Phi_0^\tau + \left[\sum_{l=0}^{k-1}(1-\eta\tau)^l\right] \times \left[\eta\sqrt{\epsilon_{\mathrm{bias}}} + \eta R_0\exp\left(-\frac{\mu_F}{40G^2}H\right) + \eta^2(R_1 + R_2)\right] \tag{92}$$
$$\overset{(b)}{\leq} \exp(-\eta\tau k)\Phi_0^\tau + \frac{1}{\tau}\sqrt{\epsilon_{\mathrm{bias}}} + \frac{1}{\tau}R_0\exp\left(-\frac{\mu_F}{40G^2}H\right) + \frac{\eta}{\tau}(R_1 + R_2)$$

where (a) is obtained by repeatedly applying the recursion and (b) uses the fact that $\sum_{l=0}^\infty(1-\eta\tau)^l = 1/\eta\tau$ and $(1-\eta\tau)^k \leq \exp(-\eta\tau k)$ whenever $\eta\tau < 1$. This establishes the corollary.

I Proof of Theorem 1

Observe the following chain of inequalities for a given $k \in \{0, \cdots, K\}$ and $g \in \{r, c\}$.

$$
\begin{aligned}
J_g^{\pi_\tau^*} - J_g^{\pi_{\theta_k}} &\overset{(a)}{=} \frac{1}{1-\gamma} \sum_{s \in \mathcal{S}} d^{\pi_\tau^*}(s) \sum_{a \in \mathcal{A}} \left\{ \pi_\tau^*(a|s) - \pi_{\theta_k}(a|s) \right\} Q_g^{\pi_{\theta_k}}(s,a) \\
&\leq \frac{1}{(1-\gamma)^2} \sum_{s \in \mathcal{S}} d^{\pi_\tau^*}(s) \| \pi_\tau^*(\cdot|s) - \pi_{\theta_k}(\cdot|s) \|_1 \\
&\overset{(b)}{\leq} \frac{1}{(1-\gamma)^2} \sum_{s \in \mathcal{S}} d^{\pi_\tau^*}(s) \sqrt{2 \mathrm{KL} \left(\pi_\tau^*(\cdot|s) \| \pi_{\theta_k}(\cdot|s) \right)} \\
&\overset{(c)}{\leq} \frac{1}{(1-\gamma)^2} \sqrt{2 \sum_{s \in \mathcal{S}} d^{\pi_\tau^*}(s) \mathrm{KL} \left(\pi_\tau^*(\cdot|s) \| \pi_{\theta_k}(\cdot|s) \right)} = \frac{1}{(1-\gamma)^2} \sqrt{2 \mathrm{KL}_k^\tau}
\end{aligned}
\tag{93}
$$

where (a), (b), and (c) are consequences of Lemma 9, the Pinkster's inequality, and the Cauchy-Schwarz inequality respectively. Taking expectation on both sides of the above inequality, we obtain,

$$
\begin{aligned}
J_g^{\pi_\tau^*} - \mathbf{E}\left[J_g^{\pi_{\theta_k}} \right] &\leq \frac{1}{(1-\gamma)^2} \mathbf{E}\left[\sqrt{2\mathrm{KL}_k^\tau} \right] \\
&\overset{(a)}{\leq} \frac{1}{(1-\gamma)^2} \sqrt{2\mathbf{E}\left[\mathrm{KL}_k^\tau \right]} \leq \frac{1}{(1-\gamma)^2} \sqrt{2\Phi_k^\tau} \\
&\overset{(b)}{\leq} \frac{\sqrt{2}}{(1-\gamma)^2} \left[\exp\left(-\frac{\eta\tau k}{2} \right) \sqrt{\Phi_0^\tau} + \frac{1}{\sqrt{\tau}} \left(\epsilon_{\mathrm{bias}} \right)^{\frac{1}{4}} + \sqrt{\frac{R_0}{\tau}} \exp\left(-\frac{\mu_F}{80G^2} H \right) + \sqrt{\frac{\eta}{\tau}} \left(\sqrt{R_1} + \sqrt{R_2} \right) \right]
\end{aligned}
\tag{94}
$$

where (a) is a result of Jensen's inequality applied to $f(x) = \sqrt{x}$, $x \geq 0$ while (b) follows from Corollary 1, and the fact that for the choice of η, τ mentioned in the theorem statement, $\tau \in [0,1]$ and $\eta\tau < 1$ for sufficiently small ϵ. Lemma 10 implies

$$
J_r^{\pi^*} - J_r^{\pi_\tau^*} \leq \tau \mathcal{H}(\pi_\tau^*) \leq \frac{\tau}{(1-\gamma)} \log A
\tag{95}
$$

Combining equation 94 and equation 95, one can obtain an upper bound on the optimality gap. To derive a similar bound for the constraint violation, observe that Lemma 10 implies that,

$$
-(\lambda_{\max} - \lambda_\tau^*) J_c^{\pi_\tau^*} \leq \frac{\tau}{2} \lambda_{\max}^2
\tag{96}
$$

Additionally, since $\lambda_\tau^* = \arg\min_{\lambda \in \Lambda} \mathcal{L}_\tau(\pi_\tau^*, \lambda)$, we have,

$$
\lambda_\tau^* = \arg\min_{\lambda \in \Lambda} \left\{ \lambda J_c^{\pi_\tau^*} + \frac{\tau}{2} \lambda^2 \right\}
\tag{97}
$$

Therefore, λ_τ^* can assume three possible values.

Case 1: If $0 < -\frac{J_c^{\pi_\tau^*}}{\tau} < \frac{\lambda_{\max}}{2}$, we have $\lambda_\tau^* = -\frac{J_c^{\pi_\tau^*}}{\tau}$, and consequently $\lambda_{\max} - \lambda_\tau^* > \frac{\lambda_{\max}}{2}$.

Case 2: If $-\frac{J_c^{\pi_\tau^*}}{\tau} \leq 0$, we get $\lambda_\tau^* = 0$ which ensures $\lambda_{\max} - \lambda_\tau^* = \lambda_{\max} > \frac{\lambda_{\max}}{2}$.

For cases 1 and 2, the following result can be derived using equation 96.

$$
-J_c^{\pi_\tau^*} \leq \tau \lambda_{\max}
\tag{98}
$$

Case 3: If $\frac{\lambda_{\max}}{2} \leq -\frac{J_c^{\pi_\tau^*}}{\tau}$, either $\lambda_\tau^* = -\frac{J_c^{\pi_\tau^*}}{\tau}$ or $\lambda_\tau^* = \lambda_{\max}$. In both cases, $\lambda_\tau^* \geq \frac{\lambda_{\max}}{2}$. Applying Lemma 10, we get,

$$
\lambda_\tau^*(J_c^{\pi^*} - J_c^{\pi_\tau^*}) \leq J_r^{\pi_\tau^*} - J_r^{\pi^*} + \tau \mathcal{H}(\pi_\tau^*)
\tag{99}
$$

Moreover, using Lemma 1, we have,

$$J_r^{\pi_\tau^*} - J_r^{\pi^*} \leq \lambda^*(J_c^{\pi^*} - J_c^{\pi_\tau^*}) \tag{100}$$

Combining equation 99 and equation 100, we arrive at,

$$(\lambda_\tau^* - \lambda^*)(J_c^{\pi^*} - J_c^{\pi_\tau^*}) \leq \tau\mathcal{H}(\pi_\tau^*) \tag{101}$$

Note that $\lambda_\tau^* \geq \frac{\lambda_{\max}}{2}$, and Lemma 1 dictates that $\lambda^* \leq \frac{\lambda_{\max}}{4}$ for the specific choice of $\lambda_{\max}$ given in the theorem statement. Clearly, $\lambda_\tau^* - \lambda^* \geq \frac{\lambda_{\max}}{4}$. Also, $J_c^{\pi^*} \geq 0$ due to feasibility of the optimal solution. Hence,

$$-J_c^{\pi_\tau^*} \leq \frac{4\tau\mathcal{H}(\pi_\tau^*)}{\lambda_{\max}} \leq \frac{\tau}{(1-\gamma)}\log A \tag{102}$$

where the last inequality follows from the fact that $4/\lambda_{\max} = (1-\gamma)c_{\text{slat}} \leq 1$. Combining equation 98 and equation 102, we obtain,

$$-J_c^{\pi_\tau^*} \leq \frac{\tau}{(1-\gamma)}\max\left\{\log A, 4c_{\text{slat}}^{-1}\right\} \tag{103}$$

I.1 Choice of Parameters

If we take $\tau = \max\{(\epsilon_{\text{bias}})^{\frac{1}{6}}, \epsilon\}$, the following bounds result from equation 95 and equation 103.

$$J_r^{\pi^*} - J_r^{\pi_\tau^*} = \mathcal{O}((\epsilon_{\text{bias}})^{\frac{1}{6}} + \epsilon), \text{ and } -J_c^{\pi_\tau^*} = \mathcal{O}((\epsilon_{\text{bias}})^{\frac{1}{6}} + \epsilon) \tag{104}$$

Moreover, if $\eta = \epsilon^2\tau$, $H = 40G^2\mu_F^{-1}\log(\tau^{-1}\epsilon^{-2})$, and $K = 2\tau^{-2}\epsilon^{-2}$, then equation 94 reduces to the following $\forall g \in \{r, c\}$.

$$J_g^{\pi_\tau^*} - \mathbf{E}[J_g^{\pi_{\theta_K}}] = \mathcal{O}(\epsilon + (\epsilon_{\text{bias}})^{\frac{1}{6}}) \tag{105}$$

To prove the above relation, we utilized the facts that (a) if $\epsilon < (\epsilon_{\text{bias}})^{\frac{1}{6}}$, we have $\tau = (\epsilon_{\text{bias}})^{\frac{1}{6}}$ and $(\epsilon_{\text{bias}})^{\frac{1}{4}}/\sqrt{\tau} = (\epsilon_{\text{bias}})^{\frac{1}{6}}$, and (b) if $\epsilon \geq (\epsilon_{\text{bias}})^{\frac{1}{6}}$, we have $\tau = \epsilon$, leading to $(\epsilon_{\text{bias}})^{\frac{1}{4}}/\sqrt{\tau} \leq \epsilon$. Note that in both of these cases, $(\epsilon_{\text{bias}})^{\frac{1}{4}}/\sqrt{\tau} \leq (\epsilon_{\text{bias}})^{\frac{1}{6}} + \epsilon$. Conjoining equation 104 and equation 105, we conclude the theorem. The criterion that H is sufficiently large is automatically satisfied if ϵ is sufficiently small.

J Auxiliary Lemmas

Lemma 9. *(Agarwal et al., 2021, Lemma 2) For any two policies π_1, π_2 and a utility $g \in \mathbb{R}^{\mathcal{S} \times \mathcal{A}}$, the following holds.*

$$J_g^{\pi_1} - J_g^{\pi_2} = \frac{1}{1-\gamma}\mathbf{E}_{(s,a)\sim\nu^{\pi_1}}\left[A_g^{\pi_2}(s,a)\right] = \frac{1}{1-\gamma}\sum_{s\in\mathcal{S}}d^{\pi_1}(s)\sum_{a\in\mathcal{A}}\left\{\pi_1(a|s) - \pi_2(a|s)\right\}Q_g^{\pi_2}(s,a) \tag{106}$$

Lemma 10. *(Ding et al., 2024, Lemma 14) The primal-dual pair $(\pi_\tau^*, \lambda_\tau^*)$ defined in equation 9 always exists and is unique for $\tau > 0$. Moreover, it satisfies the following property.*

$$\max_{\pi\in\Pi}\mathcal{L}_\tau(\pi, \lambda_\tau^*) = \mathcal{L}_\tau(\pi_\tau^*, \lambda_\tau^*) = \min_{\lambda\in\Lambda}\mathcal{L}_\tau(\pi_\tau^*, \lambda) \tag{107}$$

Consequently, the following inequalities hold $\forall\pi \in \Pi$ and $\forall\lambda \in \Lambda$.

$$J_{r+\lambda_\tau^* c}^{\pi} - \tau\mathcal{H}(\pi_\tau^*) \leq J_{r+\lambda_\tau^* c}^{\pi_\tau^*} \leq J_{r+\lambda c}^{\pi_\tau^*} + \frac{\tau}{2}\lambda^2 \tag{108}$$