On Dynamic Noise Influence in Differential Private Learning

Abstract

Protecting privacy in learning while maintaining the model performance has become increasingly critical in many applications that involve sensitive data. Private Gradient Descent (PGD) is a commonly used private learning framework, which adds noise according to the Differential Privacy protocol. Recent studies show that *dynamic privacy schedules* of decreasing noise magnitudes can improve loss at the final iteration, and yet theoretical understandings of the effectiveness of such schedules and their connections to optimization algorithms remain limited. In this paper, we provide comprehensive analysis of noise influence in dynamic privacy schedules to answer these critical questions. We first present a dynamic noise schedule minimizing the utility upper bound of PGD, and show how the noise influence from each optimization step collectively impacts utility of the final model. Our study also reveals how impacts from dynamic noise influence change when momentum is used. We empirically show the connection exists for general non-convex losses, and the influence is greatly impacted by the loss curvature.

1 Introduction

In the era of big data, privacy protection in machine learning systems is becoming a crucial topic as increasing personal data involved in training models (Dwork et al., 2020) and the presence of malicious attackers (Shokri et al., 2017; Fredrikson et al., 2015). In response to the growing demand, differential-private (DP) machine learning (Dwork et al., 2006) provides a computational framework for privacy protection and has been widely studied in various settings, including both convex and non-convex optimization (Wang et al., 2017; 2019; Jain et al., 2019).

One widely used procedure for privacy-preserving learning is the (Differentially) Private Gradient Descent (PGD) (Bassily et al., 2014; Abadi et al., 2016). A typical gradient descent procedure updates its model by the gradients of losses evaluated on the training data. When the data is sensitive, the gradients should be *privatized* to prevent excess privacy leakage. The PGD privatizes a gradient by adding controlled noise. As such, the models from PGD is expected to have a lower utility as compared to those from unprotected algorithms. In the cases where strict privacy control is exercised, or equivalently, a tight *privacy budget*, accumulating effects from highly-noised gradients may lead to unacceptable model performance. It is thus critical to design effective privatization procedures for PGD to maintain a great balance between utility and privacy.

Recent years witnessed a promising privatization direction that studies how to dynamically adjust the privacy-protecting noise during the learning process, i.e., *dynamic privacy schedules*, to boost utility under a specific privacy budget. One example is (Lee & Kifer, 2018), which reduced the noise magnitude when the loss does not decrease, due to the observation that the gradients become very small when approaching convergence, and a static noise scale will overwhelm these gradients. Another example is (Yu et al., 2019), which periodically decreased the magnitude following a predefined strategy, e.g., exponential decaying or step decaying. Both approaches confirmed the empirically advantages of decreasing noise magnitudes. Intuitively, the dynamic mechanism may coordinate with certain properties of the learning task, e.g., training data and loss surface. Yet there is no theoretical analysis available and two important questions remain unanswered: 1) *What is the form of utility-preferred noise schedules?* 2) *When and to what extent such schedules improve utility?*

To answer these questions, in this paper we develop a principled approach to construct dynamic schedules and quantify their utility bounds in different learning algorithms. Our contributions

Table 1: Comparison of utility upper bound using different privacy schedules. The algorithms are T-iteration $\frac{1}{2}R$-zCDP under the PL condition (unless marked with *). The O notation in this table drops other ln terms. Unless otherwise specified, all algorithms terminate at step $T = \mathcal{O}(\ln \frac{D}{N^2 R})$. Assume loss functions are 1-smooth and 1-Lipschitz continuous, and all parameters satisfy their numeric assumptions. Key notations: $\mathcal{O}_p$ – bound occurs in probability p; D – feature dimension; N – sample size; R – privacy budget; c_i – constant; other notations can be found in Section 4. An extended table and explanation are available in Appendix A.

Algorithm	Schedule (σ_t^2)	Utility Upper Bound
GD+MA (Wang et al., 2017)	$\mathcal{O}(\frac{T}{R_{\epsilon,\delta}})$	$\mathcal{O}\left(\frac{D \ln^2 N}{N^2 R_{\epsilon,\delta}}\right)$
*SSGD+zCDP (Feldman et al., 2020)	$\frac{1}{D}$	$\mathcal{O}\left(\left(\frac{1}{\sqrt{N}} + \frac{2\sqrt{D}}{\sqrt{R}N}\right)\ln N\right)$
Adam+MA (Zhou et al., 2020)	$\mathcal{O}(\frac{T}{R_{\epsilon,\delta}})$	$\mathcal{O}_p\left(\frac{\sqrt{D}\ln(ND\epsilon/(1-p))}{NR_{\epsilon,\delta}}\right)$
GD, Non-Private	0	$\mathcal{O}\left(\frac{D}{N^2 R}\right)$
GD+zCDP, Static Schedule	$\frac{T}{R}$	$\mathcal{O}\left(\frac{D \ln N}{N^2 R}\right)$
GD+zCDP, Dynamic Schedule	$\mathcal{O}\left(\frac{\gamma^{(t-T)/2}}{R}\right)$	$\mathcal{O}\left(\frac{D}{N^2 R}\right)$
Momentum+zCDP, Static Schedule	$\frac{T}{R}$	$\mathcal{O}\left(\frac{D}{N^2 R}(c + \ln N \mathbb{I}_{T>\hat{T}})\right)$
Momentum+zCDP, Dynamic Schedule	$\mathcal{O}\left(\frac{c_1\gamma^{T+t}+c_2\gamma^{(T-t)/2}}{R}\right)$	$\mathcal{O}\left(\frac{D}{N^2 R}(1 + \frac{cD}{N^2 R}\mathbb{I}_{T>\hat{T}})\right)$

are summarized as follows. 1) For the class of loss functions satisfying the Polyak-Lojasiewicz condition (Polyak, 1963), we show that a dynamic schedule improving the utility upper bound is shaped by the influence of per-iteration noise on the final loss. As the influence is tightly connected to the loss curvature, the advantage of using dynamic schedule depends on the loss function consequently. 2) Beyond gradient descent, our results show the gradient methods with momentum implicitly introduce a dynamic schedule and result in an improved utility bound. 3) We empirically validate our results on convex and non-convex (no need to satisfy the PL condition) loss functions. Our results suggest that the preferred dynamic schedule admits the exponentially decaying form, and works better when learning with high-curvature loss functions. Moreover, dynamic schedules give more utility under stricter privacy conditions (e.g., smaller sample size and less privacy budget).

2   RELATED WORK

Differentially Private Learning. Differential privacy (DP) characterizes the chance of an algorithm output (e.g., a learned model) to leak private information in its training data when the output distribution is known. Since outputs of many learning algorithms have undetermined distributions, the probability of their privacy leakages is hard to measure. A common approach to tackle this issue is to inject randomness with known probability distribution to *privatize* the learning procedures. Examples include output perturbation (Chaudhuri et al., 2011), objective perturbation (Chaudhuri et al., 2011) and gradient perturbation (Abadi et al., 2016; Bassily et al., 2014; Wu et al., 2017). Among these approaches, the Private Gradient Descent (PGD) has attracted extensive attention in recent years because it can be flexibly integrated with variants of gradient-based iteration methods, e.g., stochastic gradient descent, momentum methods (Qian, 1999), and Adam (Kingma & Ba, 2014), for both convex and non-convex problems.

Dynamic Policies for Privacy Protection. Wang et al. (2017) studied the empirical risk minimization using dynamic variation reduction of perturbed gradients. They showed that the utility upper bound can be achieved by gradient methods under uniform noise parameters. Instead of enhancing the gradients, Yu et al. (2019); Lee & Kifer (2018) showed the benefits of using a dynamic schedule of privacy parameters or equivalently noise scales. In addition, Pichapati et al. (2019); Thakkar et al. (2019) showed that the adaptive clipping norm, which controls sensitivity of batch gradients, similarly improves the convergence.

Utility Upper Bounds. A utility upper bound is a critical metric for privacy schedules that characterizes the maximum utility that a schedule can deliver in theory. Wang et al. (2017) is the first to prove the utility bound under the PL condition. In this paper, we improve the upper bound by a more accurate estimation of the dynamic influence of step noise. Remarkably, by introducing a dynamic schedule, we further boost the sample-efficiency of the upper bound. With a similar intuition,

Feldman et al. (2020) proposed to gradually increase the batch size, which reduces the dependence on sample size accordingly. Recently, Zhou et al. proved the utility bound by using the momentum of gradients (Polyak, 1964; Kingma & Ba, 2014). Table 1 summarizes the upper bounds of methods studied in this paper (in the last block of rows) and results from state-of-the-art algorithms based on private gradients. Our work shows that considering the dynamic influence can lead to a tighter bound.

3 PRIVATE GRADIENT DESCENT

Notations. We consider a learning task by empirical risk minimization (ERM) $f(\theta) = \frac{1}{N}\sum_{n=1}^{N} f(\theta; x_n)$ on a private dataset $\{x_n\}_{n=1}^{N}$ and $\theta \in \mathbb{R}^D$. The gradient methods are defined as $\theta_{t+1} = \theta_t - \eta_t \nabla_t$, where $\nabla_t = \nabla f(\theta_t) = \frac{1}{N}\sum_n \nabla f(\theta_t; x_n)$ denotes the non-private gradient at iteration t, η_t is the step learning rate. $\nabla_t^{(n)} = \nabla f(\theta_t; x_n)$ denotes the gradient on a sample x_n. $\mathbb{I}_c$ denotes the indicator function that returns 1 if the condition c holds, otherwise 0.

Assumptions. (1) In this paper, we assume $f(\theta)$ is continuous and differentiable. Many commonly used loss functions satisfy this assumption, e.g., the logistic function. (2) For a learning task, only finite amount of privacy cost is allowed where the maximum cost is called *privacy budget* and denoted as R. (3) Generally, we assume that loss functions $f(\theta; x)$ (sample-wise loss) are G-Lipschitz continuous and $f(\theta)$ (the empirical loss) is M-smooth.

Definition 3.1 (*G*-Lipschitz continuity). A function $f(\cdot)$ is G-Lipschitz continuous if, for $G > 0$ and all x, y in the domain of $f(\cdot)$, $f(\cdot)$ satisfies $\|f(y) - f(x)\| \le G\|y - x\|^2$. .

Definition 3.2 (*m*-strongly convexity). A function $f(\cdot)$ is m-strongly convex if $f(y) \ge f(x) + \nabla f(x)^T(y - x) + \frac{m}{2}\|y - x\|^2$, for some $m > 0$ and all x, y in the domain of $f(\cdot)$.

Definition 3.3 (*M*-smoothness). A function is M-smooth w.r.t. l_2 norm if $f(y) \le f(x) + \nabla f(x)^T(y - x) + \frac{M}{2}\|y - x\|^2$, for some constant $M > 0$ and all x, y in the domain of $f(\cdot)$.

For a private algorithm $\mathcal{M}(d)$ which maps a dataset d to some output, the privacy cost is measured by the bound of the output difference on the adjacent datasets. The adjacent datasets are defined to be datasets that only differ in one sample. In this paper, we use the zero-Concentrated Differential Privacy (zCDP, see Definition 3.4) as the privacy measurement, because it provides the simplicity and possibility of adaptively composing privacy costs at each iteration. Various privacy metrics are discussed or reviewed in (Mironov, 2017; Bun et al., 2018; Abadi et al., 2016; Desfontaines & Pejó, 2019). An example is Moment Accoutant (Abadi et al., 2016), which has a tighter privacy bound but is more complicated in computing privacy cost due to the minimization problem involved. We note that alternative metrics can be adapted to our study without major impacts to the analysis.

Definition 3.4 (*ρ*-zCDP (Bun & Steinke, 2016)). Let $\rho > 0$. A randomized algorithm $\mathcal{M} : \mathcal{D}^n \to \mathbb{R}$ satisfies ρ-zCDP if, for all adjacent datasets $d, d' \in \mathcal{D}^n$, $D_\alpha(\mathcal{M}(d)\|\mathcal{M}(d')) \le \rho\alpha$, $\forall \alpha \in (1, \infty)$ where $D_\alpha(\cdot\|\cdot)$ denotes the Rényi divergence (Rényi, 1961) of order α.

zCDP provides a linear composition of privacy costs of sub-route algorithms. When the input vector is privatized by injecting Gaussian noise of $\mathcal{N}(0, \sigma_t^2 I)$ for the t-th iteration, the composed privacy cost is proportional to $\sum_t \rho_t$ where the step cost is $\rho_t = \frac{1}{\sigma_t^2}$. For simplicity, we absorb the constant coefficient into the (residual) *privacy budget* R. The formal theorems for the privacy cost computation of composition and Gaussian noising is included in Lemmas B.1 and B.2.

Generally, we define the Private Gradient Descent (PGD) method as iterations for $t = 1 \ldots T$:

$$\theta_{t+1} = \theta_t - \eta_t v_t = \theta_t - \eta_t(\tilde{\nabla}_t + \sigma_t G \nu_t / N), \tag{1}$$

where $v_t = g_t$ is the gradient privatized from ∇_t as shown in Algorithm 1, G/N is the bound of sensitivity of the gathered gradient excluding one sample gradient, and $\nu_t \sim \mathcal{N}(0, I)$ is a vector element-wisely subject to Gaussian distribution. We use σ_t to denote the noise scale at step t and use σ to collectively represents the schedule $(\sigma_1, \ldots, \sigma_T)$ if not confusing. When the Lipschitz constant is unknown, we can control the upper bound by scaling the gradient if it is over some constant. The scaling operation is often called *clipping* in literatures since it clips the gradient norm at a threshold. After the gradient is noised, we apply a modification, $\phi(\cdot)$, to enhance its utility. In this paper, we consider two types of $\phi(\cdot)$:

$$\phi(m_t, g_t) = g_t \text{ (GD)}, \quad \phi(m_t, g_t) = [\beta(1 - \beta^{t-1})m_t + (1 - \beta)g_t]/(1 - \beta^t) \text{ (Momentum)}$$

Algorithm 1 Privatizing Gradients

Input: Raw gradients $[\nabla_t^{(1)}, \ldots, \nabla_t^{(n)}]$, v_t, residual privacy budget R_t assuming the full budget is R and $R_1 = R$.

1: $\rho_t \leftarrow 1/\sigma_t^2$ ▷ Budget request
2: **if** $\rho_t < R_t$ **then**
3:      $R_{t+1} \leftarrow R_t - \rho_t$
4:      $g_t \leftarrow \tilde{\nabla}_t + G\sigma_t\nu_t/N, \nu_t \sim \mathcal{N}(0, I)$ ▷ Privacy noise
5:      $m_{t+1} \leftarrow \phi(m_t, g_t)$ or g_1 if $t = 1$
6:      **return** $\eta_t m_{t+1}, R_{t+1}$ ▷ Utility projection
7: **else**
8:      Terminate

We now show that the PGD using Algorithm 1 guarantees a privacy cost less than R:

Theorem 3.1. *Suppose $f(\theta; x)$ is G-Lipschitz continuous and the PGD algorithm with privatized gradients defined by Algorithm 1, stops at step T. The PGD algorithm outputs θ_T and satisfies ρ-zCDP where $\rho \leq \frac{1}{2}R$.*

Note that Theorem 3.1 allows σ_t to be different throughout iterations. Next we present a principled approach for deriving dynamic schedules optimized for the final loss $f(\theta_T)$.

4 DYNAMIC POLICIES BY MINIMIZING UTILITY UPPER BOUNDS

To characterize the utility of the PGD, we adopt the Expected Excess Risk (EER), which notion is widely used for analyzing the convergence of random algorithms, e.g., (Bassily et al., 2014; Wang et al., 2017). Due to the presence of the noise and the limitation of learning iterations, optimization using private gradients is expected to reach a point with a higher loss (i.e., excess risk) as compared to the optimal solution without private protection. Define $\theta^* = \arg\min_\theta f(\theta)$, after Algorithm 1 is iterated for T times in total, the EER gives the expected utility degradation:

$$\text{EER} = \mathbb{E}_\nu[f(\theta_{T+1})] - f(\theta^*).$$

Due to the variety of loss function and complexity of recursive iterations, an exact EER with noise is intractable for most functions. Instead, we study the worst case scenario, i.e., the upper bound of the EER, and our goal is to minimize the upper bound. For consistency, we call the upper bound of EER divided by the initial error as ERUB, and $\text{ERUB} = \gamma^T + \sum_{t=1}^T \gamma^{T-t} R\alpha\sigma_t^2$ for the GD. Since the analytical form of EER is either intractable or complicated due to the recursive iterations of noise, studying the ERUB is a convenient and tractable alternative. The upper bound often has convenient functional forms which are (1) sufficiently simple, such that we can directly minimize it, and (2) closely related to the landscape of the objective depending on both the training dataset and the loss function. As a consequence, it is also used in previous PGD literature (Pichapati et al., 2019; Wang et al., 2017) for choosing proper parameters. Moreover, we let $\text{ERUB}_{\min}$ be the achievable optimal upper bound by a specific choice of parameters, e.g., the σ and T.

In this paper, we consider the class of loss functions satisfying the Polyak-Lojasiewicz (PL) condition which bounds losses by corresponding gradient norms. It is more general than the m-strongly convexity. If f is differentiable and M-smooth, then m-strongly convexity implies the PL condition.

Definition 4.1 (Polyak-Lojasiewicz condition (Polyak, 1963))**.** For $f(\theta)$, there exists $\mu > 0$ and for every θ, $\|\nabla f(\theta)\|^2 \geq 2\mu(f(\theta) - f(\theta^*))$.

The PL condition helps us to reveal how the influence of step noise propagates to the final excess error, i.e., EER. Though the assumption was also used previously in Wang et al. (2017); Zhou et al. (2020), neither did they discuss the propagated influence of noise. In the following sections, we will show how the influence can tighten the upper bound in gradient descent and its momentum variant.

4.1 GRADIENT DESCENT METHODS

For the brevity of our discussion, we first define the following constants:

$$\frac{1}{\alpha} \triangleq \frac{2RMN^2}{DG^2}(f(\theta_1) - f(\theta^*)), \ \kappa \triangleq \frac{M}{\mu}, \ \text{and} \ \gamma \triangleq 1 - \frac{1}{\kappa}, \tag{2}$$

which satisfy $\kappa \geq 1$ and $\gamma \in [0, 1)$. Note that κ is the condition number of $f(\cdot)$ if $f(\cdot)$ is strongly convex. κ tends to be large if the function is sensitive to small differences in inputs, and $1/\alpha$ tends to be large if more samples are provided and with a less strict privacy budget. The convergence of PGD under the PL condition has been studied for private (Wang et al., 2017) and non-private (Karimi et al., 2016; Nesterov & Polyak, 2006; Reddi et al., 2016) ERM. Below we extend the bound in (Wang et al., 2017) by considering dynamic influence of noise and relax σ_t to be dynamic:

Theorem 4.1. *Let α, κ and γ be defined in Eq. (2), and $\eta_t = \frac{1}{M}$. Suppose $f(\theta; x_i)$ is G-Lipschitz and $f(\theta)$ is M-smooth satisfying the Polyak-Lojasiewicz condition. For PGD, the following holds:*

$$\text{ERUB}^{uniform} = \gamma^T + R \sum\nolimits_{t=1}^{T} q_t \sigma_t^2, \ \text{where} \ q_t \triangleq \gamma^{T-t} \alpha. \tag{3}$$

In Eq. (3), the step noise magnitude σ_t^2 has an exponential *influence*, q_t, on the EER. The dynamic characteristic of the influence is the key to prove a tighter bound. Plus, on the presence of the dynamic influence, it is natural to choose a dynamic σ_t^2. When relaxing q_t to a static 1, a static σ_t^2 was studied by Wang et al.. They proved a bound which is nearly optimal except a $\ln^2 N$ factor. To get the optimal bound, in the following sections, we look for the σ and T that minimize the upper bound.

4.1.1 UNIFORM SCHEDULE

The uniform setting of σ_t has been previously studied in Wang et al. (2017). Here, we show that, by choosing a proper T, the upper bound can be further improved.

Theorem 4.2. *Suppose conditions in Theorem 4.1 are satisfied. When $\sigma_t^2 = T/R$, let α, γ and κ be defined in Eq. (2) and let T be:*

$$T = \left\lceil \mathcal{O}\left(\kappa \ln \left(1 + \frac{1}{\kappa\alpha} \right) \right) \right\rceil. \tag{4}$$

Meanwhile, if $\kappa \geq \frac{1}{1-c} > 1$, $1/\alpha > 1/\alpha_0$ for some constant $c \in (0, 1)$ and $\alpha_0 > 0$, the corresponding bound is:

$$\text{ERUB}_{\min}^{uniform} = \Theta\left(\frac{\kappa^2}{\kappa + 1/\alpha} \ln \left(1 + \frac{1}{\kappa\alpha} \right) \right). \tag{5}$$

Sketch of proof. The key of proof is to find a proper T to minimize

$$\text{ERUB} = E = \gamma^T + \sum\nolimits_{t=1}^{T} \gamma^{T-t} \alpha R \sigma^2 = \gamma^T + \alpha T \frac{1 - \gamma^T}{1 - \gamma} = \gamma^T + \alpha\kappa(1 - \gamma^T)T$$

where we use $\sigma_t = \sqrt{T/R}$. Vanishing its gradient is to solve $\gamma^T \ln \gamma + \alpha\kappa(1 - \gamma^T) - \alpha\kappa T\gamma^T \ln \gamma = 0$, which however is intractable. In (Wang et al., 2017), T is chosen to be $\mathcal{O}(\ln(1/\alpha))$ and ERUB is relaxed as $\gamma^T + \alpha\kappa T^2$. The approximation results in a less tight bound as $\mathcal{O}(\alpha(1 + \kappa \ln^2(1/\alpha)))$ which explodes as $\kappa \to \infty$.

We observe that for a super sharp loss function, i.e., a large κ, any minor perturbation may result in tremendously fluctuating loss values. In this case, not-stepping-forward will be a good choice. Thus, we choose $T = \frac{1}{\ln(1/\gamma)} \ln \left(1 + \frac{\ln(1/\gamma)}{\alpha} \right) \leq \mathcal{O}\left(\kappa \ln \left(1 + \frac{1}{\kappa\alpha} \right) \right)$ which converges to 0 as $\kappa \to +\infty$. The full proof is deferred to the appendix. $\qquad\square$

4.1.2 DYNAMIC SCHEDULE

A dynamic schedule can improve the upper bound delivered by the uniform schedule. First, we observe that the excess risk in Eq. (3) is upper bounded by two terms: the first term characterizes the error due to the finite iterations of gradient descents; the second term, a weighted sum, comes from error propagated from noise at each iteration. Now we show for any $\{q_t | q_t > 0, t = 1, \ldots, T\}$ (not limited to the q_t defined in Eq. (3)), there is a unique σ_t minimizing the weighted sum:

Lemma 4.1 (Dynamic schedule). *Suppose σ_t satisfy $\sum_{t=1}^{T} \sigma^{-2} = R$. Given a positive sequence $\{q_t\}$, the following equation holds:*

$$\min_{\sigma} R \sum_{t=1}^{T} q_t \sigma_t^2 = \left(\sum_{t=1}^{T} \sqrt{q_t} \right)^2, \text{ when } \sigma_t^2 = \frac{1}{R} \sum_{i=1}^{T} \sqrt{\frac{q_i}{q_t}}. \tag{6}$$

We see that the dynamics in σ_t come from the non-uniform nature of the weight q_t. Since q_t presents the impact of the σ_t on the final error, we denote it as *influence*. Given the dynamic schedule in Eq. (6), it is of our interest to which extent the ERUB can be improved. First, we present Theorem 4.3 to show the optimal T and ERUB.

Theorem 4.3. *Suppose conditions in Theorem 4.1 are satisfied. Let α, κ and γ be defined in Eq. (2). When $\eta_t = \frac{1}{M}$, σ_t (based on Eqs. (3) and (6)) and the T minimizing ERUB are, i.e.,*

$$\sigma_t^2 = \frac{1}{R} \frac{\sqrt{(1/\gamma)^T} - 1}{1 - \sqrt{\gamma}} \sqrt{\gamma^t}, \ T = \left\lceil \left(2\kappa \ln \left(1 + \frac{1}{\kappa\alpha} \right) \right) \right\rceil. \tag{7}$$

Meanwhile, when $\kappa \geq 1$ and $1/\alpha \geq 1/\alpha_0$ for some positive constant α_0, the minimal bound is:

$$\text{ERUB}_{\min}^{dynamic} = \Theta \left(\frac{\kappa^2}{\kappa^2 + 1/\alpha} \right). \tag{8}$$

4.1.3   DISCUSSION

In Theorems 4.2 and 4.3, we present the tightest bounds for functions satisfying the PL condition, to our best knowledge. We further analyze the advantages of our bounds from two aspects: sample efficiency and robustness to sharp losses.

Sample efficiency. Since dataset cannot be infinitely large, it is critical to know how accurate the model can be trained privately with a limited number of samples. Formally, it is of interest to study when κ is fixed and N is large enough such that $\alpha \gg 1$. Then we have the upper bound in Eq. (5) as

$$\text{ERUB}_{\min}^{\text{uniform}} \leq \mathcal{O} \left(\kappa^2 \alpha \ln \left(\frac{1}{\kappa\alpha} \right) \right) \leq \tilde{\mathcal{O}} \left(\frac{DG^2 \ln(N)}{MN^2 R} \right), \tag{9}$$

where we ignore κ and other logarithmic constants with $\tilde{\mathcal{O}}$ as done in Wang et al. (2017). As a result, we get a bound very similar to (Wang et al., 2017), except that R is replaced by $R_{MA} = \epsilon^2 / \ln(1/\delta)$ using Moment Accountant. In comparison, based on Lemma B.3, $R = 2\rho = 2\epsilon + 4\ln(1/\delta) + 4\sqrt{\ln(1/\delta)(\epsilon + \ln(1/\delta))}$ if θ_T satisfies ρ-zCDP. Because $\ln(1/\delta) > 1$, it is easy to see $R_{zCDP} > R_{MA}$ when $\epsilon \leq 2\ln(1/\delta)$. Because our bound saved a factor of $\ln N$ and is thus tighter, as compared to the one reported in (Wang et al., 2017). Remarkably, the saving is due to the maintaining of the influence terms as shown in the proof of Theorem 4.2.

Using the dynamic schedule, we have $\text{ERUB}_{\min}^{\text{dynamic}} \leq \mathcal{O}(\alpha) = \mathcal{O}\left(\frac{DG^2}{MN^2 R} \right)$, which saved another $\ln N$ factor in comparison to the one using the uniform schedule Eq. (9). As shown in Table 1, such advantage maintains when comparing with other baselines.

Robustness. Besides sample efficiency, we are also interested in robustness of the convergence. Specifically, when the samples are noisy or have noisy labels, the loss curvature may be sharp. A sharp loss function will be less smooth, i.e., a small M or has a very small PL parameter. Thus, gradients may change tremendously at some steps especially in the presence of privacy noise. Such changes have more critical impact when only a less number of iterations can be executed due to the privacy constraint. Assume α is some constant while $\kappa \gg 1/\alpha$, we immediately get:

$$\text{ERUB}_{\min}^{\text{uniform}} = \Theta \left(\kappa \ln \left(1 + \frac{1}{\kappa\alpha} \right) \right) = \Theta \left(\frac{1}{\alpha} \right) \leq \mathcal{O} \left(\frac{MN^2 R}{DG^2} \right), \ \text{ERUB}_{\min}^{\text{dynamic}} = \Theta(1).$$

Both are robust, but the dynamic schedule has a smaller factor since $1/\alpha$ could be a large number. In addition, the factor implies that when more samples are used, the uniform schedule is less robust.

4.2 Gradient Descent Methods with Momentum

Section 4.1 shows that the step noise has an exponentially increasing influence on the final loss, and therefore a decreasing noise magnitude improves the utility upper bound by a $\ln N$ factor. On the other hand, it also suggests that noise in the beginning iterations has very minor influence on the final iterations, because the accumulated noise overwhelmed the propagated information. In addition, the proper magnitude of decreasing noise can be hard to find.

The momentum method was originally proposed to stabilize the gradient estimation in Stochastic Gradient Descent (SGD) (Polyak, 1964). In this section, we show that momentum can be used to reduce the variance of privacy noise for full-batch gradient descent, and improve the utility upper bound. The momentum method or the exponentially moving average was also used in previous private learning literature. Pichapati et al. used the momentum as an estimation of gradient mean, without discussions of convergence improvements. Zhou et al. gave a bound for the Adam with DP. However, the derivation is based on gradient norm, which results in a looser bound as shown in Table 1.

The momentum method stabilizes gradients by moving average history coordinate values and thus greatly reduces the variance. The $\phi(m_t, g_t)$ can be rewritten as:

$$m_{t+1} = \phi(m_t, g_t) = \frac{v_{t+1}}{1 - \beta^t}, \ v_{t+1} = \beta v_t + (1 - \beta)g_t = (1 - \beta)\sum\nolimits_{i=1}^{t} \beta^{t-i}g_t, \ v_1 = 0, \quad (10)$$

where $\beta \in [0, 1]$. Note v_{t+1} is a biased estimation of the gradient expectation while m_{t+1} is unbiased.

Theorem 4.4 (Convergence under PL condition). *Suppose $f(\theta; x_i)$ is G-Lipschitz, and $f(\theta)$ is M-smooth and satisfies the Polyak-Lojasiewicz condition. Assume $\beta \neq \gamma$ and $\beta \in (0, 1)$. Let $\eta_t = \frac{\eta_0}{2M}$ and $\eta_0 \leq 8 \left(\sqrt{1 + 64\beta\gamma(\gamma - \beta)^{-2}(1 - \beta)^{-3}} + 1 \right)^{-1}$. Then the following holds:*

$$\text{EER} \leq \left(\gamma^T + 2R\eta_0\alpha \underbrace{U_3(\sigma, T)}_{\text{noise varinace}} \right)(f(\theta_1) - f(\theta^*)) - \zeta\frac{\eta_0}{2M} \underbrace{\sum\nolimits_{t=1}^{T} \gamma^{T-t}\mathbb{E}\|v_{t+1}\|^2}_{\text{momentum effect}} \quad (11)$$

$$\text{where} \quad \gamma = 1 - \frac{\eta_0}{\kappa}, \ \zeta = 1 - \frac{1}{\beta(1 - \beta)^3}\eta_0^2 - \frac{1}{4}\eta_0 \geq 0, \quad (12)$$

$$U_3 = \sum\nolimits_{t=1}^{T} \gamma^{T-t}\frac{(1 - \beta)^2}{(1 - \beta^t)^2} \sum\nolimits_{i=1}^{t} \beta^{2(t-i)}\sigma_i^2. \quad (13)$$

The upper bound includes three parts that influence the bound differently: *(1) Convergence.* The convergence term is mainly determined by η_0 and κ. η_0 should be in $(0, \kappa)$ such that the upper bound can converge. A large η_0 will be preferred to speed up convergence if it does not make the rest two terms worse. *(2) Noise Variance.* The second term compressed in U_3 is the effect of noise. One difference introduced by the momentum is the factor $(1 - \beta)/(1 - \beta^t)$ which decreases by t. $(1 - \beta)/(1 - \beta^t)$ decreases faster than γ^t at the beginning and converges to a non-zero constant $1 - \beta$. Therefore, the influence factor in U_3, γ^{T-t}, will be reduced to γ^T which maintains constant. Furthermore, when $t > \hat{T}$, the moving average of σ_t^2 reweighted the γ^{T-t} which also makes the quickly changing influence term smoother. *(3) Momentum Effect.* The momentum effect term can improve the upper bound when η_0 is small. For example, when $\beta = 0.9$ and $\gamma = 0.99$, then $\eta_0 \leq 0.98/M$ which is a rational value. Following the analysis, when M is large which means the gradient norms will significantly fluctuate, the momentum term may take the lead. Adjusting the noise scale in this case may be less useful for improving utility.

Theorem 4.5 (Uniform schedule). *Suppose the assumptions in Theorem 4.4 are satisfied. Let $\sigma_t^2 = T/R$, and let:*

$$\hat{T} = \max t \ s.t. \ \gamma^{t-1} \geq \frac{1 - \beta}{1 - \beta^t}, \ T = \left\lceil \mathcal{O}\left(\frac{\kappa}{\eta_0} \ln\left(1 + \frac{\eta_0}{\kappa\alpha}\right) \right) \right\rceil.$$

If some positive constant c, $1/\alpha > 1/\alpha_0$ and $\alpha_0 > 0$, the following inequality holds:

$$\text{ERUB}_{\min} \leq \mathcal{O}\left(\frac{\kappa^2}{\kappa + \eta_0/\alpha} \left[\mathbb{I}_{T \leq \hat{T}} + \gamma^{\hat{T}-1} \ln\left(1 + \frac{\eta_0}{\kappa\alpha}\right) \mathbb{I}_{T > \hat{T}} \right] \right).$$

Theorem 4.6 (Dynamic schedule). *Suppose the assumptions in Theorem 4.4 are satisfied. Let* $\alpha' = \frac{2\eta_0\alpha}{\gamma(1-\gamma\beta^2)}$, $\beta < \gamma$ *and* $\hat{T} = \max t$ *s.t.* $\gamma^{t-1} \geq \frac{1-\beta}{1-\beta^t}$. *Use the following schedule and iteration:*

$$\sigma_t^2 = \frac{1}{R}\sum_{i=1}^{T}\sqrt{\frac{q_i}{q_t}}, \ T^{dyn} = \left\lceil \mathcal{O}\left(\frac{2\kappa}{\eta_0}\ln\left(1+\frac{\eta_0}{\kappa\alpha}\right)\right)\right\rceil,$$

where $q_t = c_1\gamma^{T+t}\mathbb{I}_{T\leq\hat{T}} + \gamma^{\hat{T}-1}c_2\gamma^{T-t}\mathbb{I}_{T>\hat{T}}$ *for some positive constants* c_1 *and* c_2. *The following inequality holds:*

$$\mathrm{ERUB} \leq \gamma^T + 2\eta_0\alpha\sum_{t=1}^{T}Rq_t\sigma_t^2, \ \mathrm{ERUB_{min}} \leq \mathcal{O}\left(\frac{\kappa\alpha}{\kappa\alpha+\eta_0}\left(\frac{\kappa\alpha}{\kappa\alpha+\eta_0}\mathbb{I}_{T\leq\hat{T}}+\mathbb{I}_{T>\hat{T}}\right)\right).$$

Notably, the dynamic schedule should not be decreasing when $t \leq \hat{T}$, which is counterintuitive.

Discussion. Theoretically, the dynamic schedule is more influential in vanilla gradient descent methods than the momentum variant. The result is mainly attributed to the averaging operation. The moving averaging, $(1-\beta)\sum_{i=1}^{t}\beta^{t-i}g_i/(1-\beta^t)$, increase the influence of the under-presented initial steps and decrease the one of the over-sensitive last steps.

5 EXPERIMENTS

We empirically validate the properties of privacy schedules and their connections to learning algorithms. In this section, we briefly review the schedule behavior on logistic and quadratic losses under varying data sensitivity. More discussions and empirical results are available in Appendix D. We first show the estimated influence of step noise q_t (by retraining the private learning algorithms) in Fig. 1 Left. We see the trends of influence are in an exponential form. We then show the trends on the variance of influence in the Middle Pane, where *uni* denotes the uniform schedule baseline, *exp* is an exponential schedule, *dyn* denotes the dynamic schedule minimizing the ERUB. Dynamic schedules shows improvements of influence variance during the entire optimization span. We lastly evaluate the impacts from momentum in the Right Pane, using a Deep Neural Network (DNN) with 2 layers and 100 hidden units. Because of time costs of training deep networks, we do not estimate the influence by retraining and then compute schedules. Instead, we grid-search for the schedule hyper-parameters to find the best one. We see that influence modeled by an exponential function (*expinfl*) has comparable performance of the influence modeled by linear combination of two reverse exponential functions (*momexpinfl*). The latter only shows advantage in the setting that data scale is 25 and the number of iteration is only 100, which is expected by our analysis.

Figure 1: Comparison of dynamic schedule and uniform schedule on different data scale. Left pane is the influence by iteration estimated by retraining. The rest two are the relative loss by varying data scale (left axis with solid lines) and the variance of influence (right axis with dashed lines). The middle is on the MNIST35 dataset consisting of 1000 digit 3 and 5 images using logistic regression. The right is the final loss on subsampled MNIST dataset of 1000 training samples and 50, 000 test samples when using DNN and momentum methods.

6 CONCLUSION

When a privacy budget is provided for a certain learning task, one has to carefully schedule the privacy usage through the learning process. Uniformly scheduling the budget has been widely used in literature whereas increasing evidence suggests that dynamically schedules could empirically outperform the uniform one. This paper provided a principled analysis on the problem of optimal budget allocation and connected the advantages of dynamic schedules to both the loss structure and the learning behavior. We further validated our results through empirical studies.

REFERENCES

Martin Abadi, Andy Chu, Ian Goodfellow, H. Brendan McMahan, Ilya Mironov, Kunal Talwar, and Li Zhang. Deep Learning with Differential Privacy. In *CCS: Proceedings of the 2016 ACM SIGSAC Conference on Computer and Communications Security*, CCS '16, pp. 308–318, New York, NY, USA, 2016. ACM.

R. Bassily, A. Smith, and A. Thakurta. Private Empirical Risk Minimization: Efficient Algorithms and Tight Error Bounds. In *2014 IEEE 55th Annual Symposium on Foundations of Computer Science*, pp. 464–473, October 2014.

Mark Bun and Thomas Steinke. Concentrated Differential Privacy: Simplifications, Extensions, and Lower Bounds. In *Theory of Cryptography*, volume 9985, pp. 635–658. Springer Berlin Heidelberg, Berlin, Heidelberg, 2016.

Mark Bun, Cynthia Dwork, Guy N. Rothblum, and Thomas Steinke. Composable and Versatile Privacy via Truncated CDP. In *Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing*, STOC 2018, pp. 74–86, New York, NY, USA, 2018. ACM.

Kamalika Chaudhuri, Claire Monteleoni, and Anand D. Sarwate. Differentially Private Empirical Risk Minimization. *Journal of Machine Learning Research*, 12(Mar):1069–1109, 2011.

Rachel Cummings, Sara Krehbiel, Kevin A Lai, and Uthaipon Tantipongpipat. Differential Privacy for Growing Databases. In *Advances in Neural Information Processing Systems 31*, pp. 8864–8873. Curran Associates, Inc., 2018.

Damien Desfontaines and Balázs Pejó. SoK: Differential Privacies. *arXiv:1906.01337 [cs]*, June 2019.

Cynthia Dwork, Frank McSherry, Kobbi Nissim, and Adam Smith. Calibrating Noise to Sensitivity in Private Data Analysis. In *Theory of Cryptography*, Lecture Notes in Computer Science, pp. 265–284. Springer Berlin Heidelberg, 2006.

Cynthia Dwork, Alan Karr, Kobbi Nissim, and Lars Vilhuber. On Privacy in the Age of COVID-19. *Journal of Privacy and Confidentiality*, 10(2), June 2020.

Vitaly Feldman, Tomer Koren, and Kunal Talwar. Private stochastic convex optimization: Optimal rates in linear time. In *Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing*, STOC 2020, pp. 439–449, New York, NY, USA, June 2020. Association for Computing Machinery.

Matt Fredrikson, Somesh Jha, and Thomas Ristenpart. Model Inversion Attacks That Exploit Confidence Information and Basic Countermeasures. In *CCS: Proceedings of the 22Nd ACM SIGSAC Conference on Computer and Communications Security*, CCS '15, pp. 1322–1333, New York, NY, USA, 2015. ACM.

Prateek Jain, Dheeraj Nagaraj, and Praneeth Netrapalli. Making the Last Iterate of SGD Information Theoretically Optimal. In *Conference on Learning Theory*, pp. 1752–1755, June 2019.

Hamed Karimi, Julie Nutini, and Mark Schmidt. Linear Convergence of Gradient and Proximal-Gradient Methods Under the Polyak-Łojasiewicz Condition. In *Machine Learning and Knowledge Discovery in Databases*, Lecture Notes in Computer Science, pp. 795–811, Cham, 2016. Springer International Publishing.

Diederik P. Kingma and Jimmy Ba. Adam: A Method for Stochastic Optimization. *arXiv:1412.6980 [cs]*, December 2014.

Y. Lecun, L. Bottou, Y. Bengio, and P. Haffner. Gradient-based learning applied to document recognition. *Proceedings of the IEEE*, 86(11):2278–2324, November 1998.

Jaewoo Lee and Daniel Kifer. Concentrated Differentially Private Gradient Descent with Adaptive per-Iteration Privacy Budget. In *Proceedings of the 24th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining*, KDD '18, pp. 1656–1665, New York, NY, USA, 2018. ACM.

Ilya Mironov. Rényi Differential Privacy. In *2017 IEEE 30th Computer Security Foundations Symposium (CSF)*, pp. 263–275, Santa Barbara, CA, USA, August 2017. IEEE.

Yurii Nesterov and B.T. Polyak. Cubic regularization of Newton method and its global performance. *Mathematical Programming*, 108(1):177–205, August 2006.

Venkatadheeraj Pichapati, Ananda Theertha Suresh, Felix X. Yu, Sashank J. Reddi, and Sanjiv Kumar. AdaCliP: Adaptive Clipping for Private SGD. *arXiv:1908.07643 [cs, stat]*, October 2019.

B. T. Polyak. Gradient methods for the minimisation of functionals. *USSR Computational Mathematics and Mathematical Physics*, 3(4):864–878, January 1963.

B. T. Polyak. Some methods of speeding up the convergence of iteration methods. *USSR Computational Mathematics and Mathematical Physics*, 4(5):1–17, January 1964.

Ning Qian. On the momentum term in gradient descent learning algorithms. *Neural Networks*, 12(1): 145–151, January 1999.

Sashank J. Reddi, Ahmed Hefny, Suvrit Sra, Barnabas Poczos, and Alex Smola. Stochastic Variance Reduction for Nonconvex Optimization. In *International Conference on Machine Learning*, pp. 314–323, June 2016.

Alfréd Rényi. On measures of entropy and information. In *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability, Volume 1: Contributions to the Theory of Statistics*. The Regents of the University of California, 1961.

R. Shokri, M. Stronati, C. Song, and V. Shmatikov. Membership Inference Attacks Against Machine Learning Models. In *2017 IEEE Symposium on Security and Privacy (SP)*, pp. 3–18, May 2017.

Om Thakkar, Galen Andrew, and H. Brendan McMahan. Differentially Private Learning with Adaptive Clipping. *arXiv:1905.03871 [cs, stat]*, May 2019.

Di Wang, Minwei Ye, and Jinhui Xu. Differentially Private Empirical Risk Minimization Revisited: Faster and More General. In *Advances in Neural Information Processing Systems 30*, pp. 2722–2731. Curran Associates, Inc., 2017.

Di Wang, Changyou Chen, and Jinhui Xu. Differentially Private Empirical Risk Minimization with Non-convex Loss Functions. In *International Conference on Machine Learning*, pp. 6526–6535, May 2019.

Xi Wu, Fengan Li, Arun Kumar, Kamalika Chaudhuri, Somesh Jha, and Jeffrey Naughton. Bolt-on Differential Privacy for Scalable Stochastic Gradient Descent-based Analytics. In *Proceedings of the 2017 ACM International Conference on Management of Data*, SIGMOD '17, pp. 1307–1322, New York, NY, USA, 2017. ACM.

Lei Yu, Ling Liu, Calton Pu, Mehmet Emre Gursoy, and Stacey Truex. Differentially Private Model Publishing for Deep Learning. *proceedings of 40th IEEE Symposium on Security and Privacy*, April 2019.

Yingxue Zhou, Xiangyi Chen, Mingyi Hong, Zhiwei Steven Wu, and Arindam Banerjee. Private Stochastic Non-Convex Optimization: Adaptive Algorithms and Tighter Generalization Bounds. *arXiv:2006.13501 [cs, stat]*, August 2020.

A   COMPARISON OF ALGORITHMS

Table 2: The algorithms are T-iteration $\frac{1}{2}R$-zCDP or equivalently (ϵ, δ)-DP under the PL condition (unless marked with *). The O notation in this table drops other $\ln$ terms. Unless otherwise specified, all algorithms terminate at step $T = \mathcal{O}(\ln \frac{D}{N^2 R})$. Assume loss functions are 1-smooth and 1-Lipschitz continuous, and all parameters satisfy their numeric assumptions. Key notations: $\mathcal{O}_p$ – bound occurs in probability p; D – feature dimension; N – sample size; R – privacy budget; c_i – constant.

Algorithm	Schedule (σ_t^2)	Utility Upper Bd.
*GD+Adv (Bassily et al., 2014)	$\mathcal{O}\left(\frac{\ln(N/\delta)}{R_{\epsilon,\delta}}\right)$	$\mathcal{O}\left(\frac{D \ln^3 N}{N R_{\epsilon,\delta}}\right)$
GD+MA (Wang et al., 2017)	$\mathcal{O}(\frac{T}{R_{\epsilon,\delta}})$	$\mathcal{O}\left(\frac{D \ln^2 N}{N^2 R_{\epsilon,\delta}}\right)$
*GD+Adv+BBImp (Cummings et al., 2018)	$\mathcal{O}\left(\frac{n^2 \ln(n/\delta)}{R_{\epsilon,\delta}}\right)$	$\mathcal{O}_p\left(\frac{D^2 \ln^2(1/p)}{R_{\epsilon,\delta} N^{1-c}}\right)$
*SSGD+zCDP (Feldman et al., 2020)	$\frac{1}{D}$	$\mathcal{O}\left(\left(\frac{1}{\sqrt{N}} + \frac{2\sqrt{D}}{\sqrt{R}N}\right)\ln N\right)$
Adam+MA (Zhou et al., 2020)	$\mathcal{O}(\frac{T}{R_{\epsilon,\delta}})$	$\mathcal{O}_p\left(\frac{\sqrt{D}\ln(ND\epsilon/(1-p))}{N R_{\epsilon,\delta}}\right)$
GD, Non-Private	0	$\mathcal{O}\left(\frac{D}{N^2 R}\right)$
GD+zCDP, Static Schedule	$\frac{T}{R}$	$\mathcal{O}\left(\frac{D \ln N}{N^2 R}\right)$
GD+zCDP, Dynamic Schedule	$\mathcal{O}\left(\frac{\gamma^{(t-T)/2}}{R}\right)$	$\mathcal{O}\left(\frac{D}{N^2 R}\right)$
Momentum+zCDP, Static Schedule	$\frac{T}{R}$	$\mathcal{O}\left(\frac{D}{N^2 R}(c + \ln N \mathbb{I}_{T>\hat{T}})\right)$
Momentum+zCDP, Dynamic Schedule	$\mathcal{O}\left(\frac{c_1 \gamma^{T+t} + c_2 \gamma^{(T-t)/2}}{R}\right)$	$\mathcal{O}\left(\frac{D}{N^2 R}(1 + \frac{cD}{N^2 R}\mathbb{I}_{T>\hat{T}})\right)$

We present Table 2 as an sumpplementary to the Table 1. Asymptotic upper bounds are achieved when sample size N approaches infinity. Both R and $R_{\epsilon,\delta}$ with $R_{\epsilon,\delta} < R$ are the privacy budgets of corresponding algorithms. Specifically, $R_{\epsilon,\delta} = \epsilon^2 / \ln(1/\delta) < R$ when the private algorithm is (ϵ, δ)-DP with $\epsilon \le 2\ln(1/\delta)$.

PGD+Adv. Adv denotes the Advanced Composition method (Bassily et al., 2014). The method assumes that loss function is 1-strongly convex which implies the PL condition and optimized variable is in a convex set of diameter 1 w.r.t. l_2 norm.

PGD+MA. MA denotes the Moment Accoutant (Abadi et al., 2016) which improve the composed privacy bound versus the Advanced Composition. The improvement on privacy bound lead to a enhanced utility bound, as a result.

PGD+Adv+BBImp. The dynamic method assumes that the loss is 1-strongly convex and data comes in stream with $n \le N$ samples at each round. Their utility upper bound is achieved at some probability p with any positive c.

SSGD+zCDP. SSGD is short for Snowball Stochastic Gradient Descent where the batch size is gradually increased. The method assumes that loss functions are convex within a convex body of diameter 1 w.r.t. l_2 norm.

Adam+MA. The authors prove a convergence bound for the gradient norms which is extended to loss bound by using PL condition. They also presents the results for AdaGrad and GD which are basically of the same upper bound. Out theorems improve their bound by using the recursive derivation based on the PL condition, while their bound is a simple application of the condition on the gradient norm bound.

GD, Non-Private. This method does not inject noise into gradients but limit the number of iterations. With the bound, we can see that our utility bound are optimal with dynamic schedule.

GD+zCDP. We discussed the static and dynamic schedule for the gradient descent method where the dynamic noise influence is the key to tighten the bound.

Momemtum+zCDP. Different from the GD+zCDP, momentum methods will have two phase of utility upper bound. When T is small than some positive constant $\hat{T}$, the bound is as tight as the non-private one. Afterwards, the momentum has a bound degraded as the GD bound.

B PRELIMINARIES

B.1 PRIVACY

Lemma B.1 (Composition & Post-processing). *Let two mechanisms be $M : \mathcal{D}^n \to \mathcal{Y}$ and $M' : \mathcal{D}^n \times \mathcal{Y} \to \mathcal{Z}$. Suppose M satisfies (ρ_1, a)-zCDP and $M'(\cdot, y)$ satisfies (ρ_2, a)-zCDP for $\forall y \in \mathcal{Y}$. Then, mechanism $M'' : \mathcal{D}^n \to \mathcal{Z}$ (defined by $M''(x) = M'(x, M(x)))$ satisfies $(\rho_1 + \rho_2)$-zCDP.*

Definition B.1 (Sensitivity). The sensitivity of a gradient query ∇_t to the dataset $\{x_i\}_{i=1}^N$ is

$$\Delta_2(\nabla_t) = \max_n \left\| \frac{1}{N} \sum_{j=1, j \neq n}^N \nabla_t^{(j)} - \frac{1}{N} \sum_{j=1}^N \nabla_t^{(j)} \right\|_2 = \frac{1}{N} \max_n \left\| \nabla_t^{(n)} \right\|_2 \qquad (14)$$

where $\nabla_t^{(n)}$ denotes the gradient of the n-th sample.

Lemma B.2 (Gaussian mechanism (Bun & Steinke, 2016)). *Let $f : \mathcal{D}^n \to \mathcal{Z}$ have sensitivity Δ. Define a randomized algorithm $M : \mathcal{D}^n \to \mathcal{Z}$ by $M(x) \leftarrow f(x) + \mathcal{N}(0, \Delta^2 \sigma^2 I)$. Then M satisfies $\frac{1}{2\sigma^2}$-zCDP.*

Lemma B.3 ((Bun & Steinke, 2016)). *If M is a mechanism satisfying ρ-zCDP, then M is $(\rho + 2\sqrt{\rho \ln(1/\delta)}, \delta)$-DP for any $\delta > 0$.*

By solving $\rho + 2\sqrt{\rho \ln(1/\delta)} = \epsilon$, we can get $\rho = \epsilon + 2\ln(1/\delta) + 2\sqrt{\ln(1/\delta)(\epsilon + \ln(1/\delta))}$.

B.2 AUXILIARY LEMMAS

Lemma B.4. *If $\max_n \|x_n\|_2 = 1$ and $\frac{1}{N} \sum_n x_n = 0$, then the gradient sensitivity of the squared loss will be*

$$\Delta_2(\nabla) = \max_i \frac{1}{N} \sqrt{2f(\theta; x_i)} \|x_i\|_2 \leq \frac{1}{2}(DM \|\theta\|^2 + 1),$$

where $\Theta_{\mathcal{M}}$ is the set of all possible parameters θ_t generated by the learning algorithm $\mathcal{M}$.

Proof. According to the definition of sensitivity in Eq. (14), we have

$$\Delta_2(\nabla) = \max_i \left\| \nabla^{(i)} \right\|_2 = \max_n \frac{1}{n} \left\| A^{(i)} \theta - x_i \right\|_2$$

where we use i denotes the index of sample in the dataset. Here, we assume it is constant 1. We may get

$$\left\| A^{(i)} \theta - x_i \right\|_2^2 = \left\| x_i(x_i^\top \theta - 1) \right\|_2^2 = (x_i^\top \theta - 1)^2 \|x_i\|_2^2 = 2f(\theta; x_i) \|x_i\|_2^2$$

where $f(\theta; x_i) = \frac{1}{2}(x_i^\top \theta - 1)^2$. Thus,

$$\Delta_2(\nabla) = \max_i \frac{1}{N} \sqrt{2f(\theta; x_i)} \|x_i\|_2$$

Since $\|x_n\|_2 \leq 1$ and $\frac{1}{N} \sum_{n=1}^N x_n = 0$,

$$f(\theta) = \frac{1}{2N} \sum_{n=1}^N [(x_n^\top \theta)^2 - 2x_n^\top \theta + 1] \leq \frac{1}{2N} \sum_{n=1}^N [(\|x_n\| \|\theta\|)^2 + 1] \leq \frac{1}{2}(DM \|\theta\|^2 + 1)$$

$\square$

Lemma B.5. *Assume assumptions in Theorem 4.4 are satisfied. Given variables defined in Theorem 4.4, the following inequality holds true:*

$$\sum_{t=1}^T \gamma^{T-t} \frac{2(1-\beta)\eta_t}{b_t} \sum_{i=1}^t \beta^{t-i} \|\nabla_t - \nabla_i\|^2 \leq \frac{\eta_0^3 \beta \gamma}{2M(1-\beta)^3(\gamma-\beta)^2} \sum_{i=1}^{T-1} \gamma^{T-i} \|v_{i+1}\|^2 .$$

Proof. We first handle the inner summation. By smoothness, the inequality $\|\nabla f(x) - \nabla f(y)\| \leq M \|x - y\|$ holds true. Thus,

$$
\begin{aligned}
\sum\nolimits_{i=1}^{t} \beta^{t-i} \|\nabla_t - \nabla_i\|^2 &\leq M^2 \sum\nolimits_{i=1}^{t} \beta^{t-i} \|\theta_t - \theta_i\|^2 \\
&= M^2 \sum\nolimits_{k=0}^{t-1} \beta^k \|\theta_t - \theta_{t-k}\|^2 \\
&= M^2 \sum\nolimits_{k=0}^{t-1} \beta^k \left\| \sum\nolimits_{i=t-k}^{t-1} \eta_i v_{i+1}/b_i \right\|^2 \\
&\leq M^2 \sum\nolimits_{k=0}^{t-1} \beta^k \left(\sum\nolimits_{j=t-k}^{t-1} \eta_j^2/b_j^2 \right) \left(\sum\nolimits_{i=t-k}^{t-1} \|v_{i+1}\|^2 \right)
\end{aligned}
$$

where the last inequality is by Cauchy-Schwartz inequality. Because $\frac{1}{b_t} = \frac{1}{1-\beta^t} \leq \frac{1}{1-\beta}$ and $\eta_t = \frac{\eta_0}{2M}$,

$$
\begin{aligned}
\sum\nolimits_{i=1}^{t} \beta^{t-i} \|\nabla_t - \nabla_i\|^2 &\leq \frac{\eta_0^2}{4(1-\beta)^2} \sum\nolimits_{k=0}^{t-1} \beta^k k \sum\nolimits_{i=t-k}^{t-1} \|v_{i+1}\|^2 \\
&= \frac{\eta_0^2}{4(1-\beta)^2} \sum\nolimits_{k=0}^{t-1} \beta^k k \sum\nolimits_{i=1}^{t-1} \|v_{i+1}\|^2 \, \mathbb{I}(i \geq t-k) \\
&= \frac{\eta_0^2}{4(1-\beta)^2} \sum\nolimits_{i=1}^{t-1} \|v_{i+1}\|^2 \sum\nolimits_{k=0}^{t-1} \beta^k k \mathbb{I}(k \geq t-i) \\
&= \frac{\eta_0^2}{4(1-\beta)^2} \sum\nolimits_{i=1}^{t-1} \|v_{i+1}\|^2 \sum\nolimits_{k=t-i}^{t-1} \beta^k k
\end{aligned} \tag{15}
$$

where $\mathbb{I}(\cdot)$ is the indicating function which output 1 if the condition holds true, otherwise 0.

Denote the left-hand-side of the conclusion as LHS. We plug Eq. (15) into LHS to get

$$
\begin{aligned}
\text{LHS} &\leq \sum_{t=1}^{T} \gamma^{T-t} \frac{1}{b_t} \frac{\eta_0^3}{4M(1-\beta)} \sum\nolimits_{i=1}^{t-1} \|v_{i+1}\|^2 \sum\nolimits_{k=t-i}^{t-1} \beta^k k \\
&\leq \frac{\eta_0^3}{4M(1-\beta)^2} \sum_{t=1}^{T} \gamma^{T-t} \sum\nolimits_{i=1}^{t-1} \|v_{i+1}\|^2 \sum\nolimits_{k=t-i}^{t-1} \beta^k k
\end{aligned}
$$

where we relax the upper bound by $\frac{1}{b_t} = \frac{1}{1-\beta^t} \leq \frac{1}{1-\beta}$. Using Lemma B.6 can directly lead to the conclusion:

$$
\text{LHS} \leq \frac{\eta_0^3 \beta \gamma}{2M(1-\beta)^3(\gamma-\beta)^2} \sum_{i=1}^{T-1} \gamma^{T-i} \|v_{i+1}\|^2 .
$$

$\square$

Lemma B.6. *Given variables defined in Theorem 4.4, the following inequality holds true:*

$$
\sum_{t=1}^{T} \gamma^{T-t} \sum\nolimits_{i=1}^{t-1} \|v_{i+1}\|^2 \sum\nolimits_{k=t-i}^{t-1} k\beta^k \leq \frac{2\beta\gamma}{(\gamma-\beta)^2(1-\beta)} \sum_{i=1}^{T-1} \gamma^{T-i} \|v_{i+1}\|^2 .
$$

Proof. We first derive the summation:

$$
\begin{aligned}
U_1(t,i) \triangleq \sum_{k=t-i}^{t-1} \beta^k k &= \sum_{k=t-i}^{t-1} \sum_{j=1}^{k} \beta^k \\
&= \sum_{k=t-i}^{t-1} \sum_{j=1}^{t-1} \beta^k \mathbb{I}(j \le k) \\
&= \sum_{j=1}^{t-1} \sum_{k=\max(t-i,j)}^{t-1} \beta^k \\
&= \sum_{j=1}^{t-1} \frac{\beta^{\max(t-i,j)} - \beta^t}{1 - \beta} \\
&= \frac{1}{1-\beta} \left((t-i)\beta^{t-i} + \frac{\beta^{t-i+1} - \beta^t}{1-\beta} - \frac{\beta - \beta^t}{1-\beta} \right) \\
&= \frac{1}{1-\beta} \left((t-i)\beta^{t-i} + \frac{\beta^{t-i+1} - \beta}{1-\beta} \right)
\end{aligned}
$$

Now, we substitute $U_1(t,i)$ into LHS and replace $t - i$ by j, i.e., $t = j + i$, to get

$$
\begin{aligned}
\text{LHS} &= \sum_{t=1}^{T} \gamma^{T-t} \sum_{i=1}^{t-1} \|v_{i+1}\|^2 \frac{1}{1-\beta} \left((t-i)\beta^{t-i} + \frac{\beta^{t-i+1} - \beta}{1-\beta} \right) \\
&= \sum_{i=1}^{T-1} \|v_{i+1}\|^2 \sum_{t=i+1}^{T} \gamma^{T-t} \frac{1}{1-\beta} \left((t-i)\beta^{t-i} + \frac{\beta^{t-i+1} - \beta}{1-\beta} \right) \\
&= \sum_{i=1}^{T-1} \|v_{i+1}\|^2 \sum_{j=1}^{T-i} \gamma^{T-(j+i)} \frac{1}{1-\beta} \left(j\beta^j + \frac{\beta^{j+1} - \beta}{1-\beta} \right) \\
&= \sum_{i=1}^{T-1} \gamma^{T-i} \|v_{i+1}\|^2 \sum_{j=1}^{T-i} \gamma^{-j} \frac{1}{1-\beta} \left(j\beta^j + \frac{\beta^{j+1} - \beta}{1-\beta} \right) \\
&\le \frac{1}{1-\beta} \sum_{i=1}^{T-1} \gamma^{T-i} \|v_{i+1}\|^2 \sum_{j=1}^{T-i} \left(j \left(\frac{\beta}{\gamma} \right)^j + \frac{\beta}{1-\beta} \left(\frac{\beta}{\gamma} \right)^j \right)
\end{aligned}
$$

Let $a = \beta/\gamma$, we show

$$
\begin{aligned}
\sum_{j=1}^{T-i} j a^j &= \sum_{j=1}^{T-i} \sum_{o=1}^{j} a^j \\
&= \sum_{o=1}^{T-i} \sum_{j=o}^{T-i} a^j \\
&= \sum_{o=1}^{T-i} \left(\frac{a^o - a^{T-i+1}}{1-a} \right) \\
&= \frac{a - a^{T-i+1}}{(1-a)^2} - (T-i) \frac{a^{T-i+1}}{1-a} \\
&\le \frac{a}{(1-a)^2}.
\end{aligned}
$$

Thus,

$$\text{LHS} \le \frac{1}{1-\beta} \sum_{i=1}^{T-1} \gamma^{T-i} \|v_{i+1}\|^2 \left(\frac{a}{(1-a)^2} + \frac{\beta}{1-\beta} \sum_{j=1}^{T-i} a^j \right)$$

$$\le \frac{1}{1-\beta} \sum_{i=1}^{T-1} \gamma^{T-i} \|v_{i+1}\|^2 \left(\frac{a}{(1-a)^2} + \frac{\beta}{1-\beta} \frac{a}{1-a} \right)$$

$$\le \frac{a}{(1-a)^2(1-\beta)} \sum_{i=1}^{T-1} \gamma^{T-i} \|v_{i+1}\|^2$$

Because $\gamma < 1$, $\beta < a = \beta/\gamma$ and

$$\frac{a}{(1-a)^2} + \frac{\beta}{1-\beta} \frac{a}{1-a} \le \frac{2a}{(1-a)^2}.$$

Therefore,

$$\text{LHS} \le \frac{2a}{(1-a)^2(1-\beta)} \sum_{i=1}^{T-1} \gamma^{T-i} \|v_{i+1}\|^2 = \frac{2\beta\gamma}{(\gamma-\beta)^2(1-\beta)} \sum_{i=1}^{T-1} \gamma^{T-i} \|v_{i+1}\|^2$$

$\square$

Lemma B.7. *Suppose $\gamma \in (0,1)$ and $\beta \in (0,1)$. Define*

$$\hat{T} = \max t \text{ s.t. } \gamma^{t-1} \ge \frac{1-\beta}{1-\beta^t}.$$

If $t \le \hat{T}$, $\frac{1-\beta}{1-\beta^t} \le \gamma^{t-1}$ for $t = 1, \dots, T$. If $t > \hat{T}$, $\frac{1-\beta}{1-\beta^t} < \gamma^{\hat{T}-1}$.

Proof. Define $h(t) = \gamma^{t-1}(1-\beta^t)$ whose derivatives are

$$h'(t) = \gamma^{t-1}(1-\beta^t)\ln\gamma + \gamma^{t-1}(-\beta^t)\ln\beta$$

$$= \gamma^{t-1}\left[\ln\gamma - \beta^t(\ln\gamma + \ln\beta)\right]$$

$$= \gamma^{t-1}\left[1 - \beta^t(1 + \log_\gamma \beta)\right]\ln\gamma.$$

Simple calculation shows $1 - \beta^t(1 + \log_\gamma \beta)\big|_{t=0} = -\log_\gamma \beta < 0$ and $\lim_{t\to+\infty} 1 - \beta^t(1 + \log_\gamma \beta) = 1$. When $t = -\log_\beta(1 + \log_\gamma \beta)$ denoted as t_0, $1 - \beta^t(1 + \log_\gamma \beta) = 0$. Because $1 - \beta^t(1 + \log_\gamma \beta)$ is monotonically increasing by t and $\gamma^{t-1}\ln\gamma$ is negative, $h'(t) \ge 0$ if $t \le t_0$. Otherwise, $h'(t) < 0$. Therefore, $h(t)$ is a concave function. Because $h(1) = 1-\beta$ and $h(\hat{T}) = \gamma^{\hat{T}-1}(1-\beta^{\hat{T}}) \ge 1 - \beta > 0$, $h(t) \ge 1 - \beta$ for $t = 1, \dots, \hat{T}$. Thus, for all $t \in [1, \hat{T}]$, we have $\frac{1-\beta}{1-\beta^t} \le \gamma^{t-1}$.

For $t > \hat{T}$, because $\frac{1-\beta}{1-\beta^t}$ monotonically increases by t, we have $\frac{1-\beta}{1-\beta^t} < \frac{1-\beta}{1-\beta^{\hat{T}}} \le \gamma^{\hat{T}-1}$. $\square$

C PROOFS

Proof of Theorem 3.1. Because all sample gradient are G-Lipschitz continuous, the sensitivity of the averaged gradient is upper bounded by G/N. Based on Lemma B.2, the privacy cost of g_t is $\frac{1}{2\sigma_t^2}$[1].

Here, we make the output of each iteration a tuple of $(\theta_{t+1}, v_{t=1})$. For the 1st iteration, because θ_1 does not embrace private information by random initialization, the mapping,

$$\begin{bmatrix} v_2 \\ \theta_2 \end{bmatrix} = \begin{bmatrix} g_1 \\ \theta_1 - \eta_1 g_1 \end{bmatrix},$$

---

[1]For brevity, when we say the privacy cost of some value, e.g., gradient, we actually refer to the cost of mechanism that output the value.

is $\hat{\rho}_1$-zCDP where $\hat{\rho}_1 = \frac{1}{2\sigma_t^2}$.

Suppose the output of the t-th iteration, (θ_t, v_t), is $\hat{\rho}_t$-zCDP. At each iteration, we have the following mapping $(\theta_t, v_t) \to (\theta_{t+1}, v_{t+1})$ defined as

$$
\begin{bmatrix} v_{t+1} \\ \theta_{t+1} \end{bmatrix} = \begin{bmatrix} \phi(v_t, g_t) \\ \theta_t - \eta_t \phi(v_t, g_t) \end{bmatrix}.
$$

Thus, the output tuple (θ_{t+1}, v_{t+1}) is $(\hat{\rho}_t + \frac{1}{2\sigma_t^2})$-zCDP by Lemma B.1.

Thus, the recursion implies that (θ_{T+1}, v_{T+1}) has privacy cost as

$$
\hat{\rho}_{T+1} = \hat{\rho}_T + \frac{1}{2\sigma_T^2} = \cdots = \sum_{t=1}^{T} \frac{1}{2\sigma_t^2} = \frac{1}{2} \sum_{t=1}^{T} \rho_t \leq \frac{1}{2}(R - R_T) \leq \frac{1}{2}R.
$$

Let $\rho = \hat{\rho}_{T+1}$. Then we can get the conclusion. $\qquad\square$

C.1 GRADIENT DESCENTS

Proof of Theorem 4.1. With the definition of smoothness in Definition 3.3 and Eq. (1), we have

$$
\begin{aligned}
f(\theta_{t+1}) - f(\theta_t) &\leq -\eta_t \nabla_t^\top (\nabla_t + G\sigma_t \nu_t/N) + \frac{1}{2} M\eta_t^2 \left\| \nabla_t + G\sigma_t \nu_t/N \right\|^2 \\
&= -\eta_t (1 - \frac{1}{2} M\eta_t) \left\| \nabla_t \right\|^2 - (1 - M\eta_t)\eta_t \nabla_t^\top G\sigma_t \nu_t/N + \frac{1}{2} M\eta_t^2 \left\| G\sigma_t \nu_t/N \right\|^2 \\
&\leq -2\mu\eta_t (1 - \frac{1}{2} M\eta_t)(f(\theta_t) - f(\theta^*)) - (1 - M\eta_t)\eta_t \nabla_t^\top G\sigma_t \nu_t/N \\
&\quad + \frac{1}{2} M\eta_t^2 \left\| G\sigma_t \nu_t/N \right\|^2.
\end{aligned}
$$

where the last inequality is due to the Polyak-Lojasiewicz condition. Taking expectation on both sides, we can obtain

$$
\mathbb{E}[f(\theta_{t+1})] - \mathbb{E}[f(\theta_t)] \leq -2\mu\eta_t (1 - \frac{M}{2}\eta_t)(\mathbb{E}[f(\theta_t)] - f(\theta^*)) + \frac{M}{2}(\eta_t G\sigma_t/N)^2 \mathbb{E}\left\| \nu_t \right\|^2
$$

which can be reformulated by substacting $f(\theta^*)$ on both sides and re-arranged as

$$
\mathbb{E}[f(\theta_{t+1})] - f(\theta^*) \leq \left(1 - 2\mu\eta_t (1 - \frac{M}{2}\eta_t) \right) (\mathbb{E}[f(\theta_t)] - f(\theta^*)) + \frac{M}{2}(\eta_t G\sigma_t/N)^2 D
$$

Recursively using the inequality, we can get

$$
\begin{aligned}
\mathbb{E}[f(\theta_{T+1})] - f(\theta^*) &\leq \prod_{t=1}^{T} \left(1 - 2\mu\eta_t (1 - \frac{M}{2}\eta_t) \right) (\mathbb{E}[f(\theta_1)] - f(\theta^*)) \\
&\quad + \frac{MD}{2} \sum_{t=1}^{T} \prod_{i=t+1}^{T} \left(1 - 2\mu\eta_i (1 - \frac{M}{2}\eta_i) \right) (\eta_t G\sigma_t/N)^2.
\end{aligned}
$$

Let $\eta_t \equiv 1/M$. Then the above inequality can be simplified as

$$
\begin{aligned}
\mathbb{E}[f(\theta_{T+1})] - f(\theta^*) &\leq \gamma^T (\mathbb{E}[f(\theta_1)] - f(\theta^*)) + R \sum_{t=1}^{T} \gamma^{T-t} \frac{MD}{2R} \left(\frac{\eta_t G}{N} \right)^2 \sigma_t^2 \\
&= \gamma^T (\mathbb{E}[f(\theta_1)] - f(\theta^*)) + R \sum_{t=1}^{T} \gamma^{T-t} \alpha\sigma_t^2 (\mathbb{E}[f(\theta_1)] - f(\theta^*)) \\
&= \left(\gamma^T + R \sum_{t=1}^{T} q_t \sigma_t^2 \right) (f(\theta_1) - f(\theta^*))
\end{aligned}
$$

$\qquad\square$

Proof of Theorem 4.2. The minimizer of the upper bound of Eq. (3) can be written as

$$T^* = \arg\min_T \gamma^T + \alpha\kappa(1 - \gamma^T)T \tag{16}$$

where we substitute $\sigma^2 = T/R$ in the second line. To find the convex minimization problem, we need to vanishing its gradient which involves an equation like $T\gamma^T = c$ for some real constant c. However, the solution is $W_k(c)$ for some integer k where W is Lambert W function which does not have a simple analytical form. Instead, because $\gamma^T > 0$, we can minimize a surrogate upper bound as following

$$T^* = \arg\min_T \gamma^T + \alpha\kappa T = \frac{1}{\ln(1/\gamma)} \ln\left(\frac{\ln(1/\gamma)}{\kappa\alpha}\right), \text{ if } \kappa\alpha + \ln\gamma < 0 \tag{17}$$

where we use the surrogate upper bound in the second line and utilize $\gamma = 1 - \frac{1}{\kappa}$. However, the minimizer of the surrogate objective is not optimal for the original objective. When κ is large, the term, $-\alpha\kappa\gamma^T T$, cannot be neglected as we expect. On the other hands, T suffers from explosion if $\kappa \to \infty$ and meanwhile $1/\gamma \to_+ 1$. The tendency is counterintuitive since a small T should be taken for sharp losses. To fix the issue, we change the form of T^* as

$$T^* = \frac{1}{\ln(1/\gamma)} \ln\left(1 + \frac{\ln(1/\gamma)}{\alpha}\right), \tag{18}$$

which gradually converges to 0 as $\kappa \to \infty$.

Now we substitute Eq. (18) into the original objective function, Eq. (16), to get

$$\text{ERUB}^{\text{uniform}} = \frac{1}{1 + \ln(1/\gamma)/\alpha} \left[1 + \kappa\ln\left(1 + \frac{\ln(1/\gamma)}{\alpha}\right)\right]. \tag{19}$$

Notice that

$$\ln(1/\gamma) = \ln(\kappa/(\kappa - 1)) = \ln(1 + 1/(\kappa - 1)) \leq \frac{1}{\kappa - 1} \leq \frac{1}{c\kappa}$$

because $\kappa \geq \frac{1}{1-c} > 1$ for some constant $c \in (0, 1)$. In addition,

$$\ln(1/\gamma) = -\ln(1 - 1/\kappa) \geq 1/\kappa.$$

Now, we can get the upper bound of Eq. (19) as

$$\text{ERUB}^{\text{uniform}} \leq \frac{\kappa}{\kappa + 1/\alpha}\left[1 + \kappa\ln\left(1 + \frac{1}{c\kappa\alpha}\right)\right]$$

$$\leq c_1 \frac{\kappa}{\kappa + 1/\alpha}\kappa\left[\ln\left(1 + \frac{1}{\kappa\alpha}\right) + \ln(\frac{1}{c}))\right]$$

$$\leq c_1 c_2 \frac{\kappa^2}{\kappa + 1/\alpha}\ln\left(1 + \frac{1}{\kappa\alpha}\right)$$

for some constants c_1, c_2 and large enough $\frac{1}{\alpha}$. Also, we can get the lower bound

$$\text{ERUB}^{\text{uniform}} \geq \frac{c\kappa}{c\kappa + 1/\alpha}\left[1 + \kappa\ln\left(1 + \frac{1}{\kappa\alpha}\right)\right] \geq c\frac{\kappa^2}{\kappa + 1/\alpha}\ln\left(1 + \frac{1}{\kappa\alpha}\right).$$

where we use the condition $c \in (0, 1)$. Thus, $\text{ERUB}^{\text{uniform}} = \Theta\left(\frac{\kappa^2}{\kappa + 1/\alpha}\ln\left(1 + \frac{1}{\kappa\alpha}\right)\right)$. □

Proof of Lemma 4.1. By $\sum_{t=1}^T \sigma^{-2} = R$ and Cauchy-Schwarz inequality, we can derive the achievable lower bound as

$$R\sum_t q_t\sigma_t^2 = \sum_t \frac{1}{\sigma_t^2}\sum_t q_t\sigma_t^2 \geq \left(\sum_{t=1}^T \sqrt{q_t}\right)^2$$

where the inequality becomes equality if and only if $s/\sigma_t^2 = q_t\sigma_t^2$, i.e., $\sigma_t = (s/q_t)^{1/4}$, for some positive constant s. The equality $\sum_{t=1}^T \sigma_t^{-2} = R$ immediately suggests $\sqrt{s} = \frac{1}{R}\sum_{t=1}^T \sqrt{q_t}$. Thus, we get the σ_t. □

Proof of Theorem 4.3. The upper bound of Eq. (3) can be written as

$$\text{ERUB}^{\text{dyn}} = \gamma^T + \sum_{t=1}^{T} \gamma^{T-t} \alpha R \sigma^2$$

$$= \gamma^T + \alpha \left(\sum_{t=1}^{T} \sqrt{\gamma^{T-t}} \right)^2$$

$$= \gamma^T + \alpha \left(\frac{1 - \gamma^{T/2}}{1 - \sqrt{\gamma}} \right)^2$$

where we make use of Lemma 4.1. Then, the minimizer of the ERUB is

$$T^* = \arg\min_{T} \gamma^T + \alpha \left(\frac{1 - \gamma^{T/2}}{1 - \sqrt{\gamma}} \right)^2$$

$$= 2 \log_\gamma \left(\frac{\alpha}{\alpha + (1 - \sqrt{\gamma})^2} \right). \tag{20}$$

We can substitute Eq. (20) into ERUB^{dyn} to get

$$\text{ERUB}^{\text{dyn}}_{\text{min}} = \left(\frac{\alpha}{\alpha + (1 - \sqrt{\gamma})^2} \right)^2 + \alpha \left(\frac{1}{1 - \sqrt{\gamma}} \right)^2 \left(1 - \frac{\alpha}{\alpha + (1 - \sqrt{\gamma})^2} \right)^2$$

$$= \left(\frac{\alpha(1 - \sqrt{\gamma})^{-2}}{\alpha(1 - \sqrt{\gamma})^{-2} + 1} \right)^2 + \frac{\alpha(1 - \sqrt{\gamma})^{-2}}{\left(\alpha(1 - \sqrt{\gamma})^{-2} + 1 \right)^2}$$

$$= \frac{\alpha(1 - \sqrt{\gamma})^{-2}}{\alpha(1 - \sqrt{\gamma})^{-2} + 1}$$

Notice that $\left(1 - \sqrt{\gamma} \right)^{-2} = \kappa^2 + \kappa^2 - \kappa + 2\kappa\sqrt{\kappa(\kappa - 1)} = \kappa(2\kappa - 1 + 2\sqrt{\kappa(\kappa - 1)})$ and it is bounded by

$$\kappa(2\kappa - 1 + 2\sqrt{\kappa(\kappa - 1)}) \leq 4\kappa^2,$$

$$\kappa(2\kappa - 1 + 2\sqrt{\kappa(\kappa - 1)}) \geq \kappa(2\kappa - (3\kappa - 2) + 2\sqrt{(\kappa - 1)(\kappa - 1)}) = \kappa(-\kappa + 2 + 2\kappa - 2) = \kappa^2.$$

Therefore, $\kappa \leq \left(1 - \sqrt{\gamma} \right)^{-1} \leq 2\kappa$, with which we can derive

$$\text{ERUB}^{\text{dyn}}_{\text{min}} \leq 4 \frac{\kappa^2 \alpha}{\kappa^2 \alpha + 1},$$

$$\text{ERUB}^{\text{dyn}}_{\text{min}} \geq \frac{\kappa^2 \alpha}{4\kappa^2 \alpha + 1} \geq \frac{1}{4} \frac{\kappa^2 \alpha}{\kappa^2 \alpha + 1}.$$

Thus, $\text{ERUB}^{\text{dyn}}_{\text{min}} = \Theta \left(\frac{\kappa^2 \alpha}{\kappa^2 \alpha + 1} \right)$. □

C.2 GRADIENT DESCENTS WITH MOMENTUM

Proof of Theorem 4.4. Without loss of generality, we absorb the $C\sigma_t/N$ into the variance of ν_t such that $\nu_t \sim \mathcal{N}(0, \frac{C\sigma_t^2}{N} I)$ and $g_t \leftarrow \tilde{\nabla}_t + \nu_t$. Define $b_t = 1 - \beta^t$.

By smoothness and Eq. (1), we have

$$f(\theta_{t+1}) - f(\theta_t) \leq \nabla_t^\top (\theta_{t+1} - \theta_t) + \frac{1}{2} M \|\theta_{t+1} - \theta_t\|^2$$

$$= -\frac{\eta_t}{b_t^2} b_t \nabla_t^\top v_{t+1} + \frac{1}{2} M \frac{\eta_t^2}{b_t^2} \|v_{t+1}\|^2$$

$$= \frac{\eta_t}{b_t^2} \left(\|b_t \nabla_t - v_{t+1}\|^2 - \|b_t \nabla_t\|^2 - \|v_{t+1}\|^2 \right) + \frac{1}{2} M \frac{\eta_t^2}{b_t^2} \|v_{t+1}\|^2$$

$$= \frac{\eta_t}{b_t^2} \underbrace{\|b_t \nabla_t - v_{t+1}\|^2}_{U_1(t)} - \eta_t \|\nabla_t\|^2 - \frac{\eta_t}{b_t^2} (1 - \frac{1}{2} M \eta_t) \|v_{t+1}\|^2, \tag{21}$$

where only the $U_1(t)$ is non-negative. Specifically, $U_1(t)$ describes the difference between current gradient and the average. We can expand v_{t+1} to get an upper bound:

$$
\begin{aligned}
U_1(t) &= \|b_t \nabla_t - v_{t+1}\|^2 \\
&= \left\| (1-\beta) \sum_{i=1}^{t} \beta^{t-i} \nabla_t - (1-\beta) \sum_{i=1}^{t} \beta^{t-i} g_i \right\|^2 \\
&= (1-\beta)^2 \left\| \sum_{i=1}^{t} \beta^{t-i} (\nabla_t - g_i) \right\|^2 \\
&= (1-\beta)^2 \left\| \sum_{i=1}^{t} \beta^{t-i} (\nabla_t - \nabla_i) + \sum_{i=1}^{t} \beta^{t-i} (\nabla_i - g_i) \right\|^2 \\
&\leq 2(1-\beta)^2 \left[\left\| \sum_{i=1}^{t} \beta^{t-i} (\nabla_t - \nabla_i) \right\|^2 + \left\| \sum_{i=1}^{t} \beta^{t-i} (\nabla_i - g_i) \right\|^2 \right] \\
&\leq 2(1-\beta) \left[b_t \underbrace{\sum_{i=1}^{t} \beta^{t-i} \|\nabla_t - \nabla_i\|^2}_{U_2(t)\ \text{(gradient variance)}} + (1-\beta) \underbrace{\left\| \sum_{i=1}^{t} \beta^{t-i} \nu_i \right\|^2}_{\text{noise variance}} \right]
\end{aligned}
$$

where we use $\|x + y\|^2 \leq (\|x\| + \|y\|)^2 \leq 2(\|x\|^2 + \|y\|^2)$. The last inequality can be proved by Cauchy-Schwartz inequality for each coordinate.

We plug the $U_1(t)$ into Eq. (21) and use the PL condition to get

$$
\begin{aligned}
f(\theta_{t+1}) - f(\theta_t) &\leq \frac{\eta_t}{b_t^2} U_1(t) - \eta_t \|\nabla_t\|^2 - \frac{\eta_t}{b_t^2} \left(1 - \frac{1}{2} M \eta_t\right) \|v_{t+1}\|^2 \\
&\leq -\eta_t \|\nabla_t\|^2 + \frac{\eta_t}{b_t^2} 2(1-\beta) \left[b_t U_2(t) + (1-\beta) \left\| \sum_{i=1}^{t} \beta^{t-i} \nu_i \right\|^2 \right] \\
&\quad - \frac{\eta_t}{b_t^2} \left(1 - \frac{1}{2} M \eta_t\right) \|v_{t+1}\|^2 \\
&\leq -2\mu\eta_t (f(\theta_t) - f(\theta^*)) + \frac{2(1-\beta)\eta_t}{b_t} U_2(t) + \frac{2(1-\beta)^2 \eta_t}{b_t^2} \left\| \sum_{i=1}^{t} \beta^{t-i} \nu_i \right\|^2 \\
&\quad - \frac{\eta_t}{b_t^2} \left(1 - \frac{1}{2} M \eta_t\right) \|v_{t+1}\|^2 .
\end{aligned}
$$

Rearranging terms and taking expectation to show

$$
\begin{aligned}
\mathbb{E}[f(\theta_{t+1})] - f(\theta^*) &\leq \gamma(\mathbb{E}[f(\theta_t)] - f(\theta^*)) + \frac{2(1-\beta)^2 \eta_t}{b_t^2} \sum_{i=1}^{t} \beta^{t-i} \mathbb{E} \|\nu_i\|^2 \\
&\quad + \frac{2(1-\beta)\eta_t}{b_t} \mathbb{E}[U_2(t)] - \frac{\eta_t}{b_t^2} \left(1 - \frac{1}{2} M \eta_t\right) \mathbb{E} \|v_{t+1}\|^2 \\
&= \gamma(\mathbb{E}[f(\theta_t)] - f(\theta^*)) + \frac{2(1-\beta)^2 \eta_t}{b_t^2} \sum_{i=1}^{t} \beta^{2(t-i)} \frac{C^2 D \sigma_t^2}{N^2} \\
&\quad + \frac{2(1-\beta)\eta_t}{b_t} \mathbb{E}[U_2(t)] - \frac{\eta_t}{b_t^2} \left(1 - \frac{1}{2} M \eta_t\right) \mathbb{E} \|v_{t+1}\|^2
\end{aligned}
$$

where $\gamma = 1 - \eta_0/\kappa = 1 - 2\mu\eta_t$. The recursive inequality implies

$$\mathbb{E}[f(\theta_{T+1})] - f(\theta^*) \leq \gamma^T(f(\theta_1) - f(\theta^*)) + \sum_{t=1}^{T} \gamma^{T-t} \frac{2(1-\beta)^2 \eta_t}{b_t^2} \sum_{i=1}^{t} \beta^{2(t-i)} \frac{C^2 D \sigma_t^2}{N^2}$$

$$+ \sum_{t=1}^{T} \gamma^{T-t} \frac{2(1-\beta)\eta_t}{b_t} \mathbb{E}[U_2(t)] - \sum_{t=1}^{T} \gamma^{T-t} \frac{\eta_t}{b_t^2} (1 - \frac{1}{2} M \eta_t) \mathbb{E} \|v_{t+1}\|^2$$

$$= \left(\gamma^T + 2\eta_0 \alpha R \underbrace{\sum_{t=1}^{T} \gamma^{T-t} \frac{(1-\beta)^2}{b_t^2} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_t^2}_{U_3} \right) (f(\theta_1) - f(\theta^*))$$

$$+ \underbrace{\sum_{t=1}^{T} \gamma^{T-t} \frac{2(1-\beta)\eta_t}{b_t} \mathbb{E}[U_2(t)] - \sum_{t=1}^{T} \gamma^{T-t} \frac{\eta_t}{b_t^2} (1 - \frac{1}{2} M \eta_t) \mathbb{E} \|v_{t+1}\|^2}_{U_4(t)}.$$

where we utilize $\alpha = \frac{DC^2}{2MN^2 R} \frac{1}{f(\theta_1) - f(\theta^*)}$ and $\eta_t = \frac{\eta_0}{2M}$.

By [Lemma B.5](#), we have

$$\sum_{t=1}^{T} \gamma^{T-t} \frac{2(1-\beta)\eta_t}{b_t} U_2(t) \leq \frac{\eta_0^3 \beta \gamma}{2M(1-\beta)^3(\gamma-\beta)^2} \sum_{i=1}^{T-1} \gamma^{T-i} \|v_{i+1}\|^2.$$

Thus, by $\frac{1}{b_t} \geq 1$,

$$U_4(t) \leq \frac{\eta_0^3 \beta \gamma}{2M(1-\beta)^3(\gamma-\beta)^2} \sum_{i=1}^{T-1} \gamma^{T-i} \mathbb{E} \|v_{i+1}\|^2 - \frac{\eta_0}{2M} (1 - \frac{\eta_0}{4}) \sum_{t=1}^{T} \gamma^{T-t} \mathbb{E} \|v_{t+1}\|^2$$

$$= -\frac{\eta_0}{2M} \zeta \sum_{t=1}^{T} \gamma^{T-t} \mathbb{E} \|v_{t+1}\|^2$$

where

$$\zeta = 1 - \frac{1}{4}\eta_0 - \frac{\beta\gamma}{(\gamma-\beta)^2(1-\beta)^3} \eta_0^2 = 1 - \frac{1}{4}\eta_0 - \frac{\beta/\gamma}{(1-\beta/\gamma)^2(1-\beta)^3} \eta_0^2$$

When a small enough η_0, e.g., Specifically,

$$\eta_0 \leq \frac{(\gamma-\beta)^2(1-\beta)^3}{8\beta\gamma} \left[\sqrt{1 + \frac{64\beta\gamma}{(\gamma-\beta)^2(1-\beta)^3}} - 1 \right]$$

$$= \frac{8}{\sqrt{1 + 64\beta\gamma(\gamma-\beta)^{-2}(1-\beta)^{-3}} + 1}$$

We can have $\zeta \geq 0$.

By the definition of $U_3(T, \sigma)$, we can get

$$\mathbb{E}[f(\theta_{T+1})] - f(\theta^*) \leq \left(\gamma^T + 2\eta_0 \alpha R U_3(T, \sigma) \right) (f(\theta_1) - f(\theta^*)) - \frac{\eta_0}{2M} \zeta \sum_{t=1}^{T} \gamma^{T-t} \mathbb{E} \|v_{t+1}\|^2.$$

$\square$

Proof of Theorem 4.5. Since σ_t is static, by definition of U_3 in Theorem 4.4,

$$U_3 = \sum_{t=1}^{T} \gamma^{T-t} \frac{(1-\beta)^2}{(1-\beta^t)^2} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma^2$$

$$= \sigma^2 \sum_{t=1}^{T} \gamma^{T-t} \frac{(1-\beta)^2}{(1-\beta^t)^2} \sum_{i=1}^{t} \beta^{2(t-i)}$$

$$= \sigma^2 \sum_{t=1}^{T} \gamma^{T-t} \frac{(1-\beta)^2}{(1-\beta^t)^2} \frac{1-\beta^{2t}}{1-\beta^2}$$

$$= \sigma^2 \sum_{t=1}^{T} \gamma^{T-t} \frac{1-\beta}{1-\beta^t} \frac{1+\beta^t}{1+\beta}.$$

Because $\frac{1-\beta}{1-\beta^t} \frac{1+\beta^t}{1+\beta} \leq 1$, the U_3 will be smaller than the corresponding summation in GD with uniform schedule.

By Lemma B.7, when $T > \hat{T}$, we can rewrite U_3 as

$$U_3 \leq \sigma^2 \sum_{t=1}^{T} \gamma^{T-t} \frac{1-\beta}{1-\beta^t}$$

$$= \sigma^2 \sum_{t=1}^{\hat{T}} \gamma^{T-t} \frac{1-\beta}{1-\beta^t} + \sigma^2 \sum_{t=\hat{T}+1}^{T} \gamma^{T-t} \frac{1-\beta}{1-\beta^t}$$

$$\leq \sigma^2 \sum_{t=1}^{\hat{T}} \gamma^{T-t} \gamma^{t-1} + \sigma^2 \sum_{t=\hat{T}+1}^{T} \gamma^{T-t} \gamma^{\hat{T}-1}$$

$$= \sigma^2 \gamma^{T-1} \hat{T} + \sigma^2 \gamma^{\hat{T}-1} \sum_{t=1}^{T-\hat{T}} \gamma^{T-\hat{T}-t}$$

$$= \sigma^2 \gamma^{T-1} \hat{T} + \sigma^2 \frac{\gamma^{\hat{T}-1} - \gamma^{T-1}}{1-\gamma}$$

$$= \frac{T}{\gamma R} \gamma^T \left(\hat{T} + \frac{\gamma^{\hat{T}-T} - 1}{1-\gamma} \right)$$

where we use $\sigma^2 = T/R$ in the last line. Without assuming $T > \hat{T}$, we can generally write the upper bound as

$$U_3 \leq \frac{T}{\gamma R} \gamma^T \left(\min\{\hat{T}, T\} + \max\{\frac{\gamma^{\hat{T}-T} - 1}{1-\gamma}, 0\} \right).$$

By Theorem 4.4, because $\zeta \geq 0$, we have

$$\text{ERUB} \leq \gamma^T + 2R\eta_0 \alpha U_3$$

$$= \gamma^T (1 + \frac{\alpha'}{\gamma} T \left(\min\{\hat{T}, T\} + \max\{\frac{\gamma^{\hat{T}-T} - 1}{1-\gamma}, 0\} \right))$$

where $\alpha' = 2\eta_0 \alpha$.

First, we consider $T \leq \hat{T}$. Use $T = \frac{1}{\ln(1/\gamma)} \ln \left(1 + \frac{\eta_0}{\kappa \alpha}\right) = \left\lceil \mathcal{O} \left(\frac{\kappa}{\eta_0} \ln \left(1 + \frac{\eta_0}{\kappa \alpha}\right) \right) \right\rceil$ to get

$$\text{ERUB} \leq \left(\frac{\alpha}{\alpha + \eta_0/\kappa} \right) \left(1 + \alpha' \gamma^{-1} (\frac{2}{\ln(1/\gamma)} \ln \left(1 + \frac{\eta_0}{\kappa \alpha}\right))^2 \right)$$

$$\leq \left(\frac{\alpha}{\alpha + \eta_0/\kappa} \right) \left(1 + \frac{8\kappa^2 \alpha}{\eta_0 \gamma} \ln^2 \left(1 + \frac{\eta_0}{\kappa \alpha}\right) \right)$$

$$\leq \mathcal{O} \left(\frac{\kappa}{\kappa + \eta_0/\alpha} \left(1 + \frac{8\kappa^2 \alpha}{\eta_0 \gamma} \ln^2 \left(1 + \frac{\eta_0}{\kappa \alpha}\right) \right) \right)$$

$$= \mathcal{O} \left(\frac{\kappa}{\kappa + \eta_0/\alpha} \left(1 + \frac{4\kappa}{\gamma} \right) \right)$$

$$= \mathcal{O} \left(\frac{\kappa^2}{\kappa + \eta_0/\alpha} \right)$$

where we used $\ln(1/\gamma) \geq \eta_0/\kappa$ and $\ln(1+x) \leq \sqrt{x}$ for any $x > 0$.

Second, when $T > \hat{T}$,

$$\text{ERUB} \leq \gamma^T (1 + \frac{\alpha'}{\gamma} T \left(\hat{T} + \frac{\gamma^{\hat{T}-T} - 1}{1 - \gamma} \right))$$

$$\leq \mathcal{O} \left(\gamma^T + \frac{2\alpha'}{\gamma} T \kappa (\gamma^{\hat{T}} - \gamma^T) \right).$$

Make use of $T = \left\lceil \frac{1}{\ln(1/\gamma)} \ln \left(1 + \frac{\eta_0}{\kappa\alpha} \right) \right\rceil$ to show

$$\text{ERUB} \leq \mathcal{O} \left(\frac{\kappa}{\kappa + \eta_0/\alpha} + \frac{4\kappa^2\alpha}{\eta_0\gamma} (\gamma^{\hat{T}} - \frac{\kappa}{\kappa + \eta_0/\alpha}) \ln \left(1 + \frac{\eta_0}{\kappa\alpha} \right) \right)$$

$$\leq \mathcal{O} \left(\frac{\kappa^2}{\kappa + \eta_0/\alpha} \gamma^{\hat{T}-1} \ln \left(1 + \frac{\eta_0}{\kappa\alpha} \right) \right).$$

$\square$

Proof of Theorem 4.6. By Lemma B.7, we can rewrite U_3 as

$$U_3 = \sum_{t=1}^{T} \gamma^{T-t} \frac{(1-\beta)^2}{(1-\beta^t)^2} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2$$

$$\leq \sum_{t=1}^{\hat{T}} \gamma^{T-t} \gamma^{2(t-1)} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2 + \sum_{t=\hat{T}+1}^{T} \gamma^{T-t} \gamma^{2(\hat{T}-1)} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2$$

$$\leq \gamma^{T-\hat{T}} \underbrace{\sum_{t=1}^{\hat{T}} \gamma^{\hat{T}-t} \gamma^{2(t-1)} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2}_{V_1} + \gamma^{2(\hat{T}-1)} \underbrace{\sum_{t=\hat{T}+1}^{T} \gamma^{T-t} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2}_{V_2}$$

We derive V_1 and V_2 separately.

For V_1, we can obtain the upper bound by

$$V_1 = \sum_{t=1}^{\hat{T}} \gamma^{\hat{T}-t} \gamma^{2(t-1)} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2$$

$$= \gamma^{\hat{T}-2} \sum_{t=1}^{\hat{T}} \gamma^t \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2$$

$$= \gamma^{\hat{T}-2} \sum_{i=1}^{\hat{T}} \beta^{-2i} \sigma_i^2 \sum_{t=i}^{\hat{T}} \left(\gamma\beta^2 \right)^t$$

$$= \gamma^{\hat{T}-2} \sum_{i=1}^{\hat{T}} \beta^{-2i} \sigma_i^2 \frac{\left(\gamma\beta^2 \right)^i - \left(\gamma\beta^2 \right)^{\hat{T}+1}}{1 - \gamma\beta^2}$$

$$= \gamma^{2\hat{T}-3} \sum_{i=1}^{\hat{T}} \frac{\gamma^{i-\hat{T}-1} - \beta^{2(\hat{T}+1-i)}}{1 - \gamma\beta^2} \sigma_i^2$$

$$= \gamma^{2\hat{T}-3} \sum_{i=1}^{\hat{T}} \frac{1 - (\gamma\beta^2)^{\hat{T}+1-i}}{1 - \gamma\beta^2} \gamma^{i-\hat{T}-1} \sigma_i^2$$

$$\leq \frac{\gamma^{\hat{T}}}{\gamma^2 (1 - \gamma\beta^2)} \sum_{i=1}^{\hat{T}} \gamma^i \sigma_i^2$$

$$\leq \frac{\gamma^{\hat{T}}}{\gamma(\gamma - \beta^2)} \sum_{i=1}^{\hat{T}} \gamma^i \sigma_i^2$$

For V_2, we can derive

$$V_2 = \sum_{t=\hat{T}+1}^{T} \gamma^{T-t} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2$$

$$= \sum_{t=1}^{T} \gamma^{T-t} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2 - \sum_{t=1}^{\hat{T}} \gamma^{T-t} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2$$

$$= \sum_{t=1}^{T} \gamma^{T-t} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2 - \gamma^{T-\hat{T}} \sum_{t=1}^{\hat{T}} \gamma^{\hat{T}-t} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2.$$

We first consider the first term

$$
\sum_{t=1}^{T} \gamma^{T-t} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2
$$
$$
= \sum_{i=1}^{T} \sigma_i^2 \sum_{t=i}^{T} \gamma^{T-t} \beta^{2(t-i)}
$$
$$
= \sum_{i=1}^{T} \gamma^T \beta^{-2i} \sigma_i^2 \sum_{t=i}^{T} \gamma^{-t} \beta^{2t}
$$
$$
= \sum_{i=1}^{T} \gamma^T \beta^{-2i} \sigma_i^2 \frac{(\beta^2/\gamma)^i - (\beta^2/\gamma)^{T+1}}{1 - (\beta^2/\gamma)}
$$
$$
= \sum_{i=1}^{T} \frac{\gamma^{T+1-i} - \beta^{2(T+1-i)}}{\gamma - \beta^2} \sigma_i^2.
$$

Similarly, we have

$$
\gamma^{T-\hat{T}} \sum_{t=1}^{\hat{T}} \gamma^{\hat{T}-t} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2
$$
$$
= \gamma^{T-\hat{T}} \sum_{i=1}^{\hat{T}} \frac{\gamma^{\hat{T}+1-i} - \beta^{2(\hat{T}+1-i)}}{\gamma - \beta^2} \sigma_i^2
$$
$$
= \sum_{i=1}^{\hat{T}} \frac{\gamma^{T+1-i} - \gamma^{T-\hat{T}} \beta^{2(\hat{T}+1-i)}}{\gamma - \beta^2} \sigma_i^2.
$$

Thus,

$$
V_2 = \sum_{i=1}^{T} \frac{\gamma^{T+1-i} - \beta^{2(T+1-i)}}{\gamma - \beta^2} \sigma_i^2 - \sum_{i=1}^{\hat{T}} \frac{\gamma^{T+1-i} - \gamma^{T-\hat{T}} \beta^{2(\hat{T}+1-i)}}{\gamma - \beta^2} \sigma_i^2
$$
$$
= \sum_{i=\hat{T}+1}^{T} \frac{\gamma^{T+1-i} - \beta^{2(T+1-i)}}{\gamma - \beta^2} \sigma_i^2 + \sum_{i=1}^{\hat{T}} \frac{\gamma^{T-\hat{T}} - \beta^{2(T-\hat{T})}}{\gamma - \beta^2} \beta^{2(\hat{T}+1-i)} \sigma_i^2
$$
$$
\leq \sum_{i=\hat{T}+1}^{T} \frac{\gamma^{T+1-i} - \beta^{2(T+1-i)}}{\gamma - \beta^2} \sigma_i^2 + \sum_{i=1}^{\hat{T}} \frac{\gamma^{T-\hat{T}}}{\gamma - \beta^2} \beta^{2(\hat{T}+1-i)} \sigma_i^2.
$$

Substitute V_1 and V_2 into U_3 to get

$$
U_3 \leq \gamma^T \frac{1}{\gamma(\gamma - \beta^2)} \sum_{i=1}^{\hat{T}} \gamma^i \sigma_i^2 + \gamma^{2\hat{T}-2} \sum_{i=\hat{T}+1}^{T} \frac{\gamma^{T+1-i} - \beta^{2(T+1-i)}}{\gamma - \beta^2} \sigma_i^2
$$
$$
+ \sum_{i=1}^{\hat{T}} \frac{\gamma^{T+\hat{T}-2}}{\gamma - \beta^2} \beta^{2(\hat{T}+1-i)} \sigma_i^2
$$
$$
\leq \left(\frac{\gamma^T}{\gamma(\gamma - \beta^2)} \sum_{i=1}^{\hat{T}} (\gamma^i + \gamma^{\hat{T}-1} \beta^{2(\hat{T}+1-i)}) \sigma_i^2 + \gamma^{2\hat{T}-2} \sum_{i=\hat{T}+1}^{T} \frac{\gamma^{T+1-i} - \beta^{2(T+1-i)}}{\gamma - \beta^2} \sigma_i^2 \right)
$$
$$
\leq \left(\frac{2\gamma^T}{\gamma(\gamma - \beta^2)} \sum_{i=1}^{\hat{T}} \gamma^i \sigma_i^2 + \gamma^{2\hat{T}-2} \sum_{i=\hat{T}+1}^{T} \frac{\gamma^{T+1-i} - \beta^{2(T+1-i)}}{\gamma - \beta^2} \sigma_i^2 \right)
$$
$$
= \sum_{t=1}^{T} q_t \sigma_t^2
$$

where

$$
q_t = \frac{2}{\gamma(\gamma - \beta^2)} \gamma^{T+t} \mathbb{I}_{T \leq \hat{T}} + \gamma^{2(\hat{T}-1)} \frac{\gamma^{T+1-i} - \beta^{2(T+1-i)}}{\gamma - \beta^2} \gamma^{T-t} \mathbb{I}_{T > \hat{T}}
$$
$$
\leq c_1 \gamma^{T+t} \mathbb{I}_{T \leq \hat{T}} + \gamma^{\hat{T}-1} c_2 \gamma^{T-t} \mathbb{I}_{T > \hat{T}}
$$

where $c_1 = \frac{2}{\gamma(\gamma - \beta^2)}$ and $c_2 = \frac{\gamma^{2\hat{T}}}{\gamma - \beta^2}$.

When $T > \hat{T}$, by Lemma 4.1, the lower bound of $R \sum_{t=1}^{T} q_t \sigma_t^2$ is

$$\left(\sum_{t=1}^{T} \sqrt{q_t} \right)^2 = \gamma^T \left(\sum_{t=1}^{\hat{T}} \sqrt{c_1 \gamma^t} + \sum_{t=\hat{T}+1}^{T} \sqrt{\gamma^{\hat{T}-1} c_2 \gamma^{-t}} \right)^2$$

$$= \gamma^T \left(\sqrt{c_1 \gamma} \frac{1 - \gamma^{\hat{T}/2}}{1 - \sqrt{\gamma}} + \sqrt{c_2} \frac{1 - \gamma^{(\hat{T}-T-1)/2}}{\sqrt{\gamma} - 1} \right)^2$$

$$= \gamma^T \left(\sqrt{c_1 \gamma} \frac{1 - \gamma^{\hat{T}/2}}{1 - \sqrt{\gamma}} + \sqrt{c_2} \frac{\gamma^{(\hat{T}-T-1)/2} - 1}{1 - \sqrt{\gamma}} \right)^2$$

$$\leq \mathcal{O} \left(c_2 \left\{ \frac{\gamma^{(\hat{T}-1)/2} - \gamma^{T/2}}{1 - \sqrt{\gamma}} \right\}^2 \right)$$

which is achieved when

$$\sigma_t^2 = \frac{1}{R} \sum_{i=1}^{T} \sqrt{\frac{q_i}{q_t}}.$$

By Theorem 4.4, because $\zeta \geq 0$, we have

$$\text{ERUB} \leq \gamma^T + 2R\eta_0 \alpha U_3$$

$$= \gamma^T + 2\eta_0 \alpha \sum_{t=1}^{T} R q_t \sigma_t^2.$$

And the minimum of the upper bound is

$$\text{ERUB}_{\min} = \gamma^T + \alpha' \mathcal{O} \left(\left\{ \frac{\gamma^{(\hat{T}-1)/2} - \gamma^{T/2}}{1 - \sqrt{\gamma}} \right\}^2 \right)$$

where $\alpha' = 2\eta_0 c_2 \alpha$. Let $T = \frac{2}{\ln(1/\gamma)} \ln \left(1 + \frac{\eta_0}{\kappa \alpha} \right)$. Then,

$$\text{ERUB}_{\min} = \mathcal{O} \left(\left(\frac{\kappa \alpha}{\kappa \alpha + \eta_0} \right)^2 + \frac{\alpha'}{(1 - \sqrt{\gamma})^2} \left\{ \frac{\gamma^{(\hat{T}-1)/2} - (1 - \gamma^{(\hat{T}-1)/2})\kappa \alpha}{\kappa \alpha + \eta_0} \right\}^2 \right)$$

$$\leq \mathcal{O} \left(\left(\frac{\kappa \alpha}{\kappa \alpha + \eta_0} \right)^2 + \frac{2\eta_0 c_2 \alpha}{(1 - \sqrt{\gamma})^2} \left\{ \frac{\gamma^{(\hat{T}-1)/2}}{\kappa \alpha + \eta_0} \right\}^2 \right)$$

$$= \mathcal{O} \left(\frac{\kappa \alpha}{(\kappa \alpha + \eta_0)^2} \left(\kappa \alpha + \frac{2\eta_0 c_2/\kappa}{(1 - \sqrt{\gamma})^2} \gamma^{(\hat{T}-1)} \right) \right)$$

$$= \mathcal{O} \left(\frac{\kappa \alpha}{(\kappa \alpha + \eta_0)^2} (\kappa \alpha + c_3 \eta_0) \right)$$

$$\leq \mathcal{O} \left(\frac{\kappa \alpha}{\kappa \alpha + \eta_0} \right)$$

where c_3 is some constant.

When $T \leq \hat{T}$,

$$U_3 \leq \gamma^{T-T} \underbrace{\sum_{t=1}^{T} \gamma^{T-t} \gamma^{2(t-1)} \sum_{i=1}^{t} \beta^{2(t-i)} \sigma_i^2}_{V_1}$$

$$\leq \frac{\gamma^{T-2}}{1 - \gamma \beta^2} \sum_{i=1}^{T} \gamma^i \sigma_i^2$$

with which we obtain

$$\text{ERUB} \leq \gamma^T + 2R\eta_0 \alpha U_3$$

$$\leq \gamma^T + 2\eta_0 \alpha \frac{\gamma^{-2}}{1 - \gamma \beta^2} \sum_{t=1}^{T} R q_t \sigma_t^2.$$

where we let $q_t = \gamma^{T+t}$. By Lemma 4.1,

$$\sum_{i=1}^{T} R q_t \sigma_i^2 \geq \left(\sum_{t=1}^{T} \sqrt{q_t} \right)^2$$

$$= \gamma^T \left(\sum_{t=1}^{T} \gamma^{t/2} \right)^2$$

$$= \gamma^{T+1} \left(\frac{1 - \gamma^{T/2}}{1 - \sqrt{\gamma}} \right)^2.$$

Thus,

$$\text{ERUB}_{\min} \leq \gamma^T + 2\eta_0 \alpha \frac{\gamma^{T-1}}{1 - \gamma\beta^2} \left(\frac{1 - \gamma^{T/2}}{1 - \sqrt{\gamma}} \right)^2$$

$$= \gamma^T \left(1 + 2\eta_0 \gamma c_1 \alpha \left(\frac{1 - \gamma^{T/2}}{1 - \sqrt{\gamma}} \right)^2 \right)$$

Let $T = \left\lceil \frac{2}{\ln(1/\gamma)} \ln \left(1 + \frac{\eta_0}{\kappa\alpha} \right) \right\rceil$. Then,

$$\text{ERUB}_{\min} \leq \left(\frac{\kappa\alpha}{\kappa\alpha + \eta_0} \right)^2 \left(1 + \frac{2\eta_0 \gamma c_1 \alpha}{(1 - \sqrt{\gamma})^2} (\frac{1}{\kappa\alpha + 1})^2 \right)$$

$$\leq \left(\frac{\kappa\alpha}{\kappa\alpha + \eta_0} \right)^2 \left(1 + \mathcal{O}(\frac{1}{\kappa\alpha + 1}) \right)$$

$$\leq \mathcal{O} \left(\frac{\kappa\alpha}{\kappa\alpha + \eta_0} \right)^2.$$

In summary,

$$\text{ERUB}_{\min} \leq \mathcal{O} \left(\frac{\kappa\alpha}{\kappa\alpha + \eta_0} \left(\mathbb{I}_{T \leq \hat{T}} \frac{\kappa\alpha}{\kappa\alpha + \eta_0} + \mathbb{I}_{T > \hat{T}} \right) \right)$$

$\square$

D   Experiments

Dataset. **(1) Synthetic data**. We generate a 100-dimensional dataset including linearly separable data points using `sckit-learn` package. The data points are distributed in two points in the hyper-cubic with Euclidean distance of 10. In total, 1000 samples are generated for training with the logistic loss. **(2) Real data**. We create a subset of the MNIST dataset Lecun et al. (1998) including 1000 handwritten images of digit 3 and 5 (MNIST35). Compared to the original dataset (70, 000 samples), the small set will be more vulnerable to attack and the private learning will require larger noise (see the $1/N$ factor in Eq. (1)). Following the preprocessing in Abadi et al. (2016), we project the vectorized images into a 60-dimensional subspace extracted by PCA.

Setup. The samples are first normalized so that $\sum_{n=1}^{N} x_n = 0$ and the standard deviation is 1. Then the sample norms are scaled such that $\max_n \|x_n\| = 10$ (i.e., *data scales*). We fix the learning rate to 0.1 based on the corresponding experiments of non-private training (same setting without noise). The total privacy budget is 0.1963-zCDP (equal to $(4, 10^{-8})$-DP) which implies $R = 0.3927$. To control the sensitivity of the gradients, we clip gradients by a clipping norm fixed at 4. Formally, we scale down the sample gradients to length 4 if its norm is larger than 4. Because the schedule highly depends on the iteration number T, we grid search the best T for compared methods. Therefore, we ignore the privacy cost of such tuning in our experiments which protocol is also used in previous work Abadi et al. (2016); Wu et al. (2017). All the experiments are repeated 100 times and metrics are averaged afterwards.

Results. In the left pane of Fig. 2, the squared final loss (i.e., $f(\theta_{T+1})$) is approximately a quadratic function of the σ_t with small relative variance (i.e., the variance divided by the mean

Figure 2: Experiments with Logistic loss on synthetic data and squared, Logistic loss and DNN on MNIST35 by rows. (**left**) The final loss is fitted by a quadratic function formulated as $c_2\sigma_t^2 + c_0$. (**middle**) The influence values are estimated using the Hessian eigenvalues (squared loss) and by retraining (logistic loss). The larger is the data scale, the larger the influence variance. (**right**) The relative final training losses versus the data scale where uniform schedule (uni), dynamic schedule (dyn) and exponential-decaying schedule (exp) are compared. The relative loss is computed w.r.t. the losses of the uniform schedule. For example, if the dynamic loss is e_1 and the uniform loss is e_0, then the relative loss is $(e_1 - e_0)/e_0$. The dashed lines are with the right axis to present influence-related term.

value, $0.14, 0.032, 0.016$ for $t = 10, 60, 90$, respectively). When σ_t increases, the frequency of clipping increases as well which leads to more variance in the final losses. We use the least square method to fit the relation shown as the solid lines. The final logistic loss is more sensitive to the noise because of the additional uncertainty from the changing Hessian. We still fit a quadratic function on σ_t. It turns out the relative variances of the quadratic coefficient (approximately the influence) is small which are $0.025, 0.024, 0.027$ for $t = 10, 60, 90$, respectively.

The middle pane shows the relationship between the estimated q_t and learning steps. The q_t of squared loss is computed by analytical solution using the Hessian eigen values. The q_t of logistic loss is computed by retraining based on uniform schedule. Both loss functions show an increased influence as learning continues, which indicates that the dynamic schedule should be decreasing accordingly. The squared loss has a rather steep trend while the logistic has a relative flatten one. The reason is that the logistic loss has a larger variance in the gradient norm and therefore clipping happens more frequently (approximately more than 80% gradients are clipped). As a result, the variance of influence will be relative small for logistic losses. Moreover, we vary the data scale (scale all samples uniformly such that all sample norms are less than a specific value), changing the variance of influence, as seen in the figures.

The last pane compares uniform, dynamic and exponential decay schedules Yu et al. (2019) using final losses relative to the uniform schedule. We set the exponential decay schedule to approximate the dynamic schedule by fitting $\hat{\sigma}_t = \sigma_0 \exp(-kt)$ using the least squares method. The final losses are picked by grid searching the best $T \in [1, 100]$. We see that the advantage of the dynamic schedule over the uniform one increases when data scales less than 15. But we also notice some loss gaps decrease, suggesting that the data scale is not the inherent reason for the dynamic advantage. According to Lemma 4.1, the advantage should be proportional to $T^2 \operatorname{Var}(q_t)$ which is shown to be decreasing when the data scale is larger than 15. When the data scale continues to increase, gradient clipping will change the curvature of the loss function. Therefore, a increasing $\operatorname{Var}(Tq_t)$ is witnessed. Meanwhile, the loss gap decreases.

In the last row of Fig. 2, the DNN is experimented with the same setting. Though the influence increases by t, the variance is small and less dependent on the data scale in comparison to shallow models. Though the dynamic schedule estimated by retraining influences does not performs stably, the *exp* method still decreasingly depends on the variance of influence as expectation.