Beyond Classification: A Cough Regression Benchmark for Respiratory Acoustic Foundation Models

Mayur Sanap [1] [2] **Prasanna Desikan** [1] **Edgar Lobaton** [2]

Abstract

Respiratory acoustic foundation models (FMs) excel at cough classification, yet their ability to predict continuous health quantities from cough audio remains largely unexplored, despite the clinical value of passive age, BMI, and disease-probability estimation in settings where physical measurements are unavailable. We introduce the multi-model, multi-target cough regression benchmark evaluating five FMs (OPERA-CT, OPERA-CE, OPERA-GT, HEAR, M2D+RESP) across six targets on three datasets under subject-disjoint protocols, comparing linear, MLP-small, and full MLP regression heads. MLP-small beats the mean-predictor baseline on all tasks and linear probing in 23 of 30 model × task cases, with full MLP overfitting on small clinical data but recovering on larger sets, revealing a dataset-size × head-capacity trade-off. HEAR leads within-dataset age regression on Coswara (9.12 yr MAE); its CIDRZ result is excluded from headline claims owing to possible HEAR–CIDRZ pretraining overlap. OPERA-GT is favored over OPERA-CT on age in all three datasets, with the CIDRZ margin within seed variance, extending a generative-pretraining advantage from breath to cough. HEAR and M2D+RESP reach near-full performance at $N = 50$ samples while OPERA models require $N = 400$. Cross-dataset transfer is strongly asymmetric as large diverse data generalises to small clinical populations (CoughVID $\rightarrow$ CIDRZ: -0.17 yr) but not vice versa (CIDRZ $\rightarrow$ Coswara: $+2.43$ yr, $+26.6\%$).

[1]Centific Global Solutions Inc. [2]Department of Electrical and Computer Engineering, North Carolina State University, Raleigh, NC, USA. Correspondence to: Mayur Sanap <msanap@ncsu.edu>.

Proceedings of the Workshop on Structured Data for Health at the 43^{rd} International Conference on Machine Learning, Seoul, South Korea.

1. Introduction

Cough acoustics encode physiological state beyond categorical disease labels, with spectrotemporal features reflecting airway geometry, respiratory muscle strength, and mucosal viscosity, all of which covary quantitatively with age, body composition, and disease severity (Sharan et al., 2018; Xu et al., 2022; Rudraraju et al., 2020). In low- and middle-income countries (LMICs) where respiratory disease burden is highest (World Health Organization, 2021), birth records, weighing equipment, and radiological capacity are often unavailable, making passive cough-based estimates of age and BMI actionable proxies for triage in settings where physical measurements are unavailable.

Foundation models (FMs) pretrained on large unlabeled audio corpora learn task-agnostic embeddings that transfer efficiently via linear probing (Zhang et al., 2024; Baur et al., 2024; Niizumi et al., 2025), reducing the labeled-data burden in clinical audio AI. The three leading respiratory FM families, OPERA (Zhang et al., 2024), HEAR (Baur et al., 2024), and M2D+RESP (Niizumi et al., 2025), have been benchmarked extensively on classification, but their regression capability is almost entirely uncharacterised.

The HEAR paper (Baur et al., 2024) evaluates age and BMI regression from cough on the CIDRZ and Coswara (Bhattacharya et al., 2023) datasets, but restricts the analysis to a single model with a fixed linear probe and per-device splits, with no multi-model comparison, non-linear head evaluation, or cross-dataset generalisation. The OPERA benchmark (Zhang et al., 2024) includes regression tasks (spirometry estimation) exclusively from deep-breath and vowel sounds, not from cough. M2D+RESP (Niizumi et al., 2025) has never been evaluated on cough regression task.

We address these gaps with five contributions.

1. **Multi-model, multi-target cough regression benchmark** evaluating five FMs across six targets (age, BMI, X-ray abnormality, TB probability) on three datasets under subject-disjoint protocols, with MAE reported alongside the mean-predictor baseline (MAD).

2. **Regression head comparison** showing MLP-small wins 23 of 30 model × task combinations. Full

Figure 1. Cough regression benchmark pipeline. All audio is resampled to 16 kHz mono and padded/trimmed to 2 s. Five frozen encoders feed three regression probes and outputs cover three evaluation regimes. OPERA† comprises OPERA-CT, OPERA-CE, and OPERA-GT.

MLP overfits on small clinical data (+0.53 yr for M2D+RESP on CIDRZ) but recovers on larger ones.

3. **Generative pretraining advantage** where OPERA-GT outperforms OPERA-CT on age regression in all three datasets (CIDRZ 10.49 vs. 10.52 yr, Coswara 10.16 vs. 10.25 yr, CoughVID 9.62 vs. 9.79 yr; the CIDRZ gap lies within seed variance), extending the breath-regression finding of Zhang et al. (2024) to cough.

4. **Cross-dataset transfer asymmetry** where large web-collected data generalises to small clinical populations (CoughVID → CIDRZ −0.17 yr) while the reverse fails (CIDRZ → Coswara +2.43 yr, +26.6%).

5. **Low-data regime analysis** showing HEAR and M2D+RESP reach near-full performance at $N = 50$ samples while OPERA models require $N = 400$, indicating pretraining corpus diversity determines low-data regression performance.

The rest of this paper is organized as follows. Section 2 describes the benchmark design. Section 3 presents results and Section 4 summarises findings and future directions.

2. Benchmark Design

We evaluate five frozen FMs on three cough datasets under a unified pipeline (Figure 1), resampling all audio to 16 kHz mono and padding or trimming to 2 s, with embeddings extracted once and shared across all regimes.

2.1. Datasets and Tasks

Table 1 summarises the three datasets and six regression targets. Subject-disjoint splits (64/16/20%) are used for CIDRZ and Coswara; CoughVID reuses the official UUID-level split (train = 3050, val = 1019, test = 2789).

CIDRZ (Baur et al., 2024; Google Health AI, 2024) ($N = 1049$) contains smartphone-recorded volitional coughs from patients at a TB clinic in Zambia, with labels from clinical assessments covering age (37.1 ± 12.9 yr), BMI (21.6 ± 5.3 kg/m^2), chest X-ray abnormality (0.46 ± 0.35),

Table 1. Dataset and task summary. Prob. = prob. in [0, 1].

Dataset	Task	Unit	N
CIDRZ (Google Health AI, 2024) Zambia	Age BMI X-ray Abn. TB probability	yr kg/m^2 prob. prob.	1049
Coswara (Bhattacharya et al., 2023) India	Age	yr	2560
CoughVID (Orlandic et al., 2021) Global	Age	yr	6858

Table 2. Foundation models. All receive 2 s at 16 kHz. RC = respiratory clips. † 8.18 s positional grid; padded to 2 s.

Model	Architecture	Dim	Pretraining
OPERA-CT	Contrastive Trans.	768	136K RC
OPERA-CE	Contrastive CNN	1280	136K RC
OPERA-GT	Generative MAE†	384	136K RC
HEAR	ViT-L MAE	512	313M health
M2D+RESP	Masked Mod. + Resp	3840	AudioSet+RC

and TB probability (0.62 ± 0.24). The X-ray abnormality and TB targets are continuous derived scores, not the binary radiological and microbiological diagnoses; we therefore treat them as score-reproduction tasks, not clinical-endpoint prediction (Google Health AI, 2024).

Coswara (Bhattacharya et al., 2023) ($N = 2560$) consists of shallow-cough recordings collected remotely via a web application across India during the COVID-19 pandemic, with self-reported age labels (35.1 ± 13.9 yr, range 7–87).

CoughVID (Orlandic et al., 2021) ($N = 6858$) comprises volitional cough recordings submitted globally via smartphone with continuous age labels (34.5 ± 12.7 yr, range 5–97) and the largest training set (3,050 samples).

2.2. Foundation Models

We evaluate five frozen FMs spanning three pretraining paradigms: respiratory-specific SSL (OPERA), large-scale health audio masked autoencoding (HEAR), and gen-

Table 3. Within-dataset MAE (MLP-small, 5-seed mean ± std). MAD = mean absolute deviation (naive mean-predictor baseline); best/MAD = ratio of best MAE to MAD (lower is better; <0.90 indicates meaningful signal above chance). **Bold** = best MAE per row, excluding [†]-flagged cells. [†]HEAR CIDRZ results are potentially contaminated (CIDRZ may appear in HEAR pretraining) and are excluded from leader claims.

Task	Unit	MAD	OPERA-CT	OPERA-CE	OPERA-GT	HEAR	M2D+RESP	best/MAD
CIDRZ Age	yr	10.35	10.52 ±0.08	10.51 ±0.09	10.49 ±0.07	10.29[†] ±0.04	**10.40** ±0.05	0.99
CIDRZ BMI	kg/m²	3.74	**3.60** ±0.01	**3.60** ±0.01	3.67 ±0.01	3.60[†] ±0.02	3.63 ±0.02	0.96
CIDRZ Abn	prob	0.325	0.327 ±0.001	0.325 ±0.001	**0.316** ±0.001	0.328[†] ±0.001	0.320 ±0.004	0.97
CIDRZ TB	prob	0.205	**0.189** ±0.001	0.191 ±0.000	0.190 ±0.000	0.188[†] ±0.001	0.192 ±0.001	0.92
Coswara Age	yr	11.31	10.25 ±0.02	10.44 ±0.01	10.16 ±0.04	**9.12** ±0.07	9.58 ±0.06	**0.81**
CoughVID Age	yr	10.29	9.79 ±0.01	9.88 ±0.02	9.62 ±0.03	**9.61** ±0.02	9.79 ±0.02	0.93

eral audio masked modelling with respiratory fine-tuning (M2D+RESP). Table 2 summarises the five encoders.

OPERA-CT and OPERA-CE (Zhang et al., 2024) are contrastive models on 136K respiratory clips, differing in architecture (Transformer vs. EfficientNet CNN) and dimension (768-d vs. 1280-d). OPERA-GT (Zhang et al., 2024) is a generative masked autoencoder on the same corpus with an 8.18 s positional grid and zero-padded inputs. HEAR (Baur et al., 2024) is a ViT-L masked autoencoder pretrained on 313M health audio clips, the largest corpus and the only one covering clinical audio beyond respiratory sounds. M2D+RESP (Niizumi et al., 2025) combines masked spectrogram prediction on AudioSet with respiratory fine-tuning, with features mean-pooled to a single 3840-d vector per clip. All encoders are kept frozen and embeddings are extracted once, reused across all heads and evaluation regimes.

2.3. Regression Heads and Training

We compare three heads applied to frozen embeddings. Linear ($d_{feat} \rightarrow 1$) is the standard linear probing baseline (Zhang et al., 2024; Baur et al., 2024). MLP-small uses a 256-unit bottleneck with ReLU and 0.3 dropout ($d_{feat} \rightarrow 256 \rightarrow 1$), making it embedding-size-agnostic and comparable across all FMs. MLP is full-width ($d_{feat} \rightarrow d_{feat} \rightarrow 1$), yielding ~15M hidden parameters for M2D+RESP, expected to overfit on small datasets and recover on larger ones. All heads use Adam (lr = 10^{-4}, L2 = 10^{-5}, batch 64), LR decay 0.97/epoch, MSE loss, and early stopping (patience 10) on validation MAE for up to 64 epochs. Results are mean ± std over 5 seeds.

2.4. Evaluation Regimes and Metrics

We report MAE in natural units (years, kg/m², probability) alongside the mean absolute deviation (MAD) of each label distribution as a naive mean-predictor baseline. MAE < MAD confirms learning above chance and best MAE/MAD quantifies signal strength, since a model collapsing to the population mean can appear effective with no

patient-level prediction. Lower MAE is better throughout.

Within-dataset regression evaluates all six targets on held-out test splits (Table 3). Regression head comparison covers 90 model × task × head combinations to characterise the dataset-size × head-capacity interaction (Table 4). Cross-dataset transfer trains MLP-small on one dataset and evaluates on all others without adaptation across six age transfer conditions (Table 5)

3. Results

3.1. Within-Dataset Regression

Table 3 reports within-dataset MAE for MLP-small alongside the mean-predictor baseline (MAD, computed directly from each label distribution). All models nominally exceed the baseline on every task, but the margins vary sharply. Coswara age is the only target with clear signal (best/MAD = 0.81, HEAR −0.46 yr over M2D+RESP), with CoughVID age weaker (best/MAD = 0.93). By contrast, all four CIDRZ targets sit at or near the chance floor (best/MAD between 0.92 and 0.99; age within 1% of the baseline), so the FM embeddings carry no usable patient-level signal for this clinical cohort under linear probing. The clinical-score targets additionally show near-zero inter-model spread (TB: ≤0.004, X-ray: ≤0.012 MAE), consistent with a shared representational ceiling rather than any model recovering true clinical variation.

3.2. Generative vs. Contrastive Pretraining

OPERA-GT is favored over OPERA-CT on age regression in all three datasets (CIDRZ 10.49 vs. 10.52 yr, Coswara 10.16 vs. 10.25 yr, CoughVID 9.62 vs. 9.79 yr; 3/3 directions). The CIDRZ gap (0.03 yr) lies within one seed std, whereas the Coswara and CoughVID gaps (0.09 and 0.17 yr) exceed seed variance, so the trend is driven by the two larger datasets. This is consistent with the breath-regression finding of Zhang et al. (2024), suggesting the advantage extends beyond breath audio.

Table 4. Full three-head comparison (MAE, 5-seed mean). Age/BMI in natural units; abnormality and TB in probability units $[0, 1]$. oCT/oCE/oGT = OPERA-CT/OPERA-CE/OPERA-GT; M2D = M2D+RESP. **Bold** = best head per model × task.

| Task | Linear | | | | | MLP-small | | | | | MLP | | | | |
	oCT	oCE	oGT	HeAR	M2D	oCT	oCE	oGT	HeAR	M2D	oCT	oCE	oGT	HeAR	M2D
CIDRZ Age	**10.47**	10.55	**10.44**	10.58	10.63	10.52	**10.51**	10.49	**10.29**	**10.40**	10.57	10.70	10.54	10.39	10.93
CIDRZ BMI	3.64	3.62	**3.67**	3.64	3.68	**3.60**	**3.60**	**3.67**	**3.60**	**3.63**	**3.60**	**3.60**	**3.67**	3.61	3.91
CIDRZ Abn	**0.326**	**0.324**	0.317	0.331	0.316	0.327	0.325	**0.316**	0.328	0.320	0.331	0.329	**0.316**	0.332	**0.315**
CIDRZ TB	**0.189**	**0.190**	0.191	0.191	0.194	**0.189**	0.191	**0.190**	**0.188**	**0.192**	0.191	0.193	0.191	0.189	0.199
Coswara Age	10.25	10.46	10.25	9.50	9.98	10.25	**10.44**	10.16	**9.12**	**9.58**	**10.24**	10.49	**10.14**	9.26	9.90
CoughVID Age	9.81	**9.88**	9.71	9.78	9.86	**9.79**	**9.88**	9.62	**9.61**	**9.79**	**9.79**	9.89	**9.53**	9.67	9.95

Table 5. Cross-dataset age generalisation (MLP-small, best model per row). Gap = cross − within MAE of the same model. **Bold** = negative gap (cross-dataset outperforms in-domain).

Train → Test	Model	Cross	Within	Gap
CoughVID → CIDRZ	OPERA-CE	**10.34**	10.51	−0.17
Coswara → CIDRZ	OPERA-CE	10.54	10.51	+0.03
Coswara → CoughVID	OPERA-CT	10.42	9.79	+0.63
CoughVID → Coswara	HEAR	10.05	9.12	+0.94
CIDRZ → CoughVID	HEAR	10.54	9.61	+0.94
CIDRZ → Coswara	HEAR	11.55	9.12	+2.43

Figure 2. CIDRZ age MAE vs. training set size (MLP-small, 3-seed mean ± std). HEAR and M2D+RESP plateau by $N = 100$; OPERA models stabilise at $N = 400$.

3.3. Regression Head Comparison

MLP-small wins in 23 of 30 model × task combinations (Table 4), outperforming linear probing by up to 0.38 yr (HEAR on Coswara). Full MLP overfits on CIDRZ ($N_{\text{train}} = 669$): M2D+RESP degrades $+0.53$ yr vs. MLP-small, driven by a ~22,000:1 parameter-to-sample ratio in the 3840-d hidden layer. On larger CoughVID ($N_{\text{train}} = 3050$), full MLP recovers and OPERA-GT achieves its best result (9.53 yr), revealing a dataset-size × head-capacity trade-off as a deployment guideline.

3.4. Cross-Dataset Generalisation

Table 5 reports cross-dataset age MAE for the best model per transfer direction. Transfer succeeds only when CIDRZ is the *target*: CoughVID → CIDRZ achieves a negative gap ($−0.17$ yr) and Coswara → CIDRZ is essentially flat ($+0.03$ yr), both led by OPERA-CE, showing that large-scale web-collected data can substitute for scarce clinical training data. The reverse fails: CIDRZ → Coswara degrades $+2.43$ yr ($+26.6\%$) and CIDRZ → CoughVID $+0.94$ yr, indicating that small clinical populations do not generalise to large crowdsourced ones. HEAR is best only in the three degrading directions (gaps $\geq +0.94$ yr), where it degrades least rather than transferring well, so we do not read this as a pretraining-driven advantage; the only non-degrading transfers are led by OPERA-CE.

3.5. Low-Data Regime

Figure 2 shows CIDRZ age MAE as a function of training set size (MLP-small, 3 seeds). M2D+RESP, whose pretraining corpus does not include CIDRZ, shows flat curves across all training sizes, indicating that large-scale pretraining can produce embeddings that require minimal labeled data for regression. HEAR behaves identically, reaching within 0.02 yr of its $N = 669$ performance at $N = 50$, consistent with this account; on CIDRZ, however, we cannot exclude a contribution from possible pretraining overlap. OPERA models, also CIDRZ-clean, show substantially higher variance at $N = 50$ (std up to ± 0.22 yr) and continue improving to $N = 400$, consistent with their smaller, respiratory-only pretraining corpus.

4. Conclusion

We presented the multi-model, multi-target cough regression benchmark evaluating five respiratory acoustic FMs across six targets, three head architectures, and six cross-dataset transfer conditions. Four findings carry practical implications. First, MLP-small is the preferred head for frozen FM embeddings, outperforming linear probing in 23

of 30 cases while avoiding overfitting of full-width MLPs on small clinical datasets, though signal strength varies substantially across tasks (best/MAD ranging from 0.81 to 0.99), indicating that head choice matters most when the task has learnable structure. Second, generative pretraining (OPERA-GT) is favored over contrastive pretraining (OPERA-CT) on age regression in all three datasets (3/3 directions), though the effect is driven by Coswara and CoughVID, as the CIDRZ margin lies within seed variance; this extends a pattern observed for breath audio and is consistent with the hypothesis that reconstruction-based objectives capture more physiological variation, though the mechanism warrants further study. Third, cross-dataset transfer is strongly asymmetric and large web-collected data generalises to small clinical populations with no performance loss while the reverse fails severely ($+2.43$ yr, $81\times$ larger gap), likely reflecting the narrow demographic range of clinical cohorts relative to global crowdsourced corpora. Fourth, M2D+RESP reaches near-full performance at only $N = 50$ labeled samples (with HEAR behaving similarly) while OPERA models require $N = 400$, indicating that pretraining-corpus diversity rather than architecture drives low-data regression; we anchor this on the CIDRZ-clean M2D+RESP, as HEAR's low-data CIDRZ result may benefit from pretraining overlap. **Limitations and future work.** This study uses 2 s clips. Cross-dataset transfer is reported for age only because age is the sole target shared across two or more datasets (BMI, X-ray abnormality, and TB probability appear only in CIDRZ), so transfer is undefined for these targets rather than omitted. All conclusions are likewise scoped to the frozen-embedding, shallow-probe regime: the benchmark characterises what FM embeddings encode, not the performance ceiling achievable with fine-tuning, which was not evaluated and may further reduce the transfer gap on small clinical datasets. Future work should extend to additional targets, fine-tuning regimes, and attention-pooling heads.

Impact Statement

This paper presents work whose goal is to advance the field of Machine Learning. There are many potential societal consequences of our work, none of which we feel must be specifically highlighted here.

References

Baur, S., Nabulsi, Z., Weng, W.-H., Garrison, J., Blankemeier, L., Fishman, S., Chen, C., Kakarmath, S., Maimbolwa, M., Sanjase, N., Shuma, B., Matias, Y., Corrado, G. S., Patel, S., Shetty, S., Prabhakara, S., Muyoyeta, M., and Ardila, D. HeAR – health acoustic representations. *arXiv preprint arXiv:2403.02522*, 2024.

Bhattacharya, D., Sharma, N. K., Dutta, D., Chetupalli, S. R., Mote, P., Ganapathy, S., Chandrakiran, C., Nori, S., Suhail, K. K., Gonuguntla, S., and Alagesan, M. Coswara: A respiratory sounds and symptoms dataset for remote screening of SARS-CoV-2 infection. *Scientific Data*, 10 (1):397, 2023.

Google Health AI. Google + CIDRZ health AI evaluation zambia. Kaggle dataset, 2024. Validation dataset for AI-enabled tuberculosis screening; CIDRZ tuberculosis cough cohort. https://www.kaggle.com/datasets/googlehealthai/google-health-ai.

Niizumi, D., Takeuchi, D., Yasuda, M., Nguyen, B. T., Ohishi, Y., and Harada, N. Towards pre-training an effective respiratory audio foundation model. In *Proc. Interspeech 2025*, 2025.

Orlandic, L., Teijeiro, T., and Atienza, D. The CoughVID crowdsourcing dataset, a corpus for the study of large-scale cough analysis algorithms. *Scientific Data*, 8(1): 156, 2021.

Rudraraju, G., Palreddy, S., Mamidgi, B., Sripada, N. R., Sai, Y. P., Vodnala, N. K., and Haranath, S. P. Cough sound analysis and objective correlation with spirometry and clinical diagnosis. *Informatics in Medicine Unlocked*, 19:100319, 2020.

Sharan, R. V., Abeyratne, U. R., Swarnkar, V. R., Claxton, S., Hukins, C., and Porter, P. Predicting spirometry readings using cough sound features and regression. *Physiological Measurement*, 39(9):095001, 2018.

World Health Organization. Global tuberculosis report 2021. Technical report, World Health Organization, Geneva, 2021.

Xu, W., He, G., Pan, C., Shen, D., Zhang, N., Jiang, P., Liu, F., and Chen, J. A forced cough sound based pulmonary function assessment method by using machine learning. *Frontiers in Public Health*, 10:1015876, 2022.

Zhang, Y., Xia, T., Han, J., Wu, Y., Rizos, G., Liu, Y., Mosuily, M., Chauhan, J., and Mascolo, C. Towards open respiratory acoustic foundation models: Pretraining and benchmarking. In *Advances in Neural Information Processing Systems Datasets and Benchmarks Track*, 2024.