Online Learning of Optimal Control Signals in Stochastic Linear Dynamical Systems

Abstract

Among the most canonical systems are linear time-invariant dynamics governed by differential equations and stochastic disturbances. An interesting problem in this class of systems is learning to minimize a quadratic cost function when system matrices are unknown. This work initiates theoretical analysis of implementable reinforcement learning policies for balancing exploration versus exploitation in such systems. We present an online policy that learns the optimal control actions fast by carefully randomizing the parameter estimates to explore. More precisely, we establish performance guarantees for the presented policy showing that the regret grows as the *square-root of time* multiplied by the *number of parameters*. Implementation of the policy for a flight control task shows its efficacy. Further, we prove tight results that ensure stability under inexact system matrices and fully specify unavoidable performance degradations caused by a non-optimal policy. To obtain the results, we conduct a novel analysis for matrix perturbation, bound comparative ratios of stochastic integrals, and introduce the new method of policy differentiation. These technical novelties are trusted to provide a useful cornerstone for continuous-time reinforcement learning.

1 Introduction

State-space models are widely-used for decision-making in dynamic environments. A popular model is the one that represents the continuous-time dynamics of the environment by linear stochastic differential equations. In this setting, the multidimensional state of the system is driven by the control action and the Brownian noise, according to an Ito stochastic differential equation. The range of applications is extensive, including chemistry, biology, finance, and engineering [23, 36, 34, 28].

In many applications, uncertainties about the true dynamics necessitate adaptive methods that learn optimal actions, probably relying on parameter estimation. Accordingly, reinforcement learning policies are extensively studied for decision-making under uncertainty in discrete time systems. The existing literature is fairly rich, including efficient algorithms that use optimism in the face of uncertainty and posterior sampling [1, 2, 33, 22]. Further, regret bounds are available in the presence of domain knowledge, partial observation, and multiple agents [10, 40, 41, 3, 27].

On the other hand, the existing literature for continuous-time systems is still immature. Early papers focus on estimating unknown parameters after an infinitely long interaction with the environment [31, 30, 18, 17, 16]. Recently, sub-optimal policies with linear regrets are proposed and consistency is shown for systems with full-rank input matrices [9]. Ensuing papers study offline algorithms for learning to control continuous-time systems from a batch of data. It is shown that computation of optimal policies can be performed using dynamic programming, temporal difference, entropy regularization, and system identification [15, 35, 38, 5].

However, *online* reinforcement learning policies that can learn optimal actions from a *single* state trajectory without imposing additional costs to the system are currently unavailable. The existing results rely on asymptotic estimation, require strong assumptions, and provide sub-optimal performance [16, 9]. A fundamental challenge compared to batch learning is that an online policy needs to *simultaneously* minimize the cost and estimate the unknown dynamics parameters. These two goals are contradictory and constitute the important exploration-exploitation dilemma. That is, estimation

Submitted to *A causal view on dynamical systems* workshop at NeurIPS 2022. Do not distribute.

accuracy is necessary for optimal decision-making, while sub-optimal actions are required for obtaining accurate estimates. We provide the first online reinforcement learning policy that balances the exploration-exploitation trade-off in a provably efficient manner and under minimal assumptions.

The main contributions of this paper can be summarized as follows. In Algorithm 1, we propose an efficient online reinforcement learning policy based on randomized estimates of the unknown system matrices. Then, we establish the rates for both the error in learning the dynamics matrices, as well as scaling of the regret that the algorithm incurs. Algorithm 1 is easy to implement, yet, it learns the optimal control actions fast so that its regret at time T is $\mathcal{O}\left(\sqrt{T}\log T\right)$ (Theorem 4). Furthermore, we study stability of linear systems for inaccurate system matrices and establish the minimal stabilizability margin (Theorem 2). Finally, a reciprocal expression is provided for regret that fully captures sub-optimalities due to uncertainty or inaccuracy in approximating the optimal control action (Theorem 3). The results provide both the average-case as well as worst-case analyses, the presented bounds are tight, and the assumptions are minimal.

To study online policies, one needs to address important challenges. First, sensitivity analysis of (complex) eigenvalues of matrices with perturbed entries is needed. Further, anti-concentration bounds for smallest singular values of partially-random matrices are required. Finally, we need to fully characterize the cumulative sub-optimalities in cost function in terms of model uncertainties. Thus, we develop multiple novel techniques for *matrix-perturbation* analysis, for comparative ratios of *stochastic integrals*, and for spectral properties of *random matrices*. We also introduce the method of *policy differentiation* to precisely capture the cost of sub-optimal control actions. En route, different tools from stochastic control, Ito calculus, and stochastic analysis are used, including Hamilton-Jacobi-Bellman equations, Ito Isometry, as well as martingale convergence theorems [39, 32, 4].

This paper is organized as follows. In Section 2, we discuss the problem and the preliminary materials. Then, in Section 3, we study system stability when the control action is designed based on dynamics matrices *other than* the true ones and establish stabilizability guarantees. Next, effects of sub-optimal actions and the regret they cause are examined in Section 4. Section 5 contains the randomized-estimates policy of Algorithm 1, as well as the theoretical analysis showing its efficiency. Because of space limitations, all proofs are delegated to the appendices, as outlined on page 13.

Notation. The smallest (largest) eigenvalue of A, in magnitude, is $\boldsymbol{\lambda}_{\min}(A)$ $(\boldsymbol{\lambda}_{\max}(A))$. For $v \in \mathbb{C}^d$, its norm is defined as $\|v\|^2 = \sum_{i=1}^d |v_i|^2$. Moreover, we write $\|A\|$ for the operator norm of matrices; $\|A\| = \sup_{\|v\|=1} \|Av\|$, and $A^\dagger$ for Moore-Penrose generalized inverse. The sigma-field generated by the stochastic process $\{Y_s\}_{0 \le s \le t}$ is denoted by $\sigma(Y_{0:t})$. A multivariate normal distribution with mean μ and covariance matrix Σ is shown by $\mathcal{N}(\mu, \Sigma)$. For $\lambda \in \mathbb{C}$, we use $\Re(\lambda), \Im(\lambda)$ to denote the real and imaginary parts of λ, respectively. The symbol $\vee$ (resp., $\wedge$) is used to show the maximum (resp., minimum). Finally, $\mathcal{O}(\cdot)$ refers to the order of magnitude.

2   Problem Statement

We study reinforcement learning policies for a multidimensional Ito stochastic differential equation with unknown drift matrices. That is, the state vector at time t is $X_t \in \mathbb{R}^{d_X}$, which follows

$$\mathrm{d}X_t = (A_\star X_t + B_\star U_t)\,\mathrm{d}t + C\mathrm{d}W_t. \tag{1}$$

Above, the vector $U_t \in \mathbb{R}^{d_U}$ is the control action at time t, and the disturbance $\{W_t\}_{t \ge 0}$ is a standard Brownian motion in a d_W dimensional space. Technically, by fixing the probability space $(\Omega, \mathbb{F}, \mathbb{P})$ which is completed by adding the null-sets of $\mathbb{P}$, let all stochastic objects belong to this probability space, and let $\mathbb{E}[\cdot]$ be expectation with respect to $\mathbb{P}$ (unless otherwise explicitly stated). The Brownian motion $\{W_t\}_{t \ge 0}$ starts from the origin and has independent normally distributed increments. That is, $W_0 = 0$, for all $0 \le t_1 \le t_2 \le t_3 \le t_4$, the vectors $W_{t_2} - W_{t_1}$ and $W_{t_4} - W_{t_3}$ are statistically independent, and for all non-negative reals $s < t$, it holds that $W_t - W_s \sim \mathcal{N}(0, (t-s)I_{d_W})$. Furthermore, $C \in \mathbb{R}^{d_X \times d_W}$ reflects the effect of W_t on the state evolution.

We aim to design computationally tractable and provably efficient reinforcement learning policies for the system in (1). The transition matrix $A_\star$, the input matrix $B_\star$, and the noise-coefficient matrix C,

all are unknown. The goal is to minimize the expected average cost

$$\mathcal{J}_{\boldsymbol{\pi}} = \limsup_{T \to \infty} \frac{1}{T} \, \mathbb{E} \left[\int_0^T c_{\boldsymbol{\pi}} \left(X_t, U_t \right) \mathrm{d}t \right],$$

where $c_{\boldsymbol{\pi}} \left(X_t, U_t \right) = X_t^\top Q X_t + U_t^\top R U_t$ is the cost at t. The value of $c_{\boldsymbol{\pi}} \left(X_t, U_t \right)$ is determined by the positive definite matrices Q, R of proper dimensions and by the policy $\boldsymbol{\pi}$, as defined below.

Technically, $\boldsymbol{\pi}$ is a non-anticipative closed-loop policy: At every time t, $\boldsymbol{\pi}$ determines U_t according to the information available at the time. More precisely, $\boldsymbol{\pi}$ maps the state observations (i.e., X_s for $s \in [0, t]$) and the previously taken actions (i.e., U_s for s in the open interval $[0, t[$) to the current control action U_t. This mapping can be stochastic or deterministic. Importantly, $\boldsymbol{\pi}$ faces the fundamental exploration-exploitation dilemma for minimizing the expected average cost, because the dynamics matrices $A_\star, B_\star$ are unknown and need to be learned based on the state and action observations. The details of this dilemma is discussed in Section 5. We assume that Q, R are known to the policy, the rationale being that the decision-makers are aware of the objective they aims to achieve, while their uncertainty about the true dynamics impedes them from deciding optimally.

The benchmark for assessing reinforcement learning policies is the optimal policy $\boldsymbol{\pi}^\star$ that designs U_t *having access* to $A_\star, B_\star$. To define $\boldsymbol{\pi}^\star$, let $\Phi_{A_\star, B_\star} \left(\cdot \right) : \mathbb{R}^{d_X \times d_X} \to \mathbb{R}^{d_X \times d_X}$ be

$$\Phi_{A_\star, B_\star} \left(M \right) = A_\star^\top M + M A_\star - M B_\star R^{-1} B_\star^\top M + Q.$$

The above mapping is vital for finding $\boldsymbol{\pi}^\star$. To see the intuition, first note that a control action U_t directly influences the current cost $c_{\boldsymbol{\pi}} \left(X_t, U_t \right)$, and indirectly affects the future cost values according to (1). So, future consequences of decisions need to be considered for minimizing $\mathcal{J}_{\boldsymbol{\pi}}$, and $\Phi_{A_\star, B_\star} \left(\cdot \right)$ is employed for this purpose [12, 39]. To proceed towards identifying $\boldsymbol{\pi}^\star$, let $M = \mathcal{K} \left(A_\star, B_\star \right)$ solve the equation $\Phi_{A_\star, B_\star} \left(M \right) = 0$. In order to investigate existence and uniqueness of $\mathcal{K} \left(A_\star, B_\star \right)$, we need the following definition and assumption.

Definition 1 (Notations $\overline{\boldsymbol{\lambda}} \left(\cdot \right), \mathcal{E} \left(\cdot \right)$) *Let $\overline{\boldsymbol{\lambda}} \left(D \right)$ be the largest real-part of the eigenvalues of an arbitrary square matrix D: $\overline{\boldsymbol{\lambda}} \left(D \right) = \max \left\{ \Re \left(\lambda \right) : \det \left(D - \lambda I \right) = 0 \right\}$. Further, for arbitrary matrices $A \in \mathbb{R}^{d_X \times d_X}$, $B \in \mathbb{R}^{d_X \times d_U}$, define $\mathcal{E} \left(A, B \right) = \| A - A_\star \| + \| B - B_\star \|$. So, $\mathcal{E} \left(A, B \right)$ measures the deviation of A, B from the true dynamics matrices $A_\star, B_\star$.*

Note that unlike $\boldsymbol{\lambda}_{\min} \left(\cdot \right), \boldsymbol{\lambda}_{\max} \left(\cdot \right)$ that consider only *magnitudes* of the eigenvalues, $\underline{\boldsymbol{\lambda}} \left(\cdot \right)$ reflects the signs of the eigenvalues as well, and so can be either positive, zero, or negative. However, they are related according to $\boldsymbol{\lambda}_{\max} \left(e^D \right) = e^{\overline{\boldsymbol{\lambda}}(D)}$.

We assume that the true dynamics matrices $A_\star, B_\star$ are stabilizable:

Assumption 1 (Stabilizability) *There exists some $L \in \mathbb{R}^{d_U \times d_X}$, such that $\overline{\boldsymbol{\lambda}} \left(A_\star + B_\star L \right) < 0$.*

Assumption 1 expresses that by applying $U_t = L X_t$, the system can operate without any explosion. To see that, solve the differential equation (1) under the feedback policy $U_t = L X_t$ to obtain

$$X_t = e^{(A_\star + B_\star L)t} X_0 + \int_0^t e^{(A_\star + B_\star L)(t-s)} C \mathrm{d}W_s. \tag{2}$$

So, because of $\overline{\boldsymbol{\lambda}} \left(A_\star + B_\star L \right) < 0$, X_t does not grow unbounded with t. Importantly, existence of a stabilizing matrix L is *necessary* for the problem to be well-defined. Otherwise, state explosion renders the average cost infinite for *all* decision-making policies [12, 39]. Now, recall that $\Phi_{A_\star, B_\star} \left(\mathcal{K} \left(A_\star, B_\star \right) \right) = 0$, let $\mathcal{L} \left(A_\star, B_\star \right) = -R^{-1} B_\star^\top \mathcal{K} \left(A_\star, B_\star \right)$ and define the feedback policy

$$\boldsymbol{\pi}^\star : \qquad U_t = \mathcal{L} \left(A_\star, B_\star \right) X_t, \quad \forall t \geq 0. \tag{3}$$

We show that Assumption 1 suffices for unique existence of $\mathcal{K} \left(A_\star, B_\star \right)$ and for optimality of $\boldsymbol{\pi}^\star$.

Theorem 1 (Optimal policy) *The matrix $\mathcal{K} \left(A_\star, B_\star \right)$ uniquely exists, and $\boldsymbol{\pi}^\star$ in (3) gives*

$$\mathcal{J}_{\boldsymbol{\pi}^\star} = \inf_{\boldsymbol{\pi}} \mathcal{J}_{\boldsymbol{\pi}} = \mathbf{tr} \left(\mathcal{K} \left(A_\star, B_\star \right) C C^\top \right), \qquad \overline{\boldsymbol{\lambda}} \left(A_\star + B_\star \mathcal{L} \left(A_\star, B_\star \right) \right) < 0.$$

126 To compute $\mathcal{K}(A_\star, B_\star)$, it suffices to solve the differential equation $\dot{M}_t = \Phi_{A_\star, B_\star}(M_t)$ starting from

127 a positive semidefinite M_0. Note that it is equivalent to the integration $M_t = M_0 + \int_0^t \Phi_{A_\star, B_\star}(M_s)\mathrm{d}s$.

128 In the proof of Theorem 1, we show that $\lim_{t \to \infty} M_t = \mathcal{K}(A_\star, B_\star)$.

129 Next, we focus on learning $A_\star, B_\star$ and the additional cost compared to the cost of $\boldsymbol{\pi}^\star$. To that end,
130 we formulate sub-optimalities of decision-making policies and the increase in cost due to lack of
131 knowledge about the optimal actions $U_t = \mathcal{L}(A_\star, B_\star) X_t$. For a general policy $\boldsymbol{\pi}$, the *regret* of $\boldsymbol{\pi}$ at
132 time T is denoted by $\mathcal{R}_{\boldsymbol{\pi}}(T)$ and is defined as the cumulative increase in the cost by T. That is, the
133 difference between the instantaneous costs of $\boldsymbol{\pi}$ and $\boldsymbol{\pi}^\star$ in (3) is integrated over $0 \le t \le T$:

$$\mathcal{R}_{\boldsymbol{\pi}}(T) = \int_0^T [c_{\boldsymbol{\pi}}(X_t, U_t) - c_{\boldsymbol{\pi}^\star}(X_t, U_t)]\mathrm{d}t. \tag{4}$$

134 Clearly, randomness of state and action leads to that of regret. So, regret analyses for reinforcement
135 learning policies include worst-case analyses that establish upper-bounds for $\mathcal{R}_{\boldsymbol{\pi}}(T)$, as well as
136 average-case analyses that provide bounds for $\mathbb{E}[\mathcal{R}_{\boldsymbol{\pi}}(T)]$. Further, for unknown $A_\star, B_\star$, we hope
137 that the increasing observations of state and action over time will be effectively leveraged so that
138 eventually, the policy makes near-optimal decisions. So, as t grows, we expect $c_{\boldsymbol{\pi}}(X_t, U_t) -$
139 $c_{\boldsymbol{\pi}^\star}(X_t, U_t)$ to shrink. Accordingly, $\mathcal{R}_{\boldsymbol{\pi}}(T)$ is desired to scale sub-linearly with T. In the rest of
140 this paper, we study $\mathcal{R}_{\boldsymbol{\pi}}(T), \mathbb{E}[\mathcal{R}_{\boldsymbol{\pi}}(T)]$, and their dependence on T and the problem parameters.

141 Another quantity of interest is the *accuracy* of estimating the unknown dynamics. So, letting A_t, B_t
142 be estimates of $A_\star, B_\star$ based on the state-action observations by time t; i.e., $\{X_s, U_s\}_{0 \le s \le t}$, we
143 study the decay rate of the estimation error $\mathcal{E}(A_t, B_t)$, as defined in Definition 1.

3 Stability Analysis for Perturbed Dynamics Matrices

145 In this section, we study the effects of uncertainties about the dynamical model on the system stability.
146 We specify the minimal information one needs to possess in order to ensure stabilization, and show
147 that a coarse-grained approximation of the truth is sufficient for that purpose. Results of this section
148 will be used later in the design of randomized-estimates policy in Section 5. Importantly, the stability
149 analysis in the sequel is general, captures effects of all involved quantities, and provides tight results
150 in the sense that the conditions of Theorem 2 are required for guaranteeing stabilization. In addition,
151 the results presented here are of independent interests, because stability is required for letting the
152 system operate for a reasonable time period.

153 To proceed, note that if hypothetically the optimal linear feedback in (3) is applied to the system
154 in (1), then stability is guaranteed. More precisely, applying $\mathcal{L}(A_\star, B_\star)$, the resulting closed-loop
155 transition matrix $D_\star = A_\star + B_\star \mathcal{L}(A_\star, B_\star)$ has all its eigenvalues on the open left half-plane of
156 the complex plane, as stated in Theorem 1. The issue is that the true dynamics matrices $A_\star, B_\star$ are
157 *unknown* and need to be learned. However, if some matrices A, B meet the conditions we shortly
158 discuss, one can stabilize the system by applying the linear feedback $U_t = \mathcal{L}(A, B) X_t$.

159 To proceed, let $D = A + B\mathcal{L}(A, B)$ be the closed-loop transition matrix of a system with dynamics
160 matrices A, B, under the feedback $U_t = \mathcal{L}(A, B) X_t$. Then, let $\rho > 0$ and $\zeta < \infty$ satisfy

$$\overline{\boldsymbol{\lambda}}(D) \le -\rho, \qquad \|\mathcal{K}(A, B)\| \le \zeta. \tag{5}$$

161 The quantities in (5) both are required for studying stability of the matrix $A_\star + B_\star \mathcal{L}(A, B)$, as
162 follows. Remember that we have a closed-loop stability result in Theorem 1: $\overline{\boldsymbol{\lambda}}(D) < 0$. So, the first
163 inequality in (5) quantifies the extent to which $\mathcal{L}(A, B)$ is able to stabilize the system of parameters
164 A, B. Intuitively speaking, $\overline{\boldsymbol{\lambda}}(D)$ is the smallest (i.e., most negative) upper-bound one can hope for
165 the eigenvalues of $A_\star + B_\star \mathcal{L}(A, B)$, since the optimal feedback $\mathcal{L}(A, B)$ is purposefully designed
166 for the certainly known matrices A, B. It can be shown that the positive quantity ρ enjoys a uniform
167 positive lower-bound as long as A, B live in some neighborhoods of $A_\star, B_\star$, as expressed in (31).
168 The second inequality in (5) is somewhat guaranteed by the first one. Technically, it is shown in (22)
169 in the proof of Theorem 1 that

$$\mathcal{K}(A, B) = \int_0^\infty e^{D^\top t} \left(Q + \mathcal{L}(A, B)^\top R \mathcal{L}(A, B) \right) e^{Dt}\mathrm{d}t. \tag{6}$$

170 Therefore, $\overline{\boldsymbol{\lambda}}(D) \leq \rho < 0$ implies that for some $\zeta < \infty$, we have $\|\mathcal{K}(A, B)\| \leq \zeta$. So, ζ is merely
171 used for simplifying the expressions.

172 Towards stability analysis, we need further information of $D = A + B\mathcal{L}(A, B)$ that the Jordan
173 form of this matrix provides. Suppose that eigenvalues of D are $\lambda_1, \cdots, \lambda_k$, and let the Jordan
174 decomposition be $D = P^{-1}\Lambda P$. So, $\Lambda = \mathrm{diag}(\Lambda_1, \cdots, \Lambda_k)$ is a block-diagonal matrix, and all
175 diagonal entries of Λ_i are λ_i, the immediate off-diagonal entries above the diagonal of Λ_i are 1,
176 and all other entries of Λ_i are 0 (as shown in (24)). Now, a consequential quantity for determining
177 the stability margin is the largest block-size $\boldsymbol{\mu}_D$ defined in Definition 2. It depends on the above-
178 mentioned blocks Λ_i, $i = 1, \cdots, k$, by letting $\boldsymbol{\mu}_i$ denote the dimension of the square matrix Λ_i, and
179 referring to the largest among $\boldsymbol{\mu}_1, \cdots, \boldsymbol{\mu}_k$ by $\boldsymbol{\mu}_D$.

180 **Definition 2 (Largest block-size $\boldsymbol{\mu}_D$)** *Letting P and $\Lambda_i \in \mathbb{C}^{\boldsymbol{\mu}_i \times \boldsymbol{\mu}_i}$ be as in the Jordan decomposi-*
181 *tion $D = P^{-1}\Lambda P$ explained above, define $\boldsymbol{\mu}_D = \max\limits_{1 \leq i \leq k} \boldsymbol{\mu}_i$.*

182 The quantity $\boldsymbol{\mu}_D$ is the largest size of the blocks $\Lambda_1, \cdots, \Lambda_k$ in the Jordan form and crucially
183 determines the *order* of stability margin, as established in the following theorem.

184 **Theorem 2 (Stability margin)** *Let $P, \boldsymbol{\mu}_D$ and ρ, ζ be as in Definition 2 and (5), respectively. Then,*
185 *we have $\overline{\boldsymbol{\lambda}}(A_\star + B_\star\mathcal{L}(A, B)) < -\delta$, as long as*

$$\mathcal{E}(A, B) < \left(1 \wedge \frac{1}{\|\mathcal{L}(A, B)\|}\right) \frac{(\rho - \delta) \wedge (\rho - \delta)^{\boldsymbol{\mu}_D}}{\boldsymbol{\mu}_D^{1/2}\|P\|\|P^{-1}\|}. \tag{7}$$

186 Note that in (7), we can upper-bound $\|\mathcal{L}(A, B)\|^{-1}$ with $\boldsymbol{\lambda}_{\min}(R)\zeta^{-1}\|B\|^{-1}$. Theorem 2 states
187 that if $\mathcal{E}(A, B)$ is sufficiently small to satisfy (7), then $A_\star + B_\star\mathcal{L}(A, B)$ is stable and all of its
188 eigenvalues in the complex plane lie on the left-hand-side of the vertical line $\Re = -\delta$. In addition, (7)
189 reflects effects of different factors, as follows. First, the reason for the decrease in the stability margin
190 on the right-hand-side of (7) as $\|\mathcal{L}(A, B)\|$ increases, is that the difference between the closed-loop
191 matrices is the product of $B_\star - B$ and $\mathcal{L}(A, B)$. Further, as can be seen in Definition 2, the quantity
192 $\boldsymbol{\mu}_D^{1/2}\|P\|\|P^{-1}\| \geq 1$ quantifies non-diagonality of D, and it is 1 for fully diagonal matrices D
193 where P is the identity matrix and $\boldsymbol{\mu}_D = 1$. Therefore, more non-diagonal closed-loop matrices D
194 lead to smaller stability margins and make stabilization harder to learn.

195 The dependence on ρ corroborates the intuition that systems whose optimal closed-loop matrices has
196 eigenvalues of larger real-parts (i.e., smaller ρ), are harder to stabilize. Moreover, the expression
197 $(\rho - \delta) \wedge (\rho - \delta)^{\boldsymbol{\mu}_D}$ indicates that $\boldsymbol{\mu}_D$ is very important for determining the rates of bounding the
198 eigenvalues of $A_\star + B_\star\mathcal{L}(A, B)$. The rates for $\rho - \delta < 1$ and $\rho - \delta > 1$ are different, because of
199 a similar phenomena in sensitivity of eigenvalues of matrices to perturbations in their entries. This
200 result is of independent interest as it is a generalization of Bauer-Fike Theorem [6] to asymmetric
201 matrices we establish in the proof of Theorem 2. Intuitively, we prove that larger blocks in the Jordan
202 decomposition can lead to drastic fluctuations in eigenvalues.

203 To close this section, we provide uniform lower and upper bounds for $\rho > 0$ and $\zeta < \infty$, respectively.
204 For that purpose, similar to Definition 2, define the largest block size $\boldsymbol{\mu}_\star = \boldsymbol{\mu}_{D_\star}$ based on the
205 Jordan decomposition $D_\star = A_\star + B_\star\mathcal{L}(A_\star, B_\star) = P_\star^{-1}\Lambda_\star P_\star$. Then, we show in the proof of
206 Theorem 2 that $\mathcal{E}(A, B) \leq \epsilon_0$ is sufficient for satbilization, $\rho \geq \boldsymbol{\lambda}_{\min}(Q)4^{-1}\|\mathcal{K}(A_\star, B_\star)\|^{-1}$, and
207 $\zeta \leq 2\|\mathcal{K}(A_\star, B_\star)\|$, where

$$(1 \vee \|\mathcal{L}(A_\star, B_\star)\|)\,\epsilon_0 = \frac{\left(-\overline{\boldsymbol{\lambda}}(D_\star)\right) \wedge \left(-\overline{\boldsymbol{\lambda}}(D_\star)\right)^{\boldsymbol{\mu}_\star}}{\boldsymbol{\mu}_\star^{1/2}\|P_\star^{-1}\|\|P_\star\|} \wedge \left[4\int_0^\infty \|e^{D_\star t}\|^2 \mathrm{d}t\right]^{-1}. \tag{8}$$

4 Reciprocal Regret Bounds and Policy Differentiation

209 In this section, we investigate sub-optimalities and provide a tight expression for the regret that
210 non-optimal control actions cause. Such investigation is vital since reinforcement learning policies
211 need to learn the unknown dynamics $A_\star, B_\star$ and so their non-optimal actions are unavoidable.

To proceed, let U_t be the control action of the policy π at time t. In the following theorem, we quantify $\mathcal{R}_{\pi}(T)$ in terms of deviations $U_t - \mathcal{L}(A_\star, B_\star) X_t$, and introduce α_T that fully assesses the regret of π. In fact, α_T unifies average-case and worst-case analyses by capturing both $\mathbb{E}[\mathcal{R}_{\pi}(T)]$ and $\mathcal{R}_{\pi}(T)$. Further, Theorem 3 provides scalings with different problem parameters. For example, it shows that the difference between $\mathcal{R}_{\pi}(T)$ and $\mathbb{E}[\mathcal{R}_{\pi}(T)]$ scales linearly with the dimension of the Brownian motion. After the statement of Theorem 3, we discuss its intuitions and implications.

To establish Theorem 3, we utilize the theory of continuous-time martingales and (in Lemma 2) develop novel results on comparative ratios of stochastic integrals. More importantly, we construct the new framework of *policy differentiation* for finding tight and accurate regret bounds. Broadly speaking, policy differentiation precisely evaluates the regret in terms of infinitesimal sub-optimalities. Then, we integrate these infinitesimal deviations and obtain α_T, in which the integrand $\left\| R^{1/2}\left(L_t - \mathcal{L}(A_\star, B_\star)\right) X_t \right\|^2$ plays a role similar to the *derivative* of the regret. This framework can be used for an analogous regret analysis in other continuous-time reinforcement problems.

Theorem 3 (Regret analysis) *Suppose that L_t is a bounded piecewise continuous function of t, and π is the policy $U_t = L_t X_t$. Then, we have $\mathbb{E}[\mathcal{R}_{\pi}(T)] = \mathbb{E}[\alpha_T]$, and*

$$\mathcal{R}_{\pi}(T) \quad = \quad \alpha_T + \mathcal{O}\left(\omega_{\mathcal{R}}\, \alpha_T^{1/2} \log \alpha_T\right),$$

where $D_\star = A_\star + B_\star \mathcal{L}(A_\star, B_\star)$, $E_t = e^{D_\star^\top t} \mathcal{K}(A_\star, B_\star) e^{D_\star t}$, $\omega_{\mathcal{R}} = \dfrac{\|C\|\|\mathcal{K}(A_\star, B_\star)\|^{3/2} d_W}{\lambda_{\min}(Q)^{1/2} \lambda_{\min}(R)^{1/2}}$, and

$$\alpha_T = \int\limits_0^T \left\| R^{1/2}\left(L_t - \mathcal{L}(A_\star, B_\star)\right) X_t \right\|^2 \mathrm{d}t - 2 \int\limits_0^T \left(X_t^\top E_{T-t} B_\star \left(L_t - \mathcal{L}(A_\star, B_\star)\right) X_t \right) \mathrm{d}t.$$

The boundedness and piecewise continuity conditions are somewhat natural to ensure that the state process in (2) is well-defined. Because the optimal policy is a time-invariant feedback, violating these conditions does not lead to smaller regret. Furthermore, since by Theorem 1 we have $\overline{\lambda}(D_\star) < 0$, the matrix E_t exponentially decays as t grows. Therefore, the second integral in the definition of α_T is dominated by the first one. So, Theorem 3 shows that the sub-optimality π incurs at time t scales as *square* of the deviation $L_t - \mathcal{L}(A_\star, B_\star)$. Finally, the constant $\omega_{\mathcal{R}}$ reflects effects of different problem parameters and indicates that $\mathcal{R}_{\pi}(T) - \alpha_T$ scales linearly with d_W.

The results of Theorem 3 are interesting along different directions. First, the exact equality $\mathbb{E}[\mathcal{R}_{\pi}(T)] = \mathbb{E}[\alpha_T]$ can be used for establishing regret *lower-bounds* by studying the fastest rates $\pi^\star$ is learned and the deviation $L_t - \mathcal{L}(A_\star, B_\star)$ shrinks. Further, since $\alpha_T^{1/2} \log \alpha_T = \mathcal{O}(\alpha_T)$, not only the average-case criteria $\mathbb{E}[\mathcal{R}_{\pi}(T)]$, but also the worst-case regret $\mathcal{R}_{\pi}(T)$ are captured by α_T. In other words, Theorem 3 indicates that the fluctuations of $\mathcal{R}_{\pi}(T)$ around its expectation $\mathbb{E}[\mathcal{R}_{\pi}(T)]$ are in magnitude smaller than the expected value itself. Thus, studying α_T is sufficient and necessary for regret analysis, and it provides a general reciprocal relationship for *all* policies.

Moreover, when $A_\star, B_\star$ are unknown, a reinforcement learning policy becomes *progressively* more capable of narrowing down the sub-optimality gap as time goes by. More precisely, as soon as having sufficiently long trajectories to learn $A_\star, B_\star$ to satisfy $\left\| R^{1/2}\left(L_t - \mathcal{L}(A_\star, B_\star)\right) X_t \right\| < 1$, the regret grows much slower since α_T integrates the *squares* of these deviations. For example, if the estimation accuracy satisfies the ideal square-root rate $\mathcal{E}(A_t, B_t) = \mathcal{O}\left(t^{1/2}\right)$, then the regret scales logarithmic with time; $\alpha_T = \mathcal{O}(\log T)$. However, due to the trade-off between the exploration and exploitation, this is not the case (see Proposition 1). Technically, to obtain the above error rate, U_t needs to persistently deviate from $\mathcal{L}(A_\star, B_\star) X_t$, which causes a *linearly* growing regret.

Theorem 3 provides both a general result for analyzing online policies, as well as a useful insight on how to design them. We utilize this insight to design Algorithm 1 and to establish Theorem 4 in the next section. Indeed, we randomize the parameter estimates so that U_t appropriately deviates from $\pi^\star$, leading to $\mathcal{E}(A_t, B_t) = \mathcal{O}\left(t^{-1/4}\right)$. So, we obtain the efficient regret bound $\alpha_T = \widetilde{\mathcal{O}}\left(T^{1/2}\right)$.

5   Randomized-Estimates Policy

In this section, we discuss fast and tractable algorithms with efficient performances for cost minimization subject to uncertainties about the dynamics matrices $A_\star, B_\star$. First, we explain the fundamental

exploration-exploitation dilemma. Then, we investigate a procedure for estimating the unknown dynamics using the data of state-action trajectory. Based on that, an online reinforcement learning policy that employs randomizations of the parameter estimates for balancing exploration versus exploitation is presented in Algorithm 1. Next, a regret bound is established in Theorem 4 indicating that Algorithm 1 efficiently minimizes the cost function so that the regret scales as the square-root of the time. We also specify the rates of identifying the dynamics matrices.

First, by Theorem 3, the policy needs to ensure that $U_t \approx \mathcal{L}(A_\star, B_\star) X_t$ to incur a small regret. Further, since $A_\star, B_\star$ are unknown, the policy needs to estimate them based on $\{X_s, U_s\}_{0 \le s \le t}$ by time t. However, if $U_s \approx \mathcal{L}(A_\star, B_\star) X_s$, then some of the coordinates of the data point X_s, U_s become (almost) uninformative as they are (approximately) linear transformations of the rest of them. This defeats the purpose and renders accurate estimation of $A_\star, B_\star$ infeasible. Note that accurate approximations of $A_\star, B_\star$ are needed for taking near-optimal actions. This, known as the exploration-exploitation dilemma, is the main obstacle in online reinforcement learning and shows that a low-regret policy needs to *diversify* the actions $\{U_s\}_{0 \le s \le t}$ by *deviating* from $\{\mathcal{L}(A_\star, B_\star) X_s\}_{0 \le s \le t}$.

Now, we discuss the learning procedure in Algorithm 1, based on extensions of the least-squares estimates. Suppose that instead of the full data X_s, U_s for real values of $s \ge 0$, one has access to samples at a discrete set of time points; $X_{k\epsilon}, U_{k\epsilon}$ for $k = 0, 1, \cdots$. Then, (1) for a small ϵ gives the approximate data generation mechanism $X_{(k+1)\epsilon} - X_{k\epsilon} = (A_\star X_{k\epsilon} + B_\star U_{k\epsilon}) \epsilon + C(W_{(k+1)\epsilon} - W_{k\epsilon})$. So, a natural approach is to form a linear regression problem and estimate $A_\star, B_\star$ by a minimizer of $\sum_k \left\| X_{(k+1)\epsilon} - X_{k\epsilon} - (AX_{k\epsilon} + BU_{k\epsilon}) \epsilon \right\|^2$ over A, B. Denoting $Y_s = \left[X_s^\top, U_s^\top \right]^\top$, this gives the estimates $[A, B] = \sum_k \left(X_{(k+1)\epsilon} - X_{k\epsilon} \right) Y_{k\epsilon}^\top \epsilon \left(\sum_k Y_{k\epsilon} Y_{k\epsilon}^\top \epsilon^2 \right)^\dagger$. Therefore, letting $\epsilon \to 0$, the summations become integrations and we get the continuous-time estimate based on the full trajectory $\{X_s, U_s\}_{0 \le s \le t}$. The result is shown in (9) below and will be used by Algorithm 1.

To ensure that the system evolves stable, Algorithm 1 projects the parameter estimates on the stabilization oracle $\mathcal{S}_0$ in lights of Theorem 2. In the sequel, we explain how one can learn $\mathcal{S}_0$ fast.

Definition 3 *For a fixed $\delta_0 > 0$, let $\mathcal{S}_0$ be a set containing matrices A, B that satisfy (7) for $\delta = \delta_0$.*

Intuitively, the system is stabilized by having access to $\mathcal{S}_0$, despite uncertainties about the true dynamics matrices $A_\star, B_\star$. Note that the condition in (7) is verifiable since ρ, ζ depend on the *known* parameter estimates A, B. Availability of a stabilization set is a common assumption in the literature of online reinforcement learning policies for linear systems [31, 30, 8, 2, 33, 22, 40, 41]. For example, if an initial stabilizing feedback L_0 is available, we can devote a short time period to explore by applying sub-optimal control actions and use the resulting observations to learn $\mathcal{S}_0$. In systems operating prior to running Algorithm 1 or in open-loop-stable systems, this condition automatically holds (in the latter case, $L_0 = 0$ is an initial stabilizer). Similarly, in systems with a *reset* option that can immediately steer the system-state to small values, $\mathcal{S}_0$ can be learned fast [9, 5, 19].

In Section 3 we saw that an ϵ_0 neighborhood of $A_\star, B_\star$ is sufficient for bounding ρ, ζ, for ϵ_0 in (8). That is, a coarse-grained approximation of $A_\star, B_\star$ suffices for $\mathcal{S}_0$. So, if $\mathcal{S}_0$ is not available in advance, it can be learned from the state-input trajectories of short lengths [16, 9, 20, 21, 26, 13, 19, 24]. In addition, in Theorem 4 we show that the algorithm learns $A_\star, B_\star$ with the rate $t^{-1/4}$. So, the projection on $\mathcal{S}_0$ will be automatically performed after the time $t = \widetilde{\mathcal{O}}\left(\epsilon_0^{-4} \right)$.

The algorithm proceeds as follows. For some fixed $\gamma > 1$, the sequence $\{\gamma^n\}_{n=0}^\infty$ contains the time instants at which the algorithm updates the parameter estimates. In fact, Algorithm 1 applies control actions $U_t = \mathcal{L}(A_n, B_n) X_t$ during the time period $\gamma^n \le t < \gamma^{n+1}$, where A_n, B_n are estimates of $A_\star, B_\star$, based on the trajectory up to time γ^n. Further, to ensure that the policy commits sufficiently to explore the environment, a random matrix Θ_n is added to the parameter estimates at time $t = \gamma^n$. Then, Algorithm 1 projects the resulting $d_X \times (d_X + d_U)$ matrix onto $\mathcal{S}_0$. Formally, letting $\Pi_{\mathcal{S}_0}(\cdot)$ denote projection on $\mathcal{S}_0$, define

$$[A_n, B_n] = \Pi_{\mathcal{S}_0} \left(\left[\int_0^{\gamma^n} Y_s \mathrm{d}X_s^\top \right]^\top \left[\int_0^{\gamma^n} Y_s Y_s^\top \mathrm{d}s \right]^\dagger + \Theta_n \right), \tag{9}$$

where $Y_s = \left[X_s^\top, U_s^\top \right]^\top$ and the $d_X \times (d_X + d_U)$ matrices $\{\Theta_n\}_{n=0}^\infty$ are independent of everything else and of each others. Further, the random matrix Θ_n that is used at time $t = \gamma^n$ has independent

Gaussian entries of mean zero and standard deviation $\sigma_n = \sigma_0 \left(\gamma^{-n} n\right)^{1/4} = \sigma_0 \left(t^{-1} \log_\gamma t\right)^{1/4}$, for some fixed σ_0. This decay rate of σ_n is delicately adjusted for two purposes. On one hand, Θ_n is sufficiently large for randomizing the parameter estimates to ensure that effective exploration occurs and the current data is diverse enough so that we obtain accurate estimates in the future. On the other hand, Θ_n is sufficiently small to let the current estimates remain accurate and prevent significant deviations. Otherwise, large values of Θ_n deteriorate the current efficient exploitation.

Algorithm 1 : Randomized Estimates Policy

Select $\gamma > 1$ and $A_0, B_0 \in \mathcal{S}_0$ arbitrarily
For $0 \le t < 1 = \gamma^0$, apply $U_t = \mathcal{L}\left(A_0, B_0\right) X_t$
for $n = 0, 1, 2, \cdots$ **do**
    Update parameter estimates A_n, B_n by (9)
    while $\gamma^n \le t < \gamma^{n+1}$ **do**
        Take action $U_t = \mathcal{L}\left(A_n, B_n\right) X_t$
    end while
end for

The memory that Algorithm 1 occupies is remarkably small since in (9) it can update the involved matrices by continuously integrating the stream of the observations $Y_s, \mathrm{d}X_s$. Furthermore, calculations can be done fast, making update of the parameter estimates at time γ^n immediately effective. The rationale for freezing the parameter estimates for exponentially growing time intervals $\gamma^n \le t < \gamma^{n+1}$ is that Algorithm 1 can defer the learning step until collecting enough observations Y_s so that a new update of parameter estimates is effectively more accurate than the previous one. The following result provides performance guarantees for the randomized estimates policy.

Theorem 4 (Analysis of Algorithm 1) *Let the policy π and the estimates A_n, B_n be those in Algorithm 1. Assume that n satisfies $\gamma^n \le T < \gamma^{n+1}$. Then, using Definition 1 and (4), we have*

$$\mathcal{E}\left(A_n, B_n\right)^2 = \mathcal{O}\left(\boldsymbol{\omega}_{\mathcal{E}} T^{-1/2} \log T\right), \qquad \mathcal{R}_{\boldsymbol{\pi}}\left(T\right) = \mathcal{O}\left(\boldsymbol{\omega}_{\boldsymbol{\pi}} T^{1/2} \log T\right),$$

where

$$\boldsymbol{\omega}_{\mathcal{E}} = \left(d_X + d_U\right)\left(\frac{d_X}{\log \gamma} + \frac{d_W \|C\|^2}{\boldsymbol{\lambda}_{\min}\left(CC^\top\right)}\right), \qquad \boldsymbol{\omega}_{\boldsymbol{\pi}} = \frac{(\gamma - 1)\|C\|^2 \|\mathcal{K}\left(A_\star, B_\star\right)\|^6 \|R\|}{\boldsymbol{\lambda}_{\min}\left(Q\right)^2 \boldsymbol{\lambda}_{\min}\left(R\right)^4} \; \boldsymbol{\omega}_{\mathcal{E}}.$$

Theorem 4 indicates efficiency of Algorithm 1: At time T, the sub-optimality gap is as small as $\mathcal{O}\left(\boldsymbol{\omega}_{\boldsymbol{\pi}} T^{-1/2} \log T\right)$. It also provides $\boldsymbol{\omega}_{\boldsymbol{\pi}}, \boldsymbol{\omega}_{\mathcal{E}}$ that reflect the dependence of estimation error and regret on different parameters in the problem. So, the regret scales linearly with the number of unknown parameters in $A_\star, B_\star$, while the estimation error dwindles linearly with the dimension.

To establish Theorem 4, we study the learning step in (9) and determine the rates Algorithm 1 estimates $A_\star, B_\star$. For that purpose, we prove concentration bounds for the empirical covariance matrices of the state vectors and also show anti-concentration of the Gram matrix $\int_0^t Y_s Y_s^\top \mathrm{d}s$ of the signal $Y_s = \left[X_s^\top, U_s^\top\right]^\top$, as t grows. Then, we establish bounds on the comparative ratios of stochastic integrals and use that for controlling the estimation error. Furthermore, we show that the optimal feedback matrices have a Lipschitz property with respect to the dynamics matrices and leverage that for finding the deviation rates from the optimal feedback. Finally, we utilize policy differentiation and Theorem 3 for getting the regret bounds in Theorem 4.

Note that for obtaining descending estimation errors and sub-linear regret bounds we need $\boldsymbol{\lambda}_{\min}\left(CC^\top\right) > 0$. This is a standard requirement in estimation and control of stochastic linear systems and expresses that all coordinates of the state vectors are randomized by the Brownian motion $\{W_t\}_{t \ge 0}$ in a relatively short time [29, 37, 9]. So, all state variables have significant roles in the dynamics. From a modeling point of view, $\boldsymbol{\lambda}_{\min}\left(CC^\top\right) > 0$ indicates that the stochastic differential equation in (1) is irreducible in the sense that a smaller subset of state variables is *insufficient* for capturing the stochastic dynamical behavior of the environment.

To close this section observe that the estimation error of Algorithm 1 shrinks as $T^{-1/4}$. So, it does not decay with the ideal square-root rate because the main priority of Algorithm 1 is to minimize the

Figure 1: **Left:** Normalized estimation error $T^{1/2}\mathcal{E}\left(A_n, B_n\right)^2$ is plotted vs T, such that $\gamma^n \leq T < \gamma^{n+1}$, for some integer n. Multiple replicates of the normalized estimation error are reported in the graph, which clearly remain bounded as time grows. Therefore, the graph depicts Theorem 4 about the rates Algorithm 1 learns the dynamics matrices.

Right: The graph presents curves of the normalized regret $T^{-1/2}\mathcal{R}_{\boldsymbol{\pi}}\left(T\right)$ versus time T, while $\boldsymbol{\pi}$ is the policy in Algorithm 1. Multiple replicates of the system are simulated, all corroborating Theorem 4 that the normalized regret remains (almost) bounded.

regret by exploring minimally. However, if the randomization matrices Θ_n are persistent and their standard deviations do not diminish as n grows, then we obtain the square-root consistency. This is formalized in Proposition 1. Of course, the compromise is that $\mathcal{R}_{\boldsymbol{\pi}}\left(T\right)$ grows *linearly* with T if Θ_n does not dwindle as n grows.

Numerical Illustrations. Here, we provide numerical analyses of Algorithm 1. The details of the experiments are provided in Appendix A. The normalized rates of the estimation error are plotted in Figure 1, versus the continuous time T. To illustrate the rates in Theorem 4, the figure contains multiple trajectories of $T^{1/2}\mathcal{E}\left(A_n, B_n\right)^2$, while n, T satisfy $\gamma^n \leq T < \gamma^{n+1}$. The normalized estimation errors in Figure 1 are (almost) bounded as time grows, corroborating Theorem 4. Figure 1 depicts normalized regret versus time. That is, the horizontal axis is T, while the vertical one presents the values of $T^{-1/2}\mathcal{R}_{\boldsymbol{\pi}}\left(T\right)$, where $\boldsymbol{\pi}$ is the online reinforcement learning policy in Algorithm 1. Again, it is clear that Theorem 4 holds, as the normalized regret remains bounded as time grows.

6   Concluding Remarks

We studied online reinforcement learning policies for unknown continuous-time stochastic linear systems and presented algorithms that learn to minimize quadratic costs. Three important problems are fully investigated, followed by the intuitions and implications of the presented analyses.

First, we studied stabilization of stochastic linear systems based on incorrect dynamics matrices and proved Theorem 2 that specifies the coarse-grained approximations for guaranteeing stability. Then, proposing the novel approach of policy differentiation, we established a reciprocal result in Theorem 3 for the regret that policies cause by taking sub-optimal actions. More importantly, we presented the online reinforcement learning Algorithm 1 and established its performance guarantees. Indeed, Theorem 4 expresses that the estimation rate of Algorithm 1 is $dT^{-1/4}\log^{1/2} T$ and it enjoys the efficient regret bound $\mathcal{O}\left(d^2 T^{1/2}\log T\right)$, where T is the time and d is the dimension.

As an initiating paper on design and analysis of online reinforcement learning policies for continuous-time stochastic systems, this study introduces interesting directions for future work. That includes establishing anytime regret bounds that hold uniformly over time, investigating high-dimensional systems with structured dynamics such as low-rank or sparse matrices, designing efficient policies under imperfect state-observations, and extending the analysis to non-linear systems.

References

[1] Yasin Abbasi-Yadkori and Csaba Szepesvári. Regret bounds for the adaptive control of linear quadratic systems. In *Proceedings of the 24th Annual Conference on Learning Theory*, pages 1–26. JMLR Workshop and Conference Proceedings, 2011.

[2] Marc Abeille and Alessandro Lazaric. Improved regret bounds for thompson sampling in linear quadratic control problems. In *International Conference on Machine Learning*, pages 1–9. PMLR, 2018.

[3] Seyed Mohammad Asghari, Yi Ouyang, and Ashutosh Nayyar. Regret bounds for decentralized learning in cooperative multi-agent dynamical systems. In *Conference on Uncertainty in Artificial Intelligence*, pages 121–130. PMLR, 2020.

[4] Paolo Baldi. *Stochastic Calculus: An Introduction Through Theory and Exercises*. Springer, 2017.

[5] Matteo Basei, Xin Guo, Anran Hu, and Yufei Zhang. Logarithmic regret for episodic continuous-time linear-quadratic reinforcement learning over a finite-time horizon. *Available at SSRN 3848428*, 2021.

[6] Friedrich L Bauer and Charles T Fike. Norms and exclusion theorems. *Numerische Mathematik*, 2(1):137–141, 1960.

[7] John T Bosworth. *Linearized aerodynamic and control law models of the X-29A airplane and comparison with flight data*, volume 4356. National Aeronautics and Space Administration, Office of Management . . . , 1992.

[8] PE Caines. Continuous time stochastic adaptive control: non-explosion, ε-consistency and stability. *Systems & control letters*, 19(3):169–176, 1992.

[9] Peter E Caines and David Levanony. Stochastic ε-optimal linear quadratic adaptation: An alternating controls policy. *SIAM Journal on Control and Optimization*, 57(2):1094–1126, 2019.

[10] Asaf Cassel, Alon Cohen, and Tomer Koren. Logarithmic regret for learning linear quadratic regulators efficiently. In *International Conference on Machine Learning*, pages 1328–1337. PMLR, 2020.

[11] Siew Chan, GC Goodwin, and Kwai Sin. Convergence properties of the riccati difference equation in optimal filtering of nonstabilizable systems. *IEEE Transactions on Automatic Control*, 29(2):110–118, 1984.

[12] Goong Chen, Guanrong Chen, and Shih-Hsun Hsu. *Linear stochastic control systems*, volume 3. CRC press, 1995.

[13] Xinyi Chen and Elad Hazan. Black-box control for linear dynamical systems. In *Conference on Learning Theory*, pages 1114–1143. PMLR, 2021.

[14] C De Souza, M Gevers, and G Goodwin. Riccati equations in optimal filtering of nonstabilizable systems having singular state transition matrices. *IEEE Transactions on Automatic control*, 31(9):831–838, 1986.

[15] Kenji Doya. Reinforcement learning in continuous time and space. *Neural computation*, 12(1):219–245, 2000.

[16] Tyrone E Duncan, Lei Guo, and Bozenna Pasik-Duncan. Adaptive continuous-time linear quadratic gaussian control. *IEEE Transactions on automatic control*, 44(9):1653–1662, 1999.

[17] Tyrone E Duncan, Petr Mandl, and Bożenna Pasik-Duncan. On least squares estimation in continuous time linear stochastic systems. *Kybernetika*, 28(3):169–180, 1992.

[18] Tyrone E Duncan and Bozenna Pasik-Duncan. Adaptive control of continuous-time linear stochastic systems. *Mathematics of Control, signals and systems*, 3(1):45–60, 1990.

[19] Mohamad Kazem Shirani Faradonbeh and Mohamad Sadegh Shirani Faradonbeh. Bayesian algorithms learn to stabilize unknown continuous-time systems. *IFAC-PapersOnLine*, 55(12):377–382, 2022.

[20] Mohamad Kazem Shirani Faradonbeh, Ambuj Tewari, and George Michailidis. Finite-time adaptive stabilization of linear systems. *IEEE Transactions on Automatic Control*, 64(8):3498–3505, 2018.

[21] Mohamad Kazem Shirani Faradonbeh, Ambuj Tewari, and George Michailidis. Randomized algorithms for data-driven stabilization of stochastic linear systems. In *2019 IEEE Data Science Workshop (DSW)*, pages 170–174. IEEE, 2019.

[22] Mohamad Kazem Shirani Faradonbeh, Ambuj Tewari, and George Michailidis. On adaptive linear–quadratic regulators. *Automatica*, 117:108982, 2020.

[23] Daniel T Gillespie. Stochastic simulation of chemical kinetics. *Annu. Rev. Phys. Chem.*, 58:35–55, 2007.

[24] Dennis Gramlich and Christian Ebenbauer. Fast identification and stabilization of unknown linear systems. *arXiv preprint arXiv:2208.10392*, 2022.

[25] Panqanamala Ramana Kumar and Pravin Varaiya. *Stochastic systems: Estimation, identification, and adaptive control*. SIAM, 2015.

[26] Sahin Lale, Kamyar Azizzadenesheli, Babak Hassibi, and Anima Anandkumar. Explore more and improve regret in linear quadratic regulators. *arXiv preprint arXiv:2007.12291*, 2020.

[27] Sahin Lale, Kamyar Azizzadenesheli, Babak Hassibi, and Anima Anandkumar. Logarithmic regret bound in partially observable linear dynamical systems. *arXiv preprint arXiv:2003.11227*, 2020.

[28] Neil D Lawrence, Mark Girolami, Magnus Rattray, and Guido Sanguinetti. *Learning and inference in computational systems biology*. MIT press, 2010.

[29] David Levanony and Peter E Caines. On persistent excitation for linear systems with stochastic coefficients. *SIAM journal on control and optimization*, 40(3):882–897, 2001.

[30] Petr Mandl. Consistency of estimators in controlled systems. In *Stochastic Differential Systems*, pages 227–234. Springer, 1989.

[31] Petr Mandl, Tyrone E Duncan, and Bożenna Pasik-Duncan. On the consistency of a least squares identification procedure. *Kybernetika*, 24(5):340–346, 1988.

[32] Bernt Oksendal. *Stochastic differential equations: an introduction with applications*. Springer Science & Business Media, 2013.

[33] Yi Ouyang, Mukul Gagrani, and Rahul Jain. Posterior sampling-based reinforcement learning for control of unknown linear systems. *IEEE Transactions on Automatic Control*, 65(8):3600–3607, 2019.

[34] Huyên Pham. *Continuous-time stochastic control and optimization with financial applications*, volume 61. Springer Science & Business Media, 2009.

[35] Syed Ali Asad Rizvi and Zongli Lin. Output feedback reinforcement learning control for the continuous-time linear quadratic regulator problem. In *2018 Annual American Control Conference (ACC)*, pages 3417–3422. IEEE, 2018.

[36] Hanspeter Schmidli. *Stochastic control in insurance*. Springer Science & Business Media, 2007.

[37] Allamaraju Subrahmanyam and Ganti Prasada Rao. *Identification of Continuous-time Systems: Linear and Robust Parameter Estimation*. CRC Press, 2019.

[38] Haoran Wang, Thaleia Zariphopoulou, and Xun Yu Zhou. Reinforcement learning in continuous time and space: A stochastic control approach. *J. Mach. Learn. Res.*, 21:198–1, 2020.

[39] Jiongmin Yong and Xun Yu Zhou. *Stochastic controls: Hamiltonian systems and HJB equations*, volume 43. Springer Science & Business Media, 1999.

[40] Ingvar Ziemann and Henrik Sandberg. On a phase transition of regret in linear quadratic control: The memoryless case. *IEEE Control Systems Letters*, 5(2):695–700, 2020.

[41] Ingvar Ziemann and Henrik Sandberg. Regret lower bounds for unbiased adaptive control of linear quadratic regulators. *IEEE Control Systems Letters*, 4(3):785–790, 2020.

Contents

A Numerical Illustrations of Estimation Error and Regret

Now, we provide numerical analyses for showcasing the performance of Algorithm 1 for estimating the unknown dynamics matrices and learning the optimal policy. For this purpose, we assume that the true continuous-time system matrices are lateral-directional state-space matrices of X-29A airplane at 4000 ft altitude [7]. The system is of dimension $d_X = 4$, is controlled by two dimensional commands; $d_U = 2$, and the transition and input matrices in (1) are

$$
A_\star = \begin{bmatrix} -0.1850 & 0.1475 & -0.9825 & 0.1120 \\ -0.3467 & -1.710 & 0.9029 & -0.5843 \times 10^{-6} \\ 1.174 & -0.0825 & -0.1826 & -0.4428 \times 10^{-7} \\ 0.0 & 1.0 & 0.1429 & 0.0 \end{bmatrix}, B_\star = \begin{bmatrix} -0.4470 \times 10^{-3} & 0.4020 \times 10^{-3} \\ 0.3715 & 0.0549 \\ 0.0265 & -0.0135 \\ 0.0 & 0.0 \end{bmatrix}.
$$

Note that the above dimensions of the control action and the state vector, as well as open-loop instability of $A_\star$ and the notably small entries of $B_\star$, render control of the above-mentioned airplane challenging. Further, we let the coefficient matrix of the Brownian disturbance W_t be $C = 0.2 \times I_4$, and employ Algorithm 1 to learn to control a quadratic cost with the weight matrices $Q = 10 \times I_{d_X}, R = I_{d_U}$. The online reinforcement learning policy of Algorithm 1 is run for 500 seconds, while the parameter estimates are updated at times $t = \gamma^n$, for integer values of n and $\gamma = 1.2$. In order to find an initial stabilizing feedback, we run the Bayesian-learning stabilization algorithm for 25 seconds [19].

B Proof of Theorem 1 (Optimal policy)

Fixing $\epsilon > 0$, suppose that the control inputs U_t are frozen in intervals of length ϵ and can change only at times $k\epsilon$, for $k = 0, 1, \cdots$. That is, for all times t satisfying $k\epsilon \leq t < (k+1)\epsilon$, the action vector is fixed; $U_t = U_{k\epsilon}$. Next, we proceed towards finding a decision-making policy for minimizing the expected average cost. Note that due to the above-mentioned freezing during ϵ-length intervals, resulting decision-making policies can be sub-optimal, and indeed provide an upper bound for the optimal cost value.

Next, fix an arbitrary time horizon T, and denote the minimum cost-to-go at time t by

$$
\mathcal{V}_t (X_t) = \inf \mathbb{E} \left[\int_t^T c (X_s, U_s) \mathrm{d}s \middle| \mathcal{F}_t \right],
$$

where the infimum is taken over non-anticipating policies that freeze the control action in ϵ-length intervals, as elaborated above, and the information at time t is $\mathcal{F}_t = \sigma (X_{0:t}, U_{0:t})$; the sigma-field generated by the state and action vectors up to the time. Now, finding an optimal policy is equivalent to applying dynamic programming principle and writing Bellman optimality equations [25, 12]. So, we have

$$
\mathcal{V}_{k\epsilon} (X_{k\epsilon}) = \min_{U_{k\epsilon}} \mathbb{E} \left[\int_{k\epsilon}^{(k+1)\epsilon} c (X_t, U_{k\epsilon}) \mathrm{d}t + \mathcal{V}_{(k+1)\epsilon} \left(X_{(k+1)\epsilon} \right) \middle| \mathcal{F}_{k\epsilon} \right], \tag{10}
$$

subject to the dynamics equation in (1).

For the sake of simplicity, suppose that T/ϵ is an integer. Solving (10) for $k = T/\epsilon - 1$, we get the optimal control action $U_{k\epsilon}^\star = 0$. Accordingly, this gives

$$
\mathcal{V}_{(k+1)\epsilon} \left(X_{(k+1)\epsilon} \right) = X_{(k+1)\epsilon}^\top Q X_{(k+1)\epsilon} \epsilon,
$$

for $k = T/\epsilon - 2$, which, after substituting in (10), becomes

$$
\mathcal{V}_{k\epsilon} (X_{k\epsilon}) = \min_{U_{k\epsilon}} \int_{k\epsilon}^{(k+1)\epsilon} \mathbb{E} \left[X_t^\top Q X_t \middle| \mathcal{F}_{k\epsilon} \right] \mathrm{d}t + U_{k\epsilon}^\top R U_{k\epsilon} \epsilon + \mathbb{E} \left[X_{(k+1)\epsilon}^\top Q X_{(k+1)\epsilon} \middle| \mathcal{F}_{k\epsilon} \right] \epsilon, \tag{11}
$$

where we applied Fubini's Theorem to derive

$$
\mathbb{E} \left[\int_{k\epsilon}^{(k+1)\epsilon} c (X_t, U_{k\epsilon}) \mathrm{d}t \middle| \mathcal{F}_{k\epsilon} \right] = \int_{k\epsilon}^{(k+1)\epsilon} \mathbb{E} \left[X_t^\top Q X_t \middle| \mathcal{F}_{k\epsilon} \right] \mathrm{d}t + U_{k\epsilon}^\top R U_{k\epsilon} \epsilon. \tag{12}
$$

However, solving the dynamics (1) for $k\epsilon \leq t \leq (k+1)\epsilon$, we obtain

$$X_t = e^{A_\star(t-k\epsilon)}X_{k\epsilon} + \int_{k\epsilon}^{t} e^{A_\star(t-s)}C\mathrm{d}W_s + \int_{k\epsilon}^{t} e^{A_\star(t-s)}\mathrm{d}s B_\star U_{k\epsilon},$$

which together with Ito's Lemma, $\mathrm{d}W_s\mathrm{d}W_s^\top = I_{d_W}\mathrm{d}s$ [32], yields to

$$\mathbb{E}\left[X_t^\top Q X_t \big| \mathcal{F}_{k\epsilon}\right] = \int_{k\epsilon}^{t} \mathbf{tr}\left(e^{A_\star^\top(t-s)}Qe^{A_\star(t-s)}CC^\top\right)\mathrm{d}s$$

$$+ \left(e^{A_\star(t-k\epsilon)}X_{k\epsilon} + \int_{k\epsilon}^{t} e^{A_\star(t-s)}\mathrm{d}s B_\star U_{k\epsilon}\right)^\top Q \left(e^{A_\star(t-k\epsilon)}X_{k\epsilon} + \int_{k\epsilon}^{t} e^{A_\star(t-s)}\mathrm{d}s B_\star U_{k\epsilon}\right).$$

Plugging these results in the dynamic programming equation in (11), the expression in front of the minimum becomes the following quadratic function of $U_{k\epsilon}$:

$$X_{k\epsilon}^\top \widetilde{Q} X_{k\epsilon} + 2 X_{k\epsilon}^\top \widetilde{G} U_{k\epsilon} + U_{k\epsilon}^\top \widetilde{R} U_{k\epsilon}$$

$$+ \left(\widetilde{A}X_{k\epsilon} + \widetilde{B}U_{k\epsilon}\right)^\top P_{k+1}\left(\widetilde{A}X_{k\epsilon} + \widetilde{B}U_{k\epsilon}\right) + \mathbf{tr}\left(\widetilde{P}_{k+1}CC^\top\right),$$

where $P_{k+1} = Q\epsilon$, and

$$\widetilde{A} = e^{A_\star\epsilon},$$

$$\widetilde{B} = \int_0^\epsilon e^{A_\star s}\mathrm{d}s B_\star,$$

$$\widetilde{Q} = \int_0^\epsilon e^{A_\star^\top t}Qe^{A_\star t}\mathrm{d}t,$$

$$\widetilde{G} = \int_0^\epsilon e^{A_\star^\top t}Q\left(\int_0^t e^{A_\star(t-s)}\mathrm{d}s\right)B_\star\mathrm{d}t,$$

$$\widetilde{R} = R\epsilon + \int_0^\epsilon B_\star^\top\left(\int_0^t e^{A_\star^\top(t-s)}Qe^{A_\star(t-s)}\mathrm{d}s\right)B_\star\mathrm{d}t,$$

$$\widetilde{P}_{k+1} = P_{k+1}\int_0^\epsilon e^{A_\star^\top s}e^{A_\star s}\mathrm{d}s + \int_0^\epsilon\left(\int_0^t e^{A_\star^\top s}Qe^{A_\star s}\mathrm{d}s\right)\mathrm{d}t.$$

Note that in the last equation above, we used Ito Isometry [4] to find $\widetilde{P}_{k+1}$. Now, performing the minimization the optimal control action is

$$U_{k\epsilon}^\star = -\left(\widetilde{B}^\top P_{k+1}\widetilde{B} + \widetilde{R}\right)^{-1}\left(\widetilde{B}^\top P_{k+1}\widetilde{A} + \widetilde{G}^\top\right)X_{k\epsilon},$$

and (11) leads to

$$\mathcal{V}_{k\epsilon}(X_{k\epsilon}) = X_{k\epsilon}^\top P_k X_{k\epsilon} + \mathbf{tr}\left(CC^\top\left[P_{k+1}\int_0^\epsilon e^{A_\star^\top s}e^{A_\star s}\mathrm{d}s + \int_0^\epsilon\left(\int_0^t e^{A_\star^\top s}Qe^{A_\star s}\mathrm{d}s\right)\mathrm{d}t\right]\right), \quad (13)$$

where P_k is calculated according to the discrete time Riccati equation

$$P_k = \widetilde{Q} + \widetilde{A}^\top P_{k+1}\widetilde{A} - \left(\widetilde{G} + \widetilde{A}^\top P_{k+1}\widetilde{B}\right)\left(\widetilde{B}^\top P_{k+1}\widetilde{B} + \widetilde{R}\right)^{-1}\left(\widetilde{B}^\top P_{k+1}\widetilde{A} + \widetilde{G}^\top\right). \quad (14)$$

It is shown that if there is some matrix L such that $\boldsymbol{\lambda}_{\max}\left(\widetilde{A} + \widetilde{B}L\right) < 1$, then as $k \to -\infty$, the matrix P_k in the above discrete time Riccati equation converges to a uniquely existing matrix P that solves the algebraic Riccati equation

$$P = \widetilde{Q} + \widetilde{A}^\top P \widetilde{A} - \left(\widetilde{G} + \widetilde{A}^\top P \widetilde{B}\right)\left(\widetilde{B}^\top P \widetilde{B} + \widetilde{R}\right)^{-1}\left(\widetilde{B}^\top P \widetilde{A} + \widetilde{G}^\top\right), \tag{15}$$

regardless of the terminal matrix for the largest value $k + 1$, which here corresponds to $P_{T/\epsilon}$ [11, 14, 20].

Next, we show that if ϵ is sufficiently small, then the matrix L mentioned above exists. To that end, write

$$\widetilde{A} = e^{A_\star \epsilon} = \sum_{n=0}^{\infty} \frac{A_\star^n \epsilon^n}{n!} = I_{d_X} + \epsilon M(\epsilon) A_\star,$$

$$\widetilde{B} = \sum_{n=0}^{\infty} \int_0^\epsilon \frac{A_\star^n s^n}{n!} \mathrm{d}s B_\star = \sum_{n=0}^{\infty} \frac{A_\star^n \epsilon^{n+1}}{(n+1)!} B_\star = \epsilon M(\epsilon) B_\star,$$

where

$$M(\epsilon) = \sum_{n=1}^{\infty} \frac{A_\star^{n-1} \epsilon^{n-1}}{n!} = I_{d_X} + \epsilon \sum_{n=2}^{\infty} \frac{A_\star^{n-1} \epsilon^{n-2}}{n!}.$$

Then, letting L be as in Assumption 1, if ϵ is small enough, it holds that

$$\overline{\boldsymbol{\lambda}}\left(M(\epsilon)\left(A_\star + B_\star L\right)\right) < 0. \tag{16}$$

That is because the eigenvalues of the matrix $M(\epsilon)\left(A_\star + B_\star L\right)$ are continuous functions of ϵ, and for $\epsilon = 0$ we have $\overline{\boldsymbol{\lambda}}\left(M(0)\left(A_\star + B_\star L\right)\right) = \overline{\boldsymbol{\lambda}}\left((A_\star + B_\star L)\right) < 0$, according to Assumption 1. Hence, $\widetilde{A} + \widetilde{B}L = I_{d_X} + M(\epsilon)\left(A_\star + B_\star L\right)\epsilon$ implies that eigenvalues of $\widetilde{A} + \widetilde{B}L$ are exactly one plus the eigenvalues of $M(\epsilon)\left(A_\star + B_\star L\right)\epsilon$. So, it holds that

$$\boldsymbol{\lambda}_{\max}\left(\widetilde{A} + \widetilde{B}L\right)^2 \leq 1 + 2\epsilon\overline{\boldsymbol{\lambda}}\left(M(\epsilon)\left(A_\star + B_\star L\right)\right) + \boldsymbol{\lambda}_{\max}\left(M(\epsilon)\left(A_\star + B_\star L\right)\right)^2 \epsilon^2. \tag{17}$$

Now, putting (16) and (17) together, if ϵ is small enough, then $\boldsymbol{\lambda}_{\max}\left(\widetilde{A} + \widetilde{B}L\right) < 1$. Henceforth, suppose that ϵ is sufficiently small so that the latter inequality holds true.

As long as $\epsilon > 0$ is small enough as described above, letting the time horizon T tend to infinity, the ϵ-length frozen optimal policy for minimizing the expected average cost is

$$U_{k\epsilon}^\star = -\left(\widetilde{B}^\top P \widetilde{B} + \widetilde{R}\right)^{-1}\left(\widetilde{B}^\top P \widetilde{A} + \widetilde{G}^\top\right) X_{k\epsilon}, \tag{18}$$

where P is the unique solution of (15). On the other hand, for a fixed time horizon T, as ϵ shrinks the discrete-time Riccati equation in (14) becomes a continuous-time Riccati equation as follows. First, we have

$$\lim_{\epsilon \to 0} \frac{\widetilde{A} - I_{d_X}}{\epsilon} = A_\star,$$

$$\lim_{\epsilon \to 0} \frac{\widetilde{B}}{\epsilon} = B_\star,$$

$$\lim_{\epsilon \to 0} \frac{\widetilde{Q}}{\epsilon} = Q,$$

$$\lim_{\epsilon \to 0} \frac{\widetilde{G}}{\epsilon} = 0,$$

$$\lim_{\epsilon \to 0} \frac{\widetilde{R}}{\epsilon} = R.$$

553 Using these limits, letting $\epsilon \to 0$ in (14) leads to

$$
\begin{aligned}
\lim_{\epsilon \to 0} \frac{P_k - P_{k+1}}{\epsilon} &= \lim_{\epsilon \to 0} \frac{\widetilde{Q}}{\epsilon} + \lim_{\epsilon \to 0} \frac{\widetilde{A}^\top P_{k+1} \widetilde{A} - P_{k+1}}{\epsilon} \\
&\quad - \lim_{\epsilon \to 0} \left(\frac{\widetilde{G} + \widetilde{A}^\top P_{k+1} \widetilde{B}}{\epsilon} \right) \left(\frac{\widetilde{B}^\top P_{k+1} \widetilde{B} + \widetilde{R}}{\epsilon} \right)^{-1} \left(\frac{\widetilde{B}^\top P_{k+1} \widetilde{A} + \widetilde{G}^\top}{\epsilon} \right) \\
&= Q + A_\star^\top P_{k+1} + P_{k+1} A_\star - P_{k+1} B_\star R^{-1} B_\star^\top P_{k+1}.
\end{aligned}
$$

554 That is, the backward differential equation

$$
-\frac{\mathrm{d}P_t}{\mathrm{d}t} = \Phi_{A_\star, B_\star}(P), \tag{19}
$$

555 with the terminal condition $P_T = 0$. Thus, as $\epsilon \to 0$, the optimal policy becomes

$$
U_t^\star = -R^{-1} B_\star^\top P_t X_t,
$$

556 where P_t is the solution of (19). Similarly, letting $\epsilon \to 0$ in (15), we get the optimal policy
557 $U_t^\star = \mathcal{L}(A_\star, B_\star) X_t$ for minimizing the infinite horizon expected average cost, where

$$
\mathcal{L}(A_\star, B_\star) = -R^{-1} B_\star^\top \mathcal{K}(A_\star, B_\star),
$$

558 and $\mathcal{K}(A_\star, B_\star)$ solves $\Phi_{A_\star, B_\star}(P) = 0$. Equivalently, letting $P_{t,T}$ be the solution of (19) when the
559 time horizon is T, it holds that $\lim_{T \to \infty} P_{0,T} = \mathcal{K}(A_\star, B_\star)$, where $\mathcal{K}(A_\star, B_\star)$ solves $\Phi_{A_\star, B_\star}(P) = 0$.
560 Note that all these relationships rely on the convergence of discrete time Riccati equation (14) to the
561 algebraic Riccati equation (15), as $T \to \infty$.

562 Next, subtracting $\mathcal{V}_{(k+1)\epsilon}(X_{k\epsilon})$ from both sides of (10), dividing by ϵ, and letting $\epsilon \to 0$, Ito Isomery
563 implies that

$$
-\frac{\partial \mathcal{V}_t(X_t)}{\partial t} \mathrm{d}t = \min_{U_t} c(X_t, U_t)\, \mathrm{d}t + \mathbb{E} \left[\mathrm{d}X_t^\top \frac{\partial \mathcal{V}_t(X_t)}{\partial X_t} + \frac{1}{2} \mathrm{d}X_t^\top \frac{\partial^2 \mathcal{V}_t(X_t)}{\partial X_t \partial X_t^\top} \mathrm{d}X_t \,\Big|\, \mathcal{F}_t \right],
$$

564 where we used the limits of the matrices $\widetilde{Q}, \widetilde{G}, \widetilde{R}$ as $\epsilon \to 0$ to find the expression on the right-hand-
565 side of the above equality. Note that the above partial derivatives exist according to (13) together
566 with Dominated Convergence Theorem. Hence, substituting for $\mathrm{d}X_t$ from the dynamics (1), and
567 leveraging Ito's Lemma, we obtain the Hamilton-Jacobi-Bellman [39] equation

$$
-\frac{\partial \mathcal{V}_t(X_t)}{\partial t} = \min_{U_t} c(X_t, U_t) + \frac{\partial \mathcal{V}_t(X_t)}{\partial X_t}^\top (A_\star X_t + B_\star U_t) + \frac{1}{2} \mathbf{tr} \left(\frac{\partial^2 \mathcal{V}_t(X_t)}{\partial X_t \partial X_t^\top} CC^\top \right). \tag{20}
$$

568 Further, letting $\epsilon \to 0$, the expression in (13) gives

$$
\mathcal{V}_t(X_t) = X_t^\top P_t X_t + \int_t^T \mathbf{tr}\left(CC^\top P_s \right) \mathrm{d}s, \tag{21}
$$

569 where P_t solve (19). This can be equivalently obtained using the fact that a quadratic function of the
570 form $\mathcal{V}_t(X_t) = X_t^\top F_t X_t + \varphi_t$ solves the partial differential equation (20), as long as

$$
\begin{aligned}
-\frac{\mathrm{d}\varphi_t}{\mathrm{d}t} - X_t^\top \frac{\mathrm{d}F_t}{\mathrm{d}t} X_t &= \min_{U_t} X_t^\top Q X_t + U_t^\top R U_t \\
&\quad + 2 X_t^\top F_t (A_\star X_t + B_\star U_t) + \mathbf{tr}\left(F_t CC^\top \right),
\end{aligned}
$$

571 which after solving for U_t gives the optimal policy $U_t^\star = -R^{-1} B_\star^\top F_t X_t$, as well as

$$
\begin{aligned}
-\frac{\mathrm{d}\varphi_t}{\mathrm{d}t} - X_t^\top \frac{\mathrm{d}F_t}{\mathrm{d}t} X_t &= X_t^\top Q X_t + 2 X_t^\top F_t (A_\star X_t) \\
&\quad - X_t^\top F_t B_\star R^{-1} B_\star^\top F_t X_t + \mathbf{tr}\left(F_t CC^\top \right).
\end{aligned}
$$

572 Because the equation above needs to hold for an arbitrary X_t, it splits to

$$
-\frac{\mathrm{d}F_t}{\mathrm{d}t} = \Phi_{A_\star, B_\star}(F_t), \quad \frac{\mathrm{d}\varphi_t}{\mathrm{d}t} = -\mathbf{tr}\left(F_t CC^\top \right),
$$

573 that is, F_t solves (19). Further, note that cost-to-go at time T is zero because time-to-go is zero, which

574 provides the terminal condition $\mathcal{V}_T(X_T) = 0$, implying that $\varphi_t = \int_t^T \mathbf{tr}\left(CC^\top F_s\right)\mathrm{d}s$. Therefore, the

575 solutions F_t, φ_t of (20) lead to the same expression as in (21).

576 Finally, the expected average cost of the policy $U_t = \mathcal{L}(A_\star, B_\star)X_t$ is the limit of the expected

577 average cost of the policy $U_t = -R^{-1}B_\star^\top P_{t,T}X_t$, as $T \to \infty$;

$$
\begin{aligned}
& \limsup_{T\to\infty} \frac{1}{T}\mathbb{E}\left[\int_0^T c(X_s, U_s)\mathrm{d}s\right] \\
=\ & \limsup_{T\to\infty} \frac{1}{T}\int_0^T \mathbf{tr}\left(CC^\top P_{s,T}\right)\mathrm{d}s \\
=\ & \mathbf{tr}\left(CC^\top \lim_{T\to\infty} P_{s,T}\right) \\
=\ & \mathbf{tr}\left(CC^\top \mathcal{K}(A_\star, B_\star)\right).
\end{aligned}
$$

578 Moreover, suppose that $C = 0$, and apply the policy $U_t = \mathcal{L}(A_\star, B_\star)X_t$. Then, the state trajectory

579 becomes $X_t = e^{D_\star t}X_0$, where $D_\star = A_\star + B_\star\mathcal{L}(A_\star, B_\star)$. So, by (21), we have

$$
\begin{aligned}
& X_0^\top \mathcal{K}(A_\star, B_\star) X_0 \\
=\ & \int_0^\infty X_t^\top \left(Q + \mathcal{L}(A_\star, B_\star)^\top R\mathcal{L}(A_\star, B_\star)\right) X_t \mathrm{d}t \\
=\ & X_0^\top \int_0^\infty e^{D_\star^\top t} \left(Q + \mathcal{L}(A_\star, B_\star)^\top R\mathcal{L}(A_\star, B_\star)\right) e^{D_\star t} \mathrm{d}t X_0,
\end{aligned}
$$

580 for an arbitrary initial state X_0. Thus, (6) holds:

$$
\mathcal{K}(A_\star, B_\star) = \int_0^\infty e^{D_\star^\top t}\left(Q + \mathcal{L}(A_\star, B_\star)^\top R\mathcal{L}(A_\star, B_\star)\right) e^{D_\star t}\mathrm{d}t. \tag{22}
$$

581 Since Q is positive definite, the above equality implies that $\overline{\boldsymbol{\lambda}}(D_\star) < 0$, as well as

$$
\begin{aligned}
& D_\star^\top \mathcal{K}(A_\star, B_\star) + \mathcal{K}(A_\star, B_\star) D_\star \\
+\ & Q + \mathcal{L}(A_\star, B_\star)^\top R\mathcal{L}(A_\star, B_\star) = 0.
\end{aligned} \tag{23}
$$

582 So far, we have shown that by restricting our search for an optimal decision-making policy to the

583 class of policies that the control action is frozen during intervals of length ϵ, and then letting ϵ decay

584 to vanish, we obtain optimal policies given by (19). Next, we show that these policies are optimal in

585 the larger class of all control policies satisfying the information criteria at every time. That is, for all

586 t, the control action U_t can be determined using $\mathcal{F}_t = \sigma(X_{0:t}, U_{0:t})$. For this purpose, first note that

587 the decision-making policy $U_t = R^{-1}B_\star^\top \mathcal{K}(A_\star, B_\star)X_t$ provides an upper-bound for the optimal

588 expected average cost. That is,

$$
\inf_{\boldsymbol{\pi}} \mathcal{J}_{\boldsymbol{\pi}} \le \mathbf{tr}\left(CC^\top \mathcal{K}(A_\star, B_\star)\right).
$$

589 Now, suppose that there is another policy, denoted by $\widetilde{\pi}$, that satisfies

590 $\mathcal{J}_{\widetilde{\pi}} \le \mathbf{tr}\left(CC^\top \mathcal{K}(A_\star, B_\star)\right)$. Define cost-to-go of the policy $\widetilde{\pi}$ by

$$
\widetilde{\mathcal{V}}_t(X_t) = \mathbb{E}\left[\int_t^T c_{\widetilde{\pi}}(X_s, U_s)\mathrm{d}s \,\middle|\, \mathcal{F}_t\right],
$$

591 where T is large enough to satisfy

592 $\widetilde{\mathcal{V}}_t(X_t) \le 2X_t^\top \mathcal{K}(A_\star, B_\star)X_t + 2T\mathbf{tr}\left(CC^\top \mathcal{K}(A_\star, B_\star)\right)$, for all $0 \le t \le 1$. Note that such T

exists since $\widetilde{\pi}$ provides a smaller expected average cost than the policy $U_t = R^{-1}B_\star^\top \mathcal{K}(A_\star, B_\star) X_t$, and the desired upper-bound for $\widetilde{\mathcal{V}}_t(X_t)$ is $2\mathcal{V}_t(X_t)$; two times the cost-to-go of the policy $U_t = R^{-1}B_\star^\top \mathcal{K}(A_\star, B_\star) X_t$. Next, writing

$$\widetilde{\mathcal{V}}_t(X_t) = \mathbb{E}\left[\int_t^{t+\epsilon} c_{\widetilde{\pi}}(X_s, U_s)\mathrm{d}s + \widetilde{\mathcal{V}}_{t+\epsilon}(X_{t+\epsilon}) \,\middle|\, \mathcal{F}_t \right],$$

subtract $\widetilde{\mathcal{V}}_{t+\epsilon}(X_t)$ from both sides, and divide by ϵ. Letting ϵ decay to zero, the upper-bound for $\widetilde{\mathcal{V}}_t(X_t)$ in terms of $\mathcal{V}_t(X_t)$ implies that according to Dominated Convergence Theorem, the following derivatives exist and it holds that

$$-\frac{\partial \widetilde{\mathcal{V}}_t(X_t)}{\partial t} = c(X_t, \widetilde{\pi}(\mathcal{F}_t))$$
$$+ \frac{\partial \widetilde{\mathcal{V}}_t(X_t)^\top}{\partial X_t}(A_\star X_t + B_\star \widetilde{\pi}(\mathcal{F}_t)) + \frac{1}{2}\mathbf{tr}\left(\frac{\partial^2 \widetilde{\mathcal{V}}_t(X_t)}{\partial X_t \partial X_t^\top} CC^\top \right).$$

Now, note that since $c(X_t, U_t)$ as well as $B_\star U_t$ are continuous functions of U_t, the above partial differential equation for $\widetilde{\mathcal{V}}_t(X_t)$ indicates that $\widetilde{\pi}(\mathcal{F}_t)$ is a continuous function of X_t. This, together with the fact that W_t is an almost surely continuous function of time t, in lights of the dynamics equation in (1), leads to continuity of state trajectory X_t; i.e., $U_t = \widetilde{\pi}(\mathcal{F}_t)$ is continuous as t varies. Thus, decision-making policies that freeze for ϵ-length intervals provide accurate approximations of $U_t = \widetilde{\pi}(\mathcal{F}_t)$ in a sense that there exists a sequence $\left\{U_t^{(n)}\right\}_{n=1}^\infty$ such that $U_t^{(n)}$ freezes during intervals of the length $1/n$, and it holds that

$$\limsup_{n\to\infty} \limsup_{T\to\infty} \frac{1}{T} \int_0^T \mathbb{E}\left[\left\| U_t^{(n)} - \widetilde{\pi}(\mathcal{F}_t)\right\|\right]\mathrm{d}t = 0.$$

Therefore, we have

$$\mathcal{J}_{\widetilde{\pi}} \geq \inf_{\epsilon>0} \inf \mathcal{J}_{\boldsymbol{\pi}} = \mathbf{tr}\left(\mathcal{K}(A_\star, B_\star) CC^\top \right),$$

where the inner infimum is taken over all policies that freeze during ϵ-length intervals. This shows that the policy $U_t = -R^{-1}B_\star^\top \mathcal{K}(A_\star, B_\star) X_t$ is an optimal one, which completes the proof.

C Proof of Theorem 2 (Stability margin)

First, we study eigenvalues of the sum of two matrices. Suppose that M, Δ are arbitrary square matrices of the same size, and let $M = P^{-1}\Lambda P$ be the Jordan decomposition. That is, $\lambda_1, \cdots, \lambda_k$ are eigenvalues of M, $\Lambda \in \mathbb{C}^{d_X \times d_X}$ is a block diagonal matrix with blocks $\Lambda_1, \cdots, \Lambda_k$, and

$$
\Lambda_i = \begin{bmatrix} \lambda_i & 1 & 0 & \cdots & 0 & 0 \\ 0 & \lambda_i & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & \lambda_i & 1 \\ 0 & 0 & 0 & \cdots & 0 & \lambda_i \end{bmatrix} \in \mathbb{C}^{\boldsymbol{\mu}_i \times \boldsymbol{\mu}_i}. \tag{24}
$$

Further, similar to Definition 2, let $\boldsymbol{\mu}_M = \max_{1 \le i \le k} \boldsymbol{\mu}_i$. We prove that $\overline{\boldsymbol{\lambda}}(M - \Delta)$ is at most

$$
\overline{\boldsymbol{\lambda}}(M) + \boldsymbol{\mu}_M^{1/2} \||P\Delta P^{-1}\|| \vee \left(\boldsymbol{\mu}_M^{1/2} \||P\Delta P^{-1}\||\right)^{1/\boldsymbol{\mu}_M}. \tag{25}
$$

To show the above inequality, first let λ be an eigenvalue of $M - \Delta$ that satisfies $\Re(\lambda) > \overline{\boldsymbol{\lambda}}(M)$. So, $M - \lambda I$ is an invertible matirx, and there exists at least one vector v, such that $v \ne 0$ and $(M - \Delta - \lambda I)P^{-1}v = 0$. Then, $(M - \lambda I)P^{-1}v = \Delta P^{-1}v$ implies that

$$
v = (\Lambda - \lambda I)^{-1} P\Delta P^{-1} v. \tag{26}
$$

Because $\Lambda = \mathrm{diag}(\Lambda_1, \cdots, \Lambda_k)$, the matrix $\Lambda - \lambda I$ is block diagonal as well, and we have $(\Lambda - \lambda I)^{-1} = \mathrm{diag}\left(\left(\Lambda_1 - \lambda I_{\boldsymbol{\mu}_1}\right)^{-1}, \cdots, \left(\Lambda_k - \lambda I_{\boldsymbol{\mu}_k}\right)^{-1}\right)$. Further, it is straightforward to see that $\left(\Lambda_i - \lambda I_{\boldsymbol{\mu}_i}\right)^{-1}$ is

$$
- \begin{bmatrix} (\lambda - \lambda_i)^{-1} & (\lambda - \lambda_i)^{-2} & \cdots & (\lambda - \lambda_i)^{-\boldsymbol{\mu}_i} \\ 0 & (\lambda - \lambda_i)^{-1} & \cdots & (\lambda - \lambda_i)^{-\boldsymbol{\mu}_i + 1} \\ \vdots & \vdots & \vdots & \vdots \\ 0 & \cdots & 0 & (\lambda - \lambda_i)^{-1} \end{bmatrix}.
$$

Therefore, we have

$$
\left\||\left(\Lambda_i - \lambda I_{\boldsymbol{\mu}_i}\right)^{-1}\right\|| \le \boldsymbol{\mu}_i^{1/2} \left(|\lambda - \lambda_i| \wedge |\lambda - \lambda_i|^{\boldsymbol{\mu}_i}\right)^{-1}.
$$

Using this bound for the operator norms of blocks of the block-diagonal matrix $(\Lambda - \lambda I)^{-1}$, since $\boldsymbol{\mu}_i \le \boldsymbol{\mu}_M$ and $\Re(\lambda) > \overline{\boldsymbol{\lambda}}(M)$, the equation in (26) implies

$$
\begin{aligned}
1 &\le \left\||(\Lambda - \lambda I)^{-1} P\Delta P^{-1}\right\|| \le \left\||(\Lambda - \lambda I)^{-1}\right\|| \||P\Delta P^{-1}\|| \\
&\le \boldsymbol{\mu}_M^{1/2} \||P\Delta P^{-1}\|| \left(\left(\Re(\lambda) - \overline{\boldsymbol{\lambda}}(M)\right) \wedge \left(\Re(\lambda) - \overline{\boldsymbol{\lambda}}(M)\right)^{\boldsymbol{\mu}_M}\right)^{-1}.
\end{aligned}
$$

So, letting λ be an eigenvalue of $M - \Delta$ that satisfies $\Re(\lambda) = \overline{\boldsymbol{\lambda}}(M - \Delta)$, we obtain (25).

Now, using (25), we compare $A_\star + B_\star \mathcal{L}(A, B)$ and $D = A + B\mathcal{L}(A, B)$. Since $A_\star + B_\star \mathcal{L}(A, B) - D$ is

$$
\Delta_\star = A_\star - A - (B_\star - B) R^{-1} B \mathcal{K}(A, B), \tag{27}
$$

using (5), and letting $M = D$ in (25), we have

$$
\overline{\boldsymbol{\lambda}}(A_\star + B_\star \mathcal{L}(A, B)) \le -\rho + \boldsymbol{\mu}_D^{1/2} \||P^{-1}\|| \||P\|| \||\Delta_\star\|| \vee \left(\boldsymbol{\mu}_D^{1/2} \||P^{-1}\|| \||P\|| \||\Delta_\star\||\right)^{1/\boldsymbol{\mu}_D}.
$$

So, in order to have $\overline{\boldsymbol{\lambda}}(A_\star + B_\star \mathcal{L}(A, B)) < -\delta$, it suffices to show that

$$
\boldsymbol{\mu}_D^{1/2} \||P^{-1}\|| \||P\|| \||\Delta_\star\|| < \rho - \delta \wedge (\rho - \delta)^{\boldsymbol{\mu}_D}. \tag{28}
$$

However, since $\||\Delta_\star\|| \le \mathcal{E}(A, B)\left(1 \vee \frac{\|B\|\zeta}{\lambda_{\min}(R)}\right)$, (7) provides (28), which leads to the desired result.

 ## C.1   Proof of sufficiency of (8) for stabilization bounds

630   Next, we show that $\mathcal{E}(A, B) \leq \epsilon_0$ is sufficient for satbilization and express uniform bounds for ρ, ζ
631   in (5). Let $D_\star = A_\star + B_\star \mathcal{L}(A_\star, B_\star) = P_\star^{-1} \Lambda_\star P_\star$ be the Jordan decomposition as defined in the
632   beginning of the proof, and define the largest block size $\boldsymbol{\mu}_\star = \boldsymbol{\mu}_{D_\star}$, similar to Definition 2. Further,
633   suppose that the following is satisfied:

$$\epsilon_0 \leq \frac{1}{1 \vee \|\mathcal{L}(A_\star, B_\star)\|} \left(\frac{\left(-\overline{\boldsymbol{\lambda}}(D_\star)\right) \wedge \left(-\overline{\boldsymbol{\lambda}}(D_\star)\right)^{\boldsymbol{\mu}_\star}}{\boldsymbol{\mu}_\star^{1/2} \|\!|P_\star^{-1}|\!\|\|\!|P_\star|\!\|} \wedge \left[4 \int_0^\infty \|\!|e^{D_\star t}|\!\|^2 \mathrm{d}t \right]^{-1} \right). \tag{29}$$

634   The inequality in (29) implies that if we write $D_1 = A + B\mathcal{L}(A_\star, B_\star) = A_\star + B_\star \mathcal{L}(A_\star, B_\star) + \Delta_1 =$
635   $D_\star + \Delta_1$, then, the matrix $\Delta_1 = A - A_\star + (B - B_\star) \mathcal{L}(A_\star, B_\star)$ satisfies

$$\|\!|\Delta_1|\!\| < \frac{\left(-\overline{\boldsymbol{\lambda}}(D_\star)\right) \wedge \left(-\overline{\boldsymbol{\lambda}}(D_\star)\right)^{\boldsymbol{\mu}_\star}}{\boldsymbol{\mu}_\star^{1/2} \|\!|P_\star|\!\|\|\!|P_\star^{-1}|\!\|}.$$

So, taking $M = D_\star$, the bound in (25) implies that $\overline{\boldsymbol{\lambda}}(D_1) < 0$. Hence, we can employ Lemma 4 to
study consequences of applying the linear feedback $\mathcal{L}(A_\star, B_\star)$ to a system of dynamics matrices
A, B, and get

$$\mathcal{K}(A, B) \leq M = \mathcal{K}(A, B) + \int_0^\infty e^{D_1^\top t} F e^{D_1 t} \mathrm{d}t,$$

where

$$F = \left[\mathcal{L}(A_\star, B_\star) - \mathcal{L}(A, B)\right]^\top R \left[\mathcal{L}(A_\star, B_\star) - \mathcal{L}(A, B)\right].$$

Above, we used the fact that the initial state $X_0 = x$ in Lemma 4 is arbitrary, and so, the involved
matrices are themselves equal. Further, similar to Lemma 4, it is straightforward to see that

$$M = \int_0^\infty e^{D_1^\top t} \left[Q + \mathcal{L}(A_\star, B_\star)^\top R\mathcal{L}(A_\star, B_\star) \right] e^{D_1 t} \mathrm{d}t.$$

636   This leads to

$$Q + \mathcal{L}(A_\star, B_\star)^\top R\mathcal{L}(A_\star, B_\star) = -D_1^\top M - MD_1$$
$$= -D_\star^\top M - MD_\star - \Delta_1^\top M - M\Delta_1.$$

637   Because $\overline{\boldsymbol{\lambda}}(D_\star) < 0$, the latter equation and (6) provide

$$\begin{aligned} M &= \int_0^\infty e^{D_\star^\top t} \left[Q + \mathcal{L}(A_\star, B_\star)^\top R\mathcal{L}(A_\star, B_\star) + \Delta_1^\top M + M\Delta_1 \right] e^{D_\star t} \mathrm{d}t \\ &= \int_0^\infty e^{D_\star^\top t} \left[Q + \mathcal{L}(A_\star, B_\star)^\top R\mathcal{L}(A_\star, B_\star) \right] e^{D_\star t} \mathrm{d}t + \int_0^\infty e^{D_\star^\top t} \left[\Delta_1^\top M + M\Delta_1 \right] e^{D_\star t} \mathrm{d}t \\ &= \mathcal{K}(A_\star, B_\star) + \int_0^\infty e^{D_\star^\top t} \left[\Delta_1^\top M + M\Delta_1 \right] e^{D_\star t} \mathrm{d}t. \end{aligned}$$

638   Therefore, it holds that $\|\!|M|\!\| \leq \|\!|\mathcal{K}(A_\star, B_\star)|\!\| + 2\|\!|\Delta_1|\!\|\|\!|M|\!\| \int_0^\infty \|\!|e^{D_\star t}|\!\|^2 \mathrm{d}t$, which, according to (29)
639   and $\mathcal{K}(A, B) \leq M$, yields to

$$\|\!|\mathcal{K}(A, B)|\!\| \leq \|\!|M|\!\| \leq 2\|\!|\mathcal{K}(A_\star, B_\star)|\!\|. \tag{30}$$

To proceed, suppose that $v \in \mathbb{C}^{d_X}$ satisfies $\|v\| = 1$ and $Dv = \lambda v$. Now, (6) implies that

$$
\begin{aligned}
v^* \mathcal{K}(A, B) v &= \int_0^\infty v^* e^{D^\top t} \left[Q + \mathcal{L}(A, B)^\top R \mathcal{L}(A, B) \right] e^{Dt} v \mathrm{d}t \\
&= \int_0^\infty \left\| \left[Q + \mathcal{L}(A, B)^\top R \mathcal{L}(A, B) \right]^{\frac{1}{2}} e^{\lambda t} v \right\|^2 \mathrm{d}t,
\end{aligned}
$$

where v^* is the transposed complex conjugate of v. Thus, maximizing the left-hand-side above while taking minimum on the right-hand-side, it holds that

$$
\|\mathcal{K}(A, B)\| \geq \boldsymbol{\lambda}_{\min}(Q) \int_0^\infty e^{2\Re(\lambda)t} \mathrm{d}t \geq \frac{\boldsymbol{\lambda}_{\min}(Q)}{2\Re(-\lambda)}. \tag{31}
$$

Putting (30) and (31) together, we obtain $\overline{\boldsymbol{\lambda}}(D) \leq -\boldsymbol{\lambda}_{\min}(Q) \left(4\|\mathcal{K}(A_\star, B_\star)\|\right)^{-1}$. This and (30) imply that $\mathcal{E}(A, B) \leq \epsilon_0$ is sufficient for (5), with $\rho = \boldsymbol{\lambda}_{\min}(Q) 4^{-1} \|\mathcal{K}(A_\star, B_\star)\|^{-1}$, $\zeta = 2\|\mathcal{K}(A_\star, B_\star)\|$. $\blacksquare$

D Proof of Theorem 3 (Regret analysis)

Let $M = Q + \mathcal{L}(A_\star, B_\star)^\top R \mathcal{L}(A_\star, B_\star)$. Recall that $\boldsymbol{\pi}$ applies $U_t = L_t X_t$ at time t. Now, for a given T, suppose that $\epsilon > 0$ is a fixed small real, and let $N = \lceil T/\epsilon \rceil$. Then, define the sequence of policies $\{\boldsymbol{\pi}_i\}_{i=0}^N$:

$$\boldsymbol{\pi}_i = \begin{cases} U_t = L_t X_t & t < i\epsilon \\ U_t = \mathcal{L}(A_\star, B_\star) X_t & t \geq i\epsilon \end{cases}.$$

Note that as long as one concerns about times $t \leq T$, it holds that $\boldsymbol{\pi}^\star = \boldsymbol{\pi}_0, \boldsymbol{\pi}_N = \boldsymbol{\pi}$. Clearly, since $\mathcal{R}_{\boldsymbol{\pi}_0}(T) = 0$, we have $\mathcal{R}_{\boldsymbol{\pi}}(T) = \sum_{i=0}^{N-1}\left(\mathcal{R}_{\boldsymbol{\pi}_{i+1}}(T) - \mathcal{R}_{\boldsymbol{\pi}_i}(T)\right)$. Thus, Lemma 1 gives

$\mathcal{R}_{\boldsymbol{\pi}}(T) = \sum_{i=0}^{N-1}\left(X_{i\epsilon}^\top F_{i\epsilon} X_{i\epsilon} + 2 X_{i\epsilon}^\top g_{i\epsilon} + \beta_{i\epsilon}\right)$, where the matrix $F_{i\epsilon}$, the vector $g_{i\epsilon}$, and the scalar $\beta_{i\epsilon}$ are defined in (41), (42), and (43), respectively. Now, letting $\epsilon \to 0$, since L_t is piecewise continuous, we have

$$\mathcal{R}_{\boldsymbol{\pi}}(T) = \int_0^T \left(X_t^\top \widetilde{F}_t X_t + 2 X_t^\top \widetilde{g}_t + \widetilde{\beta}_t\right) \mathrm{d}t, \tag{32}$$

where $\widetilde{F}_t = \lim_{\epsilon \to 0, i\epsilon \to t} \epsilon^{-1} F_{i\epsilon}$, $\widetilde{g}_t = \lim_{\epsilon \to 0, i\epsilon \to t} \epsilon^{-1} g_{i\epsilon}$, and $\widetilde{\beta}_t = \lim_{\epsilon \to 0, i\epsilon \to t} \epsilon^{-1} \beta_{i\epsilon}$. Note that the above limits exist, since $F_{i\epsilon}, g_{i\epsilon}, \beta_{i\epsilon}$ are continuous. To calculate $\widetilde{F}_t, \widetilde{g}_t, \widetilde{\beta}_t$, using Lemma 1 and the piecewise continuity of L_t, we obtain $\widetilde{\beta}_t = 0$,

$$\widetilde{F}_t = S_t + 2 H_t^\top \int_t^T e^{D_\star^\top (s-t)} M e^{D_\star(s-t)} \mathrm{d}s,$$

$$\widetilde{g}_t = \int_t^T \left(H_t^\top e^{D_\star^\top (s-t)} M \int_t^s e^{D_\star(s-u)} C \mathrm{d}W_u\right) \mathrm{d}s,$$

where $S_t = L_t^\top R L_t - \mathcal{L}(A_\star, B_\star)^\top R \mathcal{L}(A_\star, B_\star)$, and

$$H_t = \lim_{\epsilon \to 0} \frac{e^{(A_\star + B_\star L_t)\epsilon} - e^{D_\star \epsilon}}{\epsilon} = B_\star (L_t - \mathcal{L}(A_\star, B_\star)).$$

Now, by (6) and $\int_T^\infty e^{D_\star^\top (s-t)} M e^{D_\star(s-t)} \mathrm{d}s = E_{T-t}$, the expression for $\widetilde{F}_t$ becomes

$$S_t + H_t^\top \mathcal{K}(A_\star, B_\star) + \mathcal{K}(A_\star, B_\star) H_t - H_t^\top E_{T-t} - E_{T-t} H_t. \tag{33}$$

So, after doing some algebra (see (51)), we get

$$\begin{aligned} S_t &+ H_t^\top \mathcal{K}(A_\star, B_\star) + \mathcal{K}(A_\star, B_\star) H_t \\ &= (L_t - \mathcal{L}(A_\star, B_\star))^\top R (L_t - \mathcal{L}(A_\star, B_\star)). \end{aligned} \tag{34}$$

Since W_u has independent increments and in $\widetilde{g}_t$ we have $u \geq t$, Fubini's Theorem gives

$$\mathbb{E}\left[X_t^\top \widetilde{g}_t\right] = \mathbb{E}\left[\mathbb{E}\left[X_t^\top \widetilde{g}_t \big| \sigma(W_{0:t})\right]\right] = \mathbb{E}\left[X_t^\top \mathbb{E}\left[\widetilde{g}_t \big| \sigma(W_{0:t})\right]\right] = 0.$$

Hence, (32), (33), (34), and Fubini's Theorem imply that $\mathbb{E}[\mathcal{R}_{\boldsymbol{\pi}}(T)] = \mathbb{E}[\alpha_T]$.

To proceed towards establishing the second result, apply Stochastic Fubini Theorem [32, 4] to get

$$\begin{aligned} \int_0^T X_t^\top \widetilde{g}_t \mathrm{d}t &= \int_0^T \int_t^T \int_t^s \left(X_t^\top H_t^\top e^{D_\star^\top (s-t)} M e^{D_\star(s-u)} C\right) \mathrm{d}W_u \mathrm{d}s \mathrm{d}t \\ &= \int_0^T \int_0^u \int_u^T \left(X_t^\top H_t^\top e^{D_\star^\top (s-t)} M e^{D_\star(s-u)} C\right) \mathrm{d}s \mathrm{d}t \mathrm{d}W_u = \int_0^T Y_u^\top \mathrm{d}W_u, \end{aligned}$$

where, using the expression for H_t, the vector Y_u can be written as

$$Y_u^\top = \int\limits_0^u \int\limits_u^T \left(X_t^\top H_t^\top e^{D_\star^\top (s-t)} M e^{D_\star(s-u)} C \right) \mathrm{d}s \mathrm{d}t = \int\limits_0^u \left(X_t^\top \left(L_t - \mathcal{L}\left(A_\star, B_\star \right) \right)^\top P_{t,u}^\top \right) \mathrm{d}t,$$

for $P_{t,u}^\top = \int\limits_u^T B_\star^\top e^{D_\star^\top (s-t)} M e^{D_\star(s-u)} C \mathrm{d}s$. Now, letting $V_T = \int\limits_0^T \|Y_u\|^2 \mathrm{d}u$, for $V_T < 1$, Ito Isometry [4], and for $V_T \geq 1$, Lemma 2, imply that

$$\int\limits_0^T Y_u^\top \mathrm{d}W_u = \mathcal{O}\left(d_W V_T^{1/2} \log^{1/2} V_T \right). \tag{35}$$

However, by using the triangle inequality and Fubini's Theorem, we obtain

$$
\begin{aligned}
V_T &\leq \int\limits_0^T \int\limits_0^u \left\| P_{t,u} \left(L_t - \mathcal{L}\left(A_\star, B_\star \right) \right) X_t \right\|^2 \mathrm{d}t \mathrm{d}u \\
&= \int\limits_0^T \left(X_t^\top \left(L_t - \mathcal{L}\left(A_\star, B_\star \right) \right)^\top \left[\int\limits_t^T P_{t,u}^\top P_{t,u} \mathrm{d}u \right] \left(L_t - \mathcal{L}\left(A_\star, B_\star \right) \right) X_t \right) \mathrm{d}t \\
&\leq \int\limits_0^T \boldsymbol{\lambda}_{\max}\left(\int\limits_t^T R^{-1/2} P_{t,u}^\top P_{t,u} R^{-1/2} \mathrm{d}u \right) \left\| R^{1/2} \left(L_t - \mathcal{L}\left(A_\star, B_\star \right) \right) X_t \right\|^2 \mathrm{d}t.
\end{aligned}
$$

The second part of the integrand above appears in $\boldsymbol{\alpha}_T$. So, we proceed by finding an upper-bound for the first part. For this purpose, we use the triangle inequality and (6) to get the equation

$$
\begin{aligned}
\boldsymbol{\lambda}_{\max}\left(\int\limits_t^T P_{t,u}^\top P_{t,u} \mathrm{d}u \right) &\leq \int\limits_t^T \left\|\left| B_\star^\top e^{D_\star^\top (u-t)} \right|\right\|^2 \left\|\left| \int\limits_u^T e^{D_\star^\top (s-u)} M e^{D_\star(s-u)} \mathrm{d}s \right|\right\|^2 \|C\|^2 \mathrm{d}u \\
&\leq \|B_\star\|^2 \|\mathcal{K}\left(A_\star, B_\star \right)\|^2 \|C\|^2 \int\limits_0^\infty \left\|\left| e^{D_\star^\top u} \right|\right\|^2 \mathrm{d}u.
\end{aligned}
$$

Therefore, by using (31), we get

$$V_T \leq \frac{\|B_\star\|^2 \|\mathcal{K}\left(A_\star, B_\star \right)\|^3 \|C\|^2}{\boldsymbol{\lambda}_{\min}\left(Q \right) \boldsymbol{\lambda}_{\min}\left(R \right)} \boldsymbol{\alpha}_T,$$

since E_t decays exponentially with t. So, (35) gives the desired result. ∎

 # E   Proof of Theorem 4 (Analysis of Algorithm 1)

672 In order to establish Theorem 4, we study the estimation procedure in (9) and specify the accuracy at
673 which the algorithm is able to estimate $A_\star, B_\star$. To that end, Lemma 5 and Lemma 6 are utilized to
674 study the Gram matrix $V_n = \int_0^{\gamma^n} Y_s Y_s^\top \mathrm{d}s$ in (9), while Lemma 2 is used for bounding the estimation
675 error. Then, by leveraging Lemma 3, we find the rates of deviating from the optimal policy in (3).
676 Finally, the resulting regret of Algorithm 1 is investigated in lights of Theorem 3.

677 By using (1) to substitute for $\mathrm{d}X_t$, $\left[\int_0^{\gamma^n} Y_s \mathrm{d}X_s^\top \right]^\top V_n^\dagger$ is

$$
\left[\int_0^{\gamma^n} Y_s Y_s^\top [A_\star, B_\star]^\top \mathrm{d}s + \int_0^{\gamma^n} Y_s \mathrm{d}W_s^\top C^\top \right]^\top V_n^\dagger.
$$

678 In (38), we show that V_n is non-singular. So, we have

$$
\left[\int_0^{\gamma^n} Y_s \mathrm{d}X_s^\top \right]^\top V_n^{-1} = [A_\star, B_\star] + \left[V_n^{-1} \int_0^{\gamma^n} Y_s \mathrm{d}W_s^\top C^\top \right]^\top . \tag{36}
$$

679 Because $[A_\star, B_\star] \in \mathcal{S}_0$, (9) and (36) lead to

$$
\mathcal{E}(A_n, B_n) \leq \left\| V_n^{-1} \int_0^{\gamma^n} Y_s \mathrm{d}W_s^\top C^\top \right\| + \|\Theta_n\|.
$$

Since entries of Θ_n are $\mathcal{N}\left(0, \gamma^{-n/2} n^{1/2}\right)$, we have

$$
\mathbb{P}\left(\|\Theta_n\| \geq d_X^{1/2} (d_X + d_U)^{1/2} \gamma^{-n/4} n^{1/2} \right) = \mathcal{O}\left(e^{-n^{1/2}} \right).
$$

This, by Borel-Cantelli Lemma, leads to

$$
\|\Theta_n\| = \mathcal{O}\left(d_X^{1/2} (d_X + d_U)^{1/2} \gamma^{-n/4} n^{1/2} \right).
$$

680 Thus, letting $d = d_X + d_U$, according to Lemma 2, $\mathcal{E}(A_n, B_n)$ is at most

$$
d^{1/2} \mathcal{O}\left(d_W^{1/2} \|C\| \left(\frac{\log \boldsymbol{\lambda}_{\max}(V_n)}{\boldsymbol{\lambda}_{\min}(V_n)} \right)^{1/2} + d_X^{1/2} \gamma^{-n/4} n^{1/2} \right). \tag{37}
$$

681 Now, Lemma 5 provides $\mathcal{O}(\log \boldsymbol{\lambda}_{\max}(V_n)) = n \log \gamma$. Further, we will shortly show that

$$
\liminf_{n \to \infty} \gamma^{-n/2} \boldsymbol{\lambda}_{\min}(V_n) \geq \boldsymbol{\lambda}_{\min}\left(CC^\top \right). \tag{38}
$$

682 Thus, (37) and (38) yield to the upper-bound

$$
\mathcal{E}(A_n, B_n) = \mathcal{O}\left(d^{1/2} \left(d_X^{1/2} + \frac{d_W^{1/2} \|C\| \log^{1/2} \gamma}{\boldsymbol{\lambda}_{\min}(CC^\top)^{1/2}} \right) \gamma^{-n/4} n^{1/2} \right).
$$

This gives the first result in Theorem 4. To prove the other statement, let $\beta_\star$ be as defined in Lemma 3.
So, Lemma 3 implies that

$$
\|\mathcal{L}(A_n, B_n) - \mathcal{L}(A_\star, B_\star)\|^2 = \mathcal{O}\left((d_X + d_U) \beta_\star^2 \left(d_X + \frac{d_W \|C\|^2 \log \gamma}{\boldsymbol{\lambda}_{\min}(CC^\top)} \right) \gamma^{-n/2} n \right).
$$

Now, since during the time period $\gamma^{n-1} \leq t < \gamma^n$ the feedback matrix is frozen to $\mathcal{L}(A_{n-1}, B_{n-1})$,
according to Lemma 5, we have

$$
\int_0^{\gamma^n} \left\| R^{1/2} (L_t - \mathcal{L}(A_\star, B_\star)) X_t \right\|^2 \mathrm{d}t = \mathcal{O}\left(\sum_{k=1}^n \beta_L \gamma^{k-1} \gamma^{-(k-1)/2} k \right),
$$

683 where

$$\beta_L = d\beta_\star^2 \left(d_X + \frac{d_W \|C\|^2 \log \gamma}{\boldsymbol{\lambda}_{\min}(CC^\top)} \right) (\gamma - 1) \|R\| \|C\|^2.$$

Moreover, since by Theorem 1 we have $\overline{\lambda}(D_\star) < 0$, the matrix E_t in Theorem 3 decays exponentially with t. So, it holds that

$$\int_0^T \left(X_t^\top E_{T-t} B_\star (L_t - \mathcal{L}(A_\star, B_\star)) X_t \right) \mathrm{d}t = \mathcal{O}\left(\log^2 T \right).$$

684 Therefore, according to Theorem 3, we have the following:

$$\boldsymbol{\mathcal{R}}_\pi(T) = \mathcal{O}\left(\sum_{k=1}^{\lceil (\log T)/(\log \gamma) \rceil} \gamma^{(k-1)/2} k \right) = \mathcal{O}\left(\frac{\beta_L}{\log \gamma} T^{1/2} \log T \right).$$

685 This, according to $\beta_\star$ in Lemma 3, completes the proof.

686 To prove (38), let $D_{k-1} = A_\star + B_\star \mathcal{L}(A_{k-1}, B_{k-1})$. Then, by Lemma 5, we have

$$\liminf_{k \to \infty} \gamma^{-k} \boldsymbol{\lambda}_{\min}\left(\int_{\gamma^{k-1}}^{\gamma^k} X_t X_t^\top \mathrm{d}t \right) \geq \eta_k \boldsymbol{\lambda}_{\min}(CC^\top), \tag{39}$$

687 where $\eta_k = \left(1 - \gamma^{-1}\right) \left(\int_0^1 \left\| e^{-D_{k-1} s} \right\|^2 \mathrm{d}s \right)^{-1}$. Hence, (39) implies that to establish (38), it suffices

688 to show that the following inequality holds for some $0 \leq \ell < n - 1$:

$$\liminf_{n \to \infty} \boldsymbol{\lambda}_{\min}\left(\sum_{k=\ell}^{n-1} \gamma^{k-n/2} \begin{bmatrix} I_{d_X} \\ \mathcal{L}(A_k, B_k) \end{bmatrix} \begin{bmatrix} I_{d_X} \\ \mathcal{L}(A_k, B_k) \end{bmatrix}^\top \right) \geq \max_{\ell \leq k \leq n-1} \frac{1}{\eta_k}. \tag{40}$$

689 For an arbitrary fixed $\epsilon > 0$, consider the event that the above-mentioned smallest eigenvalue is less
690 than ϵ, and let $\mathcal{M}_n(\epsilon)$ be the set of matrices $[A_k, B_k]_{k=\ell}^{n-1}$ for which this event occurs:

$$\mathcal{M}_n(\epsilon) = \left\{ [A_\ell, B_\ell, \cdots, A_{n-1}, B_{n-1}] : \boldsymbol{\lambda}_{\min}\left(P_{\ell,n} P_{\ell,n}^\top \right) \leq \epsilon \right\},$$

691 where the $(d_X + d_U) \times d_X(n - \ell)$ matrix $P_{\ell,n}$ is

$$\left[\gamma^{\frac{\ell}{2} - \frac{n}{4}} \begin{bmatrix} I_{d_X} \\ \mathcal{L}(A_\ell, B_\ell) \end{bmatrix}, \cdots, \gamma^{\frac{n-1}{2} - \frac{n}{4}} \begin{bmatrix} I_{d_X} \\ \mathcal{L}(A_{n-1}, B_{n-1}) \end{bmatrix} \right].$$

692 Now, note that the set of all matrices

$$F_n = \begin{bmatrix} \gamma^{\ell/2 - n/4} I_{d_X} & \cdots & \gamma^{(n-1)/2 - n/4} I_{d_X} \\ \gamma^{\ell/2 - n/4} L_\ell & \cdots & \gamma^{(n-1)/2 - n/4} L_{n-1} \end{bmatrix},$$

693 that there exists $v \in \mathbb{R}^{d_X + d_U}$ satisfying $\|v\| = 1$ and $F_n^\top v = 0$, is of dimension $d_X + d_U - 1 + (n -$
694 $\ell)(d_U - 1)$. To show that, on one hand, the set of unit $d_X + d_U$ dimensional vectors is (a sphere) of
695 dimension $d_X + d_U - 1$. On the other hand, by writing $v = \left[v_1^\top, v_2^\top \right]^\top$, for $v_1 \in \mathbb{R}^{d_X}$ and $v_2 \in \mathbb{R}^{d_U}$,
696 clearly, $F_n^\top v = 0$ is equivalent to $L_k^\top v_2 = -v_1$, for all $k = \ell, \cdots, n - 1$. The latter enforces every
697 column of L_k to be in a certain hyperplane in $\mathbb{R}^{d_U}$.

698 Thus, according to Lemma 6, the dimension of $\mathcal{M}_n(0)$ is at most $d_X + (d_U - 1)(n - \ell + 1) + (n - \ell)d_X^2$.
699 Further, if ℓ is sufficiently large so that $\gamma^{-\ell + n/2} \epsilon < 1$, then for every $[A_k, B_k]_{k=\ell}^{n-1} \in \mathcal{M}_n(\epsilon)$, there
700 exists some $\left[\widetilde{A}_k, \widetilde{B}_k \right]_{k=\ell}^{n-1} \in \mathcal{M}_n(0)$, such that for all $k = \ell, \cdots, n - 1$, it holds that

$$\left\| [A_k, B_k] - \left[\widetilde{A}_k, \widetilde{B}_k \right] \right\| = \mathcal{O}\left(\gamma^{-k/2 + n/4} \epsilon^{1/2} \right).$$

The random matrices $\{\Theta_k\}_{k=0}^{n-1}$ are independent, and entries of Θ_k are independent identically distributed $\mathcal{N}\left(0, \gamma^{-k/2}k^{1/2}\right)$ random variables. Hence, we have

$$\mathbb{P}(\mathcal{M}_n(\epsilon)) = \left[\mathcal{O}\left(\gamma^{\ell/4}\ell^{-1/4}\gamma^{-\ell/2+n/4}\epsilon^{1/2}\right) \wedge 1\right]^m,$$

where $m = (d_X d_U - d_U + 1)(n - \ell) - d_X - d_U + 1$. To see that, note that $\mathcal{M}_n(0)$ is a $d_X + (d_U - 1)(n - \ell + 1) + (n - \ell)d_X^2$ dimensional object in a $d_X(d_X + d_U)(n - \ell)$ dimensional space. So, the exponent is at least m. Letting $\ell = n - 5$, we have $m \geq 5$. Further, as n grows, $\mathcal{O}\left(\ell^{-1/4}\gamma^{(n-\ell)/4}\epsilon^{1/2}\right) < 1$ holds for $\epsilon = \max_{\ell \leq k \leq n-1} \eta_k^{-1}$. So, we have $\sum_{n=5}^{\infty} \mathbb{P}(\mathcal{M}_n(\epsilon)) = \sum_{n=5}^{\infty} \mathcal{O}\left(n^{-1/4}\right)^5 < \infty$, which by Borel-Cantelli Lemma implies (40). ∎

F Estimation Rates under Persistent Randomization

Proposition 1 *Assume that in Algorithm 1 the variance of entries of Θ_n is σ_n^2, where*

$$\liminf_{n\to\infty} \sigma_n > 0.$$

709   *Then, letting $\boldsymbol{\omega}_{\mathcal{E}}$ be as in Theorem 4, $Y_s = \left[X_s^\top, U_t^\top\right]^\top$, and $V_n = \int\limits_0^{\gamma^n} Y_s Y_s^\top \mathrm{d}s$, we have*

$$\left\|\left\|\left(\int\limits_0^{\gamma^n} Y_s \mathrm{d}X_s^\top\right)^\top V_n^\dagger - [A_\star, B_\star]\right\|\right\|^2 = \mathcal{O}\left(\boldsymbol{\omega}_{\mathcal{E}}\gamma^{-n}n^2\right).$$

Proof. By (36), it suffices to study $\Delta = V_n^\dagger \int\limits_0^{\gamma^n} Y_s \mathrm{d}W_s^\top C^\top$. In the sequel, we show that

$$\liminf_{n\to\infty} n\gamma^{-n}\boldsymbol{\lambda}_{\min}\left(V_n\right) \geq \boldsymbol{\lambda}_{\min}\left(CC^\top\right).$$

So, putting Lemma 2 and Lemma 5 together, we obtain the desired result, since they give

$$\|\Delta\|^2 = \mathcal{O}\left((d_X + d_U)\, d_W \|C\|^2 \frac{\gamma^{-n}n^2\log\gamma}{\boldsymbol{\lambda}_{\min}\left(CC^\top\right)}\right).$$

710   Thus, by (39), it is enough to show that for some $0 \leq \ell < n - 1$,

$$\liminf_{n\to\infty} \boldsymbol{\lambda}_{\min}\left(\sum_{k=\ell}^{n-1} \gamma^{k-n}n \begin{bmatrix} I_{d_X} \\ \mathcal{L}\left(A_k, B_k\right) \end{bmatrix} \begin{bmatrix} I_{d_X} \\ \mathcal{L}\left(A_k, B_k\right) \end{bmatrix}^\top\right)$$

711   is at least $\epsilon = \max\limits_{\ell \leq k \leq n-1} \eta_k^{-1}$. Let $\mathcal{M}_n(\epsilon)$ be the set of $[A_k, B_k]_{k=\ell}^{n-1}$ that the above does not hold:

712   $\mathcal{M}_n(\epsilon) = \left\{[A_\ell, B_\ell, \cdots, A_{n-1}, B_{n-1}] : \boldsymbol{\lambda}_{\min}\left(P_{\ell,n}P_{\ell,n}^\top\right) \leq \epsilon\right\}$, where $P_{\ell,n}$ is

$$\left[\gamma^{\frac{\ell-n}{2}}n^{\frac{1}{2}} \begin{bmatrix} I_{d_X} \\ \mathcal{L}\left(A_\ell, B_\ell\right) \end{bmatrix}, \cdots, \gamma^{-\frac{1}{2}}n^{\frac{1}{2}} \begin{bmatrix} I_{d_X} \\ \mathcal{L}\left(A_{n-1}, B_{n-1}\right) \end{bmatrix}\right].$$

713   Similar to the proof of Theorem 4, for $[A_k, B_k]_{k=\ell}^{n-1} \in \mathcal{M}_n(\epsilon)$, there is $\left[\widetilde{A}_k, \widetilde{B}_k\right]_{k=\ell}^{n-1} \in \mathcal{M}_n(0)$,

714   such that $\left\|\left\|[A_k, B_k] - \left[\widetilde{A}_k, \widetilde{B}_k\right]\right\|\right\|^2 = \mathcal{O}\left(\gamma^{n-k}n^{-1}\epsilon\right)$. Thus, $\liminf\limits_{n\to\infty} \sigma_n > 0$, together with the

715   dimension of $\mathcal{M}_n(0)$ that we calculated in the proof of Theorem 4, leads to

$$\mathbb{P}(\mathcal{M}_n(\epsilon)) = \left[\mathcal{O}\left(\gamma^{(n-\ell)/2}n^{-1/2}\epsilon^{1/2}\right) \wedge 1\right]^m,$$

716   for $m = (d_X d_U - d_U + 1)\left(n - \ell\right) - d_X - d_U + 1$. Finally, $\ell = n - 4$ gives $\sum\limits_{n=4}^{\infty} \mathbb{P}(\mathcal{M}_n(\epsilon)) < \infty$.

717   Therefore, Borel-Cantelli Lemma implies the desired result.   ∎

G Auxiliary Lemmas

In this section, we state the auxiliary lemmas used for establishing the main results and provide their proofs, each subsection corresponding to one lemma.

First, in Lemma 1 in Subsection G.1, we provide expressions for the difference between the regrets of two policies. Study of self-normalized stochastic integrals is the content of Lemma 2, while Lemma 3 on Lipschitz continuity of the optimal feedback with respect to the dynamics matrices is established in Subsection G.3.

Next, in Lemma 4, we consider the total cumulative cost for the case of applying a sub-optimal time-invariant linear feedback policy to a deterministic system. Then, Lemma 5 focuses on explicit calculation of the empirical covariance matrix of the state vectors. Finally, in Lemma 6 in Subsection G.6 we specify the set of dynamics matrices that possess the same optimal linear feedback matrix.

G.1 Difference in regrets of two policies

Lemma 1 *For fixed $0 \leq t_1 \leq t_2 \leq T$, define the policies π_1, π_2 according to*

$$\boldsymbol{\pi}_i = \begin{cases} U_t = LX_t & t < t_i \\ U_t = \mathcal{L}(A_\star, B_\star) X_t & t \geq t_i \end{cases}.$$

Further, let $D_\star = A_\star + B_\star \mathcal{L}(A_\star, B_\star)$, $D = A_\star + B_\star L$, $M_\star = Q + \mathcal{L}(A_\star, B_\star)^\top RL \mathcal{L}(A_\star, B_\star)$, $M = Q + LRL$, $\Delta_t = e^{D(t-t_1)} - e^{D_\star(t-t_1)}$, $Z_t = \int_{t_1}^{t} \left[e^{D(t-s)} - e^{D_\star(t-s)} \right] C dW_s$, and $S = M - M_\star = L^\top RL - \mathcal{L}(A_\star, B_\star)^\top RL \mathcal{L}(A_\star, B_\star)$.

Then, we have $\mathcal{R}_{\pi_2}(T) - \mathcal{R}_{\pi_1}(T) = X_{t_1}^\top F_{t_1} X_{t_1} + 2X_{t_1}^\top g_{t_1} + \beta_{t_1}$, where F_{t_1}, g_{t_1}, and β_{t_1} are

$$
\begin{aligned}
F_{t_1} &= \int_{t_1}^{t_2} \left(e^{D_\star^\top(t-t_1)} S e^{D_\star(t-t_1)} + 2\Delta_t^\top M e^{D_\star(t-t_1)} + \Delta_t^\top M \Delta_t \right) dt \\
&+ \int_{t_2}^{T} \left(2\Delta_{t_2}^\top e^{D_\star^\top(t-t_2)} M_\star e^{D_\star(t-t_1)} + \Delta_{t_2}^\top e^{D_\star^\top(t-t_2)} M_\star e^{D_\star(t-t_2)} \Delta_{t_2} \right) dt, \quad (41)
\end{aligned}
$$

$$
\begin{aligned}
g_{t_1} &= \int_{t_1}^{t_2} \left(S \int_{t_1}^{t} e^{D_\star(t-s)} C dW_s + \Delta_t^\top M \int_{t_1}^{t} e^{D_\star(t-s)} C dW_s + e^{D_\star^\top(t-t_1)} M Z_t \right) dt \\
&+ \int_{t_1}^{t_2} \Delta_t^\top M Z_t dt + \int_{t_2}^{T} \Delta_{t_2}^\top e^{D_\star^\top(t-t_2)} M_\star \left(e^{D_\star(t-t_2)} Z_{t_2} + \int_{t_1}^{t} e^{D_\star(t-s)} C dW_s \right) dt \\
&+ \int_{t_2}^{T} \left(e^{D_\star^\top(t-t_1)} M_\star e^{D_\star(t-t_2)} Z_{t_2} \right) dt, \quad (42)
\end{aligned}
$$

$$
\begin{aligned}
\beta_{t_1} &= \int_{t_1}^{t_2} \left(\left\| S^{1/2} \int_{t_1}^{t} e^{D_\star(t-s)} C dW_s \right\|^2 + 2Z_t^\top M \int_{t_1}^{t} e^{D_\star(t-s)} C dW_s + Z_t^\top M Z_t \right) dt \\
&+ \int_{t_2}^{T} \left(2Z_{t_2}^\top e^{D_\star^\top(t-t_2)} M_\star \int_{t_1}^{t} e^{D_\star(t-s)} C dW_s + Z_{t_2}^\top e^{D_\star^\top(t-t_2)} M_\star e^{D_\star(t-t_2)} Z_{t_2} \right) dt. (43)
\end{aligned}
$$

738 Proof. Letting $X_t^{\boldsymbol{\pi}_i}$ be the state of the system under the policy $\boldsymbol{\pi}_i$, clearly, for $t \leq t_1$, it holds that
739 $X_t^{\boldsymbol{\pi}_1} = X_t^{\boldsymbol{\pi}_2}$. So, we use X_{t_1} for both states at time t_1. Moreover, for $t_1 \leq t \leq t_2$, we have

$$X_t^{\boldsymbol{\pi}_1} = e^{D_\star(t-t_1)} X_{t_1} + \int_{t_1}^t e^{D_\star(t-s)} C \mathrm{d}W_s,$$

$$X_t^{\boldsymbol{\pi}_2} = e^{D(t-t_1)} X_{t_1} + \int_{t_1}^t e^{D(t-s)} C \mathrm{d}W_s,$$

740 where $Y_t = X_t^{\boldsymbol{\pi}_2} - X_t^{\boldsymbol{\pi}_1}$. So, by denoting the instantaneous cost of policy $\boldsymbol{\pi}_i$ at time t by $c_{\boldsymbol{\pi}_i}(t)$, we
741 get $Y_t = \Delta_t X_{t_1} + Z_t$, as well as

$$\int_{t_1}^{t_2} (c_{\boldsymbol{\pi}_2}(t) - c_{\boldsymbol{\pi}_1}(t))\mathrm{d}t = \int_{t_1}^{t_2} \left[(X_t^{\boldsymbol{\pi}_1} + Y_t)^\top M (X_t^{\boldsymbol{\pi}_1} + Y_t) - X_t^{\boldsymbol{\pi}_1\top} M_\star X_t^{\boldsymbol{\pi}_1} \right] \mathrm{d}t$$

$$= \int_{t_1}^{t_2} \left[X_t^{\boldsymbol{\pi}_1\top} S X_t^{\boldsymbol{\pi}_1} + 2Y_t^\top M X_t^{\boldsymbol{\pi}_1} + Y_t^\top M Y_t \right] \mathrm{d}t. \qquad (44)$$

742 On the other hand, for $t \geq t_2$, we have

$$\int_{t_2}^{T} (c_{\boldsymbol{\pi}_2}(t) - c_{\boldsymbol{\pi}_1}(t))\mathrm{d}t = \int_{t_2}^{T} \left[(X_t^{\boldsymbol{\pi}_1} + Y_t)^\top M_\star (X_t^{\boldsymbol{\pi}_1} + Y_t) - X_t^{\boldsymbol{\pi}_1\top} M_\star X_t^{\boldsymbol{\pi}_1} \right] \mathrm{d}t$$

$$= \int_{t_2}^{T} \left[2Y_t^\top M_\star X_t^{\boldsymbol{\pi}_1} + Y_t^\top M_\star Y_t \right] \mathrm{d}t. \qquad (45)$$

743 and

$$X_t^{\boldsymbol{\pi}_i} = e^{D_\star(t-t_2)} X_{t_2}^{\boldsymbol{\pi}_i} + \int_{t_2}^t e^{D_\star(t-s)} C \mathrm{d}W_s,$$

$$Y_t = e^{D_\star(t-t_2)} \left[X_{t_2}^{\boldsymbol{\pi}_2} - X_{t_2}^{\boldsymbol{\pi}_1} \right] = e^{D_\star(t-t_2)} \left[\Delta_{t_2} X_{t_1} + Z_{t_2} \right].$$

744 Thus, putting (44) and (45) together, we obtain the desired results. ∎

G.2 Scaling of self-normalized stochastic integrals

Lemma 2 *Suppose that $Y_t \in \mathbb{R}^m$ is a vector-valued stochastic process such that Y_t is $\mathcal{F}_t$-measurable for the natural filtration $\mathcal{F}_t = \sigma\left(\{W_s\}_{0 \leq s \leq t}\right)$. Then, letting $V_t = \int_0^t Y_s Y_s^\top \mathrm{d}s$, we have*

$$\left\| (I + V_t)^{-1/2} \int_0^t Y_s \mathrm{d}W_s^\top \right\|^2 = \mathcal{O}\left(m d_W \log \boldsymbol{\lambda}_{\max}(V_t) \right).$$

Proof. First, fix $t > 0$, and for an arbitrary $\epsilon > 0$, let $n = \lfloor t/\epsilon \rfloor$. So, for $k = 0, 1, \cdots, n$, consider the sequence of matrices $M_k = \epsilon^{-1} I + \sum_{i=0}^{k} Y_{i\epsilon} Y_{i\epsilon}^\top$. Then, for $k = 1, \cdots, n$, consider the sequence of scalars β_k defined according to $\beta_k = Y_{k\epsilon}^\top M_{k-1}^{-1} Y_{k\epsilon}$. Using the formula for determinants of the products of matrices, we have

$$\det M_k = \det \left[M_{k-1} \left(I + M_{k-1}^{-1} Y_{k\epsilon} Y_{k\epsilon}^\top \right) \right] = \det (M_{k-1}) \det \left(I + M_{k-1}^{-1} Y_{k\epsilon} Y_{k\epsilon}^\top \right).$$

746 Since all eigenvalues of $I + M_{k-1}^{-1} Y_{k\epsilon} Y_{k\epsilon}^\top$ are unit, except one of them which is $1 + \beta_k$, we have
747 $(1 + \beta_k) \det M_{k-1} = \det M_k$. On the other hand, matrix inversion formula gives

$$M_k^{-1} = M_{k-1}^{-1} - \frac{1}{1 + Y_{k\epsilon}^\top M_{k-1}^{-1} Y_{k\epsilon}} M_{k-1}^{-1} Y_{k\epsilon} Y_{k\epsilon}^\top M_{k-1}^{-1},$$

which leads to

$$Y_{k\epsilon}^\top M_k^{-1} Y_{k\epsilon} = Y_{k\epsilon}^\top \left(M_{k-1} + Y_{k\epsilon} Y_{k\epsilon}^\top \right)^{-1} Y_{k\epsilon} = \beta_k - \frac{\beta_k^2}{1 + \beta_k} = 1 - \frac{1}{1 + \beta_k} = 1 - \frac{\det M_{k-1}}{\det M_k}.$$

748 Further, by using the inequality $1 - \beta \le -\log \beta$ for $\beta > 0$, the latter equality gives

$$Y_{k\epsilon}^\top M_k^{-1} Y_{k\epsilon} \le \log \det M_k - \log \det M_{k-1}. \tag{46}$$

749 Now, let $F_k = \sum_{i=0}^{k} Y_{i\epsilon} \left(W_{(i+1)\epsilon} - W_{i\epsilon} \right)^\top$. Using the facts that $Y_{k\epsilon}, F_{k-1}$, and M_k all are $\mathcal{F}_{k\epsilon}$-
750 measurable, the Brownian motion W_t has independent increments, and its covariance matrix is a
751 multiple of identity, properties of conditional expectations give

$$\mathbb{E}\left[F_k^\top M_k^{-1} F_k \right] = \mathbb{E}\left[\mathbb{E}\left[F_k^\top M_k^{-1} F_k \middle| \mathcal{F}_{k\epsilon} \right] \right]$$

$$= \mathbb{E}\left[\mathbb{E}\left[\left(F_{k-1} + Y_{k\epsilon} \left(W_{(k+1)\epsilon} - W_{k\epsilon} \right)^\top \right)^\top M_k^{-1} \left(F_{k-1} + Y_{k\epsilon} \left(W_{(k+1)\epsilon} - W_{k\epsilon} \right)^\top \right) \middle| \mathcal{F}_{k\epsilon} \right] \right]$$

$$= \mathbb{E}\left[F_{k-1}^\top M_k^{-1} F_{k-1} + \mathbb{E}\left[\left(W_{(k+1)\epsilon} - W_{k\epsilon} \right) Y_{k\epsilon}^\top M_k^{-1} Y_{k\epsilon} \left(W_{(k+1)\epsilon} - W_{k\epsilon} \right)^\top \middle| \mathcal{F}_{k\epsilon} \right] \right]$$

$$= \mathbb{E}\left[F_{k-1}^\top M_k^{-1} F_{k-1} + \left(Y_{k\epsilon}^\top M_k^{-1} Y_{k\epsilon} \right) \epsilon I \right].$$

So, using (46) together with the fact that (as positive semidefinite matrices) the order $M_{k-1} \le M_k$
holds, we get the telescopic relationships

$$\boldsymbol{\lambda}_{\max}\left(\mathbb{E}\left[F_k^\top M_k^{-1} F_k \right] \right) - \boldsymbol{\lambda}_{\max}\left(\mathbb{E}\left[F_{k-1}^\top M_{k-1}^{-1} F_{k-1} \right] \right) \le \epsilon \left(\log \frac{\det \left(\epsilon M_k \right)}{\det \left(\epsilon M_{k-1} \right)} \right).$$

Since $F_k^\top \left(M_k \right)^{-1} F_k$ is positive semidefinite, its trace is larger than its largest eigenvalue. Hence,
adding up for $k = 0, 1, \cdots, n$, by interchanging trace and expectation, we obtain

$$\mathbb{E}\left[\boldsymbol{\lambda}_{\max}\left(F_n^\top \left(M_n \right)^{-1} F_n \right) \right] \le \mathbb{E}\left[\mathbf{tr}\left(F_n^\top \left(M_n \right)^{-1} F_n \right) \right] \le d_W \boldsymbol{\lambda}_{\max}\left(\mathbb{E}\left[F_n^\top \left(M_n \right)^{-1} F_n \right] \right),$$

which, by $\epsilon M_0 \ge I$, leads to

$$\mathbb{E}\left[\boldsymbol{\lambda}_{\max}\left(F_n^\top \left(\epsilon M_n \right)^{-1} F_n \right) \right] \le m d_W \log \boldsymbol{\lambda}_{\max}\left(\epsilon M_n \right).$$

752 Thus, according to Doob's Martingale Convergence Theorem [32, 4], we have

$$\left\| \left(\epsilon M_n \right)^{-1/2} F_n \right\|^2 = \mathcal{O}\left(m d_W \log \boldsymbol{\lambda}_{\max}\left(\epsilon M_n \right) \right).$$

753 Finally, letting $\epsilon \to 0$, we obtain the desired result, because $\epsilon M_n, F_n$ are ϵ-approximations of the
754 corresponding integrals. ∎

G.3 Lipschitz continuity of optimal feedback

756 **Lemma 3** *Using the Jordan decomposition $D_\star = A_\star + B_\star \mathcal{L}\left(A_\star, B_\star \right) = P_\star^{-1} \Lambda_\star P_\star$, define $\boldsymbol{\mu}_\star =$*
757 *$\boldsymbol{\mu}_{D_\star}$, similar to Definition 2, and suppose that $\mathcal{E}\left(A, B \right) \le \kappa_\star$, for*

$$\kappa_\star = \frac{1}{1 \vee \|\mathcal{L}\left(A_\star, B_\star \right)\|} \left(\frac{\left(-\overline{\boldsymbol{\lambda}}\left(D_\star \right) \right) \wedge \left(-\overline{\boldsymbol{\lambda}}\left(D_\star \right) \right)^{\boldsymbol{\mu}_\star}}{\boldsymbol{\mu}_\star^{1/2} \|\| P_\star^{-1} \|\| \|\| P_\star \|\|} \wedge \left[4 \int_0^\infty \|\| e^{D_\star t} \|\|^2 \mathrm{d}t \right]^{-1} \right).$$

758 *Then, letting*

$$\beta_\star = \frac{2\|\mathcal{K}\left(A_\star, B_\star\right)\|}{\lambda_{\min}\left(R\right)}\left[1 + \frac{4\|B_\star\|}{\lambda_{\min}\left(Q\right)}\|\mathcal{K}\left(A_\star, B_\star\right)\|\left(1 \vee \frac{2\left(\|B_\star\| + \kappa_\star\right)\|\mathcal{K}\left(A_\star, B_\star\right)\|}{\lambda_{\min}\left(R\right)}\right)\right],$$

759 *we have*

$$\|\mathcal{L}\left(A, B\right) - \mathcal{L}\left(A_\star, B_\star\right)\| \leq \beta_\star \mathcal{E}\left(A, B\right).$$

760 *In general, without the condition $\mathcal{E}\left(A, B\right) \leq \kappa_\star$, the constant $\beta_\star$ is replaced with*

$$\beta = \frac{\|\mathcal{K}\left(A, B\right)\|}{\lambda_{\min}\left(R\right)} + \frac{2\|B_\star\|\|\mathcal{K}\left(A_0, B_0\right)\|^2}{\lambda_{\min}\left(Q\right)\lambda_{\min}\left(R\right)}\left(1 \vee \frac{\left(\|B_\star\| + \mathcal{E}\left(A, B\right)\right)\|\mathcal{K}\left(A_0, B_0\right)\|}{\lambda_{\min}\left(R\right)}\right),$$

761 *for some convex combination $[A_0, B_0] = \eta\left[A, B\right] + (1 - \eta)\left[A_\star, B_\star\right]$, and $0 \leq \eta \leq 1$.*

Proof. Fix the matrices A, B, and consider the matrix-valued curve

$$\varphi = \left\{(1 - \eta)\left[A_\star, B_\star\right] + \eta\left[A, B\right]\right\}_{0 \leq \eta \leq 1}.$$

762 For an arbitrary $A_0, B_0 \in \varphi$, we find the derivative of the matrix $\mathcal{K}\left(A_0, B_0\right)$ at A_0, B_0, assuming
763 that the matrices A_0, B_0 vary along φ. For this purpose, letting $\Delta_A = A - A_\star$, $\Delta_B = B - B_\star$, we
764 first calculate $\mathcal{K}\left(A_1, B_1\right)$ for $A_1 = A_0 + \eta\Delta_A$, $B_1 = B_0 + \eta\Delta_B$, and then let $\eta \to 0$. First, letting
765 $P = \mathcal{K}\left(A_1, B_1\right) - \mathcal{K}\left(A_0, B_0\right)$, we get

$$
\begin{aligned}
&\mathcal{K}\left(A_0, B_0\right) B_1 R^{-1} B_1^\top \mathcal{K}\left(A_0, B_0\right) \\
={} & \eta\mathcal{K}\left(A_0, B_0\right) \Delta_B R^{-1} B_1^\top \mathcal{K}\left(A_0, B_0\right) + \mathcal{K}\left(A_0, B_0\right) B_0 R^{-1} B_1^\top \mathcal{K}\left(A_0, B_0\right) \\
={} & \eta^2\mathcal{K}\left(A_0, B_0\right) \Delta_B R^{-1}\Delta_B^\top \mathcal{K}\left(A_0, B_0\right) + \eta\mathcal{K}\left(A_0, B_0\right) \Delta_B R^{-1} B_0^\top \mathcal{K}\left(A_0, B_0\right) \\
+{} & \eta\mathcal{K}\left(A_0, B_0\right) B_0 R^{-1}\Delta_B^\top \mathcal{K}\left(A_0, B_0\right) \\
+{} & \mathcal{K}\left(A_0, B_0\right) B_0 R^{-1} B_0^\top \mathcal{K}\left(A_0, B_0\right).
\end{aligned}
$$

766 The above expression, because of

$$
\begin{aligned}
&\mathcal{K}\left(A_1, B_1\right) B_1 R^{-1} B_1^\top \mathcal{K}\left(A_1, B_1\right) \\
={} & \mathcal{K}\left(A_1, B_1\right) B_1 R^{-1} B_1^\top P + \mathcal{K}\left(A_1, B_1\right) B_1 R^{-1} B_1^\top \mathcal{K}\left(A_0, B_0\right) \\
={} & P B_1 R^{-1} B_1^\top P + \mathcal{K}\left(A_0, B_0\right) B_1 R^{-1} B_1^\top P \\
+{} & P B_1 R^{-1} B_1^\top \mathcal{K}\left(A_0, B_0\right) + \mathcal{K}\left(A_0, B_0\right) B_1 R^{-1} B_1^\top \mathcal{K}\left(A_0, B_0\right),
\end{aligned}
$$

767 implies that the followings hold true:

$$
\begin{aligned}
&\mathcal{K}\left(A_1, B_1\right) B_1 R^{-1} B_1^\top \mathcal{K}\left(A_1, B_1\right) - \mathcal{K}\left(A_0, B_0\right) B_0 R^{-1} B_0^\top \mathcal{K}\left(A_0, B_0\right) \\
={} & P B_1 R^{-1} B_1^\top P + \mathcal{K}\left(A_0, B_0\right) B_1 R^{-1} B_1^\top P + P B_1 R^{-1} B_1^\top \mathcal{K}\left(A_0, B_0\right) \\
+{} & \eta^2\mathcal{K}\left(A_0, B_0\right) \Delta_B R^{-1}\Delta_B^\top \mathcal{K}\left(A_0, B_0\right) + \eta\mathcal{K}\left(A_0, B_0\right) \Delta_B R^{-1} B_0^\top \mathcal{K}\left(A_0, B_0\right) \\
+{} & \eta\mathcal{K}\left(A_0, B_0\right) B_0 R^{-1}\Delta_B^\top \mathcal{K}\left(A_0, B_0\right). \tag{47}
\end{aligned}
$$

768 By plugging (47) and

$$
\begin{aligned}
&A_1^\top \mathcal{K}\left(A_1, B_1\right) + \mathcal{K}\left(A_1, B_1\right) A_1 = A_1^\top \mathcal{K}\left(A_0, B_0\right) + A_1^\top P + \mathcal{K}\left(A_0, B_0\right) A_1 + P A_1 \\
={} & A_0^\top \mathcal{K}\left(A_0, B_0\right) + \eta\Delta_A^\top \mathcal{K}\left(A_0, B_0\right) + A_1^\top P + \mathcal{K}\left(A_0, B_0\right) A_0 + \eta\mathcal{K}\left(A_0, B_0\right) \Delta_A + P A_1,
\end{aligned}
$$

769 in $\Phi_{A_i, B_i}\left(\mathcal{K}\left(A_i, B_i\right)\right) = 0$ for $i = 0, 1$, we obtain

$$
\begin{aligned}
0 ={} & \left[A_1^\top - \mathcal{K}\left(A_0, B_0\right) B_1 R^{-1} B_1^\top\right] P + P\left[A_1 - B_1 R^{-1} B_1^\top \mathcal{K}\left(A_0, B_0\right)\right] - P B_1 R^{-1} B_1^\top P \\
+{} & \eta\Delta_A^\top \mathcal{K}\left(A_0, B_0\right) + \eta\mathcal{K}\left(A_0, B_0\right) \Delta_A - \eta^2\mathcal{K}\left(A_0, B_0\right) \Delta_B R^{-1}\Delta_B^\top \mathcal{K}\left(A_0, B_0\right) \\
-{} & \eta\mathcal{K}\left(A_0, B_0\right) \Delta_B R^{-1} B_0^\top \mathcal{K}\left(A_0, B_0\right) - \eta\mathcal{K}\left(A_0, B_0\right) B_0 R^{-1}\Delta_B^\top \mathcal{K}\left(A_0, B_0\right),
\end{aligned}
$$

770 or equivalently,

$$0 = \widetilde{A}^\top P + P\widetilde{A} - P B_1 R^{-1} B_1^\top P + \widetilde{Q}, \tag{48}$$

for $\widetilde{A} = A_1 - B_1 R^{-1} B_1^\top \mathcal{K}(A_0, B_0)$, and

$$
\begin{aligned}
\widetilde{Q} &= \eta \Delta_A^\top \mathcal{K}(A_0, B_0) + \eta \mathcal{K}(A_0, B_0) \Delta_A - \eta^2 \mathcal{K}(A_0, B_0) \Delta_B R^{-1} \Delta_B^\top \mathcal{K}(A_0, B_0) \\
&= \eta \mathcal{K}(A_0, B_0) \left[\Delta_A + \Delta_B \mathcal{L}(A_0, B_0) \right] + \eta \left[\mathcal{L}(A_0, B_0)^\top \Delta_B^\top + \Delta_A^\top \right] \mathcal{K}(A_0, B_0) \\
&\quad - \eta^2 \mathcal{K}(A_0, B_0) \Delta_B R^{-1} \Delta_B^\top \mathcal{K}(A_0, B_0).
\end{aligned}
$$

Suppose that η is sufficiently small so that $\overline{\boldsymbol{\lambda}}\left(\widetilde{A}\right) < 0$. Note that it is possible thanks to stabilizability of A_0, B_0, Theorem 1, and $\lim_{\eta \to 0} \widetilde{A} = A_0 + B_0 \mathcal{L}(A_0, B_0) = D_0$. So, since $PB_1 R^{-1} B_1^\top P$ is a positive semidefinite matrix, (48) implies that

$$
\begin{aligned}
P &= \int_0^\infty e^{\widetilde{A}^\top t} \left(-PB_1 R^{-1} B_1^\top P + \widetilde{Q} \right) e^{\widetilde{A} t} \mathrm{d}t \\
&\leq \int_0^\infty e^{\widetilde{A}^\top t} \widetilde{Q} e^{\widetilde{A} t} \mathrm{d}t \leq \left(\left\| \widetilde{Q} \right\| \int_0^\infty \left\| e^{\widetilde{A} t} \right\|^2 \mathrm{d}t \right) I_{d_X}.
\end{aligned}
$$

This, because of $\lim_{\eta \to 0} \widetilde{Q} = 0$, leads to $\lim_{\eta \to 0} P = 0$. Thus, letting $M = \Delta_A + \Delta_B \mathcal{L}(A_0, B_0)$ and $\Delta_{\mathcal{K}(A_0, B_0)} = \lim_{\eta \to 0} \eta^{-1} P$, (48) gives the following for $\Delta_{\mathcal{K}(A_0, B_0)}$:

$$
\int_0^\infty e^{D_0^\top t} \left(\mathcal{K}(A_0, B_0) M + M^\top \mathcal{K}(A_0, B_0) \right) e^{D_0 t} \mathrm{d}t. \tag{49}
$$

By

$$
\mathcal{K}(A, B) - \mathcal{K}(A_\star, B_\star) = \int_0^1 \Delta_{(1-\eta)[A_\star, B_\star] + \eta[A, B]} \mathrm{d}\eta,
$$

(31), (49), and the Cauchy-Schwarz inequality provide

$$
\begin{aligned}
&\left\| \mathcal{K}(A, B) - \mathcal{K}(A_\star, B_\star) \right\| \\
&\leq \ \mathcal{E}(A, B) \sup_{[A_0, B_0] \in \varphi} 2 \left\| \mathcal{K}(A_0, B_0) \right\| \left(1 \vee \left\| \mathcal{L}(A_0, B_0) \right\| \right) \int_0^\infty \left\| e^{D_0 t} \right\|^2 \mathrm{d}t \\
&\leq \ \mathcal{E}(A, B) \frac{2}{\boldsymbol{\lambda}_{\min}(Q)} \sup_{[A_0, B_0] \in \varphi} \left\| \mathcal{K}(A_0, B_0) \right\|^2 \left(1 \vee \left\| \mathcal{L}(A_0, B_0) \right\| \right).
\end{aligned}
$$

Next, note that $\mathcal{E}(A, B) \leq \kappa_\star$, together with (29) and (30), implies that

$$
\left\| \mathcal{K}(A, B) - \mathcal{K}(A_\star, B_\star) \right\| \leq \mathcal{E}(A, B) \frac{8 \left\| \mathcal{K}(A_\star, B_\star) \right\|^2}{\boldsymbol{\lambda}_{\min}(Q)} \left(1 \vee \frac{2 \left(\|B_\star\| + \kappa_\star \right) \left\| \mathcal{K}(A_\star, B_\star) \right\|}{\boldsymbol{\lambda}_{\min}(R)} \right).
$$

Therefore, using (30), and putting the above inequality together with

$$
\begin{aligned}
&\left\| \mathcal{L}(A, B) - \mathcal{L}(A_\star, B_\star) \right\| \\
&= \ \left\| R^{-1} \left[(B_\star - B_1) \mathcal{K}(A, B) + B_\star (\mathcal{K}(A_\star, B_\star) - \mathcal{K}(A, B)) \right] \right\| \\
&\leq \ \left\| R^{-1} \right\| \left[\|B_\star - B_1\| \left\| \mathcal{K}(A, B) \right\| + \|B_\star\| \left\| \mathcal{K}(A_\star, B_\star) - \mathcal{K}(A, B) \right\| \right],
\end{aligned}
$$

we get the first desired result. To establish the second result, it suffices to let A_0, B_0 be the one for which the above supremum over φ is achieved. ∎

 G.4 Effects of sub-optimal linear feedback policies

783 **Lemma 4** *Consider a noiseless linear dynamical system with the stabilizable dynamics matrices*
784 A, B. *That is,* $\mathrm{d}X_t = (AX_t + BU_t)\,\mathrm{d}t$, *starting from* $X_0 = x$. *Then, if we apply the linear feedback*
785 $U_t = LX_t$, *as long as* $\overline{\lambda}(A + BL) < 0$, *it holds that*

$$
\int_0^\infty c\,(X_t, U_t)\mathrm{d}t = x^\top \mathcal{K}(A, B)\, x + \int_0^\infty \left\| R^{1/2}\left(L - \mathcal{L}(A, B)\right) e^{(A+BL)t}x \right\|^2 \mathrm{d}t.
$$

786 Proof. Denote $D_1 = A + B\mathcal{L}(A, B)$ and $D_2 = A + BL$. So, the dynamics equation $\mathrm{d}X_t =$
787 $(AX_t + BLX_t)\,\mathrm{d}t$ implies that $X_t = e^{D_2 t}x$, which leads to

$$
\int_0^\infty c\,(X_t, U_t)\mathrm{d}t = \int_0^\infty X_t^\top \left(Q + L^\top RL\right) X_t \mathrm{d}t
$$

$$
= \int_0^\infty x^\top e^{D_2^\top t}\left(Q + L^\top RL\right) e^{D_2 t}x\mathrm{d}t = x^\top Px,
$$

788 where

$$
P = \int_0^\infty e^{D_2^\top t}\left(Q + L^\top RL\right) e^{D_2 t}\mathrm{d}t
$$

$$
= \int_0^\epsilon e^{D_2^\top t}\left(Q + L^\top RL\right) e^{D_2 t}\mathrm{d}t
$$

$$
+ \; e^{D_2^\top \epsilon}\left(\int_0^\infty e^{D_2^\top t}\left(Q + L^\top RL\right) e^{D_2 t}\mathrm{d}t\right) e^{D_2 \epsilon}
$$

$$
= \int_0^\epsilon e^{D_2^\top t}\left(Q + L^\top RL\right) e^{D_2 t}\mathrm{d}t + e^{D_2^\top \epsilon}Pe^{D_2 \epsilon},
$$

789 which yields to

$$
Q + L^\top RL = \lim_{\epsilon \to 0}\frac{1}{\epsilon}\int_0^\epsilon e^{D_2^\top t}\left(Q + L^\top RL\right) e^{D_2 t}\mathrm{d}t
$$

$$
= \lim_{\epsilon \to 0}\frac{1}{\epsilon}\left[P - e^{D_2^\top \epsilon}P + e^{D_2^\top \epsilon}P - e^{D_2^\top \epsilon}Pe^{D_2 \epsilon}\right]
$$

$$
= -D_2^\top P - PD_2.
$$

790 Similar to (23), it holds that $D_1^\top \mathcal{K}(A, B) + \mathcal{K}(A, B)D_1 + Q + \mathcal{L}(A, B)^\top R\mathcal{L}(A, B)$. So, sub-
791 tracting the latter two equalities, we get

$$
(D_2 - D_1)^\top \mathcal{K}(A, B) + \mathcal{K}(A, B)(D_2 - D_1) \tag{50}
$$
$$
+ \; D_2^\top (P - \mathcal{K}(A, B)) + (P - \mathcal{K}(A, B))D_2 + S = 0,
$$

792 where
$$
S = L^\top RL - \mathcal{L}(A, B)^\top R\mathcal{L}(A, B).
$$

793 Because $\overline{\lambda}(D_2) < 0$, solving (50) for $P - \mathcal{K}(A, B)$, and using the fact $D_2 - D_1 = B\left[L - \mathcal{L}(A, B)\right]$,
794 we have

$$
P - \mathcal{K}(A, B) = \int_0^\infty e^{D_2^\top t}Fe^{D_2 t}\mathrm{d}t,
$$

where

$$F = S + [L - \mathcal{L}(A, B)]^\top B^\top \mathcal{K}(A, B) + \mathcal{K}(A, B) B [L - \mathcal{L}(A, B)].$$

Then, using $B^\top \mathcal{K}(A, B) = -R\mathcal{L}(A, B)$, after doing some algebra we obtain

$$\begin{aligned} S \quad + \quad &[L - \mathcal{L}(A, B)]^\top B^\top \mathcal{K}(A, B) + \mathcal{K}(A, B) B [L - \mathcal{L}(A, B)] \\ = \quad &[L - \mathcal{L}(A, B)]^\top R [L - \mathcal{L}(A, B)]. \end{aligned} \tag{51}$$

Thus, $P - \mathcal{K}(A, B)$ is

$$\int_0^\infty e^{D_2^\top t} [L - \mathcal{L}(A, B)]^\top R [L - \mathcal{L}(A, B)] e^{D_2 t} dt,$$

which implies the desired result. ∎

G.5 Convergence of empirical covariance matrix of the state vectors

Lemma 5 *Suppose that for $t \geq \gamma$, the linear feedback L is applied to the system* (1) *such that $\overline{\boldsymbol{\lambda}}(D) < 0$, where $D = A_\star + B_\star L$. Then, we have*

$$\lim_{T \to \infty} \frac{1}{T} \int_\gamma^{\gamma+T} X_t X_t^\top dt = \int_0^\infty e^{Ds} CC^\top e^{D^\top s} ds.$$

Proof. First, denote

$$V_T = \frac{1}{T} \int_\gamma^{\gamma+T} X_t X_t^\top dt.$$

Then, define the matrix $Y_t = X_t X_t^\top$, and apply Ito's Formula [4] to find dY_t:

$$dY_t = dX_t X_t^\top + X_t dX_t^\top + dX_t dX_t^\top.$$

Plugging in for dX_t from (1), we obtain

$$\begin{aligned} dY_t \quad = \quad &(DX_t dt + CdW_t) X_t^\top \\ + \quad &X_t (DX_t dt + CdW_t)^\top + CC^\top dt, \end{aligned}$$

where we used the facts $dtdt = 0$, $dW_t dt = 0$, and Ito Isometry $dW_t dW_t^\top = dt I_{d_W}$ [4]. Thus, we have

$$Y_{\gamma+T} - Y_\gamma = \int_\gamma^{\gamma+T} dY_t dt \quad = \quad \int_\gamma^{\gamma+T} \left(DX_t X_t^\top + X_t X_t^\top D^\top + CC^\top \right) dt + TM_{\gamma,T},$$

where

$$M_{\gamma,T} = \frac{1}{T} \int_\gamma^{\gamma+T} X_t dW_t^\top C^\top + \frac{1}{T} \left(\int_\gamma^{\gamma+T} X_t dW_t^\top C^\top \right)^\top.$$

This can equivalently be written as

$$\frac{1}{T} \left(X_{\gamma+T} X_{\gamma+T}^\top - X_\gamma X_\gamma^\top \right) \quad = \quad DV_T + V_T D^\top + CC^\top + M_{\gamma,T}.$$

Since $\overline{\boldsymbol{\lambda}}(D) < 0$, the latter equality implies that V_T is

$$\int_0^\infty e^{Ds} \left(CC^\top + M_{\gamma,T} + \frac{1}{T} X_\gamma X_\gamma^\top - \frac{1}{T} X_{\gamma+T} X_{\gamma+T}^\top \right) e^{D^\top s} ds.$$

Now, according to the following statements, the above leads to the desired result, because the terms corresponding to $M_{\gamma,T}, X_\gamma, X_{\gamma+T}$ vanish as T grows.

812   1. Clearly, it holds that $\lim\limits_{T\to\infty} T^{-1/2}\|X_\gamma\| = 0$.

813   2. Since $\overline{\boldsymbol{\lambda}}(D) < 0$, the expression

$$X_{\gamma+T} = e^{DT}X_\gamma + \int_\gamma^{\gamma+T} e^{D(\gamma+T-s)}C\mathrm{d}W_s$$

814   implies that $\lim\limits_{T\to\infty} T^{-1/2}\|X_{\gamma+T}\| = 0$.

815   3. Putting $\overline{\boldsymbol{\lambda}}(D) < 0$ together with Doob's Martingale Convergence Theorem [32, 4], we get
816     $\lim\limits_{T\to\infty} M_{\gamma,T} = 0$.

817   ■

G.6   Manifolds of dynamical systems with equal optimal feedback matrices

818

819 **Lemma 6** *Consider the set of dynamics matrices A, B that share optimal feedback with A_0, B_0:*

$$\mathcal{M}_0 = \left\{[A, B] \in \mathbb{R}^{d_X \times (d_X + d_U)} : \mathcal{L}(A, B) = \mathcal{L}(A_0, B_0)\right\}.$$

820 *Then, $\mathcal{M}_0$ is a manifold of dimension d_X^2.*

821 Proof. Suppose that for the matrix $[A, B] = [A_0, B_0] + \epsilon[M, N]$, it holds that $\mathcal{L}(A, B) = \mathcal{L}(A_0, B_0)$.
822 We find the derivative of $\mathcal{L}(A_0, B_0)$ along the direction $[M, N]$. First, using the expressions in (23)
823 for A, B and for A_0, B_0, we get

$$
\begin{aligned}
&(D_0 + \epsilon M + \epsilon N\mathcal{L}(A_0, B_0))^\top \mathcal{K}(A, B) \\
&+ \quad \mathcal{K}(A, B)(D_0 + \epsilon M + \epsilon N\mathcal{L}(A_0, B_0)) \\
&= \quad -Q - \mathcal{L}(A_0, B_0)^\top R\mathcal{L}(A_0, B_0) \\
&= \quad D_0^\top \mathcal{K}(A_0, B_0) + \mathcal{K}(A_0, B_0)D_0,
\end{aligned}
$$

824 where $D_0 = A_0 + B_0\mathcal{L}(A_0, B_0)$. Simplifying the above expressions and letting $\epsilon \to 0$, for the
825 matrix

$$\Delta = \lim_{\epsilon\to 0}\epsilon^{-1}(\mathcal{K}(A, B) - \mathcal{K}(A_0, B_0)),$$

826 we have

$$
\begin{aligned}
&D_0^\top\Delta + \Delta D_0 + (M + N\mathcal{L}(A_0, B_0))^\top \mathcal{K}(A_0, B_0) \\
&+ \quad \mathcal{K}(A_0, B_0)(M + N\mathcal{L}(A_0, B_0)) = 0.
\end{aligned}
$$

827 Thus, since according to Theorem 1, $\overline{\boldsymbol{\lambda}}(D_0) < 0$, it yields to

$$\Delta = \int_0^\infty e^{D_0^\top t}F e^{D_0 t}\mathrm{d}t,$$

828 where

$$
\begin{aligned}
F &= (M + N\mathcal{L}(A_0, B_0))^\top \mathcal{K}(A_0, B_0) \\
&+ \quad \mathcal{K}(A_0, B_0)(M + N\mathcal{L}(A_0, B_0)).
\end{aligned}
$$

829 On the other hand, $\mathcal{L}(A, B) = -R^{-1}B^\top\mathcal{K}(A, B)$ gives

$$
\begin{aligned}
0 &= \lim_{\epsilon\to 0}\frac{1}{\epsilon}\left(B^\top\mathcal{K}(A, B) - B_0^\top\mathcal{K}(A_0, B_0)\right) \\
&= \lim_{\epsilon\to 0}\frac{1}{\epsilon}\left[\left(B^\top - B_0^\top\right)\mathcal{K}(A, B)\right. \\
&\quad \left. - B_0^\top(\mathcal{K}(A_0, B_0) - \mathcal{K}(A, B))\right] \\
&= N^\top\mathcal{K}(A_0, B_0) + B_0^\top\Delta.
\end{aligned}
$$

830 So, $\mathcal{M}_0$ is a manifold, and its tangent space consists of matrices M, N satisfying the above equation.
831 To find the dimension, select a $d_X \times d_X$ matrix P arbitrarily, and let N be

$$N = -\mathcal{K}\left(A_0, B_0\right)^{-1} \int_0^\infty e^{D_0^\top t} \left[P^\top \mathcal{K}\left(A_0, B_0\right) + \mathcal{K}\left(A_0, B_0\right) P\right] e^{D_0 t} B_0 \mathrm{d}t = 0. \tag{52}$$

832 Note that since $\boldsymbol{\lambda}_{\min}\left(Q\right) > 0$, the inverse $\mathcal{K}\left(A_0, B_0\right)^{-1}$ exists. Then, solve for M according to
833 $M + N\mathcal{L}\left(A_0, B_0\right) = P$. Therefore, the matrices M, N satisfy in $N^\top \mathcal{K}\left(A_0, B_0\right) + B_0^\top \Delta = 0$, and
834 so correspond to a member of $\mathcal{M}_0$. Conversely, every matrices M, N in the tangent space of $\mathcal{M}_0$
835 provide a $d_X \times d_X$ matrix $P = M + N\mathcal{L}\left(A_0, B_0\right)$ such that $N^\top \mathcal{K}\left(A_0, B_0\right) + B_0^\top \Delta = 0$. Thus,
836 $\mathcal{M}_0$ is of dimension d_X^2, which is the desired result. ∎