A single gradient step finds adversarial examples on random two-layers neural networks

Abstract

Daniely and Schacham [2020] recently showed that gradient descent finds adversarial examples on random undercomplete two-layers ReLU neural networks. The term "undercomplete" refers to the fact that their proof only holds when the number of neurons is a vanishing fraction of the ambient dimension. We extend their result to the overcomplete case, where the number of neurons is larger than the dimension (yet also subexponential in the dimension). In fact we prove that a single step of gradient descent suffices. We also show this result for any subexponential width random neural network with smooth activation function.

1 Introduction

We study the random two-layers neural network model, $f : \mathbb{R}^d \to \mathbb{R}$ defined by

$$f(x) = \frac{1}{\sqrt{k}} \sum_{\ell=1}^{k} a_\ell \psi(w_\ell \cdot x) \,, \tag{1}$$

where $\psi : \mathbb{R} \to \mathbb{R}$ is a fixed non-linearity, the weight vectors $w_\ell \in \mathbb{R}^d$ are i.i.d. from a Gaussian $\mathcal{N}\left(0, \frac{1}{d}\mathrm{I}_d\right)$ (so they are roughly unit norm vectors), and the coefficients $a_\ell \in \mathbb{R}$ are independent from the weight vectors and i.i.d. uniformly distributed in $\{-1, +1\}$. With this parametrization, the central limit theorem says that, for $x \in \sqrt{d} \cdot \mathbb{S}^{d-1}$ (so that $w_\ell \cdot x \sim \mathcal{N}(0,1)$) and large width k, the distribution of $f(x)$ is approximately a centered Gaussian with variance $\mathbb{E}_{X \sim \mathcal{N}(0,1)}[\psi(X)^2]$.

Our goal is to study the concept of *adversarial examples* in this random model. We say that $\delta \in \mathbb{R}^d$ is an *adversarial perturbation* at $x \in \mathbb{R}^d$ if $\|\delta\| \ll \|x\|$ and $\mathrm{sign}(f(x)) \neq \mathrm{sign}(f(x + \delta))$. In this case, we call $x + \delta$ an *adversarial example*. Our main result is that, while $|f(x)| \approx 1$ with high probability, a *single* gradient step on f (i.e., a perturbation of the form $\delta = \eta \nabla f(x)$ for some $\eta \in \mathbb{R}$) suffices to find such adversarial examples, with roughly $\|\delta\| \simeq \frac{\|x\|}{\sqrt{d}} = 1$. We prove this statement for networks of subexponential width (e.g., $k \ll \exp(o(d))$) with both smooth and ReLU activation functions. We first state our result for smooth activation functions in the following theorem.

Theorem 1. *Let $\gamma \in (0, 1)$ and ψ be non-constant, Lipschitz and with Lipschtiz derivative. There exists constants C_1, C_2, C_3, C_4 depending on ψ such that the following holds true. Assume $k \geq C_1 \log^3(1/\gamma)$ and $d \geq C_2 \log(k/\gamma) \log(1/\gamma)$, and let $\eta \in \mathbb{R}$ such that $|\eta| = C_3 \frac{\sqrt{\log(1/\gamma)}}{\|\nabla f(x)\|^2}$ and $\mathrm{sign}(\eta) = -\mathrm{sign}(f(x))$. Then, with probability at least $1 - \gamma$, one has:*

$$\mathrm{sign}(f(x)) \neq \mathrm{sign}(f(x + \eta \nabla f(x))) \,.$$

Moreover we have $\|\eta \nabla f(x)\| \leq C_4 \sqrt{\log(1/\gamma)}$.

Submitted to 35th Conference on Neural Information Processing Systems (NeurIPS 2021). Do not distribute.

29 Note that our proof of Theorem 1 in Section 2 easily gives explicit values for C_1, C_2, C_3, C_4. Also
30 note that the subexponential width condition in the above Theorem is of the form $k \ll \exp(o(d))$.

31 Our second main result establishes similar behavior for the *non-smooth* ReLU activation unit.

32 **Theorem 2.** *Let $\gamma \in (0,1)$ and $\psi(t) = \max(0,t)$. There exist constants C_1, C_2, C_3, C_4, C_5 such*
33 *that the following holds true. Assume*

$$C_1 \log^6(d) \log(1/\gamma) \leq k, \quad C_2 \log^3(d) \log(1/\gamma) \leq d, \quad C_3 \log^4(k) \log(1/\gamma) \leq \frac{d}{\log(d)}$$

34 *and let $\eta \in \mathbb{R}$ such that $|\eta| = C_4 \frac{\sqrt{\log(1/\gamma)}}{\|\nabla f(x)\|^2}$ and $\mathrm{sign}(\eta) = -\mathrm{sign}(f(x))$. Then, with probability at*
35 *least $1 - \gamma$, one has:*

$$\mathrm{sign}(f(x)) \neq \mathrm{sign}(f(x + \eta \nabla f(x))).$$

36 *Moreover, we have $\|\eta \nabla f(x)\| \leq C_5 \sqrt{\log(1/\gamma)}$.*

37 Note that the subexponential condition on the width in the above Theorem is of the form $k \ll$
38 $\exp(d^{0.24})$. In fact by modifying a bit the proof we can get a condition of the form $k \ll \exp(d^\rho)$ for
39 any $\rho < 1/2$, but for the sake of clarity we only prove the weaker version stated above. The proof for
40 the ReLU activation is broken into two separate cases focusing on the overlapping regimes:

Case 1: $k \gtrsim d \log^2(d)$ and **Case 2:** $\log^6(d) \log(1/\gamma) \lesssim k \lesssim d \log^3 d$,

41 as the proofs for the two settings use distinct arguments. The proof for the first regime is similar to the
42 proof for Theorem 1 while the second uses a refinement of an argument by Daniely and Schacham.
43 These arguments are carried out in Section 3.

1.1 Related works

45 The existence of adversarial examples in neural network architectures was first evidenced in the
46 seminal paper of Szegedy et al. [2014], where the authors found adversarial examples by using
47 the L-BFGS optimization procedure. Shortly after this work, it was hypothesized in Goodfellow
48 et al. [2015] that the existence of adversarial examples stems from an excessive "linearity" of neural
49 network models. This hypothesis was experimentally confirmed by showing that a *single step* of
50 gradient descent suffices to find adversarial perturbations (the so-called *fast gradient sign method*
51 -FGSM). Our theorems can be thought of as a theoretical confirmation of the hypothesis in Goodfellow
52 et al. [2015]. In fact, as explained in Section 1.2, our proofs proceed exactly by showing that "most"
53 two-layers neural networks behave "mostly" linearly over "vast" regions of input space.

54 We note that not *all* networks are susceptible to one-step gradient attacks to find adversarial examples.
55 Indeed, in Goodfellow et al. [2015], it was shown that adversarial training can be used to build
56 networks that are somewhat robust to one-step gradient attacks. Interestingly, Madry et al. [2018]
57 showed that such models remain susceptible to *multi-steps* gradient attacks, and empirically demon-
58 strated that better robustness can be achieved with adversarial training using multi-steps gradient
59 attacks. Understanding this phenomenon theoretically remains a challenge, see for example Allen-
60 Zhu and Li [2020] for a proposed approach, and Moosavi-Dezfooli et al. [2019], Qin et al. [2019] for
61 discussion/algorithmic consequences of the relation with the phenomenon of *gradient obfuscation*
62 (Papernot et al. [2017], Athalye et al. [2018]).

63 Our work is a direct follow-up of Daniely and Schacham [2020] (which itself follows Shamir et al.
64 [2019]). Daniely and Schacham prove that multi-steps gradient descent finds adversarial examples
65 for ReLU random networks of the form (1), as long as the number of neurons is much smaller than
66 the dimension (i.e., $k = o(d)$). They explicitly conjecture that this condition is not necessary, and
67 indeed we exponentially improve it to $k = \exp(o(d))$ in Theorem 2 (see below for a discussion of k
68 exponential in the dimension). We note that Daniely and Schacham went beyond two-layers neural
69 networks, and conjectured (and proved for shrinking layers) that gradient descent finds adversarial
70 examples on random multi-layers neural networks. We give some experimental confirmation of this
71 multi-layer conjecture in Section 4.

The ultra-wide case $k = \exp(\Omega(d))$ remains open. This exponential size case seems of a different nature than the polynomial size we tackle here, at least for the ReLU activation function. In particular, it is likely that the behavior with exponential width would be closely tied to the actual limit case $k = +\infty$, where the random model (1) yields a Gaussian process. Namely, for $k = +\infty$, f is a Gaussian process indexed by the sphere (say if we restrict to inputs $x \in \sqrt{d} \cdot \mathbb{S}^{d-1}$), with $f(x) \sim \mathcal{N}(0, \mathbb{E}_{X \sim \mathcal{N}(0,1)}[\psi(X)])$ and $\mathbb{E}[f(x)f(y)] = \mathbb{E}_{X,Y \sim \mathcal{N}(0,1):\mathbb{E}[XY]=x \cdot y}[\psi(X)\psi(Y)]$. For example if the activation function is a Hermite polynomial of degree p, then f would be a spherical p-spin glass model. This polynomial case is particularly well-understood, and in fact the landscape we describe below in Section 1.2 was already described in this case by Ben Arous et al. [2020] (in particular Corollary 59). It would be interesting to see if the p-spin glass landscape literature can be extended to non-polynomial activation functions, and to a finite (but possibly exponential in d) k. A step in this latter direction was recently taken in Eldan et al. [2021], where convergence rates to the Gaussian process limit where given both for polynomial activations and for the ReLU. Finally we note that for a smooth activation it might be that there is a more direct argument to remove the subexponential width condition in Theorem 1 (in technical terms, the proof of Proposition 2 could leverage a better argument than our naive upper bound on Lip(Φ)).

Finally, we note that, in practice, it has been found that there exists "universal" adversarial perturbations that generalize across both inputs and neural networks, Moosavi-Dezfooli et al. [2017]. For the case of ReLU activation (Theorem 2), we could in fact prove our result by replacing the gradient step with a step in the direction $\sum_{\ell=1}^{k} a_\ell w_\ell$, which is indeed a direction *independent* of the input x, thus proving the existence of "universal" perturbations (generalizing across inputs) for our model.

1.2 The landscape of random two-layers neural networks

For a smooth non-linearity ψ, we have

$$\nabla f(x) = \frac{1}{\sqrt{k}} \sum_{\ell=1}^{k} a_\ell w_\ell \psi'(w_\ell \cdot x) \qquad \text{and} \qquad \nabla^2 f(x) = \frac{1}{\sqrt{k}} \sum_{\ell=1}^{k} a_\ell w_\ell w_\ell^\top \psi''(w_\ell \cdot x) \,.$$

We already claimed in the introduction that, with high probability the value of $f(x)$ was bounded:

$$\textbf{Upper-Bounded Value}: \qquad |f(x)| = O(1) \,. \tag{2}$$

We alluded to the CLT for this claim, but it is also easy to guess it intuitively by noting that:

$$\mathbb{E}[f(x)^2] = \mathbb{E}\left[\frac{1}{k} \sum_{\ell,\ell'=1}^{k} a_\ell a_{\ell'} \psi(w_\ell \cdot x)\psi(w_{\ell'} \cdot x) \right] = \mathbb{E}_{X \sim \mathcal{N}(0,1)}[\psi(X)^2] \,,$$

as $\mathbb{E}[a_\ell a_{\ell'}] = \mathbb{1}\{\ell = \ell'\}$. The formal proof of (2) (and all other claims we make here) will eventually be a simple application of the classical Bernstein concentration inequality. Similarly, it is easy to see that (noticing that $\mathbb{E}[\|\nabla f(x)\|^2] = \mathbb{E}_{X \sim \mathcal{N}(0,1)}[\psi'(X)^2]$), with high probability, the norm of the gradient of $f(x)$ is bounded below (note that we can also show that it is bounded above).

$$\textbf{Lower-Bounded Gradient}: \qquad \|\nabla f(x)\| = \Omega(1) \,. \tag{3}$$

A slightly more difficult calculation, although classical too, is that

$$\textbf{Upper-Bounded Hessian}: \qquad \|\nabla^2 f(x)\|_{\text{op}} = \widetilde{O}\left(\frac{1}{\sqrt{d}} \right) \,. \tag{4}$$

Indeed one can simply note that, for any $u \in \mathbb{S}^{d-1}$, $u^\top \nabla^2 f(x) u = \frac{1}{\sqrt{k}} \sum_{\ell=1}^{k} a_\ell (w_\ell \cdot u)^2 \psi''(w_\ell \cdot x)$ is approximately distributed as a centered Gaussian with variance

$$\mathbb{E}_{X,Y \sim \mathcal{N}(0,1):\mathbb{E}[XY]=x \cdot u}\left[\left(\frac{X}{\sqrt{d}} \right)^4 \psi''(Y)^2 \right] \,,$$

so that with probability at least $1 - \gamma$ one can expect $u^\top \nabla^2 f(x) u$ to be of order $\frac{\sqrt{\log(1/\gamma)}}{d}$, and thus by taking a union bound over a discretization of the sphere $\mathbb{S}^{d-1}$, one can show inequality (4). In

fact, interestingly, one can even hope that (4) holds true for an entire ball around x: with appropriate smoothness over ψ, this could be obtained by doing another union bound over a second discretization of a d-dimensional ball. In other words, we can expect that with high probability:

$$\forall x \in \mathbb{R}^d : \|x\| = \text{poly}(d), \text{ one has } \|\nabla^2 f(x)\|_{\text{op}} = \widetilde{O}\left(\frac{1}{\sqrt{d}}\right). \tag{5}$$

Equations (2), (3), and (5) paint a rather clear geometric picture. There are essentially two scales around a fixed $x \in \sqrt{d} \cdot \mathbb{S}^{d-1}$: The *macroscopic scale*, where one considers a perturbation $x + \delta$ with $\|\delta\| = \Omega(\sqrt{d})$, and the *mesoscopic scale* where $\|\delta\| = o(\sqrt{d})$ (we use this term because for the ReLU network there will also be a *microscopic scale*, with $\|\delta\| = o(1)$). At the macroscopic scale the landscape of f might be very complicated, but our crucial observation is that the picture at the mesoscopic scale is dramatically simpler. Namely, at the mesoscopic scale, the function f is essentially linear, since one has (thanks to (3) and (5))

Approximate Linearity : $\quad \|\nabla f(x) - \nabla f(x + \delta)\| = o(\|\nabla f(x)\|), \forall \delta : \|\delta\| = o(\sqrt{d})$. (6)

Moreover, since the height of the function is constant (by (2)) and the norm of the gradient is constant, it suffices to step at a constant distance in the direction of the gradient (or negative gradient) to change the sign of f. Assuming without any loss of generality that $f(x) > 0$, we combine (2), (3), and (6) using a standard descent lemma (Lemma 3):

$$f(x - \eta \nabla f(x)) \leq \underbrace{f(x)}_{O(1) \text{ by } (2)} - \eta \underbrace{\|\nabla f(x)\|}_{\Omega(1) \text{ by } (3)} \left(\underbrace{\|\nabla f(x)\|}_{\Omega(1) \text{ by } (3)} - \underbrace{\sup_{\frac{\|\delta\|}{\|\nabla f(x)\|} \leq \eta} \|\nabla f(x) - \nabla f(x + \delta)\|}_{o(1) \text{ by } (4) \text{ or } (6)} \right) \tag{7}$$

$$\leq C_1 - C_2 \eta < 0 \quad (\text{for } C_1/C_2 < \eta < C_3\sqrt{d}) \tag{8}$$

where C_1, C_2 and C_3 are constants that *do not* depend on k and d. In words: a single step of gradient descent (or ascent) with a $O(1)$ step-size suffices to find an adversarial example, and moreover the adversarial perturbation δ satisfies $\|\delta\| = O(1) = O(\|x\|/\sqrt{d})$.

1.3 Proof strategy

The starting point of the proof for both the smooth and ReLU case is to show (2) and (3), which we essentially do below in Section 1.4. In the smooth case, one could then prove formally (4) and conclude as indicated in the last paragraph of Section 1.2. Of course, (4) is simply ill-defined for the ReLU case, so one has to take a different route. Instead, we propose to directly prove (6), that is we study the difference of gradients at the mesoscopic scale. Using that $\|h\| = \sup_{v \in \mathbb{S}^{d-1}} v \cdot h$, we thus need to control (for some $R = o(\sqrt{d})$):

$$\sup_{\delta \in \mathbb{R}^d : \|\delta\| \leq R} \|\nabla f(x) - \nabla f(x + \delta)\| = \sup_{\substack{v \in \mathbb{S}^{d-1}, \\ \delta \in \mathbb{R}^d : \|\delta\| \leq R}} \frac{1}{\sqrt{k}} \sum_{\ell=1}^k a_\ell(w_\ell \cdot v)(\psi'(w_\ell \cdot x) - \psi'(w_\ell \cdot (x + \delta))).$$

$$\tag{9}$$

We execute this strategy first for the smooth case in Section 2. We then prove the ReLU case in Section 3, where we face an extraneous difficulty since the gradient is *not* Lipschitz at very small scale, which introduces a third scale (the *microscopic scale*) that has to be dealt with differently. Technically, this issue appears when we try to move from the discretization over v and δ in (9) to the whole space (a so-called ε-net argument).

1.4 Scaling of value and gradient

Here we show how to prove (2) and (3) (in fact, for our purpose, we only need the one-sided inequality $\|\nabla f(x)\| = \Omega(1)$) under very mild conditions on ψ which will be satisfied for both ReLU and smooth activations. We will repeatedly use Bernstein's inequality which we restate here for convenience (see e.g., Theorem 2.10 in Boucheron et al. [2013]):

Theorem 3 (Bernstein's inequality). *Let (X_ℓ) be i.i.d. centered random variables such that for all integers $q \geq 2$, $\mathbb{E}[|X_\ell|^q] \leq \frac{q!}{2}\sigma^2 c^{q-2}$ for fixed $\sigma, c > 0$. Then, with probability at least $1 - \gamma$:*

$$\sum_{\ell=1}^{k} X_\ell \leq \sqrt{2\sigma^2 k \log(1/\gamma)} + c \log(1/\gamma). \tag{10}$$

We will also use repeatedly that $\mathbb{E}_{X \sim \mathcal{N}(0,1)}[|X|^q] \leq (q-1)!! \leq \frac{q!}{2}$, as well as the following concentration of χ^2 random variables (see e.g., (2.19) in Wainwright [2019]): let $X_1, \ldots, X_k$ be i.i.d. standard Gaussians, then with probability at least $1 - \gamma$, one has:

$$\left|\sum_{\ell=1}^{k} X_\ell^2 - k\right| \leq 4\sqrt{k \log(2/\gamma)}. \tag{11}$$

We can now proceed to our various results.

Lemma 1 (Bounded Value). *Assume that there exists $\sigma, c > 0$ such that for all integers $q \geq 2$, $\mathbb{E}_{X \sim \mathcal{N}(0,1)}[|\psi(X)|^q] \leq \frac{q!}{2}\sigma^2 c^{q-2}$. Then with probability at least $1 - \gamma$ one has*

$$|f(x)| \leq \sqrt{2\sigma^2 \log(1/\gamma)} + \frac{c \log(1/\gamma)}{\sqrt{k}}.$$

Proof sketch. Since, $\frac{1}{\sqrt{k}} \sum_{\ell=1}^{k} a_\ell \psi(w_\ell \cdot x)$, we use Bernstein's inequality on $X_\ell = a_\ell \psi(w_\ell \cdot x)$. $\quad\square$

Lemma 2 (Lower-Bounded Gradient). *Let ψ be differentiable almost everywhere, and assume that there exists $\sigma', c' > 0$ such that for all integers $q \geq 2$, $\mathbb{E}_{X \sim \mathcal{N}(0,1)}[|\psi'(X)|^{2q}] \leq \frac{q!}{2}\sigma'^2 c'^{q-2}$. Then with probability at least $1 - \gamma$,*

$$\|\nabla f(x)\| \geq \left(\mathbb{E}_{X \sim \mathcal{N}(0,1)}[|\psi'(X)|^2] - \left(\sqrt{\frac{2\sigma'^2 \log(2/\gamma)}{k}} + \frac{c' \log(2/\gamma)}{k}\right)\right)^{1/2} \left(1 - 5\sqrt{\frac{\log(4/\gamma)}{d}}\right).$$

Proof sketch. Let $P = \mathrm{I}_d - \frac{xx^\top}{d}$ be the projection on the orthogonal complement of the span of x. We have $\|\nabla f(x)\| \geq \|P\nabla f(x)\| = \|\frac{1}{\sqrt{k}} \sum_{\ell=1}^{k} P a_\ell w_\ell \psi'(w_\ell \cdot x)\|$ where $a_\ell P w_\ell$ is independent of $w_\ell \cdot x$ and is distributed as $\mathcal{N}\left(0, \frac{1}{d}\mathrm{I}_{d-1}\right)$. We conclude by conditioning on the values $(w_\ell \cdot x)_{\ell \in [k]}$ and using the concentration results (10) and (11). $\quad\square$

2 Smooth Non-Linearity (Theorem 1)

In this section, we consider a 1-Lipschitz and L-smooth activation function, that is for all $s, t \in \mathbb{R}$,

$$|\psi(s) - \psi(t)| \leq |s - t| \text{ and } |\psi'(s) - \psi'(t)| \leq L|s - t|. \tag{12}$$

We also assume $\psi(0) = 0$ and denote $c_\psi^2 = \mathbb{E}_{X \sim \mathcal{N}(0,1)}[(\psi'(X))^2]$ which we assume to be non-zero (that is ψ is not a constant function).

Proposition 1 (Upper-Bounded Value and Lower-Bounded Gradient). *Under the above assumptions, one has with probability at least $1 - \gamma$,*

$$|f(x)| \leq \sqrt{2\log\left(\frac{1}{\gamma}\right)}\left(1 + \sqrt{\frac{\log\left(\frac{2}{\gamma}\right)}{k}}\right)$$

$$\text{and} \quad \|\nabla f(x)\| \geq \left(c_\psi^2 - \sqrt{\frac{2\log\left(\frac{4}{\gamma}\right)}{k}\left(1 + \sqrt{\frac{\log\left(\frac{4}{\gamma}\right)}{k}}\right)}\right)^{1/2}\left(1 - 5\sqrt{\frac{\log\left(\frac{8}{\gamma}\right)}{d}}\right).$$

Particularly, there exists $C > 0$ such that for $k \geq C \log(2/\gamma)$ and $d \geq C \log(8/\gamma)$ we have

$$|f(x)| \leq 2\sqrt{\log(1/\gamma)} \qquad \text{and} \qquad \|\nabla f(x)\| \geq c_\psi/2. \tag{13}$$

163 *Proof.* With (12) we have $|\psi(X)| \leq |X|$ and thus in Lemma 1 we can take $\sigma = c = 1$ which yields
164 the first claimed equation. For the second equation we use that $|\psi'(X)| \leq 1$ (since ψ is 1-Lipschitz)
165 and thus, in Lemma 2, we can alsotake $\sigma' = c' = 1$ yielding the second claimed equation. $\qquad\square$

166 Next, we need to control (9) where we use crucially the smoothness of the activation function.

167 **Proposition 2** (Bounded Variations). *Let $R \geq 1$. With probability at least $1 - \gamma$ one has*

$$\sup_{\delta \in \mathbb{R}^d \,:\, \|\delta\| \leq R} \|\nabla f(x) - \nabla f(x + \delta)\| \leq 20 R L \left(\sqrt{\frac{\log(Rk/\gamma)}{d}} + \frac{\log(1/\gamma)}{\sqrt{k}} \right) .$$

168 *Particularly, for any $c > 0$, there exists C_1, C_2 such that if $k \geq C_1 R^2 \log^2(1/\gamma)$ and $d \geq$*
169 *$C_2 R^2 \log(Rk/\gamma)$ then*

$$\sup_{\delta \in \mathbb{R}^d \,:\, \|\delta\| \leq R} \|\nabla f(x) - \nabla f(x + \delta)\| \leq c . \tag{14}$$

170 *Proof sketch.* Our plan is to use the fact that $\|h\| = \sup_{v \in \mathbb{S}^{d-1}} v \cdot h$. Thus, we first show, by using
171 Berstein's inequality on $X_\ell = \frac{a_\ell}{L}(w_\ell \cdot v)(\psi'(w_\ell \cdot x) - \psi'(w_\ell \cdot (x + \delta)))$, that if we fix $\delta \in \mathbb{R}^d$ such
172 that $\|\delta\| \leq R$ and $v \in \mathbb{S}^{d-1}$, then with probability at least $1 - \gamma$ one has:

$$\Phi(v, \delta) := \langle \nabla f(x) - \nabla f(x + \delta), v \rangle = \frac{L}{\sqrt{k}} \sum_{\ell=1}^{k} X_l \leq \frac{4 R L}{d} \sqrt{\log(1/\gamma)} \left(1 + \sqrt{\frac{\log(1/\gamma)}{k}} \right) .$$

173 Let $\Omega := \{(v, \delta) : \|v\| = 1, \|\delta\| \leq R\}$ and N_ε be an ε-net over Ω with $\varepsilon = 1/k$. By a union bound
174 over N_ε (whose size is at most $(10kR)^{2d}$), we obtain with probability at least $1 - \gamma$:

$$\sup_{(v, \delta) \in \Omega} \Phi(v, \delta) \leq \sup_{(v, \delta) \in N_\varepsilon} \Phi(v, \delta) + \sup_{(v, \delta), (v', \delta') \in \Omega : \|v - v'\| + \|\delta - \delta'\| \leq \varepsilon} |\Phi(v, \delta) - \Phi(v', \delta')|$$

$$\leq \frac{4 R L}{d} \sqrt{2 d \log(Rk) + \log(1/\gamma)} \left(1 + \sqrt{\frac{2 d \log(Rk) + \log(1/\gamma)}{k}} \right) + \frac{\mathrm{Lip}(\Phi)}{k} .$$

175 Then we show that the variations of Φ can be upper-bounded by a RL times a χ^2 random variable
176 and use (11) to show that with probability at least $1 - \gamma$,

$$\mathrm{Lip}(\Phi) \leq R L \left(\sqrt{k} + 4 \sqrt{\frac{\log(1/\gamma)}{d}} \right) . \qquad\square$$

177 Finally we can turn to the proof of Theorem 1. Let us first recall this theorem.

178 **Theorem 1.** *Let $\gamma \in (0, 1)$ and ψ be non-constant, Lipschitz and with Lipschtiz derivative.*
179 *There exists constants C_1, C_2, C_3, C_4 depending on ψ such that the following holds true. As-*
180 *sume $k \geq C_1 \log^3(1/\gamma)$ and $d \geq C_2 \log(k/\gamma) \log(1/\gamma)$, and let $\eta \in \mathbb{R}$ such that $|\eta| =$*
181 *$C_3 \sqrt{\log(1/\gamma)} \|\nabla f(x)\|^{-2}$ and $\mathrm{sign}(\eta) = -\mathrm{sign}(f(x))$. Then with probability at least $1 - \gamma$ one*
182 *has:*

$$\mathrm{sign}(f(x)) \neq \mathrm{sign}(f(x + \eta \nabla f(x))) .$$

183 *Moreover we have $\|\eta \nabla f(x)\| \leq C_4 \sqrt{\log(1/\gamma)}$.*

184 *Proof.* We make the following claims which hold with probability at least $1 - \gamma$. Without any loss of
185 generality we can assume that $f(x) > 0$, we will use a standard descent lemma (Lemma 3) and the
186 previous propositions to get that,

$$f(x - \eta \nabla f(x)) \leq \underbrace{f(x)}_{O(1) \text{ by } (13)} - \eta \underbrace{\|\nabla f(x)\|}_{\Omega(1) \text{ by } (13)} \left(\underbrace{\|\nabla f(x)\|}_{\Omega(1) \text{ by } (13)} - \underbrace{\sup_{\frac{\|\delta\|}{\|\nabla f(x)\|} \leq \eta} \|\nabla f(x) - \nabla f(x + \delta)\|}_{o(1) \text{ by } (14)} \right) . \tag{15}$$

187 Formally, let us set $\eta = \frac{32}{c_\psi^2 \|\nabla f(x)\|^2} \sqrt{\log(1/\gamma)}$, $R = \frac{64}{c_\psi^3} \sqrt{\log(1/\gamma)}$, $k \geq C_1 R^2 \log^2(1/\gamma)$ and $d \geq$
188 $C_2 R^2 \log(Rk/\gamma)$ where C_1 and C_2 are large enough such that (13) and (14) are valid with $c = \frac{c_\psi}{4}$.

189 By Proposition 1 we have that $\|\nabla f(x)\| \geq c_\psi/2$ and $\eta\|\nabla f(x)\|^2 = \frac{32}{c_\psi^2}\sqrt{\log(1/\gamma)} \geq \frac{16}{c_\psi^2}|f(x)|$.

190 Moreover, Proposition 2 shows that for all δ such that $\|\delta\| \leq |\eta|\|\nabla f(x)\| \leq \frac{64}{c_\psi^3}\sqrt{\log(1/\gamma)} = R$, we

191 thus have $\|\nabla f(x) - \nabla f(x+\delta)\| \leq c := \frac{c_\psi}{4}$. Consequently,

$$f(x - \eta\nabla f(x)) \leq f(x) - \eta\|\nabla f(x)\|\left(\frac{c_\psi}{2} - \frac{c_\psi}{4}\right) \leq f(x) - \eta\frac{c_\psi^2}{8} \leq -f(x). \tag{16}$$

192 Thus, with a single gradient step of size at most $R = \frac{64}{c_\psi^3}\sqrt{\log(1/\gamma)}$, we switched the sign of f. $\square$

193 3 ReLU Non-Linearity (Theorem 2)

194 In this section, we consider the ReLU non-linearity $\psi(t) = \max(0, t)$. We start by showing that $f(x)$
195 is upper bounded and that $\|\nabla f(x)\|$ is lower-bounded.

196 **Proposition 3** (Upper-Bounded Value and Lower-Bounded Gradient). *With probability at least $1 - \gamma$,*

$$|f(x)| \leq \sqrt{2\log(2/\gamma)}\left(1 + \sqrt{\frac{\log(2/\gamma)}{k}}\right)$$

$$and \quad \|\nabla f(x)\| \geq \left(\frac{1}{2} - \sqrt{\frac{2\log(4/\gamma)}{k}}\left(1 + \sqrt{\frac{\log(1/\gamma)}{k}}\right)\right)^{1/2}\left(1 - 5\sqrt{\frac{\log(4/\gamma)}{d}}\right).$$

197 *Particularly, there exists $C > 0$ such that for $k \geq C\log(1/\gamma)$ and $d \geq C\log(4/\gamma)$ we have*

$$|f(x)| \leq 2\sqrt{\log(2/\gamma)} \qquad and \qquad \|\nabla f(x)\| \geq \frac{1}{4}. \tag{17}$$

198 *Proof.* In Lemma 1 and Lemma 2, we can take $\sigma = c = \sigma' = c' = 1$ (since $|\psi(X)| \leq |X|$ and
199 $|\psi'(X)| \leq 1$), which concludes the proof. $\square$

200 Now we split the control of the gradient variation into two cases: the large width case ($Cd\log^2 d \lesssim k$)
201 and the small width case ($\log^6(d)\log(1/\gamma) \lesssim k \lesssim d\log^3 d$).

202 **Proposition 4.** *(Bounded Variations – Large Width Case) Let $1 \leq R \leq \sqrt{d}/2$, $\sqrt{k} \geq 52$ and*
203 *$d \geq \log(1/\gamma)$. Then, with probability at least $1 - \gamma$, one has*

$$\sup_{\delta\in\mathbb{R}^d : \|\delta\|\leq R} \|\nabla f(x) - \nabla f(x+\delta)\| \leq 20\left(R\log^2(Rk)\sqrt{\frac{\log d}{d}}\right)^{1/4} + 40\sqrt{\frac{d}{k}}\log(Rk).$$

204 *Particularly, for any $c > 0$, there exists C_1, C_2 such that if $C_1 d\log^2(Rk) \leq k$ and $C_2\log^4(Rk)R^2 \leq$*
205 *$\frac{d}{\log(d)}$, then we have,*

$$\sup_{\delta\in\mathbb{R}^d : \|\delta\|\leq R} \|\nabla f(x) - \nabla f(x+\delta)\| \leq c. \tag{18}$$

206 *Proof sketch.* In the smooth case (Proposition 2) we did so by using crucially the smoothness of the
207 activation function. Here, instead of smoothness, we will use that only few activations can change
208 when you make *microscopic move* (i.e., between $x + \delta$ and $x + \delta'$ with $\|\delta - \delta'\| = o(1)$). The key
209 observation that for any δ such that $\|\delta\| \leq R$,

$$\mathbb{P}(\text{sign}(w_\ell \cdot x) \neq \text{sign}(w_\ell \cdot (x+\delta))) \leq R\sqrt{\frac{2\log(d)}{d}} + \frac{1}{d}. \tag{19}$$

210 We can then use this result to bound the variation of $\nabla f(x)$ along a fixed direction. If we fix $\delta \in \mathbb{R}^d$
211 such that $\|\delta\| \leq R$ (with $R \geq 1$) and $v \in \mathbb{S}^{d-1}$, then with probability at least $1 - \gamma$ one has:

$$\Phi(v, \delta) := \langle\nabla f(x) - \nabla f(x+\delta), v\rangle = \frac{1}{\sqrt{k}}\sum_{\ell=1}^{k} X_l \leq 2\sqrt{\frac{\log(\frac{1}{\gamma})}{d}}\left(\left(2R\sqrt{\frac{\log(d)}{d}}\right)^{\frac{1}{4}} + \sqrt{\frac{\log(\frac{1}{\gamma})}{k}}\right).$$

This inequality is proven via Berstein inequality on $X_\ell := a_\ell(w_\ell \cdot v)(\psi'(w_\ell \cdot x) - \psi'(w_\ell \cdot (x + \delta)))$ where we use (19) to prove

$$\mathbb{E}[|X_\ell|^q] \leq \sqrt{\mathbb{E}[|w_\ell \cdot v|^{2q}] \times \mathbb{P}(\text{sign}(w_\ell \cdot x) \neq \text{sign}(w_\ell \cdot (x + \delta)))} \leq \frac{q!}{2}c^{q-2} \times \sigma^2, \qquad (20)$$

with $\sigma = \frac{2}{\sqrt{d}} \times \left(2R\sqrt{\frac{\log(d)}{d}}\right)^{1/4}$ and $c = \frac{2}{\sqrt{d}}$.

Then, similarly as in the proof of Proposition 2, we use a covering argument (but this time with an ε-Net of size $\varepsilon = R^{-4/3}k^{-4}$) to show that,

$$\sup_{(v,\delta)\in\Omega} \Phi(v,\delta) \leq \sup_{(v,\delta)\in N_\varepsilon} \Phi(v,\delta) + \sup_{(v,\delta),(v',\delta')\in\Omega:\|v-v'\|+\|\delta-\delta'\|\leq\varepsilon} |\Phi(v,\delta) - \Phi(v',\delta')|$$

$$\leq 2\sqrt{\frac{10d\log(Rk) + \log(2/\gamma)}{d}} \left(\left(2R\sqrt{\frac{\log(d)}{d}}\right)^{1/4} + \sqrt{\frac{10d\log(Rk) + \log(2/\gamma)}{k}}\right)$$

$$+ \sup_{(v,\delta),(v',\delta')\in\Omega:\|v-v'\|+\|\delta-\delta'\|\leq\varepsilon} |\Phi(v,\delta) - \Phi(v',\delta')|. \qquad (21)$$

By using χ^2 concentration, it is easy to see that with probability at least $1 - \gamma$:

$$|\Phi(\delta, v) - \Phi(\delta, v')| \leq \|v - v'\|\sqrt{k + 4k\sqrt{\frac{\log(k/\gamma)}{d}}}. \qquad (22)$$

By using (19) we are able to show that with probability $1 - \delta$ for any $(v, \delta) \in N_\varepsilon$ there is *at most* $4d$ different activated neurons between $\Phi(v, \delta)$ and $\Phi(v, \delta')$ and thus by applying a concentration result of Lipschitz function of Gaussians we get,

$$|\Phi(\delta, v) - \Phi(\delta', v)| \leq 18\sqrt{\frac{d}{k}}\sqrt{\log 4k + \frac{\log 8/\gamma}{d}}. \qquad (23)$$

Combining (21), (22), and (23), we get

$$\sup_{\delta\in\mathbb{R}^d:\|\delta\|\leq R} \|\nabla f(x) - \nabla f(x + \delta)\| \leq 20\left(R\log^2(Rk)\sqrt{\frac{\log d}{d}}\right)^{1/4} + 40\sqrt{\frac{d}{k}}\log(Rk). \qquad \square$$

Proposition 5 (Bounded Variations – Small Width Case). *Let* $20\log^3 d \cdot \log 1/\gamma \leq k \leq d\log^3 d$ *and* $d \geq 20 \cdot \log^3 d \cdot \log 1/\gamma$. *Then, with probability at least* $1 - \gamma$, *we have:*

$$\sup_{\delta\in\mathbb{R}^d,\|\delta\|\leq\frac{\sqrt{k}}{\log^6 d}} \|\nabla f(x) - \nabla f(x + \delta)\| \leq \frac{60}{\log d}.$$

Proof sketch. In the small width regime we assume $20\log^3 d \cdot \log 1/\delta \leq k \leq d\log^3 d$. The idea of the proof is to control the operator norm of W (where $f(x) = \frac{1}{\sqrt{k}}a^\top\psi(W^\top x)$) as well as the number of neurons with a too small activation. With high probability, we show that we have,

$$\|W\| \leq 3 \cdot \left(1 + \sqrt{\frac{10(k + \log 1/\delta)}{d}}\right), \quad \#\left\{i : |\langle w_i, x\rangle| \leq \frac{1}{\log^3 d}\right\} \leq 2 \cdot \frac{k}{\log^3 d}.$$

Thus, uniformly on $\left\{y : \|x - y\| \leq \frac{\sqrt{k}}{\log^6 d}\right\}$ the cardinal of $T_y := \{i : \text{sgn}\langle w_i, x\rangle \neq \text{sgn}\langle w_i, y\rangle\}$ is uppper-bounded:

$$|T_y| \leq 300k/\log^3 d := B. \qquad (24)$$

We can conclude the proof by using an union bound and a concentration inequality on Gaussians.

$$\|\nabla f(x) - \nabla f(y)\| = \frac{1}{\sqrt{k}}\left\|\sum_{\ell\in T} a_\ell w_\ell\right\| \leq \sup_{\substack{A\subset[k]\\|A|\leq B}} \frac{1}{\sqrt{k}}\left\|\sum_{\ell\in T} a_\ell w_\ell\right\| \leq \sqrt{\frac{B}{k}}\left(1 + \sqrt{\frac{B\log k + \log\frac{4}{\delta}}{d}}\right).$$

$$\square$$

Figure 1: Fraction of inputs with an adversarial example found after a single gradient step, for various input dimensions d, hidden layer widths k and number of hidden layers L. The colored areas represent one standard deviation.

We can now prove our final result.

Theorem 2. *Let $\gamma \in (0,1)$ and $\psi(t) = \max(0,t)$. There exist constants C_1, C_2, C_3, C_4, C_5 such that the following holds true. Assume*

$$C_1 \log^6(d) \log(1/\gamma) \le k, \quad C_2 \log^3(d) \log(1/\gamma) \le d, \quad C_3 \log^4(k) \log(1/\gamma) \le \frac{d}{\log(d)} \,,$$

and let $\eta \in \mathbb{R}$ such that $|\eta| = C_4 \frac{\sqrt{\log(1/\gamma)}}{\|\nabla f(x)\|^2}$ and $\text{sign}(\eta) = -\text{sign}(f(x))$. Then with probability at least $1 - \gamma$ one has:

$$\text{sign}(f(x)) \neq \text{sign}(f(x + \eta \nabla f(x))) \,.$$

Moreover we have $\|\eta \nabla f(x)\| \le C_5 \sqrt{\log(1/\gamma)}$.

Proof. The proof is the same as for Theorem 1 using Proposition 3 instead of Proposition 1 and Proposition 4 or Proposition 5 replacing Proposition 2. $\qquad\square$

4 Experiments

Setting   To verify our theoretical findings, we run some experiments to measure empirically the probability of finding an adversarial example in the direction $\nabla f(x)$. More precisely, we take a random point x of norm $\sqrt{d}$ and initialize a network using the procedure described in Section 1. We then find the smallest η such that a gradient step $\eta \nabla f(x)$ changes the sign of the function. η is of the opposite sign of $f(x)$ and we limit our search to $|\eta| < 20$. We explore various values of d and k, as well as deeper networks with $L = 1$ through $L = 6$ hidden layers. All hidden layers are of width k.

Results   In Figure 1, we show the fraction of examples (out of $10,000$ samples) whose sign is switched. We see that with $L = 1$ and values of d and k larger than 50, 100% of samples are switched. This confirms our theoretical results. We also observe the same statement holds for deeper networks. The values of d and k at which 100% switching is reached appears to grow with L[1]. Additionally, we track the value of the smallest η required to switch the sign of the function, and the gradient norm w.r.t x. Both are approximately constant in d and in k. This finding also holds for deeper networks (see Appendix D for the curves).

---

[1] Due to GPU memory limitations, k could not reach 1,000,000 for deeper networks.

References

Zeyuan Allen-Zhu and Yuanzhi Li. Feature purification: How adversarial training performs robust deep learning. *arXiv preprint arXiv:2005.10190*, 2020.

Anish Athalye, Nicholas Carlini, and David Wagner. Obfuscated gradients give a false sense of security: Circumventing defenses to adversarial examples. In *International Conference on Machine Learning*, 2018.

Gérard Ben Arous, Eliran Subag, and Ofer Zeitouni. Geometry and temperature chaos in mixed spherical spin glasses at low temperature: the perturbative regime. *Communications on Pure and Applied Mathematics*, 73(8):1732–1828, 2020.

Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities: A nonasymptotic theory of independence*. Oxford university press, 2013.

Amit Daniely and Hadas Schacham. Most ReLU networks suffer from ℓ^2 adversarial perturbations. In *NeurIPS*, 2020.

Ronen Eldan, Dan Mikulincer, and Tselil Schramm. Non-asymptotic approximations of neural networks by gaussian processes. *arXiv preprint arXiv:2102.08668*, 2021.

Ian Goodfellow, Jonathon Shlens, and Christian Szegedy. Explaining and harnessing adversarial examples. In *International Conference on Learning Representations*, 2015.

Aleksander Madry, Aleksandar Makelov, Ludwig Schmidt, Dimitris Tsipras, and Adrian Vladu. Towards deep learning models resistant to adversarial attacks. In *International Conference on Learning Representations*, 2018.

Michael Mitzenmacher and Eli Upfal. *Probability and computing: Randomization and probabilistic techniques in algorithms and data analysis*. Cambridge university press, 2017.

Seyed-Mohsen Moosavi-Dezfooli, Alhussein Fawzi, Omar Fawzi, and Pascal Frossard. Universal adversarial perturbations. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pages 1765–1773, 2017.

Seyed-Mohsen Moosavi-Dezfooli, Alhussein Fawzi, Jonathan Uesato, and Pascal Frossard. Robustness via curvature regularization, and vice versa. In *2019 IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*, 2019.

Nicolas Papernot, Patrick McDaniel, Ian Goodfellow, Somesh Jha, Z Berkay Celik, and Ananthram Swami. Practical black-box attacks against machine learning. In *Proceedings of the 2017 ACM on Asia conference on computer and communications security*, 2017.

Chongli Qin, James Martens, Sven Gowal, Dilip Krishnan, Krishnamurthy Dvijotham, Alhussein Fawzi, Soham De, Robert Stanforth, and Pushmeet Kohli. Adversarial robustness through local linearization. In *Advances in Neural Information Processing Systems*, 2019.

Adi Shamir, Itay Safran, Eyal Ronen, and Orr Dunkelman. A simple explanation for the existence of adversarial examples with small hamming distance. *arXiv preprint arXiv:1901.10861*, 2019.

Christian Szegedy, Wojciech Zaremba, Ilya Sutskever, Joan Bruna, Dumitru Erhan, Ian Goodfellow, and Rob Fergus. Intriguing properties of neural networks. In *International Conference on Learning Representations*, 2014.

Roman Vershynin. Introduction to the non-asymptotic analysis of random matrices. In *Compressed Sensing: Theory and Practice*, pages 210–268. Cambridge University Prteess, 2012.

Martin J Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge University Press, 2019.