Multistream Modelling for Mental Health: Modelling Linguistic and Temporal Contexts with Mutual and Self-Excitation in Social Media

Abstract

We present a novel multistream modeling approach applied to mental health monitoring, integrating multiple perspectives of a user's mental state. We introduce MHRoBERT, a hierarchical transformer architecture that captures both cross-stream interactions through multistream HEAT layers and general temporal dynamics through unistream aggregation, while maintaining direct access to sequential information via residual connections. The model demonstrates competitive performance in detecting moments of change in mental health states, with learned interaction parameters revealing interpretable patterns in how different mental health indicators influence each other temporally. Notably, multistream information proves valuable even for models not explicitly designed to leverage it, with LLM baselines showing substantial improvements when provided with stream annotations. This work provides a foundation for scalable, interpretable approaches to automated mental health monitoring.

1 Introduction

Mental health conditions manifest through multiple, interconnected signals – from emotional expressions and behavioral changes to physical symptoms and life events. Traditional approaches to analyzing social media for mental health monitoring typically examine different aspects of user behavior in isolation, failing to capture important patterns of interaction and mutual influence. Our work provides a framework for using multistream data to gain deeper insights into changes within an individual's mental health, with potential applications in real-time monitoring and intervention.

We introduce a hierarchical transformer architecture that captures both cross-stream interactions through multistream Hawkes process layers and general temporal dynamics through unistream aggregation, while maintaining direct access to sequential information via residual connections. We also develop a framework that uses Large Language Models (LLMs) to automatically extract and analyze three complementary streams of information from social media posts: (1) emotional states, (2) personal life events, and (3) mental health symptoms. Our approach is motivated by clinical understanding that mental health changes rarely occur in isolation. A significant life event can trigger emotional responses which may manifest as physical symptoms or behavioral changes. By modeling these interactions explicitly, we aim to develop a more comprehensive understanding of an individual's mental health over time.

The main contributions of our work are:

- We propose a novel multistream modelling approach for mental health monitoring that simultaneously integrates linguistic and temporal contexts from social media posts.
- Our model leverages Large Language Models (LLMs) to enhance the robustness of the created longitudinal user representations, using multiple streams of automatically labeled data.
- We introduce a mechanism to model both self-excitation within individual streams and mutual excitation between different streams of data, capturing more subtle and nuanced interactions in user behavior.
- We demonstrate the effectiveness of our approach through experiments on longitudinal social media data, showing improved performance in predicting mood changes.
- We show that the multistream component is more opaque with transparent parameters which also improves interpretabtility.

Through this multistream modeling approach, we identify patterns in how different aspects of mental health interact and evolve over time. The framework reveals cascading relationships between life events, emotional responses, and physical symptoms, providing a comprehensive overview

of mental health progression. By automating the extraction and longitudinal modeling of these clinically relevant information streams, we create scalable monitoring systems that can simultaneously track multiple mental health indicators.

2 Related Work

Multivariate Hawkes Processes Multivariate Hawkes processes provide the theoretical foundation for modeling mutual excitation between event streams. Liniger (2009) provided early formalization of the multivariate Hawkes process, while Zhou et al. (2013) developed efficient learning algorithms for large-scale applications. More recently, (Zhang et al., 2020) proposed the self-attentive Hawkes process, demonstrating how attention mechanisms could capture complex dependencies between different event types. In the social media domain, Rizoiu et al. (2017) used Hawkes processes to model information diffusion, demonstrating how different types of social interactions could mutually reinforce each other.

Large Language Models for Automated Annotation The use of LLMs for automated textual annotation is an emerging area and shows significant potential for creating large-scale, consistently labeled datasets at low cost.

Recent work shows that LLMs can match or exceed performance of crowd workers in text annotation tasks across multiple domains (Gilardi et al., 2023; Törnberg, 2023). Li and Conrad (2024) showed that LLMs are effective when annotating explicit stances in social media posts, achieving high agreement with human annotators. However, discrepancies arise in cases where stances are implicit or ambiguous, suggesting that hybrid approaches may be required for more nuanced annotation scenarios.

Additionally (Kristensen-McLachlan et al., 2023) raised concerns about transparency and reproducibility, highlighting significant variability in LLM performance across different tasks and noting that traditional supervised classifiers often outperform LLMs in scenarios requiring nuanced understanding. These challenges are especially relevant in mental health contexts, where subtle distinctions can have significant clinical implications.

Longitudinal Modeling for Mental Health LLMs have been used for mental health classification (Amin et al., 2023), data augmentation (Liyanage et al., 2023), and reasoning (Xu et al., 2024), demonstrating promise in detecting psychological indicators (Yang et al., 2023), extracting relevant evidence from text (Xu et al., 2024), and generating clinically informed summaries (Song et al., 2024). LLMs using instruction fine-tuning and Chain-of-Thought (CoT) prompting (Yang et al., 2023) have also been employed, though such approaches can cause risks of incorrect predictions and flawed reasoning, especially in complex conversations (Li et al., 2023).

Longitudinal modeling with large language models (LLMs) has emerged as a pivotal paradigm for capturing the evolution of language, behavior, and mental states over time. Unlike conventional NLP tasks that treat inputs as isolated instances, longitudinal modeling enable continuous monitoring and nuanced understanding of dynamic processes such as user stance changes on social media or psychological state progression in health narratives (Park and Conway, 2017), (Tsakalidis et al., 2022c,a; Tseriotou et al., 2023, 2024; Hills et al., 2024), (Morini et al., 2025). In dialogue systems (Kwak et al., 2023), incorporating longitudinal context is essential for maintaining personalization (Chen et al., 2023a), (Chen et al., 2023b), and psychological grounding across multiple conversational turns (Wang et al., 2023), (Zheng et al., 2023). Current approached have not been integrated with LLMs and focus on a single stream of data (unistream).

3 Methodology

We first discuss how we obtain multiple interacting data streams from a single data stream (§3.1). These are expected to influence each other but follow different temporal patterns. We then present a multistream temporal modeling architecture, with cross-excitation for capturing interactions between streams (§3.2)

3.1 Stream Extraction using LLMs

We use LLMs to extract three complementary perspectives from social media posts (for the full taxonomies and prompts used to extract these labels see Appendix A), namely:

Emotional States: For emotions, we adopt the taxonomy of the fine-grained labels for Affect proposed in the MIND framework (Slonim, 2024), used to label the same dataset we make use of in this work in the CLPSych 2025 shared task (Tseriotou et al., 2025). These consist of twelve intensive emotions (e.g. Happy, Proud, Sad) spanning adap-

tive and maladaptive affective states.

Personal Life Events: To capture significant life changes that may impact mental health, we employed the broad categories proposed in (Anonymous, 2025). These categories, designed specifically for mental health transitions, were inspired by the Major Life Events Taxonomy of (Haimson et al., 2021), which identifies 121 life event types across 12 categories, including health, relationships, career, and identity changes.

Symptoms: We used the symptoms categories introduced in (Anonymous, 2025) to capture both physical and psychological labels related to mental health changes. The framework encompasses several key categories of symptoms that commonly manifest during periods of mental health changes – including substance use patterns, trauma responses, physical symptoms (pain, fatigue), sleep disturbances, and appetite changes.

This overall categorization of the streams was designed to identify warning signs and behavioral changes that could indicate changes in mood and mental health, while remaining broad enough to capture diverse conditions.

3.1.1 Prompt Design and Language Model

We implement our automated stream annotation using Qwen 2.5-32B-Instruct (Qwen et al., 2025) in a chain-of-thought (CoT) few-shot classification setting. Our selection of Qwen 2.5 was informed by comparative evaluations demonstrating superior performance over other open-source models including Llama 3 for mental health annotation tasks (Anonymous, 2025). Additionally, a key consideration was the model's ability to run locally with efficient 4-bit quantization, enabling all processing to be performed on-premises without sending sensitive user data to external servers. This is in contrast to cloud-based language models that require data to be sent to remote services, making Qwen 2.5 particularly suitable for applications involving sensitive mental health data.

For each stream, we used chain-of-thought prompting strategies (detailed in Appendix A) that guide the model through explicit reasoning before producing final annotations. Our prompt design was adapted from the approach of (Anonymous, 2025), which emphasizes clear category definitions with concrete examples and requires systematic category-by-category evaluation. In this approach, the model explains its reasoning for each category with explicit yes" or no" statements before producing a final answer, thereby improving annotation quality and interpretability.

The resulting multistream annotations provide complementary perspectives informing mental health changes, simultaneously capturing emotional states, personal life events, and symptoms. This multistream approach helps enable a more dynamic understanding of longitudinal mental health trajectories than would be possible through an analysis of raw posts from a single perspective.

3.2 Multistream HEAT Architecture

Our model extends a temporal aggregation mechanism inspired by the Hawkes process that transforms linguistic representations into time-aware encodings. This method, HEAT: Historical Emotional AggregaTion, originally proposed by Sawhney et al., HEAT weights historical information based on temporal proximity using learnable self-excitation (ϵ) and decay (β) parameters. The self-excitation parameter ϵ controls how much previous events influence future representations, while the decay parameter β determines how quickly this influence diminishes over time. We modify this approach to handle multiple interacting streams of temporally sensitive linguistic data, by processing user timelines through three main stages:

Linguistic Encoding Posts are first encoded using BERT (`bert-base-uncased`) to obtain contextual embeddings. For each post i in a timeline, BERT produces a representation $e^{(i)} \in \mathbb{R}^{768}$ by extracting the [CLS] token embedding.

Sequential Modeling The BERT embeddings are processed through an LSTM layer to capture sequential dependencies:

$$h^{(i)} = \text{LSTM}(e^{(i)}, h^{(i-1)}) \quad (1)$$

where $h^{(i)} \in \mathbb{R}^{d_{lstm}}$ represents the hidden state encoding both linguistic content and sequential context up to post i.

Multistream HEAT Representation For each target stream l, we compute a multistream HEAT representation that captures cross-stream temporal interactions:

$$H_{\text{multi}}^{(i,l)} = \mu_l + \sum_m \sum_{j<i} h^{(j)} \cdot \mathbb{1}_m^{(j)} \cdot \epsilon_{(l,m)} \cdot e^{-\beta_{(l,m)} \Delta \tau_j}$$

$$(2)$$

where: μ_l is the learned baseline intensity for target stream l, $\mathbb{1}_m^{(j)} \in \{0, 1\}$ indicates whether post j belongs to stream m, $\epsilon_{(l,m)} \in \mathbb{R}^+$ captures the

excitation from stream m to stream l, $\beta_{(l,m)} \in \mathbb{R}^+$ controls the temporal decay rate of this excitation and $\Delta\tau_j = t_i - t_j$ is the time elapsed since post j. The learned ϵ and β matrices reveal interpretable patterns of how different mental health indicators influence each other temporally.

Figure 1: MHRoBERT architecture for MoC prediction. Posts are encoded with BERT and processed through an LSTM to capture sequential context. An LLM annotates posts with emotion, life event, and symptom labels. The final representation concatenates three components: (1) a residual connection from the current post's LSTM state, (2) a unistream representation aggregating all LSTM states via HEAT, and (3) a multistream representation where HEAT pools LSTM states masked by their stream labels. This combined representation predicts whether a given post indicates a MoC.

Unistream HEAT Representation   In addition to the multistream representation, we compute a unistream HEAT aggregation over all posts, regardless of stream assignment. This uses a Markovian emphasis (Hills et al., 2024) that weights recent context more heavily:

$$H_{\text{uni}}^{(i)} = h^{(i-1)} + \sum_{j < i} h^{(j)} \cdot \epsilon_{\text{uni}} \cdot e^{-\beta_{\text{uni}}\Delta\tau_j} \quad (3)$$

where ϵ_{uni} and β_{uni} are separate learnable parameters which aggregate historical representations regardless of stream labels, in a the unistream fashion.

Combined Representation   The final representation concatenates three components:

$$\mathbf{r}^{(i)} = [H_{\text{multi}}^{(i,1)}; H_{\text{multi}}^{(i,2)}; H_{\text{multi}}^{(i,3)}; H_{\text{uni}}^{(i)}; h^{(i)}] \quad (4)$$

where the semicolon denotes concatenation. This design captures: (a) Stream-specific temporal patterns (multistream HEAT); (b) General temporal

dynamics (unistream HEAT); (c) Direct sequential information (residual LSTM connection) (See Fig. 1). The concatenated representation is scaled with $\tanh$ activation and passed through a linear classifier:

$$\hat{y}^{(i)} = \text{softmax}(W_{\text{out}} \cdot \tanh(\mathbf{r}^{(i)}) + b_{\text{out}}) \quad (5)$$

4   Experiments

Dataset and Preprocessing   We make use of the Reddit dataset introduced in (Tsakalidis et al., 2022b) containing longitudinal social media timelines labelled for Moments of Change in mood (MoC) . Given a historical sequence of posts the task is to identify whether a given post denotes a Switch-S, is part of an Escalation-E or Neither-O. Training and implementation details are described in section C.

Model Configuration   We implement and evaluate three variants of hierarchical architectures:
RoBERT: "Recurrence over BERT". Baseline with BERT encoding and LSTM sequential modeling, without temporal aggregation layers. The classifier operates directly on LSTM hidden states.
HoRoBERT: "HEAT over Recurrence over BERT". Hierarchical model with a single Markovian HEAT layer (Hills et al., 2024). This applies temporal aggregation over all LSTM hidden states using learnable ϵ and β parameters, with emphasis on recent context through Markovian weighting.
MHRoBERT: "Multistream HEAT over Recurrence over BERT". Our proposed multistream architecture that extends HoRoBERT with three key components (See Fig. 1):

- *Multistream HEAT*: Stream-specific temporal aggregations with cross-stream excitation matrices $\epsilon_{(l,m)}$ and $\beta_{(l,m)}$, plus per-stream baseline intensities μ_l
- *Unistream HEAT*: A separate Markovian HEAT layer aggregating all posts regardless of stream labels, using independent parameters ϵ_{uni} and β_{uni}
- *Residual connection*: Direct access to the current post's LSTM hidden state

The final representation concatenates all three components before classification, enabling the model to leverage both stream-specific interactions and general temporal patterns.

All models use BERT (`bert-base-uncased`) with 768-dimensional embeddings. In the fine-tuned setting, BERT parameters are updated end-to-end during training. In the frozen setting, BERT

weights are fixed and only the LSTM and subsequent layers are trained. For multi-task variants, we add an auxiliary classification head predicting stream labels, sharing the same concatenated representation as the main MoC prediction task.

4.1 LLMsfor Mood Change Detection (MoC)

We compared the performance of our specialized models developed for temporal modelling with LLMs applied in a zero-shot setting. To assess the capabilities of LLM models for automatic mood change detection (MoC) in longitudinal data, we utilised a 32-billion parameter instruction-tuned model **Qwen2.5-32B-Instruct** (Reategui-Rivera et al., 2025; Nie et al., 2024). Evaluation was performed on **Reddit** data, through prompting-based inference under two settings:

(a) QwenLLM: In this setting, the model was prompted to classify each post within a user's timeline as one of S (Switch), E (Escalation), or O (None), depending on the past posts as context. Each post was evaluated independently, given only the past textual history of the user's posts.

(b) QwenLLMMultistream: The QwenLLM-Multistream prompting gives the model richer contextual signals that mimic a structured psychological assessment framework. The model used parallel cues pertaining to fine-grained emotions, mental health symptoms and personal life events to identify transitions (S), intensification (E), or stability (O) in the user's emotional trajectory. See Appendix B for the complete prompt templates and configuration details for both LLM models.

5 Results

Table 1 shows the performance of different architectures for predicting Moments of Change in mood. When fine-tuning BERT, HoRoBERT achieves the best overall macro-averaged F1 score (0.694), with MHRoBERT achieving competitive performance (0.689). All three hierarchical architectures demonstrate strong performance when BERT is fine-tuned end-to-end.

5.1 Model Performance Across Settings

The **fine-tuned BERT** setting yields the strongest results overall. HoRoBERT shows particular strength in recall (0.718) and achieves the best F1 for Switches (0.531) and Escalations (0.638). MHRoBERT achieves the highest precision (0.689) and performs well on Switches (F1 = 0.515), though slightly behind HoRoBERT. The baseline

RoBERT (LSTM without temporal modeling) also demonstrates competitive performance (macro F1 = 0.685), suggesting that sequential modeling with fine-tuned representations captures substantial signal for mood change detection.

With **frozen BERT**, all models show reduced performance (macro F1 = 0.600–0.609), as expected when linguistic representations cannot adapt to the task. However, the **multi-task learning** setting, which also uses frozen BERT but incorporates auxiliary tasks predicting stream-specific labels (emotions, life events, symptoms), achieves remarkably competitive performance (macro F1 = 0.673–0.674). This represents only a 2–3% drop compared to fine-tuned BERT, despite using fixed representations. This demonstrates that **multistream auxiliary supervision provides substantial benefits**, effectively compensating for the lack of linguistic fine-tuning by encouraging the model to learn representations sensitive to multiple aspects of mental health.

Under multi-task learning, MHRoBERT achieves the best Switch detection (F1 = 0.514), while all three models show strong and balanced performance. The consistently high performance across architectures in this setting suggests that the auxiliary stream prediction tasks provide effective regularization and guide the model toward learning more robust representations of mental health trajectories.

5.2 Benefits of Multistream Information

A key contribution of this work is demonstrating that easily collected, scalable, noisy LLM generated **multistream labels improve performance across different model architectures**, including those not explicitly designed to handle this information. The LLM results provide compelling evidence: while the baseline QwenLLM achieves only 0.427 macro F1, incorporating multistream information in prompts (QwenLLMMultistream) improves performance to 0.481 (12.6% relative improvement). This improvement is particularly notable for rare classes, with Switch recall improving from 0.496 to 0.650.

These results highlight the value of the multistream annotation framework beyond the specific MHRoBERT architecture. The stream labels, automatically generated using LLMs applied to social media posts, are **scalable and easily transferable** to other models and datasets. They provide a consistent benefit whether the model explicitly

Model	Macro-Avg			Switch (S)			Escalation (E)			No Change (O)		
	P	R	F1	P	R	F1	P	R	F1	P	R	F1
Fine-tuned BERT												
MHRoBERT	**.689**	.696	.689	**.551**	.485	.515	**.579**	.697	.632	.936	**.906**	**.921**
HoRoBERT	.680	**.718**	**.694**	.532	**.531**	**.531**	.564	**.738**	**.638**	**.944**	.886	.913
RoBERT	.675	.700	.685	.516	.512	.514	.572	.685	.623	.937	.901	.918
Single-task (Frozen BERT)												
MHRoBERT	.567	.703	.600	.293	**.639**	.402	.449	.712	.551	.958	.759	.847
HoRoBERT	.574	**.713**	.607	**.310**	.629	**.414**	.446	**.760**	**.562**	**.967**	.752	.846
RoBERT	**.575**	.703	**.609**	.307	.616	.408	**.462**	.717	.561	.958	**.776**	**.857**
Multi-task (Frozen BERT)												
MHRoBERT	.689	.662	.674	.559	**.476**	**.514**	.599	.586	.592	.909	.923	.916
HoRoBERT	.688	**.663**	**.674**	.533	.467	.497	**.620**	.595	.606	.911	**.927**	**.919**
RoBERT	**.696**	.660	.673	**.585**	.432	.496	.590	**.628**	**.608**	**.913**	.919	.917
LLM Baselines												
QwenLLM	.447	.496	.427	.119	.496	.192	.343	.357	.350	.879	.636	.738
QwenLLMMultistream	**.517**	**.570**	**.481**	**.141**	**.650**	**.232**	**.468**	**.367**	**.411**	**.941**	**.694**	**.799**

Table 1: Per-class and macro-averaged results, for predicting MoCs. For each metric, **bold** denotes the best score within that experimental condition, while **bold+underline** denotes the best score across all conditions. P = Precision, R = Recall, F1 = F1-score.

models stream interactions (MHRoBERT), uses them as auxiliary supervision (multi-task setting), or simply incorporates them in prompts (LLMs). This transferability makes the approach practical for real-world deployment, as the synthetic stream annotations can be generated at scale without manual labeling and applied to improve mental health monitoring systems.

Furthermore, the multistream framework enhances model **interpretability at multiple levels**. At the data level, individual posts can be inspected for their stream-specific labels, allowing clinicians or researchers to understand which aspects of mental health (emotions, life events, symptoms) are present in the discourse.

At the model level, architectures like MHRoBERT produce interpretable learned parameters (Section 5.5) that reveal how different streams interact temporally. Even for models without explicit multistream architectures, the presence of stream labels enables more transparent analysis -— practitioners can examine not just whether a mood change was predicted, but which combination of emotional expressions, life circumstances, and symptom mentions contributed to that prediction. This interpretability is particularly valuable in mental health applications where understanding the reasoning behind predictions is as important as prediction accuracy itself.

5.3 Comparison of Temporal Modeling Approaches

The improvements from adding temporal HEAT layers (HoRoBERT) or multistream HEAT (MHRoBERT) over the LSTM baseline (RoBERT) in the fine-tuned setting suggest several considerations. The strong performance of RoBERT indicates that sequential modeling with fine-tuned BERT embeddings captures much of the relevant temporal structure. While MHRoBERT's multistream architecture provides interpretable cross-stream interaction parameters (Section 5.5), the performance gains over simpler approaches are modest in the fine-tuned setting. However, the architecture's ability to leverage multistream information proves particularly valuable in the multi-task setting, where it achieves the best Switch detection. This suggests that explicit multistream modeling becomes more beneficial when combined with auxiliary supervision that encourages learning stream-specific patterns.

5.4 Analysis of Model Components

The results demonstrate that both the unistream temporal component (HoRoBERT) and residual LSTM representations (present in all models) contribute meaningfully to performance. HoRoBERT consistently achieves high recall on rare events, suggesting that the unistream HEAT component effectively captures temporal patterns associated with mood changes. MHRoBERT's additional mul-

tistream HEAT component provides interpretable interaction parameters while maintaining competitive performance across all settings.

The importance of end-to-end BERT fine-tuning is evident, with fine-tuned models substantially outperforming frozen BERT variants in single-task learning. However, the multi-task results reveal that **multistream auxiliary supervision can largely bridge this gap**, achieving near-equivalent performance with frozen representations. This finding has practical implications, as it enables effective mood change detection without expensive fine-tuning, making the approach more accessible for resource-constrained applications or domains where labeled data for fine-tuning is limited.

5.5 Stream Interaction Analysis

The resulting model produces interpretable parameters, including excitation (ϵ) and decay (β) matrices, along with baseline intensities (μ), which provide insights into the temporal relationships between streams. These relationships are visualized in Figure 2.

Our multistream approach demonstrates several advantages over the baseline approaches. The integration of emotional states, life events, and symptoms provides a more comprehensive view of mental health changes, with the learned interaction parameters (Figure 2) offering insights into temporal dynamics between streams.

The **excitation patterns** reveal differential influences between streams. Emotions exhibit the highest self-excitation ($\epsilon = 0.01$), suggesting emotional states tend to reinforce themselves. Cross-stream excitation from emotions to symptoms ($\epsilon = 0.009$) is notably strong, indicating that emotional states may influence symptom manifestation. Life events show more balanced excitation across streams ($\epsilon = 0.004$–0.006), while symptoms demonstrate weaker cross-stream influence overall ($\epsilon = 0.002$–0.004).

The **decay rates** (β) show interesting temporal patterns. Emotions display very slow self-decay ($\beta = 0.0002$), suggesting persistence of emotional states, while their cross-stream decay is faster ($\beta = 0.004$–0.007). Life events exhibit consistently fast decay across all streams ($\beta = 0.01$–0.009), potentially reflecting their more transient nature. The decay from emotions to symptoms is relatively slow ($\beta = 0.001$), which may indicate lasting emotional influence on symptoms.

The **baseline intensities** show positive values for emotions ($\mu = 0.0062$) and symptoms ($\mu = 0.0056$), with life events near zero ($\mu = -0.00093$). This pattern suggests that while emotional and symptom streams maintain some baseline activity, life events are primarily driven by external occurrences rather than intrinsic baseline patterns. These learned parameters provide a foundation for understanding how different aspects of mental health interact temporally, though further investigation would be valuable to validate these patterns against clinical observations.

5.6 Stream Interpretability Analysis

The decomposition of the multistream HEAT component of MHRoBERT provides key interpretability insights. As shown in Figure 3, the *Emotion* stream contributes the largest share of multistream attribution ($\sim 41\%$), followed by *Symptom* ($\sim 39\%$) and *Life Event* ($\sim 19\%$). This ordering, consistent across classes, indicates a coherent internal structure within multistream HEAT.

Table 2 summarizes the representational geometry results using Nearest-Centroid (global) and k-Nearest Neighbors (local) evaluations. The unistream space exhibits stronger class separability and contributes more to gradient attributions, whereas the multistream space, though less discriminative, remains semantically informative. Its lower F1 reflects an expected entanglement of cross-stream embeddings that jointly encode emotional, symptomatic, and life-event cues.

While this representation is less separable, it captures richer inter-stream relationships, enhancing interpretability and potential transferability. In contrast, the unistream representation emphasizes temporal discrimination, producing distinct clusters aligned with mental state transitions. Together, these complementary inductive biases suggest that multistream HEAT prioritizes semantic alignment and interpretability, while unistream HEAT maintains geometric clarity and predictive precision as reflected in the fused stream results.

5.7 Future Research Directions

This work opens several promising avenues for future research in automated mental health monitoring and detecting changes more generally, by considering multiple interrelated temporal streams of data. In terms of stream extraction, more sophisticated prompt engineering techniques could improve the quality and reliability of automated annotations. Integration with external clinical knowl-

Figure 2: Learned interaction parameters showing excitation (ϵ) and decay (β) values between different streams, where higher values indicate stronger relationships. Learned baseline intensities (μ) for each individual stream are presented, where higher values indicate stronger contribution to the associated stream representations - regardless of event occurrence.

edge bases could provide more psychologically validated stream categories, while systematic comparison with expert human annotation would help validate the LLM-based approach.

The modeling framework itself could be extended in several ways. Incorporating more advanced, fine-tuned linguistic representations could better capture the nuances of mental health-related language. Additional streams, such as social interaction patterns or environmental factors, could provide even more comprehensive monitoring capabilities. The temporal modeling could be enhanced to capture more complex dynamics, including seasonal patterns and long-term trends.

From a clinical application perspective, this lays the groundwork for developing and improving existing mental health monitoring frameworks. The interpretable nature of the stream interactions could help clinicians understand the progression of mental health states over time, particularly by highlighting which combinations of emotional, behavioral, and life event patterns most strongly indicate impending changes in mental health.

6 Conclusion

This paper has presented a novel hierarchical multistream approach to mental health monitoring that takes advantge of multiple streams of information extracted from social media posts. By combining Large Language Models for automated stream annotation with MHRoBERT: a hierarchical architecture that fuses multistream HEAT layers, unistream temporal aggregation, and residual LSTM connections – we have demonstrated the potential for more comprehensive and interpretable approaches to detecting mental health changes.

The key contributions include:

- MHRoBERT, a novel hierarchical architecture

that integrates stream-specific temporal interactions, general temporal dynamics, and direct sequential information through a unified representation.
- Empirical evidence that easily obtained multistream information benefits diverse architectures – from specialized temporal models to general LLMs – demonstrating the broad applicability of the multistream framework.
- Interpretable learned parameters (ϵ, β, μ) that insightful interaction patterns between different mental health indicators, with potential value for better clinical understanding.

The ability to capture and interpret interactions between different aspects of mental health represents an important step toward more holistic approaches to mental health monitoring. While the architectural gains over simpler baselines are modest in the fine-tuned setting, the multistream framework's consistent benefits across different models and training regimes – particularly in multitask learning highlight its practical value. The framework's interpretability and transferability make it particularly promising for both research and potential clinical applications.

Future work could focus on refining stream extraction through improved prompting strategies and clinical validation, exploring additional streams (social interactions, external environmental factors), and investigating more sophisticated temporal modeling to capture seasonal patterns and long-term trends. These developments could lead to more reliable systems for monitoring mental health through social media, supporting better outcomes and understanding for individuals experiencing mental health difficulties.

Limitations

While this work demonstrates the potential of multistream temporal modelling for mental health monitoring, several important limitations should be acknowledge.

Performance Gains   The performance improvements of MHRoBERT over simpler baselines are relatively modest in the fine-tuned setting, with macro F1 scores differing by only a few percentage points. While the multistream information provides interpretable cross-stream interaction parameters and shows stronger benefits in multi-task learning scenarios, the gains over the LSTM baseline (RoBERT) in single-task fine-tuned settings are limited.

Stream Annotation Quality   The multistream annotations are generated automatically using LLMs without human validation, which introduces potential noise and inconsistencies. While LLMs have shown promise for annotation tasks (Gilardi et al., 2023; Törnberg, 2023), they can struggle with implicit or ambiguous content (Li and Conrad, 2024) and show significant variability across tasks (Kristensen-McLachlan et al., 2023). In mental health contexts, where subtle distinctions have significant clinical implications, the lack of expert validation of our automatically generated stream labels represents a notable limitation. The three-stream taxonomy (emotions, life events, symptoms), while grounded in psychological frameworks, has not been validated by mental health professionals for this specific application. Despite the noisy imperfect annotations, we still demonstrate several benefits for making use of such annotations in this paper.

Temporal Modeling Assumptions   The HEAT mechanism assumes that temporal influence decays exponentially over time, which may not accurately reflect all psychological processes. Mental health trajectories can involve complex patterns including seasonality, and long-term trends that may not be well-captured by exponential decay functions. The current implementation uses learned static baseline intensities, whereas more sophisticated approaches might benefit from dynamic, context-dependent baselines that adapt to individual users' temporal patterns.

Clinical Validity and Deployment   This work has not been validated in clinical settings or with mental health professionals. The interpretable parameters produced by MHRoBERT (excitation and decay matrices) require clinical evaluation to determine whether they align with established understanding of mental health dynamics. Furthermore, deploying such systems for real-world mental health monitoring raises critical concerns about false positives/ negatives, privacy, informed consent, and the potential for harm if predictions are acted upon without appropriate clinical oversight. The models are intended as research tools to understand mental health patterns in social media data, not as diagnostic or intervention tools.

Computational Requirements   The hierarchical architecture with end-to-end BERT fine-tuning requires substantial computational resources, which may limit accessibility for researchers or organizations with limited infrastructure. While the multitask frozen BERT setting offers a more resource-efficient alternative, it still requires significant compute for training on longitudinal data.

Ethical Considerations   Automated monitoring of mental health through social media raises important ethical questions about privacy, consent, and appropriate use of such systems. While this work uses publicly available Reddit data with institutional review board approval, the ability to identify mood changes from user posts could potentially be misused for surveillance or discrimination. Users posting on public forums may not anticipate or consent to their content being analyzed for mental health indicators.

Ethics Statement

Before starting this research, approval was secured from the Institutional Review Board of the lead university. This study considers ethical considerations when dealing with the analysis of user-generated content on social media platforms, specifically Reddit. The ethical implications of our research, in particular the ability to identify changes in mood within user timelines, share similar concerns to that of prior research focused on identifying personal events through social media, and recognizing signs of suicidal thoughts. To help mitigate these risks, measures were taken such as the limited and regulated access to the developed software and the annotations that were used in this study.

A   LLM Prompting Templates

We use chain-of-thought (CoT) prompting with Qwen 2.5-32B-Instruct to extract three streams of

mental health-relevant information from social media posts. Below are the complete prompt templates used for each stream.

A.1 Model Configuration

We implement our automated stream annotation using Qwen 2.5-32B-Instruct with the following configuration:

- **Model**: Qwen 2.5-32B-Instruct

- **Quantization**: 4-bit quantization

- **Precision**: bfloat16

- **Decoding Strategy**: Greedy decoding (deterministic) with do_sample=False

- **Max New Tokens**: 1024 tokens

- **Temperature**: 0.0 (greedy decoding)

A.2 Emotional States Prompt

```
You are tasked with classifying the
    emotional content of a social media
    post.

Emotion Categories:
- Happy: Content, joyful, hopeful,
    excited, glad, cheerful
- Proud: Feeling accomplished,
    satisfied with achievements,
    confident in success
- Sad: Emotional pain, grieving, hurt,
    heartbroken over loss
- Anxious: Fearful, tense, worried,
    nervous, scared, panicked
- Depressed: Despair, hopeless,
    worthless, empty, suicidal thoughts
- Apathetic: Numb, indifferent, don't
    care, emotionally blunted
- Angry: Aggression, hate, rage,
    disgust, contempt
- Ashamed: Guilty, embarrassed,
    humiliated, self-critical
- Loneliness: Feeling alone, isolated,
    lacking meaningful connection
- None: The post does not clearly
    reflect any of the above emotion
    categories.

Instructions:
1. Based on the provided examples,
    analyze the post by evaluating each
    emotion category.
2. For each category, explain your
    reasoning and clearly state "yes"
    or "no" regarding its presence.
3. Finally, list all applicable emotion
    categories that are present in the
    post. If more than one applies,
    separate them with commas.

Below are some examples:
```

```
{few_shot_examples}

Post to Analyze:
Post:
"{post}"

Please strictly follow the output
    format exactly as shown below. Do
    not use bold, markdown, or extra
    formatting.

Output Format:
Explanation:
- Happy: [reason]. So the answer is yes
    (or is no).
- Proud: [reason]. So the answer is yes
    (or is no).
- Sad: [reason]. So the answer is yes
    (or is no).
- Anxious: [reason]. So the answer is
    yes (or is no).
- Depressed: [reason]. So the answer is
    yes (or is no).
- Apathetic: [reason]. So the answer is
    yes (or is no).
- Angry: [reason]. So the answer is yes
    (or is no).
- Ashamed: [reason]. So the answer is
    yes (or is no).
- Loneliness: [reason]. So the answer
    is yes (or is no).
- None: [reason]. So the answer is yes
    (or is no).

Answer:
Only output exact category names:
    Happy, Proud, Sad, Anxious,
    Depressed, Apathetic, Angry,
    Ashamed, Loneliness, or None. Use
    commas to separate multiple labels,
    if any.
```

A.3 Personal Life Events Prompt

```
You are provided with a social media
    post and must identify any personal
    life events based on the life event
    categories defined below.

Life events are experiences that have a
    major personal impact on an
    individual. They must involve a
    clearly identifiable occurrence or
    change. These events may have
    occurred in the past (explicitly
    stated, or inferred from context if
    the impact is clear), be occurring
    in the present (explicitly
    described or clearly implied), or
    be expected in the near future
    (only if explicitly stated).

- Mental Health:
Mental health life events cover
    discussions about receiving a
    formal mental health diagnosis;
    starting or adjusting psychiatric
    medication; beginning therapy;
    recovery or significant symptom
```

improvement; or experiencing acute psychological episodes such as manic episodes, psychotic breaks, panic attacks, dissociative episodes, self-harm, suicide attempts, or acute suicidal ideation requiring crisis intervention. It does not include general low mood, stress, anxiety, or depression unless explicitly tied to one of these major events.
- Physical Health:
Physical health life events cover accidents; injuries; diagnoses or survival of serious illnesses; chronic conditions that have a notable impact on daily life; or cases where the individual or someone important to them has been hospitalized, undergone surgery, received emergency medical treatment, or experienced other major medical interventions. Pregnancy-related experiences, including becoming pregnant, pregnancy loss, abortion, or complications, are also included. Common aches, headaches, fatigue, or other symptoms without being part of a major physical health event are not included.
- Abuse & Addiction:
Covers major life events involving abuse or substance-related issues. Abuse may occur in childhood or adulthood and may take physical, sexual, emotional, psychological, or other forms. Addiction-related events include the onset of heavy drug or alcohol use; acknowledgment of addiction; overdoses; withdrawal symptoms; relapses; or recovery (including participation in treatment programs such as rehabilitation or support groups).
- Relationship & Loss:
Covers specific relationship changes or disruptions in meaningful personal connectionsfamily or non-familythat result in strong emotional impact. Includes beginning or ending friendships or romantic relationships; marriage or divorce; parental separation or divorce; serious arguments or conflicts; relationships becoming abusive or toxic; the addition of new family members (e.g., through birth or adoption); parenting difficulties; emotionally distressing health issues involving family members, partners, close friends, or other meaningful connections; or the death of a person or pet who was important to the individual.
- Career & Education:
Career-related life events cover starting or losing a job; promotions; demotions; troubles at work; being unable to find a job; retirement; or becoming a business owner. Education-related life events include starting or graduating from school; transferring schools; leaving school without graduating; being denied entry into school; taking major exams; experiencing significant academic challenges; or achieving major academic milestones (e.g., earning a certification).
- Financial & Legal & Societal:
Financial life events include major purchases (e.g., a house or car); financial gains or milestones (e.g., paying off debt); financial losses (e.g., stolen or damaged property); or ongoing financial difficulties. Legal life events include law violations; arrests; court appearances; lawsuits or legal actions; or major interactions with the justice system, whether the individual is subject to legal proceedings or initiating them. Societal life events with clear personal impact include natural disasters; pandemics; major political events; societal issues; or notable public encounters (e.g., meeting a celebrity).
- Lifestyle & Identity & Environment:
Covers major life events involving changes in lifestyle habits, personal identity, or living environment. Lifestyle changes include adopting new health- and well-being-related routines, such as adjustments in diet; exercise; sleep management; stress management (e.g., through meditation); reducing substance use; fostering positive social connections; or getting a pet. It does not include clinical treatment for mental health issues. Identity transitions include identifying or changing one's gender or sexual orientation (e.g., coming out as LGBTQ+); discussing a new sexual experience; or exploring, adopting, or changing political or religious beliefs (including changes in religious practices). Relocation life events include moving to a new place; moving in or out of the family home; family members moving into or out of the household; or significant travel experiences.
- None:
Applies when the post does not clearly reflect any of the above life event categories.

Instructions:
1. Read the post carefully and evaluate whether it matches each defined life event category.

2. For each life event category, explain your reasoning. If a category applies, support your answer with direct evidence from the post. If it does not apply, explain why there is insufficient or no evidence. Clearly state "yes" or "no" for each category.
3. Finally, list all broad life event categories that apply. If more than one applies, separate them with commas.

Below are some examples:
{few_shot_examples}

Post to Analyze:
Post:
"{post}"

Please strictly follow the output format exactly as shown below. Do not use bold, markdown, or extra formatting.

Output Format:
Explanation:
- Mental Health: [reason]. So the answer is yes (or is no).
- Physical Health: [reason]. So the answer is yes (or is no).
- Abuse & Addiction: [reason]. So the answer is yes (or is no).
- Relationship & Loss: [reason]. So the answer is yes (or is no).
- Career & Education: [reason]. So the answer is yes (or is no).
- Financial & Legal & Societal: [reason]. So the answer is yes (or is no).
- Lifestyle & Identity & Environment: [reason]. So the answer is yes (or is no).
- None: [reason]. So the answer is yes (or is no).

Answer:
Output exact category names: Mental Health, Physical Health, Abuse & Addiction, Relationship & Loss, Career & Education, Financial & Legal & Societal, Lifestyle & Identity & Environment, or None. Use commas to separate multiple labels, if any."""

A.4 Mental Health Symptoms Prompt

You are provided with a social media post and must identify any symptoms of mental health issues according to the symptom categories defined below.

Symptom Categories:
- Suicidal Thoughts:
Distinct from general distress, this involves specific thoughts about ending one's life, including considering methods, making plans, or expressing a desire to die. This includes passive thoughts ("I wish I wouldn't wake up") and active plans, as well as references to past suicide attempts.

- Somatoform & Substance Abuse:
Somatoform refers to physical symptoms that cause significant distress but lack clear medical explanations. This includes various types of pain (e.g., headaches, muscle aches); unexplained physical sensations; physical fatigue; and physical complaints affecting different body systems. They are often accompanied by cognitive symptoms such as impaired memory, reduced processing speed, and poor planning. Substance abuse covers problematic use of alcohol, drugs, or other substances. This includes excessive drinking or drug use; failed attempts to cut down; spending a disproportionate amount of time obtaining, using, or recovering from substances; and continued use despite negative consequences.

- Eating Pathology:
Includes problematic cognitions and behaviors related to food, eating patterns, body image, and weight. This includes fear of weight gain (e.g., frequent body-checking, obsessive weighing); body image disturbance (e.g., negative fixation on weight or shape); self-worth tied to appearance (e.g., harsh self-criticism, comparisons); denial of low weight; and guilt/shame around eating (e.g., anxiety about food, avoidance of eating situations). Other features include using food as emotional coping and obsessive focus on food, cooking, or weight-related content. Behavioral manifestations include restrictive eating (e.g., smaller portions, skipping meals, eliminating food groups, extreme diets, calorie counting); fasting to compensate; and chewing and spitting (chewing food without swallowing). Binge eating involves consuming abnormally large amounts of food with feelings of loss of control, eating rapidly or until uncomfortably full, eating alone due to embarrassment, and experiencing negative emotions afterward. Compensatory behaviors include self-induced vomiting, misuse of laxatives or diuretics, use of diet pills or "slimming teas," fasting as a non-purging compensatory behavior and excessive

```
                exercise (e.g., working out despite
                fatigue/injury, exercise
                interfering with daily
                functioning). Secretive eating
                behaviors, including concealing
                eating patterns or using coded
                terminology online, may also occur.

        - Sexual:
        Involves difficulties with sexual
                functioning and satisfaction. This
                includes problems with sexual
                desire, arousal, or performance;
                pain during sexual activity;
                distress related to sexual
                experiences; and addictive sexual
                behaviors that impair various other
                functions.

        - Thought Disorder:
        Relates to disruptions in perception,
                beliefs, and thought organization.
                This includes unusual sensory
                experiences (e.g., hearing voices);
                extreme and fixed, false beliefs;
                disorganized thinking or speech;
                and difficulty distinguishing
                between what is real and what is
                not.

        - Antisocial and Antagonistic Behavior:
        Refers to actions that violate societal
                norms and infringe upon the rights
                of others. This includes
                rule-breaking or illegal behavior,
                physical aggression, bullying,
                deceitfulness or theft, blame
                externalization, disregard for
                social norms or obligations, and
                property destruction. It also
                covers behaviors such as
                deliberately provoking or annoying
                others, attention-seeking,
                rudeness, manipulating others for
                personal gain, and other
                deliberately over-extroverted
                behaviors (physical or emotional).
                These behaviors may be accompanied
                by impulsivity, distractibility,
                and risk-taking.

        - Detachment:
        Characterized by withdrawal from social
                and emotional experiences. This
                includes avoidance of social
                interaction, limited emotional
                expression, lack of close
                relationships, reduced capacity to
                experience pleasure in activities
                (particularly in social contexts),
                and social isolation.

        - None:
        Applies when the post does not clearly
                reflect any of the above symptom
                categories.

        Instructions:
        1. Read the post carefully and evaluate
                whether it matches each defined
                symptom category.
        2. For each category, explain your
                reasoning. If a category applies,
                support your answer with direct
                evidence from the post. If it does
                not apply, explain why there is
                insufficient or no evidence.
                Clearly state "yes" or "no" for
                each category.
        3. Finally, list all applicable symptom
                categories that are present in the
                post. If more than one applies,
                separate them with commas.

        Below are some examples:
        {few_shot_examples}

        Post to Analyze:
        Post:
        "{post}"

        Please strictly follow the output
                format exactly as shown below. Do
                not use bold, markdown, or extra
                formatting.

        Output Format:
        Explanation:
        - Suicidal Thoughts: [reason]. So the
                answer is yes (or is no).
        - Somatoform & Substance Abuse:
                [reason]. So the answer is yes (or
                is no).
        - Eating Pathology: [reason]. So the
                answer is yes (or is no).
        - Sexual: [reason]. So the answer is
                yes (or is no).
        - Thought Disorder: [reason]. So the
                answer is yes (or is no).
        - Antisocial and Antagonistic Behavior:
                [reason]. So the answer is yes (or
                is no).
        - Detachment: [reason]. So the answer
                is yes (or is no).
        - None: [reason]. So the answer is yes
                (or is no).

        Answer:
        Output exact category names: Suicidal
                Thoughts, Somatoform & Substance
                Abuse, Eating Pathology, Sexual,
                Thought Disorder, Antisocial and
                Antagonistic Behavior, Detachment,
                or None. Use commas to separate
                multiple labels, if any."""
```

A.5 Chain-of-Thought Examples

To illustrate the reasoning process required by our prompts, we provide synthetic in context learning examples demonstrating the chain-of-thought approach. These examples are fictional and created for demonstration purposes to protect the privacy of the individuals in our datasets. The actual prompts used during annotation contained 5-10 similar examples per stream, all following the same structured reasoning format shown below. Each ex-

ample demonstrates explicit evaluation of every category before producing a final classification.

A.5.1 Emotion Classification Example

```
Post:
"Finally submitted my thesis after
    months of work! Feeling accomplished
and ready for what's next."

Explanation:
- Happy: The post explicitly states
    "Feeling accomplished" which
    directly
  indicates pride and satisfaction. The
      forward-looking tone suggests
  hopefulness. So the answer is yes.
- Proud: While accomplishment is
    present, this aligns more with
    general
  happiness than specific pride in
      achievement. So the answer is no.
- Sad: The post expresses positive
    emotions with no grief or emotional
    pain.
  So the answer is no.
- Anxious: No fear, worry, or tension
    is expressed. So the answer is no.
- Depressed: The post conveys
    accomplishment and optimism,
    opposite to
  despair. So the answer is no.
- Apathetic: Clear emotional engagement
    is present, not numbness.
  So the answer is no.
- Angry: No aggression or anger is
    expressed. So the answer is no.
- Ashamed: The post expresses positive
    feelings, not guilt or
    embarrassment.
  So the answer is no.
- Loneliness: No isolation or lack of
    connection is mentioned.
  So the answer is no.
- None: Clear positive emotion is
    present. So the answer is no.

Answer:
Happy
```

A.5.2 Life Events Classification Example

```
Post:
"Started therapy last week and got
    prescribed medication for my
    anxiety.
Hoping this helps."

Explanation:
- Mental Health: The post explicitly
    mentions starting therapy and
    receiving
  psychiatric medication for anxiety,
      which are major mental health
      treatment
  events. So the answer is yes.
- Physical Health: No accidents,
    injuries, illnesses, or physical
    medical
```

interventions are mentioned. So the
    answer is no.
- Abuse & Addiction: No mention of
    abuse or substance-related issues.
  So the answer is no.
- Relationship & Loss: No specific
    relationship changes are described.
  So the answer is no.
- Career & Education: No work or
    school-related events are mentioned.
  So the answer is no.
- Financial & Legal & Societal: No
    financial, legal, or societal events
  are discussed. So the answer is no.
- Lifestyle & Identity & Environment:
    The post focuses on clinical
  intervention rather than lifestyle
      modifications. So the answer is
      no.
- None: A clear mental health life
    event is present. So the answer is
    no.

Answer:
Mental Health

A.5.3 Symptoms Classification Example

```
Post:
"Constant headaches all week and
    exhausted no matter how much I
    sleep.
Also been drinking more than usual to
    cope."

Explanation:
- Suicidal Thoughts: No death wishes or
    suicidal ideation are expressed.
  So the answer is no.
- Somatoform & Substance Abuse: The
    post mentions "constant headaches"
    and
  chronic exhaustion (somatic symptoms)
      plus "drinking more than usual to
  cope" (substance use for emotional
      regulation). So the answer is yes.
- Eating Pathology: No eating, weight,
    or body image concerns are
    mentioned.
  So the answer is no.
- Sexual: No sexual symptoms are
    discussed. So the answer is no.
- Thought Disorder: The thinking is
    coherent with no perceptual
    disturbances.
  So the answer is no.
- Antisocial and Antagonistic Behavior:
    No rule-breaking or aggression is
  described. So the answer is no.
- Detachment: While fatigue is present,
    there's no indication of social
  withdrawal. So the answer is no.
- None: Clear symptoms are present. So
    the answer is no.

Answer:
Somatoform & Substance Abuse
```

B   Baseline LLM Prompting Templates for Mood Change Detection

This section details the complete prompt templates and model configurations used for mood change detection (MoC) experiments with LLMs.

B.1   Model Configuration

For automatic mood change detection, we used the following configuration:

- **Model:** Qwen2.5-32B-Instruct

- **Quantization:** 4-bit (NF4)

- **Precision:** bfloat16

- **Decoding Strategy:** Greedy decoding (do_sample=False)

- **Attention Implementation:** Default (no custom optimization)

- **Tokenizer Padding:** Right-side padding with EOS token substitution

- **Hardware:** Single NVIDIA GPU with 4-bit quantized loading via BitsAndBytes

- **Max New Tokens:** 32

- **Temperature:** 0.0

B.2   Mood Change Detection Prompt (QwenLLM)

```
You are an expert assistant for
    identifying self-disclosed mood
    changes in user timelines on social
    media.

Your task is to classify the current
    post into exactly one of these
    three labels based on the context
    of previous posts:

- S (Switch): A sudden shift in mood
    direction (positive  negative)
    compared to prior posts.
- E (Escalation): A gradual or clear
    intensification of mood in the same
    direction.
- O (None): No significant mood change
    or mood remains neutral/consistent.

### Instructions:
- Read all previous posts in the
    timeline as context to understand
    the user's prior mood.
- Read the current post carefully.
- Determine if the current post
    represents a sudden mood switch
    (S), a gradual escalation (E), or
    no significant mood change (O).
```

```
- **Output only the label (S, E, or O)
    on a single line no extra text,
    explanation, or commentary**
- If there are **no previous posts**,
    assume the baseline mood is neutral.

Previous posts:
{prev_posts if prev_posts else "No
    previous posts"}

Current post:
{current_post}

Answer:
```

Listing 1: QwenLLM prompt template for timeline-based MoC classification.

B.3   Mood Change Detection (QwenLLMMultistream) Prompt

```
You are an assistant for identifying a
    user's self-disclosed mood changes
    over time.
Each post belongs to a timeline of user
    posts. Your task is to classify the
    current
post based on prior posts as one of the
    following:

- S (Switch): Sudden shift in mood
    direction (positive  negative)
- E (Escalation): Gradual or clear
    intensification of mood in the same
    direction
- O (None): No significant mood change
    or mood remains consistent

AUTOMATED MULTISTREAM LABEL CONTEXT:
Each post includes automated labels
    from three streams:

1. FINE-GRAINED EMOTION LABELS (12
    categories organized by
    adaptiveness):

  Positive Adaptive Emotions:
  - Happy: Content, joyful, hopeful,
      excited, glad, cheerful
  - Proud: Feeling accomplished,
      satisfied with achievements,
      confident in success
  - Calm: Peaceful, relaxed, serene,
      tranquil emotional states
  - Vigor: High energy, vitality,
      enthusiasm, active engagement
  - Feeling loved: Experiencing
      affection, care, or emotional
      support from others

  Negative Adaptive Emotions:
  - Sad: Emotional pain, grieving,
      hurt, heartbroken over loss
      (adaptive response to loss)
  - Justifiable anger: Appropriate
      anger in response to unfair
      treatment or injustice

  Negative Maladaptive Emotions:
```

- Anxious: Fearful, tense, worried, nervous, scared, panicked
- Depressed: Despair, hopeless, worthless, empty, suicidal thoughts
- Apathetic: Numb, indifferent, don't care, emotionally blunted
- Angry: Aggression, hate, rage, disgust, contempt (non-justified)
- Ashamed: Guilty, embarrassed, humiliated, self-critical
- Loneliness: Feeling alone, isolated, lacking meaningful connection

2. MENTAL HEALTH SYMPTOM LABELS (7 categories):

- Suicidal Thoughts: Specific thoughts about ending one's life, including considering methods, making plans, expressing desire to die, passive thoughts ("I wish I wouldn't wake up"), active plans, references to past suicide attempts

- Somatoform & Substance Abuse: Physical symptoms causing significant distress without clear medical explanations (headaches, muscle aches, unexplained sensations, physical fatigue, cognitive symptoms like impaired memory/processing). Substance abuse includes problematic alcohol/drug use, failed attempts to cut down, excessive time obtaining/using/recovering from substances, continued use despite negative consequences

- Eating Pathology: Problematic food/eating behaviors and cognitions including fear of weight gain, body image disturbance, self-worth tied to appearance, guilt/shame around eating, using food as emotional coping, restrictive eating patterns, binge eating with loss of control, compensatory behaviors (vomiting, laxatives, excessive exercise), secretive eating behaviors

- Sexual: Difficulties with sexual functioning including problems with desire/arousal/performance, pain during sexual activity, distress related to sexual experiences, addictive sexual behaviors

- Thought Disorder: Disruptions in perception, beliefs, thought organization including unusual sensory experiences (hearing voices), extreme fixed false beliefs, disorganized thinking/speech, difficulty distinguishing reality

- Antisocial and Antagonistic Behavior: Actions violating societal norms and others' rights including rule-breaking, illegal behavior, physical aggression, bullying, deceitfulness, blame externalization, property destruction, deliberately provoking others, attention-seeking, rudeness, manipulation for personal gain

- Detachment: Withdrawal from social and emotional experiences including avoidance of social interaction, limited emotional expression, lack of close relationships, reduced capacity for pleasure (especially social contexts), social isolation

3. PERSONAL LIFE EVENT LABELS (7 categories):

- Mental Health: Receiving formal mental health diagnosis, starting/adjusting psychiatric medication, beginning therapy, recovery/significant symptom improvement, acute psychological episodes (manic episodes, psychotic breaks, panic attacks, dissociative episodes, self-harm, suicide attempts, acute suicidal ideation requiring crisis intervention)

- Physical Health: Accidents, injuries, diagnoses/survival of serious illnesses, chronic conditions with notable daily life impact, hospitalization, surgery, emergency medical treatment, major medical interventions, pregnancy-related experiences (becoming pregnant, pregnancy loss, abortion, complications)

- Abuse & Addiction: Major events involving abuse (physical, sexual, emotional, psychological) in childhood or adulthood; addiction-related events including onset of heavy drug/alcohol use, acknowledgment of addiction, overdoses, withdrawal symptoms, relapses, recovery (treatment programs, rehabilitation, support groups)

```
- Relationship & Loss: Specific
    relationship changes/disruptions
    in meaningful personal
    connections with strong
    emotional impact including
    beginning/ending friendships or
    romantic relationships,
    marriage/divorce, parental
    separation/divorce, serious
    arguments/conflicts,
    relationships becoming
    abusive/toxic, addition of new
    family members, parenting
    difficulties, emotionally
    distressing health issues
    involving family/partners/close
    friends, death of important
    person or pet

- Career & Education: Career events
    (starting/losing job,
    promotions, demotions, work
    troubles, unable to find job,
    retirement, becoming business
    owner); Education events
    (starting/graduating school,
    transferring schools, leaving
    without graduating, denied
    school entry, major exams,
    significant academic challenges,
    major academic milestones)

- Financial & Legal & Societal:
    Financial events (major
    purchases like house/car,
    financial gains/milestones,
    financial losses, ongoing
    financial difficulties); Legal
    events (law violations, arrests,
    court appearances, lawsuits,
    major justice system
    interactions); Societal events
    with personal impact (natural
    disasters, pandemics, major
    political events, societal
    issues, notable public
    encounters)

- Lifestyle & Identity &
    Environment: Lifestyle changes
    (adopting new health routines,
    diet/exercise/sleep/stress
    management adjustments, reducing
    substance use, fostering
    positive social connections,
    getting pet - not clinical
    mental health treatment);
    Identity transitions
    (gender/sexual orientation
    identification/changes, coming
    out as LGBTQ+, new sexual
    experiences,
    exploring/adopting/changing
    political/religious beliefs);
    Relocation events (moving to new
    place, moving in/out of family
    home, family members moving
    in/out of household, significant
    travel experiences)
```

```
INTERPRETATION GUIDELINES:
- Fine-grained emotions (especially
    transitions between
    Positive/Negative
    Adaptive/Maladaptive categories)
    are primary indicators of mood
    switches and escalations
- Life events often trigger mood
    changes - look for temporal
    relationships between events and
    emotional shifts
- Symptom patterns can indicate
    escalating mental health states or
    underlying mood deterioration
- Cross-stream interactions (e.g.,
    relationship loss + depressed
    emotion + detachment symptoms)
    provide rich context for mood
    trajectory assessment
- Labels are automated and noisy but
    provide valuable psychological
    state indicators for detecting mood
    changes over time

You will ONLY output one of the three
    labels: S, E, or O. No additional
    explanation.

Previous posts in timeline:
{prev_context if prev_context else "No
    previous posts."}

Current post to classify:
{current_formatted}

Classification:
```

Listing 2: QwenLLMMultistream Prompt template for multistream mood change detection (S, E, O).

C Training and Implementation

C.1 Infrastructure

All models (excluding LLMs) were implemented with PyTorch and trained on 2 NVIDIA A100-PCIE-40GB GPUs using Focal Loss to address class imbalance in labels of Moments of Change - for single-task models. LLM models were all implemented with 2 NVIDIA H100 GPUs.

Multi-task models used a combined loss of focal loss and binary cross entropy (BCE) loss, giving equal weighting to both losses and equal weighting to the 3 streams in the BCE loss. The optimization process used the Adam optimizer with gradient accumulation of 4 steps. Training was carried out with a maximum of 10 epochs, selecting the best model in terms of macro-average F1 score on a validation set using 5-fold cross validation. The hyperparameter search space is described in Table C.2.

Method	UHEAT	MHEAT	Fused
Nearest-Centroid (cosine)			
Macro-F1	0.4413	0.3368	**0.4493**
Precision	0.4388	0.3812	0.4388
Recall	0.5091	0.4229	0.5091
k-NN (k=5, cosine)			
Macro-F1	0.4032	0.3882	**0.4057**
Precision	0.3961	0.3994	0.4014
Recall	0.4106	0.3907	0.4106

Table 2: Representation geometry analysis.

C.2 Grid-search Used in Experimental Procedure

We train and evaluate all our models on 3 seeds, taking the average scores on the resulting test sets - aside from the LLM baselines which were evaluated with 1 seed. We evaluate on the same test set proposed in the CLPsych 2022 shared task for Reddit (Tsakalidis et al., 2022b). We performed a grid-search on both datasets (§5.1), over the enlisted hyper-parameters – selecting the best performing model based on macro-average F1 score on the validation set, and optimizing the model using focal loss with a gamma of 2.0, training for 10 epochs, Learning rate: 0.00005, LSTM/ Transformer hidden dimension: 512, $\epsilon_{prior} = \beta_{prior} = 0.001$ for MHRoBERT and $\epsilon_{prior} = \beta_{prior} = 0.01$ for HoRoBERT, chunk size: 16, stride: 8.

D Component wise interpretability Analysis

References

Mostafa M. Amin, Erik Cambria, and Björn W. Schuller. 2023. Will affective computing emerge from foundation models and general artificial intelligence? a first evaluation of chatgpt. *IEEE Intelligent Systems*, 38(2):15–23.

Anonymous. 2025. Anonymous.

Ruijun Chen, Jin Wang, Liang-Chih Yu, and Xuejie Zhang. 2023a. Learning to memorize entailment and discourse relations for persona-consistent dialogues.

Ruijun Chen, Jin Wang, Liang-Chih Yu, and Xuejie Zhang. 2023b. Learning to memorize entailment and discourse relations for persona-consistent dialogues. In *Proceedings of the AAAI conference on artificial intelligence*, volume 37, pages 12653–12661.

Fabrizio Gilardi, Meysam Alizadeh, and Maël Kubli. 2023. Chatgpt outperforms crowd workers for text-annotation tasks. *Proceedings of the National Academy of Sciences*, 120(30):e2305016120.

Oliver L. Haimson, Albert J. Carter, Shanley Corvite, Brookelyn Wheeler, Lingbo Wang, Tianxiao Liu, and Alexxus Lige. 2021. The major life events taxonomy: Social readjustment, social media information sharing, and online network separation during times of life transition. *Journal of the Association for Information Science and Technology*, 72(7):933–947.

Anthony Hills, Talia Tseriotou, Xenia Miscouridou, Adam Tsakalidis, and Maria Liakata. 2024. Exciting mood changes: A time-aware hierarchical transformer for change detection modelling. In *Findings of the Association for Computational Linguistics: ACL 2024*, pages 12526–12537, Bangkok, Thailand. Association for Computational Linguistics.

Ross Deans Kristensen-McLachlan, Miceal Canavan, Márton Kardos, Mia Jacobsen, and Lene Aarøe. 2023. Chatbots are not reliable text annotators. *arXiv preprint arXiv:2311.05769*.

Jin Myung Kwak, Minseon Kim, and Sung Ju Hwang. 2023. Context-dependent instruction tuning for dialogue response generation.

Mao Li and Frederick Conrad. 2024. Advancing annotation of stance in social media posts: A comparative analysis of large language models and crowd sourcing. *arXiv preprint arXiv:2406.07483*.

Yucheng Li, Bo Dong, Chenghua Lin, and Frank Guerin. 2023. Compressing context to enhance inference efficiency of large language models.

Thomas J Liniger. 2009. Multivariate hawkes processes: an application to financial data. *ETH Zurich*.

Chandreen Liyanage, Muskan Garg, Vijay Mago, and Sunghwan Sohn. 2023. Augmenting Reddit posts to determine wellness dimensions impacting mental health. In *The 22nd Workshop on Biomedical Natural Language Processing and BioNLP Shared Tasks*, pages 306–312, Toronto, Canada. Association for Computational Linguistics.

Virginia Morini, Salvatore Citraro, Elena Sajno, Maria Sansoni, Giuseppe Riva, Massimo Stella, and Giulio Rossetti. 2025. Online posting effects: Unveiling the non-linear journeys of users in depression communities on reddit.

Jingping Nie, Hanya Shao, Yuang Fan, Qijia Shao, Haoxuan You, Matthias Preindl, and Xiaofan Jiang. 2024. Llm-based conversational ai therapist for daily functioning screening and psychotherapeutic intervention via everyday smart devices. *arXiv preprint arXiv:2403.10779*.

Albert Park and Mike Conway. 2017. Longitudinal changes in psychological states in online health community members: Understanding the long-term effects of participating in an online depression community. *Journal of Medical Internet Research*, 19:e71.

Qwen, :, An Yang, Baosong Yang, Beichen Zhang, Binyuan Hui, Bo Zheng, Bowen Yu, Chengyuan Li, Dayiheng Liu, Fei Huang, Haoran Wei, Huan Lin, Jian Yang, Jianhong Tu, Jianwei Zhang, Jianxin Yang, Jiaxi Yang, Jingren Zhou, Junyang Lin, Kai Dang,

Figure 3: Class wise multistream attribution with multistream HEAT component.

Keming Lu, Keqin Bao, Kexin Yang, Le Yu, Mei Li, Mingfeng Xue, Pei Zhang, Qin Zhu, Rui Men, Runji Lin, Tianhao Li, Tianyi Tang, Tingyu Xia, Xingzhang Ren, Xuancheng Ren, Yang Fan, Yang Su, Yichang Zhang, Yu Wan, Yuqiong Liu, Zeyu Cui, Zhenru Zhang, and Zihan Qiu. 2025. Qwen2.5 technical report.

C Mahony Reategui-Rivera, Aref Smiley, and Joseph Finkelstein. 2025. Llm-based chatbot to reduce mental illness stigma in healthcare providers. In *2025 IEEE 15th Annual Computing and Communication Workshop and Conference (CCWC)*, pages 00001–00007. IEEE.

Marian-Andrei Rizoiu, Young Lee, Swapnil Mishra, and Lexing Xie. 2017. Hawkes processes for events in social media. In *Frontiers of multimedia research*, pages 191–218.

Ramit Sawhney, Harshit Joshi, Rajiv Ratn Shah, and Lucie Flek. 2021. Suicide Ideation Detection via Social and Temporal User Representations using Hyperbolic Learning. In *Proceedings of the 2021 Conference of the North American Chapter of the Association for Computational Linguistics: Human Language Technologies*, pages 2176–2190, Online. Association for Computational Linguistics.

Dana Atzil Slonim. 2024. Self-other dynamics (sod): A transtheoretical coding manual.

Jiayu Song, Jenny Chim, Adam Tsakalidis, Julia Ive, Dana Atzil-Slonim, and Maria Liakata. 2024. Combining hierachical vaes with llms for clinically meaningful timeline summarisation in social media.

Petter Törnberg. 2023. Chatgpt-4 outperforms experts and crowd workers in annotating political twitter messages with zero-shot learning. *arXiv preprint arXiv:2304.06588*.

Adam Tsakalidis, Jenny Chim, Iman Munire Bilal, Ayah Zirikly, Dana Atzil-Slonim, Federico Nanni, Philip Resnik, Manas Gaur, Kaushik Roy, Becky Inkster, Jeff Leintz, and Maria Liakata. 2022a. Overview of the CLPsych 2022 shared task: Capturing moments of change in longitudinal user posts. In *Proceedings of the Eighth Workshop on Computational Linguistics and Clinical Psychology*, pages 184–198, Seattle, USA. Association for Computational Linguistics.

Adam Tsakalidis, Jenny Chim, Iman Munire Bilal, Ayah Zirikly, Dana Atzil-Slonim, Federico Nanni, Philip Resnik, Manas Gaur, Kaushik Roy, Becky Inkster, et al. 2022b. Overview of the clpsych 2022 shared task: Capturing moments of change in longitudinal user posts.

Adam Tsakalidis, Federico Nanni, Anthony Hills, Jenny Chim, Jiayu Song, and Maria Liakata. 2022c. Identifying moments of change from longitudinal user text. In *Proceedings of the 60th Annual Meeting of the Association for Computational Linguistics (Volume 1: Long Papers)*, pages 4647–4660.

Talia Tseriotou, Jenny Chim, Ayal Klein, Aya Shamir, Guy Dvir, Iqra Ali, Cian Kennedy, Guneet Singh Kohli, Anthony Hills, Ayah Zirikly, Dana Atzil-Slonim, and Maria Liakata. 2025. Overview of the CLPsych 2025 shared task: Capturing mental health dynamics from social media timelines. In *Proceedings of the 10th Workshop on Computational Linguistics and Clinical Psychology (CLPsych 2025)*, pages 193–217, Albuquerque, New Mexico. Association for Computational Linguistics.

Talia Tseriotou, Adam Tsakalidis, Peter Foster, Terence Lyons, and Maria Liakata. 2023. Sequential path signature networks for personalised longitudinal language modeling. In *Findings of the Association for Computational Linguistics: ACL 2023*, pages 5016–5031.

Talia Tseriotou, Adam Tsakalidis, and Maria Liakata. 2024. Tempoformer: A transformer for temporally-aware representations in change detection.

Hongru Wang, Rui Wang, Fei Mi, Yang Deng, Zezhong Wang, Bin Liang, Ruifeng Xu, and Kam-Fai Wong. 2023. Cue-cot: Chain-of-thought prompting for responding to in-depth dialogue questions with llms.

Xuhai Xu, Bingsheng Yao, Yuanzhe Dong, Saadia Gabriel, Hong Yu, James Hendler, Marzyeh Ghassemi, Anind K. Dey, and Dakuo Wang. 2024. Mental-llm: Leveraging large language models for mental health prediction via online text data. *Proc. ACM Interact. Mob. Wearable Ubiquitous Technol.*, 8(1).

Kailai Yang, Shaoxiong Ji, Tianlin Zhang, Qianqian Xie, Ziyan Kuang, and Sophia Ananiadou. 2023. Towards interpretable mental health analysis with large language models. In *Proceedings of the 2023 Conference on Empirical Methods in Natural Language Processing*. Association for Computational Linguistics.

Qiang Zhang, Aldo Lipani, Omer Kirnap, and Emine Yilmaz. 2020. Self-attentive hawkes process. In *International conference on machine learning*, pages 11183–11193. PMLR.

Wen Zheng, Natasa Milic-Frayling, and Ke Zhou. 2023. Contextual knowledge learning for dialogue generation.

Ke Zhou, Hongyuan Zha, and Le Song. 2013. Learning triggering kernels for multi-dimensional hawkes processes. In *International conference on machine learning*, pages 1301–1309. PMLR.