General Low-rank Matrix Optimization: Geometric Analysis and Sharper Bounds

Abstract

This paper considers the global geometry of general low-rank minimization problems via the Burer-Monterio factorization approach. For the rank-1 case, we prove that there is no spurious second-order critical point for both symmetric and asymmetric problems if the rank-2 RIP constant δ is less than $1/2$. Combining with a counterexample with $\delta = 1/2$, we show that the derived bound is the sharpest possible. For the arbitrary rank-r case, the same property is established when the rank-$2r$ RIP constant δ is at most $1/3$. We design a counterexample to show that the non-existence of spurious second-order critical points may not hold if δ is at least $1/2$. In addition, for any problem with δ between $1/3$ and $1/2$, we prove that all second-order critical points have a positive correlation to the ground truth. Finally, the strict saddle property, which can lead to the polynomial-time global convergence of various algorithms, is established for both the symmetric and asymmetric problems when the rank-$2r$ RIP constant δ is less than $1/3$. The results of this paper significantly extend several existing bounds in the literature.

1 Introduction

Given the natural numbers n, m and r, consider the low-rank matrix optimization problems

$$\min_{M \in \mathbb{R}^{n \times n}} f_s(M) \quad \text{s.t. } \operatorname{rank}(M) \leq r, \quad M^T = M, \quad M \succeq 0 \tag{1}$$

and

$$\min_{M \in \mathbb{R}^{n \times m}} f_a(M) \quad \text{s.t. } \operatorname{rank}(M) \leq r, \tag{2}$$

where the functions $f_s(\cdot)$ and $f_a(\cdot)$ are twice continuously differentiable. Problems (1)-(2) are referred to as the *symmetric* and the *asymmetric* problems, respectively. In addition, we call these problems *linear* if the objective function is induced by a linear measurement operator, i.e.,

$$f(M) = \tfrac{1}{2}\|\mathcal{A}(M) - b\|_F^2$$

for some vector $b \in \mathbb{R}^p$ and linear operator $\mathcal{A}$ mapping each matrix M to a vector in $\mathbb{R}^p$, where $f(M)$ denotes either $f_s(M)$ or $f_a(M)$. Those problems not fitting into the above model are called *nonlinear*. Low-rank optimization problems arise in a wide range of applications, e.g., matrix completion (Candès & Recht, 2009; Recht et al., 2010), phase synchronization (Singer, 2011; Boumal, 2016), and phase retrieval (Shechtman et al., 2015); see Chen & Chi (2018); Chi et al. (2019) for an overview of the topic. To overcome the non-convex rank constraint, one may resort to convex relaxations. The approach of replacing the rank constraint with a nuclear norm regularization is proven to provide the optimal sample complexity (Candès & Recht, 2009; Recht et al., 2010; Candès & Tao, 2010). However, solving the convexified problems involves computing a Singular Value Decomposition (SVD) in each iteration and results in heavy computational burdens. Along with the issue of large space complexities, the convexification approach is impractical for large-scale problems. Therefore, it is important to design efficient alternative methods with similar theoretical guarantees.

Submitted to 35th Conference on Neural Information Processing Systems (NeurIPS 2021). Do not distribute.

1.1 Burer–Monterio factorization and basic properties

Instead of directly solving convex relaxations of problems (1)-(2), we consider a computationally efficient approach, namely the Burer–Monterio factorization (Burer & Monteiro, 2003). The factorization approach is based on the observation that any matrix $M \in \mathbb{R}^{n \times m}$ with rank at most r can be written in the form of UV^T, where $U \in \mathbb{R}^{n \times r}$ and $V \in \mathbb{R}^{m \times r}$. Then, the asymmetric problem (2) is equivalent to

$$\min_{U \in \mathbb{R}^{n \times r}, V \in \mathbb{R}^{m \times r}} h_a(U, V), \tag{3}$$

where $h_a(U, V) := f_a(UV^T)$. Similarly, the symmetric problem (1) is equivalent to

$$\min_{U \in \mathbb{R}^{n \times r}} h_s(U), \tag{4}$$

where $h_s(U) := f_s(UU^T)$. The Burer-Monteiro factorization provides a natural parameterization of the low-rank structure of the unknown solution, and reformulates problems (1)-(2) as unconstrained optimization problems. In addition, the number of variables reduces from $O(n^2)$ or $O(nm)$ to as low as $O(n)$ or $O(n + m)$ when $r \ll \min\{n, m\}$. However, the reformulated problems are highly non-convex, and $\mathcal{NP}$-hard to solve in general. On the other hand, these problems share a specific non-convex structure, which makes it possible to utilize the structure and design efficient algorithms to find a global optimum under some conditions. In addition to the special structure, a regularity condition, named the Restricted Isometry Property, is required to guarantee the convergence of common iterative algorithms. We state the following two definitions only in the context of the symmetric problem since the corresponding definitions for the asymmetric problem are similar.

Definition 1 (Recht et al. (2010); Zhu et al. (2018)). Given natural numbers r and t, the function $f_s(\cdot)$ is said to satisfy the **Restricted Isometry Property** (RIP) of rank $(2r, 2t)$ for a constant $\delta \in [0, 1)$, denoted as δ-$\mathrm{RIP}_{2r,2t}$, if

$$(1 - \delta)\|K\|_F^2 \leq \left[\nabla^2 f_s(M)\right](K, K) \leq (1 + \delta)\|K\|_F^2$$

holds for all matrices $M, K \in \mathbb{R}^{n \times n}$ such that $\mathrm{rank}(M) \leq 2r, \mathrm{rank}(K) \leq 2t$, where $\left[\nabla^2 f_s(M)\right](\cdot, \cdot)$ is the curvature of the Hessian at point M.

The RIP property is satisfied in a wide range of applications, either on the entire low-rank manifold or on a part of the manifold that is of interest; examples include matrix sensing (Candès & Recht, 2009; Recht et al., 2010), matrix completion (Candès & Tao, 2010), and matrix sensing with non-Gaussian noise (Davenport et al., 2014). For instance, in the case of linear measurements with a Gaussian model, Candès & Recht (2009) showed that $O\left(nr/\delta^2\right)$ samples are enough to ensure the δ-$\mathrm{RIP}_{2r,2r}$ property with high probability. Since the analysis of the statistical behavior of problems (3)-(4) is not in the scope of this work, we focus on the case when the RIP property is satisfied by the objective function. We note that the RIP property is equivalent to the restricted strongly convex and smooth property defined in Wang et al. (2017); Park et al. (2018); Zhu et al. (2021) with the condition number $(1 + \delta)/(1 - \delta)$. Intuitively, the RIP property implies that the Hessian matrix is close to the identity tensor when the perturbation is restricted to be low-rank. This intuition naturally leads to the following definition.

Definition 2 (Bi & Lavaei (2021)). Given a natural number r, the function $f_s(\cdot)$ is said to satisfy the **Bounded Difference Property** (BDP) of rank $2r$ for a constant $\kappa \geq 0$, denoted as κ-BDP_{2r}, if

$$\left|\left[\nabla^2 f_s(M) - \nabla^2 f_s(M')\right](K, L)\right| \leq \kappa \|K\|_F \|L\|_F$$

holds for all matrices $M, M', K, L \in \mathbb{R}^{n \times n}$ such that $\mathrm{rank}(M), \mathrm{rank}(M'), \mathrm{rank}(K), \mathrm{rank}(L) \leq 2r$.

It has been proven in (Bi & Lavaei, 2021, Theorem 1) that those functions satisfying the δ-$\mathrm{RIP}_{2r,2r}$ property also satisfy the 4δ-BDP_{2r} property. With the RIP property, there are basically two categories of algorithms that can solve the factorized problem in polynomial time. Algorithms in the first category require a careful initialization so that the initial point is already in a small neighbourhood of a global optimum, and a certain local regularity condition in the neighbourhood ensures that local search algorithms will converge linearly to a global optimum; see Tu et al. (2016); Bhojanapalli et al. (2016a); Park et al. (2018) for a detailed discussion. The other class of algorithms is able

to converge globally from a random initialization. The convergence of these algorithms is usually established via the geometric analysis of the landscape of the objective function. One of the important geometric properties is the strict saddle property (Sun et al., 2018), which combined with the smoothness properties can guarantee the global polynomial-time convergence for various saddle-escaping algorithms (Jin et al., 2017, 2018; Sun et al., 2018; Huang & Becker, 2019). For the linear case, Ge et al. (2016, 2017) proved the strict saddle property for both problems (3)-(4) when the RIP constant is sufficiently small. More recently, Zhu et al. (2021) extended the results to the nonlinear asymmetric case. Moreover, a weaker geometric property, namely the non-existence of *spurious (non-global) second-order critical points*, has been established for both problems when the RIP constant is small (Li et al., 2019; Ha et al., 2020). We note that second-order critical points are points that satisfy the first-order and the second-order necessary optimality conditions, and thus the result of second-order critical points implies the non-existence of spurious local minima. Under certain regularity conditions, this weaker property is also able to guarantee the global convergence from a random initialization without an explicit convergence rate (Lee et al., 2016; Panageas & Piliouras, 2016). Please refer to Table 1 for a summary of the state-of-the-art results.

Most of the aforementioned papers are based on the following assumption on the low-rank critical points of the functions $f_s(\cdot)$ and $f_a(\cdot)$:

Assumption 1. The function $f_a(\cdot)$ has a first-order critical point M_a^* such that $\mathrm{rank}(M_a^*) \leq r$. Similarly, the function $f_s(\cdot)$ has a first-order critical point M_s^* that is symmetric, positive semi-definite and of rank at most r.

This assumption is inspired by the noiseless matrix sensing problem in the linear case for which the non-negative objective function becomes zero (the lowest value possible) at the true solution. This is a natural property of the matrix sensing problem for nonlinear measurement models as well. Under the above assumption and the RIP property, Zhu et al. (2018) proved that M_s^* and M_a^* are the unique global minima of problems (1)-(2).

Theorem 1 (Zhu et al. (2018)). *If the functions $f_s(\cdot)$ and $f_a(\cdot)$ satisfy the δ-RIP$_{2r,2r}$ property, then the critical points M_s^* and M_a^* are the* unique global minima *of problems* (1)-(2).

Given a solution (U^*, V^*) to problem (3), we observe that (U^*P, V^*P^{-T}) is also a solution for any invertible $P \in \mathbb{R}^{r \times r}$. This redundancy may induce an extreme non-convexity on the landscape of the objective function. To reduce this redundancy, Tu et al. (2016) considered the regularized problem

$$\min_{U \in \mathbb{R}^{n \times r}, V \in \mathbb{R}^{m \times r}} \rho(U, V), \tag{5}$$

where

$$\rho(U, V) := h_a(U, V) + \frac{\mu}{4} \cdot g(U, V)$$

with a constant $\mu > 0$ and the regularization term

$$g(U, V) := \|U^T U - V^T V\|_F^2.$$

The regularization term is introduced to balance the magnitudes of U^* and V^*. Zhu et al. (2018) showed that the regularization term does not introduce bias and thus problem (5) is equivalent to the original problem (2).

Theorem 2 (Zhu et al. (2018)). *Any first-order critical point (U^*, V^*) of problem* (5) *satisfies $(U^*)^T U^* = (V^*)^T V^*$. Moreover, problems* (2) *and* (5) *are equivalent.*

Detailed optimality conditions for problems (1)-(5) are provided in the appendix.

1.2 Contributions

In this work, we analyze the geometric properties of problems (4)-(5). Novel analysis methods are developed to obtain less conservative conditions for guaranteeing benign landscapes for both problems. We note that, unlike the linear measurements case, the RIP constant of nonlinear problems may not concentrate to 0 as the number of samples increases. Therefore, a sharper RIP bound leads to theoretical guarantees that hold under less stringent statistical requirements. In addition, even if the RIP constant concentrates to 0 when more samples are included, there may only be a limited number of samples available, either due to the constraints of specific applications or to the great expense of

Table 1: Comparison of the state-of-the-art results and our results. Here $\delta_{2r,2t}$ and κ are the RIP$_{2r,2t}$ and BDP$_{2r}$ constants of $f_s(\cdot)$ or $f_a(\cdot)$, respectively. Constant $\alpha(M_a^*) \in (0,1)$ only depends on M_a^*.

Problem Setups		No Spurious Second-order Critical Pts.		Strict Saddle Property	
		Existing	Ours	Existing	Ours
Rank-1 Sym.	Linear	$\delta_{2,2} < \frac{1}{2}$ (Zhang et al., 2019)	$\delta_{2,2} < \frac{1}{2}$	-	-
	Nonlinear	$\delta_{2,2} < \frac{2-O(\kappa)}{4+O(\kappa)}$ (Bi & Lavaei, 2021)	$\delta_{2,2} < \frac{1}{2}$	-	-
Rank-1 Asym.	Linear & Nonlinear	-	$\delta_{2,2} < \frac{1}{2}$	-	-
Rank-r Sym.	Linear	$\delta_{2r,2r} < \frac{1}{5}$ (Ge et al., 2016)	$\delta_{2r,2r} \le \frac{1}{3}$	$\delta_{2r,2r} < \frac{1}{10}$ (Ge et al., 2017)	$\delta_{2r,2r} < \frac{1}{3}$
	Nonlinear	$\delta_{2r,4r} < \frac{1}{5}$ (Li et al., 2019)	$\delta_{2r,2r} \le \frac{1}{3}$	-	$\delta_{2r,2r} < \frac{1}{3}$
Rank-r Asym.	Linear	$\delta_{2r,2r} < \frac{1}{3}$ (Ha et al., 2020)	$\delta_{2r,2r} \le \frac{1}{3}$	$\delta_{2r,2r} < \frac{1}{20}$ (Ge et al., 2017)	$\delta_{2r,2r} < \frac{1}{3}$
	Nonlinear	$\delta_{2r,2r} < \frac{1}{3}$ (Ha et al., 2020)	$\delta_{2r,2r} \le \frac{1}{3}$	$\delta_{2r,4r} < \frac{\alpha(M_a^*)}{100}$ (Zhu et al., 2021)	$\delta_{2r,2r} < \frac{1}{3}$

taking more samples. Hence, obtaining a sharper RIP bound is essential for many applications. We summarize our results in Table 1. More concretely, the contributions of this paper are three-folds.

First, we derive necessary conditions and sufficient conditions for the existence of spurious second-order critical points for both symmetric and asymmetric problems. Using our necessary conditions, we show that the δ-RIP$_{2r,2r}$ property with $\delta \le 1/3$ is enough to guarantee the non-existence of such points. This result provides a marginal improvement to the previous work (Ha et al., 2020), which developed the sufficient condition $\delta < 1/3$ for asymmetric problems, and is a major improvement over Ge et al. (2016) and Li et al. (2019), which requires $\delta < 1/5$ for symmetric problems. With this non-existence property and under some common regularity conditions, Lee et al. (2016); Panageas & Piliouras (2016) showed that the vanilla gradient descent method with a small enough step size and a random initialization almost surely converges to a global minimum. We note that the convergence rate was not studied and could theoretically be exponential in the worst case. In addition, by studying our necessary conditions, we show that every second-order critical point has a positive correlation to the global minimum when $\delta \in (1/3, 1/2)$. When $\delta = 1/2$, a counterexample with spurious second-order critical points is given by utilizing the sufficient conditions. We note that the sufficient conditions can greatly simplify the construction of counterexamples.

Second, we separately study the rank-1 case to further strengthen the bounds. In particular, we utilize the necessary conditions to prove that the δ-RIP$_{2,2}$ property with $\delta < 1/2$ is enough for the non-existence of spurious second-order critical points. Combining with a counterexample in the $\delta = 1/2$ case, we conclude that the bound $\delta < 1/2$ is the sharpest bound for the rank-1 case. Our results significantly extend the bounds in Zhang et al. (2019) derived for the linear symmetric case to the linear asymmetric case and the general nonlinear case. It also improves the bound in Bi & Lavaei (2021) by dropping the BDP constant.

Third, we prove that in the exact parametrization case, problems (4)-(5) both satisfy the strict saddle property (Sun et al., 2018) when the δ-RIP$_{2r,2r}$ property is satisfied with $\delta < 1/3$. This result greatly improves the bounds in Ge et al. (2017); Zhu et al. (2021) and extends the result in Ha et al. (2020) to approximate second-order critical points. With the strict saddle property and certain smoothness properties, a wide range of algorithms guarantee a global polynomial-time convergence with a random initialization; see Jin et al. (2017, 2018); Sun et al. (2018); Huang & Becker (2019). Due to the special non-convex structure of our problems and the RIP property, it is possible to prove the boundedness of the trajectory of the perturbed gradient descent method using a similar method as in Jin et al. (2017). Since the smoothness properties are satisfied over a bounded region, combining

with the strict saddle property, it follows that the perturbed gradient descent method (Jin et al., 2017) achieves a polynomial-time global convergence when $\delta < 1/3$.

1.3 Notation and organization

The operator 2-norm and the Frobenius norm of a matrix M are denoted as $\|M\|_2$ and $\|M\|_F$, respectively. The trace of matrix M is denoted as $\text{tr}(M)$. The inner product between two matrices is defined as $\langle M, N \rangle := \text{tr}(M^T N)$. For any matrix $M \in \mathbb{R}^{n \times m}$, we denote its singular values by $\sigma_1(M) \geq \cdots \geq \sigma_k(M)$, where $k := \min\{n, m\}$. For any symmetric matrix $M \in \mathbb{R}^{n \times n}$, we denote its eigenvalues by $\lambda_1(M) \geq \cdots \geq \lambda_n(M)$. The minimal eigenvalue is denoted as $\lambda_{min}(\cdot)$. For any matrix U, we use $\mathcal{P}_U$ to denote the orthogonal projection onto the column space of U. For any matrices $A, B \in \mathbb{R}^{n \times m}$, we use $A \otimes B$ to denote the fourth-order tensor whose (i, j, k, ℓ) element is $A_{i,j} B_{k,\ell}$. The identity tensor is denoted as $\mathcal{I}$. The notation $M \succeq 0$ means that the matrix M is symmetric and positive semi-definite. The sub-matrix $R_{i:j,k:\ell}$ consists of the i-th to the j-th rows and the k-th to the ℓ-th columns of matrix R. The action of the Hessian $\nabla^2 f(M)$ on any two matrices K and L is given by $[\nabla^2 f(M)](K, L) := \sum_{i,j,k,\ell} [\nabla^2 f(M)]_{i,j,k,\ell} K_{ij} L_{k,\ell}$.

In Section 2, the Singular Value Projection algorithm is analyzed as an enlightening example for our main results. Sections 3 and 4 are devoted to the non-existence of spurious second-order critical points and the strict saddle property of the low-rank optimization problem in both symmetric and asymmetric cases, respectively.

2 Motivating Example: Singular Value Projection Algorithm

Before providing theoretical results for problems (4)-(5), we first consider the Singular Value Projection Method (SVP) algorithm (Algorithm 1) as a motivating example, which is proposed in Jain et al. (2010). The SVP algorithm is basically the projected gradient method of the original low-rank problems (1)-(2) via the truncated SVD. For the asymmetric problem (2), the low-rank manifold is

$$\mathcal{M}_{asym} := \{M \in \mathbb{R}^{n \times m} \mid \text{rank}(M) \leq r\}$$

and the projection is given by only keeping components corresponding to the r largest singular values. For the symmetric problem (1), the low-rank manifold is

$$\mathcal{M}_{sym} := \{M \in \mathbb{R}^{n \times n} \mid \text{rank}(M) \leq r, \quad M^T = M, \quad M \succeq 0\}.$$

We assume without loss of generality that the gradient $\nabla f(\cdot)$ is symmetric; see Appendix A for a discussion. The projection is given by only keeping components corresponding to the r largest eigenvalues and dropping all components with negative eigenvalues. Since both low-rank manifolds are non-convex, the projection solution may not be unique and we choose an arbitrary solution when it is not unique. We note that the above projections are orthogonal in the sense that

$$\|M_+ - M\|_F = \min_{K \in \mathcal{M}} \|K - M\|_F,$$

where M_+ is the projection of a matrix M. Henceforth, $\mathcal{M}$ stands for $\mathcal{M}_{sym}$ or $\mathcal{M}_{asym}$, which should be clear from the context. Although each truncated SVD operation can be computed within $O(nmr)$ operations, the constant hidden in the $O(\cdot)$ notation is considerably larger than 1. Thus, the truncated SVD operation is significantly slower than matrix multiplication, which makes the SVP algorithm impractical for large-scale problems. However, the analysis of the SVP algorithm, combining with the equivalence property given in Ha et al. (2020), provides some insights into how to develop proof techniques for problems (4)-(5).

We extend the proof in Jain et al. (2010) and show that Algorithm 1 converges linearly to the global minimum under the δ-RIP$_{2r,2r}$ property with $\delta < 1/3$.

Theorem 3. *If function $f_s(\cdot)$ (resp. $f_a(\cdot)$) satisfies the δ-RIP$_{2r,2r}$ property with $\delta < 1/3$ and the step size is chosen to be $\eta = (1 + \delta)^{-1}$, then Algorithm 1 applied to problem* (1) *(resp.* (2)*) returns a solution M_T such that $M_T \in \mathcal{M}$ and $f(M_T) - f(M^*) \leq \epsilon$ within*

$$T := \left\lceil \frac{1}{\log[(1-\delta)/(2\delta)]} \cdot \log \left[\frac{f(M_0) - f(M^*)}{\epsilon} \right] \right\rceil$$

iterations, where $f(\cdot) := f_s(\cdot)$ (resp. $f(\cdot) := f_a(\cdot)$), M^* is the global minimum, M_0 is the initial point and $\lceil \cdot \rceil$ is the ceiling function.

Algorithm 1 Singular Value Projection (SVP) Algorithm

Input: Low-rank manifold $\mathcal{M}$, initial point M_0, number of iterations T, step size η, objective
function $f(\cdot)$.
Output: Low-rank solution M_T.
 1: **for** $t = 0, \ldots, T - 1$ **do**
 2:      Update $\tilde{M}_{t+1} \leftarrow M_t - \eta \nabla f(M_t)$.
 3:      Set M_{t+1} to be the projection of $\tilde{M}_{t+1}$ onto $\mathcal{M}$ via truncated SVD.
 4: **end for**
 5: **return** M_T.

We note that the above proof can be applied to other low-rank optimization problems with a suitable definition of the orthogonal projection. In Ha et al. (2020), it is proved that the unique global minimum is the only fixed point of the SVP algorithm if the RIP constant δ is less than $1/3$. However, the above paper has not proven the linear convergence (as done in Theorem 3). This difference leads to a strengthened inequality in the following analysis, which further serves as an essential step in proving the strict saddle property.

3   No Spurious Second-order Critical Points

In this section, we develop necessary conditions and sufficient conditions for the existence of spurious second-order critical points of problems (4)-(5). Besides the non-existence of spurious local minima, the non-existence of spurious second-order critical points also guarantees the global convergence of many first-order algorithms with random initialization under certain regularity conditions (Lee et al., 2016; Panageas & Piliouras, 2016). More precisely, we require the iteration points of the algorithm to converge to a single point and the objective function to have a Lipschitz-continuous gradient. The first condition is satisfied by the gradient descent method applied to a large class of functions known as the KŁ-functions (Attouch et al., 2013). For the second condition, many objective functions that appear in applications, e.g., the ℓ_2-loss function, do not satisfy this condition. However, if the step size is small enough, the special non-convex structure of the Burer-Monterio decomposition and the RIP property ensure that the trajectory of the gradient descent method stays in a compact set, where the Lipschitz condition is satisfied due to the second-order continuity of the functions $f_s(\cdot)$ and $f_a(\cdot)$. The proof of this claim is similar to Theorem 8 in Jin et al. (2017) and is omitted here. Therefore, the non-existence of spurious second-order critical points can ensure the global convergence of the gradient descent method for many applications.

The non-existence of spurious second-order critical points has been proved in Ge et al. (2017); Zhu et al. (2018) for problems with linear and nonlinear measurements, respectively. Recently, Ha et al. (2020) proved a relation between the second-order critical points of problem (3) or (5) and the fixed points of the SVP algorithm on problem (2). Using this relation, they showed that problems (3) and (5) have no spurious second-order critical points when the δ-RIP$_{2r,2r}$ property is satisfied with $\delta < 1/3$. In this work, we take a different approach to show that $\delta \leq 1/3$ is enough for the general case in both symmetric and asymmetric scenarios, and that $\delta < 1/2$ is enough for the rank-1 case. Moreover, we prove that there exists a positive correlation between every second-order critical point and the global minimum when $\delta \in (1/3, 1/2)$. We also show that there may exist spurious second-order critical points when $\delta = 1/2$ for both the symmetric and asymmetric problems, which extends the construction of such examples for the linear symmetric rank-1 problem in Zhang et al. (2018) to general cases. We first give necessary conditions and sufficient conditions for the existence of spurious second-order critical points below.

Theorem 4. *Let $\ell := \min\{m, n, 2r\}$. For a given $\delta \in [0, 1)$, there exists a function $f_a(\cdot)$ with the δ-RIP$_{2r,2r}$ property such that problems (3) and (5) have a spurious second-order critical point only if there exist a constant $\alpha \in (1 - \delta, (1 + \delta)/2]$, a diagonal matrix $\Sigma \in \mathbb{R}^{r \times r}$, a diagonal matrix $\Lambda \in \mathbb{R}^{(\ell-r) \times (\ell-r)}$ and matrices $A \in \mathbb{R}^{r \times r}, B \in \mathbb{R}^{r \times r}, C \in \mathbb{R}^{(\ell-r) \times r}, D \in \mathbb{R}^{(\ell-r) \times r}$ such that*

$$(1 + \delta) \min_{1 \leq i \leq r} \Sigma_{ii} \geq \max_{1 \leq i \leq \ell-r} \Lambda_{ii}, \quad \Sigma \succ 0, \ \Lambda \succeq 0,$$

$$\langle \Lambda, CD^T \rangle = \alpha \left[\text{tr}(\Sigma^2) - 2\langle \Sigma, AB^T \rangle + \|AB^T\|_F^2 + \|AD^T\|_F^2 + \|CB^T\|_F^2 + \|CD^T\|_F^2 \right], \quad (6)$$

$$\text{tr}(\Lambda^2) \leq \alpha^{-1}(2\alpha - 1 + \delta^2) \cdot \langle \Lambda, CD^T \rangle, \quad \langle \Lambda, CD^T \rangle \neq 0.$$

239 *If $CB^T = 0$ and $AD^T = 0$, then there exists a function $f_a(\cdot)$ with the δ-RIP$_{2r,2r}$ property such that*
240 *problems* (3) *and* (5) *have a spurious second-order critical point.*

241 The original problem of the non-existence of spurious second-order critical points can be viewed as
242 a property of the set of functions satisfying the RIP property, which is a convex set in an infinite-
243 dimensional functional space. The conditions in (6) reduce the infinite-dimensional problem to
244 a finite-dimensional problem by utilizing the optimality conditions and the RIP property, which
245 provides a basis of solving these conditions numerically. We note that the conditions in the third line
246 of (6) are novel and serve as an important step in developing strong theoretical guarantees. Although
247 the conditions in (6) seem complicated, they lead to strong results on the non-existence of spurious
248 second-order critical points. We provide two corollaries below to illustrate the power of the above
249 theorem. The first corollary focuses on the rank-1 case. In this case, we can simplify condition (6)
250 through suitable relaxations to obtain a sharper bound on δ that ensures the non-existence of spurious
251 second-order critical points.

252 **Corollary 1.** *Consider the case $r = 1$, and suppose that the function $f_a(\cdot)$ satisfies the δ-RIP$_{2,2}$*
253 *property with $\delta < 1/2$. Then, problems* (3) *and* (5) *have no spurious second-order critical points.*

254 The following example shows that the counterexample in Zhang et al. (2019) designed for the
255 symmetric case also works for the asymmetric rank-1 case.

256 **Example 1.** We note that Example 12 in Zhang et al. (2019) shows that problem (4) may have
257 spurious second-order critical points when $\delta = 1/2$. In general, a second-order critical point for
258 problem (4) is not a second-order critical point for problem (5), since the asymmetric manifold
259 $\mathcal{M}_{asym}$ has a larger second-order critical cone than the symmetric manifold $\mathcal{M}_{sym}$. However, it can
260 be verified that the same example also has a spurious second-order critical point in the asymmetric
261 case. For completeness, we verify the claim in the appendix.

262 It follows from Corollary 1 and Example 1 that the bound $1/2$ is the *sharpest* bound for the rank-1
263 asymmetric case. The next corollary provides a marginal improvement to the state-of-the-art result
264 for the general rank case, which derives the RIP bound $\delta < 1/3$ (Ha et al., 2020). In addition, we
265 prove that there exists a positive correlation between every second-order critical point and the global
266 minimum when $\delta < 1/2$.

267 **Corollary 2.** *Given an arbitrary r, suppose that the function $f_a(\cdot)$ satisfies the δ-RIP$_{2r,2r}$ property.*
268 *If $\delta \leq 1/3$, then both problems* (3) *and* (5) *have no spurious second-order critical points. In addition,*
269 *if $\delta \in [0, 1/2)$, then every second-order critical point $\tilde{M}$ has a positive correlation with the ground*
270 *truth M_a^*. Namely, there exists a universal function $C(\delta) : (0, 1/2) \mapsto (0, 1]$ such that*

$$\langle \tilde{M}, M_a^* \rangle \geq C(\delta) \cdot \|\tilde{M}\|_F \|M_a^*\|_F.$$

271 For the general rank-r case, we construct a counterexample with spurious second-order critical points
272 when $\delta = 1/2$.

273 **Example 2.** Let $n = m = 2r$. Now, we use the sufficiency part of Theorem 4 to construct a
274 counterexample. We choose

$$\delta := \frac{1}{2}, \quad \alpha := \frac{3}{5}, \quad \Sigma := \frac{1}{2}I_r, \quad \Lambda := \frac{3}{4}I_r, \quad A = B := 0_r, \quad C = D := I_r.$$

275 It can be verified that the conditions in (6) are satisfied and $CB^T = AD^T = 0$, which means that
276 there exists a function $f_a(\cdot)$ satisfying the δ-RIP$_{2r,2r}$ property for which problems (3) and (5) have
277 spurious second-order critical points. We also give a direct construction with linear measurements in
278 the appendix. This example illustrates that Theorem 4 can be used to systematically design instances
279 of the problem with spurious second-order critical points.

280 Before closing this section, we note that similar conditions can be obtained for the symmetric problem
281 (4). Although there exists a natural transformation of symmetric problems to asymmetric problems
282 (see the appendix), the approach requires the objective function $f_s(\cdot)$ to have the δ-RIP$_{4r,2r}$ property,
283 which provides sub-optimal RIP bounds compared to a direct analysis. We give the results of the
284 direct analysis below and omit the proof due to the similarity to the asymmetric case.

285 **Theorem 5.** *Let $\ell := \min\{n, 2r\}$. For a given $\delta \in [0, 1)$, there exists a function $f_s(\cdot)$ with the*
286 *δ-RIP$_{2r,2r}$ property such that problem* (4) *has a spurious second-order critical point only if there exist*

a constant $\alpha \in (1-\delta, (1+\delta)/2]$, a diagonal matrix $\Sigma \in \mathbb{R}^{r \times r}$, a diagonal matrix $\Lambda \in \mathbb{R}^{(\ell-r) \times (\ell-r)}$ and matrices $A \in \mathbb{R}^{r \times r}, C \in \mathbb{R}^{(\ell-r) \times r}$ such that

$$(1+\delta) \min_{1 \le i \le r} \Sigma_{ii} \ge \max_{1 \le i \le \ell-r} \Lambda_{ii}, \quad \Sigma \succ 0,$$

$$\langle \Lambda, CC^T \rangle = \alpha \left[\operatorname{tr}(\Sigma^2) - 2\langle \Sigma, AA^T \rangle + \|AA^T\|_F^2 + 2\|AC^T\|_F^2 + \|CC^T\|_F^2 \right], \quad (7)$$

$$\operatorname{tr}(\Lambda^2) \le \alpha^{-1}(2\alpha - 1 + \delta^2) \cdot \langle \Lambda, CC^T \rangle, \quad \langle \Lambda, CC^T \rangle \ne 0.$$

If $AC^T = 0$, then there exists a function $f_s(\cdot)$ with the δ-RIP$_{2r,2r}$ property for which problem (4) *has a spurious second-order critical point.*

Compared to Theorem 4, the diagonal matrix Λ is not enforced to be positive semi-definite. The reason is that the eigenvalue decomposition is used instead of the singular value decomposition in the symmetric case, and therefore some eigenvalues can be negative. Similarly, we can obtain the non-existence and the positive correlation results for the symmetric problem.

Corollary 3. *If function $f_s(\cdot)$ satisfies the δ-RIP$_{2r,2r}$ property, then the following statements hold:*

- *If $\delta \le 1/3$, then there are no spurious second-order critical points;*

- *If $\delta < 1/2$, then there exists a positive correlation between every second-order critical point and the ground truth;*

- *If $\delta = 1/2$, then there exists a counterexample with spurious second-order critical points;*

- *If $\delta < 1/2$ and $r = 1$, then there are no spurious second-order critical points.*

We note that the last statement serves as a generalization of the results in Zhang et al. (2019) to the nonlinear measurement case, and improves upon the bound in Bi & Lavaei (2021) by dropping the BDP constant.

4 Global Landscape: Strict Saddle Property

Although the non-existence of spurious second-order critical points can ensure the global convergence under certain regularity conditions, it cannot guarantee a fast convergence rate in general. Saddle-point escaping algorithms may become stuck at approximate second-order critical points for exponentially long time. To guarantee the global polynomial-time convergence, the following strict saddle property is commonly considered in the literature:

Definition 3 (Sun et al. (2018)). Consider an arbitrary optimization problem $\min_{x \in \mathcal{X} \subset \mathbb{R}^d} F(x)$ and let $\mathcal{X}^*$ denote the set of its global minima. It is said that the problem satisfies the (α, β, γ)-**strict saddle property** for $\alpha, \beta, \gamma > 0$ if at least one of the following conditions is satisfied for every $x \in \mathcal{X}$:

$$\operatorname{dist}(x, \mathcal{X}^*) \le \alpha; \quad \|\nabla F(x)\|_F \ge \beta; \quad \lambda_{min}[\nabla^2 F(x)] \le -\gamma.$$

For the low-rank problems, we choose the distance to be the Frobenius norm in the factorization space. This distance is equivalent to the Frobenius norm in the matrix space in the sense that there exist constants $c_1(\mathcal{X}^*) > 0$ and $c_2(\mathcal{X}^*) > 0$ such that

$$c_1(\mathcal{X}^*) \cdot \|U - U^*\|_F \le \|UU^T - U^*(U^*)^T\|_F \le c_2(\mathcal{X}^*) \cdot \|U - U^*\|_F$$

holds for all $U \in \mathcal{X}$ as long as $\|U - U^*\|_F$ is small (Tu et al., 2016). A similar relation holds for the asymmetric case.

In Jin et al. (2017), it has been proved that the perturbed gradient decent method can find an ϵ-approximate second-order critical point in $\tilde{O}(\epsilon^{-2})$ iterations with high probability if the Hessian of the objective function is Lipschitz. Namely, the algorithm returns a point $x \in \mathcal{X}$ such that

$$\|\nabla F(x)\|_F \le O(\epsilon), \quad \lambda_{min}[\nabla^2 F(x)] \ge -O(\sqrt{\epsilon})$$

in $\tilde{O}(\epsilon^{-2})$ iterations with high probability. If we choose $\epsilon > 0$ to be small enough such that $O(\epsilon) < \beta$ and $-O(\sqrt{\epsilon}) > -\gamma$, then the strict saddle property ensures that the returned point satisfies $\operatorname{dist}(x, \mathcal{X}^*) \le \alpha$ with high probability. We note that the Lipschitz continuity of the Hessian can be similarly guaranteed by the boundedness of trajectories of the perturbed gradient method, which can be proved similarly as Theorem 8 in Jin et al. (2017). Since the smoothness properties are satisfied

over a bounded region, we may apply the perturbed gradient descent method (Jin et al., 2017) to achieve the polynomial-time global convergence with random initialization.

In this section, we prove that problems (4) and (5) satisfy the strict saddle property with an arbitrary $\alpha > 0$ in the exact parameterization case, i.e., when the global optimum has rank r.

Assumption 2. The global optimum M_a^* or M_s^* has rank r.

It has been proved in Zhu et al. (2021) that the regularized problem (5) satisfies the strict saddle property if the function $f_a(\cdot)$ has the δ-RIP$_{2r,4r}$ property with

$$\delta < \frac{\sigma_r(M_a^*)^{3/2}}{100\|M_a^*\|_F\|M_a^*\|_2^{1/2}}.$$

Our results improve upon their bounds by allowing a larger problem-free RIP constant and requiring only the RIP$_{2r,2r}$ property (note that there are problems with RIP$_{2r,2r}$ property for which the RIP$_{2r,4r}$ property does not hold (Bi & Lavaei, 2021)). Our result can also be viewed as a robust version of the results in Ha et al. (2020).

Theorem 6. *Suppose that the function $f_a(\cdot)$ satisfies the δ-RIP$_{2r,2r}$ property with $\delta < 1/3$. Given an arbitrary constant $\alpha > 0$, if μ is selected to belong to the interval $[(1-\delta)/3, 1-\delta)$, then there exist positive constants*

$$\epsilon_1 := \epsilon_1(\delta, r, \mu, \sigma_r(M_a^*), \|M_a^*\|_F, \alpha), \quad \lambda_1 := \lambda_1(\delta, r, \mu, \sigma_r(M_a^*), \|M_a^*\|_F, \alpha)$$

such that for every $\epsilon \in (0, \epsilon_1]$ and $\lambda \in (0, \lambda_1]$, problem (5) satisfies the (α, β, γ)-strict saddle property with

$$\beta := \min\left\{\mu(\epsilon/r)^{3/2}, \lambda\right\}, \quad \gamma := \mu\epsilon.$$

We note that the constraint $\mu \in [(1-\delta)/3, 1-\delta)$ is not optimal and it can be similarly proved that $\mu \in (\delta, 1-\delta)$ also guarantees the strict saddle property. The key step in the proof is to show that for every point (U, V) at which the gradient of $f_a(UV^T)$ is small, it holds that

$$\|\nabla f_a(UV^T)\|_2 \geq (1+\delta)\sigma_r(UV^T) + C \cdot (1-3\delta)[f_a(UV^T) - f_a(M_a^*)],$$

where $C > 0$ is a constant independent of (U, V). This inequality can be viewed as a major extension of the non-existence of spurious second-order critical points when $\delta < 1/3$ (Ha et al., 2020), which shows that every spurious second-order critical point (U, V) satisfies

$$\|\nabla f_a(UV^T)\|_2 > (1+\delta)\sigma_r(UV^T).$$

We emphasize that our proof requires a new framework and is not a standard revision of the existing methods, which is the reason why sharper bounds can be established. By replacing $\|\nabla f_a(M)\|_2$ with $-\lambda_{min}(\nabla f_s(M))$, the analysis for the asymmetric case can be extended to the symmetric case with minor modifications and the same bound follows.

Theorem 7. *Suppose that the function $f_s(\cdot)$ satisfies the δ-RIP$_{2r,2r}$ property with $\delta < 1/3$. Given an arbitrary constant $\alpha > 0$, there exists a positive constant $\lambda_1 := \lambda_1(\delta, r, \sigma_r(M_s^*), \|M_s^*\|_F, \alpha)$ such that for every $\lambda \in (0, \lambda_1]$, problem (4) satisfies the (α, β, γ)-strict saddle property with*

$$\beta := \lambda, \quad \gamma := 2\lambda.$$

The above bound is the first theoretical guarantee of the strict saddle property for the nonlinear symmetric problem.

5 Conclusion

In this work, we analyze the geometric properties of low-rank optimization problems via the non-convex factorization approach. We prove novel necessary conditions and sufficient conditions for the non-existence of spurious second-order critical points in both symmetric and asymmetric cases. We show that these conditions lead to sharper bounds and greatly simplify the construction of counterexamples needed to study the sharpness of the bounds. The developed bounds significantly generalize several of the existing results. In the rank-1 case, the bound is proved to be the sharpest possible. In the general rank case, we show that there exists a positive correlation between second-order critical points and the global minimum for problems whose RIP constants are higher than the developed bound but lower than the fundamental limit obtained by the counterexamples. Finally, the strict saddle property is proved with a weaker requirement on the RIP constant for asymmetric problems. The paper develops the first strict saddle property in the literature for nonlinear symmetric problems.

References

Hedy Attouch, Jérôme Bolte, and Benar Fux Svaiter. Convergence of descent methods for semi-algebraic and tame problems: proximal algorithms, forward–backward splitting, and regularized Gauss–Seidel methods. *Mathematical Programming*, 137(1):91–129, 2013.

Srinadh Bhojanapalli, Anastasios Kyrillidis, and Sujay Sanghavi. Dropping convexity for faster semi-definite optimization. In *Conference on Learning Theory*, pp. 530–582. PMLR, 2016a.

Srinadh Bhojanapalli, Behnam Neyshabur, and Nathan Srebro. Global optimality of local search for low rank matrix recovery. In *Proceedings of the 30th International Conference on Neural Information Processing Systems*, pp. 3880–3888, 2016b.

Yingjie Bi and Javad Lavaei. On the absence of spurious local minima in nonlinear low-rank matrix recovery problems. In *International Conference on Artificial Intelligence and Statistics*, pp. 379–387. PMLR, 2021.

Nicolas Boumal. Nonconvex phase synchronization. *SIAM Journal on Optimization*, 26(4):2355–2377, 2016.

Samuel Burer and Renato DC Monteiro. A nonlinear programming algorithm for solving semidefinite programs via low-rank factorization. *Mathematical Programming*, 95(2):329–357, 2003.

Emmanuel J Candès and Benjamin Recht. Exact matrix completion via convex optimization. *Foundations of Computational mathematics*, 9(6):717–772, 2009.

Emmanuel J Candès and Terence Tao. The power of convex relaxation: Near-optimal matrix completion. *IEEE Transactions on Information Theory*, 56(5):2053–2080, 2010.

Yudong Chen and Yuejie Chi. Harnessing structures in big data via guaranteed low-rank matrix estimation: Recent theory and fast algorithms via convex and nonconvex optimization. *IEEE Signal Processing Magazine*, 35(4):14–31, 2018.

Yuejie Chi, Yue M Lu, and Yuxin Chen. Nonconvex optimization meets low-rank matrix factorization: An overview. *IEEE Transactions on Signal Processing*, 67(20):5239–5269, 2019.

Mark A Davenport, Yaniv Plan, Ewout Van Den Berg, and Mary Wootters. 1-bit matrix completion. *Information and Inference: A Journal of the IMA*, 3(3):189–223, 2014.

Rong Ge, Jason D Lee, and Tengyu Ma. Matrix completion has no spurious local minimum. *Advances in Neural Information Processing Systems*, pp. 2981–2989, 2016.

Rong Ge, Chi Jin, and Yi Zheng. No spurious local minima in nonconvex low rank problems: A unified geometric analysis. In *International Conference on Machine Learning*, pp. 1233–1242. PMLR, 2017.

Wooseok Ha, Haoyang Liu, and Rina Foygel Barber. An equivalence between critical points for rank constraints versus low-rank factorizations. *SIAM Journal on Optimization*, 30(4):2927–2955, 2020.

Zhishen Huang and Stephen Becker. Perturbed proximal descent to escape saddle points for non-convex and non-smooth objective functions. In *INNS Big Data and Deep Learning conference*, pp. 58–77. Springer, 2019.

Prateek Jain, Raghu Meka, and Inderjit Dhillon. Guaranteed rank minimization via singular value projection. In *Proceedings of the 23rd International Conference on Neural Information Processing Systems-Volume 1*, pp. 937–945, 2010.

Chi Jin, Rong Ge, Praneeth Netrapalli, Sham M Kakade, and Michael I Jordan. How to escape saddle points efficiently. In *International Conference on Machine Learning*, pp. 1724–1732. PMLR, 2017.

Chi Jin, Praneeth Netrapalli, and Michael I Jordan. Accelerated gradient descent escapes saddle points faster than gradient descent. In *Conference On Learning Theory*, pp. 1042–1085. PMLR, 2018.

Jason D Lee, Max Simchowitz, Michael I Jordan, and Benjamin Recht. Gradient descent only converges to minimizers. In *Conference on learning theory*, pp. 1246–1257. PMLR, 2016.

Qiuwei Li, Zhihui Zhu, and Gongguo Tang. The non-convex geometry of low-rank matrix optimization. *Information and Inference: A Journal of the IMA*, 8(1):51–96, 2019.

Ioannis Panageas and Georgios Piliouras. Gradient descent only converges to minimizers: Non-isolated critical points and invariant regions. *arXiv preprint arXiv:1605.00405*, 2016.

Dohyung Park, Anastasios Kyrillidis, Constantine Caramanis, and Sujay Sanghavi. Finding low-rank solutions via nonconvex matrix factorization, efficiently and provably. *SIAM Journal on Imaging Sciences*, 11(4):2165–2204, 2018.

Benjamin Recht, Maryam Fazel, and Pablo A Parrilo. Guaranteed minimum-rank solutions of linear matrix equations via nuclear norm minimization. *SIAM review*, 52(3):471–501, 2010.

Yoav Shechtman, Yonina C Eldar, Oren Cohen, Henry Nicholas Chapman, Jianwei Miao, and Mordechai Segev. Phase retrieval with application to optical imaging: a contemporary overview. *IEEE signal processing magazine*, 32(3):87–109, 2015.

Amit Singer. Angular synchronization by eigenvectors and semidefinite programming. *Applied and computational harmonic analysis*, 30(1):20–36, 2011.

Ju Sun, Qing Qu, and John Wright. A geometric analysis of phase retrieval. *Foundations of Computational Mathematics*, 18(5):1131–1198, 2018.

Stephen Tu, Ross Boczar, Max Simchowitz, Mahdi Soltanolkotabi, and Ben Recht. Low-rank solutions of linear matrix equations via Procrustes flow. In *International Conference on Machine Learning*, pp. 964–973. PMLR, 2016.

Lingxiao Wang, Xiao Zhang, and Quanquan Gu. A unified computational and statistical framework for nonconvex low-rank matrix estimation. In *Artificial Intelligence and Statistics*, pp. 981–990. PMLR, 2017.

Richard Y Zhang, Cédric Josz, Somayeh Sojoudi, and Javad Lavaei. How much restricted isometry is needed in nonconvex matrix recovery? In *NeurIPS*, 2018.

Richard Y Zhang, Somayeh Sojoudi, and Javad Lavaei. Sharp restricted isometry bounds for the inexistence of spurious local minima in nonconvex matrix recovery. *Journal of Machine Learning Research*, 20(114):1–34, 2019.

Zhihui Zhu, Qiuwei Li, Gongguo Tang, and Michael B Wakin. Global optimality in low-rank matrix optimization. *IEEE Transactions on Signal Processing*, 66(13):3614–3628, 2018.

Zhihui Zhu, Qiuwei Li, Gongguo Tang, and Michael B Wakin. The global optimization geometry of low-rank matrix optimization. *IEEE Transactions on Information Theory*, 67(2):1308–1331, 2021.