The price of unfairness in linear bandits with biased feedback

Solenne Gaucher
Département de Mathématiques d'Orsay, Université Paris-Saclay, Orsay, France
`solenne.gaucher@math.u-psud.fr`

Alexandra Carpentier
Institut für Mathematik, Universität Potsdam, Potsdam, Germany
`carpentier@uni-potsdam.de`

Christophe Giraud
Département de Mathématiques d'Orsay, Université Paris-Saclay, Orsay, France
`christophe.giraud@universite-paris-saclay.fr`

Abstract

In this paper, we study the problem of fair sequential decision making with biased linear bandit feedback. At each round, a player selects an action described by a covariate and by a sensitive attribute. The perceived reward is a linear combination of the covariates of the chosen action, but the player only observes a biased evaluation of this reward, depending on the sensitive attribute. To characterize the difficulty of this problem, we design a phased elimination algorithm that corrects the unfair evaluations, and establish upper bounds on its regret. We show that the worst-case regret is smaller than $\mathcal{O}(\kappa_*^{1/3} \log(T)^{1/3} T^{2/3})$, where κ_* is an explicit geometrical constant characterizing the difficulty of bias estimation. We prove lower bounds on the worst-case regret for some sets of actions showing that this rate is tight up to a possible sub-logarithmic factor. We also derive gap-dependent upper bounds on the regret, and matching lower bounds for some problem instance. Interestingly, these results reveal a transition between a regime where the problem is as difficult as its unbiased counterpart, and a regime where it can be much harder.

1 Introduction

Artificial intelligence is increasingly used in a wide range of decision making scenarii with higher and higher stakes, with application in online advertisement [31], credit [3], health care [11], education [28] and job interviews [35], in the hope of improving accuracy and efficiency. Recent works have shown that the decisions made by algorithms can be dangerously biased against certain categories of people, and have endeavored to mitigate this behavior [22, 14, 6, 27]. Studies have underlined that the main cause of algorithmic unfairness is the presence of bias in the training set [27], which led to the development of methods aiming to guarantee the fairness of the algorithms. This paper, in lines with these works, addresses the problem of online decision making under biased feedback.

Linear bandits have become a very popular tool in online decision making problems, when side information on the actions is available in the form of covariates. In the present paper, we consider a variant of this problem, where the agent only has access to an unfair assessment of the action taken, that is systematically biased against a group of actions. For example, examiners may be prejudiced against people from a minority group, and give them lower grades; similarly, algorithms trained on

36th Conference on Neural Information Processing Systems (NeurIPS 2022).

biased data may produce unfair assessments of the credit risk of individuals belonging to a minority group. Note that not correcting biased evaluation can have adverse effects for all parties: on the one hand, actions disadvantaged by the evaluation mechanism will be unfairly discriminated against; on the other hand, the agent may spend his budget on an unfairly advantaged action that is actually sub-optimal. The problem of sequential decision making under biased feedback can be formalized as follows.

Biased linear bandit problem   A player is presented with a set of k distinct actions characterized by covariates $x \in \mathcal{X} \subset \mathbb{R}^d$, and by known sensitive attributes $z_x \in \{-1, 1\}$ indicating the group of the action. For the sake of clarity, we consider here a two-group model (respectively privileged or discriminated against), and we defer to Appendix D discussions on how to extend this model and our algorithm to more than two groups. At each round $t \leq T$, the player chooses the action x_t and receives an unobserved reward $x_t^\top \gamma^*$, where $\gamma^* \in \mathbb{R}^d$ is the regression parameter specifying the true value of the action. The regret of the player is given by

$$R_T = \mathbb{E}\Big[\sum_{t \leq T} (x^* - x_t)^\top \gamma^* \Big], \quad \text{where} \quad x^* \in \operatorname*{argmax}_{x \in \mathcal{X}} x^\top \gamma^*. \tag{1}$$

By contrast to the classical linear bandit, the player does not observe a noisy version of the unbiased reward $x_t^\top \gamma^*$. Instead, she observes an unfair evaluation y_t of the value of the action $x_t^\top \gamma^*$, given by the following biased linear model:

$$y_t = x_t^\top \gamma^* + z_{x_t} \omega^* + \xi_t$$

 where $\xi_t \overset{i.i.d}{\sim} \mathcal{N}(0, 1)$ is a noise term. The evaluation are systematically biased against a certain group: this unequal treatment of the groups is captured by the bias parameter $\omega^* \in \mathbb{R}$.

Preliminary discussion   The biased linear bandit is a variant of the linear bandit. By contrast, in the classical linear bandit model, the agent observes a noisy version of the reward. Obviously, applying directly an algorithm designed for linear bandit to biased linear bandits without correcting the evaluations would lead to a linear regret if the evaluation mechanism is prejudiced against the group of the best action in terms of reward, and if the best action in terms of feedback belongs to the advantaged group. To avoid this pitfall, one must estimate the bias in order to correct the evaluations. This implies a change in the exploration-exploitation trade-off, as exploration becomes more expensive. Indeed, in classical bandit problems, one can compare the rewards of two actions by repeatedly sampling them - or, to put it differently, one can find the best action by sampling only those actions that seem optimal. This does not hold in the biased linear bandit: if, at some point, the set of potentially optimal actions contains representatives from both groups, and does not span $\mathbb{R}^d$, one is forced to sample sub-optimal actions to estimate the bias and improve the estimation of the unbiased rewards. For this reason, classical algorithm for linear bandit that only sample actions considered as potentially optimal, such as OFUL [1] or Phase Elimination [24], can suffer linear regret. This underlines the necessity to ensure sufficient estimation of the bias parameter, even when it implies sampling sub-optimal actions.

1.1   Related work

Fairness in bandit problems has mostly been studied from the perspective of fair budget allocation between actions. This problem is motivated by the fact that classical bandit algorithms select sub-optimal actions only a vanishing fraction of the time, which may be undesirable in many situations. To mitigate this problem and guarantee diversity in the actions selected, some papers [4, 29, 9, 16, 43] have proposed new algorithms ensuring fairness of the selection frequency of each action. The framework studied in this paper is different: we consider here that the mechanism for observing the rewards is unfair, and we aim at correcting it in order to maximize a (fair) true cumulative reward.

In this work, we consider that the agent knows the sensible attributes, and that she can treat actions differently according to their sensible attributes, in order to correct the prejudice caused by the unfair bias in the evaluation. This situation falls into the awareness framework, by contrast to the unawareness one, where using the sensitive attributes is prohibited. Whether or not it is preferable to treat different groups differently remains a controversial question. While using sensitive attributes at the time of prediction is sometimes forbidden by law, some recent works have highlighted critical

issues related to unawareness. For example, empirical evidence [26] have shown that classification algorithms based on disparate learning processes use non-sensitive features correlated with the sensitive attribute as a proxy for the later. These empirical findings have recently been supported by theoretical results established in [15] in the case of demographic parity. Similarly, the authors of [7] study a problem of fair online learning, and show that some problems feasible in the awareness framework become infeasible in the unawareness one (such as no-regret learning under demographic parity constraints). These examples advocate for the use of the sensitive attribute, as it allows for better fairness guarantees while preventing unfair discrimination based on (possibly irrelevant) non-sensitive features correlated with the sensitive attribute. Without taking a position in this debate, we underline that, in practice, this attribute (gender or minority status) is often known to the decision-maker, and that its use is in some cases allowed or even encouraged (e.g. for affirmative action).

By contrast to a line of work on statistical fairness, the aim of our model is not to correct for the possibly unequal distribution of features x and values $x^\top \gamma^*$ across the different groups. Our approach is instead related to causal fairness [18]: in the causal fairness framework, the dependencies between prediction, sensitive attributes and non-sensitive attributes are captured by a causal model. The goal is then to ensure that the sensitive attribute does not *directly* influence the prediction (in other words, that conditionally on selected resolving variables, the prediction is independent of the sensitive attribute). Here, the resolving variables may depend on the sensitive attribute in a manner that is considered as non-discriminatory. For example, one group may have, on average, more physical strength than the other one, and this skill can be considered as fair when it comes to recruit a piano mover. The biased linear model studied in this paper is a simple example of causal model with linear structural model equations $x = f(z, \xi')$ and $y = x^\top \gamma^* + z\omega^* + \xi$, where ξ and ξ' are noise terms: the covariates x may depend on the sensitive attribute z, and the biased evaluation y depends on both. In our work, we treat $x^\top \gamma^*$ as a fair evaluation of the value of action x, since it is independent of z conditionally on the resolving variable x.

The biased linear model has been studied in the batch setting in [8], where the authors investigate the optimal trade-off between minimax risk and Demographic Parity. Detection of systematic bias, interpreted as a treatment effect, has been investigated in a batch setting in [18]. In [2], the authors consider a similar model, with unobserved sensitive attribute z and known bias parameter ω^*, under additional assumption that the sensitive attribute z is independent from the covariate x. By contrast, we show that bias estimation is one of the main difficulties of the biased bandit problem.

The linear bandit with biased feedback can be viewed as a stochastic partial monitoring game. With the terminology of partial monitoring, the biased problem considered in the present paper is globally observable but not locally observable: in this case, the optimal worst-case regret rate typically increases as $\tilde{O}(T^{2/3})$. This regret rate is for example achieved in the related problem of partial linear monitoring with linear feedback and linear reward using an Information Directed Sampling algorithm [20]. However, the dependence of the regret on the geometry of the action set and on the dimension d remains in most cases an open question [25, 5, 20]. In this paper, we characterize the geometry of the biased linear bandit problem, and we investigate dependence of the regret on the gaps.

1.2 Contribution and outline

In this paper, we introduce the linear bandit problem with biased feedback. We design a new algorithm based on optimal design for this problem. We derive an upper bound on the worst case regret of this algorithm of order $\kappa_*^{1/3} \log(T)^{1/3} T^{2/3}$ for large T, where κ_* is an explicit constant depending on the geometry of the action set. We provide matching lower bounds on some problem instances, showing that the constant κ_* characterizes the difficulty of the action set. Note that this regret is higher than the classical rates of order $\tilde{O}(dT^{1/2})$ obtained for d-dimensional linear bandits: this increase corresponds to the price to pay for debiasing the unfair evaluations.

We also characterize the gap-depend regret, showing that it is of order $\left(d/\Delta_{\min} \vee \kappa(\Delta)/\Delta_{\neq}^2\right)\log(T)$, where $\Delta_{\min}$ is the minimum gap, $\Delta_{\neq}$ is the gap between the best actions of the two groups, and $\kappa(\Delta)$ corresponds to the minimum regret to pay for estimating the bias with a given variance. This bound underlines the relative difficulties of the d-dimensional linear bandit and of the bias estimation. When $d/\Delta_{\min} \geq \kappa(\Delta)/\Delta_{\neq}^2$, i.e. when one group contains all near-optimal actions, the difficulty is dominated by that of the corresponding linear bandit problem. When both groups contain near-optimal actions, and $d/\Delta_{\min} \leq \kappa(\Delta)/\Delta_{\neq}^2$, the regret corresponds to the price of debiasing the rewards.

The rest of the paper is organized as follows. In Section 2, we present the FAIR PHASED ELIMI-NATION algorithm: we first discuss parameter estimation in Section 2.1, before presenting a sketch of the algorithm in Section 2.2 (a detailed version of this algorithm is provided in Appendix B). Then, in Section 3, we establish an upper bound on its worst-case regret. In Section 4, we derive a gap-dependent upper bound on the regret of our algorithm. In Section 5, we establish lower bounds on some action sets for both the worst-case and the gap-dependent regret, showing that these rates are sharp respectively up to a sub-logarithmic factor and an absolute multiplicative constant. Additional discussions on the geometry of bias estimation are postponed to Appendix A.

1.3 Notations and additional assumptions

We assume that all covariates $x \in \mathcal{X}$ are distinct, which implies that the group z_x of action x is well defined. We also assume that no group is empty, that the set $\left\{ \begin{pmatrix} x \\ z_x \end{pmatrix} : x \in \mathcal{X} \right\}$ spans $\mathbb{R}^{d+1}$ (which guarantees identifiability of the parameters), and that the rewards are bounded: $\max_{x \in \mathcal{X}} |x^\top \gamma^*| \leq 1$.

When necessary, we underline the dependence of the regret on the parameter θ by denoting it R_T^θ. We denote by $a_x = \begin{pmatrix} x \\ z_x \end{pmatrix}$ the vector describing an action and its group, by $\theta^* = \begin{pmatrix} \gamma^* \\ \omega^* \end{pmatrix} \in \mathbb{R}^{d+1}$ the unknown parameter, and by $\mathcal{A} = \{a_x : x \in \mathcal{X}\}$ the set of actions and of corresponding sensitive attributes. We denote by $\Delta = (\Delta_x)_{x \in \mathcal{X}}$ the vector of gaps $\Delta_x = \max_{x' \in \mathcal{X}} (x' - x)^\top \gamma^*$, and by $\mathcal{C}(\mathcal{X}) = \left\{ \gamma \in \mathbb{R}^d : \forall x \in \mathcal{X}, |x^\top \gamma| \leq 1 \right\}$ the set of admissible parameters. Note that for all $x \in \mathcal{C}(\mathcal{X})$, $\Delta_x \leq 2$. For $i \leq d+1$, let e_i be the i-th vector of the canonical basis of $\mathbb{R}^{d+1}$, and for any matrix M, let M^+ be a generalized inverse of M. We denote by $\mathcal{P}^\mathcal{X}$ the set of probability measures on $\mathcal{X}$, and $\mathcal{M}^\mathcal{X} = \{\mu : \mathcal{X} \mapsto \mathbb{R}_+\}$. For any $\mu \in \mathcal{P}^\mathcal{X}$ or $\mu \in \mathcal{M}^\mathcal{X}$, we denote $V(\mu) = \sum_{x \in \mathcal{X}} \mu(x) a_x a_x^\top$ the covariance matrix corresponding to this allocation. For $u \in \mathbb{R}^{d+1}$ (resp. $\mathcal{U} \in \mathbb{R}^{d+1}$), we denote by $\mathcal{P}_u^\mathcal{X}$ (resp. $\mathcal{M}_u^\mathcal{X}$) the measures μ in $\mathcal{P}^\mathcal{X}$ (resp. in $\mathcal{M}^\mathcal{X}$) such that $u \in \text{Range}(V(\mu))$. For $\mathcal{U} \subset \mathbb{R}^{d+1}$, we denote by $\mathcal{P}_\mathcal{U}^\mathcal{X}$ (resp. $\mathcal{M}_\mathcal{U}^\mathcal{X}$) the measures μ such that $\mu \in \mathcal{P}_u^\mathcal{X}$ (resp. $\mathcal{M}_u^\mathcal{X}$) for all $u \in \mathcal{U}$.

2 Fair Phased Elimination algorithm

The Fair Phased Elimination algorithm belongs to the category of sequential elimination algorithms. Classical sequential elimination algorithms typically proceed by phases, indexed by $l = 1, 2, \ldots$. At phase l, these algorithms consider a set of potentially optimal actions $\mathcal{X}_l$. The rewards of all actions $x \in \mathcal{X}_l$ are then estimated with a given precision $O(\epsilon_l)$, typically chosen as $\epsilon_l = 2^{2-l}$, by sampling actions in $\mathcal{X}_l$. Actions sub-optimal by a gap larger than the precision level are then removed from the set $\mathcal{X}_{l+1}$ of potentially optimal actions for the phase $l+1$.

As underlined previously, classical sequential elimination algorithms may suffer linear regret in the biased linear bandit problem if actions allowing to estimate the bias are discarded by the algorithm before the best group is identified (this happens for example if at a phase l, less that $d+1$ action remains, with at least one action in each group). To mitigate this problem, we first estimate the biased evaluations of the potentially optimal actions, using ordinary least squares estimation. We then debias the estimations using an estimator for the bias relying on independent observations, which may be obtained by sampling sub-optimal actions. Before presenting the algorithm, let us discuss the estimation of the evaluations and of the bias parameter.

2.1 Optimal design for parameter estimation in the biased linear bandit

G-optimal design for biased evaluation estimation   As in the Phased Elimination algorithm [24], we rely on G-optimal design to estimate the biased evaluations $a_x^\top \theta^*$ with small error uniformly over a set of actions $\mathcal{X}_l$. More precisely, for a given set of potentially optimal actions $\mathcal{X}_l$, we compute the G-optimal design solution to the problem

$$\underset{\pi \in \mathcal{P}_{\mathcal{X}_l}^{\mathcal{X}_l}}{\text{minimize}} \max_{x \in \mathcal{X}_l} a_x^\top (V(\pi))^+ a_x . \qquad \text{(G-optimal design)} \qquad (2)$$

This can be done using polynomial-time algorithms, relying for example on interior points method [40], or on mixed integer second-order cone programming [37]. The celebrated General Equivalence theorem of Kiefer [19] and Pukelsheim [33] states that the value of Equation (2) is bounded by $d+1$.

Let π^* denote any design solution to the G-optimal design problem (2), and let $\widehat{\theta}$ denote the ordinary least square estimator obtained by sampling each action $x \in \mathcal{X}_l$ exactly $\lceil n\pi^*(x) \rceil$ times for a given $n > 0$. Then, for all $x \in \mathcal{X}_l$, the General Equivalence theorem implies that the variance of the estimate $a_x^\top \widehat{\theta}$ is smaller than $(d+1)/n$. Moreover, the G-optimal design π^* can be chosen so that it is supported by at most $(d+1)(d+2)/2$ points, so the total number of samples is at most $n + (d+1)(d+2)/2$.

Δ-optimal design for bias evaluation   In this paragraph, we introduce the Δ-optimal design, which is discussed in greater depth in Appendix A. Δ-optimal design aims at estimating a parameter with a given accuracy and with minimal regret. Similar ideas have recently been used in [42] to solve classical linear bandit problems. To estimate the bias parameter ω^*, we use the estimator $\widehat{\omega} = e_{d+1}^\top \widehat{\theta}$, where $\widehat{\theta}$ is the ordinary least square estimator for the full parameter θ^*. Now, if we sample each action $x \in \mathcal{X}$ exactly $\mu(x)$ time, the variance of $\widehat{\omega}$ is equal to $e_{d+1}^\top V(\mu)^+ e_{d+1}$. Given the vector of gaps Δ, the design μ minimizing the regret of this exploration phase, while ensuring that the variance of $\widehat{\omega}$ is smaller than 1, is solution of the problem

$$\underset{\mu \in \mathcal{M}_{\mathcal{X}}^{e_{d+1}}}{\text{minimize}} \sum_x \mu(x)\Delta_x \quad \text{such that} \quad e_{d+1}^\top V(\mu)^+ e_{d+1} \leq 1. \quad (\Delta\text{-optimal design}) \quad (3)$$

In the following, we denote μ^Δ a minimizer of (3), and $\kappa(\Delta) = \sum_{x \in \mathcal{X}} \mu^\Delta(x)\Delta_x$. Lemma 9 in Appendix A explains how to compute the design μ^Δ in polynomial time by adapting tools from c-optimal design. This lemma also shows that the support of μ^Δ can be chosen to be of cardinality at most $d + 1$. Then, choosing each action exactly $\lceil n\mu^\Delta(x) \rceil$ times for a given $n > 0$ allows us to estimate the bias with variance lower than n^{-1} and a regret no larger than $n\kappa(\Delta) + 2(d + 1)$. Obviously, we do not know the gap vector Δ beforehand, so we must estimate it as we go.

2.2   Outline of the Fair Phased Elimination algorithm

The Fair Phased Elimination algorithm, sketched in Algorithm 3, relies on the following key ideas. First, note that within a group, the order of the true rewards and of the biased evaluations are the same. Hence, within a group, we can use classical algorithms for linear bandits to choose the actions and estimate the biased evaluations with a controlled within-group regret: this is done using **G-exploration and elimination**. Second, to compare actions belonging to different groups, we independently estimate the bias parameter ω^*, using **Δ-exploration and elimination**. Finally, we underline that bias estimation may require to sample very sub-optimal actions. Therefore, it can be overly costly to estimate the bias up to the precision level required to identify the best group. To prevent this, we use a **stopping criterion**.

G-exploration and elimination   At each phase $l = 1, 2, ...$, we keep two sets of potentially optimal actions belonging to the groups $+1$ and -1, denoted respectively $\mathcal{X}_l^{(+1)}$ and $\mathcal{X}_l^{(-1)}$. If we have not identified the group containing best action, we run a G-EXP-ELIM routine 1 on each set $\mathcal{X}_l^{(z)}$ for $z = 1$ and $z = -1$. This routine samples actions according to a rounded G-optimal design on $\mathcal{X}_l^{(z)}$, with a total number of observations chosen so that the biased evaluations of all actions in $\mathcal{X}_l^{(z)}$ are known with an error at most ϵ_l. The set $\mathcal{X}_{l+1}^{(z)}$ is obtained by removing from $\mathcal{X}_l^{(z)}$ actions whose estimated evaluations are sub-optimal by a gap larger than $3\epsilon_l$, compared to the empirical best action in the group. This allows to ensure that only actions sub-optimal by a gap $\mathcal{O}(\epsilon_l)$ remain in $\mathcal{X}_{l+1}^{(z)}$, and to estimate the gap vector Δ with a precision sufficient for Δ-optimal estimation.

If the group containing the best action has been identified, we discard the other group, and run a G-EXP-ELIM routine 1 on the set of potentially optimal actions in this group.

Δ-exploration and elimination   If the group of the best action has not been found before phase l, we run the Δ-EXP-ELIM routine 2. More precisely, relying on a previous estimate $\widehat{\Delta}^l$ of the gap vector Δ, we compute the $\widehat{\Delta}^l$-optimal design $\widehat{\mu}$. We then estimate the bias using actions sampled according to a rounded version of this design, with a total number of observations chosen so that the error of bias estimation is smaller than ϵ_l, and use it to debias the reward estimation. If the debiased evaluation of the best action of each group are separated by a gap larger than $4\epsilon_l$, we consider that the best group is the one containing the empirical best action in terms of biased evaluation, and we discard the other group.

Routine 1 G-EXP-ELIM $(\mathcal{X}, n, \epsilon)$

1: Compute G-optimal design π solution of (2) on $\mathcal{X}$, with $|\operatorname{supp}(\pi)| \leq (d+1)(d+2)/2$
2: Sample $\lceil n\pi(x) \rceil$ times each action a_x for $x \in \mathcal{X}$ ▷ G-optimal parameter estimation
3: Compute the ordinary least square estimator $\widehat{\theta}$
4: $\mathcal{X}' \leftarrow \left\{ x \in \mathcal{X} : \max_{x' \in \mathcal{X}} (x' - x)^\top \widehat{\theta} \leq 3\epsilon \right\}$ ▷ Suboptimal actions elimination
5: **return** $\widehat{\theta}$ and $\mathcal{X}'$

If we cannot find the best group, we rely on estimates of the bias and of the biased evaluations obtained during the previous round to update the estimate of the gap vector $\widehat{\Delta}^{l+1}$.

Routine 2 Δ-EXP-ELIM $(\mathcal{X}, (\mathcal{X}^{(z)}, \widehat{\theta}^{(z)})_{z \in \{-1,1\}}, \widehat{\Delta}, n, \epsilon)$

1: Compute $\widehat{\Delta}$-optimal design $(\hat{\mu}, \kappa(\hat{\Delta}))$ solution of (3) on $\mathcal{X}$, with $|\operatorname{supp}(\hat{\mu})| \leq d+1$
2: Sample $\lceil n\hat{\mu}(x) \rceil$ times each action a_x for $x \in \mathcal{X}$ ▷ $\widehat{\Delta}$-optimal bias estimation
3: Compute $\widehat{\omega} = e_{d+1}^\top \widehat{\theta}$, where $\widehat{\theta}$ is the ordinary least square estimator
4: **for** $z \in \{-1, 1\}$ and $x \in \mathcal{X}^{(z)}$ **do** $\widehat{m}_x \leftarrow a_x^\top \widehat{\theta}^{(z)} - z\widehat{\omega}$ ▷ Debiased rewards estimation
5: **if** $\exists z \in \{-1, 1\}$ such that $\max_{x \in \mathcal{X}^{(z)}} \widehat{m}_x \geq \max_{x \in \mathcal{X}^{(-z)}} \widehat{m}_x + 4\epsilon$ **then** $\mathcal{Z} \leftarrow \{z\}$ ▷ Group elimination
6: **else** $\widehat{\Delta}_x \leftarrow 2 \wedge (\max_{x' \in \mathcal{X}^{(-1)} \cup \mathcal{X}^{(1)}} \widehat{m}_{x'} - \widehat{m}_x + 4\epsilon)$ for all $x \in \mathcal{X}^{(-1)} \cup \mathcal{X}^{(1)}$
7: **return** $\mathcal{Z}$ and $\widehat{\Delta}$

Stopping criterion   As underlined previously, the Δ-EXP-ELIM routine samples actions that can be very sub-optimal. As a consequence, when the gap between the best two actions of each group is small, finding the best group can be overly costly in terms of regret. To prevent this, if the best group has not been found at stage l fulfilling $\epsilon_l \leq \left(\kappa(\widehat{\Delta}^l) \log(T)/T \right)^{1/3}$, the bias estimation is stopped and the empirical best action in $\mathcal{X}_{l+1}^{(1)} \cup \mathcal{X}_{l+1}^{(-1)}$ is sampled for the remaining time (see Algorithm 3)

Algorithm 3 FAIR PHASED ELIMINATION (sketched)

1: **input:** $\delta, T, \mathcal{X}, k = |\mathcal{X}|, \epsilon_l = 2^{2-l}$ for $l \geq 1$
2: **initialize:** $\mathcal{X}_1^{(+1)} \leftarrow \{x : z_x = 1\}, \mathcal{X}_1^{(-1)} \leftarrow \{x : z_x = -1\}$,
3:        $\mathcal{Z}_1 \leftarrow \{-1, +1\}, \widehat{\Delta}^1 \leftarrow (2, ..., 2), l \leftarrow 0$
4: **while** the budget is not spent **do** $l \leftarrow l + 1$
5:     **for** $z \in \mathcal{Z}_l$ **do**
6:          $\left(\widehat{\theta}^{(z)}, \mathcal{X}_{l+1}^{(z)} \right) \leftarrow$ G-EXP-ELIM $\left(\mathcal{X}_l^{(z)}, \frac{2(d+1)}{\epsilon_l^2} \log \left(\frac{kl(l+1)}{\delta} \right), \epsilon_l \right)$
7:     **if** $\mathcal{Z}_l = \{-1, +1\}$ **then**
8:         **if** $\epsilon_l \leq \left(\kappa(\widehat{\Delta}^l) \log(T)/T \right)^{1/3}$ **then**   ▷ Stop bias estimation
9:              Sample best action in $\mathcal{X}_{l+1}^{(-1)} \cup \mathcal{X}_{l+1}^{(+1)}$ for the remaining time
10:         **else**
11:              $\left(\mathcal{Z}_{l+1}, \widehat{\Delta}^{l+1} \right) \leftarrow \Delta$-EXP-ELIM $\left(\mathcal{X}, \left(\mathcal{X}_{l+1}^{(z)}, \widehat{\theta}_l^{(z)} \right)_{z \in \{-1,1\}}, \widehat{\Delta}^l, \frac{2}{\epsilon_l^2} \log \left(\frac{l(l+1)}{\delta} \right), \epsilon_l \right)$

---

3   Upper bound on the worst-case regret of FAIR PHASED ELIMINATION

The regret of the FAIR PHASED ELIMINATION depends on the difficulty of estimating the bias parameter, captured by $\kappa(\Delta)$. Lemma 7 in Appendix A.6 shows that for all parameter $\gamma^* \in \mathcal{X}$, $\kappa(\Delta)$ is upper bounded by $2\kappa_*$, where κ_* is the *minimal variance of the bias estimator* given by

$$\kappa_* = \min_{\pi \in \mathcal{P}_{e_{d+1}}^{\mathcal{X}}} e_{d+1}^\top (V(\pi))^+ e_{d+1}.$$

The following theorem provides a bound on the worst case regret depending on κ_*. Proofs are postponed to Appendix C.2.

Theorem 1. *For the choice $\delta = T^{-1}$, there exists two numerical constants $C, C' > 0$ such that the following bound on the regret of the* FAIR PHASED ELIMINATION *algorithm 4 holds*

$$R_T \leq C \left(\kappa_*^{1/3} T^{2/3} \log(T)^{1/3} + (d \vee \kappa_*) \log(T) + d^2 + d\kappa_*^{-1/3} T^{1/3} \log(kT) \log(T)^{-1/3} \right)$$

$$\leq C' \kappa_*^{1/3} T^{2/3} \log(T)^{1/3} \quad for \quad T \geq \frac{((d \vee \kappa_*)^{3/2} \log(T)) \vee d^3}{\sqrt{\kappa_*}} \vee \frac{(d \log(kT))^3}{(\kappa_* \log(T))^2}.$$

In Section 5.1, we show that the upper bound obtained in Theorem 1 is sharp in some settings, up to the sub-logarithmic factor $\log(T)^{1/3}$.

Theorem 1 shows that the worst-case regret of the Fair Phased Elimination algorithm asymptotically grows as $C\kappa_*^{1/3} T^{2/3} \log(T)^{1/3}$. This worst-case regret rate is higher than the typical rate $Cd \log(T)T^{1/2}$ obtained under unbiased feedback on the rewards (see, e.g., [1]). This increase in the regret corresponds to the cost of learning from unfair evaluations. It is due to the fact that the algorithm may need to sample actions that are sub-optimal in order to estimate the bias parameter. Note that this rate $\widetilde{\mathcal{O}}(T^{2/3})$ is typical for globally observable bandit problems with partial linear monitoring, and can be obtained by applying results established in [20] for in the partial linear monitoring setting to the biased linear bandit problem.

By contrast to previous results, Theorem 1 characterizes precisely the dependence of the worst-case regret on the geometry of the action set. The relevant constant κ_* is the minimal variance for estimating the bias, which appears when considering the related c-optimal design problem. While the connection between G-optimal design and the linear bandit problem has already been exploited, it is to the best of our knowledge the first time that c-optimal design is related to partial monitoring.

The constant κ_* corresponds to the minimum number of samples required for estimating the bias with a variance equal to 1 (up to rounding issues). Intuitively, if the actions are very correlated with their sensitive attributes, more samples will be needed to estimate the bias with the same precision. This situation corresponds to cases where κ_* is large, and leads to a higher regret. Lemma 1, illustrated in Figure 1, relates κ_* to the margin between the two groups of actions.

Lemma 1. *κ_* is the largest constant $\kappa \geq 0$ such that, there exists an hyperplane $\mathcal{H}$ containing zero and separating the two groups, and such that, the margin to $\mathcal{H}$ is at least $\sqrt{\kappa}-1/\sqrt{\kappa}+1$ times the maximum distance of all points to the hyperplane (see Figure 1). When no such hyperplane exists, then $\kappa_* = 1$.*

(a) The margin m is equal to $\sqrt{\kappa_*}-1/\sqrt{\kappa_*}+1$ times the maximum distance M of any action to the hyperplane.

(b) $\kappa_* = 1$: the groups cannot be separated by a hyperplane containing 0.

Figure 1: Interpretation of κ_* in terms of separation of the groups.

Interestingly, Lemma 1 underlines that under reasonable assumptions, the constant κ_* may not depend on the ambient dimension d, and it can even be equal to 1. By contrast, while the Information Directed Sampling algorithm can be applied to the biased linear bandit problem, the regret bounds established in [20] are of order $\alpha^{1/3} d^{1/2} T^{2/3} \log(kT)^{1/2}$, where α is a measure of the complexity of the action set called the worst-case alignment constant. Lemma 6 in Appendix A shows that α is equivalent to the minimal variance of the bias estimator κ_*. Hence, our bound improves over previous results by a factor $d^{1/2} \log(T)^{1/6} (\log(kT)/\log(T))^{1/2}$.

The gaps are not involved in the definition of the minimal variance of bias estimation κ_*. The reader may have expected to get, instead of κ_*, the minimax regret for estimating the bias

$$\widetilde{\kappa} = \max_{\gamma \in \mathcal{C}(\mathcal{X}), x' \in \mathcal{X}} \sum_{x \in \mathcal{X}} \widetilde{\mu}(x)(x'-x)^\top \gamma, \quad \text{where}$$

$$\widetilde{\mu} = \operatorname*{argmin}_{\mu} \max_{x' \in \mathcal{X}, \gamma \in \mathcal{C}(\mathcal{X})} \sum_{x \in \mathcal{X}} \mu(x)(x'-x)^\top \gamma, \text{ such that } \mu \in \mathcal{M}_{e_{d+1}}^{\mathcal{X}} \text{ and } e_{d+1}^\top V(\mu)^+ e_{d+1} \leq 1.$$

Next lemma shows that κ_* and $\widetilde{\kappa}$ are in equivalent up to a factor 2. We refer the interested reader to Appendix A, where further discussions on the geometry of bias estimation are postponed, due to space constraints.

Lemma 2. $\widetilde{\kappa}/2 \leq \kappa_* \leq 2\widetilde{\kappa}$.

4 Upper bound on the gap-depend regret of FAIR PHASED ELIMINATION

In this section, we provide an upper bound on the worst-case regret that depends on the gap between the two best actions, and on the gap between the best actions of the two groups. Compared to instance-dependent bounds, established in the linear bandit problem in [23, 21], gap-dependent bounds characterize the dependence of the regret on a small number of parameters. They are typically less sharp than instance-dependent bounds, but allow to better highlight the influence of the parameters on the difficulty of the problem. The bound established in the following theorem relates the difficulty of the biased linear bandit to that of bias estimation, and to that of the corresponding d-dimensional linear bandit. Proofs are postponed to Appendix C.2.

Theorem 2. *Assume that* $x^* \in \operatorname{argmax}_{x \in \mathcal{X}} x^\top \gamma^*$ *is unique. Then, there exists two numerical constants* $C, C' > 0$ *such that, for the choice* $\delta = T^{-1}$, *the following bound on the regret of the* FAIR PHASED ELIMINATION *algorithm 4 holds*

$$R_T \leq C\left(\left(\frac{d}{\Delta_{\min}} \vee \frac{\kappa(\Delta \vee \Delta_{\neq} \vee \varepsilon_T)}{\Delta_{\neq}^2}\right) \log(T) + d^2 + \frac{d}{\Delta_{\min}} \log(k)\right)$$

$$\leq C'\left(\frac{d}{\Delta_{\min}} \vee \frac{\kappa(\Delta \vee \Delta_{\neq} \vee \varepsilon_T)}{\Delta_{\neq}^2}\right) \log(T) \quad \text{for} \quad T \geq k \vee e^{d\Delta_{\min}}$$

where $\Delta_{\min} = \min_{x \in \mathcal{X} \backslash x^*} \Delta_x$, $\Delta_{\neq} = \min_{x \in \mathcal{X}: z_x = -z_{x^*}} \Delta_x$, *and* $\varepsilon_T = \left(\kappa_* \log(T)/T\right)^{1/3}$.

The term $d/\Delta_{\min} \vee \kappa(\Delta \vee \Delta_{\neq} \vee \varepsilon_T)/\Delta_{\neq}^2$ highlights the two sources of difficulty of the problem. On the one hand, the term $d/\Delta_{\min}$ is unavoidable: even if the algorithm knew beforehand the group containing the best action, it would still need to play a game of d-dimensional linear bandits in this group, and suffer, in the worst-case, the corresponding gap-dependent regret [1]. Note that lower bounds on gap-depend regret of classical linear bandits follow from considering a setting with one near-optimal action with gap $\Delta_{\min}$ in each of the d dimensions. Then, any algorithm needs to explore each dimension up to $\Delta_{\min}^{-2} \log(T)$ times in order to find the best action, but can do so by choosing the near-optimal actions, thus having a regret $\Delta_{\min}^{-1} \log(T)$ in each direction. By contrast, the term $\kappa(\Delta \vee \Delta_{\neq} \vee \varepsilon_T)/\Delta_{\neq}^2$ is characteristic of the biased linear bandit problem: it is due to the fact that the algorithm may need to sample very sub-optimal actions in order to find the group containing the best action. Indeed, to identify this group, one must estimate the bias with a precision $\Delta_{\neq}$, i.e. sample sub-optimal actions with average regret $\kappa(\Delta)$ approximately $\Delta_{\neq}^{-2} \log(T)$ times.

When $d/\Delta_{\min} \leq \kappa(\Delta \vee \Delta_{\neq} \vee \varepsilon_T)/\Delta_{\neq}^2$, the regret corresponds to the regret of this bias estimation phase. In other words, when both groups contain near-optimal actions, the difficulty of the problem is dominated by the price to pay for debiasing the unfair evaluations. Interestingly, when $d/\Delta_{\min} > \kappa(\Delta \vee \Delta_{\neq} \vee \varepsilon_T)/\Delta_{\neq}^2$, the difficulty of the linear bandit with systematic bias is dominated by that of the classical d-linear bandit. In this case, the algorithm is able to find the group containing the best action, and the problem reduces to a linear bandit in dimension d. Thus, the linear bandit with systematic bias is a non trivial example of a globally observable game that can be locally observable around the best action.

Finally, we underline that the magnitude of the bias does not appear in the regret: intuitively, no matter its magnitude, the algorithm always need to estimate it up to the same precision (of order $\Delta_{\neq}$) in order to find the best group and to be optimal in terms of gap-depend regret. This indicates that our algorithm is robust against important discriminations in the evaluation mechanism.

5 Lower bounds on the regret

In this section, we derive lower bounds on the worst-case regret and the gap-dependent regret that respectively match the upper bounds established in Theorems 1 and 2 up to sub-logarithmic factors or numerical constants.

5.1 Lower bound on the worst-case regret

Theorems 1 and 2 underline the dependence of the regret on the geometry of the action set. Before stating our result, we begin by introducing the notion of κ_*-correlated action set.

Definition 1 (κ_*-correlated action set). *For $\kappa_* \geq 1$, a set of actions $\mathcal{A}$ is κ_*-correlated if $\mathcal{A} \in \mathbf{A}_{\kappa_*,d}$, where*

$$
\mathbf{A}_{\kappa_*,d} = \left\{ \begin{array}{l} \mathcal{A} = \{a_1,...,a_k\} \subset \left(\mathbb{R}^d \times \{-1,+1\}\right)^k : \\[2mm] k \in \mathbb{N}^*, \; \min_{\pi \in \mathcal{P}^{\mathcal{A}}_{e_{d+1}}} \left\{ e_{d+1}^\top \Big(\sum_{a \in \mathcal{A}} \pi(a) aa^\top\Big)^+ e_{d+1} \right\} \geq \kappa_* \end{array} \right\}
$$

is the set of actions sets such that the minimal variance of the bias estimator is larger than κ_*.

In the following theorem, we establish a lower bound on the regret valid for all $\kappa_* \geq 1$ by designing κ_*-correlated sets of actions $\mathcal{A} \in \mathbf{A}_{\kappa_*,d}$, and obtaining lower bounds on the regret of any algorithm on these sets of actions.

Theorem 3. *Let $\kappa_* \geq 1$, $d \geq 2$ and $T \geq 4^3 \kappa_*$. There exists an action set $\mathcal{A} \in \mathbf{A}_{\kappa_*,d}$ such that for any algorithm, there exists a bandit problem with parameter $\theta_T \in \mathbb{R}^{d+1}$ such that the regret of this algorithm on the problem characterized by θ_T satisfies $R_T^{\theta_T} \geq \kappa_*^{1/3} T^{2/3}/8e$.*

Previous lower bounds on the regret of linear bandits with partial monitoring, established in [20], state that the regret must be at least $c_{\mathcal{A}} T^{2/3}$ for some parameter $\theta_T \in \mathbb{R}^{d+1}$, where $c_{\mathcal{A}} > 0$ is a constant depending (not explicitly) on $\mathcal{A}$. By contrast, Theorem 3 provides an explicit characterization of the dependence of the regret rate on the geometry of the problem, which matches the upper bound of Theorem 1 up to a sub-logarithmic factor. Note that the assumption $d \geq 2$ is necessary here: if $d = 1$, there are at most two potentially optimal actions (namely, $\max\{x : x \in \mathcal{X}\}$ and $\min\{x : x \in \mathcal{X}\}$). Then, the problem becomes locally observable, and regret of order $\widetilde{O}(T^{1/2})$ can be achieved [20].

5.2 Lower bound on the gap-dependent regret

We now present a lower bound on the gap-dependent regret. More precisely, for given values of $\Delta_{\min}$ and $\Delta_{\neq}$, we establish a lower bound on the worst case regret among parameters θ verifying $\Delta_{\min} \leq \min_{x \in \mathcal{X} \setminus x^*} \Delta_x$, and $\Delta_{\neq} \leq \min_{x \in \mathcal{X}:z_x = -z_{x^*}} \Delta_x$. Before stating formally the result, let us define the corresponding parameter set. For an action set $\mathcal{A} \in \mathbf{A}_{\kappa_*,d}$, and for $(\Delta_{\min}, \Delta_{\neq}) \in (0,1)^2$ such that $\Delta_{\min} \leq \Delta_{\neq}$, we denote

$$
\Theta^{\mathcal{A}}_{\Delta_{\min}, \Delta_{\neq}} = \left\{ \begin{array}{l} \theta = \binom{\gamma}{\omega} : \; \gamma \in \mathcal{C}(\mathcal{X}), \; \exists ! \; \binom{x^*}{z_{x^*}} \in \mathrm{argmax}_{\binom{x}{z_x} \in \mathcal{A}}\{x^\top \gamma\}, \\[2mm] \forall \binom{x'}{z_{x'}} \in \mathcal{A} \text{ such that } x' \neq x^*, (x^* - x')^\top \gamma \geq \Delta_{\min}, \\[2mm] \forall \binom{x'}{z_{x'}} \in \mathcal{A} \text{ such that } z_{x'} \neq z_{x^*}, (x^* - x')^\top \gamma \geq \Delta_{\neq} \end{array} \right\}
$$

the set of parameters with minimum gap $\Delta_{\min}$, and minimum between-group-gap $\Delta_{\neq}$.

The upper bounds established in Theorem 2 underline the dependence of the gap-dependent regret on the minimal regret $\kappa(\Delta)$ for estimating the bias. Before stating our results, we define a class of problems $\Theta^{\mathcal{A}}_{\Delta_{\min}, \Delta_{\neq}, \kappa}$ such that $\kappa(\Delta) \leq \kappa$. For a parameter $\gamma \in \mathcal{C}(\mathcal{X})$, let us denote $\Delta(\gamma)_x = \max_{x' \in \mathcal{X}} (x' - x)^\top \gamma$, and $\Delta(\gamma) = (\Delta(\gamma)_x)_{x \in \mathcal{X}}$. Moreover, for a given set $\mathcal{A}$, let us denote

$$
\Theta^{\mathcal{A}}_{\Delta_{\min}, \Delta_{\neq}, \kappa} = \Theta^{\mathcal{A}}_{\Delta_{\min}, \Delta_{\neq}} \cap \left\{ \theta = \binom{\gamma}{\omega} : \; \gamma \in \mathcal{C}(\mathcal{X}), \; \kappa(\Delta(\gamma)) \leq \kappa \right\}.
$$

Theorem 4. *For all $\kappa \geq 2$ and all $d \geq 4$, there exists a set of actions $\mathcal{A} \in \mathbb{R}^{d+1}$ such that for all $(\Delta_{\min}, \Delta_{\neq}) \in (0, 1/8)^2$ with $\Delta_{\min} \leq \Delta_{\neq}$,*

$$\liminf_{T \to \infty} \sup_{\theta \in \Theta^{\mathcal{A}}_{\Delta_{\min}, \Delta_{\neq}, \kappa}} \frac{R_T^{\theta}}{\log(T)} \geq \left[\frac{d}{10\Delta_{\min}}\right] \vee \left[\frac{\kappa + 2}{8\Delta_{\neq}^2}\right]. \tag{4}$$

Theorem 4 shows that for some action sets $\mathcal{A}$, the gap-depend regret of the FAIR PHASED ELIM-
INATION algorithm is asymptotically optimal up to a numerical constant. Note that the assump-
tion $d \geq 4$ is necessary in our proof to design an action set $\mathcal{A}$ such that Equation (4) holds for
all $\Delta_{\min}, \Delta_{\neq} \in (0, 1/8)$. On the other hand, as discussed in Appendix C.6, for $d \geq 2$, for all
$\Delta_{\min}, \Delta_{\neq} \in (0, 1/8)$, we can show that there exists action sets $\mathcal{A}$ and $\theta \in \Theta^{\mathcal{A}}_{\Delta_{\min}, \Delta_{\neq}}$ such that
the lower bound in Equation (4) still holds, by considering separately the cases $d/\Delta_{\min} > \kappa/\Delta_{\neq}^2$ and
$d/\Delta_{\min} \leq \kappa/\Delta_{\neq}^2$.

6   Conclusion

In this paper, we addressed the problem of online decision making under biased bandit feedback.
We designed a new algorithm based on Δ- and G-optimal design, and obtained worst-case and
gap-dependent upper bounds on its regret. We obtained lower bounds on the regret for some problem
instances showing that these rates are tight up to sub-logarithmic factors in some settings. These rates
highlight two behaviors: on the one hand, the worst case rate $\mathcal{O}(\kappa_*^{1/3} \log(T)^{1/3} T^{2/3})$ highlights the
cost induced by the biased feedback, and the need to select sub-optimal actions in order to debias
it. On the other hand, the gap-dependent bound shows that for some instance, the problem can be
locally observable around the best action: then, the difficulty of the problem is dominated by the
difficulty of the corresponding linear bandit problem, and is no more difficult than this problem.
When this is not the case, the regret scales as $\kappa(\Delta)\Delta_{\neq}^{-2}\log(T)$, where $\Delta_{\neq}$ is the gap between the
best actions of the two groups, and $\kappa(\Delta)$ is the minimum regret for estimating the bias with a given
precision. In Appendix D, we discuss the extension of the biased linear model and of the Fair Phased
Elimination algorithm to multiple groups with different biases. This work paves the way for studying
other bandit models with unfair feedback, considering for example continuous, multi-dimensional
sensitive attributes.

Broader impact

In this work, we propose a model for sequential decision making under biased feedback. Our goal
is primarily to provide a good strategy for sequential learning in an unfair environment, and to
characterize the difficulty of this problem by bounding the regret. On the one hand, our results reveal
that maximizing the fair rewards instead of unfair evaluations may be more difficult in terms of regret,
which may discourage practitioners from correcting unfair feedbacks. On the other hand, we believe
that as fairness is an important long-term key objective, rather than discouraging the practitioner, it
will inform them to better plan the adaptation of their methods toward this aim.

Acknowledgements.

The authors would like to thank Evgenii Chzhen and Nicolas Verzelen for their valuable discussions
and suggestions.

The work of A. Carpentier is partially supported by the Deutsche Forschungsgemeinschaft (DFG)
Emmy Noether grant MuSyAD (CA 1488/1-1), by the DFG - 314838170, GRK 2297 MathCoRe, by
the FG DFG, by the DFG CRC 1294 'Data Assimilation', Project A03, by the Forschungsgruppe
FOR 5381 "Mathematical Statistics in the Information Age - Statistical Efficiency and Computational
Tractability", Project TP 02, by the Agence Nationale de la Recherche (ANR) and the DFG on the
French-German PRCI ANR ASCAI CA 1488/4-1 "Aktive und Batch-Segmentierung, Clustering und
Seriation: Grundlagen der KI" and by the UFADFH through the French-German Doktorandenkolleg
CDFA 01-18 and by the SFI Sachsen-Anhalt for the project RE-BCI. Christophe Giraud received
partial support by grant ANR-19-CHIA-0021-01 ("BiSCottE", Agence Nationale de la Recherche)
and by the ANR and the DFG on the French-German PRCI ANR-21-CE23-0035 (ASCAI).

References

[1] Y. Abbasi-Yadkori, D. Pál, and C. Szepesvári. Improved algorithms for linear stochastic bandits. In J. Shawe-Taylor, R. Zemel, P. Bartlett, F. Pereira, and K. Q. Weinberger, editors, *Advances in Neural Information Processing Systems*, volume 24. Curran Associates, Inc., 2011.

[2] A. Barik and J. Honorio. Fair sparse regression with clustering: An invex relaxation for a combinatorial problem. In *Thirty-Fifth Conference on Neural Information Processing Systems*, 2021.

[3] A. Byanjankar, M. Heikkilä, and J. Mezei. Predicting credit risk in peer-to-peer lending: A neural network approach. In *2015 IEEE Symposium Series on Computational Intelligence*, pages 719–725, 2015.

[4] L. E. Celis, S. Kapoor, F. Salehi, and N. K. Vishnoi. An algorithmic framework to control bias in bandit-based personalization, 2018.

[5] S. Chaudhuri and A. Tewari. Phased exploration with greedy exploitation in stochastic combinatorial partial monitoring games. In D. Lee, M. Sugiyama, U. Luxburg, I. Guyon, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 29. Curran Associates, Inc., 2016.

[6] S. Chawla and M. Jagadeesan. Individual Fairness in Advertising Auctions Through Inverse Proportionality. In M. Braverman, editor, *13th Innovations in Theoretical Computer Science Conference (ITCS 2022)*, volume 215 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 42:1–42:21, Dagstuhl, Germany, 2022. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.

[7] E. Chzhen, C. Giraud, and G. Stoltz. A unified approach to fair online learning via blackwell approachability. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P. Liang, and J. W. Vaughan, editors, *Advances in Neural Information Processing Systems*, volume 34, pages 18280–18292. Curran Associates, Inc., 2021.

[8] E. Chzhen and N. Schreuder. A minimax framework for quantifying risk-fairness trade-off in regression, 2020.

[9] H. Claure, Y. Chen, J. Modi, M. F. Jung, and S. Nikolaidis. Multi-armed bandits with fairness constraints for distributing resources to human teammates. *2020 15th ACM/IEEE International Conference on Human-Robot Interaction (HRI)*, pages 299–308, 2020.

[10] G. Elfving. Optimum Allocation in Linear Regression Theory. *The Annals of Mathematical Statistics*, 23(2):255 – 262, 1952.

[11] J. Fauw, J. Ledsam, B. Romera-Paredes, S. Nikolov, N. Tomasev, S. Blackwell, H. Askham, X. Glorot, B. O'Donoghue, D. Visentin, G. Driessche, B. Lakshminarayanan, C. Meyer, F. Mackinder, S. Bouton, K. Ayoub, R. Chopra, D. King, A. Karthikesalingam, and O. Ronneberger. Clinically applicable deep learning for diagnosis and referral in retinal disease. *Nature Medicine*, 24, 09 2018.

[12] V. V. Fedorov. Theory of optimal experiments. *New York : Academic Press*, 1972.

[13] J. Fellman. *On the Allocation of Linear Observations*. Commentationes physico-mathematicae. Societas Scientiarum Fennica, 1974.

[14] A. Fuster, P. Goldsmith-Pinkham, T. Ramadorai, and A. Walther. Predictably unequal? the effects of machine learning on credit markets. *The Journal of Finance*, 77(1):5–47, 2022.

[15] S. Gaucher, N. Schreuder, and E. Chzhen. Fair learning with wasserstein barycenters for non-decomposable performance measures, 2022.

[16] H. Hadiji, S. Gerchinovitz, J.-M. Loubes, and G. Stoltz. Diversity-Preserving K-Armed Bandits, Revisited. working paper or preprint, Oct. 2020.

[17] R. Harman and T. Jurík. Computing c-optimal experimental designs using the simplex method of linear programming. *Computational Statistics & Data Analysis*, 53(2):247–254, dec 2008.

[18] A. Khademi, S. Lee, D. Foley, and V. Honavar. Fairness in algorithmic decision making: An excursion through the lens of causality. In *The World Wide Web Conference*, WWW '19, page 2907–2914, New York, NY, USA, 2019. Association for Computing Machinery.

[19] J. Kiefer. General Equivalence Theory for Optimum Designs (Approximate Theory). *The Annals of Statistics*, 2(5):849 – 879, 1974.

[20] J. Kirschner, T. Lattimore, and A. Krause. Information directed sampling for linear partial monitoring. In J. D. Abernethy and S. Agarwal, editors, *Conference on Learning Theory*, volume 125 of *Proceedings of Machine Learning Research*, pages 2328–2369. PMLR, 2020.

[21] J. Kirschner, T. Lattimore, C. Vernade, and C. Szepesvari. Asymptotically optimal information-directed sampling. In M. Belkin and S. Kpotufe, editors, *Proceedings of Thirty Fourth Conference on Learning Theory*, volume 134 of *Proceedings of Machine Learning Research*, pages 2777–2821. PMLR, 15–19 Aug 2021.

[22] A. Köchling and M. C. Wehner. Discriminated by an algorithm: a systematic review of discrimination and fairness by algorithmic decision-making in the context of hr recruitment and hr development. *Business Research*, pages 1–54, 2020.

[23] T. Lattimore and C. Szepesvari. The End of Optimism? An Asymptotic Analysis of Finite-Armed Linear Bandits. In A. Singh and J. Zhu, editors, *Proceedings of the 20th International Conference on Artificial Intelligence and Statistics*, volume 54 of *Proceedings of Machine Learning Research*, pages 728–737. PMLR, 20–22 Apr 2017.

[24] T. Lattimore and C. Szepesvári. *Bandit Algorithms*. Cambridge University Press, 2020.

[25] T. Lin, B. Abrahao, R. Kleinberg, J. Lui, and W. Chen. Combinatorial partial monitoring game with linear feedback and its applications. In E. P. Xing and T. Jebara, editors, *Proceedings of the 31st International Conference on Machine Learning*, number 2, pages 901–909, Bejing, China, 22–24 Jun 2014. PMLR.

[26] Z. Lipton, J. McAuley, and A. Chouldechova. Does mitigating ml's impact disparity require treatment disparity? In S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 31. Curran Associates, Inc., 2018.

[27] N. Mehrabi, F. Morstatter, N. Saxena, K. Lerman, and A. Galstyan. A survey on bias and fairness in machine learning. *ACM Computing Surveys*, 54(6), jul 2021.

[28] Z. Papamitsiou and A. A. Economides. Learning analytics and educational data mining in practice: A systematic literature review of empirical evidence. *Journal of Educational Technology & Society*, 17(4):49–64, 2014.

[29] V. Patil, G. Ghalme, V. Nair, and Y. Narahari. Achieving fairness in the stochastic multi-armed bandit problem. *Proceedings of the AAAI Conference on Artificial Intelligence*, 34(04):5379–5386, Apr. 2020.

[30] A. Pázman. *Foundations of Optimum Experimental Design*. Mathematics and its Applications. Springer Netherlands, 1986.

[31] C. Perlich, B. Dalessandro, T. Raeder, O. Stitelman, and F. Provost. Machine learning for targeted display advertising: transfer learning in action. *Machine Learning*, 95(1):103–127, 2014.

[32] L. Pronzato and G. Sagnol. Removing inessential points in c-and A-optimal design. *Journal of Statistical Planning and Inference*, 213:233–252, 2021.

[33] F. Pukelsheim. On linear regression designs which maximize information. *Journal of statistical planning and inferrence*, 4:339–364, 1980.

[34] H. Qi. A semidefinite programming study of the elfving theorem. *Journal of Statistical Planning and Inference*, 141:3117–3130, 2011.

[35] M. Raghavan, S. Barocas, J. M. Kleinberg, and K. E. C. Levy. Mitigating bias in algorithmic hiring: evaluating claims and practices. *Proceedings of the 2020 Conference on Fairness, Accountability, and Transparency*, 2020.

[36] G. Sagnol. Plans d'expériences optimaux et application à l'estimation des matrices de trafic dans les grands réseaux : programmation conique du second ordre et sous-modularité. 2010.

[37] G. Sagnol and R. Harman. Computing exact d-optimal designs by mixed integer second order cone programming. *The Annals of Statistics*, 43, 07 2013.

[38] G. Sagnol and M. Stahlberg. Picos: A python interface to conic optimization solvers. *Journal of Open Source Software*, 7(70):3915, 2022.

[39] D. M. Titterington. Algorithms for computing D-optimal design on finite design spaces. *Proceedings of the 1976 Conf. on Information Science and Systems*, page 213–216, 1976.

[40] L. Vandenberghe, S. Boyd, and S.-P. Wu. Determinant maximization with linear matrix inequality constraints. *SIAM Journal on Matrix Analysis and Applications*, 19(2):499–533, 1998.

[41] M. Černý and M. Hladík. Two complexity results on c-optimality in experimental design. *Computational Optimization and Applications*, 51(3):1397–1408, apr 2012.

[42] A. Wagenmaker, J. Katz-Samuels, and K. Jamieson. Experimental design for regret minimization in linear bandits. In A. Banerjee and K. Fukumizu, editors, *Proceedings of The 24th International Conference on Artificial Intelligence and Statistics*, volume 130 of *Proceedings of Machine Learning Research*, pages 3088–3096. PMLR, 13–15 Apr 2021.

[43] L. Wang, Y. Bai, W. Sun, and T. Joachims. Fairness of exposure in stochastic bandits. In M. Meila and T. Zhang, editors, *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pages 10686–10696. PMLR, 18–24 Jul 2021.

[44] H. P. Wynn. The Sequential Generation of D-Optimum Experimental Designs. *The Annals of Mathematical Statistics*, 41(5):1655 – 1664, 1970.

Appendix

The Appendix is organized as follows. In Section A, we further discuss the geometry of bias estimation, and provide additional results on the constants κ_* and $\kappa(\Delta)$. Then, we provide in Section B a detailed version of the FAIR PHASED ELIMINATION algorithm 3. In Section C, we prove the main results of this paper. Finally, in Section D, we discuss the extension of the biaised linear bandits to more than 2 groups.

A   On the geometry of bias estimation

We begin in Section A.1 by highlighting the relationship of the constant κ_* with the problem of e_{d+1}-*optimal design*. Then, in Section A.2, we show that the geometrical constant κ_* can be expressed in terms of separation of the two groups. In Section A.3 and Section A.4, we relate κ_* to classical geometrical measures of the difficulty of a set of actions such as the condition.ing number and the worst-case alignment constant of [20]. In Section A.5, we show that κ_* is equivalent to the variance of the optimal design for estimation the bias against the worst parameter θ^*. In Section A.6, we provide further results on $\kappa(\Delta)$, the Δ-optimal regret for estimation the bias with variance 1 when the gap vector is Δ. Finally, in Section A.7, we propose guidance for computing the G-optimal and Δ-optimal designs.

A.1   Bias estimation as a $\mathbf{e_{d+1}}$-optimal design problem

Recall that κ_* is the minimal variance of the bias estimator related to the problem of e_{d+1}-optimal design.

$\mathbf{e_{d+1}}$**-optimal design**   Optimal design theory addresses the following problem: a scientist must design a set of n experiments $\{x_1, ..., x_n\} \in \mathcal{X}^n$ so as to estimate at best a parameter of interest, where each experiment $x \in \mathcal{X}$ corresponds to a point $a_x \in \mathbb{R}^{d+1}$. The aim of the scientist is to choose a design, i.e. a function $\mu : \mathcal{X} \mapsto \mathbb{N}$ indicating the budget $\mu(x)$ to be allocated to each experiment $x \in \mathcal{X}$. Each experiment x is then repeated exactly $\mu(x)$ times, and the corresponding observations $y_{x,1}, ..., y_{x,\mu(x)}$ are collected for each $x \in \mathcal{X}$. The law of the observations corresponding to experiment x at point a_x is given by

$$y_{x,i} = a_x^\top \theta^* + \xi_{x,i},$$

where $\xi_{x,i} \sim \mathcal{N}(0, 1)$ are independent noise terms, and $\theta^* \in \mathbb{R}^{d+1}$ is an unknown parameter. The aim of the scientist is to choose the design μ so as to best estimate (some features of) the parameter θ^*, under a constraint on the total number of experiments $\sum_{x \in \mathcal{X}} \mu(x) \leq n$ for some $n \in \mathbb{N}$.

Different criteria can be used to characterize the optimality of a design μ. For example, one may need to estimate the full parameter θ^*, in order to predict the outcomes of the experiments $x \in \mathcal{X}$ with a small uniform error: this leads to the G-optimal design problem (2). Alternatively, for c a vector in $\mathbb{R}^{d+1}$, one may aim at finding the best design $\mu \in \mathcal{N}^{\mathcal{X}}$ for estimating the scalar product $c^\top \theta^*$ under a budget constraint $\sum_{x \in \mathcal{X}} \mu(x) \leq n$, where $\mathcal{N}^{\mathcal{X}} = \{\mu : \mathcal{X} \to \mathbb{N}\}$. This problem is known as *c-optimal design*. Unbiased linear estimation of $c^\top \theta^*$ is possible only when c belongs to the image of $V(\mu)$, and in this case the best linear unbiased estimator of the scalar product $c^\top \theta^*$ is given by $c^\top \widehat{\theta}$, where $\widehat{\theta}$ is the least-square estimator defined as

$$\widehat{\theta} = V(\mu)^+ \sum_{x \in \mathcal{X}} a_x \left(\sum_{i \leq \mu(x)} y_{x,i} \right) \quad \text{for} \quad V(\mu) = \sum_{x \in \mathcal{X}} \mu(x) a_x a_x^\top. \tag{5}$$

The variance of the estimator $c^\top \widehat{\theta}$ is then equal to $c^\top V(\mu)^+ c$.

Exact c-optimal design aims at choosing the allocation $\mu \in \mathcal{N}^{\mathcal{X}}$ minimizing the variance of $c^\top \widehat{\theta}$ for a given budget $\sum_x \mu(x) \leq n$, under the constraint that $c \in \text{Range}(V(\mu))$. Let us define the normalized design $\pi : x \in \mathcal{X} \mapsto \mu(x)/n$, and let us underline that π defines a probability on $\mathcal{X}$. The variance of $c^\top \widehat{\theta}$ is then equal to $n^{-1} c^\top V(\pi)^+ c$. In the limit $n \to +\infty$, the problem is equivalent to

the problem of approximate c-optimal design (sometimes simply referred to as c-optimal design), that aims at finding a probability measure $\pi \in \mathcal{P}_c^{\mathcal{X}} := \{\pi \in \mathcal{P}^{\mathcal{X}} : c \in \mathrm{Range}(V(\pi))\}$ solution to the following problem

$$\min_{\pi \in \mathcal{P}_c^{\mathcal{X}}} c^\top V(\pi)^+ c. \qquad \text{(c-optimal design)}$$

Note that when $\{a_x : x \in \mathcal{X}\}$ spans $\mathbb{R}^{d+1}$, for any $c \in \mathbb{R}^{d+1}$, there exists a design π such that $c \in \mathrm{Range}(V(\pi))$, and hence the c-optimal design problem admits a solution.

Computation of the $\mathbf{e_{d+1}}$-optimal design Finding an exact optimal allocation $\mu \in \mathcal{N}^{\mathcal{X}}$ under the constraint that $\sum_{x \in \mathcal{X}} \mu(x) \le n$ is unfortunately NP-complete. However, finding an approximate optimal design $\pi \in \mathcal{P}_c^{\mathcal{X}}$ can be done in polynomial time [41]. Several algorithms, including multiplicative algorithms [13] and a simplex method of linear programming [17], have been proposed to iteratively approximate the optimal design. More recently, [32] suggested using screening tests to remove inessential points to accelerate optimization algorithms.

Classical results from e_{d+1}-optimal design show that there exists a c-optimal design supported by at most $d+1$ points (see, e.g., [30, 17] for a proof of this result). The following Lemma indicates how to obtain an exact design by rounding an approximate design supported by at most $d+1$ points.

Lemma 3. *For any $\pi \in \mathcal{M}_{e_{d+1}}^{\mathcal{X}}$ and any $m > 0$, the estimator $e_{d+1}^\top \widehat{\theta}_\mu$ computed from the design μ : $x \mapsto \lceil m\pi(x) \rceil$ is an unbiased estimator of $e_{d+1}^\top \theta$ and it has a variance at most $m^{-1} e_{d+1}^\top V(\pi)^+ e_{d+1}$.*

Obviously, similar results also hold for G-optimal design.

Lemma 4. *Let π be a solution of the G-optimal design problem (2). Then, for any $m > 0$ and any $x \in \mathcal{X}$, the estimator $a_x^\top \widehat{\theta}_\mu$ computed from the design $\mu : x \mapsto \lceil m\pi(x) \rceil$ is an unbiased estimator of the evaluation $a_x^\top \theta$, and it has a variance*

$$a_x^\top V(\mu)^+ a_x \le m^{-1}(d+1).$$

A.2 Interpretation of κ_* in terms of separation of the groups

Next theorem, due to Elfving, characterizes solutions to the c-optimal design problem.

Theorem 5 ([10]). *Let $\mathcal{S} = convex\ hull\ \{+a_x, -a_x : x \in \mathcal{X}\}$ be the Elfving's set of $\{a_x : x \in \mathcal{X}\} \subset \mathbb{R}^{d+1}$, and let $\partial \mathcal{S}$ denote the boundary of $\mathcal{S}$. A design $\pi \in \mathcal{P}_c^{\mathcal{X}}$ is c-optimal for $c \in \mathbb{R}^{d+1}$ if and only if there exists $\zeta \in \{-1, +1\}^{\mathcal{X}}$ and $t > 0$ such that*

$$tc = \sum_{x \in \mathcal{X}} \pi(x) \zeta_x a_x \in \partial \mathcal{S}.$$

Moreover, $t^{-2} = c^\top (V(\pi))^+ c$ is value of the c-optimal design problem.

Elfving's characterization of the e_{d+1}-optimal design allows us to derive the following equivalent characterization of κ_*.

Lemma 5. $\kappa_* = \max_{u \in \mathbb{R}^d} \dfrac{1}{\max_{x \in \mathcal{X}} \left(x^\top u + z_x\right)^2}.$

Lemma 1 follows from the characterization in Lemma 5. When $\kappa_* > 1$, the vector $\tilde{u}$ defined as

$\tilde{u} = \mathrm{argmax}_{u \in \mathbb{R}^d} \frac{1}{\max_{x \in \mathcal{X}} (x^\top u + z_x)^2}$ is a normal vector of the separating hyperplane $\mathcal{H}$ in Figure 1. Moreover, as shown in the proof of Lemma 1, the margin is in this case equal to $1 - \kappa_*^{-1/2}$, while the maximum distance of all points to the hyperplane is $1 + \kappa_*^{-1/2}$.

Application to the action set $\mathcal{A}$ of Lemma 10 To provide the reader with intuition on κ_*, we analyze here the set of actions used to derive the lower bound in Theorem 3. Let $\mathcal{A} = \left\{ \begin{pmatrix} x_1 \\ z_{x_1} \end{pmatrix}, ..., \begin{pmatrix} x_{d+1} \\ z_{x_{d+1}} \end{pmatrix} \right\}$, where $\begin{pmatrix} x_i \\ z_{x_i} \end{pmatrix} = e_i + e_{d+1}$, for $i \in \{2, ..., \lfloor d/2 \rfloor\}$, $\begin{pmatrix} x_i \\ z_{x_i} \end{pmatrix} = e_i - e_{d+1}$ for $i \in \{\lfloor d/2 \rfloor + 1, ..., d\}$, and $\begin{pmatrix} x_{d+1} \\ z_{x_{d+1}} \end{pmatrix} = -\left(1 - \frac{2}{\sqrt{\kappa_*}+1}\right) e_1 - e_{d+1}$. We show in Lemma 10 that the minimal variance for estimating the bias on $\mathcal{A}$ is indeed κ_*.

Figure 2: Illustration of Lemma 1 on the action set $\mathcal{A}$ described above for $d = 2$.

The set of actions $\mathcal{A}$ spans $\mathbb{R}^{d+1}$, however it is easy to see that only x_1 and x_{d+1} can be used to estimate the bias. On the one hand, when $\kappa = 1$, $\begin{pmatrix} x_{d+1} \\ z_{x_{d+1}} \end{pmatrix} = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$, so the bias can be evaluated just by sampling x_{d+1}. In the other hand, in the limit where $\kappa_* \to \infty$, the problems becomes more difficult as $\begin{pmatrix} x_{d+1} \\ z_{x_{d+1}} \end{pmatrix}$ tends to $- \begin{pmatrix} x_1 \\ z_{x_1} \end{pmatrix}$. In the limit $\kappa_* = \infty$, it is impossible to distinguish between the contribution of $\gamma^\top e_1$ and ω in the evaluations of actions 1 and $d + 1$: the problem becomes not identifiable. We represent this setting for an intermediate value of κ_* in Figure 2. We also represent the separating hyperplane, margin m and distance M of Lemma 1.

A.3   Comparison to the conditioning number

By contrast to classical complexity measures such as conditioning numbers that give equal weight to all observations, optimal design gives flexibility to choose $d + 1$ best actions to estimate the bias, and therefore allows for sharper bounds.

Indeed, by definition of κ_*,

$$\kappa_* \le e_{d+1} V(\pi^u)^+ e_{d+1},$$

where π^u is the uniform measure giving the same weight $1/k$ to all actions. Now, $V(\pi^u)$ is the classical covariance matrix associated with the design points $a_x \in \mathcal{A}$, so the condition number CN of this design is given by

$$CN = \frac{\lambda_{\max}(V(\pi^u))}{\lambda_{\min}(V(\pi^u))}.$$

We see that $e_{d+1} V(\pi^u)^+ e_{d+1} \le \lambda_{\min}(V(\pi^u))^{-1}$. When the actions a_x are bounded (for example $\|a_x\| \le M$), this implies that $\kappa_* \le CN/M$.

We provide an example showing that κ_* can be much smaller than the conditioning number. Consider the following example in dimension $d = 2$ with $k \ge 4$ actions, where $x_1 = (1, 0)$ and $x_2 = (-1, 0)$ belong to group 1, and $x_3, ..., x_k$ are identical, equal to $(0, 1)$, and in group -1. Then, Lemma 1 shows that the minimal variance for estimating the bias is indeed 1, and that the optimal design puts equal mass on x_1 and x_2. On the other hand, straightforward computations show that the conditioning number of the covariance matrix is $\frac{1 + (k-2)^{-1} + \sqrt{1 + (k-2)^{-2}}}{1 + (k-2)^{-1} - \sqrt{1 + (k-2)^{-2}}}$. Thus, on this example, CN/κ_* is of order k.

A.4   Comparison to the worst-case alignment constant

Lemma 5 also allows us to compare the bound in Theorem 1 with previous results on linear bandit with partial monitoring, expressed in terms of the worst-case alignment constant.

Previous work on linear bandit with partial linear monitoring measures the difficulty of the bandit game using the *worst-case alignment constant* α, defined as

$$\alpha = \max_{u \in \mathbb{R}^d} \frac{\max_{x,x' \in \mathcal{X}}((x - x')^\top u)^2}{\max_{x \in \mathcal{X}}(z_x x^\top u + 1)^2}.$$

The following Lemma shows that this constant is essentially equivalent to the minimal variance of the bias estimator κ_*.

Lemma 6. $\frac{\kappa_*}{3} \leq \alpha \leq 16\kappa_*$.

On the one hand, Lemma 6 shows that κ_* and α are essentially equivalent. In particular, Theorem 3 implies that the large T regret is of order $\alpha^{1/3} \log(T)^{1/3} T^{2/3}$. This improves over previous known rates, obtained in [20], by a factor $d^{1/2} \log(T)^{1/6} (\log(kT)/\log(T))^{1/2}$.

On the other hand, as underlined, the constant κ_* appears when considering the well-studied problem of c-optimal design. Therefore, classical results and algorithms for optimal design can be used to characterize and compute this constant.

A.5   Optimal bias estimation against the worst parameter

The constant κ_* also appears naturally when considering the related problem of optimal bias estimation against the worst parameter.

Regret of e_{d+1}-optimal design   Recall that κ_* denotes the *minimal variance of the bias estimator*, i.e. the value of the solution of the e_{d+1}-optimal design problem

$$\kappa_* = \min_{\pi \in \mathcal{P}^{\mathcal{X}}_{e_{d+1}}} e_{d+1}^\top (V(\pi))^+ e_{d+1},$$

The e_{d+1}-optimal design can be equivalently defined as the solution of the problem

$$\text{minimize} \sum_{x \in \mathcal{X}} \mu(x) \quad \text{such that } \mu \in \mathcal{M}^{\mathcal{X}}_{e_{d+1}} \text{ and } e_{d+1}^\top V(\mu)^+ e_{d+1} \leq \kappa_*. \tag{6}$$

The characterization given in Equation (6) underlines that the e_{d+1}-optimal design provides (up to discretization issues) the minimal number of samples required for estimating ω^* with a variance κ_*. Let us denote by μ^* the optimal design for estimating ω^* with a variance 1, defined as

$$\mu^* = \underset{\mu}{\text{argmin}} \sum_{x \in \mathcal{X}} \mu(x) \quad \text{such that } \mu \in \mathcal{M}^{\mathcal{X}}_{e_{d+1}} \text{ and } e_{d+1}^\top V(\mu)^+ e_{d+1} \leq 1.$$

Note that from the definition of κ_*, we have $\sum_x \mu^*(x) = \kappa_*$.

A first (naive) approach to obtain an estimate of the bias parameter ω^* with precision level $\epsilon > 0$ would consist in sampling actions according to $\epsilon^{-2}\mu^*$, rounded according to the procedure defined in Lemma 3. Let us denote by Δ_x the gap $\Delta_x = \max_{x' \in \mathcal{X}}(x' - x)^\top \gamma^*$ between the (non-observed) reward of the best action and the reward of the action x. The regret corresponding to this estimation phase would then be

$$\epsilon^{-2} \sum_{x \in \mathcal{X}} \mu^*(x) \Delta_x,$$

which can be as large as $\kappa_* \epsilon^{-2} \max_x \Delta_x$. Interestingly, we show that the regret corresponding to the e_{d+1}-optimal design is equivalent (up to a small multiplicative constant) to the minimax regret.

Optimal worst-case estimation   The minimax regret corresponds to the regret of the best sampling scheme against the worst admissible parameter γ. Note that, for a given design μ, this worst-case regret is given by

$$\max_{x' \in \mathcal{X}, \gamma \in \mathcal{C}(\mathcal{X})} \sum_x \mu(x)(x' - x)^\top \gamma,$$

where we recall that $\mathcal{C}(\mathcal{X}) = \{\gamma \in \mathbb{R}^d : \forall x \in \mathcal{X}, |x^\top \gamma| \leq 1\}$ is the set of admissible parameters. To achieve the lowest regret against the worst parameter, we must use the minimax optimal design $\widetilde{\mu}$ solution to the problem

$$\widetilde{\mu} = \underset{\mu}{\text{argmin}} \max_{x' \in \mathcal{X}, \gamma \in \mathcal{C}(\mathcal{X})} \sum_{x \in \mathcal{X}} \mu(x)(x' - x)^\top \gamma \text{ such that } \mu \in \mathcal{M}^{\mathcal{X}}_{e_{d+1}} \text{ and } e_{d+1}^\top V(\mu)^+ e_{d+1} \leq 1.$$

Lemma 2 underlines that the regret corresponding to the e_{d+1}-optimal design is no larger than twice the minimax regret.

A.6 Additionnal results the Δ-optimal design

Recall that for a vector of gaps $\Delta = (\Delta_x)_{x\in\mathcal{X}}$, μ^Δ denotes the Δ-optimal design, defined as the solution of the following problem

$$\mu^\Delta = \operatorname*{argmin}_\mu \sum_{x\in\mathcal{X}} \mu(x)\Delta_x \quad \text{such that } \mu \in \mathcal{M}^{\mathcal{X}}_{e_{d+1}} \text{ and } e_{d+1}^\top V(\mu)^+ e_{d+1} \le 1. \quad \text{(Δ-optimal design)}$$

If we knew the gaps Δ_x, we could sample the actions according to the Δ-optimal design μ^Δ, and pay the regret $\epsilon^{-2}\kappa(\Delta)$ (up to rounding error) for estimating ω^* with an error smaller than ϵ, where

$$\kappa(\Delta) = \sum_{x\in\mathcal{X}} \mu^\Delta(x)\Delta_x.$$

Lemma 7. *If $\gamma^* \in \mathcal{C}(\mathcal{X})$, then $\kappa(\Delta) \le 2\kappa_*$*

Proof. Be definition of $\mathcal{C}(\mathcal{X})$, for all $\gamma^* \in \mathcal{C}(X)$, all $x, x' \in \mathcal{X}$, we have

$$(x - x')^\top \gamma^* \le |x^\top \gamma^*| + |x'^\top \gamma^*| \le 2.$$

Then,

$$\kappa(\Delta) \le 2\min_\mu \sum_{x\in\mathcal{X}} \mu(x) \quad \text{such that } \mu \in \mathcal{M}^{\mathcal{X}}_{e_{d+1}} \text{ and } e_{d+1}^\top V(\mu)^+ e_{d+1} \le 1.$$

Let μ_* be the solution of the e_{d+1}-optimal design problem

$$\underset{\mu}{\text{minimize }} e_{d+1}^\top V(\mu)^+ e_{d+1} \text{ such that } \mu \in \mathcal{P}^{\mathcal{X}}_{e_{d+1}}.$$

By definition of κ_*, we see that $e_{d+1}^\top V(\mu_*)^+ e_{d+1} = \kappa_*$. This implies that the measure $\kappa_* \times \mu_*$ verifies the constraints $e_{d+1}^\top V(\kappa_* \times \mu_*)^+ e_{d+1} \le 1$ and $\kappa_* \mu_* \in \mathcal{M}^{\mathcal{X}}_{e_{d+1}}$. Thus,

$$\kappa(\Delta) \le 2\sum_{x\in\mathcal{X}} \kappa_* \mu_*(x) = 2\kappa_*.$$

$\square$

On the regret $\kappa(\Delta)$ The function κ verifies the following properties.

Lemma 8. *For two vectors of gaps Δ, Δ', denote by $\Delta \wedge \Delta'$ (respectively $\Delta \vee \Delta'$) the vector of gaps given by $(\Delta \wedge \Delta')_x = \Delta_x \wedge \Delta'_x$ (respectively $(\Delta \vee \Delta')_x = \Delta_x \vee \Delta'_x$) for all $x \in \mathcal{X}$. Moreover, denote $\Delta \le \Delta'$ if $\Delta_x \le \Delta'_x$ for all $x \in \mathcal{X}$. Then, the following properties hold :*

   i) for all $c > 0$, $\kappa(c\Delta) = c\kappa(\Delta)$;

   ii) if $\Delta \le \Delta'$, then $\kappa(\Delta) \le \kappa(\Delta')$;

   iii) $\kappa(\Delta \vee \Delta') \ge \kappa(\Delta) \vee \kappa(\Delta')$;

   iv) the function $\epsilon \mapsto \kappa(\Delta \vee \epsilon)$ is continuous at 0.

A.7 Computation of G- and Δ-optimal design

Computing the optimal design is a convex problem, for which many algorithms have been proposed. The first method to compute G-optimal design is due to [12] and [44]; later, [39] proposed a multiplicative weight update algorithm. More recently, [40] suggested to use a Semi-Definite Programming approach to solve the G-optimal design problem. Linear programming was used in [17] to compute c-optimal design, while [34] studied a SDP formulation of this problem. Reducing

the G-optimal problem to a Mixed-Integer, Second Order Cone Programming, [37] proposed a new algorithm based on interior point methods. We refer the interested reader to the review in [36].

In practice, one can rely on the R package OptimalDesign or the Python Package PICOS [38] to compute G- and c-optimal design.

The following Lemma allows us to reduce the problem of finding a Δ-optimal design to that of a c-optimal design for some rescaled features.

Lemma 9. *For any vector $\Delta \in (0, +\infty)^{\mathcal{X}}$, let π^{Δ} be the e_{d+1}-optimal design relative to the set $\mathcal{A}^{\Delta} = \left\{ \Delta_x^{-1/2} \begin{pmatrix} x \\ z_x \end{pmatrix} : x \in \mathcal{X} \right\}$ and let $\kappa^{\Delta} = e_{d+1}^{\top} V(\pi^{\Delta})^{+} e_{d+1}$ be the e_{d+1}-optimal variance relative to $\mathcal{A}^{\Delta}$. Then, the Δ-optimal design μ^{Δ} is given by $\mu^{\Delta}(x) = \kappa^{\Delta} \pi^{\Delta}(x) \Delta_x^{-1}$ for all $x \in \mathcal{X}$. In addition, the support of μ^{Δ} can be chosen to be of cardinnality at most $d + 1$.*

Thus, Lemma 9 shows that to compute the Δ-optimal design, one should follow these steps :

1. Compute the rescaled features $\mathcal{A}^{\Delta}$;

2. Compute the e_{d+1}-optimal design π^{Δ} on $\mathcal{A}^{\Delta}$, as well as the variance term $\kappa^{\Delta} = e_{d+1}^{\top} \left(\sum_{x \in \mathcal{X}} \frac{\pi^{\Delta}(x)}{\Delta_x} a_x a_x^{\top} \right)^{+} e_{d+1}$;

3. Compute the Δ-optimal design μ^{Δ} given by $\mu^{\Delta}(x) = \kappa^{\Delta} \pi^{\Delta}(x) \Delta_x^{-1}$ for all $x \in \mathcal{X}$.

B   Detailed Fair Phased Elimination algorithm

We present the notations used in Algorithm 4. The phases are indexed by $l \in \mathbb{N}^*$. The sets $\mathcal{X}_l^{(z)}$ for $z \in \{-1, +1\}$ corresponds to actions in group z that are considered as potentially optimal in phase l. The variable $\widehat{z}_l^*$ encodes the group determined as optimal: it is 0 as long as this group has not been determined. The subscript (z) refer to the group z when $z \in \{-1, +1\}$, and otherwise to the estimation of the bias ω^*: for example, the probability $\pi_l^{(z)}$ for $z \in \{-1, +1\}$ and $l > 1$ corresponds to the approximate G-optimal design on $\mathcal{X}_l^{(z)}$. Then, for $z \in \{-1, +1\}$, allocations $\mu^{(z)}$ (resp. $\mu^{(0)}$) correspond to allocation of samples in the exploration phase $\text{Exp}_l^{(z)}$ (resp. $\text{Exp}_l^{(0)}$). Similarly, $V_l^{(z)}$ (resp $V_l^{(0)}$) denotes the variance matrix of the estimator $\begin{pmatrix} \widehat{\gamma}_l^{(z)} \\ \widehat{\omega}_l^{(z)} \end{pmatrix}$ (resp. $\widehat{\omega}_l^{(0)}$) obtained from observations made during phase $\text{Exp}_l^{(z)}$ (resp. $\text{Exp}_l^{(0)}$). Finally, $\text{Explore}_l^{(z)}$ (resp. $\text{Explore}_l^{(0)}$) is a Boolean variable indicating whether the exploration at phase l for group z (resp. for the bias parameter) has been performed. It is used in the proofs to ensure that the corresponding estimators are well defined.

Algorithm 4 Fair Phased Elimination (detailed version)

1: **Input:** δ, T, $k = |\mathcal{X}|$

2: **Initialize:** Recovery $\leftarrow \emptyset$, $t \leftarrow 0$, $l \leftarrow 1$ $\widehat{z^*_1} \leftarrow 0$,

3: $\qquad \mathcal{X}_1^{(+1)} \leftarrow \{x : z_x = 1\}$, $\mathcal{X}_1^{(-1)} \leftarrow \{x : z_x = -1\}$, $\widehat{\Delta}_x^1 \leftarrow 2$ for $x \in \mathcal{X}$

4: **while** $t < T$ **do**

5: $\quad$ **Initialize:** $\epsilon_l \leftarrow 2^{2-l}$, $\widehat{z^*}_{l+1} \leftarrow \widehat{z^*}_l$, $\widehat{\Delta}^{l+1} \leftarrow \widehat{\Delta}^l$, $\text{Explore}_l^{(z)} \leftarrow$ False for $z \in \{-1, 0, +1\}$

6: $\quad$ **for** $z \in \{-1, +1\}$ such that $z \neq -\widehat{z^*}_l$ **do** $\qquad \qquad \triangleright$ G-optimal Exploration and Elimination

7: $\qquad \pi_l^{(z)} \leftarrow \operatorname*{argmin}_{\pi} \left\{ \max_{x \in \mathcal{X}_l^{(z)}} a_x^\top V(\pi)^+ a_x : \pi \in \mathcal{P}_{\mathcal{X}_l^{(z)}}^{\mathcal{X}_l^{(z)}}, \, |\operatorname{supp}(\pi)| \leq \frac{(d+1)(d+2)}{2} \right\}$

8: $\qquad \mu_l^{(z)}(x) \leftarrow \left\lceil \frac{2(d+1)\pi_l^{(z)}(x)}{\epsilon_l^2} \log\left(\frac{kl(l+1)}{\delta}\right) \right\rceil$ for all $x \in \mathcal{X}_l^{(z)}$

9: $\qquad n_l^{(z)} \leftarrow \sum_{x \in \mathcal{X}_l^{(z)}} \mu_l^{(z)}(x)$, $\text{Exp}_l^{(z)} \leftarrow \left\{ t+1, ..., T \wedge (t + n_l^{(z)}) \right\}$

10: $\qquad$ **if** $t + n_l^{(z)} \leq T$ **then**

11: $\qquad \qquad \text{Explore}_l^{(z)} \leftarrow$ True, choose each action $x \in \mathcal{X}_l^{(z)}$ exactly $\mu_l^{(z)}(x)$ times

12: $\qquad \qquad V_l^{(z)} \leftarrow \sum_{t \in \text{Exp}_l^{(z)}} a_{x_t} a_{x_t}^\top$, $\widehat{\theta}_l^{(z)} \leftarrow \left(V_l^{(z)} \right)^+ \sum_{t \in \text{Exp}_l^{(z)}} y_t a_{x_t}$

13: $\qquad \qquad \mathcal{X}_{l+1}^{(z)} \leftarrow \left\{ x \in \mathcal{X}_l^{(z)} : \max_{x' \in \mathcal{X}_l^{(z)}} (a_{x'} - a_x)^\top \widehat{\theta}_l^{(z)} \leq 3\epsilon_l \right\}$

14: $\qquad$ **else** for $t \in \text{Exp}_l^{(z)}$, sample empirical best action in $\mathcal{X}_l^{(z)}$

15: $\qquad t \leftarrow t + n_l^{(z)}$

16: $\quad$ **if** $\widehat{z^*}_l = 0$ **then**

17: $\qquad$ compute the $\widehat{\Delta}^l$-optimal design $\widehat{\mu}_l$ and the corresponding regret $\kappa(\widehat{\Delta}^l)$

18: $\qquad$ **if** $\epsilon_l \leq \left(\kappa(\widehat{\Delta}^l) \log(T)/T \right)^{1/3}$ **then** $\qquad \qquad \qquad \triangleright$ Recovery phase

19: $\qquad \qquad$ Recovery $\leftarrow \{t, ..., T\}$

20: $\qquad \qquad$ sample empirical best action in $\mathcal{X}_{l+1}^{(-1)} \cup \mathcal{X}_{l+1}^{(1)}$ until the end of the budget, $t \leftarrow T$

21: $\qquad$ **else** $\qquad \qquad \qquad \qquad \qquad \qquad \qquad \triangleright \widehat{\Delta}^l$-optimal Exploration and Elimination

22: $\qquad \qquad \mu_l^{(0)}(x) \leftarrow \left\lceil \frac{2\widehat{\mu}_l(x)}{\epsilon_l^2} \log\left(\frac{l(l+1)}{\delta}\right) \right\rceil$ for all $x \in \mathcal{X}$

23: $\qquad \qquad n_l^{(0)} \leftarrow \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x)$, $\text{Exp}_l^{(0)} \leftarrow \left\{ t, ..., T \wedge (t + n_l^{(0)}) \right\}$

24: $\qquad \qquad$ **if** $t + n_l^{(0)} \leq T$ **then**

25: $\qquad \qquad \qquad \text{Explore}_l^{(0)} \leftarrow$ True, choose each action $x \in \mathcal{X}$ exactly $\mu_l^{(0)}(x)$ times

26: $\qquad \qquad \qquad V_l^{(0)} \leftarrow \sum_{t \in \text{Exp}_l^{(0)}} a_{x_t} a_{x_t}^\top$, $\widehat{\omega}_l^{(0)} \leftarrow e_{d+1}^\top \left(V_l^{(0)} \right)^+ \sum_{t \in \text{Exp}_l^{(0)}} y_t a_{x_t}$

27: $\qquad \qquad \qquad$ **for** $x \in \mathcal{X}_{l+1}^{(-1)} \cup \mathcal{X}_{l+1}^{(1)}$ **do**

28: $\qquad \qquad \qquad \qquad \widehat{m}_{l,x} \leftarrow a_x^\top \widehat{\theta}_l^{(z_x)} - z_x \widehat{\omega}_l^{(0)}$

29: $\qquad \qquad \qquad \qquad \widehat{\Delta}_x^{l+1} \leftarrow \left(\max_{x' \in \mathcal{X}_{l+1}^{(-1)} \cup \mathcal{X}_{l+1}^{(1)}} \widehat{m}_{l,x'} - \widehat{m}_{l,x} + 4\epsilon_l \right) \wedge 2$

30: $\qquad \qquad \qquad$ **for** $z \in \{-1, +1\}$ **do**

31: $\qquad \qquad \qquad \qquad$ **if** $\max_{x \in \mathcal{X}_{l+1}^{(z)}} \widehat{m}_{l,x} - 2\epsilon_l \geq \max_{x \in \mathcal{X}_{l+1}^{(-z)}} \widehat{m}_{l,x} + 2\epsilon_l$ **then** $\widehat{z^*}_{l+1} \leftarrow z$

32: $\qquad \qquad$ **else** sample empirical best action in $\mathcal{X}_{l+1}^{(-1)} \cup \mathcal{X}_{l+1}^{(1)}$ until the end of the budget, $t \leftarrow T$

33: $\qquad \qquad t \leftarrow t + n_l^{(0)}$

34: $\quad l \leftarrow l + 1$

C Proofs

Before proving the main results our this paper, we begin by outlining in Section C.1 the main ideas used to obtain upper and lower bounds on the regret. Then, Theorem 1 is proved in Section C.2, Theorem 2 is proved in Section C.3, Theorem 3 is proved in Section C.4, and Theorem 4 is proved in Section C.5. Extension of Theorem 4 to $d = 2$ and $d = 3$ is discussed in Section C.6. Finally, auxiliary lemmas are proved in Appendix C.7.

For an event $\mathcal{F}$ such that $\mathbb{P}(\mathcal{F}) > 0$, we denote by $\mathbb{E}_{|\mathcal{F}}$ (resp. $\mathbb{P}_{|\mathcal{F}}$) the expectation (resp. the probability) conditionally on $\mathcal{F}$.

C.1 Outline of the proofs

C.1.1 Outline of the proof of Theorem 1

The proof of Theorem 1 can be found in Appendix C.2. We outline here the keys ingredients to this proofs. We begin by introducing some notations.

Notations We denote by L_T the largest integer l such that $\epsilon_l \geq \kappa_*^{1/3} T^{-1/3} \log(T)^{1/3}$. We denote by $L^{(0)}$ the last phase where $\widehat{\Delta}^l$-optimal Exploration and Elimination happens. We denote by $\mathrm{Exp}_l^{(z)}$ the time indices where G-exploration is performed on $\mathcal{X}_l^{(z)}$ and by $\mathrm{Exp}_l^{(0)}$ the time indices where Δ-exploration is performed at phase l. We also denote by Recovery the time indices subsequent to the stopping criterion, this set being empty when the stopping criterion is not activated.

We define a "good" event $\overline{\mathcal{F}}$ such that the errors $\left| a_x^\top \left(\theta^* - \widehat{\theta}_l \right) \right|$ and $|\omega^* - \widehat{\omega}_l^{(0)}|$ are smaller than ϵ_l for all l such that these quantities are defined, and all $x \in \mathcal{X}_l^{(-1)}$ and $\mathcal{X}_l^{(+1)}$. In the following, we use c, c' to denote positive absolute constants, which may vary from line to line. With these notations, we decompose the regret as follows :

$$
\begin{aligned}
R_T \leq\ & 2T\mathbb{P}(\mathcal{F}) + \mathbb{E}_{|\overline{\mathcal{F}}}\underbrace{\left[\sum_{l \leq L_T} \sum_{z \in \{-1,+1\}} \sum_{t \in \mathrm{Exp}_l^{(z)}} (x^* - x_t)^\top \gamma^* \right]}_{R_T^G} + \mathbb{E}_{|\overline{\mathcal{F}}}\underbrace{\left[\sum_{l \leq L^{(0)}} \sum_{t \in \mathrm{Exp}_l^{(0)}} (x^* - x_t)^\top \gamma^* \right]}_{R_T^\Delta} \\
& + \mathbb{E}_{|\overline{\mathcal{F}}}\underbrace{\left[\sum_{l \geq L_T + 1} \sum_{z \in \{-1,+1\}} \sum_{t \in \mathrm{Exp}_l^{(z)}} (x^* - x_t)^\top \gamma^* + \sum_{t \in \mathrm{Recovery}} (x^* - x_t)^\top \gamma^* \right]}_{R_T^{Rec}}.
\end{aligned}
$$

Bound on $T\mathbb{P}(\mathcal{F})$. Using arguments based on concentration of Gaussian variables, we show that $\mathbb{P}(\mathcal{F}) \leq 2T^{-1}$.

Bound on R_T^G. We show that on $\overline{\mathcal{F}}$, only actions with gaps smaller than $c\epsilon_l$ remain in the sets $\mathcal{X}_l^{(-1)}$ and $\mathcal{X}_l^{(+1)}$. The length of each G-optimal Exploration and Elimination phase is of the order $d\log(klT)/\epsilon_l^2$, so the regret of each phase is of the order $d\log(klT)/\epsilon_l$. Summing over the different phases, we find that

$$ R_T^G \leq cd\log(kL_T T)/\epsilon_{L_T}. \tag{7} $$

Using the definition of L_T, we find that $R_T^G \leq cd\log(kL_T T)\kappa_*^{-1/3}\log(T)^{-1/3}T^{1/3}$.

Bound on R_T^Δ. We show that on $\mathcal{F}$, $\widehat{\Delta}^l \geq \Delta$ for all $l \geq 1$. Then, our choice of design $\mu_l^{(0)}$ ensures that for $l \leq L^{(0)}$, on $\overline{\mathcal{F}}$,

$$
\sum_{t \in \mathrm{Exp}_l^{(0)}} (x^* - x_t)^\top \gamma^* \leq c \left(\frac{\log(l(l+1)T)}{\epsilon_l^2} \kappa(\widehat{\Delta}^l) + d + 1 \right)
$$

for some constant $c > 0$. Summing over the different phases, we find that

$$R_T^\Delta \le c\kappa(\widehat{\Delta}^{L^{(0)}}) \log(L^{(0)}T)/\epsilon_{L^{(0)}}^2. \tag{8}$$

Now, the algorithm does not enter the Recovery phase before phase $L^{(0)} + 1$, so we must have $\epsilon_{L^{(0)}}^{-2} \le T^{2/3} \log(T)^{-2/3} \kappa(\widehat{\Delta}^{L^{(0)}})^{-2/3}$. This implies that $R_T^\Delta \le c\kappa(\widehat{\Delta}^{L^{(0)}})^{1/3} \log(T)^{1/3} T^{2/3}$. Since $\kappa(\widehat{\Delta}^l) \le 2\kappa_*$, we find that $R_T^\Delta \le c'\kappa_*^{1/3} \log(T)^{1/3} T^{2/3}$.

Bound on $\mathbf{R_T^{Rec}}$. On the one hand, the actions selected during the Phases $\text{Exp}_l^{(-1)}$ and $\text{Exp}_l^{(+1)}$ for $l \ge L_T + 1$ are sub-optimal by a gap at most $c\epsilon_{L_T}$ on the event $\overline{\mathcal{F}}$. On the other hand, if the algorithm enters the Recovery phase at a phase l, then

$$\epsilon_l \le \kappa(\widehat{\Delta}^{L^{(0)}})^{1/3} T^{-1/3} \log(T)^{1/3} \le \kappa_*^{1/3} T^{-1/3} \log(T)^{1/3},$$

so we must have $l = L^{(0)} + 1 \ge L_T + 1$. Therefore, all actions selected during the Recovery phase are sub-optimal by a gap at most $c\epsilon_{L_T}$. Then, R_T^{Rec} can be bounded as $R_T^{Rec} \le c\epsilon_{L_T}T$. This implies in particular that $R_T^{Rec} \le c'\kappa_*^{1/3} \log(T)^{1/3} T^{2/3}$.

When $T \ge T_{\kappa_*,d,k}$ for some $T_{\kappa,d,k}$ large enough, we find that $\mathbb{R}_T \le c'\kappa_*^{1/3} \log(T)^{1/3} T^{2/3}$.

C.1.2 Outline of the Proof of Theorem 2

The proof of Theorem 2 is close to that of Theorem 1, and we adopt the same notations as in the proof sketch above.

Notations We denote by $L^{(0)}$ the last phase where $\widehat{\Delta}^l$-optimal Exploration and Elimination happens. We denote $\overline{\mathcal{F}}$ some "good" event such that the errors $|a_x^\top(\theta^* - \widehat{\theta}_l^{(z_x)})|$ and $|\omega^* - \widehat{\omega}_l^{(0)}|$ are smaller than ϵ_l for all l such that these quantities are defined, and all $x \in \mathcal{X}_l^{(-1)} \cup \mathcal{X}_l^{(+1)}$. We denote by $\text{Exp}_l^{(z)}$ the time indices where G-exploration is performed on $\mathcal{X}_l^{(z)}$ and by $\text{Exp}_l^{(0)}$ the time indices where Δ-exploration is performed at phase l. We also denote by Recovery the time indices subsequent to the stopping criterion, this set being empty when the stopping criterion is not activated. In the following, we use c, c' to denote positive absolute constants, which may vary from line to line.

Fact 1 Let $l_{\Delta_{\min}}$ be the largest integer such that $\epsilon_{l_{\Delta_{\min}}} \ge C\Delta_{\min}$ for some well-chosen absolute constant $C > 0$. We show that on the good event $\overline{\mathcal{F}}$, no more than $l_{\Delta_{\min}}$ G-optimal Exploration and Elimination phases are needed to find the best action. For all phases $l \ge l_{\Delta_{\min}}$, the algorithm always chooses x^*, and suffers no regret.

Fact 2 We show that on the good event $\overline{\mathcal{F}}$, for each phase l, $\widehat{\Delta}^l \le c(\Delta \vee \epsilon_l)$ for some constant c. Lemma 8 then implies that for all $l \le L^{(0)}$ and all $\tau > 0$, $\kappa(\widehat{\Delta}^l) \le c\kappa(\Delta \vee \epsilon_l) \le c(1 + \epsilon_l\tau^{-1})\kappa(\Delta \vee \tau)$.

Fact 3 Let $l_{\Delta_{\ne}}$ be the largest integer such that $\epsilon_{l_{\Delta_{\ne}}} \ge C\Delta_{\ne}$ for some well-chosen absolute constant $C > 0$. On the good event $\overline{\mathcal{F}}$, if the algorithm enters the $\widehat{\Delta}^l$-optimal Exploration and Elimination phase at round $l \ge l_{\Delta_{\ne}}$, we show that the algorithm finds the best group at this phase. This implies that $L^{(0)} \le l_{\Delta_{\ne}}$.

Fact 4 We denote by L_T the largest integer l such that $\epsilon_l \ge (\kappa_* \log(T)/T)^{1/3}$. Since $\kappa_* \ge \kappa(\widehat{\Delta}^l)$ for all $l \ge 1$, we see that if the algorithm enters the Recovery phase, we must have $L_T \le L^{(0)}$, and $\epsilon_{L^{(0)}} \le \epsilon_{L_T} \approx \varepsilon_T$.

Using **Fact 1**, we find that the regret can be written as

$$
R_T \leq 2T\mathbb{P}(\mathcal{F}) + \mathbb{E}_{|\overline{\mathcal{F}}}\underbrace{\left[\sum_{l \leq l_{\Delta_{\min}}} \sum_{z \in \{-1,+1\}} \sum_{t \in \mathrm{Exp}_l^{(z)}} (x^* - x_t)^\top \gamma^* \right]}_{R_T^G}
$$

$$
+ \mathbb{E}_{|\overline{\mathcal{F}}}\underbrace{\left[\sum_{l \leq L^{(0)}} \sum_{t \in \mathrm{Exp}_l^{(0)}} (x^* - x_t)^\top \gamma^* \right]}_{R_T^\Delta} + \mathbb{E}_{|\overline{\mathcal{F}}}\underbrace{\left[\sum_{t \in \mathrm{Recovery}} (x^* - x_t)^\top \gamma^* \right]}_{R_T^{Rec}}.
$$

Bound on R_T^G. We rely on arguments similar to those used in Equation (7) to show that $R_T^G \leq c(d+1)\log(kl_{\Delta_{\min}}T)\epsilon_{l_{\Delta_{\min}}}^{-1}$. Since $\epsilon_{l_{\Delta_{\min}}} \geq C\Delta_{\min}$, this implies that

$$
R_T^G \leq \frac{c(d+1)\log(kl_{\Delta_{\min}}T)}{\Delta_{\min}} \leq \frac{c'd\log(T)}{\Delta_{\min}}
$$

if $T \geq k$.

Bound on $R_T^\Delta + R_T^{Rec}$. We begin by bounding R_T^Δ. Recall that Equation (8) states that $R_T^\Delta \leq c\kappa(\widehat{\Delta}^{L^{(0)}})\log(l_{L^{(0)}}T)\epsilon_{L^{(0)}}^{-2}$. Using **Fact 2**, we find that for any $\tau > 0$,

$$
R_T^\Delta \leq c\kappa(\Delta \vee \tau)\log(l_{L^{(0)}}T)\left(\epsilon_{L^{(0)}}^{-2} + \epsilon_{L^{(0)}}^{-1}\tau^{-1}\right). \tag{9}
$$

Let us now consider two cases, corresponding to Recovery$= \emptyset$ and Recovery$\neq \emptyset$.

Case 1: Recovery$= \emptyset$. On the one hand, our case assumption implies that

$$
R_T^{Rec} = 0.
$$

On the other hand, by **Fact 3**, we know that on $\overline{\mathcal{F}}$, $L^{(0)} \leq l_{\Delta_{\neq}}$. Then, using the definition of $l_{\Delta_{\neq}}$ and Equation (9) with $\tau = \Delta_{\neq}$, we find that

$$
R_T^\Delta \leq c\kappa(\Delta \vee \Delta_{\neq})\log(L^{(0)}T)\Delta_{\neq}^{-2}.
$$

Case 2: Recovery$\neq \emptyset$. All actions selected during the Recovery phase belong to $\mathcal{X}_{L^{(0)}+1}^{(-1)} \cup \mathcal{X}_{L^{(0)}+1}^{(+1)}$, so on $\overline{\mathcal{F}}$ these actions are sub-optimal by a gap at most $c\epsilon_{L^{(0)}+1}$, so $R_T^{Rec} \leq cT\epsilon_{L^{(0)}+1}$. Now, since the algorithm enters the Recovery phase, we must have $\epsilon_{L^{(0)}+1} \leq (\kappa(\Delta^{L^{(0)}+1})\log(T)/T)^{1/3}$, which implies that

$$
R_T^{Rec} \leq \frac{c\kappa(\widehat{\Delta}^{L^{(0)}+1})\log(T)}{\epsilon_{L^{(0)}+1}^2}.
$$

Using **Fact 2** with $\tau = \epsilon_{L^{(0)}}$ together with Equation (9), we find that

$$
R_T^\Delta + R_T^{Rec} \leq \frac{c\kappa(\Delta \vee \epsilon_{L^{(0)}})\log(T)}{\epsilon_{L^{(0)}}^2}.
$$

On the one hand, **Fact 3** guarantees that, since we entered the Recovery phase before finding the best group, we must have $\epsilon_{L^{(0)}} \geq \epsilon_{l_{\Delta_{\neq}}}$. On the other hand, **Fact 4** ensures that $\epsilon_{L^{(0)}} \leq \varepsilon_T$. Thus,

$$
R_T^{Rec} \leq \frac{c\kappa(\Delta \vee \varepsilon_T)\log(T)}{\Delta_{\neq}^2}.
$$

Conclusion Combining these results, we find that

$$
R_T \leq c\left(\frac{d}{\Delta_{\min}} \vee \frac{\kappa(\Delta \vee \Delta_{\neq})}{\Delta_{\neq}^2} \vee \frac{\kappa(\Delta \vee \varepsilon_T)}{\Delta_{\neq}^2}\right)\log(T)
$$

when $T \geq k$. Using Lemma 8, we get that $\kappa(\Delta \vee \Delta_{\neq}) \vee \kappa(\Delta \vee \varepsilon_T) \leq \kappa(\Delta \vee \Delta_{\neq} \vee \varepsilon_T)$, which concludes the proof of the results.

C.1.3 Outline of the Proof of Theorem 4

We outline the main ingredients used to prove Theorem 4. Theorem 3 relies on similar arguments.

To prove the lower bounds, we need to construct two close problem instances with optimal actions belonging to different groups - to obtain the part of the lower bound involving $\Delta_{\neq}$ - and in addition we must also create confusing instances with different optimal actions belonging to a same group - to obtain the part of the lower bound involving $\Delta_{\min}$. This is done by considering the following set of actions and of problems.

Lemma 10. *Set* $\mathcal{A} = \left\{ \begin{pmatrix} x_1 \\ z_{x_1} \end{pmatrix}, ..., \begin{pmatrix} x_{d+1} \\ z_{x_{d+1}} \end{pmatrix} \right\}$, *where* $\begin{pmatrix} x_i \\ z_{x_i} \end{pmatrix} = e_i + e_{d+1}$, *for* $i \in \{2, ..., \lfloor d/2 \rfloor\}$, $\begin{pmatrix} x_i \\ z_{x_i} \end{pmatrix} = e_i - e_{d+1}$ *for* $i \in \{\lfloor d/2 \rfloor + 1, ..., d\}$, *and* $\begin{pmatrix} x_{d+1} \\ z_{x_{d+1}} \end{pmatrix} = -\left(1 - \frac{2}{\sqrt{\kappa_*}+1}\right) e_1 - e_{d+1}$. *It holds that*

$$\min_{\pi \in \mathcal{P}_{e_{d+1}}^{\mathcal{A}}} \left\{ e_{d+1}^\top \left(\sum_{\left(\begin{smallmatrix} x \\ z \end{smallmatrix} \right) \in \mathcal{A}} \pi(x) \begin{pmatrix} x \\ z_x \end{pmatrix} \begin{pmatrix} x \\ z_x \end{pmatrix}^\top \right)^+ e_{d+1} \right\} = \kappa_*.$$

We also define the following parameters:

$$\gamma^{(1)} = \frac{1 + \Delta_{\neq} - \Delta_{\min}}{2} \left(\sum_{1 \le j \le \lfloor d/2 \rfloor} e_j \right) + \frac{1 - \Delta_{\neq} - \Delta_{\min}}{2} \left(\sum_{\lfloor d/2 \rfloor + 1 \le j \le d} e_j \right)$$
$$+ \Delta_{\min} e_1 + \Delta_{\min} e_{\lfloor d/2 \rfloor + 1}$$

$$\gamma^{(i)} = \gamma^{(1)} + 2\Delta_{\min} e_i + 2\Delta_{\min} e_{\lfloor d/2 \rfloor + i} \quad \forall i \in \{2, ..., \lfloor d/2 \rfloor\}$$

$$\gamma^{(\lfloor d/2 \rfloor + 1)} = \frac{1 - \Delta_{\neq} - \Delta_{\min}}{2} \left(\sum_{1 \le j \le \lfloor d/2 \rfloor} e_j \right) + \frac{1 + \Delta_{\neq} - \Delta_{\min}}{2} \left(\sum_{\lfloor d/2 \rfloor + 1 \le j \le d} e_j \right)$$
$$+ \Delta_{\min} e_1 + \Delta_{\min} e_{\lfloor d/2 \rfloor + 1}.$$

The bias parameters are given by $\omega^{(i)} = -\frac{\Delta_{\neq}}{2} \; \forall i \in \{1, ..., \lfloor d/2 \rfloor\}$, and $\omega^{(\lfloor d/2 \rfloor + 1)} = \frac{\Delta_{\neq}}{2}$. The parameters $\theta^{(i)} = \begin{pmatrix} \gamma^{(i)} \\ \omega^{(i)} \end{pmatrix}$ characterize $\lfloor d/2 \rfloor + 1$ problems, with noise distribution i.i.d. $\mathcal{N}(0,1)$. We write **Problem i** for the problem characterized by $\theta^{(i)}$. Note that by construction and for any $i \in \{1, ..., \lfloor d/2 \rfloor + 1\}$, we have that $\theta^{(i)} \in \Theta_{\Delta_{\min}, \Delta_{\neq}}^{\mathcal{A}}$.

The following facts hold:

- For any $i \in \{1, ..., \lfloor d/2 \rfloor + 1\}$, action x_i is the unique optimal action in **Problem i**. Since $1/2 \ge \Delta_{\neq} \ge \Delta_{\min}$, sampling any other (sub-optimal) action leads to an instantaneous regret of at least $\Delta_{\min}$. Moreover, choosing an action in the group $-z_i$ leads to an instantaneous regret of at least $\Delta_{\neq}$.

- In **Problem i** for any $i \in \{1, ..., \lfloor d/2 \rfloor + 1\}$, action $d+1$ is very sub-optimal and sampling it leads to an instantaneous regret higher than $(1 - 2/(\sqrt{\kappa_*} + 1))(1 - \Delta_{\neq} + \Delta_{\min}) + (1 + \Delta_{\neq} + \Delta_{\min})/2 \ge 1/2$, since $\kappa_* \ge 1$ and $1/2 \ge \Delta_{\neq} \ge \Delta_{\min}$. This action is the worst action in all problems.

- Many actions are such that their distributions are the same across problems. More specifically:
  - For any $i \in \{2, ..., \lfloor d/2 \rfloor\}$, between **Problem 1** and **Problem i**, the only actions that provide different evaluations when sampled are action i and action $\lfloor d/2 \rfloor + i$, and the mean difference between the evaluations in both cases is $2\Delta_{\min}$.
  - Between **Problem 1** and **Problem** $\lfloor d/2 \rfloor + 1$, the only actions that provide different evaluations when sampled is action $d+1$, and the mean gap in this case is $\frac{2}{\sqrt{\kappa_*}+1} \Delta_{\neq} := \alpha \Delta_{\neq}$.

The proof is then divided in two parts, one part for proving the part of the bound depending on $\Delta_{\min}$ and one part for proving the part of the bound depending on $\Delta_{\neq}$.

Part of the bound depending on $\Delta_{\min}$. This part of the proof is obtained using classical arguments for K-armed bandit problems. For $i \in \{2, ..., \lfloor d/2 \rfloor\}$, all actions but x_i and $x_{\lfloor d/2 \rfloor + i}$ have the same feedback under **Problem 1** and **Problem i**. On the other hand, the average feedback for actions x_i and $x_{\lfloor d/2 \rfloor + i}$ differs by $2\Delta_{\min}$, so either action needs to be selected approximately $\frac{\log(T)}{\Delta_{\min}^2}$ times in order to identify the problem at hand with high enough probability. In **Problem 1**, the simple regret for choosing x_i or $x_{\lfloor d/2 \rfloor + i}$ is larger than $\Delta_{\min}$, so the total regret obtained when doing this is at least of the order $\frac{\log(T)}{\Delta_{\min}}$. Summing over the different actions i leads to a lower bound of the order $\frac{d \log(T)}{\Delta_{\min}}$.

Part of the bound depending on $\Delta_{\neq}$. To obtain the second part of the lower bound, we note that all actions but x_{d+1} have the same feedback under **Problem 1** and **Problem** $\lfloor d/2 \rfloor + 1$. The average feedback for actions x_{d+1} differs by $\alpha \Delta_{\neq}$ under these parameters, so action x_{d+1} needs to be selected approximately $\frac{\log(T)}{\alpha^2 \Delta_{\neq}^2} \gtrsim \frac{\log(T) \kappa_*}{\Delta_{\neq}^2}$ times to identify the problem at hand with high enough probability. Since selecting action x_{d+1} leads to an simple regret larger than $1/2$ under **Problem 1**, this implies that the regret must be at least of the order $\frac{\kappa_* \log(T)}{\Delta_{\neq}^2}$.

Bounds on $\kappa(\Delta)$ Finally, the following lemma allows to express $\kappa(\Delta)$ as a function of κ_*.

Lemma 11. *For any $i \in \{1, ..., \lfloor d/2 \rfloor + 1\}$, the gap vector Δ verifies*

$$\kappa(\Delta) = \frac{(1 + \sqrt{\kappa_*})^2 \Delta_{d+1}}{4}$$

where $\Delta_{d+1} = \max_i (x_i - x_{d+1})^\top \gamma^{(i)}$.

On the one hand, since $\kappa_* \geq 1$, we see that $\kappa_* \leq (1 + \sqrt{\kappa_*})^2 \leq 4\kappa_*$. On the other hand, $1/2 \leq \Delta_{d+1} \leq 2$, so $\kappa(\Delta) \in \left[\frac{\kappa_*}{8}, 2\kappa_*\right]$.

C.2 Proof of Theorem 1

We begin by defining for $z \in \{-1, 0, +1\}$

$$L^{(z)} = \max \left\{ l \geq 1 : \text{Explore}_l^{(z)} = \text{True} \right\}$$

the largest integer l such that $\text{Explore}_l^{(z)} = \text{True}$. Recall that κ_* is the e_{d+1}-optimal variance. By definition of the algorithm, for all $l \leq L^{(0)} + 1$, $\widehat{\Delta}^l \leq 2$, so $\kappa(\widehat{\Delta}^l) \leq 2\kappa_*$. Now, let us also define

$$L_T = \max \left\{ l \geq 1 : \epsilon_l > \left(\frac{2\kappa_* \log(T)}{T} \right)^{1/3} \right\}.$$

Then, if Recovery$\neq \emptyset$, we must have $L^{(0)} \geq L_T$. Moreover, we see that since $\epsilon_{L_T} = 2^{2 - L_T}$, we have $L_T \leq 2 + \frac{\log_2(T/(2\kappa_* \log(T)))}{3} \leq 3 \log_2(T)$ when $T > 1$.

We define a "bad" event $\mathcal{F}$, such that, on $\overline{\mathcal{F}}$, our estimators $\widehat{\gamma}_l^{(z)}$ and $\widehat{\omega}_l^{(z)}$ are close to the true parameters γ^* and ω^* for all rounds l. More precisely, let

$$\mathcal{F} = \bigcup_{l \geq 1} \mathcal{F}_l, \tag{10}$$

where for $l \geq 1$

$$\mathcal{F}_l = \left\{ \exists z \in \{-1, 1\} \text{ such that Explore}_l^{(z)} = \text{True, and } x \in \mathcal{X}_l^{(z)} \text{ such that } \left| \begin{pmatrix} \widehat{\gamma}_l^{(z)} - \gamma^* \\ \widehat{\omega}_l^{(z)} - \omega^* \end{pmatrix}^\top \begin{pmatrix} x \\ z_x \end{pmatrix} \right| \geq \epsilon_l \right\}$$

$$\bigcup \left\{ \text{Explore}_l^{(0)} = \text{True and } \left| \widehat{\omega}_l^{(0)} - \omega^* \right| \geq \epsilon_l \right\}.$$

Then, the regret decomposes as

$$R_T \leq \sum_{t \leq T} \mathbb{E}_{|\overline{\mathcal{F}}} \left[(x^* - x_t)^\top \gamma^* \right] + 2T \mathbb{P}\left[\mathcal{F} \right]. \tag{11}$$

The following lemma relies on concentration of Gaussian variables to bound the probability of the event $\mathcal{F}$.

Lemma 12. $\mathbb{P}(\mathcal{F}) \leq 2\delta$.

Now, the first term of (11) can be decomposed as

$$\sum_{t \leq T}(x^* - x_t)^\top \gamma^* \leq \sum_{z \in \{-1,0,+1\}} \sum_{l=1}^{L^{(z)}+1} \sum_{t \in \text{Exp}_l^{(z)}} (x^* - x_t)^\top \gamma^* + \sum_{t \in \text{Recovery}} (x^* - x_t)^\top \gamma^*,$$

where we use as convention that the sum over an empty set is null. Note that for $z \in \{-1, +1\}$, during the phase $\text{Exp}_l^{(z)}$ the algorithm only samples actions from $\mathcal{X}_l^{(z)}$. By contrast, during the phase $\text{Exp}_l^{(0)}$, even actions eliminated from the sets $\mathcal{X}_l^{(z)}$ can be sampled. Finally, if the algorithm stops during phase $\text{Exp}_{L^{(0)}+1}^{(0)}$, but does not have enough budget to complete the last $\widehat{\Delta}^l$-optimal Exploration and Elimination Phase, it samples the remaining actions in the set $\mathcal{X}_{L^{(0)}+2}^{(-1)} \cup \mathcal{X}_{L^{(0)}+2}^{(+1)}$. Hence, the first term of (11) can be upper-bounded by

$$\sum_{t \leq T}(x^* - x_t)^\top \gamma^* \leq \sum_{z \in \{-1,+1\}} \sum_{l=1}^{L_T} \left(\sum_{x \in \mathcal{X}_l^{(z)}} \mu_l^{(z)}(x) \right) \max_{x \in \mathcal{X}_l^{(z)}} (x^* - x)^\top \gamma^* \tag{12}$$

$$+ \sum_{z \in \{-1,+1\}} \sum_{l=L_T+1}^{L^{(z)}+1} \sum_{t \in \text{Exp}_l^{(z)}} (x^* - x_t)^\top \gamma^* + \sum_{t \in \text{Recovery}} (x^* - x_t)^\top \gamma^*$$

$$+ \sum_{l=1}^{L^{(0)}} \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x)\Delta_x + \mathbb{1}\left\{\text{Explore}_{L^{(0)}+1}^{(0)} = \text{False}\right\} \sum_{t \in \text{Exp}_{L^{(0)}+1}^{(0)}} \max_{x \in \mathcal{X}_{L^{(0)}+2}^{(-1)} \cup \mathcal{X}_{L^{(0)}+2}^{(+1)}} (x^* - x)^\top \gamma^*.$$

We begin by bounding the sum of the regret corresponding to the Recovery phase and to the phases $\text{Exp}_L^{(z)}$ for $z \in \{-1, +1\}$ and $l > L_T$ on the event $\overline{\mathcal{F}}$.

Bound on $\displaystyle\sum_{z \in \{-1,+1\}} \sum_{l=L_T+1}^{L^{(z)}+1} \sum_{t \in \text{Exp}_l^{(z)}} (x^* - x_t)^\top \gamma^* + \sum_{t \in \text{Recovery}} (x^* - x_t)^\top \gamma^*.$

Lemma 13. *Let $x^* \in \text{argmax}_{x \in \mathcal{X}} x^\top \gamma^*$ be an optimal action. Then, on the event $\overline{\mathcal{F}}$ defined in Equation (10), for $l \geq 1$ such that $\text{Explore}_l^{(z_{x^*})} = $ True,*

$$\mathcal{X}_{l+1}^{(z_{x^*})} \subset \left\{ x \in \mathcal{X}_1^{(z_{x^*})} : (x^* - x)^\top \gamma^* < 10\epsilon_{l+1} \right\}. \tag{13}$$

Moreover, for $l \geq 1$ such that $\text{Explore}_l^{(-z_{x^*})} = $ True,

$$\mathcal{X}_{l+1}^{(-z_{x^*})} \subset \left\{ x \in \mathcal{X}_1^{(-z_{x^*})} : (x^* - x)^\top \gamma^* < 42\epsilon_{l+1} \right\}.$$

Recall that if Recovery $\neq \emptyset$, $L^{(0)} \geq L_T$. Then, all actions sampled during the Recovery phase belong to $\mathcal{X}_{l+1}^{(-1)} \cup \mathcal{X}_{l+1}^{(+1)}$ for some $l \geq L_T$. Lemma 13 shows that, on $\overline{\mathcal{F}}$, for $l \geq L_T$, the actions in $\mathcal{X}_{l+1}^{(z)}$ are sub-optimal by at most $42\epsilon_{L_T+1}$. Then, we get that on the event $\overline{\mathcal{F}}$,

$$\sum_{z \in \{-1,+1\}} \sum_{l=L_T+1}^{L^{(z)}+1} \sum_{t \in \text{Exp}_l^{(z)}} (x^* - x_t)^\top \gamma^* + \sum_{t \in \text{Recovery}} (x^* - x_t)^\top \gamma^* \leq T \times 42\epsilon_{L_T+1}$$

$$\leq 53\kappa_*^{1/3} T^{2/3} \log(T)^{1/3}. \tag{14}$$

Bound on $\sum_{l=1}^{L^{(0)}} \sum_{x\in\mathcal{X}} \mu_l^{(0)}(x)\Delta_x + \mathbb{1}\left\{\text{Explore}_{L^{(0)}+1}^{(0)} = \text{False}\right\} \sum_{t\in\text{Exp}_{L^{(0)}+1}^{(0)}} \max_{x\in\mathcal{X}_{L^{(0)}+2}^{(-1)}\cup\mathcal{X}_{L^{(0)}+2}^{(+1)}} (x^* -$

$x)^\top\gamma^*$.

We begin by bounding $\mathbb{1}\left\{\text{Explore}_{L^{(0)}+1}^{(0)} = \text{False}\right\} \sum_{t\in\text{Exp}_{L^{(0)}+1}^{(0)}} \max_{x\in\mathcal{X}_{L^{(0)}+2}^{(-1)}\cup\mathcal{X}_{L^{(0)}+2}^{(+1)}} (x^*-x)^\top\gamma^*$. Recall

that $n_{L^{(0)}+1}^{(0)} = \sum_{x\in\mathcal{X}} \mu_{L^{(0)}+1}^{(0)}(x)$ is the budget that would be necessary to complete the $\widehat{\Delta}^l$-optimal

Exploration and Elimination phase at phase $L^{(0)} + 1$. On the one hand, Lemma 13 implies that on the event $\overline{\mathcal{F}}$,

$$\mathbb{1}\left\{\text{Explore}_{L^{(0)}+1}^{(0)} = \text{False}\right\} \sum_{t\in\text{Exp}_{L^{(0)}+1}^{(0)}} \max_{x\in\mathcal{X}_{L^{(0)}+2}^{(-1)}\cup\mathcal{X}_{L^{(0)}+2}^{(+1)}} (x^* - x)^\top\gamma^* \leq 42 n_{L^{(0)}+1}^{(0)}\epsilon_{L^{(0)}+2} \leq 21 n_{L^{(0)}+1}^{(0)}\epsilon_{L^{(0)}+1}.$$

On the other hand, for all $l \leq L^{(0)} + 1$, the definition of $\widehat{\Delta}^l$ implies that $\widehat{\Delta}_x^l \geq \epsilon_l$ for all $x \in \mathcal{X}$. Therefore, $21 n_{L^{(0)}+1}^{(0)}\epsilon_{L^{(0)}+1} \leq 21 n_{L^{(0)}+1}^{(0)} \min_x \widehat{\Delta}_x^{L^{(0)}+1}$. This implies that on $\overline{\mathcal{F}}$,

$$\mathbb{1}\left\{\text{Explore}_{L^{(0)}+1}^{(0)} = \text{False}\right\} \sum_{t\in\text{Exp}_{L^{(0)}+1}^{(0)}} \max_{x\in\mathcal{X}_{L^{(0)}+2}^{(-1)}\cup\mathcal{X}_{L^{(0)}+2}^{(+1)}} (x^* - x)^\top\gamma^* \leq 21 \sum_{x\in\mathcal{X}} \mu_{L^{(0)}+1}^{(0)}(x)\widehat{\Delta}_x^{L^{(0)}+1}. \quad (15)$$

Next, to bound the remaining terms of Equation (12), we bound the regret $\sum_{x\in\mathcal{X}} \mu_l^{(0)}(x)\Delta_x$ of exploration phase $\text{Exp}_l^{(0)}$ using the following lemma.

Lemma 14. *For all $l > 0$, and $z \in \{-1, +1\}$, we have*

$$\sum_{x\in\mathcal{X}_l^{(z)}} \mu_l^{(z)}(x) \leq \frac{2(d+1)}{\epsilon_l^2} \log\left(\frac{kl(l+1)}{\delta}\right) + \frac{(d+1)(d+2)}{2}.$$

and on $\overline{\mathcal{F}}$, we have

$$\sum_{x\in\mathcal{X}} \mu_l^{(0)}(x)\Delta_x \leq \sum_{x\in\mathcal{X}} \mu_l^{(0)}(x)\widehat{\Delta}_x^l \leq \frac{2\kappa(\widehat{\Delta}^l)}{\epsilon_l^2} \log\left(\frac{l(l+1)}{\delta}\right) + 2(d+1).$$

Then, Equation (15) and Lemma 14 imply that on $\overline{\mathcal{F}}$

$$\sum_{l=1}^{L^{(0)}} \sum_{x\in\mathcal{X}} \mu_l^{(0)}(x)\Delta_x + \mathbb{1}\left\{\text{Explore}_{L^{(0)}+1}^{(0)} = \text{False}\right\} \sum_{t\in\text{Exp}_{L^{(0)}+1}^{(0)}} \max_{x\in\mathcal{X}_{L^{(0)}+2}^{(-1)}\cup\mathcal{X}_{L^{(0)}+2}^{(+1)}} (x^* - x)^\top\gamma^*$$

$$\leq 21 \sum_{l=1}^{L^{(0)}+1} \sum_{x\in\mathcal{X}} \mu_l^{(0)}(x)\widehat{\Delta}_x^l$$

$$\leq 42 \sum_{l=1}^{L^{(0)}+1} \frac{\kappa(\widehat{\Delta}^l)}{\epsilon_l^2} \log\left(\frac{l(l+1)}{\delta}\right) + 42(d+1)(L^{(0)}+1)$$

$$(16)$$

We rely on the following Lemma to bound $\kappa(\widehat{\Delta}^l)$.

Lemma 15. *On $\overline{\mathcal{F}}$, we have for any $l \geq 1$ and any $\tau > 0$*

$$\kappa(\widehat{\Delta}^l) \leq 513 \left(1 + \frac{\epsilon_l}{\tau}\right) \kappa(\Delta \vee \tau).$$

and

$$\kappa(\widehat{\Delta}^l) \geq \kappa(\Delta \vee \epsilon_l).$$

Lemma 14 and Lemma 15 with $\tau = \epsilon_{L^{(0)}}$ imply that on $\overline{\mathcal{F}}$,

$$
\sum_{l=1}^{L^{(0)}+1} \frac{\kappa(\widehat{\Delta}^l)}{\epsilon_l^2} \log\left(\frac{l(l+1)}{\delta}\right) \leq 513\kappa(\Delta \vee \epsilon_{L^{(0)}}) \log\left(\frac{(L^{(0)}+1)(L^{(0)}+2)}{\delta}\right) \left(\sum_{l=1}^{L^{(0)}+1} \frac{1}{\epsilon_l^2} + \sum_{l=1}^{L^{(0)}+1} \frac{1}{\epsilon_l \epsilon_{L^{(0)}}}\right)
$$

$$
\leq 513\kappa(\Delta \vee \epsilon_{L^{(0)}}) \log\left(\frac{6L^{(0)}}{\delta}\right) \left(\frac{16}{\epsilon_{L^{(0)}}^2} + \frac{4}{\epsilon_{L^{(0)}}^2}\right)
$$

$$
\leq 10260 \log\left(\frac{6L^{(0)}}{\delta}\right) \frac{\kappa(\widehat{\Delta}^{L^{(0)}})}{\epsilon_{L^{(0)}}^2} \tag{17}
$$

where the last line follows from the second claim of Lemma 15. Now, by definition of $L^{(0)}$, $\epsilon_{L^{(0)}} \geq \left(\kappa(\widehat{\Delta}^{L^{(0)}}) \log(T)/T\right)^{1/3}$. Then, Equation (17) implies that

$$
\sum_{l=1}^{L^{(0)}+1} \frac{\kappa(\widehat{\Delta}^l)}{\epsilon_l^2} \log\left(\frac{l(l+1)}{\delta}\right) \leq 10260 \log\left(\frac{6L^{(0)}}{\delta}\right) \kappa(\widehat{\Delta}^{L^{(0)}})^{1/3} \log(T)^{-2/3} T^{2/3}. \tag{18}
$$

Moreover, we observe that during each phase l, but the last one, we sample at least

$$
\max_{z \in \{-1,1\}} \sum_{x \in \mathcal{X}_l^{(z)}} \tau_{l,x}^{(z)} \geq \frac{2(d+1)}{\delta_l^2} \log(kl(l+1)/\delta)
$$

actions during the G-optimal explorations, so the number of phases $L^{(0)}$ is never larger than

$$
\ell_T = 1 \vee \log_4(T).
$$

Using this remark, together with Equations (16) and (18), we find that on $\overline{\mathcal{F}}$

$$
\sum_{l=1}^{L^{(0)}} \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x) \widehat{\Delta}_x^l + \mathbb{1}\left\{\text{Explore}_{L^{(0)}+1}^{(0)} = \text{False}\right\} \sum_{t \in \text{Exp}_{L^{(0)}+1}^{(0)}} \max_{x \in \mathcal{X}_{L^{(0)}+2}^{(-1)} \cup \mathcal{X}_{L^{(0)}+2}^{(+1)}} (x^* - x)^\top \gamma^*
$$

$$
\leq 2^{19} \log\left(\frac{6L^{(0)}}{\delta}\right) \kappa(\widehat{\Delta}^{L^{(0)}}) T^{2/3} \log(T)^{-2/3} + 42\ell_T. \tag{19}
$$

Bound on $\sum_{z \in \{-1,+1\}} \sum_{l=1}^{L_T} \left(\sum_{x \in \mathcal{X}_l^{(z)}} \mu_l^{(z)}(x)\right) \max_{x \in \mathcal{X}_l^{(z)}} (x^* - x)^\top \gamma^*$. We bound the remaining term in Equation (12) using the first claim in Lemma 14 and Lemma 13. On $\overline{\mathcal{F}}$,

$$
\sum_{z \in \{-1,+1\}} \sum_{l=1}^{L_T} \left(\sum_{x \in \mathcal{X}_l^{(z)}} \mu_l^{(z)}(x)\right) \max_{x \in \mathcal{X}_l^{(z)}} (x^* - x)^\top \gamma^* \leq 2\sum_{l=1}^{L_T} \left(\frac{2(d+1)}{\epsilon_l^2} \log\left(\frac{kl(l+1)}{\delta}\right) + \frac{(d+1)(d+2)}{2}\right) 42\epsilon_l
$$

$$
\leq \frac{336(d+1)}{\epsilon_{L_T}} \log\left(\frac{kL_T(1+L_T)}{\delta}\right) + 168(d+1)(d+2)
$$

$$
\leq 267(d+1)\kappa_*^{-1/3} T^{1/3} \log(T)^{-1/3} \log\left(\frac{kL_T(1+L_T)}{\delta}\right)
$$

$$
+ 168(d+1)(d+2). \tag{20}
$$

Combing Equations (11), (12), (14), (19), and (20), and using $\delta = T^{-1}$, $\kappa(\widehat{\Delta}^{L^{(0)}}) \leq \kappa_*$ and $L_T \leq 4T/\log(2)$, we get for all $T \geq 1$

$$
R_T \leq C\left(\kappa_*^{1/3} T^{2/3} \log(T)^{1/3} + (d \vee \kappa_*) \log(T) + d^2 + d\kappa_*^{-1/3} T^{1/3} \log(kT) \log(T)^{-1/3}\right)
$$

for some absolute constant $C > 0$. Finally, for

$$
T \geq \frac{((d \vee \kappa_*)^{3/2} \log(T)) \vee d^3}{\sqrt{\kappa_*}} \vee \frac{(d \log(kT))^3}{(\kappa_* \log(T))^2},
$$

we get

$$
R_T \leq C' \kappa_*^{1/3} T^{2/3} \log(T)^{1/3}.
$$

C.3 Proof of Theorem 2

The beginning of the proof of Theorem 2 follows the same lines as the proof of Theorem 1. We begin by decomposing the regret as

$$R_T \leq \sum_{t \leq T} \mathbb{E}_{|\overline{\mathcal{F}}} \left[(x^* - x_t)^\top \gamma^* \right] + 2T\mathbb{P}\left[\mathcal{F}\right]. \tag{21}$$

where $\mathcal{F}$ is defined in Equation (10). On the one hand, Lemma 12 implies $T\mathbb{P}\left[\mathcal{F}\right] \leq 2\delta T$. Then, Equation (21) implies

$$R_T \leq 4\delta T + \mathbb{E}_{|\overline{\mathcal{F}}} \left[\sum_{z \in \{-1,+1\}} \sum_{l \geq 1}^{L^{(z)}+1} \sum_{t \in \text{Exp}_l^{(z)}} (x^* - x_t)^\top \gamma^* \right] + \mathbb{E}_{|\overline{\mathcal{F}}} \left[\sum_{t \in \text{Recovery}} (x^* - x_t)^\top \gamma^* \right] \tag{22}$$

$$+ \mathbb{E}_{|\overline{\mathcal{F}}} \left[\sum_{l=1}^{L^{(0)}} \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x)\Delta_x \right] + \mathbb{E}_{|\overline{\mathcal{F}}} \left[\mathbb{1}\left\{ \text{Explore}_{L^{(0)}+1}^{(0)} = \text{False} \right\} \sum_{t \in \text{Exp}_{L^{(0)}+1}^{(0)}} \max_{x \in \mathcal{X}_{L^{(0)}+2}^{(-1)} \cup \mathcal{X}_{L^{(0)}+2}^{(+1)}} (x^* - x)^\top \gamma^* \right]$$

where $\mathcal{F}$ is defined in Equation (10), and where we used the convention that the sum over an empty set is null.

Bound on $\mathbb{1}\left\{ \text{Explore}_{L^{(0)}+1}^{(0)} = \text{False} \right\} \sum_{t \in \text{Exp}_{L^{(0)}+1}^{(0)}} \max_{x \in \mathcal{X}_{L^{(z)}+1}} (x^* - x)^\top \gamma^*$.

Similarly to the proof of Theorem 1, we use Lemma 13 and Lemma 15 to show that on $\overline{\mathcal{F}}$

$$\mathbb{1}\left\{ \text{Explore}_{L^{(0)}+1}^{(0)} = \text{False} \right\} \sum_{t \in \text{Exp}_{L^{(0)}+1}^{(0)}} \max_{x \in \mathcal{X}_{L^{(z)}+1}} (x^* - x)^\top \gamma^* \leq 21 \sum_{x \in \mathcal{X}} \mu_{L^{(0)}+1}^{(0)}(x)\widehat{\Delta}_x^{L^{(0)}+1}. \tag{23}$$

Bound on $\sum_{z \in \{-1,+1\}} \sum_{l \geq 1}^{L^{(z)}+1} \sum_{t \in \text{Exp}_l^{(z)}} (x^* - x_t)^\top \gamma^*$.

Lemma 13 shows that for $l \leq L^{(z)}$, the actions in $\mathcal{X}_{l+1}^{(z)}$ are sub-optimal by at most an additional factor at most $21\epsilon_l$. Let us set $l_{\Delta_{\min}} = \lceil -\log_2(\Delta_{\min}/21)\rceil$, so that

$$\frac{\Delta_{\min}}{42} \leq \epsilon_{l_{\Delta_{\min}}} \leq \frac{\Delta_{\min}}{21}.$$

For $l \geq l_{\Delta_{\min}}$, we have $\mathcal{X}_{l+1}^{(-1)} \cup \mathcal{X}_{l+1}^{(+1)} = \{x_{z^*}\}$. Thus, $l^{(-z_{x^*})} \leq l_{\Delta_{\min}}$, and for $l \geq l_{\Delta_{\min}}$, the algorithm selects only x^* during the phase $\text{Exp}_l^{(z^*)}$. Then, combining Lemmas 14 and 13, and the fact that $L^{(z)} + 1 \leq \ell_T$, we find that, on $\overline{\mathcal{F}}$,

$$\sum_{z \in \{-1,+1\}} \sum_{l=1}^{L^{(z)}+1} \sum_{t \in \text{Exp}_l^{(z)}} (x^* - x_t)^\top \gamma^* \leq \sum_{z \in \{-1,+1\}} \sum_{l=1}^{l_{\Delta_{\min}}+1 \wedge \ell_T} \left(\sum_{x \in \mathcal{X}_l^{(z)}} \mu_l^{(z)}(x) \right) \max_{x \in \mathcal{X}_l^{(z)}} (x^* - x)^\top \gamma^*$$

$$\leq 2 \sum_{l=1}^{l_{\Delta_{\min}}+1 \wedge \ell_T} \left(\frac{2(d+1)}{\epsilon_l^2} \log\left(\frac{kl(l+1)}{\delta} \right) + \frac{(d+1)(d+2)}{2} \right) 42\epsilon_l$$

$$\leq 84(d+1)(d+2) + \epsilon_{l_{\Delta_{\min}}}^{-1} \times 672(d+1)\log\left(\frac{k(1+\ell_T)(2+\ell_T)}{\delta} \right)$$

$$\leq 84(d+1)(d+2) + \frac{28224(d+1)}{\Delta_{\min}} \log\left(\frac{k(1+\ell_T))(2+\ell_T))}{\delta} \right) \tag{24}$$

Bound on $\sum_{t \in \text{Recovery}} (x^* - x_t)^\top \gamma^* + \sum_{l=1}^{L^{(0)}} \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x)\Delta_x + \sum_{x \in \mathcal{X}} \mu_{L^{(0)}+1}^{(0)}(x)\widehat{\Delta}_x^{L^{(0)}+1}$.

We use the following lemma to bound the number of phases necessary to eliminate the sub-optimal group.

Lemma 16. *On the event $\overline{\mathcal{F}}$ defined in Equation* (10), *for $l \geq 1$ such that $\epsilon_l \leq \frac{\Delta_{\neq}}{8}$ and Explore$_L^{(0)} =$ True, $\widehat{z^*}_{l+1} = z_{x^*}$.*

Let $l_{\Delta_{\neq}} = \lceil -\log(\Delta_{\neq}/8)/\log(2) \rceil$ be such that

$$\frac{\Delta_{\neq}}{16} \leq \epsilon_{l_{\Delta_{\neq}}} \leq \frac{\Delta_{\neq}}{8}. \tag{25}$$

Lemma 16 implies that on $\overline{\mathcal{F}}$, $L^{(0)} \leq l_{\Delta_{\neq}}$.

To bound the remaining terms, we consider two cases, corresponding to Recovery$= \emptyset$ and Recovery$\neq \emptyset$.

Case 1: Recovery$= \emptyset$. Our case assumption implies that

$$\sum_{t \in \text{Recovery}} (x^* - x_t)^\top \gamma^* = 0. \tag{26}$$

Lemma 15 implies that

$$\sum_{l=1}^{L^{(0)}} \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x) \Delta_x + \sum_{x \in \mathcal{X}} \mu_{L^{(0)}+1}^{(0)}(x) \widehat{\Delta}_x^{L^{(0)}+1} \leq \sum_{l=1}^{L^{(0)}+1} \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x) \widehat{\Delta}_x^l.$$

Moreover, $L^{(0)} \leq l_{\Delta_{\neq}} \wedge \ell_T$, so on $\overline{\mathcal{F}}$

$$\sum_{l=1}^{L^{(0)}+1} \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x) \widehat{\Delta}_x^l \leq \sum_{l=1}^{(l_{\Delta_{\neq}} \wedge \ell_T)+1} \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x) \widehat{\Delta}_x^l.$$

Using Lemma 14, we find that on $\overline{\mathcal{F}}$

$$\sum_{l=1}^{(l_{\Delta_{\neq}} \wedge \ell_T)+1} \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x) \widehat{\Delta}_x^l \leq \sum_{l=1}^{(l_{\Delta_{\neq}} \wedge \ell_T)+1} \frac{2\kappa(\widehat{\Delta}^l)}{\epsilon_l^2} \log \left(\frac{l(l+1)}{\delta} \right) + 2(d+1)(\ell_T + 1)$$

$$\leq 2 \log \left(\frac{(\ell_T + 1)(\ell_T + 2)}{\delta} \right) \sum_{l=1}^{l_{\Delta_{\neq}}+1} \frac{\kappa(\widehat{\Delta}^l)}{\epsilon_l^2} + 2(d+1)(\ell_T + 1).$$

Using Lemma 15 with $\tau = \Delta_{\neq}$ and (25), we have on $\overline{\mathcal{F}}$

$$\sum_{l=1}^{l_{\Delta_{\neq}}+1} \frac{\kappa(\widehat{\Delta}^l)}{\epsilon_l^2} \leq 513\kappa(\Delta \vee \Delta_{\neq}) \sum_{l=1}^{l_{\Delta_{\neq}}+1} \left(\epsilon_l^{-2} + \epsilon_l^{-1}/\Delta_{\neq} \right)$$

$$\leq \frac{2^{18}\kappa(\Delta \vee \Delta_{\neq})}{\Delta_{\neq}^2}.$$

We obtain on $\overline{\mathcal{F}}$

$$\sum_{l=1}^{L^{(0)}+1} \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x) \widehat{\Delta}_x^l \leq 2^{19} \log \left(\frac{(\ell_T + 1)(\ell_T + 2)}{\delta} \right) \frac{\kappa(\Delta \vee \Delta_{\neq})}{\Delta_{\neq}^2} + 2(d+1)(\ell_T + 1) \tag{27}$$

Combining Equations (24), (23), (26), and (27), we find that on $\overline{\mathcal{F}}$, when Recovery= $\emptyset$, there exsists an absolute constant $c > 0$ such that for $\delta = T^{-1}$,

$$
\sum_{z \in \{-1,+1\}} \sum_{l \geq 1}^{L^{(z)}+1} \sum_{t \in \mathrm{Exp}_l^{(z)}} (x^* - x_t)^\top \gamma^* + \sum_{t \in \mathrm{Recovery}} (x^* - x_t)^\top \gamma^* + \sum_{l=1}^{L^{(0)}} \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x) \Delta_x \qquad (28)
$$

$$
+ \mathbb{1}\{\mathrm{Explore}_{L^{(0)}+1}^{(0)} = \mathrm{False}\} \sum_{t \in \mathrm{Exp}_{L^{(0)}+1}^{(0)}} \max_{x \in \mathcal{X}_{L^{(0)}+2}^{(-1)} \cup \mathcal{X}_{L^{(0)}+2}^{(+1)}} (x^* - x)^\top \gamma^*
$$

$$
\leq c \left(d^2 + \left(\frac{d}{\Delta_{\min}} \vee \frac{\kappa(\Delta \vee \Delta_{\neq})}{\Delta_{\neq}^2} \right) \log(T) + \frac{d}{\Delta_{\min}} \log(k) \right).
$$

Case 2: Recovery$\neq \emptyset$. In this case, the algorithm enters Recovery at phase $L^{(0)}$, so $\mathrm{Explore}_{L^{(0)}+1}^{(0)} = \mathrm{False}$ and $\mathrm{Exp}_{L^{(0)}+1}^{(0)} = \emptyset$, and

$$
\mathbb{1}\{\mathrm{Explore}_{L^{(0)}+1}^{(0)} = \mathrm{False}\} \sum_{t \in \mathrm{Exp}_{L^{(0)}+1}^{(0)}} \max_{x \in \mathcal{X}_{L^{(0)}+2}^{(-1)} \cup \mathcal{X}_{L^{(0)}+2}^{(+1)}} (x^* - x)^\top \gamma^* = 0. \qquad (29)
$$

Using Lemma 13, we see that

$$
\sum_{t \in \mathrm{Recovery}} (x^* - x_t)^\top \gamma^* \leq 21 T \epsilon_{L^{(0)}+1}.
$$

On the other hand, in the Recovery phase, $\epsilon_{L^{(0)}+1} \leq \left(\kappa(\widehat{\Delta}^{L^{(0)}+1}) \log(T)/T \right)^{1/3}$. Thus,

$$
\sum_{t \in \mathrm{Recovery}} (x^* - x_t)^\top \gamma^* \leq \frac{21 \kappa(\widehat{\Delta}^{L^{(0)}+1}) \log(T)}{\epsilon_{L^{(0)}+1}^2}.
$$

Now, Lemma 14 show that

$$
\sum_{l=1}^{L^{(0)}} \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x) \Delta_x \leq 4 \log(2 L^{(0)} \delta^{-1}) \sum_{l=1}^{L^{(0)}} \frac{\kappa(\widehat{\Delta}^l)}{\epsilon_l^2} + 4 d L^{(0)}.
$$

Combining these results, and using $L^{(0)} \leq \ell_T$, we see that

$$
\sum_{t \in \mathrm{Recovery}} (x^* - x_t)^\top \gamma^* + \sum_{l=1}^{L^{(0)}} \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x) \Delta_x \leq 4 d L^{(0)} + \left(4 \log(2 \ell_T \delta^{-1}) \vee 21 \log(T) \right) \sum_{l=1}^{L^{(0)}+1} \frac{\kappa(\widehat{\Delta}^l)}{\epsilon_l^2}.
$$

$$(30)$$

Using Lemma 15 with $\tau = \epsilon_{L^{(0)}}$, we see that

$$
\sum_{l=1}^{L^{(0)}+1} \frac{\kappa(\widehat{\Delta}^l)}{\epsilon_l^2} \leq 513 \sum_{l=1}^{L^{(0)}+1} \frac{\kappa(\Delta \vee \epsilon_{L^{(0)}})}{\epsilon_l^2} + 513 \sum_{l=1}^{L^{(0)}+1} \frac{\kappa(\Delta \vee \epsilon_{L^{(0)}})}{\epsilon_{L^{(0)}} \epsilon_l}
$$

$$
\leq 10260 \frac{\kappa(\Delta \vee \epsilon_{L^{(0)}})}{\epsilon_{L^{(0)}}^2}.
$$

Now, the algorithm enters the Recovery phase before finding the best group, so we must have $L^{(0)} \leq l_{\Delta_{\neq}}$. This implies that

$$
\sum_{l=1}^{L^{(0)}+1} \frac{\kappa(\widehat{\Delta}^l)}{\epsilon_l^2} \leq 2^{18} \frac{\kappa(\Delta \vee \epsilon_{L^{(0)}})}{\Delta_{\neq}^2}.
$$

Finally, note that $L^{(0)} \geq L_T$, so $\epsilon_{L^{(0)}} \leq \epsilon_{L_T} = \varepsilon_T$, and

$$\sum_{l=1}^{L^{(0)}+1} \frac{\kappa(\widehat{\Delta}^l)}{\epsilon_l^2} \leq 2^{18} \frac{\kappa(\Delta \vee \varepsilon_T)}{\Delta_{\neq}^2}. \tag{31}$$

Combining Equations (24), (29), (30), and (31), we find that on $\overline{\mathcal{F}}$, when Recovery$\neq \emptyset$, there exists an absolute constant $c > 0$ such that for $\delta = T^{-1}$,

$$\sum_{z \in \{-1,+1\}} \sum_{l \geq 1}^{L^{(z)}+1} \sum_{t \in \text{Exp}_l^{(z)}} (x^* - x_t)^\top \gamma^* + \sum_{t \in \text{Recovery}} (x^* - x_t)^\top \gamma^* + \sum_{l=1}^{L^{(0)}} \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x) \Delta_x \tag{32}$$

$$+ \mathbb{1}\{\text{Explore}_{L^{(0)}+1}^{(0)} = \text{False}\} \sum_{t \in \text{Exp}_{L^{(0)}+1}^{(0)}} \max_{x \in \mathcal{X}_{L^{(0)}+2}^{(-1)} \cup \mathcal{X}_{L^{(0)}+2}^{(+1)}} (x^* - x)^\top \gamma^*$$

$$\leq c \left(d^2 + \left(\frac{d}{\Delta_{\min}} \vee \frac{\kappa(\Delta \vee \varepsilon_T)}{\Delta_{\neq}^2} \right) \log(T) + \frac{d \log(k)}{\Delta_{\min}} \right).$$

Conclusion We conclude the proof of Theorem 2 by combining Equations (22), (28) and (32).

C.4 Proof of Theorem 3

Consider the actions $\mathcal{A}$ defined in the following lemma.

Lemma 17. *Let the action set be given by* $\mathcal{A} = \left\{ \begin{pmatrix} x_1 \\ z_{x_1} \end{pmatrix}, ..., \begin{pmatrix} x_{d+1} \\ z_{x_{d+1}} \end{pmatrix} \right\}$, *where* $\begin{pmatrix} x_1 \\ z_{x_1} \end{pmatrix} = e_1 + e_{d+1}$, $\begin{pmatrix} x_i \\ z_{x_i} \end{pmatrix} = e_i - e_{d+1}$ *for* $i \in \{2, ..., d\}$, *and* $\begin{pmatrix} x_{d+1} \\ z_{x_{d+1}} \end{pmatrix} = -\left(1 - \frac{2}{\sqrt{\kappa_*}+1} \right) e_1 - e_{d+1}$. *It holds that*

$$\min_{\pi \in \mathcal{P}^{\mathcal{A}}} \left\{ e_{d+1}^\top \left(\sum_{\binom{x}{z} \in \mathcal{A}} \pi_x \begin{pmatrix} x \\ z_x \end{pmatrix} \begin{pmatrix} x \\ z_x \end{pmatrix}^\top \right)^+ e_{d+1} \right\} = \kappa.$$

By Lemma 17, $\mathcal{A} \in \mathbf{A}_{\kappa_*,d}$. We will introduce two bandit problems characterized by two parameters $\theta_T^{(1)}$ and $\theta_T^{(2)}$ - assuming that the noise ξ_t is Gaussian and i.i.d. - and we prove that for any algorithm, the regret for one of those two problems must be of larger order than $\kappa_*^{1/3} T^{2/3}$.

We also consider the following two alternative problems. For a small $1/4 > \rho_T > 0$ where $\rho_T = T^{-1/3} \kappa_*^{1/3}$ (satisfied since $T > 4^3 \kappa_*$), the two alternative action parameters are defined as:

$$\gamma_T^{(1)} = \frac{1+\rho_T}{2} e_1 + \frac{1-\rho_T}{2} e_2 - \frac{\rho_T}{2} \left(\sum_{3 \leq j \leq d} e_j \right)$$

$$\gamma_T^{(2)} = \frac{1-\rho_T}{2} e_1 + \frac{1+\rho_T}{2} e_2 + \frac{\rho_T}{2} \left(\sum_{3 \leq j \leq d} e_j \right).$$

On top of this, two bias parameters are defined as $\omega_T^{(1)} = -\frac{\rho_T}{2}$ and $\omega_T^{(2)} = \frac{\rho_T}{2}$. Through this, we define the two bandit problems of the sketch of proof of Lemma 17 characterized by $\theta_T^{(1)} = \begin{pmatrix} \gamma_T^{(1)} \\ \omega_T^{(1)} \end{pmatrix}$ and $\theta_T^{(2)} = \begin{pmatrix} \gamma_T^{(2)} \\ \omega_T^{(2)} \end{pmatrix}$ - and where the distribution of the noise ξ_t is supposed to be Gaussian and i.i.d. We refer to these two problems respectively as **Problem 1** and **Problem 2**. We write $R_T^{(1)}$, $\mathbb{P}^{(1)}$ and

$\mathbb{E}^{(1)}$ (respectively $R_T^{(2)}$, $\mathbb{P}^{(2)}$ and $\mathbb{E}^{(2)}$) for the regret, probability and expectation for the first bandit problem, when the parameter is $\theta_T^{(1)}$ (respectively the second bandit problem with $\theta_T^{(2)}$). We also write $\mathbb{P}_j^{(i)}$ for the distribution of a sample received in **Problem i** when sampling action x_j at any given time t - note that by definition of the bandit problems, this distribution does not depend on t and on the past samples given that action x_j is sampled.

The three following facts hold on these two bandit problems:

Fact 1 The parameters $\gamma_T^{(1)}$ and $\gamma_T^{(2)}$ are chosen so that x_1 is the unique best action for **Problem 1**, and x_2 is the unique best action for **Problem 2**. Choosing any sub-optimal action induces an instantaneous regret of at least ρ_T, and choosing the very sub-optimal action x_{d+1} induces an instantaneous regret of at least $1/2$.

Fact 2 Because of the chosen bias parameters, the distributions of the evaluations of all actions but x_{d+1} are exactly the same under the two bandit problems characterized by $\theta^{(1)}$ and $\theta_T^{(2)}$ - i.e. exactly the same data is observed under the two alternative bandit problems defined by the two alternative parameters for all actions but x_{d+1}. More precisely, for $i \in \{1, 2\}$, in **Problem i** and at any time t, when sampling action x_i where $i \leq 2$, we observe a sample distributed according to $\mathcal{N}(1/2, 1)$ - i.e. $\mathbb{P}_j^{(i)}$ is $\mathcal{N}(1/2, 1)$ - and when sampling action x_i where $2 < i \geq d + 1$, we observe a sample distributed according to $\mathcal{N}(0, 1)$ - i.e. $\mathbb{P}_j^{(i)}$ is $\mathcal{N}(0, 1)$.

Fact 3 The distributions of the outcomes of the evaluation of action x_{d+1} differs in the two bandit problems. Set $\alpha = 2/(\sqrt{\kappa_*} + 1)$. In **Problem 1**, $\mathbb{P}_{d+1}^{(1)}$ is $\mathcal{N}(-\frac{1-\alpha-\rho_T\alpha}{2}, 1)$. In **Problem 2**, $\mathbb{P}_{d+1}^{(2)}$ is $\mathcal{N}(-\frac{1-\alpha+\rho_T\alpha}{2}, 1)$. So that the difference between the means of the evaluations of action x_{d+1} in the two bandit problems is $\bar{\Delta} = \rho_T\alpha = \frac{2\rho_T}{\sqrt{\kappa_*}+1} \leq \frac{2\rho_T}{\sqrt{\kappa_*}}$.

For $i \leq d + 1$, we write $N_i(T)$ for the number of times that action x_i has been selected before time T. In **Problem 1**, choosing the action x_{d+1} leads to an instantaneous regret larger than $\frac{1}{2}$ (**Fact 1**), so that

$$R_T^{(1)} \geq \frac{\mathbb{E}^{(1)}\left[N_{x_{d+1}}(T)\right]}{2}.$$

If $\mathbb{E}^{(1)}\left[N_{d+1}(T)\right] \geq \frac{T^{2/3}\kappa_*^{1/3}}{2}$, then Theorem 1 follows immediately; we therefore consider from now on the case when

$$\mathbb{E}^{(1)}\left[N_{d+1}(T)\right] \leq \frac{T^{2/3}\kappa_*^{1/3}}{2}. \tag{33}$$

Now, let us define the event

$$F = \left\{N_1(T) \geq \frac{T}{2}\kappa_*^{1/3}\right\}.$$

Note that action x_1 is optimal for **Problem 1** and that action x_2 is optimal for **Problem 2** (**Fact 1**). Since choosing an action that is sub-optimal leads to an instantaneous regret larger than ρ_T (**Fact 1**), we also have

$$R_T^{(1)} \geq \frac{T\rho_T}{2}\mathbb{P}^{(1)}\left(\overline{F}\right)$$

and

$$R_T^{(2)} \geq \frac{T\rho_T}{2}\mathbb{P}^{(2)}\left(F\right).$$

Then, Bretagnolle-Huber inequality (see, e.g., Theorem 14.2 in [24]) implies that

$$R_T^{(1)} + R_T^{(2)} \geq \frac{T\rho_T}{4}\exp\left(-KL\left(\mathbb{P}^{(1)}, \mathbb{P}^{(2)}\right)\right).$$

For the choice $\rho_T = T^{-1/3}\kappa_*^{1/3}$, this implies that

$$R_T^{(1)} + R_T^{(2)} \geq \frac{T^{2/3}\kappa_*^{1/3}}{4}\exp\left(-KL\left(\mathbb{P}^{(1)}, \mathbb{P}^{(2)}\right)\right). \tag{34}$$

Now, the Kullback-Leibler divergence between $\mathbb{P}^{(1)}$ and $\mathbb{P}^{(2)}$ can be rewritten as follows (see, e.g., Lemma 15.1 in [24]) :

$$KL(\mathbb{P}^{(1)}, \mathbb{P}^{(2)}) \quad = \quad \frac{1}{2} \sum_{j \leq d+1} \mathbb{E}^{(1)} \left[N_j(T) \right] KL(\mathbb{P}_j^{(1)}, \mathbb{P}_j^{(2)}).$$

By **Fact 2**, we have that for any $j \leq d$, $\mathbb{P}_j^{(1)} = \mathbb{P}_j^{(2)}$. So that

$$KL(\mathbb{P}^{(1)}, \mathbb{P}^{(2)}) \quad = \quad \frac{1}{2} \mathbb{E}^{(1)} \left[N_{d+1}(T) \right] KL(\mathbb{P}_{d+1}^{(1)}, \mathbb{P}_{d+1}^{(2)}).$$

By the characterization of $\mathbb{P}_{d+1}^{(1)}, \mathbb{P}_{d+1}^{(2)}$ in **Fact 3**, and recalling that the Kullback-Leibler divergence between two normalized Gaussian distributions is given by the squared distance between their means, we find that

$$KL(\mathbb{P}^{(1)}, \mathbb{P}^{(2)}) \quad = \quad \frac{1}{2} \mathbb{E}^{(1)} \left[N_{d+1}(T) \right] \bar{\Delta}^2.$$

Thus, by the definition of $\bar{\Delta}$ in **Fact 3** and by Equation (33)

$$KL \left(\mathbb{P}^{(1)}, \mathbb{P}^{(2)} \right) = \frac{1}{2} \mathbb{E}^{(1)} \left[N_{d+1}(T) \right] \left(\frac{2\rho_T}{\sqrt{\kappa_*} + 1} \right)^2 \leq \frac{T^{2/3} \kappa_*^{1/3}}{4} \times \frac{4\rho_T^2}{\kappa_*} = 1, \qquad (35)$$

reminding that $\rho_T = T^{-1/3} \kappa_*^{1/3}$.

Combining Equations (34) and (35) implies that

$$\max \left\{ R_T^{(1)}, R_T^{(2)} \right\} \geq \frac{T^{2/3} \kappa^{1/3}}{8} \exp(-1),$$

which concludes the proof of Theorem 3.

C.5   Proof of Theorems 4

Theorems 4 follows directly from the next Theorem.

Theorem 6. *For all $\kappa_* \geq 1$ and all $d \geq 4$, there exists an action set $\mathcal{A} \in \mathbf{A}_{\kappa_*, d}$, such that for all bandit algorithms, for all $(\Delta_{\min}, \Delta_{\neq}) \in (0, 1/8)^2$ with $\Delta_{\min} \leq \Delta_{\neq}$, and for all budget $T \geq 2$, there exists a problem characterized by $\theta \in \Theta_{\Delta_{\min}, \Delta_{\neq}}^{\mathcal{A}}$ such that the regret of the algorithm on the problem satisfies*

$$R_T^\theta \quad \geq \quad \left[\frac{d}{10\Delta_{\min}} \log(T) \left[1 - \frac{\log \left(\frac{8d \log(T)}{\Delta_{\min}^2} \right)}{\log(T)} \right] \right] \vee \left[\frac{\kappa_* + 1}{4\Delta_{\neq}^2} \log(T) \left[1 - \frac{\log \left(\frac{8\kappa_* \log(T)}{\Delta_{\neq}^3} \right)}{\log(T)} \right] \right]$$

$$\vee \left[\frac{\kappa_*}{4\Delta_{\neq}^2} \left[1 \wedge \log \left(\frac{T\Delta_{\neq}^3}{8\kappa_*} \right) \right] \right]. \qquad (36)$$

Moreover, on this problem, $\kappa(\Delta) \in [\kappa_*/8, 2\kappa_*]$.

Remark 1. *Note that Theorem 6 allows us to recover a lower bound similar to that of Theorem 3 by choosing $\Delta_{\neq}$ and $\Delta_{\min}$ of the order $\kappa_*^{1/3} T^{-1/3}$, however this bound only holds for d larger than 4.*

We prove Theorem 6 for the following set of actions $\mathcal{A}$: $\mathcal{A} = \left\{ \left(\begin{smallmatrix} x_1 \\ z_{x_1} \end{smallmatrix} \right), ..., \left(\begin{smallmatrix} x_{d+1} \\ z_{x_{d+1}} \end{smallmatrix} \right) \right\}$, where $\left(\begin{smallmatrix} x_i \\ z_{x_i} \end{smallmatrix} \right) = e_i + e_{d+1}$, for $i \in \{2, ..., \lfloor d/2 \rfloor\}$, $\left(\begin{smallmatrix} x_i \\ z_{x_i} \end{smallmatrix} \right) = e_i - e_{d+1}$ for $i \in \{\lfloor d/2 \rfloor + 1, ..., d\}$, and $\left(\begin{smallmatrix} x_{d+1} \\ z_{x_{d+1}} \end{smallmatrix} \right) = - \left(1 - \frac{2}{\sqrt{\kappa_*} + 1} \right) e_1 - e_{d+1}$. Then, by Lemma 10, for this choice of action set, we have $\mathcal{A} \in \mathbf{A}_{\kappa_*, d}$.

We consider the following set of bandit problems: for $i \in \{1, ..., \lfloor d/2 \rfloor + 1\}$ **Problem i** is characterized by the parameter $\theta^{(i)}$, where $\theta^{(i)} = \begin{pmatrix} \gamma^{(i)} \\ \omega^{(i)} \end{pmatrix}$ is defined as:

$$\gamma^{(1)} = \frac{1 + \Delta_{\neq} - \Delta_{\min}}{2} \left(\sum_{1 \leq j \leq \lfloor d/2 \rfloor} e_j \right) + \frac{1 - \Delta_{\neq} - \Delta_{\min}}{2} \left(\sum_{\lfloor d/2 \rfloor + 1 \leq j \leq d} e_j \right) + \Delta_{\min} e_1 + \Delta_{\min} e_{\lfloor d/2 \rfloor + 1}$$

$$\gamma^{(i)} = \gamma^{(1)} + 2\Delta_{\min} e_i + 2\Delta_{\min} e_{\lfloor d/2 \rfloor + i} \quad \forall i \in \{2, ..., \lfloor d/2 \rfloor\}$$

$$\gamma^{(\lfloor d/2 \rfloor + 1)} = \frac{1 - \Delta_{\neq} - \Delta_{\min}}{2} \left(\sum_{1 \leq j \leq \lfloor d/2 \rfloor} e_j \right) + \frac{1 + \Delta_{\neq} - \Delta_{\min}}{2} \left(\sum_{\lfloor d/2 \rfloor + 1 \leq j \leq d} e_j \right) + \Delta_{\min} e_1 + \Delta_{\min} e_{\lfloor d/2 \rfloor + 1},$$

and the bias parameters are defined as $\omega^{(i)} = -\frac{\Delta_{\neq}}{2} \forall i \in \{1, ..., \lfloor d/2 \rfloor\}$, and otherwise $\omega^{(\lfloor d/2 \rfloor + 1)} = \frac{\Delta_{\neq}}{2}$. We write $\mathbb{E}^{(i)}, \mathbb{P}^{(i)}, R_T^{(i)}$ for resp. the probability, expectation, and regret, in **Problem i**. Note that this choice of parameters ensures that $\forall i \in \{1, ..., \lfloor d/2 \rfloor + 1\}, \theta^{(i)} \in \Theta_{\Delta_{\min}, \Delta_{\neq}}^{\mathcal{A}}$.

Set $\mathcal{A} = \left\{ \begin{pmatrix} x_1 \\ z_{x_1} \end{pmatrix}, ..., \begin{pmatrix} x_{d+1} \\ z_{x_{d+1}} \end{pmatrix} \right\}$, where $\begin{pmatrix} x_i \\ z_{x_i} \end{pmatrix} = e_i + e_{d+1}$, for $i \in \{2, ..., \lfloor d/2 \rfloor\}$, $\begin{pmatrix} x_i \\ z_{x_i} \end{pmatrix} = e_i - e_{d+1}$ for $i \in \{\lfloor d/2 \rfloor + 1, ..., d\}$, and $\begin{pmatrix} x_{d+1} \\ z_{x_{d+1}} \end{pmatrix} = -\left(1 - \frac{2}{\sqrt{\kappa_*} + 1} \right) e_1 - e_{d+1}$. Then, Lemma 10 shows that $\mathcal{A} \in \mathbf{A}_{\kappa_*, d}$.

The following facts hold:

Fact 1 For any $i \in \{1, ..., \lfloor d/2 \rfloor + 1\}$, action x_i is the unique optimal action in **Problem i**. Since $1/2 \geq \Delta_{\neq} \geq \Delta_{\min}$, sampling any other (sub-optimal) action leads to an instantaneous regret of at least $\Delta_{\min}$. Moreover, choosing an action in the group $-z_i$ leads to an instantaneous regret of at least $\Delta_{\neq}$.

Fact 2 In **Problem i** for any $i \in \{1, ..., \lfloor d/2 \rfloor + 1\}$, action $d + 1$ is very sub-optimal and sampling it leads to an instantaneous regret higher than $(1 - 2/(\sqrt{\kappa_*} + 1))(1 - \Delta_{\neq} + \Delta_{\min}) + (1 + \Delta_{\neq} + \Delta_{\min})/2 \geq 1/2$, since $\kappa_* \geq 1$ and $1/2 \geq \Delta_{\neq} \geq \Delta_{\min}$.

Fact 3 In **Problem i**, for $i \in \{1, ..., \lfloor d/2 \rfloor + 1\}$, when sampling action x_j at time, t the distribution of the observation does not depend on t or on the past (except through the choice of x_j) and is $\mathbb{P}_j^{(i)}$. It is characterized as:

$$\forall i \in \{1, ..., \lfloor d/2 \rfloor + 1\}, \mathbb{P}_1^{(i)}, \mathbb{P}_{\lfloor d/2 \rfloor + 1}^{(i)} \text{ are } \mathcal{N}((1 + \Delta_{\min})/2, 1)$$

$$\forall i \in \{1, ..., \lfloor d/2 \rfloor + 1\}, \forall j \in \{2, ..., d\} \setminus \{\lfloor d/2 \rfloor + 1, i, \lfloor d/2 \rfloor + i\}, \mathbb{P}_j^{(i)} \text{ is } \mathcal{N}((1 - \Delta_{\min})/2, 1),$$

$$\forall i \in \{2, \lfloor d/2 \rfloor\}, \mathbb{P}_i^{(i)} \text{ is } \mathcal{N}((1 + 3\Delta_{\min})/2, 1) \quad \mathbb{P}_{\lfloor d/2 \rfloor + i}^{(i)} \text{ is } \mathcal{N}((1 + 3\Delta_{\min})/2, 1)$$

$$\forall i \in \{1, \lfloor d/2 \rfloor\}, \mathbb{P}_{d+1}^{(i)} \text{ is } \mathcal{N}(-(1 - \alpha)(1 + \Delta_{\neq} + \Delta_{\min})/2 + \Delta_{\neq}/2, 1),$$

$$\mathbb{P}_{d+1}^{(\lfloor d/2 \rfloor + 1)} \text{ is } \mathcal{N}(-(1 - \alpha)(1 - \Delta_{\neq} + \Delta_{\min})/2 - \Delta_{\neq}/2, 1) \text{ where } \alpha = 2/(\sqrt{\kappa_*} + 1).$$

So that:

Fact 3.1 For any $i \in \{2, ..., \lfloor d/2 \rfloor\}$, between **Problem 1** and **Problem i**, the only actions that provide different evaluations when sampled are action i and action $\lfloor d/2 \rfloor + i$, and the mean gaps in both cases is $2\Delta_{\min}$.

Fact 3.2 Between **Problem 1** and **Problem** $\lfloor d/2 \rfloor + 1$, the only action that provide different evaluation when sampled is action $d + 1$, and the mean gap in this case is $\alpha \Delta_{\neq}$.

For $j \leq d + 1$, we write $N_j(T)$ for the total number of times action x_j has been selected before time T. Then, for $j \in \{1, ..., \lfloor d/2 \rfloor\}$, let $E^{(j)} = \{N_i(T) \leq T/2\}$. Note that for $i \in \{1, ..., \lfloor d/2 \rfloor\}$, in **Problem i** the action x_i is the optimal action. Therefore, for any efficient algorithm, for all $i \in \{1, ..., \lfloor d/2 \rfloor\}$ the event $E^{(i)}$ should have a low probability under $\mathbb{P}^{(i)}$. Indeed, for $i \in \{1, ..., \lfloor d/2 \rfloor\}$, the regret of the algorithm under **Problem i** can be lower-bounded as follows - see **Facts 1 and 2**:

$$R_T^{(i)} \geq \sum_{j \leq \lfloor d/2 \rfloor, \, j \neq i} \mathbb{E}^{(i)} [N_j(T)] \Delta_{\min} + \sum_{\lfloor d/2 \rfloor + 1 \leq j \leq d} \mathbb{E}^{(i)} [N_j(T)] \Delta_{\neq} + \frac{\mathbb{E}^{(i)} [N_{d+1}(T)]}{2} \quad (37)$$

Since $\sum_j \mathbb{E}^{(i)}[N_j(T)] = T$ and $\Delta_{\min} \le \Delta_{\ne} \le \frac{1}{2}$, this implies together with **Facts 1**:

$$R_T^{(i)} \ge \left(T - \mathbb{E}^{(i)}[N_i(T)]\right)\Delta_{\min}$$

Using the definition of $E^{(i)}$, we find that

$$R_T^{(i)} \ge \frac{T\Delta_{\min}}{2}\mathbb{P}^{(i)}\left(E^{(i)}\right). \tag{38}$$

In particular for **Problem 1**, for any $i \in \{1, ..., \lfloor d/2 \rfloor\}$,

$$R_T^{(1)} \ge \frac{T\Delta_{\min}}{2}\mathbb{P}^{(1)}\left(\overline{E^{(i)}}\right). \tag{39}$$

since $E^{(1)} \supset \overline{E^{(i)}}$.

Similarly, let us also define the event $F = \left\{\sum_{i \le \lfloor d/2 \rfloor} N_i(T) \ge T/2\right\}$. Then, in **Problem 1**, the group 1 contains the optimal action, and so for any efficient algorithm, the event F should have a low probability under $\mathbb{P}^{(1)}$. Indeed, Equation (37) also implies

$$R_T^{(1)} \ge \left(T - \mathbb{E}^{(1)}\left[\sum_{i \le \lfloor d/2 \rfloor} N_i(T)\right]\right)\Delta_{\ne} \ge \frac{T\Delta_{\ne}}{2}\mathbb{P}^{(1)}\left(\overline{F}\right). \tag{40}$$

On the other hand, for any efficient algorithm, the event F should have high probability under $\mathbb{P}^{(\lfloor d/2 \rfloor + 1)}$. Indeed, under problem **Problem $\lfloor d/2 \rfloor + 1$**, the regret can be lower-bounded as follows - see **Facts 1 and 2**:

$$R_T^{(\lfloor d/2 \rfloor + 1)} \ge \sum_{j \le \lfloor d/2 \rfloor} \mathbb{E}^{(\lfloor d/2 \rfloor + 1)}[N_j(T)]\Delta_{\ne} + \sum_{\lfloor d/2 \rfloor + 2 \le j \le d} \mathbb{E}^{(\lfloor d/2 \rfloor + 1)}[N_j(T)]\Delta_{\min} + \frac{\mathbb{E}^{(\lfloor d/2 \rfloor + 1)}[N_{d+1}(T)]}{2}.$$

which implies that

$$R_T^{(\lfloor d/2 \rfloor + 1)} \ge \sum_{j \le \lfloor d/2 \rfloor} \mathbb{E}^{(\lfloor d/2 \rfloor + 1)}[N_j(T)]\Delta_{\ne} \ge \frac{T\Delta_{\ne}}{2}\mathbb{P}^{(\lfloor d/2 \rfloor + 1)}(F). \tag{41}$$

Now, Bretagnolle-Huber inequality (see, e.g., Theorem 14.2 in [24]) implies that for all $i \in \{2, ..., \lfloor d/2 \rfloor\}$,

$$\frac{1}{2}\exp\left(-KL\left(\mathbb{P}^{(1)}, \mathbb{P}^{(i)}\right)\right) \le \mathbb{P}^{(i)}\left(E^{(i)}\right) + \mathbb{P}^{(1)}\left(\overline{E^{(i)}}\right) \tag{42}$$

and that

$$\frac{1}{2}\exp\left(-KL\left(\mathbb{P}^{(1)}, \mathbb{P}^{(\lfloor d/2 \rfloor + 1)}\right)\right) \le \mathbb{P}^{(\lfloor d/2 \rfloor + 1)}(F) + \mathbb{P}^{(1)}\left(\overline{F}\right). \tag{43}$$

On the one hand, Equation (42) implies that for any $i \in \{2, ..., \lfloor d/2 \rfloor\}$,

$$\begin{aligned} KL\left(\mathbb{P}^{(1)}, \mathbb{P}^{(i)}\right) &\ge -\log\left(2\mathbb{P}^{(i)}\left(E^{(i)}\right) + 2\mathbb{P}^{(1)}\left(\overline{E^{(i)}}\right)\right) \\ &\ge \log(T) - \log\left(2T\mathbb{P}^{(i)}\left(E^{(i)}\right) + 2T\mathbb{P}^{(1)}\left(\overline{E^{(i)}}\right)\right). \end{aligned} \tag{44}$$

Combining Equations (38), (39), and (44), we find that

$$KL\left(\mathbb{P}^{(1)}, \mathbb{P}^{(i)}\right) \ge \log(T) - \log\left(\frac{4(R_T^{(i)} + R_T^{(1)})}{\Delta_{\min}}\right). \tag{45}$$

On the other hand, Equation (43) implies that

$$\begin{aligned} KL\left(\mathbb{P}^{(1)}, \mathbb{P}^{(\lfloor d/2 \rfloor + 1)}\right) &\ge -\log\left(2\mathbb{P}^{(\lfloor d/2 \rfloor + 1)}(F) + 2\mathbb{P}^{(1)}\left(\overline{F}\right)\right) \\ &\ge \log(T) - \log\left(2T\mathbb{P}^{(\lfloor d/2 \rfloor + 1)}(F) + 2T\mathbb{P}^{(1)}\left(\overline{F}\right)\right). \end{aligned} \tag{46}$$

Combining Equations (38), (39), and (46), we find that

$$KL\left(\mathbb{P}^{(1)}, \mathbb{P}^{(\lfloor d/2 \rfloor+1)}\right) \geq \log(T) - \log\left(\frac{4(R_T^{(\lfloor d/2 \rfloor+1)} + R_T^{(1)})}{\Delta_{\neq}}\right). \tag{47}$$

Also, note that for all $i \in \{2, ..., \lfloor d/2 \rfloor + 1\}$, the Kullback-Leibler divergence between $\mathbb{P}^{(1)}$ and $\mathbb{P}^{(i)}$ can be decomposed as follows (see, e.g., Lemma 15.1 in [24]) :

$$KL(\mathbb{P}^{(1)}, \mathbb{P}^{(i)}) = \sum_{j \leq d+1} \mathbb{E}^{(1)}[N_j(T)] \, KL(\mathbb{P}_j^{(1)}, \mathbb{P}_j^{(i)}). \tag{48}$$

Lower bound in $d\Delta_{\min}^{-1} \log T$**.** By design, for $i \in \{2, ..., \lfloor d/2 \rfloor\}$, all actions but x_i and $x_{\lfloor d \rfloor + i}$ have the same distribution under $\mathbb{P}^{(1)}$ and $\mathbb{P}^{(i)}$ - see **Fact 3.1**. Then, Equation (48) becomes from **Fact 3.1** and from the expression of KL divergence between standard Gaussian distributions:

$$KL(\mathbb{P}^{(1)}, \mathbb{P}^{(i)}) = \frac{4\Delta_{\min}^2}{2} \mathbb{E}^{(1)}[N_i(T)] + \frac{4\Delta_{\min}^2}{2} \mathbb{E}^{(1)}[N_{\lfloor d \rfloor + i}(T)].$$

So that, summing over $i \in \{2, ..., \lfloor d/2 \rfloor\}$, and by **Fact 1**:

$$\sum_{i \in \{2, ..., \lfloor d/2 \rfloor\}} KL(\mathbb{P}^{(1)}, \mathbb{P}^{(i)}) \leq 2\Delta_{\min} R_T^{(1)}.$$

So that by Equation (45) (summing over $i \in \{2, ..., \lfloor d/2 \rfloor\}$):

$$2\Delta_{\min} R_T^{(1)} \geq \sum_{i \in \{2, ..., \lfloor d/2 \rfloor\}} \left[\log(T) - \log\left(\frac{4(R_T^{(i)} + R_T^{(1)})}{\Delta_{\min}}\right)\right]$$

$$= (\lfloor d/2 \rfloor - 1)\log(T) - \sum_{i \in \{2, ..., \lfloor d/2 \rfloor\}} \log\left(\frac{4(R_T^{(i)} + R_T^{(1)})}{\Delta_{\min}}\right).$$

Let us assume that our algorithm satisfies $\max_{i \leq \lfloor d/2 \rfloor} R_T^{(i)} \leq \frac{d\log(T)}{\Delta_{\min}}$ - otherwise the bound immediately follows for this algorithm. Then

$$R_T^{(1)} \geq \frac{1}{2\Delta_{\min}}(\lfloor d/2 \rfloor - 1)\log(T) - \frac{1}{2\Delta_{\min}} \sum_{i \in \{2, ..., \lfloor d/2 \rfloor\}} \log\left(\frac{8d\log T}{\Delta_{\min}^2}\right)$$

$$\geq \frac{1}{2\Delta_{\min}}(\lfloor d/2 \rfloor - 1)\left[\log(T) - \log\left(\frac{8d\log(T)}{\Delta_{\min}^2}\right)\right]. \tag{49}$$

Sine $d \geq 4$, we note that $\lfloor d/2 \rfloor - 1 \geq d/5$. This concludes the proof for this part of the bound.

Lower bound in $\kappa_* \Delta_{\neq}^{-2} \log T$**.** By design, all actions but x_{d+1} have the same evaluation under **Problem 1** and **Problem** $\lfloor d/2 \rfloor + 1$ - see **Fact 3.2**. Then, by **Fact 3.2** and the expression between the KL divergence of standard Gaussians, Equation (48) becomes

$$KL(\mathbb{P}^{(1)}, \mathbb{P}^{(\lfloor d/2 \rfloor+1)}) = \mathbb{E}^{(1)}[N_{d+1}(T)] \frac{(\alpha\Delta_{\neq})^2}{2} = \frac{1}{2}\mathbb{E}^{(1)}[N_{d+1}(T)]\left(\frac{2\Delta_{\neq}}{\sqrt{\kappa_*}+1}\right)^2.$$

Combined with equation (47), this implies that

$$\frac{1}{2}\mathbb{E}^{(1)}[N_{d+1}(T)]\left(\frac{2\Delta_{\neq}}{\sqrt{\kappa_*}+1}\right)^2 \geq \log(T) - \log\left(\frac{4(R_T^{(\lfloor d/2 \rfloor+1)} + R_T^{(1)})}{\Delta_{\neq}}\right). \tag{50}$$

Let us assume that our algorithm satisfies $\max_{i \leq \lfloor d/2 \rfloor+1} R_T^{(i)} \leq \frac{\kappa_* \log(T)}{\Delta_{\neq}^2}$ - otherwise the bound immediately follows for this algorithm. We then have

$$\frac{1}{2}\mathbb{E}^{(1)}[N_{d+1}(T)]\left(\frac{2\Delta_{\neq}}{\sqrt{\kappa_*}+1}\right)^2 \geq \log(T) - \log\left(\frac{8\kappa_* \log(T)}{\Delta_{\neq}^3}\right).$$

Using Equation (37), we find that

$$R_T^{(1)} \geq \frac{\kappa_*+1}{4\Delta_{\neq}^2}\left[\log(T) - \log\left(\frac{8\kappa_* \log(T)}{\Delta_{\neq}^3}\right)\right]. \tag{51}$$

Lower bound in $\kappa_* \Delta_{\neq}^{-2}$. Let us assume that our algorithm satisfies $\max_{i \leq \lfloor d/2 \rfloor + 1} R_T^{(i)} \leq \frac{\kappa_*}{\Delta_{\neq}^2}$ - otherwise the bound immediately follows for this algorithm. Then, Equation (50) implies

$$\frac{1}{2} \mathbb{E}^{(1)} [N_{d+1}(T)] \left(\frac{2\Delta_{\neq}}{\sqrt{\kappa_*}} \right)^2 \geq \log(T) - \log\left(\frac{8\kappa_*}{\Delta_{\neq}^3} \right).$$

Using again Equation (37), we find that

$$R_T^{(1)} \geq \frac{\kappa_* + 1}{4\Delta_{\neq}^2} \log\left(\frac{T\Delta_{\neq}^3}{8\kappa_*} \right). \tag{52}$$

We conclude the proof of Theorem 6 by combining Equations (49), (51) and (52).

Bounds on $\kappa(\Delta)$ Finally, Lemma 11 allows us to express $\kappa(\Delta)$ as a function of κ_*. On the one hand, since $\kappa_* \geq 1$, we see that $\kappa_* \leq (1 + \sqrt{\kappa_*})^2 \leq 4\kappa_*$. On the other hand, $1/2 \leq \Delta_{d+1} \leq 2$, so $\kappa(\Delta) \in \left[\frac{\kappa_*}{8}, 2\kappa_* \right]$.

C.6 Extension of the gap-dependent lower bounds to $d = 2, 3$

Theorem 4 can be extended to $d \in \{2, 3\}$ by considering separately the cases $\frac{d}{\Delta_{\min}} \geq \frac{\kappa}{\Delta_{\neq}^2}$ and $\frac{d}{\Delta_{\min}} < \frac{\kappa}{\Delta_{\neq}^2}$.

Case 1 : $\frac{d}{\Delta_{\min}} \geq \frac{\kappa}{\Delta_{\neq}^2}$ Let us consider the set of actions defined by $\mathcal{A} = \left\{ \begin{pmatrix} x_1 \\ z_{x_1} \end{pmatrix}, ..., \begin{pmatrix} x_{d+1} \\ z_{x_{d+1}} \end{pmatrix} \right\}$, where $\begin{pmatrix} x_i \\ z_{x_i} \end{pmatrix} = e_1 + e_{d+1}$ for $i \in \{1, ..., d\}$, and $\begin{pmatrix} x_{d+1} \\ z_{x_{d+1}} \end{pmatrix} = - \left(1 - \frac{2}{\sqrt{\kappa_*}+1} \right) e_1 - e_{d+1}$. Using the same proof as in Lemma 17, we see that

$$\min_{\pi \in \mathcal{P}^{\mathcal{A}}} \left\{ e_{d+1}^\top \left(\sum_{\left(\begin{smallmatrix} x \\ z \end{smallmatrix} \right) \in \mathcal{A}} \pi_x \begin{pmatrix} x \\ z_x \end{pmatrix} \begin{pmatrix} x \\ z_x \end{pmatrix}^\top \right)^+ e_{d+1} \right\} = \kappa.$$

Then, we consider the following problems : for $i \leq d$, **Problem i** is characterized by the parameter $\theta^{(i)}$, where $\theta^{(i)} = \begin{pmatrix} \gamma^{(i)} \\ \omega^{(i)} \end{pmatrix}$ is defined as:

$$\gamma^{(1)} = \frac{1 - \Delta_{\min}}{2} \sum_{i \leq d} e_i + \Delta_{\min} e_1$$

$$\gamma^{(i)} = \frac{1 - \Delta_{\min}}{2} \sum_{i \leq d} e_i + \Delta_{\min} e_1 + \Delta_{\min} e_i \quad \text{for i >1}$$

and the bias parameters are defined as $\omega^{(i)} = 0$ for $i \leq d$. The following facts hold:

Fact 1 For any $i \in \{1, ..., d\}$, action x_i is the unique optimal action in **Problem i**. Sampling any other (sub-optimal) action leads to an instantaneous regret of at least $\Delta_{\min}$.

Fact 2 In **Problem i**, for $i \in \{1, ..., d\}$, when sampling action x_j at time, t the distribution of the observation does not depend on t or on the past (except through the choice of x_j) and is $\mathbb{P}_j^{(i)}$. It is characterized as:

$$\forall i \in \{1, ..., d\}, \mathbb{P}_1^{(i)} \text{ is } \mathcal{N}((1 + \Delta_{\min})/2, 1)$$

$$\forall i \in \{1, ..., d\}, \mathbb{P}_{d+1}^{(1)} \text{ is } \mathcal{N}(-(1 - \frac{2}{\sqrt{\kappa_*}+1})(1 + \Delta_{\min})/2, 1)$$

$$\forall i \in \{2, ..., d\}, \mathbb{P}_i^{(i)} \text{ is } \mathcal{N}((1 + 3\Delta_{\min})/2, 1)$$

$$\forall i, j \in \{2, ..., d\}, i \neq j : \mathbb{P}_j^{(i)} \text{ is } \mathcal{N}((1 - \Delta_{\min})/2, 1)$$

So that for any $i \in \{2, ..., d\}$, between **Problem 1** and **Problem i**, the only action that provides different evaluations when sampled is action i, and the mean gap is $2\Delta_{\min}$.

Since $\Delta_{\neq} \leq \frac{1}{8}$, this choice of parameters ensures that $\forall i \in \{1, ..., d\}, \theta^{(i)} \in \Theta^{\mathcal{A}}_{\Delta_{\min}, \Delta_{\neq}, \kappa_*}$. Adapting the proof of Lemma 17, we note that the minimal variance of bias estimation is at least κ_*. This proves that $\mathcal{A} \in \Theta^{\mathcal{A}}_{\Delta_{\min}, \Delta_{\neq}, \kappa_*}$. Now, the lower bound

$$R_T \geq \frac{d-1}{2\Delta_{\min}} \left[\log(T) - \log\left(\frac{8d \log(T)}{\Delta^2_{\min}} \right) \right]$$

follows directly using arguments from the proof of Theorem 6.

Case 2 : $\frac{d}{\Delta_{\min}} > \frac{\kappa}{\Delta^2_{\neq}}$ Let the action set be given by $\mathcal{A} = \left\{ \begin{pmatrix} x_1 \\ z_{x_1} \end{pmatrix}, ..., \begin{pmatrix} x_{d+1} \\ z_{x_{d+1}} \end{pmatrix} \right\}$, where $\begin{pmatrix} x_1 \\ z_{x_1} \end{pmatrix} = e_1 + e_{d+1}$, $\begin{pmatrix} x_i \\ z_{x_i} \end{pmatrix} = e_i - e_{d+1}$ for $i \in \{2, ..., d\}$, and $\begin{pmatrix} x_{d+1} \\ z_{x_{d+1}} \end{pmatrix} = -\left(1 - \frac{2}{\sqrt{\kappa_*}+1} \right) e_1 - e_{d+1}$. By Lemma 17, $\mathcal{A} \in \mathbf{A}_{\kappa_*, d}$. We consider two bandit problems characterized by two parameters $\theta^{(1)}$ and $\theta^{(2)}$, defined as:

$$
\begin{aligned}
\gamma^{(1)} &= \frac{1+\Delta_{\neq}}{2} e_1 + \frac{1-\Delta_{\neq}}{2} e_2 - \frac{\Delta_{\neq}}{2} e_3 \\
\gamma^{(2)} &= \frac{1-\Delta_{\neq}}{2} e_1 + \frac{1+\Delta_{\neq}}{2} e_2 + \frac{\Delta_{\neq}}{2} e_3.
\end{aligned}
$$

On top of this, two bias parameters are defined as $\omega^{(1)} = -\frac{\Delta_{\neq}}{2}$ and $\omega^{(2)} = \frac{\Delta_{\neq}}{2}$.

The following facts hold:

Fact 1 For any $i \in \{1, 2\}$, action x_i is the unique optimal action in **Problem i**. Since $1/2 \geq \Delta_{\neq}$, sampling any other (sub-optimal) action leads to an instantaneous regret of at least $\Delta_{\neq}$.

Fact 2 In **Problem i**, for $i \in \{1, ..., d\}$, when sampling action x_j at time, t the distribution of the observation does not depend on t or on the past (except through the choice of x_j) and is $\mathbb{P}^{(i)}_j$. It is characterized as:

$$\forall i \in \{1, 2\}, \forall j \in \{1, 2\}, \mathbb{P}^{(i)}_j \text{ is } \mathcal{N}(1/2, 1)$$

$$\forall i \in \{1, 2\}, \mathbb{P}^{(1)}_3 \text{ is } \mathcal{N}(0, 1)$$

$$\mathbb{P}^{(1)}_{d+1} \text{ is } \mathcal{N}\left(\left(1 - \frac{2}{\sqrt{\kappa_*}+1} \right) \left(\frac{1+\Delta_{\neq}}{2} \right) + \frac{\Delta_{\neq}}{2}, 1 \right)$$

$$\mathbb{P}^{(2)}_{d+1} \text{ is } \mathcal{N}\left(\left(1 - \frac{2}{\sqrt{\kappa_*}+1} \right) \left(\frac{1-\Delta_{\neq}}{2} \right) - \frac{\Delta_{\neq}}{2}, 1 \right)$$

So that, between **Problem 1** and **Problem 2**, the only action that provides different evaluations when sampled is action 1, and the mean gaps in both cases is $\frac{2\Delta_{\neq}}{\sqrt{\kappa_*}+1}$.

Note that the minimum gap for these parameters is $\Delta_{\neq} \geq \Delta_{\min}$. Thus, this choice of parameters ensures that $\forall i \in \{1, ..., d\}, \theta^{(i)} \in \Theta^{\mathcal{A}}_{\Delta_{\min}, \Delta_{\neq}, \kappa_*}$. Adapting the proof of Lemma 17, we note that the minimal variance of bias estimation is at least κ_*. This proves that $\mathcal{A} \in \Theta^{\mathcal{A}}_{\Delta_{\min}, \Delta_{\neq}, \kappa_*}$. Then, the lower bound

$$R_T \geq \frac{\kappa_* + 1}{4\Delta^2_{\neq}} \left[\log(T) - \log\left(\frac{8\kappa_* \log(T)}{\Delta^3_{\neq}} \right) \right].$$

follows directly using arguments from the proof of Theorem 6.

C.7 Auxiliary Lemmas

C.7.1 Proof of Lemma 1

Lemma 1 follows from the characterization of κ_* given in Lemma 5. We begin by proving the first statement. Assume that $\kappa_* > 1$ (otherwise the first statement is void). Note that for all $u \in \mathbb{R}^d$, $\lim_{\lambda \to +\infty} (\max_{x \in \mathcal{X}} \left(x^\top (\lambda u) + z_x \right)^2)^{-1} = 0$, so the minimum over $u \in \mathbb{R}^d$ of $(\max_{x \in \mathcal{X}} \left(x^\top (\lambda u) + z_x \right)^2)^{-1}$ is attained for some vector $\tilde{u} \in \mathbb{R}^d$. Since $\kappa_* > 1$, $\tilde{u}$ is not null.

Moreover, $\max_{x\in\mathcal{X}}(1 + z_x x^\top \tilde{u})^2 < 1$, so $\max_{x\in\mathcal{X}} z_x x^\top \tilde{u} < 0$. Thus, for all $x \in \mathcal{X}$, $x^\top \tilde{u}$ and z_x are of opposite sign, and $x^\top \tilde{u} \neq 0$. This implies that the hyperplane containing 0 with normal vector $\tilde{u}$ contains no action, and separates the two groups. Moreover,

$$\kappa_*^{-1/2} = \max_{x\in\mathcal{X}} |z_x x^\top \tilde{u} + 1|.$$

We denote $x^{(1)} \in \mathrm{argmax}_{x\in\mathcal{X}}\, z_z x^\top \tilde{u}$, and $x^{(2)} \in \mathrm{argmin}_{x\in\mathcal{X}}\, z_z x^\top \tilde{u}$. Let us show that $(z_{x^{(1)}} x^{(1)\top} \tilde{u} + 1) = -\left(1 + z_{x^{(2)}} x^{(2)\top} \tilde{u}\right)$, i.e that $z_{x^{(1)}} x^{(1)\top} \tilde{u} + z_{x^{(2)}} x^{(2)\top} \tilde{u} = -2$. Indeed, note that

$$\kappa_*^{-1/2} = (z_{x^{(1)}} x^{(1)\top} \tilde{u} + 1) \vee -(1 + z_{x^{(2)}} x^{(2)\top} \tilde{u}).$$

Then, for $u' = \dfrac{-2}{\left(z_{x^{(1)}} x^{(1)} + z_{x^{(2)}} x^{(2)}\right)^\top \tilde{u}} \tilde{u}$, we see that

$$z_{x^{(1)}} x^{(1)\top} u' + 1 = -\left(1 + z_{x^{(2)}} x^{(2)\top} u'\right) = \max_{x\in\mathcal{X}} |z_x x^\top u' + 1|.$$

By contradiction, let us first assume that $z_{x^{(1)}} x^{(1)\top} \tilde{u} + z_{x^{(2)}} x^{(2)\top} \tilde{u} < -2$. Then,

$$\max_{x\in\mathcal{X}} |z_x x^\top u' + 1| = z_{x^{(1)}} x^{(1)\top} u' + 1 < z_{x^{(1)}} x^{(1)\top} \tilde{u} + 1 = \kappa_*^{-1/2}$$

which contradicts the definition of κ_*.

Similarly, if we assume that $z_{x^{(1)}} x^{(1)\top} \tilde{u} + z_{x^{(2)}} x^{(2)\top} \tilde{u} > -2$, then

$$\max_{x\in\mathcal{X}} |z_x x^\top u' + 1| = -(z_{x^{(2)}} x^{(2)\top} u' + 1) < -(z_{x^{(2)}} x^{(2)\top} \tilde{u} + 1) = \kappa_*^{-1/2}$$

which contradicts again the definition of κ_*. Therefore,

$$(z_{x^{(1)}} x^{(1)\top} \tilde{u} + 1) = -\left(1 + z_{x^{(2)}} x^{(2)\top} \tilde{u}\right) = \kappa_*^{-1/2}.$$

Then, the hyperplane containing 0 with normal vector $\tilde{u}$ separates the actions of the two groups. Moreover, the margin is $-z_{x^{(1)}} x^{(1)\top} \tilde{u} = 1 - \kappa_*^{-1/2}$, while the maximum distance of all points is $-z_{x^{(2)}} x^{(2)\top} \tilde{u} = 1 + \kappa_*^{-1/2}$. Thus, there exists $\tilde{u}$ such that the hyperplane containing 0 with normal vector $\tilde{u}$ separates the actions of the two groups, with margin equal to $\frac{\sqrt{\kappa_*}-1}{\sqrt{\kappa_*}+1}$ times the maximum distance of all points to the hyperplane.

Conversely, assume that there exists $\kappa > \kappa_*$ such that there exists $u \in \mathbb{R}^d$ such that the hyperplane containing 0 with normal vector u separates the actions of the two groups, with margin equal to $\frac{\sqrt{\kappa}-1}{\sqrt{\kappa}+1} = \frac{1-\kappa^{-1/2}}{1+\kappa^{-1/2}}$ times the maximum distance of all points to the hyperplane, denoted hereafter d. Since the hyperplane separates the points, we can assume without loss of generality that for all $x \in \mathcal{X}$, $z_x x^\top u < 0$. Similarly, up to a renormalization, we can assume without loss of generality that $d = 1 + \kappa^{-1/2}$. Then,

$$\begin{aligned}
\max_{x\in\mathcal{X}} |z_x x^\top u + 1| &= (\max_{x\in\mathcal{X}} z_x x^\top u + 1) \vee -(\min_{x\in\mathcal{X}} z_x x^\top u + 1) \\
&= \left(-\frac{1-\kappa^{-1/2}}{1+\kappa^{-1/2}} \times (1+\kappa^{-1/2}) + 1\right) \vee -(1 - \kappa^{-1/2} - 1) = \kappa^{-1/2} < \kappa_*^{-1/2}
\end{aligned}$$

which contradicts the definition of κ_*. This concludes the proof of the first statement.

To prove the second statement, let us assume that no separating hyperplane containing zero exists. Then, for all $u \in \mathbb{R}^d$, there exists $x \in \mathcal{X}$ such that $z_x x^\top u \geq 0$. This implies that $\min_{u\in\mathbb{R}^d} \max_{x\in\mathcal{X}}(z_x x^\top u + 1) \geq 1$, so $\kappa_* \leq 1$. Choosing $u = 0$, we see that $\kappa_* \geq 1$, which implies that $\kappa_* = 1$.

C.7.2 Proof of Lemma 2

Since for all $\gamma \in \mathcal{X}$ and all $x \in \mathcal{X}$, $|x^\top \gamma| \leq 1$, it is easy to see that the gaps are bounded by 2, and that $\widetilde{\kappa} \leq 2\kappa_*$.

Let us now show that $\widetilde{\kappa} \geq \kappa_*/2$.

$$
\left(x^{(1)}, x^{(2)}, \widetilde{\gamma}\right) \in \underset{(x,x')\in\mathcal{X}, \gamma\in\mathcal{C}(\mathcal{X})}{\operatorname{argmax}} (x - x')^\top \gamma
$$

$$
\overline{x} = \frac{1}{2}(x^{(1)} + x^{(2)})
$$

$$
\widetilde{n} = \sum_{x\in\mathcal{X}} \widetilde{\mu}(x)
$$

$$
\text{and} \quad \widetilde{x} = \frac{1}{\widetilde{n}} \sum_{x\in\mathcal{X}} \widetilde{\mu}(x)x.
$$

Recall that κ_* can equivalently be defined as the budget necessary to estimate the bias with a variance smaller than 1. Therefore, we have

$$
\widetilde{n} \geq \kappa_*. \tag{53}
$$

Let us define $\Delta_{\max}$ as $\Delta_{\max} = (x^{(1)} - x^{(2)})^\top \widetilde{\gamma} = \max_{(x,x')\in\mathcal{X}, \gamma\in\mathcal{C}(\mathcal{X})} (x - x')^\top \gamma$. By definition of $\widetilde{\kappa}$ and $\widetilde{\mu}$,

$$
\widetilde{\kappa} \geq \sum_{x\in\mathcal{X}} \widetilde{\mu}(x)(x^{(1)} - x)^\top \widetilde{\gamma}
$$

$$
= \widetilde{n}(x^{(1)} - \widetilde{x})^\top \widetilde{\gamma}.
$$

Using Equation (53), we find that

$$
\frac{\widetilde{\kappa}}{\kappa_*} \geq (x^{(1)} - \overline{x})^\top \widetilde{\gamma} + (\overline{x} - \widetilde{x})^\top \widetilde{\gamma}
$$

$$
= \frac{\Delta_{\max}}{2} + (\overline{x} - \widetilde{x})^\top \widetilde{\gamma}. \tag{54}
$$

Now, since $\widetilde{\gamma} \in \mathcal{C}(\mathcal{X})$, we also have $-\widetilde{\gamma} \in \mathcal{C}(\mathcal{X})$, and therefore

$$
\widetilde{\kappa} \geq \sum_{x\in\mathcal{X}} \widetilde{\mu}(x)(x^{(2)} - x)^\top (-\widetilde{\gamma})
$$

$$
= \widetilde{n}(\widetilde{x} - x^{(2)})^\top \widetilde{\gamma}
$$

Using again Equation (53), we find that

$$
\frac{\widetilde{\kappa}}{\kappa_*} \geq (\widetilde{x} - \overline{x})^\top \widetilde{\gamma} + (\overline{x} - x^{(2)})^\top \widetilde{\gamma}
$$

$$
= (\widetilde{x} - \overline{x})^\top \widetilde{\gamma} + \frac{\Delta_{\max}}{2}. \tag{55}
$$

Combining Equations (54) and (55), we find that

$$
\frac{\widetilde{\kappa}}{\kappa_*} \geq \frac{\Delta_{\max}}{2} + |(\overline{x} - \widetilde{x})^\top \widetilde{\gamma}|.
$$

This implies in particular that $\widetilde{\kappa} \geq \frac{\Delta_{\max}\kappa_*}{2}$.

To conclude the proof of the Lemma, we show that $\Delta_{\max} \geq 1$. By contradiction, assume that $\Delta_{\max} < 1$.

For all non-zero vector $u \in \mathbb{R}^d$, let us denote $x_u = \operatorname{argmax}_{x\in\mathcal{X}} |x^\top u|$. Since $\mathcal{X}$ spans $\mathbb{R}^d$, we necessarily have $|x_u^\top u| > 0$, so we can define the normalized vector $\tilde{u} = u/|x_u^\top u|$ such that $\tilde{u}$ belongs to the set $\mathcal{C}(\mathcal{X})$. Finally, denote $x_u^{(1)}, x_u^{(2)} \in \operatorname{argmax}_{x,x'\in\mathcal{X}} (x_u^{(1)} - x_u^{(2)})^\top \tilde{u}$. Note that by definition of $\Delta_{\max}$, we always have $(x_u^{(1)} - x_u^{(2)})^\top \tilde{u} \leq \Delta_{\max} < 1$.

Case 1 : $x_u^\top \tilde{u} > 0$ Then, by definition of x_u and $x_u^{(1)}$, we see that $x_u^{(1)^\top} \tilde{u} = x_u^\top \tilde{u} = 1$. Then, $(x_u^{(1)} - x_u^{(2)})^\top \tilde{u} < 1$ implies that $1 - x_u^{(2)^\top} \tilde{u} < 1$, so $x_u^{(2)^\top} \tilde{u} > 0$, and in particular $x_u^{(2)^\top} u > 0$.

Case 2 : $x_u^\top u < 0$ Then, by definition of x_u and $x_u^{(2)}$, we see that $x_u^{(2)^\top} \tilde{u} = x_u^\top u = -1$. Then $(x_u^{(1)} - x_u^{(2)})^\top \tilde{u} < 1$ implies that $x_u^{(1)^\top} \tilde{u} + 1 < 1$, so $x_u^{(1)^\top} \tilde{u} < 0$, and in particular $x_u^{(1)^\top} u < 0$.

Putting together Case 1 and Case 2, we see that $x_u^{(1)^\top} u$ and $x_u^{(2)^\top} u$ are of the same sign and are not null. By definition of $x_u^{(1)}$ and $x_u^{(2)}$, we conclude that for all $x \in \mathcal{X}$, the sign of $x^\top u$ is the same, and that $x^\top u$ is not 0. Since this is true for all non-zero vector u, this implies in particular that no hyperplane containing the origin can separate the actions, which contradicts the assumption that $\mathcal{X}$ spans $\mathbb{R}^d$.

C.7.3 Proof of Lemmas 3 and 4

We begin by proving Lemma 4. Recall that π is a G-optimal design for the set $\{a_x : x \in \mathcal{X}\}$, and that μ is defined as $\mu(x) = \lceil m\pi(x) \rceil$ for all $x \in \mathcal{X}$.

We first observe that $V(\pi) = A_\pi^\top A_\pi$, where A_π is the matrix with lines given by $[\sqrt{\pi(x)} a_x^\top]_{x \in \mathcal{X}}$. Since the supports of μ and π are the same, we get that $\text{Range}(A_\pi^\top) = \text{Range}(A_\mu^\top)$. As a consequence

$$\text{Range}(V(\pi)) = \text{Range}(A_\pi^\top) = \text{Range}(A_\mu^\top) = \text{Range}(V(\mu)),$$

and $x \in \text{Range}(V(\mu))$ for all $x \in \mathcal{X}$. This ensures that $a_x^\top \widehat{\theta}_\mu$ is an unbiased estimator of $a_x^\top \theta^*$.

Furthermore $V(\mu) \succcurlyeq mV(\pi)$, so the variance $a_x^\top V(\mu)^+ a_x$ of $a_x^\top \widehat{\theta}_\mu$ is upper-bounded by $a_x^\top V(\mu)^+ a_x \leq m^{-1} a_x^\top V(\pi)^+ a_x$. Now, the General Equivalence Theorem of Kiefer and Pukelshein shows that $\max_{x \in \mathcal{X}} a_x^\top V(\pi)^+ a_x \leq d+1$. Thus, $a_x^\top V(\pi)^+ a_x \leq m^{-1}(d+1)$.

We now prove Lemma 3. Recall that $\pi \in \mathcal{M}_{e_{d+1}}^{\mathcal{X}}$ is such that $e_{d+1} \in \text{Range } V(\pi)$, and that μ is defined as $\mu(x) = \lceil m\pi(x) \rceil$ for all $x \in \mathcal{X}$. Using similar arguments, we can show that $e_{d+1} \in \text{Range}(V(\mu))$, which ensures that $e_{d+1}^\top \widehat{\theta}_\mu$ is an unbiased estimator of $e_{d+1}^\top \theta^*$. The second part of the Lemma follows directly using that $V(\mu) \succcurlyeq mV(\pi)$.

C.7.4 Proof of Lemma 5

Elfving's set $\mathcal{S}$ for estimating the bias in the biased linear bandit problem is given by

$$\mathcal{S} = \text{convex hull} \left\{ \begin{pmatrix} x \\ z_x \end{pmatrix}, \begin{pmatrix} -x \\ -z_x \end{pmatrix} : x \in \mathcal{X} \right\},$$

or equivalently by

$$\mathcal{S} = \text{convex hull} \left\{ \pm \begin{pmatrix} z_x x \\ 1 \end{pmatrix} : x \in \mathcal{X} \right\}.$$

Now, Theorem 5 indicates that $\kappa_*^{-1/2} e_{d+1}$ belongs to a supporting hyperplane of $\mathcal{S}$. We first show that when $\mathcal{A}$ spans $\mathbb{R}^{d+1}$, any normal vector $w \in \mathbb{R}^{d+1}$ to this hyperplane is such that $w^\top e_{d+1} \neq 0$.

By contradiction, let us assume that $\kappa_*^{-1/2} e_{d+1}$ belongs to some supporting hyperplane $\mathcal{H}$ of $\mathcal{S}$ parametrized as $\mathcal{H} = \{a \in \mathbb{R}^{d+1} : a^\top w = b\}$, where the normal vector w is of the form $w = \begin{pmatrix} u \\ 0 \end{pmatrix}$. Then, $\kappa_*^{-1/2} e_{d+1} \in \mathcal{H}$, so $\kappa_*^{-1/2} e_{d+1}^\top w = b$, and thus $b = 0$. Now, $\mathcal{H}$ is a supporting hyperplane of $\mathcal{S}$, so for all $a \in \mathcal{S}$ we see that $a^\top w \leq b$. In particular, for all $x \in \mathcal{X}$, $x^\top u \leq 0$ and $-x^\top u \leq 0$, so $x^\top u = 0$. This implies that $\mathcal{X}$ is supported by an hyperplane in $\mathbb{R}^d$ with normal vector u, which contradicts our assumption that $\mathcal{A}$ spans $\mathbb{R}^{d+1}$. Thus, the supporting hyperplane of $\mathcal{S}$ containing $\kappa_*^{-1/2} e_{d+1}$ has a normal vector $w \in \mathbb{R}^{d+1}$ such that $w^\top e_{d+1} \neq 0$. In particular, we can parameterize this hyperplane as $\mathcal{H}_{u,b} = \{a \in \mathbb{R}^{d+1} : a^\top \begin{pmatrix} u \\ 1 \end{pmatrix} = b\}$ for some $b \in \mathbb{R}$ and $u \in \mathbb{R}^d$.

Now, if $\mathcal{H}_{u,b}$ is a supporting hyperplane of $\mathcal{S}$, then, by definition, $\mathcal{S}$ is contained in the half space $\{a \in \mathbb{R}^{d+1} : a^\top \begin{pmatrix} u \\ 1 \end{pmatrix} \leq b\}$. In particular, for all $x \in \mathcal{X}$, one must have $z_x x^\top u + 1 \leq b$ and

$-z_x x^\top u - 1 \le b$: therefore, for all $x \in \mathcal{X}$, $|z_x x^\top u + 1| \le b$. Moreover, $\mathcal{H}_{u,b}$ is a supporting hyperplane of $\mathcal{S}$, so there exists an extreme point $a \in \mathcal{S}$ such that $a \in \mathcal{H}_{u,b}$. Note that $\mathcal{S}$ is the convex hull of $\left\{ \pm \left(\begin{smallmatrix} z_x x \\ 1 \end{smallmatrix} \right) : x \in \mathcal{X} \right\}$, so the extreme points of $\mathcal{S}$ are in $\left\{ \pm \left(\begin{smallmatrix} z_x x \\ 1 \end{smallmatrix} \right) : x \in \mathcal{X} \right\}$. In particular, this implies that $b = \max \left\{ |z_x x^\top u + 1| : x \in \mathcal{X} \right\}$. Thus, the supporting hyperplane of $\mathcal{S}$ containing $\kappa_*^{-1/2} e_{d+1}$ is necessarily of the form $\mathcal{H}_{u,\max\{|z_x x^\top u+1|:x\in\mathcal{X}\}}$.

On the one hand, $\kappa_*^{-1/2}$ belongs to the boundary of $\mathcal{S}$ and therefore to a supporting hyperplane $\mathcal{H}_{u,\max\{|z_x x^\top u+1|:x\in\mathcal{X}\}}$ of $\mathcal{S}$. Then, there exists $u \in \mathbb{R}^d$ such that $\kappa_*^{-1/2} = \max \left\{ |z_x x^\top u + 1| : x \in \mathcal{X} \right\}$.

On the other hand, it is easy to verify that for all $u \in \mathbb{R}^d$, $\mathcal{H}_{u,\max\{|z_x x^\top u+1|:x\in\mathcal{X}\}}$ is a supporting hyperplane of $\mathcal{S}$. Now, $\kappa_*^{-1/2} e_{d+1}$ belongs to $\mathcal{S}$, so $\kappa_*^{-1/2} e_{d+1}^\top \left(\begin{smallmatrix} u \\ 1 \end{smallmatrix} \right) \le \max \left\{ |z_x x^\top u + 1| : x \in \mathcal{X} \right\}$. These two results imply that

$$\kappa_*^{-1/2} = \min_{u \in \mathbb{R}^d} \max_{x \in \mathcal{X}} |z_x x^\top u + 1|$$

which proves the Lemma.

C.7.5 Proof of Lemma 6

We prove that $2(\sqrt{\kappa_*} - 1)^2 \vee 1 \le \alpha \le 8(\kappa_* + 1)$. Lemma 6 follows directly by noticing that $\alpha \ge 1$ and $\kappa_* \ge 1$.

Let us begin by proving that $2(\sqrt{\kappa_*} - 1)^2 \le \alpha$ for $\kappa_* > 1$ (otherwise this inequality is automatically verified). Note that for all $u \in \mathbb{R}^d$, $\lim_{\lambda \to +\infty} \frac{1}{\max_{x \in \mathcal{X}} (x^\top (\lambda u) + z_x)^2} = 0$, so the minimum over $u \in \mathbb{R}^d$ of $\frac{1}{\max_{x \in \mathcal{X}} (x^\top u + z_x)^2} = 0$ is attained for some vector $\tilde{u} \in \mathbb{R}^d$. Let us also denote $\tilde{x} \in \mathrm{argmax}_{x \in \mathcal{X}} (z_x x^\top \tilde{u} + 1)^2$, such that

$$\kappa_* = \frac{1}{\left(z_{\tilde{x}} \tilde{x}^\top \tilde{u} + 1\right)^2}.$$

With these notations, we see that for all $x \in \mathcal{X}$,

$$(z_x x^\top \tilde{u} + 1)^2 \le (z_{\tilde{x}} \tilde{x}^\top \tilde{u} + 1)^2 = \kappa_*^{-1} < 1.$$

This implies that for all $x \in \mathcal{X}$,

$$z_x x^\top \tilde{u} \le -1 + \kappa_*^{-1/2} < 0.$$

Now, let us denote $x^{(1)}, x^{(2)} \in \mathrm{argmax}_{x,x' \in \mathcal{X}} (x - x')^\top \tilde{u}$. By definition of α, we see that

$$\alpha \ge \frac{\left((x^{(1)} - x^{(2)})^\top \tilde{u}\right)^2}{\left(z_{\tilde{x}} \tilde{x}^\top \tilde{u} + 1\right)^2} = \left((x^{(1)} - x^{(2)})^\top \tilde{u}\right)^2 \times \kappa_*.$$

Since $z_x x^\top \tilde{u} < 0$ for all $x \in \mathcal{X}$, and since no group is empty, we can conclude that there exists $x, x' \in \mathcal{X}$ such that $x^\top \tilde{u} > 0$ and $x'^\top \tilde{u} < 0$. In particular, by definition of $x^{(1)}$ and $x^{(2)}$, we see that $(x^{(1)})^\top \tilde{u} > 0$ and $(x^{(2)})^\top \tilde{u} < 0$. Then,

$$\left((x^{(1)} - x^{(2)})^\top \tilde{u}\right)^2 \ge \left((x^{(1)})^\top \tilde{u}\right)^2 + \left((x^{(2)})^\top \tilde{u}\right)^2 \ge 2(1 - \kappa_*^{-1/2})^2.$$

This implies that

$$\alpha \ge 2(1 - \kappa_*^{-1/2})^2 \times \kappa_* = 2(\sqrt{\kappa_*} - 1)^2.$$

Let us now prove that $\alpha \ge 1$. Note that by assumption, $\mathcal{X}$ spans $\mathbb{R}^d$, and in particular there exists $\tilde{u} \in \mathbb{R}^d$ and $x, x' \in \mathcal{X}$ such that $\max_{x \in \mathcal{X}} x^\top \tilde{u} > 0$ and $\min_{x \in \mathcal{X}} x^\top \tilde{u} \le 0$. Thus, $\max_{x,x' \in \mathcal{X}} ((x - x')^\top \tilde{u})^2 \ge \max_{x \in \mathcal{X}} (x^\top \tilde{u})^2$. For any $\lambda > 0$, choosing $u = \lambda \tilde{u}$ in the definition of α implies that

$$\alpha \ge \frac{\lambda^2 \max_{x \in \mathcal{X}} (x^\top u)^2}{\max_{x \in \mathcal{X}} (\lambda z_x x^\top u + 1)^2}.$$

Letting λ go to infinity, we find that $\alpha \geq 1$.

Finally, we prove that $\alpha \leq 8(\kappa_* + 1)$. For all $u \in \mathbb{R}^d$, we see that

$$\frac{\max_{x,x' \in \mathcal{X}}((x - x')^\top u)^2}{\max_{x \in \mathcal{X}}(z_x x^\top u + 1)^2} \leq \frac{4 \max_{x \in \mathcal{X}}(z_x x^\top u)^2}{\max_{x \in \mathcal{X}}(z_x x^\top u + 1)^2}.$$

Now, we see that

$$\frac{\max_{x \in \mathcal{X}}(z_x x^\top u)^2}{\max_{x \in \mathcal{X}}(z_x x^\top u + 1)^2} \leq \frac{2 \max_{x \in \mathcal{X}}(z_x x^\top u + 1)^2 + 2}{\max_{x \in \mathcal{X}}(z_x x^\top u + 1)^2} \leq 2 + \frac{2}{\max_{x \in \mathcal{X}}(z_x x^\top u + 1)^2}.$$

This in turn implies that for all $u \in \mathbb{R}^d$,

$$\frac{\max_{x,x' \in \mathcal{X}}((x - x')^\top u)^2}{\max_{x \in \mathcal{X}}(z_x x^\top u + 1)^2} \leq 8(1 + \kappa_*),$$

which finally implies that $\alpha \leq 8(1 + \kappa_*)$.

C.7.6 Proof of Lemma 8

Proof of Claim i) The proof of the first claim is immediate by definition of κ. Indeed, let $\widetilde{\mathcal{M}} = \left\{ \mu \in \mathcal{M}_{e_{d+1}}^{\mathcal{X}} : e_{d+1}^\top V(\mu)^+ e_{d+1} \leq 1 \right\}$ be the set of measures μ admissible for estimating ω^* with a precision level 1. Then,

$$\kappa(c\Delta) \quad = \quad \min_{\mu \in \widetilde{\mathcal{M}}} \sum_x \mu(x) c\Delta_x = c \min_{\mu \in \widetilde{\mathcal{M}}} \sum_x \mu(x)\Delta_x = c\kappa(\Delta).$$

Proof of Claim ii) The proof of the second claim is also straightforward. If $\Delta \leq \Delta'$, then for all $\mu \in \widetilde{\mathcal{M}}$, $\sum_x \mu(x)\Delta_x \leq \sum_x \mu(x)\Delta'_x$. Recall that $\mu^{\Delta'} = \operatorname{argmin}_{\mu \in \widetilde{\mathcal{M}}} \sum_x \mu(x)\Delta'_x$. Then,

$$\kappa(\Delta') \quad = \quad \sum_x \mu^{\Delta'}(x)\Delta'_x \geq \sum_x \mu^{\Delta'}(x)\Delta_x \geq \min_{\mu \in \widetilde{\mathcal{M}}} \sum_x \mu(x)\Delta_x = \kappa(\Delta).$$

Proof of Claim iii) To prove the third claim, note that

$$\kappa(\Delta \vee \Delta') \quad = \quad \min_{\mu \in \widetilde{\mathcal{M}}} \sum_x \mu(x) \left(\Delta_x \vee \Delta_x\right)$$

$$\geq \quad \min_{\mu \in \widetilde{\mathcal{M}}} \left(\sum_x \mu(x)\Delta_x \vee \sum_x \mu(x)\Delta'_x \right)$$

$$\geq \quad \left(\min_{\mu \in \widetilde{\mathcal{M}}} \sum_x \mu(x)\Delta_x \right) \vee \left(\min_{\mu \in \widetilde{\mathcal{M}}} \sum_x \mu(x)\Delta'_x \right)$$

$$\geq \quad \kappa(\Delta) \vee \kappa(\Delta').$$

Proof of Claim iv) Recall that

$$\kappa(\Delta) = \min_{\mu \in \widetilde{\mathcal{M}}} \sum_x \mu(x)\Delta_x.$$

Let us define a sequence $(\mu_n)_{n \in \mathbb{N}} \in \widetilde{\mathcal{M}}^{\mathbb{N}}$ such that $\sum_x \mu_n(x)\Delta_x \xrightarrow[n \to \infty]{} \kappa(\Delta)$, and let us denote $\kappa_n = \sum_x \mu_n(x)\Delta_x$. According to Claim ii), we have

$$\kappa(\Delta) \leq \kappa(\Delta \vee \epsilon) = \min_{\mu \in \widetilde{\mathcal{M}}} \sum_x \mu(x) \left(\Delta_x \vee \epsilon\right) \leq \sum_x \mu_n(x)\Delta_x + \epsilon \sum_x \mu_n(x).$$

It follows that for all n,

$$\kappa(\Delta) \leq \liminf_{\epsilon \to 0^+} \kappa(\Delta \vee \epsilon) \leq \limsup_{\epsilon \to 0^+} \kappa(\Delta \vee \epsilon) \leq \kappa_n.$$

Letting n go to infinity, we get that $\lim_{\epsilon \to 0^+} \kappa(\Delta \vee \epsilon) = \kappa(\Delta)$.

C.7.7 Proof of Lemma 9

Setting $\mu \cdot \Delta = (\mu(x)\Delta_x)_{x\in\mathcal{X}}$ and

$$
V_\Delta(\lambda) = \sum_{x\in\mathcal{X}} \lambda_x \begin{pmatrix} \Delta_x^{-1/2}x \\ \Delta_x^{-1/2}z_x \end{pmatrix} \begin{pmatrix} \Delta_x^{-1/2}x \\ \Delta_x^{-1/2}z_x \end{pmatrix}^\top ,
$$

we observe that $V_\Delta(\mu \cdot \Delta) = V(\mu)$. Hence,

$$
\kappa(\Delta) = \min_{\substack{\mu\in\mathcal{M}^+ \\ e_{d+1}^\top V_\Delta(\mu\cdot\Delta)^+ e_{d+1}\leq 1}} \sum_{x\in\mathcal{X}} (\mu \cdot \Delta)_x.
$$

We observe that $e_{d+1} \in \mathrm{Range}(V(\mu))$ is equivalent to $e_{d+1} \in \mathrm{Range}(V_\Delta(\mu\cdot\Delta))$. Hence, $\mu^\Delta \cdot \Delta = \lambda^\Delta$ where

$$
\lambda^\Delta \in \operatorname*{argmin}_{\substack{\lambda\in\mathbb{R}_+^{\mathcal{X}} \\ e_{d+1}\in\mathrm{Range}(V_\Delta(\lambda)) \\ e_{d+1}^\top V_\Delta(\lambda)^+ e_{d+1}\leq 1}} \sum_{x\in\mathcal{X}} \lambda_x.
$$

The conclusion then follows by noticing that by homogeneity, $\lambda^\Delta = \kappa^\Delta \pi^\Delta$.

C.7.8 Proof of Lemma 12

Lemma 12 follows directly from Lemmas 18 and 19.

Lemma 18.

$$
\mathbb{P}\left(\exists l \geq 1, z \in \{-1,1\} \text{ such that } \mathit{Explore}_l^{(z)} = \mathit{True}, \text{ and } x \in \mathcal{X}_l^{(z)} \text{ such that } \left| \begin{pmatrix} \widehat{\gamma}_l^{(z)} - \gamma^* \\ \widehat{\omega}_l^{(z)} - \omega^* \end{pmatrix}^\top \begin{pmatrix} x \\ z_x \end{pmatrix} \right| \geq \epsilon_l \right) \leq \delta.
$$

Lemma 19.

$$
\mathbb{P}\left(\exists l \geq 1 \text{ such that } \mathit{Explore}_l^{(0)} = \mathit{True} \text{ and } \left| \widehat{\omega}_l^{(0)} - \omega^* \right| \geq \epsilon_l \right) \leq \delta.
$$

C.7.9 Proof of Lemma 13

To prove Lemma 13, we rely on the following key lemma. This lemma proves that on $\overline{\mathcal{F}}$, i.e. when the error bounds hold, the algorithm never eliminates the best action or the best group.

Lemma 20. *On the event $\overline{\mathcal{F}}$, for all $x^* \in \operatorname{argmax}_{x\in\mathcal{X}} x^\top \gamma^*$ and all l such that $\mathit{Explore}_l^{(z_{x^*})} = True$, $x^* \in \mathcal{X}_{l+1}^{(z_{x^*})}$. Moreover, on the event $\overline{\mathcal{F}}$, for all l such that $\mathit{Explore}_l^{(0)} = True$, there exists $x^* \in \operatorname{argmax}_{x\in\mathcal{X}} x^\top \gamma^*$ such that $\widehat{z}^*_{l+1} \neq -z_{x^*}$.*

Let $l \geq 1$ be such that $\mathit{Explore}_l^{(z_{x^*})} = \text{True}$. Then, on $\overline{\mathcal{F}}$, $x^* \in \mathcal{X}_{l+1}^{(z_{x^*})}$ by Lemma 20. Moreover, for all $x \in \mathcal{X}_{l+1}^{(z_{x^*})}$, by definition of $\mathcal{X}_{l+1}^{(z_{x^*})}$, we have that on $\overline{\mathcal{F}}$

$$
\left(\begin{pmatrix} x^* \\ z_{x^*} \end{pmatrix} - \begin{pmatrix} x \\ z_{x^*} \end{pmatrix} \right)^\top \begin{pmatrix} \widehat{\gamma}_l^{(z)} \\ \widehat{\omega}_l^{(z)} \end{pmatrix} \leq 3\epsilon_l.
$$

which implies that

$$
\left(\begin{pmatrix} x^* \\ z_{x^*} \end{pmatrix} - \begin{pmatrix} x \\ z_{x^*} \end{pmatrix} \right)^\top \begin{pmatrix} \gamma^* \\ \omega^* \end{pmatrix} \leq 3\epsilon_l + \left| \begin{pmatrix} x^* \\ z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(z)} - \gamma^* \\ \widehat{\omega}_l^{(z)} - \omega^* \end{pmatrix} \right| + \left| \begin{pmatrix} x \\ z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(z)} - \gamma^* \\ \widehat{\omega}_l^{(z)} - \omega^* \end{pmatrix} \right|.
$$

Thus, on the event $\overline{\mathcal{F}}$, for all $x \in \mathcal{X}_{l+1}^{(z_{x^*})}$

$$
(x^* - x)^\top \gamma^* < 5\epsilon_l,
$$

which proves Equation (13). To prove the second claim of Lemma 13, assume that for all $x' \in \operatorname{argmax}_{x\in\mathcal{X}} x^\top \gamma^*$, $z_{x'} = z_{x^*}$ (when this does not hold, the second claim follows from Equation

(13)). Now, let $l \geq 1$ be such that $\text{Explore}_l^{(-z_{x^*})} = \text{True}$. By Lemma 20, on $\overline{\mathcal{F}}$, $x^* \in \mathcal{X}_l^{(z_{x^*})}$ and $\widehat{z}_l^* = 0$. Then, the algorithm is unable to determine the group containing the best set during the phase $\text{Exp}_{l-1}^{(0)}$, so there must exist $x' \in \mathcal{X}_l^{(-z_{x^*})}$ such that

$$
\begin{pmatrix} x^* \\ z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_{l-1}^{(z_{x^*})} \\ \widehat{\omega}_{l-1}^{(z_{x^*})} \end{pmatrix} \leq \begin{pmatrix} x' \\ -z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_{l-1}^{(-z_{x^*})} \\ \widehat{\omega}_{l-1}^{(-z_{x^*})} \end{pmatrix} + 2z_{x^*}\widehat{\omega}_{l-1}^{(0)} + 4\epsilon_{l-1}.
$$

It follows that

$$
\begin{pmatrix} x^* - x' \\ 2z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \gamma^* \\ \omega^* \end{pmatrix} \leq \begin{pmatrix} x^* \\ z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \gamma^* - \widehat{\gamma}_{l-1}^{(z_{x^*})} \\ \omega^* - \widehat{\omega}_{l-1}^{(z_{x^*})} \end{pmatrix} + \begin{pmatrix} x' \\ -z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_{l-1}^{(-z_{x^*})} - \gamma^* \\ \widehat{\omega}_{l-1}^{(-z_{x^*})} - \omega^* \end{pmatrix} + 2z_{x^*}\widehat{\omega}_{l-1}^{(0)} + 4\epsilon_{l-1}.
$$

On $\overline{\mathcal{F}}$, this implies that

$$
\begin{pmatrix} x^* - x' \\ 2z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \gamma^* \\ \omega^* \end{pmatrix} < 2z_{x^*}\widehat{\omega}_{l-1}^{(0)} + 6\epsilon_{l-1}
$$

so

$$
(x^* - x')^\top \gamma^* \leq 2z_{x^*}\left(\widehat{\omega}_{l-1}^{(0)} - \omega^*\right) + 6\epsilon_{l-1} < 8\epsilon_{l-1} = 16\epsilon_l. \tag{56}
$$

Moreover, for all $x \in \mathcal{X}_{l+1}^{(-z_{x^*})}$ we have $(a_{x'} - a_x)^\top \theta_l^{(-z_{x^*})} \leq 3\epsilon_l$, so following the same lines as for the first claim, we get $(x' - x)^\top \gamma^* < 5\epsilon_l$. Combining this bound with (56), we get

$$
\max_{x \in \mathcal{X}_{l+1}^{(-z_{x^*})}} (x^* - x)^\top \gamma^* < 21\epsilon_l.
$$

This concludes the proof of Lemma 13.

C.7.10   Proof of Lemma 14

For $z \in \{-1, +1\}$ and $l > 0$,

$$
\sum_x \mu_l^{(z)}(x) \leq \sum_x \frac{2(d+1)\pi_l^{(z)}(x)}{\epsilon_l^2} \log\left(\frac{kl(l+1)}{\delta}\right) + |\text{supp}(\pi_l^{(z)})|.
$$

Now, $\text{supp}(\pi_l^{(z)}) \leq \frac{(d+1)(d+2)}{2}$ and $\sum_x \pi_l^{(z)}(x) = 1$, so

$$
\sum_x \mu_l^{(z)}(x) \leq \frac{2(d+1)}{\epsilon_l^2} \log\left(\frac{kl(l+1)}{\delta}\right) + \frac{(d+1)(d+2)}{2}
$$

which proves the first claim of Lemma 14.

To prove the second claim, we bound the regret for bias estimation at stage l as follows. On $\overline{\mathcal{F}}$, we have $\Delta_x \leq \widehat{\Delta}_x^l$ for all $x \in \mathcal{X}$ and $l \geq 1$, so

$$
\sum_{x \in \mathcal{X}} \mu_l^{(0)}(x)\Delta_x \leq \sum_{x \in \mathcal{X}} \mu_l^{(0)}(x)\widehat{\Delta}_x^l.
$$

Recall that $\hat{\mu}_l$ is the $\widehat{\Delta}^l$-optimal design, and that for all $x \in \mathcal{X}$, $\mu_l^{(0)}(x) = \lceil \frac{2\hat{\mu}_l(x)}{\epsilon_l^2} \log\left(\frac{l(l+1)}{\delta}\right)\rceil$. Since $\widehat{\Delta}_x^l \leq 2$ for all $x \in \mathcal{X}$, we have

$$
\sum_{x \in \mathcal{X}} \mu_l^{(0)}(x)\widehat{\Delta}_x^l \leq \sum_{x \in \mathcal{X}} \frac{2\hat{\mu}_l(x)}{\epsilon_l^2} \log\left(\frac{l(l+1)}{\delta}\right)\widehat{\Delta}_x^l + 2|\text{supp}(\mu_l^{(0)})|
$$

and $|\text{supp}(\mu_l^{(0)})| \leq d+1$, so

$$
\sum_x \mu_l^{(0)}(x)\Delta_x \leq \frac{2}{\epsilon_l^2} \log\left(\frac{l(l+1)}{\delta}\right) \sum_{x \in \mathcal{X}} \hat{\mu}_l(x)\widehat{\Delta}_x^l + 2(d+1).
$$

By definition of $\hat{\mu}_l(x)$, we have that

$$
\sum_{x \in \mathcal{X}} \hat{\mu}_l(x)\widehat{\Delta}_x^l = \kappa(\widehat{\Delta}^l).
$$

It follows that, on $\overline{\mathcal{F}}$,

$$
\sum_x \mu_l^{(0)}(x)\Delta_x \leq \sum_x \mu_l^{(0)}(x)\widehat{\Delta}_x^l \leq \frac{2}{\epsilon_l^2} \log\left(\frac{l(l+1)}{\delta}\right)\kappa(\widehat{\Delta}^l) + 2(d+1).
$$

C.7.11 Proof of Lemma 15

For the first claim, we rely on the next lemma.

Lemma 21. *Let us set $\ell_x = \max\left\{l \geq 1 : x \in \mathcal{X}_l^{(-1)} \cup \mathcal{X}_l^{(1)}\right\}$. On $\overline{\mathcal{F}}$, we have for any $l \geq 1$*

1. *$\widehat{\Delta}_x^l \leq \Delta_x + 16\epsilon_l$ for all $x \in \mathcal{X}_l^{(-1)} \cup \mathcal{X}_l^{(1)}$ (i.e. for all x such that $l \leq \ell_x$);*

2. *if $\Delta_x \geq 21\epsilon_l$ then $\ell_x \leq l$;*

3. *$\epsilon_{\ell_x} < \Delta_x$ for all $x \in \mathcal{X}$.*

Lemma 15 relies on the following remarks : if Δ, Δ' are such that $\Delta_x \leq \Delta_x'$ for all $x \in \mathcal{X}$, then by Lemma 8 (ii)), $\kappa(\Delta) \leq \kappa(\Delta')$. Let us now prove that for all $l \geq 1$ and all $x \in \mathcal{X}$, $\widehat{\Delta}_x^l \leq 513(\Delta \vee \epsilon_l)$.

Case $\epsilon_l \geq \Delta_x$. On $\overline{\mathcal{F}}$, we have $l \leq \ell_x - 1$ according to the third claim of Lemma 21. So, on $\overline{\mathcal{F}}$,

$$\widehat{\Delta}_x^l \leq \Delta_x + 16\epsilon_l \leq 17(\Delta_x \vee \epsilon_l).$$

Case $\epsilon_l < \Delta_x$. Then, on $\overline{\mathcal{F}}$, we have $32\epsilon_{l+5} < \Delta_x$ and so $l + 5 \geq \ell_x$ according to the second claim of Lemma 21. Hence, on $\overline{\mathcal{F}}$, according to Lemma 21, we have

$$\widehat{\Delta}_x^l \leq \max_{k=0,\ldots,5} \widehat{\Delta}_x^{\ell_x - k} \leq \Delta_x + 16\epsilon_{\ell_x - 5}$$

$$\leq \Delta_x + 512\epsilon_{\ell_x} \leq 513\Delta_x.$$

Thus, for all $l \geq 1$ and all $x \in \mathcal{X}$,

$$\widehat{\Delta}_x^l \leq 513(\Delta \vee \epsilon_l).$$

Now, let $\widetilde{\mathcal{M}} = \left\{\mu \in \mathcal{M}_{e_{d+1}}^{\mathcal{X}} : e_{d+1}^\top V(\mu)^+ e_{d+1} \geq 1\right\}$ the measures μ admissible for estimating ω^* with a precision level 1. Note that for all $a, b, c > 0$,

$$(1 + ab^{-1})(c \vee b) = (c + cab^{-1}) \vee (a + b) \geq c \vee (a + b) \geq c \vee a. \tag{57}$$

Using Equation (57) with $a = \Delta_x$, $b = \tau$ and $c = \epsilon$, we see that

$$\kappa(\Delta \vee \epsilon) \;\; = \;\; \min_{\mu \in \widetilde{\mathcal{M}}} \sum_x \mu(x)(\Delta_x \vee \epsilon) \leq (1 + \epsilon/\tau) \min_{\mu \in \widetilde{\mathcal{M}}} \sum_x \mu(x)(\Delta_x \vee \tau) = (1 + \epsilon/\tau)\kappa(\Delta \vee \tau).$$

Using Lemma 8 together with $\widehat{\Delta}_x^l \leq 513(\Delta \vee \epsilon_l)$, we find that

$$\kappa(\widehat{\Delta}_x^l) \leq 513\kappa(\Delta \vee \epsilon_l) \leq 513(1 + \epsilon_l/\tau)\kappa(\Delta \vee \tau).$$

This proves the first claim of Lemma 15.

To prove the second claim, we use Lemma 8 and the fact that for all x, $\widehat{\Delta}_x^l \geq \epsilon_l$. Moreover, on $\overline{\mathcal{F}}$, $\widehat{\Delta}_x^l \geq \Delta_x$ for all $x \in \mathcal{X}$. Then, $\kappa(\widehat{\Delta}) \geq \kappa(\epsilon_l \vee \Delta)$ by Lemma 8 (iii)).

C.7.12 Proof of Lemmas 16

To prove Lemma 16, let us consider l such that $\epsilon_l \leq \frac{\Delta_{\neq}}{8}$. According to Lemma 20, on $\overline{\mathcal{F}}$ we know that $\widehat{z^*}_l \neq -z_{x^*}$. When $\widehat{z^*}_l = z_{x^*}$, then we also have $\widehat{z^*}_{l+1} = z_{x^*}$ and the conclusion follows immediately. Let us consider now the case where $\widehat{z^*}_l = 0$. By definition of $\Delta_{\neq}$, for all $x' \in \mathcal{X}_{l+1}^{(-z_{x^*})}$,

$$(x^* - x')^\top \gamma^* \geq \Delta_{\neq}.$$

This implies that

$$\begin{pmatrix} x^* \\ z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(z_{x^*})} \\ \widehat{\omega}_l^{(z_{x^*})} \end{pmatrix} - z_{x^*} \widehat{\omega}_l^{(0)} \;\; \geq \;\; \max_{x \in \mathcal{X}_{l+1}^{(-z_{x^*})}} \begin{pmatrix} x \\ -z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(-z_{x^*})} \\ \widehat{\omega}_l^{(-z_{x^*})} \end{pmatrix} + z_{x^*} \widehat{\omega}_l^{(0)}$$

$$+ \begin{pmatrix} x^* \\ z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(z_{x^*})} - \gamma^* \\ \widehat{\omega}_l^{(z_{x^*})} - \omega^* \end{pmatrix} + \min_{x \in \mathcal{X}_{l+1}^{(-z_{x^*})}} \begin{pmatrix} x \\ -z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \gamma^* - \widehat{\gamma}_l^{(-z_{x^*})} \\ \omega^* - \widehat{\omega}_l^{(-z_{x^*})} \end{pmatrix}$$

$$+ \Delta_{\neq} + 2z_{x^*} \left(\omega^* - \widehat{\omega}_l^{(0)}\right).$$

On $\overline{\mathcal{F}}$, it follows that

$$\begin{pmatrix} x^* \\ z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(z_{x^*})} \\ \widehat{\omega}_l^{(z_{x^*})} \end{pmatrix} - z_{x^*}\widehat{\omega}_l^{(0)} - 2\epsilon_l \;\geq\; \max_{x \in \mathcal{X}_{l+1}^{(-z_{x^*})}} \begin{pmatrix} x \\ -z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(-z_{x^*})} \\ \widehat{\omega}_l^{(-z_{x^*})} \end{pmatrix} + z_{x^*}\widehat{\omega}_l^{(0)} - 6\epsilon_l + \Delta_{\neq}.$$

When $\Delta_{\neq} \geq 8\epsilon_l$, this implies that $\widehat{z_{l+1}^*} = z_{x^*}$.

C.7.13 Proof of Lemmas 10 and 17

We prove Lemma 10. The proof of Lemma 17 follows by noticing that the two actions sets are equal up to a permutation of the direction of some basis vectors. To prove Lemma 17, we rely on Elfving's characterization of c-optimal design, given in Theorem 5. Theorem 5 shows that for $\pi \in \mathcal{P}^{\{1,...,d+1\}}$ to be e_{d+1}-optimal, there must exist $t > 0$ and $\zeta \in \{-1,+1\}^{d+1}$ such that

$$\sum_{1 \leq i \leq d+1} \pi_i \;=\; 1$$

$$0 \;=\; \pi_1 \zeta_1 - (1 - \frac{2}{\sqrt{\kappa_*}+1})\pi_{d+1}\zeta_{d+1}$$

$$\forall i \in \{2,...,d\},\; 0 \;=\; \pi_i \zeta_i$$

$$t \;=\; \sum_{1 \leq i \leq \lfloor d/2 \rfloor} \pi_i \zeta_i - \sum_{\lfloor d/2 \rfloor + 1 \leq i \leq d+1} \pi_i \zeta_i.$$

Solving this system, we find that $t^{-2} = \kappa_*$. Note that the unicity of the solution for the corresponding probability measure π guarantees that $t e_{d+1}$ belongs to the boundary of $\mathcal{S}$.

C.7.14 Proof of Lemma 11

For a given parameter γ^*, let us denote by Δ_i the gap corresponding to the action i. To compute $\kappa(\Delta)$, we could want to rely on Lemma 9 to find the Δ-optimal design, corresponding to the e_{d+1}-optimal design on the rescaled features $\Delta_x^{-1/2}\begin{pmatrix} x \\ z_x \end{pmatrix}$. Theorem 5 indeed allows us to compute such a design, as seen in the proof of Lemma 10. Unfortunately, we cannot rescale the features using the true gaps, since $\Delta_{x^*} = 0$. To circumvent this problem, we rely on the following reasoning :

1. We use Lemma 9 and Theorem 5 to compute the design $\mu^{\Delta \vee \epsilon}$ for $\epsilon \in (0, \Delta_{\min})$; and the corresponding regret $\kappa(\Delta \vee \epsilon)$;
2. We find the value of $\kappa(\Delta)$ by noticing that $\epsilon \mapsto \kappa(\Delta \vee \epsilon)$ is continuous at 0.

For $\epsilon \in (0, \Delta_{\min})$, define $\overline{\Delta} = \Delta \vee \epsilon$, and $\overline{x} = \overline{\Delta}_x^{-1/2} x$. Let $\overline{\pi}$ denote the e_{d+1}-optimal design for the rescaled features $\overline{x}$, and let $\overline{\kappa_*}$ denote its variance. Then, Lemma 9 ensures that $\kappa(\overline{\Delta}) = \overline{\kappa_*}$.

Now, Theorem 5 shows that there exists $\zeta \in \{-1,+1\}^{d+1}$ such that

$$\sum_{1 \leq i \leq d+1} \overline{\pi}_i \;=\; 1$$

$$0 \;=\; \overline{\pi}_1 \zeta_1 \overline{\Delta}_1^{-1/2} - (1 - \frac{2}{\sqrt{\kappa_*}+1})\overline{\pi}_{d+1}\zeta_{d+1}\overline{\Delta}_{d+1}^{-1/2}$$

$$\forall i \in \{2,...,d\},\; 0 \;=\; \overline{\pi}_i \zeta_i \overline{\Delta}_i^{-1/2}$$

$$\overline{\kappa_*}^{-1/2} \;=\; \sum_{1 \leq i \leq \lfloor d/2 \rfloor} \overline{\pi}_i \zeta_i \overline{\Delta}_i^{-1/2} - \sum_{\lfloor d/2 \rfloor + 1 \leq i \leq d+1} \overline{\pi}_i \zeta_i \overline{\Delta}_i^{-1/2}$$

and $\overline{\kappa_*}^{-1/2} e_{d+1}$ belongs to the boundary of $\mathcal{S}$. Solving this system, we find that

$$\kappa(\overline{\Delta})^{-1/2} = \overline{\kappa_*}^{-1/2} = \frac{\left(\frac{2}{\sqrt{\kappa_*}+1}\right)\overline{\Delta}_{d+1}^{-1/2}}{1 + \left(1 - \frac{2}{\sqrt{\kappa_*}+1}\right)\overline{\Delta}_{d+1}^{-1/2}\overline{\Delta}_1^{1/2}}.$$

As in Lemma 10, the unicity of the solution for the corresponding probability measure $\overline{\pi}$ guarantees that $\overline{\kappa_*}^{-1/2}e_{d+1}$ belongs to the boundary of the Elfving's set. Now, $\epsilon \leq \Delta_{\min}$, so

$$\kappa(\overline{\Delta})^{-1/2} = \kappa(\Delta \vee \epsilon)^{-1/2} = \frac{\left(\frac{2}{\sqrt{\kappa_*}+1}\right)\Delta_{d+1}^{-1/2}}{1 + \left(1 - \frac{2}{\sqrt{\kappa_*}+1}\right)\Delta_{d+1}^{-1/2}\epsilon^{1/2}}.$$

The fourth claim of Lemma 8 ensures that $\kappa(\Delta \vee \epsilon) \underset{\epsilon \to 0}{\to} \kappa(\Delta)$. Therefore,

$$\kappa(\Delta) = \lim_{\epsilon \to 0} \left(\frac{\left(\frac{2}{\sqrt{\kappa_*}+1}\right)\Delta_{d+1}^{-1/2}}{1 + \left(1 - \frac{2}{\sqrt{\kappa_*}+1}\right)\Delta_{d+1}^{-1/2}\epsilon^{1/2}}\right)^{-2} = \frac{(\sqrt{\kappa_*}+1)^2\Delta_{d+1}}{4}.$$

C.7.15 Proof of Lemma 18

Recall that $\xi_t = y_t - x_t^\top\gamma^* - z_{x_t}\omega^*$. For $l \geq 0$ and $z \in \{-1, +1\}$, when $\text{Explore}_l^{(z)} = \text{True}$, the least square estimator $\begin{pmatrix}\widehat{\gamma}_l^{(z)} \\ \widehat{\omega}_l^{(z)}\end{pmatrix}$ is given by

$$\begin{pmatrix}\widehat{\gamma}_l^{(z)} \\ \widehat{\omega}_l^{(z)}\end{pmatrix} = \left(V_l^{(z)}\right)^+ \sum_{t \in \text{Exp}_l^{(z)}} \left(\begin{pmatrix}x_t \\ z_{x_t}\end{pmatrix}^\top \begin{pmatrix}\gamma^* \\ \omega^*\end{pmatrix} + \xi_t\right)\begin{pmatrix}x_t \\ z_{x_t}\end{pmatrix}$$

$$= \left(V_l^{(z)}\right)^+ \left(V_l^{(z)}\right)\begin{pmatrix}\gamma^* \\ \omega^*\end{pmatrix} + \left(V_l^{(z)}\right)^+ \sum_{t \in \text{Exp}_l^{(z)}} \xi_t\begin{pmatrix}x_t \\ z_{x_t}\end{pmatrix},$$

where $\left(V_l^{(z)}\right)^+$ is a generalized inverse of $V_l^{(z)}$. Since $V_l^{(z)}\left(V_l^{(z)}\right)^+ V_l^{(z)} = V_l^{(z)}$, multiplying the left and right hand side of the last equation by $V_l^{(z)}$, we find that

$$V_l^{(z)}\begin{pmatrix}\widehat{\gamma}_l^{(z)} - \gamma^* \\ \widehat{\omega}_l^{(z)} - \omega^*\end{pmatrix} = V_l^{(z)}\left(V_l^{(z)}\right)^+ \sum_{t \in \text{Exp}_l^{(z)}} \xi_t\begin{pmatrix}x_t \\ z_{x_t}\end{pmatrix}. \tag{58}$$

By Lemma 4, for all $x \in \mathcal{X}_l^{(z)}$, $\begin{pmatrix}x \\ z_x\end{pmatrix} \in \text{Range}\left(V_l^{(z)}\right)$, so

$$V_l^{(z)}\left(V_l^{(z)}\right)^+ \begin{pmatrix}x \\ z_x\end{pmatrix} = \begin{pmatrix}x \\ z_x\end{pmatrix}. \tag{59}$$

Then,

$$\begin{pmatrix}\widehat{\gamma}_l^{(z)} - \gamma^* \\ \widehat{\omega}_l^{(z)} - \omega^*\end{pmatrix}^\top \begin{pmatrix}x \\ z_x\end{pmatrix} = \begin{pmatrix}\widehat{\gamma}_l^{(z)} - \gamma^* \\ \widehat{\omega}_l^{(z)} - \omega^*\end{pmatrix}^\top V_l^{(z)}\left(V_l^{(z)}\right)^+ \begin{pmatrix}x \\ z_x\end{pmatrix}$$

$$= \sum_{t \in \text{Exp}_l^{(z)}} \begin{pmatrix}x_t \\ z_{x_t}\end{pmatrix}^\top \left(V_l^{(z)}\right)^+ V_l^{(z)}\left(V_l^{(z)}\right)^+ \begin{pmatrix}x \\ z_x\end{pmatrix}\xi_t$$

$$= \sum_{t \in \text{Exp}_l^{(z)}} \begin{pmatrix}x_t \\ z_{x_t}\end{pmatrix}^\top \left(V_l^{(z)}\right)^+ \begin{pmatrix}x \\ z_x\end{pmatrix}\xi_t,$$

where the first and third lines follow from Equation (59), and the second line follows from Equation (58). By definition of our algorithm, conditionally on $\mathcal{X}_l^{(z)}$ and $\text{Explore}_l^{(z)} = \text{True}$, the variables $(\xi_t)_{t \in \text{Exp}_l^{(z)}}$ are independent centered normal gaussian variables. Then,

$$\mathbb{P}_{|\mathcal{X}_l^{(z)}, \text{Explore}_l^{(z)}=\text{True}}\left(\left|\begin{pmatrix}\widehat{\gamma}_l^{(z)} - \gamma^* \\ \widehat{\omega}_l^{(z)} - \omega^*\end{pmatrix}^\top \begin{pmatrix}x \\ z_x\end{pmatrix}\right| \geq \sqrt{2\sum_{t \in \text{Exp}_l^{(z)}}\left(\begin{pmatrix}x_t \\ z_{x_t}\end{pmatrix}^\top \left(V_l^{(z)}\right)^+ \begin{pmatrix}x \\ z_x\end{pmatrix}\right)^2 \log\left(\frac{kl(l+1)}{\delta}\right)}\right) \leq \frac{\delta}{kl(l+1)}.$$

Expanding $\left(\left(\begin{smallmatrix} x_t \\ z_{x_t} \end{smallmatrix}\right)^\top \left(V_l^{(z)}\right)^+ \left(\begin{smallmatrix} x \\ z_x \end{smallmatrix}\right)\right)^2 = \left(\begin{smallmatrix} x \\ z_x \end{smallmatrix}\right)^\top \left(V_l^{(z)}\right)^+ \left(\begin{smallmatrix} x_t \\ z_{x_t} \end{smallmatrix}\right)\left(\begin{smallmatrix} x_t \\ z_{x_t} \end{smallmatrix}\right)^\top \left(V_l^{(z)}\right)^+ \left(\begin{smallmatrix} x \\ z_x \end{smallmatrix}\right)$, and using the definition of $V_l^{(z)}$, we find that

$$\mathbb{P}_{|\mathcal{X}_l^{(z)},\,\text{Explore}_l^{(z)}=\text{True}}\left(\left|\left(\begin{smallmatrix} \widehat{\gamma}_l^{(z)}-\gamma^* \\ \widehat{\omega}_l^{(z)}-\omega^* \end{smallmatrix}\right)^\top \left(\begin{smallmatrix} x \\ z_x \end{smallmatrix}\right)\right| \geq \sqrt{2\left(\begin{smallmatrix} x \\ z_x \end{smallmatrix}\right)^\top \left(V_l^{(z)}\right)^+ V_l^{(z)} \left(V_l^{(z)}\right)^+ \left(\begin{smallmatrix} x \\ z_x \end{smallmatrix}\right)\log\left(\frac{kl(l+1)}{\delta}\right)}\right) \leq \frac{\delta}{kl(l+1)}$$

which in turn implies (using Equation (59))

$$\mathbb{P}_{|\mathcal{X}_l^{(z)},\,\text{Explore}_l^{(z)}=\text{True}}\left(\left|\left(\begin{smallmatrix} \widehat{\gamma}_l^{(z)}-\gamma^* \\ \widehat{\omega}_l^{(z)}-\omega^* \end{smallmatrix}\right)^\top \left(\begin{smallmatrix} x \\ z_x \end{smallmatrix}\right)\right| \geq \sqrt{2\left\|\left(\begin{smallmatrix} x \\ z_x \end{smallmatrix}\right)\right\|^2_{\left(V_l^{(z)}\right)^+}\log\left(\frac{kl(l+1)}{\delta}\right)}\right) \leq \frac{\delta}{kl(l+1)}$$

Now, using Lemma 4 and the definition of μ_l^z, we see that for all $x \in \mathcal{X}_l^{(z)}$,

$$\left(\begin{smallmatrix} x \\ z_x \end{smallmatrix}\right)^\top \left(V_l^{(z)}\right)^+ \left(\begin{smallmatrix} x \\ z_x \end{smallmatrix}\right) \leq \frac{\epsilon_l^2}{2\log\left(kl(l+1)/\delta\right)}.$$

Finally, for all $x \in \mathcal{X}_l^{(z)}$,

$$\mathbb{P}_{|\mathcal{X}_l^{(z)},\,\text{Explore}_l^{(z)}=\text{True}}\left(\left|\left(\begin{smallmatrix} \widehat{\gamma}_l^{(z)}-\gamma^* \\ \widehat{\omega}_l^{(z)}-\omega^* \end{smallmatrix}\right)^\top \left(\begin{smallmatrix} x \\ z_x \end{smallmatrix}\right)\right| \geq \epsilon_l\right)$$

$$\leq \mathbb{P}_{|\mathcal{X}_l^{(z)},\,\text{Explore}_l^{(z)}=\text{True}}\left(\left|\left(\begin{smallmatrix} \widehat{\gamma}_l^{(z)}-\gamma^* \\ \widehat{\omega}_l^{(z)}-\omega^* \end{smallmatrix}\right)^\top \left(\begin{smallmatrix} x \\ z_x \end{smallmatrix}\right)\right| \geq \sqrt{2\left\|\left(\begin{smallmatrix} x \\ z_x \end{smallmatrix}\right)\right\|^2_{\left(V_l^{(z)}\right)^+}\log\left(\frac{kl(l+1)}{\delta}\right)}\right) \leq \frac{\delta}{kl(l+1)}.$$

Integrating out the conditioning on the value of $\mathcal{X}_l^{(z)}$ and $\text{Explore}_l^{(z)}$ and using a union bound yields the desire result.

C.7.16   Proof of Lemma 19

The proof is similar to that of Lemma 18. If $\text{Explore}_l^{(0)} = \text{True}$, then $\widehat{\omega}_l$ is defined as

$$\widehat{\omega}_l^{(0)} = e_{d+1}^\top \left(V_l^{(0)}\right)^+ \sum_{t\in\text{Exp}_l^{(0)}}\left(\left(\begin{smallmatrix} x_t \\ z_{x_t} \end{smallmatrix}\right)^\top \left(\begin{smallmatrix} \gamma^* \\ \omega^* \end{smallmatrix}\right) + \xi_t\right)\left(\begin{smallmatrix} x_t \\ z_{x_t} \end{smallmatrix}\right).$$

Since $\left(\begin{smallmatrix} x \\ z_x \end{smallmatrix}\right)_{x\in\mathcal{X}}$ spans $\mathbb{R}^{d+1}$, μ is finite and $e_{d+1} \in \text{Range}\left(V(\hat{\mu}_l)\right)$. Then, according to Lemma 3, for every round l, we have $e_{d+1} \in \text{Range}\left(V_l^{(0)}\right)$, so $V_l^{(0)}\left(V_l^{(0)}\right)^+ e_{d+1} = e_{d+1}$. This implies that

$$\widehat{\omega}_l^{(0)} - \omega^* = \sum_{t\in\text{Exp}_l^{(0)}} e_{d+1}^\top \left(V_l^{(0)}\right)^+ \left(\begin{smallmatrix} x_t \\ z_{x_t} \end{smallmatrix}\right)\xi_t.$$

By definition of our algorithm, conditionally on $\text{Explore}_l^{(0)} = \text{True}$, the variables $(\xi_t)_{t\in\text{Exp}_l^{(0)}}$ are independent centered normal gaussian variables. Then,

$$\mathbb{P}_{|\text{Explore}_l^{(0)}=\text{True}}\left(\left|\widehat{\omega}_l^{(0)} - \omega^*\right| \geq \sqrt{2\sum_{t\in\text{Exp}_l^{(z)}}\left(e_{d+1}^\top \left(V_l^{(0)}\right)^+ \left(\begin{smallmatrix} x_t \\ z_{x_t} \end{smallmatrix}\right)\right)^2\log\left(\frac{l(l+1)}{\delta}\right)}\right) \leq \frac{\delta}{l(l+1)}.$$

Using again $V_l^{(0)}\left(V_l^{(0)}\right)^+ e_{d+1} = e_{d+1}$ and the definition of $V_l^{(0)}$, we find that

$$\mathbb{P}_{|\text{Explore}_l^{(0)}=\text{True}}\left(\left|\widehat{\omega}_l^{(0)} - \omega^*\right| \geq \sqrt{2e_{d+1}^\top \left(V_l^{(0)}\right)^+ e_{d+1}\log\left(\frac{l(l+1)}{\delta}\right)}\right) \leq \frac{\delta}{l(l+1)}. \qquad (60)$$

Now, Lemma 3 and the definition of $\mu_l^{(0)}$ imply that

$$e_{d+1}^\top \left(V_l^{(0)}\right)^+ e_{d+1} \le \frac{\epsilon_l^2}{2\log\left(l(l+1)/\delta\right)}.$$

Finally, Equation (60) implies that

$$\mathbb{P}_{|\text{Explore}_l^{(0)}=\text{True}}\left(\left|\widehat{\omega}_l^{(0)} - \omega^*\right| \ge \epsilon_l\right) \le \frac{\delta}{l(l+1)}.$$

Using a union bound over the phases $\text{Exp}_l^{(0)}$ yields the result.

C.7.17   Proof of Lemma 20

To prove Lemma 20, we begin by showing that it is enough to prove that for $l \ge 1$,

$$\mathcal{F}_l \supset \quad \left\{\exists x^* \in \operatorname*{argmax}_{x \in \mathcal{X}} x^\top \gamma^* : \text{Explore}_l^{(z_{x^*})} = \text{True and } x^* \notin \mathcal{X}_{l+1}^{(z_{x^*})}\right\} \tag{61}$$

$$\cup \quad \left\{\bigcap_{l' \le l} \overline{\left\{\exists x^* \in \operatorname*{argmax}_{x \in \mathcal{X}} x^\top \gamma^* : \text{Explore}_{l'}^{(z_{x^*})} = \text{True and } x^* \notin \mathcal{X}_{l'+1}^{(z_{x^*})}\right\}}\right.$$

$$\left.\bigcap \left\{\text{Explore}_l^{(0)} = \text{True and } \forall x^* \in \operatorname*{argmax}_{x \in \mathcal{X}} x^\top \gamma^*, \widehat{z^*}_{l+1} = -z_{x^*}\right\}\right\}.$$

Indeed, denoting $\mathcal{F}_l^{(1)} = \left\{\exists x^* \in \operatorname{argmax}_{x \in \mathcal{X}} x^\top \gamma^* : \text{Explore}_l^{(z_{x^*})} = \text{True and } x^* \notin \mathcal{X}_{l+1}^{(z_{x^*})}\right\}$ and $\mathcal{F}_l^{(2)} = \left\{\text{Explore}_l^{(0)} = \text{True and } \forall x^* \in \operatorname{argmax}_{x \in \mathcal{X}} x^\top \gamma^*, \widehat{z^*}_{l+1} = -z_{x^*}\right\}$, we see that Equation (61) would then be rewritten as

$$\mathcal{F}_l \quad \supset \quad \mathcal{F}_l^{(1)} \bigcup \left\{\bigcap_{l' \le l} \overline{\mathcal{F}_{l'}^{(1)}} \bigcap \mathcal{F}_l^{(2)}\right\}$$

which implies

$$\bigcup_{l \ge 1} \mathcal{F}_l \supset \bigcup_{l \ge 1} \left\{\mathcal{F}_l^{(1)} \bigcup \left\{\left\{\bigcap_{l' \le l} \overline{\mathcal{F}_{l'}^{(1)}} \bigcap \mathcal{F}_l^{(2)}\right\} \bigcup_{l' \le l} \mathcal{F}_{l'}^{(1)}\right\}\right\} \quad \supset \quad \bigcup_{l \ge 1} \left\{\mathcal{F}_l^{(1)} \cup \mathcal{F}_l^{(2)}\right\}.$$

Then, Equation (61) would imply that

$$\overline{\mathcal{F}} = \overline{\bigcup_{l \ge 1} \mathcal{F}_l} \subset \overline{\bigcup_{l \ge 1} \left\{\mathcal{F}_l^{(1)} \bigcup \mathcal{F}_l^{(2)}\right\}} = \bigcap_{l \ge 1} \left\{\overline{\mathcal{F}_l^{(1)}} \bigcap \overline{\mathcal{F}_l^{(2)}}\right\},$$

thus proving Lemma 20. To prove Equation (61), we show that both $\mathcal{F}_l^{(1)}$ and $\bigcap_{l' \le l} \overline{\mathcal{F}_{l'}^{(1)}} \bigcap \mathcal{F}_l^{(2)}$ imply $\mathcal{F}_l$.

If $\mathcal{F}_l^{(1)}$ is true: then $\exists x^* \in \operatorname{argmax}_{x \in \mathcal{X}} : \text{Explore}_l^{(z_{x^*})} = \text{True and } x^* \notin \mathcal{X}_{l+1}^{(z_{x^*})}$.
Without loss of generality, assume that $l > 1$ is the smallest integer such that $\text{Explore}_l^{(z_{x^*})} = \text{True}$ and $x^* \notin \mathcal{X}_{l+1}^{(z_{x^*})}$. Then, necessarily $x^* \in \mathcal{X}_l^{(z_{x^*})}$ (because either $l = 1$, or $\text{Explore}_{l-1}^{(z_{x^*})} = \text{True}$). Now, because $x^* \in \mathcal{X}_l^{(z_{x^*})} \setminus \mathcal{X}_{l+1}^{(z_{x^*})}$, there exists $x \in \mathcal{X}_l^{(z_{x^*})}$ such that

$$(x - x^*)^\top \widehat{\gamma}_l^{(z_{x^*})} \ge 3\epsilon_l$$

and in particular

$$x^\top \widehat{\gamma}_l^{(z_{x^*})} - \epsilon_l > (x^*)^\top \widehat{\gamma}_l^{(z_{x^*})} + \epsilon_l.$$

Recall that by definition of x^*, $(\gamma^*)^\top (x^* - x) \ge 0$. This in turn implies that

$$\begin{pmatrix} x \\ z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(z_{x^*})} - \gamma^* \\ \widehat{\omega}_l^{(z_{x^*})} - \omega^* \end{pmatrix} - \epsilon_l > \begin{pmatrix} x^* \\ z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(z_{x^*})} - \gamma^* \\ \widehat{\omega}_l^{(z_{x^*})} - \omega^* \end{pmatrix} + \epsilon_l.$$

The last equation implies that either $\begin{pmatrix} x \\ z_x \end{pmatrix}^\top \begin{pmatrix} \gamma_l^{(z)} - \gamma^* \\ \widehat{\omega}_l^{(z)} - \omega^* \end{pmatrix} > \epsilon_l$ or $\begin{pmatrix} x^* \\ z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \gamma_l^{(z)} - \gamma^* \\ \widehat{\omega}_l^{(z)} - \omega^* \end{pmatrix} < -\epsilon_l$, which in turn implies $\mathcal{F}_l$.

If $\bigcap_{l' \leq l} \overline{\mathcal{F}_{l'}^{(1)}} \bigcap \mathcal{F}_l^{(2)}$ is true: then $\text{Explore}_l^{(0)} = \text{True}$ and $\forall x^* \in \text{argmax}_{x \in \mathcal{X}} x^\top \gamma^*, \widehat{z_{l+1}^*} = -z_{x^*}$.
Moreover, for all $l' \leq l$, $\text{Explore}_{l'}^{(z_{x^*})} = \text{False}$ or $x^* \in \mathcal{X}_{l'+1}^{(z_{x^*})}$.
Note that this case can only hold if all optimal actions x^* belong to the same group z_{x^*}. Without loss of generality, assume that $l > 1$ is the smallest integer such that $\text{Explore}_l^{(0)} = \text{True}$ and $\widehat{z_{l+1}^*} = -z_{x^*}$, and for all $l' \leq l$, $\text{Explore}_{l'}^{(z_{x^*})} = \text{False}$ or $x^* \in \mathcal{X}_{l'+1}^{(z_{x^*})}$. Note that because $\text{Explore}_l^{(0)} = \text{True}$, necessarily $\text{Explore}_{l'}^{(z_{x^*})} = \text{True}$ for all $l' \leq l$, and in particular $x^* \in \mathcal{X}_{l+1}^{(z_{x^*})}$.

Then, there exists $x \in \mathcal{X}_{l+1}^{(-z_{x^*})}$ such that

$$\begin{pmatrix} x \\ -z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(-z_{x^*})} \\ \widehat{\omega}_l^{(-z_{x^*})} \end{pmatrix} - \begin{pmatrix} x^* \\ z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(z_{x^*})} \\ \widehat{\omega}_l^{(z_{x^*})} \end{pmatrix} + 2 z_{x^*} \widehat{\omega}_l^{(0)} \geq 4\epsilon_l.$$

Recall that all optimal actions x^* are in the same group z_{x^*}, so $(\gamma^*)^\top (x^* - x) > 0$. This in turn implies that

$$\begin{pmatrix} x \\ -z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(-z_{x^*})} - \gamma^* \\ \widehat{\omega}_l^{(-z_{x^*})} - \omega^* \end{pmatrix} - \begin{pmatrix} x^* \\ z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(z_{x^*})} - \gamma^* \\ \widehat{\omega}_l^{(z_{x^*})} - \omega^* \end{pmatrix} + 2 z_{x^*} (\widehat{\omega}_l^{(0)} - \omega^*) \geq 4\epsilon_l.$$

The last equation implies that either $\begin{pmatrix} x \\ -z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(-z_{x^*})} - \gamma^* \\ \widehat{\omega}_l^{(-z_{x^*})} - \omega^* \end{pmatrix} \geq \epsilon_l$, or $\begin{pmatrix} x^* \\ z_{x^*} \end{pmatrix}^\top \begin{pmatrix} \widehat{\gamma}_l^{(z_{x^*})} - \gamma^* \\ \widehat{\omega}_l^{(z_{x^*})} - \omega^* \end{pmatrix} \leq -\epsilon_l$, or $z_{x^*} (\widehat{\omega}_l^{(0)} - \omega^*) \geq \epsilon_l$, which in turn implies $\mathcal{F}_l$.

C.7.18 Proof of Lemma 21

The first claim holds for $l = 1$. For $l \geq 1$, for any $x \in \mathcal{X}_{l+1}^{(-1)} \cup \mathcal{X}_{l+1}^{(1)}$, we have $\widehat{\Delta}_x^{l+1} \leq \Delta_x + 8\epsilon_l$ on $\overline{\mathcal{F}}$ according to the definition of $\widehat{\Delta}^{l+1}$ and $\mathcal{F}$. The first claim then follows.

For the second claim, Lemma 13 gives that, on $\overline{\mathcal{F}}$, $\Delta_x < 21\epsilon_l$ for any $x \in \mathcal{X}_{l+1}^{(-1)} \cup \mathcal{X}_{l+1}^{(1)}$. So $\Delta_x \geq 21\epsilon_l$ implies $x \notin \mathcal{X}_{l+1}^{(-1)} \cup \mathcal{X}_{l+1}^{(1)}$ and hence $l \geq \ell_x$ on $\overline{\mathcal{F}}$.

For the third claim, we notice that

$$\max_{x' \in \mathcal{X}_{\ell_x}^{(z_x)}} (a_{x'} - a_x)^\top \widehat{\theta}_{\ell_x}^{(z_x)} > 3\epsilon_{\ell_x},$$

since $x \notin \mathcal{X}_{\ell_x + 1}$. Since the left-hand side is smaller than $\Delta_x + 2\epsilon_{\ell_x}$ on $\overline{\mathcal{F}}$, we get $\Delta_x > \epsilon_{\ell_x}$.

D   Extension to M groups

Model   We extend the biased linear bandit to Z groups, denoted $\mathcal{Z} = \{1, ..., Z\}$. The evaluations are given by

$$y_t = x_t^\top \gamma + Z_{x_t}^\top \omega + \xi_t,$$

where Z_x is the z_x-th vector of the canonical basis in $\mathbb{R}^Z$, and $\omega = \{\omega_1, ..., \omega_Z\} \in \mathbb{R}^Z$ is the vector of biases. Note that for the model to be identifiable, we must assume it does not contain an intercept. For $x \in \mathcal{X}$, we denote $a_x = \begin{pmatrix} x \\ Z_x \end{pmatrix}$. To ensure identifiability of the model, we further assume that the set $\mathcal{A} = \{a_x : x \in \mathcal{X}\}$ spans $\mathbb{R}^{d+Z}$.

Estimation of the biased evaluations   Adapting the G-EXP-ELIM routine to the multiple group framework is rather straightforward. Note that this routine can be used as is to eliminate within-group sub-optimal actions. The actions of each group span a sub-space of dimension $d+1$, so the G-optimal measure is still supported by $O(d^2)$ points. Moreover, the variance corresponding to this G-optimal design is still $d + 1$.

Estimation of the bias By contrast, the bias elimination routine must be modified in order to handle Z groups. At each phase l, we denote by $\mathcal{Z}_l$ the set of groups that have not been eliminated yet. If more than one group remain in $\mathcal{Z}_l$, we compute the difference $\omega_1 - \omega_z$ for all group z remaining in $\mathcal{Z}_l$ with precision $\epsilon_l/2$ using a modified Δ-EXP-ELIM routine, which we call Δ-MULT-EXP-ELIM, described in 5. This routine samples action according to the distribution μ_z, where for any groups $z \neq 1$, we defined μ_z as the solution of the problem

$$\underset{\mu \in \mathcal{M}_{\mathcal{X}}^{e_{d+1}-e_{d+z}}}{\text{minimize}} \sum_x \mu(x)\Delta_x \quad \text{such that} \quad (e_{d+1} - e_{d+z})^\top V(\mu)^+ (e_{d+1} - e_{d+z}) \leq 1 \tag{62}$$

We also define $\widetilde{\kappa}_z(\Delta)$ as the corresponding regret :

$$\widetilde{\kappa}_z(\Delta) \quad = \quad \sum_x \mu_z(x)\Delta_x.$$

Note that the support of the distribution μ_z is at most of size $d + Z$. This two-by-two comparison allows us to compute, for each $z, z' \in \mathcal{Z}_l$, the difference of bias $\omega_z - \omega_{z'} = \omega_1 - \omega_{z'} - (\omega_1 - \omega_z)$ with precision level ϵ_l. Then, we can use these bias estimates to eliminate groups that are sub-optimal by a gap larger than $4\epsilon_l$. Again, we rely on estimates of the biases and of the biased evaluations obtained during the previous round to update the estimate of the gap vector $\widehat{\Delta}^{l+1}$.

Algorithm 5 Δ-MULT-EXP-ELIM $(\mathcal{X}, \mathcal{Z}, (\mathcal{X}^{(z)}, \widehat{\theta}^{(z)})_{z \in \mathcal{Z}}, \widehat{\Delta}, n, \epsilon)$

1: **for** $z \in \mathcal{Z}, z \neq 1$ **do**
2:      Compute $\widehat{\Delta}$-optimal design $\hat{\mu}_z$ solution of (62) on $\mathcal{X}$, with $|\operatorname{supp}(\hat{\mu}_z)| \leq d + Z$
3:      Sample $\lceil n\hat{\mu}_z(x) \rceil$ times each action a_x for $x \in \mathcal{X}$
4:      Compute $\widehat{\omega}_1 - \widehat{\omega}_z = (e_{d+1} - e_{d+z})^\top \widehat{\theta}$, where $\widehat{\theta}$ is the ordinary least square estimator
5: **for** $z \in \mathcal{Z}$ and $x \in \mathcal{X}^{(z)}$ **do** $\widehat{m}_x \leftarrow a_x^\top \widehat{\theta}^{(z)} + (\widehat{\omega}_1 - \widehat{\omega}_z)$
6: **for** $z \in \mathcal{Z}$ and $x \in \mathcal{X}^{(z)}$ **do** $\widehat{\Delta}_x \leftarrow 2 \wedge \left(\max_{z' \in \mathcal{Z}, x' \in \mathcal{X}^{(z')}} \widehat{m}_{x'} - \widehat{m}_x + 4\epsilon \right)$
7: **for** $z \in \mathcal{Z}$ **do**
8:      **if** $\max_{z' \in \mathcal{Z}} \max_{x \in \mathcal{X}^{(z')}} a_x^\top \widehat{\theta}^{(z')} + (\widehat{\omega}_1 - \widehat{\omega}_{z'}) \geq \max_{x \in \mathcal{X}^{(z)}} a_x^\top \widehat{\theta}^{(z)} + (\widehat{\omega}_1 - \widehat{\omega}_z) + 4\epsilon$ **then** $\mathcal{Z} \leftarrow \mathcal{Z} \setminus \{z\}$
9: **return** $\mathcal{Z}$ and $\widehat{\Delta}$

Stopping criterion We denote by $\widetilde{\kappa}_{\mathcal{Z}_l}(\widehat{\Delta}^l) = \sum_{z \in \mathcal{Z}_l, z \neq 1} \widetilde{\kappa}_z(\widehat{\Delta}^l)$ the regret for estimating the biases at phase l. If $\epsilon_l \leq \left(\widetilde{\kappa}_{\mathcal{Z}_l}(\widehat{\Delta}^l) \log(T)/T \right)^{1/3}$, bias estimation becomes too costly, so we sample the empirical best action for the remaining time. The FAIR PHASED ELIMINATION FOR MULTIPLE GROUPS algorithm is presented in 6.

D.1 Worst case regret

Before analyzing the worst case regret of Algorithm 6, we introduce a new quantity, $\widetilde{\kappa}_*$, defined as

$$\widetilde{\kappa}_* = \sum_{z \in \mathcal{Z}, z \neq 1} \min_{\pi \in \mathcal{P}_{e_{d+1}-e_{d+z}}^{\mathcal{X}}} (e_{d+1} - e_{d+z})^\top (V(\pi))^+ (e_{d+1} - e_{d+z}).$$

Note that for all $z \in \mathcal{Z}, z \neq 1$, and $l \geq 1$, we have $\widetilde{\kappa}_{\mathcal{Z}_l}(\widehat{\Delta}^l) \leq 2\widetilde{\kappa}_*$.

Claim 1. *For the choice $\delta = T^{-1}$, there exists an absolute constant $C > 0$ and a constant $T_{\widetilde{\kappa}_*,k,Z,d,k}$ depending on $\widetilde{\kappa}_*, k, Z, d,$ and k such that the following bound on the regret of the* FAIR PHASED ELIMINATION FOR MULTIPLE GROUPS *algorithm 6 holds*

$$R_T \leq CZ \left(\widetilde{\kappa}_* \log(T) \right)^{1/3} T^{2/3} \quad \text{for} \quad T \geq T_{\widetilde{\kappa}_*,k,Z,d,k}.$$

Sketch of Proof. We sketch here a proof of Claim 1, highlighting the main differences with the two-groups setting. We begin by introducing some notations.

Algorithm 6 FAIR PHASED ELIMINATION FOR MULTIPLE GROUPS

1: **input:** $\delta, T, \mathcal{X}, k = |\mathcal{X}|, \epsilon_l = 2^{2-l}$ for $l \geq 1$
2: **initialize:** $\widehat{\Delta}^1 \leftarrow (2, ..., 2), l \leftarrow 0, \mathcal{Z}_1 = \mathcal{Z}$
3: **for** $z \in \mathcal{Z}_1$ **do** $\mathcal{X}_1^{(z)} \leftarrow \{x : z_x = z\}$
4: **while** the budget is not spent **do** $l \leftarrow l + 1$
5:     **for** $z \in \mathcal{Z}_l$ **do**
6:         $\left(\widehat{\theta}^{(z)}, \mathcal{X}_{l+1}^{(z)}\right) \leftarrow$ G-EXP-ELIM $\left(\mathcal{X}_l^{(z)}, \frac{2(d+1)}{\epsilon_l^2} \log\left(\frac{kl(l+1)}{\delta}\right), \epsilon_l\right)$
7:     **if** $|\mathcal{Z}_l| > 1$ **then**
8:         Compute $\widetilde{\kappa}_{\mathcal{Z}_l}(\widehat{\Delta}^l) = \sum\limits_{z \in \mathcal{Z}_l, z \neq 1} \widetilde{\kappa}_z(\widehat{\Delta}^l)$.
9:         **if** $\epsilon_l \leq \left(\widetilde{\kappa}_{\mathcal{Z}_l}(\widehat{\Delta}^l) \log(T)/T\right)^{1/3}$ **then**   ▷ Stop bias estimation
10:             Sample best action in $\cup_{z \in \mathcal{Z}_l} \mathcal{X}_{l+1}^{(z)}$ for the remaining time
11:     **else**
12:         $\left(\mathcal{Z}_{l+1}, \widehat{\Delta}^{l+1}\right) \leftarrow \Delta$-MULT-EXP-ELIM $\left(\mathcal{X}, \mathcal{Z}_l, \left(\mathcal{X}_{l+1}^{(z)}, \widehat{\theta}_l^{(z)}\right)_{z \in \mathcal{Z}_l}, \widehat{\Delta}^l, \frac{8}{\epsilon_l^2} \log\left(\frac{Zl(l+1)}{\delta}\right), \epsilon_l\right)$

Notations We denote by L_T the largest integer l such that $\epsilon_l \geq \left(2\widetilde{\kappa}_*^{1/3} \log(T)/T\right)^{1/3}$. For $z \in \mathcal{Z}$, we denote by L^Δ the last phase where $\widehat{\Delta}^l$-optimal Exploration and Elimination is performed. We denote by Exp-G$_l^{(z)}$ the time indices where G-exploration is performed on $\mathcal{X}_l^{(z)}$ and by Exp-D$_l^{(z)}$ the time indices where Δ-exploration is performed at phase l for estimating the difference $\omega_1 - \omega_z$. We also denote by Recovery the time indices subsequent to the stopping criterion, this set being empty when the stopping criterion is not activated.

We define a "good" event $\overline{\mathcal{F}}$ such that for all $z, z' \in \mathcal{Z}$ and all $x \in \mathcal{X}_1^{(z)}$, the errors $\left| a_x^\top \left(\theta^* - \widehat{\theta}_l^{(z)}\right) \right|$ and $|(\omega_z^* - \omega_{z'}^*) - ((\widehat{\omega}_l)_z - (\widehat{\omega}_l)_{z'})|$ are smaller than ϵ_l for all l such that these quantities are defined. In the following, we use c, c' to denote positive absolute constants, which may vary from line to line. With these notations, we decompose the regret as follows :

$$
R_T \leq 2T\mathbb{P}(\mathcal{F}) + \mathbb{E}_{|\overline{\mathcal{F}}}\left[\underbrace{\sum_{l \leq L_T} \sum_{z \in \mathcal{Z}_l} \sum_{t \in \text{Exp-G}_l^{(z)}} (x^* - x_t)^\top \gamma^*}_{R_T^G}\right] + \mathbb{E}_{|\overline{\mathcal{F}}}\left[\underbrace{\sum_{l \leq L^\Delta} \sum_{z \in \mathcal{Z}_l, z \neq 1} \sum_{t \in \text{Exp-D}_l^{(z)}} (x^* - x_t)^\top \gamma^*}_{R_T^\Delta}\right]
$$

$$
+ \mathbb{E}_{|\overline{\mathcal{F}}}\left[\underbrace{\sum_{l \geq L_T+1} \sum_{z \in \mathcal{Z}_l} \sum_{t \in \text{Exp-G}_l^{(z)}} (x^* - x_t)^\top \gamma^* + \sum_{t \in \text{Recovery}} (x^* - x_t)^\top \gamma^*}_{R_T^{Rec}}\right].
$$

Bound on $T\mathbb{P}(\mathcal{F})$. Using arguments based on concentration of Gaussian variables, we can show that $\mathbb{P}(\mathcal{F}) \leq 2T^{-1}$.

Bound on R_T^G. The analysis is similar to the two-groups setting. We can show that on $\overline{\mathcal{F}}$, only actions with gaps smaller than $c\epsilon_l$ remain in the sets $\mathcal{X}_l^{(z)}$ for $z \in \mathcal{Z}_l$. The length of each G-optimal Exploration and Elimination phase for one group is of the order $(d+1)\log(klT)/\epsilon_l^2$, so the regret corresponding to phase l is of the order $Z(d+1)\log(klT)/\epsilon_l$. Summing over the different phases, we find that

$$
R_T^G \leq c(d+1)Z \log(kL_TT)/\epsilon_{L_T}. \tag{63}
$$

Using the definition of L_T, we find that $R_T^G \leq c(d+1)Z \log(kL_TT)\widetilde{\kappa}_*^{-1/3} \log(T)^{-1/3}T^{1/3}$.

Bound on R_T^{Rec}. On the one hand, the actions selected during the Phases Exp-G$_l^{(z)}$ for $l \geq L_T + 1$ are sub-optimal by a gap at most $c\epsilon_{L_T}$ on the event $\overline{\mathcal{F}}$. On the other hand, if the algorithm enters the Recovery phase at a phase l, then

$$\epsilon_l \leq \widetilde{\kappa}_{\mathcal{Z}_{L^\Delta}}(\widehat{\Delta}^{L^\Delta})^{1/3} T^{-1/3} \log(T)^{1/3} \leq 2\widetilde{\kappa}_*^{1/3} T^{-1/3} \log(T)^{1/3},$$

so we must have $l = L^\Delta + 1 \geq L_T + 1$. Therefore, all actions selected during the Recovery phase are sub-optimal by a gap at most $c\epsilon_{L_T}$. Then, R_T^{Rec} can be bounded as $R_T^{Rec} \leq c\epsilon_{L_T} T$. This implies in particular that $R_T^{Rec} \leq c'\widetilde{\kappa}_*^{1/3} \log(T)^{1/3} T^{2/3}$.

Bound on R_T^Δ. To bound R_T^Δ, we introduce further notations. Let us denote by $l_1, ..., l_R$ the phases at which at least one group is eliminated, by $S_1 i$ the sets of groups remaining at the beginning of phase l_i, and by S_{R+1} the set of groups that are never eliminated. We also write $l_{R+1} = L^\Delta$. We abuse notations and denote Exp-D$_l^{(S)} = \cup_{z \in S}$ Exp-D$_l^{(z)}$. Then, we see that

$$R_T^\Delta \leq \sum_{i \leq R+1} \sum_{l \leq l_i} \sum_{t \in \text{Exp-D}_l^{(S_i)}} (x^* - x_t)^\top \gamma^*.$$

The rest of the proof is similar to that in the two-communities setting. We show that on $\mathcal{F}$, $\widehat{\Delta}^l \geq \Delta$ for all $l \geq 1$. Then, our choice of design $\widehat{\mu}_{z_l,z}$ at phase l ensures that for $i \leq R + 1$, on $\overline{\mathcal{F}}$,

$$\sum_{t \in \text{Exp-D}_l^{(S_i)}} (x^* - x_t)^\top \gamma^* \leq c \sum_{z \in S_i} \left(\frac{\log(Zl(l+1)T)}{\epsilon_l^2} \widetilde{\kappa}_z(\widehat{\Delta}^l) + d + 1 \right)$$

for some constant $c > 0$. Using arguments similar to the two-groups setting, we can sum over the different phases $l \leq l_i$, and find that

$$\sum_{l \leq l_i} \sum_{t \in \text{Exp-D}_l^{(S_i)}} (x^* - x_t)^\top \gamma^* \leq c\widetilde{\kappa}_{S_i}(\widehat{\Delta}^{l_i}) \log(Zl_iT)/\epsilon_{l_i}^2. \tag{64}$$

By definition of S_i we have that $\widetilde{\kappa}_{\mathcal{Z}_{l_i}}(\widehat{\Delta}^{l_i}) = \widetilde{\kappa}_{S_i}(\widehat{\Delta}^{l_i})$. Now, the algorithm does not enter the Recovery phase before phase $l_i + 1$, so we must have
$\epsilon_{l_i}^{-2} \leq T^{2/3} \log(T)^{-2/3} \widetilde{\kappa}_{\mathcal{Z}_{l_i}}(\widehat{\Delta}^{l_i})^{-2/3}$. This implies that

$$\sum_{l \leq l_i} \sum_{t \in \text{Exp-D}_l^{(S_i)}} (x^* - x_t)^\top \gamma^* \leq c\widetilde{\kappa}_{\mathcal{Z}_{l_i}}(\widehat{\Delta}^{l_i})^{1/3} \left(\log(T)^{1/3} + \log(Z) \right) T^{2/3}.$$

We use that $\widetilde{\kappa}_{\mathcal{Z}_{l_i}}(\widehat{\Delta}^{l_i}) \leq \widetilde{\kappa}_*$ and sum over $i \leq R + 1 < Z$, and we find that
$R_T^\Delta \leq CZ\widetilde{\kappa}_*^{1/3} \log(T)^{1/3} T^{2/3}$ for T large enough.

When $T \geq T_{\widetilde{\kappa}_*,k,Z,d,k}$ for some $T_{\widetilde{\kappa}_*,k,Z,d,k}$ large enough, we find that $\mathbb{R}_T \leq c'Z\widetilde{\kappa}_*^{1/3} \log(T)^{1/3} T^{2/3}$. $\qquad\square$

D.2 Gap-dependent regret

Before stating the bound on the gap-dependent regret, we introduce further notations. For $z \in \mathcal{Z}$, we denote $\Delta_{\neq,z} = \min_{x:z_x=z} \Delta_x$, $\Delta_{\neq} = \min_{x:z\neq z^*} \Delta_{\neq,z}$, $\Delta_{\min} = \min_{x \in \mathcal{X} \setminus x^*} \Delta_x$, and $\varepsilon_T = (\widetilde{\kappa}_* \log(T)/T)^{1/3}$.
Then, we claim that the following gap-dependent regret bound on the regret of Algorithm 5 holds.

Claim 2. *Assume that $x^* \in \text{argmax}_{x \in \mathcal{X}} x^\top \gamma^*$ is unique. Then, there exists an absolute constant $C > 0$ and a constant $T_{\widetilde{\kappa}_*,k,Z,d,k,\Delta_{\neq},\Delta_{\min}}$ depending on $\widetilde{\kappa}_*, k, Z, d, k, \Delta_{\min}$, and $(\Delta_{\neq,z})_{z \neq z^*}$ such that the following bound on the regret of the* FAIR PHASED ELIMINATION FOR MULTIPLE GROUPS *algorithm 6 holds for $T \geq T_{\widetilde{\kappa}_*,k,Z,d,k,\Delta_{\neq},\Delta_{\min}}$*

$$R_T \leq C \left(\frac{d}{\Delta_{\min}} \vee \sum_{z \neq z^*, z \neq 1} \frac{\kappa_z (\Delta \vee \Delta_{\neq,z} \vee \varepsilon_T)}{(\Delta_{\neq,z})^2} + \frac{\kappa_{z^*} (\Delta \vee \Delta_{\neq} \vee \varepsilon_T)}{(\Delta_{\neq})^2} \right) \log(T).$$

Sketch of Proof. We sketch here a proof of Claim 2. We begin by introducing some notations.

Notations We define a "good" event $\overline{\mathcal{F}}$ such that for all $z, z' \in \mathcal{Z}$ and all $x \in \mathcal{X}_1^{(z)}$, the errors $\left| a_x^\top \left(\theta^* - \widehat{\theta}_l^{(z)} \right) \right|$ and $|(\omega_z^* - \omega_{z'}^*) - ((\widehat{\omega}_l)_z - (\widehat{\omega}_l)_{z'})|$ are smaller than ϵ_l for all l such that these quantities are defined. For each group $z \in \mathcal{Z}$, we denote by Exp-G$_l^{(z)}$ the time indices where G-exploration is performed on $\mathcal{X}_l^{(z)}$. For $z \in \mathcal{Z}$, $z \neq 1$, we denote by Exp-D$_l^{(z)}$ the time indices where Δ-exploration is performed at phase l to estimate the difference $\omega_1 - \omega_z$, and by $L^{(z)}$ the last phase l such that $z \in \mathcal{Z}_l$ and bias exploration is performed at this phase. We denote by L^Δ the last phase l where bias estimation is performed. Moreover, we denote by S the sets of groups eliminated before the stopping criterion is activated, and write $\overline{S} = \mathcal{Z} \setminus S$. We abuse notations and denote Exp-D$_l^{(S)} = \cup_{z \in S}$ Exp-D$_l^{(z)}$. We also denote by Recovery the time indices subsequent to the stopping criterion, this set being empty when the stopping criterion is not activated. In the following, we use c, c' to denote positive absolute constants, which may vary from line to line.

Fact 1 Let $l_{\Delta_{\min}}$ be the largest integer such that $\epsilon_{l_{\Delta_{\min}}} \geq C\Delta_{\min}$ for some well-chosen absolute constant $C > 0$. Similarly to the two-groups setting, we can show that on the good event $\overline{\mathcal{F}}$, no more than $l_{\Delta_{\min}}$ G-optimal Exploration and Elimination phases are needed to find the best action. For all phases $l \geq l_{\Delta_{\min}}$, the algorithm always chooses x^*, and suffers no regret.

Fact 2 Similarly to the two-groups setting, we can show that on the good event $\overline{\mathcal{F}}$, for each phase l, $\widehat{\Delta}^l \leq c (\Delta \vee \epsilon_l)$ for some constant c. Moreover, for all $l \leq L^\Delta$, all groups $z \neq 1$, and all $\tau > 0$, $\widetilde{\kappa}_z(\widehat{\Delta}^l) \leq c\widetilde{\kappa}_z(\Delta \vee \epsilon_l) \leq c(1 + \epsilon_l \tau^{-1})\widetilde{\kappa}_z(\Delta \vee \tau)$.

Fact 3 For $z \in \mathcal{Z} \setminus \{z^*\}$, let $l_{\Delta_{\neq,z}}$ be the largest integer such that $\epsilon_{l_{\Delta_{\neq,z}}} \geq C\Delta_{\neq,z}$ for some well-chosen absolute constant $C > 0$. On the good event $\overline{\mathcal{F}}$, if $\widehat{\Delta}^l$-optimal Exploration and Elimination is performed at phase $l \geq l_{\Delta_{\neq,z}}$, and $z \in \mathcal{Z}_l$, then the algorithm eliminates z at this phase. This implies that $L^{(z)} \leq l_{\Delta_{\neq,z}}$, and that $L^\Delta \leq l_{\Delta_{\neq}}$.

Fact 4 We denote by L_T the largest integer l such that $\epsilon_l \geq (2\widetilde{\kappa}_* \log(T)/T)^{1/3}$. Since $2\widetilde{\kappa}_* \geq \widetilde{\kappa}(\widehat{\Delta}^l)$ for all $l \geq 1$ and all $z \in \mathcal{Z}$, we see that if the algorithm enters the Recovery phase, we must have $L_T \leq L^\Delta$, and $\epsilon_{L^\Delta} \leq \epsilon_{L_T} \approx \varepsilon_T$.

Using **Fact 1**, we find that the regret can be written as

$$
R_T \leq 2T\mathbb{P}(\mathcal{F}) + \mathbb{E}_{|\overline{\mathcal{F}}}\left[\underbrace{\sum_{l \leq l_{\Delta_{\min}}} \sum_{z \in \mathcal{Z}_l} \sum_{t \in \text{Exp-G}_l^{(z)}} (x^* - x_t)^\top \gamma^*}_{R_T^G} \right] + \mathbb{E}_{|\overline{\mathcal{F}}}\left[\underbrace{\sum_{z \in S} \sum_{l \leq L^{(z)}} \sum_{t \in \text{Exp-D}_l^{(z)}} (x^* - x_t)^\top \gamma^*}_{R_T^{\Delta,S}} \right]
$$

$$
+ \mathbb{E}_{|\overline{\mathcal{F}}}\left[\underbrace{\sum_{l \leq L^\Delta} \sum_{t \in \text{Exp-D}_l^{(\overline{S})}} (x^* - x_t)^\top \gamma^*}_{R_T^{\Delta,\overline{S}}} \right] + \mathbb{E}_{|\overline{\mathcal{F}}}\left[\underbrace{\sum_{t \in \text{Recovery}} (x^* - x_t)^\top \gamma^*}_{R_T^{Rec}} \right].
$$

Bound on R_T^G. We rely on arguments similar to those used in Equation (63) to show that $R_T^G \leq c(d+1) \log(kl_{\Delta_{\min}}T)\epsilon_{l_{\Delta_{\min}}}^{-1}$. Since $\epsilon_{l_{\Delta_{\min}}} \geq C\Delta_{\min}$, this implies that

$$
R_T^G \leq \frac{c(d+1) \log(kl_{\Delta_{\min}}T)}{\Delta_{\min}} \leq \frac{c'd \log(T)}{\Delta_{\min}}
$$

if $T \geq k$.

Bound on $R_T^{\Delta,S}$. Using arguments similar to the two-groups settings, we can show that for all $z \neq 1$

$$
\sum_{l \leq L^{(z)}} \sum_{t \in \text{Exp-D}_l^{(z)}} (x^* - x_t)^\top \gamma^* \leq c\widetilde{\kappa}_z(\widehat{\Delta}^{L^{(z)}}) \log(l_{L^{(z)}}T)\epsilon_{L^{(z)}}^{-2}. \tag{65}
$$

Using **Fact 2** with $\tau = \Delta_{\neq,z}$ together with **Fact 3**, we find that

$$R_T^{\Delta,S} \le c \sum_{z \in \mathcal{S}} \widetilde{\kappa}_z(\Delta \vee \Delta_{\neq,z}) \log(L^{(z)}T)(\Delta_{\neq,z})^{-2}.$$

Bound on $\mathbf{R_T^{\Delta,\overline{S}} + R_T^{Rec}}$. If the algorithm does not enter the Recovery phase, then $R_T^{Rec} = 0$ and $\overline{S} = \{z^*\}$. Then, the algorithms finds the best group, and the last bias exploration phase is performed at phase $\max_{z \neq z^*} L^{(z)} \le \max_{z \neq z^*} l_{\Delta_{\neq,z}} = l_{\Delta_{\neq}}$. Then, Equation (65) implies that

$$R_T^{\Delta,\overline{S}} \le c\widetilde{\kappa}_{z^*}(\Delta \vee \Delta_{\neq}) \log(L^{(z)}T)(\Delta_{\neq})^{-2}.$$

If the algorithms enters the Recovery phase, we can use again the same arguments to show that $R_T^{\Delta,\overline{S}} \le c \sum_{z \in \overline{S}} \widetilde{\kappa}_z(\widehat{\Delta}^{L^\Delta}) \log(l_{L^{(z)}}T)\epsilon_{L^{(z)}}^{-2}$. Using **Fact 2** and Equation (65), we find that for $\tau = \epsilon_{L^\Delta}$,

$$R_T^{\Delta,\overline{S}} \le c \sum_{z \in \overline{S}} \widetilde{\kappa}_z(\Delta \vee \epsilon_{L^\Delta}) \log(l_{L^\Delta}T)\epsilon_{L^\Delta}^{-2} = c \frac{\widetilde{\kappa}_{\overline{S}}(\Delta \vee \epsilon_{L^\Delta}) \log(l_{L^\Delta}T)}{\epsilon_{L^\Delta}^2}.$$

Since all actions selected during the Recovery phase belong to $\cup_{z \in \overline{S}} \mathcal{X}_l^{(z)}$, on $\overline{\mathcal{F}}$ these actions are sub-optimal by a gap at most $c\epsilon_{L^\Delta+1}$, so $R_T^{Rec} \le cT\epsilon_{L^\Delta+1}$. Now, since the algorithm enters the Recovery phase, we must have $\epsilon_{L^\Delta+1} \le (\widetilde{\kappa}_{\overline{S}}(\Delta^{L^\Delta+1}) \log(T)/T)^{1/3}$, which implies that

$$R_T^{Rec} \le \frac{c\widetilde{\kappa}_{\overline{S}}(\widehat{\Delta}^{L^\Delta+1}) \log(T)}{\epsilon_{L^\Delta+1}^2}.$$

Together with **Fact 2**, this implies that

$$R_T^{\Delta,\overline{S}} + R_T^{Rec} \le \frac{c\widetilde{\kappa}_{\overline{S}}(\Delta \vee \epsilon_{L^\Delta}) \log(T)}{\epsilon_{L^\Delta}^2}.$$

On the one hand, **Fact 3** guarantees that, since we entered the Recovery phase before eliminating any group in $\overline{S}$, we must have $L^\Delta \le \min_{z \in \overline{S} \setminus \{z^*\}} l_{\Delta_{\neq,z}}$, so $\epsilon_{L^\Delta} \ge c \max_{z \in \overline{S}} \Delta_{\neq,z}$. On the other hand, **Fact 4** ensures that $\epsilon_{L^\Delta} \le \varepsilon_T$. Thus,

$$R_T^\Delta + R_T^{Rec} \le \sum_{s \in \overline{S} \setminus \{z^*\}} \frac{c\widetilde{\kappa}_z(\Delta \vee \varepsilon_T) \log(T)}{(\Delta_{\neq,z})^2} + \frac{c\widetilde{\kappa}_{z^*}(\Delta \vee \varepsilon_T) \log(T)}{(\Delta_{\neq})^2}.$$

Conclusion Combining these results, we find that

$$R_T \le c \left(\frac{d}{\Delta_{\min}} \vee \sum_{z \neq z^*, z \neq 1} \frac{\widetilde{\kappa}_z(\Delta \vee \Delta_{\neq,z}) \vee \widetilde{\kappa}_z(\Delta \vee \varepsilon_T)}{(\Delta_{\neq,z})^2} + \frac{\widetilde{\kappa}_{z^*}(\Delta \vee \Delta_{\neq}) \vee \widetilde{\kappa}_{z^*}(\Delta \vee \varepsilon_T)}{(\Delta_{\neq})^2} \right) \log(T)$$

when $T \ge k$. Using Lemma 8, we get that $\widetilde{\kappa}_z(\Delta \vee \Delta_{\neq}) \vee \widetilde{\kappa}_z(\Delta \vee \varepsilon_T) \le \widetilde{\kappa}_z(\Delta \vee \Delta_{\neq} \vee \varepsilon_T)$, which concludes the proof of the results. $\qquad\square$