Determinantal point processes based on orthogonal polynomials for sampling minibatches in SGD

Abstract

Stochastic gradient descent (SGD) is a cornerstone of machine learning. When the number N of data items is large, SGD relies on constructing an unbiased estimator of the gradient of the empirical risk using a small subset of the original dataset, called a minibatch. Default minibatch construction involves uniformly sampling a subset of the desired size, but alternatives have been explored for variance reduction. In particular, experimental evidence suggests drawing minibatches from determinantal point processes (DPPs), a tractable distribution over minibatches that favours diversity among selected items. However, like in recent work on DPPs for coresets, providing a systematic and principled understanding of how and why DPPs help has been difficult. In this work, we contribute an orthogonal polynomial-based determinantal point process paradigm for performing minibatch sampling in SGD. Our approach leverages the specific data distribution at hand, which endows it with greater sensitivity and power over existing data-agnostic methods. We substantiate our method via a detailed theoretical analysis of its convergence properties, interweaving between the discrete data set and the underlying continuous domain. In particular, we show how specific DPPs and a string of controlled approximations can lead to gradient estimators with a variance that decays faster with the batchsize than under uniform sampling. Coupled with existing finite-time guarantees for SGD on convex objectives, this entails that, for a large enough batchsize and a fixed budget of item-level gradients to evaluate, DPP minibatches lead to a smaller bound on the mean square approximation error than uniform minibatches. Moreover, our estimators are amenable to a recent algorithm that directly samples linear statistics of DPPs (i.e., the gradient estimator) without sampling the underlying DPP (i.e., the minibatch), thereby reducing computational overhead. We provide detailed synthetic experiments to substantiate our theoretical claims.

1   Introduction

Consider minimizing an empirical loss

$$\min_{\theta \in \Theta} \frac{1}{N} \cdot \sum_{i=1}^{N} \mathcal{L}(\boldsymbol{z}_i, \theta) + \lambda(\theta), \tag{1}$$

with some penalty $\lambda : \Theta \mapsto \mathbb{R}_+$. Many learning tasks, such as regression and classification, are usually framed that way [1]. When $N \gg 1$, computing the gradient of the objective in (1) becomes a bottleneck, even if individual gradients $\nabla_\theta \mathcal{L}(\boldsymbol{z}_i, \theta)$ are cheap to evaluate. For a fixed computational budget, it is thus tempting to replace vanilla gradient descent by more iterations but using an approximate gradient, obtained using only a few datapoints. Stochastic gradient descent (SGD; [2]) follows this template. In its simplest form, SGD builds an unbiased estimator at each

iteration of gradient descent, independently from past iterations, using a minibatch of random samples from the data set. Theory [3] and practice suggest that the variance of the gradient estimators in SGD should be kept as small as possible. It is thus natural that variance reduction for SGD has been a rich area of research; see for instance the detailed references in [Section 2][4].

In a related vein, determinantal point processes (DPPs) are probability distributions over subsets of a (typically large or infinite) ground set that are known to yield samples made of collectively diverse items, while being tractable both in terms of sampling and inference. Originally introduced in electronic optics [5], they have been turned into generic statistical models for repulsion in spatial statistics [6] and machine learning [7]. In ML, DPPs have also been shown to be efficient sampling tools; see Section 2. Importantly for us, there is experimental evidence that minibatches in (1) drawn from DPPs and other repulsive point processes can yield gradient estimators with low variance for advanced learning tasks [4, 8], though a conclusive theoretical result has remained elusive. In particular, it is hard to see the right candidate DPP when, unlike linear regression, the objective function does not necessarily have a geometric interpretation.

Our contributions are as follows. We combine continuous DPPs based on orthogonal polynomials [9] and kernel density estimators built on the data to obtain two gradient estimators; see Section 3. We prove that their variance is $\mathcal{O}(1/p^{1+1/d})$, where p is the size of the minibatch; see Section 4. This provides theoretical backing to the claim that DPPs yield variance reduction [4] over, say, sampling without replacement. In passing, the combination of analytic tools –orthogonal polynomials–, and an essentially discrete subsampling task –minibatch sampling– sheds light on new ways to build discrete DPPs for subsampling. Finally, we demonstrate our theoretical results on simulated data in Section 5.

Notation. Let data be denoted by $\mathfrak{D} := \{z_1, \ldots, z_N\}$, and assume that the z_i's are drawn i.i.d. from a distribution γ on $\mathbb{R}^d$. Assume γ is compactly supported, with support $\mathcal{D} \subset [-1, 1]^d$ bounded away from the border of $[-1, 1]^d$. Assume also that γ is continuous with respect to the Lebesgue measure, and that its density is bounded away from zero on its support. While our assumptions exclude learning problems with discrete labels, such as classification, we later give experimental support that our estimators yield variance reduction in that case too. We define the *empirical measure* $\hat{\gamma}_N := N^{-1} \cdot \sum_{i=1}^N \delta_{z_i}$, where δ_{z_i} is the delta measure at the point $z_i \in \mathbb{R}^d$. Clearly, $\hat{\gamma}_N \to \gamma$ in $\mathcal{P}(\mathbb{R}^d)$; under our operating assumption of compact support, this amounts to convergence in $\mathcal{P}(\mathcal{D})$.

For simplicity, we assume that no penalty is used in (1), but our results will extend straightforwardly. We denote the gradient of the empirical loss by $\Xi_N = \Xi_N(\theta) := N^{-1} \cdot \sum_{i=1}^N \nabla_\theta \mathcal{L}(z_i, \theta)$. A minibatch is a (random) subset $A \subset [N]$ of size $|A| = p \ll N$ such that the random variable

$$\Xi_A = \Xi_A(\theta) := \sum_{i \in A} w_i \nabla_\theta \mathcal{L}(z_i, \theta), \tag{2}$$

for suitable weights (w_i), provides a good approximation for Ξ_N.

2 Background and relevant literature

Stochastic gradient descent. Going back to [2], SGD has been a major workhorse for machine learning; see e.g. [1, Chapter 14]. The basic version of the algorithm, applied to the empirical risk minimization (1), is to repeatedly update

$$\theta_t \leftarrow \theta_t - \gamma_t \Xi_A(\theta_t), \quad t \in \mathbb{N}, \tag{3}$$

where γ_t is a (typically decreasing) stepsize, and Ξ_A is a minibatch-based estimator (2) of the gradient of the penalized empirical risk function (1), possibly depending on past iterations. Most theoretical analyses assume that for any θ, $\Xi_A(\theta)$ in the t-th update (3) is unbiased, conditionally on the history of the Markov chain (θ_t) so far. For simplicity, we henceforth make the assumption that Ξ_A does not depend on the past of the chain. In particular, using such unbiased gradients, one can derive a nonasymptotic bound [3] on the mean square distance of θ_t to the minimizer of (1), for strongly convex and smooth loss functions like linear regression and logistic regression with ℓ_2 penalization. More precisely, for $\gamma_t \propto t^{-\alpha}$ and $0 < \alpha < 1$, [3, Theorem 1] yields

$$\mathbb{E}\|\theta_t - \theta_\star\|^2 \leq f(t) \left(\mathbb{E}\|\theta_0 - \theta_\star\|^2 + \frac{\sigma^2}{L} \right) + C\frac{\sigma^2}{t^\alpha}, \tag{4}$$

where $C, L > 0$ are problem-dependent constants, $f(t) = \mathcal{O}(e^{-t^{\alpha}})$, and $\sigma^2 = \mathbb{E}[\|\Xi_A(\theta_\star)\|^2|\mathfrak{D}]$ is the trace of the covariance matrix of the gradient estimator, evaluated at the optimizer $\theta_\star$ of (1). The initialization bias is thus forgotten subexponentially fast, while the asymptotically leading term is proportional to σ^2/t^α. Combined with practical insight that variance reduction for gradients is key, theoretical results like (4) have motivated methodological research on efficient gradient estimators [4, Section 2], i.e., constructing minibatches so as to minimize σ^2. In particular, repulsive point processes such as determinantal point processes have been empirically demonstrated to yield variance reduction and overall better performance on ML tasks [4, 8]. Our paper is a stab at a theoretical analysis to support these experimental claims.

The Poissonian benchmark. The default approach to sample a minibatch $A \subset [N]$ is to sample p data points from $\mathfrak{D}$ uniformly, with or without replacement, and take $w_i = 1/p$ constant in (2). Both sampling with or without replacement lead to unbiased gradient estimators. A similar third approach is Poissonian random sampling. This simply consists in starting from $A = \emptyset$, and independently adding each element of $\mathfrak{D}$ to the minibatch A with probability p/N. The Poisson estimator $\Xi_{A,\mathrm{Poi}}$ is then (2), with constant weights again equal to $1/p$. When $p \ll N$, which is the regime of SGD, the cardinality of A is tightly concentrated around p, and $\Xi_{A,\mathrm{Poi}}$ has the same fluctuations as the two default estimators, while being easier to analyze. In particular, $\mathbb{E}[\Xi_{A,\mathrm{Poi}}|\mathfrak{D}] = \Xi_N$ and $\mathrm{Var}[\Xi_{A,\mathrm{Poi}}|\mathfrak{D}] = O_P(p^{-1})$; see Appendix S1 for details.

DPPs as (sub)sampling algorithms. As distributions over subsets of a large ground set that favour diversity, DPPs are intuitively good candidates at subsampling tasks, and one of their primary applications in ML has been as summary extractors [7]. Since then, DPPs or mixtures thereof have been used, e.g., to generate experimental designs for linear regression, leading to strong theoretical guarantees [10, 11, 12]; see also [13] for a survey of DPPs in randomized numerical algebra, or [14] for feature selection in linear regression with DPPs.

When the objective function of the task has less structure than linear regression, it has been more difficult to prove that finite DPPs significantly improve over i.i.d. sampling. For coreset construction, for instance, [15] manage to prove that a projection DPP necessarily improves over i.i.d. sampling with the same marginals [15, Corollary 3.7], but the authors stress the disappointing fact that current concentration results for strongly Rayleigh measures (such as DPPs) do not allow yet to prove that DPP coresets need significantly fewer points than their i.i.d. counterparts [15, Section 3.2]. Even closer to our motivation, DPPs for minibatch sampling have shown promising experimental performance [4], but reading between the lines of the proof of [4, Theorem 1], a bad choice of DPP can even yield a *larger* variance than i.i.d. sampling!

For such unstructured problems (compared to, say, linear regression) as coreset extraction or loss-agnostic minibatch sampling, we propose to draw inspiration from work on continuous DPPs, where faster-than-i.i.d. error decays have been proven in similar contexts. For instance, orthogonal polynomial theory motivated [9] to introduce a particular DPP, called a multivariate orthogonal polynomial ensemble, and prove a faster-than-i.i.d. central limit theorem for Monte Carlo integration of smooth functions. While resorting to continuous tools to study a discrete problem such as minibatch sampling may be unexpected, we shall see that the assumption that the size of the dataset is large compared to the ambient dimension crucially allows to transfer variance reduction arguments.

3 DPPs, and two gradient estimators

Since we shall need both discrete and continuous DPPs, we assume that $\mathcal{X}$ is either $\mathbb{R}^d$ or $\mathfrak{D}$, and follow [16] in introducing DPPs in an abstract way that encompasses both cases. After that, we propose two gradient estimators for SGD that build on a particular family of continuous DPPs introduced in [9] for Monte Carlo integration, called multivariate orthogonal polynomial ensembles.

3.1 DPPs: The kernel machine of point processes

A point process on $\mathcal{X}$ is a distribution on finite subsets S of $\mathcal{X}$; see [17] for a general reference. Given a reference measure μ on X, a point process is said to be determinantal (DPP) if there exists a

function $K : \mathcal{X} \times \mathcal{X} \to \mathbb{R}$, the *kernel* of the DPP, such that, for every $n \geq 1$,

$$\mathbb{E}\left[\sum_{\neq} f(x_{i_1}, \ldots, x_{i_n})\right] = \int_{(\mathcal{X})^n} f(x_1, \ldots, x_n) \cdot \mathrm{Det}\left[K(x_i, x_\ell)\right]_{i,\ell=1}^n \mathrm{d}\mu^{\otimes n}(x_1, \ldots, x_n) \quad (5)$$

for every bounded Borel function $f : \mathcal{X}^n \to \mathbb{R}$, where the sum in the LHS of (5) ranges over all pairwise distinct n-uplets of the random finite subset S. A few remarks are in order. First, satisfying (5) for every n is a strong constraint on K, so that not every kernel yields a DPP. A set of sufficient conditions on K is given by the Macchi-Soshnikov theorem [5, 18]. In words, if $K(x, y) = K(y, x)$, and if K further corresponds to an integral operator

$$f \mapsto \int K(x, y) f(y) \mathrm{d}\mu(y), \quad f \in L^2(\mathcal{X}, \mu)$$

that is trace-class with spectrum in $[0, 1]$, then the corresponding DPP exists. Second, note in (5) that the kernel of a DPP encodes how the points in the random configurations interact. A strong point if favour of DPPs is that, unlike most interacting point processes, sampling and inference are tractable [16, 6]. Third, (5) yields simple formulas for the mean and variance of linear statistics of a DPP.

Proposition 1 (See e.g. [19, 20]). *Let $S \sim DPP(K, \mu)$ and $\Phi : \mathcal{X} \to \mathbb{R}$ be a bounded Borel function,*

$$\mathbb{E}\left[\sum_{x \in S} \Phi(x)\right] = \int \Phi(x) K(x, x) \mathrm{d}\mu(x) \quad (6)$$

$$\mathrm{Var}\left[\sum_{x \in S} \Phi(x)\right] = \iint \|\Phi(x) - \Phi(y)\|_2^2 |K(x, y)|^2 \mathrm{d}\mu(x) \mathrm{d}\mu(y)$$

$$+ \int \|\Phi(x)\|_2^2 \left(K(x, x) - \int |K(x, y)|^2 \mathrm{d}\mu(y)\right) \mathrm{d}\mu(x). \quad (7)$$

Since the seminal paper [7], the case $\mathcal{X} = \mathfrak{D}$ has been the most common in machine learning. Taking μ to be $\hat{\gamma}_N$, any kernel is given by its restriction to $\mathfrak{D}$, usually given as an $N \times N$ matrix $\mathbf{K} = K|_{\mathfrak{D}}$. Equation (6) with $n = p$, A a subset of size p of $\mathfrak{D}$, and Φ the indicator of A yields $\mathbb{P}(A \subset S) = N^{-p} \mathrm{Det} \mathbf{K}_A$. This is the usual way finite DPPs are introduced [7], except maybe for the factor N^{-p}, which comes from using $\hat{\gamma}_N$ as the reference measure instead of $N\hat{\gamma}_N$. In this finite setting, a careful implementation of the general DPP sampler of [16] yields m DPP samples of average cardinality $\mathrm{Tr}(\mathbf{K})$ in $\mathcal{O}(N^\omega + mN\mathrm{Tr}(\mathbf{K})^2)$ operations [21].

We now go back to a general $\mathcal{X}$ and fix $p \in \mathbb{N}$. A canonical way to construct DPPs generating configurations of p points almost surely, i.e. $S = \{x_1, \ldots, x_p\}$, is the following. Consider p orthonormal functions $\phi_0, \ldots, \phi_{p-1}$ in $L^2(\mu)$, and take for kernel

$$K(x, y) = \sum_{k=0}^{p-1} \phi_k(x) \phi_k(y). \quad (8)$$

In this setting, the (permutation invariant) random variables $x_1, \ldots, x_p$ with joint distribution

$$\frac{1}{p!} \mathrm{Det}\left[K(x_i, x_\ell)\right]_{i,\ell=1}^p \prod_{i=1}^p \mathrm{d}\mu(x_i) \quad (9)$$

generate a DPP $\{x_1, \ldots, x_p\}$ with kernel $K(x, y)$, called a *projection* DPP. For further information on determinantal point processes, we refer the reader to [16, 7].

3.2 Multivariate orthogonal polynomial ensembles

This section paraphrases [9] in their definition of a particular projection DPP on $\mathcal{X} = \mathbb{R}^d$, called a multivariate orthogonal polynomial ensemble (OPE). Fix some reference measure $q(x)\mathrm{d}x$ on $[-1, 1]^d$, and assume that it puts positive mass on some open subset of $[-1, 1]^d$. Now choose an ordering of the monomial functions $(x_1, \ldots, x_d) \mapsto x_1^{\alpha_1} \cdots x_d^{\alpha_d}$; in this work we use the graded lexical order. Then apply the Gram-Schmidt algorithm in $L^2(q(x)\mathrm{d}x)$ to these ordered monomials. This yields a sequence of orthonormal polynomial functions $(\phi_k)_{k \in \mathbb{N}}$, the multivariate orthonormal polynomials w.r.t. q. Finally, plugging the first p multivariate orthonormal polynomials $\phi_0, \ldots, \phi_{p-1}$ into the projection kernel (8), we obtain a projection DPP with kernel denoted as $K_q^{(p)}$, referred to as the *multivariate OPE* associated with the measure $q(x)\mathrm{d}x$.

3.3 Our first estimator: reweight, restrict, and saturate an OPE kernel

Let $h > 0$ and

$$\tilde{\gamma}(\boldsymbol{z}) = \frac{1}{Nh^d} \sum_{i=1}^{N} k\left(\frac{\boldsymbol{z} - \boldsymbol{z}_i}{h}\right) \tag{10}$$

be a single-bandwith kernel density estimator of the pdf of the data-generating distribution γ; see [22, Section 4.2], In particular, k is chosen so that $\int k(x)\mathrm{d}x = 1$. Note that the approximation kernel k is unrelated to any DPP kernel in this paper. Let now $q(x) = q_1(x_1)\ldots q_d(x_d)$ be a separable pdf on $[-1,1]^d$, where each q_i is Nevai-class[1]. Let $K_q^{(p)}$ be the multivariate OPE kernel defined in Section 3.2, and form a new kernel

$$K_{q,\tilde{\gamma}}^{(p)}(x,y) := \sqrt{\frac{q(x)}{\tilde{\gamma}(x)}} K_q^{(p)}(x,y) \sqrt{\frac{q(y)}{\tilde{\gamma}(y)}}.$$

The form of $K_{q,\tilde{\gamma}}^{(p)}$ is reminiscent of importance sampling [23], which is no accident. Indeed, while the positive semidefinite matrix

$$K_{q,\tilde{\gamma}}^{(p)}|_{\mathfrak{D}} := \left(K_{q,\tilde{\gamma}}^{(p)}(\boldsymbol{z}_i, \boldsymbol{z}_j)\right)_{1 \leq i,j \leq N}$$

is not necessarily the kernel matrix of a DPP on $(\{1,\ldots,N\}, \hat{\gamma}_N)$, see Section 3.1, we built it to be close to a projection of rank p. More precisely,

$$\int K_{q,\tilde{\gamma}}^{(p)}(\boldsymbol{z}_k, y) K_{q,\tilde{\gamma}}^{(p)}(y, \boldsymbol{z}_\ell)\mathrm{d}\hat{\gamma}_N(y) = \sqrt{\frac{q(\boldsymbol{z}_k)}{\tilde{\gamma}(\boldsymbol{z}_k)}} \left[\frac{1}{N} \sum_{n=1}^{N} K_q^{(p)}(\boldsymbol{z}_k, \boldsymbol{z}_n) K_q^{(p)}(\boldsymbol{z}_n, \boldsymbol{z}_\ell) \frac{q(\boldsymbol{z}_n)}{\tilde{\gamma}(\boldsymbol{z}_n)}\right] \sqrt{\frac{q(\boldsymbol{z}_\ell)}{\tilde{\gamma}(\boldsymbol{z}_\ell)}}.$$

If N is large compared to p and d, so that in particular $\tilde{\gamma} \approx \gamma$, the term within brackets will be close to $\int K_q^{(p)}(\boldsymbol{z}_k, \boldsymbol{z}) K_q^{(p)}(\boldsymbol{z}, \boldsymbol{z}_\ell) q(\boldsymbol{z})\mathrm{d}\boldsymbol{z} = K_q^{(p)}(\boldsymbol{z}_k, \boldsymbol{z}_\ell)$, so that $K_{q,\tilde{\gamma}}^{(p)}|_{\mathfrak{D}}$ is almost a projection in $L^2(\hat{\gamma}_N)$.

Let us actually consider the orthogonal projection matrix $\widetilde{\mathbf{K}}$ with the same eigenvectors as $K_{q,\tilde{\gamma}}^{(p)}|_{\mathfrak{D}}$, but with the p largest eigenvalues replaced by 1, and the rest of the spectrum set to 0. By the Macchi-Soshnikov theorem, $\widetilde{\mathbf{K}}$ is the kernel matrix of a DPP; see Section 3.1. We thus consider a minibatch $A \sim \mathrm{DPP}(\widetilde{\mathbf{K}})$. Coming from a projection DPP, $|A| = p$ almost surely, and we define the gradient estimator

$$\Xi_{A,\mathrm{DPP}} := \sum_{i \in A} \frac{\nabla_\theta \mathcal{L}(\boldsymbol{z}_i, \theta)}{\widetilde{\mathbf{K}}_{ii}}. \tag{11}$$

In Section 4, we shall prove that $\Xi_{A,\mathrm{DPP}}$ is unbiased, and examine under what assumptions its variance decreases faster than $1/p$. In terms of computational cost, the bottleneck is computing the p largest eigenvalues of $K_{q,\tilde{\gamma}}^{(p)}|_{\mathfrak{D}}$, along with the corresponding eigenvectors. This can be done once before running SGD, as a preprocessing step. Note that storing the kernel in diagonalized form only requires $\mathcal{O}(Np)$ storage. Each iteration of SGD then only requires sampling a rank-p projection DPP with diagonalized kernel, which takes $\mathcal{O}(Np^2)$ elementary operations [21]. Alternately, (11) can be sampled directly, without sampling the underlying DPP. Indeed the Laplace transform of (11) is a Fredholm determinant, and is is shown in [24] that Nyström-type approximations of that determinant, followed by Laplace inversion, yield an accurate inverse CDF sampler.

Finally, we stress the unusual way in which our finite DPP kernel $\widetilde{\mathbf{K}}$ is constructed, through a reweighted continuous OPE kernel, restricted to the actual dataset. This construction is interesting *per se*, as it is key to leveraging analytic techniques from the continuous case in Section 4.

3.4 Our second estimator: sample the OPE, but smooth the gradient

In Section 3.3, we smoothed the empirical distribution of the data and restricted a continuous kernel to the dataset $\mathfrak{D}$, to make sure that the drawn minibatch would be a subset of $\mathfrak{D}$. But one could actually define another gradient estimator, directly from an OPE sample $A = \{\boldsymbol{w}_1, \ldots, \boldsymbol{w}_p\} \sim \mathrm{DPP}(K_q^{(p)}, q)$.

---

[1]See [9, Section 4] for details. It suffices that each q_i is positive on $[-1,1]^d$.

183 Note that in that case, the "generalized minibatch" $A \subset [-1, 1]^d$ is not necessarily a subset of the
184 dataset $\mathfrak{D}$. Defining a kernel density estimate of the gradient,

$$\widehat{\nabla_\theta \mathcal{L}}(\boldsymbol{z}, \theta) := \frac{1}{Nh^d} \sum_{i=1}^{N} \nabla_\theta \mathcal{L}(\boldsymbol{z}_i, \theta) \cdot k\left(\frac{\boldsymbol{z} - \boldsymbol{z}_i}{h}\right),$$

185 we consider the estimator

$$\Xi_{A,\mathrm{s}} = \sum_{w_j \in A} \frac{\widehat{\nabla_\theta \mathcal{L}}(\boldsymbol{w}_j, \theta)}{q(\boldsymbol{w}_j) K_q^{(p)}(\boldsymbol{w}_j, \boldsymbol{w}_j)}.$$

186 Since each evaluation of this estimator is at least as costly as evaluating the actual gradient $\Xi_N(\theta)$, its
187 use is mostly theoretical: the analysis of the fluctuations of $\Xi_{A,\mathrm{s}}$ is easier than that of $\Xi_{A,\mathrm{DPP}}$, while
188 requiring the same key steps. Moreover, the computation of all pairwise distances in $\Xi_{A,\mathrm{s}}$ could be
189 efficiently approximated, possibly using random projection arguments [25], so that the limited scope
190 of $\Xi_{A,\mathrm{s}}$ might be overcome in future work. Note also that, like $\Xi_{A,\mathrm{DPP}}$, inverse Laplace sampling
191 [24] applies to $\Xi_{A,\mathrm{s}}$.

192 4   Analysis of determinantal sampling of SGD gradients

193 We first analyze the bias, and then the fluctuations, of the gradient estimators introduced in Section 3.
194 By Proposition 1 with $\Phi = \nabla_\theta \mathcal{L}(\cdot, \theta)$, it comes

$$\mathbb{E}[\Xi_{A,\mathrm{DPP}} | \mathfrak{D}] = \int_D \nabla_\theta \mathcal{L}(\boldsymbol{z}, \theta) \underline{K_{q,\gamma}^{(p)}(\boldsymbol{z}, \boldsymbol{z})^{-1}} \cdot \underline{K_{q,\gamma}^{(p)}(\boldsymbol{z}, \boldsymbol{z})} \mathrm{d}\hat\gamma_N(z) = \frac{1}{N} \sum_{i=1}^{N} \nabla_\theta \mathcal{L}(\boldsymbol{z}_i, \theta),$$

195 so that we immediatly get the following result.

196 **Proposition 2.** $\mathbb{E}[\Xi_{A,\mathrm{DPP}} | \mathfrak{D}] = \Xi_N$.

197 Thus, $\Xi_{A,\mathrm{DPP}}$ is unbiased, like the classical Poissonian benchmark in Section 2. However, the
198 smoothed estimator $\Xi_{A,\mathrm{s}}$ from Section 3.4 is slightly biased. Note that while the results of [3], like
199 (4), do not apply to biased estimators, SGD can be analyzed in the small-bias setting [26].

200 **Proposition 3.** *Assume that k in (10) has compact support and q is bounded on D. Then $\mathbb{E}[\Xi_{A,\mathrm{s}} | \mathfrak{D}] =$*
201 *$\Xi_N + O_P(ph/N)$.*

202 *Proof.* Using the first part of Proposition 1 again, it comes

$$\mathbb{E}[\Xi_{A,\mathrm{s}} | \mathfrak{D}] = \mathbb{E}_{A \sim \mathrm{DPP}(K_q^{(p)}, q)} \left[\sum_{w_j \in A} \frac{\widehat{\nabla_\theta \mathcal{L}}(\boldsymbol{w}_j, \theta)}{q(\boldsymbol{w}_j) K_q^{(p)}(\boldsymbol{w}_j, \boldsymbol{w}_j)} \right]$$

$$= \int_D \frac{\widehat{\nabla_\theta \mathcal{L}}(\boldsymbol{w}, \theta)}{\underline{q(\boldsymbol{w}) K_q^{(p)}(\boldsymbol{w}, \boldsymbol{w})}} \cdot \underline{K_q^{(p)}(\boldsymbol{w}, \boldsymbol{w}) q(\boldsymbol{w})} \mathrm{d}\boldsymbol{w} = \int_D \left(\frac{1}{Nh^d} \sum_{i=1}^{N} \nabla_\theta \mathcal{L}(\boldsymbol{z}_i, \theta) k\left(\frac{\boldsymbol{z} - \boldsymbol{z}_i}{h}\right) \right) \mathrm{d}\boldsymbol{w}$$

$$= \frac{1}{N} \sum_{i=1}^{N} \left(\int_D \frac{1}{h^d} \cdot k\left(\frac{\boldsymbol{w} - \boldsymbol{z}_i}{h}\right) \mathrm{d}\boldsymbol{w} \right) \cdot \nabla_\theta \mathcal{L}(\boldsymbol{z}_i, \theta) = \frac{1}{N} \sum_{i=1}^{N} \nabla_\theta \mathcal{L}(\boldsymbol{z}_i, \theta) + O_P(ph/N), \quad (12)$$

203 where, in the last step, we have used the fact that $h^{-d} \cdot k\left(h^{-1}(\boldsymbol{w} - \boldsymbol{z}_i)\right)$ is a probability density.
204 The $O_P(ph/N)$ error term arises because we integrate that pdf on D rather than $\mathrm{Supp}(k)$. But since
205 k is a compactly supported kernel, for points $\boldsymbol{z}_i$ that are within a distance $O(h)$ from the boundary
206 of D, we incur an error of $O_P(1)$. By Proposition 1, the expected number of such points $\boldsymbol{z}_i$ is
207 $\int_{D_h} K_q^{(p)}(\boldsymbol{w}, \boldsymbol{w}) q(\boldsymbol{w}) \mathrm{d}\boldsymbol{w}$, where D_h is the h-neighbourhood of the boundary of D. By a classical
208 asymptotic result of Totik, see [9, Theorem 4.8], $\boldsymbol{w} \mapsto K_q^{(p)}(\boldsymbol{w}, \boldsymbol{w})$ is $O(p)$ on D_h; whereas q is a
209 bounded density, implying that $\int_{D_h} q(\boldsymbol{w}) \mathrm{d}\boldsymbol{w} = O(\mathrm{Vol}(D_h)) = O(h)$. Putting together all of these,
210 we obtain the $O_P(ph/N)$ error term in (12). ∎

211 The rest of this section is devoted to the more involved task of analyzing the fluctuations of $\Xi_{A,\mathrm{DPP}}$
212 and $\Xi_{A,\mathrm{s}}$. For the purposes of our analysis, we discuss certain desirable regularity behaviour for our
213 kernels and loss functions in Section 4.1. Then, we tackle the main ideas behind the study of the
214 fluctuations of our estimators in Section 4.2. Details can be found in Appendix S3.

4.1 Some regularity phenomena

Assume that $(\frac{1}{p} \cdot K_q^{(p)}(\boldsymbol{z}, \boldsymbol{z}))^{-1} \nabla_\theta \mathcal{L}(\boldsymbol{z}, \theta)$ is bounded on the domain D uniformly in N (note that p possibly depends on N). Furthermore, assume that $\boldsymbol{z} \mapsto (p^{-1} \cdot K_q^{(p)}(\boldsymbol{z}, \boldsymbol{z}) q(\boldsymbol{z}))^{-1} \nabla_\theta \mathcal{L}(\boldsymbol{z}, \theta) \cdot \tilde{\gamma}(\boldsymbol{z})$ is 1-Lipschitz, with Lipschitz constant $O_P(1)$ and bounded in θ.

Such properties are natural in the context of various known asymptotic phenomena, in particular asymptotic results on OPE kernels and the convergence of the KDE $\tilde{\gamma}$ to the distribution γ. The detailed discussion is deferred to Appendix S2; we record here in passing that the above asymptotic phenomena imply that

$$\frac{\nabla_\theta \mathcal{L}(\boldsymbol{z}, \theta)}{\frac{1}{p} \cdot K_q^{(p)}(\boldsymbol{z}, \boldsymbol{z}) q(\boldsymbol{z})} \cdot \tilde{\gamma}(\boldsymbol{z}) \longrightarrow \nabla_\theta \mathcal{L}(\boldsymbol{z}, \theta) \cdot \gamma(\boldsymbol{z}) \prod_{j=1}^{d} \sqrt{1 - \boldsymbol{z}[j]^2}. \tag{13}$$

Our desired regularity properties may therefore be understood in terms of closeness to this limit, which itself has similar regularity. At the price of these assumptions, we can use analytic tools to derive fluctuations for $\Xi_{A,\mathrm{DPP}}$ by working on the limit in (13). For the fluctuation analysis of the smoothed estimator $\Xi_{A,\mathrm{s}}$, we similarly assume that $(q(\boldsymbol{w}) \cdot \frac{1}{p} K_q^{(p)}(\boldsymbol{w}, \boldsymbol{w}))^{-1} \widehat{\nabla_\theta \mathcal{L}}(\boldsymbol{w}, \theta)$ is $O_P(1)$ and 1-Lipschitz with an $O_P(1)$ Lipschitz constant that is bounded in θ. These are once again motivated by the convergence of $(q(\boldsymbol{w}) \cdot \frac{1}{p} K_q^{(p)}(\boldsymbol{w}, \boldsymbol{w}))^{-1} \widehat{\nabla_\theta \mathcal{L}}(\boldsymbol{w}, \theta)$ to $\nabla_\theta \mathcal{L}(\boldsymbol{w}, \theta) \prod_{j=1}^{d} \sqrt{1 - \boldsymbol{w}[j]^2}$.

4.2 Reduced fluctuations for determinantal samplers

For succinctness of presentation, we focus on the theoretical analysis of the smoothed estimator $\Xi_{A,\mathrm{s}}$, leaving the analysis of $\Xi_{A,\mathrm{DPP}}$ to Appendix S3, while still capturing the main ideas of our approach.

Proposition 4. *Under the assumptions in Section 4.1,* $\mathrm{Var}[\Xi_{A,\mathrm{s}}|\mathfrak{D}] = O_P(p^{-(1+1/d)})$.

Proof Sketch. Invoking (7) in the case of a projection kernel,

$$\mathrm{Var}[\Xi_{A,\mathrm{s}}|\mathfrak{D}] = \iint \left\| \frac{\widehat{\nabla_\theta \mathcal{L}}(\boldsymbol{z}, \theta)}{q(\boldsymbol{z}) K_q^{(p)}(\boldsymbol{z}, \boldsymbol{z})} - \frac{\widehat{\nabla_\theta \mathcal{L}}(\boldsymbol{w}, \theta)}{q(\boldsymbol{w}) K_q^{(p)}(\boldsymbol{w}, \boldsymbol{w})} \right\|_2^2 |K_q^{(p)}(\boldsymbol{z}, \boldsymbol{w})|^2 \mathrm{d}q(\boldsymbol{z}) \mathrm{d}q(\boldsymbol{w})$$

$$= \frac{1}{p^2} \iint \left\| \frac{\widehat{\nabla_\theta \mathcal{L}}(\boldsymbol{z}, \theta)}{q(\boldsymbol{z}) \cdot \frac{1}{p} K_q^{(p)}(\boldsymbol{z}, \boldsymbol{z})} - \frac{\widehat{\nabla_\theta \mathcal{L}}(\boldsymbol{w}, \theta)}{q(\boldsymbol{w}) \cdot \frac{1}{p} K_q^{(p)}(\boldsymbol{w}, \boldsymbol{w})} \right\|_2^2 |K_q^{(p)}(\boldsymbol{z}, \boldsymbol{w})|^2 \mathrm{d}q(\boldsymbol{z}) \mathrm{d}q(\boldsymbol{w})$$

$$\lesssim \mathcal{M}_\theta \cdot \frac{1}{p^2} \int \int \|\boldsymbol{z} - \boldsymbol{w}\|_2^2 |K_q^{(p)}(\boldsymbol{z}, \boldsymbol{w})|^2 \mathrm{d}q(\boldsymbol{z}) \mathrm{d}q(\boldsymbol{w}), \tag{14}$$

where we used the 1-Lipschitzianity of $\frac{\widehat{\nabla_\theta \mathcal{L}}(\boldsymbol{z}, \theta)}{q(\boldsymbol{z}) K_q^{(p)}(\boldsymbol{z}, \boldsymbol{z})}$, with $\mathcal{M}_\theta = O_P(1)$ the Lipschitz constant.

We control the integral in (14) by invoking the renowned Christoffel-Darboux formula for the OPE kernel $K_q^{(p)}$ [27]. For clarity of presentation, we outline here the main ideas for $d = 1$; the details for general dimensions are available in Appendix S3. Broadly speaking, since $K_q^{(p)}$ is an orthogonal projection of rank p in $L_2(q)$, we may observe that $\iint |K_q^{(p)}(\boldsymbol{z}, \boldsymbol{w})|^2 \mathrm{d}q(\boldsymbol{z}) \mathrm{d}q(\boldsymbol{w}) = p$; so that without the $\|z - w\|_2^2$ term in (14), we would have a $O_P(p^{-1})$ behaviour in total, which would be similar to the Poissonian estimator. However, it turns out that the leading order contribution to $\int \int \|z - w\|_2^2 |K_q^{(p)}(\boldsymbol{z}, \boldsymbol{w})|^2 \mathrm{d}q(\boldsymbol{z}) \mathrm{d}q(\boldsymbol{w})$ comes from near the diagonal $z = w$, and this turns out to be neutralised in an extremely precise manner by the $\|\boldsymbol{z} - \boldsymbol{w}\|_2^2$ factor that vanishes on the diagonal.

This neutralisation is systematically captured by the Christoffel-Darboux formula [27], which implies

$$K_q^{(p)}(x, y) = (x - y)^{-1} A_p \cdot (\phi_q(x) \phi_{q-1}(y) - \phi_q(y) \phi_{q-1}(x)),$$

where $A_p = O(1)$ and ϕ_q, ϕ_{q-1} are two orthonormal functions in $L_2(q)$. Substituting this into (14), a simple computation shows that $\iint \|z - w\|_2^2 |K_q^{(p)}(\boldsymbol{z}, \boldsymbol{w})|^2 \mathrm{d}q(\boldsymbol{z}) \mathrm{d}q(\boldsymbol{w}) = O(1)$. This leads to a variance bound $\mathrm{Var}[\Xi_{A,\mathrm{s}}|\mathfrak{D}] = O_P(p^{-2})$, which is the desired rate for $d = 1$. For general d, we use the fact that $q = \bigotimes_{i=1}^{d} q_i$ is a product measure, and apply the Christoffel-Darboux formula for each q_i, leading to a variance bound of $\mathrm{Var}[\Xi_{A,\mathrm{s}}|\mathfrak{D}] = O_P(p^{-(1+1/d)})$, as desired. ∎

248 The theoretical analysis of $\mathrm{Var}[\Xi_{A,\mathrm{DPP}}|\mathfrak{D}]$ follows the broad contours of the argument for
249 $\mathrm{Var}[\Xi_{A,\mathrm{s}}|\mathfrak{D}]$ as above, with additional difficulties introduced by the spectral truncation from $K_{q,\tilde{\gamma}}^{(p)}$ to
250 $\widetilde{\mathbf{K}}$; see Section 3. This is addressed by an elaborate spectral approximation analysis in Appendix S3,
251 which combines with the above ideas to indicate a fluctuation bound of $\mathrm{Var}[\Xi_A|\mathfrak{D}] = O_P(p^{-(1+1/d)})$,
252 as desired. We defer the detailed arguments to the Section S3.

5 Experiments

254 In this section, we compare the empirical performance and the variance decay of our gradient
255 estimator $\Xi_{A,\mathrm{DPP}}$ to the default $\Xi_{A,\mathrm{Poi}}$. We do not to include $\Xi_{A,\mathrm{s}}$, whose interest is mostly
256 theoretical; see Section 4. Moreover, while we focus on simulated data to illustrate our theoretical
257 analysis, we provide experimental results on a real dataset in Appendix S4. Throughout this section,
258 the pdf q introduced in Section 3.2 is taken to be $q(\boldsymbol{w}) \propto \prod_{j=1}^d (1 + \boldsymbol{w}[j])^{\alpha_j}(1 - \boldsymbol{w}[j])^{\beta_j}$, with
259 $\alpha_j, \beta_j \in [-1/2, 1/2]$ tuned to match the first two moments of the jth marginal of $\hat{\gamma}_N$. All DPPs are
260 sampled using the Python package DPPy [28], which is under MIT licence.

Experimental setup. We consider the ERM setting (1) for linear regression $\mathcal{L}_{\mathrm{lin}}((x,y),\theta) =$
262 $0.5(\langle x,\theta \rangle - y)^2$ and logistic regression $\mathcal{L}_{\mathrm{log}}((x,y),\theta) = \log[1 + \exp(-y\langle x,\theta \rangle)]$, both with an
263 additional ℓ_2 penalty $\lambda(\theta) = (\lambda_0/2)\|\theta\|_2^2$. Here the features are $x \in [-1,1]^{d-1}$, and the labels are,
264 respectively, $y \in [-1,1]$ and $y \in \{-1,1\}$. Note that our proofs currently assume that the law γ of
265 $z = (x,y)$ is continuous w.r.t. Lebesgue, which in all rigour excludes logistic regression. However,
266 we demonstrate below that if we draw a minibatch using our DPP but on features only, and then
267 deterministically associate each label to the drawn features, we observe the same gains for logistic
268 regression as for linear regression, where the DPP kernel takes into account both features and labels.

269 For each experiment, we generate $N = 1000$ data points x_i with either the uniform distribution or a
270 mixture of 2 well-separated Gaussian distributions on $[-1,1]^{d-1}$. The variable $y_i \in \mathbb{R}$ is generated
271 as $\langle x_i, \theta_{\mathrm{true}} \rangle + \varepsilon_i$, where $\theta_{\mathrm{true}} \in \mathbb{R}^{d-1}$ is given, and $\varepsilon \in \mathbb{R}^N$ is a white Gaussian noise vector. In the
272 linear regression case, we scale y_i by $\|y\|_\infty$ to make sure that $y_i \in [-1,1]$. In the logistic regression
273 case, we replace each y_i by its sign. The regularization parameter λ_0 is manually set to be 0.1 and
274 the stepsize for the t-th iteration is set as $1/t^{0.9}$, so that (4) applies. In each experiment, performance
275 metrics are averaged over 1000 independent runs of each SGD variant.

Performance evaluation of sampling strategies in SGD. Figure 1 summarizes the experimental
277 results of $\Xi_{A,\mathrm{Poi}}$ and $\Xi_{A,\mathrm{DPP}}$, with $p = 5$ and $p = 10$. The top row shows how the norm of the
278 complete gradient $\|\Xi_N(\theta_t)\|$ decreases with the number of individual gradient evaluations $t \times p$,
279 called here *budget*. The bottom row shows the decrease of $\|\theta_t - \theta_\star\|$. Note that using $t \times p$ on all
280 x-axes allows comparing different batchsizes. Error bars on the bottom row are $\pm$ one standard
281 deviation of the mean. On all experiments, using a DPP consistently improves the performance of
282 Poisson minibatches of the same size, and the DPP with batchsize 5 sometimes even outperforms
283 Poisson sampling with batchsize 10, showing that smaller but more diverse batches can be a better
284 way of spending a fixed number of gradient evaluations. This is particularly true for mixture data
285 (middle column), where forcing diversity with our DPP brings the biggest improvement. Maybe less
286 intuitively, the gains for the logistic regression in $d = 11$ (last column) are also significant, while
287 the case of discrete labels is not covered yet by our theoretical analysis, and the moderately large
288 dimension makes the improvement in the decay rate of the variance minor. This indicates that there is
289 variance reduction beyond the change of the rate.

Variance decay. For a given dimension d, we want to infer the rate of decay of the variance
291 $\sigma^2 = \mathbb{E}[\|\Xi_A(\theta_\star)\|^2|\mathfrak{D}]$, to confirm the $\mathcal{O}(p^{-1-1/d})$ rate discussed in Section 4. We take $\mathcal{L} = \mathcal{L}_{\mathrm{lin}}$
292 as an example, with $N = 1000$ i.i.d. samples $\mathfrak{D}$ from the uniform distribution on $[-1,1]^d$. For
293 $d = 1, 2, 3$, we show in Figure 2 the sample variance of 1000 realizations of the variance of
294 $\|\Xi_{A,\mathrm{Poi}}(\theta_\star)\|^2$ (white dots) and $\|\Xi_{A,\mathrm{DPP}}(\theta_\star)\|^2$ (black dots), conditionally on $\mathfrak{D}$. Blue and red
295 dots indicate standard 95% marginal confidence intervals, for indication only. The slope found by
296 maximum likelihood in a linear regression in the log-log scale are indicated as legend. The experiment
297 confirms that σ^2 is smaller for the DPP, decays as $p^{-1-1/d}$, and that this decay starts at a batchsize p
298 that increases with dimension.

Figure 1: Summary of the performance of two sampling strategies in SGD.

Figure 2: Summary of the variance decay results.

6 Discussion

In this work, we introduced an orthogonal polynomial-based DPP paradigm for sampling minibatches in SGD that entails variance reduction in the resulting gradient estimator. We substantiated our proposal by detailed theoretical analysis and numerical experiments. Our work raises natural questions and leaves avenues for improvement in several directions. These include the smoothed estimator $\Xi_{A,\mathrm{s}}$, which calls for further investigation in order to be deployed as a computationally attractive procedure; improvement in the dimension dependence of the fluctuation exponent when the gradients are smooth enough, like [29, 30] did for [9]; sharpening of the regularity hypotheses for our theoretical investigations to obtain a more streamlined analysis. While our estimators were motivated by a continuous underlying data distribution, our experiments suggest notably good performance in situations like logistic regression, where the data is at least partially discrete. Extensions to account for discrete settings in a principled manner, via discrete OPEs or otherwise, would be a natural topic for future research. Finally, while our results already partly apply to more sophisticated gradient estimators like Polyak-Rupert averaged gradients [3, Section 4.2], it would be interesting to introduce repulsiveness across consecutive SGD iterations to further minimize the variance of averaged estimators. In summary, we believe that the ideas put forward in the present work will motivate a new perspective on improved minibatch sampling for SGD, more generally on estimators based on linear statistics (e.g. in coreset sampling), and beyond.

References

[1] S. Shalev-Shwartz and S. Ben-David. *Understanding machine learning: From theory to algorithms*. Cambridge university press, 2014.

[2] H. Robbins and S. Monro. A stochastic approximation method. *The Annals of Mathematical Statistics*, pages 400–407, 1951.

[3] É. Moulines and F. Bach. Non-asymptotic analysis of stochastic approximation algorithms for machine learning. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 24, pages 451–459, 2011.

[4] C. Zhang, H. Kjellstrom, and S. Mandt. Determinantal point processes for mini-batch diversification. In *Uncertainty in Artificial Intelligence (UAI)*, 2017.

[5] O. Macchi. The coincidence approach to stochastic point processes. *Advances in Applied Probability*, 7:83–122, 1975.

[6] F. Lavancier, J. Møller, and E. Rubak. Determinantal point process models and statistical inference. *Journal of the Royal Statistical Society, Series B*, 2014.

[7] A. Kulesza and B. Taskar. *Determinantal Point Processes for Machine Learning*, volume 5 of *Foundations and Trends in Machine Learning*. Now Publishers Inc., 2012.

[8] C. Zhang, H. Kjellstrom, and S. Mandt. Active mini-batch sampling using repulsive point processes. In *Proceedings of the AAAI conference on Artificial Intelligence*, 2019.

[9] R. Bardenet and A. Hardy. Monte Carlo with determinantal point processes. *Annals of Applied Probability*, 2020.

[10] A. Nikolov, M. Singh, and U. T. Tantipongpipat. Proportional volume sampling and approximation algorithms for a-optimal design. *arXiv:1802.08318*, 2018.

[11] M. Derezinski and M. K. Warmuth. Unbiased estimates for linear regression via volume sampling. In *Advances in Neural Information Processing Systems (NeurIPS)*. Curran Associates, Inc., 2017.

[12] A. Poinas and R. Bardenet. On proportional volume sampling for experimental design in general spaces. *arXiv preprint arXiv:2011.04562*, 2020.

[13] M. Derezinski and M. W. Mahoney. Determinantal point processes in randomized numerical linear algebra. *Notices of the American Mathematical Society*, 68(1), 2021.

[14] A. Belhadji, R. Bardenet, and P. Chainais. A determinantal point process for column subset selection. *Journal of Machine Learning Research (JMLR)*, 2020.

[15] N. Tremblay, S. Barthelmé, and P.-O. Amblard. Determinantal point processes for coresets. *Journal of Machine Learning Research*, 20(168):1–70, 2019.

[16] J. B. Hough, M. Krishnapur, Y. Peres, and B. Virág. Determinantal processes and independence. *Probability surveys*, 2006.

[17] D. J. Daley and D. Vere-Jones. *An Introduction to the Theory of Point Processes*, volume 1. Springer, 2nd edition, 2003.

[18] A. Soshnikov. Determinantal random point fields. *Russian Mathematical Surveys*, 55:923–975, 2000.

[19] A. Soshnikov. Gaussian limit for determinantal random point fields. *Annals of Probability*, 30(1):171–187, 2002.

[20] S. Ghosh. Determinantal processes and completeness of random exponentials: the critical case. *Probability Theory and Related Fields*, 163(3):643–665, 2015.

[21] J. A. Gillenwater. *Approximate inference for determinantal point processes*. PhD thesis, University of Pennsylvania, 2014.

[22] M. P. Wand and M. C. Jones. *Kernel smoothing*. CRC press, 1994.

[23] C. P. Robert and G. Casella. *Monte Carlo statistical methods*. Springer, 2004.

[24] R. Bardenet and S. Ghosh. Learning from DPPs via Sampling: Beyond HKPV and symmetry. *arXiv preprint arXiv:2007.04287*, 2020.

[25] S. S. Vempala. *The random projection method*, volume 65. American Mathematical Society, 2005.

[26] A. Benveniste, M. Métivier, and P. Priouret. *Adaptive algorithms and stochastic approximations*. Springer Science & Business Media, 1990.

[27] B. Simon. The christoffel–darboux kernel. In *Perspectives in PDE, Harmonic Analysis and Applications*, volume 79 of *Proceedings of Symposia in Pure Mathematics*, pages 295–335, 2008.

[28] G. Gautier, R. Bardenet, G. Polito, and M. Valko. DPPy: Sampling determinantal point processes with Python. *Journal of Machine Learning Research; Open Source Software (JMLR MLOSS)*, 2019.

[29] A. Belhadji, R. Bardenet, and P. Chainais. Kernel quadrature with determinantal point processes. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2019.

[30] A. Belhadji, R. Bardenet, and P. Chainais. Kernel interpolation with continuous volume sampling. In *International Conference on Machine Learning (ICML)*, 2020.