Convergence of score-based generative modeling for general data distributions

Abstract

We give polynomial convergence guarantees for denoising diffusion models that do not rely on the data distribution satisfying functional inequalities or strong smoothness assumptions. Assuming a L^2-accurate score estimate, we obtain Wasserstein distance guarantees for any distributions of bounded support or sufficiently decaying tails, as well as TV guarantees for distributions with further smoothness assumptions.

1 Introduction

Diffusion models have gained huge popularity in recent years in machine learning, as a method to learn and generate new samples from a data distribution. Score-based generative modeling (SGM), as a particular kind of diffusion model, uses a learned score function to transform white noise to the data distribution by following a stochatic differential equation. While SGM's have achieved state-of-the-art performance for artificial image and audio generation [SE19; Dat+19; Gra+19; SE20; Son+20; Men+21; Son+21b; Son+21a; Ram+22; Jin+22], our theoretical understanding of these models is still under development.

There have been some recent progress in theoretical guarantees of SGM [LLT22; De +21]; however, these works either suffer from exponential dependence on parameters or rely on strong assumptions on the data distribution such as smoothness or functional inequalities, which are rarely satisfied in practical situations. Indeed, one of the most successful applications of diffusion model is conditional generation of images based on given text, for which the data distribution is multimodal and hence does not fit into the existing theory. In this work, we aim to develop theoretical guarantees (with polynomial complexity) for SGM for a more general class of distributions.

1.1 Prior work on convergence guarantees

We highlight two recent works. [LLT22] is the first to give polynomial convergence guarantees in TV distance under L^2-accurate score error for a reasonable family of distributions. They introduce a framework to reduce the analysis under L^2-accurate score error to L^∞-score error. However, they rely on the data distribution satisfying smoothness conditions and a log-Sobolev inequality—a strong assumption which essentially limits the guarantees to unimodal distributions. [De 22] gives convergence in Wasserstein distance for distributions with bounded support $\mathcal{M}$. In particular, this covers the case of distributions supported on lower-dimensional manifolds, where guarantees in TV distance are unattainable. However, their guarantees have dependence $\exp(O(\operatorname{diam}(\mathcal{M})^2/\varepsilon))$ for general distributions, and an improved, but still exponential dependence $\exp(\widetilde{O}(\Gamma))$ for distributions with Hessian bounded by Γ/σ_t^2.

We note that other works also analyze the generalization error of a learned score estimate [BMR20; De 22], and without further assumptions, learning an L^2-accurate score estimate requires a number

35 of samples exponential in the dimension. This is beyond the scope of our paper, as we assume that an
36 L^2-accurate score estimate is obtainable.

37 Upon finishing our paper, we learned of a concurrent and independent work [Che+22] which obtained
38 theoretical guarantees for score-based generative modeling under similarly general assumptions on
39 the data distribution.

1.2 Our contributions

41 In this work, we analyze convergence in the most general setting of distributions of bounded support,
42 as in [De 22]. We give Wasserstein bounds for *any* distribution of bounded support (or sufficiently
43 decaying tails), and TV bounds for distributions under smoothness assumptions, that are polynomial
44 in all parameters, and do not rely on the data distribution satisfying any functional inequality.

45 We streamline the analysis of [LLT22] and make a key improvement to their χ^2-based analysis by
46 bounding a certain KL divergence without relying on a global functional inequality. We use a key
47 lemma that distriutions which are close in χ^2-divergence have score functions that are close in L^2,
48 which may be of independent interest.

2 Problem setting and main results

50 A general framework for many score-based generative models is where noise is injected into a data
51 distribution P_{data} via a forward SDE [Son+20]

$$d\widetilde{x}_t = f(\widetilde{x}_t, t)\, dt + g(t)\, dw_t, \quad t \in [0, T], \tag{1}$$

52 where $\widetilde{x}_0 \sim \widetilde{P}_0 := P_{\text{data}}$. Let $\widetilde{p}_t$ denote the density of $\widetilde{x}_t$. Remarkably, $\widetilde{x}_t$ also satisfies a reverse-time
53 SDE,

$$d\widetilde{x}_t = [f(\widetilde{x}_t, t) - g(t)^2 \nabla \ln \widetilde{p}_t(\widetilde{x}_t)]\, dt + g(t)\, d\widetilde{w}_t, \quad t \in [0, T], \tag{2}$$

54 where $\widetilde{w}_t$ is a backward Brownian motion [And82].

55 Of course, in practice the score function $\nabla \ln \widetilde{p}_t$ is not available, and has to be learned from em-
56 pirical samples $\widetilde{x}_t$ from the forward SDE starting from the data samples. Denoising auto-encoders
57 are often used for the score function estimate, which will result in a guarantee that the L^2 error
58 $\mathbb{E}_{\widetilde{p}_t}[\|\nabla \ln \widetilde{p}_t(x) - s(x, t)\|^2]$ is small [Vin11]. The natural question is the following:

59   Given L^2-error bounds of the score function, how close is the distribution generated
60   by (2) to the data distribution P_{data}?

61 We note it is more realistic to consider L^2 rather than L^∞ error, and this makes the analysis more
62 challenging. In addition, we would hope to get a result with as few structural assumptions as possible
63 on P_{data}, so that it can be useful in realistic scenarios where SGM is applied.

64 To state the problem more precisely, we will consider specifically denoising diffusion probabilis-
65 tic modeling (DDPM) [Son+20], where in the forward SDE, $f(x, t) = -\frac{1}{2}g(t)^2 x$ for some non-
66 decreasing function g to be chosen. The forward process is an Ornstein-Uhlenbeck process with time
67 rescaling: $\widetilde{x}_t$ has the same distribution as

$m_t \widetilde{x}_0 + \sigma_t z$, where

$$m_t = \exp\left[-\frac{1}{2}\int_0^t g(s)^2\, ds\right], \ \sigma_t^2 = 1 - \exp\left[-\int_0^t g(s)^2\, ds\right], \ \text{and } z \sim N(0, I). \tag{3}$$

68 Given an estimate score function $s(x, t)$ approximating $\nabla \ln \widetilde{p}_t(x)$, we can simulate the reverse
69 process (reparameterizing $t \leftarrow T - t$ and denoting $p_t := \widetilde{p}_{T-t}$)

$$dx_t = \frac{1}{2}g(T - t)^2\, (x_t + 2\nabla \ln p_t(x_t))\, dt + g(T - t)\, dw_t \tag{4}$$

70 with the exponential integrator discretization [ZC22], where $h_k = t_{k+1} - t_k$ and $\eta_{k+1} \sim N(0, I_d)$:

$$z_{t_{k+1}} = z_{t_k} + \gamma_{1,k}(z_{t_k} + 2s(T - t_k, z_{t_k})) + \sqrt{\gamma_{2,k}} \cdot \eta_{k+1}, \tag{5}$$

$$\text{where } \gamma_{1,k} = \exp\left[\frac{1}{2}G_{t_k, t_{k+1}}\right] - 1, \ \gamma_{2,k} = \exp\left[G_{t_k, t_{k+1}}\right] - 1, \ \text{and } G_{t', t} := \int_{t'}^t g(T - s)^2\, ds. \tag{6}$$

71 We initialize z_0 with a prior distribution that approximates $p_0 = \widetilde{p}_T$ for sufficiently large T:

$$z_0 \sim p_{\text{prior}} := N(0, \sigma_T^2 I_d) \approx N(0, I_d). \tag{7}$$

72 While we focus on DDPM, we note that the continuous process underlying DDPM is equivalent to
73 that of score-matching Langevin diffusion (SMLD) under reparameterization in time and space (see
74 [LLT22, §C.2]). We will further take $g \equiv 1$ for convenience in stating our results.

75 Our goal is to obtain a quantitative guarantee for the distance between the distribution q_{t_K} for z_{t_K}
76 (for appropriate $t_K \approx T$) and P_{data}, under a L^2-score error guarantee. In the following, we assume a
77 sequence of discretization points $0 = t_0 < t_1 < \cdots < t_K \leq T$ has been chosen.

78 **Assumption 1** (L^2 score error)**.** *For any $t \in \{T - t_0, \ldots, T - t_K\}$, the error in the score estimate*
79 *is bounded in $L^2(\widetilde{p}_t)$:*

$$\|\nabla \ln \widetilde{p}_t - s(\cdot, t)\|_{L^2(\widetilde{p}_t)}^2 = \mathbb{E}_{\widetilde{p}_t}[\|\nabla \ln \widetilde{p}_t(x) - s(x, t)\|^2] \leq \varepsilon_t^2 := \frac{\varepsilon_\sigma^2}{\sigma_t^4}.$$

80 We note that the gradient $\nabla \ln \widetilde{p}_t$ grows as $\frac{1}{\sigma_t^2}$ as $t \to 0$, so this is a reasonable assumption, and
81 quantitatively weaker than a uniform bound over t.

82 **Assumption 2** (Bounded support)**.** *P_{data} is supported on $B_R(0) := \{x \in \mathbb{R}^d : \|x\| \leq R\}$.*

83 For simplicity, we assume bounded support in the main text, but note that our results generalize to
84 distributions with sufficiently fast power decay. In the application of image generation, pixel values
85 are bounded, so Assumption 2 is satisfied with R typically on the order of $\sqrt{d}$.

86 These are the only assumptions we need to obtain a polynomial complexity guarantee. We also
87 consider the following stronger smoothness assumption, which is Assumption A.6 in [De 22] and
88 will give better dependencies. Note that [De 22, Theorem I.8] shows a (nonuniform) version of
89 Assumption 3 holds when p_0 is a smooth density on a convex submanifold.

90 **Assumption 3.** *The following bound of the Hessian of the log-pdf holds for any $t > 0$ and x:*

$$\left\|\nabla^2 \ln p_t(x)\right\| \leq \frac{C}{\sigma_t^2},$$

91 *for some constant $C > 0$.*

92 Finally, the following smoothness assumption on $\widetilde{p}_0$ will allow us to obtain TV guarantees.

93 **Assumption 4.** *P_{data} admits a density $\widetilde{p}_0 \propto e^{-V(x)}$ where $V(x)$ is L-smooth.*

Algorithm 1 DDPM with exponential integrator [Son+20; ZC22]

INPUT: Time T; discretization points $0 = t_0 < t_1 < \cdots < t_N \leq T$; score estimates $s(\cdot, T - t_k)$;
radius R; function g (default: $g \equiv 1$)
Draw $z_0 \sim p_{\text{prior}}$ from the prior distribution p_{prior} given by (7).
for k from 1 to N **do**
    Compute z_{t_k} from $z_{t_{k-1}}$ using (5).
end for
Let $\widehat{z}_{t_N} = z_{t_N}$ if $z_{t_N} \in B_R(0)$; otherwise, let $\widehat{z}_{t_N} = 0$.

---

94 **Theorem 2.1** (Wasserstein+TV error for distributions with bounded support)**.** *Suppose that As-*
95 *sumption 1 and 2 hold with $R \geq \sqrt{d}$. Then there is a sequence of discretization points*
96 *$0 = t_0 < t_1 < \cdots < t_N < T$ with $N = O(\text{poly}(d, R, 1/\varepsilon_W))$ such that if $\varepsilon_\sigma = \widetilde{o}\left(\frac{\varepsilon_{\text{TV}}^{6.5} \varepsilon_W^5}{R^9 d^{2.25}}\right)$,*
97 *then the distribution q_{t_N} of the output z_{t_N} of DDPM is ε_{TV}-close in TV distance to a distribution*
98 *that is ε_W in W_2 distance from P_{data}. If in addition Assumption 3 holds with $C \geq R^2$, it suffices for*
99 *$\varepsilon_\sigma = \widetilde{o}\left(\frac{\varepsilon_{\text{TV}}^4}{C^2 d}\right)$ (note that the $\widetilde{o}(\cdot)$ hides logarithmic dependence on ε_W).*

100 This result is perhaps surprising at first glance, as it is well known that for sampling algorithms such
101 as Langevin Monte Carlo, structural assumptions on the target distribution—such as a log-Sobolev
102 inequality—are required to obtain similar theoretical guarantees, even with the knowledge of the

103 exact score function. The key reason that we can do better is that we utilize a *sequence* of score
104 functions s_t along the reverse SDE, which is not available in standard sampling settings. To some
105 extent, this result shows the power of DDPM and other reverse SDE-based methods compared with
106 generative modeling based on standard Langevin Monte Carlo.

107 A statement with more precise dependencies, and which works for unbounded distributions with
108 sufficiently decaying tails, can be found as Theorem D.2. We note that under the Hessian bound
109 (Assumption 3), up to logarithmic factors, the same score error bound suffices to obtain a fixed TV
110 distance to a distribution arbitrarily close in W_2 distance. By truncating the resulting distribution, we
111 can also obtain purely Wasserstein error bounds.

112 **Theorem 2.2** (Wasserstein error for distributions with bounded support). *In the same setting as*
113 *Theorem 2.1, consider the distribution $\widehat{q}_{t_N}$ of the truncated output $\widehat{x}_{t_N}$ of DDPM. If Assumptions 1*
114 *and 2 hold with $R \geq \sqrt{d}$ and $\varepsilon_\sigma = \widetilde{o}\left(\frac{\varepsilon_W^{18}}{R^{22}d^{2.25}}\right)$, then with appropriate (polynomial) choice*
115 *of parameters, $W_2(\widehat{q}_{t_K}, P_{\text{data}}) \leq \varepsilon_W$. If in addition Assumption 3 holds with $C \geq R^2$, then*
116 $\varepsilon_\sigma = \widetilde{o}\left(\frac{\varepsilon_W^8}{C^2R^8d}\right)$ *suffices.*

117 With an extra assumption on the smoothness of P_{data}, we can also obtain purely TV error bounds:

118 **Theorem 2.3** (TV error for distributions under smoothness assumption). *Suppose that Assumptions 1*
119 *and 4 hold, P_{data} is subexponential (with a fixed constant), and denote $R = \max\left\{\sqrt{d}, \mathbb{E}_{P_{data}}\|X\|\right\}$.*
120 *Then there is a sequence of discretization points $0 = t_0 < t_1 < \cdots < t_N < T$ with $N =$*
121 $O(\text{poly}(d, R, 1/\varepsilon_{\text{TV}}))$ *such that if $\varepsilon_\sigma = \widetilde{o}\left(\frac{\varepsilon_{\text{TV}}^{11.5}}{R^{14}d^{2.25}L^5}\right)$, then $TV(q_{t_N}, P_{\text{data}}) \leq \varepsilon_{\text{TV}}$. If in addition*
122 *Assumption 3 holds with $C \geq R^2$, then $\varepsilon_\sigma = \widetilde{o}\left(\frac{\varepsilon_{\text{TV}}^4}{C^2d}\right)$ suffices.*

123 A more precise statement can be found as Theorem D.3, which also works more generally with
124 sufficient tail decay. We note that this result can be derived directly by combining Theorem D.2 and a
125 TV error bound between P_{data} and p_{t_N} (Lemma C.4) depending on the smoothness of P_{data}.

3 Proof overview

127 Our proof uses the framework by [LLT22] to convert guarantees under L^∞-accurate score function to
128 under L^2-accurate score function. For the analysis under L^∞-accurate score function, we interpolate
129 the discrete process with estimated score, $z_t \sim q_t$, and derive a differential inequality

$$\frac{d}{dt}\chi^2(q_t\|p_t) = -g(T-t)^2\mathscr{E}_{p_t}\left(\frac{q_t}{p_t}\right) + 2\mathbb{E}\left[\left\langle g(T-t)^2(s(z_{t_k}, T-t_k) - \nabla\ln p_t(z_t), \nabla\frac{q_t(x)}{p_t(x)}\right\rangle\right].$$

130 We bound resulting error terms, making ample use of the Donsker-Varadhan variational principle to
131 convert expectations to be under p_t. Under small enough step sizes, this shows that $\chi^2(q_t\|p_t)$ grows
132 slowly (Theorem A.10), which suffices as χ^2-divergence decays exponentially in the forward process.

133 Our main innovation over the analysis of [LLT22] is bounding a KL divergence term $\text{KL}(q_t\psi_t\|p_t)$
134 without a global log-Sobolev inequality for p_t. We note that it suffices for p_t to be a mixture of
135 distributions each satisfying a log-Sobolev inequality, with the logarithm of the minimum mixture
136 weight bounded below, and in Lemma B.2, we show that we can decompose any distributed of
137 bounded support in this manner if we move a small amount of its mass. In Section C, we show that
138 this does not significantly affect the estimate of the score estimate, by interpreting the score function
139 as solving a Bayesian inference problem. We note the key Lemma C.5 may be of independent interest.

140 Finally, we reduce from the L^2 to L^∞ setting by bounding the probabilities of hitting a bad set where
141 the score error is large, and carefully choose parameters (Section D). This gives a TV error bound to
142 $\widetilde{p}_\delta$—the forward distribution at small positive time. Finally, we can bound the Wasserstein distance
143 of $\widetilde{p}_\delta$ to $\widetilde{P}_0$ (in the general case) or the TV distance (under additional smoothness of $\widetilde{P}_0$.)

144 In Section E we show that the Hessian is always bounded by $O\left(\frac{d}{\sigma_t^2}\right)$ with high probability (cf.
145 Assumption 3). We speculate that a high-probability rather than uniform bound on the Hessian (as in
146 Lemma A.13) can be used to obtain better dependencies, and leave this as an open problem.

References

[And82] Brian DO Anderson. "Reverse-time diffusion equation models". In: *Stochastic Processes and their Applications* 12.3 (1982), pp. 313–326.

[BMR20] Adam Block, Youssef Mroueh, and Alexander Rakhlin. "Generative modeling with denoising auto-encoders and Langevin sampling". In: *arXiv preprint arXiv:2002.00107* (2020).

[CCN21] Hong-Bin Chen, Sinho Chewi, and Jonathan Niles-Weed. "Dimension-free log-Sobolev inequalities for mixture distributions". In: *Journal of Functional Analysis* 281.11 (2021), p. 109236.

[Che+21] Sinho Chewi, Murat A Erdogdu, Mufan Bill Li, Ruoqi Shen, and Matthew Zhang. "Analysis of Langevin Monte Carlo from Poincaré to Log-Sobolev". In: *arXiv preprint arXiv:2112.12662* (2021).

[Che+22] Sitan Chen, Sinho Chewi, Jerry Li, Yuanzhi Li, Adil Salim, and Anru R. Zhang. *Sampling is as easy as learning the score: theory for diffusion models with minimal data assumptions*. arXiv:2209.11215. 2022.

[Dat+19] Sumanth Dathathri, Andrea Madotto, Janice Lan, Jane Hung, Eric Frank, Piero Molino, Jason Yosinski, and Rosanne Liu. "Plug and play language models: A simple approach to controlled text generation". In: *arXiv preprint arXiv:1912.02164* (2019).

[De +21] Valentin De Bortoli, James Thornton, Jeremy Heng, and Arnaud Doucet. "Diffusion Schrödinger bridge with applications to score-based generative modeling". In: *Advances in Neural Information Processing Systems* 34 (2021).

[De 22] Valentin De Bortoli. "Convergence of denoising diffusion models under the manifold hypothesis". In: *arXiv preprint arXiv:2208.05314* (2022).

[EHZ21] Murat A. Erdogdu, Rasa Hosseinzadeh, and Matthew S. Zhang. "Convergence of Langevin Monte Carlo in Chi-Squared and Renyi Divergence". In: *arXiv preprint arXiv:2007.11612* (2021).

[Gra+19] Will Grathwohl, Kuan-Chieh Wang, Jörn-Henrik Jacobsen, David Duvenaud, Mohammad Norouzi, and Kevin Swersky. "Your classifier is secretly an energy based model and you should treat it like one". In: *arXiv preprint arXiv:1912.03263* (2019).

[Jin+22] Bowen Jing, Gabriele Corso, Renato Berlinghieri, and Tommi Jaakkola. "Subspace Diffusion Generative Models". In: *arXiv preprint arXiv:2205.01490* (2022).

[LLT22] Holden Lee, Jianfeng Lu, and Yixin Tan. "Convergence for score-based generative modeling with polynomial complexity". In: *arXiv preprint arXiv:2206.06227* (2022).

[LM00] Beatrice Laurent and Pascal Massart. "Adaptive estimation of a quadratic functional by model selection". In: *Annals of Statistics* (2000), pp. 1302–1338.

[Men+21] Chenlin Meng, Yutong He, Yang Song, Jiaming Song, Jiajun Wu, Jun-Yan Zhu, and Stefano Ermon. "SDEdit: Guided image synthesis and editing with stochastic differential equations". In: *International Conference on Learning Representations*. 2021.

[Ram+22] Aditya Ramesh, Prafulla Dhariwal, Alex Nichol, Casey Chu, and Mark Chen. "Hierarchical text-conditional image generation with clip latents". In: *arXiv preprint arXiv:2204.06125* (2022).

[SE19] Yang Song and Stefano Ermon. "Generative Modeling by Estimating Gradients of the Data Distribution". In: *Proceedings of the 33rd Annual Conference on Neural Information Processing Systems*. 2019.

[SE20] Yang Song and Stefano Ermon. "Improved techniques for training score-based generative models". In: *arXiv preprint arXiv:2006.09011* (2020).

[Son+20] Yang Song, Jascha Sohl-Dickstein, Diederik P Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. "Score-Based Generative Modeling through Stochastic Differential Equations". In: *International Conference on Learning Representations*. 2020.

[Son+21a] Yang Song, Conor Durkan, Iain Murray, and Stefano Ermon. "Maximum likelihood training of score-based diffusion models". In: *Advances in Neural Information Processing Systems* 34 (2021).

[Son+21b] Yang Song, Liyue Shen, Lei Xing, and Stefano Ermon. "Solving Inverse Problems in Medical Imaging with Score-Based Generative Models". In: *arXiv preprint arXiv:2111.08005* (2021).

202 [Ver18]   Roman Vershynin. *High-dimensional probability: An introduction with applications in*
203   *data science*. Vol. 47. Cambridge university press, 2018.
204 [Vin11]   Pascal Vincent. "A connection between score matching and denoising autoencoders".
205   In: *Neural computation* 23.7 (2011), pp. 1661–1674.
206 [ZC22]   Qinsheng Zhang and Yongxin Chen. "Fast Sampling of Diffusion Models with Expo-
207   nential Integrator". In: *arXiv preprint arXiv:2204.13902* (2022).

Notation and definitions

We let $\widetilde{p}_t$ denote the density of $\widetilde{x}_t$ under the forward process (1). Note that $x_0 \sim \widetilde{P}_0$ may not admit a density, but $\widetilde{x}_t$ will for $t > 0$. For the reverse process, we use the notation $p_t = \widetilde{p}_{T-t}$, $x_t = \widetilde{x}_{T-t}$. We defined m_t and σ_t in (3),

$$m_t = \exp\left[-\frac{1}{2}\int_0^t g(s)^2\,ds\right], \qquad \sigma_t^2 = 1 - \exp\left[-\int_0^t g(s)^2\,ds\right],$$

and note that $\widetilde{p}_t = (M_{m_t \sharp} \widetilde{P}_0) * \varphi_{\sigma_t^2}$, where $M_m(x) = mx$ denotes multiplication by m, $F_\sharp P$ denotes the pushforward of the measure P by F, and φ_{σ^2} is the density of $N(0, \sigma^2 I_d)$. When $g \equiv 1$, we note the bound $\sigma_t^2 \le \min\{1, t\}$ and $\sigma_t^2 = \Theta(\min\{1, t\})$.

We will let z_t denote the (interpolated) discrete process (see (12)) and let q_t be the density of z_t. We define

$$\phi_t(x) = \frac{q_t(x)}{p_t(x)}, \qquad \psi_t(x) = \frac{\phi_t(x)}{\mathbb{E}_{p_t}\phi_t^2}, \tag{8}$$

and note that $q_t\psi_t$ is a probability density. We defined $G_{t',t} = \int_{t'}^t g(T-s)^2\,ds$ in (6).

We denote the estimated score function by either $s(x,t)$ and $s_t(x)$ interchangeably.

A random variable X is subgaussian with constant C if

$$C = \|X\|_{\psi_2} := \inf\left\{t > 0 : \mathbb{E}\exp(X^2/t^2) \le 2\right\} < \infty.$$

A $\mathbb{R}^d$-valued random variable X is subgaussian with constant C if for all $v \in \mathbb{S}^{d-1}$, $\langle X, v\rangle$ is subgaussian. We also define

$$\|X\|_{2,\psi_2} := \|\|X\|_2\|_{\psi_2}.$$

Given a probability measure P on $\mathbb{R}^d$ with density p, the associated Dirichlet form is $\mathscr{E}_p(f,g) := \int_{\mathbb{R}^d} \langle \nabla f, \nabla g\rangle\, P(dx) = \int_{\mathbb{R}^d} \langle \nabla f, \nabla g\rangle\, p(x)\,dx$; denote $\mathscr{E}_p(f) = \mathscr{E}_p(f,f)$. we say that a log-Sobolev inequality (LSI) holds with constant C_{LS} if for any probability density q,

$$\mathrm{KL}(q\|p) \le \frac{C_{\mathrm{LS}}}{2}\mathscr{E}_p\left(\frac{q}{p}, \ln\frac{q}{p}\right) = \frac{C_{\mathrm{LS}}}{2}\int_{\mathbb{R}^d}\left\|\nabla\ln\frac{q(x)}{p(x)}\right\|^2 q(x)\,dx. \tag{9}$$

Note $\int_{\mathbb{R}^d}\left\|\nabla\ln\frac{q(x)}{p(x)}\right\|^2 q(x)\,dx$ is also known as the Fisher information of q with respect to p. Alternatively, defining the entropy by $\mathrm{Ent}_p(f) = \mathbb{E}_p f(x)\ln f(x) - \mathbb{E}_p f(x)\ln\mathbb{E}_p f(x)$, for any $f \ge 0$,

$$\mathrm{Ent}_p(f) \le \frac{C_{\mathrm{LS}}}{2}\mathscr{E}_p(f, \ln f) = \frac{C_{\mathrm{LS}}}{2}\int_{\mathbb{R}^d}\|\nabla\ln f(x)\|^2 f(x)p(x)dx. \tag{10}$$

A   DDPM with L^∞-accurate score estimate

We consider the error between the exact backwards SDE (4) and the exponential integrator with estimated score (5). In this section, we bound the error assuming that the score estimate s is accurate in L^∞.

Assumption 5 (L^∞ score error). *For any $t \in \{T - t_0, \ldots, T - t_K\}$, the error in the score estimate is bounded:*

$$\|\nabla\ln\widetilde{p}_t - s(\cdot, t)\|_\infty = \sup_{x\in\mathbb{R}^d}\|\nabla\ln\widetilde{p}_t(x) - s(x,t)\| \le \varepsilon_{\infty,t}^2 \tag{11}$$

for some non-decreasing function $\varepsilon_{\infty,t}^2$.

234 In Section D, we will relax this condition to score function being accurate in L^2.

235 First, we construct the following continuous-time process which interpolates the discrete-time
236 process (5), for $t \in [t_k, t_{k+1}]$:

$$dz_t = g(T-t)^2 \left(\frac{1}{2} z_t + s(z_{t_k}, T - t_k) \right) dt + g(T-t) \, dw_t. \tag{12}$$

237 Integration gives that

$$z_t - z_{t_k} = \left(\exp\left(\frac{1}{2} G_{t_k,t} \right) - 1 \right) (z_{t_k} + 2s(z_{t_k}, T - t_k))$$

$$+ \int_{t_k}^{t} \exp\left(\frac{1}{2} \int_{t_k}^{t'} g(T - t'')^2 \, dt'' \right) g(t') \, dw_{t'}, \tag{13}$$

238 where $G_{t',t}$ is defined in (6).

239 Letting q_t be the distribution of z_t and p_t be the distribution of x_t, we have by [LLT22, Lemma A.2]
240 that

$$\frac{d}{dt} \chi^2(q_t \| p_t) = -g(T-t)^2 \mathscr{E}_{p_t}\left(\frac{q_t}{p_t} \right) + 2\mathbb{E}\left[\left\langle g(T-t)^2(s(z_{t_k}, T - t_k) - \nabla \ln p_t(z_t), \nabla \frac{q_t(x)}{p_t(x)} \right\rangle \right]. \tag{14}$$

241 (Note that in our case, $\widehat{f}$ also depends on z_t rather than just z_{t_k}, but this does not change the
242 calculation.) Define ϕ_t, ψ_t as in (8): $\phi_t(x) = \frac{q_t(x)}{p_t(x)}$, $\psi_t(x) = \frac{\phi_t(x)}{\mathbb{E}_{p_t} \phi_t^2}$.

243 To bound (14), we use the following lemma.

244 **Lemma A.1** (cf. [EHZ21, Lemma 1], [LLT22, Lemma A.3]). *For any $C > 0$ and any $\mathbb{R}^d$-valued*
245 *random variable u, we have*

$$\mathbb{E}\left[\left\langle u, \nabla \frac{q_t(z_t)}{p_t(z_t)} \right\rangle \right] \leq C \cdot (\chi^2(q_t \| p_t) + 1) \cdot \mathbb{E}\left[\|u\|^2 \, \psi_t(z_t) \right] + \frac{1}{4C} \mathscr{E}_{p_t}\left(\frac{q_t}{p_t} \right).$$

246 *Proof.* By Young's inequality,

$$\mathbb{E}\left[\left\langle u, \nabla \frac{q_t(z_t)}{p_t(z_t)} \right\rangle \right] = \mathbb{E}\left[\left\langle u \sqrt{\frac{q_t(z_t)}{p_t(z_t)}}, \sqrt{\frac{p_t(z_t)}{q_t(z_t)}} \nabla \frac{q_t(z_t)}{p_t(z_t)} \right\rangle \right]$$

$$\leq C\mathbb{E}\left[\|u\|^2 \frac{q_t(z_t)}{p_t(z_t)} \right] + \frac{1}{4C} \mathbb{E}_{p_t}\left[\left\| \nabla \frac{q_t(x)}{p_t(x)} \right\|^2 \right]$$

$$= C\mathbb{E}_{p_t} \phi_t^2 \cdot \mathbb{E}\left[\|u\|^2 \, \psi_t(z_t) \right] + \frac{1}{4C} \mathscr{E}_{p_t}\left(\frac{q_t}{p_t} \right)$$

$$= C(\chi^2(q_t \| p_t) + 1) \cdot \mathbb{E}\left[\|u\|^2 \, \psi_t(z_t) \right] + \frac{1}{4C} \mathscr{E}_{p_t}\left(\frac{q_t}{p_t} \right). \qquad \square$$

247 **Lemma A.2.** *Suppose that (11) holds for $t = T - t_k$, $\nabla \ln p_{t_k}(x)$ is L_{T-t_k}-Lipschitz, g is non-*
248 *decreasing, and that $h_k \leq \frac{1}{20 L_{T-t_k} g(T-t_k)^2}$. Then we have for $t \in [t_k, t_{k+1}]$ that*

$$\frac{d}{dt} \chi^2(q_t \| p_t) \leq -\frac{1}{2} g(T-t)^2 \mathscr{E}_{p_t}\left(\frac{q_t}{p_t} \right) + 12 g(T-t)^2 (\chi^2(q_t \| p_t) + 1) \cdot$$

$$\left[\varepsilon_{\infty, T-t_k}^2 + 16 G_{t_k,t}^2 L_{T-t_k}^2 \left[\mathbb{E}[\psi_t(z_t) \|z_t\|^2] + 16\mathbb{E}[\psi_t(z_t) \|\nabla \ln p_t(x_t)\|^2] \right] \right.$$

$$\left. + 64 G_{t_k,t} L_{T-t_k}^2 (8 \, \mathrm{KL}(\psi_t q_t \| p_t) + 2d + 16 \ln 2) + \mathbb{E}\left[\|\nabla \ln p_{t_k}(z_t) - \nabla \ln p_t(z_t)\|^2 \, \psi_t(z_t) \right] \right].$$

249 *Proof.* We bound the second term on the RHS of (14). By Lemma A.1,

$$\mathbb{E}\left[\left\langle s(z_{t_k}, T - t_k) - \nabla \ln p_t(z_t), \nabla \frac{q_t(z_t)}{p_t(z_t)} \right\rangle\right]$$

$$\leq (\chi^2(q_t\|p_t) + 1)\mathbb{E}\left[\|s(z_{t_k}, T - t_k) - \nabla \ln p_t(z_t)\|^2 \psi_t(z_t)\right] + \frac{1}{4}\mathcal{E}_{p_t}\left(\frac{q_t}{p_t}\right). \tag{15}$$

250 Now

$$\|s(z_{t_k}, T - t_k) - \nabla \ln p_t(z_t)\|^2$$

$$\leq 3\left[\|s(z_{t_k}, T - t_k) - \nabla \ln p_{t_k}(z_{t_k})\|^2 + \|\nabla \ln p_{t_k}(z_{t_k}) - \nabla \ln p_{t_k}(z_t)\|^2 + \|\nabla \ln p_{t_k}(z_t) - \nabla \ln p_t(z_t)\|^2\right]$$

$$\leq 3\left[\|s(z_{t_k}, T - t_k) - \nabla \ln p_{t_k}(z_{t_k})\|^2 + L_{T-t_k}^2 \|z_{t_k} - z_t\|^2 + \|\nabla \ln p_{t_k}(z_t) - \nabla \ln p_t(z_t)\|^2\right]$$

251 and

$$\mathbb{E}\left[\|s(z_{t_k}, T - t_k) - \nabla \ln p_{t_k}(z_{t_k})\|^2 \psi_t(z_t)\right] \leq \varepsilon_{\infty, T-t_k}^2$$

252 by definition of $\varepsilon_{\infty,t}$, so by Lemma A.3,

$$\mathbb{E}\left[\|s(z_{t_k}, T - t_k) - \nabla \ln p_t(z_t)\|^2 \psi_t(z_t)\right]$$

$$\leq 3\left[\varepsilon_{\infty, T-t_k}^2 + L_{T-t_k}^2 \mathbb{E}\left[\|z_t - z_{t_k}\|^2 \psi_t(z_t)\right] + \mathbb{E}\left[\|\nabla \ln p_{t_k}(z_t) - \nabla \ln p_t(z_t)\|^2 \psi_t(z_t)\right]\right]$$

$$\leq 3\left[\varepsilon_{\infty, T-t_k}^2 + 16 G_{t_k,t}^2 L_{T-t_k}^2 \left[\mathbb{E}[\psi_t(z_t)\|z_t\|^2] + 4\mathbb{E}[\psi_t(z_t)\|s(z_{t_k}, T - t_k) - \nabla \ln p_t(z_t)\|^2]\right.\right.$$

$$+ 16\mathbb{E}[\psi_t(z_t)\|\nabla \ln p_t(x_t)\|^2]\right] + 64 G_{t_k,t} L_{T-t_k}^2 (8\,\mathrm{KL}(\psi_t q_t\|p_t) + 2d + 16\ln 2)$$

$$+ \mathbb{E}\left[\|\nabla \ln p_{t_k}(z_t) - \nabla \ln p_t(z_t)\|^2 \psi_t(z_t)\right]\Bigg]$$

253 The condition on h_k and the fact that g is non-decreasing implies $192 G_{t_k,t}^2 L_{T-t_k}^2 \leq \frac{1}{2}$. Rearranging
254 gives

$$\mathbb{E}\left[\|s(z_{t_k}, T - t_k) - \nabla \ln p_t(z_t)\|^2 \psi_t(z_t)\right]$$

$$\leq 6\left[\varepsilon_{\infty, T-t_k}^2 + 16 G_{t_k,t}^2 L_{T-t_k}^2 \left[\mathbb{E}[\psi_t(z_t)\|z_t\|^2] + 16\mathbb{E}[\psi_t(z_t)\|\nabla \ln p_t(x_t)\|^2]\right.\right.$$

$$+ 64 G_{t_k,t} L_{T-t_k}^2 (8\,\mathrm{KL}(\psi_t q_t\|p_t) + 2d + 16\ln 2) + \mathbb{E}\left[\|\nabla \ln p_{t_k}(z_t) - \nabla \ln p_t(z_t)\|^2 \psi_t(z_t)\right]\Bigg]$$

255 Substituting into (15) and that inequality into (14) give the conclusion. $\qquad\square$

256 **Lemma A.3.** *Suppose that $h_k \leq \frac{1}{2g(T-t_k)^2}$. Then for $t \in [t_{k+1}, t_k]$,*

$$\mathbb{E}\left[\|z_t - z_{t_k}\|^2 \psi_t(z_t)\right] \leq 16 G_{t_k,t}^2 \left[\mathbb{E}[\psi_t(z_t)\|z_t\|^2] + 4\mathbb{E}[\psi_t(z_t)\|s(z_{t_k}, T - t_k) - \nabla \ln p_t(z_t)\|^2]\right.$$

$$\left.+ 16\mathbb{E}[\psi_t(z_t)\|\nabla \ln p_t(z_t)\|^2]\right] + 64 G_{t_k,t}(8\,\mathrm{KL}(\psi_t q_t\|p_t) + 2d + 16\ln 2).$$

257 *Proof.* Consider (13). The assumption on h_k implies $G_{t_k,t} \leq G_{t_k,t_{k+1}} \leq \frac{1}{2}$, so $\exp\left(\frac{1}{2}G_{t_k,t}\right) - 1 \leq$
258 $G_{t_k,t}$. Let Y denote the last term of (13). Then

$$\|z_t - z_{t_k}\| \leq G_{t_k,t}[\|z_{t_k}\| + 2\|s(z_{t_k}, T - t_k)\|] + \|Y\|$$
$$\leq G_{t_k,t}[\|z_t\| + \|z_{t_k} - z_t\| + 2\|s(z_{t_k}, T - t_k) - \nabla \ln p_t(z_t)\| + 2\|\nabla \ln p_t(z_t)\|] + \|Y\|.$$

259 Again using $G_{t_k,t} \leq \frac{1}{2}$, rearranging gives

$$\|z_t - z_{t_k}\| \leq 2 G_{t_k,t}[\|z_t\| + 2\|s(z_{t_k}, T - t_k) - \nabla \ln p_t(z_t)\| + 4\|\nabla \ln p_t(z_t)\|] + 2\|Y\|,$$

260 and

$$\mathbb{E}\left[\|z_t - z_{t_k}\|^2 \psi_t(z_t)\right] \leq 16 G_{t_k,t}^2 \Big[\mathbb{E}[\psi_t(z_t)\|z_t\|^2] + 4\mathbb{E}[\psi_t(z_t)\|s(z_{t_k}, T - t_k) - \nabla \ln p_t(z_t)\|^2]$$
$$+ 16\mathbb{E}[\psi_t(z_t)\|\nabla \ln p_t(x_t)\|^2]\Big] + 16\mathbb{E}[\psi_t(z_t)\|Y\|^2].$$

261 By Lemma A.4,

$$\mathbb{E}[\psi_t(z_t)\|Y\|^2] \leq 4 G_{t_k,t}(8\,\mathrm{KL}(\psi_t q_t\|p_t) + 2d + 16\ln 2).$$

262 The lemma follows. □

263 **Lemma A.4.** *For $t \in [t_k, t_{k+1}]$,*

$$\mathbb{E}\left[\psi_t(z_t)\left\|\int_{t_k}^t \exp\left(\frac{1}{2}\int_{t_k}^{t'} g(T - t'')^2\, dt''\right) g(t')\, dw_{t'}\right\|^2\right]$$
$$\leq 2(\exp(G_{t_k,t}) - 1)\big(8\,\mathrm{KL}(\psi_t q_t\|p_t) + 2d + 16\ln 2\big).$$

264 *Proof.* Note that $Y := \int_{t_k}^t \exp\left(\frac{1}{2}\int_{t_k}^{t'} g(T - t'')^2\, dt''\right) g(t')\, dw_{t'}$ is a Gaussian random vector with
265 variance

$$\int_{t_k}^t \exp\left(\int_{t_k}^{t'} g(T - t'')^2\, dt''\right) g(t')^2\, dt' \cdot I_d = \exp\left(\int_{t_k}^{t'} g(T - t'')^2\, dt''\right)\Big|_{t'=t_k}^{t'=t} \cdot I_d$$
$$= (\exp(G_{t_k,t}) - 1) \cdot I_d.$$

266 (Note that this calculation shows that the continuous-time process (12) does agree with the discrete-
267 time process (5) at $t = t_{k+1}$.) Using the Donsker-Varadhan variational principle, for any random
268 variable X,

$$\tilde{\mathbb{E}}X \leq \mathrm{KL}(\tilde{\mathbb{P}}\|\mathbb{P}) + \ln \mathbb{E}\exp X.$$

269 Applying this to $X = c\left(\|Y\| - \mathbb{E}\|Y\|\right)^2$ for a constant $c > 0$ to be chosen later, and $\tilde{\mathbb{P}}$ such that
270 $\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}}(z_t) = \psi_t(z_t)$, we can bound

$$\tilde{\mathbb{E}}\|Y\|^2 \leq 2\mathbb{E}\left[\|Y\|^2\right] + 2\tilde{\mathbb{E}}\left[(Y - \mathbb{E}\|Y\|)^2\right]$$
$$\leq 2\mathbb{E}\left[\|Y\|^2\right] + \frac{2}{c}\left[\mathrm{KL}(\tilde{\mathbb{P}}\|\mathbb{P}) + \ln\mathbb{E}\exp\left(c\left(\|Y\| - \mathbb{E}\|Y\|\right)^2\right)\right] \qquad (16)$$
$$\leq 2d(\exp(G_{t_k,t}) - 1) + \frac{2}{c}\left[\mathrm{KL}(\tilde{\mathbb{P}}\|\mathbb{P}) + \ln\mathbb{E}\exp\left(c\left(\|Y\| - \mathbb{E}\|Y\|\right)^2\right)\right]. \qquad (17)$$

271 Now following [Che+21, Theorem 4], we set $c = \frac{1}{8(\exp(G_{t_k,t}) - 1)}$, so that

$$\mathbb{E}\left[\frac{(\|Y\| - \mathbb{E}\|Y\|)^2}{8(\exp(G_{t_k,t}) - 1)}\right] \leq 2.$$

272 Next, we have

$$\mathrm{KL}(\tilde{\mathbb{P}}\|\mathbb{P}) = \mathbb{E}_{\psi_t q_t}\ln\psi_t = \mathbb{E}_{\psi_t q_t}\ln\frac{\phi_t}{\mathbb{E}_{p_t}\phi_t^2} = \frac{1}{2}\mathbb{E}_{\psi_t q_t}\ln\frac{\phi_t^2}{(\mathbb{E}_{p_t}\phi_t^2)^2}$$
$$= \frac{1}{2}\left[\mathbb{E}_{\psi_t q_t}\ln\frac{\phi_t^2}{\mathbb{E}_{p_t}\phi_t^2} - \ln\mathbb{E}_{p_t}\phi_t^2\right] = \frac{1}{2}\left[\mathbb{E}_{\psi_t q_t}\ln\frac{\psi_t q_t}{p_t} - \ln\mathbb{E}_{p_t}\phi_t^2\right].$$

273 Noting that $\mathbb{E}_{p_t}\phi_t^2 = \chi^2(q_t\|p_t) + 1 \geq 1$, we have that

$$\mathrm{KL}(\tilde{\mathbb{P}}\|\mathbb{P}) \leq \frac{1}{2}\,\mathrm{KL}(\psi_t q_t\|p_t).$$

274 Substituting everything into (17) gives the desired inequality. □

Let

$$K_z = \mathbb{E}\left[\psi_t(z_t)\,\|z_t\|^2\right] \tag{18}$$

$$K_V = \mathbb{E}\left[\psi_t(z_t)\,\|\nabla\ln p_t(z_t)\|^2\right] \tag{19}$$

$$K_{\Delta V} = \mathbb{E}\left[\psi_t(z_t)\,\|\nabla\ln p_{t_k}(z_t) - \nabla\ln p_t(z_t)\|^2\right] \tag{20}$$

$$K = \mathrm{KL}(\psi_t q_t \| p_t). \tag{21}$$

In order to bound the RHS in Lemma A.2, we need to bound all four of these quantities, which we do in Lemma A.5, A.6, A.8, and Section B, respectively. The main innovation in our analysis compared to [LLT22] is a new way to bound K, which we present in a separate section.

First we bound K_z. Recall the norm

$$\|X\|_{2,\psi_2} = \inf\left\{L > 0 : \mathbb{E}e^{\frac{\|X\|_2^2}{L^2}} \le 2\right\}.$$

(In other words, this is the usual Orlicz norm applied to $\|X\|_2$.)

Lemma A.5. *For $t \in [t_k, t_{k+1}]$,*

$$\mathbb{E}\left[\psi_t(z_t)\,\|z_t\|^2\right] \le \|x_t\|_{2,\psi_2}^2 \cdot \left[\mathrm{KL}(\psi_t q_t\|p_t) + \ln 2\right].$$

Proof. By the Donsker-Varadhan variational principle,

$$\mathbb{E}\left[\psi_t(z_t)\,\|z_t\|^2\right] = \frac{2}{s}\mathbb{E}_{\psi_t q_t}\left[\frac{s}{2}\,\|x\|^2\right] \le \frac{2}{s}\left[\mathrm{KL}(\psi_t q_t\|p_t) + \ln\mathbb{E}_{p_t}\left[e^{\frac{s}{2}\|x\|^2}\right]\right]$$

for any $s > 0$. Choosing $s = 2\|x_t\|_{2,\psi_2}^{-2}$, we have $\mathbb{E}_{p_t}\left[e^{\frac{s}{2}\|x\|^2}\right] \le 2$, which gives the desired inequality. $\qquad\square$

The following bounds K_V; note that the proof does not depend on the definition of q_t, only that it is a probability density.

Lemma A.6 ([LLT22, Corollary C.7], [Che+21, Lemma 16]).

$$\mathbb{E}\left[\psi_t(z_t)\,\|\nabla\ln p_t(z_t)\|^2\right] \le \frac{4}{\chi^2(q_t\|p_t)+1}\cdot\mathscr{E}_{p_t}\left(\frac{q_t}{p_t}\right) + 2dL.$$

We use the following lemma to bound $K_{\Delta V}$ in Lemma A.8.

Lemma A.7 ([LLT22, Lemma C.12]). *Suppose that $p(x) \propto e^{-V(x)}$ is a probability density on $\mathbb{R}^d$, where $V(x)$ is L-smooth. Let $p_\alpha(x) = \alpha^d p(\alpha x)$ and $\varphi_{\sigma^2}(x)$ denote the density function of $N(0, \sigma^2 I_d)$. Then for $\sigma^2 \le \frac{1}{2\alpha^2 L}$,*

$$\left\|\nabla\ln\frac{p(x)}{(p_\alpha * \varphi_{\sigma^2})(x)}\right\| \le 6\alpha^2 L\sigma d^{1/2} + (\alpha + 2\alpha^3 L\sigma^2)(\alpha-1)L\,\|x\| + (\alpha - 1 + 2\alpha^3 L\sigma^2)\,\|\nabla V(x)\|.$$

Lemma A.8. *Suppose that $h_k \le \frac{1}{4Lg(T-t_k)^2}$ where $\nabla\ln p_t$ is L_{T-t}-smooth ($L_{T-t} \ge 1$) and $L = \max_{t \in [t_k, t_{k+1}]} L_{T-t}$. For $t \in [t_k, t_{k+1}]$,*

$$\mathbb{E}\left[\psi_t(z_t)\,\|\nabla\ln p_{t_k}(z_t) - \nabla\ln p_t(z_t)\|^2\right]$$
$$\le 25L_{T-t}^2\left(8G_{t_k,t}d + G_{t_k,t}^2\mathbb{E}\left[\psi_t(z_t)\,\|z_t\|^2\right]\right) + 100L_{T-t}^2 G_{t_k,t}^2\mathbb{E}\left[\psi_t(z_t)\,\|\nabla\ln p_t(z_t)\|^2\right]$$

Proof. We have the following relationship for $t \in [t_k, t_{k+1}]$:

$$p_{t_k} = (p_t)_\alpha * \varphi_{\sigma^2}.$$

294 where $p_\alpha(x) = \alpha^d p(\alpha x)$, $\alpha = e^{\frac{1}{2}\int_{t_k}^t g(T-s)^2\,ds}$ and $\sigma^2 = 1 - e^{-\int_{t_k}^t g(T-s)^2\,ds}$. Observe that since
295 $h_k \leq \frac{1}{4g(T-t_k)^2}$,

$$\alpha \leq 1 + \int_{t_k}^t g(T-s)^2 ds \leq 1 + h_k g(T-t_k)^2 \leq 1 + \frac{1}{4}$$

$$\sigma^2 = 1 - e^{-\int_{t_k}^t g(T-s)^2 ds} \leq \int_{t_k}^t g(T-s)^2 ds \leq h_k g(T-t_k)^2 \leq \frac{1}{4}.$$

296 We note that

$$\sigma^2 \leq h_k g(T-t_k)^2 \leq \frac{1}{4L_t} \leq \frac{1}{2\alpha^2 L_t}$$

297 so the hypothesis of Lemma A.7 is satisfied. Using Lemma A.7, we obtain

$$\mathbb{E}\left[\psi_t(z_t)\left\|\nabla \ln p_{t_k}(z_t) - \nabla \ln p_t(z_t)\right\|^2\right]$$

$$\leq 72\alpha^4 L_{T-t}^2 \sigma^2 d + 4(\alpha + 2\alpha^3 L_{T-t}\sigma^2)^2(\alpha-1)^2 L_{T-t}^2 \mathbb{E}\left[\psi(z_t)\left\|z_t\right\|^2\right]$$

$$+ 4(\alpha - 1 + 2\alpha^3 L_{T-t}\sigma^2)^2 \mathbb{E}\left[\psi_t(z_t)\left\|\nabla \ln p_t(z_t)\right\|^2\right]$$

$$\leq 72(5/4)^4 L_{T-t}^2 G_{t_k,t} d + 4(2\alpha)^2 G_{t_k,t}^2 L_{T-t}^2 \mathbb{E}\left[\psi(z_t)\left\|z_t\right\|^2\right]$$

$$+ 4(G_{t_k,t} + 4L_{T-t}G_{t_k,t})^2 \mathbb{E}\left[\psi_t(z_t)\left\|\nabla \ln p_t(z_t)\right\|^2\right]$$

$$\leq 200 L_{T-t}^2 d G_{t_k,t} + 25 L_{T-t}^2 G_{t_k,t}^2 \mathbb{E}\left[\psi(z_t)\left\|z_t\right\|^2\right] + 100 L_{T-t}^2 G_{t_k,t}^2 \mathbb{E}\left[\psi_t(z_t)\left\|\nabla \ln p_t(z_t)\right\|^2\right].$$

298 $\hfill\square$

299 Now we put everything together. Write $G_t = G_{t_k,t}$ for short. Suppose L_t is non-increasing. By
300 Lemma A.2,

$$\frac{d}{dt}\chi^2(q_t\|p_t) \leq -\frac{1}{2}g(T-t)^2 \mathscr{E}_{p_t}\left(\frac{q_t}{p_t}\right) + 12g(T-t)^2(\chi^2(q_t\|p_t) + 1)\cdot E$$

where $E \leq 16 G_t^2 L_{T-t_k}^2(K_z + 16 K_V) + 64 G_t L_{T-t_k}^2(8K + 2d + 16\ln 2) + \varepsilon_{\infty,T-t_k}^2 + K_{\Delta V}$.

301 By Lemma A.8, $K_{\Delta V} \leq 25 L_{T-t}^2(8G_t d + G_t^2 K_z) + 100 L_{T-t}^2 G_t^2 K_V$, so

$$E \leq 41 G_t^2 L_{T-t}^2 K_z + 356 G_t^2 L_{T-t}^2 K_V + 64 G_t L_{T-t}^2(8K + 6d + 16\ln 2) + \varepsilon_{\infty,T-t_k}^2.$$

302 By Lemma A.5, $K_z \leq \|x_t\|_{2,\psi_2}^2(K + \ln 2)$, and by Corollary A.6, $K_V \leq \frac{4}{\chi^2(q_t\|p_t)+1}\cdot \mathscr{E}_{p_t}\left(\frac{q_t}{p_t}\right) +$
303 $2dL$, so

$$E \leq 41 G_t^2 L_{T-t}^2\left(\|x_t\|_{2,\psi_2}^2(K + \ln 2)\right) + 356 G_t^2 L_{T-t}^2\left(\frac{4}{\chi^2(q_t\|p_t)+1}\cdot \mathscr{E}_{p_t}\left(\frac{q_t}{p_t}\right) + 2dL\right)$$

$$+ 64 G_t L_{T-t}^2(8K + 6d + 16\ln 2) + \varepsilon_{\infty,T-t_k}^2.$$

304 Now, if $h_k \leq \frac{\varepsilon'_{h_k}}{20g(T-t_k)^2 L_{T-t_{k+1}}}$, then

$$E \leq {\varepsilon'_{h_k}}^2\left[\|x_t\|_{2,\psi_2}^2(K + \ln 2) + \left(\frac{4}{\chi^2(q_t\|p_t)+1}\cdot \mathscr{E}_{p_t}\left(\frac{q_t}{p_t}\right) + 2dL_{T-t}\right)\right]$$

$$+ 4\varepsilon'_{h_k} L_{T-t}(8K + 2d + 16\ln 2) + \varepsilon_{\infty,T-t_k}^2.$$

305 Let $M_{T-t} := \|x_t\|_{2,\psi_2}^2$. Assume that $K \leq \frac{A_{T-t}}{\chi^2(q_t\|p_t)+1} + B_{T-t}$. Then we obtain

$$12g(T-t)^2(\chi^2(q_t\|p_t) + 1)\cdot E$$

$$\leq 12g(T-t)^2\Big[\mathscr{E}_{p_t}\left(\frac{q_t}{p_t}\right)({\varepsilon'_{h_k}}^2\cdot(A_{T-t}M_{T-t} + 4) + \varepsilon'_{h_k}\cdot 32 L_{T-t}A_{T-t})$$

$$+ (\chi^2(q_t\|p_t) + 1)({\varepsilon'_{h_k}}^2\cdot((B_{T-t} + \ln 2)M_{T-t} + 2dL)$$

$$+ \varepsilon'_{h_k}\cdot L_{T-t}(8B_{T-t} + 6d + 16\ln 2)) + \varepsilon_{\infty,T-t_k}^2\Big].$$

306 If $\varepsilon'_{h_k} \leq \min\left\{\frac{1}{\sqrt{48(A_{T-t}M_{T-t}+4)}}, \frac{1}{128L_{T-t}A_{T-t}}\right\}$, then

$$\frac{d}{dt}\chi^2(q_t\|p_t) \leq 12g(T-t)^2\Big[(\chi^2(q_t\|p_t)+1)(\varepsilon'^2_{h_k} \cdot ((B_{T-t}+\ln 2)M_{T-t}+2dL_{T-t})$$
$$+ \varepsilon'_{h_k} \cdot L_{T-t}(8B_{T-t}+6d+16\ln 2)) + \varepsilon^2_{\infty,T-t_k}\Big].$$

307 If $\varepsilon'_{h_k} \leq \min\left\{\frac{\sqrt{\varepsilon'}}{g(T-t)\sqrt{24(T-t_k)((B_{T-t}+\ln 2)M_{T-t}+2dL_{T-t})}}, \frac{\varepsilon'}{24g(T-t)^2(T-t)L_{T-t}(8B_{T-t}+6d+16\ln 2)}\right\}$,

308 we get

$$\frac{d}{dt}\chi^2(q_t\|p_t) \leq \frac{\varepsilon'}{T-t}(\chi^2(q_t\|p_t)+1) + \varepsilon^2_{\infty,T-t_k}g(T-t)^2.$$

309 Integration gives

$$\chi^2(q_{t_k}\|p_{t_k}) \leq e^{\varepsilon'\int_0^{t_k}\frac{1}{T-t}dt}(\chi^2(q_0\|p_0)+1) + \int_0^{t_k}e^{\int_t^{t_k}\frac{\varepsilon'}{T-s}ds}\varepsilon^2_{T-t}g(T-t)^2\,dt$$

$$\leq \left(\frac{T}{T-t_k}\right)^{\varepsilon'}\chi^2(q_0\|p_0) + \left(\left(\frac{T}{T-t_k}\right)^{\varepsilon'}-1\right) + \int_0^{t_k}\left(\frac{T-t}{T-t_k}\right)^{\varepsilon'}\varepsilon^2_{T-t}g(T-t)^2\,dt.$$

310 Taking $\varepsilon' = \frac{\varepsilon}{\ln\left(\frac{T}{T-t_N}\right)}$ then gives the following Theorem A.10. We first introduce a technical

311 assumption.

312 **Definition A.9.** *Let $f : \mathbb{R}_{>0} \to \mathbb{R}_{>0}$. We say that f has at most power growth and decay (with some*

313 *constant $c > 0$) if $\max_{u\in[\frac{t}{2},t]} f(u) \in \left[\frac{f(t)}{c}, cf(t)\right]$.*

314 **Theorem A.10.** *Suppose that the following hold.*

315   *1. Assumption 5 holds for $\varepsilon_{\infty,t}$.*

316   *2. $\|\widetilde{x}_t\|^2_{2,\psi_2} \leq M_t$.*

317   *3. The KL bound $\mathrm{KL}(\psi_t q_t\|p_t) \leq \frac{A_{T-t}}{\chi^2(q_t\|p_t)+1} + B_{T-t}$ holds for any density q_t and $t < t_N$,*

318   *where $\psi_t(x) = \frac{q_t(x)/p_t(x)}{\chi^2(q_t\|p_t)+1}$.*

319   *4. $g(t), A_t, B_t, L_t, M_t$ have at most polynomial growth and decay (with some constant c).*

320 *Then there is some constant c' (depending on c) such that if the step sizes satisfy*

$$h_k \leq \min\left\{\frac{T-t_k}{2}, \frac{c'\varepsilon'_{h_k}}{g(T-t_k)^2 L_{T-t_k}}\right\},$$

$$\text{where } \varepsilon'_{h_k} = \min\left\{\frac{1}{\sqrt{A_{T-t_k}M_{T-t_k}+1}}, \frac{1}{L_{T-t_k}A_{T-t_k}},\right.$$

$$\frac{\sqrt{\varepsilon/\ln\left(\frac{T}{T-t_N}\right)}}{g(T-t_k)\sqrt{(T-t_k)((B_{T-t_k}+1)M_{T-t_k}+dL_{T-t_k})}},$$

$$\left.\frac{\varepsilon/\ln\left(\frac{T}{T-t_N}\right)}{g(T-t_k)^2(T-t_k)L_{T-t_k}(B_{T-t_k}+d)}\right\},$$

321 *then for $0 \leq k \leq N$,*

$$\chi^2(q_{t_k}\|p_{t_k}) \leq e^\varepsilon \chi^2(q_0\|p_0) + (e^\varepsilon-1) + e^\varepsilon\int_0^{t_k}\varepsilon^2_{\infty,T-t}g(T-t)^2\,dt.$$

322 *Proof.* This follows from the above calculations and the observation that if we replace $F(T-t)$

323 by $F(T-t_k)$, for some F satisfying the power growth and decay assumption, then we change the

324 bound by at most a constant factor, because the step size satisfies $h_k = t_{k+1} - t_k \leq \frac{T-t_k}{2}$. □

We specialize this theorem in the case of distributions with bounded support. Note that although not every initial distribution $\widetilde{p}_t$ may satisfy a KL inequality as required by condition 3 of Theorem 30, Lemma B.2 will give the existence of a distribution that does, and is close in TV-error. Later in Section C, we show that this will have a small effect on the score function, and hence allow us to prove our main theorems.

Corollary A.11. *Suppose that Assumptions 5 and 2 hold, $R^2 \geq d$, $g \equiv 1$, and that $\widetilde{P}_0$ is such that the KL inequality* (30) *holds. Let $\delta = T - t_N$. If $0 < \delta, \varepsilon < \frac{1}{2}$, $h_k =$*
$$O\left(\frac{\varepsilon}{\max\{T-t_k, (T-t_k)^{-3}\}R^4 d \ln\left(\frac{T}{\delta}\right)\ln\left(\frac{R}{\delta\varepsilon_K}\right)}\right), \text{ then for any } 0 \leq k \leq N,$$

$$\chi^2(q_{t_k}\|p_{t_k}) \leq e^\varepsilon \chi^2(q_0\|p_0) + \varepsilon + e^\varepsilon \int_0^{t_k} \varepsilon_{\infty, T-t}^2 \, dt.$$

Proof. For $g \equiv 1$, note that $\sigma_{T-t}^2 = \Theta(\min\{T - t, 1\})$. From Lemma A.13, we can choose
$$L_t = \frac{R^2}{\sigma_t^4} = O\left(\frac{R^2}{\min\{(T-t)^2, 1\}}\right).$$

From Lemma A.15, we can choose
$$M_t = \max\{R^2, d\}.$$

The KL inequality (30) gives us
$$A_t = 6(e+1)\sigma_t^2 = O(\min\{T - t, 1\})$$
$$B_t = \ln\left(\frac{1}{\varepsilon}\right) + d\ln\left(1 + O\left(\frac{R}{\sqrt{T - t_N}}\right)\right)$$

We now check the requirements on h_k. We need

$$\varepsilon'_{h_k} = O\left(\frac{1}{\sqrt{A_{T-t_k}M_{T-t_k} + 1}}\right) \qquad \Longleftarrow \varepsilon'_{h_k} = O\left(\frac{1}{\max\{R, \sqrt{d}\}}\right) \tag{22}$$

$$\varepsilon'_{h_k} = O\left(\frac{1}{L_{T-t_k}A_{T-t_k}}\right) \qquad \Longleftarrow \varepsilon'_{h_k} = O\left(\frac{T - t_k}{R^2}\right) \tag{23}$$

$$\varepsilon'_{h_k} = O\left(\frac{\sqrt{\varepsilon/\ln\left(\frac{T}{\delta}\right)}}{\sqrt{(T - t_k)((B_{T-t_k} + 1)M_{T-t_k} + dL_{T-t_k})}}\right). \tag{24}$$

For $T - t_k \leq 1$, (24) is implied by

$$\varepsilon'_{h_k} = O\left(\frac{\sqrt{\varepsilon/\ln\left(\frac{T}{\delta}\right)}}{\sqrt{(T - t_k)\left(\ln\left(\frac{1}{\varepsilon_K}\right) + d\ln\left(\frac{R}{\delta}\right)\right)\max\{R^2, d\} + \frac{dR^2}{T-t_k}}}\right)$$

$$\Longleftarrow \varepsilon'_{h_k} = O\left(\sqrt{\frac{\varepsilon(T - t_k)}{d\max\{R^2, d\}\ln\left(\frac{T}{\delta}\right)\ln\left(\frac{R}{\delta\varepsilon_K}\right)}}\right),$$

and for $T - t_k > 1$,

$$\varepsilon'_{h_k} = O\left(\frac{\sqrt{\varepsilon/\ln\left(\frac{T}{\delta}\right)}}{\sqrt{T\left(\ln\left(\frac{1}{\varepsilon}\right) + d\ln\left(\frac{R}{\delta}\right)\right)\max\{R^2, d\} + dR^2}}\right)$$

$$\Longleftarrow \varepsilon'_{h_k} = O\left(\sqrt{\frac{\varepsilon}{Td\max\{R^2, d\}\ln\left(\frac{T}{\delta}\right)\ln\left(\frac{R}{\delta\varepsilon_K}\right)}}\right).$$

Finally, the last requirement is

$$\varepsilon'_{h_k} = O\left(\frac{\varepsilon/\ln\left(\frac{T}{\delta}\right)}{(T-t_k)L_{T-t_k}(B_{T-t_k}+d)}\right)$$

$$\Longleftarrow \varepsilon'_{h_k} = O\left(\frac{\varepsilon}{R^2\max\{T-t_k,(T-t_k)^{-1}\}d\ln\left(\frac{T}{\delta}\right)\ln\left(\frac{R}{\delta\varepsilon_K}\right)}\right).$$

As long as $R^2 = \Omega(d)$ and $\varepsilon < 1$, the last equation implies all the others. Plugging this into Theorem A.10 gives the result. □

Above, we use the Hessian bound $\left\|\nabla^2\ln p_t(x)\right\| \le \frac{R^2}{\sigma_t^4}$ given in Lemma A.13. Under the stronger smoothness assumption given by Assumption 3, we can take the step sizes to be larger.

Corollary A.12. *Suppose that Assumptions 5, 2, 3 hold, $C \ge R^2 \ge d$, $g \equiv 1$, and that $\widetilde{P}_0$ is such that the KL inequality (30) holds. Let $\delta = T - t_N$. If $0 < \delta, ep < \frac{1}{2}$ and $\varepsilon < 1/\sqrt{T}$,*

$$h_k = O\left(\frac{\varepsilon}{\max\{T-t_k,(T-t_k)^{-1}\}C^2 d\ln\left(\frac{T}{\delta}\right)\ln\left(\frac{R}{\delta\varepsilon_K}\right)}\right), \text{ then for any } 0 \le k \le N,$$

$$\chi^2(q_{t_k}\|p_{t_k}) \le e^\varepsilon\chi^2(q_0\|p_0) + \varepsilon + e^\varepsilon\int_0^{t_k}\varepsilon_{\infty,T-t}^2\,dt.$$

Proof. We instead have the bound $L_t = \frac{C}{\sigma_t^2}$. The requirement (22) stays the same, while (23) is implied by $\varepsilon'_{h_k} = O(1/C)$. Inequality (24), for $T - t_k \le 1$, is implied by

$$\varepsilon'_{h_k} = O\left(\sqrt{\frac{1}{d\max\{C,R^2\}\ln\left(\frac{T}{\delta}\right)\ln\left(\frac{R}{\delta\varepsilon_K}\right)}}\right).$$

and for $T - t_k > 1$,

$$\varepsilon'_{h_k} = O\left(\sqrt{\frac{\varepsilon}{Td\max\{C,R^2\}\ln\left(\frac{T}{\delta}\right)\ln\left(\frac{R}{\delta\varepsilon_K}\right)}}\right).$$

Finally, the last requirement is implied by

$$\varepsilon'_{h_k} = O\left(\frac{\varepsilon}{Cd\ln\left(\frac{T}{\delta}\right)\ln\left(\frac{R}{\delta\varepsilon_K}\right)}\right),$$

and for $C \ge R^2, \varepsilon \le 1/\sqrt{T}$, implies all the others. □

A.1 Auxiliary bounds

In this section we give bounds on the Hessian (L_t, Lemma A.13), initial χ^2 divergence $\chi^2(q_0\|p_0)$ (Lemma A.14), and Orlicz norm (M_t, Lemma A.15).

Lemma A.13 (Hessian bound). *Suppose that μ is a probability measure supported on a bounded set $\mathcal{M} \subset \mathbb{R}^d$ with radius R. Then letting φ_{σ^2} denote the density of $N(0, \sigma^2 I_d)$,*

$$\left\|\nabla^2\ln(\mu * \varphi_{\sigma^2}(x))\right\| \le \frac{R^2}{\sigma^4}. \tag{25}$$

Therefore, for $\widetilde{P}_0$ supported on $B_R(0)$, we have

$$\left\|\nabla^2\ln\widetilde{p}_t(x)\right\| \le \frac{R^2}{\sigma_t^4}. \tag{26}$$

Proof. Let μ_{x,σ^2} denote the density $\mu(du)$ weighted with the gaussian $\varphi_{\sigma^2}(u-x)$, that is,

$\mu_{x,\sigma^2}(du) = \dfrac{e^{-\frac{\|x-u\|^2}{2\sigma^2}}\mu(du)}{\int_{\mathbb{R}^d} e^{-\frac{\|x-u\|^2}{2\sigma^2}}\mu(du)}$. We note the following calculations:

$$\nabla \ln(\mu * \varphi_{\sigma^2}(x)) = \frac{\nabla \int_{\mathbb{R}^d} e^{-\frac{\|x-u\|^2}{2\sigma^2}}\mu(du)}{\int_{\mathbb{R}^d} e^{-\frac{\|x-u\|^2}{2\sigma^2}}\mu(du)} = \frac{\int_{\mathbb{R}^d} -\frac{x-u}{\sigma^2}e^{-\frac{\|x-u\|^2}{2\sigma^2}}\mu(du)}{\int_{\mathbb{R}^d} e^{-\frac{\|x-u\|^2}{2\sigma^2}}\mu(du)} = -\frac{1}{\sigma^2}\mathbb{E}_{\mu_{x,\sigma^2}}(x-u) \tag{27}$$

$$\nabla^2 \ln(\mu * \varphi_{\sigma^2}(x)) = \frac{1}{\sigma^4}\operatorname{Cov}_{\mu_{x,\sigma^2}}(x-u) = \frac{1}{\sigma^4}\operatorname{Cov}_{\mu_{x,\sigma^2}}(x). \tag{28}$$

The covariance of a distribution supported on a set of radius R is bounded by R^2 in operator norm. Inequality (25) then follows from (28).

For (26), note that $\widetilde{p}_t = M_{m_t\sharp}\widetilde{P}_0 * \varphi_{\sigma_t^2}$, where m_t is given by (3) and M_m denotes multiplication by m. Since $M_{m_t\sharp}\widetilde{P}_0$ is supported on $B_{m_tR}(0) \subset B_R(0)$, the result follows. $\qquad\square$

Lemma A.14 (Bound on initial χ^2-divergence). *Suppose that $\widetilde{P}_0$ is supported on $B_R(0)$. Let $p_{\mathrm{prior}} = N(0, (1 - e^{G_{0,t}})I_d)$. Then*

$$\chi^2(p_{\mathrm{prior}}\|\widetilde{p}_T) \leq \exp\left[\frac{R^2 \exp(-G_{0,T})}{1 - \exp(-G_{0,T})}\right]$$

and for $0 < \varepsilon < \frac{1}{2}$ and $G_{0,T} \geq \ln\left(\frac{4R^2}{\varepsilon^2}\right) \vee 1$, we have $\chi^2(p_{\mathrm{prior}}\|\widetilde{p}_T) \leq \varepsilon^2$.

Proof. We have for $x_0 \sim \widetilde{P}_0$ that

$$\chi^2\left(N(0, (1 - e^{-G_{0,T}})I_d)\|N(x_0 \exp\left(-\frac{1}{2}G_{0,T}\right), (1 - \exp(-G_{0,T}))I_d)\right)$$

$$\leq \exp\left[\frac{\|x_0\|^2 \exp(-G_{0,T})}{1 - \exp(-G_{0,T})}\right] \leq \exp\left(\frac{R^2 \exp(-G_{0,T})}{1 - \exp(-G_{0,T})}\right)$$

Using convexity of χ^2-divergence then gives the result. For $G_{0,T} \geq \ln\left(\frac{4R^2}{\varepsilon^2}\right) \vee 1$, we have

$$\exp\left[\frac{R^2 \exp(-G_{0,T})}{1 - \exp(-G_{0,T})}\right] \leq \exp\left[\frac{\varepsilon^2/4}{1/2}\right] \leq \varepsilon^2. \qquad\square$$

Lemma A.15 (Subgaussian bound). *Suppose $\widetilde{P}_0$ is supported on $B_R(0)$. Then for $X \sim \widetilde{p}_t$,*

$$\|X\|_{2,\psi_2} \leq \sqrt{\frac{e}{\ln 2}} \cdot \left(4m_tR + 6C_1\sigma_t\sqrt{d}\right) = O(\max\{R, \sqrt{d}\}),$$

where m_t, σ_t are as in (3) and C_1 is an absolute constant.

Proof. Let $Y \sim \widetilde{P}_0$ s.t. $X = m_tY + \sigma_t\xi$ for some $\xi \sim N(0, I_d)$ independent of Y. Define $U = \|X\|_2 := \left(\sum_{i=1}^d X_i^2\right)^{1/2}$, then for $p \geq 1$,

$$\begin{aligned}
\mathbb{E}|U|^p = \mathbb{E}\|X\|_2^p &\leq \mathbb{E}\left(\|m_tY\|_2 + \|\sigma_t\xi\|_2\right)^p \\
&\leq 2^{p-1}\mathbb{E}\left[\|m_tY\|_2^p + \|\sigma_t\xi\|_2^p\right] \\
&\leq 2^{p-1}\left[(m_tR)^p + \sigma_t^p \cdot 2^{p/2}\frac{\Gamma((d+p)/2)}{\Gamma(d/2)}\right] \\
&\leq 2^{p-1}\left[(m_tR)^p + C_1(\sqrt{2}\sigma_t)^p \cdot \left(d^{p/2} + p^{p/2}\right)\right]
\end{aligned}$$

where Γ is the commonly used gamma function and C_1 is an absolute constant. Therefore

$$(\mathbb{E}|U|^p)^{1/p} \leq 2m_t R + \sqrt{2}C_1\sigma_t(\sqrt{d} + \sqrt{p}) \leq K\sqrt{p},$$

where $K = 2m_t R + 3C_1\sigma_t\sqrt{d}$. Now consider $V = U/K$, then for some $\lambda > 0$ small enough, by Taylor expansion,

$$\mathbb{E}\left[e^{\lambda^2 V^2}\right] = \mathbb{E}\left[1 + \sum_{p=1}^{\infty} \frac{(\lambda^2 V^2)^p}{p!}\right] = 1 + \sum_{p=1}^{\infty} \frac{\lambda^{2p}\mathbb{E}\left[V^{2p}\right]}{p!}.$$

Note that $\mathbb{E}\left[V^{2p}\right] \leq (2p)^p$, while Stirling's approximation yields $p! \geq (p/e)^p$. Substituting these two bounds, we get

$$\mathbb{E}e^{\lambda^2 V^2} \leq 1 + \sum_{p=1}^{\infty} \left(\frac{2\lambda^2 p}{p/e}\right) = \sum_{p=0}^{\infty}(2e\lambda^2)^p = \frac{1}{1 - 2e\lambda^2},$$

provided that $2e\lambda^2 < 1$, in which case the geometric series above converges. To bound this quantity further, we can use the numeric inequality $1/(1-x) \leq e^{2x}$ which is valid for $x \in [0, 1/2]$. It follows that

$$\mathbb{E}e^{\lambda^2 V^2} \leq e^{4e\lambda^2} \quad \text{for all } \lambda \text{ satisfying } |\lambda| \leq 1/2\sqrt{e}.$$

Now set $4e\lambda^2 = \ln 2$, then

$$\mathbb{E}\left[e^{\frac{\ln 2}{4eK^2}\|X\|_2^2}\right] \leq 2,$$

which implies that

$$\|X\|_{2,\psi_2} \leq \sqrt{\frac{4e}{\ln 2}}K = \sqrt{\frac{e}{\ln 2}} \cdot \left(4m_t R + 6C_1\sigma_t\sqrt{d}\right). \qquad \square$$

B Bounding the KL divergence

In this section, we bound the quantity $K = \mathrm{KL}(\psi_t q_t \| p_t)$, where ψ_t is as in (8). While p_t is defined by the DDPM process, in this section we do *not* assume q_t is the density of the discretized process; rather, it is any density for which $\mathscr{E}_{p_t}\left(\frac{q_t}{p_t}\right)$ and $\chi^2(q_t\|p_t)$ are finite.

Lemma B.1. *Suppose that $\widetilde{P}_0$ is a probability measure on $\mathbb{R}^d$ such that*

$$\widetilde{P}_0 = \sum_{j=1}^{m} w_j \widetilde{P}_{j,0}, \tag{29}$$

where $w_j > 0$, $\sum_{j=1}^{m} w_j = 1$, and each $\widetilde{P}_{j,0}$ is a probability measure. For $t > 0$, let $\widetilde{p}_t$ and $\widetilde{p}_{j,t}$ be the densities obtained by running the forward DDPM process (1) for time t, and $p_t = \widetilde{p}_{T-t}$, $p_{j,t} = \widetilde{p}_{j,T-t}$. Let $w_{\min} = \min_{1 \leq j \leq m} w_j$ and suppose all the $\widetilde{P}_{j,t}$ satisfy a log-Sobolev constant with constant C_t. Then for any q_t, where ψ_t is as in (8)

$$\mathrm{KL}(\psi_t q_t \| p_t) \leq \frac{2C_{T-t}}{\chi^2(q_t\|p_t) + 1} \cdot \mathscr{E}_{p_t}\left(\frac{q_t}{p_t}\right) + \ln\left(\frac{1}{w_{\min}}\right).$$

While we need p_t to satisfy a log-Sobolev inequality to get a bound of the form $\frac{C}{\chi^2(q_t\|p_t)+1}\mathscr{E}_{p_t}\left(\frac{q_t}{p_t}\right)$ ([LLT22, Lemma C.8]), we note that if we allow additive slack, it suffices for p_t to be a mixture of distributions satisfying a log-Sobolev inequality, with the *logarithm* of the minimum mixture weight bounded below. In Lemma B.2 we will see that we can almost decompose any distribution of bounded support in this manner, if we move a small amount of the mass.

Proof. Let $\overline{f}_t : [m] \to \mathbb{R}$ be the function

$$\overline{f}_t(j) = \int_{\mathbb{R}^d} \frac{\psi_t(x)q_t(x)}{p_t(x)}P_{j,t}(x)\,dx.$$

By decomposition of entropy and the fact that each $P_{i,t}$ satisfies LSI with constant C_{T-t},

$$
\begin{aligned}
\mathrm{KL}(\psi_t q_t \| p_t) &\leq \mathrm{Ent}_{p_t}\left(\frac{\psi_t q_t}{p_t}\right) \\
&= \sum_{i=1}^{m} \int_{\mathbb{R}^d} w_i \, \mathrm{Ent}_{P_{i,t}}\left(\frac{\psi_t q_t}{p_t}\right) + \mathrm{Ent}_w(\overline{f}_t) \\
&\leq \frac{C_t}{2} \sum_{i=1}^{m} w_i \mathscr{E}_{P_{i,t}}\left(\ln \frac{\psi_t q_t}{p_t}, \frac{\psi_t q_t}{p_t}\right) + \mathrm{Ent}_w(\overline{f}_t) \\
&\leq \frac{C_t}{2} \mathscr{E}_{p_t}\left(\ln \frac{\psi_t q_t}{p_t}, \frac{\psi_t q_t}{p_t}\right) + \mathrm{Ent}_w(\overline{f}_t) \\
&= \frac{C_t}{2} \int_{\mathbb{R}^d} \left\| \nabla \ln \frac{\psi_t(x) q_t(x)}{p_t(x)} \right\|^2 \psi_t(x) q_t(x) \, dx + \mathrm{Ent}_w(\overline{f}_t) \\
&= 2C_t \int_{\mathbb{R}^d} \left\| \nabla \ln \frac{q_t(x)}{p_t(x)} \right\|^2 \psi_t(x) q_t(x) \, dx + \mathrm{Ent}_w(\overline{f}_t) \\
&= 2C_t \int_{\mathbb{R}^d} \left\| \nabla \frac{q_t(x)}{p_t(x)} \right\|^2 \frac{\psi_t(x) p_t(x)^2}{q_t(x)} \, dx + \mathrm{Ent}_w(\overline{f}_t) \\
&= \frac{2C_t}{\chi^2(q_t \| p_t) + 1} \cdot \int \left\| \nabla \frac{q_t(x)}{p_t(x)} \right\|^2 p_t(x) \, dx + \mathrm{Ent}_w(\overline{f}_t) \\
&\leq \frac{2C_t}{\chi^2(q_t \| p_t) + 1} \cdot \mathscr{E}_{p_t}\left(\frac{q_t}{p_t}\right) + \ln\left(\frac{1}{w_{\min}}\right),
\end{aligned}
$$

where the last inequality follows from noting $w_j \overline{f}_t(j)$ is a probability mass function on $[m]$, so that $\overline{f}_t(j) \leq \frac{1}{w_j}$ and

$$
\mathrm{Ent}_w(\overline{f}_t) = \sum_{j=1}^{m} w_j \overline{f}_t(j) \ln(\overline{f}_t(j)) \leq \sum_{j=1}^{m} w_j \overline{f}_t(j) \ln\left(\frac{1}{w_{\min}}\right) = \ln\left(\frac{1}{w_{\min}}\right). \qquad \square
$$

Lemma B.2. *Suppose $0 < \varepsilon_K < \frac{1}{2}$, and that $\overline{P}_0$ is a probability measure such that $\overline{P}_0(\mathcal{M}) \geq 1 - \frac{\varepsilon_K}{8}$. Let $\mathcal{N}\left(\mathcal{M}, \frac{\sigma_t}{2}\right)$ denote the covering number of $\mathcal{M}$ with balls of radius σ_t. Given $\delta > 0$, there exists a distribution $\widetilde{P}_0$ such that $\chi^2(\widetilde{P}_0 \| \overline{P}_0) \leq \varepsilon_K$ and considering the DDPM process started with $\widetilde{P}_0$, for all $0 \leq t \leq T - \delta$,*

$$
\mathrm{KL}(\psi_t q_t \| p_t) \leq \left(\frac{6(1+e)\sigma_{T-t}^2}{\chi^2(q_t \| p_t) + 1} \mathscr{E}_{p_t}\left(\frac{q_t}{p_t}\right) + \ln\left(\frac{\mathcal{N}(\mathcal{M}, \sigma_\delta/2)}{\varepsilon_K}\right)\right).
$$

In particular, for $\mathcal{M} = B_R(0)$ in $\mathbb{R}^d$,

$$
\mathrm{KL}(\psi_t q_t \| p_t) \leq \left(\frac{6(1+e)\sigma_{T-t}^2}{\chi^2(q_t \| p_t) + 1} \mathscr{E}_{p_t}\left(\frac{q_t}{p_t}\right) + \ln\left(\frac{1}{\varepsilon_K}\right) + d \ln\left(1 + \frac{4R}{\sigma_\delta}\right)\right). \tag{30}
$$

Proof. Partition $\mathcal{M}$ into disjoint subsets $\mathcal{M}_j$, $1 \leq j \leq N := \mathcal{N}(\mathcal{M}, \sigma_\delta/2)$ of diameter at most σ_δ, and decompose

$$
\overline{P}_0 = w_* P_* + \sum_{j=1}^{n} \overline{w}_j \widetilde{P}_{j,0}
$$

where p_j is supported on $\mathcal{M}_j$ and $P_* = \overline{P}_0(\cdot | \mathcal{M}^c)$. We will zero out the coefficients of all small components: let $Z = \sum_{j: w_j \geq \frac{\varepsilon_K}{8N}} w_j$ and

$$
w_j = \begin{cases} \frac{\overline{w}_j}{Z}, & j \in [n], \overline{w}_j \geq \frac{\varepsilon_K}{8N} \\ 0, & \text{otherwise,} \end{cases}
$$

and define

$$\widetilde{P}_0 = \sum_{j=1}^{n} w_j \widetilde{P}_{j,0}.$$

Note that $Z \geq 1 - \frac{\varepsilon_K}{8} - \sum_{j:w_j \leq \frac{\varepsilon_K}{8N}} \geq 1 - \frac{\varepsilon_K}{4}$. As probability distributions on $[m] \cup \{*\}$,

$$\chi^2(w||\overline{w}) \leq \left(\frac{1}{1 - \frac{\varepsilon_K}{4}}\right)^2 - 1 \leq \varepsilon_K,$$

and hence the same bound holds for $\chi^2(\widetilde{P}_0||\overline{P}_0)$. Note each $M_{m_t\sharp}\widetilde{P}_{j,0}$ is supported on a set of diameter $m_t\sigma \leq \sigma$. By Theorem 1 of [CCN21], noting that

$$\chi^2(N(\mu_2, \Sigma)||N(\mu_1, \Sigma)) = \exp\left[(\mu_2 - \mu_1)^\top \Sigma^{-1}(\mu_2 - \mu_1)\right] \leq e$$

when $\Sigma = \sigma^2 I$ and $\|\mu_2 - \mu_1\| \leq \sigma$, $\widetilde{P}_{j,t} = (M_{m_t\sharp}\widetilde{P}_{j,0}) * \varphi_{\sigma^2}$ satisfies a log-Sobolev inequality with constant $6(1 + e)\sigma_t^2$. The result then follows from Lemma B.1. For $\mathcal{M} = B_R(0)$, we use the bound $\mathcal{N}(B_R(0), \sigma_\delta/2) \leq \left(1 + \frac{4R}{\sigma_\delta}\right)^d$ [Ver18, Corollary 4.2.13]. $\qquad\square$

In the next section, we show that we can move a small amount of mass ε without significantly affecting the score function. This is necessary, as our guarantees on the score estimate are for the original distribution and not the perturbed one in Lemma B.2.

C The effect of perturbing the data distribution on the score function

In this section we consider the effect of perturbing the data distribution on the score function. The key observation is that the score function can be interpreted as the solution to an inference problem, that of recovering the original data point from a noisy sample, with data distribution as the given prior distribution. We show through a coupling argument that we can bound the difference between the score functions in terms of the distance between the two data distributions. This will allow us to "massage" the data distribution in order to optimally bound $\mathrm{KL}(\psi_t q_t||p_t)$ in Section B.

C.1 Perturbation under χ^2 error and truncation

We first give a general lemma on denoising error from a mismatched prior.

Lemma C.1. *Let φ be a probability density on $\mathbb{R}^d$, and $P_{0,x}, P_{1,x}$ be measures on $\mathbb{R}^d$. For $i = 0, 1$, let P_i denote the joint distribution of $x_i \sim P_{i,x}$ and $y_i = x_i + \xi_i$ where $\xi_i \sim \varphi$, and let $P_{i,y}$ denote the marginal distribution of y. Let*

$$m^{(k)}(\varepsilon) := \sup_{0 \leq f \leq 1, \int_{\mathbb{R}^d} f\varphi \, dx \leq \varepsilon} \int_{\mathbb{R}^d} f(x) \|x\|^k \varphi(x) \, dx.$$

Let $\varepsilon_{\mathrm{TV}} = \mathrm{TV}(P_{0,x}, P_{1,x})$ and $\varepsilon_\chi^2 = \chi^2(P_{0,x}||P_{1,x})$. Then

$$\int_{\mathbb{R}^d} P_{0,y}(dy_0) \left\| \int_{\mathbb{R}^d} x_0 P_0(dx_0|y_0) - \int_{\mathbb{R}^d} x_1 P_1(dx_1|y_0) \right\|^2$$
$$\leq 8m^{(2)}(\varepsilon_{\mathrm{TV}}) + \varepsilon_\chi \sqrt{m^{(4)}(\varepsilon_{\mathrm{TV}})}$$

For $\varphi = \varphi_{\sigma^2}$, the upper bound is $O\left(\sigma^2 \varepsilon_\chi \left(d + \ln\left(\frac{1}{\varepsilon_{\mathrm{TV}}}\right)\right)\right)$.

Note the tricky part of the proof is to deal with $P_1(dx_1|y_0)$, which can be thought of as inferring x assuming the incorrect prior $P_{1,x}$, rather than the actual prior $P_{0,x}$.

Proof. For notational clarity, we will denote draws from the conditional distribution as $\widehat{x}_0$ and $\widehat{x}_1$, for example $P_0(d\widehat{x}_0|y_0)$. Let $r_i(y) = \int_{\mathbb{R}^d}(\widehat{x}_i - y)P_i(d\widehat{x}_i|y)$. Let $P_{0,1}$ be a coupling of $(x_0, y_0 = $

438 $x_0 + \xi_0, x_1, y_1 = y_1 + \xi_1)$ such that $x_0 = x_1$ with probability $1 - \varepsilon_{\mathrm{TV}}$ and $\xi_0 = \xi_1$ with probability
439 1. We have

$$\int_{\mathbb{R}^d} P_{0,y}(dy_0) \|r_0(y_0) - r_1(y_0)\|^2 = \underbrace{\int_{\{y_0 = y_1\}} P_{0,1,y}(dy_0, dy_1) \|r_0(y_0) - r_1(y_0)\|^2}_{(I)}$$

$$+ \underbrace{\int_{\{y_0 \neq y_1\}} P_{0,1,y}(dy_0, dy_1) \|r_0(y_0) - r_1(y_0)\|^2}_{(II)}.$$

440 Define a measure Q (not necessarily a probability measure) on $\mathbb{R}^d$ by

$$Q(A) := P_{0,1}(y_0 \in A \text{ and } y_0 = y_1).$$

441 Note that

$$Q(A) \leq \min\{P_{0,y}(A), P_{1,y}(A)\},$$

442 so Q is absolutely continuous with respect to $P_{0,y}$ and $P_{1,y}$, and by assumption on the coupling,

$$Q(\mathbb{R}^d) \geq 1 - \varepsilon_{\mathrm{TV}}. \tag{31}$$

443 Under $P_{0,1}$, when $y_0 = y_1$, we can couple $P_0(d\widehat{x}_0|y_0)$ and $P_1(d\widehat{x}_1|y_0)$ so that $x_0 = x_1$ with
444 probability $\min\left\{\frac{dQ}{dP_{0,y}}, \frac{dQ}{dP_{1,y}}\right\}$. Let $\widehat{P}(d\widehat{x}_0, d\widehat{x}_1|y_0)$ denote this coupled distribution. Then as in
445 Lemma C.5,

$$(I) \leq \int_{\{y_0 = y_1\}} P_{0,1,y}(dy_0, dy_1) \left\| \int_{\{\widehat{x}_0 \neq \widehat{x}_1\}} ((\widehat{x}_0 - y_0) - (\widehat{x}_1 - y_0)) \widehat{P}(d\widehat{x}_0, d\widehat{x}_1|y_0) \right\|^2$$

$$\leq 2 \int_{\mathbb{R}^d} P_{0,1,y}(dy_0, dy_1) \left(\int_{\{\widehat{x}_0 \neq \widehat{x}_1\}} \|\xi_0\|^2 \, \widehat{P}(d\widehat{x}_0, d\widehat{x}_1|y_0) + \int_{\{\widehat{x}_0 \neq \widehat{x}_1\}} \|\xi_1\|^2 \, \widehat{P}(d\widehat{x}_0, d\widehat{x}_1|y_1) \right)$$

446 We bound this by first bounding

$$\int_{\mathbb{R}^d} P_{0,1,y}(dy_1, dy_2) \widehat{P}(\widehat{x}_0 \neq \widehat{x}_1) \leq \int_{\mathbb{R}^d} P_{0,y}(dy) \max\left\{ 1 - \frac{dQ}{dP_{0,y}}, 1 - \frac{dQ}{dP_{1,y}} \right\} \leq 2\varepsilon_{\mathrm{TV}}, \tag{32}$$

447 which follows from the two inequalities (using (31))

$$\int_{\mathbb{R}^d} P_{0,y}(dy) \left(1 - \frac{dQ}{dP_{0,y}} \right) = 1 - Q(\mathbb{R}^d) \leq \varepsilon_{\mathrm{TV}}$$

$$\int_{\mathbb{R}^d} P_{0,y}(dy) \left(1 - \frac{dQ}{dP_{1,y}} \right) \leq \int_{\mathbb{R}^d} P_{1,y}(dy) \left(1 - \frac{dQ}{dP_{1,y}} \right) + \mathrm{TV}(P_{0,y}, P_{1,y})$$

$$\leq (1 - Q(\mathbb{R}^d)) + \varepsilon_{\mathrm{TV}} \leq 2\varepsilon_{\mathrm{TV}}.$$

448 From (32), and the fact that the distribution of (x_i, y_i) is the same as $(\widehat{x}_i, y_i)$ by Nishimori's identity,
449 we obtain

$$(I) \leq 2(m^{(2)}(2\varepsilon_{\mathrm{TV}}) + m^{(2)}(2\varepsilon_{\mathrm{TV}})) = 4m^{(2)}(\varepsilon_{\mathrm{TV}}).$$

450 Now for the second term (II),

$$(II) \leq 2 \int_{\{y_0 \neq y_1\}} P_{0,1,y}(dy_0, dy_1)(\|r_0(y_0)\|^2 + \|r_1(y_0)\|^2).$$

451 The first term satisfies $\int_{\{y_0 \neq y_1\}} P_{0,1,y}(dy_0, dy_1) \|r_0(y_0)\|^2 \leq m^{(2)}(\varepsilon_{\mathrm{TV}})$. For the second term, we
452 note that Cauchy-Schwarz gives for any measures P and Q that

$$\int_\Omega f(x) P(dx) \leq \int_\Omega f(x) Q(dx) + \int_\Omega \left(\frac{dP}{dQ} - 1 \right) f(x) Q(dx)$$

$$\leq \int_\Omega f(x) Q(dx) + \sqrt{\chi^2(P\|Q) \int_\Omega f(x)^2 Q(dx)}$$

453 to switch from the measure $P_{0,y}$ to $P_{1,y}$:

$$\int_{\{y_0 \neq y_1\}} P_{0,1,y}(dy_0) \left\| r_1(y_0) \right\|^2 = \int_{\mathbb{R}^n} P_{0,y}(dy_0) P_{0,1,y}(y_0 \neq y_1 | y_0) \left\| r_1(y_0) \right\|^2$$

$$\leq \int_{\mathbb{R}^n} P_{1,y}(dy_0) P_{0,1,y}(y_0 \neq y_1 | y_0) \left\| r_1(y_0) \right\|^2 + \sqrt{\chi^2(P_{0,y} \| P_{1,y}) \int P_{1,y}(dy_0) P_{0,1,y}(y_0 \neq y_1 | y_0) \left\| r_1(y_0) \right\|^4}$$

454 (Note that intentionally, the measure is $P_{1,y}$, though we use y_0 for the variable.) Hence,

$$\int_{\mathbb{R}^n} P_{1,y}(dy_0) P_{0,1,y}(y_0 \neq y_1 | y_0) \leq \mathrm{TV}(P_{0,y}, P_{1,y}) + \int_{\mathbb{R}^n} P_{0,y}(dy_0) P_{0,1,y}(y_0 \neq y_1 | y_0) \leq 2\varepsilon_{\mathrm{TV}}$$

455 so

$$\int_{\{y_0 \neq y_1\}} P_{0,1,y}(dy_0) \left\| r_1(y_0) \right\|^2 \leq m^{(2)}(2\varepsilon_{\mathrm{TV}}) + \sqrt{\chi^2(P_{0,x} \| P_{1,x}) m^{(4)}(2\varepsilon_{\mathrm{TV}})},$$

456 where we used the data processing inequality.

457 For $\varphi = \varphi_{\sigma^2}$, we obtain by Lemma C.6 that the bound is

$$O\left(\sigma^2 (\varepsilon_{\mathrm{TV}} + \varepsilon_\chi \varepsilon_{\mathrm{TV}}^{1/2}) \left(d + \ln\left(\frac{1}{\varepsilon_{\mathrm{TV}}} \right) \right) \right) = O\left(\sigma^2 \varepsilon_\chi \left(d + \ln\left(\frac{1}{\varepsilon_{\mathrm{TV}}} \right) \right) \right). \qquad \square$$

458 **Lemma C.2.** *Let $\widetilde{P}^{(0)} = \widetilde{P}_0^{(0)}$ and $\widetilde{P}^{(1)} = \widetilde{P}_0^{(1)}$ be two probability distributions on $\mathbb{R}^d$ such that*
459 $\chi^2(\widetilde{P}^{(1)} \| \widetilde{P}^{(0)}) \leq \varepsilon_\chi^2 \leq 1$.

460   *1. For any $\sigma > 0$,*

$$\int \left\| \nabla \ln(\widetilde{P}^{(0)} * \varphi_{\sigma^2})(x) - \nabla \ln(\widetilde{P}^{(1)} * \varphi_{\sigma^2})(x) \right\|^2 (\widetilde{P}^{(1)} * \varphi_{\sigma^2})(dx) = O\left(\frac{\varepsilon_\chi \left(d + \ln\left(\frac{1}{\varepsilon_\chi} \right) \right)}{\sigma^2} \right).$$

461   *2. Let $\widetilde{p}_t^{(i)}$ be the density resulting from running (1) starting from $\widetilde{P}^{(i)}$, and let σ_t be as in (3).*
462   *Then for any $t > 0$,*

$$\int \left\| \nabla \ln \widetilde{p}_t^{(0)}(x) - \nabla \ln \widetilde{p}_t^{(1)}(x) \right\|^2 \widetilde{p}_t^{(1)}(x) \, dx = O\left(\frac{\varepsilon_\chi \left(d + \ln\left(\frac{1}{\varepsilon_\chi} \right) \right)}{\sigma_t^2} \right).$$

463 *Proof.* For part 1, note by (27) that

$$\nabla \ln(\widetilde{P}^{(i)} * \varphi_{\sigma^2})(y) = \frac{1}{\sigma^2} \mathbb{E}_{\widetilde{P}_{y,\sigma^2}^{(i)}} (x - y),$$

464 where $\widetilde{P}_{y,\sigma^2}^{(i)}$ is the "tilted" probability distribution defined by

$$\frac{d\widetilde{P}_{y,\sigma^2}^{(i)}}{d\widetilde{P}^{(i)}}(x) \propto e^{-\frac{\|x-y\|^2}{2\sigma^2}}.$$

465 By Bayes's rule, this can be viewed as the conditional probability that $x_0 = x$ given $x_t = y$, where
466 $x_0 \sim \widetilde{P}^{(i)}$ and $y = x_0 + \sigma\xi, \xi \sim N(0, I_d)$. Hence this fits in the framework of Lemma C.1 and

$$\int \left\| \nabla \ln(\widetilde{P}^{(0)} * \varphi_{\sigma^2})(y) - \nabla \ln(\widetilde{P}^{(1)} * \varphi_{\sigma^2})(y) \right\|^2 (\widetilde{P}^{(1)} * \varphi_{\sigma^2})(dy)$$

$$= \frac{1}{\sigma^4} \int_{\mathbb{R}^d} \left\| \mathbb{E}_{\widetilde{P}_{y,t}^{(0)}}[x] - \mathbb{E}_{\widetilde{P}_{y,t}^{(1)}}[x] \right\|^2 (\widetilde{P}^{(1)} * \varphi_{\sigma^2})(dy)$$

$$= O\left(\frac{1}{\sigma^4} \sigma^2 \varepsilon_\chi \left(d + \ln\left(\frac{1}{\varepsilon_{\mathrm{TV}}} \right) \right) \right),$$

467 giving the result.

468 For part 2, note that $\widetilde{p}_t^{(i)} = (M_{m_t \sharp} \widetilde{P}^{(i)}) * \varphi_{\sigma_t^2}$. Applying part 1 with $\widetilde{P}^{(i)} \hookleftarrow M_{m_t \sharp} \widetilde{P}^{(i)}$ (which
469 preserves χ^2-divergence) and $\sigma = \sigma_t$ gives the result.

470 $\square$

471 Finally, we argue that a score estimate that is accurate with respect to $\widehat{p}_t^{(1)}$ will still be accurate with
472 respect to $\widehat{p}_t^{(0)}$, with high probability. When using this lemma, we will substitute in the bound from
473 Lemma C.2.

474 **Lemma C.3.** *Let $\widetilde{P}_0^{(0)}$ and $\widetilde{P}_0^{(1)}$ be two probability distributions on $\mathbb{R}^d$ with TV distance ε. Suppose*
475 *the estimated score function $s_t(x)$ satisfies*

$$\left\| \nabla \ln \widetilde{p}_t^{(0)} - s_t \right\|_{L^2(\widetilde{p}_t^{(0)})}^2 = \mathbb{E}_{\widetilde{p}_t^{(0)}} \left[\left\| \nabla \ln \widetilde{p}_t^{(0)}(x) - s_t(x) \right\|^2 \right] \leq \varepsilon_t^2$$

476 *for $t \in (0, T]$, and $\nabla \ln \widetilde{p}_t^{(0)}$ is L_t-Lipschitz. Then for $t \in (0, T]$ and any $\varepsilon_\infty > 0$,*

$$P_{\widetilde{p}_t^{(1)}} \left(\left\| s_t - \nabla \ln \widetilde{p}_t^{(1)} \right\| \geq \varepsilon_\infty \right) \leq \varepsilon + \frac{4}{\varepsilon_\infty^2} \cdot \left[\varepsilon_t^2 + \int \left\| \nabla \ln \widetilde{p}_t^{(1)}(x) - \nabla \ln \widetilde{p}_t^{(0)}(x) \right\|^2 \widetilde{p}_t^{(1)}(x) \, dx \right].$$

477 *Proof.* We have

$$P_{\widetilde{p}_t^{(1)}} \left(\left\| s_t - \nabla \ln \widetilde{p}_t^{(1)} \right\| \geq \varepsilon_\infty \right)$$
$$\leq P_{\widetilde{p}_t^{(1)}} \left(\left\| s_t - \nabla \ln \widetilde{p}_t^{(0)} \right\| \geq \varepsilon_\infty/2 \right) + P_{\widetilde{p}_t^{(1)}} \left(\left\| \nabla \ln \widetilde{p}_t^{(0)} - \nabla \ln \widetilde{p}_t^{(1)} \right\| \geq \varepsilon_\infty/2 \right)$$
$$\leq \mathrm{TV}(\widetilde{p}_t^{(0)}, \widetilde{p}_t^{(1)}) + P_{\widetilde{p}_t^{(0)}} \left(\left\| s_t - \nabla \ln \widetilde{p}_t^{(0)} \right\| \geq \varepsilon_\infty/2 \right) + P_{\widetilde{p}_t^{(1)}} \left(\left\| \nabla \ln \widetilde{p}_t^{(0)} - \nabla \ln \widetilde{p}_t^{(1)} \right\| \geq \varepsilon_\infty/2 \right).$$

478 The first term is bounded by $\mathrm{TV}(\widetilde{P}^{(0)}, \widetilde{P}^{(1)}) \leq \varepsilon$. For the second term, by Chebyshev's Inequality,

$$P_{\widetilde{p}_t^{(0)}} \left(\left\| s_t - \nabla \ln \widetilde{p}_t^{(0)} \right\| \geq \varepsilon_1/2 \right) \leq \frac{4}{\varepsilon_\infty^2} \mathbb{E}_{\widetilde{p}_t^{(0)}} \left[\left\| s_t - \nabla \ln \widetilde{p}_t^{(0)} \right\|^2 \right] \leq \frac{4\varepsilon_t^2}{\varepsilon_\infty^2};$$

479 For the last term, again by Chebyshev's Inequality,

$$P_{\widetilde{p}_t^{(1)}} \left(\left\| \nabla \ln \widetilde{p}_t^{(0)} - \nabla \ln \widetilde{p}_t^{(1)} \right\| \geq \varepsilon_\infty/2 \right) \leq \frac{4}{\varepsilon_1^2} \int \left\| \nabla \ln \widetilde{p}_t^{(1)}(x) - \nabla \ln \widetilde{p}_t^{(0)}(x) \right\|^2 \widetilde{p}_t^{(1)}(x) dx.$$

480 We conclude the proof by combining the these three inequalities. $\square$

481 We will need the following to obtain a TV error bound to $\widetilde{p}_0$.

482 **Lemma C.4.** *Suppose that $\widetilde{p}_0 \propto e^{-V(x)}$ is a probability density on $\mathbb{R}^d$ with bounded first moment*
483 $\mathbb{E}_{\widetilde{p}_0} \|X\|$, *and V is L-smooth. Then for $t > 0$ such that $\alpha_t \sigma_t \leq \frac{1}{2L}$, we have*

$$\mathrm{TV}(\widetilde{p}_t, \widetilde{p}_0) \leq 2(\alpha_t - 1) \cdot (L \mathbb{E}_{\widetilde{p}_0} \|X\| + d) + \frac{3}{2} d L \alpha_t \sigma_t.$$

484 *Here $\alpha_t = 1/m_t$ and σ_t are defined in (3). In particular, $TV(\widetilde{p}_\delta, \widetilde{p}_0) \leq \varepsilon_{\mathrm{TV}}$ if $\delta = O(\frac{\varepsilon_{TV}^2}{R^2 L^2})$ and*
485 $R = \max \left\{ \sqrt{d}, \mathbb{E}_{\widetilde{p}_0} \|X\| \right\}$.

486 *Proof.* Without loss of generality, we assume that $\widetilde{p}_0(x) = e^{-V(x)}$. Note that $\widetilde{p}_t(x) =$
487 $\int \alpha_t^d \widetilde{p}_0(\alpha_t y) \varphi_{\sigma_t^2}(x - y) \, dy$. Let $\widetilde{q}_t(x) := \alpha_t^d \widetilde{p}_0(\alpha_t x)$, which is also a probability density on $\mathbb{R}^d$.
488 Then by the triangle inequality,

$$\mathrm{TV}(\widetilde{p}_t, \widetilde{p}_0) \leq \mathrm{TV}(\widetilde{p}_t, \widetilde{q}_t) + \mathrm{TV}(\widetilde{q}_t, \widetilde{p}_0).$$

489 For the second term,

$$\begin{aligned}
|\widetilde{q}_t(x) - \widetilde{p}_0(x)| &= \left| \alpha_t^d \widetilde{p}_0(\alpha_t x) - \widetilde{p}_0(x) \right| \\
&= \left| e^{-V(\alpha_t x) + d \ln \alpha_t} - e^{-V(x)} \right| \\
&\leq \max \left\{ e^{-V(x)}, e^{-V(\alpha_t x) + d \ln \alpha_t} \right\} \cdot \left(1 - e^{-|V(x) - V(\alpha_t x) + d \ln \alpha_t|} \right) \\
&\leq (\widetilde{p}_0(x) + \widetilde{q}_t(x)) \cdot (|V(x) - V(\alpha_t x)| + d \ln \alpha_t) \\
&\leq (\widetilde{p}_0(x) + \widetilde{q}_t(x)) \cdot [L \|x\| (\alpha_t - 1) + d \ln \alpha_t],
\end{aligned}$$

where in the second inequality, we use the fact that $1 - e^x \le |x|$ for all $x \le 0$. Thus

$$
\begin{aligned}
\mathrm{TV}(\widetilde{q}_t(x), \widetilde{p}_0(x)) &= \frac{1}{2} \int |\widetilde{q}_t(x) - \widetilde{p}_0(x)|\, dx \\
&\le \int \left[L\,(\alpha_t - 1)\,\|x\| + d\ln\alpha_t \right] \widetilde{p}_0(x)\, dx + \int \left[L\,(\alpha_t - 1)\,\|x\| + d\ln\alpha_t \right] \widetilde{q}_t(x)\, dx \\
&\le L(\alpha_t - 1) \left(\int \|x\|\,\widetilde{p}_0(x)dx + \int \|x\|\,\widetilde{q}_t(x)dx \right) + 2d\ln\alpha_t \\
&\le 2L(\alpha_t - 1) \int \|x\|\,\widetilde{p}_0(x)dx + 2d\ln\alpha_t.
\end{aligned}
$$

Now for the first term,

$$
\widetilde{p}_t(x) - \widetilde{q}_t(x) = \int \widetilde{q}_t(x - y)\varphi_{\sigma_t^2}(y)\, dy - \widetilde{q}_t(x) = \int \left(\widetilde{q}_t(x - \sigma_t y) - \widetilde{q}_t(x) \right) \varphi(y)dy,
$$

where $\varphi(y)$ is the density of the d-dimensional standard Gaussian distribution. Apply Minkowski's inequality for integrals:

$$
\begin{aligned}
\int |\widetilde{p}_t(x) - \widetilde{q}_t(x)|\, dx &= \int \left| \int \left(\widetilde{q}_t(x - \sigma_t y) - \widetilde{q}_t(x) \right) \varphi(y)dy \right| dx \\
&\le \int \left[\int |\widetilde{q}_t(x - \sigma_t y) - \widetilde{q}_t(x)|\, dx \right] \varphi(y)\, dy \\
&\le \int \left[\int \left(e^{L\|\alpha_t \sigma_t y\|} - 1 \right) \widetilde{q}_t(x)dx \right] \varphi(y)\, dy \\
&= \int \left(e^{L\|\alpha_t \sigma_t y\|} - 1 \right) \varphi(y)\, dy \\
&= (2\pi)^{-d/2} \int e^{L\alpha_t \sigma_t \|y\| - \frac{\|y\|^2}{2}}\, dy - 1 \\
&\le (2\pi)^{-d/2} \int e^{\left[-\frac{1}{2} + (L\alpha_t \sigma_t)^2 \right]\|y\|^2}\, dy + L\alpha_t \sigma_t \int \|y\|\,\varphi(y)\, dy - 1 \\
&\le \left[\frac{1}{1 - 2\,(L\alpha_t \sigma_t)^2} \right]^{d/2} + \sqrt{d} L\alpha_t \sigma_t - 1 \\
&\le e^{2d(L\alpha_t \sigma_t)^2} - 1 + \sqrt{d} L\alpha_t \sigma_t \\
&\le 4d\,(L\alpha_t \sigma_t)^2 + \sqrt{d} L\alpha_t \sigma_t,
\end{aligned}
$$

where in the third inequality, we use the elementary inequality $e^x \le x + e^{x^2}$, which is valid for all $x \in \mathbb{R}$, and in the fifth inequality, we use $\frac{1}{1-2x} \le e^{4x}$, which holds for $x \in [0, 1/3]$. Hence if $L\alpha_t \sigma_t \le 1/2$, we have

$$
\mathrm{TV}(\widetilde{p}_t, \widetilde{q}_t) \le \frac{1}{2} \int |\widetilde{p}_t(x) - \widetilde{q}_t(x)|\, dx \le \frac{3}{2} dL\alpha_t \sigma_t.
$$

Now we conclude the proof by combining the bounds for $\mathrm{TV}(\widetilde{p}_t, \widetilde{q}_t)$ and $\mathrm{TV}(\widetilde{p}_0, \widetilde{q}_t)$:

$$
\begin{aligned}
\mathrm{TV}(\widetilde{p}_t, \widetilde{p}_0) &\le \mathrm{TV}(\widetilde{p}_t, \widetilde{q}_t) + \mathrm{TV}(\widetilde{q}_t, \widetilde{p}_0) \\
&\le 2L(\alpha_t - 1) \int \|x\|\,\widetilde{p}_0(x)dx + 2d\ln\alpha_t + \frac{3}{2} dL\alpha_t \sigma_t \\
&\le 2\,(\alpha_t - 1) \cdot (L\mathbb{E}_{\widetilde{p}_0} \|X\| + d) + \frac{3}{2} dL\alpha_t \sigma_t,
\end{aligned}
$$

where we use the fact that $\ln x \le x - 1$ for all $x \ge 1$. Recall that $\alpha_t = 1/m_t = e^{t/2}$ and $\sigma_t^2 = 1 - e^{-t}$ when $g \equiv 1$. It suffices for

$$
\max \left\{ 2\,(L\mathbb{E}_{\widetilde{p}_0} \|X\| + d)\,(\alpha_\delta - 1),\ \frac{3}{2} dL\alpha_\delta \sigma_\delta \right\} \le \frac{\varepsilon_{\mathrm{TV}}}{2},
$$

which is implied by

$$\delta \precsim \min\left\{ \frac{\varepsilon_{\mathrm{TV}}}{L\mathbb{E}_{\widetilde{p}_0}\|X\| + d}, \frac{\varepsilon_{\mathrm{TV}}^2}{d^2 L^2}\right\} \asymp \frac{\varepsilon_{\mathrm{TV}}^2}{R^2 L^2},$$

for appropriate constants, as $R \geq \max\left\{\sqrt{d}, \mathbb{E}_{\widetilde{p}_0}\|X\|\right\}$. □

C.2 Perturbation under TV error

Although we will not need it in our proof, we note that we can derive a similar perturbation result under TV error, which might be of independent interest.

Lemma C.5. *Let $K(x, dy)$ be a probability kernel on $\mathbb{R}^d$, let $P_{0,x}, P_{1,x}$ be measures on $\mathbb{R}^d$. Let P_i denote the joint distribution of $x_i \sim P_{i,x}$ and $y_i \sim K(x_i, \cdot)$, and let $P_{i,y}$ denote the marginal distribution of y. Suppose there is a coupling $P_{0,1}$ of $(x_0, y_0) \sim P_0$ and $(x_1, y_1) \sim P_1$ such that*

- *$x_0 = x_1$ with probability $1 - \varepsilon$,*

- *$x_0 = x_1$ implies $y_0 = y_1$, and*

- *$\mathbb{E}[\|y_0 - y_1\|^2] \leq \varepsilon_W^2$.*

Define the tail error by

$$m_i(\varepsilon) := \sup_{0 \leq f \leq 1, \int_{\mathbb{R}^d} f\varphi\, dx \leq \varepsilon} \int_{\mathbb{R}^d} f(x)\|x\|^2 P_i(dx).$$

Let $r_i(y) = \int_{\mathbb{R}^d} x_i P_i(dx_i|y)$, and suppose that $r_1(y) = \int_{\mathbb{R}^d} x_1 P_1(dx_1|y)$ is L_1-Lipschitz. Then

$$\int_{\mathbb{R}^d} P_{0,y}(dy_0)\left\| \int_{\mathbb{R}^d} x_0 P_0(dx_0|y_0) - \int_{\mathbb{R}^d} x_1 P_1(dx_1|y_0)\right\|^2$$
$$\leq 4(m_0(2\varepsilon) + m_0(\varepsilon) + m_1(2\varepsilon) + m_1(\varepsilon)) + 2L_1^2 \varepsilon_W^2$$
$$\leq 4(m_0(2\varepsilon) + m_1(2\varepsilon)) + 4(1 + L_1^2)(m_0(\varepsilon) + m_1(\varepsilon)).$$

Proof. For notational clarity, we will denote draws from the conditional distribution as $\widehat{x}_0$ and $\widehat{x}_1$, for example $P_0(d\widehat{x}_0|y_0)$. We have

$$\int_{\mathbb{R}^d} P_{0,y}(dy_0)\|r_0(y_0) - r_1(y_0)\|^2 \leq 2\underbrace{\int_{\mathbb{R}^d \times \mathbb{R}^d} P_{0,1,y}(dy_0, dy_1)\|r_0(y_0) - r_1(y_1)\|^2}_{(I)}$$
$$+ 2\underbrace{\int_{\mathbb{R}^d \times \mathbb{R}^d} P_{0,1,y}(dy_0, dy_1)\|r_1(y_1) - r_1(y_0)\|^2}_{(II)}.$$

For the first term (I), we split it as

$$(I) \leq \underbrace{\int_{\{y_0 = y_1\}} P_{0,1,y}(dy_0, dy_1)\|r_0(y_0) - r_1(y_0)\|^2}_{(i)} + \underbrace{\int_{\{y_0 \neq y_1\}} P_{0,1,y}(dy_0, dy_1)\|r_0(y_0) - r_1(y_1)\|^2}_{(ii)}.$$

Define the measure Q on $\mathbb{R}^d$ by

$$Q(A) := P_{0,1}(y_0 \in A \text{ and } y_0 = y_1).$$

As in Lemma C.2, under $P_{0,1}$, when $y_0 = y_1$, we can couple $P_0(d\widehat{x}_0|y_0)$ and $P_1(d\widehat{x}_1|y_0)$ so that $x_0 = x_1$ with probability $\min\left\{\frac{dQ}{dP_{0,y}}, \frac{dQ}{dP_{1,y}}\right\}$. Let $\widehat{P}(d\widehat{x}_0, d\widehat{x}_1|y_0)$ denote this coupled distribution.

519 Then

$$(i) \le \int_{\{y_0 = y_1\}} P_{0,1,y}(dy_0, dy_1) \left\| \int_{\{\widehat{x}_0 \ne \widehat{x}_1\}} (\widehat{x}_0 - \widehat{x}_1) \widehat{P}(d\widehat{x}_0, d\widehat{x}_1 | y_0) \right\|^2$$

$$\le 2 \int_{\mathbb{R}^d} P_{0,1,y}(dy_0, dy_1) \left(\int_{\{\widehat{x}_0 \ne \widehat{x}_1\}} \|\widehat{x}_0\|^2 \, \widehat{P}(d\widehat{x}_0, d\widehat{x}_1 | y_0) + \int_{\{\widehat{x}_0 \ne \widehat{x}_1\}} \|\widehat{x}_1\|^2 \, \widehat{P}(d\widehat{x}_0, d\widehat{x}_1 | y_1) \right)$$

$$\le 2(m_0(2\varepsilon) + m_1(2\varepsilon))$$

520 as in Lemma C.2. Now

$$(ii) \le 2 \int_{\{y_0 \ne y_1\}} P_{0,1,y}(dy_0, dy_1)(\|r_0(y_0)\|^2 + \|r_1(y_1)\|^2) \le 2(m_0(\varepsilon) + m_1(\varepsilon)).$$

521 Finally, for the second term (II), we use the fact that r_1 is L_1 Lipschitz and the coupling to conclude

$$(II) \le \int_{\mathbb{R}^d} P_{0,1,y}(dy_0, dy_1) L_1^2 \|y_0 - y_1\|^2 \le L_1^2 \varepsilon_W^2.$$

522 We conclude the proof by combining the inequalities for (i), (ii), and (II).

523 For the second upper bound, we note that

$$\mathbb{E}[\|y_0 - y_1\|^2] \le 2(\mathbb{E}[\|y_0\|^2] + \mathbb{E}[\|y_1\|^2]) \le 2(m_0(\varepsilon) + m_1(\varepsilon)). \qquad \square$$

524 ### C.3 Gaussian tail calculation

525 **Lemma C.6.** *Let μ be the standard Gaussian measure on $N(0, I_d)$. Then*

$$\sup_{\mu(A) \le \varepsilon} \int_A \|x\|^2 \mu(dx) \le \varepsilon \left(2d + 3 \ln \left(\frac{1}{\varepsilon} \right) + 3 \right) = O\left(\varepsilon \left(d + \ln \left(\frac{1}{\varepsilon} \right) \right) \right)$$

$$\sup_{\mu(A) \le \varepsilon} \int_A \|x\|^4 \mu(dx) \le \varepsilon \left(2d + 3 \ln \left(\frac{1}{\varepsilon} \right) \right)^2 + 3\varepsilon \left(2d + 3 \ln \left(\frac{1}{\varepsilon} \right) \right) + 9\varepsilon = O\left(\varepsilon \left(d^2 + \ln \left(\frac{1}{\varepsilon} \right)^2 \right) \right).$$

526 *Proof.* By the χ^2 tail bound in [LM00], for $t \ge 0$,

$$\mu(\|X\|^2 \ge 2d + 3t) \le \mathbb{P}(\|X\|^2 \ge d + 2\sqrt{dt} + 2t) \le e^{-t}, \tag{33}$$

527 so $\|X\|^2$ is stochastically dominated by a random variable with cdf $F(y) = 1 - e^{-\frac{y-2d}{3}}$. Then letting
528 P_Y be the measure corresponding to F,

$$\sup_{\mu(A) \le \varepsilon} \int_A \|x\|^2 \mu(dx) \le \sup_{P_Y(A) \le \varepsilon} \int_A y P_Y(dy) = \int_{2d+3\ln\left(\frac{1}{\varepsilon}\right)}^{\infty} y dF(y)$$

$$= \varepsilon \left(2d + 3 \ln \left(\frac{1}{\varepsilon} \right) \right) + \int_{2d+3\ln\left(\frac{1}{\varepsilon}\right)}^{\infty} e^{-\frac{y-2d}{3}} dy = \varepsilon \left(2d + 3 \ln \left(\frac{1}{\varepsilon} \right) \right) + 3\varepsilon$$

529 and

$$\sup_{\mu(A) \le \varepsilon} \int_A \|x\|^4 \mu(dx) \le \sup_{P_Y(A) \le \varepsilon} \int_A y^2 P_Y(dy) = \int_{2d+3\ln\left(\frac{1}{\varepsilon}\right)}^{\infty} y^2 dF(y)$$

$$= \varepsilon \left(2d + 3 \ln \left(\frac{1}{\varepsilon} \right) \right)^2 + \int_{2d+3\ln\left(\frac{1}{\varepsilon}\right)}^{\infty} 2y e^{-\frac{y-2d}{3}} dy$$

$$= \varepsilon \left(2d + 3 \ln \left(\frac{1}{\varepsilon} \right) \right)^2 - \left[3y e^{-\frac{y-2d}{3}} \right] \Big|_{2d+3\ln\left(\frac{1}{\varepsilon}\right)}^{\infty} + \int_{2d+3\ln\left(\frac{1}{\varepsilon}\right)}^{\infty} 3e^{-\frac{y-2d}{3}} dy$$

$$= \varepsilon \left(2d + 3 \ln \left(\frac{1}{\varepsilon} \right) \right)^2 + 3\varepsilon \left(2d + 3 \ln \left(\frac{1}{\varepsilon} \right) \right) + 9\varepsilon. \qquad \square$$

D Guarantees under L^2-accurate score estimate

We will state our results under a more general tail bound assumption.

Assumption 6 (Tail bound). $R : [0, 1] \to [0, \infty)$ *is a function such that* $P_{\text{data}}(B_{R(\varepsilon)}(0)) \geq 1 - \varepsilon$.

Our result will require $R(\varepsilon)$ to grow at most as a sufficiently small power of ε^{-1} as $\varepsilon \to 0$; in particular, this holds for subexponential distributions. By taking R to be a constant function, this contains the assumption of bounded support (Assumption 2) as a special case.

D.1 TV error guarantees

We follow the framework of [LLT22] to convert guarantees under L^∞-accurate score estimate, to guarantees under L^2-accurate score estimate.

Theorem D.1 ([LLT22, Theorem 4.1]). *Let* $(\Omega, \mathcal{F}, \mathbb{P})$ *be a probability space and* $\{\mathcal{F}_n\}$ *be a filtration of the sigma field* $\mathcal{F}$. *Suppose* $X_n \sim p_n$, $Z_n \sim q_n$, *and* $\overline{Z}_n \sim \overline{q}_n$ *are* $\mathcal{F}_n$-*adapted random processes taking values in* Ω, *and* $B_n \subseteq \Omega$ *are sets such that the following hold for every* $n \in \mathbb{N}_0$.

    1. If $Z_k \in B_k^c$ *for all* $0 \leq k \leq n - 1$, *then* $Z_n = \overline{Z}_n$.

    2. $\chi^2(\overline{q}_n \| p_n) \leq D_n^2$.

    3. $\mathbb{P}(X_n \in B_n) \leq \delta_n$.

Then the following hold.

$$\text{TV}(q_n, \overline{q}_n) \leq \sum_{k=0}^{n-1}(D_k^2 + 1)^{1/2}\delta_k^{1/2} \qquad \text{TV}(p_n, q_n) \leq D_n + \sum_{k=0}^{n-1}(D_k^2 + 1)^{1/2}\delta_k^{1/2} \qquad (34)$$

Theorem D.2 (DDPM with L^2-accurate score estimate). *Let* $0 < \varepsilon_\chi, \varepsilon_{\text{TV}}, \delta < \frac{1}{2}$. *Suppose that Assumption 6 for a sufficiently small value of* c *that* R_0 *is such that* $R\left(\frac{c\varepsilon_{\text{TV}}^3 \delta^6 \varepsilon_\chi^{12}}{R_0^{19} d^5}\right) \leq R_0$, *and* $R_0^2 \geq d$. *Suppose one of the following cases holds.*

    1. Let $P_{\text{data}}, s(\cdot, t)$ *be such that Assumption 1 holds, with* $R_0^2 \geq d$. *Suppose that*

$$\varepsilon_\sigma = O\left(\frac{\varepsilon_{\text{TV}} \delta^{5/2} \varepsilon_\chi^{11/2}}{B^{9/4}}\right),$$

    where $B = R_0^4 d \ln\left(\frac{T}{\delta}\right) \ln\left(\frac{R_0 d}{\delta \varepsilon_{\text{TV}} \varepsilon_\chi}\right)$, *and we run* (5) *starting from* p_{prior} *for time* $T = \ln\left(\frac{16 R_0^2}{\varepsilon_\chi^2}\right)$, $N = O\left(\frac{B\left(T + \frac{1}{\delta^2}\right)}{\varepsilon_\chi^2}\right)$ *steps with step sizes satisfying* $h_k = O\left(\frac{\varepsilon_\chi^2}{B \max\{T - t_k, (T - t_k)^{-3}\}}\right)$.

    2. Let $P_{\text{data}}, s(\cdot, t)$ *be such that Assumptions 1 and 3 hold, with* $C \geq R_0^2$. *Suppose*

$$\varepsilon_\sigma = O\left(\frac{\varepsilon_{\text{TV}} \varepsilon_\chi^3}{T^{5/2} B}\right),$$

    where $B = C^2 d \ln\left(\frac{T}{\delta}\right) \ln\left(\frac{R_0 d}{\delta \varepsilon_{\text{TV}} \varepsilon_\chi}\right)$, *and we run* (5) *starting from* p_{prior} *for time* $T = \ln\left(\frac{16 R_0^2}{\varepsilon_\chi^2}\right)$, $N = O\left(\frac{B\left(T + \ln\left(\frac{1}{\delta}\right)\right)}{\varepsilon_\chi^2}\right)$ *steps with step sizes satisfying* $h_k = O\left(\frac{\varepsilon_\chi^2}{B \max\{T - t_k, (T - t_k)^{-1}\}}\right)$.

Then the resulting distribution q_{t_N} *is such that* q_{t_N} *is* ε_{TV}-*far in TV distance from a distribution* $\overline{q}_{t_N}$, *where* $\overline{q}_{t_N}$ *satisfies* $\chi^2(\overline{q}_{t_N} \| p_{t_N}) \leq \varepsilon_\chi^2$. *In particular, taking* $\varepsilon_\chi = \varepsilon_{\text{TV}}$, *we have* $\text{TV}(q_T, P_{\text{data}}) \leq 2\varepsilon_{\text{TV}}$.

Note that the condition on R can be satisfied if $R(\varepsilon) = o(R^{-1/19})$ (no effort has been made to optimize the exponent).

Proof. We invoke Lemma B.2 for a ε_K to be chosen, to obtain a distribution $\widetilde{P}_0$ on $B_{R_0}(0)$, where $R_0 \geq R(\varepsilon_K/8)$. Let $B = R_0^4 d \ln\left(\frac{T}{\delta}\right) \ln\left(\frac{R_0}{\delta \varepsilon_K}\right)$ and $B = C^2 d \ln\left(\frac{T}{\delta}\right) \ln\left(\frac{R_0}{\delta \varepsilon_K}\right)$ in case 1 and case 2, respectively; our choice of $\varepsilon_K = O\left(\frac{\varepsilon_{\text{TV}}^2 \delta^6}{n^2 R_0^6}\right)$ will give the definition of B in the theorem statement. In the following, we define $\widetilde{p}_t$ with $\widetilde{P}_0$, rather than P_{data}, as the initial distribution. Note that since $\text{TV}(P_{\text{data}}, \widetilde{P}_0) \leq \sqrt{\varepsilon_K} = o(\varepsilon_{\text{TV}})$ (and the same holds for their evolutions under (1)), it suffices to consider convergence to $\widetilde{p}_\delta$.

We first define the bad sets where the error in the score estimate is large,

$$B_t := \{\|\nabla \ln \widetilde{p}_t(x) - s(x,t)\| > \varepsilon_{\infty,t}\} \tag{35}$$

for some $\varepsilon_{\infty,t}$ to be chosen.

Given $t \geq 0$, let $t_- = t_k$ where k is such that $t \in [t_k, t_{k+1})$. Given bad sets B_t, define the interpolated process on $[t_k, t_{k+1})$ by

$$d\bar{z}_t = g(T-t)^2 \left(\frac{1}{2} z_t + b(z_-, T - t_-)\right) dt + g(T-t)\, dw_t \tag{36}$$

$$\text{where } b(z,t) = \begin{cases} s(z,t), & z \notin B_t \\ \nabla \ln \widetilde{p}_t(z), & z \in B_t \end{cases}.$$

In other words, simulate the reverse SDE using the score estimate as long as the point is in the good set at the previous discretization timepoint t_k, and otherwise use the actual gradient $\nabla \ln p_t$. Let $\bar{q}_t$ denote the distribution of $\bar{z}_t$ when $\bar{z}_0 \sim q_0$. Note that this process is defined only for purposes of analysis, as we do not have access to $\nabla \ln p_t$. As before, we let denote q_t the distribution of z_t defined by (12).

We can couple this process with the exponential integrator (5) using s so that as long as $x_{t_m} \notin B_{T-t_m}$, the processes agree, thus satisfying condition 1 of Theorem D.1.

Then by Lemma C.3,

$$\widetilde{P}_t^{(0)}(B_t) = \varepsilon_K + \frac{4}{\varepsilon_{\infty,t}^2}\left(\varepsilon_t^2 + O\left(\frac{\varepsilon_K\left(d + \ln\left(\frac{1}{\varepsilon_K}\right)\right)}{\sigma_t^2}\right)\right),$$

Then by choice of h_k and either Corollary A.11 or A.12, when $\int_0^{t_n} \varepsilon_t^2\, dt = O(1)$,

$$\chi^2(\bar{q}_{t_k}\|p_{t_k}) = e^\varepsilon \chi^2(q_0\|p_0) + \varepsilon + e^\varepsilon \int_0^{t_n} \varepsilon_{\infty, T-t}^2\, dt \tag{37}$$

$$\leq 2\chi^2(q_0\|p_0) + O(1),$$

where $\varepsilon = \frac{\varepsilon_\chi^2}{4}$. For $\chi^2(\bar{q}_{t_k}\|p_{t_k})$ to be bounded by ε_χ^2, it suffices for the terms in (37) to be bounded by $\frac{\varepsilon_\chi^2}{2}, \frac{\varepsilon_\chi^2}{4}, \frac{\varepsilon_\chi^2}{4}$; this is implied by

$$T = \ln\left(\frac{16R^2}{\varepsilon_\chi^2}\right) \text{ by Lemma A.14}$$

$$\int_0^{t_n} \varepsilon_{\infty, T-t}^2\, dt = O(\varepsilon_\chi^2). \tag{38}$$

By Theorem D.1,

$$\text{TV}(q_{t_n}, \bar{q}_{t_n}) \leq \sum_{k=0}^{n-1} (1 + \chi^2(q_{t_k} \| p_{t_k}))^{1/2} P(B_{t_k})^{1/2}$$

$$\leq \sum_{k=0}^{n-1} \left(2\chi^2(q_0\|p_0)^{1/2} + O(1)\right) \delta_t^{1/2} \tag{39}$$

$$= O\left(\sum_{k=0}^{n-1} \frac{\varepsilon_{t_k}}{\varepsilon_{\infty,t_k}} + \sqrt{\varepsilon_K}\left(1 + \frac{\sqrt{d + \ln(1/\varepsilon_K)}}{\varepsilon_{\infty,t_k}\sigma_{T-t_k}}\right)\right). \tag{40}$$

For this to be bounded by ε_{TV}, it suffices for

$$\sum_{k=0}^{n-1} \frac{\varepsilon_t}{\varepsilon_{\infty,t}} = O(\varepsilon_{\text{TV}}) \tag{41}$$

$$\varepsilon_K = O\left(\frac{\min_k \varepsilon_{t_k}^2 \sigma_{T-t_k}^2}{d + \ln(1/\varepsilon_K)}\right). \tag{42}$$

We bound (42) crudely, as the dependence on ε_K will be logarithmic. Using $\varepsilon_{t_k}^2 = \varepsilon_\sigma^2/\sigma_{t_k}^4$, it suffices that

$$\varepsilon_K = O\left(\frac{\varepsilon_\sigma^2}{d + \ln(1/\varepsilon_K)}\right). \tag{43}$$

We will return to this after deriving a condition on ε_σ. It remains to bound (38) and (41). We break up the timepoints depending on whether $T - t > 1$. Let

$$(t_0, t_1, \ldots, t_N) = (t_0, \ldots, t_{n^{\text{coarse}}-1}, t'_0, \ldots, t'_{n^{\text{fine}}})$$

and $u_k = T - t'_k$, where $t_{n^{\text{coarse}}-1} \leq T - 1 \leq t'_1$. Let $h'_k = t'_{k+1} - t'_k$. Note the "fine" timepoints will be closer together than the "coarse" timepoints. We break up the integral (38) and the sum (41) into the parts involving the coarse and fine timepoints. For (38), it suffices to have

$$\text{(38), coarse:} \quad \int_0^{t'_0} \varepsilon_{\infty,T-t}^2 \, dt \leq T \max_{0 \leq k \leq n^{\text{coarse}}} \varepsilon_{\infty,T-t_k}^2 = O(\varepsilon_\chi^2)$$

so it suffices to take $\varepsilon_{\infty,T-t_k}^2 \asymp \frac{\varepsilon_\chi^2}{T}$. Let $\alpha = 3$ in case 1 and $\alpha = 1$ in case 2. For the fine part, recalling our choice of h'_k, it suffices to have (note we can redefine $\varepsilon_t = \varepsilon_{t_k}$ when $t \in [t_k, t_{k+1})$ without any harm)

$$\text{(38), fine:} \quad \int_{t'_0}^{t'_{n^{\text{fine}}}} \varepsilon_{\infty,T-t}^2 \, dt = \sum_{k=0}^{n^{\text{fine}}-1} h'_k \varepsilon_{\infty,T-t'_k}^2 = O(\varepsilon_\chi^2)$$

$$\Longleftarrow \sum_{k=0}^{n^{\text{fine}}-1} \frac{\varepsilon_\chi^2 u_k^\alpha}{B} \varepsilon_{\infty,u_k}^2 = O(\varepsilon_\chi^2)$$

$$\Longleftarrow \sum_{k=0}^{n^{\text{fine}}-1} \frac{u_k^\alpha \varepsilon_{\infty,u_k}^2}{B} = O(1). \tag{44}$$

For (41), it suffices to have

$$\text{(41), coarse:} \quad \sum_{k=0}^{n^{\text{coarse}}-1} \frac{\varepsilon_{T-t_k}}{\varepsilon_{\infty,T-t_k}} \asymp n^{\text{coarse}} \frac{\varepsilon_\sigma}{\varepsilon_\chi/\sqrt{T}} = O(\varepsilon_{\text{TV}})$$

$$\Longleftarrow \varepsilon_\sigma = O\left(\frac{\varepsilon_{\text{TV}}\varepsilon_\chi}{n^{\text{coarse}}\sqrt{T}}\right) \tag{45}$$

and

$$\text{(41), fine:} \quad \sum_{k=0}^{n^{\text{fine}}-1} \frac{\varepsilon_{u_k}}{\varepsilon_{\infty,u_k}} \asymp \sum_{k=0}^{n^{\text{fine}}-1} \frac{\varepsilon_\sigma}{u_k \varepsilon_{\infty,u_k}} = O(\varepsilon_{\text{TV}}). \tag{46}$$

597 Note that in light of the required step sizes, we can take $n^{\text{coarse}} \asymp \frac{T^2 B}{\varepsilon_\chi^2}$. Considering the equality case

598 of Hölder's inequality on $(44)^{1/3}(46)^{2/3}$ suggests that we take

$$\varepsilon_\sigma \asymp \frac{\varepsilon_{\text{TV}} B^{1/2}}{\left(\sum_{k=0}^{n^{\text{fine}}-1} u_k^{\frac{\alpha-2}{3}}\right)^{3/2}} \tag{47}$$

$$\varepsilon_{\infty,u_k} \asymp \frac{B^{1/2}}{u_k^{\frac{\alpha+1}{3}} \left(\sum_{k=0}^{n^{\text{fine}}-1} u_k^{\frac{\alpha-2}{3}}\right)^{1/2}} \tag{48}$$

599 Note that the number of steps needed in the fine part is $O\left(\frac{B}{\varepsilon_\chi^2 \delta^2}\right)$ in the first case and $O\left(\frac{B}{\varepsilon_\chi^2}\right) \ln\left(\frac{1}{\delta}\right)$

600 in the second case. We can check that (47) and (48) make (44) and (46) satisfied.

601 Finally, we calculate the denominator for ε_σ. In case 1, note that starting from $T - t_0' = O(1)$ and

602 taking steps of size $h_k' \asymp \frac{\varepsilon_\chi^2}{B(T-t_k')^3}$, it takes $n^{\text{fine}} = \Theta\left(\frac{B}{\varepsilon_\chi^2 \delta^2}\right)$ steps to reach $T - t = \delta$.

$$u_k = T - t_k' = \left(1 + \Theta\left(\frac{k\varepsilon_\chi^2}{B}\right)\right)^{-\frac{1}{2}}$$

$$\sum_{k=0}^{n^{\text{fine}}-1} u_k^{1/3} \asymp \sum_{k=0}^{n^{\text{fine}}-1} \left(1 + \Theta\left(\frac{k\varepsilon_\chi^2}{B}\right)\right)^{-\frac{1}{6}} \asymp \frac{B}{\varepsilon_\chi^2}(n^{\text{fine}})^{\frac{5}{6}} \asymp \left(\frac{B}{\varepsilon_\chi^2}\right)^{11/6} \frac{1}{\delta^{5/3}}.$$

603 Then we obtain

$$\varepsilon_\sigma \asymp \varepsilon_{\text{TV}} B^{1/2} \frac{\varepsilon_\chi^{11/2}}{B^{11/4}} \delta^{5/2} = \frac{\varepsilon_{\text{TV}} \delta^{5/2} \varepsilon_\chi^{11/2}}{B^{9/4}}.$$

604 In case 1, our requirement is

$$\varepsilon_\sigma \asymp O\left(\frac{\varepsilon_{\text{TV}} \delta^{5/2} \varepsilon_\chi^{11/2}}{B^{9/4}} \wedge \frac{\varepsilon_{\text{TV}} \varepsilon_\chi^3}{T^{5/2} B}\right),$$

605 but note that the first bound is more stringent. Now, returning to (43), we see that it suffices to take

606 $\varepsilon_K = O\left(\frac{1}{d}\left(\frac{\varepsilon_{\text{TV}} \delta^{5/2} \varepsilon_\chi^{11/2}}{R_0^9 d^{9/4}}\right)^{2+\beta}\right)$ for any $\beta > 0$ (this will "solve" the $\log(1/\varepsilon_K)$ appearing in B.)

607 In case 2, we have instead $u_k = \exp\left(-\Theta\left(\frac{\varepsilon_\chi^2}{B} k\right)\right)$ so

$$\varepsilon_\sigma \asymp \varepsilon_{\text{TV}} B^{1/2} \left(\frac{\varepsilon_\chi^2}{B}\right)^{3/2} = \frac{\varepsilon_{\text{TV}} \varepsilon_\chi^3}{B}. \qquad \square$$

608 **Theorem D.3** (TV error for DDPM with L^2-accurate score estimate and smoothness)**.** *Let* $0 <$

609 $\varepsilon_{\text{TV}} < \frac{1}{2}$*. Suppose that Assumption 4 and 6 for a sufficiently small value of c that* R_0 *is such that*

610 $R\left(\frac{c\varepsilon_{\text{TV}}^{15}}{R_0^{31} d^5 L^{12}}\right) \leq R_0$*, and* $R_0^2 \geq \max\left\{d, \mathbb{E}_{P_{data}}\left[\|X\|^2\right]\right\}$*, and one of the following cases holds.*

611 *1. Let* $P_{\text{data}}, s(\cdot, t)$ *be such that Assumption 1 holds. Suppose that*

$$\varepsilon_\sigma = O\left(\frac{\varepsilon_{\text{TV}}^{11.5}}{B^{9/4} R_0^5 L^5}\right),$$

612 *where* $B = R_0^4 d \ln\left(\frac{TR_0^2 L^2}{\varepsilon_{\text{TV}}^2}\right) \ln\left(\frac{R_0^3 dL^2}{\varepsilon_{\text{TV}}^3 \varepsilon_\chi}\right)$*, and we run* (5) *starting from* p_{prior} *for time*

613 $T = \ln\left(\frac{16R_0^2}{\varepsilon_{\text{TV}}^2}\right)$, $N = O\left(\frac{B\left(T + \left(\frac{R_0 L}{\varepsilon_{\text{TV}}}\right)^4\right)}{\varepsilon_{\text{TV}}^2}\right)$ *steps with step sizes satisfying* $h_k = $

614 $O\left(\frac{\varepsilon_\chi^2}{B \max\{T-t_k, (T-t_k)^{-3}\}}\right)$*.*

2. *Let $P_{\text{data}}, s(\cdot, t)$ be such that Assumptions 1 and 3 hold, with $C \geq R_0^2$. Suppose*

$$\varepsilon_\sigma = O\left(\frac{\varepsilon_{\text{TV}}^4}{T^{5/2}B}\right),$$

where $B = C^2 d \ln\left(\frac{TR_0^2 L^2}{\varepsilon_{\text{TV}}^2}\right) \ln\left(\frac{R_0^3 dL^2}{\varepsilon_{\text{TV}}^4}\right)$, and we run (5) starting from p_{prior} for time

$$T = \ln\left(\frac{16R_0^2}{\varepsilon_{\text{TV}}^2}\right), \quad N = O\left(\frac{B\left(T + \ln\left(\frac{R_0 L}{\varepsilon_{\text{TV}}}\right)\right)}{\varepsilon_{\text{TV}}^2}\right) \text{ steps with step sizes satisfying } h_k =$$

$$O\left(\frac{\varepsilon_\chi^2}{B \max\{T - t_k, (T - t_k)^{-1}\}}\right).$$

Then the resulting distribution q_{t_N} is such that q_{t_N} is ε_{TV}-far in TV distance from the data distribution P_{data}.

Proof. With the result of Theorem D.2, we see that $\text{TV}(q_{t_N}, p_{t_N}) \leq 2\varepsilon_{\text{TV}}$. Now by Lemma C.4, if we further assume

$$\delta = O\left(\frac{\varepsilon_{\text{TV}}^2}{R_0^2 L^2}\right),$$

then $\text{TV}(p_{t_N}, P_{\text{data}}) \leq \varepsilon_{\text{TV}}$. We conclude the proof by triangle inequality and replacing the δ-dependence with $O(\frac{\varepsilon_{\text{TV}}^2}{R_0^2 L^2})$ in the previous theorem. $\square$

Proof of Theorem 2.3. If P_{data} is subexponential with a fixed constant, note that Assumption 6 holds with $R(\varepsilon) = O\left(\ln\left(\frac{1}{\varepsilon}\right)\right)$ and hence R_0 is logarithmic in all parameters. $\square$

D.2 Wasserstein error guarantees

Proof of Theorem 2.1. If $T - t_N = \delta$, then $W_2(\widetilde{p}_0, \widetilde{p}_\delta) \leq \sigma_\delta \leq \sqrt{\delta}$. Choosing $\delta = \varepsilon_W^2$, we see by Theorem D.2 it suffices to take

$$\varepsilon_\sigma = O\left(\frac{\varepsilon_{\text{TV}}^{13/2}(\varepsilon_W^2)^{5/2}}{\left(R^4 d \ln\left(\frac{T}{\delta}\right) \ln\left(\frac{RN}{\delta \varepsilon_W}\right)\right)^{9/4}}\right).$$

Simplifying gives $\varepsilon_\sigma = \widetilde{o}\left(\frac{\varepsilon_{\text{TV}}^{6.5}\varepsilon_W^5}{R^9 d^{2.25}}\right)$. If Assumption 3 also holds, then it suffices to take

$$\varepsilon_\sigma = O\left(\frac{\varepsilon_{\text{TV}}^4}{T^{5/2}C^2 d \ln\left(\frac{T}{\delta}\right) \ln\left(\frac{RN}{\delta \varepsilon_W}\right)}\right).$$

Simplifying gives $\varepsilon_\sigma = \widetilde{o}\left(\frac{\varepsilon_{\text{TV}}^4}{C^2 d}\right)$. $\square$

Proof of Theorem 2.2. To obtain purely Wasserstein error guarantees, we include an extra step of replacing any sample $z_{t_N} \sim q_{t_N}$ falling outside $B_R(0)$ by 0. Suppose $T - t_N = \delta$. Let $\widehat{q}_{t_N}$ be the resulting distribution. Then

$$W_2(\widetilde{p}_0, \widehat{q}_{t_N}) \leq W_2(\widetilde{p}_0, \widetilde{p}_\delta) + W_2(\widetilde{p}_\delta, \widehat{q}_{t_N})$$
$$\leq \sigma_\delta + W_2(\widetilde{p}_\delta, \widehat{q}_{t_N}) \leq \sqrt{\delta} + W_2(\widetilde{p}_\delta, \widehat{q}_{t_N}).$$

We choose $\delta = \frac{\varepsilon_W^2}{4}$ so the first term is $\leq \frac{\varepsilon_W}{2}$. It suffices to bound the second term $W_2(\widetilde{p}_\delta, \widehat{q}_{t_N})$ also by $\frac{\varepsilon_W}{2}$. We bound it in terms of $\text{TV}(\widetilde{p}_\delta, \widehat{q}_{t_N})$ using the fact that $\widehat{q}_{t_N}$ is supported on $B_R(0)$ and using a Gaussian tail calculation for $\widetilde{p}_\delta$. Consider a coupling of $x_{t_N} = \widetilde{x}_\delta \sim \widetilde{p}_\delta$ and $\widehat{z}_{t_N} \sim \widehat{q}_{t_N}$ such that $x_\delta \neq \widehat{z}_{t_N}$ with probability ε_{TV}. Express $\widetilde{x}_\delta = m_\delta \widetilde{x}_0 + \sigma_\delta \xi$ where $\widetilde{x}_0 \sim \widetilde{p}_0$. Now

$$\mathbb{E}[\|\widetilde{x}_\delta - \widehat{z}_{t_N}\|^2] \leq \sup_{P(A) \leq \varepsilon_{\text{TV}}} 2\left(\mathbb{E}[\|m_\delta \widetilde{x}_0 - z_{t_N}\|^2 \, \mathbb{1}_A] + \sigma_\delta^2 \mathbb{E}[\|\xi\|^2 \, \mathbb{1}_A]\right)$$
$$= 2\left(4R^2 \varepsilon_{\text{TV}} + \sigma_\delta^2 \varepsilon_{\text{TV}} \cdot O\left(d + \ln\left(\frac{1}{\varepsilon_{\text{TV}}}\right)\right)\right),$$

639 where the bound on the second term uses Lemma C.6. Using $R^2 \geq d$, we see that it suffices to choose
640 $\varepsilon_{\mathrm{TV}} = O\left(\frac{\varepsilon_W^2}{R^2}\right)$ for appropriate choice of constants. By Theorem D.2, it suffices to take

$$\varepsilon_\sigma = O\left(\frac{(\varepsilon_W^2/R^2)^{13/2}\left(\varepsilon_W^2\right)^{5/2}}{\left(R^4 d\ln\left(\frac{T}{\delta}\right)\ln\left(\frac{RN}{\delta\varepsilon_W}\right)\right)^{9/4}}\right).$$

641 Simplifying gives $\widetilde{o}\left(\frac{\varepsilon_W^{18}}{R^{22}d^{2.25}}\right)$.

642 In case 2, it suffices to take

$$\varepsilon_\sigma = O\left(\frac{(\varepsilon_W^2/R^2)^4}{T^{5/2}(C^2 d\ln\left(\frac{T}{\delta}\right)\ln\left(\frac{RN}{\delta\varepsilon_W}\right))}\right).$$

643 Simplifying gives $\varepsilon_\sigma = \widetilde{o}\left(\frac{\varepsilon_W^8}{C^2 R^8 d}\right)$. $\qquad\square$

E High-probability bound on the Hessian

645 In this section we obtain a high-probability bound on the Hessian of $\ln\widetilde{p}_t$, i.e., the Jacobian of the
646 score function.

647 To see why we expect Hessian to usually be smaller than the worst-case bound given by Lemma A.13,
648 note that we can express (27) and (28) as

$$\nabla \ln(\mu * \varphi_{\sigma^2}(y)) = -\frac{1}{\sigma^2}\mathbb{E}[Y - X|Y = y] \tag{49}$$

$$\nabla^2 \ln(\mu * \varphi_{\sigma^2}(y)) = -\frac{1}{\sigma^4}\operatorname{Cov}[Y - X|Y = y] \tag{50}$$

649 where $X \sim \mu$ and $Y = X + \sigma\xi$, $\xi \sim N(0, I_d)$. We expect that the random variable $Y - X$ is
650 distributed as $N(0, \sigma^2 I_d)$, which suggests that the covariance (50) may be bounded by $\frac{1}{\sigma^2}$ rather
651 than $\frac{1}{\sigma}$ with high probability. Indeed, we can easily construct an example where the worst case of
652 Lemma A.13 is attained—for example, $\mu = \frac{1}{2}(\delta_{-v} + \delta_v)$ for $\|v\|_2 = R$, at $x = 0$—but this point has
653 exponentially small probability density under $\mu * \varphi_{\sigma^2}$.

654 The following lemma uses a ε-net argument to bound the operator norm of the variance of a conditional
655 distribution, with high probability.

656 **Lemma E.1.** *Suppose X is a $\mathbb{R}^d$-valued random variable over the probability space $(\Omega, \mathcal{G}, P)$, and*
657 $\mathcal{F} \subseteq \mathcal{G}$ *is a σ-subalgebra. If X is subgaussian, then*

$$\mathbb{P}\left(\mathbb{E}\left[\left\|XX^\top\right\| \mid \mathcal{F}\right] \geq 2\|X\|_{\psi_2}^2 \ln\left(\frac{2 \cdot 5^d}{\varepsilon}\right)\right) \leq \varepsilon.$$

658 *Proof.* By Jensen's inequality and Markov's inequality, for any $v \in \mathbb{S}^{d-1}$,

$$\begin{aligned}
\mathbb{P}\left(\mathbb{E}[v^\top XX^\top v|\mathcal{F}] \geq \lambda^2\right) &= \mathbb{P}\left(e^{\mathbb{E}[v^\top XX^\top v|\mathcal{F}]/c^2} \geq e^{\lambda^2/c^2}\right) \\
&\leq \mathbb{P}\left(\mathbb{E}\left[e^{\langle X,v\rangle^2/c^2}|\mathcal{F}\right] \geq e^{\lambda^2/c^2}\right) \\
&\leq \frac{\mathbb{E}\left[\mathbb{E}[e^{\langle X,v\rangle^2/c^2}|\mathcal{F}]\right]}{e^{\lambda^2/c^2}} = \frac{\mathbb{E}\left[e^{\langle X,v\rangle^2/c^2}\right]}{e^{\lambda^2/c^2}} \leq 2e^{-\lambda^2/\|X\|_{\psi_2}},
\end{aligned}$$

659 where the last inequality follows from taking $c = \|X\|_{\psi_2}$. Now take a $\frac{1}{2}$-net $\mathcal{N}$ of $\mathbb{S}^{d-1}$ of size $\leq 5^d$
660 [Ver18, Cor. 4.2.13]. By a union bound,

$$\mathbb{P}\left(\exists v \in \mathcal{N} : \mathbb{E}[v^\top XX^\top v|\mathcal{F}] \geq \lambda^2\right) \leq 5^d \cdot 2 \cdot e^{-\lambda^2/\|X\|_{\psi_2}^2} = \varepsilon$$

when we take $\lambda = \|X\|_{\psi_2} \sqrt{\ln\left(\frac{2\cdot 5^d}{\varepsilon}\right)}$. By [Ver18, Lemma 4.4.1], the operator norm can be bounded by the norm on an ε-net,

$$\|A\| \leq 2 \sup_{v \in A} \|\langle A, v \rangle\| = 2 \sup_{v \in A} |v^\top A v|.$$

where the second inequality holds when A is symmetric. The result follows from applying this to $\mathbb{E}[v^\top X X^\top v | \mathcal{F}]$. $\qquad\square$

From this we obtain the desired high-probability bound.

Lemma E.2. *There is a universal constant C such that the following holds. For any starting distribution $\widetilde{P}_0$, letting $\widetilde{P}_t$ be the law of the DDPM process* (1) *at time t, we have*

$$\widetilde{P}_t\left(\left\|\nabla^2 \ln \widetilde{p}_t(x)\right\| \leq \frac{C(d + \ln\left(\frac{1}{\varepsilon}\right))}{\sigma_t^2}\right) \geq 1 - \varepsilon.$$

Note that there is no dependence on the radius.

Proof. Apply (50) with $\mu = M_{m_t \sharp} \widetilde{P}_0$ to obtain $\nabla^2 \ln \widetilde{p}_t$. Noting that $Y - X \sim N(0, \sigma^2 I_d)$ is subgaussian with $\|Y - X\|_{\psi_2} \leq C_2 \sigma$ for some universal constant C_2, the result follows from Lemma E.1. $\qquad\square$