Mutation-Driven Follow the Regularized Leader for Last-Iterate Convergence in Zero-Sum Games

Abstract

In this study, we consider a variant of the Follow the Regularized Leader (FTRL) dynamics in two-player zero-sum games. FTRL is guaranteed to converge to a Nash equilibrium when time-averaging the strategies, while many variants suffer from the issue of limit cycling behavior, i.e., lacks the last-iterate convergence guarantee. To resolve this issue, we propose a mutation-driven FTRL (M-FTRL), an algorithm that introduces mutation for the perturbation of action probabilities. We then investigate the continuous-time dynamics of M-FTRL and provide the strong convergence guarantees toward stationary points which approximate Nash equilibria under full-information feedback. Furthermore, our simulation demonstrates that M-FTRL can enjoy faster convergence rates than FTRL and optimistic FTRL under full-information feedback and surprisingly exhibits clear convergence under bandit feedback.

1 INTRODUCTION

Our study focuses on the problem of learning an equilibrium in two-player zero-sum games. In order to find an equilibrium in two-player zero-sum games, we need to solve a minimax optimization (or saddle-point optimization) in the form of $\min_x \max_y f(x, y)$. Motivated by advances of multi-agent reinforcement learning [Busoniu et al., 2008] and Generative Adversarial Networks (GANs) [Goodfellow et al., 2014], the development of algorithms that efficiently approximate the solution of the minimax optimization is attracting considerable interest [Blum and Monsour, 2007, Daskalakis et al., 2018].

There are many studies focusing on developing no-regret learning algorithms where the iterate-average strategy profile converges to a Nash equilibrium of two-player zero-sum

games [Banerjee and Peng, 2005, Zinkevich et al., 2007, Daskalakis et al., 2011]. However, well-known no-regret learning algorithms such as Follow the Regularized Leader (FTRL) are shown to cycle and fail to converge without time-averaging [Mertikopoulos et al., 2018, Bailey and Piliouras, 2018]. In recent years, several studies have developed and analyzed algorithms whose last strategy converges, such as optimistic FTRL (O-FTRL) [Daskalakis et al., 2018, Daskalakis and Panageas, 2019, Mertikopoulos et al., 2019, Wei et al., 2021, Lei et al., 2021]. This convergence property is known as *last-iterate convergence*. However, the convergence results for optimistic multiplicative weights update, which is tantamount to O-FTRL with entropy regularization, require that the equilibrium in underlying games must be unique [Daskalakis and Panageas, 2019, Wei et al., 2021].

In this study, as an alternative, we propose a mutation-driven FTRL (M-FTRL), an algorithm that introduces mutation for the perturbation of action probabilities. We first identify the discrete-time version of the M-FTRL dynamics and then modify it to the continuous-time version to provide the theoretical analysis. We prove the followings: 1) M-FTRL dynamics induced by the entropy regularizer is equivalent to replicator-mutator dynamics (RMD) [Hofbauer et al., 2009, Zagorsky et al., 2013, Bauer et al., 2019]; 2) for general regularization functions, the strategy trajectory of M-FTRL converges to a stationary point of the RMD; 3) the trajectory of M-FTRL with the entropy regularizer converges to an approximate Nash equilibrium at an exponentially fast rate. To the best of our knowledge, we are the first to provide the convergence result for RMD in two-player zero-sum games.

Furthermore, our simulation demonstrates that M-FTRL can enjoy faster convergence rates than FTRL and optimistic FTRL under full-information feedback, i.e., M-FTRL converges to a stationary point, which approximates a Nash equilibrium, faster. It also exhibits clear convergence under partial-information or bandit feedback, where each player takes the feedback about the payoffs from his or her chosen actions. We empirically observe the last-iterate convergence behavior in the M-FTRL dynamics, as well as under full-

Submitted to the 38th Conference on Uncertainty in Artificial Intelligence (UAI 2022). **To be used for reviewing only**.

Figure 1: Learning dynamics of FTRL and M-FTRL in biased Rock-Paper-Scissors. The red dot represents the Nash equilibrium point of the game.

information feedback, while neither FTRL nor O-FTRL reveals such behavior. This is surprising because it is an open question if a last-iterate convergence guarantee is provided under bandit feedback.

2 RELATED LITERATURE

Average-iterate convergence There are many previous studies focusing on developing no-regret learning algorithms enjoy average-iterate convergence in zero-sum games [Cesa-Bianchi and Lugosi, 2006, Zinkevich et al., 2007, Hofbauer et al., 2009, Syrgkanis et al., 2015]. FTRL is one of the most widely studied no-regret learning algorithm and has been shown to be convergent if the equilibrium is deterministic or strict [Mertikopoulos et al., 2018, Giannou et al., 2021]. If the equilibrium strategy is a mixed strategy with full support, FTRL's trajectory can be recurrent [Mertikopoulos et al., 2018]. For extensive-form games, counterfactual regret minimization [Zinkevich et al., 2007] and its variants have been developed as a no-regret learning algorithm [Gibson et al., 2012, Tammelin, 2014, Lanctot et al., 2017, Schmid et al., 2019, Brown and Sandholm, 2019, Davis et al., 2020]. However, most of these algorithms have not been proven that the last-iterate strategy converges.

Last-iterate convergence In recent years, various algorithms using an optimistic online learning framework [Rakhlin and Sridharan, 2013a,b] have been proposed for last-iterate convergence in minimax optimization. Optimistic gradient descent ascent [Daskalakis et al., 2018, Mertikopoulos et al., 2019, Wei et al., 2021] and optimistic multiplicative weights update [Daskalakis and Panageas, 2019, Wei et al., 2021, Lei et al., 2021] are the variants of O-FTRL, and they have been shown to enjoy the last-iterate convergence guarantee in games with a unique equilibrium. Furthermore, Nguyen et al. [2021] have proposed the no-regret learning algorithm, which exhibits the last-iterate convergence in asymmetric repeated games. In contrast to their optimistic modification of FTRL, which boosts updates for expected utitilies, our method is motivated by replicator-mutator dynamics and provides an alternative way to enjoy the last-iterate convergence guarantee.

Replicator-Mutator Dynamics Evolutionary game theory has been strongly related to learning dynamics. In fact, it is well-known that cross learning converges to the replicator dynamics (RD) in the continuous-time limit [Börgers and Sarin, 1997, Bloembergen et al., 2015], similarly to FTRL. On the other hand, RMD [Hofbauer and Sigmund, 1998] has been overlooked in the context of learning. Introducing mutation empirically makes numerical errors in computation small [Zagorsky et al., 2013]. However, it makes difficult to analyze the properties. Some notable exceptions report that mutation stabilizes the dynamics [Bomze and Burger, 1995, Bauer et al., 2019]. Let π^μ be an interior stationary point of RMD with mutation rate μ, then π^μ is ε-Nash equilibrium of the underlying game for $\varepsilon = \mu$ [Bauer et al., 2019]. Also, evolutionary game dynamics such as RD typically exhibits continua of stationary points and is unlikely to converge to a unique, stable stationary point. Mutation dissolves continua of neutrally stable equilibria into isolated, asymptotically stable ones [Bomze and Burger, 1995].

3 PRELIMINARIES

3.1 ZERO-SUM NORMAL-FORM GAME

A two-player normal-form game is defined by utility functions $u_i \in [-u_{\max}, u_{\max}]^{A_1 \times A_2}$, where A_i is the finite action space for player $i \in \{1, 2\}$. In a two-player zero-sum normal-form game, u_i satisfies $u_1(a_1, a_2) + u_2(a_1, a_2) = 0$ for all $a_1 \in A_1$ and $a_2 \in A_2$. In this game, each player i selects action $a_i \in A_i$ simultaneously. Then, player i receives utility $u_i(a_1, a_2)$. Let us denote $\pi_i \in \Delta(A_i)$ as a *mixed strategy* for player i. We define a *strategy profile* as $\pi = (\pi_1, \pi_2)$. For a given strategy profile π, the expected utility for player i is given by $v_i^\pi = \mathbb{E}_{a \sim \pi}[u_i(a_1, a_2)]$. We further define the conditional expected utility of taking ac-

tion $a_i \in A_i$ as $q_i^\pi(a_i) = \mathbb{E}_{a_{-i} \sim \pi_{-i}}[u_i(a_i, a_{-i})]$, where $-i$ represents the opponent of player i. Finally, we denote the expected utility vector as $q_i^\pi = (q_i^\pi(a_i))_{a_i \in A_i}$.

3.2 NASH EQUILIBRIUM AND EXPLOITABILITY

A common solution concept for two-player games is a *Nash equilibrium*, where no player cannot improve his/her expected utility by deviating from his/her specified strategy. In two-player zero-sum normal-form games, a Nash equilibrium $\pi^* = (\pi_1^*, \pi_2^*)$ ensures the following condition: $\forall \pi_1 \in \Delta(A_1), \forall \pi_2 \in \Delta(A_2)$,

$$v_1^{\pi_1^*, \pi_2} \geq v_1^{\pi_1^*, \pi_2^*} \geq v_1^{\pi_1, \pi_2^*}.$$

An ϵ-*Nash equilibrium* (π_1, π_2) is an approximation of a Nash equilibrium, which satisfies the following inequality:

$$\max_{\tilde{\pi}_1 \in \Delta(A_1)} v_1^{\tilde{\pi}_1, \pi_2} + \max_{\tilde{\pi}_2 \in \Delta(A_2)} v_2^{\pi_1, \tilde{\pi}_2} \leq \epsilon.$$

Furthermore, we call $\mathrm{exploit}(\pi) := \max_{\tilde{\pi}_1 \in \Delta(A_1)} v_1^{\tilde{\pi}_1, \pi_2} + \max_{\tilde{\pi}_2 \in \Delta(A_2)} v_2^{\pi_1, \tilde{\pi}_2}$ as *exploitability* of a given strategy profile π. Exploitability is a metric for measuring how close π is to a Nash equilibrium π^* in two-player zero-sum games [Johanson et al., 2011, 2012, Lockhart et al., 2019, Timbers et al., 2020, Abe and Kaneko, 2021]. From the definition, a Nash equilibrium π^* has the lowest exploitability of 0.

3.3 PROBLEM SETTING

In this study, we consider the setting where the game is played repeatedly for T iterations. At each iteration $t \in [T]$, each player i determines the (mixed) strategy $\pi_i^t \in \Delta(A_i)$ based on the past-observed feedback. Then, each player i observes the new feedback. In this study, we focus on two feedback cases: *full-information feedback* and *bandit feedback*. At the end of the iteration t under full-information feedback, player i observes the expected utility vector $(q_i^{\pi^t}(a_i))_{a_i \in A_i}$ as feedback. Under bandit feedback, each player i chooses an action a_i^t according to π_i^t. Then, each player observes the realized utility $u_i(a_1^t, a_2^t)$.

FTRL is a widely used learning algorithm in the repeated game setting. For player i, FTRL methods are defined with *regularization function* $\psi_i : \Delta(A_i) \to \mathbb{R}$, which is strictly convex and continuously differentiable on $\Delta(A_i)$. In FTRL, each player i determines her strategy π_i^t at iteration t as follows:

$$\pi_i^t = \arg\max_{p \in \Delta(A_i)} \left\{ \langle y_i^t, p \rangle - \psi_i(p) \right\},$$

$$y_i^t(a_i) = \sum_{s=1}^{t-1} q_i^{\pi^s}(a_i).$$

3.4 OTHER NOTATIONS

For a strictly convex and continuously differentiable function ψ, the associated *Bregman divergence* is defined as $D_\psi(x, x') = \psi(x) - \psi(x') - \langle \nabla\psi(x'), x - x' \rangle$. The *Kullback-Leibler divergence*, which is the Bregman divergence with the entropy regularizer $\psi(x) = \sum_i x_i \ln x_i$, is denoted by $\mathrm{KL}(x, x') = \sum_i x_i \ln \frac{x_i}{x_i'}$. Besides, we define the sum of Bregman divergences and sum of Kullback-Leibler divergences as $D_\psi(\pi, \pi') = \sum_{i=1}^{2} D_{\psi_i}(\pi_i, \pi_i')$ and $\mathrm{KL}(\pi, \pi') = \sum_{i=1}^{2} \mathrm{KL}(\pi_i, \pi_i')$, respectively.

4 MUTANT FOLLOW THE REGULARIZED LEADER

In this section, we introduce *Mutant Follow the Regularized Leader* (M-FTRL), which is inspired by the replicator-mutator dynamics [Hofbauer and Sigmund, 1998, Zagorsky et al., 2013]. Let us see what happens in a biased version of the Rock-Paper-Scissors game, see Table 1. Figure 1 compares trajectories of FTRL and M-FTRL with varying mutation parameters. Figure 1a shows that the trajectories form a cycle and never converge to the Nash equilibrium because the game is intransitive. Note, however, that the time-averaged trajectory of FTRL converges to interior Nash equilibria in zero-sum games [Hofbauer et al., 2009]. In contrast, Figures 1b and 1c exhibit a clear convergence to the unique stationary point, which is almost equivalent to the interior Nash equilibrium (the red dot) without taking the time average. As the mutation parameter increases to 1.0, although the stationary point becomes far from the Nash equilibrium, it is still asymptotically stable in Figure 1d. Thus, mutation is expected to ensure that the trajectory of a learning dynamics reaches an approximated equilibrium.

4.1 ALGORITHM

We propose a discrete-time version of the M-FTRL algorithm under two feedback cases: full-information feedback and bandit feedback. First, we provide the strategy update rule under full-information feedback:

$$\pi_i^t = \arg\max_{p \in \Delta(A_i)} \left\{ \eta \left\langle \sum_{s=1}^{t-1} q_i^{\mu, t}, p \right\rangle - \psi_i(p) \right\}, \quad (1)$$

$$q_i^{\mu, s}(a_i) = q_i^{\pi^s}(a_i) + \frac{\mu}{\pi_i^s(a_i)} \left(c_i(a_i) - \pi_i^s(a_i) \right),$$

where $\eta \in \mathbb{R}_{>0}$ is the learning rate, $\mu \in \mathbb{R}_{>0}$ is the *mutation parameter*, and $c_i \in \Delta(A_i) \cap \mathbb{R}_{>0}^{|A_i|}$ is the *relative mutation probabilities*.

As shown in Figure 1a, strategies π_i^t updated by (1) would converge to the stationary point, which is different from the Nash equilibrium of the original game. From Lemma 3.5 of [Bauer et al., 2019], the stationary point of (RMD)

Algorithm 1 Mutant Follow the Regularized Leader with adaptive relative mutation probabilities for player i.

Require: Time horizon T, learning rate η, regularization function ψ_i, mutation parameter μ, update frequency N, initial strategy π_i^0
1: $c_i \leftarrow \left(\frac{1}{|A_i|} \right)_{a_i \in A_i}$
2: $\tau \leftarrow 0$
3: Initialize z_i^0 so that $\pi_i^0 = \arg\max\limits_{p \in \Delta(A_i)} \left\{ \langle z_i^0, p \rangle - \psi_i(p) \right\}$
4: **for** $t = 1, 2, \cdots, T$ **do**
5:   Compute strategy π_i^t by
$$\pi_i^t = \arg\max_{p \in \Delta(A_i)} \left\{ \langle z_i^t, p \rangle - \psi_i(p) \right\}$$
6:   **for** $a \in A_i$ **do**
7:     $z_i^{t+1}(a) \leftarrow z_i^t(a) + \eta \left(q_i^{\pi^t}(a) + \frac{\mu}{\pi_i^t(a)} (c_i(a) - \pi_i^t(a)) \right)$
8:   **end for**
9:   $\tau \leftarrow \tau + 1$
10: **if** $\tau = N$ **then**
11:    $c_i \leftarrow \pi_i^t$
12:    $\tau \leftarrow 0$
13: **end if**
14: **end for**

is a 2μ-Nash equilibrium of the original game, and the stationary point is not Nash equilibrium unless (c_1, c_2) is a Nash equilibrium. Therefore, for convergence to a Nash equilibrium of the original game, we introduce a technique to adapt the relative mutation probabilities. That is, we copy probabilities from π_i^t into c_i every $N(\leq T)$ iterations. This technique is similar to the direct convergence method by [Perolat et al., 2021]. The pseudo-code of our algorithm with adaptive relative mutation probabilities is presented in Algorithm 1.

Under bandit feedback, each player i needs to estimate $q_i^{\mu,t} = \left(q_i^{\pi^t}(a_i) + \frac{\mu}{\pi_i^t(a_i)} (c_i(a_i) - \pi_i^t(a_i)) \right)_{a_i \in A_i}$ from the realized utility $u_i(a_1^t, a_2^t)$. Similarly to [Wei and Luo, 2018, Ito, 2021], we construct the following estimator $\hat{q}_i^{\mu,t}$:

$$\hat{q}_i^{\mu,t}(a_i) = \frac{u_i(a_1^t, a_2^t)}{\pi_i^t(a_i^t)} \mathbb{1}[a_i = a_i^t] + \frac{\mu}{\pi_i^t(a_i)} \left(c_i(a_i) - \pi_i^t(a_i) \right). \tag{2}$$

It is easy to confirm that $\hat{q}_i^{\mu,t}$ is an unbiased estimator of $q_i^{\mu,t}$. Under bandit feedback, we use $\hat{q}_i^{\mu,t}$ instead of $q_i^{\mu,t}$ in (1). Note that M-FTRL does not require any information about the opponent's strategy π_{-i}^t under bandit feedback.

5 THEORETICAL ANALYSIS

In this section, we provide the theoretical relationship to the stationary point of RMD and the bound of the exploitability. Instead of the discrete-time version of M-FTRL algorithm, we analyze the theoretical properties of the following continuous-time version of M-FTRL dynamics:

$$\pi_i^t = \arg\max_{p \in \Delta(A_i)} \left\{ \langle z_i^t, p \rangle - \psi_i(p) \right\}, \tag{3}$$

$$z_i^t(a_i) = \int_0^t \left(q_i^{\pi^s}(a_i) + \frac{\mu}{\pi_i^s(a_i)} \left(c_i(a_i) - \pi_i^s(a_i) \right) \right) ds.$$

First, we show that this dynamics is a generalization of RMD [Bauer et al., 2019]. That is, the dynamics of M-FTRL with the entropy regularizer $\psi_i(p) = \sum_{a_i \in A_i} p(a_i) \ln p(a_i)$ induces replicator-mutator dynamics:

Theorem 5.1. *The dynamics defined by (3) with the entropy regularizer $\psi_i(p) = \sum_{a_i \in A_i} p(a_i) \ln p(a_i)$ is equivalent to replicator-mutator dynamics:*

$$\frac{d}{dt}\pi_i^t(a_i) = \pi_i^t(a_i) \left(q_i^{\pi^t}(a_i) - \langle \pi_i^t, q_i^{\pi^t} \rangle \right) + \mu \left(c_i(a_i) - \pi_i^t(a_i) \right). \tag{RMD}$$

The proof of this theorem is shown in Appendix C.1.

From here, we derive the relationship between the stationary point π^μ of (RMD) and the updated strategy profile π^t. Note that, from Lemma 3.3 in [Bauer et al., 2019], for any $\mu \in \mathbb{R}_{>0}$ there exists $\pi^\mu \in \prod_{i=1}^2 \left(\Delta(A_i) \cap \mathbb{R}_{>0}^{|A_i|} \right)$ such that π^μ is a stationary point of (RMD). Thus, π^μ is well-defined. We first derive the time derivative of the (sum of) Bregman divergence between π^μ and π^t:

Theorem 5.2. *Let $\pi_i^\mu \in \Delta(A_i)$ be a stationary point of (RMD) for all $i \in \{1, 2\}$. Then, π^t updated by M-FTRL satisfies that:*

$$\frac{d}{dt} D_\psi(\pi^\mu, \pi^t)$$
$$= -\mu \sum_{i=1}^2 \sum_{a_i \in A_i} c_i(a_i) \left(\sqrt{\frac{\pi_i^t(a_i)}{\pi_i^\mu(a_i)}} - \sqrt{\frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)}} \right)^2.$$

Furthermore, if the regularizer is entropy $\psi_i(p) = \sum_{a_i \in A_i} p(a_i) \ln p(a_i)$, then π^t satisfies that:

$$\frac{d}{dt} \text{KL}(\pi^\mu, \pi^t) \leq -\mu\xi \text{KL}(\pi^\mu, \pi^t),$$

where $\xi = \min_{i \in \{1,2\}, a_i \in A_i} \frac{c_i(a_i)}{\pi_i^\mu(a_i)}$.

The first statement implies that $\frac{d}{dt} D_\psi(\pi^\mu, \pi^t) = 0$ holds if and only if $\pi^t = \pi^\mu$, and $\forall \pi^t \neq \pi^\mu, \frac{d}{dt} D_\psi(\pi^\mu, \pi^t) < 0$. Thus, by Lyapunov arguments [Khalil, 2015], the Bregman

divergence between π^μ and π^t converges to 0, and then π^t converges to π^μ. Note that Theorem 5.2 holds for all stationary points of (RMD). This means that for a fixed μ and $(c_i)_{i=1}^2$, the stationary point is unique. From the second statement, we can show that exponential convergence rates can be achieved when using the entropy regularizer:

Corollary 5.3. *Assume that the regularizer is entropy* $\psi_i(p) = \sum_{a_i \in A_i} p(a_i) \ln p(a_i)$. *Then, M-FTRL's trajectory converges to a stationary point of (RMD) exponentially fast, i.e.,*

$$\mathrm{KL}(\pi^\mu, \pi^t) \le \mathrm{KL}(\pi^\mu, \pi^0) \exp(-\mu\xi t).$$

Finally, combining this corollary and Lemma 3.5 in [Bauer et al., 2019], we can derive the exploitability bound of π^t.

Theorem 5.4. *Assume that the regularizer is entropy* $\psi_i(p) = \sum_{a_i \in A_i} p(a_i) \ln p(a_i)$. *Then, the exploitability for M-FTRL is bounded as:*

$$\mathrm{exploit}(\pi^t) \le 2\mu$$
$$+ 2u_{\max}\sqrt{(\ln 2)\mathrm{KL}(\pi^\mu, \pi^0)} \exp\left(-\frac{\mu\xi}{2}t\right).$$

Theorem 5.4 means that π^t converges to a 2μ-Nash equilibrium exponentially fast. The proof of the theorem is shown in Section 5.2.

5.1 PROOF SKETCH OF THEOREM 5.2

We sketch below the proof of Theorem 5.2. Complete proof and associated Lemmas are presented in Appendix C.2-C.4.

Proof of the first part of Theorem 5.2. First, we derive the time derivative of the Bregman divergence between $\pi \in \Delta(A_1) \times \Delta(A_2)$ and π^t:

Lemma 5.5. *For any* $\pi \in \Delta(A_1) \times \Delta(A_2)$, π^t *updated by M-FTRL satisfies that:*

$$\frac{d}{dt}D_\psi(\pi, \pi^t) = \sum_{i=1}^2 \sum_{a \in A} \pi_i^t(a_i)\pi_{-i}(a_{-i})u_i(a)$$
$$+ 2\mu - \mu \sum_{i=1}^2 \sum_{a_i \in A_i} c_i(a_i)\frac{\pi_i(a_i)}{\pi_i^t(a_i)}.$$

The proof of Lemma 5.5 stems from the fact that $D_\psi(\pi, \pi^t) = \sum_{i=1}^2 \left(\max_{p \in \Delta(A_i)}\{\langle z_i^t, p\rangle - \psi_i(p)\} - \langle z_i^t, \pi_i\rangle + \psi_i(\pi_i)\right)$.

Next, we derive the relationship between the expected utilities v^{π^μ} and v^{π_i', π_{-i}^μ} for any $\pi_i' \in \Delta(A_i)$:

Lemma 5.6. *Let* $\pi_i^\mu \in \Delta(A_i)$ *be a stationary point of (RMD) for all* $i \in \{1, 2\}$. *Then, for any* $\pi_i' \in \Delta(A_i)$:

$$\sum_{a \in A} \pi_i'(a_i)\pi_{-i}^\mu(a_{-i})u_i(a)$$
$$= \sum_{a \in A} \pi_i^\mu(a_i)\pi_{-i}^\mu(a_{-i})u_i(a) + \mu - \mu \sum_{a_i \in A_i} c_i(a_i)\frac{\pi_i'(a_i)}{\pi_i^\mu(a_i)}.$$

This result can be shown by the fact that π_i^μ is the stationary point of (RMD), i.e., $\pi_i^\mu(a_i)\left(q_i^{\pi^\mu}(a_i) - \langle\pi_i^\mu, q_i^{\pi^\mu}\rangle\right) + \mu\left(c_i(a_i) - \pi_i^\mu(a_i)\right) = 0$.

By combining Lemmas 5.5 and 5.6, we can obtain:

$$\frac{d}{dt}D_\psi(\pi^\mu, \pi^t) = \sum_{i=1}^2 \sum_{a \in A} \pi_i^t(a_i)\pi_{-i}^\mu(a_{-i})u_i(a)$$
$$+ 2\mu - \mu\sum_{i=1}^2 \sum_{a_i \in A_i} c_i(a_i)\frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)}$$
$$= \sum_{i=1}^2 \sum_{a \in A} \pi_i^\mu(a_i)\pi_{-i}^\mu(a_{-i})u_i(a)$$
$$+ 4\mu - \mu\sum_{i=1}^2 \sum_{a_i \in A_i} c_i(a_i)\left(\frac{\pi_i^t(a_i)}{\pi_i^\mu(a_i)} + \frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)}\right)$$
$$= 4\mu - \mu\sum_{i=1}^2 \sum_{a_i \in A_i} c_i(a_i)\left(\frac{\pi_i^t(a_i)}{\pi_i^\mu(a_i)} + \frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)}\right)$$
$$= -\mu\sum_{i=1}^2 \sum_{a_i \in A_i} c_i(a_i)\left(\sqrt{\frac{\pi_i^t(a_i)}{\pi_i^\mu(a_i)}} - \sqrt{\frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)}}\right)^2,$$

where we use $\sum_{i=1}^2 \sum_{a \in A} \pi_i^\mu(a_i)\pi_{-i}^\mu(a_{-i})u_i(a) = \sum_{i=1}^2 v_i^{\pi^\mu} = 0$. This concludes the first statement of the theorem.

Proof of the second part of Theorem 5.2. Let us define $\xi_i = \min_{a_i \in A_i} \frac{c_i(a_i)}{\pi_i^\mu(a_i)}$. From the first part of the theorem, we have:

$$\frac{d}{dt}D_\psi(\pi^\mu, \pi^t)$$
$$= -\mu\sum_{i=1}^2 \sum_{a_i \in A_i} c_i(a_i)\left(\frac{\pi_i^t(a_i)}{\pi_i^\mu(a_i)} + \frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)} - 2\right)$$
$$= -\mu\sum_{i=1}^2 \sum_{a_i \in A_i} \frac{c_i(a_i)}{\pi_i^\mu(a_i)}\frac{(\pi_i^t(a_i) - \pi_i^\mu(a_i))^2}{\pi_i^t(a_i)}$$
$$\le -\mu\sum_{i=1}^2 \xi_i \sum_{a_i \in A_i} \frac{(\pi_i^t(a_i) - \pi_i^\mu(a_i))^2}{\pi_i^t(a_i)}$$
$$\le -\mu\sum_{i=1}^2 \xi_i \ln\left(1 + \sum_{a_i \in A_i} \frac{(\pi_i^t(a_i) - \pi_i^\mu(a_i))^2}{\pi_i^t(a_i)}\right)$$
$$= -\mu\sum_{i=1}^2 \xi_i \ln\left(\sum_{a_i \in A_i} \pi_i^\mu(a_i)\frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)}\right)$$

$$\leq -\mu \sum_{i=1}^{2} \xi_i \sum_{a_i \in A_i} \pi_i^{\mu}(a_i) \ln \left(\frac{\pi_i^{\mu}(a_i)}{\pi_i^t(a_i)} \right)$$

$$= -\mu \sum_{i=1}^{2} \xi_i \mathrm{KL}(\pi_i^{\mu}, \pi_i^t) \leq -\mu \xi \sum_{i=1}^{2} \mathrm{KL}(\pi_i^{\mu}, \pi_i^t), \quad (4)$$

where the second inequality follows from $x \geq \ln(1+x)$ for all $x > 0$, and the third inequality follows from the concavity of the $\ln(\cdot)$ function and Jensen's inequality for concave functions. On the other hand, if $\psi_i(p) = \sum_{a_i \in A_i} p(a_i) \ln p(a_i)$, then $D_{\psi_i}(\pi_i^{\mu}, \pi_i^t) = \mathrm{KL}(\pi_i^{\mu}, \pi_i^t)$. Thus, we have $D_{\psi}(\pi^{\mu}, \pi^t) = \sum_{i=1}^{2} \mathrm{KL}(\pi_i^{\mu}, \pi_i^t)$. From this fact and (4), we have:

$$\frac{d}{dt} \mathrm{KL}(\pi^{\mu}, \pi^t) \leq -\mu \xi \mathrm{KL}(\pi^{\mu}, \pi^t).$$

This concludes the second statement of the theorem. $\quad\square$

5.2 PROOF OF THEOREM 5.4

From the definition of exploitability, we have:

$$\mathrm{exploit}(\pi^t) = \sum_{i=1}^{2} \max_{\tilde{\pi}_i \in \Delta(A_i)} v_i^{\tilde{\pi}_i, \pi_{-i}^t}$$

$$= \sum_{i=1}^{2} \left(\max_{\tilde{\pi}_i \in \Delta(A_i)} v_i^{\tilde{\pi}_i, \pi_{-i}^{\mu}} \right.$$
$$\left. + \max_{\tilde{\pi}_i \in \Delta(A_i)} v_i^{\tilde{\pi}_i, \pi_{-i}^t} - \max_{\tilde{\pi}_i \in \Delta(A_i)} v_i^{\tilde{\pi}_i, \pi_{-i}^{\mu}} \right)$$

$$\leq \sum_{i=1}^{2} \left(\max_{\tilde{\pi}_i \in \Delta(A_i)} v_i^{\tilde{\pi}_i, \pi_{-i}^{\mu}} + \max_{\tilde{\pi}_i \in \Delta(A_i)} \left(v_i^{\tilde{\pi}_i, \pi_{-i}^t} - v_i^{\tilde{\pi}_i, \pi_{-i}^{\mu}} \right) \right)$$

$$\leq \sum_{i=1}^{2} \left(\max_{\tilde{\pi}_i \in \Delta(A_i)} v_i^{\tilde{\pi}_i, \pi_{-i}^{\mu}} \right.$$
$$\left. + \|\pi_i^{\mu} - \pi_i^t\|_1 \max_{\tilde{\pi}_{-i} \in \Delta(A_{-i})} \|q_i^{\pi_i^t, \tilde{\pi}_{-i}}\|_{\infty} \right)$$

$$\leq \sum_{i=1}^{2} \left(\max_{\tilde{\pi}_i \in \Delta(A_i)} v_i^{\tilde{\pi}_i, \pi_{-i}^{\mu}} + u_{\max} \sqrt{2(\ln 2)\mathrm{KL}(\pi_i^{\mu}, \pi_i^t)} \right),$$
$$(5)$$

where the second inequality follows from Hölder's inequality, and the last inequality follows from Lemma 11.6.1 in [Cover and Thomas, 2006].

From Lemma 3.5 of [Bauer et al., 2019], a stationary point π^{μ} of (RMD) satisfies that for all $i \in \{1, 2\}$ and $a_i \in A_i$, $q_i^{\pi^{\mu}}(a_i) - \langle \pi_i^t, q_i^{\pi^{\mu}} \rangle \leq \mu$. Therefore, the term of $\max_{\tilde{\pi}_i \in \Delta(A_i)} v_i^{\tilde{\pi}_i, \pi_{-i}^{\mu}}$ can be bounded as:

$$\sum_{i=1}^{2} \max_{\tilde{\pi}_i \in \Delta(A_i)} v_i^{\tilde{\pi}_i, \pi_{-i}^{\mu}} = \sum_{i=1}^{2} \left(\max_{\tilde{\pi}_i \in \Delta(A_i)} v_i^{\tilde{\pi}_i, \pi_{-i}^{\mu}} - v_i^{\pi_i^{\mu}, \pi_{-i}^{\mu}} \right)$$

$$= \sum_{i=1}^{2} \left(\max_{a_i \in A_i} q_i^{\pi^{\mu}}(a_i) - \langle \pi_i^{\mu}, q_i^{\pi^{\mu}} \rangle \right) \leq 2\mu. \quad (6)$$

By combining (5), (6), and Corollary 5.3, we have:

$$\mathrm{exploit}(\pi^t) \leq 2\mu + u_{\max} \sum_{i=1}^{2} \sqrt{2(\ln 2)\mathrm{KL}(\pi_i^{\mu}, \pi_i^t)}$$

$$\leq 2\mu + u_{\max} \sqrt{2(\ln 2)} \sqrt{2 \sum_{i=1}^{2} \mathrm{KL}(\pi_i^{\mu}, \pi_i^t)}$$

$$\leq 2\mu + 2\sqrt{\ln 2} u_{\max} \sqrt{\mathrm{KL}(\pi^{\mu}, \pi^0) \exp(-\mu\xi t)}$$

$$= 2\mu + 2u_{\max} \sqrt{(\ln 2)\mathrm{KL}(\pi^{\mu}, \pi^0)} \exp\left(-\frac{\mu\xi}{2} t\right),$$

where the second inequality follows from $\sqrt{a} + \sqrt{b} \leq \sqrt{2(a+b)}$ for $a, b > 0$. This concludes the statement. $\quad\square$

6 EXPERIMENTS

In this section, we empirically evaluate M-FTRL. We compare its performance to those of FTRL and O-FTRL.

We conduct experiments on the following games: biased rock-paper-scissors (BRPS), a normal-form game with multiple Nash equilibria (M-Eq), and random utility games. BRPS and M-Eq have the following utility matrix, respectively:

Table 1: Biased RPS utilities

	R	P	S
R	0	-0.1	0.3
P	0.1	0	-0.1
S	-0.3	0.1	0

Table 2: M-Eq utilities

	y_1	y_2
x_1	0.1	-0.2
x_2	-0.4	0.3
x_3	-1	0.9

The set of Nash equilibria in M-Eq is given by:

$$\Pi_1^* = \left\{ x \in \Delta^3 \mid x_2 = -\frac{22}{12}x_1 + \frac{19}{12}; \ x_3 = \frac{10}{12}x_1 - \frac{7}{12} \right\},$$

$$\Pi_2^* = \left\{ \left(\frac{1}{2}, \frac{1}{2} \right) \right\}.$$

For random utility games, we generate each component in a utility matrix uniformly at random in $[0, 1]$. We consider random utility games with action sizes $|A_1| = |A_2| = 10$ and $|A_1| = |A_2| = 50$. For each game, we average the results for 100 instances. We generate the initial strategy profile π^0 uniformly at random in $\Delta(A_1) \times \Delta(A_2)$ for each instance. We use the entropy regularizer $\phi_i(p) = \sum_{a_i \in A_i} p(a_i) \ln p(a_i)$ in all experiments.

6.1 FULL-INFORMATION FEEDBACK

First, we provide the results under full-information feedback. In these experiments, we analyze the performance of M-FTRL with fixed relative mutation probabilities $c_i = \frac{1}{|A_i|}$

Figure 2: Exploitability of π^t for M-FTRL, FTRL, and O-FTRL under full-information feedback.

(a) M-FTRL with fixed relative mutation probabilities

(b) M-FTRL with adaptive relative mutation probabilities

(c) FTRL

(d) O-FTRL

Figure 3: Trajectories of π^t for M-FTRL, FTRL and O-FTRL in BRPS under full-information feedback. We set the initial strategy profile to $\pi_i^0 = \frac{1}{|A_i|}$ for $i \in \{1, 2\}$. The black point represents the equilibrium strategy. The blue/red points represent the initial/final points, respectively.

(a) M-FTRL with fixed relative mutation probabilities

(b) O-FTRL

Figure 4: Initial strategies and final strategies for player 1 in 100 instances (M-Eq under full-information feedback). The green line represents the set of equilibrium strategies for player 1. The blue/red points represent the initial/final points, respectively.

and one with adaptive relative mutation probabilities (Algorithm 1). We set the learning rate to $\eta = 10^{-1}$ for all algorithms, and set the mutation parameter to $\mu = 10^{-2}$ for M-FTRL. For M-FTRL with adaptive relative mutation probabilities, we set $N = 4,000$ in BRPS and M-Eq, and $N = 20,000$ in the random utility games.

Figure 2 shows the average exploitability of π^t updated by each algorithm. We find that the exploitability of M-FTRL converges to a constant value faster than FTRL and O-FTRL. Furthermore, by adapting the relative mutation probabilities, the exploitability of M-FTRL's strategy profile quickly converges to 0.

Next, we compare the trajectories of strategies updated by each algorithm. Figure 3 shows the trajectories of π^t updated by each algorithm from an instance of RBPS. Note that in this figure, we set the initial strategy to $\pi_i^0 = \frac{1}{|A_i|}$ for $i \in \{1, 2\}$. We can observe that FTRL's strategies cycle around the Nash equilibrium strategy, and O-FTRL's strategies gradually approach the Nash equilibrium strategy. Unlike these methods, M-FTRL's strategies quickly approach the stationary point. Figure 4 shows the initial strategies and final strategies for player 1 in M-Eq. We find that M-FTRL's strategy profile converges to a unique stationary point regardless of the setting of the initial point, while O-FTRL's strategy profile converges to a different Nash equilibrium for each instance. This result highlights the uniqueness property of the stationary point from Theorem 5.2.

6.2 BANDIT FEEDBACK

Next, we provide the results under bandit feedback. We set the learning rate to $\eta = 10^{-4}$ for all algorithms, and set the mutation parameter to $\mu = 10^{-2}$ for M-FTRL. In the bandit feedback experiments, we focus on the performance of M-FTRL with fixed relative mutation probabilities $c_i = \frac{1}{|A_i|}$. In FTRL and O-FTRL algorithms, we use the unbiased estimator by [Lattimore and Szepesvári, 2020] as the estimator of $q_i^{\pi^t}$ so that the estimator takes values in

Figure 5: Exploitability of π^t for M-FTRL, FTRL, and O-FTRL under bandit feedback.

(a) M-FTRL with fixed relative mutation probabilities

(b) FTRL

(c) O-FTRL

Figure 6: Trajectories of π^t for M-FTRL, FTRL and O-FTRL in BRPS under bandit feedback. We set the initial strategy profile to $\pi_i^0 = \frac{1}{|A_i|}$ for $i \in \{1, 2\}$. The black point represents the equilibrium strategy. The blue/red points represent the initial/final points, respectively.

$(-\infty, u_{\max}]$ for computational stability. We provide further details on the estimator in Appendix A. Note that M-FTRL does not need this estimator, but it is sufficient to use the importance-weighted estimator in (2).

Figure 5 shows the average exploitability of π^t updated by each algorithm, and Figure 6 shows the trajectories of π^t updated by each algorithm from an instance of RBPS. We can see that unlike the experimental results under full-information feedback, O-FTRL's trajectory does not converge to a Nash equilibrium. On the other hand, M-FTRL's trajectory converges near a stationary point. These results suggest that M-FTRL has the last-iterate convergence property even under bandit feedback.

7 CONCLUSION

In this study, we proposed M-FTRL, a simple FTRL algorithm that incorporates mutation for last-iterate convergence to a stationary point. We proved that the M-FTRL dynamics induced by the entropy regularizer is equivalent to RMD. Besides, we showed that the trajectory of M-FTRL with general regularization functions converges to a stationary point of the RMD. The numerical simulation reveals that M-FTRL outperforms the state-of-the-art FTRL and O-FTRL

in a variety of two-person zero-sum games. In future studies, we will extend M-FTRL algorithm and provide a theoretical analysis to more complex games, such as extensive-form games and Markov games.

References

Kenshi Abe and Yusuke Kaneko. Off-policy exploitability-evaluation in two-player zero-sum markov games. In *AAMAS*, 2021.

James P Bailey and Georgios Piliouras. Multiplicative weights update in zero-sum games. In *Economics and Computation*, pages 321–338, 2018.

Bikramjit Banerjee and Jing Peng. Efficient no-regret multi-agent learning. In *AAAI*, pages 41–46, 2005.

Johann Bauer, Mark Broom, and Eduardo Alonso. The stabilization of equilibria in evolutionary game dynamics through mutation: mutation limits in evolutionary games. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 475(2231):20190355, 2019.

Daan Bloembergen, Karl Tuyls, Daniel Hennes, and Michael Kaisers. Evolutionary dynamics of multi-agent

learning: A survey. *Journal of Artificial Intelligence Research*, 53:659–697, 2015.

Avrim Blum and Yishay Monsour. Learning, regret minimization, and equilibria. In *Algorithmic game theory*, pages 79–101. Cambridge University Press, 2007.

Immanuel M Bomze and Reinhard Burger. Stability by mutation in evolutionary games. *Games and Economic Behavior*, 11(2):146–172, 1995.

Noam Brown and Tuomas Sandholm. Solving imperfect-information games via discounted regret minimization. In *AAAI*, pages 1829–1836, 2019.

Lucian Busoniu, Robert Babuska, and Bart De Schutter. A comprehensive survey of multiagent reinforcement learning. *IEEE Transactions on Systems, Man, and Cybernetics, Part C (Applications and Reviews)*, 38(2):156–172, 2008.

Tilman Börgers and Rajiv Sarin. Learning through reinforcement and replicator dynamics. *Journal of Economic Theory*, 77(1):1–14, 1997.

Nicolo Cesa-Bianchi and Gábor Lugosi. *Prediction, learning, and games*. Cambridge university press, 2006.

Thomas M. Cover and Joy A. Thomas. *Elements of information theory 2nd Edition*. Wiley-Interscience, 2006.

Constantinos Daskalakis and Ioannis Panageas. Last-iterate convergence: Zero-sum games and constrained min-max optimization. *Innovations in Theoretical Computer Science*, 2019.

Constantinos Daskalakis, Alan Deckelbaum, and Anthony Kim. Near-optimal no-regret algorithms for zero-sum games. In *Symposium on Discrete Algorithms*, pages 235–254, 2011.

Constantinos Daskalakis, Andrew Ilyas, Vasilis Syrgkanis, and Haoyang Zeng. Training gans with optimism. In *ICLR*, 2018.

Trevor Davis, Martin Schmid, and Michael Bowling. Low-variance and zero-variance baselines for extensive-form games. In *ICML*, pages 2392–2401, 2020.

Angeliki Giannou, Emmanouil Vlatakis-Gkaragkounis, and Panayotis Mertikopoulos. The convergence rate of regularized learning in games: From bandits and uncertainty to optimism and beyond. In *NeurIPS*, 2021.

Richard G Gibson, Marc Lanctot, Neil Burch, Duane Szafron, and Michael Bowling. Generalized sampling and variance in counterfactual regret minimization. In *AAAI*, pages 1355–1361, 2012.

Ian Goodfellow, Jean Pouget-Abadie, Mehdi Mirza, Bing Xu, David Warde-Farley, Sherjil Ozair, Aaron Courville, and Yoshua Bengio. Generative adversarial nets. In *NeurIPS*, 2014.

Josef Hofbauer and Karl Sigmund. *Evolutionary Games and Population Dynamics*. Cambridge University Press, Cambridge, 1998.

Josef Hofbauer, Sylvain Sorin, and Yannick Viossat. Time average replicator and best-reply dynamics. *Mathematics of Operations Research*, 34(2):263–269, 2009.

Shinji Ito. Parameter-free multi-armed bandit algorithms with hybrid data-dependent regret bounds. In *COLT*, pages 2552–2583, 2021.

Michael Johanson, Kevin Waugh, Michael Bowling, and Martin Zinkevich. Accelerating best response calculation in large extensive games. In *IJCAI*, 2011.

Michael Johanson, Nolan Bard, Neil Burch, and Michael Bowling. Finding optimal abstract strategies in extensive-form games. In *AAAI*, pages 1371–1379, 2012.

Hassan K. Khalil. *Nonlinear Control, Global Edition*. Pearson Education, 2015.

Marc Lanctot, Vinicius Zambaldi, Audrunas Gruslys, Angeliki Lazaridou, Karl Tuyls, Julien Perolat, David Silver, and Thore Graepel. A unified game theoretic approach to multiagent reinforcement learning. In *NeurIPS*, page 4190–4203, 2017.

Tor Lattimore and Csaba Szepesvári. *Bandit algorithms*. Cambridge University Press, 2020.

Qi Lei, Sai Ganesh Nagarajan, Ioannis Panageas, et al. Last iterate convergence in no-regret learning: constrained min-max optimization for convex-concave landscapes. In *AISTATS*, pages 1441–1449, 2021.

Edward Lockhart, Marc Lanctot, Julien Pérolat, Jean-Baptiste Lespiau, Dustin Morrill, Finbarr Timbers, and Karl Tuyls. Computing approximate equilibria in sequential adversarial games by exploitability descent. *arXiv preprint arXiv:1903.05614*, 2019.

Panayotis Mertikopoulos, Christos Papadimitriou, and Georgios Piliouras. Cycles in adversarial regularized learning. In *Symposium on Discrete Algorithms*, pages 2703–2717, 2018.

Panayotis Mertikopoulos, Bruno Lecouat, Houssam Zenati, Chuan-Sheng Foo, Vijay Chandrasekhar, and Georgios Piliouras. Optimistic mirror descent in saddle-point problems: Going the extra (gradient) mile. In *ICLR*, 2019.

Tri-Dung Nguyen, Alain B Zemhoho, Long Tran-Thanh, et al. Last round convergence and no-dynamic regret in asymmetric repeated games. In *ALT*, pages 553–577, 2021.

Julien Perolat, Remi Munos, Jean-Baptiste Lespiau, Shayegan Omidshafiei, Mark Rowland, Pedro Ortega, Neil Burch, Thomas Anthony, David Balduzzi, Bart De Vylder, et al. From poincaré recurrence to convergence in imperfect information games: Finding equilibrium via regularization. In *ICML*, pages 8525–8535, 2021.

Alexander Rakhlin and Karthik Sridharan. Online learning with predictable sequences. In *COLT*, pages 993–1019, 2013a.

Sasha Rakhlin and Karthik Sridharan. Optimization, learning, and games with predictable sequences. In *NeurIPS*, 2013b.

Martin Schmid, Neil Burch, Marc Lanctot, Matej Moravcik, Rudolf Kadlec, and Michael Bowling. Variance reduction in monte carlo counterfactual regret minimization (vr-mccfr) for extensive form games using baselines. In *AAAI*, pages 2157–2164, 2019.

Shai Shalev-Shwartz et al. Online learning and online convex optimization. *Foundations and trends in Machine Learning*, 4(2):107–194, 2011.

Vasilis Syrgkanis, Alekh Agarwal, Haipeng Luo, and Robert E Schapire. Fast convergence of regularized learning in games. In *NeurIPS*, 2015.

Oskari Tammelin. Solving large imperfect information games using cfr+. *arXiv preprint arXiv:1407.5042*, 2014.

Finbarr Timbers, Edward Lockhart, Marc Lanctot, Martin Schmid, Julian Schrittwieser, Thomas Hubert, and Michael Bowling. Approximate exploitability: Learning a best response in large games. *arXiv preprint arXiv:2004.09677*, 2020.

Chen-Yu Wei and Haipeng Luo. More adaptive algorithms for adversarial bandits. In *COLT*, pages 1263–1291, 2018.

Chen-Yu Wei, Chung-Wei Lee, Mengxiao Zhang, and Haipeng Luo. Linear last-iterate convergence in constrained saddle-point optimization. In *ICLR*, 2021.

Benjamin M. Zagorsky, Johannes G. Reiter, Krishnendu Chatterjee, and Martin A. Nowak. Forgiver triumphs in alternating prisoner's dilemma. *PLOS ONE*, pages 1–8, 2013.

Martin Zinkevich, Michael Johanson, Michael Bowling, and Carmelo Piccione. Regret minimization in games with incomplete information. In *NeurIPS*, pages 1729–1736, 2007.

A UNBIASED ESTIMATOR FOR FTRL AND O-FTRL UNDER BANDIT FEEDBACK

For FTRL and O-FTRL under bandit feedback, we use the following unbiased estimator of $q_i^{\pi^t}$ which is proposed by [Lattimore and Szepesvári, 2020]:

$$\hat{q}_i^{\pi^t}(a_i) = u_{\max} - \frac{u_{\max} - u_i(a_1^t, a_2^t)}{\pi_i^t(a_i^t)}\mathbb{1}[a_i = a_i^t].$$

This estimator takes values in $(-\infty, u_{\max}]$ while the standard importance-weighted estimator takes values in $(-\infty, \infty)$.

B ADDITIONAL LEMMAS

Lemma B.1. *For any $\pi \in \Delta(A_1) \times \Delta(A_2)$, π^t updated by M-FTRL satisfies that:*

$$D_\psi(\pi, \pi^t) = \sum_{i=1}^{2}\left(\max_{p \in \Delta(A_i)}\left\{\langle z_i^t, p\rangle - \psi_i(p)\right\} - \langle z_i^t, \pi_i\rangle + \psi_i(\pi_i)\right).$$

Lemma B.2. *Let $\pi_i^\mu \in \Delta(A_i)$ be a stationary point of (RMD). If $c_i \in \Delta(A_i) \cap \mathbb{R}_{>0}^{A_i}$ and $\mu > 0$, then we also have $\pi_i^\mu \in \Delta(A_i) \cap \mathbb{R}_{>0}^{A_i}$.*

C PROOFS

C.1 PROOF OF THEOREM 5.1

Proof of Theorem 5.1. By the method of Lagrange multiplier, we have:

$$\pi_i^t(a_i) = \frac{\exp\left(z_i^t(a_i)\right)}{\sum_{a_i' \in A_i}\exp\left(z_i^t(a_i')\right)}.$$

Therefore, the time derivative of $\pi_i^t(a_i)$ is given as follows:

$$\begin{aligned}
\frac{d}{dt}\pi_i^t(a_i) &= \frac{\frac{d}{dt}\exp\left(z_i^t(a_i)\right)}{\sum_{a_i' \in A_i}\exp\left(z_i^t(a_i')\right)} - \frac{\exp\left(z_i^t(a_i)\right)\frac{d}{dt}\left(\sum_{a_i' \in A_i}\exp\left(z_i^t(a_i')\right)\right)}{\left(\sum_{a_i' \in A_i}\exp\left(z_i^t(a_i')\right)\right)^2}\\
&= \frac{\exp\left(z_i^t(a_i)\right)\frac{d}{dt}z_i^t(a_i)}{\sum_{a_i' \in A_i}\exp\left(z_i^t(a_i')\right)} - \frac{\exp\left(z_i^t(a_i)\right)\left(\sum_{a_i' \in A_i}\exp\left(z_i^t(a_i')\right)\frac{d}{dt}z_i^t(a_i')\right)}{\left(\sum_{a_i' \in A_i}\exp\left(z_i^t(a_i')\right)\right)^2}\\
&= \pi_i^t(a_i)\frac{d}{dt}z_i^t(a_i) - \pi_i^t(a_i)\sum_{a_i' \in A_i}\pi_i^t(a_i')\frac{d}{dt}z_i^t(a_i').
\end{aligned}$$

From the definition of $z_i^t(a_i)$, we have:

$$\frac{d}{dt}z_i^t(a_i) = q_i^{\pi^t}(a_i) + \frac{\mu}{\pi_i^t(a_i)}\left(c_i(a_i) - \pi_i^t(a_i)\right).$$

By combining these equalities, we get:

$$\begin{aligned}
\frac{d}{dt}\pi_i^t(a_i) =&\, \pi_i^t(a_i)\left(q_i^{\pi^t}(a_i) + \frac{\mu}{\pi_i^t(a_i)}\left(c_i(a_i) - \pi_i^t(a_i)\right) - \sum_{a_i' \in A_i}\pi_i^t(a_i')\left(q_i^{\pi^t}(a_i') + \frac{\mu}{\pi_i^t(a_i')}\left(c_i(a_i') - \pi_i^t(a_i')\right)\right)\right)\\
=&\, \pi_i^t(a)\left(q_i^{\pi^t}(a_i) - \langle\pi_i^t, q_i^{\pi^t}\rangle\right) + \mu\left(c_i(a_i) - \pi_i^t(a_i)\right) - \mu\pi_i^t(a_i)\sum_{a_i' \in A_i}\left(c_i(a_i') - \pi_i^t(a_i')\right)\\
=&\, \pi_i^t(a)\left(q_i^{\pi^t}(a_i) - \langle\pi_i^t, q_i^{\pi^t}\rangle\right) + \mu\left(c_i(a_i) - \pi_i^t(a_i)\right).
\end{aligned}$$

$\square$

C.2 PROOF OF LEMMA 5.5

Proof of Lemma 5.5. Let us define $\psi_i^*(z_i) = \max_{p \in \Delta(A_i)} \{\langle z_i, p \rangle - \psi_i(p)\}$. Then, from Lemma B.1, the time derivative of $D_\psi(\pi, \pi^t)$ is given as:

$$
\begin{aligned}
\frac{d}{dt} D_\psi(\pi, \pi^t) &= \sum_{i=1}^2 \frac{d}{dt} \left(\max_{p \in \Delta(A_i)} \{\langle z_i^t, p \rangle - \psi_i(p)\} - \langle z_i^t, \pi_i \rangle + \psi_i(\pi_i) \right) \\
&= \sum_{i=1}^2 \frac{d}{dt} \left(\psi_i^*(z_i^t) - \langle z_i^t, \pi_i \rangle \right) \\
&= \sum_{i=1}^2 \left(\left\langle \frac{d}{dt} z_i^t, \nabla \psi_i^*(z_i^t) \right\rangle - \left\langle \frac{d}{dt} z_i^t, \pi_i \right\rangle \right) \\
&= \sum_{i=1}^2 \left\langle \frac{d}{dt} z_i^t, \nabla \psi_i^*(z_i^t) - \pi_i \right\rangle.
\end{aligned}
$$

From maximizing argument of [Shalev-Shwartz et al., 2011], we have $\nabla \psi_i^*(z_i) = \arg \max_{p \in \Delta(A_i)} \{\langle z_i, p \rangle - \psi_i(p)\}$ and then $\nabla \psi_i^*(z_i^t) = \pi_i^t$. Furthermore, from the definition of $z_i^t(a_i)$, we have $\frac{d}{dt} z_i^t(a_i) = q_i^{\pi^t}(a_i) + \frac{\mu}{\pi_i^t(a_i)} (c_i(a_i) - \pi_i^t(a_i))$. Then,

$$
\begin{aligned}
\frac{d}{dt} D_\psi(\pi, \pi^t) &= \sum_{i=1}^2 \left\langle \frac{d}{dt} z_i^t, \pi_i^t - \pi_i \right\rangle \\
&= \sum_{i=1}^2 \sum_{a_i \in A_i} \left(q_i^{\pi^t}(a_i) + \frac{\mu}{\pi_i^t(a_i)} (c_i(a_i) - \pi_i^t(a_i)) \right) (\pi_i^t(a_i) - \pi_i(a_i)) \\
&= \sum_{i=1}^2 \sum_{a_i \in A_i} (\pi_i^t(a_i) - \pi_i(a_i)) \sum_{a_{-i} \in A_{-i}} \pi_{-i}^t(a_{-i}) \left(u_i(a_i, a_{-i}) + \frac{\mu}{\pi_i^t(a_i)} (c_i(a_i) - \pi_i^t(a_i)) \right) \\
&= \sum_{i=1}^2 \sum_{a_i \in A_i} (\pi_i^t(a_i) - \pi_i(a_i)) \sum_{a_{-i} \in A_{-i}} \pi_{-i}^t(a_{-i}) \left(u_i(a_i, a_{-i}) + \mu \left(\frac{c_i(a_i)}{\pi_i^t(a_i)} - 1 \right) \right) \\
&= \sum_{i=1}^2 \sum_{a_i \in A_i} (\pi_i^t(a_i) - \pi_i(a_i)) \sum_{a_{-i} \in A_{-i}} \pi_{-i}^t(a_{-i}) \left(u_i(a_i, a_{-i}) + \mu \frac{c_i(a_i)}{\pi_i^t(a_i)} \right),
\end{aligned}
$$

where the last equality follows from $\sum_{a_i \in A_i} \pi_i(a_i) \sum_{a_{-i} \in A_{-i}} \mu \pi_{-i}^t(a_{-i}) = \mu$ for all $\pi_i \in \Delta(A_i)$. Furthermore,

$$
\begin{aligned}
\frac{d}{dt} D_\psi(\pi, \pi^t) &= \sum_{i=1}^2 \sum_{a_i \in A_i} (\pi_i^t(a_i) - \pi_i(a_i)) \sum_{a_{-i} \in A_{-i}} \pi_{-i}^t(a_{-i}) \left(u_i(a_i, a_{-i}) + \mu \frac{c_i(a_i)}{\pi_i^t(a_i)} \right) \\
&= \sum_{i=1}^2 \sum_{a \in A} \pi_i^t(a_i) \pi_{-i}^t(a_{-i}) u_i(a) - \sum_{i=1}^2 \sum_{a \in A} \pi_i(a_i) \pi_{-i}^t(a_{-i}) u_i(a) \\
&\quad + \sum_{i=1}^2 \sum_{a \in A} \pi_i^t(a_i) \pi_{-i}^t(a_{-i}) \mu \frac{c_i(a_i)}{\pi_i^t(a_i)} - \sum_{i=1}^2 \sum_{a \in A} \pi_i(a_i) \pi_{-i}^t(a_{-i}) \mu \frac{c_i(a_i)}{\pi_i^t(a_i)} \\
&= -\sum_{i=1}^2 \sum_{a \in A} \pi_i(a_i) \pi_{-i}^t(a_{-i}) u_i(a) + 2\mu - \mu \sum_{i=1}^2 \sum_{a \in A} \pi_i(a_i) \pi_{-i}^t(a_{-i}) \frac{c_i(a_i)}{\pi_i^t(a_i)} \\
&= -\sum_{i=1}^2 \sum_{a \in A} \pi_i(a_i) \pi_{-i}^t(a_{-i}) u_i(a) + 2\mu - \mu \sum_{i=1}^2 \sum_{a_i \in A_i} c_i(a_i) \frac{\pi_i(a_i)}{\pi_i^t(a_i)} \\
&= \sum_{i=1}^2 \sum_{a \in A} \pi_i^t(a_i) \pi_{-i}(a_{-i}) u_i(a) + 2\mu - \mu \sum_{i=1}^2 \sum_{a_i \in A_i} c_i(a_i) \frac{\pi_i(a_i)}{\pi_i^t(a_i)},
\end{aligned}
$$

where the third equality follows from $\sum_{i=1}^{2}\sum_{a\in A}\pi_i^t(a_i)\pi_{-i}^t(a_{-i})u_i(a_i,a_{-i}) = \sum_{i=1}v_i^{\pi^t} = 0$ and $\sum_{a\in A}\pi_i^t(a_i)\pi_{-i}^t(a_{-i})\mu\frac{c_i(a_i)}{\pi_i^t(a_i)} = \mu\sum_{a_i\in A_i}c_i(a_i)\frac{\pi_i(a_i)}{\pi_i^t(a_i)}\sum_{a_{-i}\in A_{-i}}\pi_{-i}^t(a_{-i}) = \mu\sum_{a_i\in A_i}c_i(a_i)\frac{\pi_i(a_i)}{\pi_i^t(a_i)}$. $\qquad\square$

C.3 PROOF OF LEMMA 5.6

Proof of Lemma 5.6. By using the ordinary differential equation (RMD), we have for all $i \in \{1,2\}$ and $a_i \in A_i$:

$$\pi_i^\mu(a_i)\left(q_i^{\pi^\mu}(a_i) - \langle\pi_i^\mu, q_i^{\pi^\mu}\rangle\right) + \mu\left(c_i(a_i) - \pi_i^\mu(a_i)\right)$$

$$= \pi_i^\mu(a_i)\left(\sum_{a_{-i}\in A_{-i}}\pi_{-i}^\mu(a_{-i})u_i(a_i,a_{-i}) - \sum_{a_i'\in A_i}\sum_{a_{-i}\in A_{-i}}\pi_i^\mu(a_i')\pi_{-i}^\mu(a_{-i})u_i(a_i',a_{-i})\right) + \mu\left(c_i(a_i) - \pi_i^\mu(a_i)\right) = 0.$$

Then, from Lemma B.2, we get:

$$\sum_{a_{-i}\in A_{-i}}\pi_{-i}^\mu(a_{-i})u_i(a_i,a_{-i}) = \sum_{a_i'\in A_i}\sum_{a_{-i}\in A_{-i}}\pi_i^\mu(a_i')\pi_{-i}^\mu(a_{-i})u_i(a_i',a_{-i}) - \frac{\mu}{\pi_i^\mu(a_i)}\left(c_i(a_i) - \pi_i^\mu(a_i)\right).$$

Note that from the following Lemma B.2, $\frac{1}{\pi_i^\mu(a_i)}$ is well-defined. Then, for any $\pi_i' \in \Delta(A_i)$ we have:

$$\sum_{a\in A}\pi_i'(a_i)\pi_{-i}^\mu(a_{-i})u_i(a_i,a_{-i}) = \sum_{a\in A}\pi_i^\mu(a_i)\pi_{-i}^\mu(a_{-i})u_i(a_i,a_{-i}) - \mu\sum_{a_i\in A_i}\frac{\pi_i'(a_i)}{\pi_i^\mu(a_i)}\left(c_i(a_i) - \pi_i^\mu(a_i)\right)$$

$$= \sum_{a\in A}\pi_i^\mu(a_i)\pi_{-i}^\mu(a_{-i})u_i(a_i,a_{-i}) + \mu - \mu\sum_{a_i\in A_i}c_i(a_i)\frac{\pi_i'(a_i)}{\pi_i^\mu(a_i)}.$$

$\qquad\square$

C.4 PROOF OF THEOREM 5.2

Proof of Theorem 5.2. First, we prove the first part of the theorem. By setting $\pi = \pi^\mu$ in Lemma 5.5 and $\pi' = \pi^t$ in Lemma 5.6, we have:

$$\frac{d}{dt}D_\psi(\pi^\mu,\pi^t) = \sum_{i=1}^{2}\sum_{a\in A}\pi_i^t(a_i)\pi_{-i}^\mu(a_{-i})u_i(a_i,a_{-i}) + 2\mu - \mu\sum_{i=1}^{2}\sum_{a_i\in A_i}c_i(a_i)\frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)}$$

$$= \sum_{i=1}^{2}\sum_{a\in A}\pi_i^\mu(a_i)\pi_{-i}^\mu(a_{-i})u_i(a_i,a_{-i}) + 4\mu - \mu\sum_{i=1}^{2}\sum_{a_i\in A_i}c_i(a_i)\frac{\pi_i^t(a_i)}{\pi_i^\mu(a_i)} - \mu\sum_{i=1}^{2}\sum_{a_i\in A_i}c_i(a_i)\frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)}$$

$$= 4\mu - \mu\sum_{i=1}^{2}\sum_{a_i\in A_i}c_i(a_i)\left(\frac{\pi_i^t(a_i)}{\pi_i^\mu(a_i)} + \frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)}\right)$$

$$= 4\mu - \mu\sum_{i=1}^{2}\sum_{a_i\in A_i}c_i(a_i)\left(\left(\sqrt{\frac{\pi_i^t(a_i)}{\pi_i^\mu(a_i)}} - \sqrt{\frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)}}\right)^2 + 2\right)$$

$$= -\mu\sum_{i=1}^{2}\sum_{a_i\in A_i}c_i(a_i)\left(\sqrt{\frac{\pi_i^t(a_i)}{\pi_i^\mu(a_i)}} - \sqrt{\frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)}}\right)^2,$$

where the third equality follows from $\sum_{i=1}^{2}\sum_{a\in A}\pi_i^\mu(a_i)\pi_{-i}^\mu(a_{-i})u_i(a_i,a_{-i}) = \sum_{i=1}^{2}v_i^{\pi^\mu} = 0$.

Next, we prove the second part of the theorem. From the first part of the theorem, we have:

$$\frac{d}{dt}D_\psi(\pi^\mu, \pi^t) = -\mu \sum_{i=1}^{2} \sum_{a_i \in A_i} c_i(a_i) \left(\frac{\pi_i^t(a_i)}{\pi_i^\mu(a_i)} + \frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)} - 2 \right)$$

$$\leq -\mu \sum_{i=1}^{2} \left(\min_{a_i \in A_i} \frac{c_i(a_i)}{\pi_i^\mu(a_i)} \right) \sum_{a_i \in A_i} \pi_i^\mu(a_i) \left(\frac{\pi_i^t(a_i)}{\pi_i^\mu(a_i)} + \frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)} - 2 \right)$$

$$= -\mu \sum_{i=1}^{2} \left(\min_{a_i \in A_i} \frac{c_i(a_i)}{\pi_i^\mu(a_i)} \right) \sum_{a_i \in A_i} \frac{(\pi_i^t(a_i) - \pi_i^\mu(a_i))^2}{\pi_i^t(a_i)}$$

$$\leq -\mu \sum_{i=1}^{2} \left(\min_{a_i \in A_i} \frac{c_i(a_i)}{\pi_i^\mu(a_i)} \right) \ln \left(1 + \sum_{a_i \in A_i} \frac{(\pi_i^t(a_i) - \pi_i^\mu(a_i))^2}{\pi_i^t(a_i)} \right)$$

$$= -\mu \sum_{i=1}^{2} \left(\min_{a_i \in A_i} \frac{c_i(a_i)}{\pi_i^\mu(a_i)} \right) \ln \left(\sum_{a_i \in A_i} \pi_i^\mu(a_i) \frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)} \right)$$

$$\leq -\mu \sum_{i=1}^{2} \left(\min_{a_i \in A_i} \frac{c_i(a_i)}{\pi_i^\mu(a_i)} \right) \sum_{a_i \in A_i} \pi_i^\mu(a_i) \ln \left(\frac{\pi_i^\mu(a_i)}{\pi_i^t(a_i)} \right)$$

$$= -\mu \sum_{i=1}^{2} \left(\min_{a_i \in A_i} \frac{c_i(a_i)}{\pi_i^\mu(a_i)} \right) \mathrm{KL}(\pi_i^\mu, \pi_i^t) \leq -\mu \left(\min_{i \in \{1,2\}, a_i \in A_i} \frac{c_i(a_i)}{\pi_i^\mu(a_i)} \right) \sum_{i=1}^{2} \mathrm{KL}(\pi_i^\mu, \pi_i^t), \quad (7)$$

where the second inequality follows from $x \geq \ln(1+x)$ for all $x > 0$, and the third inequality follows from the concavity of the $\ln(\cdot)$ function and Jensen's inequality for concave functions. On the other hand, when $\psi_i(p) = \sum_{a_i \in A_i} p(a_i) \ln p(a_i)$, $D_{\psi_i}(\pi_i^\mu, \pi_i^t) = \mathrm{KL}(\pi_i^\mu, \pi_i^t)$. Thus, we have $D_\psi(\pi^\mu, \pi^t) = \sum_{i=1}^{2} \mathrm{KL}(\pi_i^\mu, \pi_i^t)$. From this fact and (7), we have:

$$\frac{d}{dt}\mathrm{KL}(\pi^\mu, \pi^t) \leq -\mu \left(\min_{i \in \{1,2\}, a_i \in A_i} \frac{c_i(a_i)}{\pi_i^\mu(a_i)} \right) \mathrm{KL}(\pi^\mu, \pi^t).$$

$\square$

D PROOFS OF ADDITIONAL LEMMAS

D.1 PROOF OF LEMMA B.1

Proof of Lemma B.1. First, for any $\pi \in \Delta(A_1) \times \Delta(A_2)$,

$$D_\psi(\pi, \pi^t) = \sum_{i=1}^{2} D_{\psi_i}(\pi_i, \pi_i^t) = \sum_{i=1}^{2} \left(\psi_i(\pi_i) - \psi_i(\pi_i^t) - \langle \nabla \psi_i(\pi_i^t), \pi_i - \pi_i^t \rangle \right). \quad (8)$$

From the assumptions on ψ_i and the first-order necessary conditions for the optimization problem of $\arg\max_{p \in \Delta(A_i)} \{\langle z_i^t, p \rangle - \psi_i(p)\}$, for $\pi_i^t = \arg\max_{p \in \Delta(A_i)} \{\langle z_i^t, p \rangle - \psi_i(p)\}$, there exists $\lambda \in \mathbb{R}$ such that

$$z_i^t - \nabla \psi_i(\pi_i^t) = \lambda \mathbf{1}.$$

Therefore, we have:

$$\langle z_i^t, \pi_i - \pi_i^t \rangle = \langle \lambda \mathbf{1} + \nabla \psi_i(\pi_i^t), \pi_i - \pi_i^t \rangle = \langle \nabla \psi_i(\pi_i^t), \pi_i - \pi_i^t \rangle. \quad (9)$$

By combining (8) and (9):

$$
\begin{aligned}
D_\psi(\pi, \pi^t) &= \sum_{i=1}^{2} \left(\psi_i(\pi_i) - \psi_i(\pi_i^t) - \left\langle z_i^t, \pi_i - \pi_i^t \right\rangle \right) \\
&= \sum_{i=1}^{2} \left(\left\langle z_i^t, \pi_i^t \right\rangle - \psi_i(\pi_i^t) - \left\langle z_i^t, \pi_i \right\rangle + \psi_i(\pi_i) \right) \\
&= \sum_{i=1}^{2} \left(\max_{p \in \Delta(A_i)} \left\{ \left\langle z_i^t, p \right\rangle - \psi_i(p) \right\} - \left\langle z_i^t, \pi_i \right\rangle + \psi_i(\pi_i) \right).
\end{aligned}
$$

$\square$

D.2   PROOF OF LEMMA B.2

Proof of Lemma B.2. We assume that there exists $i \in \{1, 2\}$ and $a_i \in A_i$ such that $\pi_i^\mu(a_i) = 0$. Then, for such i and a_i, we have:

$$
\frac{d}{dt}\pi_i^\mu(a_i) = \pi_i^\mu(a_i) \left(q_i^{\pi^\mu}(a_i) - \left\langle \pi_i^\mu, q_i^{\pi^\mu} \right\rangle \right) + \mu \left(c_i(a_i) - \pi_i^\mu(a_i) \right) = \mu c_i(a_i) > 0.
$$

This contradicts that $\frac{d}{dt}\pi_i^\mu(a_i) = 0$ since π_i^μ is a stationary point. Therefore, for all $i \in \{1, 2\}$ and $a_i \in A_i$, we have $\pi_i^\mu(a_i) > 0$. $\square$