Generalization of Spectral Graph Neural Networks

Abstract

Spectral graph neural networks (GNNs) have achieved remarkable success across various applications, yet their generalization properties remain poorly understood. This paper bridges this gap by analyzing the impact of graph homophily and architectural choices on the generalization of spectral GNNs. We derive a general form of uniform transductive stability for spectral GNNs and provide an explicit stability analysis for graphs with two node classes, providing a comprehensive framework to understand their generalization. Based on this stability analysis, we establish a generalization error bound, demonstrating that better stability leads to improved generalization. Our theoretical findings reveal that spectral GNNs generalize well on graphs with strong homophily or heterophily but struggle on graphs with weaker structural properties. We also identify conditions under which increasing the polynomial order in spectral GNN architectures may degrade generalization. Empirical results on synthetic and real-world benchmark datasets align closely with our theoretical findings.

1 Introduction

Generalization is a fundamental challenge in machine learning, crucial for understanding both the theoretical limits and practical performance of algorithms. Researchers have developed various measures to study generalization, including Vapnik–Chervonenkis (VC) dimension (Cherkassky et al., 1999), PAC-Bayes bound (McAllester, 1998), Rademacher complexity (Bartlett & Mendelson, 2002), and the stability of learning algorithms (Bousquet & Elisseeff, 2002). These measures provide insights into a model's ability to generalize beyond its training data. Graph neural networks (GNNs) have achieved remarkable success across various practical applications (Zhou et al., 2020), yet their generalization capabilities remain poorly understood. Unlike traditional machine learning models that operate on independent and identically distributed (i.i.d.) data, GNNs work on interdependent data where the graph topology and node/edge features are inherently linked. This interconnected structure makes it difficult to evaluate how well a GNN trained on one graph or a set of graphs can generalize to unseen graphs.

Research on GNN generalization primarily focuses on two tasks: graph classification and node classification. In graph classification, where graphs are typically i.i.d., generalization has been studied through connections with WL algorithms (Morris et al., 2023; D'Inverno et al., 2024; Franks et al., 2024) and data-dependent PAC-Bayes bounds (Liao et al., 2021; Ju et al., 2023). In node classification, which involves transductive learning where node features are known during training, approaches like Transductive Rademacher complexity and uniform transductive stability are more common. These methods explore the impact of factors such as graph matrix representations and GNN depth on generalization (Oono & Suzuki, 2020; El-Yaniv & Pechyony, 2007; Tang & Liu, 2023b; Zhou & Wang, 2021; Cong et al., 2021).

In this work, we focus on the generalization of GNNs for node classification. Unlike graph classification, node classification performance is influenced by the distribution of node classes, which is closely tied to graph homophily. In homophilic graphs, connected nodes tend to belong to the same class, whereas in heterophilic graphs, connected nodes are often from different classes. Empirical evidence shows that the edge homophilic ratio (Zhu et al., 2020) significantly affects GNN performance. For example, models like GCN and GAT excel on homophilic graphs but underperform on heterophilic graphs (Kipf & Welling, 2017; Velickovic et al., 2018). This motivates us to explore the relationship between graph homophily and the generalization of GNNs in node classification tasks,

raising the question: *how does GNN generalization depend on node class distributions?* Notably, this question has remained largely unexplored in previous research.

We examine the generalization of GNNs through a spectral perspective, as spectral GNNs have demonstrated strong performance on both homophilic and heterophilic graphs. Spectral GNNs operate in the spectral domain, applying frequency-domain convolutions to extract structural information (Balcilar et al., 2021). Formally, a spectral GNN is defined as:

$$\Psi(M, X) = \sigma(g_\Theta(M) f_W(X)), \tag{1}$$

where $M \in \mathbb{R}^{n \times n}$ is a graph matrix (e.g., Laplacian or adjacency matrix), $g_\Theta(M) = \sum_{k=0}^{K} \theta_k T_k(M)$ represents graph convolution using $\Theta = \{\theta_k\}_{k=0}^{K}$ and the k-th polynomial basis $T_k(\cdot)$, $f_W(X)$ is a feature transformation function parameterized by W, and σ is a non-linear function such as softmax.

One notable observation about spectral GNNs is that, although the frequency response function can theoretically be approximated by a sum of polynomial basis with negligible error as the order K increases (Powell, 1981), empirical results show that higher polynomial orders do not consistently lead to better performance. This discrepancy raises the question: *is the gap between theoretical study and empirical performance, particularly with respect to polynomial order K, related to the generalization of spectral GNNs?* To date, no work has provided a clear explanation for this phenomenon.

We address the aforementioned questions by focusing on the generalization of spectral GNNs. In transductive learning, the relationships between labeled and unlabeled nodes are critical for generalization. To measure this, we employ uniform transductive stability (El-Yaniv & Pechyony, 2006), which evaluates the stability of spectral GNNs under perturbations to individual training samples.

To study the interplay between graph structure and GNN performance, we use the contextual stochastic block model (cSBM) (Baranwal et al., 2023; Deshpande et al., 2018), a widely used generative model that captures both homophilic and heterophilic graph structures in a controlled and analytically tractable manner. Previous studies have demonstrated that cSBM models real-world datasets such as Citeseer, Cora, and Polblogs, which are frequently used in GNN research (Deshpande et al., 2018; LEI, 2016; Dreveton et al., 2023; Kipf & Welling, 2017; Zhang et al., 2021). Specifically, cSBM generates graphs with well-defined block structures, where nodes within the same block are more likely to be connected (homophilic graphs) or nodes between blocks have a higher connection probability (heterophilic graphs). Leveraging cSBM allows us to systematically vary graph homophily and examine its impact on GNN generalization properties.

Contributions. Our main contributions are summarized as follows:

- We analyze the γ-uniform transductive stability of spectral GNNs by decomposing it into two factors: the Lipschitz continuity and smoothness of the loss function and the spectral GNN, and the gradient norm bound (Theorem 6). This enables us to study the effects of node class distribution and spectral GNN architecture on training gradients through *an explicit gradient norm bound* (Theorem 8).

- We establish the generalization error bound of spectral GNNs based on their stability, where good stability indicates strong generalization capability (Theorem 9). To further explicitly analyze the effects of graph homophily and polynomial order on generalization, we derive an explicit form of the gradient norm bound for two node classes (Theorem 13).

- We prove that spectral GNNs generalize well on graphs that are strong homophilic or heterophilic, but perform poorly on graphs that are moderately homophilic or heterophilic. We identify conditions under which increasing the polynomial order in spectral GNN architectures may degrade generalization, providing insights into architectural design (Theorems 14 and 15; Proposition 16).

To validate our theoretical findings, we conduct experiments on nine synthetic datasets and nine real-world benchmark datasets for node classification. The experimental results align closely with our theoretical analysis.

2 RELATED WORKS

Previous work. We review prior studies on GNN generalization, typically categorized into graph classification and node classification, with a primary focus on the latter.

Generalization Bound Methods		Analysis		Key factors in Bounds				
		Lipschitz	Gradient	Graph Matrix	Tsize	Depth	Hom	Poly
VC bound	(Scarselli et al., 2018)	–	n/a	n	$\sqrt{}$	–	–	–
RC bound (*transductive*)	(Esser et al., 2021)	$\sqrt{}$	n/a	$\|M\|_\infty, \|MX\|_{2\to\infty}$	$\sqrt{}$	$\sqrt{}$	–	–
	(Tang & Liu, 2023b)	$\sqrt{}$	n/a	$\|M\|_\infty$	$\sqrt{}$	$\sqrt{}$	–	–
	(Oono & Suzuki, 2020)	$\sqrt{}$	n/a	$\|MX\|_F$	$\sqrt{}$	$\sqrt{}$	–	–
US bound (*Inductive*)	(Verma & Zhang, 2019)	–	$\sqrt{}$	$\|M\|_2$	$\sqrt{}$	–	–	–
	(Zhou & Wang, 2021)	–	$\sqrt{}$	$\|M\|_2, \|MX\|_2$	$\sqrt{}$	$\sqrt{}$	–	–
US bound (*transductive*)	(Cong et al., 2021)	$\sqrt{}$	$\sqrt{}$	d_{max}	$\sqrt{}$	$\sqrt{}$	–	–
	Our work	$\sqrt{}$	$\sqrt{}$	M_{ij}	$\sqrt{}$	–	$\sqrt{}$	$\sqrt{}$

Table 1: Comparison of generalization bounds for GNNs on node classification. Here, $\|\cdot\|_2, \|\cdot\|_F$, $\|\cdot\|_\infty, \|\cdot\|_{2\to\infty}$ denote the spectral norm, Frobenius norm, infinity norm and maximum column ℓ_2-norm. n is graph node number. $d_{\max}$ denotes the maximum node degree. The other factors Tsize, Depth, Hom, and Poly refer to training sample size, model depth, graph homophily, and polynomial order, respectively. $\sqrt{}$ indicates "discussed", while – indicates "not discussed".

Graph classification is typically considered as an inductive learning task. One prominent method is the Vapnik–Chervonenkis (VC) bound, which relates the VC dimension of a GNN to the number of colors generated by the 1-WL algorithm, reflecting the number of graphs the 1-WL algorithm can distinguish (Morris et al., 2023). The PAC-Bayes bound is another approach. Liao et al. (2021) connects generalization bound to factors like maximum node degree and GNN depth, while Ju et al. (2023) refines this by tying the bound to the largest singular value of the graph matrix. Behboodi et al. (2022) further extends the PAC-Bayes framework to equivariant networks, highlighting the influence of group properties on generalization. Rademacher complexity and uniform convergence have also been explored. Garg et al. (2020) shows that a GNN's computational tree complexity bounds its overall complexity, and Maskey et al. (2022) shows that generalization bound increases with model complexity but decreases with higher average node degrees.

Generalization analysis for node classification is more challenging than for graph classification due to its transductive nature (Tang & Liu, 2023b). Recent studies focus on how graph matrices and GNN depth influence generalization. Rademacher complexity (RC) is commonly used for node classification tasks due to its strong theoretical foundation in transductive learning (El-Yaniv & Pechyony, 2007). It has been shown that the transductive RC of a GNN is proportional to the infinity norm of its graph matrix, with generalization bounds provided for several classic GNNs (Tang & Liu, 2023b). Esser et al. (2021) uses a planted model to illustrate the relationship between GNN generalization and graph matrix compatibility. Oono & Suzuki (2020) shows that gradient boosting reduces RC in multi-scale GNNs. Uniform stability (US) offers another key approach for analyzing generalization. Verma & Zhang (2019) relates the generalization error bound of single-layer GCNs to the largest absolute eigenvalue of the graph matrix, while Cong et al. (2021) demonstrates that increasing GNN depth improves stability and lowers generalization error bounds. Other works deviate from transductive assumptions, addressing non-i.i.d. settings. For example, Ma et al. (2021) derives PAC-Bayesian bounds by assuming i.i.d. node classes given fixed node features.

Our work. In this work, we focus on node classification tasks, investigating how node class distribution and the architecture of spectral GNNs influence generalization. While prior studies have examined factors like graph size, training set size, graph matrix norms, and node features, they have largely overlooked the role of graph homophily in generalization and the impact of increasing the polynomial order of spectral GNNs. To our knowledge, this is the first study to analyze the effects of graph homophily and polynomial order on the generalization of spectral GNNs.

Table 1 compares our work with other methods for node classification, highlighting several key aspects: (1) **Analysis settings** (inductive or transductive): The VC bound (Scarselli et al., 2018) is data-independent and thus agnostic to inductive or transductive settings. While Verma & Zhang (2019); Zhou & Wang (2021) derive bounds for GNNs in inductive settings, others (Esser et al., 2021; Tang & Liu, 2023b; Oono & Suzuki, 2020; Cong et al., 2021) and our work address the more complex transductive setting. (2) **Analysis frameworks**: Rademacher complexity estimates a model's capacity to fit noise based on graph structure and node features but does not account for node labels. As a result, methods such as (Esser et al., 2021; Tang & Liu, 2023b; Oono & Suzuki, 2020) are unable to analyze the effect of graph homophily, which depends on both graph structure and node

labels. In contrast, uniform stability enables analysis of the relationship between generalization and graph homophily using gradient-based analysis. It is worth noting that while (Cong et al., 2021) employs uniform transductive stability, their analysis focuses solely on how GNN depth impacts Lipschitz continuity and gradient. (3) **Key factors in bounds**: Training sample size is a critical factor in all bounds except the VC bound (Scarselli et al., 2018). Model depth (number of GNN layers) is addressed in (Esser et al., 2021; Tang & Liu, 2023b; Zhou & Wang, 2021; Cong et al., 2021). Our work examines spectral GNNs, where the architecture comprises only one layer of K-order polynomials. Unlike prior studies that focus on various graph matrix norms, our analysis takes a finer-grained approach by considering the expectation of individual graph matrix elements. Moreover, we explore the effects of graph homophily and the polynomial order on the generalization of spectral GNNs, aspects not previously investigated.

3 PROBLEM SETUP

An undirected, unweighted graph is denoted as $G = (V, E)$, where V is the set of nodes, E is the set of edges, and $|V| = n$. In this work, we model graphs with controlled properties using the generalized multi-class contextual stochastic block model (cSBM) (Baranwal et al., 2023). A graph $G \sim cSBM(n, f, \Pi, Q)$ consists of n nodes, partitioned into C distinct classes. Here, f is the dimension of node features, $\Pi = \{\Pi_i\}_{i \in [C]}$ is a set of C continuous distributions, where $\pi_i \in \mathbb{R}^f$ and $\Sigma_i \in \mathbb{R}^{f \times f}$ are the mean and variance of Π_i respectively, and $Q \in \mathbb{R}^{C \times C}$ is a symmetric edge-probability matrix. Each node v_i is assigned a class y_i sampled uniformly from a set of C classes, and its feature vector $x_i \in \mathbb{R}^f$ is drawn from the distribution Π_{y_i}. This generates the node feature matrix $X \in \mathbb{R}^{n \times f}$. For the adjacency matrix $A \in \mathbb{R}^{n \times n}$, an edge between node v_i and v_j (i.e., $A_{ij} = 1$) is generated according to the edge-probability matrix Q based on the following probability:

$$P(A_{ij} = 1 \mid y_i, y_j) = Q_{y_i y_j},$$

where $Q_{y_i y_j}$ gives the edge formation probability between class y_i and class y_j.

For a spectral GNN Ψ, following (Wang & Zhang, 2022; Lu et al., 2024), we first consider $f_W(X) = XW$, and then discuss the effect of non-linear feature transformation. Here $W \in \mathbb{R}^{f \times C}$. Let $S_m = (X, \{y_i\}_{i=1}^m)$ be a training set containing m labelled nodes, randomly sampled form the graph G, and $\mathcal{D}_u = (X, \{y_i\}_{i=m+1}^n)$ be the testing set containing the other nodes in the graph G. We define a loss function $\ell(y_i, \hat{y}_i|_{\Theta, W})$ to measure the discrepancy between the truth class y_i and the prediction $\hat{y}_i$ when a spectral GNN is parameterized by Θ, W. The empirical loss $\mathcal{L}_{S_m}(\Theta, W)$ and the expected loss $\mathcal{L}_{\mathcal{D}_u}(\Theta, W)$ are defined as:

$$\mathcal{L}_{S_m}(\Theta, W) = \frac{1}{m} \sum_{i=1}^m \ell(y_i, \hat{y}_i|_{\Theta, W}), \quad \mathcal{L}_{\mathcal{D}_u}(\Theta, W) = \frac{1}{n-m} \sum_{i=m+1}^n \ell(y_i, \hat{y}_i|_{\Theta, W}).$$

Following (El-Yaniv & Pechyony, 2006), testing datasets are randomly sampled from real data and we treat the loss on these testing datasets as the expected loss. Given that the optimal parameters Θ^*, W^* minimize the empirical loss $\mathcal{L}_{S_m}(\Theta^*, W^*)$, our goal is to bound the generalization error:

$$\mathcal{L}_{\mathcal{D}_U}(\Theta^*, W^*) - \mathcal{L}_{S_m}(\Theta^*, W^*).$$

A small generalization error bound indicates that spectral GNNs can perform well on testing data.

3.1 ASSUMPTIONS

We first introduce assumptions used in the generalization analysis of spectral GNNs.

Assumption 1 (Lipschitz Continuity and Smoothness). The loss function ℓ and the spectral GNN Ψ are both Lipschitz continuous and smooth.

Assumption 1 is commonly used in the analysis of neural networks trained with gradient descent (Ghadimi & Lan, 2013). They are necessary conditions ensuring that the neural network training converges (Arfken et al., 2011; Liao et al., 2021). We use $\mathrm{Lip}(\bullet)$ and $\mathrm{Smt}(\bullet)$ to denote the *Lipschitz* constant and smoothness of a function, respectively, and $\| \bullet \|_F$ denotes the Frobenius norm.

Assumption 2 (Bounded Gradients). The gradients of both the spectral GNN and the loss function $\ell(y_i, \hat{y}_i)$ w.r.t. any parameters Θ, W, and for any node v_i with class y_i and prediction $\hat{y}_i$, are bounded:

$$\|\nabla_\Theta \hat{y}_i\|_F^2 + \|\nabla_W \hat{y}_i\|_F^2 \leq \beta_2^2; \quad \|\nabla_{\hat{y}_i} \ell(y_i, \hat{y}_i)\|_F \leq \beta_1. \tag{2}$$

Here, $\nabla_\Theta \hat{y}_i$ and $\nabla_W \hat{y}_i$ represent the gradients of $\hat{y}_i$ with respect to the parameters Θ and W, respectively. $\nabla_{\hat{y}_i} \ell(y_i, \hat{y}_i)$ refers to the gradient of the loss function with respect to $\hat{y}_i$.

Remark. The loss surface of neural networks often contains many local minima (Dinh et al., 2017). The bounded gradient assumption ensures convergence during optimization (Li & Liu, 2021).

Unlike margin loss, the cross-entropy loss function is unbounded. For theoretical analysis, it is typically assumed that the sample loss is bounded.

Assumption 3 (Bounded Sample Loss). For any sample (x_i, y_i), the maximum loss between the ground truth class y_i and the prediction $\hat{y}_i$ satisfies $\ell(\hat{y}_i, y_i) \leq B_\ell$.

Following the previous work (Zhang et al., 2019; Kuzborskij & Lampert, 2018), we assume that parameter norms are bounded during training.

Assumption 4 (Bounded Parameters). Each parameter $\theta_k \in \Theta$ and the parameter matrix W in a spectral GNN are bounded, i.e., $|\theta_k| \leq B_\Theta$ and $\|W\|_F \leq B_W$.

4 GENERAL RESULTS

In this section, we examine the generalization of spectral GNNs through *uniform transductive stability* (El-Yaniv & Pechyony, 2006). We define γ-uniform transductive stability for spectral GNNs, analyze the key factors influencing γ by deriving the gradient norm bound, and use these insights to establish the generalization error bound.

Definition 5 (γ-Uniform Transductive Stability). Let $S_m^{ij} = (X, \{y_t\}_{t=1, t \neq i}^m \cup \{y_j\})$ be a perturbed dataset obtained by replacing the i-th sample in S_m with the j-th sample from $\mathcal{D}_u$. Let $\{\Theta^*, W^*\}$ and $\{\Theta', W'\}$ be the optimal parameters of a spectral GNN Ψ trained on S_m and S_m^{ij}, respectively. Denote the predictions for node v_i by $\hat{y}_i|_{\Theta^*, W^*}$ and $\hat{y}_i|_{\Theta', W'}$. The spectral GNN Ψ is γ-*uniform transductive stable* if for any $i \in [n]$:

$$\max_{1 \leq i \leq n} |\ell(y_i, \hat{y}_i|_{\Theta^*, W^*}) - \ell(y_i, \hat{y}_i|_{\Theta', W'})| \leq \gamma.$$

A larger γ indicates worse stability of spectral GNNs. Below, we decompose γ into two terms: r and β. Here, r accounts for the Lipschitz continuity and smoothness of the loss function and spectral GNNs, while β bounds the gradient norm during training.

Theorem 6 (Stability and Gradient Norm). *Let Ψ be a spectral GNN trained using gradient descent for T iterations with a learning rate η on a training dataset S_m, and evaluated on a testing set $\mathcal{D}_u$. Under Assumption 1, for all iterations $t \in [1, T]$ and any sample (x_i, y_i) in S_m or $\mathcal{D}_u$, if the gradient norm satisfies $\|\nabla \ell(y_i, \hat{y}_i|_{\Theta^t, W^t})\|_F \leq \beta$, where $\{\Theta^t, W^t\}$ are the parameters at the t-th iteration, then Ψ satisfies γ-uniform transductive stability with:*

$$\gamma = r\beta, \quad r = \frac{2\eta \alpha_1}{m} \sum_{t=1}^T (1 + \eta \alpha_2)^{t-1},$$

where $\alpha_1 = Lip(\ell) \cdot Lip(\Psi)$ and $\alpha_2 = Smt(\Psi)\beta_1 + Smt(\ell)Lip(\Psi)\beta_2$.

Remark. α_1 and α_2 represent the Lipschitz constant and smoothness of the loss function ℓ and the spectral GNN Ψ, respectively. They determine how parameter updates Θ^t and W^t during training affect the loss of a sample (x_i, y_i) through the term r. As described in Eq. (2), β_1 and β_2 are the bounds for the loss and its gradient, respectively. The proof is provided in Appendix B.

Unlike previous work (Cong et al., 2021), which assumes a fixed gradient norm β, we explicitly derive β to analyze how graph homophily and the polynomial order of spectral GNNs influence the gradient norm and, in turn, stability. To begin, we introduce the concept of a walk on a graph and its expectation, both critical for analyzing the stability of spectral GNNs. A k-*length walk* on a graph G is defined as a sequence of k edges $\{e_1, e_2, \ldots, e_k\}$, where $e_i \in E$, and the endpoint of e_i is the starting point of e_{i+1} for $i \in \{1, \ldots, k-1\}$. The expectation of k-length walks is defined as follows.

Definition 7 ($\mathbb{E}[A_{ij}^k]$). For a graph $G \sim cSBM(n, f, \Pi, Q)$ with adjacency matrix A, the expectation of the element A_{ij}^k in the k-th power of A is:

$$\mathbb{E}[A_{ij}^k] = \sum_{p \in P_{ij}^k} \prod_{(v,v') \in p} Q_{yy'},$$

where P_{ij}^k is the set of all k-length walks between v_i and v_j, and $Q_{yy'}$ is the transition probability between the classes y of v and y' of v'.

Remark. Intuitively, A_{ij}^k represents the number of k-length walks between nodes v_i and v_j. The first moment $\mathbb{E}[A_{ij}^k]$ gives the expected number of such walks in the random graph generated by the cSBM. Since $Q_{yy'}$ represents the probability of an edge between nodes of classes y and y', a larger $Q_{yy'}$ increases the likelihood of edges in walks involving transitions between these classes, resulting in higher expected counts of such walks.

The following theorem reveals how the gradient norm bound β depends on the expectation of k-length walks and their node class distributions within the graph.

Theorem 8. *Consider a spectral GNN Ψ with polynomial order K trained using full-batch gradient descent for T iterations with a learning rate η on a training dataset S_m sampled from a graph $G \sim cSBM(n, f, \Pi, Q)$ with average node degree $d \ll n$. When $n \to \infty$ and $K \ll n$, under Assumptions 1, 2, and 4, for any node v_i, $i \in [n]$, and for a constant $\epsilon \in (0, 1)$, with probability at least $1 - \epsilon$, Ψ satisfies γ-uniform transductive stability, where $\gamma = r\beta$ and*

$$\beta = \frac{1}{\epsilon} \Bigg[O\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2 \right] \right) + O\left(\|\pi_{y_i}^\top \pi_{y_i} + \Sigma_{y_i}\|_F \right)$$

$$+ O\left(\sum_{k=1}^K \sum_{j=1}^n \mathbb{E}[A_{ij}^k] \left\| \sum_{t=1}^n \mathbb{E}[A_{it}^k] \pi_{y_j}^\top \pi_{y_t} + \mathbb{E}[A_{ij}^k] \Sigma_{y_j} \right\|_F \right) \Bigg].$$

Proof sketch. The main idea is to first derive the gradient of a sample loss and the expected gradient norm $\mathbb{E}\left[\|\nabla \ell(y_i, \hat{y}_i|_{\Theta^t, W^t})\|_F \right]$ for node (x_i, y_i) with fixed parameters (n, f, Π, Q). Then the gradient norm bound β is derived using Markov's Inequality (Evans & Rosenthal, 2004). When parameters Θ, W are bounded (Assumption 4), the main factors in $\mathbb{E}\left[\|\nabla \ell(y_i, \hat{y}_i|_{\Theta^t, W^t})\|_F \right]$ are $\|\hat{y}_i - y_i\|_F^2$ and moments of $A^k X$. When graph structure and node features are independent given node labels, we have $\mathbb{E}\left[A^k X \right] = \mathbb{E}\left[A^k \right] \mathbb{E}\left[X \right]$. While A^k depends on the graph structure, governed by the parameter Q, node feature X is controlled by parameter Π, shown as the mean π_{y_i} and variance Σ_{y_i} of nodes belong to class y_i. The full proof is provided in Appendix C. $\square$

Remark. Theorem 8 shows that the gradient norm bound β is primarily influenced by two factors:

(1) **Expected prediction error** $\mathbb{E}[\|\hat{y}_i - y_i\|_F^2]$ quantifies the difference between the truth node class y_i and the predicted label $\hat{y}_i$ for a node v_i by a spectral GNN. A well-performing spectral GNN is characterized by a low expected prediction error.

(2) **Expectation of k-length walks** $\mathbb{E}[A_{ij}^k]$ measures the interaction between nodes v_i and v_j through k-length walks. The existence of these walks depends on the labels of all nodes along the walk, with edge probabilities in k-length walks determined by Q in cSBM.

When a γ-transductive learning algorithm is trained on two nearly identical training sets, differing by just one sample, the expected generalization error equals the expected increase in sample loss (El-Yaniv & Pechyony, 2006). Based on this known result, Theorem 6, and β obtained from Theorem 8, we have the following generalization error bound for spectral GNNs.

Theorem 9 (Generalization Error Bound). *Let $H_2(n) \triangleq \sum_{i=1}^n \frac{1}{i^2}$ and $\Omega(m, n - m) \triangleq (n - m)^2 (H_2(n) - H_2(n - m))$. For $\epsilon \in (0, 1)$, if a spectral GNN is γ-uniform transductive stability with probability $1 - \epsilon$, then under Assumption 3, for $\delta \in (0, 1)$, with probability at least $(1 - \delta)(1 - \epsilon)$, the generalization error $\mathcal{L}_{\mathcal{D}_u}(\Theta, W) - \mathcal{L}_{S_m}(\Theta, W)$ is upper-bounded by:*

$$\gamma + \left(2\gamma + \left(\frac{1}{n - m} + \frac{1}{m} \right) (B_\ell - \gamma) \right) \sqrt{2\Omega(m, n - m) \log \frac{1}{\delta}}. \tag{3}$$

Remark. The generalization error bound of a spectral GNN is closely related to its uniform transductive stability γ, the number of training samples m, and the total number of nodes n in the graph. A smaller γ indicates stronger stability, which in turn suggests better generalization performance. Factors such as graph homophily and the polynomial order of a spectral GNN affect γ, thereby impacting the generalization error. From Theorem 6, we observe that $\gamma = O(1/m)$. When n is sufficiently large, the term $1/(n - m)$ becomes negligible, and $\Omega(m, n - m)$ increases as $O(m^{1/2})$.

The following lemma shows that increasing the number of training samples m improves the generalization. The proof is provided in Appendix D.

Lemma 10. *Consider a spectral GNN trained with m samples as $n \to \infty$. As the sample size m increases, the generalization error bound decreases at the rate $O(1/m) + O\left(\sqrt{2\log(1/\delta)/m}\right)$.*

Thus far, we have considered only linear feature transformation functions f_W in spectral GNNs. We now establish a result on how a non-linear f_W influences uniform transductive stability.

Proposition 11. *For a spectral GNN $\Psi_{\tilde{\sigma}}$ with a non-linear feature transformation function $f_W(X) = \tilde{\sigma}(XW)$, assume the gradient norm bound β in Theorem 9 is the same for Ψ and $\Psi_{\tilde{\sigma}}$. If $Lip(\tilde{\sigma}) \leq 1$ and $Smt(\tilde{\sigma}) \leq 1$, then $\gamma_{\tilde{\sigma}} \leq \gamma$, where $\gamma_{\tilde{\sigma}}$ is the stability of $\Psi_{\tilde{\sigma}}$.*

Remark. The $\gamma_{\tilde{\sigma}}$-uniform transductive stability of spectral GNNs $\Psi_{\tilde{\sigma}}$ with a non-linear transformation is bounded by the stability of its linear counterpart Ψ, provided that the activation function satisfies $Lip(\tilde{\sigma}) \leq 1$ and $Smt(\tilde{\sigma}) \leq 1$. These conditions ensure that the non-linear transformation does not excessively amplify inputs or gradients, thus maintaining stability. Common activation functions like ReLU, Sigmoid, and Tanh satisfy these criteria, indicating that they may contribute to stabilizing the training of spectral GNNs by preventing large output fluctuations in response to small input changes. The proof is provided in Appendix D.

5 FURTHER ANALYSIS

In this section, we analyze the impact of node class distribution and spectral GNN architecture on the generalization error bound. To derive an explicit form for property analysis, we consider $cSBM(n, f, \mu, u, \lambda, d)$, a well-studied specialization of the general multi-class cSBM (Deshpande et al., 2018), widely used in prior studies on graph analysis (Esser et al., 2021; Ma et al., 2022; Baranwal et al., 2021; Baranwal et al.). Specifically, for a node v_i with label $y_i \in \{\pm 1\}$, its feature is sampled from a Gaussian distribution:

$$x_i \sim \mathcal{N}(y_i \sqrt{\mu/n}u, I_f/f).$$

Two nodes of the same class are connected with probability $c_{\text{in}} = d + \lambda\sqrt{d}$, while nodes of different classes are connected with probability $c_{\text{out}} = d - \lambda\sqrt{d}$. In this simplified 2-class cSBM, the distribution Π reduces to $\{\Pi_{\pm}\}$, and the edge-probability matrix simplifies to a 2×2 matrix with diagonal elements c_{in}/n and off-diagonal elements c_{out}/n.

By adjusting the parameter λ in cSBM, we can generate graphs with varying node class distributions. One way to quantify the node class distribution is the edge homophilic ratio (Zhu et al., 2020),

$$H_{edge} = \frac{|\{e_{ij} \mid v_i, v_j \in V, e_{ij} \in E, y_i = y_j\}|}{|E|}.$$

The relationship between the parameters d, λ in cSBM and the edge homophilic ratio is as follows.

Proposition 12. *For a graph $G \sim cSBM(n, \mu, u, \lambda, d)$, the expected edge homophily ratio is:*

$$\mathbb{E}[H_{edge}] = \frac{d + \lambda\sqrt{d}}{2d}; \quad \mathbb{E}[H_{edge}] = \frac{c_{in}}{c_{in} + c_{out}}. \tag{4}$$

When $\lambda > 0$, the graph tends to be homophilic as $\mathbb{E}[H_{edge}] > 0.5$. Conversely, when $\lambda < 0$, the graph tends to be heterophilic. The proof is provided in Appendix F.

5.1 UNIFORM TRANSDUCTIVE STABILITY

We now establish stability for graphs with two node classes and Gaussian-distributed node features.

Theorem 13. *Consider a spectral GNN Ψ parameterized by Θ, W trained using full-batch gradient descent for T iterations with a learning rate η on a training dataset containing m samples drawn from nodes on a graph $G \sim cSBM(n, f, \mu, u, \lambda, d)$. When $n \to \infty$, $k \ll n$, and $d \ll n$, under Assumptions 1, 2, and 4, for any node v_i on the graph, with probability at least $1 - \epsilon$ for a constant $\epsilon \in (0, 1)$, Ψ satisfies γ-uniform transductive stability, where $\gamma = r\beta$ and*

$$\beta = \frac{1}{\epsilon}\left[O\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]\right) + O\left(\sum_{k=2}^{K}\left(\mathbb{E}\left[\left(A_{ij}^k \mid y_i = y_j\right)^2\right] + \mathbb{E}\left[\left(A_{ij}^k \mid y_i \neq y_j\right)^2\right]\right)\right)\right].$$

Proof sketch. The proof follows the same structure as Theorem 8. The gradient norm bound β can be explicitly expressed as the expected prediction error $\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]$ and the homophily-aware walk variance $\zeta_k = \mathbb{E}\left[(A_{ij}^k \mid y_i = y_j)^2\right] + \mathbb{E}\left[(A_{ij}^k \mid y_i \neq y_j)^2\right]$. The connection between ζ_k and H_{edge} can be analyzed in a tractable manner. The full proof is provided in Appendix E.3. □

Remark. The theorem derives the explicit form of k-length walks A_{ij}^k. Notably, $(A_{ij}^k|y_i = y_j)$ and $(A_{ij}^k|y_i \neq y_j)$ follow distinct distributions based on whether nodes v_i and v_j share the same label. When $k = 1$, A_{ij}^k follows a Bernoulli distribution. For $n \to \infty$ with $d \ll n$ and $2 \leq k \ll n$, A_{ij}^k follows a Poisson distribution. The term $\zeta_k = \mathbb{E}[(A_{ij}^k|y_i = y_j)^2] + \mathbb{E}[(A_{ij}^k|y_i \neq y_j)^2]$ captures the *homophily-aware walk variance*, reflecting the variance in k-length walks between same-class or different-class nodes. This depends on the edge probabilities c_{in} and c_{out}: (1) $c_{in} = c_{out}$: the graph is essentially an Erdős-Rényi graph, lacking clusters or multipartite structure, leading to higher variance in k-length walks. (2) $c_{in} > c_{out}$: the graph is homophilic with cluster patterns, and walks are concentrated within clusters, reducing variance. (3) $c_{in} < c_{out}$: the graph is heterophilic with multipartite patterns, and walks are concentrated along edges connecting different classes, affecting the variance. In general, the absence of clear cluster or multipartite structures increases randomness in k-length walks, resulting in higher ζ_k.

5.2 MAIN FACTORS IN STABILITY

We first analyze how exactly the expected prediction error $\mathbb{E}[\|\hat{y}_i - y_i\|_F^2]$ and the homophily-aware walk variance ζ_k vary with the parameters λ and K, and then examine the combined effects of λ and K on the stability and generalization of spectral GNNs.

Theorem 14 ($\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]$ and λ, K). *Given a graph $G \sim cSBM(n, \mu, u, \lambda, d)$ and a spectral GNN of order K, $\mathbb{E}[\|\hat{y}_i - y_i\|_F^2]$ for any node v_i satisfies the following: it increases with $\lambda \in [-\sqrt{d}, 0]$, decreases with $\lambda \in [0, \sqrt{d}]$, and reaches its maximum at $\lambda = 0$; it increases with K if $\sum_{k=2}^K \theta_k \frac{(k-1)!}{2^{k-1}}$ grows more slowly than $\sum_{k=2}^K \theta_k^2 \frac{(k-1)!}{2^k}$ as K increases.*

Remark. When $\lambda = 0$, the graph is neither homophilic nor heterophilic, resulting in the maximum expected error. When $\lambda = \pm\sqrt{d}$, the expected error is minimized. This implies that spectral GNNs perform well on strong homophilic or heterophilic graphs but poorly on graphs that are neither. The relationship between the expected norm $\mathbb{E}[\|\hat{y}_i - y_i\|_F^2]$ and the order K is nonetheless intricate, depending on $\Theta = \{\theta_k\}_{k=0}^K$. The proof is provided in Appendix F.

We observe that ζ_k exhibits the same trend as $\mathbb{E}[\|\hat{y}_i - y_i\|_F^2]$ with respect to changes in λ; however, their behavior diverges with respect to K, as characterized in the following theorem.

Theorem 15 (ζ_k and λ, K). *Given a graph $G \sim cSBM(n, \mu, u, \lambda, d)$ and a spectral GNN of order K, ζ_k has the following properties: (1) it increases with $\lambda \in [-\sqrt{d}, 0]$, decreases with $\lambda \in [0, \sqrt{d}]$, and achieves its maximum value at $\lambda = 0$; (2) it increases with k as k grows, for $k \in [0, K]$.*

Remark. When d is fixed, $\lambda \to \sqrt{d}$, nodes with the same class form clusters, and when $\lambda \to -\sqrt{d}$, they form a bipartite structure. In both cases, the graph structure exhibits clear patterns, leading to a small variance $\mathbb{V}\left[A_{ij}^k\right] = \mathbb{E}\left[(A_{ij}^k)^2\right] - (\mathbb{E}\left[A_{ij}^k\right])^2$ and, consequently, a small ζ_k. When $\lambda \to 0$, the graph lacks simple patterns, resulting in a large variance and a correspondingly large ζ_k. When $k \in [0, K]$ increases, more walks between two nodes exist and thus the variance $\mathbb{V}\left[A_{ij}^k\right]$ increases. Larger variance corresponds to a larger ζ_k. The proof is provided in Appendix F.

Based on Theorems 14 and 15, the following proposition summarizes how λ and K influence the γ-uniform transductive stability of spectral GNNs. The proof is provided in Appendix F.

Proposition 16. *For a fixed K, γ-uniform transductive stability and generalization error bound strictly increase as λ moves from $-\sqrt{d}$ to 0, and decreases as λ moves from 0 to $\sqrt{d}$. For a fixed λ, if $\sum_{k=2}^K \theta_k \frac{(k-1)!}{2^{k-1}}$ grows more slowly than $\sum_{k=2}^K \theta_k^2 \frac{(k-1)!}{2^k}$ as K increases, then γ-uniform transductive stability and generalization error bound increase with K.*

5.3 PRACTICAL IMPLICATIONS

We discuss two practical implications of our theoretical findings.

Rewiring graphs: Our analysis establishes a strong connection between graph homophily and the generalization error bound, offering practical insights for rewiring graphs to enhance the performance

Figure 1: Testing accuracy, accuracy gap, and loss gap of five spectral GNNs on synthetic and real-world datasets: **(a)-(c)** *Syn-Varying-H_{edge}*: synthetic datasets with varying $H_{edge} \in [0.1, 0.9]$ (step 0.1) and $m = 3,000$; **(d)-(f)** *Syn-Varying-m*: synthetic datasets with varying training sample sizes $m \in [500, 4500]$ (step 500) of fixed H_{edge}; **(g)-(i)** *Real-m-100*: real-world datasets with varying H_{edge} (shown on the right) and $m = 100$.

of spectral GNNs. Graphs with strong homophilic or heterophilic structures exhibit low variance in k-length walks, which reveals clearer structural patterns. This reduction in variance decreases the gradient norm bound β (Theorems 8 and 13), leading to improved γ-uniform transductive stability (Theorem 6). Enhanced stability, in turn, reduces the generalization error bound (Theorem 9).

Constrained graph convolution: Our theoretical results indicate that constraining the graph convolution parameters $0 \leq \theta_k \leq 1$ prevents the generalization error bound from increasing with the polynomial order K. This is because the constraint ensures that the condition in Proposition 16, where $\sum_{k=2}^{K} \theta_k \frac{(k-1)!}{2^{k-1}}$ increases slower than $\sum_{k=2}^{K} \theta_k^2 \frac{(k-1)!}{2^k}$, is violated, as $\theta_k \geq \theta_k^2$. Previous work (He et al., 2021) reports that constraining θ_k to non-negative values with Bernstein polynomial basis leads to valid polynomial filters. Our analysis further suggests adding the constraint $\theta_k \leq 1$ to maintain stable generalization error as K increases.

6 EXPERIMENTS

Synthetic and real-wrold datasets. We use the following datasets: (1) *Synthetic datasets* consist of nine graphs generated using $G \sim cSBM(n, f, \mu, u, \lambda, d)$ following (Chien et al., 2021). Each graph contains 5,000 nodes with 2,000-dimensional features and an average degree of 5. The edge homophily ratios (H_{edge}) range from 0.1 to 0.9 in steps of 0.1. (2) *Real-world datasets* consist of ten benchmark node classification datasets (Texas, Wisconsin, Actor, Chameleon, Squirrel, Citeseer, Pubmed, Cora, Ogbn-Arxiv, Ogbn-Products) with H_{edge} varying between 0.11 and 0.81. Following previous work (He et al., 2021; 2022; Chien et al., 2021), we randomly split each dataset into 60% for training, 20% for validation, and 20% for testing.

Spectral GNNs. We select five widely recognized spectral GNNs for our experiments: ChebNet (Defferrard et al., 2016), GPRGNN (Chien et al., 2021), BernNet (He et al., 2021), JacobiConv (Wang & Zhang, 2022), and ChebNetII (He et al., 2022). For consistency with Eq. (1), we use a single-layer ChebNet rather than the typical two-layer version.

Further details about the dataset statistics, spectral GNNs, hyper-parameter settings used in our experiments and additional experimental results are provided in Appendix G. Below, we discuss the effects of node class distribution and polynomial order on the accuracy and loss gaps of spectral GNNs.

Figure 2: Accuracy gap and loss gap of JacobiConv and BernNet when the polynomial order K increases, where $\rho_1 = \sum_{k=2}^{K} \theta_k \frac{(k-1)!}{2^{k-1}}$ and $\rho_2 = \sum_{k=2}^{K} \theta_k^2 \frac{(k-1)!}{2^k}$: (a-b),(e-f) show the results on a synthetic dataset of $H_{edge} = 0.2$; (c-d),(g-h) show the results on Chameleon dataset of $H_{edge} = 0.24$.

Figure 1(a)-(c) illustrates that as H_{edge} of synthetic datasets varies from 0.1 to 0.9, accuracy and loss gaps increase monotonically for $H_{edge} \in [0.1, 0.5]$ and decrease for $H_{edge} \in [0.5, 0.9]$, reaching their maximum at $H_{edge} = 0.5$ across all spectral GNNs. This aligns with our theoretical analysis in Proposition 16, which states that the generalization error bound increases as H_{edge} moves from 0 to 0.5 and decreases as H_{edge} moves from 0.5 to 1. In Figure 1(d)-(f), when the training sample number m increases from 500 to 4,500 on synthetic datasets, both accuracy and loss gaps decrease consistently. This is consistent with Lemma 10, which shows that the generalization error bound decreases with increasing m. Figure 1(g)-(i) shows that when training all real-world datasets with the same sample size $m = 100$, spectral GNNs exhibit a similar trend to synthetic datasets. Loss gaps are small at the extreme ends of the H_{edge} range but increase as H_{edge} transitions from Texas (0.11) to Ogbn-Products (0.81). These results align with Proposition 16.

Figure 2 shows that when the slope of ρ_1 is smaller than that of ρ_2, accuracy and loss gaps increase. Conversely, gaps decrease when ρ_1 grows faster than ρ_2, aligning with Proposition 16. Intuitively, this condition can be understood as follows: (1). *Non-negative θ_k*: Spectral GNNs constrained to $0 \le \theta_k \le 1$ demonstrate strong generalization. In this case, $\theta_k \ge \theta_k^2$, ensuring ρ_1 grows faster than ρ_2, violating the condition of Proposition 16, and preventing the generalization error bound from increasing with K. For instance, BernNet enforces non-negative θ_k, and as shown in Figure 2(e-h), its accuracy and loss gaps remain stable with increasing K. (2). *Unrestricted θ_k*: Spectral GNNs allowing both positive and negative θ_k may exhibit poor generalization. If $\theta_k < 0$, $\rho_1 \le \rho_2$. When $\theta_{k_1} \le 0$ and $\theta_{k_2} \ge 0$, ρ_1 typically grows slower than ρ_2, satisfying the condition of Proposition 16 and leading to increasing generalization error bounds with K. For example, JacobiConv does not restrict the sign of θ_k, and its accuracy and loss gaps increase with K in Figure 2(a-b).

7 CONCLUSION, LIMITATIONS, AND FUTURE WORK

This work investigates how node class distribution and architectural choices impact the generalization of spectral GNNs. Our findings show that spectral GNNs generalize well on graphs with strong homophilic or heterophilic structures, where node class distributions exhibit clear patterns, and the generalization error of spectral GNNs increases with polynomial order under certain conditions.

We derive the uniform transductive stability of spectral GNNs on graphs generated by the general multi-class cSBM, providing insights into the relationships between graph homophily, polynomial order, and generalization error bounds. However, this analysis is limited to a specialized cSBM, leaving room for further exploration of more diverse graph generation models. Another limitation lies in architectural choices. While these choices, such as the selection of the graph matrix (e.g., Laplacian vs. adjacency matrix) and polynomial basis (e.g., Chebyshev vs. Bernstein), are critical to generalization performance, we do not explore their specific impacts on generalization bounds. Future work could investigate how these design decisions influence the theoretical and practical performance of spectral GNNs. Finally, our theoretical analysis assumes training with gradient descent, whereas Adam is the optimizer most commonly used in practice. This discrepancy between theoretical assumptions and practical applications highlights an important direction for future research to bridge the gap and improve the relevance of theoretical findings to real-world scenarios.

REFERENCES

George B Arfken, Hans J Weber, and Frank E Harris. *Mathematical methods for physicists: a comprehensive guide*. Academic press, 2011.

Muhammet Balcilar, Guillaume Renton, Pierre Héroux, Benoit Gaüzère, Sébastien Adam, and Paul Honeine. Analyzing the expressive power of graph neural networks in a spectral perspective. In *International Conference on Learning Representations*, 2021.

Aseem Baranwal, Kimon Fountoulakis, and Aukosh Jagannath. Effects of graph convolutions in multi-layer networks. In *The Eleventh International Conference on Learning Representations*.

Aseem Baranwal, Kimon Fountoulakis, and Aukosh Jagannath. Graph convolution for semi-supervised classification: Improved linear separability and out-of-distribution generalization. In *International Conference on Machine Learning*, pp. 684–693. PMLR, 2021.

Aseem Baranwal, Kimon Fountoulakis, and Aukosh Jagannath. Optimality of message-passing architectures for sparse graphs. *Advances in Neural Information Processing Systems*, 36:40320–40341, 2023.

Peter L Bartlett and Shahar Mendelson. Rademacher and gaussian complexities: Risk bounds and structural results. *Journal of Machine Learning Research*, 3(Nov):463–482, 2002.

Arash Behboodi, Gabriele Cesa, and Taco Cohen. A pac-bayesian generalization bound for equivariant networks. *ArXiv*, abs/2210.13150, 2022. URL https://api.semanticscholar.org/CorpusID:253097790.

Olivier Bousquet and André Elisseeff. Stability and generalization. *The Journal of Machine Learning Research*, 2:499–526, 2002.

Vladimir Cherkassky, Xuhui Shao, Filip M Mulier, and Vladimir N Vapnik. Model complexity control for regression using vc generalization bounds. *IEEE transactions on Neural Networks*, 10 (5):1075–1089, 1999.

Eli Chien, Jianhao Peng, Pan Li, and Olgica Milenkovic. Adaptive universal generalized pagerank graph neural network. *arXiv: Learning*, 2021.

Weilin Cong, Morteza Ramezani, and Mehrdad Mahdavi. On provable benefits of depth in training graph convolutional networks. *Advances in Neural Information Processing Systems*, 34:9936–9949, 2021.

Michaël Defferrard, Xavier Bresson, and Pierre Vandergheynst. Convolutional neural networks on graphs with fast localized spectral filtering. *Advances in neural information processing systems*, 29, 2016.

Yash Deshpande, Subhabrata Sen, Andrea Montanari, and Elchanan Mossel. Contextual stochastic block models. *Advances in Neural Information Processing Systems*, 31, 2018.

Laurent Dinh, Razvan Pascanu, Samy Bengio, and Yoshua Bengio. Sharp minima can generalize for deep nets. In *International Conference on Machine Learning*, 2017.

Maximilien Dreveton, Felipe S. Fernandes, and Daniel R. Figueiredo. Exact recovery and bregman hard clustering of node-attributed stochastic block model. *ArXiv*, abs/2310.19854, 2023. URL https://api.semanticscholar.org/CorpusID:264824545.

Rick Durrett. *Probability: theory and examples*, volume 49. Cambridge university press, 2019.

Giuseppe Alessio D'Inverno, Monica Bianchini, and Franco Scarselli. Vc dimension of graph neural networks with pfaffian activation functions. 2024. URL https://api.semanticscholar.org/CorpusID:267095159.

Ran El-Yaniv and Dmitry Pechyony. Stable transductive learning. In *Learning Theory: 19th Annual Conference on Learning Theory, COLT 2006, Pittsburgh, PA, USA, June 22-25, 2006. Proceedings 19*, pp. 35–49. Springer, 2006.

Ran El-Yaniv and Dmitry Pechyony. Transductive rademacher complexity and its applications. In *Annual Conference Computational Learning Theory*, 2007. URL https://api.semanticscholar.org/CorpusID:6444242.

Pascal Esser, Leena Chennuru Vankadara, and Debarghya Ghoshdastidar. Learning theory can (sometimes) explain generalisation in graph neural networks. *Advances in Neural Information Processing Systems*, 34:27043–27056, 2021.

Michael J Evans and Jeffrey S Rosenthal. *Probability and statistics: The science of uncertainty*. Macmillan, 2004.

Billy J Franks, Christopher Morris, Ameya Velingker, and Floris Geerts. Weisfeiler-leman at the margin: When more expressivity matters. *arXiv preprint arXiv:2402.07568*, 2024.

Vikas Garg, Stefanie Jegelka, and Tommi Jaakkola. Generalization and representational limits of graph neural networks. In *International Conference on Machine Learning*, pp. 3419–3430. PMLR, 2020.

Saeed Ghadimi and Guanghui Lan. Stochastic first- and zeroth-order methods for nonconvex stochastic programming. *SIAM J. Optim.*, 23:2341–2368, 2013. URL https://api.semanticscholar.org/CorpusID:14112046.

Mingguo He, Zhewei Wei, Zengfeng Huang, and Hongteng Xu. Bernnet: Learning arbitrary graph spectral filters via bernstein approximation. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2021.

Mingguo He, Zhewei Wei, and Ji rong Wen. Convolutional neural networks on graphs with chebyshev approximation, revisited. *ArXiv*, abs/2202.03580, 2022. URL https://api.semanticscholar.org/CorpusID:246652363.

Haotian Ju, Dongyue Li, Aneesh Sharma, and Hongyang R Zhang. Generalization in graph neural networks: Improved pac-bayesian bounds on graph diffusion. In *International Conference on Artificial Intelligence and Statistics*, 2023.

Thomas Kipf and M. Welling. Semi-supervised classification with graph convolutional networks. In *International Conference on Learning Representations*, 2017.

Ilja Kuzborskij and Christoph Lampert. Data-dependent stability of stochastic gradient descent. In *International Conference on Machine Learning*, pp. 2815–2824. PMLR, 2018.

JING LEI. A goodness-of-fit test for stochastic block models. *The Annals of Statistics*, 44(1):401–424, 2016.

Shaojie Li and Yong Liu. Improved learning rates for stochastic optimization: Two theoretical viewpoints. *ArXiv*, abs/2107.08686, 2021. URL https://api.semanticscholar.org/CorpusID:236087657.

Renjie Liao, Raquel Urtasun, and Richard Zemel. A pac-bayesian approach to generalization bounds for graph neural networks. In *International Conference on Learning Representations*, 2021.

Kangkang Lu, Yanhua Yu, Hao Fei, Xuan Li, Zixuan Yang, Zirui Guo, Meiyu Liang, Mengran Yin, and Tat-Seng Chua. Improving expressive power of spectral graph neural networks with eigenvalue correction. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 38, pp. 14158–14166, 2024.

Jiaqi Ma, Junwei Deng, and Qiaozhu Mei. Subgroup generalization and fairness of graph neural networks. In *Neural Information Processing Systems*, 2021. URL https://api.semanticscholar.org/CorpusID:235669723.

Yao Ma, Xiaorui Liu, Neil Shah, and Jiliang Tang. Is homophily a necessity for graph neural networks? In *10th International Conference on Learning Representations, ICLR 2022*, 2022.

Sohir Maskey, Ron Levie, Yunseok Lee, and Gitta Kutyniok. Generalization analysis of message passing neural networks on large random graphs. *Advances in neural information processing systems*, 35:4805–4817, 2022.

David A. McAllester. Some pac-bayesian theorems. *Machine Learning*, 37:355–363, 1998. URL https://api.semanticscholar.org/CorpusID:11417123.

Christopher Morris, Floris Geerts, Jan Tönshoff, and Martin Grohe. Wl meet vc. In *International Conference on Machine Learning*, pp. 25275–25302. PMLR, 2023.

Kenta Oono and Taiji Suzuki. Optimization and generalization analysis of transduction through gradient boosting and application to multi-scale graph neural networks. *Advances in Neural Information Processing Systems*, 33:18917–18930, 2020.

Oleg Platonov, Denis Kuznedelev, Michael Diskin, Artem Babenko, and Liudmila Prokhorenkova. A critical look at the evaluation of gnns under heterophily: Are we really making progress? *arXiv preprint arXiv:2302.11640*, 2023.

Michael James David Powell. *Approximation theory and methods*. Cambridge university press, 1981.

Franco Scarselli, Ah Chung Tsoi, and Markus Hagenbuchner. The vapnik–chervonenkis dimension of graph and recursive neural networks. *Neural Networks*, 108:248–259, 2018.

Huayi Tang and Y. Liu. Towards understanding the generalization of graph neural networks. *ArXiv*, abs/2305.08048, 2023a. URL https://api.semanticscholar.org/CorpusID:258685824.

Huayi Tang and Yong Liu. Towards understanding the generalization of graph neural networks. In *International Conference on Machine Learning*, 2023b.

Petar Velickovic, Guillem Cucurull, Arantxa Casanova, Adriana Romero, Pietro Lio', and Yoshua Bengio. Graph attention networks. *ArXiv*, abs/1710.10903, 2018.

Saurabh Verma and Zhi-Li Zhang. Stability and generalization of graph convolutional neural networks. In *Proceedings of the 25th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining*, pp. 1539–1548, 2019.

Xiyuan Wang and Muhan Zhang. How powerful are spectral graph neural networks. *ArXiv*, abs/2205.11172, 2022.

Jingzhao Zhang, Tianxing He, Suvrit Sra, and Ali Jadbabaie. Why gradient clipping accelerates training: A theoretical justification for adaptivity. *arXiv preprint arXiv:1905.11881*, 2019.

Yingxue Zhang, Florence Regol, Soumyasundar Pal, Sakif Khan, Liheng Ma, and Mark Coates. Detection and defense of topological adversarial attacks on graphs. In *International Conference on Artificial Intelligence and Statistics*, pp. 2989–2997, 2021.

Jie Zhou, Ganqu Cui, Shengding Hu, Zhengyan Zhang, Cheng Yang, Zhiyuan Liu, Lifeng Wang, Changcheng Li, and Maosong Sun. Graph neural networks: A review of methods and applications. *AI open*, 1:57–81, 2020.

Xianchen Zhou and Hongxia Wang. The generalization error of graph convolutional networks may enlarge with more layers. *Neurocomputing*, 424:97–106, 2021.

Jiong Zhu, Yujun Yan, Lingxiao Zhao, Mark Heimann, Leman Akoglu, and Danai Koutra. Beyond homophily in graph neural networks: Current limitations and effective designs. *Advances in neural information processing systems*, 33:7793–7804, 2020.

A BOUND OF SAMPLE LOSS AND GRADIENT CHANGE WRT. PARAMETERS

In this section, we prove the maximum change of sample loss and maximum change of gradient of loss function with respect to parameters $\{\Theta, W\}$ of spectral GNNs in Eq. (1). These two properties are widely used in the subsequent analysis.

Based on Assumption 1, we have the following lemmas.

Lemma 17 (Bound of Loss function to Parameters). *Under Assumption 1, given a loss function ℓ and a spectral GNN, for parameters $\bar{\Theta}, \bar{W}, \Theta', W'$ and any node v_i with truth class y_i we have*

$$\|\ell(y_i, \hat{y}_i|_{\Theta=\bar{\Theta}, W=\bar{W}}) - \ell(y_i, \hat{y}_i|_{\Theta', W'})\|_F \leq \alpha_1 \sqrt{\|\bar{\Theta} - \Theta'\|_F^2 + \|\bar{W} - W'\|_F^2}$$

where $\alpha_1 = Lip(\ell)Lip(\Psi)$.

Proof. Under Assumption 1, we have

$$\|\ell(y_i, \hat{y}_i|_{\tau=\bar{\tau}}) - \ell(y_i, \hat{y}_i|_{\tau=\tau'})\| \leq Lip(\ell)\|\hat{y}_i|_{\tau=\bar{\tau}} - \hat{y}_i|_{\tau=\tau'}\|_F$$

and

$$\|Lip(\ell)\|\hat{y}_i|_{\tau=\bar{\tau}} - \hat{y}_i|_{\tau=\tau'}\|_F \leq Lip(\Psi)\|\bar{\tau} - \tau'\|_F.$$

This leads to

$$\|\ell(y_i, \hat{y}_i|_{\tau=\bar{\tau}}) - \ell(y_i, \hat{y}_i|_{\tau=\tau'})\| \leq Lip(\ell)Lip(\Psi)\|\bar{\tau} - \tau'\|_F.$$

$\square$

Lemma 18 (Bound of Gradient to Parameters). *Under Assumption 1, Assumption 2, for parameters $\bar{\Theta}, \bar{W}, \Theta', W'$ of a spectral GNN, the following holds for any node v_i with truth class y_i*

$$\|\nabla\ell(y_i, \hat{y}_i|_{\Theta=\bar{\Theta}, W=\bar{W}}) - \nabla\ell(y_i, \hat{y}_i|_{\Theta', W'})\|_F \leq \alpha_2 \sqrt{\|\bar{\Theta} - \Theta'\|_F^2 + \|\bar{W} - W'\|_F^2}$$

where $\alpha_2 = (Smt(\Psi)\beta_1 + Smt(\ell)Lip(\Psi)\beta_2)$.

Proof. Since we know that

$$\nabla\ell(y_i, \hat{y}_i|_{\tau=\bar{\tau}}) = \nabla_{\hat{y}_i}\ell(y, \hat{y}_i)|_{\tau=\bar{\tau}} \cdot \nabla\hat{y}_i|_{\tau=\bar{\tau}}$$

and

$$\nabla\ell(y_i, \hat{y}_i|_{\tau=\tau'}) = \nabla_{\hat{y}_i}\ell(y, \hat{y}_i)|_{\tau=\tau'} \cdot \nabla\hat{y}_i|_{\tau=\tau'}.$$

this gives

$$\nabla\ell(y_i, \hat{y}_i|_{\tau=\bar{\tau}}) - \nabla\ell(y_i, \hat{y}_i|_{\tau=\tau'}) = \nabla_{\hat{y}_i}\ell(y, \hat{y}_i)|_{\tau=\bar{\tau}}(\nabla\hat{y}_i|_{\tau=\bar{\tau}} - \nabla\hat{y}_i|_{\tau=\tau'}) \\ + (\nabla_{\hat{y}_i}\ell(y, \hat{y}_i)|_{\tau=\bar{\tau}} - \nabla_{\hat{y}_i}\ell(y, \hat{y}_i)|_{\tau=\tau'})\nabla\hat{y}_i|_{\tau=\tau'}.$$

Hence, we obtain the following

$$\|\nabla\ell(y_i, \hat{y}_i|_{\tau=\bar{\tau}}) - \nabla\ell(y_i, \hat{y}_i|_{\tau=\tau'})\|_F \leq \|\nabla_{\hat{y}_i}\ell(y, \hat{y}_i)|_{\tau=\bar{\tau}}\|_F \cdot \|\nabla\hat{y}_i|_{\tau=\bar{\tau}} - \nabla\hat{y}_i|_{\tau=\tau'}\|_F \\ + \|\nabla_{\hat{y}_i}\ell(y, \hat{y}_i)|_{\tau=\bar{\tau}} - \nabla_{\hat{y}_i}\ell(y, \hat{y}_i)|_{\tau=\tau'}\|_F \cdot \|\nabla\hat{y}_i|_{\tau=\tau'}\|_F. \tag{5}$$

Under Assumption 1 and Assumption 2, we have:

$$\|\nabla\hat{y}_i|_{\tau=\bar{\tau}} - \nabla\hat{y}_i|_{\tau=\tau'}\|_F \leq Smt(\Psi)\|\bar{\tau} - \tau'\|_F \\ \|\nabla_{\hat{y}_i}\ell(y, \hat{y}_i)|_{\tau=\bar{\tau}}\|_F \leq \beta_1. \tag{6}$$

Under Assumption 1, we have

$$\|\nabla_{\hat{y}_i}\ell(y, \hat{y}_i)|_{\tau=\bar{\tau}} - \nabla_{\hat{y}_i}\ell(y, \hat{y}_i)|_{\tau=\tau'}\|_F \leq Smt(\ell)\|\hat{y}_i|_{\tau=\bar{\tau}} - \hat{y}_i|_{\tau=\tau'}\|_F \\ \leq Smt(\ell)Lip(\Psi)\|\bar{\tau} - \tau'\|_F. \tag{7}$$

Under Assumption 2, we have:

$$\|\nabla\hat{y}_i|_{\tau=\tau'}\|_F \leq \beta_2. \tag{8}$$

Substitute Eq. (6), Eq. (7), Eq. (8) into Eq. (5), we have

$$\|\nabla\ell(y_i, \hat{y}_i|_{\tau=\bar{\tau}}) - \nabla\ell(y_i, \hat{y}_i|_{\tau=\tau'})\|_F \leq Smt(\Psi)\|\bar{\tau} - \tau'\|_F \cdot \beta_1 + Smt(\ell)Lip(\Psi)\|\bar{\tau} - \tau'\|_F \cdot \beta_2 \\ = (Smt(\Psi)\beta_1 + Smt(\ell)Lip(\Psi)\beta_2)\|\bar{\tau} - \tau'\|_F$$

$\square$

B UNIFORM TRANSDUCTIVE STABILITY OF SPECTRAL GNNs

Theorem 6 (Stability and Gradient Norm). *Let Ψ be a spectral GNN trained using gradient descent for T iterations with a learning rate η on a training dataset S_m, and evaluated on a testing set $\mathcal{D}_u$. Under Assumption 1, for all iterations $t \in [1, T]$ and any sample (x_i, y_i) in S_m or $\mathcal{D}_u$, if the gradient norm satisfies $\|\nabla \ell(y_i, \hat{y}_i|_{\Theta^t, W^t})\|_F \leq \beta$, where $\{\Theta^t, W^t\}$ are the parameters at the t-th iteration, then Ψ satisfies γ-uniform transductive stability with:*

$$\gamma = r\beta, \quad r = \frac{2\eta\alpha_1}{m}\sum_{t=1}^{T}(1 + \eta\alpha_2)^{t-1},$$

where $\alpha_1 = Lip(\ell) \cdot Lip(\Psi)$ and $\alpha_2 = Smt(\Psi)\beta_1 + Smt(\ell)Lip(\Psi)\beta_2$.

Proof. We denote $\tau = [\Theta; W]$ as the concatenation of parameters Θ, W. According to Lemma 17, Lemma 18, we have

$$\|\ell(y_i, \hat{y}_i|_\tau) - \ell(y_i, \hat{y}_i|_{\tau'})\|_F \leq \alpha_1 \|\tau - \tau'\|_F$$

where $\alpha_1 = Lip(\ell)Lip(\Psi)$. and

$$\|\nabla\ell(y_i, \hat{y}_i|_\tau) - \nabla\ell(y_i, \hat{y}_i|_{\tau'})\|_F \leq \alpha_2 \|\tau - \tau'\|_F$$

where $\alpha_2 = (Smt(\Psi)\beta_1 + Smt(\ell)Lip(\Psi)\beta_2)$. The updation rule of gradient descent is:

$$\tau^{t+1} = \tau^t - \eta\nabla\mathcal{L}_{S_m}(\tau^t)$$
$$\tau_{ij}^{t+1} = \tau_{ij}^t - \eta\nabla\mathcal{L}_{S_m^{ij}}(\tau_{ij}^t)$$

where

$$\mathcal{L}_{S_m}(\tau^t) = \frac{1}{m}\sum_{r=1}^{m}\ell(y_r, \hat{y}_r|_{\tau^t})$$

$$\mathcal{L}_{S_m^{ij}}(\tau_{ij}^t) = \frac{1}{m}\sum_{r=1}^{m}\ell(y_r, \hat{y}_r|_{\tau_{ij}^t})$$

are the empirical loss on training dataset S_m and S_m^{ij} respectively. The difference between empirical loss is:

$$\mathcal{L}_{S_m^{ij}}(\tau_{ij}^t) - \mathcal{L}_{S_m}(\tau^t) = \frac{1}{m}\left[\sum_{r=1, r \neq i,j}^{m}\left(\ell(y_r, \hat{y}_r|_{\tau_{ij}^t}) - \ell(y_r, \hat{y}_r|_{\tau^t})\right) + \ell(y_j, \hat{y}_j|_{\tau_{it}^t}) - \ell(y_i, \hat{y}_i|_{\tau^t})\right].$$

We derive the parameter difference:

$$\|\tau_{ij}^{t+1} - \tau^{t+1}\|_F = \left\|\tau_{ij}^t - \eta\nabla\mathcal{L}_{S_m^{ij}}(\tau_{ij}^t) - \tau^t + \eta\nabla\mathcal{L}_{S_m}(\tau^t)\right\|_F$$

$$\leq \|\tau_{ij}^t - \tau^t\|_F + \eta\|\nabla(\mathcal{L}_{S_m}(\tau^t) - \mathcal{L}_{S_m^{ij}}(\tau_{ij}^t))\|_F$$

$$= \|\tau_{ij}^t - \tau^t\|_F + \frac{\eta}{m}\left\|\nabla\left[\sum_{\substack{r=1 \\ r \neq i,j}}^{m}\left(\ell(y_r, \hat{y}_r|_{\tau_{ij}^t}) - \ell(y_r, \hat{y}_r|_{\tau^t})\right) + \ell(y_j, \hat{y}_j|_{\tau_{ij}^t}) - \ell(y_i, \hat{y}_i|_{\tau^t})\right]\right\|_F$$

$$\leq \|\tau_{ij}^t - \tau^t\|_F + \frac{\eta}{m}\left\|\sum_{\substack{r=1 \\ r \neq i,j}}^{m}\alpha_2\|\tau_{ij}^t - \tau^t\|_F + \nabla\left[\ell(y_j, \hat{y}_j|_{\tau_{ij}^t}) - \ell(y_i, \hat{y}_i|_{\tau^t})\right]\right\|_F \quad (Assumption\ 1)$$

$$\leq \|\tau_{ij}^t - \tau^t\|_F + \frac{\eta}{m}(m-1)\alpha_2\|\tau_{ij}^t - \tau^t\|_F + \frac{\eta}{m}\left\|\nabla\left[\ell(y_j, \hat{y}_j|_{\tau_{ij}^t}) - \ell(y_i, \hat{y}_i|_{\tau^t})\right]\right\|_F$$

$$\leq \|\tau_{ij}^t - \tau^t\|_F + \frac{\eta}{m}(m-1)\alpha_2\|\tau_{ij}^t - \tau^t\|_F + \frac{2\eta\beta}{m} \quad (Theorem\ 13)$$

$$= \left(1 + \frac{m-1}{m}\eta\alpha_2\right)\|\tau_{ij}^t - \tau^t\|_F + \frac{2\eta\beta}{m}$$

$$\leq (1 + \eta\alpha_2)\|\tau_{ij}^t - \tau^t\|_F + \frac{2\eta\beta}{m}$$

After T iterations, we obtain

$$\left\|\tau_{ij}^T - \tau^T\right\|_F \leq (1+\eta\alpha_2)\left\|\tau_{ij}^{T-1} - \tau^{T-1}\right\|_F + \frac{2\eta\beta}{m}$$

$$\leq (1+\eta\alpha_2)[(1+\eta\alpha_2)\left\|\tau_{ij}^{T-2} - \tau^{T-2}\right\|_F + \frac{2\eta\beta}{m}]$$

$$\leq (1+\eta\alpha_2)^T \left\|\tau_{ij}^0 - \tau^0\right\|_F + \sum_{t=1}^{T}(1+\eta\alpha_2)^{t-1}\frac{2\eta\beta}{m}$$

$$= \sum_{t=1}^{T}(1+\eta\alpha_2)^{t-1}\frac{2\eta\beta}{m}$$

If the loss function ℓ is α_1 Lipschitz continuous, then for the loss function ℓ on any sample (x_i, y_i) with parameter $\tau^T = [\Theta^T; W^T], \tau_{ij}^T = [\Theta_{ij}^T; W_{ij}^T]$, we have:

$$\left|\ell(\hat{y}_i, y_i; \tau^T) - \ell(\hat{y}_i, y_i; \tau_{ij}^T)\right| \leq \alpha_1\left|\tau^T - \tau_{ij}^T\right|$$

$$\leq \alpha_1\sum_{t=1}^{T}(1+\eta\alpha_2)^{t-1}\frac{2\eta\beta}{m}$$

$\square$

C UNIFORM TRANSDUCTIVE STABILITY ON GENERAL MULTI-CLASS CSBM

We derive the uniform transductive stability of spectral GNNs defined in Eq. (1) on graphs generated by $G \sim cSBM(n, f, \Pi, Q)$. Then we discuss how the non-linear feature transformation function affect the stability.

We first give a brief introduction to inequalities and lemmas used in this proof.

Lemma 19 (Jensen's Inequality). *Let X be an arbitrary random variable, and let $f : \mathbb{R}^1 \to \mathbb{R}^1$ be a convex function such that $\mathbb{E}\left[f(X)\right]$ is finite. Then $f(\mathbb{E}\left[f(X)\right]) \leq \mathbb{E}\left[f(X)\right]$.*

Lemma 20 (Markov's Inequality). *If X is a non-negative random variable, then for all $a > 0$,*

$$P(X \geq a) \leq \frac{\mathbb{E}\left[X\right]}{a}.$$

That is, the probability that X exceeds any given value a is no more than the expectation of X divided by a.

Remark. Lemma 19, Lemma 20 are important inequalities about a variable and its expectation. Details can be found in (Evans & Rosenthal, 2004).

Lemma 21 (Cauchy-Schwarz Inequality (Arfken et al., 2011)).

$$(\sum_{k=1}^{n} a_k b_k)^2 \leq (\sum_{k=1}^{n} a_k^2)(\sum_{k=1}^{n} b_k^2)$$

The square of the ℓ_2-norm of product of two vectors is smaller than the product of the square of ℓ_2-norm of each vector.

Lemma 22 (Trace and Frobenius Norm). *For any matrix $A \in \mathbb{R}^{n \times n}$, the relation between its trance and its Frobenius norm is*

$$Tr(A) \leq \sqrt{n} \cdot \|A\|_F.$$

Proof.

$$Tr(A) = \sum_{i=1}^{n} a_{ii} \leq \sum_{i=1}^{n} |a_{ii}| \leq \sqrt{n \cdot \sum_{i=1}^{n}|a_{ii}|^2} \quad (Lemma\ 21) = \sqrt{n} \cdot \sqrt{\sum_{i=1}^{n} a_{ii}^2} = \|A\|_F$$

$\square$

Lemma 23 (Partial Derivatives). *For spectral graph neural networks $\hat{Y} = softmax(\sum_{k=0}^{K} \theta_k \tilde{A}^K XW)$, node feature matrix $X \in \mathbb{R}^{n \times f}$ and ground truth node label matrix $Y \in \mathbb{R}^{n \times C}$, the cross-entropy loss of sample (x_i, y_i) is $\ell(\hat{y}_i, y_i; \Theta, W) = -\sum_{c=1}^{C} Y_{ic} \log\left(\hat{Y}_{ic}\right)$, then the partial derivative of $\ell(\hat{y}_i, y_i; \Theta, W)$ with respect to θ_k and W is*

$$\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial \theta_k} = \sum_{c=1}^{C} \left(\hat{Y}_{ic} - Y_{ic}\right) \left(\tilde{A}^k XW\right)_{ic}$$

and

$$\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial W_{pq}} = \left(\hat{Y}_{iq} - Y_{iq}\right) \left(\sum_{k=0}^{K} \theta_k \tilde{A}^k X\right)_{ip}$$

Proof. We begin with

$$Z = \sum_{k=0}^{K} \theta_k \tilde{A}^k XW, \quad \hat{Y}_{ic} = \frac{e^{Z_{ic}}}{\sum_{c'=1}^{C} e^{Z_{ic'}}}, \quad \ell(\hat{y}_i, y_i; \Theta, W) = -\sum_{c=1}^{C} Y_{ic} \log(\hat{Y}_{ic})$$

Then, we have

$$\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial \hat{Y}_{ic}} = -\sum_{c=1}^{C} \frac{Y_{ic}}{\hat{Y}_{ic}},$$

$$\frac{\partial \hat{Y}_{ic}}{\partial Z_{ic'}} = \hat{Y}_{ic}(\delta_{cc'} - \hat{Y}_{ic'}),$$

where δ_{cq} is the Kronecker delta, which is 1 if $c = c'$ and 0 otherwise.

(1) Gradient w.r.t. θ_k

We have

$$\frac{\partial Z_{ic}}{\partial \theta_k} = (\tilde{A}^k XW)_{ic}$$

By the chain rule of gradient, we have:

$$\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial \theta_k} = -\sum_{c=1}^{C} \frac{\ell(\hat{y}_i, y_i; \Theta, W)}{\partial \hat{Y}_{ic}} \cdot \left(\sum_{c'=1}^{C} \frac{\partial \hat{Y}_{ic}}{\partial Z_{ic'}} \cdot \frac{\partial Z_{ic'}}{\partial \theta_k}\right)$$

$$= -\sum_{c=1}^{C} \frac{Y_{ic}}{\hat{Y}_{ic}} \cdot \left(\sum_{c'=1}^{C} \hat{Y}_{ic} \left(\delta_{cc'} - \hat{Y}_{ic'}\right) \cdot \left(\tilde{A}^k XW\right)_{ic'}\right)$$

$$= -\sum_{c=1}^{C} Y_{ic} \cdot \left(\sum_{c'=1}^{C} \left(\delta_{cc'} - \hat{Y}_{ic'}\right) \cdot \left(\tilde{A}^k XW\right)_{ic'}\right)$$

$$= -\sum_{c=1}^{C} Y_{ic} \cdot \left(\left(\tilde{A}^k XW\right)_{ic} - \sum_{c'=1}^{C} \hat{Y}_{ic'} \left(\tilde{A}^k XW\right)_{ic'}\right)$$

$$= -\sum_{c=1}^{C} Y_{ic} \left(\tilde{A}^k XW\right)_{ic} + \sum_{c'=1}^{C} \hat{Y}_{ic'} \left(\tilde{A}^k XW\right)_{ic'}$$

$$= \sum_{c=1}^{C} \left(\hat{Y}_{ic} - Y_{ic}\right) \left(\tilde{A}^k XW\right)_{ic}$$

(2) Gradient w.r.t. W

Based on the following

$$Z_{ic} = \sum_{k=0}^{K} \theta_k \sum_{j=1}^{n} (\tilde{A}^k)_{ij} \sum_{r=1}^{f} X_{jr} W_{rc},$$

we have

$$\frac{\partial Z_{ic}}{\partial W_{pq}} = \sum_{k=0}^{K} \theta_k \sum_{j=1}^{n} (\tilde{A}^k)_{ij} X_{jp} \delta_{cq} = \delta_{cq} \sum_{k=0}^{K} \theta_k \left(\tilde{A}^k X \right)_{ip}$$

where δ_{cq} is the Kronecker delta, which is 1 if $c = q$ and 0 otherwise. Then, by the chain rule of gradient, we have:

$$
\begin{aligned}
\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial W_{pq}} &= -\sum_{c=1}^{C} \frac{\ell(\hat{y}_i, y_i; \Theta, W)}{\partial \hat{Y}_{ic}} \cdot \left(\sum_{c'=1}^{C} \frac{\partial \hat{Y}_{ic}}{\partial Z_{ic'}} \cdot \frac{\partial Z_{ic'}}{\partial W_{pq}} \right) \\
&= -\sum_{c=1}^{C} \frac{Y_{ic}}{\hat{Y}_{ic}} \cdot \left(\sum_{c'=1}^{C} \hat{Y}_{ic} \left(\delta_{cc'} - \hat{Y}_{ic'} \right) \cdot \left(\delta_{c'q} \sum_{k=0}^{K} \theta_k \left(\tilde{A}^k X \right)_{ip} \right) \right) \\
&= -\sum_{c=1}^{C} Y_{ic} \cdot \left(\sum_{c'=1}^{C} \left(\delta_{cc'} - \hat{Y}_{ic'} \right) \cdot \left(\delta_{c'q} \sum_{k=0}^{K} \theta_k \left(\tilde{A}^k X \right)_{ip} \right) \right) \\
&= -\sum_{c=1}^{C} Y_{ic} \cdot \left(\left(\delta_{cq} \sum_{k=0}^{K} \theta_k \left(\tilde{A}^k X \right)_{ip} \right) - \sum_{c'=1}^{C} \hat{Y}_{ic'} \left(\delta_{c'q} \sum_{k=0}^{K} \theta_k \left(\tilde{A}^k X \right)_{ip} \right) \right) \\
&= -\sum_{c=1}^{C} Y_{ic} \left(\delta_{cq} \sum_{k=0}^{K} \theta_k \left(\tilde{A}^k X \right)_{ip} \right) + \sum_{c'=1}^{C} \hat{Y}_{ic'} \left(\delta_{c'q} \sum_{k=0}^{K} \theta_k \left(\tilde{A}^k X \right)_{ip} \right) \\
&= \sum_{c=1}^{C} \left(\hat{Y}_{ic} - Y_{ic} \right) \left(\delta_{cq} \sum_{k=0}^{K} \theta_k \left(\tilde{A}^k X \right)_{ip} \right) \\
&= \sum_{c=1}^{C} \sum_{k=0}^{K} \theta_k \delta_{cq} \left(\hat{Y}_{ic} - Y_{ic} \right) \left(\tilde{A}^k X \right)_{ip} \\
&= \left(\hat{Y}_{iq} - Y_{iq} \right) \left(\sum_{k=0}^{K} \theta_k \tilde{A}^k X \right)_{ip}
\end{aligned}
$$

$\square$

Theorem 8. *Consider a spectral GNN Ψ with polynomial order K trained using full-batch gradient descent for T iterations with a learning rate η on a training dataset S_m sampled from a graph $G \sim cSBM(n, f, \Pi, Q)$ with average node degree $d \ll n$. When $n \to \infty$ and $K \ll n$, under Assumptions 1, 2, and 4, for any node v_i, $i \in [n]$, and for a constant $\epsilon \in (0, 1)$, with probability at least $1 - \epsilon$, Ψ satisfies γ-uniform transductive stability, where $\gamma = r\beta$ and*

$$
\begin{aligned}
\beta = \frac{1}{\epsilon} \Bigg[&O\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2 \right] \right) + O\left(\|\pi_{y_i}^\top \pi_{y_i} + \Sigma_{y_i}\|_F \right) \\
&+ O\left(\sum_{k=1}^{K} \sum_{j=1}^{n} \mathbb{E}[A_{ij}^k] \left\| \sum_{t=1}^{n} \mathbb{E}[A_{it}^k] \pi_{y_j}^\top \pi_{y_t} + \mathbb{E}[A_{ij}^k] \Sigma_{y_j} \right\|_F \right) \Bigg].
\end{aligned}
$$

Proof. Any spectral GNNs in Eq. (1) with linear feature transformation function, and polynomial basis expanded on normalized graph matrix can be transformed into the format:

$$\hat{Y} = softmax(\sum_{k=0}^{K} \theta_k \tilde{A}^k X W) \tag{9}$$

where $\tilde{A} = D^{-\frac{1}{2}} A D^{-\frac{1}{2}}$ is the normalized graph adjacency matrix, D is the diagonal degree matrix. We denotes $Y \in \mathbb{R}^{n \times C}$ as the ground truth node label matrix.

(1) Walk counting
According to Definition 7, we have

$$\mathbb{E}[A_{ij}^k] = \sum_{p \in P_{ij}^k} \prod_{(v,v') \in p} Q_{yy'}$$

(2) Feature expectation

Since we have $G \sim cSBM(n, f, \Pi, Q)$, node classes have a uniform prior $y_i \sim \mathcal{U}(1, C)$. Thus,

$$
\begin{aligned}
\mathbb{E}\left[XW\right]_{ij} &= \frac{1}{n} \sum_{u=1}^{n} (\pi_{y_u} W)_j \\
&= \frac{1}{n} \sum_{u=1}^{n} \sum_{c=1}^{C} p(y_u = c)(\pi_c W)_j \\
&= \frac{1}{n} \sum_{u=1}^{n} \sum_{c=1}^{C} \frac{1}{C} (\pi_c W)_j \\
&= \frac{1}{C} \sum_{c=1}^{C} (\pi_c W)_j
\end{aligned}
\tag{10}
$$

When $k \geq 1$, we have

$$
\begin{aligned}
\mathbb{E}[(\tilde{A}^k XW)_{ij}] &= \mathbb{E}\left[\tilde{A}_{i:}^k\right] \mathbb{E}\left[(XW)_{:j}\right] \\
&= \sum_{s=1}^{n} \mathbb{E}\left[\tilde{A}_{is}^k\right] \mathbb{E}\left[(XW)_{sj}\right] \\
&= \sum_{s=1}^{n} \mathbb{E}\left[\tilde{A}_{is}^k\right] \cdot \frac{1}{C} \sum_{c=1}^{C} (\pi_c W)_j
\end{aligned}
$$

when $k = 0$, we have

$$
\begin{aligned}
\mathbb{E}[(IXW)_{ij}] &= \mathbb{E}\left[(XW)_{ij}\right] \\
&= \frac{1}{C} \sum_{c=1}^{C} (\pi_c W)_j
\end{aligned}
$$

Thus,

$$
\mathbb{E}[(\tilde{A}^k XW)_{ij}] = \begin{cases} \frac{1}{C} \sum_{c=1}^{C} (\pi_c W)_j, & k = 0 \\ \sum_{s=1}^{n} \mathbb{E}\left[\tilde{A}_{is}^k\right] \cdot \frac{1}{C} \sum_{c=1}^{C} (\pi_c W)_j, & k \geq 1 \end{cases}
\tag{11}
$$

(3) Gradient Norm.

The gradient norm can be relaxed as:

$$
\begin{aligned}
\mathbb{E}\left[\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F\right] &\leq \mathbb{E}\left[\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_{\ell_1}\right] \\
&= \sum_{k=0}^{K} \mathbb{E}\left[\|\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial \theta_k}\|_{\ell_1}\right] + \mathbb{E}\left[\|\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial W}\|_{\ell_1}\right]
\end{aligned}
\tag{12}
$$

According to Eq. (9) and Lemma 23, we get the partial derivatives $\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial \theta_k}$ and $\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial W_{pq}}$.

Specially, when $m = 1$, we get the partial derivatives of empirical loss on training sample (x_i, y_i)

$$
\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial \theta_k} = \sum_{c=1}^{C} \left(\hat{Y}_{ic} - Y_{ic}\right) \left(\tilde{A}^k XW\right)_{ic}
\tag{13}
$$

$$
\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial W_{pq}} = \left(\hat{Y}_{iq} - Y_{iq}\right) \left(\sum_{k=0}^{K} \theta_k \tilde{A}^k X\right)_{ip}
\tag{14}
$$

Thus, we have:

$$
\begin{aligned}
\mathbb{E}\left[\|\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial \theta_k}\|_{\ell_1}\right] &= \mathbb{E}\left[|\sum_{c=1}^{C}\left(\hat{Y}_{ic} - Y_{ic}\right)\left(\tilde{A}^k XW\right)_{ic}|\right] \\
&\leq \sum_{c=1}^{C} \mathbb{E}\left[|\left(\hat{Y}_{ic} - Y_{ic}\right)\left(\tilde{A}^k XW\right)_{ic}|\right] \\
&= \sum_{c=1}^{C} \mathbb{E}\left[|\left(\hat{Y}_{ic} - Y_{ic}\right)| \cdot |\left(\tilde{A}^k XW\right)_{ic}|\right] \\
&\leq \sum_{c=1}^{C} \frac{1}{2}\left(\mathbb{E}\left[\left(\hat{Y}_{ic} - Y_{ic}\right)^2\right] + \mathbb{E}\left[\left(\tilde{A}^k XW\right)_{ic}^2\right]\right), \quad (Lemma\ 28) \\
&= \frac{1}{2}\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \mathbb{E}\left[\|\tilde{A}_{i:}^k XW\|_F^2\right]\right)
\end{aligned}
\tag{15}
$$

and

$$
\begin{aligned}
\mathbb{E}\left[\|\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial W}\|_{\ell_1}\right] &= \sum_{p=1}^{f}\sum_{q=1}^{C} \mathbb{E}\left[\|\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial W_{pq}}\|_{\ell_1}\right] \\
&= \sum_{p=1}^{f}\sum_{q=1}^{C} \mathbb{E}\left[|\left(\hat{Y}_{iq} - Y_{iq}\right)\left(\sum_{k=0}^{K}\theta_k \tilde{A}^k X\right)_{ip}|\right] \\
&\leq \sum_{p=1}^{f}\sum_{k=0}^{K}|\theta_k|\left(\sum_{q=1}^{C} \mathbb{E}\left[|\left(\hat{Y}_{iq} - Y_{iq}\right)| \cdot |\left(\tilde{A}^k X\right)_{ip}|\right]\right) \\
&\leq \sum_{p=1}^{f}\sum_{k=0}^{K}|\theta_k|\left(\mathbb{E}\left[\sum_{q=1}^{C}\left(\hat{Y}_{iq} - Y_{iq}\right)^2\right] + \mathbb{E}\left[\sum_{q=1}^{C}\left(\tilde{A}^k X\right)_{ip}^2\right]\right), \quad (Lemma\ 28) \\
&= \sum_{p=1}^{f}\sum_{k=0}^{K}|\theta_k|\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\mathbb{E}\left[\left(\tilde{A}^k X\right)_{ip}^2\right]\right) \\
&= \sum_{k=0}^{K}|\theta_k|\left(f \cdot \mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\mathbb{E}\left[\|\tilde{A}_{i:}^k X\|_F^2\right]\right)
\end{aligned}
\tag{16}
$$

(4) Expectation $\mathbb{E}\left[\|\tilde{A}_{i:}^k XW\|_F^2\right]$ **and** $\mathbb{E}\left[\|\tilde{A}_{i:}^k X\|_F^2\right]$

For sparse graphs G and its adjacency matrix A, when $d \ll n$ and $k \ll n$, A_{ia}^k, A_{ib}^k can be treated as independent variables due to following reasons: (1) The overlap between walks of different lengths is limited due to the sparsity; (2) there exist k-length walk between two nodes is rare event when $k \ll n$ and the joint occurrences of two rare event can be neglected. (3) when $d \ll n$, the variance of A_{ij}^k can be neglected compared with $\left(\mathbb{E}\left[A_{ij}^k\right]\right)^2$. Thus, by Eq. (11), we have the following for the case $k \geq 1$:

$$\mathbb{E}[\|\tilde{A}_{i:}^k XW\|_F^2] = \mathbb{E}\left[\sum_{c=1}^{C}\left(\sum_{s=1}^{n}\tilde{A}_{is}^k (XW)_{sc}\right)^2\right]$$

$$= \mathbb{E}\left[\sum_{c=1}^{C}\sum_{s=1}^{n}\sum_{t=1}^{n}\tilde{A}_{is}^k \tilde{A}_{it}^k (XW)_{sc}(XW)_{tc}\right]$$

$$= \sum_{c=1}^{C}\sum_{s,t=1}^{n}\mathbb{E}\left[\tilde{A}_{is}^k \tilde{A}_{it}^k (XW)_{sc}(XW)_{tc}\right]$$

$$= \sum_{c=1}^{C}\sum_{s,t=1}^{n}\mathbb{E}\left[\tilde{A}_{is}^k\right]\cdot\mathbb{E}\left[\tilde{A}_{it}^k\right]\cdot\mathbb{E}[(XW)_{sc}(XW)_{tc}]$$

$$= \sum_{c=1}^{C}\sum_{s=1}^{n}\mathbb{E}\left[\tilde{A}_{is}^k\right]\left[\sum_{t=1,t\neq s}\mathbb{E}\left[\tilde{A}_{it}^k\right]\cdot\mathbb{E}[(XW)_{sc}(XW)_{tc}]\right.$$

$$\left. + \mathbb{E}\left[\tilde{A}_{is}^k\right]\cdot\mathbb{E}\left[(XW)_{sc}^2\right]\right]$$

$$= \frac{1}{d^{2k}}\sum_{c=1}^{C}\sum_{s=1}^{n}\mathbb{E}\left[\tilde{A}_{is}^k\right]\left[\sum_{t=1,t\neq s}^{n}\mathbb{E}\left[\tilde{A}_{it}^k\right]\cdot(\pi_{y_s}W)_c\cdot(\pi_{y_t}W)_c\right.$$

$$\left. + \mathbb{E}\left[\tilde{A}_{is}^k\right]\cdot W_{:c}^\top\left(\pi_{y_s}^\top\pi_{y_s}+\Sigma_{y_s}\right)W_{:c}\right]$$

when $k = 0$:

$$\mathbb{E}\left[\|\tilde{A}_{i:}^k XW\|_F^2\right] = \mathbb{E}\left[\|X_{i:}W\|_F^2\right]$$

$$= \mathbb{E}\left[\sum_{c=1}^{C}(XW)_{ic}^2\right]$$

$$= \sum_{c=1}^{C}W_{:c}^\top\left(\pi_{y_i}^\top\pi_{y_i}+\Sigma_{y_i}\right)W_{:c}$$

Thus, we have

$$\mathbb{E}\left[\|\tilde{A}_{i:}^k XW\|_F^2\right] = \begin{cases} \sum_{c=1}^{C}W_{:c}^\top\left(\pi_{y_i}^\top\pi_{y_k}+\Sigma_{y_i}\right)W_{:c}, k=0 \\ \frac{1}{d^{2k}}\sum_{c=1}^{C}\sum_{s=1}^{n}\mathbb{E}\left[\tilde{A}_{is}^k\right]\left[\sum_{t=1,t\neq s}^{n}\mathbb{E}\left[\tilde{A}_{it}^k\right]\cdot(\pi_{y_s}W)_c\cdot(\pi_{y_t}W)_c\right. \\ \left. + \mathbb{E}\left[\tilde{A}_{is}^k\right]\cdot W_{:c}^\top\left(\pi_{y_s}^\top\pi_{y_s}+\Sigma_{y_s}\right)W_{:c}\right], k\geq 1 \end{cases} \quad (17)$$

Similarly, by Eq. (10), we have

$$\mathbb{E}\left[\|\tilde{A}_{i:}^k X\|_F^2\right] = \begin{cases} \sum_{c=1}^{C}I_{:c}^\top\left(\pi_{y_i}^\top\pi_{y_k}+\Sigma_{y_i}\right)I_{:c}, k=0 \\ \frac{1}{d^{2k}}\sum_{q=1}^{f}\sum_{s=1}^{n}\mathbb{E}\left[\tilde{A}_{is}^k\right]\left[\sum_{t=1,t\neq s}^{n}\mathbb{E}\left[\tilde{A}_{it}^k\right]\cdot\pi_{y_s,q}\cdot\pi_{y_t,q}\right. \\ \left. + \mathbb{E}\left[\tilde{A}_{is}^k\right]\cdot I_{:q}^\top\left(\pi_{y_s}^\top\pi_{y_s}+\Sigma_{y_s}\right)I_{:q}\right], k\geq 1 \end{cases} \quad (18)$$

By putting Eq. (17) into Eq. (15), Eq. (18) into Eq. (16), and Eq. (15), Eq. (16) into Eq. (12), we have the following

$$\mathbb{E}\left[\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F\right] \leq \frac{1}{2}\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \sum_{c=1}^{C} W_{:c}^{\top}\left(\pi_{y_i}^{\top}\pi_{y_i} + \Sigma_{y_i}\right)W_{:c}\right)$$

$$+ \sum_{k=1}^{K}\frac{1}{2}\bigg[\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \frac{1}{d^{2k}}\sum_{c=1}^{C}\sum_{s=1}^{n}\mathbb{E}\left[\tilde{A}_{is}^k\right]$$

$$\cdot\bigg[\sum_{t=1,t\neq s}^{n}\mathbb{E}\left[\tilde{A}_{it}^k\right]\cdot\left(\pi_{y_t}W\right)_c\cdot\left(\pi_{y_t}W\right)_c + \mathbb{E}\left[\tilde{A}_{is}^k\right]\cdot W_{:c}^{\top}\left(\pi_{y_s}^{\top}\pi_{y_s} + \Sigma_{y_s}\right)W_{:c}\bigg]\bigg]$$

$$+ |\theta_0|\left(f\cdot\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\sum_{c=q}^{f}I_{:q}^{\top}\left(\pi_{y_i}^{\top}\pi_{y_i} + \Sigma_{y_i}\right)I_{:q}\right)$$

$$+ \sum_{k=1}^{K}|\theta_k|\bigg[f\cdot\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\frac{1}{d^{2k}}\sum_{c=1}^{C}\sum_{s=1}^{n}\mathbb{E}\left[\tilde{A}_{is}^k\right]$$

$$\cdot\bigg[\sum_{t=1,t\neq s}^{n}\mathbb{E}\left[\tilde{A}_{it}^k\right]\cdot\pi_{y_s,q}\cdot\pi_{y_t,q} + \mathbb{E}\left[\tilde{A}_{is}^k\right]\cdot I_{:q}^{\top}\left(\pi_{y_s}^{\top}\pi_{y_s} + \Sigma_{y_s}\right)I_{:q}\bigg]\bigg]$$

$$= \left(\frac{K+1}{2} + f\sum_{k=0}^{K}|\theta_k|\right)\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]$$

$$+ \frac{1}{2}\sum_{c=1}^{C}W_{:c}^{\top}\left(\pi_{y_i}^{\top}\pi_{y_i} + \Sigma_{y_i}\right)W_{:c} + |\theta_0|C\sum_{c=1}^{C}I_{:c}^{\top}\left(\pi_{y_i}^{\top}\pi_{y_i} + \Sigma_{y_i}\right)I_{:c}$$

$$+ \sum_{k=1}^{K}\frac{1}{d^{2k}}\sum_{c=1}^{C}\sum_{s=1}^{n}\mathbb{E}\left[\tilde{A}_{is}^k\right]$$

$$\cdot\bigg[\sum_{t=1,t\neq s}^{n}\mathbb{E}\left[\tilde{A}_{it}^k\right]\cdot\left(\pi_{y_t}W\right)_c\cdot\left(\pi_{y_t}W\right)_c + \mathbb{E}\left[\tilde{A}_{is}^k\right]\cdot W_{:c}^{\top}\left(\pi_{y_s}^{\top}\pi_{y_s} + \Sigma_{y_s}\right)W_{:c}\bigg]$$

$$+ \sum_{k=1}^{K}\frac{C}{d^{2k}}|\theta_k|\sum_{q=1}^{f}\sum_{s=1}^{n}\mathbb{E}\left[\tilde{A}_{is}^k\right]$$

$$\cdot\bigg[\sum_{t=1,t\neq s}^{n}\mathbb{E}\left[\tilde{A}_{it}^k\right]\cdot\pi_{y_s,q}\cdot\pi_{y_t,q} + \mathbb{E}\left[\tilde{A}_{is}^k\right]\cdot I_{:q}^{\top}\left(\pi_{y_s}^{\top}\pi_{y_s} + \Sigma_{y_s}\right)I_{:q}\bigg]$$

$$= \left(\frac{K+1}{2} + f\sum_{k=0}^{K}|\theta_k|\right)\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]$$

$$+ \frac{1}{2}\sum_{c=1}^{C}W_{:c}^{\top}\left(\pi_{y_i}^{\top}\pi_{y_i} + \Sigma_{y_i}\right)W_{:c} + |\theta_0|C\sum_{c=1}^{C}I_{:c}^{\top}\left(\pi_{y_i}^{\top}\pi_{y_i} + \Sigma_{y_i}\right)I_{:c}$$

$$+ \sum_{k=1}^{K}\frac{1}{d^{2k}}\sum_{c=1}^{C}\sum_{s=1}^{n}\mathbb{E}\left[\tilde{A}_{is}^k\right]\left[W_{:c}^{\top}\left(\sum_{\substack{t=1\\t\neq s}}^{n}\mathbb{E}\left[\tilde{A}_{it}^k\right]\pi_{y_s}^{\top}\pi_{y_t} + \mathbb{E}\left[\tilde{A}_{is}^k\right]\left(\pi_{y_s}^{\top}\pi_{y_s} + \Sigma_{y_s}\right)\right)W_{:c}\right]$$

$$+ \sum_{k=1}^{K}\frac{C|\theta_k|}{d^{2k}}\sum_{q=1}^{f}\sum_{s=1}^{n}\mathbb{E}\left[\tilde{A}_{is}^k\right]\left[\sum_{\substack{t=1\\t\neq s}}^{n}\mathbb{E}\left[\tilde{A}_{it}^k\right]\pi_{y_s,q}\pi_{y_t,q} + \mathbb{E}\left[\tilde{A}_{is}^k\right]I_{:q}^{\top}\left(\left(\pi_{y_s}^{\top}\pi_{y_s} + \Sigma_{y_s}\right)\right)I_{:q}\right]$$

We further simplify it and relax it under Assumption 4 that:

$$\mathbb{E}\left[\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F\right] \leq \left(\frac{K+1}{2} + f\sum_{k=0}^{K} B_\Theta\right) \mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]$$

$$+ \frac{1}{2}Tr\left(W^T\left(\pi_{y_i}^\top \pi_{y_i} + \Sigma_{y_i}\right)W\right) + B_\Theta C Tr\left(\pi_{y_i}^\top \pi_{y_i} + \Sigma_{y_i}\right)$$

$$+ \sum_{k=1}^{K}\frac{1}{d^{2k}}\sum_{s=1}^{n}\mathbb{E}\left[\tilde{A}_{is}^k\right]Tr\left(\sum_{\substack{t=1\\t\neq s}}^{n}\mathbb{E}\left[\tilde{A}_{it}^k\right]\pi_{y_s}^\top \pi_{y_t} + \mathbb{E}\left[\tilde{A}_{is}^k\right]\left(\pi_{y_s}^\top \pi_{y_s} + \Sigma_{y_s}\right)\right)$$

$$+ \sum_{k=1}^{K}\frac{CB_\Theta}{d^{2k}}\sum_{s=1}^{n}\mathbb{E}\left[\tilde{A}_{is}^k\right]\left[\sum_{\substack{t=1\\t\neq s}}^{n}\mathbb{E}\left[\tilde{A}_{it}^k\right]Tr\left(\pi_{y_s}^\top \pi_{y_t}\right) + \mathbb{E}\left[\tilde{A}_{is}^k\right]Tr\left(\left(\pi_{y_s}^\top \pi_{y_s} + \Sigma_{y_s}\right)\right)\right] \quad (19)$$

$$\leq \left(\frac{K+1}{2} + fB_\Theta(K+1)\right)\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]$$

$$+ \left(\frac{B_W^2}{2} + B_\Theta C\right)Tr\left(\pi_{y_i}^\top \pi_{y_i} + \Sigma_{y_i}\right)$$

$$+ \sum_{k=1}^{K}\frac{1+CB_\Theta}{d^{2k}}\sum_{j=1}^{n}\mathbb{E}\left[A_{ij}^k\right]Tr\left(\sum_{\substack{t=1\\t\neq j}}^{n}\mathbb{E}\left[A_{it}^k\right]\pi_{y_j}^\top \pi_{y_t} + \mathbb{E}\left[A_{ij}^k\right]\left(\pi_{y_j}^\top \pi_{y_j} + \Sigma_{y_j}\right)\right)$$

With Lemma 22, we rewrite it as

$$\mathbb{E}\left[\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F\right] \leq O\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]\right) + O\left(\|\pi_{y_i}^\top \pi_{y_i} + \Sigma_{y_i}\|_F\right)$$

$$+ O\left(\sum_{k=1}^{K}\sum_{j=1}^{n}\mathbb{E}\left[A_{ij}^k\right]\|\sum_{t=1}^{n}\mathbb{E}\left[A_{it}^k\right]\pi_{y_j}^\top \pi_{y_t} + \mathbb{E}\left[A_{ij}^k\right]\Sigma_{y_j}\|_F\right) \quad (20)$$

(5) Concentration Bound.

By Jensen's inequality (Lemma 19), we have:

$$\mathbb{E}[\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F]^2 \leq \mathbb{E}[\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F^2]$$

i.e.,

$$\mathbb{E}[\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F] \leq \sqrt{\mathbb{E}[\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F^2]} \quad (21)$$

By Markov's inequality (Lemma 20), for a positive constant a, we have:

$$\mathbb{P}(\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F \geq a) \leq \frac{\mathbb{E}[\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F]}{a} = \epsilon \quad (22)$$

solving for a:

$$a = \frac{\mathbb{E}[\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F]}{\epsilon} \quad (23)$$

Therefore, combining Eq. (20), Eq. (21), Eq. (22), Eq. (23), with probability at least $1 - \epsilon$, we have

$$\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F \leq \beta = \frac{1}{\epsilon}\mathbb{E}\left[\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F\right]$$

When $\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F \leq \beta$, according to Theorem 6, spectral GNNs on graphs $G \sim cSBM(n, f, \Pi, Q)$ has γ- uniform transductive stability, we rewrite it in big O format:

$$\gamma = r\beta; \beta = \frac{1}{\epsilon}\left[O\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]\right) + O\left(\|\pi_{y_i}^\top \pi_{y_i} + \Sigma_{y_i}\|_F\right)\right.$$

$$\left. + O\left(\sum_{k=1}^{K}\sum_{j=1}^{n}\mathbb{E}\left[A_{ij}^k\right]\|\sum_{t=1}^{n}\mathbb{E}\left[A_{it}^k\right]\pi_{y_j}^\top \pi_{y_t} + \mathbb{E}\left[A_{ij}^k\right]\Sigma_{y_j}\|_F\right)\right],$$

where r is the same as that in Theorem 6.

$\square$

D GENERALIZATION ERROR BOUND OF SPECTRAL GNNS

We derive the generalization error bound of spectral GNNs based on uniform transductive stability. Then we analyze how training sample number affect the generalization error bound.

We first introduce two lemmas for this proof.

Lemma 24 (Inequality for permutation (El-Yaniv & Pechyony, 2006)). *Let Z be a random permutation vector. Let $f(Z)$ be an (m, q)-symmetric permutation function satisfying $\|f(Z) - f(Z^{ij})\| \leq \beta$ for all $i \in I_1^m, j \in I_{m+1}^{m+q}$. Let $H_2(n) \triangleq \sum_{i=1}^n \frac{1}{i^2}$ and $\omega(m, q) \triangleq q^2 (H_2(m + q) - H_2(q))$. Then*

$$\mathbb{P}\left(f(Z) - \mathbb{E}[f(Z)] \geq \epsilon\right) \leq \exp\left(-\frac{\epsilon^2}{2\beta^2 \Omega(m, q)}\right)$$

Lemma 25 (Risk and uniform stability (El-Yaniv & Pechyony, 2006)). *Give any training set S_m and test set $\mathcal{D}_u$, we have:*

$$\mathbb{E}[\mathcal{L}_{\mathcal{D}_u}(\Theta, W) - \mathcal{L}_{S_m}(\Theta, W)] = \mathbb{E}[\Delta(i, j, i, i)] \quad i \in I_1^m, j \in I_{m+1, m+q}$$

where $\Delta(i, j, i, i)$ is the loss change of sample (x_i, y_i) when model is trained on two datasets exchange sample x_i, y_i in training dataset and sample (x_j, y_j) in testing dataset.

Theorem 9 (Generalization Error Bound). *Let $H_2(n) \triangleq \sum_{i=1}^n \frac{1}{i^2}$ and $\Omega(m, n - m) \triangleq (n - m)^2 (H_2(n) - H_2(n - m))$. For $\epsilon \in (0, 1)$, if a spectral GNN is γ-uniform transductive stability with probability $1 - \epsilon$, then under Assumption 3, for $\delta \in (0, 1)$, with probability at least $(1 - \delta)(1 - \epsilon)$, the generalization error $\mathcal{L}_{\mathcal{D}_u}(\Theta, W) - \mathcal{L}_{S_m}(\Theta, W)$ is upper-bounded by:*

$$\gamma + \left(2\gamma + \left(\frac{1}{n-m} + \frac{1}{m}\right)(B_\ell - \gamma)\right)\sqrt{2\Omega(m, n - m)\log\frac{1}{\delta}}. \tag{3}$$

Proof. Let $\Delta(i, j, s, t) \triangleq \ell(\hat{y}_t, y_t; \Theta_{ij}^T, W_{ij}^T) - \ell(\hat{y}_s, y_s; \Theta^T, W^T)$, where Θ_{ij}^T, W_{ij}^T are model parameters trained on dataset S_m^{ij} for T iterations and Θ^T, W^T are model parameters trained on dataset S_m.

We first derive a bound on the permutation stability of function $f(S_m, \mathcal{D}_u) \triangleq \mathcal{L}_{\mathcal{D}_u}(\Theta, W) - \mathcal{L}_{S_m}(\Theta, W)$:

$$\|(\mathcal{L}_{\mathcal{D}_u}(\Theta, W) - \mathcal{L}_{S_m}(\Theta, W)) - (\mathcal{L}_{\mathcal{D}_u}(\Theta^{ij}, W^{ij}) - \mathcal{L}_{S_m}(\Theta^{ij}, W^{ij}))\| \leq$$

$$\frac{1}{q}\sum_{r=m+1, r\neq j}^{m+q} \|\Delta(i, j, r, r)\| + \frac{1}{q}\|\Delta(i, j, i, j)\| + \frac{1}{m}\sum_{r=1, r\neq i}^m \|\Delta(i, j, r, r)\| + \frac{1}{m}\|\Delta(i, j, j, i)\| \tag{24}$$

where $q = n - m$.

According to Definition 5, Assumption 3 and Theorem 6, we have

$$\max_{1 \leq r \leq m+q} \|\Delta(i, j, r, r)\| \leq \gamma = \alpha_1 \sum_{t=1}^T (1 + \eta\alpha_2)^{t-1} \frac{2\eta\beta}{m}$$

and Eq. (24) is bounded by

$$\|(\mathcal{L}_{\mathcal{D}_u}(\Theta, W) - \mathcal{L}_{S_m}(\Theta, W)) - (\mathcal{L}_{\mathcal{D}_u}(\Theta^{ij}, W^{ij}) - \mathcal{L}_{S_m}(\Theta^{ij}, W^{ij}))\|$$

$$\leq \frac{q-1}{q}\gamma + \frac{1}{q}B_\ell + \frac{m-1}{m}\gamma + \frac{1}{m}B_\ell$$

$$= \left(\frac{q-1}{q} + \frac{m-1}{m}\right)\gamma + \left(\frac{1}{q} + \frac{1}{m}\right)B_\ell$$

Let $\tilde{\beta} = \left(\frac{q-1}{q} + \frac{m-1}{m} \right) \gamma + \left(\frac{1}{q} + \frac{1}{m} \right) B_\ell$. Then, the function $f(S_m, \mathcal{D}_u) = \mathcal{L}_{\mathcal{D}_u}(\Theta, W) - \mathcal{L}_{S_m}(\Theta, W)$ has transductive stability $\tilde{\beta}$. Apply Lemma 24 to $f(S_m, \mathcal{D}_u)$, equating the bound to δ

$$\exp\left(-\frac{\epsilon^2}{2\tilde{\beta}^2 \Omega(m, q)} \right) = \delta$$

we get

$$\epsilon = \tilde{\beta}\sqrt{2\Omega(m, q) \log \frac{1}{\delta}}$$

Therefore, we obtain that the probability at least $1 - \delta$ that

$$\mathcal{L}_{\mathcal{D}_u}(\Theta, W) - \mathcal{L}_{S_m}(\Theta, W) - \mathbb{E}\left[\mathcal{L}_{\mathcal{D}_u}(\Theta^{ij}, W^{ij}) - \mathcal{L}_{S_m}(\Theta^{ij}, W^{ij}) \right] \leq \tilde{\beta}\sqrt{2\Omega(m, q)\log\frac{1}{\delta}} \quad (25)$$

According to Lemma 25 and Theorem 6, for $1 \leq i \leq m, m+1 \leq j \leq n$, we have

$$\mathbb{E}\left[\mathcal{L}_{\mathcal{D}_u}(\Theta^{ij}, W^{ij}) - \mathcal{L}_{S_m}(\Theta^{ij}, W^{ij}) \right] = \mathbb{E}\left[\Delta(i, j, i, i) \right] \leq \gamma \quad (26)$$

Substitute Eq. (26) into Eq. (25), we get:

$$\mathcal{L}_{\mathcal{D}_u}(\Theta, W) \leq \mathcal{L}_{S_m}(\Theta, W) + \gamma + \tilde{\beta}\sqrt{2\Omega(m, q)\log\frac{1}{\delta}}$$

We rewrite it as

$$\mathcal{L}_{\mathcal{D}_u}(\Theta, W) - \mathcal{L}_{S_m}(\Theta, W) \leq \gamma + \left(2\gamma + \left(\frac{1}{n-m} + \frac{1}{m} \right)(B_\ell - \gamma) \right)\sqrt{2\Omega(m, n-m)\log\frac{1}{\delta}}$$

$\square$

Lemma 10. *Consider a spectral GNN trained with m samples as $n \to \infty$. As the sample size m increases, the generalization error bound decreases at the rate $O(1/m) + O\left(\sqrt{2\log(1/\delta)/m} \right)$.*

Proof. (1) $\frac{1}{n-m}$ is neglectable compared with $\frac{1}{m}$

As $m < n$, we have $m = o(n)$.

$\frac{m}{n-m} = \frac{m}{n} \cdot \frac{1}{1-\frac{m}{n}}$ when $n \to \infty$, we have $\frac{m}{n} \to 0$ and $\frac{1}{1-\frac{m}{n}} \to 1$ as $m = o(n)$. Therefore,

$$\lim_{n\to\infty} \frac{m}{n-m} = 0, \lim_{n\to\infty} \frac{\frac{1}{n-m}}{\frac{1}{m}} = 0;$$

which indicates

$$\frac{1}{n-m} = o(\frac{1}{m})$$

(2) $\Omega(m, n-m)$ increase with m

$\Omega(m, n-m) = (n-m)^2 \left(H_2(n) - H_2(n-m) \right)$ and $H_2(k) = \sum_{i=1}^{k} \frac{1}{i^2}$. So

$$H_2(n) - H_2(n-m) = \sum_{i=n-m+1}^{n} \frac{1}{i^2}$$

As

$$m \cdot \frac{1}{n^2} \leq \sum_{i=n-m+1}^{n} \frac{1}{i^2} \leq m \cdot \frac{1}{(n-m)^2},$$

we have

$$m \cdot \frac{1}{n^2} \leq H_2(n) - H_2(n-m) \leq m \cdot \frac{1}{(n-m)^2}.$$

Therefore,

$$(n-m)^2 \cdot m \cdot \frac{1}{n^2} \le \Omega(m, n-m) \le (n-m)^2 \cdot m \cdot \frac{1}{(n-m)^2},$$

i.e.,

$$\frac{m(n-m)^2}{n^2} \le \Omega(m, n-m) \le m$$

Thus,

$$\Omega(m, n-m) = O(m)$$

As $\gamma = O(\frac{1}{m})$, we have

$$\gamma + \left(2\gamma + \left(\frac{1}{n-m} + \frac{1}{m}\right)(B_\ell - \gamma)\right)\sqrt{2\Omega(m, n-m)\log\frac{1}{\delta}}$$

$$= O(\frac{1}{m}) + \left(O(\frac{1}{m}) + \left(o(\frac{1}{m}) + \frac{1}{m}\right)\left(B_\ell - O(\frac{1}{m})\right)\right)\sqrt{2O(m)\log\frac{1}{\delta}}$$

$$= O(\frac{1}{m}) + B_\ell O(\frac{1}{m})O(m^{1/2})\sqrt{2\log\frac{1}{\delta}}$$

$$= O\left(\frac{1}{m} + B_\ell\sqrt{\frac{2\log(\frac{1}{\delta})}{m}}\right)$$

Thus, the total effect on bound is: $O\left(\frac{1}{m} + O(\sqrt{\frac{2\log(\frac{1}{\delta})}{m}})\right)$.

$\square$

Proposition 11. *For a spectral GNN $\Psi_{\tilde{\sigma}}$ with a non-linear feature transformation function $f_W(X) = \tilde{\sigma}(XW)$, assume the gradient norm bound β in Theorem 9 is the same for Ψ and $\Psi_{\tilde{\sigma}}$. If $Lip(\tilde{\sigma}) \le 1$ and $Smt(\tilde{\sigma}) \le 1$, then $\gamma_{\tilde{\sigma}} \le \gamma$, where $\gamma_{\tilde{\sigma}}$ is the stability of $\Psi_{\tilde{\sigma}}$.*

Proof. We consider spectral GNN Ψ:

$$\Psi(M, X) = \sigma(\sum_{k=0}^{K} \tilde{A}^k XW)$$

and spectral GNN $\Psi_{\tilde{\sigma}}$:

$$\Psi_{\tilde{\sigma}}(M, X) = \sigma(\sum_{k=0}^{K} \tilde{\sigma}\left(\tilde{A}^k XW\right))$$

(1) Lipschitz Constant:

For any two sets of parameters (Θ_1, W_1) and (Θ_2, W_2):

$$\|\Psi_{\tilde{\sigma}}(\Theta_1, W_1) - \Psi_{\tilde{\sigma}}(\Theta_2, W_2)\|$$

$$= \|\sigma(\sum_{i=0}^{K} \theta_{1k} \tilde{\sigma}(\tilde{A}^k X W_1)) - \sigma(\sum_{i=0}^{K} \theta_{2k} \tilde{\sigma}(\tilde{A}^k X W_2))\|$$

$$\leq Lip(\sigma)\| \sum_{i=0}^{K} \theta_{1k} \tilde{\sigma}(\tilde{A}^k X W_1) - \sum_{i=0}^{K} \theta_{2k} \tilde{\sigma}(\tilde{A}^k X W_2)\|$$

$$\leq Lip(\sigma)\| \sum_{i=0}^{K} (\theta_{1k} - \theta_{2k}) \tilde{\sigma}(\tilde{A}^k X W_1) + \sum_{i=0}^{K} \theta_{2k}(\tilde{\sigma}(\tilde{A}^k X W_1) - \tilde{\sigma}(\tilde{A}^k X W_2))\|$$

$$\leq Lip(\sigma)(\| \sum_{i=0}^{K} (\theta_{1k} - \theta_{2k}) \tilde{\sigma}(\tilde{A}^k X W_1)\| + \| \sum_{i=0}^{K} \theta_{2k}(\tilde{\sigma}(\tilde{A}^k X W_1) - \tilde{\sigma}(\tilde{A}^k X W_2))\|)$$

$$\leq Lip(\sigma)(\|\Theta_1 - \Theta_2\|_F \cdot \max_k \|\tilde{\sigma}(\tilde{A}^k X W_1)\|_2 + \|\Theta_2\|_F \cdot Lip(\tilde{\sigma}) \cdot \max_k \|\tilde{A}^k X (W_1 - W_2)\|_2)$$

Since $Lip(\tilde{\sigma}) \leq 1$, we have:

$$\|\Psi_{\tilde{\sigma}}(\Theta_1, W_1) - \Psi_{\tilde{\sigma}}(\Theta_2, W_2)\| \leq Lip(\sigma)(\|\Theta_1 - \Theta_2\|_F \cdot C_1 + \|\Theta_2\|_F \cdot \|W_1 - W_2\|_F \cdot C_2)$$

where C_1, C_2 are constants depending on $X, \tilde{A}$.

The right hand side is identical to the bound we get for Ψ without activation function. Therefore, $Lip(\Psi_{\tilde{\sigma}}) \leq Lip(\Psi)$.

(2) Smoothness Constant:

We first get partial derivatives of Ψ:

$$\frac{\partial \Psi}{\partial \theta_k} = \nabla \sigma(\sum_{i=0}^{K} \theta_i \tilde{A}^i X W) \cdot \tilde{A}^k X W$$

$$\frac{\partial \Psi_{\tilde{\sigma}}}{\partial \theta_k} = \nabla \sigma(\sum_{i=0}^{K} \theta_i \tilde{\sigma}(\tilde{A}^i X W)) \cdot \tilde{\sigma}(\tilde{A}^k X W)$$

Partial derivatives of $\Psi_{\tilde{\sigma}}$ are:

$$\frac{\partial \Psi}{\partial W} = \nabla \sigma(\sum_{i=0}^{K} \theta_i \tilde{A}^i X W) \cdot \sum_{i=0}^{K} \theta_i \tilde{A}^i X$$

$$\frac{\partial \Psi_{\tilde{\sigma}}}{\partial W} = \nabla \sigma(\sum_{i=0}^{K} \theta_i \tilde{\sigma}(\tilde{A}^i X W)) \cdot \sum_{i=0}^{K} \theta_i \nabla \tilde{\sigma}(\tilde{A}^i X W) \cdot \tilde{A}^i X$$

The Lipschitz constant of these gradients determine the smoothness. For $\Psi_{\tilde{\sigma}}$, the additional $\tilde{\sigma}$ and $\nabla\tilde{\sigma}$ terms do not increase the Lipschitz constant of the gradient as $Lip(\tilde{\sigma}) \leq 1, Smt(\tilde{\sigma}) \leq 1$.

1) $\tilde{\sigma}$ is 1-Lipschitz, so it doesn't increase the difference between inputs. 2) $\nabla\tilde{\sigma}$ is bounded by 1 (since $Smt(\tilde{\sigma}) \leq 1$), so it doesn't amplify the gradient.

Therefore, the Lipschitz constant of the gradient of $\Psi_{\tilde{\sigma}}$ is at most equal to that of Ψ, i.e., :

$$Smt(\Psi_{\tilde{\sigma}}) \leq Smt(\Psi)$$

(3) stability $\gamma_{\tilde{\sigma}}$

According to Theorem 6, we have $\alpha_1 = Lip(\ell) \cdot Lip(\Psi)$ and $\alpha_2 = Smt(\Psi)\beta_1 + Smt(\ell)Lip(\Psi)\beta_2$. Thus, we have a smaller $\alpha_{1\tilde{\sigma}}, \alpha_{2\tilde{\sigma}}$ as $Lip(\Psi_{\tilde{\sigma}}) \leq Lip(\Psi)$ and $\Psi_{\tilde{\sigma}} \leq Smt(\Psi)$. Then, we have $r_{\tilde{\sigma}} \leq r$.

As β is the same for $\Psi_{\tilde{\sigma}}$ and Ψ and $\gamma_{\tilde{\gamma}} = \beta r_{\tilde{\sigma}}, \gamma = \beta r$, we have

$$\gamma_{\tilde{\sigma}} \leq \gamma$$

$\square$

E UNIFORM TRANSDUCTIVE STABILITY ON cSBM

We show the uniform transductive stability of spectral GNNs of architecture in Eq. (1) on graphs generated by $G \sim cSBM(n, f, \mu, u, \lambda, d)$. Theorem 13 is a specialized form of Theorem 8. The data model is specialized to be nodes of binary classes and Gaussian node features.

We first introduce lemmas that are closely related to calculating node features after graph convolution in Appendix E.1. Then we give the expectation and variance of element A_{ij}^k in adjacency matrix and the expectation and variance of node features after graph convolution in Appendix E.2. Based on that, we derive the transductive stability of spectral GNNs on specialized data model in Appendix E.3.

E.1 LEMMAS FOR THEOREM 13

Lemma 26 (Poisson Limit Theorem (Durrett, 2019))**.** *For each n, let $X_{n,m}, 1 \leq m \leq n$ be independent random variables with $\mathbb{P}\left[X_{n,m} = 1\right] = p_{n,m}, \mathbb{P}\left[X_{n,m} = 0\right] = 1 - p_{n,m}$. Suppose (1) $\sum_{m=1}^{n} p_{n,m} \to \lambda \in (0, \infty)$ and (2) $\max_{1 \leq m \leq n} p_{n,m} \to 0$, if $S_n = \sum_{m=1}^{n} X_{n,m}$, then S_n obeys the Gaussian distribution $Poisson(\lambda)$.*

Remark. The Poisson limit theorem is also know as law of rare events. It states that the total number of events will follow a Poisson distribution if the probability of occurrence of an event is small in every trail and it may occur in a large number of trials. More details can be found in (Durrett, 2019).

Lemma 27 (Binomial coefficient approximation)**.** *When $n \gg k$, the binomial coefficient $\binom{n}{k} = \frac{n^k}{k!}$.*

Proof.

$$
\begin{aligned}
\binom{n}{k} &= \frac{n!}{k!(n-k)!} \\
&= \frac{n \cdot (n-1) \cdot (n-2) \cdot \ldots \cdot (n-k+1) \cdot (n-k)!}{k! \cdot (n-k)!} \\
&= \frac{n \cdot (n-1) \cdot (n-2) \cdot \ldots \cdot (n-k+1) \cdot (n-k)!}{k! \cdot (n-k)!} \\
&= \frac{n \cdot (n-1) \cdot (n-2) \cdot \ldots \cdot (n-k+1)}{k!} \\
&= \frac{n^k}{k!}, \quad n \gg k
\end{aligned}
$$

$\square$

Lemma 28 (Expecatations of $\mathbb{E}\left[AB\right]$)**.** *For any two random variable A, B, we have the expectations*

$$
\mathbb{E}[AB] \leq \frac{1}{2}\mathbb{E}[A^2] + \frac{1}{2}\mathbb{E}[B^2]
$$

Proof. Define a function $f(t)$ for any real number t:

$$
f(t) = \mathbb{E}[(\frac{1}{\sqrt{2}}A - \frac{t}{\sqrt{2}}B)^2]
$$

Since this is an expectation of a squared term, we have $f(t) \geq 0$ for any real t.

Expand $f(t)$:

$$
f(t) = \mathbb{E}[\frac{1}{2}A^2 - tAB + \frac{t^2}{2}B^2] = \frac{1}{2}\mathbb{E}[A^2] - t\mathbb{E}[AB] + \frac{t^2}{2}\mathbb{E}[B^2]
$$

set $t = 1$, we have

$$
\frac{1}{2}\mathbb{E}[A^2] - \mathbb{E}[AB] + \frac{1}{2}\mathbb{E}[B^2] \geq 0
$$

i.e.,

$$
\mathbb{E}[AB] \leq \frac{1}{2}\mathbb{E}[A^2] + \frac{1}{2}\mathbb{E}[B^2]
$$

$\square$

Lemma 29 (Monotonicity of $g(\lambda) = \left(\left(d + \lambda\sqrt{d} \right)^k - \left(d - \lambda\sqrt{d} \right)^k \right)^2$). *The function* $g(\lambda) =$

$\left(\left(d + \lambda\sqrt{d} \right)^k - \left(d - \lambda\sqrt{d} \right)^k \right)^2, \lambda \in [-\sqrt{d}, \sqrt{d}]$

- *monotonously increases on* $\lambda \in [0, \sqrt{d}]$;
- *monotonously decreases on* $\lambda \in [-\sqrt{d}, 0]$;
- *achieves the minimum value when* $\lambda = 0$.

Proof. First, observe that $g(\lambda)$ is a symmetry function. Thus, we only need to analyze its behaviour for $\lambda \geq 0$ and then mirror the results for $\lambda < 0$.

Define:

$$A = d + \lambda\sqrt{d}, \quad B = d - \lambda\sqrt{d}$$

So, the function becomes:

$$g(\lambda) = (A^k - B^k)^2$$

Compute the derivative $g'(\lambda)$:

$$g'(\lambda) = 2k\sqrt{d}(A^k - B^k)(A^{k-1} + B^{k-1})$$

When $\lambda \geq 0$, $A \geq B \geq 0$, both $(A^k - B^k)$ and $(A^{k-1} + B^{k-1})$ are non-negative, thus, $g'(\lambda) \geq 0$. This indicates that $g(\lambda)$ monotonously increases on $\lambda \in [0, \sqrt{d}]$.

Due to the even symmetry of $g(\lambda)$, $g(\lambda)$ monotonously decreases on $\lambda \in [-\sqrt{d}, 0]$.

$\square$

Lemma 30 (Monotonicity of $g(\lambda) = \sum_{s=1}^{k} \left(d + \lambda\sqrt{d} \right)^{k-s} \left(d - \lambda\sqrt{d} \right)^s$). *The function* $g(\lambda) =$

$\sum_{s=1}^{k} \left(d + \lambda\sqrt{d} \right)^{k-s} \left(d - \lambda\sqrt{d} \right)^s, \lambda \in [-\sqrt{d}, \sqrt{d}]$

- *monotonously decreases on* $\lambda \in [0, \sqrt{d}]$;
- *monotonously increases on* $\lambda \in [-\sqrt{d}, 0]$;
- *achieves the maximum value at* $\lambda = 0$.

Proof. $g(\lambda)$ can be rewritten as

$$g(\lambda) = (2d)^k - \left(d + \lambda\sqrt{d} \right)^k - \left(d - \lambda\sqrt{d} \right)^k$$

compute the first derivative of $g(\lambda)$ with respect to λ:

$$g'(\lambda) = k\sqrt{d} \left[\left(d - \lambda\sqrt{d} \right)^{k-1} - \left(d + \lambda\sqrt{d} \right)^{k-1} \right]$$

- when $\lambda > 0$, we have $\left(d - \lambda\sqrt{d} \right) < \left(d + \lambda\sqrt{d} \right)$, i.e., $g'(\lambda) < 0$, thus, $g(\lambda)$ is strictly decreasing;
- when $\lambda < 0$, we have $\left(d - \lambda\sqrt{d} \right) > \left(d + \lambda\sqrt{d} \right)$, i.e., $g'(\lambda) > 0$, thus, $g(\lambda)$ is strictly increasing;
- when $\lambda = 0$, $g'(\lambda) > 0$, $g(\lambda)$ achieves the maximum value.

$\square$

E.2 Expectation and Variance of A_{ij}^k and $\left(\tilde{A}^k X W\right)_{ij}$

Theorem 31 (Expectation and variance of A_{ij}^k). *Given a graph generated by $G \sim cSBM(n, f, \mu, u, \lambda, d)$. When $n \to \infty, d \ll n, 2 \le k \le k^2 \ll n$, the k-length walk connecting node v_i, v_j obeys Poisson distribution $Poisson(\rho')$, where*

$$
\rho' = \begin{cases} \rho_= = \frac{(k-1)!}{n \cdot 2^{k-1}} \sum_{a=2}^{k+1} O \left(\sum_{s=\min(2, 2(a-2), 2(k+1-a))}^{\min(2(a-1), 2(k+1-a))} c_{in}^{k-s} \cdot c_{out}^s \right), & \text{if} \quad y_i = y_j \\ \rho_{\neq} = \frac{(k-1)!}{n \cdot 2^{k-1}} \sum_{a=1}^{k} O \left(\sum_{s=1}^{\min(2a-1, 2(k-a)+1)} c_{in}^{k-s} \cdot c_{out}^s \right), & \text{if} \quad y_i \neq y_j \end{cases}
$$

and expectation is $\mathbb{E}\left[A_{ij}^k\right] = \rho'$, variance is $\mathbb{V}\left[A_{ij}^k\right] = \rho'$.

When $k = 1$, the walk connecting node v_i, v_j obeys Bernoulli distribution $Ber(p)$, where

$$
p = \begin{cases} p_= = \frac{c_{in}}{n}, & \text{if} \quad y_i = y_j \\ p_{\neq} = \frac{c_{out}}{n}, & \text{if} \quad y_i \neq y_j \end{cases}
$$

and expectation is $\mathbb{E}\left[A_{ij}^k\right] = p$, variance is $\mathbb{V}\left[A_{ij}^k\right] = p(1-p)$.

Proof. According to Definition 7, we have

$$
\mathbb{E}[A_{ij}^k] = \sum_{p \in P_{ij}^k} \prod_{(v,v') \in p} Q_{yy'}
$$

When $C = 2$, we have

$$
Q_{yy'} = \begin{cases} \frac{c_{in}}{n}, & \text{if} \quad y = y' \\ \frac{c_{out}}{n}, & \text{if} \quad y \neq y' \end{cases} \tag{27}
$$

Case 1: $y_i = y_j$ **and** $k \ge 2$

For nodes v_i and v_j sharing the same class y_i, we consider walks of length k that include a nodes sharing the class y_i and $k + 1 - a$ nodes with different classes.

As we start at v_i and end at v_j, both with class y_i, we need to choose $a - 2$ nodes from the same cluster and $k - a$ nodes from the other cluster.

The total number of ways to arrange these nodes in a walk is $(k - 1)!$ as we have $k - 1$ positions to fill. The probability of each edge depends on whether it's connecting same-class or different-class nodes.

Number of ways to choose the nodes:

- Choose $a - 2$ nodes from $(\frac{n}{2} - 2)$ nodes in the same cluster: $\binom{\frac{n}{2}-2}{a-2}$;

- Choose $k - a + 1$ nodes from $\frac{n}{2}$ nodes in the other cluster: $\binom{\frac{n}{2}}{k-a+1}$.

Number of ways to arrange these nodes: $(k - 1)!$.

Consider the class change of the k-length walk, we denote s the number of walk of class change, when $2a \ge k + 1$, we have $s_{min} = \min(2, 2(k + 1 - a)), s_{max} = 2(k + 1 - a)$; when $2a \le k + 1$, we have $s_{min} = \min(2, 2(a - 2)), s_{max} = 2(a - 1)$.

Therefore, the probability that there are walk of length k and a nodes on walk sharing same class with v_i is:

$$
p_k^a(v_i, v_j \mid y_i = y_j) =
$$

$$
\begin{cases} \binom{\frac{n}{2}-2}{a-2} \cdot \binom{\frac{n}{2}}{k-a+1} \cdot (k-1)! \cdot \left(\sum_{s=\min(2, 2(k+1-a))}^{2(k+1-a)} \left(\frac{c_{in}}{n}\right)^{k-s} \cdot \left(\frac{c_{out}}{n}\right)^s \right), & \text{if} \quad 2a \ge k+1; \\ \binom{\frac{n}{2}-2}{a-2} \cdot \binom{\frac{n}{2}}{k-a+1} \cdot (k-1)! \cdot \left(\sum_{s=\min(2, 2(a-2))}^{2(a-1)} \left(\frac{c_{in}}{n}\right)^{k-s} \cdot \left(\frac{c_{out}}{n}\right)^s \right), & \text{if} \quad 2a < k+1; \end{cases}
$$

The probability that there are walk of length k connecting node v_i, v_j and $y_i = y_j$ is:

$$p_k(v_i, v_j | y_i = y_j) =$$

$$\sum_{a=2}^{\frac{k+1}{2}} \binom{\frac{n}{2} - 2}{a - 2} \cdot \binom{\frac{n}{2}}{k - a + 1} \cdot (k-1)! \cdot \left(\sum_{s=\min(2,2(a-2))}^{2(a-1)} \left(\frac{c_{in}}{n} \right)^{k-s} \cdot \left(\frac{c_{out}}{n} \right)^s \right)$$

$$+ \sum_{\frac{k+1}{2}}^{k+1} \binom{\frac{n}{2} - 2}{a - 2} \cdot \binom{\frac{n}{2}}{k - a + 1} \cdot (k-1)! \cdot \left(\sum_{s=\min(2,2(k+1-a))}^{2(k+1-a)} \left(\frac{c_{in}}{n} \right)^{k-s} \cdot \left(\frac{c_{out}}{n} \right)^s \right) \quad (28)$$

When $k \ll n$, using Lemma 27, binomial coefficients

$$\binom{\frac{n}{2} - 2}{a - 2} = \frac{\left(\frac{n}{2} - 2 \right)^{a-2}}{(a-2)!}$$

$$\binom{\frac{n}{2}}{k - a + 1} = \frac{\left(\frac{n}{2} \right)^{k-a+1}}{(k - a + 1)!}$$

Then,

$$\binom{\frac{n}{2} - 2}{a - 2} \cdot \binom{\frac{n}{2}}{k - a + 1} \cdot (k-1)! = \frac{\left(\frac{n}{2} - 2 \right)^{a-2}}{(a-2)!} \cdot \frac{\left(\frac{n}{2} \right)^{k-a+1}}{(k - a + 1)!} \cdot (k-1)!$$

$$= O\left(\left(\frac{n}{2} \right)^{k-1} \cdot \binom{k-1}{a-2} \right)$$

Then we can simplify Eq. (28) to

$$p_k(v_i, v_j | y_i = y_j) = \sum_{a=2}^{\frac{k+1}{2}} O\left(\left(\frac{n}{2} \right)^{k-1} \cdot \binom{k-1}{a-2} \right) \cdot \left(\sum_{s=\min(2,2(a-2))}^{2(a-1)} \left(\frac{c_{in}}{n} \right)^{k-s} \cdot \left(\frac{c_{out}}{n} \right)^s \right)$$

$$+ \sum_{\frac{k+1}{2}}^{k+1} O\left(\left(\frac{n}{2} \right)^{k-1} \cdot \binom{k-1}{a-2} \right) \cdot \left(\sum_{s=\min(2,2(k+1-a))}^{2(k+1-a)} \left(\frac{c_{in}}{n} \right)^{k-s} \cdot \left(\frac{c_{out}}{n} \right)^s \right)$$

$$= \frac{1}{n \cdot 2^{k-1}} \sum_{a=2}^{\frac{k+1}{2}} O\left(\binom{k-1}{a-2} \cdot \left(\sum_{s=\min(2,2(a-2))}^{2(a-1)} c_{in}^{k-s} \cdot c_{out}^s \right) \right)$$

$$+ \frac{1}{n \cdot 2^{k-1}} \sum_{\frac{k+1}{2}}^{k+1} O\left(\binom{k-1}{a-2} \cdot \left(\sum_{s=\min(2,2(k+1-a))}^{2(k+1-a)} c_{in}^{k-s} \cdot c_{out}^s \right) \right)$$

$$= \frac{(k-1)!}{n \cdot 2^{k-1}} \sum_{a=2}^{k+1} O\left(\sum_{s=\min(2,2(a-2),2(k+1-a))}^{\min(2(a-1),2(k+1-a))} c_{in}^{k-s} \cdot c_{out}^s \right)$$

$$(29)$$

Case 2: $y_i \neq y_j$ and $k \geq 2$

For node v_i, v_j, when they have different classes $y_i \neq y_j$, we count the walk number that the walk has length k and there are a nodes of same class of y_i and $k + 1 - a$ nodes of different class of y_i.

We need to choose $a - 1$ nodes from the same cluster as v_i and $k - a$ nodes from the cluster of v_j as $y_i \neq y_j$.

The total number of ways to arrange these nodes in a walk is $(k - 2)!$ as we have $k - 2$ positions to fill.

Number of ways to choose the nodes:

- Choose $a - 1$ nodes from $(\frac{n}{2} - 1)$ nodes in the same cluster as v_i: $\binom{\frac{n}{2}-1}{a-1}$;

- Choose $k - a$ nodes from $(\frac{n}{2} - 1)$ nodes in the same cluster as v_j: $\binom{\frac{n}{2}-1}{k-a}$.

Number of ways to arrange these nodes: $(k - 1)!$.

Consider the class change of the k-length walk, we denote s the number of walk of class change, when $2a \geq k + 1$, we have $s_{min} = 1, s_{max} = 2(k - a) + 1$; when $2a \leq k + 1$, we have $s_{min} = 1, s_{max} = 2a - 1$.

Therefore, the probability that there are walk of length k and a nodes on walk sharing same class with v_i is

$$p_k^a(v_i, v_j | y_i \neq y_j) =$$
$$\begin{cases} \binom{\frac{n}{2}-1}{a-1} \cdot \binom{\frac{n}{2}-1}{k-a} \cdot (k-1)! \cdot \left(\sum_{s=1}^{2(k-a)+1} \left(\frac{c_{in}}{n}\right)^{k-s} \cdot \left(\frac{c_{out}}{n}\right)^s \right), & \text{if} \quad 2a \geq k + 1 \\ \binom{\frac{n}{2}-1}{a-1} \cdot \binom{\frac{n}{2}-1}{k-a} \cdot (k-1)! \cdot \left(\sum_{s=1}^{2a-1} \left(\frac{c_{in}}{n}\right)^{k-s} \cdot \left(\frac{c_{out}}{n}\right)^s \right), & \text{if} \quad 2a < k + 1 \end{cases}$$

The probability that there are walk of length k connecting node v_i, v_j and $y_i \neq y_j$ is

$$p_k(v_i, v_j | y_i \neq y_j) =$$
$$\sum_{a=1}^{\frac{k+1}{2}} \binom{\frac{n}{2}-1}{a-1} \cdot \binom{\frac{n}{2}-1}{k-a} \cdot (k-1)! \cdot \left(\sum_{s=1}^{2a-1} \left(\frac{c_{in}}{n}\right)^{k-s} \cdot \left(\frac{c_{out}}{n}\right)^s \right)$$
$$+ \sum_{a=\frac{k+1}{2}}^{k} \binom{\frac{n}{2}-1}{a-1} \cdot \binom{\frac{n}{2}-1}{k-a} \cdot (k-1)! \cdot \left(\sum_{s=1}^{2(k-a)+1} \left(\frac{c_{in}}{n}\right)^{k-s} \cdot \left(\frac{c_{out}}{n}\right)^s \right) \tag{30}$$

When $k \ll n$, using Lemma 27, we have $\binom{\frac{n}{2}-1}{a-1} = \frac{(\frac{n}{2}-1)^{a-1}}{(a-1)!}, \binom{\frac{n}{2}-1}{k-a} = \frac{(\frac{n}{2}-1)^{k-a}}{(k-a)!}$.

Then:

$$\binom{\frac{n}{2}-1}{a-1} \cdot \binom{\frac{n}{2}-1}{k-a} \cdot (k-1)! = \frac{(\frac{n}{2}-1)^{a-1}}{(a-1)!} \cdot \frac{(\frac{n}{2}-1)^{k-a}}{(k-a)!} \cdot (k-1)!$$
$$= \left(\frac{n}{2}-1\right)^{k-1} \cdot \binom{k-1}{a-1}$$

We simplify Eq. (30) to

$$p_k(v_i, v_j | y_i \neq y_j) = \sum_{a=1}^{\frac{k+1}{2}} \left(\frac{n}{2}-1\right)^{k-1} \cdot \binom{k-1}{a-1} \cdot \left(\sum_{s=1}^{2a-1} \left(\frac{c_{in}}{n}\right)^{k-s} \cdot \left(\frac{c_{out}}{n}\right)^s \right)$$
$$+ \sum_{a=\frac{k+1}{2}}^{k} \left(\frac{n}{2}-1\right)^{k-1} \cdot \binom{k-1}{a-1} \cdot \left(\sum_{s=1}^{2(k-a)+1} \left(\frac{c_{in}}{n}\right)^{k-s} \cdot \left(\frac{c_{out}}{n}\right)^s \right)$$
$$= \frac{1}{n \cdot 2^{k-1}} \sum_{a=1}^{\frac{k+1}{2}} O\left(\binom{k-1}{a-1} \cdot \left(\sum_{s=1}^{2a-1} c_{in}^{k-s} \cdot c_{out}^s \right) \right) \tag{31}$$
$$+ \frac{1}{n \cdot 2^{k-1}} \sum_{a=\frac{k+1}{2}}^{k} O\left(\binom{k-1}{a-1} \cdot \left(\sum_{s=1}^{2(k-a)+1} c_{in}^{k-s} \cdot c_{out}^s \right) \right)$$
$$= \frac{(k-1)!}{n \cdot 2^{k-1}} \sum_{a=1}^{k} O\left(\sum_{s=1}^{\min(2a-1, 2(k-a)+1)} c_{in}^{k-s} \cdot c_{out}^s \right)$$

Case 3: $k = 1$
When $k = 1$, $A^k = A$,
$$\mathbb{E}[A_{ij}] = \begin{cases} \frac{c_{in}}{n}; \text{if} \quad y_i = y_j \\ \frac{c_{out}}{n}; \text{if} \quad y_i \neq y_j \end{cases}$$

In the following, we prove that when a graph is sparse and k is small, A_{ij}^k can be modeled with Poisson Distribution.

(1) when a sparse graph contains a large number of nodes $n \to \infty, d \ll n$, the potential k-length walk has a low probability of existing; (2) when $k \ll n$, the dependence between two different k-length walks is negligible; (3) the number of potential k-length walk is large $n^{k-1}, n \to \infty$. Thus, according to Lemma 26, the k-length walk connecting node v_i, v_j obeys the Poisson distribution $Poisson(\rho')$ when $k \geq 2$ where

$$
\rho' = \begin{cases} \rho_= = \frac{(k-1)!}{n \cdot 2^{k-1}} \sum_{a=2}^{k+1} O\left(\sum_{s=\min(2,2(a-2),2(k+1-a))}^{\min(2(a-1),2(k+1-a))} c_{in}^{k-s} \cdot c_{out}^s \right), & \text{if} \quad y_i = y_j \\ \rho_{\neq} = \frac{(k-1)!}{n \cdot 2^{k-1}} \sum_{a=1}^{k} O\left(\sum_{s=1}^{\min(2a-1,2(k-a)+1)} c_{in}^{k-s} \cdot c_{out}^s \right), & \text{if} \quad y_i \neq y_j \end{cases}
$$

When $k = 1$, $p(v_i, v_j)$ obeys the Bernoulli distribution $Ber(p)$ that

$$
p = \begin{cases} \frac{c_{in}}{n}, & \text{if} \quad y_i = y_j \\ \frac{c_{out}}{n}, & \text{if} \quad y_i \neq y_j \end{cases}
$$

$\square$

Theorem 32 (Expectation and variance of $(\tilde{A}^k XW)_{ij}$). *Given a graph generated by $G \sim cSBM(n, f, \mu, u, \lambda, d)$. The input node feature matrix is X and the normalized adjacency matrix is $\tilde{A}$. The k-th power matrix $\tilde{A}^k$ is applied to obtain a new feature matrix $\tilde{A}^k XW$, then the expectation and the variance of $(\tilde{A}^k XW)_{ij}$ are as follows:*

For $k = 1$:

$$
\mathbb{E}\left[(\tilde{A}^k XW)_{ij} \right] = \frac{1}{2d} \sqrt{\frac{\mu}{n}} (c_{in} - c_{out}) y_i u W_{:j}
$$

$$
\mathbb{V}\left[(\tilde{A}^k XW)_{ij} \right] = \frac{1}{2 \cdot d^2} \left(d - \frac{c_{in}^2 + c_{out}^2}{n} \right) \cdot \left(\frac{\mu}{n} (uW_{:j})^2 + \frac{||W_{:j}||_2^2}{f} \right)
$$

For $k \geq 2$:

$$
\mathbb{E}\left[(\tilde{A}^k XW)_{ij} \right] = \frac{(k-1)!}{d^k \cdot 2^{k-1}} O\left(c_{in}^k - c_{out}^k \right) \sqrt{\frac{\mu}{n}} y_i u W_{:j}
$$

$$
\mathbb{V}\left[(\tilde{A}^k XW)_{ij} \right] = \frac{(k-1)!}{d^{2k} \cdot 2^k} \left(\sum_{a=2}^{k+1} O\left(\sum_{s=\min(2,2(a-2),2(k+1-a))}^{\min(2(a-1),2(k+1-a))} c_{in}^{k-s} \cdot c_{out}^s \right) \right.
$$
$$
\left. + \sum_{a=1}^{k} O\left(\sum_{s=1}^{\min(2a-1,2(k-a)+1)} c_{in}^{k-s} \cdot c_{out}^s \right) \right) \left(\frac{\mu}{n} (uW_{:j})^2 + \frac{||W_{:j}||_2^2}{f} \right)
$$

Proof. Given that the node feature x_i for node v_i, generated by a conditional Stochastic Block Model (cSBM) conditioned on u and node class y_i, is distributed as:

$$
x_i \sim \mathcal{N}\left(\sqrt{\frac{\mu}{n}} y_i u, \frac{I_f}{f} \right)
$$

For a linear transformation matrix W, the transformed node feature is given by:

$$
x_i W \sim \mathcal{N}\left(\sqrt{\frac{\mu}{n}} y_i uW, \frac{W^T W}{f} \right)
$$

Feature after transformation with W and propagation with $\tilde{A}^k$ is

$$
\begin{aligned}
\left(\tilde{A}^k XW\right)_{ij} &= \sum_{r=1}^{n} \tilde{A}^k_{ir} (XW)_{rj} \\
&= \sum_{r=1}^{n} \tilde{A}^k_{ir} \left(\sqrt{\frac{\mu}{n}} y_r u W_{:j} + \frac{\epsilon_r W_{:j}}{\sqrt{f}} \right) \\
&= \sum_{r=1}^{n} \tilde{A}^k_{ir} \sqrt{\frac{\mu}{n}} \, y_r u W_{:j}
\end{aligned}
$$

and

$$
\mathbb{E}\left[\left(\tilde{A}^k XW\right)_{ij}\right] = \sqrt{\frac{\mu}{n}} \left(\sum_{r=1}^{n} \mathbb{E}\left[\tilde{A}^k_{ir}\right] y_r \right) u W_{:j} \tag{32}
$$

We now derive the expectation $\mathbb{E}[A^k_{ij}]$ of the adjacency matrix A raised to the power k.

1. Expectation $\mathbb{E}\left[\left(\tilde{A}^k XW\right)_{ij}\right]$ **when** $k \geq 2$

Two clusters generated by cSBM are in equal size. According to Theorem 31, we have

$$
\begin{aligned}
\mathbb{E}\left[\left(\tilde{A}^k XW\right)_{ij}\right] &= \sqrt{\frac{\mu}{n}} \left(\sum_{r=1}^{n} \mathbb{E}\left[\tilde{A}^k_{ir}\right] y_r \right) u W_{:j} \\
&= \frac{1}{d^k} \sqrt{\frac{\mu}{n}} \left(\sum_{r=1}^{n} \left(\mathbb{E}\left[A^k_{ir}|y_i = y_r\right] + \mathbb{E}\left[A^k_{ir}|y_i \neq y_r\right] \right) y_r \right) u W_{:j} \\
&= \frac{1}{d^k} \sqrt{\frac{\mu}{n}} \Bigg(\sum_{r=1}^{n} \bigg(\frac{(k-1)!}{n \cdot 2^{k-1}} \sum_{a=2}^{k+1} O\left(\sum_{s=\min(2,2(a-2),2(k+1-a))}^{\min(2(a-1),2(k+1-a))} c_{in}^{k-s} \cdot c_{out}^{s} \right) \\
&\qquad + \frac{(k-1)!}{n \cdot 2^{k-1}} \sum_{a=1}^{k} O\left(\sum_{s=1}^{\min(2a-1,2(k-a)+1)} c_{in}^{k-s} \cdot c_{out}^{s} \right) \bigg) y_r \Bigg) u W_{:j} \\
&= \frac{(k-1)!}{d^k \cdot 2^{k-1}} O\Bigg(\sum_{a=2}^{k+1} \sum_{s=\min(2,2(a-2),2(k+1-a))}^{\min(2(a-1),2(k+1-a))} c_{in}^{k-s} \cdot c_{out}^{s} \\
&\qquad - \sum_{a=1}^{k} \sum_{s=1}^{\min(2a-1,2(k-a)+1)} c_{in}^{k-s} \cdot c_{out}^{s} \Bigg) \sqrt{\frac{\mu}{n}} y_i u W_{:j} \\
&= \frac{(k-1)!}{d^k \cdot 2^{k-1}} O\left(c_{in}^{k} - c_{out}^{k} \right) \sqrt{\frac{\mu}{n}} y_i u W_{:j}
\end{aligned}
$$

2. Variance $\mathbb{E}\left[\left(\tilde{A}^k XW\right)_{ij}\right]$ **when** $k \geq 2$

The variance of new feature X'_{ij} given u, Y can be expressed as:

$$
\mathbb{V}\left[(\tilde{A}^k XW)_{ij}\right] = \mathbb{V}\left[\sum_{r=1}^{n} \tilde{A}_{ir}^k(\sqrt{\frac{\mu}{n}}y_r uW_{:j} + \frac{\epsilon_r W_{:j}}{\sqrt{f}})\right]
$$

$$
= \sum_{r=1}^{n} \mathbb{V}\left[\tilde{A}_{ir}^k(\sqrt{\frac{\mu}{n}}y_r uW_{:j} + \frac{\epsilon_r W_{:j}}{\sqrt{f}})\right], \quad \text{feature dimension independent}
$$

$$
= \sum_{r=1}^{n} \left[\mathbb{E}\left[(\tilde{A}^k)_{ir}^2\right]\mathbb{E}\left[(\sqrt{\frac{\mu}{n}}y_r uW_{:j} + \frac{\epsilon_r W_{:j}}{\sqrt{f}})^2\right] - \left(\mathbb{E}\left[\tilde{A}_{ir}^k\right]\right)^2\left(\mathbb{E}\left[\sqrt{\frac{\mu}{n}}y_r uW_{:j} + \frac{\epsilon_r W_{:j}}{\sqrt{f}}\right]\right)^2\right]
$$

$$
= \sum_{r=1}^{n} \left[\mathbb{E}\left[(\tilde{A}^k)_{ir}^2\right]\left(\left(\sqrt{\frac{\mu}{n}}y_r uW_{:j}\right)^2 + \frac{||W_{:j}||_2^2}{f}\right) - \left(\mathbb{E}\left[\tilde{A}_{ir}^k\right]\right)^2(\mathbb{E}\left[\sqrt{\frac{\mu}{n}}y_r uW_{:j} + \frac{\epsilon_r W_{:j}}{\sqrt{f}}\right])^2\right]
$$

$$
= \sum_{r=1}^{n} \left[\mathbb{E}\left[(\tilde{A}^k)_{ir}^2\right]\left(\left(\sqrt{\frac{\mu}{n}}y_r uW_{:j}\right)^2 + \frac{||W_{:j}||_2^2}{f}\right) - \left(\mathbb{E}\left[\tilde{A}_{ir}^k\right]\right)^2\left(\sqrt{\frac{\mu}{n}}y_r uW_{:j}\right)^2\right]
$$

$$
= \sum_{r=1}^{n} \left[\left(\left(\mathbb{E}\left[\tilde{A}_{ir}^k\right]\right)^2 + \mathbb{V}\left[\tilde{A}_{ir}^k\right]\right)\cdot\left(\left(\sqrt{\frac{\mu}{n}}y_r uW_{:j}\right)^2 + \frac{||W_{:j}||_2^2}{f}\right)\right.
$$

$$
\left. - \left(\mathbb{E}\left[\tilde{A}_{ir}^k\right]\right)^2\left(\sqrt{\frac{\mu}{n}}y_r uW_{:j}\right)^2\right]
$$

$$
= \frac{1}{d^{2k}}\sum_{r=1}^{n} \left[\left(\left(\mathbb{E}\left[A_{ir}^k\right]\right)^2 + \mathbb{V}\left[A_{ir}^k\right]\right)\cdot\left(\frac{\mu}{n}\left(uW_{:j}\right)^2 + \frac{||W_{:j}||_2^2}{f}\right)\right.
$$

$$
\left. - \left(\mathbb{E}\left[A_{ir}^k\right]\right)^2\frac{\mu}{n}\left(uW_{:j}\right)^2\right]
$$

$$
= \frac{1}{d^{2k}}\sum_{r=1}^{n} \left[\left(\mathbb{E}\left[A_{ir}^k\right]\right)^2\cdot\frac{||W_{:j}||_2^2}{f} + \mathbb{V}\left[A_{ir}^k\right]\cdot\left(\frac{\mu}{n}\left(uW_{:j}\right)^2 + \frac{||W_{:j}||_2^2}{f}\right)\right]
$$

$$
= \frac{1}{d^{2k}}\frac{n}{2}\left(\left(\mathbb{E}\left[A_{ir}^k|y_i = y_r\right]\right)^2 + \left(\mathbb{E}\left[A_{ir}^k|y_i \neq y_r\right]^2\right)\right)\cdot\frac{||W_{:j}||_2^2}{f}
$$

$$
+ \frac{1}{d^{2k}}\frac{n}{2}\left(\mathbb{V}\left[A_{ir}^k|y_i = y_r\right] + \mathbb{V}\left[A_{ir}^k|y_i \neq y_r\right]\right)\cdot\left(\frac{\mu}{n}\left(uW_{:j}\right)^2 + \frac{||W_{:j}||_2^2}{f}\right)
$$

$$
\tag{33}
$$

According to Theorem 31, when $k \geq 2$, we have

$$
\left(\mathbb{E}\left[A_{ij}^k|y_i = y_j\right]\right)^2 = \left(\frac{(k-1)!}{n\cdot 2^{k-1}}\sum_{a=2}^{k+1}O\left(\sum_{s=\min(2,2(a-2),2(k+1-a))}^{\min(2(a-1),2(k+1-a))}c_{in}^{k-s}\cdot c_{out}^s\right)\right)^2
$$

$$
\left(\mathbb{E}\left[A_{ij}^k|y_i \neq y_j\right]\right)^2 = \left(\frac{(k-1)!}{n\cdot 2^{k-1}}\sum_{a=1}^{k}O\left(\sum_{s=1}^{\min(2a-1,2(k-a)+1)}c_{in}^{k-s}\cdot c_{out}^s\right)\right)^2
$$

Two clusters generated by cSBM are in equal size. Then, Eq. (33) is written as:

$$
\mathbb{V}\left[(\tilde{A}^k XW)_{ij}\right] = \frac{1}{d^{2k}}\frac{n}{2}\left(\left(\mathbb{E}\left[A_{ir}^k|y_i=y_r\right]\right)^2 + \left(\mathbb{E}\left[A_{ir}^k|y_i\neq y_r\right]\right)^2\right)\cdot\frac{||W_{:j}||_2^2}{f}
$$

$$
+\frac{1}{d^{2k}}\frac{n}{2}\left(\mathbb{V}\left[A_{ir}^k|y_i=y_r\right] + \mathbb{V}\left[A_{ir}^k|y_i\neq y_r\right]\right)\cdot\left(\frac{\mu}{n}(uW_{:j})^2 + \frac{||W_{:j}||_2^2}{f}\right)
$$

$$
=\frac{((k-1)!)^2}{n\cdot d^{2k}\cdot 2^{2k-1}}O\Bigg(\left(\sum_{a=2}^{k+1}O\left(\sum_{s=\min(2,2(a-2),2(k+1-a))}^{\min(2(a-1),2(k+1-a))}c_{in}^{k-s}\cdot c_{out}^s\right)\right)^2
$$

$$
+\left(\sum_{a=1}^{k}O\left(\sum_{s=1}^{\min(2a-1,2(k-a)+1)}c_{in}^{k-s}\cdot c_{out}^s\right)\right)^2\Bigg)\cdot\frac{||W_{:j}||_2^2}{f}
$$

$$
+\frac{(k-1)!}{d^{2k}\cdot 2^k}\Bigg(\sum_{a=2}^{k+1}O\left(\sum_{s=\min(2,2(a-2),2(k+1-a))}^{\min(2(a-1),2(k+1-a))}c_{in}^{k-s}\cdot c_{out}^s\right)
$$

$$
+\sum_{a=1}^{k}O\left(\sum_{s=1}^{\min(2a-1,2(k-a)+1)}c_{in}^{k-s}\cdot c_{out}^s\right)\Bigg)\left(\frac{\mu}{n}(uW_{:j})^2 + \frac{||W_{:j}||_2^2}{f}\right)
$$

$$
=\frac{(k-1)!}{d^{2k}\cdot 2^k}\Bigg(\sum_{a=2}^{k+1}O\left(\sum_{s=\min(2,2(a-2),2(k+1-a))}^{\min(2(a-1),2(k+1-a))}c_{in}^{k-s}\cdot c_{out}^s\right)
$$

$$
+\sum_{a=1}^{k}O\left(\sum_{s=1}^{\min(2a-1,2(k-a)+1)}c_{in}^{k-s}\cdot c_{out}^s\right)\Bigg)\left(\frac{\mu}{n}(uW_{:j})^2 + \frac{||W_{:j}||_2^2}{f}\right),\quad n\to\infty
$$

3. Expectation and variance of $\left(\tilde{A}^k XW\right)_{ij}$ when $k=1$

$$
\mathbb{E}\left[(\tilde{A}XW)_{ij}\right] = \sqrt{\frac{\mu}{n}}\left(\sum_{r=1}^{n}\mathbb{E}\left[\tilde{A}_{ir}\right]y_r\right)uW_{:j}
$$

$$
=\frac{1}{d}\sqrt{\frac{\mu}{n}}\left(\sum_{r=1}^{n}\mathbb{E}\left[A_{ir}|y_i=y_r\right]y_i - \sum_{r=1}^{n}\mathbb{E}\left[A_{ir}|y_i\neq y_r\right]y_i\right)uW_{:j}
$$

$$
=\frac{1}{d}\sqrt{\frac{\mu}{n}}\left(\frac{n}{2}\frac{c_{in}}{n}y_i - \frac{n}{2}\frac{c_{out}}{n}y_i\right)uW_{:j}
$$

$$
=\frac{1}{2d}\sqrt{\frac{\mu}{n}}(c_{in}-c_{out})y_i uW_{:j}
$$

when $k=1$, we have

$$
\left(\mathbb{E}\left[A_{ij}^k|y_i=y_j\right]\right)^2 = \left(\frac{c_{in}}{n}\right)^2
$$

$$
\left(\mathbb{E}\left[A_{ij}^k|y_i\neq y_j\right]\right)^2 = \left(\frac{c_{out}}{n}\right)^2
$$

Eq. (33) is written as:

$$\mathbb{V}\left[(\tilde{A}XW)_{ij}\right] = \frac{1}{d^2}\frac{n}{2}\left(\left(\mathbb{E}\left[A_{ir}^k|y_i=y_r\right]\right)^2 + \left(\mathbb{E}\left[A_{ir}^k|y_i\neq y_r\right]\right)^2\right)\cdot\frac{||W_{:j}||_2^2}{f}$$

$$+ \frac{1}{d^2}\frac{n}{2}\left(\mathbb{V}\left[A_{ir}^k|y_i=y_r\right] + \mathbb{V}\left[A_{ir}^k|y_i\neq y_r\right]\right)\cdot\left(\frac{\mu}{n}\left(uW_{:j}\right)^2 + \frac{||W_{:j}||_2^2}{f}\right)$$

$$= \frac{1}{d^2}\frac{n}{2}\left(\left(\frac{c_{in}}{n}\right)^2 + \left(\frac{c_{out}}{n}\right)^2\right)\cdot\frac{||W_{:j}||_2^2}{f}$$

$$+ \frac{1}{d^2}\frac{n}{2}\left(\frac{c_{in}}{n}\left(1-\frac{c_{in}}{n}\right) + \frac{c_{out}}{n}\left(1-\frac{c_{out}}{n}\right)\right)\cdot\left(\frac{\mu}{n}\left(uW_{:j}\right)^2 + \frac{||W_{:j}||_2^2}{f}\right)$$

$$= \frac{1}{2n\cdot d^2}\left(c_{in}^2 + c_{out}^2\right)\cdot\frac{||W_{:j}||_2^2}{f}$$

$$+ \frac{1}{2\cdot d^2}\left(d - \frac{c_{in}^2 + c_{out}^2}{n}\right)\cdot\left(\frac{\mu}{n}\left(uW_{:j}\right)^2 + \frac{||W_{:j}||_2^2}{f}\right)$$

$$= \frac{1}{2\cdot d^2}\left(d - \frac{c_{in}^2 + c_{out}^2}{n}\right)\cdot\left(\frac{\mu}{n}\left(uW_{:j}\right)^2 + \frac{||W_{:j}||_2^2}{f}\right), \quad n\to\infty$$

$\square$

E.3 UNIFORM TRANSDUCTIVE STABILITY ON $G\sim cSBM(n,f,\mu,u,\lambda,d)$

We first give a lemma about the order of $\mathbb{E}\left[A_{ij}^k\right]$, which will be used in proof of Theorem 13.

Lemma 33 (order of $\mathbb{E}\left[A_{ij}^k\right]$). *The order of $\mathbb{E}\left[A_{ij}^k\right]$ is $O\left(\frac{k!\cdot d^k}{n\cdot 2^k}\right)$.*

Proof. According to Theorem 31, $A_{ij}^k|y_i=y_j$ and $A_{ij}^k|y_i\neq y_j$ obeys different Poisson distributions. As

$$c_{in}^{k-s}\cdot c_{out}^s = O\left(d^k\right),$$

then,

$$\rho_= = \frac{(k-1)!}{n\cdot 2^{k-1}}\sum_{a=2}^{k+1} O\left(\sum_{s=\min(2,2(a-2),2(k+1-a))}^{\min(2(a-1),2(k+1-a))} c_{in}^{k-s}\cdot c_{out}^s\right)$$

$$= \frac{(k-1)!}{n\cdot 2^{k-1}}\sum_{a=2}^{k+1} O\left(\sum_{s=\min(2,2(a-2),2(k+1-a))}^{\min(2(a-1),2(k+1-a))} d^k\right)$$

$$= \frac{(k-1)!}{n\cdot 2^{k-1}}\sum_{a=2}^{k+1} O\left(k\cdot d^k\right)$$

$$= \frac{(k-1)!}{n\cdot 2^{k-1}} O\left(k^2\cdot d^k\right)$$

$$= O\left(\frac{k!\cdot d^k}{n\cdot 2^k}\right)$$

similarly, we have $\rho_{\neq} = O\left(\frac{k!\cdot d^k}{n\cdot 2^k}\right)$

$\square$

According to Theorem 8, we prove Theorem 13 that a specific case that when graph $G\sim cSBM(n,f,\mu,u,\lambda,d)$.

Theorem 13. *Consider a spectral GNN Ψ parameterized by Θ, W trained using full-batch gradient descent for T iterations with a learning rate η on a training dataset containing m samples drawn from nodes on a graph $G\sim cSBM(n,f,\mu,u,\lambda,d)$. When $n\to\infty$, $k\ll n$, and $d\ll n$, under*

Assumptions 1, 2, and 4, for any node v_i on the graph, with probability at least $1 - \epsilon$ for a constant $\epsilon \in (0, 1)$, Ψ satisfies γ-uniform transductive stability, where $\gamma = r\beta$ and

$$\beta = \frac{1}{\epsilon} \left[O\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]\right) + O\left(\sum_{k=2}^K \left(\mathbb{E}\left[\left(A_{ij}^k \mid y_i = y_j\right)^2\right] + \mathbb{E}\left[\left(A_{ij}^k \mid y_i \neq y_j\right)^2\right]\right)\right)\right].$$

Proof. Any spectral GNNs in Eq. (1) with linear feature transformation function, and polynomial basis expanded on normalized graph matrix can be transformed into the format:

$$\hat{Y} = softmax(\sum_{k=0}^K \theta_k \tilde{A}^k XW) \tag{34}$$

where $\tilde{A} = D^{-\frac{1}{2}} A D^{-\frac{1}{2}}$ is the normalized graph adjacency matrix, D is the diagonal degree matrix. We denotes $Y \in \mathbb{R}^{n \times C}$ as the ground truth node label matrix.

When graph $G \sim cSBM(n, f, \mu, u, \lambda, d)$, the node feature

$$x_i \sim \mathcal{N}(y_i \sqrt{\mu/n}u, I_f/f)$$

Denote $B = XW$ and $S = BB^\top$, then we have

$$B_{ik} \sim \mathcal{N}(y_i \sqrt{\frac{\mu}{n}} uW_{:k}, \frac{\|W_{:k}\|_F^2}{f})$$

when $i \neq j$, B_{ik}, B_{jk} are independent, then

$$\mathbb{E}[S_{ij}] = \sum_{k=1}^C \mathbb{E}\left[B_{ik} B_{kj}^\top\right]$$

$$= \sum_{k=1}^C y_i y_j \frac{\mu}{n} (uW_{:k})^2$$

$$= y_i y_j \frac{\mu}{n} \|uW\|_F^2$$

when $i = j$:

$$\mathbb{E}[S_{ii}] = \frac{\mu}{n}\|uW\|_F^2 + \frac{\|W\|_F^2}{f}$$

When node number $n \to \infty$, we have

$$\sum_{q=1,q\neq j}^n \mathbb{E}[S_{jq}] = \frac{n}{2} y_j^2 \frac{\mu}{n}\|uW\|_F^2 + \frac{n}{2} y_j(-y_j) \frac{\mu}{n}\|uW\|_F^2 = 0$$

$$\sum_{j=1}^n \sum_{q=1,q\neq j}^n \mathbb{E}\left[A_{ij}^k A_{iq}^k\right] \mathbb{E}[S_{jq}]$$

$$= \frac{n^2}{4} \rho_{k=}^2 \frac{\mu}{n}\|uW\|_F^2; \quad (y_i = y_j = y_q)$$

$$+ \frac{n^2}{4} \rho_{k=}\rho_{k\neq} -\frac{\mu}{n}\|uW\|_F^2; \quad (y_i = y_j \neq y_q)$$

$$+ \frac{n^2}{4} \rho_{k\neq}\rho_{k=} -\frac{\mu}{n}\|uW\|_F^2; \quad (y_i \neq y_j = y_q) \tag{35}$$

$$+ \frac{n^2}{4} \rho_{k\neq}^2 \frac{\mu}{n}\|uW\|_F^2; \quad (y_i = y_q \neq y_j)$$

$$= \frac{n^2}{4} \cdot \frac{\mu}{n}\|uW\|_F^2 \cdot \left(\rho_{k=}^2 - 2\rho_{k\neq}\rho_{k=} + \rho_{k\neq}^2\right)$$

$$= \frac{n^2}{4} \cdot \frac{\mu}{n}\|uW\|_F^2 \cdot \left(\rho_{k=} - \rho_{k\neq}\right)^2$$

According to Theorem 31, when $k \geq 2$, $A_{ij}^k \sim Poisson(\rho_k')$, then

$$
\mathbb{E}\left[\|\tilde{A}_{i:}^k XW\|_F^2\right] = \mathbb{E}\left[\tilde{A}_{i:}^k XW (XW)^\top \left(\tilde{A}_{i:}^k\right)^\top\right]
$$

$$
= \mathbb{E}\left[\tilde{A}_{i:}^k S \left(\tilde{A}_{i:}^k\right)^\top\right]
$$

$$
= \mathbb{E}\left[\sum_{q=1}^n \sum_{j=1}^n \left(\tilde{A}_{ij}^k \tilde{A}_{iq}^k S_{jq}\right)\right]
$$

$$
= \frac{1}{d^{2k}} \mathbb{E}\left[\sum_{q=1}^n \sum_{j=1}^n \left(A_{ij}^k A_{iq}^k S_{jq}\right)\right]
$$

$$
= \frac{1}{d^{2k}} \sum_{q=1}^n \sum_{j=1}^n \mathbb{E}\left[A_{ij}^k A_{iq}^k\right] \mathbb{E}\left[S_{jq}\right]
$$

$$
= \frac{1}{d^{2k}} \sum_{j=1}^n \mathbb{E}\left[\left(A_{ij}^k\right)^2\right] \mathbb{E}\left[S_{jj}\right] + \frac{1}{d^{2k}} \sum_{j=1}^n \sum_{q=1,q\neq j}^n \mathbb{E}\left[A_{ij}^k A_{iq}^k\right] \mathbb{E}\left[S_{jq}\right]
$$

$$
= \frac{1}{d^{2k}} \frac{n}{2} \mathbb{E}\left[\left(A_{ij}^k\right)^2 \mid y_i = y_j\right] \mathbb{E}\left[S_{jj}\right] + \frac{1}{d^{2k}} \frac{n}{2} \mathbb{E}\left[\left(A_{ij}^k\right)^2 \mid y_i \neq y_j\right] \mathbb{E}\left[S_{jj}\right]
$$

$$
+ \frac{1}{d^{2k}} \frac{n^2}{4} \cdot \frac{\mu}{n} \|uW\|_F^2 \cdot (\rho_{k=} - \rho_{k\neq})^2 \quad (Eq.\ (35))
$$

$$
= \frac{1}{d^{2k}} \frac{n}{2} \left(\rho_{k=} + \rho_{k=}^2 + \rho_{k\neq} + \rho_{k\neq}^2\right) \left(\frac{\mu}{n} \|uW\|_F^2 + \frac{\|W\|_F^2}{f}\right)
$$

$$
+ \frac{1}{d^{2k}} \frac{n^2}{4} \cdot \frac{\mu}{n} \|uW\|_F^2 \cdot (\rho_{k=} - \rho_{k\neq})^2
$$

$$
= \frac{1}{2d^{2k}} \zeta_k \left(\mu \|uW\|_F^2 + \frac{n\|W\|_F^2}{f}\right) + \frac{n\mu}{4d^{2k}} \|uW\|_F^2 \cdot (\rho_{k=} - \rho_{k\neq})^2
$$

where $\zeta_k = \rho_{k=}^2 + \rho_{k=} + \rho_{k\neq}^2 + \rho_{k\neq}$

When $k = 1$, $A_{ij} \sim Ber(p)$, then

$$
\mathbb{E}\left[\|\tilde{A}_{i:}XW\|_F^2\right] = \frac{1}{d^2} \frac{n}{2} \left(p_=^2 + p_=(1 - p_=) + p_{\neq}^2 + p_{\neq}(1 - p_{\neq})\right) \left(\frac{\mu}{n} \|uW\|_F^2 + \frac{\|W\|_F^2}{f}\right)
$$

$$
= \frac{1}{d^2} \frac{n}{2} \left(p_= + p_{\neq}\right) \left(\frac{\mu}{n} \|uW\|_F^2 + \frac{\|W\|_F^2}{f}\right)
$$

$$
= \frac{1}{d^2} \frac{n}{2} \frac{2d}{n} \left(\frac{\mu}{n} \|uW\|_F^2 + \frac{\|W\|_F^2}{f}\right)
$$

$$
= \frac{1}{d} \left(\frac{\mu}{n} \|uW\|_F^2 + \frac{\|W\|_F^2}{f}\right)
$$

Substitute it into Eq. (15), we have

$$
\mathbb{E}\left[|\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial \theta_k}|\right] =
$$

$$
\begin{cases}
\frac{1}{2} \left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \left(\frac{\mu}{n} \|uW\|_F^2 + \frac{\|W\|_F^2}{f}\right)\right), & \text{if}\quad k = 0 \\
\frac{1}{2} \left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \frac{1}{d} \left(\frac{\mu}{n} \|uW\|_F^2 + \frac{\|W\|_F^2}{f}\right)\right), & \text{if}\quad k = 1 \\
\frac{1}{2} \left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \frac{1}{2d^{2k}} \zeta_k \left(\mu \|uW\|_F^2 + \frac{n\|W\|_F^2}{f}\right) + \frac{n\mu}{4d^{2k}} \|uW\|_F^2 \cdot (\rho_{k=} - \rho_{k\neq})^2\right), & \text{if}\quad k \geq 2
\end{cases}
$$

$$
(36)
$$

similarly, we have

$$\mathbb{E}\left[\|\tilde{A}_{i:}^k X\|_F^2\right] = \begin{cases} \frac{\mu}{n}\|u\|_F^2 + 1, & \text{if} \quad k = 0 \\ \frac{1}{d}\left(\frac{\mu}{n}\|u\|_F^2 + 1\right), & \text{if} \quad k = 1 \\ \frac{1}{2d^{2k}}\zeta_k\left(\mu\|u\|_F^2 + 1\right) + \frac{n\mu}{4d^{2k}}\|u\|_F^2 \cdot (\rho_{k=} - \rho_{k\neq})^2, & \text{if} \quad k \geq 2 \end{cases}$$

Substitute it into Eq. (16), we have

$$\mathbb{E}\left[\|\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial W}\|_{\ell_1}\right] = |\theta_0|\left(f \cdot \mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\left(\frac{\mu}{n}\|u\|_F^2 + 1\right)\right)$$

$$+ |\theta_1|\left(f \cdot \mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\frac{1}{d}\left(\frac{\mu}{n}\|u\|_F^2 + 1\right)\right)$$

$$+ \sum_{k=2}^{K} \frac{1}{d^{2k}}|\theta_k|\left(f \cdot \mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\frac{1}{2d^{2k}}\zeta_k\left(\mu\|u\|_F^2 + 1\right) + \frac{n\mu}{4d^{2k}}\|u\|_F^2 \cdot (\rho_{k=} - \rho_{k\neq})^2\right)$$

$$\tag{37}$$

Substitute Eq. (36), Eq. (37) into Eq. (12), we have

$$\mathbb{E}\left[\|\nabla \ell(\hat{y}_i, y_i; \Theta, W)\|_F\right] \leq \sum_{k=0}^{K} \mathbb{E}\left[\|\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial \theta_k}\|_{\ell_1}\right] + \mathbb{E}\left[\|\frac{\partial \ell(\hat{y}_i, y_i; \Theta, W)}{\partial W}\|_{\ell_1}\right]$$

$$= \frac{1}{2}\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \left(\frac{\mu}{n}\|uW\|_F^2 + \frac{\|W\|_F^2}{f}\right)\right)$$

$$+ \frac{1}{2}\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \frac{1}{d}\left(\frac{\mu}{n}\|uW\|_F^2 + \frac{\|W\|_F^2}{f}\right)\right)$$

$$+ \sum_{k=2}^{K} \frac{1}{2}\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \frac{1}{2d^{2k}}\zeta_k\left(\mu\|uW\|_F^2 + \frac{n\|W\|_F^2}{f}\right) + \frac{n\mu}{4d^{2k}}\|uW\|_F^2 \cdot \tilde{\zeta}_k^2\right) \tag{38}$$

$$+ |\theta_0|\left(f \cdot \mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\left(\frac{\mu}{n}\|u\|_F^2 + 1\right)\right)$$

$$+ |\theta_1|\left(f \cdot \mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\frac{1}{d}\left(\frac{\mu}{n}\|u\|_F^2 + 1\right)\right)$$

$$+ \sum_{k=2}^{K} \frac{1}{d^{2k}}|\theta_k|\left(f \cdot \mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\frac{1}{2d^{2k}}\zeta_k\left(\mu\|u\|_F^2 + 1\right) + \frac{n\mu}{4d^{2k}}\|u\|_F^2 \cdot \tilde{\zeta}_k^2\right)$$

where $\zeta_k = \rho_{k=}^2 + \rho_{k=} + \rho_{k\neq}^2 + \rho_{k\neq}, \tilde{\zeta}_k = \rho_{k=} - \rho_{k\neq}$.

According to Lemma 33, when $n \to \infty$, we have

$$n\left(\tilde{\zeta}_k\right)^2 = n\left(\rho_= - \rho_{\neq}\right)^2$$

$$= n\left(O\left(\frac{k! \cdot d^k}{n \cdot 2^k}\right)\right)^2$$

$$= nO\left(\frac{(k! \cdot d^k)^2}{n^2 \cdot 2^{2k}}\right)$$

$$= O\left(\frac{(k! \cdot d^k)^2}{n \cdot 2^{2k}}\right)$$

$$\to 0$$

Thus, $n\left(\tilde{\zeta}_k\right)^2$ can be neglected. Thus, we rewrite Eq. (38) as

$$
\begin{aligned}
\mathbb{E}\left[\|\nabla\ell(\hat{y}_i, y_i; \Theta, W)\|_F\right] &= \sum_{k=0}^{K} \mathbb{E}\left[\|\frac{\partial\ell(\hat{y}_i, y_i; \Theta, W)}{\partial\theta_k}\|_{\ell_1}\right] + \mathbb{E}\left[\|\frac{\partial\ell(\hat{y}_i, y_i; \Theta, W)}{\partial W}\|_{\ell_1}\right] \\
&= \frac{1}{2}\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \left(\frac{\mu}{n}\|uW\|_F^2 + \frac{\|W\|_F^2}{f}\right)\right) \\
&\quad + \frac{1}{2}\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \frac{1}{d}\left(\frac{\mu}{n}\|uW\|_F^2 + \frac{\|W\|_F^2}{f}\right)\right) \\
&\quad + \sum_{k=2}^{K}\frac{1}{2}\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \frac{1}{2d^{2k}}\zeta_k\left(\mu\|uW\|_F^2 + \frac{n\|W\|_F^2}{f}\right)\right) \\
&\quad + |\theta_0|\left(f\cdot\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\left(\frac{\mu}{n}\|u\|_F^2 + 1\right)\right) \\
&\quad + |\theta_1|\left(f\cdot\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\frac{1}{d^{2k-1}}\left(\frac{\mu}{n}\|u\|_F^2 + 1\right)\right) \\
&\quad + \sum_{k=2}^{K}\frac{1}{d^{2k}}|\theta_k|\left(f\cdot\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\frac{1}{2d^{2k}}\zeta_k\left(\mu\|u\|_F^2 + 1\right)\right) \\
&\leq \frac{1}{2}\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \left(\frac{\mu}{n}\|u\|_F^2 B_W^2 + \frac{B_W^2}{f}\right)\right) \\
&\quad + \frac{1}{2}\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \frac{1}{d}\left(\frac{\mu}{n}\|u\|_F^2 B_W^2 + \frac{B_W^2}{f}\right)\right) \\
&\quad + \sum_{k=2}^{K}\frac{1}{2}\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + \frac{1}{2d^{2k}}\zeta_k\left(\mu\|u\|_F^2 B_W^2 + \frac{nB_W^2}{f}\right)\right) \\
&\quad + B_\Theta\left(f\cdot\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\left(\frac{\mu}{n}\|u\|_F^2 + 1\right)\right) \\
&\quad + B_\Theta\left(f\cdot\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\frac{1}{d}\left(\frac{\mu}{n}\|u\|_F^2 + 1\right)\right) \\
&\quad + \sum_{k=2}^{K}\frac{1}{d^{2k}}B_\Theta\left(f\cdot\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] + C\frac{1}{2d^{2k}}\zeta_k\left(\mu\|u\|_F^2 + 1\right)\right) \\
&= \left(\frac{K+1}{2} + 2fB_\Theta + \sum_{k=2}^{K}\frac{f}{d^{2k}}B_\Theta\right)\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] \\
&\quad + \left(1 + \frac{1}{d}\right)\left(\left(\frac{B_W^2}{2} + CB_\Theta\right)\frac{\mu}{n}\|u\|_F^2 + \frac{B_W^2}{2f} + CB_\Theta\right) \\
&\quad + \sum_{k=2}^{K}\frac{\zeta_k}{d^{2k}}\left(\left(\mu\|u\|_F^2 + \frac{n}{f}\right)\frac{B_W^2}{4} + \left(\mu\|u\|_F^2 + 1\right)\frac{B_\Theta}{d^{2k}}\right)
\end{aligned}
\tag{39}
$$

We write it in big O format:

$$
\mathbb{E}\left[\|\nabla\ell(\hat{y}_i, y_i; \Theta, W)\|_F\right] = O\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]\right) + O\left(\sum_{k=2}^{K}\zeta_k\right)
$$

where $\zeta_k = \mathbb{E}\left[\left(A_{ij}^k \mid y_i = y_j\right)^2\right] + \mathbb{E}\left[\left(A_{ij}^k \mid y_i \neq y_j\right)^2\right]$

After get the upper bound of the norm of gradient, according to Theorem 6, we have the uniform transductive stability of spectral GNNs on graphs $G \sim cSBM(n, f, \mu, u, \lambda, d)$ of two classes $C = 2$ in big O format that

$$\gamma = r\beta; \beta = \frac{1}{\epsilon}\left[O\left(\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]\right) + O\left(\sum_{k=2}^{K}\left(\mathbb{E}\left[\left(A_{ij}^k \mid y_i = y_j\right)^2\right] + \mathbb{E}\left[\left(A_{ij}^k \mid y_i \neq y_j\right)^2\right]\right)\right)\right]$$

where r is the same as that in Theorem 6.

$\square$

F RELATION BETWEEN STABILITY OF SPECTRAL GNNS, NODE CLASS DISTRIBUTION AND ARCHITECTURE OF SPECTRAL GNNS

In this section, we fist derive the relation between parameter λ in cSBM and the edge homophilic ratio on graph. Then, we first study how the expected prediction error $\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]$ and ζ_k changes with λ, K. Next, we discuss how λ, K affect the uniform transductive stability and generalization of spectral GNNs.

Proposition 12. *For a graph $G \sim cSBM(n, \mu, u, \lambda, d)$, the expected edge homophily ratio is:*

$$\mathbb{E}[H_{edge}] = \frac{d + \lambda\sqrt{d}}{2d}; \quad \mathbb{E}[H_{edge}] = \frac{c_{in}}{c_{in} + c_{out}}. \tag{4}$$

Proof. As graphs generated with cSBM contains two clusters of same size. Thus, there are $\frac{n}{2}$ nodes having same class.

The expected number of edges between nodes of the same class is:

$$\mathbb{E}[E_{\text{same}}] = \binom{\frac{2}{n}}{2} \cdot \frac{c_{in}}{n} = \frac{c_{in}(n-2)}{8}$$

The expected number of edges between nodes of different class is:

$$\mathbb{E}[E_{\text{diff}}] = \frac{n}{2} \cdot \frac{n}{2} \cdot \frac{c_{out}}{n} \cdot \frac{1}{2} = \frac{c_{out}n}{8}$$

Thus, the expectation of H_{edge} is the expected number of edges between nodes with the same class to the total expected number of edges that

$$\begin{aligned}
\mathbb{E}\left[H_{edge}\right] &= \mathbb{E}[E_{\text{same}}] + \mathbb{E}[E_{\text{diff}}] \\
&= \frac{\frac{c_{in}(n-2)}{8}}{\frac{c_{in}(n-2)}{8} + \frac{c_{out}n}{8}} \\
&= \frac{(d + \lambda\sqrt{d})(n-2)}{(d + \lambda\sqrt{d})(n-2) + (d - \lambda\sqrt{d})n} \\
&= \frac{d + \lambda\sqrt{d}}{2d}, n \to \infty.
\end{aligned}$$

We also get the expectation between the H_{edge} and c_{in}, c_{out} that

$$\begin{aligned}
\mathbb{E}\left[H_{edge}\right] &= \mathbb{E}[E_{\text{same}}] + \mathbb{E}[E_{\text{diff}}] \\
&= \frac{\frac{c_{in}(n-2)}{8}}{\frac{c_{in}(n-2)}{8} + \frac{c_{out}n}{8}} \\
&= \frac{c_{in}(n-2)}{c_{in}(n-2) + c_{out}n} \\
&= \frac{c_{in}}{c_{in} + c_{out}}, n \to \infty.
\end{aligned}$$

$\square$

Theorem 14 ($\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]$ and λ, K). *Given a graph $G \sim cSBM(n, \mu, u, \lambda, d)$ and a spectral GNN of order K, $\mathbb{E}[\|\hat{y}_i - y_i\|_F^2]$ for any node v_i satisfies the following: it increases with $\lambda \in [-\sqrt{d}, 0]$, decreases with $\lambda \in [0, \sqrt{d}]$, and reaches its maximum at $\lambda = 0$; it increases with K if $\sum_{k=2}^{K} \theta_k \frac{(k-1)!}{2^{k-1}}$ grows more slowly than $\sum_{k=2}^{K} \theta_k^2 \frac{(k-1)!}{2^k}$ as K increases.*

Proof. Denote

$$Z = \sum_{k=0}^{K} \theta_k \tilde{A}^k X W; \qquad \hat{Y} = \text{softmax}(Z)$$

Then, for any node v_i with truth class y_i, we denote its prediction as

$$\hat{y}_i = \text{softmax}(Z_{i:})$$

For the binary classification $C = 2$, for a node with truth class $y_i = [1, 0]$, the predicted class

$$\hat{y}_i = [\hat{y}_1, \hat{y}_2] = \text{softmax}([Z_{i1}, Z_{i2}]) = [\sigma(Z_{i1} - Z_{i2}), 1 - \sigma(Z_{i1} - Z_{i2})]$$

where $\sigma(x) = \frac{1}{1+e^{-x}}$ is the sigmoid function.

We denote $z_i = Z_{i1} - Z_{i2}$, then

$$\hat{y}_i = [\sigma(z_i), 1 - \sigma(z_i)]$$

Thus,

$$\|\hat{y}_i - y_i\|_F^2 = (\sigma(z_i) - 1)^2 + (1 - \sigma(z_i))^2 = 2(1 - \sigma(z_i))^2$$

and

$$\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] = 2\mathbb{E}\left[(1 - \sigma(z_i))^2\right]$$

As node feature $x_i \sim \mathcal{N}(y_i \sqrt{\mu/n}u, I_f/f)$, any linear combination of Gaussian variable is still Gaussian variable, then

$$z_i \sim \mathcal{N}(\mu_{z_i}, \omega_{z_i}^2)$$

where

$$\mu_{z_i} = \mathbb{E}\left[z_i\right] = \mathbb{E}\left[Z_{i1} - Z_{i2}\right] = \mathbb{E}\left[Z_{i1}\right] - \mathbb{E}\left[Z_{i2}\right]$$

As $c_{in} = d + \lambda\sqrt{d}, c_{out} = d - \lambda\sqrt{d}, \lambda \in [-\sqrt{d}, \sqrt{d}]$, we have

$$c_{in}^k - c_{out}^k = O(d^k); \quad c_{in}^k = O(d^k); \quad c_{out}^k = O(d^k) \tag{40}$$

As $u \sim \mathcal{L}(0, I_f), d \ll f$ and Θ, W are bounded according to Assumption 4, we analyze dominant terms in μ_{z_i} and $\omega_{z_i}^2$.

According to Theorem 32, we have the expectation of $(\tilde{A}^k X W)_{ij}$, then

$$\begin{aligned}
\mu_{z_i} = \mathbb{E}\left[Z_{i1}\right] - \mathbb{E}\left[Z_{i2}\right] &= \theta_0 \sqrt{\frac{\mu}{n}} y_i u \left(W_{:1} - W_{:2}\right) \\
&+ \theta_1 \frac{1}{2d} \sqrt{\frac{\mu}{n}} \left(c_{in} - c_{out}\right) y_i u \left(W_{:1} - W_{:2}\right) \\
&+ \sum_{k=2}^{K} \theta_k \frac{(k-1)!}{d^k \cdot 2^{k-1}} O\left(c_{in}^k - c_{out}^k\right) \sqrt{\frac{\mu}{n}} y_i u \left(W_{:1} - W_{:2}\right) \\
&= O\left(\sum_{k=2}^{K} \theta_k \frac{(k-1)!}{2^{k-1}}\right) \quad (Eq.\ (40))
\end{aligned} \tag{41}$$

As $\tilde{A}^k, X$ are independent and columns of X are independent, $\left(\sum_{k=0}^{K} \theta_k \tilde{A}^k X\right)_{ij}, \left(\sum_{k=0}^{K} \theta_k \tilde{A}^k X\right)_{it}$ are independent. According to Theorem 32, we

have the variance of $(\tilde{A}^k XW)_{ij}$. Then we have

$$
\begin{aligned}
\omega_{z_i}^2 &= \mathbb{V}\left[Z_{i1} - Z_{i2}\right] \\
&= \mathbb{V}\left[\left(\sum_{k=0}^{K} \theta_k \tilde{A}^k X\right)_{i:}(W_{:1} - W_{:2})\right] \\
&= \mathbb{V}\left[\sum_{j=1}^{f}\left(\sum_{k=0}^{K} \theta_k \tilde{A}^k X\right)_{ij}(W_{j1} - W_{j2})\right] \\
&= \sum_{j=1}^{f}(W_{j1} - W_{j2})^2 \sum_{k=0}^{K} \theta_k^2 \mathbb{V}\left[\left(\tilde{A}^k X\right)_{ij}\right] \quad \text{(independent)} \\
&= \sum_{j=1}^{f}(W_{j1} - W_{j2})^2 \sum_{k=0}^{K} \theta_k^2\Bigg[\frac{1}{2 \cdot d^2}\left(d - \frac{c_{in}^2 + c_{out}^2}{n}\right) \cdot \left(\frac{\mu}{n}(uW_{:j})^2 + \frac{||W_{:j}||_2^2}{f}\right) \\
&\quad + \frac{(k-1)!}{d^{2k} \cdot 2^k}\Bigg(\sum_{a=2}^{k+1} O\Bigg(\sum_{s=\min(2,2(a-2),2(k+1-a))}^{\min(2(a-1),2(k+1-a))} c_{in}^{k-s} \cdot c_{out}^s\Bigg) \\
&\quad + \sum_{a=1}^{k} O\Bigg(\sum_{s=1}^{\min(2a-1,2(k-a)+1)} c_{in}^{k-s} \cdot c_{out}^s\Bigg)\Bigg)\left(\frac{\mu}{n}(uW_{:j})^2 + \frac{||W_{:j}||_2^2}{f}\right)\Bigg] \\
&= O\left(\sum_{k=2}^{K} \theta_k^2 \frac{(k-1)!}{2^k}\right) \quad (Eq.~(40))
\end{aligned}
\tag{42}
$$

1. $\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]$ **and** λ**.**
According to Lemma 29 and Lemma 30, we know that

- μ_{z_i} monotonously decreases and $\omega_{z_i}^2$ monotonously increases on $\lambda \in [-\sqrt{d}, 0]$;

- μ_{z_i} monotonously increases and $\omega_{z_i}^2$ monotonously decreases on $\lambda \in [0, \sqrt{d}]$;

- μ_{z_i} achieves the minimum value and $\omega_{z_i}^2$ achieves the maximum value when $\lambda = 0$.

$$
\mathbb{E}[(1 - \sigma(z_i))^2] = \int_{-\infty}^{\infty}(1 - \sigma(z_i))^2 \cdot \frac{1}{\sqrt{2\pi}\omega_{z_i}}e^{-\frac{(z - \mu_{z_i})^2}{2\omega_{z_i}^2}}\,dz_i
\tag{43}
$$

the integral decreases with μ_{z_i} and $\omega_{z_i}^2$, thus,

- $\mathbb{E}[(1 - \sigma(z_i))^2]$ increases on $\lambda \in [-\sqrt{d}, 0]$;
- $\mathbb{E}[(1 - \sigma(z_i))^2]$ decreases on $\lambda \in [0, \sqrt{d}]$;
- $\mathbb{E}[(1 - \sigma(z_i))^2]$ achieves the maximum value when $\lambda = 0$.

$\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]$ has the same trend with $\mathbb{E}\left[(1 - \sigma(z_i))^2\right]$.

2. $\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]$ **and** K**.**

we rewrite variable z that
$$
z = \mu_{z_i} + \omega_{z_i}y,
$$
where $y \sim \mathcal{N}(0, 1)$.

Thus, Eq. (43) is rewritten as

$$
\mathbb{E}\left[(1 - \sigma(z_i))^2\right] = \int_{-\infty}^{\infty}(1 - \sigma(\mu_{z_i} + \omega_{z_i}y))^2\frac{1}{\sqrt{2\pi}}e^{-\frac{y^2}{2}}\,dy
$$

(1) μ_{z_i} increases faster than $\omega_{z_i}^2$ when K increases

In this case, z will be dominated by μ_{z_i}, thus, we have

$$
\begin{aligned}
\mathbb{E}\left[(1 - \sigma(z))^2\right] &= \int_{-\infty}^{\infty} \left(1 - \sigma\left(\mu_{z_i}\right)\right)^2 \frac{1}{\sqrt{2\pi}} e^{-\frac{y^2}{2}} \, dy \\
&= \left(1 - \sigma\left(\mu_{z_i}\right)\right)^2 \\
&\leq 0.25
\end{aligned}
$$

(2) μ_{z_i} increases slower than $\omega_{z_i}^2$ when K increases

In this case, z will be dominated by $\omega_{z_i} y$, thus, we have

$$
\begin{aligned}
\mathbb{E}\left[(1 - \sigma(z))^2\right] &= \int_{-\infty}^{\infty} \left(1 - \sigma\left(\omega_{z_i} y\right)\right)^2 \frac{1}{\sqrt{2\pi}} e^{-\frac{y^2}{2}} \, dy \\
&= \int_{-\infty}^{0} (1 - 0) \frac{1}{\sqrt{2\pi}} e^{-\frac{y^2}{2}} \, dy + \int_{0}^{\infty} (1 - 1)^2 \frac{1}{\sqrt{2\pi}} e^{-\frac{y^2}{2}} \, dy \\
&= 0.5
\end{aligned}
$$

From above analysis, we know that when μ_{z_i} increases slower than $\omega_{z_i}^2$ when K increases, $\mathbb{E}\left[(1 - \sigma(z))^2\right]$ tends to have a larger value towards 0.5 compared with the case that μ_{z_i} increases faster than $\omega_{z_i}^2$ when K increases with corresponding values less or equal to 0.25.

In brief, when μ_{z_i} increases slower than $\omega_{z_i}^2$ when K increases, $\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]$ increases with K.

From Eq. (41), Eq. (42), we know the the dominant term of μ_{z_i} is $\sum_{k=2}^{K} \theta_k \frac{(k-1)!}{2^{k-1}}$, the dominant term of $\omega_{z_i}^2$ is $\sum_{k=2}^{K} \theta_k^2 \frac{(k-1)!}{2^k}$. Thus, $\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]$ increases with K if $\sum_{k=2}^{K} \theta_k \frac{(k-1)!}{2^{k-1}}$ grows slower than $\sum_{k=2}^{K} \theta_k^2 \frac{(k-1)!}{2^k}$.

$\square$

Theorem 15 (ζ_k and λ, K). *Given a graph $G \sim cSBM(n, \mu, u, \lambda, d)$ and a spectral GNN of order K, ζ_k has the following properties: (1) it increases with $\lambda \in [-\sqrt{d}, 0]$, decreases with $\lambda \in [0, \sqrt{d}]$, and achieves its maximum value at $\lambda = 0$; (2) it increases with k as k grows, for $k \in [0, K]$.*

Proof. The proof of this theorem is incorporated into the proof of Proposition 16. See the proof of Proposition 16 for details. $\square$

Proposition 16. *For a fixed K, γ-uniform transductive stability and generalization error bound strictly increase as λ moves from $-\sqrt{d}$ to 0, and decreases as λ moves from 0 to $\sqrt{d}$. For a fixed λ, if $\sum_{k=2}^{K} \theta_k \frac{(k-1)!}{2^{k-1}}$ grows more slowly than $\sum_{k=2}^{K} \theta_k^2 \frac{(k-1)!}{2^k}$ as K increases, then γ-uniform transductive stability and generalization error bound increase with K.*

Proof. According to Theorem 6 and Theorem 13, the uniform stability of spectral GNNs depends on the upper bound of the gradient norm β, and

$$
\begin{aligned}
\beta =& \left(\frac{K+1}{2} + 2f B_\Theta + \sum_{k=2}^{K} \frac{f}{d^{2k}} B_\Theta\right) \mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right] \\
&+ \left(1 + \frac{1}{d}\right) \left(\left(\frac{B_W^2}{2} + C B_\Theta\right) \frac{\mu}{n} \|u\|_F^2 + \frac{B_W^2}{2f} + C B_\Theta\right) \\
&+ \sum_{k=2}^{K} \frac{\zeta_k}{d^{2k}} \left(\left(\mu \|u\|_F^2 + \frac{n}{f}\right) \frac{B_W^2}{4} + \left(\mu \|u\|_F^2 + 1\right) \frac{B_\Theta}{d^{2k}}\right)
\end{aligned}
$$

where $\zeta_k = \rho_{\underline{=}}^2 + \rho_= + \rho_{\neq}^2 + \rho_{\neq}$, and $\rho_=$ and $\rho_{\neq}$ are the parameters of distribution in Theorem 31.

Denote

$$\psi_y = \left(\frac{K+1}{2} + 2fB_\Theta + \sum_{k=2}^{K} \frac{f}{d^{2k}} B_\Theta \right);$$

$$\psi_1 = \sum_{k=2}^{K} \frac{\zeta_k}{d^{2k}} \left(\left(\mu\|u\|_F^2 + \frac{n}{f} \right) \frac{B_W^2}{4} + \left(\mu\|u\|_F^2 + 1 \right) \frac{B_\Theta}{d^{2k}} \right).$$

We show that the terms $\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2 \right], \psi_y$, and ψ_1 can all be affected by λ, K.

(1) **Term** $\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2 \right]$

According to Theorem 14, the expected prediction error $\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2 \right]$ strictly increases with $\lambda \in [-\sqrt{d}, 0]$ and decreases with $\lambda \in [0, \sqrt{d}]$. In addition, it increases with K when $\sum_{k=2}^{K} \theta_k \frac{(k-1)!}{2^{k-1}}$ grows slower than $\sum_{k=2}^{K} \theta_k^2 \frac{(k-1)!}{2^k}$.

(2) **Term** ψ_y

As $\psi_y = \left(\frac{K+1}{2} + \sum_{k=0}^{K} |\theta_k| f \right)$ which does not contain λ, the class distribution has no effect on ψ_y. It also increases with order K.

(3) **Terms** ψ_1

ψ_1 is closely related with $\zeta_k = \rho_=^2 + \rho_= + \rho_{\neq}^2 + \rho_{\neq}$. According to Theorem 31, we have

$$\zeta_k = \rho_=^2 + \rho_= + \rho_{\neq}^2 + \rho_{\neq}$$

$$= \left(\frac{(k-1)!}{n \cdot 2^{k-1}} \sum_{a=2}^{k+1} O\left(\sum_{s=\min(2,2(a-2),2(k+1-a))}^{\min(2(a-1),2(k+1-a))} c_{in}^{k-s} \cdot c_{out}^s \right) \right)^2$$

$$+ \frac{(k-1)!}{n \cdot 2^{k-1}} \sum_{a=2}^{k+1} O\left(\sum_{s=\min(2,2(a-2),2(k+1-a))}^{\min(2(a-1),2(k+1-a))} c_{in}^{k-s} \cdot c_{out}^s \right)$$

$$+ \left(\frac{(k-1)!}{n \cdot 2^{k-1}} \sum_{a=1}^{k} O\left(\sum_{s=1}^{\min(2a-1,2(k-a)+1)} c_{in}^{k-s} \cdot c_{out}^s \right) \right)^2$$

$$+ \frac{(k-1)!}{n \cdot 2^{k-1}} \sum_{a=1}^{k} O\left(\sum_{s=1}^{\min(2a-1,2(k-a)+1)} c_{in}^{k-s} \cdot c_{out}^s \right).$$

As $c_{in} = d + \lambda\sqrt{d}$ and $c_{out} = d - \lambda\sqrt{d}$, all four terms in ρ are in the form of $g(\lambda) = \sum_{s=1}^{k} \left(d + \lambda\sqrt{d} \right)^{k-s} \cdot \left(d - \lambda\sqrt{d} \right)^s$. According to Lemma 30, all functions in the form of $g(\lambda)$ strictly increase on $\lambda \in [-\sqrt{d}, 0]$ and decreases on $\lambda \in [0, \sqrt{d}]$. Since all the other elements in ψ_1 except ζ_k are positive, ψ_1 strictly increases on $\lambda \in [-\sqrt{d}, 0]$ and decreases on $\lambda \in [0, \sqrt{d}]$.

When k increases, ζ_k contains more items and thus ψ_1 increases with order K.

According to Proposition 12, we have

$$\lambda \in [0, \sqrt{d}] \Leftrightarrow H_{edge} \in [0.5, 1] \text{ and } \lambda \in [-\sqrt{d}, 0] \Leftrightarrow H_{edge} \in [0, 0.5].$$

According to Theorem 9, any factors affecting γ affect the generalization error bound. Thus, we conclude the following cases:

(a) uniform transductive stability γ, generalization error bound and λ

From the above analysis, we know that ϕ_y is not affected by λ, and terms $\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2 \right]$, ψ_1, and ψ_2 strictly increase on $\lambda \in [-\sqrt{d}, 0]$ and decrease on $\lambda \in [0, \sqrt{d}]$. This shows that

the stability decreases and the generalization error bound increases when $H_{edge} \in (0, 0.5]$. The stability increases and the generalization error bound decreases when $H_{edge} \in [0, 5, 1)$. Spectral GNNs are stable and generalize well on strong homophilic and heterophilic graphs.

(b) uniform transductive stability γ, generalization error bound, and K

From the above analysis, we know that terms ϕ_y, ψ_1, ψ_2 increase with k. Thus, when $\sum_{k=2}^{K} \theta_k \frac{(k-1)!}{2^{k-1}}$ grows slower than $\sum_{k=2}^{K} \theta_k^2 \frac{(k-1)!}{2^k}$, the expected prediction error $\mathbb{E}\left[\|\hat{y}_i - y_i\|_F^2\right]$ increases with K and the uniform transductive stability γ also increases with K. Under this condition, the stability becomes worse and generalization error bound thus increases when K increase.

$\square$

G DETAILS OF EXPERIMENTS

G.1 DATASETS

The statistical properties of real-world datasets, including the number of nodes, edges, feature dimensions, node classes, and edge homophily ratios, are summarized in Table 2 and Table 3. We use the directed and cleaned versions of the Chameleon and Squirrel datasets provided by (Platonov et al., 2023), where repeated nodes have been removed.

Statistics	Texas	Wisconsin	Cornell	Actor	Chameleon	Squirrel	Citeseer	Pubmed	Cora
# Nodes	183	251	183	7,600	890	2,223	3,327	19,717	2,708
# Edges	295	466	295	26,752	27,168	131,436	4,676	44,327	5,278
# Features	1,703	1,703	1,703	932	2,325	2,089	3,703	500	1,433
# Classes	5	5	5	5	5	6	5	7	
Edge Homophily	0.11	0.21	0.22	0.24	0.22	0.74	0.8	0.81	

Table 2: Statistics of real-world datasets.

Statistics	OGBN-Arxiv	OGBN-Products
# Nodes	169,343	2,449,029
# Edges	2,315,598	61,859,140
# Features	128	100
# Classes	40	47
Edge Homophily	0.65	0.81

Table 3: Statistics of OGBN datasets.

G.2 SPECTRAL GNNS

We detail the spectral GNNs used in our experiments below. For a graph with adjacency matrix A, degree matrix D, and identity matrix I, we define the following matrices: the normalized Laplacian matrix $\hat{L} = I - D^{-1/2}AD^{-1/2}$, the shifted normalized Laplacian matrix $\tilde{L} = -D^{-1/2}AD^{-1/2}$, the normalized adjacency matrix $\tilde{A} = D^{-1/2}AD^{-1/2}$, and the normalized adjacency matrix with self-loops $\tilde{A}' = (D + I)^{-1/2}(A + I)(D + I)^{-1/2}$.

ChebNet (Defferrard et al., 2016): This model uses the Chebyshev basis to approximate a spectral filter:

$$\hat{Y} = \sum_{k=0}^{K} \theta_k T_k(\tilde{L}) f_W(X)$$

where X is the raw feature matrix, $\Theta = [\theta_0, \theta_1, \ldots, \theta_K]$ is the graph convolution parameter, W is the feature transformation parameter and $f_W(X)$ is usually a 2-layer MLP. $T_k(\tilde{L})$ is the k-th Chebyshev

basis expanded on the shifted normalized graph Laplacian matrix $\tilde{L}$ and is recursively calculated:

$$T_0(\tilde{L}) = I$$
$$T_1(\tilde{L}) = \tilde{L}$$
$$T_k(\tilde{L}) = 2\tilde{L}T_{k-1}(\tilde{L}) - T_{k-2}(\tilde{L})$$

ChebNetII (He et al., 2022): The model is formulated as

$$\hat{Y} = \frac{2}{K+2}\sum_{k=0}^{K}\sum_{j=0}^{K}\theta_j T_k(x_j)T_k(\tilde{L})f_W(X),$$

where X is the input feature matrix, W is the feature transformation parameter, $f_W(X)$ is usually a 2-layer MLP, $T_k(\cdot)$ is the k-th Chebyshev basis expanded on $\cdot$, $x_j = \cos\left((j+1/2)\,\pi/\,(K+1)\right)$ is the j-th Chebyshev node, which is the root of the Chebyshev polynomials of the first kind with degree $K+1$, and θ_j is a learnable parameter. Graph convolution parameter in ChebNet is reparameterized with Chebyshev nodes and learnable parameters θ_j.

JacobiConv (Wang & Zhang, 2022): This model uses the Jacobi basis to approximate a filter as:

$$\hat{Y} = \sum_{k=0}^{K}\theta_k T_k^{a,b}(\tilde{A})f_W(X),$$

where X is the input feature matrix, $\Theta = [\theta_0, \theta_1, \ldots, \theta_K]$ is the graph convolution parameter, W is the feature transformation parameter and $f_W(X)$ is usually a 2-layer MLP. $T_k^{a,b}(\tilde{A})$ is the Jacobi basis on normalized graph adjacency matrix $\tilde{A}$ and is recursively calculated as

$$T_k^{a,b}(\tilde{A}) = I$$
$$T_k^{a,b}(\tilde{A}) = \frac{1-b}{2}I + \frac{a+b+2}{2}\tilde{A}$$
$$T_k^{a,b}(\tilde{A}) = \gamma_k \tilde{A}T_{k-1}^{a,b}(\tilde{A}) + \gamma_k' T_{k-1}^{a,b}(\tilde{A}) + \gamma_k'' T_{k-2}^{a,b}(\tilde{A})$$

where $\gamma_k = \frac{(2k+a+b)(2k+a+b-1)}{2k(k+a+b)}, \gamma_k' = \frac{(2k+a+b-1)(a^2-b^2)}{2k(k+a+b)(2k+a+b-2)}, \gamma_k'' = \frac{(k+1-1)(k+b-1)(2k+a+b)}{k(k+a+b)(2k+a+b-2)}$. a and b are hyper-parameters. Usually, grid search is used to find the optimal a and b values.

GPRGNN (Chien et al., 2021): This model uses the monomial basis to approximate a filter:

$$\hat{Y} = \sum_{k=0}^{K}\theta_k \tilde{A}'^k f_W(X)$$

where X is the input feature matrix, $\Theta = [\theta_0, \theta_1, \ldots, \theta_K]$ is the graph convolution parameter, W is the feature transformation parameter and $f_W(X)$ is usually a 2-layer MLP. $\tilde{A}'$ is the normalized adjacency matrix with self-loops.

BernNet (He et al., 2021): This model uses the Bernstein basis for approximation:

$$\hat{Y} = \sum_{k=0}^{K}\theta_k \frac{1}{2^K}\binom{K}{k}(2I - \hat{L})^{K-k}\hat{L}^k f_W(X)$$

where X is the input feature matrix, $\Theta = [\theta_0, \theta_1, \ldots, \theta_K]$ is the graph convolution parameter, W is the feature transformation parameter and $f_W(X)$ is usually a 2-layer MLP. $\hat{L}$ is the normalized Laplacian matrix.

G.3 HYPER-PARAMETER SETTINGS

All experiments were conducted on an NVIDIA RTX A6000 GPU with 48GB of memory.

We employ a two-layer Multi-Layer Perceptron (MLP) with a hidden layer size of 64 for the feature transformation function f_W, using ReLU as the activation function across all spectral GNN models.

Following (Tang & Liu, 2023a; Cong et al., 2021), the dropout rate and weight decay are set to 0.0. The Adam optimizer is used for optimization. Each experiment runs for a maximum of 300 iterations and is repeated 10 times to report the mean and variance of the results. A grid search is conducted to determine the best learning rate from $\{0.05, 0.01, 0.001\}$.

G.4 DETAILED EXPERIMENTAL RESULTS

H_{edge}	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9
ChebNet	94.92±0.24	86.08±0.43	81.09±0.63	75.11±0.73	72.69±0.66	74.66±0.65	79.62±0.78	86.03±0.6	94.64±0.39
Acc Gap	5.08±0.24	13.92±0.41	18.91±0.57	24.89±0.72	27.3±0.62	25.34±0.68	20.38±0.74	13.97±0.61	5.36±0.41
Loss Gap	0.64±0.07	3.15±0.14	3.72±0.2	5.42±0.24	5.88±0.5	6.01±0.27	4.62±0.3	3.04±0.18	0.98±0.06
ChebNetII	92.19±0.51	85.03±0.58	79.83±0.43	77.55±0.64	77.34±0.54	77.7±0.57	78.22±0.73	83.68±0.41	91.43±0.48
Acc Gap	7.81±0.47	14.97±0.58	20.17±0.41	22.45±0.66	22.66±0.49	22.3±0.57	21.77±0.71	16.32±0.44	8.57±0.47
Loss Gap	0.66±0.07	1.84±0.11	3.55±0.21	4.77±0.26	4.86±0.13	4.64±0.21	4.23±0.33	2.14±0.17	0.72±0.05
JacobiConv	89.25±3.35	77.23±4.51	77.19±0.66	77.0±0.55	79.06±0.61	80.2±0.57	84.64±0.39	90.48±0.24	96.91±0.24
Acc Gap	10.71±2.86	22.73±4.36	22.8±0.67	23.0±0.54	20.94±0.61	19.8±0.6	15.36±0.41	9.51±0.24	3.09±0.25
Loss Gap	0.69±0.26	1.58±0.45	4.08±0.21	4.33±0.14	5.36±0.33	1.95±0.13	1.58±0.13	0.99±0.06	0.16±0.01
GPRGNN	90.33±0.57	87.06±0.64	81.71±0.41	77.03±0.47	77.23±0.65	79.52±0.59	82.72±0.52	89.25±0.5	96.45±0.18
Acc Gap	9.66±0.54	12.94±0.67	18.29±0.42	22.96±0.49	22.77±0.64	20.48±0.6	17.27±0.52	10.75±0.54	3.55±0.2
Loss Gap	1.42±0.08	2.21±0.14	3.27±0.2	4.72±0.19	5.17±0.13	4.7±0.25	3.7±0.47	2.4±0.32	1.05±0.11
BernNet	87.44±0.5	82.92±0.67	79.3±0.44	77.69±0.53	77.97±0.54	77.49±0.72	76.58±0.79	79.73±1.3	85.68±1.05
Acc Gap	12.55±0.5	17.08±0.76	20.7±0.44	22.31±0.54	22.03±0.55	22.51±0.64	23.41±0.8	20.27±1.39	14.32±1.06
Loss Gap	1.2±0.06	2.45±0.21	3.69±0.16	4.77±0.24	4.72±0.15	4.7±0.17	4.35±0.35	2.92±0.31	1.36±0.14

Table 4: Testing accuracy, accuracy gap, loss gap of spectral GNNs on synthetic datasets with edge homophilic ratio $H_{edge} \in [0.1, 0.9]$. Small accuracy and loss gaps imply good generalization capability.

Datasets	Texas	Wisconsin	Actor	Squirrel	Chameleon	Cornell	Citeseer	Pubmed	Cora
ChebNet	40.82±7.25	52.23±3.77	26.63±0.53	30.08±1.14	33.94±1.58	44.88±6.19	64.16±0.82	84.74±0.37	74.95±0.96
Acc Gap	59.18±6.94	47.77±3.92	73.26±0.54	69.92±1.28	66.06±1.52	55.12±5.95	35.82±0.75	15.25±0.37	25.05±0.92
Loss Gap	5.91±0.66	5.77±0.87	21.64±0.8	35.68±2.33	36.17±3.04	6.57±0.82	4.68±0.22	1.44±0.06	3.9±0.29
ChebNetII	77.55±5.71	74.38±3.08	27.94±0.36	28.1±1.82	38.45±1.63	73.69±5.12	65.85±0.52	84.7±0.3	74.0±0.8
Acc Gap	22.45±5.2	25.62±3.31	71.94±0.33	71.83±1.77	61.47±1.53	26.31±5.0	34.12±0.48	15.16±0.28	26.0±0.75
Loss Gap	1.1±0.27	1.39±0.32	20.16±0.76	27.56±2.88	19.33±1.68	1.7±0.3	2.66±0.09	1.13±0.09	2.14±0.09
JacobiConv	78.06±5.31	77.62±2.92	27.89±0.63	26.78±1.28	32.2±2.08	80.41±3.98	73.56±0.64	86.33±0.47	84.31±0.49
Acc Gap	21.94±5.41	22.38±2.85	71.97±0.66	50.85±11.88	63.82±9.46	19.59±4.18	26.41±0.65	10.87±1.45	15.69±0.5
Loss Gap	0.94±0.26	1.19±0.22	31.67±0.86	32.75±11.57	38.77±7.16	0.91±0.16	2.16±0.06	0.51±0.14	1.28±0.09
GPRGNN	46.84±6.22	72.08±3.23	26.29±0.65	29.91±1.19	34.28±1.58	61.33±6.12	72.89±0.62	85.42±0.4	84.37±0.51
Acc Gap	53.16±6.12	27.92±2.92	71.52±4.82	70.09±1.09	65.72±1.69	38.67±6.43	27.08±0.67	14.58±0.37	15.63±0.54
Loss Gap	3.35±0.83	1.6±0.31	29.22±2.69	35.34±5.58	29.88±2.22	2.2±0.53	3.32±0.16	1.24±0.09	1.54±0.1
BernNet	75.92±5.31	81.85±2.23	27.28±0.76	33.42±1.14	33.72±1.38	81.43±3.46	67.17±0.59	84.82±0.25	73.39±0.87
Acc Gap	24.08±5.41	18.15±2.16	72.61±0.71	66.58±1.11	66.28±1.33	18.57±3.57	32.8±0.57	14.95±0.45	26.61±0.87
Loss Gap	1.24±0.31	0.87±0.26	24.68±0.71	28.17±1.47	27.83±1.75	1.06±0.18	2.66±0.09	1.13±0.13	2.18±0.08

Table 5: Testing accuracy, accuracy gap, loss gap of spectral GNNs on real world datasets with edge homophilic ratio $H_{edge} \in [0.11, 0.81]$. Small accuracy and loss gaps imply good generalization capability.

Order K	1	2	3	4	5	6	7	8	9	10
ChebNet	87.31±0.3	89.11±0.31	88.48±0.49	84.19±0.9	71.3±3.0	79.58±0.52	80.77±0.62	76.21±0.51	82.94±0.48	86.08±0.41
Acc Gap	12.7±0.32	10.89±0.31	11.52±0.5	15.8±0.92	28.7±3.54	20.42±0.51	19.23±0.57	23.79±0.47	17.06±0.45	13.92±0.42
Loss Gap	2.2±0.09	1.76±0.07	1.9±0.14	2.84±0.27	7.2±1.45	3.88±0.2	3.08±0.21	3.79±0.26	3.8±0.11	3.15±0.14
ChebNetII	85.92±0.56	80.1±0.99	82.65±0.7	85.56±0.45	84.64±0.8	84.62±0.59	85.27±0.51	86.2±0.64	86.39±0.5	85.03±0.57
Acc Gap	14.07±0.53	19.9±1.02	17.35±0.73	14.44±0.45	15.36±0.87	15.38±0.6	14.73±0.5	13.79±0.6	13.61±0.49	14.97±0.58
Loss Gap	1.94±0.08	3.23±0.31	2.62±0.14	2.06±0.14	1.94±0.21	1.95±0.17	1.99±0.15	1.75±0.14	1.83±0.11	1.84±0.11
JacobiConv	77.44±0.67	80.51±0.48	49.44±1.12	39.85±1.91	48.81±2.65	47.73±7.63	60.29±7.48	67.53±7.95	68.03±9.15	77.23±4.79
Acc Gap	22.55±0.62	19.49±0.46	50.56±1.18	60.13±1.98	51.19±2.63	52.25±7.08	39.7±7.32	32.45±7.76	31.96±9.19	22.73±4.82
Loss Gap	5.72±0.19	5.8±0.26	8.81±0.79	12.63±1.22	7.3±1.01	8.23±1.77	4.98±1.23	3.42±1.39	3.33±1.32	1.58±0.48
GPRGNN	83.61±0.66	86.14±0.29	79.44±1.05	88.36±0.28	87.25±0.5	88.0±0.39	87.57±0.47	87.5±0.3	87.17±0.3	87.06±0.59
Acc Gap	16.39±0.69	13.86±0.29	20.56±1.06	11.63±0.29	12.76±0.49	12.01±0.32	12.43±0.48	12.49±0.33	12.84±0.29	12.94±0.68
Loss Gap	2.37±0.11	2.21±0.1	3.18±0.19	1.83±0.1	2.14±0.2	1.93±0.09	2.06±0.13	2.12±0.09	2.19±0.13	2.21±0.14
BernNet	82.76±0.72	81.14±0.41	81.21±0.57	81.47±0.6	81.77±0.66	82.11±0.75	82.32±0.88	82.55±0.84	82.8±0.81	82.92±0.79
Acc Gap	17.24±0.71	18.86±0.39	18.79±0.56	18.53±0.7	18.23±0.62	17.89±0.85	17.68±0.84	17.45±0.79	17.2±0.79	17.08±0.7
Loss Gap	2.45±0.17	3.02±0.11	2.95±0.21	2.84±0.2	2.75±0.21	2.65±0.21	2.59±0.22	2.54±0.2	2.49±0.21	2.45±0.21

Table 6: Testing accuracy, accuracy gap, loss gap of spectral GNNs on synthetic dataset of edge homophilic ratio $H_{edge} = 0.2$ when $K \in [1, 10]$. Small accuracy and loss gaps imply good generalization capability.

Order K	1	2	3	4	5	6	7	8	9	10
ChebNet	83.78±2.45	80.61±4.59	80.51±3.47	61.73±5.0	63.37±8.57	36.33±5.72	44.18±5.0	24.39±2.14	30.2±4.8	40.82±7.35
Acc Gap	16.22±2.45	19.39±4.8	19.49±3.78	38.27±5.0	36.63±7.86	63.67±6.12	55.82±5.0	75.61±2.24	69.8±5.0	59.18±7.15
Loss Gap	1.49±0.44	1.26±0.44	1.48±0.31	2.77±0.53	3.08±0.59	8.98±0.68	6.09±0.72	7.99±0.93	9.0±1.03	5.91±0.69
ChebNetII	80.41±3.98	75.41±5.72	76.53±4.29	76.53±4.59	76.94±5.0	78.78±5.61	78.88±5.2	77.45±4.9	76.94±5.72	77.55±5.51
Acc Gap	19.59±3.78	24.59±5.2	23.47±4.59	23.47±4.49	23.06±4.8	21.22±5.61	21.12±5.82	22.55±4.49	23.06±5.61	22.45±5.31
Loss Gap	0.74±0.14	1.2±0.44	1.15±0.29	1.28±0.3	1.23±0.33	1.11±0.29	1.16±0.26	1.21±0.29	1.24±0.27	1.1±0.27
JacobiConv	52.24±5.41	80.92±3.78	75.31±5.31	74.39±3.78	79.08±3.67	78.67±4.08	80.0±3.06	73.67±6.33	77.65±5.41	78.06±5.61
Acc Gap	47.76±5.31	19.08±3.98	24.69±5.0	25.61±3.67	20.92±3.47	21.33±3.67	20.0±3.06	26.33±6.84	22.35±5.1	21.94±5.41
Loss Gap	2.54±0.42	0.89±0.2	1.1±0.25	1.18±0.27	0.9±0.17	0.97±0.16	0.93±0.13	1.22±0.39	0.97±0.26	0.94±0.24
GPRGNN	53.88±4.8	49.18±5.1	46.73±5.82	45.82±6.64	46.12±5.41	45.61±5.2	46.43±4.59	46.12±5.0	47.55±4.8	46.84±6.22
Acc Gap	46.12±4.9	50.82±5.31	53.27±5.61	54.18±6.63	53.88±5.72	54.39±5.2	53.57±4.9	53.88±4.9	52.45±5.1	53.16±6.43
Loss Gap	2.6±0.44	3.21±0.53	3.5±0.67	3.6±0.63	3.58±0.63	3.51±0.64	3.47±0.48	3.44±0.61	3.22±0.73	3.35±0.83
BernNet	76.73±3.67	75.92±2.45	75.61±3.67	77.04±3.88	77.14±4.39	75.2±4.7	74.9±5.72	75.2±5.2	74.8±5.92	75.71±5.71
Acc Gap	23.27±3.67	24.08±2.65	24.39±3.57	22.96±3.98	22.86±4.29	24.8±4.69	25.1±5.2	24.8±5.61	25.2±6.02	24.29±5.61
Loss Gap	0.96±0.22	0.95±0.18	1.01±0.17	1.02±0.21	1.06±0.21	1.13±0.25	1.19±0.31	1.18±0.26	1.27±0.34	1.25±0.31

Table 7: Testing accuracy, accuracy gap, loss gap of spectral GNNs on Texas dataset of edge homophilic ratio $H_{edge} = 0.11$ when $K \in [1, 10]$. Small accuracy and loss gaps imply good generalization capability.