A Consolidated Cross-Validation Algorithm for Support Vector Machines via Data Reduction

Abstract

We propose a consolidated cross-validation (CV) algorithm for training and tuning the support vector machines (SVM) on reproducing kernel Hilbert spaces. Our consolidated CV algorithm utilizes a recently proposed exact leave-one-out formula for the SVM and accelerates the SVM computation via a data reduction strategy. In addition, to compute the SVM with the bias term (intercept), which is not handled by the existing data reduction methods, we propose a novel two-stage consolidated CV algorithm. With numerical studies, we demonstrate that our algorithm is about an order of magnitude faster than the two mainstream SVM solvers, kernlab and LIBSVM, with almost the same accuracy.

1 Introduction

This paper concerns one of the most successful classifiers, the kernel support vector machine (SVM) (Cortes and Vapnik, 1995; Vapnik, 1995, 1998), which has been popularly used on structured data in the past two decades. The success of the SVM is mainly attributed to its appealing geometric interpretation, solid theoretical foundation, and high predictive power. To assess the predictive accuracy of the SVM, cross-validation (CV)(Wahba and Wold, 1975; Arlot and Celisse, 2010) is perhaps the most commonly used method in practice. In a K-fold CV procedure, the training data is randomly split into K equal-sized groups. Based on data splitting, part of the data is used for training each competing model and the rest of the data is reserved for evaluating the prediction error. The model with the smallest CV error is finally elected. Typical choices of K are 5, 10, or n (sample size), where $K = n$ yields the so-called leave-one-out cross-validation (LOOCV).

LOOCV is generally less used than ten-fold and five-fold CV, largely because of the two popular arguments: (1) high computational cost of LOOCV; (2) much larger variance than five-fold or ten-fold CV. We must point out that while the first argument is true in some sense, the second argument is actually a wide-spreading misconception about LOOCV. For instance, Kohavi (1995) and Hastie et al. (2009) argue that leave-one-out is almost unbiased, but it has high variance, leading to unreliable estimates. The argument, however, is not generally true. A series of revealing works, e.g., Burman (1989); Bengio and Grandvalet (2004); Molinaro et al. (2005); Zhang and Yang (2015), have shown that, both empirically and theoretically, for modeling procedures with low instability, LOOCV often has the smallest variability. For example, in the context of the kernel SVM, Wang and Zou (2021) provided convincing numerical examples to show that (1) LOOCV has almost no bias in estimating the generalization error; (2) LOOCV does not have higher variance than ten-fold and five-fold CV. Consequently LOOCV results in a smaller overall error when estimating the prediction error as compared with ten-fold and five-fold CV.

From the aforementioned arguments, we can see the only legitimate complaint of LOOCV would arise from its expensive computation, as a typical approach needs to fit the models n times on the leave-one-out data before evaluating their performance with each of the sample removed, so the

computational cost is roughly n times as large as the cost of a single fit on the full data. To mitigate the computational burden, Golub et al. (1979) proposed a shortcut formula of LOOCV for smoothing splines such that the whole computation time is of the same order of fitting a single model, and the shortcut formula later evolved into the generalized cross-validation (GCV) for ridge regression.

Nevertheless, for the kernel classifiers, how to efficiently compute the exact LOOCV is a long-standing open problem. The shortcut cross-validation formula has been long considered as a unique property of some linear smoothers, and many works such as the generalized approximated cross-validation (GACV) (Wahba et al., 1999) resorted to approximating LOOCV, while there is no theoretical guarantee that LOOCV can always be well approximated. To solve the exact (rather than approximated) LOOCV, until very recently, Wang and Zou (2022) successfully proposed a leave-one-out lemma extending the Golub-Heath-Wahba formula to the kernel classifiers. Specifically, they showed the exact LOOCV error can be obtained by slightly varying the class labels without literally leaving out some samples during the CV procedure. Since no sample is left out, all the folds of LOOCV are using the same complete data and thus redundant computational efforts can be saved to dramatically accelerate LOOCV. Based on the leave-one-out lemma, Wang and Zou (2022) unified the training and tuning of the SVM and developed a new `magicsvm` algorithm, which often runs a magnitude faster than the state-of-art SVM solvers, e.g., `kernlab` (Karatzoglou et al., 2004) and `LIBSVM` (Chang and Lin, 2011).

In this work, the main contribution is to propose a consolidated CV algorithm via data reduction. The data reduction method was first proposed by Ghaoui et al. (2010) for the lasso method (Tibshirani, 1996) and then extended to the SVM (Ogawa et al., 2013; Wang et al., 2014; Pan and Xu, 2018; Hong et al., 2019). The key renovation of our proposal is to reduce all the cross-validated data in a consolidated manner, thereby aiming to speedup the whole SVM procedure. Our method is fundamentally different from the existing methods which isolate the model training and tuning. Moreover, the existing data reduction methods cannot handle the SVM with the bias (intercept), which is essentially useful for achieving high prediction accuracy. To handle the SVM with the bias, we propose a novel *two-stage consolidated CV*; such an extension is highly non-trivial.

We implement the consolidate CV in the `ccvsvm` algorithm. Simulations and nine benchmark data are used to demonstrate the superior performance of `ccvsvm` To give a quick demonstration, our consolidated CV algorithm reduces the run time from more than 1.5 hours (by LIBSVM) to less than one minute, when performing the exact LOOCV for the kernel SVM on a data set arrhythmia.

2 Methodology

2.1 SVM and the Exact Leave-One-Out Lemma

Since we need to work with the fundamentals of the SVM, we first review the SVM in this section.

We focus on binary classification. Let $L(u) = (1-u)_+ = \max(1-u, 0)$ be the *hinge loss*. Suppose there are n training samples, $(\mathbf{x}_i, y_i)$, $i = 1, 2, \ldots, n$, where each $\mathbf{x}_i \in I\!\!R^p$ and $y_i = \pm 1$. The SVM can be formulated as a function estimation problem in an RKHS (Wahba, 1990):

$$\hat{f}_l = \operatorname*{argmin}_{f \in \mathcal{H}_K} \left[\frac{1}{n} \sum_{i=1}^{n} (1 - y_i f(\mathbf{x}_i))_+ + \lambda_l \|f\|_{\mathcal{H}_K}^2 \right], \tag{1}$$

where $\lambda_l > 0$ is a tuning parameter chosen from a decreasing sequence $\lambda_1 > \lambda_2 > \ldots > \lambda_L$, $\mathcal{H}_K$, the reproducing kernel Hilbert space, is generated by a bivariate kernel function $K : \mathcal{X} \times \mathcal{X} \to I\!\!R$, and the classifier $\hat{f}$ is thus dubbed *kernel SVM*. Throughout this paper, we consider the *universal kernel*, whose induced RKHS $\mathcal{H}_K$ is rich enough to yield arbitrarily accurate decision boundaries (Steinwart, 2001; Micchelli et al., 2006). A commonly used universal kernel is the radial kernel $K(\mathbf{x}_i, \mathbf{x}_j) = \exp(-\sigma \|\mathbf{x}_i - \mathbf{x}_j\|_2^2)$.

By the representer theorem (Wahba, 1990), problem (1) has a finite-dimensional solution:

$$\hat{\boldsymbol{\alpha}}_l = \operatorname*{argmin}_{\boldsymbol{\alpha} \in I\!\!R^n} \left[\frac{1}{n} \sum_{i=1}^{n} (1 - y_i \mathbf{K}_i' \boldsymbol{\alpha})_+ + \lambda_l \boldsymbol{\alpha}' \mathbf{K} \boldsymbol{\alpha} \right], \tag{2}$$

where $\mathbf{K}$ is the $n \times n$ kernel matrix with $K_{ij} = K(\mathbf{x}_i, \mathbf{x}_j)$ and is assumed to be positive definite; $\mathbf{K}_i$ is its ith row. Thus problem (2) has a unique minimizer $\hat{f}(\mathbf{x}) = \sum_{i=1}^{n} \hat{\alpha}_i K(\mathbf{x}_i, \mathbf{x})$.

To tune the model, with the LOOCV procedure, the SVM is fitted on the training data with the jth sample opted out: for each $l = 1, 2, \ldots, L$ and each $j = 1, 2, \ldots, n$, let $\tilde{\boldsymbol{\alpha}}_l^{[-j]}$ be

$$\tilde{\boldsymbol{\alpha}}_l^{[-j]} = \underset{\boldsymbol{\alpha} \in \mathbb{R}^{n-1}}{\mathrm{argmin}} \left[\frac{1}{n} \sum_{i \neq j} \left(1 - y_i \mathbf{K}_i^{[-j]'} \boldsymbol{\alpha} \right)_+ + \lambda_l \boldsymbol{\alpha}' \mathbf{K}^{[-j]} \boldsymbol{\alpha} \right], \tag{3}$$

where $\mathbf{K}^{[-j]}$ is the leave-one-out kernel matrix induced by the training data without the jth sample. Problem (2) refers to *the complete data problem*, and problem (3) refers to *the LOOCV problem*.

The bottleneck of the LOOCV problem is mainly due to the computation involving n different leave-one-out kernel matrices. To reduce the computational burden, this work is based on the *exact leave-one-out lemma* (Wang and Zou, 2022) for the kernel SVM, and the key idea is to obtain the exact LOOCV from the complete kernel matrix.

Lemma 2.1. *(Exact leave-one-out lemma) For a given j, let $\tilde{y}_i^{[j]} = y_i$ if $i \neq j$ and $\tilde{y}_j^{[j]} = 0$. Define*

$$\hat{\boldsymbol{\alpha}}_l^{[-j]} = \underset{\boldsymbol{\alpha} \in \mathbb{R}^n}{\mathrm{argmin}} \left[\frac{1}{n} \sum_{i=1}^n \left(1 - \tilde{y}_i^{[j]} \mathbf{K}_i' \boldsymbol{\alpha} \right)_+ + \lambda_l \boldsymbol{\alpha}' \mathbf{K} \boldsymbol{\alpha} \right]. \tag{4}$$

Then the solution of problem (3) *can be obtained as*

$$\tilde{\boldsymbol{\alpha}}_l^{[-j]} = (\hat{\alpha}_{1,l}^{[-j]}, \ldots, \hat{\alpha}_{j-1,l}^{[-j]}, \hat{\alpha}_{j+1,l}^{[-j]}, \ldots, \hat{\alpha}_{n,l}^{[-j]})'.$$

Although problem (3) can be transformed into problem (4), the solutions of the two problems have different lengths. One can show that $\hat{\alpha}_{j,l}^{[-j]}$, the jth element of $\hat{\boldsymbol{\alpha}}_l^{[-j]}$, is zero, and the solution of problem (3) can be retrieved by knocking off the jth element from $\hat{\boldsymbol{\alpha}}_l^{[-j]}$.

As a consequence of transforming problem (3) into problem (4), the same kernel matrix $\mathbf{K}$ is used in all the folds during LOOCV, rather than the leave-one-out matrices $\mathbf{K}^{[-j]}$, while slightly different responses are crafted for different j. By sharing the same kernel matrix, some redundant calculations can be saved and Wang and Zou (2022) developed the efficient algorithm *magicsvm*.

2.2 Consolidated CV via Data Reduction

On the basis of Lemma 2.1, we propose a data reduction strategy to accelerate the LOOCV computation of the kernel SVM, which is referred to as *consolidated CV*.

For notational convenience, the complete data problem (2) can be written as a special case of problem (4) with $j = 0$, i.e., $\hat{\boldsymbol{\alpha}}_l \equiv \hat{\boldsymbol{\alpha}}_l^{[-0]}$ and

$$\hat{\boldsymbol{\alpha}}_l^{[-0]} = \underset{\boldsymbol{\alpha} \in \mathbb{R}^n}{\mathrm{argmin}} \left[\frac{1}{n} \sum_{i=1}^n \left(1 - \tilde{y}_i^{[0]} \mathbf{K}_i' \boldsymbol{\alpha} \right)_+ + \lambda_l \boldsymbol{\alpha}' \mathbf{K} \boldsymbol{\alpha} \right],$$

where we define $\tilde{y}_i^{[0]} = y_i$ for each $i = 1, 2, \ldots, n$. By solving problem (4) for all $j = 0, 1, \ldots, n$, we both train the SVM through the complete data problem (2) and tune it using LOOCV.

The idea of consolidated CV is motivated by the sparsity of the solution $\hat{\boldsymbol{\alpha}}_l^{[-j]}$ in problem (4). To see this, we check the optimality condition of problem (4) by taking the sub-differential of the objective with respect to each $\mathbf{K}_i' \boldsymbol{\alpha}$, for each $j = 0, 1, \ldots, n$:

$$0 \in \frac{1}{n} \tilde{y}_i^{[j]} \partial L \left(\tilde{y}_i^{[j]} \mathbf{K}_i' \hat{\boldsymbol{\alpha}}_l^{[-j]} \right) + 2\lambda_l \hat{\alpha}_{i,l}^{[-j]}, \ \forall i = 1, \ldots, n,$$

where $\partial L(t)$ is the subgradient of the hinge loss function: $\partial L(t) = 1$, if $t < 1$; $\partial L(t) = 0$, if $t > 1$; and $\partial L(t) \in [-1, 0]$ if $t = 1$. It follows that

$$\hat{\alpha}_{i,l}^{[-j]} = \begin{cases} \dfrac{\tilde{y}_i^{[j]}}{2n\lambda_l}, & \text{if } \tilde{y}_i^{[j]} \mathbf{K}_i' \hat{\boldsymbol{\alpha}}_l^{[-j]} < 1, \\ 0, & \text{if } \tilde{y}_i^{[j]} \mathbf{K}_i' \hat{\boldsymbol{\alpha}}_l^{[-j]} > 1. \end{cases}$$

112   By translating $\tilde{y}_i^{[j]}$ back to y_i, we see

$$\hat{\alpha}_{i,l}^{[-j]} = \begin{cases} \dfrac{y_i}{2n\lambda_l}, & \text{if } y_i\mathbf{K}_i'\hat{\alpha}_l^{[-j]} < 1 \text{ and } i \neq j, \\ 0, & \text{if } y_i\mathbf{K}_i'\hat{\alpha}_l^{[-j]} > 1 \text{ or } i = j. \end{cases} \tag{5}$$

113   Expression (5) hints on a possible data reduction strategy: before invoking the actual calculation
114   of $\hat{\alpha}_l^{[-j]}$, if we are advised that $y_i\mathbf{K}_i'\hat{\alpha}_l^{[-j]} > 1$ for some i, then we can directly set $\hat{\alpha}_{i,l}^{[-j]}$ to zero;
115   likewise, if $y_i\mathbf{K}_i'\hat{\alpha}_l^{[-j]} < 1$ is given, then $\hat{\alpha}_{i,l}^{[-j]}$ must be $y_i/(2n\lambda_l)$ unless $i = j$. We can pre-
116   determine the values of some coordinates and only need to focus on the calculation of the remaining
117   ones. Hence the dimension of problem (4) can be reduced.

118   The key to performing the data reduction through expression (5) is to know whether $y_i\mathbf{K}_i'\hat{\alpha}_l^{[-j]} < 1$
119   or > 1 for some i before $\hat{\alpha}_l^{[-j]}$ is actually computed. We present the following theorem.

Theorem 2.2. *For some $l > 1$, suppose we have solved*

$$\hat{\alpha}_{l-1} = \operatorname*{argmin}_{\alpha \in \mathbb{R}^n} \left[\frac{1}{n} \sum_{i=1}^{n} (1 - y_i\mathbf{K}_i'\alpha)_+ + \lambda_{l-1}\alpha'\mathbf{K}\alpha \right].$$

120   *For each $i = 1, 2, \ldots, n$, define*

$$a_{i,l}^+ = \frac{\lambda_{l-1} + \lambda_l}{2\lambda_l} y_i\mathbf{K}_i'\hat{\alpha}_{l-1} + \frac{\lambda_{l-1} - \lambda_l}{2\lambda_l}\sqrt{B}\sqrt{\hat{\alpha}_{l-1}'\mathbf{K}\hat{\alpha}_{l-1}} + \frac{B}{2n\lambda_l},$$

$$a_{i,l}^- = \frac{\lambda_{l-1} + \lambda_l}{2\lambda_l} y_i\mathbf{K}_i'\hat{\alpha}_{l-1} - \frac{\lambda_{l-1} - \lambda_l}{2\lambda_l}\sqrt{B}\sqrt{\hat{\alpha}_{l-1}'\mathbf{K}\hat{\alpha}_{l-1}} - \frac{B}{2n\lambda_l},$$

121   *where $B = \max_i K(\mathbf{x}_i, \mathbf{x}_i)$ and $B = 1$ for the radial kernel. Then for each $j = 0, 1, \ldots, n$, it holds*

$$a_{i,l}^- \leq y_i\mathbf{K}_i'\hat{\alpha}_l^{[-j]} \leq a_{i,l}^+, \ \forall i \neq j. \tag{6}$$

122   *Further, let $\mathcal{L} = \{i : a_{i,l}^+ < 1\}$ and $\mathcal{R} = \{i : a_{i,l}^- > 1\}$. Then the solution of problem (4) satisfies that*

$$\hat{\alpha}_{i,l}^{[-j]} = \begin{cases} \dfrac{\tilde{y}_i^{[j]}}{2n\lambda_l}, & \text{if } i \in \mathcal{L}; \\ 0, & \text{if } i \in \mathcal{R}. \end{cases}$$

123

124   The proof of Theorem 2.2 is given in Section A.1 of the appendix. Note that Theorem 2.2 holds
125   for $\hat{\alpha}_l^{[-j]}, \forall j = 0, 1, \ldots, n$. By utilizing knowledge of $\hat{\alpha}_{l-1}$, which is the solution of the complete
126   data problem with the tuning parameter λ_{l-1}, we can pre-determining certain coordinates for both
127   the solutions of the complete data problem and all the LOOCV problems with λ_l, i.e., for all
128    $j = 0, 1, \ldots, n$, through $\mathcal{L}$ and $\mathcal{R}$, thus performing data reduction in a *consolidated* fashion.

129   To solve problem (4), Theorem 2.2 implies that $\hat{\alpha}_{i,l}^{[-j]}$ for $i \in \mathcal{L}$ and $i \in \mathcal{R}$ can be pre-determined, so
130   we only need to solve $\hat{\alpha}_{i,l}^{[-j]}$, for $i \in \mathcal{S}$ where $\mathcal{S} \equiv (\mathcal{L} \cup \mathcal{R})^C$. Denote by τ the one-to-one mapping
131   from $\{1, 2, \ldots, n_s\}$ to $\mathcal{S}$, where n_s is the cardinality of $\mathcal{S}$. Let Γ be the $n \times n_s$ sub-matrix of $\mathbf{K}$ such
132   that its ith column $\Gamma_i = \mathbf{K}_{\tau(i)}$. Let Σ be the $n_s \times n_s$ matrix such that $\Sigma_{ij} = K_{\tau(i)\tau(j)}$.

133   For each $j = 0, 1, \ldots, n$, let $\bar{\mathbf{y}}^{[j]}$ be the n-vector with the ith element to be $\tilde{y}_i^{[j]}$ if $i \in \mathcal{S}$, and 0 if
134    $i \notin \mathcal{S}$. The solution of problem (4) is obtained as

$$\hat{\alpha}_{i,l}^{[-j]} = \begin{cases} \dfrac{\tilde{y}_i^{[j]}}{2n\lambda_l}, & \text{if } i \in \mathcal{L}, \\ 0, & \text{if } i \in \mathcal{R}, \\ \hat{\eta}_{\tau^{-1}(i),l}^{[-j]}, & \text{if } i \in \mathcal{S}, \end{cases} \tag{7}$$

135   where $\hat{\eta}_{\tau^{-1}(i),l}^{[-j]}$ is the $\tau^{-1}(i)$th element of

$$\hat{\eta}_l^{[-j]} = \operatorname*{argmin}_{\eta \in \mathbb{R}^{n_s}} \left[\frac{1}{n} \sum_{i=1}^{n} \left(1 - \tilde{y}_i^{[j]}\Gamma_i'\eta - \frac{1}{2n\lambda_l}\tilde{y}_i^{[j]}\mathbf{K}_i'\bar{\mathbf{y}}^{[j]} \right)_+ + \frac{1}{n}\bar{\mathbf{y}}^{[j]'}\Gamma\eta + \lambda_l\eta'\Sigma\eta \right]. \tag{8}$$

136 The dimension of problem (8) is n_s, which is lower than n – the dimension of the original problem (4).
137 The matrices $\mathbf{\Gamma}$ and $\mathbf{\Sigma}$ are the same for each $j = 0, 1, \ldots, n$. We shall introduce an optimization
138 algorithm for solving problem (8) in the next section.

2.3 A Consolidated Algorithm for Solving Problem (8)

140 Due to Theorem 2.2, we can perform LOOCV by solving problem (8), a reduced-optimization
141 problem, for each j. To overcome the computational challenge caused by non-smoothness of the
142 hinge loss, we consider a smoothed loss,

$$
L_\tau(u) = \begin{cases} 0 & u \geq 1 + \tau, \\ (u - (1 + \tau))^2/(4\tau) & 1 - \tau < u < 1 + \tau, \\ 1 - u & u \leq 1 - \tau, \end{cases}
$$

143 for some small $\tau > 0$. One can show that L_τ has a Lipschitz continuous gradient,
144 $|L'_\tau(t_1) - L'_\tau(t_2)| \leq \frac{1}{2\tau}|t_1 - t_2|, \forall t_1, t_2 \in \mathbb{R}$. Thus a smoothed surrogate of problem (8) is
145

$$
\hat{\boldsymbol{\eta}}_{\tau,l}^{[-j]} = \underset{\boldsymbol{\eta} \in \mathbb{R}^{n_s}}{\operatorname{argmin}} \left[\frac{1}{n} \sum_{i=1}^n L^\tau \left(\tilde{y}_i^{[j]} \mathbf{\Gamma}_i' \boldsymbol{\eta} - \frac{1}{2n\lambda_l} \tilde{y}_i^{[j]} \mathbf{K}_i' \bar{\mathbf{y}}^{[j]} \right) + \frac{1}{n} \bar{\mathbf{y}}^{[j]'} \mathbf{\Gamma} \boldsymbol{\eta} + \lambda_l \boldsymbol{\eta}' \mathbf{\Sigma} \boldsymbol{\eta} \right]. \tag{9}
$$

146 Problem (9) can be solved using the proximal gradient descent (PGD) algorithm (Parikh and Boyd,
147 2014). Specifically, the matrix inversion is computed first

$$
\mathbf{P}^{-1} = \left(2\lambda_l \mathbf{\Sigma} + \frac{1}{n\tau} \mathbf{\Gamma}' \mathbf{\Gamma} \right)^{-1}. \tag{10}
$$

148 Then, for each $j = 0, 1, \ldots, n$, we update

$$
\boldsymbol{\eta}^{[-j]} \leftarrow \boldsymbol{\eta}^{[-j]} - \mathbf{P}^{-1} \left(\mathbf{\Gamma}' \mathbf{z}^{(k)} + \frac{1}{n} \mathbf{\Gamma}' \bar{\mathbf{y}}^{[j]} + 2\lambda_l \mathbf{\Sigma} \boldsymbol{\eta}^{[-j]} \right) \tag{11}
$$

149 until convergence, and then let $\hat{\boldsymbol{\eta}}_{\tau,l}^{[-j]} \leftarrow \boldsymbol{\eta}^{[-j]}$. We claim the above algorithm is consolidated since
150 the same matrix inversion $\mathbf{P}^{-1}$ obtained from equation (10) can be used in equation (11) for all j (all
151 folds.) By saving huge computational efforts in inverting n matrices, the consolidated CV algorithm
152 is much more efficient than the standard CV implementation. We also include the warm-start and
153 Nesterov's acceleration to further accelerate the computation.

154 We just discussed the PGD algorithm for solving a smoothed SVM problem (9). Interestingly, the
155 *exact* SVM solution based on problem (8) can be obtained by iteratively solving problem (9) with
156 $\tau_1 > \tau_2 > \ldots$ where $\tau_1 = 1$ and $\tau_k = \tau_{k-1}/8$ for $k > 1$. The iteration is able to reach the exact
157 solution of problem (8) in a finite number of steps, following a simple projection step. To conserve
158 space, we omit details and refer interesting readers to Wang and Zou (2022).

3 two-stage Consolidated CV for the General SVM Problems

160 The consolidated CV developed in Section 2 does not include the bias; nonetheless, the SVM without
161 the bias may have lower prediction accuracy and its use is limited in certain applications. Although a
162 regularized bias can be used by adding a constant feature to the data matrix, the standard practice
163 of the SVM does not regularize the bias term. Thus our goal is to compute the SVM with the bias,
164 namely, *the general SVM problems*. In this section, we extend the consolidated CV to handle the
165 general SVM problems. Such an extension turns out to be non-trivial.

166 The general SVM problem is formulated as follows,

$$
(\hat{\beta}_{0,l}, \hat{\boldsymbol{\alpha}}_l) = \underset{\beta_0 \in \mathbb{R}, \, \boldsymbol{\alpha} \in \mathbb{R}^n}{\operatorname{argmin}} \frac{1}{n} \sum_{i=1}^n [1 - y_i(\beta_0 + \mathbf{K}_i' \boldsymbol{\alpha})]_+ + \lambda_l \boldsymbol{\alpha}' \mathbf{K} \boldsymbol{\alpha}, \tag{12}
$$

167 and the corresponding LOOCV problems are

$$
(\hat{\beta}_{0,l}^{[-j]}, \hat{\boldsymbol{\alpha}}_l^{[-j]}) = \underset{\beta_0 \in \mathbb{R}, \, \boldsymbol{\alpha} \in \mathbb{R}^n}{\operatorname{argmin}} \frac{1}{n} \sum_{i=1}^n \left[1 - \tilde{y}_i^{[j]}(\beta_0 + \mathbf{K}_i' \boldsymbol{\alpha}) \right]_+ + \lambda_l \boldsymbol{\alpha}' \mathbf{K} \boldsymbol{\alpha}. \tag{13}
$$

168 For notational convenience, we let $\tilde{y}_i^{[0]} = y_i$ and let $(\hat{\beta}_{0,l}, \hat{\boldsymbol{\alpha}}_l) = (\hat{\beta}_{0,l}^{[-0]}, \hat{\boldsymbol{\alpha}}_l^{[-0]})$, so we extend
169 problem (13) with $j = 0$ to include the complete data problem (12) as a special case.

170 The key difficulty of developing the consolidated CV procedure for the general SVM problems is
171 that $|\hat{\beta}_{0,l} - \hat{\beta}_{0,l}^{[-j]}|$ is hard to bound. To this end, we propose a *two-stage consolidated CV procedure*,
172 where we give a consolidated bound of $|\hat{\beta}_{0,l} - \hat{\beta}_{0,l}^{[-j]}|$ for all j in the first stage.

173 For $l > 1$, suppose we have found the solutions of problems (12) and (13) with the tuning parameter
174 λ_{l-1}. Denote these solutions by $(\hat{\beta}_{0,l-1}, \hat{\boldsymbol{\alpha}}_{l-1})$ and $(\hat{\beta}_{0,l-1}^{[-j]}, \hat{\boldsymbol{\alpha}}_{l-1}^{[-j]})$. In Lemma 3.1, for each i, we
175 give a consolidated bound of $y_i \mathbf{K}_i' \hat{\boldsymbol{\alpha}}_l^{[-j]}$ for all $j = 0, 1, \ldots, n$ and $j \neq i$.

176 **Lemma 3.1.** *For each $i = 1, 2, \ldots, n$, define*

$$
\begin{aligned}
c_{i,l}^+ &= \max_{\substack{j=0,1,\ldots,n \\ j \neq i}} \left\{ \frac{\lambda_{l-1} + \lambda_l}{2\lambda_l} y_i \mathbf{K}_i' \hat{\boldsymbol{\alpha}}_{l-1}^{[-j]} + \frac{\lambda_{l-1} - \lambda_l}{2\lambda_l} \sqrt{B} \sqrt{\hat{\boldsymbol{\alpha}}_{l-1}^{[-j]\prime} \mathbf{K} \hat{\boldsymbol{\alpha}}_{l-1}^{[-j]}} \right\}, \\
c_{i,l}^- &= \min_{\substack{j=0,1,\ldots,n \\ j \neq i}} \left\{ \frac{\lambda_{l-1} + \lambda_l}{2\lambda_l} y_i \mathbf{K}_i' \hat{\boldsymbol{\alpha}}_{l-1}^{[-j]} - \frac{\lambda_{l-1} - \lambda_l}{2\lambda_l} \sqrt{B} \sqrt{\hat{\boldsymbol{\alpha}}_{l-1}^{[-j]\prime} \mathbf{K} \hat{\boldsymbol{\alpha}}_{l-1}^{[-j]}} \right\},
\end{aligned}
\tag{14}
$$

177 *where $B = \max_i K(\mathbf{x}_i, \mathbf{x}_i)$ and $B = 1$ for the radial kernel. Then for any $i = 1, \ldots, n$, it holds that*
178

$$
c_{i,l}^- \leq y_i \mathbf{K}_i' \hat{\boldsymbol{\alpha}}_l^{[-j]} \leq c_{i,l}^+, \ \forall j \neq i.
\tag{15}
$$

179

180 On the basis of the bounds of $y_i \mathbf{K}_i' \hat{\boldsymbol{\alpha}}_l^{[-j]}$ given in Lemma 3.1, we next present Lemma 3.2 and
181 Lemma 3.3 to give bounds of $\hat{\beta}_{0,l}^{[-j]}$ that are consolidated for all $j = 0, 1, \ldots, n$.

182 **Lemma 3.2.** *With $c_{i,l}^-$ and $c_{i,l}^+$ from Lemma 3.1, for a given constant b, define $\mathcal{S}_1(b) = \{i : y_i b +$
183 $c_{i,l}^+ < 1\}$ and $\mathcal{S}_2(b) = \{i : y_i b + c_{i,l}^- > 1\}$. Let $n_+(b) = \sum_{i \in (\mathcal{S}_1(b) \cup \mathcal{S}_2(b))^C} I(y_i = 1)$ and
184 $n_-(b) = \sum_{i \in (\mathcal{S}_1(b) \cup \mathcal{S}_2(b))^C} I(y_i = -1)$. Define $\psi^+(b) = \sum_{i \in \mathcal{S}_1(b)} y_i + n_+(b) + 1$ and $\psi^-(b) =$
185 $\sum_{i \in \mathcal{S}_1(b)} y_i - n_-(b) - 1$. Then we have*

186   *(1) both $\psi^+(b)$ and $\psi^-(b)$ are non-increasing in b;*

187   *(2) $\psi^+(b) < 0$ implies $b > \hat{\beta}_{0,l}^{[-j]}$ for all $j = 0, 1, \ldots, n$;*

188   *(3) $\psi^-(b) > 0$ implies $b < \hat{\beta}_{0,l}^{[-j]}$ for all $j = 0, 1, \ldots, n$.*

189 Following Lemma 3.2, we develop a bi-section algorithm in Algorithm 1. As shown in Lemma 3.3,
190 Algorithm 1 yields consolidated bounds for $\hat{\beta}_{0,l}^{[-j]}$ for all $j = 0, 1, \ldots, n$.

191 **Lemma 3.3.** *Suppose the input of Algorithm 1, $c_{i,l}^-$ and $c_{i,l}^+$, satisfies inequality (15), then the output
192 of Algorithm 1, $\beta_{0,l}^+$ and $\beta_{0,l}^-$, satisfies that*

$$
\beta_{0,l}^- < \hat{\beta}_{0,l}^{[-j]} < \beta_{0,l}^+. \ \forall j = 0, 1, \ldots, n.
\tag{16}
$$

193

194 It immediately follows from Lemma 3.3 that

$$
|\hat{\beta}_{0,l} - \hat{\beta}_{0,l}^{[-j]}| < \beta_{0,l}^+ - \beta_{0,l}^-,
\tag{17}
$$

195 for any j, achieving the goal of the first stage.

196 We have constructed the bounds in inequalities (15) and (16). However, these bounds are too loose
197 to develop data reduction rules in practice. The loose bounds are mainly caused by the maximum
198 and minimum operators that are involved in equations (14). To this end, in the second stage, we give
199 refined bounds, which are presented below.

Algorithm 1 Bi-section algorithm to find $\beta_{0,l}^+$ and $\beta_{0,l}^-$

Input: $\mathbf{y}, c_{i,l}^-, c_{i,l}^+, \epsilon = 10^{-7}$.

1: Compute B^+ and B^- as

$$B^+ = \max\left\{ \max_{\{i:y_i=-1\}}\{c_{i,l}^+ - 1\}, \max_{\{i:y_i=1\}}\{1 - c_{i,l}^-\} \right\} + \epsilon,$$

$$B^- = \min\left\{ \min_{\{i:y_i=-1\}}\{c_{i,l}^- - 1\}, \min_{\{i:y_i=1\}}\{1 - c_{i,l}^+\} \right\} - \epsilon.$$

2: Let $a^+ \leftarrow B^+, c^+ \leftarrow B^-, b^+ \leftarrow (a^+ + c^+)/2$.
3: **repeat**
4:     Compute $\psi^+(b^+)$.
5:     Let $a^+ \leftarrow b^+$ and $b^+ \leftarrow (b^+ + c^+)/2$ if $\psi^+(b^+) < 0$.
6:     Let $c^+ \leftarrow b^+$ and $b^+ \leftarrow (a^+ + b^+)/2$ if $\psi^+(b^+) \geq 0$.
7: **until** $|a^+ - c^+| < \epsilon$.
8: Let $\beta_{0,l}^+ \leftarrow a^+$.
9: Let $a^- \leftarrow B^+, c^- \leftarrow B^-, b^- \leftarrow (a^- + c^-)/2$.
10: **repeat**
11:     Compute $\psi^-(b^-)$.
12:     Let $a^- \leftarrow b^-$ and $b^- \leftarrow (b^- + c^-)/2$ if $\psi^-(b^-) \leq 0$.
13:     Let $c^- \leftarrow b^-$ and $b^- \leftarrow (a^- + b^-)/2$ if $\psi^-(b^-) > 0$.
14: **until** $|a^- - c^-| < \epsilon$.
15: Let $\beta_{0,l}^- \leftarrow c^-$.

Output: $\beta_{0,l}^+$ **and** $\beta_{0,l}^-$.

Lemma 3.4. *For each $i = 1, 2, \ldots, n$, define*

$$\tilde{c}_{i,l}^+ = \frac{\lambda_{l-1} + \lambda_l}{2\lambda_l} y_i \mathbf{K}_i' \hat{\boldsymbol{\alpha}}_{l-1} + \frac{\lambda_{l-1} - \lambda_l}{2\lambda_l}\sqrt{B}\sqrt{\hat{\boldsymbol{\alpha}}_{l-1}' \mathbf{K}\hat{\boldsymbol{\alpha}}_{l-1}} + \sqrt{\frac{B^2}{16n^2\lambda_l^2} + \frac{B(\beta_{0,l}^+ - \beta_{0,l}^-)}{2n\lambda_l}} + \frac{B}{4n\lambda_l},$$

$$\tilde{c}_{i,l}^- = \frac{\lambda_{l-1} + \lambda_l}{2\lambda_l} y_i \mathbf{K}_i' \hat{\boldsymbol{\alpha}}_{l-1} - \frac{\lambda_{l-1} - \lambda_l}{2\lambda_l}\sqrt{B}\sqrt{\hat{\boldsymbol{\alpha}}_{l-1}' \mathbf{K}\hat{\boldsymbol{\alpha}}_{l-1}} - \sqrt{\frac{B^2}{16n^2\lambda_l^2} + \frac{B(\beta_{0,l}^+ - \beta_{0,l}^-)}{2n\lambda_l}} - \frac{B}{4n\lambda_l},$$

where $\beta_{0,l}^+$ and $\beta_{0,l}^-$ are produced by Algorithm 1. Then for any $j = 1, \ldots, n$, it holds that

$$\tilde{c}_{i,l}^- \leq y_i \mathbf{K}_i' \hat{\boldsymbol{\alpha}}_l^{[-j]} \leq \tilde{c}_{i,l}^+, \ \forall j = 0, 1, \ldots, n, \text{ and } j \neq i. \tag{18}$$

Hence by Lemmata 3.1 and 3.4, we have

$$\hat{c}_{i,l}^- \equiv \max\{c_{i,l}^-, \tilde{c}_{i,l}^-\} \leq y_i \mathbf{K}_i' \hat{\boldsymbol{\alpha}}_l^{[-j]} \leq \min\{c_{i,l}^+, \tilde{c}_{i,l}^+\} \equiv \hat{c}_{i,l}^+, \tag{19}$$

for any $j = 0, 1, \ldots, n$, and $j \neq i$. We then use $\max\{c_{i,l}^-, \tilde{c}_{i,l}^-\}$ and $\min\{c_{i,l}^+, \tilde{c}_{i,l}^+\}$ as the input in Algorithm 1 to yield the output $\tilde{\beta}_{0,l}^+$ and $\tilde{\beta}_{0,l}^-$. Then by inequality (18) and Lemma 3.2, we have

$$\tilde{\beta}_{0,l}^- < \hat{\beta}_{0,l}^{[-j]} < \tilde{\beta}_{0,l}^+. \ \forall j = 0, 1, \ldots, n. \tag{20}$$

Therefore, we glean inequalities (18) and (20), which are the refined bounds of inequalities (15) and (16). Using the refined bounds, we now present the main theorem.

Theorem 3.5. *The solution of problem* (12), $\hat{\boldsymbol{\alpha}}_l$, *satisfies:*

$$\hat{\alpha}_{i,l} = \begin{cases} \dfrac{y_i}{2n\lambda_l}, & \text{if } i \in \tilde{\mathcal{L}}; \\ 0, & \text{if } i \in \tilde{\mathcal{R}}, \end{cases}$$

Table 1: Run time (in second), objective value, and test error of four kernel SVM solvers under mixture Gaussian distributed data with $p = \{20, 200\}$, and $n = \{200, 400, 800, 1600\}$. The test error is assessed on independently generated test data. The numbers are the average quantities over 50 independent runs and the standard errors are presented in parentheses.

p	n	method	time (s)	objective	test error	method	time (s)	objective	test error
20	200	ccvsvm	5.1	0.814 (.005)	0.351 (.007)	kernlab	73.4	0.814 (.005)	0.351 (.007)
		magicsvm	7.7	0.814 (.005)	0.351 (.007)	LIBSVM	144.4	0.828 (.014)	0.351 (.007)
	400	ccvsvm	44.2	0.827 (.003)	0.332 (.005)	kernlab	334.3	0.827 (.003)	0.332 (.005)
		magicsvm	87.8	0.827 (.003)	0.332 (.005)	LIBSVM	879.7	0.827 (.003)	0.332 (.005)
	800	ccvsvm	446.8	0.846 (.002)	0.309 (.002)	kernlab	2220.2	0.846 (.002)	0.309 (.002)
		magicsvm	847.3	0.846 (.002)	0.309 (.002)	LIBSVM	6519.7	0.846 (.002)	0.310 (.002)
	1600	ccvsvm	3829.5	0.853 (.001)	0.297 (.001)	kernlab	25530.5	0.853 (.001)	0.297 (.001)
		magicsvm	7024.1	0.853 (.001)	0.297 (.001)	LIBSVM	63886.1	0.853 (.001)	0.297 (.001)
200	200	ccvsvm	6.8	0.780 (.006)	0.337 (.015)	kernlab	337.6	0.780 (.006)	0.339 (.015)
		magicsvm	12.9	0.780 (.006)	0.337 (.015)	LIBSVM	932.5	0.780 (.006)	0.342 (.015)
	400	ccvsvm	66.0	0.794 (.003)	0.366 (.015)	kernlab	2304.1	0.794 (.003)	0.366 (.015)
		magicsvm	150.1	0.794 (.003)	0.366 (.015)	LIBSVM	6641.9	0.794 (.003)	0.368 (.015)
	800	ccvsvm	530.4	0.811 (.001)	0.346 (.015)	kernlab	36771.4	0.811 (.001)	0.346 (.015)
		magicsvm	996.1	0.811 (.001)	0.346 (.015)	LIBSVM	109365.5	0.811 (.001)	0.346 (.015)
	1600	ccvsvm	5489.2	0.821 (.001)	0.322 (.013)	kernlab	461245.7	0.821 (.001)	0.322 (.013)
		magicsvm	10803.9	0.821 (.001)	0.322 (.013)	LIBSVM	1436416.1	0.821 (.001)	0.322 (.013)

and for any $j = 1, \ldots, n$, the solution of problem (13), $\hat{\boldsymbol{\alpha}}_l^{[-j]}$, satisfies:

$$\hat{\alpha}_{i,l}^{[-j]} = \begin{cases} \dfrac{y_i}{2n\lambda_l}, & \text{if } i \in \tilde{\mathcal{L}} \text{ and } i \neq j; \\ 0, & \text{if } i \in \tilde{\mathcal{R}} \text{ or } i = j, \end{cases}$$

where $\hat{c}_{i,l}^+$ and $\hat{c}_{i,l}^-$ are given in inequality (19) and

$$\tilde{\mathcal{L}} = \left\{ i : y_i = 1 \text{ and } \tilde{\beta}_{0,l}^+ + \hat{c}_{i,l}^+ < 1 \right\} \cup \left\{ i : y_i = -1 \text{ and } -\tilde{\beta}_{0,l}^- + \hat{c}_{i,l}^+ < 1 \right\},$$

$$\tilde{\mathcal{R}} = \left\{ i : y_i = 1 \text{ and } \tilde{\beta}_{0,l}^- + \hat{c}_{i,l}^- > 1 \right\} \cup \left\{ i : y_i = -1 \text{ and } -\tilde{\beta}_{0,l}^+ + \hat{c}_{i,l}^- > 1 \right\}.$$

Thus by Theorem 3.5, problem (13) can be solved through some reduced-dimensional optimization problems, which are similar to problem (8) where the bias is excluded. Therefore, we can follow the discussions in Section 2.3 to employ the same PGD algorithm and the exact smoothing technique to obtain the exact solution for problem (13). Details of the algorithm are omitted to conserve space.

4 Numerical Studies

In this section, we demonstrate the computational advantages of fitting the kernel SVM using `ccvsvm` over the three other competitors, `magicsvm`, `kernlab`, and `LIBSVM`, with simulations and real data applications. To pursue the high predictive accuracy, we focus on the SVM with the bias, so we do not compare `ccvsvm` with the algorithms that cannot handle the bias.

4.1 Simulations

A commonly used simulation data from mixture Gaussian distributions (Hastie et al., 2009) is used. We generate mean vectors $\boldsymbol{\mu}_{k_+}$ from $N(\boldsymbol{\mu}_+, \mathbf{I}_p)$ where $k = 1, 2, \ldots, 10$ in which $\boldsymbol{\mu}_+ = (1, 1, \ldots, 1, 0, 0, \ldots, 0)$ with half of the coordinates to be zero. Each positive-class training sample is independently generated from $N(\boldsymbol{\mu}_{k_+}, 3^2)$ where k is drawn from the discrete uniform distribution on $\{1, 2, \ldots, 10\}$. Using the same procedure, we obtain the negative-class training data from $N(\boldsymbol{\mu}_{k_-}, 3^2)$ where k is also uniform on $\{1, 2, \ldots, 10\}$ and $\boldsymbol{\mu}_- = (0, 0, \ldots, 0, 1, 1, \ldots, 1)$. For each combination of the feature dimension $p = 20$ and 200 and the sample size $n = 200, 400, 800$, and 1600, we fit the kernel SVM using the four kernel SVM solvers, `ccvsvm`, `magicsvm`, `kernlab`, and `LIBSVM`, to compute the entire solution paths at a sequence of 50 tuning parameters, uniformly distributed on the

Table 2: Run time (in second) of four SVM solvers for benchmark data, averaged over 50 runs.

data	n	p	ccvsvm	magicsvm	kernlab	LIBSVM
arrhythmia	452	191	48.076	113.099	1554.579	5061.881
australian	690	14	202.863	412.323	902.463	2178.644
chess	3196	37	21768.612	38942.348	> 240 hours	> 240 hours
heart	270	13	8.464	16.466	89.373	168.477
leuk	72	7218	0.464	0.828	1548.724	4811.612
malaria	71	22283	0.504	0.819	4804.442	13835.143
musk	476	166	62.169	127.656	1563.262	4778.779
sonar	208	60	4.736	7.033	98.080	221.505
valley	606	100	149.034	311.147	2230.010	6428.014

logarithm scale between e^{-6} and e^6. The radial kernel is used and the bandwidth is the default option of `kernlab`, which generally performs well. We compared the run time, objective function value, and test error of the four solvers, where the run time includes the whole computation process including training and tuning the model. The objective function value is computed from equation (2). Test error is assessed on $10,000$ test samples which are independently generated from the same distribution. Computations were conducted on an Intel(R) Xeon(R) Gold 6230 CPU @ 2.10 GHz.

Table 1 shows that, to reach the same objective value and the test error, our `ccvsvm` algorithm is roughly twice as fast as `magicsvm`, and it is about an order of magnitude faster than `kernlab` and `LIBSVM`. In addition, we observe that `kernlab` and `LIBSVM` significantly slow down as p increases, e.g., LIBSVM is about 20 times slower when p grows from 20 to 200, whereas the speed of `ccvsvm` and `magicsvm` is quite insensitive to the change of dimensions. Remarkably, for $p = 200$ and $n = 1600$, our `ccvsvm` algorithm finishes training and tuning the SVM using LOOCV within two hours, while with the same accuracy, LIBSVM spends about 400 hours, lasting over 16 days.

4.2 Benchmark Data Applications

We test the performance on benchmark data applications. We study nine commonly used real data applications from the UCI machine learning repository (Dua and Graff, 2017). The sample sizes range from 208 to $3,196$. Two high-dimensional data sets with the number of features $p = 7,218$ and $22,283$ are included. Each data set is split into a training set and a test set with the ratio $9 : 1$. The kernel SVM is trained and tuned by the four solvers on the training set, and the test error is assessed on the test set. We adopt the training-test split-ratio $9 : 1$ because we aim to assign most of the samples to the training set and the computation time can be evaluated using relatively large data.

Table 2 exhibits the timing comparisons, where we discover our `ccvsvm` algorithm is clearly the fastest. It is about as twice as fast as `magicsvm`, and it is significantly faster than `kernlab` and `LIBSVM`. Especially for the two high-dimensional examples, `magicsvm` is thousands or even tens of thousands faster than `kernlab` and `LIBSVM`, and `ccvsvm` further cuts the run time of `magicsvm` into half. Similar to the simulation results, we observe that all the four kernel SVM solvers deliver almost the same objective values and test errors on the real data applications; for sake of space limit, we omit the results for the accuracy.

5 Discussions

In this work, we have introduced a consolidated CV procedure and developed an algorithm called `ccvsvm` for the kernel SVM, which is one of the most successful classifiers. Our work is built on the recently proposed leave-one-out lemma and the `magicsvm` algorithm: the `ccvsvm` algorithm can even double the speed of `magicsvm`, which has already shown remarkable computational advantages over the mainstream SVM solvers, `kernlab` and `LIBSVM`.

Limitation. The proposed method is only for LOOCV and SVM since it utilizes the special structure of support vectors. However, in future works, it is interesting to explore if the consolidated CV can be generalized to other K-fold CV or the hold-out validation more broadly, and also extended to other computationally expensive machine learning methods.

Societal impact. This work does not present any foreseeable societal consequence.

References

ARLOT, S. and CELISSE, A. (2010). A survey of cross-validation procedures for model selection. *Statistics Surveys* **4** 40–79.

BENGIO, Y. and GRANDVALET, Y. (2004). No unbiased estimator of the variance of k-fold cross-validation. *Journal of Machine Learning Research* **5** 1089–1105.

BURMAN, P. (1989). A comparative study of ordinary cross-validation, v-fold cross-validation and the repeated learning-testing methods. *Biometrika* **76** 503–514.

CHANG, C.-C. and LIN, C.-J. (2011). Libsvm: A library for support vector machines. *ACM Transactions on Intelligent Systems and Technology (TIST)* **2** 1–27.

CORTES, C. and VAPNIK, V. (1995). Support-vector networks. *Machine Learning* **20** 273–297.

DUA, D. and GRAFF, C. (2017). UCI machine learning repository.
URL http://archive.ics.uci.edu/ml

GHAOUI, L. E., VIALLON, V. and RABBANI, T. (2010). Safe feature elimination for the lasso and sparse supervised learning problems. *arXiv preprint arXiv:1009.4219* .

GOLUB, G. H., HEATH, M. and WAHBA, G. (1979). Generalized cross-validation as a method for choosing a good ridge parameter. *Technometrics* **21** 215–223.

HASTIE, T., TIBSHIRANI, R. and FRIEDMAN, J. (2009). *The Elements of Statistical Learning*. Springer Series in Statistics, Springer New York Inc., New York, NY, USA.

HONG, B., ZHANG, W., LIU, W., YE, J., CAI, D., HE, X. and WANG, J. (2019). Scaling up sparse support vector machines by simultaneous feature and sample reduction. *Journal of Machine Learning Research* **20** 1–39.

KARATZOGLOU, A., SMOLA, A., HORNIK, K. and ZEILEIS, A. (2004). kernlab-an s4 package for kernel methods in r. *Journal of Statistical Software* **11** 1–20.

KOHAVI, R. (1995). A study of cross-validation and bootstrap for accuracy estimation and model selection. In *International Joint Conference on Artificial Intelligence*, vol. 14.

MICCHELLI, C. A., XU, Y. and ZHANG, H. (2006). Universal kernels. *Journal of Machine Learning Research* **7** 2651–2667.

MOLINARO, A. M., SIMON, R. and PFEIFFER, R. M. (2005). Prediction error estimation: a comparison of resampling methods. *Bioinformatics* **21** 3301–3307.

OGAWA, K., SUZUKI, Y. and TAKEUCHI, I. (2013). Safe screening of non-support vectors in pathwise svm computation. In *International Conference on Machine Learning*.

PAN, X. and XU, Y. (2018). A novel and safe two-stage screening method for support vector machine. *IEEE Transactions on Neural Networks and Learning Systems* **30** 2263–2274.

PARIKH, N. and BOYD, S. (2014). Proximal algorithms. *Foundations and Trends in Optimization* **1** 127–239.

STEINWART, I. (2001). On the influence of the kernel on the consistency of support vector machines. *Journal of Machine Learning Research* **2** 67–93.

TIBSHIRANI, R. (1996). Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society: Series B (Methodological)* **58** 267–288.

VAPNIK, V. (1995). *The Nature of Statistical Learning Theory*. Springer Science & Business Media.

VAPNIK, V. (1998). *Statistical Learning Theory*. Wiley.

WAHBA, G. (1990). *Spline Models for Observational Data*, vol. 59. SIAM.

WAHBA, G., LIN, Y. and ZHANG, H. (1999). Gacv for support vector machines. *Advances in Neural Information Processing Systems* 297–310.

WAHBA, G. and WOLD, S. (1975). Periodic splines for spectral density estimation: The use of cross validation for determining the degree of smoothing. *Communications in Statistics-Theory and Methods* **4** 125–141.

315 WANG, B. and ZOU, H. (2021). Honest leave-one-out cross-validation for estimating post-tuning generalization
316   error. *Stat* **10**.

317 WANG, B. and ZOU, H. (2022). Fast and exact leave-one-out analysis of large-margin classifiers. *Technometrics*
318   **To appear.**

319 WANG, J., WONKA, P. and YE, J. (2014). Scaling svm and least absolute deviations via exact data reduction.
320   In *International Conference on Machine Learning*.

321 ZHANG, Y. and YANG, Y. (2015). Cross-validation for selecting a model selection procedure. *Journal of*
322   *Econometrics* **187** 95–112.