Learning (Very) Simple Generative Models Is Hard

Sitan Chen
UC Berkeley
sitanc@berkeley.edu

Jerry Li
Microsoft Research
jerrl@microsoft.com

Yuanzhi Li
CMU
yuanzhil@andrew.cmu.edu

Abstract

Motivated by the recent empirical successes of deep generative models, we study the computational complexity of the following unsupervised learning problem. For an unknown neural network $F : \mathbb{R}^d \to \mathbb{R}^{d'}$, let D be the distribution over $\mathbb{R}^{d'}$ given by pushing the standard Gaussian $\mathcal{N}(0, \mathrm{Id}_d)$ through F. Given i.i.d. samples from D, the goal is to output *any* distribution close to D in statistical distance.

We show under the statistical query (SQ) model that no polynomial-time algorithm can solve this problem even when the output coordinates of F are one-hidden-layer ReLU networks with $\log(d)$ neurons. Previously, the best lower bounds for this problem simply followed from lower bounds for *supervised learning* and required at least two hidden layers and $\mathrm{poly}(d)$ neurons [CGKM22, DV21].

The key ingredient in our proof is an ODE-based construction of a compactly supported, piecewise-linear function f with polynomially-bounded slopes such that the pushforward of $\mathcal{N}(0, 1)$ under f matches all low-degree moments of $\mathcal{N}(0, 1)$.

1 Introduction

In recent years, deep generative models such as variational autoencoders, generative adversarial networks, and normalizing flows [GPAM$^+$14, KW13, RM15] have seen incredible success in modeling real world data. These work by learning a parametric transformation (e.g. a neural network) of a simple distribution, usually a standard normal random variable, into a complex and high-dimensional one. The learned distributions have been shown to be shockingly effective at modeling real world data. The success of these generative models begs the following question: when is it possible to learn such a distribution? Not only is this a very natural question from a learning-theoretic perspective, but understanding this may also lead to more direct methods to learn generative models for real data.

More formally, we consider the following problem. Let D be the unknown pushforward distribution over $\mathbb{R}^d$ given by $f(g)$, where $g \sim \mathcal{N}(0, \mathrm{Id})$ is a standard normal Gaussian, and f is an unknown feed-forward neural network with non-linear (typically ReLU) activations. Such distributions naturally arise as the output of many common deep generative models in practice. The learner is given n samples from D, and their goal is to output the description of some distribution which is close to D.

When f is a one-layer network (i.e. of the form $f(g) = \mathrm{ReLU}(Wg)$), there are efficient algorithms for learning the distribution [WDS19, LLDD20]. However, this setting is unsatisfactory in many ways, as one-layer networks lack much of the complex structure that makes these generative models so appealing in practice. Indeed, when the neural network only has a single layer, the resulting distribution is similar to a truncated Gaussian, and one can leverage techniques developed for learning from truncated samples. Notably, this structure disappears even with two-layer neural networks. Even in the two-layer case, despite significant interest, very little is known about how to learn D efficiently.

In fact, a recent line of work suggests that learning neural network pushforwards of Gaussians may be an inherently difficult computational task. Recent results of [DV21, CGKM22] show hardness of *supervised* learning from labeled Gaussian examples under cryptographic asssumptions, and the latter also demonstrates hardness for all statistical query (SQ) algorithms (see Section 1.3 for a more

36th Conference on Neural Information Processing Systems (NeurIPS 2022).

detailed description of related work). These naturally imply hardness in the unsupervised setting (see supplement). However, these lower bound constructions still have their downsides. For one, all of these constructions require at least three layers (i.e. two hidden layers), and so leave open the possibility that efficient learning is possible when the neural network only has one hidden layer. Additionally, the resulting neural networks in these constructions are quite complicated. In particular, the size of the neural networks in these hard instances, that is, the number of hidden nonlinear activations for any output coordinate, must be polynomially large. This begs the natural question:

Can we learn pushforwards of Gaussians under one-hidden-layer neural networks of small size?

1.1 Our Results

We demonstrate strong evidence that despite the simplicity of the setting, this learning task is already computationally intractable. We show there is no polynomial-time statistical query (SQ) algorithm which can learn the distribution of $f(g)$, when $g \sim \mathcal{N}(0, \mathrm{Id})$ and each output coordinate of f is a one-hidden-layer neural network of *logarithmic hidden size*. We formally define the SQ model in Definition 2; we note that it is well-known to capture almost all popular learning algorithms [FGR+17].

Theorem 1.1 (informal, see Theorem 3.1). *For any $d > 0$, and any $C \geq 1$, there exists a family of one-hidden-layer neural networks $\mathcal{F}$ from $\mathbb{R}^d$ to $\mathbb{R}^{d^C}$ so that the following properties hold. For any $f \in \mathcal{F}$, let D_f denote the distribution of $f(g)$, for $g \sim \mathcal{N}(0, \mathrm{Id})$. Then, we have that:*

- *For all $f \in \mathcal{F}$, $\mathrm{d_{TV}}(\mathcal{N}(0, \mathrm{Id}), D_f) = \Omega(1)$,[1]*

- *Every output coordinate of f is a sum of $O(\log d / \log \log d)$ ReLUs, with $\mathrm{poly}(d)$-bounded weights.*

- *Any SQ algorithm which can distinguish between D_f and $\mathcal{N}(0, \mathrm{Id})$ with high probability for all $f \in \mathcal{F}$ requires $d^{\omega(1)}$ time and/or samples.*

In other words, there is a family of one-hidden-layer ReLU networks of logarithmic size whose corresponding pushforwards are statistically very far from Gaussian, yet no efficient SQ algorithm can distinguish them from a Gaussian. Note this implies hardness even of *improperly* learning such pushforwards: not only is it hard to recover the parameters of the network or output a network close to the underlying distribution D_f, but it is hard to learn *any* distribution close to D_f. In contrast, if one ignores issues of computational efficiency, one can easily learn networks in this family with polynomial sample complexity via hypothesis selection, as there are only $\mathrm{poly}(d)$ many parameters specifying any network in this family.

Since such networks are arguably some of the simplest neural networks with more than one layer, this suggests that learning even the most basic deep generative models may already be a very difficult task, at least without additional assumptions. Still, this is by no means the last word in this direction. Given the real world success of deep generative models, a natural and important direction is to identify natural conditions under which we can efficiently learn. We view our results as a first step towards understanding the computational landscape of this important learning problem, and our result *provides evidence that (strong) assumptions need to be made on f for the pushforward $f(g)$ to be efficiently learnable, even in very simple two-layer cases.*

1.2 Our Techniques

Like many recent SQ lower bounds, ours follows the general framework which was introduced in [DKS17] and builds on [FGR+17]. Here one considers the following "non-Gaussian component analysis" task. We will consider a family of distributions, parametrized by a "hidden" direction $v \in \mathbb{S}^{d-1}$, which are Gaussian in every direction orthogonal to the hidden direction but which are very non-Gaussian along the hidden direction.

To formalize this, let D be a distribution D over $\mathbb{R}$ which is known to the learner and far in statistical distance from Gaussian. Given a unit vector v in d dimensions, let P_v^D denote the distribution over $\mathbb{R}^d$ whose projection along v is given by D and whose projection in all directions orthogonal to v is standard Gaussian. Given samples from some unknown distribution over $\mathbb{R}^d$, the goal is to

---

[1] $\mathrm{d_{TV}}$ denotes total variation distance. A lower bound for Wasserstein distance also holds, see supplement.

decide whether the unknown distribution is $\mathcal{N}(0, \mathrm{Id}_d)$ or P_v^D for some v. [DKS17] showed that if D's moments match those of $\mathcal{N}(0,1)$ up to some degree m, then under mild conditions, any SQ algorithm for this task requires at least $d^{\Omega(m)}$ queries (Lemma 3.3).

Suppose one could exhibit a one-hidden-layer ReLU network $f : \mathbb{R}^\ell \to \mathbb{R}$ such that the pushforward $D = f(\mathcal{N}(0, \mathrm{Id}))$ satisfied such properties. Then we can realize P_v^D as a pushforward as follows. Let U be a rotation mapping the first standard basis vector in $\mathbb{R}^d$ to v. Then consider the function $F : \mathbb{R}^{\ell+d-1} \to \mathbb{R}^d$ mapping z to $U \cdot (f(z_1, \ldots, z_\ell), z_{\ell+1}, \ldots, z_{\ell+d-1})$. One can check that every output coordinate of F is computed by a one-hidden-layer ReLU network with size essentially equal to that of f. By [DKS17], we would immediately get the desired SQ lower bound.

The main challenge is thus to construct such a network whose pushforward matches the low-degree moments of $\mathcal{N}(0,1)$. It is not hard to ensure the existence of such f with essentially *infinite* weights (Corollary 4.2 and Lemma 4.4). It is much less clear whether this is possible with *polynomially bounded* weights, and this is our primary technical contribution. We design and analyze a certain ODE which defines a one-parameter family of perturbations to f, such that the low-degree moments of the corresponding pushforwards remain unchanged over time. By evolving along this family over an inverse-polynomial time scale, we obtain a network with polynomially bounded weights whose pushforward matches the low-degree moments of $\mathcal{N}(0,1)$. We defer the details to Section 4.

1.3 Related Work

A full literature review on the theory of learning deep generative models is beyond the scope of this paper (see e.g. the survey of [GSW+21]). For conciseness we cover only the most relevant papers.

Upper bounds. In terms of upper bounds, much of the literature has focused on a different setting, where the goal is to understand when first order dynamics can learn toy generative models [FFGT17, DISZ17, GHP+19, LLDD20, AZL21, JMGL22], which are much simpler than the ones we consider here. For learning pushforwards of Gaussians under neural networks with ReLU activations, algorithms with provable guarantees are only known when the network has no hidden layers [WDS19, LLDD20]. This is in contrast to the supervised setting, where fixed parameter tractable algorithms are known for learning ReLU networks of arbitrary depth [CKM22].

A different line of work seeks to find efficient learning algorithms when the activations are given by low degree polynomials [FFGT17, LD20, CLLZ22]. Arguably the closest to our work is [CLLZ22], which gives polynomial-time algorithms for learning low-degree polynomial transformations of Gaussians, in a smoothed setting. It is a very interesting open question if similar smoothed assumptions can be leveraged to circumvent our lower bound when the activations are ReLU. Unfortunately, these papers heavily leverage the nice moment structure of low-degree Gaussian polynomials, and it is unclear how their techniques can generalize to different activations.

Lower bounds. Much of the literature on lower bounds for learning neural networks has focused on the supervised setting, where a learner is given labeled examples $(x, f(x))$, and the goal is to output a good predictor. There are many lower bounds known in the distribution-free setting [BR92, Vu98, KS09, LSSS14, DV20], however, these do not transfer over to our (unsupervised) setting. When x is Gaussian, the aforementioned work of [CGKM22] derives hardness for learning two-hidden-layer networks with polynomial size for all SQ algorithms, as well as under cryptographic assumptions (see also [DV21]). It is not hard to show (see supplement) that this lower bound immediately implies a lower bound for the unsupervised problem. In the supervised setting, lower bounds are also known against restricted families of SQ [VW19, GGJ+20, DKKZ20, SVWX17], when there are adversarially noisy labels [KK14, DKZ20, GGK20, SZB21], and in discrete settings [Val84, Kha95, AK95, Fel09, CGV15, DGKP20, AAK21], but to our knowledge, these results do not transfer to our setting.

The literature on lower bounds for the unsupervised problem we consider here is much sparser. Besides [DV21, CGKM22], we also mention the recent work of [CLLM22] that studies whether achieving small Wasserstein GAN loss implies distribution learning. A corollary of their results is cryptographic hardness for learning pushforwards of Gaussians under networks with constant depth and polynomial size, but only when the learner is given by a Lipschitz ReLU network discriminator. However, this does not rule out efficient algorithms which do not output such Lipschitz discriminators.

Finally, we remark that the family of hard distributions we construct can be thought of as a close cousin of the "parallel pancakes" construction of [DKS17]. This and slight modifications thereof are mixtures of Gaussians which are known to be computationally hard both in the SQ model [DKS17, BLPR19] and under cryptographic assumptions [BRST21, GVV22].

SQ lower bounds via ODEs. We remark that in a very different context, [DKZ20] also used an ODE to design a moment-matching construction. While our approach draws inspiration from theirs, an important difference is that they use their ODE as a "size reduction" trick to construct a step function $f : \mathbb{R} \to \{\pm 1\}$ with a small number of linear pieces such that $\mathbb{E}_{g \sim \mathcal{N}(0,1)}[f(g)g^k] = 0$ for all small k, while we use our ODE as a "weight reduction" trick to construct a continuous neural network $f : \mathbb{R} \to \mathbb{R}$ with bounded weights such that $\mathbb{E}_{g \sim \mathcal{N}(0,1)}[f(g)^k] = \mathbb{E}_{g \sim \mathcal{N}(0,1}[g^k]$. The form of the moments $\mathbb{E}_{g \sim \mathcal{N}(0,1)}[f(g)g^k]$ they consider is simpler than in our setting, and while they essentially run their ODE to singularity and use non-quantitative facts like the invertibility of a certain Jacobian, we only run our ODE for a finite horizon and need to carefully control the condition number of the Jacobian arising in our setting over this horizon (e.g. Lemma 4.8).

2 Technical Preliminaries

We freely abuse notation and use the same symbols to denote probability distributions, their laws, and their density functions. Given a distribution A over a domain Ω and a function $f : \Omega \to \Omega'$, we let $f(A)$ denote the *pushforward* of A through f, that is, the distribution A' of the random variable $f(z)$ for $z \sim A$. Given distributions p, q for which p is absolutely continuous with respect to q, let $\chi^2(p, q)$ denote their chi-squared divergence. Let $p \star q$ denote the convolution of p and q. Also, we use $\|\cdot\|_p$ to denote ℓ^p norm, omitting the subscript when $p = 2$. $\sigma_{\min}(\cdot)$ denotes minimum singular value. Given an invertible matrix M, we use $M^{-\top}$ to denote the matrix $(M^\top)^{-1} = (M^{-1})^\top$.

Henceforth $\mathbb{E}_g[\cdot]$ will always denote $\mathbb{E}_{g \sim \mathcal{N}(0,\mathrm{Id})}[\cdot]$, where the dimensionality is implicit from context. Let $\gamma_{\sigma^2}(x) \triangleq \frac{1}{\sigma\sqrt{2\pi}} e^{-x^2/(2\sigma^2)}$. Given $S \subset \mathbb{R}$, we use $\gamma_{\sigma^2}(S)$ to denote $\int_{-\infty}^{\infty} \gamma(x) \cdot \mathbb{1}[x \in S]\, dx$. When $\sigma = 1$, we omit the subscript σ^2. We will also use $\gamma^{(d)}(x)$ to denote the density of $\mathcal{N}(0, \mathrm{Id}_d)$.

Definition 1 (One-hidden-layer ReLU networks). *We say that $g : \mathbb{R}^d \to \mathbb{R}$ is a one-hidden-layer ReLU network with size S and W-bounded weights if there exist $w_1, \ldots, w_S \in \mathbb{R}^d$, $b_1, \ldots, b_S \in \mathbb{R}$, and $s_1, \ldots, s_S \in \{\pm 1\}$ for which $g(x) = \sum_{i=1}^{S} s_i \mathrm{ReLU}(\langle w_i, x \rangle + b_i)$ and $\|w_i\|, |b_i| \le W$ for all $i \in S$.[2] Here $\mathrm{ReLU}(z) \triangleq \max(0, z)$. Given $f : \mathbb{R}^d \to \mathbb{R}^{d'}$, together with a distribution A over $\mathbb{R}^d$, we say that $f(A)$ is a one-hidden-layer ReLU network pushforward of A with size S and W-bounded weights if each coordinate of f is of this form.*

Fact 2.1 ([GMSR20]). *If $V \in \mathbb{R}^{n \times n}$ is a Vandermonde matrix with nodes $z_1, \ldots, z_n$, that is, $V_{i,j} = z_j^{i-1}$, and $\{z_i\}$ are ζ-separated, then $\sigma_{\min}(V) \ge \frac{1}{n} \cdot \Omega(\zeta)^{n-1}$.*

Theorem 2.2 (Peano's existence theorem, see e.g. Theorem 2.1 from [Har02]). *For $T, r > 0$ and $y_0 \in \mathbb{R}^n$, let $B \subset \mathbb{R} \times \mathbb{R}^n$ be the parallelepiped consisting of (t, y) for which $0 \le t \le T$ and $\|y - y_0\|_\infty \le r$. If $f : B \to \mathbb{R}$ is continuous and satisfies $|f(t, y)| \le M$ for all $(t, y) \in B$, then the initial value problem $\{y'(t) = f(t, y)$ and $y(0) = y_0\}$ has a solution over $t \in [0, \min(T, r/M)]$.*

Definition 2 (Statistical queries). *Given distribution D over $\mathbb{R}^d$ and parameters $\tau, t > 0$, a $\mathrm{STAT}(\tau)$ oracle takes in any query of the form $f : \mathbb{R}^d \to [-1, 1]$ and outputs a value from $[\mathbb{E}_{x \sim D}[f(x)] - \tau, \mathbb{E}_{x \sim D}[f(x)] + \tau]$, while a $\mathrm{VSTAT}(t)$ oracle takes in any query of the form $f : \mathbb{R}^d \to [0, 1]$ and outputs a value from $[\mathbb{E}_{x \sim D}[f(x)] - \tau, \mathbb{E}_{x \sim D}[f(x)] + \tau]$ for $\tau = \max(1/t, \sqrt{\mathbb{V}_{x \sim D}[f(x)]/t})$.*

3 Statistical Query Lower Bound

Our main result is the following lower bound in this model for learning ReLU network pushforwards:

Theorem 3.1. *Let $d \in \mathbb{N}$ be sufficiently large. Any SQ algorithm which, given SQ access to an arbitrary one-hidden-layer ReLU network pushforward of $\mathcal{N}(0, \mathrm{Id}_d)$ of size $O(\log d / \log \log d)$ with $\mathrm{poly}(d)$-bounded weights, outputs a distribution which is $O(1)$-close in $d_{\mathrm{TV}}(\cdot)$ must make at least $d^{\Omega(\log d / \log \log d)}$ queries to either $\mathrm{STAT}(\tau)$ or $\mathrm{VSTAT}(1/\tau^2)$ for $\tau = d^{-\Omega(\log d / \log \log d)}$.*

---

[2] In practice, one can think of S as the number of channels, e.g. in a deconvolution network.

Our proof will invoke the following key technical result whose proof we defer to Section 4. It exhibits a one-hidden-layer ReLU network $f : \mathbb{R}^2 \to \mathbb{R}$ with bounded weights under which the pushforward of $\mathcal{N}(0,1)$ matches the low-degree moments of $\mathcal{N}(0,1)$ to arbitrary precision, in addition to some other technical conditions that we need to formally establish our statistical query lower bound:

Theorem 3.2. *Fix any odd m and $\nu, \sigma < 1$. There is a one-hidden-layer ReLU network $f^* : \mathbb{R}^2 \to \mathbb{R}$ of size $O(m)$ with weights at most $m^{O(m)}$ for which the pushforward $D \triangleq f^*(\mathcal{N}(0, \mathrm{Id}))$ satisfies*

1. $|\mathbb{E}_{x \sim D}[x^k] - \mathbb{E}_{g \sim \mathcal{N}(0,1)}[g^k]| < \nu$ for all $k = 1, \ldots, m$

2. $\chi^2(D, \mathcal{N}(0,1)) \leq \exp(O(m))/\sigma$

3. $\mathrm{d}_{\mathrm{TV}}(P_v^D, P_{v'}^D) \geq 1 - 2\sigma \log(1/\sigma) - m^{-\Omega(m)}$ for any $v, v' \in \mathbb{S}^{d-1}$ satisfying $|\langle v, v' \rangle| \geq 1/2$.

The rest of the proof of our lower bound will then follow the framework introduced in [DKS17] and subsequently generalized in [DK20]. First, given a distribution D over $\mathbb{R}$ and $v \in \mathbb{S}^{d-1}$, let P_v^D denote the distribution over $\mathbb{R}^d$ with density $P_v^D(x) = D(\langle v, x \rangle) \cdot \gamma^{(d-1)}(x - \langle v, x \rangle v)$, that is, the distribution which is given by D in the direction v and is given by $\mathcal{N}(0, \mathrm{Id} - vv^\top)$ orthogonal to v. We use the following generic statistical query lower bound about learning such distributions, a proof of which we include in the supplement for completeness.

Lemma 3.3. *Let $m \in \mathbb{N}$ and $0 < C < 1/2$. Let D be a distribution such that 1) $\chi^2(D, \mathcal{N}(0,1))$ is finite, and 2) $|\mathbb{E}_{x \sim D}[x^k] - \mathbb{E}_{g \sim \mathcal{N}(0,1)}[g^k]| \leq \Omega(d)^{-(m+1)(1/4 - C/2)} \sqrt{\chi^2(D, \mathcal{N}(0,1))}$ for all $k = 1, \ldots, m$.*

Consider the set of distributions $\{P_v^D\}_{v \in \mathbb{S}^{d-1}}$ for $d \geq m^{\Omega(1/C)}$. If there is some $\epsilon > 0$ for which $\mathrm{d}_{\mathrm{TV}}(P_v^D, P_{v'}^D) > 2\epsilon$ whenever $|\langle v, v' \rangle| \leq 1/2$, then any SQ algorithm which, given SQ access to P_v^D for an unknown $v \in \mathbb{S}^{d-1}$, outputs a hypothesis Q with $\mathrm{d}_{\mathrm{TV}}(Q, P_v^D) \leq \epsilon$ needs at least d^{m+1} queries to $\mathrm{STAT}(\tau)$ or to $\mathrm{VSTAT}(1/\tau^2)$ for $\tau \triangleq O(d)^{-(m+1)(1/4 - C/2)} \cdot \sqrt{\chi^2(D, \mathcal{N}(0,1))}$.

Proof of Theorem 3.1. By Theorem 3.2 applied with sufficiently large odd m and sufficiently small σ, there exists a distribution $D = f^*(\mathcal{N}(0, \mathrm{Id}_2))$ over $\mathbb{R}$ for $f^* : \mathbb{R}^2 \to \mathbb{R}$ of size $O(m)$ with $m^{O(m)}$-bounded weights satisfying the hypotheses of Lemma 3.3 for $\epsilon = 0.49$, and $\chi^2(D, \mathcal{N}(0,1)) \leq \exp(O(m))$ (note that we can absorb the $1/\sigma$ factor in Theorem 3.2 into $\exp(O(m))$ because we can take $\sigma = \exp(-\Theta(m))$). As long as $m \leq d^{O(C)}$, we conclude that an SQ algorithm for learning any distribution from $\{P_v^D\}_{v \in \mathbb{S}^{d-1}}$ to total variation distance $1/4$ must make at least d^{m+1} queries to $\mathrm{STAT}(\tau)$ or $\mathrm{VSTAT}(1/\tau^2)$ for $\tau \triangleq O(d)^{-(m+1)(1/4 - C/2)} \cdot \exp(O(m))$. By taking $m = \Theta(\log d / \log \log d)$, we ensure that $m^{O(m)} \leq \mathrm{poly}(d)$. We're done by taking C in Lemma 3.3 to be $C = 1/4$.

The proof of the theorem is complete upon noting that any distribution P_v^D can be implemented as a pushforward of $\mathcal{N}(0, \mathrm{Id}_{d+1})$ under a one-hidden-layer ReLU network $F_v : \mathbb{R}^{d+1} \to \mathbb{R}^d$ of size $O(\log d / \log \log d)$ with $\mathrm{poly}(d)$-bounded weights. Let $U \in O(d)$ be a rotation mapping the first standard basis vector in $\mathbb{R}^d$ to v. Then for $F_v(z_1, \ldots, z_{d+1}) \triangleq U(f^*(z_1, z_2), z_3, \ldots, z_{d+1})$ we have that $F_v(\mathcal{N}(0, \mathrm{Id}_{d+1})) = P_v^D$ as desired. Furthermore, note that every output coordinate of $F_v(z_1, \ldots, z_{d+1})$ is a one-hidden-layer ReLU network of the form $\alpha f^*(z_1, z_2) + \langle u, (z_3, \ldots, z_{d+1}) \rangle$ for some vector $(\alpha, u) \in \mathbb{R}^d$. Note that the size of this network is two plus that of f^*, and its weights are also upper bounded by $\mathrm{poly}(d)$, so F_v's output coordinates are of size $O(\log d / \log \log d)$. $\qquad\square$

Remark 3.4. Theorem 1.1 was stated with output dimension polynomially bigger than input dimension, whereas in our construction, the output dimension (d) is less than the input dimension ($d + 1$). One can get the former by a padding argument (i.e. by duplicating output coordinates) to give a generator with arbitrarily large polynomial stretch and such that the $d^{\log d / \log \log d}$ lower bound still applies.

4 Moment-Matching Construction

In this section we prove Theorem 3.2, the main technical ingredient in the proof of Theorem 3.1.

4.1 Moment-Matching With Unbounded Weights

We begin by making the simple observation that for one-hidden-layer networks with *unbounded weights*, it is easy to construct networks such that the pushforward of $\mathcal{N}(0,1)$ under these networks

matches the moments of $\mathcal{N}(0,1)$. The starting point for this is the following standard construction. Roughly speaking, it gives a set of well-separated points on the real line and a distribution over them which matches low-order moments with $\mathcal{N}(0,1)$.

Lemma 4.1 (Lemma 4.3 from [DKS17]). *For any $m \in \mathbb{N}$, there exist weights $\lambda_1, \ldots, \lambda_m \geq 0$ and points $h_1, \ldots, h_m \in \mathbb{R}$ for which*

1. *(Moments match) $\sum_{i=1}^m \lambda_i h_i^k = \mathbb{E}_g[g^k]$ for all $k = 0, \ldots, 2m-1$ (note that the special case of $k = 0$ implies $\sum_{i=1}^m \lambda_i = 0$).*

2. *(Points symmetric about origin) $h_1 \leq \cdots \leq h_m$ and $h_i = -h_{m-i+1}$ for all $1 \leq i \leq m$.*

3. *(Weights symmetric) $\lambda_1 \leq \cdots \leq \lambda_{\lceil m/2 \rceil}$ and $\lambda_i = \lambda_{m-i+1}$.*

4. *(Points bounded and separated) $\Omega(1/\sqrt{m}) \leq |h_i| \leq O(\sqrt{m})$ for all $1 \leq i \leq m$ and $\{h_i\}$ are $\Omega(1/\sqrt{m})$-separated.*

5. *(Weights not too small) $\min_i \lambda_i \geq e^{-cm}$ for an absolute constant $c > 0$.*

6. *(Central point and weight) If m is odd, then $h_{(m+1)/2} = 0$ and $\lambda_{(m+1)/2} = \Theta(1/\sqrt{m})$.*

This immediately implies that there exists a *discontinuous* piecewise linear function $f : \mathbb{R} \to \mathbb{R}$ for which the pushforward $f(\mathcal{N}(0,1))$ matches the low-degree moments of $\mathcal{N}(0,1)$. The reason is that we can take f to be a step function which takes on values given by the positions of the points $h_1, \ldots, h_m$ in Lemma 4.1. We can then take the steps to have lengths such that the probability that a standard Gaussian input to f falls under the step given by some h_i is precisely λ_i (see Figure 1).

Corollary 4.2. *For any $m \in \mathbb{N}$, there is a partition of $\mathbb{R}$ into disjoint intervals $I_1, \ldots, I_m$, along with a choice of scalars $h_1, \ldots, h_m$, such that the step function $f : \mathbb{R} \to \mathbb{R}$ given by $f(z) = \sum_{i=1}^m h_i \cdot \mathbb{1}[z \in I_i]$ satisfies $\mathbb{E}_{x \sim f(\mathcal{N}(0,1))}[x^k] = \mathbb{E}_{g \sim \mathcal{N}(0,1)}[g^k]$ for all $k = 0, \ldots, 2m-1$.*

Proof. Let $\{\lambda_i\}, \{h_i\}$ be as in Lemma 4.1. As $\sum_i \lambda_i = 1$, there are intervals $\mathbb{R} = I_1 \sqcup \cdots \sqcup I_m$ for which $\gamma(I_i) = \lambda_i$. As $\mathbb{E}_{x \sim f(\mathcal{N}(0,1))}[x^k] = \sum_i \lambda_i h_i^k$, the claim follows by Part 1 of Lemma 4.1. $\square$

By infinitesimally perturbing f in Corollary 4.2, we can ensure $f(\mathcal{N}(0,1))$ still *approximately* matches the low-degree moments of $\mathcal{N}(0,1)$ to arbitrary precision and that the linear pieces of f have finite slopes, though some slopes will be arbitrarily large. The new function can thus be represented as a one-hidden-layer ReLU network, but unfortunately its weights will be arbitrarily large. The key challenge in the sequel is to design a better perturbation so the resulting f has *polynomially bounded* slopes yet is such that $f(\mathcal{N}(0,1))$ matches the low-degree moments of $\mathcal{N}(0,1)$.

4.2 Bump Construction

Before we describe our perturbation scheme, we slightly modify the construction in Corollary 4.2. In place of a step function, we will consider a certain sum of *bump functions*, illustrated in Figure 2.

Definition 3 (Bump functions). *Given $w, \epsilon > 0$ and $h, c \in \mathbb{R}$, define $T_c^{w,h,\epsilon} : \mathbb{R} \to \mathbb{R}$ by*

$$
T_c^{w,h,\epsilon}(z) = \begin{cases} \frac{h}{\epsilon}(z - c + \epsilon + w) & \text{if } z \in [c - \epsilon - w, c - w] \\ h & \text{if } z \in [c - w, c + w] \\ -\frac{h}{\epsilon}(z - c - \epsilon - w) & \text{if } z \in [c + w, c + \epsilon + w] \\ 0 & \text{otherwise.} \end{cases}
$$

Because these are piecewise linear functions, we can implement them as one-hidden-layer ReLU networks:

Fact 4.3. *Given $w, \epsilon > 0$ and $h, c \in \mathbb{R}$, $T_c^{w,h,\epsilon}$ can be implemented as a one-hidden-layer ReLU network with size 4 and W-bounded weights for $W \leq \frac{h}{\epsilon} \max(1, |c| + \epsilon + w)$.*

In the next lemma, we show that we can replace the construction in Corollary 4.2, given by a step function, with a sum of well-separated bumps with infinitesimally small ϵ parameter. This will be the instance on which we will apply our perturbation scheme.

Lemma 4.4. *For any odd m and $\nu < 1$, there are centers $c_1 \leq \cdots \leq c_{m-1}$, widths $w_1, \ldots, w_{m-1} > 0$, heights $h_1 \leq \cdots \leq h_{m-1} \in \mathbb{R}$, and parameter $\bar{\epsilon} > 0$ for which the following holds. Define the function $f : \mathbb{R} \to \mathbb{R}$ (see Figure 1) by $f(z) \triangleq \sum_{i=1}^{m-1} T_{c_i}^{w_i, h_i, \epsilon}(z)$ for any $0 \leq \epsilon < \bar{\epsilon}$. Then f satisfies*

Figure 1: Left: construction from Lemma 4.1 gives rise to step function in Corollary 4.2. Right: removing central spike gives rise to sum of bumps in Lemma 4.4

1. *(Bumps are well-separated) For all $1 \leq i < m - 1$, $c_i + m^{-3/2} \leq c_{i+1}$.*

2. *(Moments match) $|\mathbb{E}_{x \sim f(\mathcal{N}(0,1))}[x^k] - \mathbb{E}_g[g^k]| < \nu$ for all $k = 1, \ldots, 2m - 1$.*

3. *(Symmetricity) $w_i = w_{m-i}$, $h_i = -h_{m-i}$, and $c_i = -c_{m-i}$ for all $1 \leq i < m$.*

4. *(Bounded, separated heights) $\Omega(1/\sqrt{m}) \leq |h_i| \leq O(\sqrt{m})$ for all $1 \leq i < m$, and $\{h_i\}$ are $\Omega(1/\sqrt{m})$-separated.*

5. *(Intervals not too thin) $\min_i \gamma([c_i - w_i, c_i + w_i]) \geq e^{-cm}$ for an absolute constant $c > 0$.*

6. *(Bounded endpoints) $|c_i| + w_i \leq O(\log m)$ for all $1 \leq i < m$.*

Proof sketch, see supplement. The idea is that when m is odd, one of the points h_i from Lemma 4.1 is zero and does not contribute to the moments. We want to design f for which $f(\mathcal{N}(0,1))$ is the mixture of point masses given by removing this point and its corresponding weight. We take f to have one linear piece for every one of the remaining point masses. By Part 6 of Lemma 4.1, these points have total weight $1 - \Theta(1/\sqrt{m})$, so by anticoncentration we can arrange these linear pieces to be $\sim m^{-3/2}$-separated, satisfying Part 1. The remaining parts follow easily from Lemma 4.1. □

Unfortunately the slopes in the function constructed in Lemma 4.4 are arbitrarily large if ν is arbitrarily small. The issue still remains of how to get a continuous piecewise-linear function whose slopes are *polynomially bounded* so that the corresponding ReLU network has polynomially bounded weights. As we illustrate next however, the $\Omega(m^{-3/2})$ spacing between the bumps in the definition of f in Lemma 4.4 gives us sufficient "room" to carefully perturb f to achieve this goal.

4.2.1 Some Estimates for Bump Moments

For convenience, define $M_{c,k}^{w,h,\epsilon} \triangleq \mathbb{E}_g[T_c^{w,h,\epsilon}(g)^k]$. We conclude this subsection by collecting some useful bounds for this quantity. As we verify in the supplement, $M_{c,k}^{w,h,\epsilon}$ is continuously differentiable with respect to ϵ when $\epsilon > 0$. Additionally, we have the following (see supplement):

Lemma 4.5. $\frac{\partial M_{x,k}^{w,h,\epsilon}}{\partial h} = \frac{k}{h} M_{x,k}^{w,h,\epsilon}$.

Lemma 4.6. *For any $w \geq 0$, $c, h, h' \in \mathbb{R}$, $\epsilon' \geq \epsilon \geq 0$, and even $k \in \mathbb{N}$, we have the bound $|M_{c,k}^{w,h',\epsilon'} - M_{c,k}^{w,h,\epsilon}| \leq h^k \left(|(h'/h)^k - 1| + \epsilon' - \epsilon\right)$. In particular, this implies that $\left|\frac{\partial M_{c,k}^{w,h,\epsilon}}{\partial \epsilon}\right| \leq h^k$.*

4.3 ODE-Driven Perturbation

Denote the parameters of the function constructed in Lemma 4.4 by $\{(h_i(0), w_i, c_i)\}_{1 \leq i < m}$. We also define $\epsilon(0)$ to be some arbitrarily small positive quantity satisfying $\epsilon(0) \leq \bar{\epsilon}$ for $\bar{\epsilon}$ from Lemma 4.4.

We will design an ODE whose solution specifies a one-parameter family of functions

$$f_t \triangleq \sum_{i=1}^{m-1} T_{c_i}^{w_i, h_i(t), \epsilon(t)} \tag{1}$$

that arise from gradually perturbing the function from Lemma 4.4. Roughly speaking, starting at $h_i(0)$ and $\epsilon(0)$ for all $1 \leq i < m$, perturbing the function along this one-parameter family will correspond to keeping the widths w_i and centers c_i of the bumps fixed, increasing the ϵ parameter of every bump at unit speed, and evolving the heights $h_i(t)$ in such a way that the moments of the pushforward of $\mathcal{N}(0,1)$ under f_t remain constant in t for all $0 \leq t \leq T$. We illustrate this evolution

in Figure 2. Here T is some horizon which is at least inverse-polynomially large but smaller than $m^{-3/2}$ so that the "edges" $c_i \pm (\epsilon(t) + w_i)$ of the bumps don't collide with each other (this is where we make crucial use of Part 1 of Lemma 4.4). At the end of this horizon, we want to show that the heights will not have changed too much, whereas the bumps now have ϵ parameter given by inverse-polynomially large T. This will imply that f_T has polynomially bounded slopes as desired.

Figure 2: Evolution of one of the $m-1$ bumps constituting f_t

As the odd moments of $f_0(\mathcal{N}(0,1))$ vanish and the parameters $\{(h_i(0), w_i, c_i)\}_{1 \leq i < m}$ satisfy the symmetry properties from Part 3 of Lemma 4.4, it is easy to ensure that the odd moments of $f_t(\mathcal{N}(0,1))$ also vanish: simply take $h_i(t) = -h_{m-i}(t)$ for all $1 \leq i \leq m-1$.

We thus focus on evolving $(h_1(t), \ldots, h_{(m-1)/2}(t))$. For convenience denote this by $\mathbf{h}(t)$. Define the *moment vector* $\mu : \mathbb{R}^{(m-1)/2} \times \mathbb{R} \to \mathbb{R}^{m-1}$ by

$$\mu(\mathbf{h}, \epsilon) \triangleq \left\{ \sum_{i=1}^{(m-1)/2} M_{c_i, 2\ell}^{w_i, h_i, \epsilon} \right\}_{1 \leq \ell \leq (m-1)/2}$$

For any fixed $(\mathbf{h}, \epsilon)$ in a small neighborhood of $(\mathbf{h}(0), \epsilon(0))$, we want to show there is a direction $v \in \mathbb{R}^{(m-1)/2}$ such that the directional derivative of μ in the direction $w \triangleq (v_1, \ldots, v_{(m-1)/2}, 1)$ is zero. The constraint that the directional derivative $\nabla_w \mu(\mathbf{h}, \epsilon)$ vanishes specifies a linear system in v:

$$\sum_{i=1}^{(m-1)/2} v_i \cdot \frac{\partial M_{c_i, 2\ell}^{w_i, h_i, \epsilon}}{\partial h_i} = -\sum_{i=1}^{(m-1)/2} \frac{\partial M_{c_i, 2\ell}^{w_i, h_i, \epsilon}}{\partial \epsilon} \quad \forall \, 1 \leq \ell \leq (m-1)/2. \tag{2}$$

Recalling Lemma 4.5, we can rewrite this as

$$\sum_{i=1}^{(m-1)/2} v_i \cdot \frac{2\ell}{h_i} M_{c_i, 2k}^{w_i, h_i, \epsilon} = -\sum_{i=1}^{(m-1)/2} \frac{\partial M_{c_i, 2\ell}^{w_i, h_i, \epsilon}}{\partial \epsilon} \quad \forall \, 1 \leq \ell \leq (m-1)/2.$$

To express this more compactly, define $\mathbf{b}(\epsilon) \in \mathbb{R}^{(m-1)/2}$ and $Z(\mathbf{h}, \epsilon)$ by

$$\mathbf{b}(\epsilon)_\ell \triangleq -\sum_{i=1}^{(m-1)/2} \frac{\partial M_{c_i, 2\ell}^{w_i, h_i, \epsilon}}{\partial \epsilon} \qquad \text{and} \qquad Z(\mathbf{h}, \epsilon)_{i,\ell} \triangleq M_{c_i, 2\ell}^{w_i, h_i, \epsilon_i}. \tag{3}$$

Also define the matrices $A(\mathbf{h}) \triangleq \operatorname{diag}(1/h_1, \ldots, 1/h_{(m-1)/2})$ and $B \triangleq \operatorname{diag}(2, 4, \ldots, m-1)$. Then (2) is equivalent to $v^\top \cdot A(\mathbf{h}) Z(\mathbf{h}, \epsilon) B = \mathbf{b}(\epsilon)^\top$. Provided $A(\mathbf{h}) Z(\mathbf{h}, \epsilon) B$ is invertible, the natural choice for v would thus be $v = B^{-1} Z(\mathbf{h}, \epsilon)^{-\top} A(\mathbf{h})^{-1} \cdot \mathbf{b}(\epsilon)$. Therefore, defining

$$w(t, \mathbf{h}) \triangleq \left(B^{-1} Z(\mathbf{h}, \epsilon(0) + t)^{-\top} A(\mathbf{h})^{-1} \cdot \mathbf{b}(\epsilon(0) + t), 1 \right), \tag{4}$$

we consider the following initial value problem

$$\mathbf{h}'(t) = w(t, \mathbf{h}(t)) \qquad \text{and} \qquad \mathbf{h}(0) = (h_1(0), \ldots, h_{(m-1)/2}(0)). \tag{5}$$

Note that if we had a solution $\mathbf{h}(t)$ to (5) for $t \in [0, T]$ for some horizon T, then we would have

$$\frac{\partial}{\partial t} \mu(\mathbf{h}(t), t) = \left(\frac{\partial}{\partial (\mathbf{h}(t), t)} \mu(\mathbf{h}(t), t) \right) \cdot \mathbf{h}'(t) = \nabla_{w(t)} \mu(\mathbf{h}(t), t) = \mathbf{0}, \tag{6}$$

implying that the low-degree moments of f_t defined in (1) are constant in t as desired.

4.4 Existence and Boundedness of $\mathbf{h}(t)$

To carry out the strategy in Section 4.3, we must establish that (1) a solution to the initial value problem (5) exists over a non-negligible horizon T, and (2) the entries of $\mathbf{h}(t)$ do not explode in t.

For both of these, we need to show that the matrix $Z(\mathbf{h}, \epsilon)$ is invertible or, more specifically, well-conditioned for $(\mathbf{h}, \epsilon)$ in a neighborhood of $(\mathbf{h}(0), \epsilon(0))$. We first establish this at time $t = 0$ by relating $Z(\mathbf{h}(0), \epsilon(0))$ to a certain Vandermonde matrix and appealing to Fact 2.1, see supplement:

Lemma 4.7. $\sigma_{\min}(Z(\mathbf{h}(0), \epsilon(0))) \geq m^{-Cm}$ for an absolute constant $C > 0$.

Thus, for $(\mathbf{h}, \epsilon)$ in a neighborhood of $(\mathbf{h}(0), \epsilon(0))$, $Z(\mathbf{h}, \epsilon)$ is well-conditioned (see supplement).

Lemma 4.8. Let $C > 0$ be from Lemma 4.7. For any $(\mathbf{h}, \epsilon)$ satisfying $\|\mathbf{h} - \mathbf{h}(0)\|_{\infty} \leq m^{-C'm}$ and $0 \leq \epsilon - \epsilon(0) \leq m^{-C'm}$ for sufficiently large absolute constant $C' > 0$, $\sigma_{\min}(Z(\mathbf{h}, \epsilon)) \geq m^{-Cm}/2$.

To establish Property (1), we must first verify that $w(t, \mathbf{h})$ is continuous (see supplement for proof):

Lemma 4.9. Let $C' > 0$ be the absolute constant from Lemma 4.8. Then the function $w(t, \mathbf{h})$ defined in (4) is continuous with respect to both t and $\mathbf{h}$ for $t \leq m^{-C'm}$ and $\|\mathbf{h} - \mathbf{h}(0)\|_{\infty} \leq m^{-C'm}$.

Lastly, we must show that under the hypotheses of Lemma 4.9, $\|w(t, \mathbf{h})\|_{\infty}$ is not too large. This implies Property (1) by Theorem 2.2 and Property (2) because $\|\mathbf{h}'(t)\|_{\infty} = \|w(t, \mathbf{h}(t))\|_{\infty}$:

Lemma 4.10. Let $C' > 0$ be the absolute constant from Lemma 4.8. If $t \leq m^{-C'm}$ and $\|\mathbf{h} - \mathbf{h}(0)\|_{\infty} \leq m^{-C'm}$, then $\|w(t, \mathbf{h})\|_{\infty} \leq m^{C''m}$ for some absolute constant $C'' > 0$.

Proof. By the second part of Lemma 4.6, every entry of $\mathbf{b}(\epsilon(0) + t)$ is at most $\frac{m-1}{2} \cdot (\max_i h_i)^{m-1} \leq \frac{m-1}{2} \cdot (O(\sqrt{m}) + m^{-C'm})^{m-1} \leq m^{O(m)}$, where in the penultimate step we used Part 4 of Lemma 4.4 and our hypothesis on $\mathbf{h}$. Note that $\sigma_{\min}(Z(\mathbf{h}, \epsilon(0) + t)) \geq m^{-Cm}/2$ by Lemma 4.8, $\sigma_{\min}(B) \geq 2$, and $\sigma_{\min}(A(\mathbf{h})) \geq \min_i 1/h_i \geq \Omega(\sqrt{m})$ by Part 4 of Lemma 4.4 and our hypothesis on $\mathbf{h}$. We conclude that $B^{-1}Z(\mathbf{h}, \epsilon(0) + t)^{-\top}A(\mathbf{h})^{-1} \cdot \mathbf{b}(\epsilon(0) + t)$ has L_{∞} norm at most $m^{C''m}$ for some absolute constant $C'' > 0$, so $\|w(t, \mathbf{h})\|_{\infty} \leq m^{C''m}$ as claimed. $\square$

We are now ready to put these ingredients together to prove the key lemma for showing Theorem 3.2:

Lemma 4.11. Fix any odd $m \in \mathbb{N}$ and any $0 < \nu < 1$. There is a one-hidden-layer ReLU network $f : \mathbb{R} \to \mathbb{R}$ of size $O(m)$ with weights at most $m^{O(m)}$ for which the pushforward $D = f(\mathcal{N}(0,1))$ satisfies $|\mathbb{E}_{x \sim D}[x^k] - \mathbb{E}_{g \sim \mathcal{N}(0,1)}[g^k]| < \nu$ for all $k = 1, \ldots, m$.

Proof. Define the parallelepiped B of pairs $(t, \mathbf{h})$ for which $0 \leq t \leq m^{-C'm}$ and $\|\mathbf{h} - \mathbf{h}(0)\|_{\infty} \leq m^{-C'm}$. By Lemma 4.9, w is continuous over B. By Lemma 4.10, $\|w(t, \mathbf{h})\|_{\infty} \leq m^{C''m}$.

By Theorem 2.2, the initial value problem in (5) has a solution $\mathbf{h}(t)$ over $t \in [0, T]$ for $T = m^{-(C'+C'')m}$. Furthermore, because $\|\mathbf{h}'(t)\|_{\infty} = \|w(t, \mathbf{h}(t))\|_{\infty} \leq m^{C''m}$, we conclude that $\frac{1}{T}\|\mathbf{h}(T)\|_{\infty} \leq m^{C''m}$. The slopes of the bumps $T_{c_i}^{w_i, h_i(T), \epsilon(0) + T}$ are therefore bounded by $m^{C''m}$.

For $(m-1)/2 < i \leq m-1$, define $h_i(T) = -h_{m-i}(T)$ and consider the one-parameter family of functions $f_t \triangleq \sum_{i=1}^{m-1} T_{c_i}^{w_i, h_i(t), \epsilon(0) + t}$. Because $c_i = c_{m-i}$ and $w_i = w_{m-i}$ for all $1 \leq i < m$ by Part 3 of Lemma 4.4, we conclude that the odd moments of $f_T(\mathcal{N}(0,1))$ all vanish. As for the even moments, because $\frac{\partial}{\partial t}\mu(\mathbf{h}(t), t) = \mathbf{0}$ by (6), we conclude that the even moments of $f_T(\mathcal{N}(0,1))$ up to degree $m-1$ agree with those of $f_0(\mathcal{N}(0,1))$. So because $D_0 = f_0(\mathcal{N}(0,1))$ satisfies $|\mathbb{E}_{x \sim D}[x^k] - \mathbb{E}_{g \sim \mathcal{N}(0,1)}[g^k]| < \nu$ for $1 \leq k \leq m$ by Part 2 of Lemma 4.4, the same holds for f_T.

As the endpoints of the intervals supporting the bumps are bounded by $O(\log m)$, Fact 4.3 implies f_T is a one-hidden-layer network with size $O(m)$ and $m^{O(m)}$-bounded weights. $\square$

4.5 Proof of Theorem 3.2

Having designed a pushforward given by a one-hidden-layer network with $m^{O(m)}$-bounded weights satisfying the first part of Theorem 3.2, we now modify this to also satisfy the remaining two parts.

To do this, we convolve a suitable scaling of the pushforward by a thin Gaussian. The following lemmas show that the moments continue to match, the chi-squared divergence between the result and $\mathcal{N}(0,1)$ is bounded, and the third part of Theorem 3.2 is satisfied (see supplement for proof):

Lemma 4.12. *Let D be any symmetric distribution for which $|\mathbb{E}_{x \sim D}[x^k] - \mathbb{E}_g[g^k]| < \nu$ for all $1 \le k < m$. For any $c \in \mathbb{R}$ let $c \cdot D$ denote the distribution obtained by rescaling D by a factor of c. Then $D' \triangleq \sqrt{1 - \sigma^2} \cdot D \star \mathcal{N}(0, \sigma^2)$ satisfies $|\mathbb{E}_{x \sim D'}[x^k] - \mathbb{E}_g[g^k]| < \nu$ for all $1 \le k < m$.*

Lemma 4.13. *For any density A on $[-R, R]$ and $\sigma \le 1/2$, $\chi^2(A \star \mathcal{N}(0, \sigma^2), \mathcal{N}(0, 1)) \le e^{O(R^2)}/\sigma$.*

Lemma 4.14. *Let $D = f(\mathcal{N}(0, 1))$ be from Lemma 4.11, and define $D' \triangleq \sqrt{1 - \sigma^2} \cdot D \star \mathcal{N}(0, \sigma^2)$. Then for any $v, v' \in \mathbb{S}^{d-1}$ satisfying $|\langle v, v' \rangle| \le 1/2$, $d_{\mathrm{TV}}(P_v^{D'}, P_{v'}^{D'}) \ge 1 - 2\sigma \log(1/\sigma) - m^{-\Omega(m)}$.*

Proof of Theorem 3.2. Let f be the function constructed in Lemma 4.11 and define $f^* : \mathbb{R}^2 \to \mathbb{R}$ by $f^*(z_1, z_2) = \sqrt{1 - \sigma^2} f(z_1) + \sigma z_2$. Note that $f^*(\mathcal{N}(0, \mathrm{Id}))$ is exactly $\sqrt{1 - \sigma^2} f(\mathcal{N}(0, 1)) \star \mathcal{N}(0, \sigma^2)$, so the three parts of Theorem 3.2 follow immediately from Lemmas 4.12, 4.13, and 4.14 respectively. $\qquad\qquad\square$

Conclusion.   In this work we established that learning one-hidden-layer ReLU network pushforwards of logarithmic size is hard for efficient SQ algorithms. Given that SQ algorithms capture a large family of efficient algorithms, this suggests that this basic learning problem, despite its simple description, is computationally intractable. Of course, our lower bound construction is worst case, and so it is unlikely to actually arise in practice. A natural open question, as mentioned previously, is to come up with natural assumptions (for instance, smoothed analysis settings like in [CLLZ22]) that allow us to circumvent this lower bound, and to achieve efficient learning algorithms.

References

[AAK21]   Naman Agarwal, Pranjal Awasthi, and Satyen Kale. A deep conditioning treatment of neural networks. In *Algorithmic Learning Theory*, pages 249–305. PMLR, 2021.

[AK95]   Dana Angluin and Michael Kharitonov. When won't membership queries help? *Journal of Computer and System Sciences*, 50(2):336–355, 1995.

[AZL21]   Zeyuan Allen-Zhu and Yuanzhi Li. Forward super-resolution: How can gans learn hierarchical generative models for real-world distributions. *arXiv preprint arXiv:2106.02619*, 2021.

[BLPR19] Sébastien Bubeck, Yin Tat Lee, Eric Price, and Ilya Razenshteyn. Adversarial examples from computational constraints. In *International Conference on Machine Learning*, pages 831–840. PMLR, 2019.

[BR92]   Avrim L Blum and Ronald L Rivest. Training a 3-node neural network is np-complete. *Neural Networks*, 5(1):117–127, 1992.

[BRST21] Joan Bruna, Oded Regev, Min Jae Song, and Yi Tang. Continuous lwe. In *Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing*, pages 694–707, 2021.

[CGKM22] Sitan Chen, Aravind Gollakota, Adam R Klivans, and Raghu Meka. Hardness of noise-free learning for two-hidden-layer neural networks. *arXiv preprint arXiv:2202.05258*, 2022.

[CGV15]   Aloni Cohen, Shafi Goldwasser, and Vinod Vaikuntanathan. Aggregate pseudorandom functions and connections to learning. In *Theory of Cryptography Conference*, pages 61–89. Springer, 2015.

[CKM22]   Sitan Chen, Adam R Klivans, and Raghu Meka. Learning deep relu networks is fixed-parameter tractable. In *2021 IEEE 62nd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 696–707. IEEE, 2022.

[CLLM22] Sitan Chen, Jerry Li, Yuanzhi Li, and Raghu Meka. Minimax optimality (probably) doesn't imply distribution learning for gans. *arXiv preprint arXiv:2201.07206*, 2022.

[CLLZ22] Sitan Chen, Jerry Li, Yuanzhi Li, and Anru R Zhang. Learning polynomial transformations. *arXiv preprint arXiv:2204.04209*, 2022.

[DGKP20] Abhimanyu Das, Sreenivas Gollapudi, Ravi Kumar, and Rina Panigrahy. On the learnability of random deep networks. In *Proceedings of the Fourteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 398–410. SIAM, 2020.

[DISZ17] Constantinos Daskalakis, Andrew Ilyas, Vasilis Syrgkanis, and Haoyang Zeng. Training gans with optimism. *arXiv preprint arXiv:1711.00141*, 2017.

[DK20] Ilias Diakonikolas and Daniel M Kane. Hardness of learning halfspaces with massart noise. *arXiv preprint arXiv:2012.09720*, 2020.

[DKKZ20] Ilias Diakonikolas, Daniel M Kane, Vasilis Kontonis, and Nikos Zarifis. Algorithms and sq lower bounds for pac learning one-hidden-layer relu networks. In *Conference on Learning Theory*, pages 1514–1539. PMLR, 2020.

[DKS17] Ilias Diakonikolas, Daniel M Kane, and Alistair Stewart. Statistical query lower bounds for robust estimation of high-dimensional gaussians and gaussian mixtures. In *2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 73–84. IEEE, 2017.

[DKZ20] Ilias Diakonikolas, Daniel Kane, and Nikos Zarifis. Near-optimal sq lower bounds for agnostically learning halfspaces and relus under gaussian marginals. *Advances in Neural Information Processing Systems*, 33:13586–13596, 2020.

[DV20] Amit Daniely and Gal Vardi. Hardness of learning neural networks with natural weights. *Advances in Neural Information Processing Systems*, 33:930–940, 2020.

[DV21] Amit Daniely and Gal Vardi. From local pseudorandom generators to hardness of learning. In *Conference on Learning Theory*, pages 1358–1394. PMLR, 2021.

[Fel09] Vitaly Feldman. On the power of membership queries in agnostic learning. *The Journal of Machine Learning Research*, 10:163–182, 2009.

[FFGT17] Soheil Feizi, Farzan Farnia, Tony Ginart, and David Tse. Understanding gans: the lqg setting. *arXiv preprint arXiv:1710.10793*, 2017.

[FGR+17] Vitaly Feldman, Elena Grigorescu, Lev Reyzin, Santosh S Vempala, and Ying Xiao. Statistical algorithms and a lower bound for detecting planted cliques. *Journal of the ACM (JACM)*, 64(2):1–37, 2017.

[GGJ+20] Surbhi Goel, Aravind Gollakota, Zhihan Jin, Sushrut Karmalkar, and Adam Klivans. Superpolynomial lower bounds for learning one-layer neural networks using gradient descent. In *International Conference on Machine Learning*, pages 3587–3596. PMLR, 2020.

[GGK20] Surbhi Goel, Aravind Gollakota, and Adam Klivans. Statistical-query lower bounds via functional gradients. *Advances in Neural Information Processing Systems*, 33:2147–2158, 2020.

[GHP+19] Gauthier Gidel, Reyhane Askari Hemmat, Mohammad Pezeshki, Rémi Le Priol, Gabriel Huang, Simon Lacoste-Julien, and Ioannis Mitliagkas. Negative momentum for improved game dynamics. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 1802–1811. PMLR, 2019.

[GMSR20] Spencer Gordon, Bijan Mazaheri, Leonard J Schulman, and Yuval Rabani. The sparse hausdorff moment problem, with application to topic models. *arXiv preprint arXiv:2007.08101*, 2020.

[GPAM+14] Ian Goodfellow, Jean Pouget-Abadie, Mehdi Mirza, Bing Xu, David Warde-Farley, Sherjil Ozair, Aaron Courville, and Yoshua Bengio. Generative adversarial nets. *Advances in neural information processing systems*, 27, 2014.

[GSW+21] Jie Gui, Zhenan Sun, Yonggang Wen, Dacheng Tao, and Jieping Ye. A review on generative adversarial networks: Algorithms, theory, and applications. *IEEE Transactions on Knowledge and Data Engineering*, 2021.

[GVV22]   Aparna Gupte, Neekon Vafa, and Vinod Vaikuntanathan. Continuous lwe is as hard as lwe & applications to learning gaussian mixtures. *arXiv preprint arXiv:2204.02550*, 2022.

[Har02]   Philip Hartman. *Ordinary differential equations*. SIAM, 2002.

[JMGL22] Samy Jelassi, Arthur Mensch, Gauthier Gidel, and Yuanzhi Li. Adam is no better than normalized SGD: Dissecting how adaptivity improves GAN performance, 2022.

[Kha95]   Michael Kharitonov. Cryptographic lower bounds for learnability of boolean functions on the uniform distribution. *Journal of Computer and System Sciences*, 50(3):600–610, 1995.

[KK14]   Adam Klivans and Pravesh Kothari. Embedding hard learning problems into gaussian space. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques (APPROX/RANDOM 2014)*. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2014.

[KS09]   Adam R Klivans and Alexander A Sherstov. Cryptographic hardness for learning intersections of halfspaces. *Journal of Computer and System Sciences*, 75(1):2–12, 2009.

[KW13]   Diederik P Kingma and Max Welling. Auto-encoding variational bayes. *arXiv preprint arXiv:1312.6114*, 2013.

[LD20]   Yuanzhi Li and Zehao Dou. Making method of moments great again?–how can gans learn distributions. *arXiv preprint arXiv:2003.04033*, 2020.

[LLDD20] Qi Lei, Jason Lee, Alex Dimakis, and Constantinos Daskalakis. SGD learns one-layer networks in wgans. In *International Conference on Machine Learning*, pages 5799–5808. PMLR, 2020.

[LSSS14] Roi Livni, Shai Shalev-Shwartz, and Ohad Shamir. On the computational efficiency of training neural networks. *Advances in neural information processing systems*, 27, 2014.

[RM15]   Danilo Rezende and Shakir Mohamed. Variational inference with normalizing flows. In *International conference on machine learning*, pages 1530–1538. PMLR, 2015.

[SVWX17] Le Song, Santosh Vempala, John Wilmes, and Bo Xie. On the complexity of learning neural networks. *Advances in neural information processing systems*, 30, 2017.

[SZB21]   Min Jae Song, Ilias Zadik, and Joan Bruna. On the cryptographic hardness of learning single periodic neurons. *Advances in Neural Information Processing Systems*, 34, 2021.

[Val84]   Leslie G Valiant. A theory of the learnable. *Communications of the ACM*, 27(11):1134–1142, 1984.

[Vu98]   Van H Vu. On the infeasibility of training neural networks with small mean-squared error. *IEEE Transactions on Information Theory*, 44(7):2892–2900, 1998.

[VW19]   Santosh Vempala and John Wilmes. Gradient descent for one-hidden-layer neural networks: Polynomial convergence and sq lower bounds. In *Conference on Learning Theory*, pages 3115–3117. PMLR, 2019.

[WDS19]   Shanshan Wu, Alexandros G Dimakis, and Sujay Sanghavi. Learning distributions generated by one-layer relu networks. *Advances in neural information processing systems*, 32, 2019.

A Deferred Proofs From Section 3

We begin by reviewing standard concepts pertaining to establishing statistical query lower bounds for unsupervised learning problems, as developed in [FGR[+]17].

Definition 4 (Distributional search problems). *Let $\mathcal{D}$ be a set of probability distributions, let $\mathcal{F}$ be a set of* solutions, *and let $\mathcal{Z} : \mathcal{D} \to 2^{\mathcal{F}}$ be a map that takes any $D \in \mathcal{D}$ to a subset of $\mathcal{F}$ corresponding to the valid solutions for D. We say that $\mathcal{Z}$ specifies a* distributional search problem *over $\mathcal{D}$ and $\mathcal{F}$: given oracle access to an unknown $D \in \mathcal{D}$, the goal of the learner is to output a valid solution from $\mathcal{Z}(D)$.*

Definition 5 (Statistical query oracles). *Given a distribution D over $\mathbb{R}^d$ and parameters $\tau, t > 0$, a $\mathrm{STAT}(\tau)$ oracle takes in any query of the form $f : \mathbb{R}^d \to [-1, 1]$ and outputs a value from the interval $[\mathbb{E}_{x \sim D}[f(x)] - \tau, \mathbb{E}_{x \sim D}[f(x)] + \tau]$, while a $\mathrm{VSTAT}(t)$ oracle takes in any query of the form $f : \mathbb{R}^d \to [0, 1]$ and outputs a value from the interval $[\mathbb{E}_{x \sim D}[f(x)] - \tau, \mathbb{E}_{x \sim D}[f(x)] + \tau]$ for $\tau = \max(1/t, \sqrt{\mathbb{V}_{x \sim D}[f(x)]/t})$.*

Definition 6 (Pairwise correlation). *Given distributions p, q over a domain Ω which are absolutely continuous with respect to a distribution r over Ω, we let $\chi_r^2(p, q)$ denote the* pairwise correlation, *that is*

$$\chi_r^2(p, q) \triangleq \int_\Omega p(x)q(x)/r(x)\,\mathrm{d}x - 1.$$

Note that when $p = q$, this is simply the chi-squared divergence between p and r.

We say that a set of m distributions $\mathcal{D} = \{D_1, \ldots, D_m\}$ is (γ, β)-correlated relative to a distribution μ over $\mathbb{R}^d$ if

$$|\chi_\mu(D_i, D_j)| \leq \begin{cases} \gamma & \text{if } i \neq j \\ \beta & \text{if } i = j \end{cases}.$$

Definition 7 (Statistical dimension). *Let $\beta, \gamma > 0$, let $\mathcal{Z}$ be a distributional search problem over distributions $\mathcal{D}$ and solutions $\mathcal{F}$, and let N be the largest integer for which there exists a distribution μ and a finite subset $\mathcal{D}_\mu \subseteq \mathcal{D}$ such that for any $f \in \mathcal{F}$, $\mathcal{D}_f \triangleq \mathcal{D}_\mu \backslash \mathcal{Z}^{-1}(f)$ is (γ, β)-correlated relative to μ and $|\mathcal{D}_f| \geq N$. We say that the* statistical dimension *with pairwise correlations (γ, β) of $\mathcal{Z}$ is N and denote it by $\mathrm{SD}(\mathcal{Z}, \gamma, \beta)$.*

Lemma A.1 (Corollary 3.12 from [FGR[+]17]). *Let $\mathcal{Z}$ be a distributional search problem over distributions $\mathcal{D}$ and solutions $\mathcal{F}$. For γ, β, if $N = \mathrm{SD}(\mathcal{Z}, \gamma, \beta)$, then any statistical query algorithm for $\mathcal{Z}$ requires at least $N\gamma/(\beta - \gamma)$ queries to $\mathrm{STAT}(\sqrt{2\gamma})$ or $\mathrm{VSTAT}(1/(6\gamma))$.*

We will use the following two lemmas from [DKS17] and [DK20]. Recall from the main body that P_v^D denotes the distribution over $\mathbb{R}^d$ with density

$$P_v^D(x) = D(\langle v, x \rangle) \cdot \gamma^{(d-1)}(x - \langle v, x \rangle v),$$

that is the distribution which is given by D in the direction v and is given by $\mathcal{N}(0, \mathrm{Id} - vv^\top)$ orthogonal to v.

Lemma A.2 (Lemma 3.5 from [DK20]). *There is an absolute constant $c > 0$ such that the following holds. Let $m \in \mathbb{N}$ and $\nu > 0$. If a distribution D over $\mathbb{R}$ is such that 1) $\chi^2(D, \mathcal{N}(0, 1))$ is finite, and 2) $|\mathbb{E}_{x \sim D}[x^k] - \mathbb{E}_{g \sim \mathcal{N}(0,1)}[g^k]| \leq \nu$ for all $k = 1, \ldots, m$, then for all $v, v' \in \mathbb{S}^{d-1}$ for which $|\langle v, v' \rangle| < c$,*

$$|\chi_{\mathcal{N}(0, \mathrm{Id}_d)}^2(P_v^D, P_{v'}^D)| \leq |\langle v, v' \rangle|^{m+1} \chi^2(D, \mathcal{N}(0, 1)) + \nu^2.$$

Fact A.3 (Lemma 3.7 from [DKS17]). *For any constant $0 < C < 1/2$, there exists a set S of 2^{d^C} unit vectors in $\mathbb{S}^{d-1}$ such that any pair of distinct $u, v \in S$ satisfies $|\langle u, v \rangle| < d^{C-1/2}$.*

We can now prove the generic statistical query lower bound of Lemma 3.3:

Proof of Lemma 3.3. Let S be the set of 2^{d^C} unit vectors from Fact A.3. In the notation of Definition 7, take $\mu = \mathcal{N}(0, \mathrm{Id})$ and let $\mathcal{D}_\mu \triangleq \{P_v^D\}_{v \in S}$. By Lemma A.2, for any distinct $v, v' \in S$ we have

$$\chi_\mu(P_v^D, P_{v'}^D) \leq |\langle v, v' \rangle|^{m+1} \chi^2(D, \mathcal{N}(0,1)) + O(\tau^2) \leq \Omega(d)^{-(m+1)(1/2-C)} \chi^2(D, \mathcal{N}(0,1)).$$

On the other hand, if $v = v' \in S$, then $\chi_\mu(P_v^D, P_v^D) = \chi^2(D, \mathcal{N}(0,1)) + O(\tau^2) \leq 2\chi^2(D, \mathcal{N}(0,1))$. So for

$$\gamma \triangleq \Omega(d)^{-(m+1)(1/2-C)} \chi^2(D, \mathcal{N}(0,1)) \qquad \text{and} \qquad \beta \triangleq 2\chi^2(D, \mathcal{N}(0,1)),$$

$\mathcal{D}_\mu$ is (γ, β)-correlated with respect to μ.

Consider the distributional search problem $\mathcal{Z}$ mapping any distribution P_v^D to the set of probability distributions which are ϵ-close in total variation distance to P_v^D. Because $\mathrm{d}_{\mathrm{TV}}(P_v, P_{v'}) > 2\epsilon$ for distinct $v, v' \in S$, for any distribution f over $\mathbb{R}^d$ we have that $|\mathcal{Z}^{-1}(f)| \leq 1$. We conclude that $\mathrm{SD}(\mathcal{Z}, \gamma, \beta) \geq 2^{\Omega(d^C)}$. By Lemma A.1, we conclude that any SQ algorithm for $\mathcal{Z}$ requires at least $2^{\Omega(d^C)} d^{-(m+1)(1/2-C)}$ calls to either $\mathrm{STAT}(\tau)$ or $\mathrm{VSTAT}(1/\tau^2)$. Note that because we are assuming that $d \geq m^{\Omega(1/C)}$, we have $2^{\Omega(d^{C/2})} \geq d^{m+1}$, so the total number of required queries is at least $2^{\Omega(d^{C/2})} \geq d^{m+1}$ as claimed. $\qquad\square$

B   Deferred Proofs From Section 4

B.1   Proof of Fact 4.3

Proof. We have for all $z \in \mathbb{R}$ that $T_c^{w,h,\epsilon}(z)$ is equal to

$$\frac{h}{\epsilon} \left(\mathrm{ReLU}(z - c + \epsilon + w) - \mathrm{ReLU}(z - c + w) - \mathrm{ReLU}(z - c - w) + \mathrm{ReLU}(z - c - \epsilon - w) \right).$$

$\qquad\square$

B.2   Proof of Lemma 4.4

To prove Lemma 4.4, we will need the following modification of the construction in Lemma 4.1.

Lemma B.1. *For any odd $m \in \mathbb{N}$, there exist weights $\lambda_1, \ldots, \lambda_{m-1} \geq 0$ and points $h_1, \ldots, h_{m-1} \in \mathbb{R}$ for which*

1. *(Sum of weights bounded away from 1)* $\sum_{i=1}^{m-1} \lambda_i = 1 - \Theta(1/\sqrt{m})$.

2. *(Moments match)* $|\sum_{i=1}^{m-1} \lambda_i h_i^k - \mathbb{E}_g[g^k]| < \nu$ *for all $k = 1, \ldots, 2m - 1$.*

3. *(Points symmetric about origin)* $h_1 \leq \cdots \leq h_{m-1}$ *and $h_i = -h_{m-i}$ for all $1 \leq i < m$.*

4. *(Weights symmetric)* $\lambda_1 \leq \cdots \leq \lambda_{(m-1)/2}$ *and $\lambda_i = \lambda_{m-i}$.*

5. *(Points bounded and separated)* $\Omega(1/\sqrt{m}) \leq |h_i| \leq O(\sqrt{m})$ *for all $1 \leq i < m$, and $\{h_i\}$ are $\Omega(1/\sqrt{m})$-separated.*

6. *(Weights not too small)* $\min_i \lambda_i \geq e^{-cm}$ *for an absolute constant $c > 0$.*

Proof. Because m is odd, we can take the weights and points to be given by Lemma 4.1 and remove the $(m+1)/2$-th weight and point– recall that the $(m+1)/2$-th point is 0 and thus does not contribute to $\sum_i \lambda_i h_i^k$. The fact that $\sum_{i=1}^{m-1} \lambda_i = 1 - \Theta(1/\sqrt{m})$ then follows from the fact that the $(m+1)/2$-th weight from Lemma 4.1 is of order $\Theta(1/\sqrt{m})$ by Part 6 of Lemma 4.1. The remaining parts of the lemma follow by the corresponding parts of Lemma 4.1. $\qquad\square$

Proof of Lemma 4.4. Let $\lambda_1, \ldots, \lambda_{m-1}, h_1, \ldots, h_{m-1}$ be as in Lemma B.1. As $\sum_i \lambda_i = 1 - \Theta(1/\sqrt{m})$, we claim there exist intervals $I_1, \ldots, I_{m-1}$ such that for any $i < j$, all points in I_i are strictly smaller than all points in I_j, such that $\gamma(I_i) = \lambda_i$ for all i, and such that the right endpoint of any I_i is at least $m^{-3/2}$ smaller than the left endpoint of I_{i+1}.

We can construct these intervals in an inductive fashion. First, let $\gamma \triangleq 1 - \sum_i \lambda_i = \Theta(1/\sqrt{m})$. Let $I_1 = [a_1, b_1]$ for $a_1 < b_1 < 0$ such that $\gamma(I_1) = \lambda_1$ and $\gamma((-\infty, a_1]) = \gamma/m$. Given $I_1, \ldots, I_i$ for $1 \leq i < (m-1)/2$, if $I_i = [a_i, b_i]$ is the right endpoint of I_i, then let $a_{i+1} > b_i$ be such that $\gamma([b_i, a_{i+1}]) = \gamma/m$, and define $I_{i+1} = [a_{i+1}, b_{i+1}]$ for $b_{i+1} > a_{i+1}$ satisfying $\gamma([a_{i+1}, b_{i+1}]) = \lambda_{i+1}$. By construction,

$$\gamma((-\infty, b_{(m-1)/2}]) = \sum_{i=1}^{(m-1)/2} \gamma(I_i) + \frac{m-1}{2} \cdot \frac{\gamma}{m} = \frac{1-\gamma}{2} + \frac{m-1}{2m} \cdot \frac{\gamma}{2} = \frac{1}{2} - \frac{\gamma}{2m},$$

so by Gaussian anticoncentration and the fact that $\gamma = \Theta(1/\sqrt{m})$, we conclude that $b_{(m-1)/2} \leq -\Omega(m^{-3/2})$. In the same way, we also conclude that because $\gamma([b_i, a_{i+1}]) = \frac{\gamma}{m}$, we must have $b_i - a_{i+1} \leq -\Omega(m^{-3/2})$. Finally, for $i = (m-1)/2 + 1, \ldots, m-1$, we can define I_i to be the reflection of I_{m-i} about the origin. Note that by our bounds on $b_i - a_{i+1}$ for $1 \leq i \leq (m-1)/2$ and on $b_{(m-1)/2}$, all of the intervals are $\Omega(m^{-3/2})$-separated from each other as claimed. And by design, $\gamma(I_i) = \lambda_i$ for all $1 \leq i \leq m-1$.

While the lemma is stated in terms of $\epsilon > 0$, let us first consider the following construction where $\epsilon = 0$. We can take the centers $c_1, \ldots, c_{m-1}$ in the lemma to be the centers of $I_1, \ldots, I_{m-1}$, and $w_1, \ldots, w_{m-1}$ to be half of the widths of $I_1, \ldots, I_{m-1}$, in which case $f \triangleq \sum_{i=1}^{m-1} T_{c_i}^{w_i, h_i, 0}$ immediately satisfies Parts 1 and 3 of the lemma. Then the pushforward of $\mathcal{N}(0, 1)$ under this choice of f is the distribution which with probability γ equals zero (when $z \sim \mathcal{N}(0, 1)$ lies outside of $I_1, \ldots, I_{m-1}$) and otherwise takes the value h_i with probability λ_i. Parts 2, 4, and 5 then follow from Lemma B.1. Finally, note that a_1 defined above is at most $O(\log m)$ in magnitude because $\gamma((-\infty, a_1]) = \gamma/m$ by Part 6 of Lemma B.1. This establishes Part 6 of the lemma.

Finally, note that by taking ϵ infinitesimally small (relative to ν) but positive, the function f defined in the lemma satisfies all of the parts of the lemma. $\qquad\square$

B.3 Proofs from Section 4.2.1

First, we give an explicit expression for $M_{c,k}^{w,h,\epsilon} \triangleq \mathbb{E}_g[T_c^{w,h,\epsilon}(g)^k]$:

Lemma B.2. *For $c, w, h, \epsilon > 0$ satisfying $c - \epsilon - w \geq 0$, we have*

$$M_{c,k}^{w,h,\epsilon} = h^k \gamma([c-w, c+w]) +$$

$$(k-1)!! \left(\frac{h}{\epsilon}\right)^k \sum_{\substack{i=0 \\ even}}^{k} \binom{k}{i} \left[(-c+\epsilon+w)^{k-i} \gamma([c-\epsilon-w, c-w]) + (c+\epsilon+w)^{k-i} \gamma([c+w, c+\epsilon+w]) \right]$$

$$- \left(\frac{h}{\epsilon}\right)^k \sum_{i=0}^{k} \binom{k}{i} \left[(-c+\epsilon+w)^{k-i} \left(p_i(c-w)\gamma(c-w) - p_i(c-\epsilon-w)\gamma(c-\epsilon-w) \right) \right.$$

$$\left. + (-1)^i(c+\epsilon+w)^{k-i} \left(p_i(c+\epsilon+w)\gamma(c+\epsilon+w) - p_i(c+w)\gamma(c+w) \right) \right]$$

for all even k.

To show this we use the form of the moments of a truncated Gaussian. Given $m, i \in \mathbb{N}$, let $m^{\Downarrow i} \triangleq m(m-2) \cdots (m-2i+2)$. Also let $m^{\Downarrow 0} = 1$. Then:

Lemma B.3. *For any $k \in \mathbb{N}$, define the polynomial*

$$p_k(x) \triangleq \sum_{i=0}^{\lfloor (k-1)/2 \rfloor} (k-1)^{\Downarrow i} x^{k-1-2i}.$$

For any $a \leq b$,

$$\mathbb{E}_g[g^k \cdot \mathbb{1}[a \leq g \leq b]] = \begin{cases} (k-1)!! \cdot \gamma([a,b]) - (p_k(b)\gamma(b) - p_k(a)\gamma(a)) & \text{if } k \text{ even} \\ -(p_k(b)\gamma(b) - p_k(a)\gamma(b)) & \text{if } k \text{ odd} \end{cases}$$

Corollary B.4. *For any $c, d \in \mathbb{R}$ and $k \in \mathbb{N}$ even,*

$$\mathbb{E}_g[(cg + d)^k \cdot \mathbb{1}[a \le g \le b]] = \sum_{\substack{i=0 \\ even}}^{k} \binom{k}{i} c^i d^{k-i} (k-1)!! \gamma([a, b])$$

$$- \sum_{i=0}^{k} \binom{k}{i} c^i d^{k-i} (p_i(b)\gamma(b) - p_i(a)\gamma(a)).$$

Proof of Lemma B.2. By Corollary B.4, the contribution from the interval $g \in [c - \epsilon - w, c - w]$ to $\mathbb{E}_g\left[\left(T_c^{w,h,\epsilon}(z)\right)^k\right]$ is given by

$$\sum_{\substack{i=0 \\ even}}^{k} \binom{k}{i} \left(\frac{h}{\epsilon}\right)^i \left(\frac{h}{\epsilon}(-c + \epsilon + w)\right)^{k-i} (k-1)!! \gamma([c - \epsilon - w, c - w]) -$$

$$\sum_{i=0}^{k} \binom{k}{i} \left(\frac{h}{\epsilon}\right)^i \left(\frac{h}{\epsilon}(-c + \epsilon + w)\right)^{k-i} (p_i(c - w)\gamma(c - w) - p_i(c - \epsilon - w)\gamma(c - \epsilon - w)).$$

Similarly, the contribution from the interval $g \in [c + w, c + \epsilon + w]$ is given by

$$\sum_{\substack{i=0 \\ even}}^{k} \binom{k}{i} \left(\frac{-h}{\epsilon}\right)^i \left(\frac{h}{\epsilon}(c + \epsilon + w)\right)^{k-i} (k-1)!! \gamma([c + w, c + \epsilon + w]) -$$

$$\sum_{i=0}^{k} \binom{k}{i} \left(\frac{-h}{\epsilon}\right)^i \left(\frac{h}{\epsilon}(c + \epsilon + w)\right)^{k-i} (p_i(c + \epsilon + w)\gamma(c + \epsilon + w) - p_i(c + w)\gamma(c + w)).$$

Finally, the contribution from the interval $g \in [c - w, c + w]$ is given by $h^k \cdot \gamma([c - w, c + w])$. $\square$

Lemma 4.5 now follows immediately from LemmaB.2. Furthermore, it is clear from Lemma B.2 that $M_{c,k}^{w,h,\epsilon}$ is continuously differentiable with respect to ϵ when $\epsilon > 0$.

Proof of Lemma 4.6. Note that $0 \le M_{c,k}^{w,h,\epsilon} \le h^k \gamma([c - \epsilon - w, c + \epsilon + w])$, so the first part of the lemma follows by upper bounding $|M_{c,k}^{w,h',\epsilon'} - M_{c,k}^{w,h,\epsilon}|$ by

$$|h'^k - h^k| \gamma([c - \epsilon' - w, c + \epsilon' + w]) + h^k \gamma([c - \epsilon' - w, c - \epsilon - w] \cup [c + \epsilon + w, c + \epsilon' + w])$$

$$\le |h'^k - h^k| + h^k(\epsilon' - \epsilon) = h^k \left(|(h'/h)^k - 1| + \epsilon' - \epsilon\right),$$

where in the last step we used that $\gamma([a, a + \eta]) \le \eta/2$ for any $a \in \mathbb{R}$, $\eta \ge 0$. The second part of the lemma then follows by taking $h = h'$ and $\epsilon' \to \epsilon$. $\square$

B.4 Proof of Lemma 4.7

Proof. For convenience, in this proof we refer to $Z(\mathbf{h}(0), \epsilon(0))$ as Z. Note that $Z_{i,\ell} = \gamma([c_i - w_i, c_i + w_i]) \cdot h_i(0)^{2\ell} + \xi_{i,\ell}$ for some $\xi_{i,\ell}$ which can be made arbitrarily small by taking $\epsilon(0)$ to be arbitrarily small. We can thus write $Z = \Lambda H + \Xi$ for $\Lambda = \text{diag}(\lambda_1 h_1(0)^2, \ldots, \lambda_{(m-1)/2} h_{(m-1)/2}(0)^2)$, $H \in \mathbb{R}^{(m-1)/2 \times (m-1)/2}$ given by $H_{i,\ell} = h_i(0)^{2\ell-2}$, and Ξ a matrix consisting of arbitrarily small positive entries. So $\sigma_{\min}(Z) \ge (\min_i \lambda_i h_i(0)^2) \cdot \sigma_{\min}(H) - \xi \ge (e^{-cm}/m) \cdot \sigma_{\min}(H) - \xi$ for arbitrarily small $\xi > 0$, where in the last step we used Parts 4 and 5 of Lemma 4.4.

Finally, note that H is a Vandermonde matrix with nodes $h_1(0)^2, \ldots, h_{(m-1)/2}(0)^2$. As $\{h_i(0)\}$ are $\Omega(1/\sqrt{m})$-separated and lie within $[\Omega(1/\sqrt{m}), O(\sqrt{m})]$, $\{h_i(0)^2\}$ are $\Omega(1/m)$-separated. So by Fact 2.1, $\sigma_{\min}(H) \ge m^{-O(m)}$, concluding the proof. $\square$

B.5 Proof of Lemma 4.8

Proof. For convenience, in this proof we refer to $Z(\mathbf{h}, \epsilon)$ and $Z(\mathbf{h}(0), \epsilon(0))$ by Z' and Z respectively. By Lemma 4.6, each entry of Z' differs from the corresponding entry of Z by at most

$$(\max_i h_i^{m-1}) \cdot \left(\left| (1 + m^{-C'm}/(\min_i h_i))^{m-1} - 1 \right| + m^{-C'm} - \epsilon(0) \right).$$

As $\max_i h_i \leq O(\sqrt{m})$ and $\min_i h_i \geq \Omega(1/\sqrt{m})$ by Part 4 of Lemma 4.4, and $\epsilon(0)$ can is an arbitrarily small positive quantity, the above is at most $m^{-C''m}$ for some absolute constant $C'' > 0$ which is increasing in C'. So $\|Z - Z'\|_{\mathsf{op}} \leq \|Z - Z'\|_F \leq (m-1)/2 \cdot m^{-C''m} \ll m^{-Cm}$ provided we take C' sufficiently large. $\qquad \square$

B.6 Proof of Lemma 4.9

Proof of Lemma 4.9. By our expression for $M_{c,k}^{w,h,\epsilon}$ in Lemma B.2 and the definition of $\mathbf{b}(\epsilon)$ in (3), $\mathbf{b}(\epsilon(0)+t)$ is clearly continuous in t whenever $t \geq 0$ (because $\epsilon(0) > 0$). Similarly, $A(\mathbf{h})Z(\mathbf{h}, \epsilon(0)+t)B$ is clearly continuous with respect to $\mathbf{h}$ and t whenever $t \geq 0$ and $h_i \neq 0$ for all i. By Lemma 4.8, if $t \leq m^{-C'm}$ and $\|\mathbf{h} - \mathbf{h}(0)\|_\infty \leq m^{-C'm}$ (which additionally implies that $h_i \neq 0$ for all i, by Part 4 of Lemma 4.4), then $A(\mathbf{h})Z(\mathbf{h}, \epsilon(0)+t)B$ is invertible. We conclude that for such $t, \mathbf{h}, w$ is continuous. $\qquad \square$

B.7 Proof of Lemma 4.12

Proof. For any $1 \leq k < m$,

$$\mathop{\mathbb{E}}_{x \sim D'}[x^k] = \mathop{\mathbb{E}}_{z \sim \sqrt{1-\sigma^2} \cdot D, g \sim \mathcal{N}(0,1)}[(z+\sigma g)^k] = \mathop{\mathbb{E}}_{g,g' \sim \mathcal{N}(0,1)}[(\sqrt{1-\sigma^2}g' + \sigma g)^k] = \mathop{\mathbb{E}}_{g \sim \mathcal{N}(0,1)}[g^k],$$

where in the second step we used that $\sqrt{1-\sigma^2} \cdot D$ matches the moments of $\mathcal{N}(0, 1-\sigma^2)$ up to degree m, and in the last step we used that $\sqrt{1-\sigma^2}g' + \sigma g$ is distributed as a draw from $\mathcal{N}(0,1)$. $\quad \square$

B.8 Proof of Lemma 4.13

Proof. Let $\widetilde{A} \triangleq A \star \mathcal{N}(0,1)$. By definition, $\widetilde{A}(x) = \int_{-\infty}^\infty A(s)\gamma_{\sigma^2}(x-s) \, \mathrm{d}s$. So

$$1 + \chi^2(\widetilde{A}, \mathcal{N}(0,1)) = \int_{-\infty}^\infty \frac{1}{\gamma(x)} \left(\int_{-\infty}^\infty \int_{-\infty}^\infty A(s)A(t)\gamma_{\sigma^2}(x-s)\gamma_{\sigma^2}(x-t) \, \mathrm{d}s\mathrm{d}t \right) \mathrm{d}x. \tag{7}$$

Note that for any $s, t \in \mathbb{R}$,

$$\frac{\gamma_{\sigma^2}(x-s)\gamma_{\sigma^2}(x-t)}{\gamma(x)} = \frac{1}{\sigma^2\sqrt{2\pi}} \exp\left(\frac{-(x-s)^2 - (x-t)^2}{2\sigma^2} + x^2/2 \right)$$

$$= \frac{1}{\sigma^2\sqrt{2\pi}} \exp\left(-\frac{2-\sigma^2}{2\sigma^2} \left(x - \frac{s+t}{2-\sigma^2} \right)^2 + \frac{2st - (s^2+t^2)(1-\sigma^2)}{2\sigma^2(2-\sigma^2)} \right),$$

so

$$\int_{-\infty}^\infty \frac{\gamma_{\sigma^2}(x-s)\gamma_{\sigma^2}(x-t)}{\gamma(x)} \, \mathrm{d}x = \frac{e^{\frac{2st-(s^2+t^2)(1-\sigma^2)}{2\sigma^2(2-\sigma^2)}}}{\sigma\sqrt{2-\sigma^2}}.$$

Eq. (7) thus becomes

$$1 + \chi^2(\widetilde{A}, \mathcal{N}(0,1)) = \int_{-\infty}^\infty \int_{-\infty}^\infty A(s)A(t) \cdot \frac{e^{\frac{2st-(s^2+t^2)(1-\sigma^2)}{2\sigma^2(2-\sigma^2)}}}{\sigma\sqrt{2-\sigma^2}} \, \mathrm{d}s\mathrm{d}t \tag{8}$$

As A is supported on $[-R, R]$,

$$\frac{e^{\frac{2st-(s^2+t^2)(1-\sigma^2)}{2\sigma^2(2-\sigma^2)}}}{\sigma\sqrt{2-\sigma^2}} \leq e^{O(R^2/(1-\sigma^2))} \leq e^{O(R^2)}.$$

Substituting this into (8), we find that

$$1 + \chi^2(\widetilde{A}, \mathcal{N}(0,1)) \leq \frac{e^{O(k)}}{\sigma\sqrt{2-\sigma^2}} \int \int A(s)A(t) \, \mathrm{d}s\mathrm{d}t = \frac{e^{O(R^2)}}{\sigma\sqrt{2-\sigma^2}} \leq e^{O(R^2)}/\sigma. \qquad \square$$

B.9 Proof of Lemma 4.14

We will use the following elementary fact about total variation distance:

Fact B.5. *Given two distributions p, q over a domain Ω, $d_{\mathrm{TV}}(p, q) = 1 - \int_\Omega \min(p(x), q(x)) \, \mathrm{d}x$.*

Proof of Lemma 4.14. By Fact B.5, it suffices to upper bound $\int_{\mathbb{R}^d} \min(P_v(z), P_{v'}(z)) \, \mathrm{d}z$. Let H denote the plane spanned by v, v'. As the component in H of a sample from either P_v or $P_{v'}$ is independent from the component in $H^\perp$, and the latter is distributed as $\mathcal{N}(0, \Pi_{H^\perp})$, it suffices to bound $\int_H \min(P_v(z), P_{v'}(z)) \, \mathrm{d}z$. Let x, y be orthogonal coordinates for H with v in the direction of the x-axis, and let x', y' be orthogonal coordinates for H with v' in the direction of the x'-axis. Let θ be the angle between v, v'. Then

$$\int_H \min(P_v(z), P_{v'}(z)) \, \mathrm{d}z = \int_{-\infty}^\infty \int_{-\infty}^\infty \min(D'(x)\gamma(y), D'(x')\gamma(y')) \, \mathrm{d}x \mathrm{d}y$$
$$= \int_{-\infty}^\infty \int_{-\infty}^\infty \min(D'(x)\gamma(y), D'(x')\gamma(y')) \csc\theta \, \mathrm{d}x \mathrm{d}x'. \tag{9}$$

For $1 \le i \le m - 1$, let D_i denote the distribution of $\sqrt{1 - \sigma^2} \cdot T_{x_i}^{w_i, \mathbf{h}_i(T), T}(g)$ for $g \sim \mathcal{N}(0, 1)$ conditioned on $g \in [x_i - w_i - T, x_i + w_i + T]$. Also let D_m denote the distribution which is a point mass at zero. Let $D_i' \triangleq D_i \star \mathcal{N}(0, 1)$. Note that there is a distribution p over $[m]$ for which $D' = \mathbb{E}_{i \sim p}[D_i']$. Then we can upper bound (9) by

$$\max_{i,j \in [m]} \int_{-\infty}^\infty \int_{-\infty}^\infty \min(D'(x)_i \gamma(y), D_j'(x')\gamma(y')) \csc\theta \, \mathrm{d}x \mathrm{d}x' \tag{10}$$

Note that for $1 \le i \le m - 1$, $\mathbb{P}_{x \sim D_i'}[|x - x_i| > a] \le \sigma + \xi_i$ for $a \triangleq 2\sigma\sqrt{\log(1/\sigma)}$ and $\xi_i \triangleq \frac{\gamma([x_i - w_i - T, x_i - w_i] \cup [x_i + w_i, x_i + w_i + T])}{\gamma([x_i - w_i - T, x_i + w_i + T])}$. And for $i = m$, $\mathbb{P}_{x \sim D_m'}[|x - x_i| > a] \le \sigma$ for $x_m \triangleq 0$. So we get an upper bound on (10) of

$$\sigma + \max_{i \in [m-1]} \xi_i + \max_{i,j \in [m]} \int_{x_i - a}^{x_i + a} \int_{x_j - a}^{x_j + a} \min(D_i'(x)\gamma(y), D_j'(x')\gamma(y')) \csc\theta \, \mathrm{d}x \mathrm{d}x'$$

As D_i' is a convolution of D_i with $\mathcal{N}(0, \sigma^2)$, $D_i'(x) \le \frac{1}{\sigma\sqrt{2\pi}}$ for all $x \in \mathbb{R}$. And $\gamma(y) \le 1/\sqrt{2\pi}$, so the above display is at most

$$\sigma + \max_{i \in [m-1]} \xi_i + a^2 \csc\theta/(2\pi\sigma) = \sigma + \max_{i \in [m-1]} \xi_i + \sigma \csc\theta \log(1/\sigma)/\pi.$$

Note that if $|\langle v, v' \rangle| \le 1/2$, then $\csc\theta/\pi \le 2/(\sqrt{3}\pi) \le 1$. Finally, to bound ξ_i, first note that for any $i \in [m-1]$, $\gamma([x_i - w_i - T, x_i + w_i + T]) = \lambda_i$, and recall that $\lambda_i \ge e^{-cm}$ for some absolute constant $c > 0$. On the other hand, $\gamma([x_i - w_i - T, x_i - w_i]) \le T/\sqrt{2\pi} = m^{-(C' + C'')m}/\sqrt{2\pi}$ by Gaussian anti-concentration and our choice of $T = m^{-(C' + C'')m}$ in the proof of Lemma 4.11. So by taking the constants C', C'' to be larger than c, we conclude that $\xi_i \le m^{-\Omega(m)}$ for all $i \in [m-1]$. $\qquad\square$

C Hardness for Estimation in Wasserstein

We now show an analogous version of Theorem 3.1 under the Wasserstein-1 metric rather than total variation distance. We begin by observing that the ODE-based evolution does not move the pushforward at time zero, i.e. the distribution constructed in Lemma 4.4, too far away in Wasserstein distance over a time horizon of T:

Lemma C.1. *Let $f_0, f_T : \mathbb{R} \to \mathbb{R}$ denote the functions from Lemmas 4.4 and 4.11 respectively. Define $D_0 \triangleq f_0(\mathcal{N}(0, 1))$ and $D_T \triangleq f_T(\mathcal{N}(0, 1))$. Then $W_1(D, D') \le m^{-\Omega(m)}$.*

Proof. Recall that $w_1, \ldots, w_{m-1}$ denote the widths of the bumps in f_0, f_T, $x_1, \ldots, x_{m-1}$ denote the centers, and the heights and ϵ parameters for the bumps in f_0, f_T are given by $\{h_i(0)\}_i, \epsilon(0)$

and $\{h_i(T)\}_i, \epsilon(0) + T$ respectively, for $T = m^{-(C'+C'')m}$ and $\epsilon(0)$ an arbitrarily small positive quantity. Also recall from the proof of Lemma 4.11 that $|h_i(0) - h_i(T)| \leq m^{-C'm}$ for all i.

Now consider any $g \in \mathbb{R}$. If $g \in [x_i - w_i, x_i + w_i]$ for some i, then $|f_0(g) - f_T(g)| = |h_i(0) - h_i(T)| \leq m^{-C'm}$. Furthermore,

$$\underset{g \sim \mathcal{N}(0,1)}{\mathbb{P}} [g \in [x_i - \epsilon_i(0) - T - w_i, x_i - w_i] \cup [x_i + w_i, x_i + \epsilon_i(0) + T + w_i] \text{ for some } i] \leq O(mT)$$

Finally, for all g that do not lie in any of the aforementioned intervals, i.e. that do not lie in the support of any bump from f_0 or f_T, note that $f_0(g) = f_T(g) = 0$ by construction. We conclude that for any 1-Lipschitz function $h : \mathbb{R} \to \mathbb{R}$,

$$\left| \underset{z \sim D}{\mathbb{E}}[h(z)] - \underset{z \sim D'}{\mathbb{E}}[h(z)] \right| = \left| \underset{g}{\mathbb{E}}[h(f_0(g)) - h(f_T(g))] \right|$$

$$\leq \underset{g}{\mathbb{E}}[|f_0(g) - f_T(g)|] \leq m^{-C'm} + O(mT) \leq m^{-\Omega(m)}$$

as claimed. $\qquad\square$

We can now show the analogue of Lemma 4.14 for Wasserstein distance:

Lemma C.2. *Let $D = f(\mathcal{N}(0,1))$ be from Lemma 4.11, and define $D' \triangleq \sqrt{1-\sigma^2} \cdot D \star \mathcal{N}(0,\sigma^2)$ for $\sigma \ll 1/\sqrt{m}$. Then for any $v, v' \in \mathbb{S}^{d-1}$ satisfying $|\langle v, v' \rangle| \leq 1/2$, $W_1(P_v^{D'}, P_{v'}^{D'}) \geq \Omega(1/\sqrt{m})$.*

Proof. Let f_0 be the function from Lemma 4.4, and let A denote $f_0(\mathcal{N}(0,1))$. We begin by lower bounding $W_1(P_v^A, P_{v'}^A)$, which we will do by showing that with $\Omega(1)$ probability, a sample x from $P_{v''}^A$ will be distance $\Omega(1/\sqrt{m})$ from the support of P_v^A. As the distance from a point x to the affine hyperplane $\Lambda_h \triangleq \{z : \langle z, v \rangle = h\}$ is $|\langle v, x \rangle - h|$, if x is of the form $h'v' + v^\perp$ for some $h' \in \mathbb{R}$, then x is at distance

$$\left| h'\langle v', v \rangle + \langle v^\perp, v \rangle - h \right|$$

from the hyperplane. Note that P_v^A is supported on the hyperplanes $\Lambda_{h_1}, \ldots, \Lambda_{h_{m-1}}$ for $h_1, \ldots, h_{m-1}$ from Lemma 4.4. And for $x \sim P_{v'}^A$, h' takes on the value h_i with probability λ_i (where $\{\lambda_i\}$ are also from Lemma 4.4), while $v^\perp$ is an independent draw from $\mathcal{N}(0, \mathrm{Id} - v'v'^\perp)$. We conclude that $h'\langle v', v \rangle + \langle v^\perp, v \rangle$ is distributed as $\mathcal{N}(h'\langle v', v \rangle, 1 - \langle v', v \rangle^2)$. Therefore, the event that x is at distance $\Omega(1/\sqrt{m})$ from the support of P_v^A is equivalent to the event that a sample from $\mathcal{N}(h'\langle v', v \rangle, 1 - \langle v', v \rangle^2)$ is $\Omega(1/\sqrt{m})$-far from every $h_1, \ldots, h_{m-1}$. But note that because $h_1, \ldots, h_{m-1}$ are $\Omega(1/\sqrt{m})$-separated, there is an absolute constant $c > 0$ such that the union of the balls of radius $c/\sqrt{m}$ around $h_1, \ldots, h_{m-1}$ cover at most a constant fraction of the interval $[h'\langle v', v \rangle - 1, h'\langle v', v \rangle + 1]$. Because $1 - \langle v', v \rangle^2 \geq 3/4$, a constant fraction of the mass of $\mathcal{N}(h'\langle v', v \rangle, 1 - \langle v', v \rangle^2)$ is located in this interval, concluding the proof that $W_1(P_v^A, P_{v'}^A) \geq \Omega(1/\sqrt{m})$.

By Lemma C.1 and the fact that scaling by $\sqrt{1-\sigma^2}$ and convolving by $\mathcal{N}(0,\sigma^2)$ incurs $O(\sigma) = o(1/\sqrt{m})$ in Wasserstein, we conclude that $W_1(P_v^{D'}, P_v^A) = W_1(D', A) = o(1/\sqrt{m})$ and similarly for $W_1(P_{v'}^{D'}, P_{v'}^A)$. So by triangle inequality for Wasserstein, $W_1(P_v^{D'}, P_{v'}^{D'}) = \Omega(1/\sqrt{m})$ as claimed. $\qquad\square$

We conclude that in Theorem 3.2, the distribution D also satisfies the Wasserstein analogue of Part 3, i.e. $W_1(P_v^D, P_{v'}^D) \geq \Omega(1/\sqrt{m})$ for any $v, v' \in \mathbb{S}^{d-1}$ satisfying $|\langle v, v' \rangle| \geq 1/2$. We can now prove an analogue of Theorem 3.1 for Wasserstein:

Theorem C.3. *Let $d \in \mathbb{N}$ be sufficiently large. Any SQ algorithm which, given SQ access to an arbitrary one-hidden-layer ReLU network pushforward of $\mathcal{N}(0, \mathrm{Id}_d)$ of size $O(\log d / \log \log d)$ with $\mathrm{poly}(d)$-bounded weights, outputs a distribution which is $O(\sqrt{\log \log d / \log d})$-close in $\mathrm{d_{TV}}(\cdot)$ must make at least $d^{\Omega(\log d / \log \log d)}$ queries to either $\mathrm{STAT}(\tau)$ or $\mathrm{VSTAT}(1/\tau^2)$ for $\tau = d^{-\Omega(\log d / \log \log d)}$.*

Proof. By Theorem 3.2 applied with sufficiently large odd m and sufficiently small σ, together with the above consequence of Lemma C.2, there exists a distribution $D = f^*(\mathcal{N}(0, \mathrm{Id}_2))$ over $\mathbb{R}$ for

$f^* : \mathbb{R}^2 \to \mathbb{R}$ of size $O(m)$ with $m^{O(m)}$-bounded weights satisfying the hypotheses of Lemma 3.3 for $\epsilon = O(1/\sqrt{m})$, and $\chi^2(D, \mathcal{N}(0,1)) \leq \exp(O(m))$ (note that while Lemma 3.3 is stated for $\mathrm{d}_{\mathrm{TV}}(\cdot)$, it is also true with $\mathrm{d}_{\mathrm{TV}}(\cdot)$ replaced with Wasserstein-1). As long as $m \leq d^{O(C)}$, we conclude that an SQ algorithm for learning any distribution from $\{P_v^D\}_{v \in \mathbb{S}^{d-1}}$ to Wasserstein-1 distance $O(1/\sqrt{m})$ must make at least d^{m+1} queries to $\mathrm{STAT}(\tau)$ or $\mathrm{VSTAT}(1/\tau^2)$ for $\tau \triangleq O(d)^{-(m+1)(1/4-C/2)} \cdot \exp(O(m))$. By taking $m = \Theta(\log d / \log\log d)$, we ensure that $m^{O(m)} \leq \mathrm{poly}(d)$. We're done by taking C in Lemma 3.3 to be $C = 1/4$. $\qquad\square$

D   Hardness From Supervised Learning

In this section we make rigorous the claim from the introduction that lower bounds for PAC learning neural networks from Gaussian labeled examples imply lower bounds for learning neural network pushforwards. Formally, consider the following distinguishing problem:

Definition 8 (Distinguishing labeled examples from Gaussian). *For $d \in \mathbb{N}$, let $\mathcal{C}_d$ be some class of functions from $\mathbb{R}^d$ to $\mathbb{R}$. The learner is given $\mathrm{poly}(d)$ many samples $(x_1, y_1), \ldots, (x_N, y_N)$ where $x_1, \ldots, x_N$ are independent draws from $\mathcal{N}(0, \mathrm{Id}_d)$ such that one of the following is true: 1) there is some $h \in \mathcal{C}$ for which $y_i = h(x_i)$ for all $i \in [N]$, or 2) every y_i is an independent sample from $\mathcal{N}(0,1)$. We say that an algorithm distinguishes between these two situations with constant advantage if the probability it outputs* YES *(resp.* NO*) under the former (resp. latter) is at least $2/3$, where the probability is with respect to the randomness of the samples and internal randomness of the algorithm.*

Here we make the simple observation that an oracle for distinguishing any given family of non-Gaussian pushforwards from $\mathcal{N}(0, \mathrm{Id})$ (an easier task than actually learning pushforwards) immediately implies an algorithm for the distinguishing task in Definition 8.

Lemma D.1. *For $d \in \mathbb{N}$, let $\mathcal{C}_d$ be any function class from $\mathbb{R}^d \to \mathbb{R}$ for which the indexing functions $f^{[j]}$, given by $f^{[j]}(x) = x_j$ for some $j \in [d]$, are elements of $\mathcal{C}$. Suppose that for any $d_1, d_2 = \mathrm{poly}(d)$, there is a $\mathrm{poly}(d)$-time algorithm $\mathcal{A}$ for the following task. Let $d_1, d_2 = \mathrm{poly}(d)$, and let $\mathcal{S}$ be a known set of functions $f : \mathbb{R}^{d_1} \to \mathbb{R}^{d_2}$ whose output coordinates are all elements of $\mathcal{C}_{d_1}$ and such that for any $f \in \mathcal{S}$, $\mathrm{d}_{\mathrm{TV}}(f(\mathcal{N}(0, \mathrm{Id}_{d_1})), \mathcal{N}(0, \mathrm{Id}_{d_2})) \geq 1/2$. Then $\mathcal{A}$ can distinguish with constant advantage whether it is given $\mathrm{poly}(d)$ samples from $f(\mathcal{N}(0, \mathrm{Id}_{d_1}))$ for some $f \in \mathcal{S}$ versus samples from $\mathcal{N}(0, \mathrm{Id}_{d_2})$.*

Under this hypothesis, there is a $\mathrm{poly}(d)$-time algorithm that solves the distinguishing problem of Definition 8 to constant advantage.

Proof. Note that in situation 1) of Definition 8, the joint distribution over (x, y) is given by the pushforward $f(\mathcal{N}(0, \mathrm{Id}))$ where $f : \mathbb{R}^{d+1} \to \mathbb{R}^{d+1}$ is as follows: the first d output coordinates are given by the d indexing functions $f^{[1]}, \ldots, f^{[d]}$, and the last output coordinate is given by h. By taking $\mathcal{S}$ in the hypothesis to consist of such f, we can thus apply the algorithm $\mathcal{A}$ to distinguish between the two situations in Definition 8 to constant advantage. $\qquad\square$

Note that the contrapositive of the above lemma implies that any lower bound for the task in Definition 8 immediately implies a lower bound for learning pushforwards. While the aforementioned lower bounds of [CGKM22, DV21], which apply when $\mathcal{C}_d$ is the family of neural networks with at least two hidden layers and polynomially bounded size and weights, do not show hardness for the task in Definition 8, note that hardness for this task immediately implies hardness for PAC learning $\mathcal{C}_d$ from Gaussian examples. Indeed, given an algorithm $\mathcal{A}$ that, given $(x_1, h(x_1)), \ldots, (x_N, h(x_N))$, outputs a predictor $\widehat{h}$ for which $\mathbb{E}_g[(h(g) - \widehat{h}(g))^2]$ is small, one can easily solve the task in Definition 8 by running $\mathcal{A}$ and estimating the square loss of the predictor from some fresh samples. In situation 2) of Definition 8, because the labels are random, no predictor can achieve low square loss. So the algorithm which outputs YES if and only if the empirical square loss on fresh samples is small will distinguish between the two situations with constant advantage.

In other words, showing hardness of Definition 8 for $\mathcal{C}_d$ would be a *stronger result* than what is already shown in [CGKM22, DV21]. Putting this and Lemma D.1 together, we conclude that even this stronger hardness result would only imply hardness for learning pushforwards given by f whose output coordinates are functions in $\mathcal{C}_d$ given by neural networks with at least two hidden layers and

polynomially bounded size and weights. In contrast, in the present work, we show hardness for *one* hidden layer, *logarithmic* size, and polynomially bounded weights.