Mean Estimation in High-Dimensional Binary Markov Gaussian Mixture Models

Abstract

We consider a high-dimensional mean estimation problem over a binary hidden Markov model, which illuminates the interplay between memory in data, sample size, dimension, and signal strength in statistical inference. In this model, an estimator observes n samples of a d-dimensional parameter vector $\theta_* \in \mathbb{R}^d$, multiplied by a random sign S_i ($1 \le i \le n$), and corrupted by isotropic standard Gaussian noise. The sequence of signs $\{S_i\}_{i \in [n]} \in \{-1, 1\}^n$ is drawn from a stationary homogeneous Markov chain with flip probability $\delta \in [0, 1/2]$. As δ varies, this model smoothly interpolates two well-studied models: the Gaussian Location Model for which $\delta = 0$ and the Gaussian Mixture Model for which $\delta = 1/2$. Assuming that the estimator knows δ, we establish a nearly minimax optimal (up to logarithmic factors) estimation error rate, as a function of $\|\theta_*\|, \delta, d, n$. We then provide an upper bound to the case of estimating δ, assuming a (possibly inaccurate) knowledge of θ_*. The bound is proved to be tight when θ_* is an accurately known constant. These results are then combined to an algorithm which estimates θ_* with δ unknown a priori, and theoretical guarantees on its error are stated.

1 Introduction

Memory between data samples are ubiquitous in practical applications, as data collected from networks or sampled time series inevitably inherit spatial or temporal statistical dependencies. Numerous examples arise in imaging, meteorology, health care, finance, social science, and so on [Glaeser et al., 1996, Bertrand et al., 2000, Sacerdote, 2001, Duflo and Saez, 2003, Christakis and Fowler, 2013]. In principle, prior knowledge of the existence of such memory can be used to improve the performance of statistical estimators compared to memoryless models. Development and analysis of statistical inference algorithms for models with memory has been extensively explored from the algorithmic perspective, in which computationally efficient algorithms such as Baum-Welch and message-passing were developed [Ephraim and Merhav, 2002, van Handel, 2008, MacKay, 2003, Wainwright and Jordan, 2008], and from a theoretical perspective in the classical, fixed-dimensional and asymptotic regime, e.g., [Györfi et al., 2002, Chapter 27 and references therein]. However, much less is understood about about the high dimensional, non-asymptotic regime, which is of paramount importance in modern applications, and the focus of current extensive research [Vershynin, 2018, Wainwright, 2019]. As we exemplify in this paper, the error in estimation problems of this form depends in an intricate way on the interplay between the number of samples, the dimension of the vector parameters to be estimated, the noise level (signal-to-noise ratio), and the level of memory between the samples.

An ever popular and fundamental model is the Gaussian mixture model, for which memory exists between samples when the latent variables determining the component of each sample are dependent. Numerous recent papers [Balakrishnan et al., 2017, Xu et al., 2016, Jin et al., 2016,

Klusowski and Brinda, 2016, Dwivedi et al., 2020b, 2018, 2020a, Zhao et al., 2018, Yan et al., 2017, Weinberger and Bresler, 2022] have focus on high-dimensional *memoryless* models, and analyzed computationally efficient estimation algorithms, most notably the expectation maximization (EM) algorithm. Specifically, the seminal [Balakrishnan et al., 2017] has provided theoretical guarantees for EM in memoryless models, though without proving minimax optimality, which was subsequently established in [Wu and Zhou, 2019]. As a notable exception, Yang et al. [2015] has generalized the analysis of [Balakrishnan et al., 2017] to a hidden Markov model (which has memory), yet again, without determining how the minimax error depends on the number of samples, dimension, noise level, and the amount of memory. In this paper, we address the question of precise characterization of the minimax error rate in terms of these parameters, in the context of a hidden Markov high-dimensional Gaussian mixture model. Our characterization illuminates the opportunities and challenges associated with optimal inference in high dimensional models with memory. We next turn to formally define this model and the estimation problem, describe known results, and then present our contributions.

1.1 Problem formulation

Let $S_0^n := (S_0, S_1, \ldots, S_n)$ be the following homogeneous binary symmetric Markov chain, $S_i \in \{-1, 1\}$, $\mathbb{P}[S_0 = 1] = 1/2$ and

$$S_i = \begin{cases} S_{i-1}, & \text{w.p. } 1 - \delta \\ -S_{i-1}, & \text{w.p. } \delta \end{cases} \tag{1}$$

for $i \in [n] := \{1, \ldots, n\}$, and where $\delta \in [0, 1]$ is the flip probability of the binary Markov chain. We also denote $\rho := 1 - 2\delta \in [-1, 1]$ which is the correlation between adjacent samples $\rho = \mathbb{E}[S_i S_{i+1}]$. At each time point $i \in [n]$, a sample of a d-dimensional Gaussian mixture model is observed

$$X_i = S_i \theta_* + Z_i, \tag{2}$$

where $Z_i \sim N(0, I_d)$ is an i.i.d. sequence, independent of S_0^n, and where $\theta_* \in \mathbb{R}^d$, $d \geq 1$. At its two extremes, this model degenerates to one of two well-studied and fundamental models. When $\delta = 0$, the memory length is infinite, and the sign $S_0 = S_1 \cdots = \cdots = S_n$ is fixed. Thus, up to this sign ambiguity, the model (2) is the standard *Gaussian location model* (GLM), which is essentially a memoryless model (and exactly so if S_0 is known). When $\delta = \frac{1}{2}$, the signs S_0^n are i.i.d. and have no memory at all. The model (2) is then a *Gaussian mixture model* (GMM) with two symmetric components, which is also a memoryless model. In all other cases, $0 < \delta < \frac{1}{2}$ (or $\frac{1}{2} < \delta < 1$), the model is a simple version of a *hidden Markov model* (HMM).

The inference problem we consider in this paper is the estimation of $\theta_* \in \mathbb{R}^d$, under the loss function

$$\text{loss}(\hat{\theta}, \theta_*) := \min\{\|\hat{\theta} - \theta_*\|, \|\hat{\theta} + \theta_*\|\}, \tag{3}$$

that is, the Euclidean distance error under a possible sign ambiguity.[1] An intermediate goal (or an additional problem) is to estimate δ, under the regular absolute error loss function $|\hat{\delta} - \delta|$. The fundamental limits of this estimation problem will be gauged by the *local* minimax rate, which is the maximal decrease rate of the loss possible for any estimator, given n samples, at dimension $d \geq 2$, for signal strength $\|\theta_*\| = t$, and under flip probability δ. Specifically, for $d \geq 2$ it is defined as

$$\mathsf{M}(n, d, \delta, t) := \inf_{\hat{\theta}(X_1^n)} \sup_{\|\theta_*\| = t} \mathbb{E}\left[\text{loss}(\theta_*, \hat{\theta}(X_1^n)) \right]. \tag{4}$$

For general $d \geq 1$, the *global* minimax rate is defined with the condition $\|\theta_*\| = t$ replaced by $\|\theta_*\| \leq t$ (this condition trivializes the estimator for $d = 1$).

1.2 Known minimax estimation errors rates for GLM and GMM

Before delving into models with memory ($0 < \delta < 1/2$), we review known results on the minimax rates in memoryless high dimensional Gaussian models – the GLM ($\delta = 0$, the GMM ($\delta = 1/2$). We refer the reader to note that the minimax error rate in these models may undergo two possible phase transitions – one as t increases, and the other one as d increases. We also remark that the regime of interest is that of low-separation ($t \lesssim 1$), in which accurately detecting the components is impossible, yet parameter estimation with vanishing loss is possible.

---

[1] Similar bounds can be derived for the squared loss by trivial extensions.

At the first extreme, the local minimax rate for the GLM ($\delta = 0$) is the usual parametric error rate

$$
\mathsf{M}_{\mathrm{GLM}}(n,d,t) := \mathsf{M}(n,d,0,t) \asymp \begin{cases} t, & t \leq \sqrt{\frac{d}{n}} \\ \sqrt{\frac{d}{n}}, & t \geq \sqrt{\frac{d}{n}} \end{cases} . \tag{5}
$$

This is achieved by the trivial estimator $\hat{\theta} = 0$ if $t \leq \sqrt{\frac{d}{n}}$ and the simple empirical average estimator $\hat{\theta} = \frac{1}{n}\sum_{i=1}^{n} X_i$ if $t \geq \sqrt{\frac{d}{n}}$ (see [Wu, 2017, Sec. III-9]). The rate $\Theta(\sqrt{\frac{d}{n}})$ is then the *global* minimax rate, i.e., the largest error over $t = \|\theta_*\| > 0$. This model does not have a phase transition with dimension. At the other extreme, the GMM model ($\delta = \frac{1}{2}$) undergoes a phase transition at the dimension $d = n$. At low dimension, $d \leq n$, the minimax rate was neatly shown in [Wu and Zhou, 2019, Appendix B] to be

$$
\mathsf{M}_{\mathrm{GMM}}(n,d,t) \equiv \mathsf{M}\left(n,d,\frac{1}{2},t\right) \asymp \begin{cases} t, & t \leq \left(\frac{d}{n}\right)^{1/4} \\ \frac{1}{t}\sqrt{\frac{d}{n}}, & \left(\frac{d}{n}\right)^{1/4} \leq t \leq 1 \\ \sqrt{\frac{d}{n}}, & t > 1 \end{cases} , \tag{6}
$$

whereas at high dimension $d \geq n$, it is as for the Gaussian location model in (5), i.e., $\mathsf{M}_{\mathrm{GMM}}(n,d,t) = \mathsf{M}_{\mathrm{GLM}}(n,d,t)$. However, at this high dimensional regime, the loss is also lower bounded by a constant even if the signal strength t is unbounded. The loss in (6) is achieved by the trivial estimator $\hat{\theta} = 0$ if $t \leq \left(\frac{d}{n}\right)^{1/4}$ and by an estimator given by a properly scaled and shifted principal component of the empirical covariance matrix of $X_1^n := (X_1, X_2, \ldots, X_n)$, if $t \geq \left(\frac{d}{n}\right)^{1/4}$. For the GMM at low dimension, $d \leq n$, the global minimax rate is $\left(\frac{d}{n}\right)^{1/4}$, which is worse than the minimax rate of the Gaussian location model $\sqrt{\frac{d}{n}}$.

Therefore, the GMM has worse estimation performance compared to the GLM from three aspects: First, at low dimension, $d \leq n$, it has a larger global minimax rate $\left(\frac{d}{n}\right)^{1/4}$ compared to the parametric error rate of the GLM, $\sqrt{\frac{d}{n}}$; Second, at low dimension, $d \leq n$, parametric error rate is achieved only for constant separation $t \geq 1$; Third, the transition to the high dimension regime occurs at $d = n$.

As is intuitively appealing from a "data-processing" reasoning, a Markov model with flip probability δ' should allow for lower estimation error of θ_* compared to a Markov model with $\delta > \delta'$. Indeed, and as a specific simple example, any Markov model with $\delta < \frac{1}{2}$ can be easily transformed to a GMM model by randomizing the signs of each of the samples by an independent Rademacher variable. Thus we may deduce, e.g., that since at high dimension ($d \geq n$) the GLM and the GMM have the same minimax rates, the minimax rates for $d \geq n$ are in fact as in (5) for *any* $\delta \in [0, 1]$. We thus henceforth exclusively focus on the regime $d \leq n$. As we show, it is generally true that the improvement in estimation error when δ is reduced is less profound as the dimension increases.

1.3 Contributions

We first consider the case in which δ is known to the estimator of θ_*. For this case, we analyze the loss of an estimator that is based on a computation of the principal component of a properly chosen empirical covariance matrix. We show (Theorem 1) that, at low dimension, $d \leq \delta n$, it achieves local minimax rate of

$$
\mathsf{M}(n,d,\delta,t) \lesssim \begin{cases} t, & t \leq \left(\frac{\delta d}{n}\right)^{1/4} \\ \frac{1}{t}\sqrt{\frac{\delta d}{n}}, & \left(\frac{\delta d}{n}\right)^{1/4} \leq t \leq \sqrt{\delta} \\ \sqrt{\frac{d}{n}}, & t \geq \sqrt{\delta} \end{cases} , \tag{7}
$$

and at high dimension $d \geq \delta n$, it is as for the Gaussian location model in (5). The rate of this estimator is then further shown to be asymptotically optimal (up to a logarithmic factor) via a minimax lower bound (Theorem 2). Evidently, the loss in (7) smoothly interpolates the rates of the GLM in (5) and the GMM in (6). Moreover, it is evident that the loss is improved with the decrease of δ from all three aspects previously mentioned. First, at low dimension, $d \leq \delta n$, the global minimax rate is $\left(\frac{\delta d}{n}\right)^{1/4}$

(obtained by equating the first and second cases in (7)). Hence, reducing the flip probability from $\delta = \Theta(1)$ to $\delta = \frac{d}{n}$ smoothly reduces the global minimax rate from the parametric error rate $\Theta(\sqrt{\frac{d}{n}})$ of the GLM to the $\Theta\left(\left(\frac{d}{n}\right)^{1/4}\right)$ of the GMM. Second, at low dimension, $d \leq \delta n$, the minimal signal strength required to obtain parametric error rate is $t = \Theta(\sqrt{\delta})$ (obtained from the third case in (7)). This, again, improves by decreasing δ, and matches the extremes $t = \Theta(\sqrt{\frac{d}{n}})$ of GLM and $t = \Theta(1)$ of GMM. Third, the transition to the high dimension regime occurs at $d = \delta n$, which is again better with lower δ, and matches the transition point of the GMM given by $d = n$. This lower transition point allows us to achieve the error rate of the GLM for any signal strength, even in a regime in which the loss $O(\sqrt{\frac{d}{n}})$ vanishes with $n \to \infty$ (unlike for GMM, $\delta = \frac{1}{2}$); specifically, this occurs whenever $d \geq \delta n$ yet $d = o(n)$. Beyond the formal proof, Appendix A provides a heuristic justification for why the minimax error is naturally expected to scale as in (7).

Second, as a step towards the removal of the assumption that δ is known to the estimator, we consider the complementary problem of estimating δ whenever an estimate $\theta_\sharp$ of θ_* is available (which can be either exact $\theta_\sharp = \theta_*$, or inaccurate $\theta_\sharp \neq \theta_*$). We propose a simple estimator for δ, and analyze its error in case of a mismatch (Theorem 4). We then specify this result to the matched case $\theta_\sharp = \theta_*$ and show that in the non-trivial regime ($\|\theta_*\| \lesssim 1$) its error rate is $\tilde{O}(\frac{1}{\|\theta_*\|^2}\sqrt{\frac{1}{n}})$. We then proceed to show an impossibility lower bound of $\Omega(\sqrt{\frac{1}{n}})$ (Proposition 6) for this error rate. The precise dependence of the estimation error rate of δ on $\|\theta_*\|$ is therefore not precisely determined, and we discuss the challenges in settling this matter.

Third, we consider the case in which the estimator of θ_* has no prior knowledge of δ. We propose a three step algorithm for this case (Algorithm 1). First, a (possibly) gross estimate $\hat{\theta}^{(A)}$ of θ_* is computed based on third of the samples, assuming the worst case of $\delta = \frac{1}{2}$. Then, an estimate $\hat{\delta}^{(B)}$ of δ is computed using another third of the samples, assuming the estimate $\hat{\theta}^{(A)}$. Finally, a refined estimate $\hat{\theta}^{(C)}$ of θ_* is obtained by (essentially) assuming that δ is $\hat{\delta}^{(B)}$. At each of the steps above, the algorithm may stop and decide to return its current estimate when it determines that no further improvement is possible by moving on to the next steps. We analyze the estimation loss of this algorithm (Theorem 7), and show that this algorithm is capable of partially achieving the gains associated with the case of known δ.

1.4 Additional related work

Both the GLM [Johnstone, 2002, Tsybakov, 2008] and GMM [Lindsay, 1995, McLachlan et al., 2019] are classic models which were well-explored from numerous perspectives. In the last few years, there is a surge of interest in the non-asymptotic performance analysis of computationally efficient estimation algorithms for this estimation task. For example, Moitra and Valiant [2010], Kalai et al. [2010], Anandkumar et al. [2014], Hardt and Price [2015], Wu and Yang [2020] have analyzed method-of-moments-based algorithms, and various other papers considered the expectation-maximization (EM) algorithm [Balakrishnan et al., 2017, Xu et al., 2016, Jin et al., 2016, Klusowski and Brinda, 2016, Weinberger and Bresler, 2022, Dwivedi et al., 2020b,a, 2018, Zhao et al., 2018, Yan et al., 2017]. Specifically, the local minimax rate for GMM in (6) was determined in [Wu and Zhou, 2019] as a benchmark for the operation of the EM algorithm.

The model (2) is a simple instance of a HMM [Ephraim and Merhav, 2002, van Handel, 2008] in high dimension. Parameter estimation in such models is practically performed via the Baum-Welch algorithm [Baum et al., 1970], which is a computationally efficient version of EM for HMMs. To the best of our knowledge, there were little attempts to characterize the minimax rates in such models with the exception of [Yang et al., 2015] previously mentioned. However, the qualifying condition for the estimation error bound of [Yang et al., 2015] on EM is that $t = \|\theta_*\| \gtrsim \log \frac{1}{1-(1-2\delta)^2}$, that is, a non-trivial separation when δ is constant, which further blows up as $\delta \downarrow 0$. By contrast, in this paper, our goal is to characterize the estimation error in the regime of δ and $t = \|\theta_*\|$ in which the minimax rate is affected by these parameters, and this requires analyzing vanishing δ and t.

More broadly, there is a growing interest in advancing the quantitative understanding of the performance of statistical learning and inference with dependent data. Bresler et al. [2020] studied

linear regression with Markovian covariates and characterized the minimax error rate in terms of the mixing time of the Markov chain. A stochastic gradient descent-style algorithm adapted to the Markov setting was shown to be minimax optimal. Statistical estimation problems including linear and logistic regression with more general network dependencies among response variables were studied by Daskalakis et al. [2019] and Kandiros et al. [2021]. Learnability and generalization bounds were derived by Dagan et al. [2019] for dependent data satisfying the so-called Dobrushin's condition.

1.5 Notation conventions

For a vector $v \in \mathbb{R}^d$, $\|v\|$ is the Euclidean norm. For a positivedefinite matrix A, $\lambda_{\max}(A)$ and $v_{\max}(A)$ are the maximal eigenvalue and the associated eigenvector (of unit norm) of A. Unless otherwise stated, the constants involved in Bachmann-Landau notation are numerical, and do not depend on the parameters (n, d, δ, t). It holds that $a \gtrsim b$ (resp. $a \lesssim b$) if there exists a constant $c > 0$ (resp. $C > 0$) such that $a \geq cb$ (resp. $a \leq Cb$). If $a \lesssim b$ and $a \gtrsim b$. then $a \asymp b$. Integer constraints (ceiling and floor) on large quantities that do not affect the results are omitted for brevity. For a real numbers a, b the shorthand notation $a \vee b := \max\{a, b\}$, $a \wedge b := \min\{a, b\}$ and $(a)_+ := a \vee 0$ is used. A sequence of objects $X_1, \cdots, X_n$ is denoted by X_1^n. Expectation, variance and probability are denoted by $\mathbb{E}, \mathbb{V}$ and $\mathbb{P}$, respectively. Equality in distribution of random variables X and Y is denoted by $X \stackrel{d}{=} Y$. All logarithms $\log$ are to the base e.

2 Mean estimation for a known flip probability

In this section, we consider the problem of estimating θ_* whenever δ is exactly known to the estimator. In that case, it may be assumed w.l.o.g. that $\delta \in [0, \frac{1}{2}]$, as otherwise one may negate each of the even samples to obtain an equivalent model with δ replaced with $1 - \delta$. Hence also $\rho \in [0, 1]$. We next describe an estimator for this task, state a bound on its performance, and then show that it matches (up to a logarithmic factor) an impossibility result.

The estimator operationally interpolates and therefore simultaneously generalizes the empirical average estimator (21) and the (properly scaled) principal component estimator (23) analyzed in [Wu and Zhou, 2019, Appendix B]. It degenerates to the latter estimators if $\delta \downarrow 0$ or $\delta \uparrow \frac{1}{2}$. Specifically, the estimator partitions the sample into blocks of equal length k each (which will later be set to $k = \frac{1}{8\delta}$, according to the mixing time of the Markov chain S_0^n). Let ℓ denote the number of blocks respectively, so that $k\ell = n$. Let $\mathcal{I}_i = \{(i-1)k + 1, (i-1)k + 2, \cdots, ik\}$ denote the indices of the ith block. Further, let $\{R_i\}_{i \in [\ell]}$ be an i.i.d. Rademacher sequence ($R_i \sim \text{Uniform}\{-1, 1\}$), and let $\overline{X}_i := R_i \cdot \frac{1}{k} \sum_{j \in \mathcal{I}_i} X_j \stackrel{d}{=} \overline{S}_i \theta_* + \overline{Z}_i$ denote the average of the samples in the ith block (randomized with a sign R_i), where $\overline{S}_i := R_i \cdot \frac{1}{k} \sum_{j \in \mathcal{I}_i} S_j$ is the *gain* (average of the signs) of the ith block, and $\overline{Z}_i := R_i \cdot \frac{1}{k} \sum_{j \in \mathcal{I}_i} Z_j \sim N\left(0, \frac{1}{k} \cdot I_d\right)$ is the average noise of the ith block. Due to the sign randomization, it holds that $\{\overline{S}_i\}_{i \in [\ell]}$ is an i.i.d. sequence. Since $\{\overline{Z}_i\}_{i \in [\ell]}$ is also an i.i.d. sequence, then so is $\{\overline{X}_i\}_{i \in [\ell]}$. For notational simplicity we will omit the block index i of a generic block. For block length k, we denote by $\xi_k := \mathbb{E}[\overline{S}^2] = \mathbb{E}[(\frac{1}{k} \sum_{j=1}^k S_j)^2]$ the second moment of the gain $\overline{S}$. Note that $\xi_k \in [\frac{1}{k}, 1]$ for any $\delta \in [0, \frac{1}{2}]$ and is in particular always positive. For a sequence of samples $X_1^n = (X_1, \ldots, X_n)$, we define by $\hat{\Sigma}_{n,k}(X_1^n)$ the empirical covariance matrix of the averaged samples over blocks $\{\overline{X}_i\}_{i \in [\ell]}$, that is $\hat{\Sigma}_{n,k}(X_1^n) := \frac{1}{\ell} \sum_{i=1}^{\ell} \overline{X}_i \overline{X}_i^\top$, whose population average is $\Sigma_{n,k}(\theta_*)$, where $\Sigma_{n,k}(\theta_*) := \mathbb{E}[\overline{X}\,\overline{X}^\top] = \xi_k \theta_* \theta_*^\top + \frac{1}{k} I_d$. We note that θ is the principal component of $\Sigma_{n,k}(\theta)$, that is, $\lambda_{\max}(\Sigma_{n,k}(\theta)) = \xi_k \|\theta\|^2 + \frac{1}{k}$ and the corresponding eigenvector is $v_{\max}(\Sigma_{n,k}(\theta)) = \theta$. We thus consider the following estimator for θ_*, from a sequence X_1^n, and with a block length of k

$$\hat{\theta}_{\text{cov}}(X_1^n; k) := \sqrt{\frac{1}{\xi_k}\left(\lambda_{\max}(\hat{\Sigma}_{n,k}(X_1^n)) - \frac{1}{k}\right)_+} \cdot v_{\max}\left(\hat{\Sigma}_{n,k}(X_1^n)\right). \tag{8}$$

The estimator is thus constructed from two types of averages: First, a coherent average of the samples at each block, to obtain ℓ block-samples $\overline{X}_i$ with gain $\overline{S}_i$ and noise variance reduced by a factor of k. Second, an incoherent average of the "square" of the ℓ block-samples $\overline{X}_i \overline{X}_i^\top$, which resolves

the remaining sign ambiguity between blocks. This balance two extreme cases: If $\delta = 0$, then this reduces the problem to the GLM (with a sign ambiguity) and $k = n$ is an optimal choice. If $\delta = \frac{1}{2}$, then this reduces the problem to the GMM, in which coherent averaging is non-beneficial and $k = 1$ is rate optimal. Generally, the optimal choice of the block length k is proportional to the mixing time of the Markov chain $\Theta(\frac{1}{\delta})$. This choice assures that the random gain $\overline{S}$ is ± 1 with a (constant) high probability. In fact, an elementary, yet crucial, part of the analysis establishes that the random gain $\overline{S}$ has constant variance for this choice of block length (see Lemma 9 in Appendix B.1). On the other hand, if $k = \Omega(\frac{1}{\delta})$, then the random gain $\overline{S}$ will not be ± 1 (or not even bounded away from zero) with high probability, and such choice is never efficient. Specifically, we consider the estimator in (8) with $k = \frac{1}{8\delta}$. The above estimation procedure is depicted in Figure 2 in Appendix B.1.

Let us denote $\beta(n, d, \delta) := \sqrt{\frac{d}{n}} \vee \left(\frac{\delta d}{n}\right)^{1/4}$, which will actually be the global minimax rate.

Theorem 1. *Assume that $\delta \geq \frac{1}{n}$ and $d \leq n$, and set $\hat{\theta} \equiv \hat{\theta}_{\mathrm{cov}}(X_1^n; k)$ with $k = \frac{1}{8\delta}$. Then, there exist numerical constants $c_0, c_1, c_2 > 0$ such that for every $\theta_* \in \mathbb{R}^d$*

$$\mathbb{E}\left[\mathsf{loss}(\hat{\theta}, \theta_*)\right] \leq c_0 \cdot \begin{cases} \beta(n, d, \delta), & \|\theta_*\| \leq \beta(n, d, \delta) \\ \sqrt{\frac{d}{n}} + \frac{1}{\|\theta_*\|}\sqrt{\frac{\delta d}{n}} + \frac{1}{\|\theta_*\|} \cdot \frac{d}{n}, & \beta(n, d, \delta) \leq \|\theta_*\| \end{cases} \tag{9}$$

and $\mathsf{loss}(\hat{\theta}, \theta_*) \leq c_1 \cdot \log(n) \cdot \mathbb{E}\left[\mathsf{loss}(\hat{\theta}, \theta_*)\right]$ with probability larger than $1 - \frac{c_2}{n}$.

Theorem 1 is proved in Appendix B.1. Evidently, Theorem 1 implies that the upper bound on the minimax rate stated in (7) above holds in low dimension, $d \leq \delta n$, whereas $\mathsf{M}(n, d, \delta, t) \asymp \mathsf{M}_{\mathrm{GLM}}(n, d, t)$ holds in high dimension $d \geq \delta n$.[2] We remark that the condition $\delta \geq \frac{1}{n}$ is mild as otherwise the model (2) is essentially equivalent to GLM. See Remark 8 in Appendix B.1. Numerical validation of the performance of the estimator $\hat{\theta}_{\mathrm{cov}}(X_1^n; k)$ is shown in Appendix F.

We next consider an impossibility result. As we have seen, at high dimension, $d \geq \delta n$, the minimax error rates achieved are the same as for the Gaussian location model, and thus clearly cannot be improved. We thus next focus on the low dimensional regime $d \leq \delta n$.

Theorem 2. *Assume that $2 \leq d \leq \delta n$ and $n \geq \frac{128}{d}$. Then the local minimax rate is bounded as*

$$\mathsf{M}(n, d, \delta, t) \gtrsim \frac{1}{\sqrt{\log(n)}} \cdot \begin{cases} t, & t \leq \left(\frac{\delta d}{n}\right)^{1/4} \\ \frac{1}{t}\sqrt{\frac{\delta d}{n}}, & \left(\frac{\delta d}{n}\right)^{1/4} \leq t \leq \sqrt{\delta} \\ \sqrt{\frac{d}{n}}, & t \geq \sqrt{\delta} \end{cases} \tag{10}$$

Hence, the minimax rates achieved by the estimator in Theorem 1 are nearly asymptotically optimal, up to a $\sqrt{\log(n)}$ factor. The full proof of Theorem 2 together with a summary of the main ideas used in the proof is presented in Appendix B.2.

Remark 3 (Relaxation of the noise distribution assumption). For the sake of concreteness and clarity of exposition, we have assumed in Theorems 1 and 2 that the noise samples $\{Z_i\}_{i=1}^n$ are i.i.d. isotropic Gaussians. There are two straightforward relaxations of this assumption. First, our minimax upper bound (Theorem 1) can be proved to any subGaussian noise distribution, simply because all the concentration bounds for Gaussian random variables used in the proof admit subGaussian analogues. The impossibility result (converse, Theorem 2) trivially holds for subGaussian noise since Gaussians are special case of subGaussians. Second, the isotropic assumption can be relaxed to anisotropic noise with known covariance Σ by simple standardization: If $Z_i \sim N(0_d, \Sigma)$ are i.i.d. for some *known* $\Sigma \succ 0$, then the estimator will multiply the samples by $\Sigma^{-1/2}$ and reduce the problem back to the isotropic setting. After applying the estimator we propose for the isotropic case, the estimator will obtain its final estimate by multiplying its isotropic estimate by $\Sigma^{1/2}$) (However, the loss of

---

[2]When $\|\theta_*\| \leq \beta(n, d, \delta)$, the estimator $\hat{\theta}_{\mathrm{cov}}(X_1^n; k)$ in Theorem 1 only achieves a rate $\beta(n, d, \delta)$ which is larger than the promised rate $\|\theta_*\|$ in (7). However, since the estimator is assumed to know t (but not the direction of θ_*; a common formulation in high-dimensional statistics), then it can output the zero vector. It then incurs loss $\|\theta\|_*$ for any $\theta_* \in \mathbb{R}^d$, matching the promised rate when $\|\theta_*\| \leq \beta(n, d, \delta)$. To summarize, for any value of t, the minimax rate is achieved by the minimum rate of $\hat{\theta}_{\mathrm{cov}}(X_1^n; k)$ and $\hat{\theta}_0(X_1^n) \equiv 0$.

249 the estimator will then be gauged by the Mahalanobis distance, parameterized by Σ). We refer the
250 reader to Appendix G for a discussion on more challenging directions in which the isotropic Gaussian
251 assumption can be relaxed.

3 Flip probability estimation for a given estimator of θ_*

253 In this section, we consider the problem of estimating δ whenever θ_* is approximately known to be
254 $\theta_\sharp$. We propose a simple estimator, and then discuss the importance of the accuracy of θ_*. We then
255 derive an impossibility result for the matched case, $\theta_\sharp = \theta_*$.

256 First note that an estimator for δ can be easily obtained from an estimator for ρ, with essentially the
257 same error rate, via $\hat{\delta} = \frac{1}{2}(1 - \hat{\rho})$. Thus we focus on estimating ρ. Assume for simplicity that n
258 is even. Observing that $\mathbb{E}[X_{2i}^\top X_{2i+1}] = \rho\|\theta_*\|^2$, we propose the following natural estimator for ρ,
259 which replaces the population average with empirical average:

$$\hat{\rho}_{\text{corr}}(X_1^n; \theta_\sharp) = \frac{1}{\|\theta_\sharp\|^2} \cdot \frac{2}{n} \sum_{i=1}^{n/2} X_{2i}^\top X_{2i-1}. \tag{11}$$

260 That is, the estimator is based on evaluating the correlation of each of two adjacent samples X_{2i} and
261 X_{2i-1}. We first state a general bound on the estimation error of this estimator. We then consider the
262 case in which θ_* is known, and show how the estimation error is improved in this case.

263 **Theorem 4.** *Assume that $d \leq n$. Let $\theta_* \in \mathbb{R}^d$ and let $\theta_\sharp$ be an estimate of θ_*. Set $\hat{\rho} \equiv \hat{\rho}_{\text{corr}}(X_1^n; \theta_\sharp)$*
264 *and $\hat{\delta} = \frac{1}{2}(1 - \hat{\rho})$. Then, it holds with probability $1 - \frac{8}{n}$ that*

$$\left|\hat{\delta} - \delta\right| = \frac{1}{2}\left|\hat{\rho} - \rho\right| \leq \frac{\left|\|\theta_*\|^2 - \|\theta_\sharp\|^2\right|}{\|\theta_\sharp\|^2} + 16\log(n)\left[\sqrt{\frac{\delta}{n}} + \frac{1}{\|\theta_\sharp\|}\sqrt{\frac{1}{n}} + \frac{1}{\|\theta_\sharp\|^2}\sqrt{\frac{d}{n}}\right]. \tag{12}$$

265 The proof of Theorem 4 appears in Appendix C.1. Note that Theorem 4 states a high-probability
266 bound, suitable to its usage later on in Section 4. A bound on the expectation of the error can be
267 obtained by standard methods (integrating tails).

268 **The effect of knowledge of θ_*** If θ_* is known up to a sign, i.e., $\theta_\sharp = \pm\theta_*$, then for the purpose of ρ
269 (or equivalently δ) estimation, the model (2) can be reduced to a one-dimensional model by rotational
270 invariance of isotropic Gaussian (See additional details in Appendix C.2). It then immediately follows
271 from Theorem 4 that:

272 **Corollary 5.** *Assume that $d \leq n$, $\|\theta_*\| \leq 1$ and $\theta_\sharp = \pm\theta_*$. Let U_1^n be defined in as $U_i :=$*
273 *$\|\theta_*\| \cdot S_i + W_i$ where $W_i \sim N(0,1)$ i.i.d., $\hat{\rho} \equiv \hat{\rho}_{\text{corr}}(U_1^n; \theta_\sharp)$ and $\hat{\delta} = \frac{1}{2}(1 - \hat{\rho})$. Then it holds with*
274 *probability $1 - \frac{8}{n}$ that*

$$\left|\hat{\delta} - \delta\right| = \frac{1}{2}\left|\hat{\rho} - \rho\right| \leq \frac{18\log(n)}{\|\theta_*\|^2}\sqrt{\frac{1}{n}}. \tag{13}$$

275 Numerical validation of the performance of the estimator $\hat{\delta}_{\text{corr}}(X_1^n; \theta_\sharp) = \frac{1}{2}(1 - \hat{\rho}_{\text{corr}}(X_1^n; \theta_\sharp))$ in the
276 mismatched (Theorem 4) and matched (Corollary 5) cases is provided in Appendix F.

277 We next consider an impossibility lower bound.

278 **Proposition 6.** *Suppose that $\theta_\sharp = \theta_*$ and $\|\theta_*\| \leq \frac{1}{\sqrt{2}}$. Then $\inf_{\hat{\delta}(U_1^n)} \sup_{\delta \in [0,1]} \mathbb{E}[|\delta - \hat{\delta}(U_1^n)|] \geq \frac{1}{32\sqrt{n}}$,*

279 *where the infimum is over any estimator $\hat{\delta}(U_1^n)$ based on the model $U_i = \|\theta_*\|S_i + W_i$ where each*
280 *W_i is i.i.d. $N(0,1)$.*

281 The proof of Proposition 6 is presented in Appendix C.2. According to Corollary 5 and Proposition
282 6, in estimating δ with a known θ_*, though the dependence $\Theta(\frac{1}{\sqrt{n}})$ of the minimax error rate on
283 the sample size is shown to be nearly optimal, it is unclear what the optimal dependence on the
284 signal strength should be. This is left as an interesting open question and we discuss the challenges
285 associated with this problem in Appendix C.2.

4 Mean estimation under an unknown flip probability

As we have seen, if an estimator for θ_* knows the value of δ, and if both $\delta \leq \frac{1}{2}$ and $d \leq \delta n$ hold, then the estimator can achieve improved error rates over the GMM case ($\delta = \frac{1}{2}$). In this section, we assume that both θ_* and δ are unknown, and so the estimator is required to estimate δ in order to use this knowledge for an estimator of θ_*. We propose an estimation procedure of three steps based on sample splitting of $3n$ samples. We mention at the outset that the regime in which improvement is possible will be for low signal strength $\|\theta_*\| \lesssim 1$ (low separation between the components), and up to a dimension which depends on δ. Of course the estimation procedure does not know (θ_*, δ) in advance, and so it is required to identify if (θ_*, δ) are in this regime during its operation.

We now begin with an overview of the steps of the estimation algorithm. At Step A, the algorithm estimates θ_* based on X_1^n assuming a Gaussian mixture model ($\delta = \frac{1}{2}$) to obtain an estimate $\hat{\theta}^{(A)}$. Then, based on $\|\hat{\theta}^{(A)}\|$, the algorithm decides whether improvement is potentially possible had δ was known. There are two cases. The first case is that $\|\hat{\theta}^{(A)}\|$ is too low, and then its estimate is not sufficiently accurate to be used in the next steps. Essentially, this happens when the norm is below the global minimax rate $(\frac{d}{n})^{1/4}$, and the estimation error of the norm on the same scale as the norm of $\|\theta_*\|$. A trivial estimator of $\hat{\theta} = 0$ is then optimal in terms of error rates. It can be already noted at this step that while the global minimax rate for the known δ case is $(\frac{\delta d}{n})^{1/4}$, here the algorithm already stops and estimates $\hat{\theta} = 0$ even if just $\|\theta_*\| \lesssim (\frac{d}{n})^{1/4}$, leading to larger global minimax rate. The second case is that $\|\theta_*\|$ is larger than a constant. In this case, the estimation based on a GMM already achieves the optimal parametric $O(\sqrt{\frac{d}{n}})$ error rate of the Gaussian location model, and so no further estimation steps are necessary. Otherwise, an improvement in the estimation is possible. The algorithm proceeds to Step B, and uses X_{n+1}^{2n} to obtain an estimate $\hat{\delta}^{(B)}$ of δ based on the mismatched $\theta_\sharp \equiv \hat{\theta}^{(A)}$. Then, based on the estimate $\hat{\delta}^{(B)}$ the algorithm decides whether the accuracy of $\hat{\delta}^{(B)}$ is sufficient to be used in an refined estimation of θ_*. If the accuracy of $\hat{\delta}^{(B)}$ is not good enough, then the algorithm outputs the estimate from Step A, that is $\hat{\theta}^{(A)}$. Otherwise, it proceeds to Step C, in which θ_* is re-estimated using X_{2n+1}^{3n}, based on a mismatched choice of k, that is $k \asymp \frac{1}{\hat{\delta}^{(B)}}$ instead of $k = \frac{1}{8\delta}$. Intuitively, the estimated value $\hat{\delta}^{(B)}$ should be larger than δ so the resulting block size $k \asymp \frac{1}{\hat{\delta}^{(B)}}$ will be such that the gain in the block is still close to 1 with high probability. On the other hand, it is desired that $\hat{\delta}^{(B)}$ will be on the same scale as δ so that the estimation rate (295) (see also (7)) – which now essentially holds with $\hat{\delta}^{(B)}$ instead of δ – would be as small as possible. Thus, if the algorithm has assured in Step B that $\hat{\delta}^{(B)} \asymp \delta$, then at Step C it will achieve the error rate indicated in (295).

The formal description of the estimation algorithm is provided in Algorithm 1. We remark that refining the estimation of δ can be easily incorporated as a fourth step of this algorithm, but we do not present this in order to keep the statement of the result simple. The error of the estimator output by Algorithm 1 is as follows:

Theorem 7. *There exist constants $c_1, c_2 \geq 0$ and $\lambda_\theta, \lambda_\delta \geq 1$ such that if $d \leq \frac{n}{4\lambda_\theta^2 \log^2(n) \wedge 16}$ then the output $\hat{\theta}$ of Algorithm 1 satisfies for any $\theta_* \in \mathbb{R}^d$, with probability $1 - O(\frac{1}{n})$: If $d \leq \frac{1}{64\lambda_\delta^2 \lambda_\theta^2 \log^2(n)} \delta^4 n$ then*

$$\mathsf{loss}(\hat{\theta}, \theta_*) \leq c_1 \log n \cdot \begin{cases} \|\theta_*\|, & \|\theta_*\| \leq \lambda_\theta \log(n) \left(\frac{d}{n}\right)^{1/4} \\ \frac{1}{\|\theta_*\|}\sqrt{\frac{d}{n}}, & \lambda_\theta \log(n) \left(\frac{d}{n}\right)^{1/4} \leq \|\theta_*\| \leq \sqrt{8\lambda_\delta \lambda_\theta \log(n)} \left(\frac{d}{\delta^2 n}\right)^{1/4} \\ \frac{1}{\|\theta_*\|}\sqrt{\frac{\delta d}{n}}, & \sqrt{8\lambda_\delta \lambda_\theta \log(n)} \left(\frac{d}{\delta^2 n}\right)^{1/4} \leq \|\theta_*\| \leq \sqrt{\delta} \\ \sqrt{\frac{d}{n}}, & \|\theta_*\| \geq \sqrt{\delta} \end{cases} \tag{17}$$

If $d \geq \frac{1}{64\lambda_\delta^2 \lambda_\theta^2 \log^2(n)} \delta^4 n$ then

$$\mathsf{loss}(\hat{\theta}, \theta_*) \leq c_2 \log n \cdot \begin{cases} \|\theta_*\|, & \|\theta_*\| \leq \lambda_\theta \log(n) \left(\frac{d}{n}\right)^{1/4} \\ \frac{1}{\|\theta_*\|}\sqrt{\frac{d}{n}}, & \lambda_\theta \log(n) \left(\frac{d}{n}\right)^{1/4} \leq \|\theta_*\| \leq \sqrt{8\lambda_\delta \lambda_\theta \log(n)} \left(\frac{d}{\delta^2 n}\right)^{1/4} \\ \sqrt{\frac{d}{n}}, & \|\theta_*\| \geq \sqrt{8\lambda_\delta \lambda_\theta \log(n)} \left(\frac{d}{\delta^2 n}\right)^{1/4} \end{cases} \tag{18}$$

Algorithm 1 Mean estimation for unknown δ

1: **input:** Parameters $\lambda_\theta, \lambda_\delta > 0$ (from (295) (296) (298)), $3n$ data samples X_1^{3n} from the model (2)

2: **step A:** Estimate θ_* assuming a Gaussian mixture model

$$\hat{\theta}^{(A)} \equiv \hat{\theta}_{\mathrm{cov}}(X_1^n; k = 1) \tag{14}$$

3: **if** $\|\hat{\theta}^{(A)}\| \leq 2\lambda_\theta \cdot \log(n) \cdot (\frac{d}{n})^{1/4}$ **then**

4:     **return** $\hat{\theta} = 0$ $\triangleright$ No further improvement can be guaranteed

5: **else if** $\|\hat{\theta}^{(A)}\| \geq \frac{1}{2}$ **then**

6:     **return** $\hat{\theta} = \hat{\theta}^{(A)}$ $\triangleright$ No further improvement is possible

7: **end if**

8: **step B:** Estimate δ assuming a mismatched mean value of $\hat{\theta}^{(A)}$

$$\hat{\delta}^{(B)} = \hat{\delta}_{\mathrm{corr}}(X_{n+1}^{2n}; \hat{\theta}^{(A)}) \tag{15}$$

9: **if** $\hat{\delta}^{(B)} \leq 64\lambda_\delta\lambda_\theta \frac{\log(n)}{\|\hat{\theta}^{(A)}\|^2}\sqrt{\frac{d}{n}}$ **then**

10:     **return** $\hat{\theta} = \hat{\theta}^{(A)}$ $\triangleright$ No further improvement can be guaranteed

11: **end if**

12: **step C**: Estimate θ_* assuming a Markov model with a mismatched flip probability $\hat{\delta}^{(B)}$

$$\hat{\theta}^{(C)} = \hat{\theta}_{\mathrm{cov}}\left(X_{2n+1}^{3n}; k = \frac{1}{16\hat{\delta}^{(B)}}\right) \tag{16}$$

13: **return** $\hat{\theta} = \hat{\theta}^{(C)}$.

Theorem 7 implies that up to logarithmic factors, the error rates of the known δ case are recovered for low enough dimension $d \lesssim \delta^4 n$ and $\|\theta_*\| \gtrsim \left(\frac{d}{\delta^2 n}\right)^{1/4}$. The analysis of Algorithm 1 appears in Appendix D. Numerical validation of the performance of Algorithm 1 can be found in Appendix F.

The impact of unknown δ on the estimation error of θ_*   Comparing Theorems 2 (known δ) and 7 (unknown δ) reveals the deterioration in the estimation of θ_* due to the lack of knowledge of δ from the three aspects mentioned in Section 1.3 (ignoring logarithmic factors): First, the global minimax rate is $(\frac{d}{n})^{1/4}$ as for the GMM, instead of the rate $(\frac{\delta d}{n})^{1/4}$ for the Markov model case with known δ. Second, at the regime $(\frac{d}{n})^{1/4} \lesssim \|\theta_*\| \lesssim (\frac{d}{\delta^2 n})^{1/4}$ the error rate is $O(\frac{1}{\|\theta_*\|}\sqrt{\frac{d}{n}})$ instead of the lower $O(\frac{1}{\|\theta_*\|}\sqrt{\frac{\delta d}{n}})$. Third, the algorithm is only effective when the dimension is as low as $d \lesssim \delta^4 n$. For higher dimensions, the rates of the GMM are achieved, which can be achieved even without the knowledge of δ.

5   Conclusion and future work

In this paper, we have considered an elementary, yet fundamental, high-dimensional model with memory. We have obtained a sharp bound on the minimax rate of estimation in case the underlying statistical dependency (flip probability) is known, and proposed a three-step estimation algorithm when it is unknown. This has revealed the gains possible in estimation rates due to the memory between the samples, and smoothly interpolated between the extreme cases of GLM and GMM. An interesting open problem is to either characterize the optimality of the algorithm or improving in the unknown δ case; this requires understanding optimal estimation of the flip probability, and the challenges associated with this problem are illuminated.

Naturally, as the model considered in this paper is basic, there is an ample of possibilities to generalize this model. These include, a larger number of components in the mixture, statistical dependency with a more complicated graphical structure between the data samples, existence of nuisance parameters such as the noise variance, sharp finite-sample/finite-iteration analysis of specific practical algorithms such as Baum-Welch, location-scale model with anisotropic noise, heavy-tailed noise, and so on.

References

Animashree Anandkumar, Rong Ge, Daniel Hsu, Sham M Kakade, and Matus Telgarsky. Tensor decompositions for learning latent variable models. *The Journal of Machine Learning Research*, 15(1):2773–2832, 2014.

Sivaraman Balakrishnan, Martin J. Wainwright, and Bin Yu. Statistical guarantees for the EM algorithm: From population to sample-based analysis. *The Annals of Statistics*, 45(1):77–120, 2017.

Leonard E. Baum, Ted Petrie, George Soules, and Norman Weiss. A maximization technique occurring in the statistical analysis of probabilistic functions of Markov chains. *The Annals of Mathematical Statistics*, 41(1):164–171, 1970.

Marianne Bertrand, Erzo FP Luttmer, and Sendhil Mullainathan. Network effects and welfare cultures. *The quarterly journal of economics*, 115(3):1019–1055, 2000.

N. M. Blachman and L. Few. Multiple packing of spherical caps. *Mathematika*, 10:84–88, 1963. ISSN 0025-5793. doi: 10.1112/S0025579300003405. URL https://doi.org/10.1112/S0025579300003405.

Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities: A nonasymptotic theory of independence*. Oxford university press, 2013.

Guy Bresler, Prateek Jain, Dheeraj Nagaraj, Praneeth Netrapalli, and Xian Wu. Least squares regression with markovian data: Fundamental limits and algorithms. In *Proceedings of the 34th International Conference on Neural Information Processing Systems*, NIPS'20, Red Hook, NY, USA, 2020. Curran Associates Inc. ISBN 9781713829546.

Yeshwanth Cherapanamjeri, Samuel B. Hopkins, Tarun Kathuria, Prasad Raghavendra, and Nilesh Tripuraneni. Algorithms for heavy-tailed statistics: regression, covariance estimation, and beyond. In Konstantin Makarychev, Yury Makarychev, Madhur Tulsiani, Gautam Kamath, and Julia Chuzhoy, editors, *Proccedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing, STOC 2020, Chicago, IL, USA, June 22-26, 2020*, pages 601–609. ACM, 2020. doi: 10.1145/3357713.3384329. URL https://doi.org/10.1145/3357713.3384329.

Nicholas A Christakis and James H Fowler. Social contagion theory: examining dynamic social networks and human behavior. *Statistics in medicine*, 32(4):556–577, 2013.

T. M. Cover and J. A. Thomas. *Elements of Information Theory*. Wiley-Interscience, 2006. ISBN 0471241954.

Yuval Dagan, Constantinos Daskalakis, Nishanth Dikkala, and Siddhartha Jayanti. Learning from weakly dependent data under dobrushin's condition. *CoRR*, abs/1906.09247, 2019. URL http://arxiv.org/abs/1906.09247.

Constantinos Daskalakis, Nishanth Dikkala, and Ioannis Panageas. Regression from dependent observations. In *STOC'19—Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing*, pages 881–889. ACM, New York, 2019.

Esther Duflo and Emmanuel Saez. The role of information and social interactions in retirement plan decisions: Evidence from a randomized experiment. *The Quarterly journal of economics*, 118(3): 815–842, 2003.

Raaz Dwivedi, Koulik Khamaru, Martin J Wainwright, Michael I. Jordan, et al. Theoretical guarantees for EM under misspecified Gaussian mixture models. In *Advances in Neural Information Processing Systems*, pages 9681–9689, 2018.

Raaz Dwivedi, Nhat Ho, Koulik Khamaru, Martin J. Wainwright, Michael I. Jordan, and Bin Yu. Challenges with EM in application to weakly identifiable mixture models. *CoRR*, abs/1902.00194, 2019. URL http://arxiv.org/abs/1902.00194.

Raaz Dwivedi, Nhat Ho, Koulik Khamaru, Martin Wainwright, Michael Jordan, and Bin Yu. Sharp analysis of expectation-maximization for weakly identifiable models. In *International Conference on Artificial Intelligence and Statistics*, pages 1866–1876. PMLR, 2020a.

Raaz Dwivedi, Nhat Ho, Koulik Khamaru, Martin J Wainwright, Michael I Jordan, and Bin Yu. Singularity, misspecification and the convergence rate of EM. *The Annals of Statistics*, 48(6): 3161–3182, 2020b.

Yariv Ephraim and Neri Merhav. Hidden Markov processes. *IEEE Transactions on information theory*, 48(6):1518–1569, 2002.

Thomas S. Ferguson. An inconsistent maximum likelihood estimate. *J. Amer. Statist. Assoc.*, 77(380):831–834, 1982. ISSN 0162-1459. URL http://links.jstor.org/sici?sici= 0162-1459(198212)77:380<831:AIMLE>2.0.CO;2-Y&origin=MSN.

Edward L Glaeser, Bruce Sacerdote, and Jose A Scheinkman. Crime and social interactions. *The Quarterly journal of economics*, 111(2):507–548, 1996.

László Györfi, Michael Kohler, Adam Krzyżak, and Harro Walk. *A distribution-free theory of nonparametric regression*. Springer Series in Statistics. Springer-Verlag, New York, 2002. ISBN 0-387-95441-4. doi: 10.1007/b97848. URL https://doi.org/10.1007/b97848.

Moritz Hardt and Eric Price. Tight bounds for learning a mixture of two Gaussians. In *Proceedings of the forty-seventh annual ACM symposium on Theory of computing*, pages 753–760. ACM, 2015.

Samuel B. Hopkins. Mean estimation with sub-Gaussian rates in polynomial time. *The Annals of Statistics*, 48(2):1193 – 1213, 2020. doi: 10.1214/19-AOS1843. URL https://doi.org/10. 1214/19-AOS1843.

Samuel B. Hopkins, Jerry Li, and Fred Zhang. Robust and heavy-tailed mean estimation made simple, via regret minimization. In *NeurIPS*, 2020. URL https://proceedings.neurips.cc/paper/ 2020/hash/8a1276c25f5efe85f0fc4020fbf5b4f8-Abstract.html.

Daniel J. Hsu and Sham M. Kakade. Learning mixtures of spherical gaussians: moment methods and spectral decompositions. In Robert D. Kleinberg, editor, *Innovations in Theoretical Computer Science, ITCS '13, Berkeley, CA, USA, January 9-12, 2013*, pages 11–20. ACM, 2013. doi: 10.1145/2422436.2422439. URL https://doi.org/10.1145/2422436.2422439.

Chi Jin, Yuchen Zhang, Sivaraman Balakrishnan, Martin J. Wainwright, and Michael I. Jordan. Local maxima in the likelihood of Gaussian mixture models: Structural results and algorithmic consequences. In *Advances in neural information processing systems*, pages 4116–4124, 2016.

Iain M. Johnstone. Function estimation and Gaussian sequence models. *Unpublished manuscript*, 2 (5.3):2, 2002.

Adam Tauman Kalai, Ankur Moitra, and Gregory Valiant. Efficiently learning mixtures of two gaussians. In *Proceedings of the forty-second ACM symposium on Theory of computing*, pages 553–562, 2010.

Vardis Kandiros, Yuval Dagan, Nishanth Dikkala, Surbhi Goel, and Constantinos Daskalakis. Statistical estimation from dependent data. In Marina Meila and Tong Zhang, editors, *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pages 5269–5278. PMLR, 18–24 Jul 2021. URL https://proceedings. mlr.press/v139/kandiros21a.html.

Jason M. Klusowski and W.D. Brinda. Statistical guarantees for estimating the centers of a two-component Gaussian mixture by EM. *arXiv preprint arXiv:1608.02280*, 2016.

Bruce G. Lindsay. Mixture models: Theory, geometry, and applications. Ims, 1995.

David J. C. MacKay. *Information theory, inference and learning algorithms*. Cambridge University Press, New York, 2003. ISBN 0-521-64298-1.

Geoffrey J McLachlan, Sharon X Lee, and Suren I Rathnayake. Finite mixture models. *Annual review of statistics and its application*, 6:355–378, 2019.

Ankur Moitra and Gregory Valiant. Settling the polynomial learnability of mixtures of gaussians. In *2010 IEEE 51st Annual Symposium on Foundations of Computer Science*, pages 93–102. IEEE, 2010.

Tongzheng Ren, Fuheng Cui, Sujay Sanghavi, and Nhat Ho. Beyond EM algorithm on over-specified two-component location-scale gaussian mixtures. *CoRR*, abs/2205.11078, 2022. doi: 10.48550/arXiv.2205.11078. URL `https://doi.org/10.48550/arXiv.2205.11078`.

Phillippe Rigollet and Jan-Christian Hütter. High dimensional statistics. *Lecture notes for course 18S997*, 2019.

Bruce Sacerdote. Peer effects with random assignment: Results for dartmouth roommates. *The Quarterly journal of economics*, 116(2):681–704, 2001.

Alexandre B. Tsybakov. *Introduction to Nonparametric Estimation*. Springer Publishing Company, Incorporated, 2008. ISBN 0387790519.

Ramon van Handel. Hidden Markov models. *Unpublished lecture notes*, 2008.

Roman Vershynin. *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge University Press, 2018.

Martin J. Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge University Press, 2019.

Martin J. Wainwright and Michael I. Jordan. Graphical models, exponential families, and variational inference. *Foundations and Trends® in Machine Learning*, 1(1–2):1–305, 2008. ISSN 1935-8237. doi: 10.1561/2200000001. URL `http://dx.doi.org/10.1561/2200000001`.

Nir Weinberger and Guy Bresler. The em algorithm is adaptively-optimal for unbalanced symmetric gaussian mixtures. *Journal of Machine Learning Research*, 23(103):1–79, 2022. URL `http://jmlr.org/papers/v23/21-0186.html`.

Yihong Wu. Lecture notes on information-theoretic methods for high-dimensional statistics. *Lecture Notes for ECE598YW (UIUC)*, 16, 2017.

Yihong Wu and Pengkun Yang. Optimal estimation of Gaussian mixtures via denoised method of moments. *The Annals of Statistics*, 48(4):1981–2007, 2020.

Yihong Wu and Harrison H. Zhou. Randomly initialized EM algorithm for two-component Gaussian mixture achieves near optimality in $o(\sqrt{n})$ iterations. *arXiv preprint arXiv:1908.10935*, 2019.

Ji Xu, Daniel J. Hsu, and Arian Maleki. Global analysis of expectation maximization for mixtures of two Gaussians. In *Advances in Neural Information Processing Systems*, pages 2676–2684, 2016.

Bowei Yan, Mingzhang Yin, and Purnamrita Sarkar. Convergence of gradient EM on multi-component mixture of Gaussians. In *Advances in Neural Information Processing Systems*, pages 6956–6966, 2017.

Fanny Yang, Sivaraman Balakrishnan, and Martin J. Wainwright. Statistical and computational guarantees for the Baum-Welch algorithm. In *2015 53rd Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pages 658–665. IEEE, 2015.

Yuhong Yang and Andrew Barron. Information-theoretic determination of minimax rates of convergence. *Annals of Statistics*, pages 1564–1599, 1999.

Ruofei Zhao, Yuanzhi Li, and Yuekai Sun. Statistical convergence of the EM algorithm on Gaussian mixture models. *arXiv preprint arXiv:1810.04090*, 2018.