Teamwork makes von Neumann work: Min-Max Optimization in Two-Team Zero-Sum Games

Abstract

Motivated by recent advances in both theoretical and applied aspects of multi-player games, spanning from e-sports to multi-agent generative adversarial networks, we focus on min-max optimization in team zero-sum games. In this class of games, players are split in two teams with payoffs equal within the same team and of opposite sign across the opponent team. Unlike the textbook two player zero-sum games, finding a Nash equilibrium in our class can be shown to be CLS-hard, i.e., it is unlikely to have a polynomial time algorithm for computing Nash equilibria. Moreover In this generalized framework, we establish that even asymptotic last iterate or time average convergence to a Nash Equilibrium is not possible using Gradient Descent Ascent (GDA), its optimistic variant and extra gradient. Specifically, we present a family of team games whose induced utility is non-multilinear with non-attractive *per-se* mixed Nash Equilibria, as strict saddle points of the underlying optimization landscape. Leveraging techniques from control theory, we complement these negative results by designing a modified GDA that converges locally to Nash equilibria. Finally, we discuss connections of our framework with AI architectures with team competition structure like multi-agent generative adversarial networks.

1 Introduction

Team competition has played a central role in the development of Game Theory (Marschak, 1955; von Stengel & Koller, 1997; Bacharach, 1999; Gold, 2005), Economics (Marschak, 1955; Gottinger, 1974) and Evolutionary Biology (Nagylaki, 1993; Nowak et al., 2004), however, the behavior of the underlying dynamics within the teams are usually sidelined. Either for reasons of mathematical convenience or bigger picture understanding, "teams" in literature are typically modeled as if they were unitary actors, i.e., single individuals without unveiling the internal decision-making of the team members (see Kim et al. (2019)).

For instance, in the biology setting of *weak selection model* (Nagylaki, 1993; Chastain et al., 2014; Mehta et al., 2015) species are modeled to compete as teams, while at the crux of the matter the genes of each species are the actual players and their alleles are the actions in the survival game. Similary, it was the social-media collaboration of the Reddit retail trading crowd as a team that touches off the last year's GameStop frenzy of short squeeze (Umar et al., 2021; Hasso et al., 2021) transforming the markets into a tug of war game against the team of Wall Street hedge funds.

Recently, these intrinsic details behind the competition among teams has attracted renewed interest in the Machine Learning community, motivated by the advent of multi-agent systems that are used for generative tasks or playing to complex games like CTF (Jaderberg et al., 2019) or Starcraft (Vinyals et al., 2019). So as to win this kind of games, self-training AI systems have to develop both collaborative attributes (coordination within each team) as well as contesting ones (competition across the teams). Moreover, following the complementary thread of multi-agent generative adversarial network research, the creation of a pool by efficient incumbent agents, either in generators (Arora et al., 2017; Hoang et al., 2017; 2018; Zhang et al., 2018; Tang, 2020), or discriminators (Hardy et al., 2019; Albuquerque et al., 2019) has been tested providing significant statistical and

computational benefits. In this direction, researchers strive to harness the efficacy of distributed processing, utilizing shallower networks that can learn all the while more diverse datasets [1].

In order to shed some light to this persistent strain of research, the main premise of the theoretical scaffolding developed in this paper is that

> *The "unitary two-players" min-max approach misses the critical component*
> *of the collective strategy making within each competing team.*

Our class of games. In this regard, we turn our attention to *Two-Team Zero-Sum games*, proposed by Schulman & Vazirani (2019b), a quite general class of min-max optimization problems that includes bilinear games as well as a wide range of non-convex non-concave games. In this class, the players fall in two teams of size k_1, k_2 and submit their own probabilistic strategy vector independently, akin to a general normal form multi-player game. Following the econometric common value assumption of Marschak (1955), what makes a group of players a *team* is that in any outcome the players of each team receive identical payoff. Thus, to build some intuition, it is easy to see that if perfect coordination existed within each team, the interaction between the teams is merely a zero-sum game between two "virtual" players. To streamline our presentation here, we defer the more precise description of our model to Section 2.

Challenges behind Two-Team Zero-Sum games. In the archetypical case of two players, i.e., $(k_1 = k_2 = 1)$, min-max stategies are typically thought of as the axiomatically correct predictions thanks to the seminal Von Neumann's minmax theorem (Von Neumann, 1928). Unfortunately min-max optimization for case $k > 1$ is a much more tenuous affair: Schulman & Vazirani (2019b) preclude the existence of unique value by presenting a family of team games where $\min\max \neq \max\min$ together with bounds about this duality gap, which quantifies exactly the effect of exchanging the order of strategy commitment either between the teams or the players thereof.

If defining the correct figure of merit for Team games is rife with frustration, what is even more demanding is understanding what kind of algorithms/dynamics are able to solve this problem when a game-theoretically meaningful solution exists: Firstly, computing local Nash Equilibria (NE) in general non-convex non-concave games are PPAD-complete (Daskalakis et al., 2009; 2021). Thus, all well-celebrated first-order methods, like gradient descent-ascent (Lin et al., 2020; Daskalakis & Panageas, 2019), its optimistic (Popov, 1980; Daskalakis & Panageas, 2018; Mertikopoulos et al., 2019) and extragradient variant (Korpelevich, 1976) would require an exponential number of steps in the parameters of the problem to find an appoximate NE under Nemirovsky-Yudin (Nemirovskij & Yudin, 1983) oracle optimization model. Secondly, even if a regret notion could be defined, no-regret methodology is guarranteed to attract only to the set of coarse correlated equilibria (CCE) (Fudenberg, 1991; Hannan, 2016; Flokas et al., 2020; Giannou et al., 2021), a weaker notion that may be exclusively supported on strictly dominated strategies, even for simple symmetric two-player games (See also Viossat & Zapechelnyuk (2013)).

Whilst the aforementioned intractability failures for the general case of non-convex non-concave min-max problems provides significant insights, they can not *a fortiori* answer the fundamental question, restricted in the model of Two-Team Zero-Sum Games:

> *Can we compute Nash equilibria in Two-Team Zero-Sum Games and*
> *ultimately are there first order methods which converge to them under tangible guarantees?*

Our results. To the best of our knowledge, the following contributions are the first-of-its kind type of results for the case of Two-Team Zero-Sum games:

- For the case of computational complexity of approximate (possibly mixed) NE we establish a sweeping negative result proving that it is CLS-hard (Theorem 3.1), i.e., is computationally harder of finding pure NE in a congestion game or finding approximate gradient descent fixed points.
- From optimization perspective, we settle these questions with a resounding "no" for all the well-known discrete gradient flow variations. Specifically, we present a simple family of two-team with two-players zero-sum games where Projected-GDA, Optimistic-GDA and Extra Gradient fail even

---

[1] Indeed, from the training perspective, it's more computationally preferable to back-propagate through two equally sized neural networks with smaller capacity rather than through a giant single one that would be twice as deep.(Tang, 2020)

to stabilize around a mixed NE, when they are initialized nearby (Theorem 3.5). Additionally, for the category GDA in the non-degenerate team games with unique mixed NE, one could acquire an even stronger result for any high-dimensional configuration of actions and players. (Theorem 3.2)

- In order to make some substantial headway under the burden of the above instability results, we shift our attention on adaptive control generalizations of the celebrated Washout filters–traditionally used for stabilizing the Dutch-roll motion of an aircraft during a flight (Hassouneh et al., 2004; Grant & Reid, 1997). Inspired by this framework, we propose the modified KPV-GDA[2] which consists a tandem combination of GDA together with a stabilizing feedback introduces by Bazanella et al. (1997).

$$\begin{cases} state^{(k+1)} = & state^{(k)} + \eta GDA(state^{(k+1)}) + \eta K(state^{(k)} - stress^{(k)}) \\ stress^{(k+1)} = & stresss^{(k)} + \eta P(state^{(k)} - stress^{(k)}) \end{cases} \quad \text{(KPV-GDA)}$$

The main linchpin of KPV-GDA method is the Simon's and Theil's (Simon, 1956; Theil, 1957) *certainty equivalence principle*, a widely used methodology in Control theory in developing applied dynamic rational expectations models. According to this principle, the feedback law is split into two optimization steps whereby $K-step$ attracts quickly the $state$ to the $stress$ while $P-step$ converges slowly to the fixed points of GDA. Comparing with the plethora of the proposed dynamics for min-max problems, the crucial advantage of the afore-described technique is that does not introduce any extra fixed points than GDA's ones. In Section 2.2, we provide some illustrative examples of KPV-GDA technique, while in Theorem 3.7 we prove the existence of such control feedback for our class of games.

- Finally, in Section 4 we provide a series of experiments in simple two-team zero-sum games showcasing both the messy behaviours of traditional methods like GDA,OGDA and the power of KPV-GDA method in these optimization environments. Additionally, we show that multi-agent GAN architectures achieve better performance than the sigle-agent ones, in terms of network capacity, when their trained in synthetic or real-world datasets like CIFAR10.

2 PRELIMINARIES

2.1 DEFINITIONS

Our setting. Formally, a *two-team game* in normal form is defined as a tuple $\Gamma = \Gamma(\mathcal{N}, \mathcal{A}, u)$ consisting of (i) a finite set of *players* $\mathcal{N}$, split into two teams A, B with k_A and k_B players correspondingly such that: $\mathcal{N} = \mathcal{N}_A \cup \mathcal{N}_B = \{A_1, \cdots, A_{k_A}, B_1, \cdots, B_{k_B}\}$; (ii) a finite set of *actions* (or *pure strategies*) $\mathcal{A}_i = \{\alpha_1, \ldots, \alpha_{n_i}\}$ per player $i \in \mathcal{N}$; (iii) each team's *payoff* function $u_A, u_B : \mathcal{A} \to \mathbb{R}$, where $\mathcal{A} := \prod_i \mathcal{A}_i$ denotes the ensemble of all possible action profiles $\alpha = (\alpha_{A_1}, \ldots, \alpha_{A_{k_A}}, \alpha_{B_1}, \ldots, \alpha_{B_{k_B}})$ while the *individual utility* of a player is identical to her teammates, i.e., $u_i = u_A$ & $u_j = u_B$ $\forall (i,j) \in \mathcal{N}_A \times \mathcal{N}_B$. In this general context, players could also adhere *mixed strategies*, i.e, probability distributions $s_k \in \Delta(\mathcal{A}_k)$ over the pure strategies $\alpha_k \in \mathcal{A}_k$. Correspondingly, we define the product distributions $\mathbf{x} = s_{A_1} \otimes \cdots \otimes s_{A_{k_A}}, \mathbf{y} = s_{B_1} \otimes \cdots \otimes s_{B_{k_B}}$ as the teams' strategies. Collectively, we will write $\mathcal{X} := \prod_{i \in \mathcal{N}_A} \mathcal{X}_i = \prod_{i \in \mathcal{N}_A} \Delta(\mathcal{A}_i), \mathcal{Y} := \prod_{i \in \mathcal{N}_A} \mathcal{Y}_i = \prod_{i \in \mathcal{N}_B} \Delta(\mathcal{A}_i)$ the space of mixed strategy profiles of teams A, B.

Similarly with the bilinear two-player games, the teams' utility functions can be expressed via the payoff-*tensors* $\mathbf{A}, \mathbf{B} \in \mathbb{R}^\tau$ with $\tau = \prod_{i \in \mathcal{N}} |\mathcal{A}_i|$ and acquire the form[3]:

$$u_A = \mathbf{A}_{\mathbf{x}}^{\mathbf{y}} \ \& \ u_B = \mathbf{B}_{\mathbf{x}}^{\mathbf{y}} \tag{2.1}$$

---

[2]The name "KPV-GDA" is an initialism from (K, P)-Vaned Gradient Descent Ascent method; Just like the tail section of an aircraft where the vanes are flight control surfaces that control the unstable yaw, (K, P) control feedback aims to stabilize the unstable arrows of Gradient flow around a mixed NE.

[3]Figuratively, the latter form denotes what is known as a tensor contraction given: $\mathbf{A}_{\mathbf{x}}^{\mathbf{y}} = \sum_{i,\ldots,j,k,\ldots,l} x_{i,\ldots,j} A_{i,\ldots,j,k\ldots,l} y_{k\ldots,l}$ If $\mathbf{x}, \mathbf{y}$ have shapes (i, j) and (k, l) that would be equivalent to $\mathtt{u} = \mathtt{einsum('ijkl,ij,kl', \ A, \ x, \ y)}$

In terms of solutions, we focus on the per player *Nash Equilibrium* (NE), i.e., a state strategy profile $s^* = (\mathbf{x}, \mathbf{y}) = \left((s^*_{A_1}, \ldots, s^*_{A_{k_A}}), (s^*_{B_1}, \ldots, s^*_{B_{k_B}}) \right)$ such that

$$u_i(s^*) \geq u_i(s_i; s^*_{-i})^4 \text{ for all } s_i \in \Delta(\mathcal{A}_i) \text{ and all } i \in \mathcal{N} \tag{NE}$$

The state strategy profile s^* is called *pure* if every player of both teams choose a single action; otherwise we say that it is mixed. Finally, a two-team game is called *two-team zero-sum* if $u_A = -u_B$ or equivalently $\mathbf{A} + \mathbf{B} = \mathbf{O}$.

Remark 2.1. A quite technical prerequisite for the rest of this work, we will assume that a succinct representation of the utility tensors of the game are available or equivalently that a payoff oracle provides efficiently both the value of the utility function and its derivatives for a specific input, which is consistent with the vast majority of the applications that are described in the literature (von Stengel & Koller, 1997).

A first approach on computing Nash equilibria in Two-Team Zero-Sum games. Given the existence of the duality-gap between the $\min \max$ and $\max \min$, in lieu of the two-player zero-sum game an equilibrium in our setting can not be computed via linear programming. For the goal of computing Nash equilibria in two-team zero-sum games, we have experimented with a selection of first order methods that have been utilized with varying success in the setting of two-person zero-sum case. Namely, for a given scalar, the real-valued utility function $f(\mathbf{x}, \mathbf{y}) = \mathbf{A}^{\mathbf{y}}_{\mathbf{x}}$, we get:

1. Gradient Descent-Ascent

$$
\begin{cases}
x_i^{(k+1)} = \Pi_{\mathcal{X}_i} \left\{ x_i^{(k)} - \eta \nabla_{x_i} f(\mathbf{x}^{(k)}, \mathbf{y}^{(k)}) \right\} \\
y_j^{(k+1)} = \Pi_{\mathcal{Y}_j} \left\{ y_j^{(k)} + \eta \nabla_{y_j} f(\mathbf{x}^{(k)}, \mathbf{y}^{(k)}) \right\}
\end{cases}
$$

2. Optimistic Gradient Descent-Ascent

$$
\begin{cases}
x_i^{(k+1)} = \Pi_{\mathcal{X}_i} \left\{ x_i^{(k)} - 2\eta \nabla_{x_i} f(\mathbf{x}^{(k)}, \mathbf{y}^{(k)}) + \eta \nabla_{x_i} f(\mathbf{x}^{(k-1)}, \mathbf{y}^{(k-1)}) \right\} \\
y_j^{(k+1)} = \Pi_{\mathcal{Y}_j} \left\{ y_j^{(k)} + 2\eta \nabla_{y_j} f(\mathbf{x}^{(k)}, \mathbf{y}^{(k)}) - \eta \nabla_{y_j} f(\mathbf{x}^{(k-1)}, \mathbf{y}^{(k-1)}) \right\}
\end{cases}
$$

3. Extra Gradient Method

$$
\begin{cases}
x_i^{(k+\frac{1}{2})} = \Pi_{\mathcal{X}_i} \left\{ x_i^{(k)} - \eta \nabla_{x_i} f(\mathbf{x}^{(k)}, \mathbf{y}^{(k)}) \right\}, & x_i^{(k+1)} = \Pi_{\mathcal{X}_i} \left\{ x_i^{(k)} - \eta \nabla_{x_i} f(\mathbf{x}^{(k+\frac{1}{2})}, \mathbf{y}^{(k+\frac{1}{2})}) \right\} \\
y_j^{(k+\frac{1}{2})} = \Pi_{\mathcal{Y}_j} \left\{ y_j^{(k)} + \eta \nabla_{y_j} f(\mathbf{x}^{(k)}, \mathbf{y}^{(k)}) \right\}, & y_j^{(k+1)} = \Pi_{\mathcal{Y}_j} \left\{ y_j^{(k)} + \eta \nabla_{y_j} f(\mathbf{x}^{(k+\frac{1}{2})}, \mathbf{y}^{(k+\frac{1}{2})}) \right\}
\end{cases}
$$

where $x_i^{(k)}$ (or $y_j^{(k)}$) denotes the strategy vector of the i-th minimizing (or j-th maximizing agent) at time-step k, the step-size is denoted by η and $\Pi_{\mathcal{X}_i}, \Pi_{\mathcal{Y}_j}$ are the projection operators to the corresponding simplices. The below remark will play a key role in the sequel.

Remark 2.2. Any fixed point of the aforementioned discrete time dynamics on the utility function corresponds necessarily to the Nash equilibria of the game.

Hence, an important testbed for the long-run behavior of GDA, OGDA and EG methods is to examine whether these methods stabilize around their fixed points, which consist the effectively Nash equilibria of the game. In Section 3.2, we show that in lack of pure Nash equilibria, all the above methods fail to stabilize on their fixed points even for a simple class of $(2, 2)$-players game, and as a consequence to the mixed Nash equilibria of the game.

The presence of these results showcases the need of a different approach that lies outside purely optimization based ideas. Inspired by the applications of washout filters to stabilize highly susceptible systems and their adaptive control generalizations, we design a new incarnation of GDA vaned by two matrices-control feedback. Surprisingly, in contrast with the aforementioned traditional methods, our proposed technique accomplishes last-iterate stabilization on its fixed point, i.e., the mixed Nash equilibria of the team game.

---

[4]We are using here the standard shorthand $(s_1, \ldots, s_i, \ldots, s_{|\mathcal{N}|})$ to highlight the strategy of a given player $i \in \mathcal{N}$ versus the rest of players $\mathcal{N} \setminus \{i\}$.

(K, P)-**Vaned GDA Method.** After concatenating the vectors of the minimizing and the maximizing agents $\mathbf{z}^{(k)} = (\mathbf{x}^{(k)}, \mathbf{y}^{(k)})$ we can write our method, for appropriate matrices K, P:

$$\begin{cases} \mathbf{z}^{(k+1)} = & \Pi_{\mathcal{Z}} \Big\{ \mathbf{z}^{(k)} + \eta \big({-\nabla_{\mathbf{x}} f(\mathbf{z}^{(k)}) \atop \nabla_{\mathbf{y}} f(\mathbf{z}^{(k)})} \big) + \eta K(\mathbf{z}^{(k)} - \boldsymbol{\theta}^{(k)}) \Big\} \\ \boldsymbol{\theta}^{(k+1)} = & \Pi_{\mathcal{Z}} \Big\{ \boldsymbol{\theta}^{(k)} + \eta P(\mathbf{z}^{(k)} - \boldsymbol{\theta}^{(k)}) \Big\} \end{cases} \tag{2.2}$$

Intuitively, the added variable $\boldsymbol{\theta}^{(k)}$ holds an estimate of the fixed point, and through the feedback $\eta K(\mathbf{z}^{(k)} - \boldsymbol{\theta}^{(k)})$ the vector $\mathbf{z}$ stabilizes around that estimate which slowly moves towards the real fixed point of the plain GDA dynamic. It is crucial to note that no additional fixed points are introduced to the system.

2.2 Two Illustrative Examples

Our first example supports a double role: Firstly, it exemplifies how our two-team min-max competition can captures the formulation of multi-agent GANs' architectures. Secondly, it hints also an early separation between the optimization methods, since as we will see GDA will not converge to the Nash Equilibrium/ground-truth distribution.

2.2.1 Learning a mixture of Gaussians with multi-agent GANs

Consider the case of $\mathcal{M}$, a mixture of gaussian distribution with two components, $C_1 \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{I})$ and $C_2 \sim \mathcal{N}(-\boldsymbol{\mu}, \boldsymbol{I})$ and mixture weights π_1, π_2 to be positive such that $\pi_1 + \pi_2 = 1$ and $\pi_1, \pi_2 \neq \frac{1}{2}$.

To learn the distribution above, we utilize an instance of a *Team*-WGAN in which there exists a generating team of agents $G_p : \mathbb{R} \to \mathbb{R}, G_{\boldsymbol{\theta}} : \mathbb{R}^n \to \mathbb{R}^n$, and a discriminating team of agents $D_{\mathbf{v}} : \mathbb{R}^n \to \mathbb{R}, D_{\mathbf{w}} : \mathbb{R}^n \to \mathbb{R}$, all described by the following equations:

$$\begin{array}{ll} \text{Generators:} & G_p(\zeta) = p + \zeta \ , \ G_\theta(\mathbf{z}) = \mathbf{z} + \boldsymbol{\theta} \\ \text{Discriminators:} & D_{\mathbf{v}}(\mathbf{y}) = \langle \mathbf{v}, \mathbf{y} \rangle \ , \ D_{\mathbf{w}}(\mathbf{y}) = \sum_i w_i y_i^2 \end{array} \tag{2.3}$$

The generating agent G_θ maps random noise $\mathbf{z} \sim \mathcal{N}(0, \boldsymbol{I})$ to samples while generating agent $G_p(\zeta)$, utilizing an independent source of randomness $\zeta \sim \mathcal{N}(0, 1)$, probabilistically controls the sign of the output of the generator G_θ. The probability of ultimately generating a sample $\mathbf{y} = \mathbf{z} + \boldsymbol{\theta}$ is equal to $\zeta + p$, while the probability of the sample being $\mathbf{y} = -\mathbf{z} - \boldsymbol{\theta}$ is equal to $1 - (p + \zeta)$.

On the other end, there stands the discriminating team of $D_{\mathbf{v}}, D_{\mathbf{w}}$. Discriminators, $D_v(\mathbf{y}), D_w(\mathbf{y})$ map any given sample $\mathbf{y}$ to a scalar value accounting for the realness or fakeness of it – negative meaning fake, positive meaning real. The discriminators are disparate in the way they measure realness of samples as seen in their definitions.

We follow the formalism of the Wasserstein GAN to form the optimization objective:

$$\max_{\mathbf{v}, \mathbf{w}} \min_{\boldsymbol{\theta}, p} \left\{ \begin{array}{c} \mathbb{E}_{\mathbf{y} \sim real} \Big[D_{\mathbf{v}}(\mathbf{y}) + D_{\mathbf{w}}(\mathbf{y}) \Big] \\ - \\ \mathbb{E}_{z \sim \mathcal{N}(0, \boldsymbol{I}), \zeta \sim \mathcal{N}(0,1)} \left[\left\{ \begin{array}{c} G_p(\zeta) \cdot \Big(D_{\mathbf{v}}(G_{\boldsymbol{\theta}}(\mathbf{y})) + D_{\mathbf{w}}(G_{\boldsymbol{\theta}}(\mathbf{y})) \Big) \\ - \\ (1 - G_p(\zeta)) \cdot \Big(D_{\mathbf{v}}(G_{\boldsymbol{\theta}}(\mathbf{y})) + D_{\mathbf{w}}(G_{\boldsymbol{\theta}}(\mathbf{y})) \Big) \end{array} \right\} \right] \end{array} \right\} \tag{2.4}$$

Equation equation 2.4 yields the simpler form:

$$\max_{\mathbf{v}, \mathbf{w}} \min_{\boldsymbol{\theta}, p} (\pi_1 - \pi_2) \mathbf{v}^T \boldsymbol{\mu} - 2p \mathbf{v}^T \boldsymbol{\theta} + \mathbf{v}^T \boldsymbol{\theta} + \sum_i^n w_i(\mu_i^2 - \theta_i^2) \tag{2.5}$$

It is easy to check that Nash equilibria of Equation equation 2.4 must satisfy:

$$\left\{ \begin{array}{llll} \boldsymbol{\theta} & = & \boldsymbol{\mu}, & p = 1 - \pi_2 = \pi_1 \\ \boldsymbol{\theta} & = & -\boldsymbol{\mu}, & p = 1 - \pi_1 = \pi_2. \end{array} \right\}$$

Figure 1 demonstrates both GDA's failure and OGDA, EG and our KPV-GDA methods' success to converge to the above Nash equilibria and tantamountly to discover the groudtruth mixture.

Figure 1: Parameter training of the configuration under different algorithms

2.2.2 MULTIPLAYER MATCHING PENNIES

Interestingly enough, there are non-trivial instances of two-team competition settings that even Optimistic GDA and EG Method fail to converge. Such is the case for a team version of the well-known game of matching pennies. The game can be shortly described as such: "*coordinate with your team-mates to play a game of matching pennies against the opposing team, coordinate not and pay a penalty*". The penalty is set to $\frac{1}{2}$. For the interested reader, we defer to appendix A.3 the precise description of the game in a contracted tensor/table. Since every player has simply two actions, their probability vector can be represented by a single variable in $[0, 1]$. Considering the minimizing team $\mathbf{x}$ its players are x_1, x_2, while the players of the maximizing team $\mathbf{y}$ are y_1, y_2. The multiplayer of matching pennies is described by the utility function:

$$u(x_1, x_2, y_1, y_2) = -x_1x_2 - x_1y_1 - x_1y_2 + 1.5(x_1 + x_2) - x_2y_1 - x_2y_2 + y_1y_2 + 0.5(y_1 + y_2) - 1 \quad (2.6)$$

As we can see in Figures 2 and 3, multiplayer matching pennies game consists an excellent benchmark where all traditional gradient flow discretizations fail under perfect competition setting. Interestingly, we are not aware of similar example in min-max literature and it has been our starting point for seeking new optimization techniques inspired by Control theory. Indeed, KPV-GDA variation with $(K, P) = (-1.1\mathbf{I}, 0.3\mathbf{I})$ achieves to converge to the unique mixed Nash Equilibrium of the game. In the following sections, we provide theorems that explained formally this long-run behavior of the examinded dynamics.

Figure 2: Multiplayer matching pennies under different algorithms

Figure 3: Projected Trajectory of Team A under different algorithms

3 OUR MAIN RESULTS

3.1 ON THE COMPLEXITY OF TWO-TEAM ZERO-SUM GAMES

We start this section by showing that computing a Nash equilibrium in two team zero-sum games is computationally hard and thus getting a polynomial time algorithm that computes a Nash equilibrium is unlikely.

Theorem 3.1 (CLS-hard). *Computing a Nash equilibrium in Two team zero-sum games is* CLS-*hard.*

The main idea of the proof of Theorem 3.1 relies on a reduction of approximating Nash equilibria in congestion games, which has been shown to be complete for the interesting class of CLS, which contains the problem of continuous optimization. For consicion, we defer the proof of the above theorem to the paper's supplement.

3.2 FIRST-ORDER METHODS FAIL TO STABILIZE

The negative computational complexity result we proved for two team zero-sum games (Theorem 3.1) does not conclude the prospect of having algorithms (learning dynamics, first-order methods) that converge to Nash equilibria and thus can approximate them well enough. Unfortunately, we can even prove negative results about convergence to Nash equilibria in two team zero-sum games of well established methods that are commonly used and have found enormous success in classic two player zero-sum games.

In this section, we are going to construct a family of two team zero-sum games with the property that GDA, OGDA and EG fail to stabilize to Nash equilibria. This result indicates how challenging and rich the setting of team zero-sum games can be and why provable guarantees about convergence have not been established yet. Before defining the family of two team zero-sum games, we prove an important Theorem which states that GDA does not stabilize around mixed Nash equilibria. This fact is a stepping stone in constructing the family of team-zero sum games later. We present the proof all of the below statements in detail in the paper's appendix.

Weakly-stable Nash equilibrium (Kleinberg et al., 2009; Mehta et al., 2015). Consider the set of Nash equilibria with the property that if any single randomizing agent of one team is forced to play any strategy in her current support with probability one, all other agents of the same team must remain indifferent between the strategies in their support. This type of Nash equilibria is called weakly-stable. Please note that trivially pure Nash equilibria are weakly-stable. It has been shown that mixed Nash equilibria are not weakly-stable in generic games[5] games (Kleinberg et al., 2009). We can show that Nash equilibria that are not weakly-stable Nash are actually unstable for GDA and moreover, using standard machinery from dynamical systems, that the set of initial conditions that converges to Nash equilibria that are not weakly-stable should be of measure zero. Formally we prove that:

Theorem 3.2 (Non weakly-stable Nash are unstable). *Consider a two team zero-sum game with utility function of Team B (**y** vector) being $U(\mathbf{x}, \mathbf{y})$ and Team A (**x** vector) being $-U(\mathbf{x}, \mathbf{y})$. Moreover, assume that $(\mathbf{x}^*, \mathbf{y}^*)$ is a Nash equilibrium of full support that is not weakly-stable. It follows that the set of initial conditions so that GDA converges to $(\mathbf{x}^*, \mathbf{y}^*)$ is of measure zero for stepsize $\eta < \frac{1}{L}$ where L is the Lipschitz constant of ∇U.*

3.3 GENERALIZED MATCHING PENNIES (GMP)

Inspired by Theorem 3.2, in this section we construct a family of team zero-sum games so that GDA, OGDA and EG methods fail to converge (if the initialization is a random point in the simplex, the probability of convergence of the aforementioned methods is zero). The intuition is to construct a family of games, each of which has only mixed Nash equilibria (that are not weakly-stable), i.e., the constructed games should lack pure Nash equilibria; using Theorem 3.2, it would immediately imply our claim for GDA. It turns out that OGDA and EG also fail to converge for the same family.

Definition of GMP. Consider a setting of two teams (Team A, Team B), each of which has $n = 2$ players. Inspired by the standard matching pennies game and the game defined in Schulman & Vazirani (2019a), we allow each agent i to have two strategies/actions that is $S = \{H, T\}$ for both teams with 2^4 possible strategy profiles. In case all the members of a Team choose the same strategy say H or T then the Team "agrees" to play H or T (otherwise the Team "does not agree").

Thus, in the case both teams "agree", the payoff of each team is actually the payoffs for the two player matching pennies. If one team "agrees" and the other does not, the team that "agrees" gets

---

[5]Roughly speaking, games in which we add small Gaussian noise on every payoff.

Figure 4: Parameter training of the configuration under different algorithms

payoff $\omega \in (0, 1)$ and the other team gets penalty ω. If both teams fail to "agree", both team get payoff zero. Let x_i with $i \in \{1, 2\}$ be the probability that agent i of Team A chooses H and $1 - x_i$ the probability that she chooses T. We also denote $\mathbf{x}$ the vector of probabilities for Team A. Similarly, we denote y_i for $i \in \{1, 2\}$ be the probability that agent i of Team B chooses H and $1 - y_i$ the probability that she chooses T and $\mathbf{y}$ the probability vector.

The first fact about the game that we defined is that for $\omega \in (0, 1)$, there is only one Nash equilibrium $(\mathbf{x}^*, \mathbf{y}^*)$, which is the uniform, i.e., $x_1^* = x_2^* = y_1^* = y_2^* = \frac{1}{2}$ for all agents i.

Lemma 3.3 (GMP has a unique Nash). *The Generalized Matching Pennies game exhibits a unique Nash equilibrium which is* $(\mathbf{x}^*, \mathbf{y}^*) = ((\frac{1}{2}, \frac{1}{2}), (\frac{1}{2}, \frac{1}{2}))$.

Remark 3.4. The fact that the game we defined has a unique Nash equilibrium that is in the interior of $[0, 1]^4$ is really crucial for our negative convergence results later in the section as we will show that it is not weakly-stable Nash equilibrium and the negative result about GDA will be a corollary due to Theorem 3.2. Please also note that if $\omega = 1$ then there are more Nash equilibria, in particular the $(\mathbf{0}, \mathbf{0}), (\mathbf{1}, \mathbf{0}), (\mathbf{0}, \mathbf{1}), (\mathbf{1}, \mathbf{1})$ are also Nash equilibria (which are pure).

The following Theorem is the main (negative) result of this section.

Theorem 3.5 (GDA, OGDA, EG fail). *Consider GMP game with* $\omega \in (0, 1)$. *Assume that* $\eta_{GDA} < \frac{1}{4}$, $\eta_{OGDA} < \frac{1}{8}$ *and* $\eta_{EG} < \frac{\omega}{2}$ *(bound on the stepsize for GDA, OGDA and EG methods respectively). It holds that the set of initial conditions so that GDA, OGDA, EG converge (stabilize to any point) is of measure zero.*

Remark 3.6 (Average iterate also fails). One might ask what happens when we consider average iterates instead of last iterate. It is well-known fact Syrgkanis et al. (2015) that the average iterate of no-regret algorithms converges to course-correlated equilibria (CCE) so we expect that the average iterate stabilizes. Nevertheless, CCE might not be Nash equilibria. Indeed we can construct examples in which the average iterate of GDA, OGDA and EG experimentally fails to stabilize to Nash equilibria. In particular we consider a slight modification of GMP; players and strategies are the same but the payoff matrix has changed and can be found below (table on the right):

	HH	HT/TH	TT		HH	HT/TH	TT
HH	$1, -1$	$\omega, -\omega$	$-1, 1$	HH	$2, -2$	$\frac{1}{2}, -\frac{1}{2}$	$-2, 2$
HT/TH	$-\omega, \omega$	$0, 0$	$-\omega, \omega$	HT/TH	$-\frac{1}{2}, \frac{1}{2}$	$0, 0$	$-\frac{1}{2}, \frac{1}{2}$
TT	$-1, 1$	$\omega, -\omega$	$1, -1$	TT	$-1, 1$	$\frac{1}{2}, -\frac{1}{2}$	$1, -1$

Table 1: GMP configurations of Theorem 3.5 (left) and Remark 3.6 (right)

Figure 4 illustrates that the average iterates of GDA, OGDA, and EG stabilizes to points that are not Nash equilibria. Note that since our method (see next subsection) converges locally, the average iterate should converge locally to a Nash equilibrium.

3.4 Wash-out Filters & Adaptive Control

The aforementioned results indicates that to answer the tantalizing question of finding NE in two-team zero-sum games, our machiner should be broaden outside the limits of textbook optimization arsenal. The mainstay on this effort and our positive result is KPV-GDA method defined in (2.2), inspired by the adaptive control toolbox and washout filters. Our main statement shows that KPV-GDA stabilizes around any Nash equilbrium for appropriate choices of matrices K, P. The formal theorem is given below:

Theorem 3.7 (KPV stabilizes). *Consider a team zero-sum game so that the utility of Team B is $U(\mathbf{x}, \mathbf{y})$ and hence the utility of Team A is $-U(\mathbf{x}, \mathbf{y})$ and a Nash equilibrium $(\mathbf{x}^*, \mathbf{y}^*)$ of the game. Moreover we assume*

$$\left(\begin{array}{cc} -\nabla_{\mathbf{xx}} U(\mathbf{x}^*, \mathbf{y}^*) & -\nabla_{\mathbf{xy}} U(\mathbf{x}^*, \mathbf{y}^*) \\ \nabla_{\mathbf{yx}} U(\mathbf{x}^*, \mathbf{y}^*) & \nabla_{\mathbf{yy}} U(\mathbf{x}^*, \mathbf{y}^*) \end{array} \right) \ is\ invertible.$$

For any fixed stepsize $\eta > 0$, we can always find matrices K, P so that GDA-KPV method defined in (2.2) converges locally to $(\mathbf{x}^*, \mathbf{y}^*)$.

4 Experiments

In this section we perform a series of numerical experiments to validate our theoretical findings. Our experiment setting includes a 2-D Gaussian Mixture Model with 8 modes. Our architecture includes 8 "shallow" generators and discriminators with 2 layers of 2-16-2 ReLUs activations, compared with a giant single-agent GAN with 4 layers of 2-128-256-1024-2 activations. Interestingly, the giant one fails in a double sense; It appears both mode-collapsing and mode-drop phenomena without stabilizing. On the other hand our architecture with a small number of neurons achieves to We defer an execution of multi-generators multi-discriminators architectures for CIFAR-10 again to the paper's supplement.

(a) Each generator of MGAN learns one mode of 8-GMM (b) Mode Collapse of single-agent GANs (c) Single-agent GAN can't discriminate between the modes

5 Conclusions and Open Problems

In this paper, we have presented a number of negative results about the problem of finding a Nash equilibrium in team zero-sum games and moreover about the inability of commonly used methods for min-max optimization such as GDA, OGDA and EG to stabilize. We also presented a method (called KPV-GDA) that manages to stabilize around Nash equilibria. Given these results, a number of interesting open questions emerge.

Open Questions. One question for future consideration is the global convergence and the rates of convergence of KPV method. We believe that KPV converges globally for appropriate choice of matrices K, P. One other possible direction is to find a systematic way to get the matrices K, P.

REPRODUCIBILITY STATEMENT

In our submission folder, we provide all the necessary additional technical materials and complete proofs of the main draft's statements in the appendix section. We also uploaded the code of our experiments (Python/Pytorch/Tensorflow).

REFERENCES

Isabela Albuquerque, João Monteiro, Thang Doan, Breandan Considine, Tiago Falk, and Ioannis Mitliagkas. Multi-objective training of generative adversarial networks with multiple discriminators. In *International Conference on Machine Learning*, pp. 202–211. PMLR, 2019.

Martin Arjovsky, Soumith Chintala, and Léon Bottou. Wasserstein generative adversarial networks. In *Proceedings of the 34th International Conference on Machine Learning-Volume 70*, pp. 214–223, 2017.

Sanjeev Arora, Rong Ge, Yingyu Liang, Tengyu Ma, and Yi Zhang. Generalization and equilibrium in generative adversarial nets (gans). In *International Conference on Machine Learning*, pp. 224–232. PMLR, 2017.

Yakov Babichenko and Aviad Rubinstein. Settling the complexity of nash equilibrium in congestion games. In Samir Khuller and Virginia Vassilevska Williams (eds.), *STOC '21: 53rd Annual ACM SIGACT Symposium on Theory of Computing, Virtual Event, Italy, June 21-25, 2021*, pp. 1426–1437. ACM, 2021.

Michael Bacharach. Interactive team reasoning: A contribution to the theory of co-operation. *Research in Economics*, 53(2):117–147, 1999. ISSN 1090-9443. doi: https://doi.org/10.1006/reec.1999.0188. URL https://www.sciencedirect.com/science/article/pii/S1090944399901886.

Nicola Basilico, Andrea Celli, Giuseppe De Nittis, and Nicola Gatti. Computing the team–maxmin equilibrium in single–team single–adversary team games. *Intelligenza Artificiale*, 11(1):67–79, 2017a.

Nicola Basilico, Andrea Celli, Giuseppe De Nittis, and Nicola Gatti. Team-maxmin equilibrium: efficiency bounds and algorithms. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 31, 2017b.

A. S. Bazanella, P. V. Kokotovic, and A. S. e Silva. On the control of dynamic systems with unknown operating point. In *1997 European Control Conference (ECC)*, pp. 3434–3439, 1997. doi: 10.23919/ECC.1997.7082644.

Andrea Celli and Nicola Gatti. Computational results for extensive-form adversarial team games. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 32, 2018.

Erick Chastain, Adi Livnat, Christos Papadimitriou, and Umesh Vazirani. Algorithms, games, and evolution. *Proceedings of the National Academy of Sciences*, 111(29):10620–10623, 2014.

Constantinos Daskalakis and Ioannis Panageas. The limit points of (optimistic) gradient descent in min-max optimization. *Advances in Neural Information Processing Systems*, 31, 2018.

Constantinos Daskalakis and Ioannis Panageas. Last-iterate convergence: Zero-sum games and constrained min-max optimization. *Innovations in Theoretical Computer Science*, 2019.

Constantinos Daskalakis, Paul W Goldberg, and Christos H Papadimitriou. The complexity of computing a nash equilibrium. *SIAM Journal on Computing*, 39(1):195–259, 2009.

Constantinos Daskalakis, Stratis Skoulakis, and Manolis Zampetakis. The complexity of constrained min-max optimization. In *Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing*, pp. 1466–1478, 2021.

Ishan Durugkar, Ian Gemp, and Sridhar Mahadevan. Generative multi-adversarial networks. *arXiv preprint arXiv:1611.01673*, 2016.

Lampros Flokas, Emmanouil Vlatakis-Gkaragkounis, Thanasis Lianeas, Panayotis Mertikopoulos, and Georgios Piliouras. No-regret learning and mixed nash equilibria: They do not mix. In *NeurIPS'20: The 34th International Conference on Neural Information Processing Systems*, 2020.

Drew Fudenberg. Jean tirole game theory, 1991.

Angeliki Giannou, Emmanouil Vasileios Vlatakis-Gkaragkounis, and Panayotis Mertikopoulos. Survival of the strictest: Stable and unstable equilibria under regularized learning with partial information. In Mikhail Belkin and Samory Kpotufe (eds.), *Proceedings of Thirty Fourth Conference on Learning Theory*, volume 134 of *Proceedings of Machine Learning Research*, pp. 2147–2148. PMLR, 15–19 Aug 2021. URL https://proceedings.mlr.press/v134/giannou21a.html.

Natalie Gold. Introduction: Teamwork in theory and in practice. In *Teamwork*, pp. 1–21. Springer, 2005.

Hans W Gottinger. J. marschak and roy radner," economic theory of teams"(book review). *Theory and Decision*, 5(3):349, 1974.

Peter R Grant and Lloyd D Reid. Motion washout filter tuning: Rules and requirements. *Journal of aircraft*, 34(2):145–151, 1997.

James Hannan. 4. approximation to rayes risk in repeated play. In *Contributions to the Theory of Games (AM-39), Volume III*, pp. 97–140. Princeton University Press, 2016.

Corentin Hardy, Erwan Le Merrer, and Bruno Sericola. Md-gan: Multi-discriminator generative adversarial networks for distributed datasets. In *2019 IEEE international parallel and distributed processing symposium (IPDPS)*, pp. 866–877. IEEE, 2019.

Tim Hasso, Daniel Müller, Matthias Pelster, and Sonja Warkulat. Who participated in the gamestop frenzy? evidence from brokerage accounts. *Finance Research Letters*, pp. 102140, 2021.

Munther A Hassouneh, Hsien-Chiarn Lee, and Eyad H Abed. Washout filters in feedback control: Benefits, limitations and extensions. In *Proceedings of the 2004 American control conference*, volume 5, pp. 3950–3955. IEEE, 2004.

Quan Hoang, Tu Dinh Nguyen, Trung Le, and Dinh Phung. Multi-generator generative adversarial nets. *arXiv preprint arXiv:1708.02556*, 2017.

Quan Hoang, Tu Dinh Nguyen, Trung Le, and Dinh Phung. Mgan: Training generative adversarial nets with multiple generators. In *International conference on learning representations*, 2018.

Max Jaderberg, Wojciech M Czarnecki, Iain Dunning, Luke Marris, Guy Lever, Antonio Garcia Castaneda, Charles Beattie, Neil C Rabinowitz, Ari S Morcos, Avraham Ruderman, et al. Human-level performance in 3d multiplayer games with population-based reinforcement learning. *Science*, 364(6443):859–865, 2019.

Jeongbin Kim, Thomas R Palfrey, and Jeffrey R Zeidel. A theory of games played by teams of players. 2019.

R. Kleinberg, G. Piliouras, and É. Tardos. Multiplicative updates outperform generic no-regret learning in congestion games. In *STOC*, 2009.

GM Korpelevich. The extragradient method for finding saddle points and other problems. *Matecon*, 12:747–756, 1976.

Jason D. Lee, Ioannis Panageas, Georgios Piliouras, Max Simchowitz, Michael I. Jordan, and Benjamin Recht. First-order methods almost always avoid strict saddle points. *Math. Program.*, 176(1-2):311–337, 2019. doi: 10.1007/s10107-019-01374-3. URL https://doi.org/10.1007/s10107-019-01374-3.

Dan Li, Dacheng Chen, Baihong Jin, Lei Shi, Jonathan Goh, and See-Kiong Ng. Mad-gan: Multi-variate anomaly detection for time series data with generative adversarial networks. In *International Conference on Artificial Neural Networks*, pp. 703–716. Springer, 2019.

Tianyi Lin, Chi Jin, and Michael Jordan. On gradient descent ascent for nonconvex-concave mini-max problems. In *International Conference on Machine Learning*, pp. 6083–6093. PMLR, 2020.

Jakob Marschak. Elements for a theory of teams. *Management science*, 1(2):127–137, 1955.

H Brendan McMahan, Geoffrey J Gordon, and Avrim Blum. Planning in the presence of cost functions controlled by an adversary. In *Proceedings of the 20th International Conference on Machine Learning (ICML-03)*, pp. 536–543, 2003.

R. Mehta, I. Panageas, and G. Piliouras. Natural selection as an inhibitor of genetic diversity: Multiplicative weights updates algorithm and a conjecture of haploid genetics. In *ITCS*, 2015.

Panayotis Mertikopoulos, Houssam Zenati, Bruno Lecouat, Chuan-Sheng Foo, Vijay Chan-drasekhar, and Georgios Piliouras. Optimistic mirror descent in saddle-point problems: Going the extra (gradient) mile. In *ICLR'19-International Conference on Learning Representations*, pp. 1–23, 2019.

Mohammad Sal Moslehian. Ky fan inequalities. *CoRR*, abs/1108.1467, 2011.

Thomas Nagylaki. The evolution of multilocus systems under weak selection. *Genetics*, 134(2): 627–647, 1993.

Arkadij Semenovič Nemirovskij and David Borisovich Yudin. Problem complexity and method efficiency in optimization. 1983.

Martin A Nowak, Akira Sasaki, Christine Taylor, and Drew Fudenberg. Emergence of cooperation and evolutionary stability in finite populations. *Nature*, 428(6983):646–650, 2004.

Leonid Denisovich Popov. A modification of the arrow-hurwicz method for search of saddle points. *Mathematical notes of the Academy of Sciences of the USSR*, 28(5):845–848, 1980.

R.W. Rosenthal. A class of games possessing pure-strategy Nash equilibria. *International Journal of Game Theory*, 2(1):65–67, 1973.

Leonard J. Schulman and Umesh V. Vazirani. The duality gap for two-team zero-sum games. *Games Econ. Behav.*, 115:336–345, 2019a. doi: 10.1016/j.geb.2019.03.011. URL https://doi.org/10.1016/j.geb.2019.03.011.

Leonard J Schulman and Umesh V Vazirani. The duality gap for two-team zero-sum games. *Games and Economic Behavior*, 115:336–345, 2019b.

Herbert A Simon. Dynamic programming under uncertainty with a quadratic criterion function. *Econometrica, Journal of the Econometric Society*, pp. 74–81, 1956.

E.D. Sontag. Mathematical control theory. In *Texts in Applied Mathematics*, 1998.

Vasilis Syrgkanis, Alekh Agarwal, Haipeng Luo, and Robert E Schapire. Fast convergence of regu-larized learning in games. In *Advances in Neural Information Processing Systems*, pp. 2989–2997, 2015.

Shichang Tang. Lessons learned from the training of gans on artificial datasets. *IEEE Access*, 8: 165044–165055, 2020.

Henri Theil. A note on certainty equivalence in dynamic planning. *Econometrica: Journal of the Econometric Society*, pp. 346–349, 1957.

Zaghum Umar, Mariya Gubareva, Imran Yousaf, and Shoaib Ali. A tale of company fundamentals vs sentiment driven pricing: The case of gamestop. *Journal of Behavioral and Experimental Finance*, 30:100501, 2021.

Oriol Vinyals, Igor Babuschkin, Wojciech M Czarnecki, Michaël Mathieu, Andrew Dudzik, Juny-oung Chung, David H Choi, Richard Powell, Timo Ewalds, Petko Georgiev, et al. Grandmaster level in starcraft ii using multi-agent reinforcement learning. *Nature*, 575(7782):350–354, 2019.

Yannick Viossat and Andriy Zapechelnyuk. No-regret dynamics and fictitious play. *Journal of Economic Theory*, 148(2):825–842, 2013.

John Von Neumann. Zur theorie der gesellschaftsspiele. *Math*, 100:295–320, 1928.

Bernhard von Stengel and Daphne Koller. Team-maxmin equilibria. *Games and Economic Behavior*, 21(1-2):309–321, 1997.

Hongyang Zhang, Susu Xu, Jiantao Jiao, Pengtao Xie, Ruslan Salakhutdinov, and Eric P Xing. Stackelberg gan: Towards provable minimax equilibrium via multi-generator architectures. *arXiv preprint arXiv:1811.08010*, 2018.

Youzhi Zhang and Bo An. Computing team-maxmin equilibria in zero-sum multiplayer extensive-form games. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 34, pp. 2318–2325, 2020a.

Youzhi Zhang and Bo An. Converging to team-maxmin equilibria in zero-sum multiplayer games. In *International Conference on Machine Learning*, pp. 11033–11043. PMLR, 2020b.

A APPENDIX

A.1 FURTHER RELATED WORK

The properties of team games have given rise to a vast corpus of literature which we cannot hope to review here; for an appetizer we refer the reader to Kim et al. (2019); Gold (2005) and reference therein.On the other had, the convergence of first order methods to NE, like GDA, OGDA or Extragradient in *team games* (even finite ones) is nowhere near as well understood.

A notable exception to this is the case of *team-maxmin equilibria* (TME), a notion introduced by von Stengel & Koller (1997). In this category of two-team zero-sum games, a team of $n-$players plays defensively against a single *adversarial* player. TME provide crucial information about the robustness since they give the team players the highest payoff they employ uncoordinated strategies. Unfortunately, TME correspond to the solution of a non–linear non–convex program. (Basilico et al., 2017b) provided heuristics with iterative LP and support enumeration methods with proved approximation guarantees. For the case of extensive or large-sized normal form games, Basilico et al. (2017a); Celli & Gatti (2018) introduced a relaxed notion of *team-maxmin equilibrium with coordination device* (TMECor). In TMECor solutions, the team of players shares their strategies *ex-ante*[6] against the adversary. While TMECor shares some properties of NEs in zero-sum two-player games (e.g., exchangeability), the proposed algorithms by (Celli & Gatti, 2018) leverages tools like Mixed-Integer Linear Program (MILP) that involves a large number of integer variables. To tackle heuristically these hurdles, in (Zhang & An, 2020a;b) had been proposed modified versions of incremental strategy generation (ISG) algorithm (See (McMahan et al., 2003)) for large sized games.

Describing the history and the literature surrounding multi-agent GANs would take us too far afield, so we do not attempt it. Scratching only the surface, we mention that: *a)* In MGAN (Hoang et al., 2018) and MAD-GAN (Li et al., 2019), we have a mixture of generative models with asymptotically infinite capacity networks. *b)* In the seminal case of MIX+GAN (Arora et al., 2017), an extra regularization term is typically added to discourage the weights of a mixture of generators being too far away from uniform. *c)* In Stackelberg GAN, (Zhang et al., 2018) exploit the leader-followers relation of discriminator vs generators to achieve smaller minimax gap and better Frechét metrics. For the case of *multiple-discriminators*: *d)* In GMAN (Durugkar et al., 2016), increased number of agents are used to achieve higher quality images, while *e)* MD-GAN (Hardy et al., 2019) are designed to generalize over different datasets and multiple tasks that are spread on multiple workers.

While the main contribution of this work is to highlight the inherent difficulties that the above heuristic architectures are cursed, our theoretical findings in team competition hint also at new optimization schemes like KPV-GDA whose performance is illustrated in Section 4 with delicate network architectures that contain even multiple generators and discriminators simultaneously.

---

[6]That is, the team members are allowed to discuss and agree on tactics before the game starts, but they cannot communicate during the game.

A.2 PROOF OF THEOREM 3.1

We will reduce the problem of finding a Nash equilibrium in congestion games to the problem of finding a Nash equilibrium in two team zero-sum games. The result then will follow since computing Nash equilibria in congestion games is CLS-hard (Babichenko & Rubinstein (2021)).

A *congestion game* is defined by the tuple $(N; E; (S_i)_{i \in N}; (c_e)_{e \in E})$ where N is the set of *agents*, E is a set of *resources* (also known as *edges* or *facilities*), and each player i has a set S_i of subsets of E. Each strategy $s_i \in S_i$ is a set of edges (a *path*), and c_e is a cost (negative utility) function associated with facility e. For a strategy profile $\mathbf{s} = (s_1, s_2, \ldots, s_N)$, the cost of player i is given by $c_i(\mathbf{s}) = \sum_{e \in s_i} c_e(\ell_e(\mathbf{s}))$, where $\ell_e(\mathbf{s})$ is the number of players using e in $\mathbf{s}$ (the load of edge e). It is a well-known result Rosenthal (1973) that congestion games exhibit a potential function $\Phi(\mathbf{s})$, that

$$\Phi(\mathbf{s}) = \sum_{e \in E} \sum_{j=1}^{\ell_e(\mathbf{s})} c_e(j)$$

with the property that if any agent i changes her strategy to s_i' it holds that

$$\Phi(s_i', \mathbf{s}_{-i}) - \Phi(s_i, \mathbf{s}_{-i}) = c_i(s_i', \mathbf{s}_{-i}) - c_i(s_i, \mathbf{s}_{-i}).$$

Reduction. Consider a congestion game $(N; E; (S_i)_{i \in N}; (c_e)_{e \in E})$ with $n = |N|$ players and potential function Φ. We define a team zero-sum game as follows: Team A has n players, in which each agent i chooses strategies from S_i. Team B has n players, with each agent j having only one possible choice (singleton set of actions) call it d, i.e., these are dummy players. If players from Team A choose strategy profile $\mathbf{s}$ (Team B has only one choice) then they get utility $u_A(\mathbf{s}, d) = -\Phi(s)$. The utility members of Team B get is $u_B(\mathbf{s}, d) = \Phi(s)$.

Let $\mathbf{x}^* \equiv (x_1^*, \ldots, x_n^*)$ and $(d, \ldots, d)$ be a (possibly mixed) Nash equilibrium in the team zero-sum game we defined. We shall show that $(x_1^*, \ldots, x_n^*)$ is a Nash equilibrium of the original congestion game and the reduction will be complete. Aiming for contradiction, suppose $(x_1^*, \ldots, x_n^*)$ is not a Nash equilibrium of the original congestion game. Then there exists an agent i that can deviate from strategy x_i^* to $\tilde{x}_i$ and decrease her expected cost. Hence we have that

$$0 < \mathbb{E}_{\mathbf{s} \sim \mathbf{x}^*}[c_i(\mathbf{s})] - \mathbb{E}_{\mathbf{s} \sim (\tilde{x}_i^*, \mathbf{x}_{-i}^*)}[c_i(\mathbf{s})]$$
$$= \mathbb{E}_{\mathbf{s} \sim \mathbf{x}^*}[\Phi(\mathbf{s})] - \mathbb{E}_{\mathbf{s} \sim (\tilde{x}_i^*, \mathbf{x}_{-i}^*)}[\Phi(\mathbf{s})] \quad \text{(Property of potential)}.$$

We conclude that $\mathbb{E}_{\mathbf{s} \sim \mathbf{x}^*}[u_A(\mathbf{s}, d)] = -\mathbb{E}_{\mathbf{s} \sim \mathbf{x}^*}[\Phi(\mathbf{s})] < -\mathbb{E}_{\mathbf{s} \sim (\tilde{x}_i^*, \mathbf{x}_{-i}^*)}[\Phi(\mathbf{s})] = \mathbb{E}_{\mathbf{s} \sim (\tilde{x}_i^*, \mathbf{x}_{-i}^*)}[u_A(\mathbf{s}, d)]$ which is a contradiction since $(x_1^*, \ldots, x_n^*)$ is a Nash equilibrium for the team zero-sum game hence if player i deviates, her payoff (i.e., the payoff of her Team) should not increase.

A.3 MULTIPLAYER MATCHING PENNIES

Below, we present the exact definition of the (2x2) two-team of two-players matching pennies that we discuss in Sec

		Team B		
		HH	HT/TH	TT
	HH	$-1, 1$	$-1/2, 1/2$	$1, -1$
Team A	HT/TH	$1/2, -1/2$	$0, 0$	$1/2, -1/2$
	TT	$1, -1$	$-1/2, 1/2$	$-1, 1$

A.4 PROOF OF THEOREM 3.2

Since $(\mathbf{x}^*, \mathbf{y}^*)$ is not weakly-stable, there exist players i, j from the same team (say B without loss of generality) and strategies k, l, l' so that so that if i is forced to play k, then j's best response

is l and that gives larger payoff than another strategy l' in her support. Formally it holds that (by multi-linearity of U)

$$\text{(payoff if } i, j \text{ choose } k, l) \; \frac{\partial^2 U(\mathbf{x}^*, \mathbf{y}^*)}{\partial y_{ik} \partial y_{jl}} > \frac{\partial^2 U(\mathbf{x}^*, \mathbf{y}^*)}{\partial y_{ik} \partial y_{jl'}} \; \text{(payoff if } i, j \text{ choose } k, l'), \quad \text{(A.1)}$$

and also $\frac{\partial U(\mathbf{x}^*, \mathbf{y}^*)}{\partial y_{jl}} = \frac{\partial U(\mathbf{x}^*, \mathbf{y}^*)}{\partial y_{jl'}}$ (*). We shall show that $\nabla^2_{\mathbf{yy}} U(\mathbf{x}^*, \mathbf{y}^*)$ has a strictly positive eigenvalue tangent in the product of simplices (note that if we were working with A, we would show that $\nabla^2_{\mathbf{xx}} U(\mathbf{x}^*, \mathbf{y}^*)$ has a strictly negative eigenvalue). Consider a vector of size $\sum_i |S_i|$, (where $|S_i|$ is the cardinality of the strategy space of agent i in Team B) which has 1 at coordinates $(i, k), (j, l)$, -1 at coordinate (j, l') and from which we subtract y_i^*; we denote by $\mathbf{v}$ the resulting vector. We shall show that $\mathbf{v}^\top \nabla^2_{\mathbf{xx}} U(\mathbf{x}^*, \mathbf{y}^*) \mathbf{v} < 0$.

By multilinearity of U it follows that $\frac{\partial^2 U}{\partial y_{js} \partial y_{js'}} = 0$ (**) for all agents j and strategies s, s' and the same is true for x variables (team A). We conclude that

$$\frac{1}{2} \mathbf{v}^\top \nabla^2_{\mathbf{yy}} U(\mathbf{x}^*, \mathbf{y}^*) \mathbf{v} = \frac{\partial^2 U(\mathbf{x}^*, \mathbf{y}^*)}{\partial y_{ik} \partial y_{jl}} - \frac{\partial^2 U(\mathbf{x}^*, \mathbf{y}^*)}{\partial y_{ik} \partial y_{jl'}} - \sum_{s \in S_i} y_{is}^* \left(\frac{\partial^2 U(\mathbf{x}^*, \mathbf{y}^*)}{\partial y_{is} \partial y_{jl}} - \frac{\partial^2 U(\mathbf{x}^*, \mathbf{y}^*)}{\partial y_{is} \partial y_{jl'}} \right)$$

$$= \frac{\partial^2 U(\mathbf{x}^*, \mathbf{y}^*)}{\partial y_{ik} \partial y_{jl}} - \frac{\partial^2 U(\mathbf{x}^*, \mathbf{y}^*)}{\partial y_{ik} \partial y_{jl'}} - \left(\frac{\partial U(\mathbf{x}^*, \mathbf{y}^*)}{\partial y_{jl}} - \frac{\partial U(\mathbf{x}^*, \mathbf{y}^*)}{\partial y_{jl'}} \right)$$

$$\overset{(*)}{=} \frac{\partial^2 U(\mathbf{x}^*, \mathbf{y}^*)}{\partial y_{ik} \partial y_{jl}} - \frac{\partial^2 U(\mathbf{x}^*, \mathbf{y}^*)}{\partial y_{ik} \partial y_{jl'}} \overset{(A.1)}{>} 0.$$

Therefore $\nabla^2_{\mathbf{yy}} U(\mathbf{x}^*, \mathbf{y}^*)$ has a positive eigenvalue and as a result

$$R := \begin{pmatrix} -\nabla^2_{\mathbf{xx}} U(\mathbf{x}^*, \mathbf{y}^*) & \mathbf{0} \\ \mathbf{0} & \nabla^2_{\mathbf{yy}} U(\mathbf{x}^*, \mathbf{y}^*) \end{pmatrix} \quad \text{(A.2)}$$

must have a positive and a negative eigenvalue (since the trace is zero).

We consider the Jacobian of the GDA dynamics at $(\mathbf{x}^*, \mathbf{y}^*)$. The corresponding matrix is the following:

$$J_{\text{GDA}} = \mathbf{I} + \eta \begin{pmatrix} -\nabla^2_{\mathbf{xx}} U(\mathbf{x}^*, \mathbf{y}^*) & -\nabla^2_{\mathbf{xy}} U(\mathbf{x}^*, \mathbf{y}^*) \\ \nabla^2_{\mathbf{yx}} U(\mathbf{x}^*, \mathbf{y}^*) & \nabla^2_{\mathbf{yy}} U(\mathbf{x}^*, \mathbf{y}^*) \end{pmatrix}, \quad \text{(A.3)}$$

We will show that J_{GDA} has an eigenvalue (possible complex) with absolute value greater than one. It suffices to show that $J_{\text{GDA}} - \mathbf{I}$ has an eigenvalue with positive real part (because then J_{GDA} would have an eigenvalue with real part greater than 1 and hence magnitude greater than one). Due to (**), we get that $J_{\text{GDA}} - \mathbf{I}$ has trace zero. To reach contradiction suppose that no eigenvalue of $J_{\text{GDA}} - \mathbf{I}$ has positive real part, then all eigenvalues of $J_{\text{GDA}} - \mathbf{I}$ should be imaginary or zero. But an imaginary eigenvalue (that is not zero) also results in an eigenvalue with magnitude greater than one for J_{GDA}, therefore all eigenvalues of $J_{\text{GDA}} - \mathbf{I}$ should be zero. We use Ky Fan inequalities which states that the sequence (in decreasing order) of the eigenvalues of $\frac{1}{2}(H + H^\top)$ majorizes the real part of the sequence of the eigenvalues of H (see Moslehian (2011), page 4) for any matrix H. We choose $H = \frac{1}{\eta} \cdot (J_{\text{GDA}} - \mathbf{I})$. Since R has both a negative and a positive eigenvalue, we get that H has an eigenvalue with negative real part. The claim follows since H has trace zero, thus it should have an eigenvalue with positive real part as well.

We conclude that J_{GDA} has an eigenvalue with absolute value greater than one. Using Theorem 2.2 in Daskalakis & Panageas (2018), it occurs that the set of initial conditions so that GDA converge to $(\mathbf{x}^*, \mathbf{y}^*)$ is of measure zero (for the particular choice of the stepsize).

B Proof of Lemma 3.3

Firstly, we start with the min-max formulation of GMP game:

$$\min_{\mathbf{x} \in [0,1]^2} \max_{\mathbf{y} \in [0,1]^2} -x_1 x_2 y_1 y_2 - (1 - x_1)(1 - x_2)(1 - y_1)(1 - y_2) + x_1 x_2 (1 - y_1)(1 - y_2) +$$

$$+ (1 - x_1)(1 - x_2) y_1 y_2 + \omega \left(1 - x_1 x_2 - (1 - x_1)(1 - x_2)\right) \left(y_1 y_2 + (1 - y_1)(1 - y_2)\right) -$$

$$- \omega \left(1 - y_1 y_2 - (1 - y_1)(1 - y_2)\right) \left(x_1 x_2 + (1 - x_1)(1 - x_2)\right)$$

or after simplification it is equivalent with

$$\min_{\mathbf{x}\in[0,1]^2} \max_{\mathbf{y}\in[0,1]^2} (\omega+1)(x_1+x_2)+(1-\omega)(y_1+y_2)-(x_1+x_2)(y_1+y_2)-2\omega x_1 x_2+2\omega y_1 y_2. \quad \text{(B.1)}$$

Remark B.1. Please note that the objective in the min-max is *multi-linear* and the degree of each variable in every summand is at most one (total degree is 2). Moreover note that due to the non convexity-concavity of the function above, the max-min is not equal to the min-max.

Let $(x_1^*, x_2^*, y_1^*, y_2^*)$ be a Nash equilibrium. Assuming $x_1^*, x_2^*, y_1^*, y_2^* \in (0,1)$ from (B.1) and first-order conditions we get the system of equations

1. $\omega + 1 - y_1^* - y_2^* - 2\omega x_2^* = 0$,
2. $\omega + 1 - y_1^* - y_2^* - 2\omega x_1^* = 0$,
3. $1 - \omega - x_1^* - x_2^* + 2\omega y_2^* = 0$,
4. $1 - \omega - x_1^* - x_2^* + 2\omega y_1^* = 0$.

Combining the first two equations, we have $x_1^* = x_2^*$ and combining the last two it follows $y_1^* = y_2^*$. Dividing equation one by ω and subtracting three, it follows that $\frac{1}{\omega} - 2\frac{y_1^*}{\omega} + \omega - 2\omega y_1^* = 0$. Hence we conclude that $y_1^* = \frac{1}{2}$. As a result $y_2^* = \frac{1}{2}$ and substituting in first equation $x_1^* = x_2^* = \frac{1}{2}$.

- Assume now that $x_1^* = 0$ and $x_2^*, y_1^*, y_2^* \in (0,1)$. Following the same idea, now only equations 2, 3, 4 hold and instead of the first we have the constraint $\omega + 1 - y_1^* - y_2^* - 2\omega x_2^* \geq 0$. From 3, 4 we conclude that $y_1^* = y_2^*$ and using 2 it holds that $y_1^* = y_2^* = \frac{1+\omega}{2}$. Using 3 follows that $1 - \omega - x_2^* + \omega(\omega+1) = 0$. Thus $x_2^* = 1 + \omega^2 > 1$ (this is not possible because $x_2^* \in [0,1]$).

- Consider the case that $x_1^* = 1$ and $x_2^*, y_1^*, y_2^* \in (0,1)$. Only equations 2, 3, 4 hold and instead of the first we have the constraint $\omega + 1 - y_1^* - y_2^* - 2\omega x_2^* \leq 0$. From 3, 4 we conclude that $y_1^* = y_2^*$ and using 2 it holds that $y_1^* = y_2^* = \frac{1-\omega}{2}$. Using 3 follows that $-\omega - x_2^* + \omega(1-\omega) = 0$. Thus $x_2^* = -\omega^2 < 0$ (this is not possible because $x_2^* \in [0,1]$). By symmetry the same happens when $x_2^* = 0$ or $x_2^* = 1$ and $x_1^*, y_1^*, y_2^* \in (0,1)$.

- Case $x_1^* = x_2^* = 0$ and $y_1^*, y_2^* \in (0,1)$. Using 3, 4 we get $y_1^* = y_2^* = \frac{\omega-1}{2\omega} < 0$ (this is not possible).

- Case $x_1^* = x_2^* = 1$ and $y_1^*, y_2^* \in (0,1)$. Using 3, 4 we get $y_1^* = y_2^* = \frac{\omega+1}{2\omega} > 1$ (this is not possible).

- Case $x_1^* = 0$ and $x_2^* = 1$ and $y_1^*, y_2^* \in (0,1)$. Using 3, 4 we get $y_1^* = y_2^* = \frac{1}{2}$. Moreover one becomes $\omega - 2\omega x_2^* \geq 0$ and two $\omega - 2\omega x_1^* \leq 0$, that is $x_1^* \geq \frac{1}{2}$ and $x_2^* \leq \frac{1}{2}$ (contradiction). The case $x_1^* = 1$ and $x_2^* = 0$ and $y_1^*, y_2^* \in (0,1)$ is symmetric.

Similarly one can consider the case where the $\mathbf{y}$ team plays pure and $x_1^*, x_2^* \in (0,1)$. One can also check that all possible pure strategy profiles are not Nash equilibria.

C PROOF OF THEOREM 3.5

We split the proof in 3 parts. Before we start the proof, note that the Hessian of U (C.3) has infinity norm less than 4 (since $\omega \in (0,1)$), so U has gradient Lipschitz with $L \leq 4$. Thus for the rest of the proof for GDA, we choose $\eta_{\text{OGDA}} < \frac{1}{4}$.

GDA. For GDA the proof will be straightforward. We will show that $(x_1^*, x_2^*, y_1^*, y_2^*)$ is a weakly Nash equilibrium. Then the claim about GDA will follow because of Lemma 3.3, Theorem 3.2 and Remark 2.2.

Assume that player x_1 fixes his strategy to $x_1 = 0$. and y_1, y_2 keep their strategy $(\frac{1}{2}, \frac{1}{2})$. We shall show that x_2 is not indifferent in his support and would like to change his mixed strategy $x_2 = \frac{1}{2}$ to pure. When $x_1 = 0$ and $y_1 = y_2 = \frac{1}{2}$ the payoff of Team A (x variables) becomes $-\omega x_2 - 1 + \frac{\omega}{2}$. Since $\omega \in (0,1)$, x_2 prefers to play $x_2 = 0$ (instead of $\frac{1}{2}$ she had). We conclude that $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ is

not a weakly-stable Nash equilibrium.

OGDA. The Jacobian of the update rule of OGDA dynamics (use the same machinery of Section 3 in Daskalakis & Panageas (2018)) is the following:

$$
J_{\text{OGDA}} = \begin{pmatrix}
\mathbf{I} - 2\eta_{\text{OGDA}}\nabla^2_{\mathbf{xx}}U & -2\eta_{\text{OGDA}}\nabla^2_{\mathbf{xy}}U & \eta_{\text{OGDA}}\nabla^2_{\mathbf{xx}}U & \eta_{\text{OGDA}}\nabla^2_{\mathbf{xy}}U \\
2\eta_{\text{OGDA}}\nabla^2_{\mathbf{yx}}U & \mathbf{I} + 2\eta_{\text{OGDA}}\nabla^2_{\mathbf{yy}}U & -\eta_{\text{OGDA}}\nabla^2_{\mathbf{yx}}U & -\eta_{\text{OGDA}}\nabla^2_{\mathbf{yy}}U \\
\mathbf{I} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\
\mathbf{0} & \mathbf{I} & \mathbf{0} & \mathbf{0}
\end{pmatrix},
$$
(C.1)

where U is the payoff of Team B (max) and $-U$ is the payoff of team A. Substituting for MPG payoff at Nash equilibrium, we have

$$
\nabla^2_{\mathbf{xx}}U = \begin{pmatrix} 0 & -2\omega \\ -2\omega & 0 \end{pmatrix}, \nabla^2_{\mathbf{yy}}U = \begin{pmatrix} 0 & 2\omega \\ 2\omega & 0 \end{pmatrix}, \nabla^2_{\mathbf{xy}}U = \begin{pmatrix} -1 & -1 \\ -1 & -1 \end{pmatrix}.
$$

The corresponding Jacobian matrix becomes:

$$
J_{\text{OGDA}} = I + \eta_{\text{OGDA}} \begin{pmatrix}
0 & 4\omega & 2 & 2 & 0 & -2\omega & -1 & -1 \\
4\omega & 0 & 2 & 2 & -2\omega & 0 & -1 & -1 \\
-2 & -2 & 0 & 4\omega & 1 & 1 & 0 & -2\omega \\
-2 & -2 & 4\omega & 0 & 1 & 1 & -2\omega & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 1 & 0 & 0 & 0 & 0
\end{pmatrix},
$$
(C.2)

It turns out that matrix (C.2) has an eigenvalue that is $1 + \eta_{\text{OGDA}} \cdot 2\omega\left(\sqrt{1 + \frac{1}{2\omega}} - 1\right) > 1$ with corresponding right eigenvector $\left(2\omega\left(1 - \sqrt{1 + \frac{1}{2\omega}}\right), 2\omega\left(-1 + \sqrt{1 + \frac{1}{2\omega}}\right), 0, 0, -1, 1, 0, 0\right)^{\top}$ for all $\eta_{\text{OGDA}} > 0$. It is also easy to see the Hessian of U, that is

$$
\begin{pmatrix} \nabla^2_{\mathbf{xx}}U & \nabla^2_{\mathbf{xy}}U \\ \nabla^2_{\mathbf{yx}}U & \nabla^2_{\mathbf{yy}}U \end{pmatrix} = \begin{pmatrix}
0 & -2\omega & -1 & -1 \\
-2\omega & 0 & -1 & -1 \\
-1 & -1 & 0 & 2\omega \\
-1 & -1 & 2\omega & 0
\end{pmatrix}
$$
(C.3)

is invertible for $\omega \in (0,1)$ (this is true as the eigenvalues are $-2\omega, 2\omega, -2\sqrt{1 + \omega^2}, 2\sqrt{1 + \omega^2}$, non of which is zero. From Theorem 3.2 in Daskalakis & Panageas (2018) follows that the initial conditions so that OGDA converges to the Nash equilibrium is of measure zero for stepsize $\eta_{\text{OGDA}} < \frac{1}{2L} \leq \frac{1}{8}$ (where L is the Lipschitz constant of ∇U).

EG. The Jacobian of EG dynamics computed at the Nash equilibrium (fixed point) is given below (made use of chain rule):

$$
J_{\text{EG}} = \mathbf{I} + \eta_{\text{EG}} \begin{pmatrix}
-\nabla^2_{\mathbf{xx}}U(\mathbf{x}^*, \mathbf{y}^*) & -\nabla^2_{\mathbf{xy}}U(\mathbf{x}^*, \mathbf{y}^*) \\
\nabla^2_{\mathbf{yx}}U(\mathbf{x}^*, \mathbf{y}^*) & \nabla^2_{\mathbf{yy}}U(\mathbf{x}^*, \mathbf{y}^*)
\end{pmatrix} + \eta^2_{\text{EG}} \begin{pmatrix}
-\nabla^2_{\mathbf{xx}}U(\mathbf{x}^*, \mathbf{y}^*) & -\nabla^2_{\mathbf{xy}}U(\mathbf{x}^*, \mathbf{y}^*) \\
\nabla^2_{\mathbf{yx}}U(\mathbf{x}^*, \mathbf{y}^*) & \nabla^2_{\mathbf{yy}}U(\mathbf{x}^*, \mathbf{y}^*)
\end{pmatrix}^2.
$$
(C.4)

We substitute with the values and we get

$$
J_{\text{EG}} = \mathbf{I} + \eta_{\text{EG}} \begin{pmatrix}
0 & 2\omega & 1 & 1 \\
2\omega & 0 & 1 & 1 \\
-1 & -1 & 0 & 2\omega \\
-1 & -1 & 2\omega & 0
\end{pmatrix} + \eta^2_{\text{EG}} \begin{pmatrix}
4\omega^2 - 2 & -2 & 4\omega & 4\omega \\
-2 & 4\omega^2 - 2 & 4\omega & 4\omega \\
-4\omega & -4\omega & -2 + 4\omega^2 & -2 \\
-4\omega & -4\omega & -2 & -2 + 4\omega^2
\end{pmatrix}.
$$
(C.5)

The eigenvalues of J_{EG} are $1 + \eta_{\text{EG}}(-2\omega + 4\eta_{\text{EG}}\omega^2), 1 + \eta_{\text{EG}}(-2\omega + 4\eta_{\text{EG}}\omega^2)$,
$1 + \eta_{\text{EG}}\left(-4\eta_{\text{EG}} + 2\omega + 4\eta_{\text{EG}}\omega^2 - 2\sqrt{-1 - 8\eta_{\text{EG}}\omega - 16\eta^2_{\text{EG}}\omega^2}\right)$, and
$1 + \eta_{\text{EG}}\left(-4\eta_{\text{EG}} + 2\omega + 4\eta_{\text{EG}}\omega^2 + 2\sqrt{-1 - 8\eta_{\text{EG}}\omega - 16\eta^2_{\text{EG}}\omega^2}\right)$. If $-4\eta_{\text{EG}} + 2\omega + 4\eta_{\text{EG}}\omega^2 > 0$

then J_{EG} will have an eigenvalue with absolute value greater than one. A sufficient condition for that is $\eta_{\text{EG}} \leq \frac{\omega}{2}$. Finally, for the same choice of stepsizes we have that J_{EG} is invertible for all $(\mathbf{x}, \mathbf{y})$, hence the EG dynamics is a local diffeomorphism. From Theorem 2 in arxiv version of Lee et al. (2019) the claim for EG method follows.

D PROOF OF THEOREM 3.7

We first compute the Jacobian of GDA-KPV dynamics (2.2) where we have eliminated one variable per agent (to avoid using the projection operator). The Jacobian has the following form:

$$
J_{\text{KPV-GDA}} = \mathbf{I} + \eta \left(K + \begin{pmatrix} -\nabla_{\mathbf{xx}}U(\mathbf{x}^*, \mathbf{y}^*) & -\nabla_{\mathbf{yx}}U(\mathbf{x}^*, \mathbf{y}^*) \\ \nabla_{\mathbf{xy}}U(\mathbf{x}^*, \mathbf{y}^*) & \nabla_{\mathbf{yy}}U(\mathbf{x}^*, \mathbf{y}^*) \end{pmatrix} \quad -K \\ P \qquad\qquad\qquad\qquad\qquad -P \right). \tag{D.1}
$$

It suffices to show that there exist matrices K, P so that J_{KPV} has all its eigenvalues in absolute value less than one (lie inside the unit sphere).

We get (trivially from identity matrix) that $[\lambda \mathbf{I} - \mathbf{I} - \eta\tilde{H} \quad \mathbf{I}]$ has full rank for all λ, thus $(\mathbf{I} + \eta\tilde{H}, B)$ is stabilizable (see Sontag (1998) for more information on control theory, controlizable and stabilizable pair of matrices). Moreover by assumption we have that $\tilde{H} := \begin{pmatrix} -\nabla_{\mathbf{xx}}U(\mathbf{x}^*, \mathbf{y}^*) & -\nabla_{\mathbf{xy}}U(\mathbf{x}^*, \mathbf{y}^*) \\ \nabla_{\mathbf{yx}}U(\mathbf{x}^*, \mathbf{y}^*) & \nabla_{\mathbf{yy}}U(\mathbf{x}^*, \mathbf{y}^*) \end{pmatrix}$ is invertible, thus Proposition 5 from Hassouneh et al. (2004) indicates that there exist matrices K, P such that $(\mathbf{x}^*, \mathbf{y}^*, \mathbf{x}^*, \mathbf{y}^*)$ (fixed point in the lifted system) is asymptotically stable. As a result we have local convergence around the Nash equilibrium and the claim follows.

E   CIFAR-10

CIFAR-10 is a well established test bed for various GAN architectures. It contains 10 balanced classes of images of different objects.

Just to further illustrate the merits of multi-agent GANs, we offer a selection of images generated by WGAN-GP Arjovsky et al. (2017) and the MGAN Hoang et al. (2017).

In the pictures provided by the WGAN-GP we see that across iterations, the generated samples tend to cover only certain classes of the dataset while it takes longer for the samples to become realistic.

On the other hand, the MGAN architecture from an early stage provides with diverse samples that tend to be more realistc from early on.

Figure 6: MGAN after 300 iterations

Figure 7: WGAN after 10000 iterations

Figure 8: WGAN after 15000 iterations

Figure 9: WGAN after 25000 iterations