Surrogate Regret Bounds for Polyhedral Losses

Abstract

Surrogate risk minimization is an ubiquitous paradigm in supervised machine learning, wherein a target problem is solved by minimizing a surrogate loss on a dataset. Surrogate regret bounds, also called excess risk bounds, are a common tool to prove generalization rates for surrogate risk minimization. While surrogate regret bounds have been developed for certain classes of loss functions, such as proper losses, general results are relatively sparse. We provide two general results. The first gives a linear surrogate regret bound for any polyhedral (piecewise-linear convex) surrogate, meaning that surrogate generalization rates translate directly to target rates. The second shows that for sufficiently non-polyhedral surrogates, the regret bound is a square root, meaning fast surrogate generalization rates translate to slow rates for the target. Together, these results suggest polyhedral surrogates are weakly optimal in many cases.

1 Introduction

In supervised learning, our goal is to learn a hypothesis that solves some target problem given labeled data. The problem is often specified by a target loss ℓ that is discrete in the sense that the prediction lies in a finite set, as with classification, ranking, and structured prediction. As minimizing the target ℓ over a data set is typically NP-hard, we instead turn to the general paradigm of *surrogate risk minimization*. Here a surrogate loss function L is chosen and minimized instead of the target ℓ. A link function ψ then maps the surrogate predictions to target predictions.

Naturally, a central question in surrogate risk minimization is how to choose the surrogate L. A minimal requirement is statistical *consistency*, which roughly says that approaching the best possible surrogate loss will eventually lead to the best possible target loss. A more precise goal is to understand the rate at which this minimization occurs, called the *generalization rate*: how quickly the target loss approaches the best possible, in terms of the number of data points. In other words, if we define the *regret* of a hypothesis to be its performance relative to the Bayes optimal hypothesis, we wish to know how quickly the ℓ-regret approaches zero.

Surrogate regret bounds are a common tool to prove generalization rates for surrogate risk minimization, by bounding the target regret as a function of the surrogate regret. This bound allows generalization rates for surrogate losses—of which there are many, often leveraging properties of L like strong convexity and smoothness—to imply rates for the target problem. Roughly speaking, a surrogate regret bound takes the form

$$\forall \text{ surrogate hypotheses } h, \quad \text{Regret}_\ell(\psi \circ h) \leq \zeta(\text{Regret}_L(h)), \tag{1}$$

where the *regret transfer function* $\zeta : \mathbb{R}_+ \to \mathbb{R}_+$ controls how surrogate regret translates to target regret. For example, if we have a sequence of surrogate hypotheses h_n achieving a fast rate $\text{Regret}_L(h_n) = O(1/n)$, and we have a regret transfer of $\zeta(\epsilon) = \sqrt{\epsilon}$, then we can only guarantee a slower target rate of $\text{Regret}_\ell(\psi \circ h_n) = O(1/\sqrt{n})$. The ideal regret transfer is *linear*, i.e. $\zeta(\epsilon) = C \cdot \epsilon$ for some $C > 0$, which would have directly translated the fast-rate guarantee to the target.

Much is known about surrogate regret bounds for particular problems, especially for binary classification and related problems like bipartite ranking; see § 1.1 for a discussion of the literature. General results which span multiple target problems, however, are sparse. Furthermore, we still lack guidance as to which surrogate will lead to the best overall target generalization rate, as this rate depends on both the surrogate generalization rate and the regret transfer function. For example, consider the choice between a *polyhedral* (piecewise-linear and convex) surrogate, like hinge loss, or a smooth surrogate like exponential or squared loss. The surrogate rate for polyhedral losses will likely be a slower $O(1/\sqrt{n})$, whereas one typically achieves a faster $O(1/n)$ rate with smooth surrogates—but what is the tradeoff when it comes to the regret transfer function ζ, which ultimately decides the target generalization rate? To answer these types of questions, we need more general results.

Our first result is a general surrogate regret bound for polyhedral surrogates. Perhaps suprisingly, we show that *every* consistent surrogate in this class achieves a linear regret transfer, meaning their surrogate generalization rates translate directly to target rates, up to a constant factor.

Theorem 1 (Linear bound for polyhedral). *Let L be any polyhedral surrogate and ψ any link function such that (L, ψ) is consistent for the target loss ℓ. Then L satisfies Equation 1 with a linear ζ.*

The recent embedding framework of Finocchiaro et al. [8] shows how to construct consistent polyhedral surrogates for any discrete target loss; Theorem 1 additionally endows these surrogates with linear regret bounds. In particular, this result immediately applies to the Weston–Watkins surrogate [24], Lovász hinge [8, 26], and other polyhedral surrogates in the literature currently lacking regret bounds.

We also give a partial converse: If the surrogate is sufficiently "non-polyhedral", i.e. smooth and locally strongly convex, then ζ cannot shrink faster than a square root. For these losses, therefore, even fast surrogate generalization rates would likely translate to slow rates for the target.

Theorem 2 (Lower bound for "non-polyhedral"). *Let L be a locally strongly convex surrogate with locally Lipschitz gradient. If L satisfies Equation 1 for a target loss ℓ, then there exists $c > 0$ such that, for all small enough $\epsilon \geq 0$, we have $\zeta(\epsilon) \geq c \cdot \sqrt{\epsilon}$.*

While it may seem intuitive that $\zeta = \sqrt{\cdot}$ for strongly convex losses, it is well-known that losses like exponential loss for binary classification also have $\zeta = \sqrt{\cdot}$, even though they are neither strongly convex nor strongly smooth [4]. Theorem 2 offers an explanation for this phenomenon, and together with Theorem 1 clarifies an intriguing trend in the literature: ζ is typically either linear or square-root.

Taken together, our results shed light on the dichotomy expressed above: slower polyhedral surrogate rates of $O(1/\sqrt{n})$ will translate to the target unchanged, while most smooth surrogates, even if achieving a fast surrogate generalization rate of $O(1/n)$, will likely yield the same target rate of $O(1/\sqrt{n})$. Our results therefore suggest an inherent tradeoff fast surrogate rates and optimal regret transfer, a phenomenon also observed by Mahdavi et al. [11] for binary classification. In particular, our results suggest that polyhedral losses may be weakly optimal in many cases.

Our proof of the upper bound, Theorem 1, is nonconstructive. After proving the main results (Sections 3 and 4), we make it constructive in Section 5. Specifically, a polyhedral loss's regret transfer function can be bounded by $\zeta(t) \leq \alpha \cdot t$ where the constant is of the form $\alpha = \alpha_\ell \cdot \alpha_L \cdot \alpha_\psi$. Each component of α is constructed as a function of its respective object ℓ, L, and ψ. A corollary of the derivation is that a polyhedral loss L and link ψ are consistent for ℓ if and only if they satisfy the weak indirect elicitation criterion and ψ is ϵ-*separated*, a condition introduced in Finocchiaro et al. [8].

1.1 Literature on surrogate regret bounds

Surrogate regret bounds were perhaps developed first by Zhang et al. [28], as a method to prove statistical consistency for various surrogate losses for binary classification. These results were then generalized by Bartlett et al. [4] for margin losses, and Reid and Williamson [19] for composite proper losses. These latter works and others give a fairly complete picture for binary classification: a characterization of calibrated surrogates, and precise closed forms for the optimal ζ for each. By extension, close variations on binary classification are also well-understood, such as cost-sensitive binary classification [21], binary classification with an abstain option [3, 27], bipartite ranking [1, 12], and ordinal regression via reduction to binary classification [15]. For multiclass classification, we have nice consistency characterizations with respect to 0-1 loss [23], with some surrogate regret bounds [7, 16].

Beyond binary classification and 0-1 loss, our knowledge drops off precipitously, with few general results. There are some works which provide partial understand for specific target problems. For example, in the problem of multiclass classification with an abstain option, Ramaswamy et al. [18] give three consistent polyhedral surrogates (two of which are commonly used as inconsistent surrogates for 0-1 loss), along with linear surrogate regret bounds. Ramaswamy et al. [17] show how to extend these surrogates to nested, hierarchical versions, again providing linear regret bounds. Motivated by these polyhedral surrogates, together with examples arising in top-k prediction [10, 25] and other structured prediction problems [26], Finocchiaro et al. [8] introduce an embedding framework for polyhedral surrogates. This framework shows that all polyhedral surrogates are consistent for some discrete target problem, and vice versa, but do not provide surrogate regret bounds. Our results fill this gap, giving linear regret bounds for all polyhedral losses, and thus all discrete target problems.

Perhaps closest in spirit to our work are several recent works on general structured prediction problems. Ciliberto et al. [6, Theorem 2] and Osokin et al. [14, Theorem 7] give surrogate regret bounds for a quadratic surrogate, for a general class of structed prediction problems. Blondel [5, Proposition 5] gives surrogate regret bounds for a tighter class of quadratic surrogates based on projection oracles. Nowak-Vila et al. [13, Theorem 4.4] give even more general bounds, using a broader class of surrogates based on estimating a vector-valued expectation. All of these works apply to a broad range of discrete target problems. The main difference to our work is that their upper bounds are all for strongly convex surrogates,[1] whereas ours are for polyhedral surrogates. Osokin et al. [14, Theeorem 8] and Nowak-Vila et al. [13, Theorem 4.5] also provide lower bounds, in the same spirit as our Theorem 2 but only for the classes of surrogates they study. In particular, our results recover their bounds up to a constant factor.

2   Preliminaries

Target problems.   We consider supervised machine learning problems where data points have features in $\mathcal{X}$ and labels in $\mathcal{Y}$, a finite set. The *target problem* is to learn a hypothesis mapping $\mathcal{X}$ to a *report* (or *prediction*) *space* $\mathcal{R}$, possibly different from $\mathcal{Y}$. For example, binary classification has $\mathcal{Y} = \mathcal{R} = \{-1, +1\}$, whereas for ranking problems $\mathcal{R}$ may be the set of permutations of $\mathcal{Y}$. In this paper, we assume $\mathcal{R}$ is a finite set. The target problem is specified by the *target loss* $\ell : \mathcal{R} \to \mathbb{R}_+^{\mathcal{Y}}$, which maps a report (prediction) $r \in \mathcal{R}$ to the vector $\ell(r)$ of nonnegative loss values for each label, i.e. $\ell(r) = (\ell(r)_y)_{y \in \mathcal{Y}}$. Given a hypothesis $g : \mathcal{X} \to \mathcal{R}$ and a data point $(x, y) \in \mathcal{X} \times \mathcal{Y}$, the loss is $\ell(g(x))_y$. We make a minimal non-redundancy assumption on ℓ, formalized in Section 2.1. We may refer to ℓ as a *discrete* target loss to emphasize our assumption that $\mathcal{R}$ and $\mathcal{Y}$ are finite.

Surrogate problems.   We recall the surrogate risk minimization framework. First, one constructs a *surrogate loss function* $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$. Then, one minimizes it to learn a hypothesis of the form $h : \mathcal{X} \to \mathbb{R}^d$. Finally, one applies a *link function* $\psi : \mathbb{R}^d \to \mathcal{R}$, which maps h to the implied target hypothesis $\psi \circ h$. In other words, if a surrogate prediction $h(x) \in \mathbb{R}^d$ implies a prediction $\psi(h(x)) \in \mathcal{R}$ for the target problem. For example, in binary classification, the target problem is given by 0-1 loss, $\ell(r)_y = \mathbb{1}\{r \neq y\}$, with surrogate losses typically defined over $\mathbb{R}$. Common surrogate losses include hinge loss and logistic loss. These surrogates are typically associated with the link $\psi(u) = \mathrm{sign}(u)$, which maps negative predictions to -1 and nonnegative predictions to $+1$.

Surrogate regret bounds.   The suitability and efficiency of a chosen surrogate is often quantified with a *surrogate regret bound*. Formally, the *surrogate regret* of h with respect to data distribution $\mathcal{D}$ is given by $R_L(h; \mathcal{D}) = \mathbb{E}_{(X,Y) \sim \mathcal{D}} L(h(X))_Y - \inf_{h' : \mathcal{X} \to \mathbb{R}^d} \mathbb{E}_{(X,Y) \sim \mathcal{D}} L(h'(X))_Y$, where the minimum is taken over all measurable functions. (One may equivalently assume the Bayes optimal hypothesis is in the function class.) Given h, the *target regret* of the implied hypothesis $\psi \circ h$ is $R_\ell(\psi \circ h; \mathcal{D}) = \mathbb{E}_{(X,Y) \sim \mathcal{D}} \ell(\psi(h(X)))_Y - \inf_{h' : \mathcal{X} \to \mathcal{R}} \mathbb{E}_{(X,Y) \sim \mathcal{D}} \ell(h'(X))_Y$, where we recall that ψ is the link function.

A surrogate regret bound answers the following question: when we minimize the surrogate regret at a rate $f(n)$, given n data points, how fast does the target regret diminish? Formally, we say that

---

[1]These works express surrogate regret bounds with ζ^{-1} appearing on the left-hand side of eq. (1), and thus the terms "upper bound" and "lower bound" are reversed from their terminology.

139 $\zeta : \mathbb{R}_+ \to \mathbb{R}_+$ is a *regret transfer function* if ζ is continuous at 0 and satisfies $\zeta(0) = 0$. Given a
140 rerget transfer function ζ, we say that (L, ψ) guarantees a *regret transfer of* ζ for ℓ if

$$\forall \mathcal{D}, \forall h : \mathcal{X} \to \mathbb{R}^d, \quad R_\ell(\psi \circ h; \mathcal{D}) \leq \zeta(R_L(h; \mathcal{D})) . \tag{2}$$

141 We note that a classic minimal requirement for (L, ψ) is that they be *consistent* for ℓ, which is the
142 case if there exists any regret transfer ζ guaranteed by (L, ψ) for ℓ [22]. In other words, consistency
143 implies that $R_L(h; \mathcal{D}) \to 0 \implies R_\ell(\psi \circ h; \mathcal{D}) \to 0$, but says nothing about the relative rates.

144 **Polyhedral surrogates.** A function $f : \mathbb{R}^d \to \mathbb{R}$ is called *polyhedral* if it can be written as a
145 pointwise maximum of a finite set of affine functions [20, § 19]. The epigraph of a polyhedral
146 function is a polyhedral set, i.e. the intersection of a finite set of closed halfspaces; thus polyhedral
147 functions are always convex. We say a surrogate loss $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$ is *polyhedral* if, for each fixed
148 $y \in \mathcal{Y}$, the loss as a function of prediction, i.e. $r \mapsto L(r)_y$, is polyhedral.

2.1 Property elicitation and the conditional approach

150 Our technical approach is based on property elicitation, which studies the structure of ℓ and L as
151 they relate to $\Delta_{\mathcal{Y}}$. This involves a *conditional* perspective. Given a data distribution $\mathcal{D}$ on $\mathcal{X} \times \mathcal{Y}$,
152 for $(X, Y) \sim \mathcal{D}$, we consider the format of possible target predictions $g(x) = r \in \mathcal{R}$; surrogate
153 predictions $h(x) = u \in \mathbb{R}^d$; and conditional distributions $\Pr[\cdot \mid X] = p \in \Delta_{\mathcal{Y}}$. By dropping the role
154 of the hypotheses and the feature space $\mathcal{X}$, we can focus on the relationships of the functions $\ell(r)_y$
155 and $L(u)_y$ and the conditional distribution p.

156 In particular, in our notation the expected L-loss of prediction u on distribution p is simply the
157 inner product $\langle p, L(u) \rangle$. The *Bayes risk* of L is $\underline{L}(p) = \inf_{u \in \mathbb{R}^d} \langle p, L(u) \rangle$, and similarly the Bayes
158 risk of ℓ is $\underline{\ell}(p) = \inf_{r \in \mathcal{R}} \langle p, \ell(r) \rangle$. We abuse notation to define the *conditional surrogate regret*
159 $R_L(u, p) = \langle p, L(u) \rangle - \underline{L}(p)$, and similarly the *conditional target regret* $R_\ell(r, p) = \langle p, \ell(r) \rangle - \underline{\ell}(p)$.
160 Given a regret transfer function ζ, we say (L, ψ) guarantees a *conditional regret transfer* of ζ for ℓ if,
161 for all $p \in \Delta_{\mathcal{Y}}$ and all $u \in \mathbb{R}^d$, $R_\ell(\psi(u), p) \leq \zeta(R_L(u, p))$.

162 The following is a direct result of Jensen's inequality, using that $R_L(h; \mathcal{D}) = \mathbb{E}_X R_L(h(X), p_X)$
163 where p_X is the conditional distribution on Y given X.

164 **Observation 1.** *If (L, ψ) guarantee a conditional regret transfer of ζ for ℓ, and ζ is concave, then*
165 *(L, ψ) guarantee a regret transfer of ζ for ℓ.*

166 **Property elicitation.** To begin, we define a *property*, which is essentially a statistic or summary of
167 a distribution, such as the mode. We then define what it means for a loss function to *elicit* a property.
168 We write $\Gamma : \Delta_{\mathcal{Y}} \rightrightarrows R$ as shorthand for $\Gamma : \Delta_{\mathcal{Y}} \to 2^R \setminus \{\emptyset\}$.

169 **Definition 1** (Property, level set). *A property is a function $\Gamma : \Delta_{\mathcal{Y}} \rightrightarrows R$, for some set R. The level*
170 *set of Γ for report r is the set $\Gamma_r = \{p \in \Delta_{\mathcal{Y}} : r \in \Gamma(p)\}$.*

171 The following definition applies both to target losses and surrogate losses.

172 **Definition 2** (Elicits, report space). *A loss function $f : R \to \mathbb{R}_+^{\mathcal{Y}}$, where the domain R is called the*
173 *report space, elicits a property $\Gamma : \Delta_{\mathcal{Y}} \rightrightarrows \mathcal{R}$ if*

$$\forall p \in \Delta_{\mathcal{Y}}, \quad \Gamma(p) = \arg \min_{r \in R} \langle p, f(r) \rangle =: \mathrm{prop}[f](p) . \tag{3}$$

174 As each loss f elicits a unique property, we refer to it as $\mathrm{prop}[f]$. In particular, we will consistently
175 use the notation $\Gamma = \mathrm{prop}[L]$ where L is a surrogate loss with report space $\mathbb{R}^d$, and we will use
176 $\gamma = \mathrm{prop}[\ell]$ where ℓ is a target loss with finite report space $\mathcal{R}$.

177 We will assume throughout that the target loss is *non-redundant*, meaning that all $r \in \mathcal{R}$ are possibly
178 uniquely optimal. Formally, we suppose that for all $r \in \mathcal{R}$, there exists $p \in \Delta_{\mathcal{Y}}$ with $\gamma(p) = \{r\}$.[2]

---

[2]Technically, this definition differs from the one given in literature [8, 9], but the two are equivalent for finite elicitable properties.

Calibration and consistency. It is known that consistency implies the weaker condition of *calibration*, which implies the still-weaker, but very useful, condition of *indirect elicitation*. To define these conditions, consider L, ψ, ℓ and $\gamma = \mathrm{prop}[\ell]$. Recall that $\gamma(p)$ is the set of target reports that are optimal for p. The typical definition of indirect elicitation, which we reformulate below, is as follows: a surrogate-link pair (L, ψ) *indirectly elicits* γ if $u \in \mathrm{prop}[L](p) \implies \psi(u) \in \gamma(p)$ for all $p \in \Delta_{\mathcal{Y}}$. Note that if $\psi(u) \notin \gamma(p)$, then u is a "bad" (suboptimal) surrogate prediction for p. Let $B_{L,\psi,\ell}(p) = \{R_L(u, p) \, : \, \psi(u) \notin \gamma(p)\}$, the set of expected surrogate losses resulting from bad predictions. Thus, we may equivalently state indirect elicitation as the requirement that suboptimal surrogate reports be bad, which clarifies the relationship to calibration: calibration further requires a nonzero gap between the expected loss of a "good" report and that of any bad one.

Definition 3 (Indirectly Elicits, Calibrated). Let ℓ be a target loss and $\gamma = \mathrm{prop}[\ell]$. A surrogate-link pair (L, ψ) *indirectly elicits* γ if $0 \notin B_{L,\psi,\ell}(p)$ for all $p \in \Delta_{\mathcal{Y}}$. The pair (L, ψ) is *calibrated* for ℓ if $0 < \inf B_{L,\psi,\ell}(p)$ for all $p \in \Delta_{\mathcal{Y}}$.

Fact 1 ([4, 22]). *Let $\ell : \mathcal{R} \to \mathbb{R}_+^{\mathcal{Y}}$ be a target loss with finite $\mathcal{R}$ and $\mathcal{Y}$. If a surrogate and link (L, ψ) are consistent for ℓ, then (L, ψ) are calibrated for L. If (L, ψ) are calibrated for ℓ, then they indirectly elicit ℓ.*

3 Upper Bound: Polyhedral Surrogates

In this section, we describe the proof of Theorem 1, that polyhedral surrogates guarantee linear regret transfers. At a high level, the proof (whose details appear in Appendix A) has two parts:

1. Fix $p \in \Delta_{\mathcal{Y}}$. Prove that there is some constant α_p such that $R_\ell(\psi(u), p) \leq \alpha_p R_L(u, p)$ for all $u \in \mathbb{R}^d$.

2. Apply this result to each p in a carefully chosen finite subset of $\Delta_{\mathcal{Y}}$. Argue that the maximum α_p from this finite set suffices.

The first part will actually hold for any calibrated surrogate. The second part relies on a crucial fact about polyhedral losses L: their properties $\Gamma = \mathrm{prop}[L]$ cover the whole simplex with only a finite number of polyhedral level sets. In other words, although the prediction space $\mathbb{R}^d$ is infinite, one can restrict to just a finite set of prediction points and always have a global minimizer of expected loss.

3.1 Linear transfer for fixed p

First, we establish a linear transfer for any fixed conditional distribution p. This holds for any calibrated surrogate, not necessarily polyhedral.

Lemma 1. *Let $\ell : \mathcal{R} \to \mathbb{R}_+^{\mathcal{Y}}$ be a discrete target loss and suppose the surrogate $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$ and link $\psi : \mathbb{R}^d \to \mathcal{R}$ are calibrated for ℓ. Then for any $p \in \Delta_{\mathcal{Y}}$, there exists $\alpha_p \geq 0$ such that, for all $u \in \mathbb{R}^d$,*

$$R_\ell(\psi(u), p) \leq \alpha_p R_L(u, p).$$

While this result is nice, it is also somewhat nonconstructive and potentially loose. In Section 5, we investigate the constant α_p for polyhedral losses in particular.

3.2 Linear overall transfer function

Given Lemma 1, we might hope to obtain that, for any calibrated surrogate, there exists $\alpha = \sup_p \alpha_p$ such that $R_\ell(\psi(u), p') \leq \alpha R_L(u, p')$ for all p'. However, we know this is false in general: not all surrogates yield linear regret transfers! Indeed, for many surrogates, the supremum will be $+\infty$.

However, polyhedral surrogates have a special structure that allows us to show α is finite. We first present some tools that hold for general surrogates, then the key implication from L being polyhedral.

Lemma 2. *If (L, ψ) indirectly elicits γ, then $\Gamma = \mathrm{prop}[L]$ refines γ in the sense that, for all $u \in \mathbb{R}^d$, there exists $r \in \mathcal{R}$ such that $\Gamma_u \subseteq \gamma_r$.*

The next lemma states that surrogate regret is linear in p on any fixed level set Γ_{u^*} of $\Gamma = \mathrm{prop}[L]$. By combining this fact with Lemma 2, we obtain that target regret is linear on these level sets as well.

Lemma 3. *Suppose (L, ψ) indirectly elicits $\gamma = \mathrm{prop}[\ell]$ and let $\Gamma = \mathrm{prop}[L]$. Then for any fixed $u, u^* \in \mathbb{R}^d$ and $r \in \mathcal{R}$, the functions $R_L(u, \cdot)$ and $R_\ell(r, \cdot)$ are linear on Γ_{u^*} in their second arguments.*

We now arrive at the key property of polyhedral losses. It requires that L be *minimizable:* for all $p \in \Delta_\mathcal{Y}$, there exists some optimal prediction u, i.e. where $\langle p, L(u) \rangle = \inf_{u' \in \mathbb{R}^d} \langle p, L(u') \rangle$. Note that if (L, ψ) indirectly elicits ℓ for some ψ and ℓ then L is minimizable.

Lemma 4. *If $L : \mathbb{R}^d \to \mathbb{R}_+^\mathcal{Y}$ is polyhedral and minimizable, then $\Gamma = \mathrm{prop}[L]$ has a finite set of level sets that union to $\Delta_\mathcal{Y}$. Also, these level sets are polytopes.*

We are now ready to restate and sketch a proof of the main upper bound.

Theorem 3 (Theorem 1 restated)**.** *Suppose the surrogate loss $L : \mathbb{R}^d \to \mathbb{R}_+^\mathcal{Y}$ and link $\psi : \mathbb{R}^d \to \mathcal{R}$ are consistent for the target loss $\ell : \mathcal{R} \to \mathbb{R}_+^\mathcal{Y}$. If L is polyhedral, then (L, ψ) guarantee a linear regret transfer for ℓ, i.e. there exists $\alpha \geq 0$ such that, for all $\mathcal{D}$ and all measurable $h : \mathcal{X} \to \mathbb{R}^d$,*

$$R_\ell(\psi \circ h; \mathcal{D}) \leq \alpha R_L(h; \mathcal{D}).$$

The key point of the proof is that, by Lemma 4, the polyhedral loss L has a finite set $U \subset \mathbb{R}^d$ of predictions such that (a) for each $u \in U$, the level set Γ_u is a polytope, and (b) $\cup_{u \in U} \Gamma_u = \Delta_\mathcal{Y}$. Let $\mathcal{Q}$ be the finite set of all vertices of all these level sets. For any vertex $q \in \mathcal{Q}$, we have a linear regret transfer for this fixed q with constant α_q from Lemma 1. Lemma 3 allows us to write the regret of a general p as a convex combination of the regrets of its containing level set's vertices. So we obtain a bound of $\alpha = \max_{q \in \mathcal{Q}} \alpha_q$.

4 Lower bound

In this section, we show that if a surrogate loss is sufficiently "non-polyhedral", then the best regret transfer it can achieve is $\zeta(\epsilon)$ on the order of $\sqrt{\epsilon}$. (Proofs are deferred to Appendix B.) Specifically, we consider a relaxation of L being both strongly smooth and strongly convex.

We next formalize this relaxation. It will assume L is strongly smooth everywhere and strongly convex around some *boundary prediction for L, ℓ*, which we define to be a $u_0 \in \mathbb{R}^d$ such that, for some $r, r' \in \mathcal{R}$, there exists a distribution p_0 such that $p_0 \in \Gamma_{u_0} \cap \gamma_r \cap \gamma_{r'}$. Here $\Gamma = \mathrm{prop}[L]$ and $\gamma = \mathrm{prop}[\ell]$. The upshot of this definition is that for the conditional distribution p_0, u_0 minimizes expected L-loss, and it may link to either r or r', as either of these minimize expected ℓ-loss. We also note that an *open neighborhood* of u_0 is an open set in $\mathbb{R}^d$ containing u_0.

Assumption 1. $L : \mathbb{R}^d \to \mathbb{R}_+^\mathcal{Y}$ is a surrogate and $\psi : \mathbb{R}^d \to \mathcal{R}$ a link, consistent with the discrete target $\ell : \mathcal{R} \to \mathbb{R}_+^\mathcal{Y}$, satisfying the following: (i) For all $y \in \mathcal{Y}$, the function $L(\cdot)_y$ is differentiable with a locally Lipschitz gradient. (ii) For some $\alpha > 0$ and some $u_0 \in \mathbb{R}^d$, a boundary prediction for L, ℓ, the expected loss function $u \mapsto \langle p, L(u) \rangle$ is α-strongly convex on an open neighborhood of u_0.

We observe that boundary points are guaranteed to exist if (L, ψ) indirectly elicit ℓ. This holds because ℓ has a finite report space $\mathcal{R}$, so for example the fact that its full-dimensional level sets union to the simplex implies the existence of $r, r' \in \mathcal{R}$ and $p_0 \in \gamma_r \cap \gamma_{r'}$; and some surrogate prediction u_0 minimizes expected surrogate loss on p_0.

The following result is our main lower bound.

Theorem 4 (Stronger version of Theorem 2.)**.** *Suppose the surrogate loss L and link ψ satisfy a regret transfer of ζ for a target loss ℓ. If L, ψ, and ℓ satisfy Assumption 1, then there exists $c > 0$ such that, for some $\epsilon^* > 0$, for all $0 \leq \epsilon < \epsilon^*$, $\zeta(\epsilon) \geq c\sqrt{\epsilon}$.*

The key idea of the proof is to fix a boundary report u_0 and consider a sequence of distributions $p_\lambda \to p_0$ as $\lambda \to 0$. Target regret $R_\ell(\psi(u_0), p_\lambda)$ shrinks linearly in λ, but surrogate regret $R_L(u_0, p_\lambda)$ shrinks quadratically. This holds roughly because expected surrogate loss is strongly convex in some neighborhood of u_0 and is strongly smooth elsewhere.[3]

---

[3]We note that a tempting approach instead is to fix a distribution p_0 and consider a sequence of predictions. But, we know from Lemma 1 that this will fail: for any consistent surrogate and any fixed p_0, there is a linear regret transfer if we restrict to p_0.

We observe that Assumption 1 captures a wide range of surrogate losses beyond those that are strongly convex and strongly smooth. For example, exponential loss is not strongly convex, but does satisfy Assumption 1. As another example, consider Huber loss for binary classification; here $\mathcal{Y} = \{\pm 1\}$ and the prediction space is $\mathbb{R}$, and Huber loss can be defined by letting the hinge loss be $t = \max\{0, 1 - ry\}$ and setting $L(r)_y = t^2$ if $t \leq 2$, else $4(t - 1)$. Theorem 4 does capture Huber, but needs the strength of Assumption 1(ii) as Huber is strongly convex on $[-1, 1]$ but linear outside that interval.

5 Unpacking the constant for polyhedral losses

Our main upper bound for polyhedral losses, Theorem 1, is mostly nonconstructive: it says polyhedral surrogates have some α such that $R_\ell(\psi \circ h; \mathcal{D}) \leq \alpha \cdot R_L(h; \mathcal{D})$. In this section, we make the result constructive by unpacking the implications of consistency for polyhedral surrogates. Proofs appear in Appendix C.

We will give a bound on the constant of the form $\alpha \leq \alpha_\ell \cdot \alpha_L \cdot \alpha_\psi$, i.e. one component each relating to the structure of the target loss, surrogate loss, and link. First, we will derive α_L using the theory of *Hoffman constants* from linear optimization. Then, we will show that *any* link for a consistent polyhedral loss must satisfy a condition called ϵ-separation, introduced in [8]. We will set $\alpha_\psi = \frac{1}{\epsilon}$. With α_ℓ simply an upper bound on the target loss, we will finally prove $\alpha \leq \alpha_\ell \cdot \alpha_L \cdot \alpha_\psi$.

Hoffman constants. The intuition we explore here is the linear growth rate of a given polyhedral function as we move away from its optimal set. In particular, consider the expected loss $u \mapsto \langle p, L(u) \rangle$. Expected loss should grow linearly with distance from the optimal set, which happens to be $\Gamma(p)$. In particular, the worst growth rate is governed roughly by the smallest slope of the function in any direction. This is captured by the *Hoffman constant* of an associated system of linear inequalities. In Appendix C, we use Hoffman constants to obtain the following result, where we define $H_{L,p}$.

Lemma 5. *Let $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$ be a polyhedral loss with $\Gamma = \mathrm{prop}[L]$. Then for any fixed p, there exists some smallest constant $H_{L,p} \geq 0$ such that $d_\infty(u, \Gamma(p)) \leq H_{L,p} R_L(u, p)$ for all $u \in \mathbb{R}^d$.*

To foreshadow the usefulness of $H_{L,p}$, recall the proof strategy of the upper bound: use Lemma 1 to obtain a constant α_p for each p; then carefully choose a finite set of p's and take the maximum constant. In the same way, we will be able to set the "loss" constant $\alpha_L = \max_p H_{L,p}$. First, however, we will need to investigate separated links.

Separated link functions. A link ψ is ϵ-separated if, when $\psi(u^*)$ is an optimal target prediction and $\psi(u)$ is not, then $\|u^* - u\|_\infty > \epsilon$. The definition was introduced in [8] and is rephrased in terms of the properties $\Gamma = \mathrm{prop}[L]$ and $\gamma = \mathrm{prop}[\ell]$.

Definition 4 (Separated Link). *Let properties $\Gamma : \Delta_{\mathcal{Y}} \rightrightarrows \mathbb{R}^d$ and $\gamma : \Delta_{\mathcal{Y}} \rightrightarrows \mathcal{R}$ be given. We say a link $\psi : \mathbb{R}^d \to \mathcal{R}$ is ϵ-separated if for all $u \in \mathbb{R}^d$ with $\psi(u) \notin \gamma(p)$, we have $d_\infty(u, \Gamma(p)) > \epsilon$, where $d_\infty(u, A) \doteq \inf_{a \in A} \|u - a\|_\infty$.*

Recall from the previous subsection that Hoffman constants allowed us to show that polyhedral expected losses grow linearly as we move away from the optimal set. Now, ϵ-separated link functions allow us to guarantee that we move at least a constant distance from the optimal set before we start linking to an "incorrect" report $r' \in \mathcal{R}$. But when does a polyhedral loss have an associated ϵ-separated link?

[8] showed that in fact, if L indirectly elicits ℓ, then there always exists an ϵ-separated link ψ such that (L, ψ) is consistent for L. Here, we strengthen this result much further, and give a converse that we believe is of independent interest: if L indirectly elicits ℓ, then *every* link making L consistent is ϵ-separated for some $\epsilon > 0$.

Lemma 6. *Let polyhedral surrogate $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$, discrete loss $\ell : \mathcal{R} \to \mathbb{R}_+^{\mathcal{Y}}$, and link $\psi : \mathbb{R}^d \to \mathcal{R}$ be given such that (L, ψ) is calibrated with respect to ℓ. Then there exists $\epsilon > 0$ such that ψ is ϵ-separated with respect to $\Gamma \doteq \mathrm{prop}[L]$ and $\gamma \doteq \mathrm{prop}[\ell]$.*

Combining the loss and link. We can now follow the general proof strategy of the main upper bound, but constructively. Given Lemma 6, we can make the following definition.

318 **Definition 5** (ϵ_ψ)**.** If (L, ψ) are calibrated for ℓ, then let $\epsilon_\psi \doteq \sup\{\epsilon \ : \ \psi$ is ϵ-separated$\}$.

319 We will also need an upper bound on the regret of ℓ.

320 **Definition 6** (C_ℓ)**.** Given a discrete target loss $\ell : \mathcal{R} \to \mathbb{R}_+^{\mathcal{Y}}$, define $C_\ell = \max_{r,r' \in \mathcal{R}, y \in \mathcal{Y}} \ell(r)_y -$
321 $\ell(r')_y$.

322 Observe that $C_\ell = \max_{r,p} R_\ell(r, p)$.

323 For the proof, we first give a bound for any fixed p.

324 **Lemma 7.** *Let $\ell : \mathcal{R} \to \mathbb{R}_+^{\mathcal{Y}}$ be a discrete target loss, $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$ be a polyhedral surrogate loss,*
325 *and $\psi : \mathbb{R}^d \to \mathcal{R}$ a link function. If (L, ψ) indirectly elicit ℓ and ψ is ϵ-separated, then for all u and*
326 *p,*

$$R_\ell(\psi(u), p) \leq \frac{C_\ell H_{L,p}}{\epsilon} R_L(u, p).$$

327 Recall that Lemma 4 gave that, if L is polyhedral, then $\Gamma = \mathrm{prop}[L]$ has a finite set of full-dimensional
328 level sets, each a polytope, that union to the simplex.

329 **Definition 7** (H_L)**.** Given a polyhedral surrogate loss $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$, let $\mathcal{Q}$ be the set of all vertices
330 of the full-dimensional level sets of $\Gamma = \mathrm{prop}[L]$, and define $H_L \doteq \max_{p \in \mathcal{Q}} H_{L,p}$.

331 **Theorem 5** (Constructive linear transfer)**.** *Let $\ell : \mathcal{R} \to \mathbb{R}_+^{\mathcal{Y}}$ be a discrete target loss, $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$*
332 *be a polyhedral surrogate loss, and $\psi : \mathbb{R}^d \to \mathcal{R}$ a link function. If (L, ψ) are consistent for ℓ, then*

$$(\forall h, \mathcal{D}) \quad R_\ell(\psi \circ h; \mathcal{D}) \leq \frac{C_\ell H_L}{\epsilon_\psi} R_L(h; \mathcal{D}) \ .$$

333 The proof closely mirrors the proof of the nonconstructive upper bound, Theorem 1, but using the
334 constructive constants $C_\ell, H_L, \epsilon_\psi$ derived above. To summarize, we have shown the following.

335 **Theorem 6.** *Let $\ell : \mathcal{R} \to \mathbb{R}_+^{\mathcal{Y}}$ be a discrete target loss, $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$ be a polyhedral surrogate loss,*
336 *and $\psi : \mathbb{R}^d \to \mathcal{R}$ a link function. The following are equivalent:*

337   *1. (L, ψ) are consistent for ℓ.*

338   *2. (L, ψ) indirectly elicit ℓ and ψ is ϵ-separated for some $\epsilon > 0$.*

339   *3. (L, ψ) guarantee a linear regret transfer for ℓ.*

340 *Furthermore, if any of the above hold, then in particular (L, ψ) guarantee the regret transfer*
341 *$\zeta(t) = \left(\frac{C_\ell H_L}{\epsilon_\psi} \right) t$.*

5.1 Further tightening

343 While our goal is not to focus on exact constants for specific problems, we remark on a few ways
344 Theorem 5 may be loose and how one could compute the tightest possible constant α^* for which
345 $R_\ell(\psi \circ h; \mathcal{D}) \leq \alpha^* R_L(h; \mathcal{D})$ for all h and $\mathcal{D}$. In general, for a fixed p, there is some smallest α_p^* such
346 that $R_\ell(\psi(u), p) \leq \alpha_p^* R_L(u, p)$ for all u. Then, it follows from our results that $\alpha^* = \max_{p \in \mathcal{Q}} \alpha_p^*$ for
347 the finite set $\mathcal{Q}$ used in the proof, i.e. the vertices of the full-dimensional level sets of $\Gamma = \mathrm{prop}[L]$.

348 Above, we bounded $\alpha_p^* \leq \frac{C_\ell H_{L,p}}{\epsilon_\psi}$. The intuition is that some u at distance $\geq \epsilon_\psi$ from $\Gamma(p)$, the
349 optimal set, may link to a "bad" report $r = \psi(u) \notin \gamma(p)$. The rate at which L grows is at least
350 $H_{L,p}$, so the surrogate loss at u may be as small as $\frac{\epsilon_\psi}{H_{L,p}}$, while the target regret may be as high as
351 $C_\ell = \max_{r',p'} R_\ell(r', p')$. The ratio of regrets is therefore bounded by $\frac{H_{L,p} C_\ell}{\epsilon_\psi}$.

352 The tightest possible bound, on the other hand, is $\alpha^* = \sup_{u:\psi(u) \notin \gamma(p)} \frac{R_\ell(\psi(u),p)}{R_L(u,p)}$. This can be
353 smaller if the values of numerator and denominator are correlated across u. For example, u may only
354 be ϵ_ψ-close to the optimal set when it links to reports $\psi(u)$ with lower target regret; or L may have a
355 smaller slope in the direction where the link's separation is larger than ϵ.

To illustrate with a concrete example, consider the *binary encoded predictions (BEP) surrogate* of [18] for the abstain target loss, $\ell(r, y) = \frac{1}{2}$ if $r = \bot$, otherwise $\ell(r, y) = \mathbb{1}[r \neq y]$. The surrogate involves an injective map $B : \mathcal{Y} \to \{-1, 1\}^d$ for $d = \lceil \log_2 |\mathcal{Y}| \rceil$. It is $L(u)_y = \max_{j=1\ldots d}(1 - u_d B(y)_d)_+$, where $(\cdot)_+$ indicates taking the maximum of the argument and zero. The associated link is $\psi(u) = \bot$ if $\min_{j=1\ldots d} |u_j| \leq \frac{1}{2}$, otherwise $\psi(u) = \arg\min_{y \in \mathcal{Y}} \|B(y) - u\|_\infty$.

One can show that for $p = \delta_y$, i.e. the distribution with full support on some $y \in \mathcal{Y}$, $L(u)_y = d_\infty(u, \Gamma(p))$ exactly, giving $H_{L,p} = 1$. It is almost immediate that $\epsilon_\psi = \frac{1}{2}$. Meanwhile, $R_\ell(r, p) \leq 1$, giving us an upper bound $\alpha_p^* \leq \frac{(1)(1)}{1/2} = 2$. In fact, this is slightly loose; the exact constant, given in [18] is 1. The looseness stems from the fact that for $p = \delta_y$, the closest reports u to the optimal set, i.e. at distance only $\epsilon_\psi = \frac{1}{2}$ away, do not link to reports maximizing target regret; they link to the abstain report $\bot$, which has regret only $\frac{1}{2}$. With this correction, and an observation that all u linking to reports $y' \neq y$ are at distance at least $\frac{3}{2}$ from $\Gamma(p)$, we restore the tight bound $\alpha_p^* \leq 1$. A similar but slightly more involved calculation can be carried out for the other vertices $p \in \mathcal{Q}$, which turn out to be all vertices of the form $\frac{1}{2}\delta_y + \frac{1}{2}\delta_{y'}$.

Different norms. This paper uses L_∞ norm for the minimum-slope H_L and the ϵ-separated link. One reason is that it is easiest to prove the former by reduction to existence of Hoffman constants; another reason is compatibility of both with the abstain loss example above. However, all definitions hold for other L_p norms and so does the main upper bound, as existence of an H_L and ϵ_ψ in L_∞ imply existence of constants in L_p for $p \geq 1$. The optimal constant may change if the distance measure changes.

6 Broader Impacts

As a theoretical work, this papers' impacts are mostly in the form of downstream research and applications. Instead of direct applications, we anticipate this work leading to more investigation of surrogate losses to improve discrete prediction tasks. It may inform practitioners' choices of which surrogate losses they use for supervised learning tasks. Of course, such machine learning tasks can be solved for ethical or unethical purposes. We do not know of particular risk of negative impacts of this work beyond risks of supervised machine learning in general.

References

[1] Shivani Agarwal. Surrogate regret bounds for bipartite ranking via strongly proper losses. *The Journal of Machine Learning Research*, 15(1):1653–1674, 2014.

[2] Franz Aurenhammer. A criterion for the affine equivalence of cell complexes in R d and convex polyhedra in R d+1. *Discrete & Computational Geometry*, 2(1):49–64, December 1987. ISSN 0179-5376, 1432-0444. doi: 10.1007/BF02187870. URL http://link.springer.com/article/10.1007/BF02187870.

[3] Peter L Bartlett and Marten H Wegkamp. Classification with a reject option using a hinge loss. *Journal of Machine Learning Research*, 9(Aug):1823–1840, 2008.

[4] Peter L. Bartlett, Michael I. Jordan, and Jon D. McAuliffe. Convexity, classification, and risk bounds. *Journal of the American Statistical Association*, 101(473):138–156, 2006. URL http://amstat.tandfonline.com/doi/abs/10.1198/016214505000000907.

[5] Mathieu Blondel. Structured prediction with projection oracles. *Advances in neural information processing systems*, 32:12145–12156, 2019.

[6] Carlo Ciliberto, Lorenzo Rosasco, and Alessandro Rudi. A consistent regularization approach for structured prediction. *Advances in neural information processing systems*, 29:4412–4420, 2016.

[7] John Duchi, Khashayar Khosravi, Feng Ruan, et al. Multiclass classification, information, divergence and surrogate risk. *The Annals of Statistics*, 46(6B):3246–3275, 2018.

[8] Jessie Finocchiaro, Rafael Frongillo, and Bo Waggoner. An embedding framework for consistent polyhedral surrogates. In *Advances in neural information processing systems*, 2019.

[9] Rafael Frongillo and Ian Kash. General truthfulness characterizations via convex analysis. In *Web and Internet Economics*, pages 354–370. Springer, 2014.

[10] Maksim Lapin, Matthias Hein, and Bernt Schiele. Loss functions for top-k error: Analysis and insights. In *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition*, pages 1468–1477, 2016.

[11] Mehrdad Mahdavi, Lijun Zhang, and Rong Jin. Binary excess risk for smooth convex surrogates. *arXiv preprint arXiv:1402.1792*, 2014.

[12] Aditya Krishna Menon and Robert C Williamson. Bipartite ranking: a risk-theoretic perspective. *The Journal of Machine Learning Research*, 17(1):6766–6867, 2016.

[13] Alex Nowak-Vila, Francis Bach, and Alessandro Rudi. A general theory for structured prediction with smooth convex surrogates. *arXiv preprint arXiv:1902.01958*, 2019.

[14] Anton Osokin, Francis Bach, and Simon Lacoste-Julien. On structured prediction theory with calibrated convex surrogate losses. In *Advances in Neural Information Processing Systems*, pages 302–313, 2017.

[15] Fabian Pedregosa, Francis Bach, and Alexandre Gramfort. On the consistency of ordinal regression methods. *Journal of Machine Learning Research*, 18:1–35, 2017.

[16] Bernardo Avila Pires, Csaba Szepesvari, and Mohammad Ghavamzadeh. Cost-sensitive multiclass classification risk bounds. In *International Conference on Machine Learning*, pages 1391–1399, 2013.

[17] Harish Ramaswamy, Ambuj Tewari, and Shivani Agarwal. Convex calibrated surrogates for hierarchical classification. In *International Conference on Machine Learning*, pages 1852–1860, 2015.

[18] Harish G Ramaswamy, Ambuj Tewari, Shivani Agarwal, et al. Consistent algorithms for multiclass classification with an abstain option. *Electronic Journal of Statistics*, 12(1):530–554, 2018.

[19] Mark D Reid and Robert C Williamson. Surrogate regret bounds for proper losses. In *Proceedings of the 26th Annual International Conference on Machine Learning*, pages 897–904, 2009.

[20] R.T. Rockafellar. *Convex analysis*, volume 28 of *Princeton Mathematics Series*. Princeton University Press, 1997.

[21] Clayton Scott et al. Calibrated asymmetric surrogate losses. *Electronic Journal of Statistics*, 6: 958–992, 2012.

[22] Ingo Steinwart and Andreas Christmann. *Support Vector Machines*. Springer Science & Business Media, September 2008. ISBN 978-0-387-77242-4. Google-Books-ID: HUnqnrpYt4IC.

[23] Ambuj Tewari and Peter L. Bartlett. On the consistency of multiclass classification methods. *The Journal of Machine Learning Research*, 8:1007–1025, 2007. URL http://dl.acm.org/citation.cfm?id=1390325.

[24] Yutong Wang and Clayton Scott. Weston-watkins hinge loss and ordered partitions. *Advances in neural information processing systems*, 2020.

[25] Forest Yang and Sanmi Koyejo. On the consistency of top-k surrogate losses. *CoRR*, abs/1901.11141, 2019. URL http://arxiv.org/abs/1901.11141.

[26] Jiaqian Yu and Matthew B Blaschko. The lovász hinge: A novel convex surrogate for submodular losses. *IEEE transactions on pattern analysis and machine intelligence*, 2018.

[27] Ming Yuan and Marten Wegkamp. Classification methods with reject option based on convex risk minimization. *Journal of Machine Learning Research*, 11(Jan):111–130, 2010.

[28] Tong Zhang et al. Statistical behavior and consistency of classification methods based on convex risk minimization. *The Annals of Statistics*, 32(1):56–85, 2004.

A Omitted Proofs - Upper Bound

This section contains omitted proofs from Section 3.

Lemma (Lemma 1). *Let $\ell : \mathcal{R} \to \mathbb{R}_+^{\mathcal{Y}}$ be a discrete target loss and suppose the surrogate $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$ and link $\psi : \mathbb{R}^d \to \mathcal{R}$ are calibrated for ℓ. Then for any $p \in \Delta_{\mathcal{Y}}$, there exists $\alpha_p \geq 0$ such that, for all $u \in \mathbb{R}^d$,*

$$R_\ell(\psi(u), p) \leq \alpha_p R_L(u, p).$$

Proof. Fix $p \in \Delta_{\mathcal{Y}}$. Let $C_p = \max_{r \in \mathcal{R}} R_\ell(r, p)$. The maximum exists because ℓ is discrete, i.e. $\mathcal{R}$ is finite. Meanwhile, recall that, when defining calibration, we let $B_{L,\psi,\ell}(p) = \{R_L(u, p) : \psi(u) \notin \gamma(p)\}$. Let $B_p = \inf B_{L,\psi,\ell}(p)$. By definition of calibration, we have $B_p > 0$.

To combine these bounds, let $\alpha_p = \frac{C_p}{B_p}$. Let $u \in \mathbb{R}^d$. There are two cases. If $\psi(u) \in \gamma(p)$, then $R_\ell(\psi(u), p) = 0 \leq R_L(u, p)$ immediately. If $\psi(u) \notin \gamma(p)$, then

$$\begin{aligned} R_\ell(\psi(u), p) &\leq C_p \\ &= \alpha_p \cdot B_p \\ &\leq \alpha_p R_L(u, p). \end{aligned}$$

$\square$

Lemma (Lemma 2). *If (L, ψ) indirectly elicits ℓ, then $\Gamma = \text{prop}[L]$ refines $\gamma = \text{prop}[\ell]$ in the sense that, for all $u \in \mathbb{R}^d$, there exists $r \in \mathcal{R}$ such that $\Gamma_u \subseteq \gamma_r$.*

Proof. For any u, let $r = \psi(u)$. By indirect elicitation, $u \in \Gamma(p) \implies r \in \gamma(p)$. So $\Gamma_u = \{p : u \in \Gamma(p)\} \subseteq \{p : r \in \gamma(p)\} = \gamma_r$. $\qquad\square$

Lemma (Lemma 3). *Suppose (L, ψ) indirectly elicits ℓ and let $\Gamma = \text{prop}[L]$. Then for any fixed $u, u^* \in \mathbb{R}^d$ and $r \in \mathcal{R}$, the functions $R_L(u, \cdot)$ and $R_\ell(r, \cdot)$ are linear in their second arguments on Γ_{u^*}.*

Proof. Let $u^* \in \mathbb{R}^d$ and $p \in \Gamma_{u^*}$. By definition, for all $p \in \Gamma_{u^*}$, $\underline{L}(p) = \langle p, L(u^*)\rangle$. So for fixed u,

$$R_L(u, p) = \langle p, L(u)\rangle - \langle p, L(u^*)\rangle = \langle p, L(u) - L(u^*)\rangle,$$

a linear function of p on Γ_{u^*}. Next, by Lemma 2, there exists r^* such that $\Gamma_{u^*} \subseteq \gamma_{r^*}$. By the same argument, for fixed r, $R_\ell(r, p) = \langle p, \ell(r) - \ell(r^*)\rangle$, a linear function of p on γ_{r^*} and thus on Γ_{u^*}. $\quad\square$

Lemma (Lemma 4). *If $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$ is polyhedral and minimizable, then $\Gamma = \text{prop}[L]$ has a finite set of level sets that union to $\Delta_{\mathcal{Y}}$. Also, these level sets are polytopes.*

Proof. This can be deduced from the embedding framework of [8]. In particular, Lemma 5 of [8] states that if L is polyhedral, then its Bayes risk $\underline{L}$ is concave polyhedral, i.e. is the pointwise minimum of a finite set of affine functions. It follows that there exists a finite set $U \subset \mathbb{R}^d$ such that

$$\underline{L}(p) = \min_{u \in \mathbb{R}^d} \langle p, L(u)\rangle = \min_{u \in U} \langle p, L(u)\rangle . \tag{4}$$

We claim the level sets of U witness the claim. First, it is known (e.g. from theory of power diagrams, [2]) that if $\underline{L}$ is a polyhedral function represented as (4) and $u \in U$, then $\Gamma_u = \{p \in \Delta_{\mathcal{Y}} : \langle p, L(u)\rangle = \underline{L}(p)\}$ is a polytope. Finally, suppose for contradiction that there exists $p \in \Delta_{\mathcal{Y}}$, $p \notin \cup_{u \in U}\Gamma_u$. Then there must be some $u' \notin U$ with $p \in \Gamma_{u'}$, implying that $\langle p, L(u')\rangle > \max_{u \in U}\langle p, L(u)\rangle$, contradicting (4). $\qquad\square$

Theorem (Theorem 3). *Suppose the surrogate loss $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$ and link $\psi : \mathbb{R}^d \to \mathcal{R}$ are consistent for the target loss $\ell : \mathcal{R} \to \mathbb{R}_+^{\mathcal{Y}}$. If L is polyhedral, then (L, ψ) guarantee a linear regret transfer for ℓ, i.e. there exists $\alpha \geq 0$ such that, for all $\mathcal{D}$ and all measurable $h : \mathcal{X} \to \mathbb{R}^d$,*

$$R_\ell(\psi \circ h; \mathcal{D}) \leq \alpha R_L(h; \mathcal{D}).$$

Proof. We first recall that by Fact 1, consistency implies that (L, ψ) are calibrated for ℓ and that (L, ψ) indirectly elicit ℓ. Next, by Observation 1, it suffices to show a linear *conditional* regret transfer, i.e. for all $p \in \Delta_{\mathcal{Y}}$ and $u \in \mathbb{R}^d$, we show $R_\ell(\psi(u), p) \leq \alpha R_L(u, p)$.

By Lemma 4, the polyhedral loss L has a finite set $U \subset \mathbb{R}^d$ of predictions such that (a) for each $u \in U$, the level set Γ_u is a polytope, and (b) $\cup_{u \in U}\Gamma_u = \Delta_{\mathcal{Y}}$. Let $\mathcal{Q}_u \subset \Delta_{\mathcal{Y}}$ be the finite set of vertices of the polytope Γ_u, and define the finite set $\mathcal{Q} = \cup_{u \in U}\mathcal{Q}_u$.

By Lemma 1, for each $q \in \mathcal{Q}$, there exists $\alpha_q \geq 0$ such that $R_\ell(\psi(u), q) \leq \alpha_q R_L(u, q)$ for all u. We choose

$$\alpha = \max_{q \in \mathcal{Q}} \alpha_q.$$

To prove the conditional regret transfer, consider any $p \in \Delta_{\mathcal{Y}}$ and any $u \in \mathbb{R}^d$. There exists $u \in U$ such that $p \in \Gamma_u$, a polytope. So we can write p as a convex combination of its vertices, i.e.

$$p = \sum_{q \in \mathcal{Q}_u} \beta(q)q$$

for some probability distribution β. Recall that $\mathcal{Q}_u \subseteq \Gamma_u$ and R_L and R_ℓ are linear in p on Γ_u by Lemma 3. So, for any u':

$$
\begin{aligned}
R_\ell(\psi(u'), p) &= R_\ell\left(\psi(u'), \sum_{q \in \mathcal{Q}_u} \beta(q)q\right) \\
&= \sum_{q \in \mathcal{Q}_u} \beta(q)R_\ell(\psi(u'), q) \\
&\leq \sum_{q \in \mathcal{Q}_u} \beta(q)\alpha_q R_L(u', q) \\
&\leq \alpha \sum_{q \in \mathcal{Q}_u} \beta(q)R_L(u', q) \\
&= \alpha R_L(u', p).
\end{aligned}
$$

$\square$

B   Omitted Proofs - Lower Bound

This section contains omitted proofs from Section 4.

Theorem (Theorem 4). *Suppose the surrogate loss L and link ψ satisfy a regret transfer of ζ for a target loss ℓ. If L, ψ, and ℓ satisfy Assumption 1, then there exists $c > 0$ such that, for some $\epsilon^* > 0$, for all $0 \leq \epsilon < \epsilon^*$, $\zeta(\epsilon) \geq c\sqrt{\epsilon}$.*

Proof outline: By assumption we have a boundary point u_0 which is L-optimal for a distribution p_0. We have some r, r' which are both optimal for p_0, and $\psi(u_0) = r'$. First, we will choose a p_1 where r is uniquely optimal, hence u_0 is a strictly suboptimal choice. We then consider a sequence of distributions $p_\lambda = (1 - \lambda)p_0 + \lambda p_1$, approaching p_0 as $\lambda \to 0$. For all such p_λ, it will happen that r is optimal while u_0 and $r' = \psi(u_0)$ are strictly suboptimal. We show that $R_\ell(r', p_\lambda) = c_\ell \lambda$ for some constant c_ℓ and all small enough λ. Meanwhile, we will show that $R_L(u_0, p_\lambda) \leq O(\lambda^2)$, proving the result. The last fact will use the properties of strong smoothness and strong convexity in a neighborhood of u_0.

Proof. Obtain α, u_0, p_0, r, r', and an open neighborhood of u_0 from Assumption 1 and the definition of boundary point. Assume without loss of generality that $\psi(u_0) = r'$; otherwise, swap the roles of r and r'.

Linearity of $R_\ell(r', p_\lambda)$.   As ℓ is non-redundant by assumption, there exists some $p_1 \in \mathring{\gamma}_r$, the relative interior of the full-dimensional level set γ_r. We therefore have $R_\ell(r', p_1) = \langle p_1, \ell(r') - \ell(r)\rangle =: c_\ell > 0$, and $R_\ell(r', p_0) = 0$. Let $p_\lambda := (1 - \lambda)p_0 + \lambda p_1$. By convexity of γ_r, we have $p_\lambda \in \gamma_r$ for all $\lambda \in [0, 1]$, which gives $R_\ell(r', p_\lambda) = \lambda c_\ell$.

Obtaining the global minimizer u_λ of L_λ.   Let $L_\lambda : \mathbb{R}^d \to \mathbb{R}_+$ be given by $L_\lambda(u) = \langle p_\lambda, L(u)\rangle = (1 - \lambda)\langle p_0, L(u)\rangle + \lambda\langle p_1, L(u)\rangle$. Let $\delta > 0$ such that the above open neighborhood of u_0 contains the Euclidean ball $B_\delta(u_0)$ of radius δ around u_0. Let $u_1 \in \Gamma(p_1)$. We argue that for all small enough λ, $L_\lambda(u)$ is uniquely minimized by some $u_\lambda \in B_\delta(u_0)$. For any $u \notin B_\delta(u_0)$, we have, using local strong convexity and the optimality of u_1,

$$
\begin{aligned}
L_\lambda(u) - L_\lambda(u_0) &= (1 - \lambda)\left(L_0(u) - L_0(u_0)\right) + \lambda\left(L_1(u) - L_1(u_0)\right) \\
&\geq (1 - \lambda)\left(\frac{\alpha}{2}\delta^2\right) + \lambda\left(L_1(u_1) - L_1(u_0)\right) \\
&> 0
\end{aligned}
$$

if $\lambda < \lambda^* := \alpha\delta^2/(2\alpha\delta^2 + 4L_1(u_0) - 4L_1(u_1))$. For the remainder of the proof, let $\lambda < \lambda^*$. Then any $u \notin B_\delta(u_0)$ has $L_\lambda(u) > L_\lambda(u_0)$, hence is suboptimal. By α-strong convexity of L_0 on $B_\delta(u_0)$, L_λ is strictly convex on $B_\delta(u_0)$. So it has a unique minimizer u_λ, and by the above argument this is the global minimizer of L_λ. Then $\underline{L}(p_\lambda) = L_\lambda(u_\lambda)$, and thus $R_L(u_0, p_\lambda) = L_\lambda(u_0) - L_\lambda(u_\lambda)$. We also observe here that $R_L(u_0, p_\lambda)$ is continuous in λ, e.g. because the Bayes risk of L is continuous in p as is $\langle p, L(u_0)\rangle$. It is also zero when $\lambda = 0$.

Showing R_L is quadratic in λ. By assumption, the gradient of L_y is locally Lipschitz for all $y \in \mathcal{Y}$. We will apply this fact to the compact set $\mathcal{C} = \{u \in \mathbb{R}^d : \|u - u_1\| \leq \|u_0 - u_1\| + \delta\}$. By compactness, we have a finite subcover of open neighborhoods; let β be the minimum Lipschitz constant over this finite set of neighborhoods. We thus have that L_y is β-strongly smooth on $\mathcal{C}$, and hence so is L_λ for any $\lambda \in [0, 1]$.

We now upper bound $\|u_\lambda - u_0\|_2$, and then apply strong smoothness to upper bound $R_L(u_0, p_\lambda) = L_\lambda(u_0) - L_\lambda(u_\lambda)$. Consider the first-order optimality condition of L_λ:

$$0 = \nabla L_\lambda(u_\lambda) = (1 - \lambda)\nabla L_0(u_\lambda) + \lambda \nabla L_1(u_\lambda)$$
$$\implies (1 - \lambda)\|\nabla L_0(u_\lambda)\|_2 = \lambda\|\nabla L_1(u_\lambda)\|_2 .$$

By optimality of u_0 and u_1, strong convexity of L_0 and strong smoothness of L_1, and the triangle inequality, we have

$$\|\nabla L_0(u_\lambda)\|_2 = \|\nabla L_0(u_\lambda) - \nabla L_0(u_0)\|_2 \geq \alpha\|u_\lambda - u_0\|_2 ,$$
$$\|\nabla L_1(u_\lambda)\|_2 = \|\nabla L_1(u_\lambda) - \nabla L_1(u_1)\|_2 \leq \beta\|u_\lambda - u_1\|_2$$
$$\leq \beta\left(\|u_\lambda - u_0\|_2 + \|u_0 - u_1\|_2\right) .$$

Combining,

$$(1 - \lambda)\alpha\|u_\lambda - u_0\|_2 \leq (1 - \lambda)\|\nabla L_0(u_\lambda)\|_2$$
$$= \lambda\|\nabla L_1(u_\lambda)\|_2$$
$$\leq \lambda\beta\left(\|u_\lambda - u_0\|_2 + \|u_0 - u_1\|_2\right) .$$

Now rearranging and taking $\lambda \leq \frac{1}{2}\frac{\alpha}{\alpha+\beta}$, we have

$$\|u_\lambda - u_0\|_2 \leq \frac{\lambda\beta}{(1 - \lambda)\alpha - \lambda\beta}\|u_0 - u_1\|_2 \leq \lambda\frac{2\beta}{\alpha}\|u_0 - u_1\|_2 .$$

Finally, from strong smoothness of L_λ and optimality of u_λ,

$$L_\lambda(u_0) - L_\lambda(u_\lambda) \leq \frac{\beta}{2}\|u_0 - u_\lambda\|_2^2 \leq \frac{\beta}{2}\left(\lambda\frac{2\beta}{\alpha}\|u_0 - u_1\|_2\right)^2 = c_L\lambda^2 ,$$

where $c_L = \frac{2\beta^3}{\alpha^2}\|u_0 - u_1\|_2^2 > 0$.

To conclude: we have found a $\lambda^* > 0$ and shown that for all $0 \leq \lambda < \lambda^*$, $R_\ell(r', p_\lambda) = c_\ell\lambda$ and $R_L(u_0, p_\lambda) \leq c_L\lambda^2$. In particular, let $\epsilon^* = \sup_{0 \leq \lambda < \lambda^*} R_L(u_0, p_\lambda)$. Then for all $0 \leq \epsilon < \epsilon^*$, by continuity, we can choose $\lambda < \lambda^*$ such that $R_L(u_0, p_\lambda) = \epsilon \leq c_L\lambda^2$. Meanwhile, $R_\ell(\psi(u_0), p_\lambda) = c_\ell\lambda \geq \frac{c_\ell}{\sqrt{c_L}}\sqrt{\epsilon}$. Recalling that $\zeta(R_L(u_0, p_\lambda)) \geq R_\ell(\psi(u_0), p_\lambda)$ by definition, this implies $\zeta(\epsilon) \geq c\sqrt{\epsilon}$ for all $\epsilon < \epsilon^*$, with $c = \frac{c_\ell}{\sqrt{c_L}}$. $\qquad\square$

C Omitted Proofs - Constant Derivation

This section contains omitted proofs from Section 5.

Hoffman constants. First we appeal to a known fact, the existence of Hoffman constants for systems of linear inequalities.

Theorem 7 (Hoffman constant,). *Given a matrix $A \in \mathbb{R}^{m \times n}$, there exists some smallest $H(A) \geq 0$, called the* Hoffman constant *(with respect to $\|\cdot\|_\infty$), such that for all $b \in \mathbb{R}^m$ and all $x \in \mathbb{R}^n$,*

$$d_\infty(x, S(A, b)) \leq H(A)\|(Ax - b)_+\|_\infty , \tag{5}$$

where $S(A, b) = \{x \in \mathbb{R}^n \mid Ax \leq b\}$ and $(u)_+ \doteq \max(u, 0)$ component-wise.

Lemma (Lemma 5). *Let $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$ be a polyhedral loss with $\Gamma = \mathrm{prop}[L]$. Then for any fixed p, there exists some smallest constant $H_{L,p} \geq 0$ such that $d_\infty(u, \Gamma(p)) \leq H_{L,p}R_L(u, p)$ for all $u \in \mathbb{R}^d$.*

Proof. Since L is polyhedral, there exist $a_1, \ldots, a_m \in \mathbb{R}^d$ and $c \in \mathbb{R}^m$ such that we may write $\langle p, L(u) \rangle = \max_{1 \leq j \leq m} a_j \cdot u + c_j$. Let $A \in \mathbb{R}^{m \times d}$ be the matrix with rows a_j, and let $b = \underline{L}(p)\mathbb{1} - c$, where $\mathbb{1} \in \mathbb{R}^m$ is the all-ones vector. Then we have

$$
\begin{aligned}
S(A, b) &\doteq \{u \in \mathbb{R}^d \mid Au \leq b\} \\
&= \{u \in \mathbb{R}^d \mid Au + c \leq \underline{L}(p)\mathbb{1}\} \\
&= \{u \in \mathbb{R}^d \mid \forall i \, (Au + c)_i \leq \underline{L}(p)\} \\
&= \{u \in \mathbb{R}^d \mid \max_i \, (Au + c)_i \leq \underline{L}(p)\} \\
&= \{u \in \mathbb{R}^d \mid \langle p, L(u) \rangle \leq \underline{L}(p)\} \\
&= \Gamma(p) \, .
\end{aligned}
$$

Similarly, we have $\max_i \, (Au - b)_i = \langle p, L(u) \rangle - \underline{L}(p) = R_L(u, p) \geq 0$. Thus,

$$
\begin{aligned}
\|(Au - b)_+\|_\infty &= \max_i \, ((Au - b)_+)_i \\
&= \max((Au - b)_1, \ldots, (Au - b)_m, 0) \\
&= \max(\max_i \, (Au - b)_i, \, 0) \\
&= \max_i \, (Au - b)_i \\
&= R_L(u, p) \, .
\end{aligned}
$$

Now applying Theorem 7, we have

$$
\begin{aligned}
d_\infty(u, \Gamma(p)) &= d_\infty(u, S(A, b)) \\
&\leq H(A)\|(Au - b)_+\|_\infty \\
&= H(A)R_L(u, p) \, .
\end{aligned}
$$

$\square$

Separated links.

Lemma (Lemma 6). *Let polyhedral surrogate $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$, discrete loss $\ell : \mathcal{R} \to \mathbb{R}_+^{\mathcal{Y}}$, and link $\psi : \mathbb{R}^d \to \mathcal{R}$ be given such that (L, ψ) is calibrated with respect to ℓ. Then there exists $\epsilon > 0$ such that ψ is ϵ-separated with respect to $\Gamma \doteq \mathrm{prop}[L]$ and $\gamma \doteq \mathrm{prop}[\ell]$.*

Proof. Suppose that ψ is not ϵ-separated for any $\epsilon > 0$. Then letting $\epsilon_i \doteq 1/i$ we have sequences $\{p_i\}_i \subset \Delta_{\mathcal{Y}}$ and $\{u_i\}_i \subset \mathbb{R}^d$ such that for all $i \in \mathbb{N}$ we have both $\psi(u_i) \notin \gamma(p_i)$ and $d_\infty(u_i, \Gamma(p_i)) \leq \epsilon_i$. First, observe that there are only finitely many values for $\gamma(p_i)$ and $\Gamma(p_i)$, as $\mathcal{R}$ is finite and L is polyhedral. Thus, there must be some $p \in \Delta_{\mathcal{Y}}$ and some infinite subsequence indexed by $j \in J \subseteq \mathbb{N}$ where for all $j \in J$, we have $\psi(u_j) \notin \gamma(p)$ and $\Gamma(p_j) = \Gamma(p)$.

Next, observe that, as L is polyhedral, the expected loss $\langle p, L(u) \rangle$ is β-Lipschitz in $\|\cdot\|_\infty$ for some $\beta > 0$. Thus, for all $j \in J$, we have

$$
\begin{aligned}
d_\infty(u_i, \Gamma(p)) \leq \epsilon_j &\implies \exists u^* \in \Gamma(p) \|u_j - u^*\|_\infty \leq \epsilon_j \\
&\implies |\langle p, L(u_j) \rangle - \langle p, L(u^*) \rangle| \leq \beta\epsilon_j \\
&\implies |\langle p, L(u_j) \rangle - \underline{L}(p)| \leq \beta\epsilon_j \, .
\end{aligned}
$$

Finally, for this p, we have

$$
\inf_{u : \psi(u) \notin \gamma(p)} \langle p, L(u) \rangle \leq \inf_{j \in J} \langle p, L(u_j) \rangle = \underline{L}(p) \, ,
$$

contradicting the calibration of ψ.

$\square$

Combining the loss and link.

Lemma (Lemma 7). *Let $\ell : \mathcal{R} \to \mathbb{R}_+^{\mathcal{Y}}$ be a discrete target loss, $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$ be a polyhedral surrogate loss, and $\psi : \mathbb{R}^d \to \mathcal{R}$ a link function. If (L, ψ) indirectly elicit ℓ and ψ is ϵ-separated, then for all u and p,*

$$
R_\ell(\psi(u), p) \leq \frac{C_\ell H_{L,p}}{\epsilon} R_L(u, p).
$$

616 *Proof.* If $\psi(u) \in \gamma(p)$, then $R_\ell(u, p) = 0$ and we are done. Otherwise, applying the definition of
617 ϵ-separated and Lemma 5,

$$\epsilon < d_\infty(u, \Gamma(p))$$
$$\leq H_{L,p} R_L(u, p).$$

618 So $R_\ell(\psi(u), p) \leq C_\ell \leq \frac{C_\ell H_{L,p}}{\epsilon} R_L(u, p)$. $\qquad\square$

619 **Theorem** (Constructive linear transfer, Theorem 5). *Let $\ell : \mathcal{R} \to \mathbb{R}_+^{\mathcal{Y}}$ be a discrete target loss,*
620 $L : \mathbb{R}^d \to \mathbb{R}_+^{\mathcal{Y}}$ *be a polyhedral surrogate loss, and $\psi : \mathbb{R}^d \to \mathcal{R}$ a link function. If (L, ψ) are*
621 *consistent for ℓ, then*

$$(\forall h, \mathcal{D}) \quad R_\ell(\psi \circ h; \mathcal{D}) \leq \frac{C_\ell H_L}{\epsilon_\psi} R_L(h; \mathcal{D}) .$$

622 The proof closely mirrors the proof of the nonconstructive upper bound, Theorem 1.

623 *Proof.* By Lemma 6, ψ is separated and ϵ_ψ well-defined. By Lemma 7, for each $p \in \mathcal{Q}$,
624 $R_\ell(\psi(u), p) \leq \frac{C_\ell H_L}{\epsilon_\psi} R_L(u, p)$ for all u. Now consider a general p, which is in some full-dimensional
625 polytope level set Γ_u. Write $p = \sum_{q \in \mathcal{Q}_u} \beta(q)q$ for some probability distribution β, where $\mathcal{Q}_u$ is the
626 set of vertices of Γ_u. By Lemma 3, R_L and R_ℓ are linear in p on Γ_u, so for any u',

$$R_\ell(\psi(u'), p) = \sum_{q \in \mathcal{Q}_u} \beta(q) R_\ell(\psi(u'), q)$$
$$\leq \sum_{q \in \mathcal{Q}_u} \beta(q) \frac{C_\ell H_{L,p}}{\epsilon_\psi} R_L(u', q)$$
$$\leq \frac{C_\ell H_L}{\epsilon_\psi} \sum_{q \in \mathcal{Q}_u} \beta(q) R_L(u', q)$$
$$\leq \frac{C_\ell H_L}{\epsilon_\psi} R_L(u', p).$$

627 This conditional regret transfer implies a full regret transfer with the same constant by Observation
628 1. $\qquad\square$