A Note on Sparse Generalized Eigenvalue Problem

Abstract

The sparse generalized eigenvalue problem (SGEP) aims to find the leading eigen-vector with sparsity structure. SGEP plays an important role in statistical learning and has wide applications including, but not limited to, sparse principal compo-nent analysis, sparse canonical correlation analysis and sparse Fisher discriminant analysis, etc. Due to the sparsity constraint, the solution of SGEP entails inter-esting properties from both numerical and statistical perspectives. In this paper, we provide a detailed sensitivity analysis for SGEP and establish the rate-optimal perturbation bound under the sparse setting. Specifically, we show that the bound is related to the perturbation/noise level and the recovery of the true support of the leading eigenvector as well. We also investigate the estimator of SGEP via imposing a non-convex regularization. Such estimator can achieve the optimal error rate and can recover the sparsity structure as well. Extensive numerical experiments corroborate our theoretical findings via using alternating direction method of multipliers (ADMM)-based computational method.

1 Introduction

The sparse generalized eigenvalue problem (SGEP) is to solve the following constrained optimization problem:

$$\max_{\mathbf{x} \in \mathbb{R}^p} \frac{\mathbf{x}^{\mathrm{T}} \widetilde{A} \mathbf{x}}{\mathbf{x}^{\mathrm{T}} \widetilde{B} \mathbf{x}}, \quad \text{subject to} \quad \|\mathbf{x}\|_0 \leq k, \tag{1}$$

where $\widetilde{A}$, $\widetilde{B}$ are both p-by-p symmetric matrices and $\widetilde{B}$ is (semi) positive definite; $\|\mathbf{x}\|_0$ denotes the ℓ_0-norm of $\mathbf{x}$, i.e., the number of nonzero entries of x; k is an integer which may be much smaller than p. For a fixed k, we let $\mathbf{x}_*$ be the solution to (1). Moreover, matrices $\widetilde{A}$ and $\widetilde{B}$ usually have the following decomposition:

$$\widetilde{A} = A + E, \qquad \widetilde{B} = B + F, \tag{2}$$

where A, B are underlying symmetric matrices and B is positive definite. However, A and B are unobserved in practice and they are contaminated by some noise/perturbation matrices E and F, respectively. Let λ_1 be the largest generalized eigenvalue of matrix pair (A, B) and $\mathbf{u}_1$ be its corresponding eigenvector which is also known as the leading eigenvector. In addition, $\mathbf{u}_1$ is always assumed to be sparse, i.e., $\|\mathbf{u}_1\|_0 \ll p$. Therefore, the SGEP is essentially to find an approximation of $\mathbf{u}_1$ based on observation $(\widetilde{A}, \widetilde{B})$.

The SGEP is challenging due to the following aspects. First, the sparsity constraint optimization problem (1) is NP-hard, since it is essentially a subset selection problem [15, 16]. To recover the underlying sparsity structure, we need to try different k which is computationally expensive. Second, due to insufficient number of samples, $\widetilde{A}$ and $\widetilde{B}$ can be both ill-conditioned or even share an approximate common null space, and perturbations E and F can be large. As a result, the leading eigenvector of pair $(\widetilde{A}, \widetilde{B})$ may not be a good approximation of $\mathbf{u}_1$.

Though challenging, especially when there is a limited number of data samples, SGEP and related problems have received more and more attentions in the past decade and various numerical methods are proposed to solve the SGEP. For example, in 20, The SGEP is framed as a d.c. (difference of convex functions) program and is solved as a sequence of convex programs by invoking the majorization-minimization method. In 19, the ℓ_0 norm constraint is approximated by a continuous surrogate function, an algorithm inspired by the majorization-minimization method is developed via iteratively majorizing the surrogate function by a quadratic separable function, and a systematic way based on smoothing is proposed to deal with the singularity issue. In 26, a two-stage computational framework is proposed to solve the SGEP, in which the first stage computes an initial guess via convex relaxation, and the second stage solves a nonconvex optimization via a fixed step size steepest ascent method, and followed by the simple truncation. In 18, a general framework called sparse estimation with linear programming is proposed for sparse canonical correlation analysis. Combining the idea of ℓ_1-penalized adaptive normalized quasi-Newton algorithm with nested orthogonal complement structure, an algorithm is proposed in 28 to solve the SGEP. 9 solves the SGEP by modifying the standard generalized orthogonal iteration with a sparsity-inducing penalty for the eigenvectors. 6 proposes a two-stage estimation method for sparse canonical correlation analysis via using semidefinite programming method.

Despite the efforts mentioned above, there is very little work investigating the optimal estimation error and no work provides theoretical guarantees whether the underlying sparsity structure of the leading eigenvector of SGEP can be recovered. In this paper, we build the theoretical gap in the literature. First, we provide both upper and lower bounds of the approximation error given noisy observations in the sparse generalized eigenvalue problem setting. To be specific, we prove that the difference between $\mathbf{u}_1$ and $\mathbf{x}_*$ is bounded by two parts, the perturbation error and error of not finding the true support of $\mathbf{u}_1$. Furthermore, we show that the error bound we obtained is rate optimal. That is, the lower bound matches the upper bound up to a multiplicative constant. Second, we consider a family of non-convex penalty functions and reformulate (1) into a regularization problem. We show that such regularized estimator enjoys the merits of sparsity and nearly unbiasedness. With these three nice properties, the estimator can be shown to achieve the optimal estimation error rate and can recover the true support of leading eigenvector as well. In addition, we also present a new computational algorithm. The estimation procedure adopts the Alternating Direction Method of Multipliers (ADMM, 33, 17). Our method can recover the sparsity structure under mild conditions.

The rest of paper is organized as follows. In Section 2, we introduce the sparse generalized eigenvalue problem with its applications and technical tools. In Section 3, we provide a sensitivity analysis for SGEP and establish its upper and lower perturbation bounds. In Section 4, we propose a regularized estimator via using non-convex penalization method. We also propose a computational algorithm (NC-SGEP) to tackle with the estimation issues. Multiple numerical results are given in Section 5. The concluding remark is given in Section 6.

Notations: The calligraphic letters $\mathcal{I}$, $\mathcal{J}$, $\mathcal{K}$, $\mathcal{L}$ and $\mathcal{S}$ are usually used to denote index sets. $|\mathcal{I}|$ denotes the cardinality of $\mathcal{I}$, e.g., $\mathcal{I} = \{i_1, i_2, \ldots, i_s\}$, where $i_1, i_2, \ldots, i_s$ are distinct integers, then $|\mathcal{I}| = s$. We use $\mathbf{x}$ to denote a generic vector in $\mathbb{R}^p$ and $\mathbf{x}^T$ to denote its transpose. $\mathbf{x}[j]$ is the jth element in $\mathbf{x}$. $\mathbf{x}[\mathcal{I}]$ stands for the subvector of $\mathbf{x}$ with indices in set $\mathcal{I}$. We use A to denote a matrix in $\mathbb{R}^{p \times p}$. $A[i, j]$ is the entry in the ith row and the jth column. $A[\mathcal{I}, \mathcal{J}]$ stands for the submatrix of A with row indices in set $\mathcal{I}$ and column indices in set $\mathcal{J}$. If $\mathcal{I} = \mathcal{J}$, $A[\mathcal{I}, \mathcal{I}]$ is also denoted by $A_{\mathcal{I}}$. We also let $(\mathcal{I}, \mathcal{J})$ be the union of sets $\mathcal{I}$ and $\mathcal{J}$. For $\mathbf{x} \in \mathbb{R}^p$, $\mathrm{supp}(\mathbf{x})$ denotes the index set of all nonzero entries of $\mathbf{x}$. For matrix A, $\mathrm{supp}(A)$ denotes the index set of all nonzero rows of A. I_p is the $p \times p$ identity matrix. $\|\mathbf{x}\|_0$ represents the number of non-zero entries in $\mathbf{x}$; $\|\mathbf{x}\|_2 := \sqrt{\sum_j \mathbf{x}^2[j]}$; $\|A\|_2$ denotes the largest singular value of A; $\|A\|_F$ and $\|A\|_*$ is the Frobenius norm and nuclear norm of matrix A respectively. $\|A\|_\infty := \max_{i,j} |A[i, j]|$. We write $a \gg b$ ($a \ll b$) if $a \geq Kb$ ($a \leq b/K$) for some sufficiently large number K.

2   Preliminary

Applications Sparse generalized eigenvalue problem has a wide application. Many high-dimensional multivariate statistical problems can be formulated as special instances of (1).

Example 1 [*Sparse Principal Component Analysis* [38, 4, 1, 31, 32]] *Given n observations with p features, sparse principal component analysis (SPCA) seeks the best low dimensional projection of the observed data for increasing interpretability, minimizing information loss and achieving sparsity structure. Let $\widetilde{\Sigma}$ be the empirical covariance matrix. SPCA aims to solve*

$$\max_{\mathbf{x} \in \mathbb{R}^p} \mathbf{x}^T \widetilde{\Sigma} \mathbf{x}, \ \ subject \ to \ \|\mathbf{x}\|_0 \leq k, \|\mathbf{x}\|_2 = 1. \tag{3}$$

Such problem is a special case of SGEP where $\widetilde{A} = \widetilde{\Sigma}$ and $\widetilde{B} = I$.

Example 2 [*Sparse Fisher's Discriminant Analysis* [27, 8, 13, 10, 12]] *Given n observations with p features from K different classes, Fisher's discriminant analysis seeks a projection for mapping observations to the low dimensional space on which the between-class variance Σ_b is large while the within-class variance Σ_w is small. Let $\widetilde{\Sigma}_b$ and $\widetilde{\Sigma}_w$ be the sample estimates of Σ_b and Σ_w, respectively. To obtain the sparse leading discriminant vector, one can solve*

$$\max_{\mathbf{x} \in \mathbb{R}^p} \mathbf{x}^T \widetilde{\Sigma}_b \mathbf{x}, \ \ subject \ to \ \mathbf{x}^T \widetilde{\Sigma}_w \mathbf{x} = 1, \ \|\mathbf{x}\|_0 \leq k. \tag{4}$$

Such a problem can be formulated as an SGEP with $\widetilde{A} = \widetilde{\Sigma}_b$ and $\widetilde{B} = \widetilde{\Sigma}_w$.

Example 3 [*Sparse Canonical Correlation Analysis* [35, 6, 7]] *Given two random vectors $X \in \mathbb{R}^p$, $Y \in \mathbb{R}^p$, let $\Sigma_{xx}, \Sigma_{yy}, \Sigma_{xy}$ be the covariance matrices for X, Y, and the cross-covariance matrix between X and Y, respectively. Let $\widetilde{\Sigma}_{xx}, \widetilde{\Sigma}_{yy}, \widetilde{\Sigma}_{xy}$ be estimators (constructed from samples) for Σ_{xx}, Σ_{yy}, Σ_{xy}, respectively. Sparse canonical correlation analysis (SCCA) aims to solve the constrained optimization problem:*

$$\max_{\mathbf{u}_x, \mathbf{u}_y} \mathbf{u}_x^{\mathrm{T}} \widetilde{\Sigma}_{xy} \mathbf{u}_y, \ \ subject \ to \ \mathbf{u}_x^{\mathrm{T}} \widetilde{\Sigma}_{xx} \mathbf{u}_x = 1, \ \mathbf{u}_y^{\mathrm{T}} \widetilde{\Sigma}_{yy} \mathbf{u}_y = 1,$$

$$\|\mathbf{u}_x\|_0 \leq k_x, \ \|\mathbf{u}_y\|_0 \leq k_y,$$

where k_x and k_y are two small integers. Such a problem can be recast as an SGEP with

$$\widetilde{A} = \begin{bmatrix} 0 & \widetilde{\Sigma}_{xy} \\ \widetilde{\Sigma}_{xy}^{\mathrm{T}} & 0 \end{bmatrix}, \quad \widetilde{B} = \begin{bmatrix} \widetilde{\Sigma}_{xx} & 0 \\ 0 & \widetilde{\Sigma}_{yy} \end{bmatrix}, \quad \mathbf{u}_1 = \begin{bmatrix} \mathbf{u}_x \\ \mathbf{u}_y \end{bmatrix}.^1$$

Strictly speaking, two problems are not exactly equivalent, because $\|\mathbf{u}\|_0 \leq k_x + k_y$ does not necessarily imply $\|\mathbf{u}_x\|_0 \leq k_x$ and $\|\mathbf{u}_y\|_0 \leq k_y$.

Technical preparation We first introduce some useful terminologies for describing the generalized eigenvalue problem (GEP). Let $A \in \mathbb{R}^{p \times p}$ be a symmetric matrix and $B \in \mathbb{R}^{p \times p}$ be a symmetric and positive definite matrix. Then GEP for the matrix pair (A, B) is defined as

$$A\mathbf{u}_i = \lambda_i B\mathbf{u}_i; \ \ i = 1, \ldots, p.$$

Without loss of generality, we can always assume that $\lambda_1 \geq \cdots \geq \lambda_p$. Here $\mathbf{u}_i$ is ith eigenvector and λ_i is ith eigenvalue; $(\lambda_i, \mathbf{u}_i)$ is called ith eigenpair of (A, B). Notice that $\mathbf{u}_i$ is only determined up to a scale. In this paper, we always assume that $\|\mathbf{u}_i\|_2 = 1$. The perturbation analysis of generalized eigenvalue problem has been extensively studied since 1970s and many perturbation bounds have been developed [22, 23, 24, 29]. For reader convenience, we present several existing fundamental results before moving to our main theory immediately.

Definition 1 *The angle between $\mathbf{x}, \mathbf{y}(\neq \mathbf{0}) \in \mathbb{R}^p$ is defined as $\theta(\mathbf{x}, \mathbf{y}) := \arccos \frac{|\mathbf{x}^{\mathrm{T}} \mathbf{y}|}{\|\mathbf{x}\|_2 \|\mathbf{y}\|_2}$.*

We assume $\mathbf{x}$ and $\mathbf{y} \in \mathbb{R}^p$ have unit l_2 norms and let $[\mathbf{x}, X_2]$ be an orthogonal matrix. Then it holds

$$|\sin \theta(\mathbf{x}, \mathbf{y})| = \|X_2^{\mathrm{T}} \mathbf{y}\|_2. \tag{5}$$

Furthermore, it can be checked that $|\sin \theta(\mathbf{x}, \mathbf{y})| \leq \|\mathbf{x} - \mathbf{y}\|_2 \leq \sqrt{2}|\sin \theta(\mathbf{x}, \mathbf{y})|$. Thus $|\sin \theta(\mathbf{x}, \mathbf{y})|$ is a measure to quantify the distance between two vectors. In generalized eigenvalue problem, Crawford number is an important quantity for characterizing the perturbation bound.

Definition 2 *The* Crawford number *of a symmetric matrix pair (A, B) is defined as*

$$c(A, B) := \min_{\|\mathbf{x}\|_2 = 1} \sqrt{(\mathbf{x}^{\mathrm{T}} A \mathbf{x})^2 + (\mathbf{x}^{\mathrm{T}} B \mathbf{x})^2}.$$

A symmetric matrix pair (A, B) is referred to as "definite" if $c(A, B) > 0$.

Obviously, the symmetric matrix pair (A, B) is definite for any positive definite B. By [22], it is known that the Crawford number is continuous with respect to the matrix pair.

3 Understanding of Perturbation Bound

3.1 On the General Setting

In order to understand the difference between the solution of (1) and the true leading eigenvector, we first describe the perturbation results without considering the sparsity. The following lemma is from 25 and is a generalization of the Davis-Kahn's $\sin \Theta$ theorem in 3 (also see 21, 11). It gives an upper bound for $|\sin \theta(\mathbf{u}_1, \widetilde{\mathbf{u}}_1)|$, where $\widetilde{\mathbf{u}}_1$ is the leading eigenvector of $(\widetilde{A}, \widetilde{B})$.

Lemma 1 *Suppose (A, B), $(\widetilde{A}, \widetilde{B}) = (A + E, B + F)$ are both symmetric-definite pairs, and their eigenvalues satisfy $\lambda_1 \geq \cdots \geq \lambda_p \geq 0$ and $\widetilde{\lambda}_1 \geq \cdots \geq \widetilde{\lambda}_p \geq 0$, respectively. Let $\phi_1 = \arctan \lambda_1 > \widetilde{\phi}_2 = \arctan \widetilde{\lambda}_2$, $\mathbf{u}_1$, $\widetilde{\mathbf{u}}_1$ be the eigenvectors corresponding to λ_1, $\widetilde{\lambda}_1$, respectively. Denote*

$$C_u = \frac{\sqrt{2(\|A\|_2^2 + \|B\|_2^2)}}{c(A, B)}, \quad \xi = \frac{\sqrt{\|E\mathbf{u}_1\|_2^2 + \|F\mathbf{u}_1\|_2^2}}{c(\widetilde{A}, \widetilde{B})},$$

then

$$|\sin \theta(\mathbf{u}_1, \widetilde{\mathbf{u}}_1)| \leq \frac{C_u \, \xi}{\sin(\phi_1 - \widetilde{\phi}_2)}.$$

Therefore, it holds asymptotically that

$$|\sin \theta(\mathbf{u}_1, \widetilde{\mathbf{u}}_1)| \lesssim \frac{C_u \, \epsilon}{c(A, B) \sin(\phi_1 - \phi_2)}. \tag{6}$$

Although perturbation upper bound has been studied extensively, there are not many results on lower bound. Next we establish the perturbation lower bound. That is, given any symmetric definite matrix pair (A, B) with $\lambda_1 > \lambda_2$ and any sufficiently small constant $\epsilon > 0$, we can always find a matrix pair (E, F) satisfying $\sqrt{\|E\|_2^2 + \|F\|_2^2} < \epsilon$ such that the distance between $\mathbf{u}_1$ and $\widetilde{\mathbf{u}}_1$ are lower bounded by a quantity in the same order of the bound in Lemma 1. The result is stated in the next lemma.

Lemma 2 *Follow the notations in Lemma 1. For any small positive constant ϵ, it holds that*

$$\sup_{(E,F) \in \mathcal{F}_\epsilon} |\sin \theta(\mathbf{u}_1, \widetilde{\mathbf{u}}_1)| \gtrsim \frac{C_l \, \epsilon}{c(A, B) \sin(\phi_1 - \phi_2)}, \tag{7}$$

where $C_l = \frac{c(A,B)}{\sqrt{2(\|A\|_2^2 + \|B\|_2^2)} \kappa^{\frac{3}{2}}}$ with $\kappa := \|B\|_2 \|B^{-1}\|_2$, $\mathcal{F}_\epsilon := \{(E, F) | \sqrt{\|E\|_2^2 + \|F\|_2^2} \leq \epsilon\}$.

Comparing (6) and (7) and noticing that C_u, C_l are two constants, we can see that upper and lower bounds of approximation error only differ up to a multiplicative constant. Quantity $\epsilon / c(A, B)$ can be seen as the relative perturbation and $\frac{1}{\sin(\phi_1 - \phi_2)}$ is a monotonically decreasing function of the gap $\phi_1 - \phi_2$. The upper and lower bounds are both proportional to $\frac{\epsilon}{c(A,B) \sin(\phi_1 - \phi_2)}$. Thus, we declare that the perturbation bound for the leading eigenvector $\mathbf{u}_1$ is *rate-optimal*.

3.2 On the Sparse Setting

Now we are ready to present the upper and lower perturbation bounds under the sparse setting. Throughout the rest of this section, the following assumptions are assumed.

A1 For any $\mathcal{K} \supset \mathrm{supp}(\mathbf{u}_1)$ with $|\mathcal{K}| \leq s + k$, it holds

$$\frac{\sqrt{\|\widetilde{A}_{\mathcal{K}} - A_{\mathcal{K}}\|_2^2 + \|\widetilde{B}_{\mathcal{K}} - B_{\mathcal{K}}\|_2^2}}{c(A_{\mathcal{K}}, B_{\mathcal{K}})} < 1,$$

where $c(A_{\mathcal{K}}, B_{\mathcal{K}})$ is the Crawford number of $(A_{\mathcal{K}}, B_{\mathcal{K}})$.

A2 For any $\mathcal{K} \supset \mathrm{supp}(\mathbf{u}_1)$ with $|\mathcal{K}| \leq s + k$, $\widetilde{A}_{\mathcal{K}}$ and $\widetilde{B}_{\mathcal{K}}$ are positive definite.

Assumption **A1** requires that the perturbation within a small superset of $\mathbf{u}_1$ is tiny. It says that one can get a good approximation for $\mathbf{u}_1$ (according to Lemma 1 and 2) when a small superset of $\mathbf{u}_1$ is available. It is, in fact, a necessary condition for computing a good approximation for $\mathbf{u}_1$. Assumption **A2** is a technical requirement to ensure positive definiteness for submatrices of $\widetilde{B}$.

156 We further adopt the following notations. The leading eigenvector $\mathbf{u}_1$ is assumed to be sparse, i.e.,
157 $\|\mathcal{S}\|_0 \ll p$ where $\mathcal{S} := \mathrm{supp}(\mathbf{u}_1)$. We also define the perturbation level

$$\epsilon := \max_{\mathcal{K}:\mathcal{K}\supset\mathrm{supp}(u_1)} \sqrt{\|\widetilde{A}_{\mathcal{K}} - A_{\mathcal{K}}\|_2^2 + \|\widetilde{B}_{\mathcal{K}} - B_{\mathcal{K}}\|_2^2},$$

158 the perturbation set $\mathcal{F}_{\epsilon,l} := \{(E,F) | \sqrt{\|E_{\mathcal{K}}\|_2^2 + \|F_{\mathcal{K}}\|_2^2} \leq \epsilon$ for any subset $\mathcal{K}$ with size less than $l\}$.

159 **Upper Bound of** $|\sin\theta(\mathbf{u}_1, \mathbf{x}_*)|$

160 **Theorem 1** *Let* $\mathcal{J} = \mathrm{supp}(\mathbf{x}_*)$, $\mathcal{K} = \mathcal{S}\cup\mathcal{J}$ *and* $\mathcal{L} = \mathcal{K}\setminus\mathcal{J}$. *Denote* $\rho_* = \frac{\mathbf{x}_*^{\mathrm{T}}\widetilde{A}\mathbf{x}_*}{\mathbf{x}_*^{\mathrm{T}}\widetilde{B}\mathbf{x}_*}$, $c_{\mathcal{K}} = c(A_{\mathcal{K}}, B_{\mathcal{K}})$
161 *and* $\widetilde{c}_{\mathcal{K}} = c(\widetilde{A}_{\mathcal{K}}, \widetilde{B}_{\mathcal{K}})$. *Let* $\delta = \frac{\|\widetilde{A}_{(\mathcal{L},\mathcal{J})}[\mathbf{x}_*]_{\mathcal{J}} - \rho_*\widetilde{B}_{(\mathcal{L},\mathcal{J})}[\mathbf{x}_*]_{\mathcal{J}}\|_2}{\widetilde{c}_{\mathcal{K}}}$. *If* $\delta < \sqrt{1 + \rho_*^2}$ *and*

$$\arctan\rho_* > \arctan\mu_2 + \arctan\epsilon + \arctan\frac{\delta}{\sqrt{1+\rho_*^2}}, \tag{8}$$

162 *then*

$$|\sin\theta(\mathbf{u}_1, \mathbf{x}_*)| \leq \frac{1}{\widetilde{c}_{\mathcal{K}}}\left(\frac{\epsilon\, C_\epsilon}{c(A_{\mathcal{K}}, B_{\mathcal{K}})\sin(\phi_1 - \widetilde{\phi}_2)} + \frac{\delta\, C_\delta}{\sin(\phi_* - \widetilde{\phi}_2)}\right),$$

163 *where* $\phi_* = \arctan\rho_*$, μ_2 *is the second largest eigenvalue for* $(A_{\mathcal{K}}, B_{\mathcal{K}})$, C_ϵ *and* C_δ *are some*
164 *constants.*

165 **Remark**: The upper bound has two terms. The first term is due to the perturbation, which is
166 approximately proportional to the perturbation level ϵ. The second term is due to the failure in finding
167 the true support set of $\mathbf{u}_1$. If $\mathrm{supp}(\mathbf{u}_1) \subset \mathrm{supp}(\mathbf{x}_*)$, then $\delta = 0$, consequently, the second term
168 vanishes. Therefore, it holds asymptotically that

$$|\sin\theta(\mathbf{u}_1, \mathbf{x}_*)| \leq \frac{C_{u,\mathcal{K}}\,\epsilon}{c(A_{\mathcal{K}}, B_{\mathcal{K}})\sin(\phi_1 - \phi_2)} \tag{9}$$

169 with $C_{u,\mathcal{K}} = \frac{\sqrt{2(\|A_{\mathcal{K}}\|_2^2 + \|B_{\mathcal{K}}\|_2^2)}}{c(A_{\mathcal{K}}, B_{\mathcal{K}})}$, when $\mathrm{supp}(\mathbf{x}_*) \supset \mathrm{supp}(\mathbf{u}_1)$ and ϵ is sufficiently small.

170 **Lower Bound of** $|\sin\theta(\mathbf{u}_1, \mathbf{x}_*)|$ Next we present the lower bound for $|\sin\theta(\mathbf{u}_1, \mathbf{x}_*)|$. The proof
171 follows immediately from the proofs of Theorem 1 and Lemma 2.

172 **Theorem 2** *Follow the notations and assumptions in Theorem 1. Then the following result holds.*

173 **Case (a)** *If* $\delta = 0$, *then*

$$\max_{(E,F)\in\mathcal{F}_{\epsilon,s+k}} |\sin\theta(\mathbf{u}_1, \mathbf{x}_*)| \gtrsim \frac{C_{l,\mathcal{K}}\,\epsilon}{c(A_{\mathcal{K}}, B_{\mathcal{K}})\sin(\phi_1 - \phi_2)},$$

174 *where* $C_{l,\mathcal{K}} = \frac{c(A_{\mathcal{K}}, B_{\mathcal{K}})}{\sqrt{2(\|A_{\mathcal{K}}\|_2^2 + \|B_{\mathcal{K}}\|_2^2)}\kappa_2^{\frac{3}{2}}(B_{\mathcal{K}})}$.

175 **Case (b)** *If* $\epsilon = 0$, *then*

$$\max_{(E,F)\in\mathcal{F}_{\epsilon,s+k}} |\sin\theta(\mathbf{u}_1, \mathbf{x}_*)| \gtrsim \frac{1}{\sqrt{2}\kappa_2^{\frac{3}{2}}(B_{\mathcal{K}})}\frac{\widehat{\xi}_{\mathcal{K}}}{\sin(\phi_* - \phi_2)},$$

176 *where* $\widehat{\xi}_{\mathcal{K}} = \sqrt{\frac{\|\widehat{E}[u_1]_{\mathcal{K}}\|_2^2 + \|\widehat{F}[u_1]_{\mathcal{K}}\|_2^2}{\|A\|_2^2 + \|B\|_2^2}}$, *and* $\widehat{E}, \widehat{F} \in \mathbb{R}^{|\mathcal{K}|\times|\mathcal{K}|}$ *are some matrices controlled by* δ.

177 In **Case (a)**, we can see that the lower bound is at the same order of the first term of the upper bound
178 in Theorem 1; In **Case (b)**, $\widehat{\xi}_{\mathcal{K}} = \mathcal{O}(\delta)$, then the lower bound is at the same order of the second term
179 of the upper bound in Theorem 1. We may declare that our perturbation bounds are *rate-optimal*.

180 **Connection to the literature** Although we do not make any statistical assumptions on E and F,
181 we still make connections to the statistical literature here. First, it can be computed that

$$\sin(\phi_1 - \phi_2) = \frac{\lambda_1 - \lambda_2}{\sqrt{\lambda_1^2 + 1}\sqrt{\lambda_2^2 + 1}}.$$

When $B = I$ is an identity matrix, $C_{u,\mathcal{K}} = O(\sqrt{\lambda_1^2 + 1})$ and $C_{l,\mathcal{K}} = \Omega(\frac{1}{\sqrt{\lambda_1^2+1}})$. Thus our lower bound can be simplified to $\frac{\sqrt{\lambda_2^2+1}}{\lambda_1 - \lambda_2}\epsilon$. In sparse PCA [32], it has been established that the minimax lower bound $\inf_{\tilde{\mathbf{u}}_1} \sup_{\mathcal{M}} (\mathbb{E}\|\mathbf{u}_1 - \tilde{\mathbf{u}}_1\|_2^2)^{1/2}$ is $\Theta(\frac{\sqrt{\lambda_1\lambda_2}}{\lambda_1-\lambda_2}\epsilon)$ with $\epsilon = \sqrt{\frac{s \log p}{n}}$ and $\mathcal{M}$ is the model space where covariance matrix A satisfies that its principle singular vector is s-sparse and λ_1 is larger than λ_2 by certain margin. When λ_1, λ_2 and λ_1/λ_2 are bounded constants, this minimax rate matches ours. In [2], they provide the minimax perturbation bound for singular subspace of non-symmetric matrices. The minimax rate of $\|\tilde{\mathbf{u}}_1 - \mathbf{u}_1\|_2$ is

$$\Theta(\frac{\alpha z_{21} + \beta z_{12}}{\alpha^2 + \beta^2 - \min\{z_{12}^2, z_{21}^2\}}),$$

where the detailed definitions of α, β, z_{12}, z_{21} can be found accordingly in 2. Especially, for symmetric A and $\tilde{A}$, the above bound can be simplified to $\frac{\epsilon}{\lambda_1 - \lambda_2}$ when $\epsilon \ll \lambda_1 - \lambda_2$. Again, when λ_1 and λ_2 are bounded constants, the rate is consistent with ours.

4 Understanding of Estimation Quality

In order to estimate the sparse leading eigenvector, it is straightforward to solve (1). It can be seen that the sparsity of solution to (1) depends on the choice of k, which may not be easily tuned. It is also computationally expensive to find the optimal solution to (1). Alternatively, such problem can be solved via regularization method. For example, 19 uses a smooth surrogate function to replace $\| \cdot \|_0$ penalty for solving SGEP.

A good estimator should have the following properties, 1. sparsity, 2. nearly unbiasedness, 3. stability. In high dimensional statistical problem, $\tilde{B}$ is always singular when the sample size is smaller than the number of features. In order to obtain a good approximation of $\mathbf{u}_1$, we consider the following restricted problem

$$(P1') \quad \min_{\mathbf{x} \in \mathbb{R}^p, \|\mathbf{x}\|_0 \leq s_n} -\mathbf{x}^T \widetilde{A}\mathbf{x} + p_\lambda(\mathbf{x}) \qquad s.t. \; \mathbf{x}^T \widetilde{B}\mathbf{x} \leq 1, \tag{10}$$

where $p_\lambda(\mathbf{x}) := \sum_{j=1}^p p_\lambda(\mathbf{x}[j])$ and $p_\lambda(x)$ is some univariate non-convex function, and s_n is the restricted dimension. In practice, s_n needn't be a small number. It can grow with sample size n. The restriction $\|\mathbf{x}\|_0 \leq s_n$ is needed for enforcing the solution to be nearly low-dimensional. Along with p_λ, the estimator can recover the sparsity structure. This method can be viewed as the counterpart of two stage methods [26, 6], where they need to find a good approximation of $\mathbf{u}_1$ in the first stage.

4.1 Penalization Function

For the choice of p_λ, we consider a family of special non-convex penalties:

$$\mathcal{P}_\lambda = \{p_\lambda(x) : p_\lambda(x) \text{ satisfies (a1) - (a3)}\}, \tag{11}$$

where (a1) Function $p_\lambda(x)$ is an even function, i.e., $p_\lambda(x) = p_\lambda(-x)$; (a2) The derivative of $p_\lambda(x)$, $p_\lambda'(x)$, exists in $(0, \infty)$; $\lim_{x \downarrow 0} p_\lambda'(x) = \lambda$ and $p_\lambda'(x) \equiv 0$ if $x \geq \gamma\lambda$ for some constant γ; (a3) On $(0, \infty)$, $p_\lambda'(x)$ is monotone decreasing and Lipschitz continuous, i.e., there exists a constant κ such that $0 \leq \frac{p_\lambda'(x_1) - p_\lambda'(x_2)}{x_2 - x_1} \leq \kappa$ for any $0 < x_1 < x_2$.

Here (a1) requires $p_\lambda(x)$ to be symmetric; (a2) specifies the local property of derivatives of $p_\lambda(x)$ around 0 and assumes the flatness of $p_\lambda(x)$ for larger x; (a3) puts continuity constraints on $p_\lambda'(x)$. Many popular penalty functions are included in $\mathcal{P}_\lambda$, for example, smoothly clipped absolute deviation (**SCAD**, 5), minimax concave penalty (**MCP**, 37) , etc. Based on the definitions, we can see that any $p_\lambda(x) \in \mathcal{P}_\lambda$ is similar to L1 norm locally around the origin. On the other hand, $p_\lambda(x)$ puts a smaller penalization on the signal compared with L1 penalty. They do not give the penalization to those large values. Intuitively speaking, the regularized estimator with non-convex penalty may outperform the L1 norm-based estimator since it is nearly unbiased.

4.2 Support Recovery

In this section, we provide the error bound and oracle properties of the non-convex estimator under suitable conditions. Some additional notations are introduced as follows. We define $\hat{\mathbf{u}}_1$ as $\arg\max_{\mathbf{x}:\|\mathbf{x}\|_0 \leq s_n, \mathbf{x}^T \widetilde{B}\mathbf{x} \leq 1} \mathbf{x}^T \widetilde{A}\mathbf{x}$ and denote $\tilde{\rho} = \hat{\mathbf{u}}_1^T \widetilde{A}\hat{\mathbf{u}}_1$. Vector $\hat{\mathbf{u}}_1$ can be viewed as the best approximation of the leading eigenvector in the restricted space. We define $c(A, B, s_n) := \max_{|\mathcal{K}| \leq s_n} c(A_\mathcal{K}, B_\mathcal{K})$ and $c(\tilde{A}, \tilde{B}, s_n) := \max_{|\mathcal{K}| \leq s_n} c(\widetilde{A}_\mathcal{K}, \widetilde{B}_\mathcal{K})$. We also define $N(A, B, s_n) := \max_{|\mathcal{K}| \leq s_n} \sqrt{\|A_\mathcal{K}^2 + B_\mathcal{K}^2\|_2}$. We define $\epsilon_f := \max_{|\mathcal{K}| \leq s_n} \|F_\mathcal{K}\|_2$ and $\epsilon_s := \max_{\mathcal{K}:\|\mathcal{K}\|_0 \leq s_n}\{\|E_\mathcal{K}\|_2, \|F_\mathcal{K}\|_2\}$. A different set of assumptions is stated as follows.

B0 (*Regularity*) $\lambda_1 - \lambda_2$ is positive; $\|B\|, \|B^{-1}\|$ are bounded by some constant.

B1 (*Signal*) $\min\{|\mathbf{u}_1[j]| : j \in \mathrm{supp}(\mathbf{u}_1)\} \gg \sqrt{S}\lambda$.

B2 (*Penalization*) $\lambda \gg \epsilon_s \cdot C_F$, where C_F is a constant depending on $c(A, B, s_n)$, $c(\tilde{A}, \tilde{B}, s_n)$ and $N(A, B, s_n)$.

B3 (*Support size*) $|\mathcal{S}|\lambda^2 \ll 1$.

B4 (*Identifiability*) $\max_{\mathcal{K}:\|\mathcal{K}\|_0 \leq s_n, \mathcal{S} \not\subset \mathcal{K}} \lambda_1(A_\mathcal{K}, B_\mathcal{K}) < \lambda_1 - 2\epsilon$.

Condition **B0** ensures that the leading eigenvector is unique and underlying matrix B is well-behaved. Condition **B1** requires that the absolute values of entries in the true support is not too small so that $\mathcal{S}$ can be identified. Condition **B2** specifies the relationship between penalty level λ and noise level ϵ. That is, penalty level λ should be at least larger than the noise level up to a multiplicative constant. Condition **B3** makes sure that the support size is not too large, i.e., $\mathbf{u}_1$ should be sparse. The following theorem guarantees that the estimator is not far away from the re-scaled leading eigenvector. Condition **B4** is for the identifiability of $|\mathcal{S}|$ to ensure that the support of $\mathbf{u}_1$ could be still identified after perturbation. Without loss of generality, we always assume $s_n \geq |\mathcal{S}|$. Then we have the following results.

Theorem 3 *Let $\hat{\mathbf{x}}$ be the optimizer of Problem (P1'). Under Conditions **B0** - **B4**, it holds that*

$$\min_{\mathrm{sgn} \in \{-1, 1\}} \|\mathrm{sgn} \cdot \hat{\mathbf{x}} - \mathbf{u}_{1s}\|_2 \leq C_1 \sin(\theta(\hat{\mathbf{u}}_1, \mathbf{u}_1)) + C_2\epsilon_f,$$

for some constants C_1 and C_2.

Here $\mathbf{u}_{1s} = \mathbf{u}_1(\mathbf{u}_1^T B\mathbf{u}_1)^{-1/2}$ is the re-scaled leading eigenvector. Furthermore, we can recover the support of $\mathbf{u}_1$ under this restricted problem.

Theorem 4 *Under the same conditions in Theorem 3, it holds that*

$$\mathrm{supp}(\hat{\mathbf{x}}) = \mathrm{supp}(\mathbf{u}_1). \tag{12}$$

Based on Theorem 4, we actually have even stronger results. When all required conditions are met, the proposed estimator is equal to the oracle estimator which is the one estimated when the true support $\mathcal{S}$ is known to us. Therefore, the estimator achieves the optimal error bound which matches the one obtained in Theorem 1. Specifically, the proposed estimator achieves the optimal error rate $(\frac{|\mathcal{S}|\log p}{n})^{1/2}$ in sparse PCA and sparse CCA problems.

Theorem 5 *Under the same set of conditions in Theorem 4, we have that*

$$|\sin\theta(\hat{\mathbf{x}}, \mathbf{u}_1)| \leq C_{u,\mathcal{S}} \frac{\epsilon}{c(A_\mathcal{S}, B_\mathcal{S})\sin(\phi_1 - \phi_2)}, \tag{13}$$

with $C_{u,\mathcal{S}} = C\frac{\sqrt{2(\|A_\mathcal{S}\|_2^2 + \|B_\mathcal{S}\|_2^2)}}{c(A_\mathcal{S}, B_\mathcal{S})}$ (C is a universal constant).

Remark 1 *To achieve the optimal statistical error rate $(\frac{|\mathcal{S}|\log p}{n})^{1/2}$, most existing methods are two-stage based. For example, 34 study sparse PCA problem and propose a "sparse orthogonal iteration pursuit" (SOAP) algorithm which involves "relax" and "tighten" stages; 6 consider sparse CCA problem and adopt the two-stage regularization approach with L1 penalty for the first stage and group LASSO penalty for the second stage. By comparison, our method does not require convex relaxation in the first stage.*

4.3 Computation

For computational purpose, we consider to solve (P1') by using the alternating direction method of multipliers (ADMM, 33, 17). Specifically, we relax the problem (P1') by reformulating it to

$$\min_{\mathbf{x},\mathbf{y},\mathbf{z}} \mathcal{L}(\mathbf{x},\mathbf{z},\mathbf{y}), \qquad s.t.\ \mathbf{x}^T \widetilde{B} \mathbf{x} = 1,\ \|\mathbf{x}\|_0 \le s_n, \tag{14}$$

where

$$\mathcal{L}(\mathbf{x},\mathbf{z},\mathbf{y}) = -\mathbf{x}^T \tilde{A}\mathbf{x} + p_\lambda(\mathbf{z}) + \mathbf{y}^T(\mathbf{x} - \mathbf{z}) + \frac{\eta}{2}\|\mathbf{x} - \mathbf{z}\|_2^2.$$

In (14), we introduce several auxiliary variables for the following reasons. We construct a new vector $\mathbf{z}$ which is a copy of $\mathbf{x}$. It can help us to split the original problem to two simple separate sub-problems. $\mathbf{y}$ is the dual variable for the constraint $\mathbf{x} = \mathbf{z}$.

By formulation (14), we can optimize the objective function with respect to each variable iteratively. The steps for updating $\mathbf{x}, \mathbf{y}, \mathbf{z}$ are described as follows.

Update $\mathbf{x}$: At $(t+1)$-th iteration, we aim to find $\mathbf{x}^{(t+1)}$ which is

$$\arg\min_{\mathbf{x} \in \mathcal{D}} \quad \frac{\eta}{2}\|\mathbf{x} - \mathbf{z}^{(t)}\|_2^2 + (\mathbf{y}^{(t)})^T(\mathbf{x} - \mathbf{z}^{(t)}) - \mathbf{x}^T \tilde{A}\mathbf{x}, \tag{15}$$

where $\mathcal{D} = \{\mathbf{x} : \mathbf{x}^T \widetilde{B}\mathbf{x} = 1, \mathbf{x}[-\mathcal{S}_t] = \mathbf{0}\}$ and $\mathcal{S}_t$ is the set of indices corresponding to first s_n largest absolute values of entries in $\mathbf{b}_1^{(t+1)}$ (see (16)). This sub-problem can be solved via the following recursive formula,

$$\mathbf{b}_m^{(t+1)} = \mathbf{z}^{(t)} - \frac{\mathbf{y}^{(t)} - \tilde{A}\mathbf{x}_{m-1}^{(t+1)}}{\eta}, \tag{16}$$

$$\mathbf{x}_m^{(t+1)}[\mathcal{S}_t] = (\beta_m^{(t+1)} \widetilde{B}_{\mathcal{S}_t} + I)^{-1}\mathbf{b}_m^{(t+1)}[\mathcal{S}_t],$$

where $\mathbf{x}_0^{(t+1)} = \mathbf{x}^{(t)}$ and $m = 1, 2, \ldots$. Scalar $\beta_m^{(t+1)}$ satisfies $1 = \sum_j \frac{d_j(\widetilde{\mathbf{b}}_m^{(t+1)}[j])^2}{(\beta_m^{(t+1)}d_j+1)^2}$ with $\widetilde{\mathbf{b}}_m^{(t+1)} = U^T \mathbf{b}_m^{(t+1)}[\mathcal{S}_t]$; $\widetilde{B}_{\mathcal{S}_t} = UDU^T$ and $D = \mathrm{diag}(d_1, \ldots, d_{s_n})$.

Update $\mathbf{z}$: We know that $\mathbf{y}^{(t+1)} = \arg\min_{\mathbf{z}} \frac{\eta}{2}\|\mathbf{z} - \mathbf{x}^{(t)}\|_2^2 + (\mathbf{y}^{(t)})^T(\mathbf{x}^{(t+1)} - \mathbf{z}) + p_\lambda(\mathbf{z})$. Each entry of $\mathbf{z}$ can be optimized separately. Specifically, if we take p_λ as the MCP penalty, then $\mathbf{y}^{(t+1)}$ has the following analytical form,

$$\mathbf{z}^{(t+1)}[j] = \begin{cases} \check{\mathbf{z}}^{(t)} & \text{if } |\check{\mathbf{z}}^{(t)}| > \gamma\lambda, \\ \frac{\mathrm{sgn}(\check{\mathbf{z}}^{(t)}[j])(|\check{\mathbf{z}}^{(t)}[j]| - \frac{\lambda}{\eta})_+}{1 - \frac{1}{\eta\gamma}} & \text{if } |\check{\mathbf{z}}^{(t)}| \le \gamma\lambda, \end{cases}$$

where $\check{\mathbf{z}}^{(t)} := \mathbf{x}^{(t+1)} + \frac{\mathbf{y}^{(t)}}{\eta}$.

Update $\mathbf{y}$: By dual variable update in 17, we have $\mathbf{y}^{(t+1)} = \mathbf{y}^{(t)} + \eta(\mathbf{x}^{(t+1)} - \mathbf{z}^{(t+1)})$.

We call the above procedure as SGEP (NC-SGEP) algorithm. Such proposed algorithm works on matrix-vector product and eigen-decomposition for submatrices of $\tilde{B}$, which is computationally efficient. Compared with other truncation methods, we do not need to make efforts to choose best s_n due to the existence of non-convex regularization term. In practice, we can choose s_n comparable to the sample size. The following theorem gives the local convergence of NC-SGEP algorithm.

Theorem 6 *Under Conditions **B0** - **B4**, if η is large enough and the initial supports of $\mathbf{x}^{(0)}$ and $\mathbf{z}^{(0)}$ include* $\mathrm{supp}(\mathbf{u}_1)$*, then it holds*

$$T(\epsilon) \le \frac{C(\mathcal{L}(\mathbf{x}^{(0)}, \mathbf{z}^{(0)}, \mathbf{y}^{(0)}) - \bar{f})}{\epsilon},$$

where $T(\epsilon) = \min\{t : \|\mathbf{x}^{(t)} - \mathbf{z}^{(t)}\| \le \epsilon\}$ and $\bar{f} = \min_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \mathbf{x}, \mathbf{y})$ for some constant C. Moreover, $\mathrm{supp}(\hat{\mathbf{u}}_1) = \mathrm{supp}(\mathbf{u}_1)$ *if $\epsilon = o(\epsilon_s)$.*

5 Numerical Experiments

Validation of Perturbation Bounds We conduct perturbation analyses of proposed estimator under different settings. The matrix pair is set as $A = 4\mathbf{u}_1\mathbf{u}_1^T + I - P_{\mathbf{u}_1}$, $B = I$. We sample n data which follows $N(0, A)$ and sample another n data which follows $N(0, B)$. The $\tilde{A}$ and $\tilde{B}$ are constructed based on the sample covariance correspondingly. The leading eigenvector $\mathbf{u}_1$ has unit norm and has non-zero entries in first $|\mathcal{S}|$ positions. We fixed dimension $p \equiv 100$ and let number of samples (n) and support size ($|\mathcal{S}|$) vary. Each setting is repeated for 100 times with fixed choice of $\lambda = 0.3$, $\eta = 1$, $s_n = 25$. The mean and standard deviation of $\|\hat{\mathbf{x}} - \mathbf{u}_1\|_2$ are reported. We also compute the oracle estimator which is the best $\hat{\mathbf{x}}$ when true support $\mathcal{S}$ is known. From Table 1, when noise level $\sqrt{\log p/n}$ is small, we can see that the estimation error is quite close to the optimal (oracle) error. In addition, we can see that estimation error is proportional to $\sqrt{|\mathcal{S}|}$ and $\sqrt{\log p/n}$. This indicates that our method can achieve the theoretical optimal error rate, i.e., $O(\sqrt{\frac{|\mathcal{S}|\log p}{n}})$.

Table 1: Estimation error $\|\hat{\mathbf{x}} - \mathbf{u}_1\|_2$ under different perturbation and sparsity level. "Oracle" / "Est": with / without knowing support $\mathcal{S}$. "Sd" is the standard deviation of "Est".

$|\mathcal{S}|$	n $\sqrt{\frac{\log p}{n}}$	100	200	400	800	1600	3200	6400
		0.230	0.162	0.115	0.081	0.057	0.041	0.029
2	Oracle	0.056	0.036	0.029	0.019	0.013	0.010	0.007
	Est	0.140	0.047	0.043	0.024	0.019	0.012	0.008
	Sd	(0.190)	(0.034)	(0.030)	(0.022)	(0.013)	(0.009)	(0.006)
4	Oracle	0.107	0.073	0.057	0.037	0.028	0.019	0.014
	Est	0.225	0.131	0.077	0.043	0.032	0.022	0.017
	Sd	(0.244)	(0.061)	(0.036)	(0.028)	(0.017)	(0.011)	(0.009)
8	Oracle	0.173	0.118	0.087	0.062	0.044	0.030	0.022
	Est	0.277	0.191	0.110	0.082	0.058	0.043	0.028
	Sd	(0.279)	(0.156)	(0.066)	(0.039)	(0.031)	(0.022)	(0.012)

Validation of Sparsity Recovery The underlying matrix pair is set as $A = 3\mathbf{u}_1\mathbf{u}_1^T + I - P_{\mathbf{u}_1}$, $B = I$. Thus, we naturally take $\tilde{B} = B = I$ and $\tilde{A}$ as the sample covariance of data. We let dimension p grow from 16 to 256, fix the sample size $n = 100$ and set $\lambda = 0.5$, $\eta = 1$ and $s_n = 50$. We compare the proposed method with the semidefinite programming method [30, 14, 36, SDP] with L1 (SDP_L1) and MCP (SDP_MCP) penalty. The estimation error and percentage of support recovery are reported in Table 2. We can see that the proposed method can have a slightly better performance. This is because that NC-SGEP optimizes objective in space $\mathbb{R}^p$ unlike those SDP methods work on space $\mathbb{R}^{p \times p}$. In addition, the method can recover the sparsity structure pretty well, while SDP methods can never recover the true support set.

Table 2: Estimation accuracy for sparse canonical correlation analysis.

	p	16	32	64	128	256
	NC-SGEP	0.074 (0.056)	0.076 (0.068)	0.072 (0.059)	0.081(0.054)	0.084 (0.054)
$\|\hat{\mathbf{x}}\hat{\mathbf{x}}^T - \mathbf{u}_1\mathbf{u}_1^T\|_\infty$	SDP_L1	0.071 (0.046)	0.080 (0.041)	0.095 (0.049)	0.094 (0.039)	0.100 (0.041)
	SDP_MCP	0.070 (0.045)	0.079 (0.040)	0.093 (0.048)	0.093 (0.038)	0.096 (0.039)
	NC-SGEP	93 %	82 %	68 %	59 %	49 %
Recovery of $|\mathcal{S}|$	SDP_L1	-	-	-	-	-
	SDP_MCP	-	-	-	-	-

6 Conclusion

In this paper, we establish the upper and lower bounds for perturbation analysis of sparse generalized eigenvalue problem. We also propose a new statistical estimation method. The proposed method gives a sparse, nearly unbiased and stable solution to SGEP. We show that the proposed estimator can achieve the optimal estimation error rate. Multiple numerical results validate our theories and also show the superior performance of the proposed method. In the future work, we may focus on extending the current results to the problem of finding multiple leading sparse eigenvectors. Although computing leading vectors one by one seems straightforward, it is still not an easy problem since it requires the orthogonalization procedure. The computational complexity may require a special care.

References

[1] Aharon Birnbaum, Iain M Johnstone, Boaz Nadler, and Debashis Paul. Minimax bounds for sparse pca with noisy high-dimensional data. *Annals of statistics*, 41(3):1055, 2013.

[2] T Tony Cai and Anru Zhang. Rate-optimal perturbation bounds for singular subspaces with applications to high-dimensional statistics. *The Annals of Statistics*, 46(1):60–89, 2018.

[3] Chandler Davis and William Morton Kahan. The rotation of eigenvectors by a perturbation. III. *SIAM J. Numer. Anal.*, 7(1):1–46, 1970.

[4] Alexandre d'Aspremont, Francis Bach, and Laurent El Ghaoui. Optimal solutions for sparse principal component analysis. *J. Machine Learning Res.*, 9(Jul):1269–1294, 2008.

[5] Jianqing Fan and Runze Li. Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of the American statistical Association*, 96(456):1348–1360, 2001.

[6] Chao Gao, Zongming Ma, and Harrison H Zhou. An efficient and optimal method for sparse canonical correlation analysis. *ArXiV e-prints*, 1409, 2014.

[7] Chao Gao, Zongming Ma, Harrison H Zhou, et al. Sparse cca: Adaptive estimation and computational barriers. *The Annals of Statistics*, 45(5):2074–2101, 2017.

[8] Yaqian Guo, Trevor Hastie, and Robert Tibshirani. Regularized linear discriminant analysis and its application in microarrays. *Biostatistics*, 8(1):86–100, 2007.

[9] Sungkyu Jung, Jeongyoun Ahn, and Yongho Jeon. Penalized orthogonal iteration for sparse estimation of generalized eigenvalue problem. *J. Comput. Graphical Statist.*, pages 1–23, 2019.

[10] Mladen Kolar and Han Liu. Optimal feature selection in high-dimensional discriminant analysis. *IEEE Trans. Inf. Theory*, 61(2):1063–1083, 2014.

[11] Ren-Cang Li. Matrix perturbation theory. In L. Hogben, R. Brualdi, and G. W. Stewart, editors, *Handbook of Linear Algebra*, chapter 21. CRC Press, Boca Raton, FL, 2nd edition, 2014.

[12] Qing Mai, Yi Yang, and Hui Zou. Multiclass sparse discriminant analysis. *arXiv:1504.05845*, 2015.

[13] Qing Mai, Hui Zou, and Ming Yuan. A direct approach to sparse discriminant analysis in ultra-high dimensions. *Biometrika*, 99(1):29–42, 2012.

[14] Jérôme Malick, Janez Povh, Franz Rendl, and Angelika Wiegele. Regularization methods for semidefinite programming. *SIAM Journal on Optimization*, 20(1):336–356, 2009.

[15] Baback Moghaddam, Yair Weiss, and Shai Avidan. Generalized spectral bounds for sparse LDA. In *ICML*, pages 641–648. ACM, 2006.

[16] Baback Moghaddam, Yair Weiss, and Shai Avidan. Spectral bounds for sparse PCA: Exact and greedy algorithms. In *Adv. Neural Inform. Process. Systems*, pages 915–922, 2006.

[17] Neal Parikh and Stephen Boyd. Proximal algorithms. *Foundations and Trends in optimization*, 1(3):127–239, 2014.

[18] Sandra E Safo, Jeongyoun Ahn, Yongho Jeon, and Sungkyu Jung. Sparse generalized eigenvalue problem with application to canonical correlation analysis for integrative analysis of methylation and gene expression data. *Biometrics*, 2018.

[19] Junxiao Song, Prabhu Babu, and Daniel P Palomar. Sparse generalized eigenvalue problem via smooth optimization. *IEEE Trans. Signal Process.*, 63(7):1627–1642, 2015.

[20] Bharath K Sriperumbudur, David A Torres, and Gert RG Lanckriet. A majorization-minimization approach to the sparse generalized eigenvalue problem. *Mach. Learn.*, 85(1-2):3–39, 2011.

[21] Gilbert W Stewart and Ji-Guang Sun. *Matrix Perturbation Theory*. Academic Press, Boston, 1990.

[22] GW Stewart. Pertubation bounds for the definite generalized eigenvalue problem. *Linear Algebra Appl.*, 23:69–85, 1979.

[23] Ji-Guang Sun. The perturbation bounds of generalized eigenvalues of a class of matrix-pairs. *Math. Numer. Sinica*, 4(1):23–29, 1982.

[24] Ji-Guang Sun. Perturbation analysis for the generalized eigenvalue and the generalized singular value problem. In *Matrix Pencils*, pages 221–244. Springer, 1983.

[25] Ji-guang Sun. The perturbation bounds for eigenspaces of a definite matrix-pair. *Numer. Math.*, 41(3):321–343, 1983.

[26] Kean Ming Tan, Zhaoran Wang, Han Liu, and Tong Zhang. Sparse generalized eigenvalue problem: Optimal statistical rates via truncated Rayleigh flow. *J. R. Statist. Soc. B*, 80(5):1057–1086, 2018.

[27] Robert Tibshirani, Trevor Hastie, Balasubramanian Narasimhan, and Gilbert Chu. Class prediction by nearest shrunken centroids, with applications to dna microarrays. *Statistical Science*, pages 104–117, 2003.

[28] Kengo Uchida and Isao Yamada. A nested ℓ_1-penalized adaptive normalized quasi-Newton algorithm for sparsity-aware generalized eigen-subspace extraction. In *2018 Asia-Pacific Signal and Information Processing Association Annual Summit and Conference (APSIPA ASC)*, pages 212–217. IEEE, 2018.

[29] Charles F Van Loan and Gene H Golub. *Matrix Computations*. Johns Hopkins University Press, Baltimore, MD, 4th edition, 2012.

[30] Lieven Vandenberghe and Stephen Boyd. Semidefinite programming. *SIAM review*, 38(1):49–95, 1996.

[31] Vincent Q Vu, Juhee Cho, Jing Lei, and Karl Rohe. Fantope projection and selection: A near-optimal convex relaxation of sparse pca. In *Advances in neural information processing systems*, pages 2670–2678, 2013.

[32] Vincent Q Vu, Jing Lei, et al. Minimax sparse principal subspace estimation in high dimensions. *The Annals of Statistics*, 41(6):2905–2947, 2013.

[33] Bo Wahlberg, Stephen Boyd, Mariette Annergren, and Yang Wang. An admm algorithm for a class of total variation regularized estimation problems. *IFAC Proceedings Volumes*, 45(16):83–88, 2012.

[34] Zhaoran Wang, Huanran Lu, and Han Liu. Nonconvex statistical optimization: Minimax-optimal sparse pca in polynomial time. *arXiv preprint arXiv:1408.5352*, 2014.

[35] Daniela M Witten, Robert Tibshirani, and Trevor Hastie. A penalized matrix decomposition, with applications to sparse principal components and canonical correlation analysis. *Biostatistics*, 10(3):515–534, 2009.

[36] Henry Wolkowicz, Romesh Saigal, and Lieven Vandenberghe. *Handbook of semidefinite programming: theory, algorithms, and applications*, volume 27. Springer Science & Business Media, 2012.

[37] Cun-Hui Zhang. Nearly unbiased variable selection under minimax concave penalty. *The Annals of statistics*, 38(2):894–942, 2010.

[38] Hui Zou, Trevor Hastie, and Robert Tibshirani. Sparse principal component analysis. *J. Comput. Graphical Statist.*, 15(2):265–286, 2006.