On the Identifiability of Nonlinear ICA: Sparsity and Beyond

Abstract

Nonlinear independent component analysis (ICA) aims to recover the underlying independent latent sources from their observable nonlinear mixtures. How to make the nonlinear ICA model identifiable up to certain trivial indeterminacies is a long-standing problem in unsupervised learning. Recent breakthroughs reformulate the standard independence assumption of sources as conditional independence given some auxiliary variables (e.g., class labels and/or domain/time indexes) as weak supervision or inductive bias. However, nonlinear ICA with unconditional priors cannot benefit from such developments. We explore an alternative path and consider only assumptions on the mixing process, such as *Structural Sparsity* or *Independent Influences*. We show that under specific instantiations of such constraints, the independent latent sources can be identified from their nonlinear mixtures up to a permutation and a component-wise transformation, thus achieving nontrivial identifiability of nonlinear ICA without auxiliary variables. We provide estimation methods and validate the theoretical results experimentally. The results on image data suggest that our conditions may hold in a number of practical data generating processes.

1 Introduction

Nonlinear independent component analysis (ICA) is fundamental in unsupervised learning. It generalizes linear ICA (Comon, 1994) to identify latent sources from observations, which are assumed to be a nonlinear mixture of the sources. For an observed vector $\mathbf{x}$, nonlinear ICA expresses it as $\mathbf{x} = \mathbf{f}(\mathbf{s})$, where $\mathbf{f}$ is an unknown invertible mixing function, and $\mathbf{s}$ is a latent random vector representing the (marginally) independent sources. The goal is to recover function $\mathbf{f}$ as well as sources $\mathbf{s}$ from the observed mixture $\mathbf{x}$ up to certain indeterminacies. While nonlinear ICA is of general interest in a variety of tasks, such as disentanglement (Lachapelle et al., 2022) and unsupervised learning (Oja, 2002), its identifiability has been a long-standing problem for decades. The key difficulty is that, without additional assumptions, there exist infinite ways to transform the observations into independent components while still mixed w.r.t. the sources (Hyvärinen and Pajunen, 1999).

To deal with this challenge, existing works introduce the auxiliary variable $\mathbf{u}$ (e.g., class label, domain index, time index) and assume that sources are conditionally independent given $\mathbf{u}$ (Hyvärinen and Morioka, 2017; Hyvärinen et al., 2019; Yao et al., 2021; Lachapelle et al., 2022). Most of them require the auxiliary variable to be observable, while clustering-based methods (Willetts and Paige, 2021) and those for time series (Hälvä and Hyvärinen, 2020; Hälvä et al., 2021) are exceptions. These works impose mild restrictions on the mixing process but require many distinct values of $\mathbf{u}$ for identifiability, which might restrict their practical applicability. Motivated by this, Yang et al. (2022) reduce the number of required distinct values by strengthening the functional assumptions on the mixing process. As described, all these results are based on conditional independence of the sources given auxiliary variable $\mathbf{u}$, instead of the standard marginal independence assumption, to achieve identifiability in specific

tasks, thanks to the weak supervision provided by the auxiliary variables. The identifiability of the original setting without auxiliary variables, which is crucial for unsupervised learning, stays unsolved.

Instead of introducing auxiliary variables, there exists the possibility to rely on further restrictions on the mixing functions to achieve identifiability, but limited results are available in the literature over the past two decades. With the assumption of conformal transformation, Hyvärinen and Pajunen (1999) proved identifiability of the nonlinear ICA solution up to rotation indeterminacy for only two sources. Another result is shown for the mixture that consists of component-wise nonlinearities added to a linear mixture (Taleb and Jutten, 1999).

In this work, we aim to show the identifiability of nonlinear ICA with unconditional priors. We prove that without any kind of auxiliary variables, the latent sources can be identified from nonlinear mixtures up to a component-wise invertible transformation and a permutation under assumptions only on the mixing process. Specifically, we mainly focus on the *structural sparsity* condition w.r.t. the structure from sources to observed variables, i.e., the support of the Jacobian matrix of the mixing function. Under structural conditions on the mixing process, one could identify the true sources by sparsity regularization during estimation (Thm. 1). By removing rotation indeterminacy, structural sparsity benefits the identifiability of linear ICA with Gaussian sources as well, which was previously thought to be unsolvable (Prop. 1, Thm. 2). Besides, we also develop another identifiability theory based on *Independent Influences*, which is an analogous concept of "sparsity" regarding the individual influencing process instead of the support (Thm. 4). We establish, to the best of our knowledge, one of the first identifiability results of the original nonlinear ICA model in a fully unsupervised setting.

Moreover, we also provide identifiability results for the underdetermined case, where the number of observed variables is larger than that of sources (Thm. 2, Thm. 3). This is of great practical interest and cannot be handled with previous conditions on nonlinear ICA, even with the help of auxiliary variables (Hyvärinen et al., 2019). We validate our theoretical claims experimentally, and the results on the image dataset suggest that our conditions appear to be reasonable for practical data generating processes.

2 Preliminaries

We consider the following data-generating process of ICA

$$p_{\mathbf{s}}(\mathbf{s}) = \prod_{i=1}^{n} p_{\mathbf{s}_i}(\mathbf{s}_i), \tag{1}$$

$$\mathbf{x} = \mathbf{f}(\mathbf{s}), \tag{2}$$

where n denotes the number of latent sources, $\mathbf{x} = (\mathbf{x}_1, \ldots, \mathbf{x}_n)$ denotes the observed random vector, $\mathbf{s} = (\mathbf{s}_1, \ldots, \mathbf{s}_n)$ is the latent random vector representing the marginally independent sources, $p_{\mathbf{s}_i}$ is the marginal probability density function (PDF) of the i-th source s_i, $p_{\mathbf{s}}$ is the joint PDF of random vector $\mathbf{s}$, and $\mathbf{f} : \mathbf{s} \to \mathbf{x}$ denotes a nonlinear mixing function. For linear ICA, Eq. (2) is restricted as

$$\mathbf{x} = \mathbf{A}\mathbf{s}, \tag{3}$$

where $\mathbf{A} := [\mathbf{A}_{:,1} \cdots \mathbf{A}_{:,n}]$ denotes the mixing matrix. The goal of ICA is to learn an estimated unmixing function $\hat{\mathbf{f}}^{-1} : \mathbf{x} \to \hat{\mathbf{s}}$ (or alternatively in the linear case, an estimated unmixing matrix $\hat{\mathbf{A}}^{-1}$), such that $\hat{\mathbf{s}} = (\hat{\mathbf{s}}_1, \ldots, \hat{\mathbf{s}}_n)$ consists of independent estimated sources.

Notations. Throughout this work, for any matrix $\mathbf{M}$, we use $\mathbf{M}_{i,:}$ to refer to its ith row, and $\mathbf{M}_{:,j}$ to indicate its jth column. For any set of indices $\mathcal{S} \subset \{1, \ldots, m\} \times \{1, \ldots, n\}$, we analogously denote $\mathcal{S}_{i,:} := \{j \mid (i,j) \in \mathcal{S}\}$ and $\mathcal{S}_{:,j} := \{i \mid (i,j) \in \mathcal{S}\}$. We denote by $|\cdot|$ the cardinality of a set. We also define the following technical notations.

Definition 1. *Given a subset $\mathcal{S} \subseteq \{1, \ldots, n\}$, the subspace $\mathbb{R}^n_{\mathcal{S}}$ is defined as*

$$\mathbb{R}^n_{\mathcal{S}} := \{z \in \mathbb{R}^n \mid i \notin \mathcal{S} \implies z_i = 0\}. \tag{4}$$

In other words, $\mathbb{R}^n_{\mathcal{S}}$ refers to the subspace of $\mathbb{R}^n$ indicated by an index set $\mathcal{S}$. In the following, we define the support of a matrix.

Definition 2. *The support of matrix $\mathbf{M} \in \mathbb{R}^{m \times n}$ is defined as*

$$\text{supp}(\mathbf{M}) := \{(i,j) \mid \mathbf{M}_{i,j} \neq 0\}. \tag{5}$$

With slight abuse of notation, we also reuse the notation supp to denote the support of a matrix-valued function, depending on the context.

Definition 3. *The support of function* $\mathbf{M} : \Theta \to \mathbb{R}^{m \times n}$ *is defined as*

$$\mathrm{supp}(\mathbf{M}(\Theta)) := \{(i, j) \mid \exists \theta \in \Theta, \mathbf{M}(\theta)_{i,j} \neq 0\}. \tag{6}$$

3 Identifiability with Structural Sparsity

3.1 Nonlinear ICA with Unconditional Priors: Structural Sparsity

In this section, we consider the identifiability of nonlinear ICA with unconditional priors, where the mixing function $\mathbf{f}$ in Eq. (2) is nonlinear. Different from recent breakthroughs in identifiable nonlinear ICA with side information (Hyvärinen et al., 2019; Lachapelle et al., 2022), in our setting, sources $\mathbf{s}$ in Eq. (1) do not need to be mutually conditionally independent given an auxiliary variable. Instead, our setting is consistent with the original ICA problem based on a general marginal independence assumption of sources. Besides, no particular form of the source distribution is assumed, which is different from most works that directly assume an exponential family (Hyvärinen et al., 2019; Lachapelle et al., 2022). The goal of identifiable nonlinear ICA is to estimate the unmixing function $\hat{\mathbf{f}}^{-1}$ so that the independent sources are identified up to a certain indeterminacy, which is usually a composition of a component-wise invertible transformation and a permutation (Hyvärinen and Pajunen, 1999). We formulate our structural sparsity condition below and present the identifiability result. The proof is provided in Appx. A.1.

Theorem 1. *Assume*

i The observed data is sampled from a nonlinear ICA model as defined in Eqs. (1) and (2).

ii Mixing function $\mathbf{f}$ is invertible and smooth. Its inverse is also smooth.

iii Denote by $\mathcal{F}$ the support of (all possible values of) $\mathbf{J_f}(\mathbf{s})$. For all $i \in \{1, \ldots, n\}$, there exist $\{\mathbf{s}^{(\ell)}\}_{\ell=1}^{|\mathcal{F}_{i,:}|}$ *such that* $\mathrm{span}\{\mathbf{J_f}(\mathbf{s}^{(\ell)})_{i,:}\}_{\ell=1}^{|\mathcal{F}_{i,:}|} = \mathbb{R}^n_{\mathcal{F}_{i,:}}$.

iv $|\mathrm{supp}(\mathbf{J_{\hat{f}}}(\hat{\mathbf{s}}))| \leq |\mathrm{supp}(\mathbf{J_f}(\mathbf{s}))|$.

v (Structural Sparsity) For all $k \in \{1, \ldots, n\}$, there exists $\mathcal{C}_k$ such that

$$\bigcap_{i \in \mathcal{C}_k} \mathrm{supp}(\mathbf{J_f}(\mathbf{s})_{i,:}) = \{k\}.$$

Then $\mathbf{h} := \hat{\mathbf{f}}^{-1} \circ \mathbf{f}$ is a composition of a component-wise invertible transformation and a permutation.

Assumption *iii* in a generic sense rules out a set of specific parameters where the Jacobian is partially constant and can be contained in a proper affine subspace aligned by its support (i.e., $\mathbb{R}^n_{\mathcal{G}_{i,:}}$). Assumption *iv* corresponds to incorporating a suitable sparsity regularization term into the estimating process with no restriction on the ground truth. It helps to find the estimated mixing function with the minimal L_0 norm among all functions that allow the model to perfectly fit the true distribution of sources. Thus, our discussion will focus on the assumption of structural sparsity, which is the core of our theory.

We start with the intuition of the proposed direction on exploiting sparsity on the mixing process for identifiability. As discussed in previous works (Hyvärinen and Pajunen, 1999), there exist infinitely many ways to preserve the independence among variables after mixing them up, such as Darmois construction (Darmois, 1951) and measure-preserving automorphism (MPA) (Hyvärinen and Pajunen, 1999). Besides, sources with non-Gaussian priors could be transferred to marginally independent Gaussians by trivial point-wise transformations (Hyvärinen et al., 2019). Thus, the existing strategy of exploiting independence and non-Gaussianity for identification fails in the nonlinear case. However, if the true structure is sparse enough, any alternative solution that produces any indeterminacy beyond component-wise transformations and permutations (e.g., mixtures or rotations) may correspond to a denser structure. In the light of that, instead of restricting the functional class (e.g., post nonlinear models (Taleb and Jutten, 1999) or conformal maps (Hyvärinen and Pajunen, 1999)) or introducing auxiliary variables for extra structural dependencies between latent sources (Hyvärinen et al., 2019; Lachapelle et al., 2022), we leverage the sparsity pattern, i.e., the support of the Jacobian of the mixing function, to identify the sources.

The structural sparsity assumption (Assumption *v* in Thm. 1) implies that, for every latent source $\mathbf{s}_i$, there exists a set of observed variable(s) such that $\mathbf{s}_i$ is the only latent source that participates in the generation of all observed variable(s) in the set. Graphically, for every latent source $\mathbf{s}_i$, there exists a set of observed variable(s) such that the intersection of their/its parent(s) is $\mathbf{s}_i$ (e.g., for $\mathbf{s}_1$ in Fig. 1, there exist $\mathbf{x}_1$ and $\mathbf{x}_4$ such that the intersection of their parents is $\mathbf{s}_1$.). This encourages the connections to be sparse enough to make the disentanglement of each source based on structural conditions possible. Together with the regularization of sparsity during the estimating process (e.g., minimizing the L_0 norm of the Jacobian as Assumption *iv*) and other mild conditions, Assumption *v* illustrates a graphical pattern of nonlinear ICA models that could be identified by exploiting structural sparsity. It indicates a structural criterion of the sparsity that is needed for identifiability. Beyond that, structural sparsity is also of great practical interest for the interpretability of results (Zhang et al., 2009). Meanwhile,

Figure 1: The structural sparsity assumption of Thm. 1, where the matrix represents $\mathrm{supp}(\mathbf{J_f}(\mathbf{s}))$.

in the literature of causality, various versions of Occam's razor (e.g., faithfulness (Spirtes et al., 2000), minimality principle (Zhang, 2013), and frugality (Raskutti and Uhler, 2018; Forster et al., 2020)) are also fundamental to the identifiability of the underlying causal structure.

With structural sparsity, Thm. 1 shows the identifiability of nonlinear ICA while maintaining the standard mutually marginal independence assumption of sources. This is consistent with the original setting of ICA and plays an important role in a more general range of unsupervised learning tasks, compared to previous works relying on conditional independence given auxiliary variables. Moreover, one may modify Thm. 1 to consider the generating process based on auxiliary variables by imposing similar restrictions on the mixing process. Different from previous works (Hyvärinen et al., 2019; Yang et al., 2022), this extension removes common restrictions on the required auxiliary variables, such as the number of distinct values (e.g., class labels), non-proportional variances, or sufficient variability. Therefore, although our main result (Thm. 1) focuses on nonlinear ICA with unconditional priors, it also provides potential insight into improving the flexibility of utilizing the auxiliary variable when it is available. As a trade-off, the additional assumptions on the sparsity might limit its usage.

Besides, under additional assumptions, we could reduce the indeterminacy in Thm. 1 and further identify sources up to a component-wise *linear* transformation. Most works in nonlinear ICA focus on the identifiability up to a component-wise invertible transformation and a permutation (Hyvärinen and Morioka, 2016; Khemakhem et al., 2020), which is also called *permutation-identifiability* in the literature of disentanglement (Lachapelle et al., 2022). This indeterminacy is trivial compared to the fundamental nonuniqueness of nonlinear ICA and is analogous to the indeterminacy involving rescaling and permutation in linear ICA (Hyvärinen and Pajunen, 1999). Recently, Yang et al. (2022) provided an identifiability result based on auxiliary variables that further reduces the indeterminacy of the component-wise nonlinear transformation to a linear one. Inspired by it but without auxiliary variables, we show conditions for the identifiability of nonlinear ICA with unconditional priors up to a component-wise linear transformation, with the proof in Appx. A.2. The reduced indeterminacy may give rise to a more informative disentanglement and open up the possibility for further improving the quality of recovery with only element-wise linear operations, such as the Hadamard product.

Corollary 1. *Assume*

  i The observed data is sampled from a nonlinear ICA model as defined in Eqs. (1) and (2).

  ii The function $\mathbf{h} := \hat{\mathbf{f}}^{-1} \circ \mathbf{f}$ is a composition of a component-wise invertible transformation and a permutation.

  iii The mixing function $\mathbf{f}$ is volume-preserving.

  iv The source distribution $p_{\mathbf{s}}(\mathbf{s})$ is a factorial multivariate Gaussian.

  Then $\mathbf{h} := \hat{\mathbf{f}}^{-1} \circ \mathbf{f}$ is a composition of a component-wise linear transformation and a permutation.

3.2 Removing Rotation Indeterminacy with Structural Sparsity

Rotation indeterminacy is one of the major obstacles to the identifiability of ICA. Linear ICA exploits the maximization of the non-Gaussianity of the estimated sources (e.g., Kurtois (Hyvärinen and Oja,

1997)). As a direct result, the typical assumption is that at most one of the sources can have Gaussian distribution. In contrast, for the nonlinear case, a trivial point-wise function could transform sources to have any marginal distribution including Gaussian thus invalidating the effect of non-Gaussianity. Therefore, removing rotation indeterminacy remains critical for the identifiability in the nonlinear case. For instance, "rotated-Gaussian" MPA produces nonuniqueness in nonlinear ICA due to rotation indeterminacy (Hyvärinen and Pajunen, 1999). It first maps the distribution to an isotropic Gaussian, then applies a rotation and maps it back without affecting its original distribution. To address it for identifiability, Yang et al. (2022) assume that there do not exist two conditionally independent Gaussians such that their variances are proportional.

In the previous section, we have introduced the structural sparsity for the identifiability of nonlinear ICA. One may wonder whether it straightforwardly leads to the identifiability of linear ICA with Gaussian sources (Gaussian ICA), which was previously perceived to be impossible due to its rotation indeterminacy. The idea is simple–any rotation of the true mixing matrix $\mathbf{A}$ will be less sparse if $\mathbf{A}$ satisfies the proposed structural sparsity assumption. While the joint distribution of Gaussian sources stays invariant across rotations, the sparsity of the mixing matrix (i.e., L_0 norm) keeps changing. We first consider Assumption *v* in Thm. 1 for Gaussian ICA as an extension. Because the Jacobian of linear mixing function is fixed, Assumption *iii* of Thm. 1 does not directly hold in the linear Gaussian case. As a result, we cannot directly apply Thm. 1 here and therefore propose some alternative conditions:

Proposition 1. *Assume*

i *The observed data is sampled from a linear ICA model defined in Eqs. (1) and (3) with Gaussian sources.*

ii *Mixing matrix $\mathbf{A}$ is invertible.*

iii *There exists a matrix $\hat{\mathbf{A}}$ such that $\forall j \in \text{supp}(\mathbf{A})_{i,:},\ \text{supp}\ (\hat{\mathbf{A}}\mathbf{A}^{-1})_{j,:} \in \mathbb{R}^n_{\text{supp}(\hat{\mathbf{A}})_{i,:}}.$*

iv $|\text{supp}(\hat{\mathbf{A}})| \leq |\text{supp}(\mathbf{A})|.$

v *(Structural Sparsity) For all $k \in \{1, \ldots, n\}$, there exists $\mathcal{C}_k$ such that*

$$\bigcap_{i \in \mathcal{C}_k} \text{supp}(\mathbf{A}_{i,:}) = \{k\}.$$

Then $\hat{\mathbf{A}} = \mathbf{ADP}$, where $\mathbf{D}$ is a diagonal matrix and $\mathbf{P}$ is a column permutation matrix.

The proof is provided in Appx. A.3. Similar to Assumption *iii* of Thm. 1, Assumption *iii* of Prop. 1 excludes a specific set of parameters that makes structure-based identification ill-posed.

The underdetermined case. For the full identifiability of nonlinear ICA, it is noteworthy that the invertibility of the mixing function is technically essential in our result (Thm. 1). This invertibility implies that the number of observed variables is equal to that of sources. Although that setting is rather common in the literature, especially for the theory of nonlinear cases (Hyvärinen and Pajunen, 1999; Hyvärinen et al., 2019), underdetermined ICA (i.e., there are more observed variables), may be of more practical interest and more challenging (Kim and Yoo, 2004). Therefore, we develop an alternative structural sparsity condition for the underdetermined case. As a trade-off, in the nonlinear case, the structural condition that removes the requirement of invertibility could only deal with the rotation indeterminacy. Thus we start from Gaussian ICA to introduce the condition.

Definition 4. *Given a matrix $\mathbf{S} \in \mathbb{R}^{m \times n}$, we define the function* overlap $: \mathbb{R}^{m \times n} \to \{0, 1\}^{m \times n}$ *as*

$$(\text{overlap}(\mathbf{S}))_{ij} = \begin{cases} 1 & \text{if } \mathbf{S}_{ij} = 1 \text{ and it is not the only nonzero entry in } \mathbf{S}_{i,:} \\ 0 & \text{otherwise.} \end{cases}$$

Theorem 2. *Assume*

i *The observed data is sampled from a linear ICA model defined in Eqs. (1) and (3) with Gaussian sources. Differently, the number of observed variables (denoted as m) could be larger than that of the sources n, i.e., $m \geq n$.*

ii *The nonzero coefficients of matrix $\mathbf{A}$ are randomly drawn from a distribution that is absolutely continuous with respect to Lebesgue measure.*

219   *iii* $|\operatorname{supp}(\hat{\mathbf{A}})| \le |\operatorname{supp}(\mathbf{A})|$.

iv (Structural Sparsity) Given $\mathcal{C} \subseteq \{1, 2, \ldots, n\}$ where $|\mathcal{C}| > 1$, let $\mathbf{A}_\mathcal{C} \in \mathbb{R}^{m \times |\mathcal{C}|}$ represents a submatrix of $\mathbf{A} \in \mathbb{R}^{m \times n}$ consisting of columns with indices $\mathcal{C}$. Then, for all $k \in \mathcal{C}$, we have

$$\left| \bigcup_{k' \in \mathcal{C}} \operatorname{supp}(\mathbf{A}_{k'}) \right| - \operatorname{rank}(\operatorname{overlap}(\mathbf{A}_\mathcal{C})) > |\operatorname{supp}(\mathbf{A}_k)|.$$

220
221 *Then $\hat{\mathbf{A}} = \mathbf{ADP}$, where $\mathbf{D}$ is a diagonal matrix and $\mathbf{P}$ is a column permutation matrix.*

222 The proof is provided in Appx. A.4. Intuitively, Assumption *iv* in Thm. 2
223 encourages the variability between the influences of each source. For each
224 source, its influence on the observations should be as distinctive as possible,
225 which is correlated with the sparsity of the mixing matrix. Fig. 2 is an illus-
226 tration of the assumption. In that case, $k = 1$ and $\mathcal{C} = \{1, 2, 3\}$. Let $\mathbf{A}_\mathcal{C}$
227 denotes the submatrix shown in the figure, where $\left| \bigcup_{k' \in \mathcal{C}} \operatorname{supp}(\mathbf{A}_{k'}) \right| = 7$. Blue
228 dotted square denotes $\operatorname{overlap}(\mathbf{A}_\mathcal{C})$ and we have $\operatorname{rank}(\operatorname{overlap}(\mathbf{A}_\mathcal{C})) = 2$. Thus,
229 $\left| \bigcup_{k' \in \mathcal{C}} \operatorname{supp}(\mathbf{A}_{k'}) \right| - \operatorname{rank}(\operatorname{overlap}(\mathbf{A}_\mathcal{C})) = 5$ (the number of black dots), which

Figure 2: Assumption *iv* of Thm. 2.

230 is larger than $|\operatorname{supp}(\mathbf{A}_1)| = 4$. For the nonlinear case, we prove that it could
231 help to distinguish spurious solutions due to the rotation indeterminacy, i.e., the
232 "rotated-Gaussian" MPA (Defn. 2.5 in Gresele et al. (2021)). Gresele et al. (2021) prove that this class
233 of MPAs could be ruled out with assumptions of conformal maps, non-Gaussianity and orthogonality
234 of the Jacobian of the mixing function. Differently, we address it with the proposed structural sparsity
235 condition (Assumption *iv* in Thm. 2). The corresponding theorem for the nonlinear case, with its
236 proof given in Appx. A.5, is as follows:

237 **Theorem 3.** *Given a nonlinear ICA model defined in Eqs. (1) and (2), where $\mathbf{f}$ is the true mixing*
238 *function. Consider $\hat{\mathbf{f}} = \mathbf{f} \circ \mathbf{G}^{-1} \circ \mathbf{U} \circ \mathbf{G}$, where $\mathbf{G}$ denotes an invertible Gaussianization[1] that*
239 *maps the distribution to an standard isotropic (rotation-invariant) Gaussian, $\mathbf{U}$ denotes a rotation,*
240 *and $\mathbf{G}^{-1}$ maps the distribution back to that before applying $\hat{\mathbf{U}} \circ \mathbf{G}$. If Assumptions ii, iii and iv of*
241 *Thm. 2 are satisfied by replacing $\mathbf{A}$ with $\mathbf{J}_\mathbf{f}(\mathbf{s})$ and $\hat{\mathbf{A}}$ with $\mathbf{J}_{\hat{\mathbf{f}}}(\mathbf{s})$, then function $\mathbf{h} := \hat{\mathbf{f}}^{-1} \circ \mathbf{f}$ is a*
242 *composition of a component-wise invertible transformation and a permutation.*

243 Besides removing rotation indeterminacy, Thm. 3 also provides an extra insight for the full identifia-
244 bility of nonlinear ICA, because all previous works assume that the nonlinear mixing function must be
245 invertible (Hyvärinen and Morioka, 2016; Yang et al., 2022) even with the help of auxiliary variables.

4   Identifiability with Independent Influences

247 In the previous section, we have presented theoretical results based on conditions of structural sparsity.
248 Alternatively, we show in this section that sparsity conditions on the support might not be necessary in
249 some specific scenarios, and that one can leverage the independence (in a non-statistical sense) among
250 the influences from different sources to the observations to achieve identifiability of nonlinear ICA.
251 This implies sparse interactions between the individual influencing processes and could be viewed as
252 an analogous concept of "sparsity" w.r.t. influences. We show that without any kind of auxiliary vari-
253 ables, the full identifiability of nonlinear ICA can also be provided given the condition of independent
254 influences, thus complementing the results shown above from a different view of "sparsity".

255 **Nonlinear ICA with unconditional priors: Independent Influences.** We now specify the setting
256 of the required estimating process. We sample the estimated sources $\hat{\mathbf{s}}$ from a factorial multivariate
257 Gaussian:

$$p_{\hat{\mathbf{s}}}(\hat{\mathbf{s}}) = \prod_{i=1}^{n} \frac{1}{Z_i} \exp\left(-\theta_{i,1}\hat{s}_i - \theta_{i,2}\hat{s}_i^2\right), \tag{7}$$

---

[1]One example is described in (Gresele et al., 2021), i.e., a composition of the element-wise CDFs of a smooth factorised density and a Gaussian, respectively.

where $Z_i > 0$ is a constant. The sufficient statistics $\theta_{i,1} = -\frac{\mu_i}{\sigma_i^2}$ and $\theta_{i,2} = \frac{1}{2\sigma_i^2}$ are assumed to be linearly independent. We constraint the variances σ_i^2 to be distinct without loss of generality.

Theorem 4. *Consider the model defined in Eqs.* (1)*,* (2)*, and* (7)*. Assume*

   i *(Independent Influences): The influence of each source on the observed variables is independent of each other, i.e., each partial derivative $\partial \mathbf{f}/\partial s_i$ is independent of the other sources and their influences in a non-statistical sense.*

   ii *The mixing function $\mathbf{f}$ and its inverse are twice differentiable.*

   iii *The Jacobian determinant of mixing function can be factorized as $\det(\mathbf{J_f}(\mathbf{s})) = \prod_{i=1}^{n} y_i(s_i)$, where y_i is a function that depends only on s_i. Note that volume-preserving transformation is a special case when $y_i(s_i) = 1, i = 1, \ldots, n$.*

   iv *During estimation, the columns of the Jacobian of the estimated unmixing function are regularized to be mutually orthogonal and with equal euclidean norm.*

Then $\mathbf{h} := \hat{\mathbf{f}}^{-1} \circ \mathbf{f}$ is a composition of a component-wise invertible transformation and a permutation.

The proof is provided in Appx. A.6. Note that some part of the proof technique is inspired by or based on Lemma 1 in Yang et al. (2022). Assumption *iv* is a regularization during estimation and thus puts no restriction on the true generating process. We achieve it by optimizing an objective function with a designed regularization term, of which the property is as follows (proof in Appx. A.7):

Proposition 2. *The following inequality holds*

$$n \log \left(\frac{1}{n} \sum_{i=1}^{n} \left\| \frac{\partial \hat{\mathbf{f}}^{-1}}{\partial x_i} \right\|_2 \right) - \log \left| \det(\mathbf{J}_{\hat{\mathbf{f}}^{-1}}(\mathbf{x})) \right| \geq 0, \tag{8}$$

with equality iff. $\mathbf{J}_{\hat{\mathbf{f}}^{-1}}(\mathbf{x}) = \mathbf{O}(\mathbf{x})\lambda(\mathbf{x})$, where $\mathbf{O}(\mathbf{x})$ is an orthogonal matrix and $\lambda(\mathbf{x})$ is a scalar.

Regarding the assumption of independent influences, to build intuition, one can consider a filmmaking process in a safari park, of which the main characters are wild animals. The task of the boom operators here is to record the sound of animals. As there are many animals and microphones, the mixing (recording) process is highly dependent on the relative positions between them. Thus, the animal (source s_i), loosely speaking, influences the recording (mixing function $\mathbf{f}$) through its position. The moving direction and speed of them can be loosely interpreted as the partial derivative w.r.t. the i-th source, i.e. $\partial \mathbf{f}/\partial s_i$. Assuming that the wild animals are not cooperative enough to fine-tune their positions and speeds for a better recording and the safari park is not crowded, the influences of animals on the recording are generated independently by them in the sense that they are not affected by the others while moving. As a result of that generating process, the column vectors of the Jacobian of the mixing function are uncorrelated with each other and the partial derivative w.r.t. to each source is independent of other sources. Other practical scenarios include fields that adopt independent influences in process like orthogonal coordinate transformations (Gresele et al., 2021), such as dynamic control (Mistry et al., 2010) and structural geology (De Paor, 1983). It is worth noting that this assumption of independent influences is different from the orthogonality condition proposed in (Gresele et al., 2021). For instance, if the sample mean is not zero, uncorrelatedness differs from orthogonality. And we argue that uncorrelatedness as a specific way to achieve independence might be more sensible as a condition on the influences. Besides, in (Gresele et al., 2021), orthogonality, together with conformal map and the others, are proposed as conditions to rule out two specific types of spurious solutions (i.e., Darmois construction (Darmois, 1951) and "rotated-Gaussian" MPAs (Locatello et al., 2019)) without an identifiability result, although its empirical results are encouraging.

Having said that, a violation of the assumption of independent influences could be ascribed to a deliberate global adjustment of sources. For example, cinema audio systems are carefully adjusted to achieve a homogeneous sound effect on every audience. The position and orientation of each speaker are fine-tuned according to the others. In this case, it may be difficult to distinguish the influences from different speakers because of the fine-tuning. This may lead to a high degree of multicollinearity across the columns of Jacobian, thus violating the assumption of independent influences.

Regarding Assumption *iii* in Thm. 3, we first note that mixing functions with factorial Jacobian determinants are of a much wider range as compared to component-wise transformations. To see this,

| (a) *SS.* | (b) *II.* | (c) *VP.* | (d) *Base.* |

Figure 3: Pearson correlation matrices between the ground-truth and the recovered sources.

a straightforward example is the volume-preserving transformation, which has been widely adopted in generative models (Zhang et al., 2021). In fact, all transformations with constant Jacobian determinant can be factorized w.r.t. the sources, which, generally speaking, are not component-wise. Besides, consistent with the empirical results in Yang et al. (2022), there exists other non-volume-preserving transformations with factorial Jacobian determinant [2]. Moreover, the volume-preserving assumption has been demonstrated to be helpful to weaken the requirement of auxiliary variables, specifically by reducing the number of required labels (Sorrenson et al., 2020; Yang et al., 2022). Thm. 4 takes this further by imposing a weaker version, i.e., Assumption (iii), which, together with the other constraints, removes the need for any auxiliary variable.

5   Experiments

To validate the proposed theory of the identifiability of nonlinear ICA with unconditional priors, we conduct experiments based on our main condition, i.e., structural sparsity (Thm. 1), as well as the condition of the independent influences (Thm. 4).

Setup.   For the required regularization during estimation, we consider a regularized maximum-likelihood approach with the following objective: $\mathcal{L}(\hat{\mathbf{f}}^{-1}; \mathbf{x}) = \mathbb{E}_{\mathbf{x}}\left[\log p_{\hat{\mathbf{f}}^{-1}}(\mathbf{x}) - \lambda \mathbf{R}(\hat{\mathbf{f}}^{-1}, \mathbf{x})\right]$, where λ is a regularization parameter and $\mathbf{R}(\hat{\mathbf{f}}^{-1}, \mathbf{x})$ is the regularization term. For Thm. 1, we use L_1 norm as an approximation of the L_0 norm for efficiency (Donoho and Elad, 2003), therefore $\mathbf{R}(\hat{\mathbf{f}}^{-1}, \mathbf{x}) := \|\mathbf{J}_{\hat{\mathbf{f}}^{-1}}(\mathbf{x})\|_1$; for Thm. 4, $\mathbf{R}(\hat{\mathbf{f}}^{-1}, \mathbf{x})$ corresponds to LHS of Eq. (8). Following (Sorrenson et al., 2020; Yang et al., 2022), we train a GIN (Sorrenson et al., 2020) to maximize the the objective function $\mathcal{L}(\hat{\mathbf{f}}^{-1}; \mathbf{x})$.

Ablation study.   We conduct an ablation study to verify the necessity of the proposed assumptions. Specifically, we focus on the following models that correspond to different assumptions: *(SS)* The assumption of structural sparsity, as well as other assumptions in Thm. 1, are satisfied; *(II)* The assumption of independent influences, as well as other assumptions in Thm. 4, are satisfied; *(VP)* Compared to *II*, only the assumption of independent influences is violated while the other assumptions (e.g., factorizable Jacobian determinant) are still satisfied; *(Base)* The vanilla baseline. Compared to *VP*, the (un)mixing function is not restricted to having factorizable Jacobian determinants. The data are generated according to the required assumptions.

Figure 4: Ablation study.

We also conduct comparison between the assumption of independent influences and orthogonality in (Gresele et al., 2021), which are presented in Appx. B together with experimental settings. Results for each model are summarised in Fig. 4. For evaluation, we use the mean correlation coefficient (MCC) between the true sources and the estimated ones (Hyvärinen and Morioka, 2016). One could observe that when the proposed assumptions are fully satisfied (*SS* or *II*), our model achieves the highest MCC on average. This indicates that it is actually possible to identify sources from highly nonlinear mixtures up to trivial indeterminacies only based on restrictions on the mixing process. Moreover, regarding Thm. 4, the importance of Assumption *iii* is supported by the higher performance of *VP* compared to that of *Base*. The visualization of the Pearson correlation matrices is shown in Fig. 3.

---

[2]A toy example: the Jacobian determinant of mixing function w.r.t. $\left(x_1 = \frac{as_1s_2}{a+bs_2}, x_2 = \frac{bs_1}{b+as_2}\right)$ is abs_1, where $a, b \neq 0$ are some constants.

Stability. To study the stability of the performance of identification w.r.t. different datasets varying the number of sources n. We test the model *SS* (Thm. 1) with different n. Visually, we find that *SS* consistently outperforms *Base* (Fig. 5). Meanwhile, when the number of sources increases, one could observe that the MCC of *Base* decreases while that of *SS* stays stable. The stable empirical performance further validates our theoretical claims about the identifiability with structural sparsity.

Figure 5: MCC w.r.t. different number of sources.

Image dataset. To study how reasonable the proposed theories are w.r.t. the practical generating process of observational data, we conduct experiments on the "Triangles" image dataset (Yang et al., 2022). The process of generating this dataset mimics the process of drawing triangles by humans: i) First, we sample the elements needed for humans to draw a monochrome triangle (i.e., rotation, width, height and grey level) from a factorial multivariate Gaussian distribution. Different from (Yang et al., 2022), we always sample from a single distribution in order to guarantee that all priors are unconditionally independent; ii) Then, for each pixel, we decide whether it locates inside the triangle based on the sampled

Figure 6: Identification results on Triangles. Each row represents a source identified by our model, with it varying from -2 to +2 SDs to illustrate its influence. The rightmost column is a heat map given by the absolute pixel difference between -1 and +1 SDs. Visually, the four rows correspond to rotation, width, height and gray level, respectively.

elements and assign its gray level accordingly. Therefore, the process is similar to human drawing triangles. Even though each image is generated from these semantic elements, the true generating process and sources are still unknown (e.g., a pixel could be (indirectly) influenced by multiple elements in a complicated way). We apply GIN with sparsity regularization as the estimating method. The visualization of the identified sources (Fig. 6) indicates that our conditions may hold in practice.

6   Discussion

We provide identifiability results for nonlinear ICA with unconditional priors, which serve as one of the first steps to solve a long-standing problem in unsupervised learning. In particular, we prove that the i.i.d. latent sources can be recovered up to a component-wise invertible transformation and a permutation with only conditions on the nonlinear mixing process (e.g., structural sparsity). Therefore it stays closer to the original notion of ICA that is based on the marginal independence assumption of latent sources, while previous works rely on conditional independence on auxiliary variables as weak supervision or inductive bias. Besides, by removing rotation indeterminacy, structural sparsity benefits the identifiability of Gaussian ICA as well, which was also thought to be unsolvable before. Moreover, the results on the underdetermined case are of great practical interest and introduce insight for extending identifiable nonlinear ICA to general real-world settings.

Our results on images illustrate the validity of the proposed conditions in practical data generating processes. In spite of this, it is possible that part of them is violated in several specific scenarios as discussed before. For example, the structural sparsity conditions do not apply to fully-connected structures, though the practical significance of such cases may be compromised by the lack of interpretability. As a complementary solution, we formulate the independent influences condition, which does not rely on the sparse structure of supports. While arguably natural in general, it could still be violated due to a deliberate global adjustment of sources for homogeneity. We argue that this is inevitably a trade-off between introducing auxiliary variables and imposing restrictions on the mixing process to achieve the identifiability, whose practical use depends on the scenario and information available. Future work includes further generalizing and validating our theory.

References

P. Comon. Independent component analysis, a new concept? *Signal processing*, 36(3):287–314, 1994.

G. Darmois. Analyse des liaisons de probabilité. In *Proc. Int. Stat. Conferences 1947*, page 231, 1951.

D. G. De Paor. Orthographic analysis of geological structures—I. deformation theory. *Journal of Structural Geology*, 5(3-4):255–277, 1983.

L. Dinh, J. Sohl-Dickstein, and S. Bengio. Density estimation using Real NVP. *arXiv preprint arXiv:1605.08803*, 2016.

D. L. Donoho and M. Elad. Optimally sparse representation in general (nonorthogonal) dictionaries via L_1 minimization. *Proceedings of the National Academy of Sciences*, 100(5):2197–2202, 2003.

H. Flanders. Liouville's theorem on conformal mapping. *Journal of Mathematics and Mechanics*, 15 (1):157–161, 1966.

M. Forster, G. Raskutti, R. Stern, and N. Weinberger. The frugal inference of causal relations. *The British Journal for the Philosophy of Science*, 2020.

L. Gresele, J. von Kügelgen, V. Stimper, B. Schölkopf, and M. Besserve. Independent mechanism analysis, a new concept? *Advances in Neural Information Processing Systems*, 2021.

H. Hälvä and A. Hyvärinen. Hidden markov nonlinear ICA: Unsupervised learning from nonstationary time series. In *Conference on Uncertainty in Artificial Intelligence*, pages 939–948. PMLR, 2020.

H. Hälvä, S. Le Corff, L. Lehéricy, J. So, Y. Zhu, E. Gassiat, and A. Hyvärinen. Disentangling identifiable features from noisy data with structured nonlinear ICA. *Advances in Neural Information Processing Systems*, 34, 2021.

A. Hyvärinen and H. Morioka. Unsupervised feature extraction by time-contrastive learning and nonlinear ICA. *Advances in Neural Information Processing Systems*, 29:3765–3773, 2016.

A. Hyvärinen and H. Morioka. Nonlinear ICA of temporally dependent stationary sources. In *International Conference on Artificial Intelligence and Statistics*, pages 460–469. PMLR, 2017.

A. Hyvärinen and E. Oja. A fast fixed-point algorithm for independent component analysis. *Neural computation*, 9(7):1483–1492, 1997.

A. Hyvärinen and P. Pajunen. Nonlinear independent component analysis: Existence and uniqueness results. *Neural networks*, 12(3):429–439, 1999.

A. Hyvärinen, H. Sasaki, and R. Turner. Nonlinear ICA using auxiliary variables and generalized contrastive learning. In *International Conference on Artificial Intelligence and Statistics*, pages 859–868. PMLR, 2019.

I. Khemakhem, D. Kingma, R. Monti, and A. Hyvärinen. Variational autoencoders and nonlinear ICA: A unifying framework. In *International Conference on Artificial Intelligence and Statistics*, pages 2207–2217. PMLR, 2020.

S. G. Kim and C. D. Yoo. Underdetermined independent component analysis by data generation. In *International Conference on Independent Component Analysis and Signal Separation*, pages 445–452. Springer, 2004.

D. P. Kingma and P. Dhariwal. Glow: Generative flow with invertible 1x1 convolutions. *Advances in neural information processing systems*, 31, 2018.

S. Lachapelle, P. R. López, Y. Sharma, K. Everett, R. L. Priol, A. Lacoste, and S. Lacoste-Julien. Disentanglement via mechanism sparsity regularization: A new principle for nonlinear ICA. *Conference on Causal Learning and Reasoning*, 2022.

F. Locatello, S. Bauer, M. Lucic, G. Raetsch, S. Gelly, B. Schölkopf, and O. Bachem. Challenging common assumptions in the unsupervised learning of disentangled representations. In *international conference on machine learning*, pages 4114–4124. PMLR, 2019.

M. Mistry, J. Buchli, and S. Schaal. Inverse dynamics control of floating base systems using orthogonal decomposition. In *2010 IEEE international conference on robotics and automation*, pages 3406–3412. IEEE, 2010.

E. Oja. Unsupervised learning in neural computation. *Theoretical computer science*, 287(1):187–207, 2002.

G. Raskutti and C. Uhler. Learning directed acyclic graph models based on sparsest permutations. *Stat*, 7(1):e183, 2018.

P. Sorrenson, C. Rother, and U. Köthe. Disentanglement by nonlinear ICA with general incompressible-flow networks (GIN). *arXiv preprint arXiv:2001.04872*, 2020.

P. Spirtes, C. N. Glymour, R. Scheines, and D. Heckerman. *Causation, prediction, and search.* MIT press, 2000.

A. Taleb and C. Jutten. Source separation in post-nonlinear mixtures. *IEEE Transactions on signal Processing*, 47(10):2807–2820, 1999.

L. N. Trefethen and D. Bau III. *Numerical linear algebra*, volume 50. Siam, 1997.

M. Willetts and B. Paige. I don't need **u**: Identifiable non-linear ICA without side information. *arXiv preprint arXiv:2106.05238*, 2021.

X. Yang, Y. Wang, J. Sun, X. Zhang, S. Zhang, Z. Li, and J. Yan. Nonlinear ICA using volume-preserving transformations. In *International Conference on Learning Representations*, 2022.

W. Yao, Y. Sun, A. Ho, C. Sun, and K. Zhang. Learning temporally causal latent processes from general temporal data. *arXiv preprint arXiv:2110.05428*, 2021.

J. Zhang. A comparison of three occam's razors for markovian causal models. *The British journal for the philosophy of science*, 64(2):423–448, 2013.

K. Zhang, H. Peng, L. Chan, and A. Hyvärinen. ICA with sparse connections: Revisited. In *International Conference on Independent Component Analysis and Signal Separation*, pages 195–202. Springer, 2009.

S. Zhang, C. Zhang, N. Kang, and Z. Li. iVPF: Numerical invertible volume preserving flow for efficient lossless compression. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pages 620–629, 2021.