Globally Convergent Policy Search for Output Estimation

Abstract

We introduce the first direct policy search algorithm which provably converges to the globally optimal *dynamic* filter for the classical problem of predicting the outputs of a linear dynamical system, given noisy, partial observations. Despite the ubiquity of partial observability in practice, theoretical guarantees for direct policy search algorithms, one of the backbones of modern reinforcement learning, have proven difficult to achieve. This is primarily due to the degeneracies which arise when optimizing over filters that maintain an internal state.

In this paper, we provide a new perspective on this challenging problem based on the notion of *informativity*, which intuitively requires that all components of a filter's internal state are representative of the true state of the underlying dynamical system. We show that informativity overcomes the aforementioned degeneracy. Specifically, we propose a *regularizer* which explicitly enforces informativity, and establish that gradient descent on this regularized objective – combined with a "reconditioning step" – converges to the globally optimal cost at a $\mathcal{O}(1/T)$ rate.

1 Introduction

Data used for prediction and control of real world dynamical systems is almost always noisy and incomplete (partially observed). Sensors and other measurement procedures inevitably introduce errors into the datasets, so designing reliable learning algorithms for these noisy or partially observed domains requires confronting fundamental questions of disturbance filtering and state estimation. Despite the ubiquity of partial observation in practice, these concerns are often underexplored in modern analyses of learning for control that assume perfect observations of the underlying dynamics.

In this work, we study the output estimation (OE) problem or learning to predict in partially observed linear dynamical systems. The output estimation problem is one of the most fundamental problems in theoretical statistics and learning theory. Both in theory and in practice, advances in predicting partially observed linear systems have led to successes in a variety of areas from controls to biology and economics, (c.f. e.g. Athans [1974], Lillacci and Khammash [2010], Gautier and Poignet [2001]). We revisit this classical problem from a modern optimization perspective, and study the possibility of learning the optimal predictor via model-free procedures and direct policy search.

Relative to model-based procedures, which first estimate the underlying dynamics and then return a policy by solving an optimization problem using the estimated model, model-free methods offer several potential advantages. For instance, direct policy search allows one to easily specify the complexity of the policy class over which one searches. In addition, model-free policy search optimizes for performance directly on the true system, rather than an approximate model. As such, there is no gap between the model used for synthesis and the system on which the policy is deployed.

In light of these advantages, there has recently been significant interest from both theoreticians and practitioners in understanding the foundations of model-free decision making. So far, this attention

has focused mostly on problems with *full-state* observation such as the linear quadratic regulator (`LQR`) or fully-observed Markov Decision Processes (`MDPs`), for which optimal policies are *static* (without memory). Progress on partially observed problems has been complicated by the difficulties associated with optimizing over *dynamic* policies (with memory) that maintain internal state to summarize past observations. In this paper, we provide the first policy search algorithm which provably converges to the globally optimal filter for the `OE` problem, and shed new light on the intricacies of the underlying optimization landscape.

The Output Estimation problem. We study one of the simplest and most basic problems with partial observability: the *output estimation* (`OE`) problem. In brief, the goal is to search for a predictor of the output $\mathbf{z}(t)$ of a linear dynamical system given partial measurements $\mathbf{y}(t)$. For the *true system* with states $\mathbf{x}(t)$ and dynamics that evolve according to,

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbf{x}(t) = \mathbf{A}\mathbf{x}(t) + \mathbf{w}(t), \quad \mathbf{y}(t) = \mathbf{C}\mathbf{x}(t) + \mathbf{v}(t), \quad \mathbf{z}(t) = \mathbf{G}\mathbf{x}(t), \quad \mathbf{x}(0) = 0,$$
$$\mathbf{w}(t) \overset{\text{i.i.d}}{\sim} \mathcal{N}(0, \mathbf{W}_1), \quad \mathbf{v}(t) \overset{\text{i.i.d}}{\sim} \mathcal{N}(0, \mathbf{W}_2), \tag{1.1}$$

the goal is to find the parameters $\mathsf{K} = (\mathbf{A}_\mathsf{K}, \mathbf{B}_\mathsf{K}, \mathbf{C}_\mathsf{K})$ of the *filter* (interchangably, *policy*),

$$\frac{\mathrm{d}}{\mathrm{d}t}\hat{\mathbf{x}}(t) = \mathbf{A}_\mathsf{K}\hat{\mathbf{x}}(t) + \mathbf{B}_\mathsf{K}\mathbf{y}(t), \quad \hat{\mathbf{z}}(t) = \mathbf{C}_\mathsf{K}\hat{\mathbf{x}}(t), \tag{1.2}$$

that minimizes the steady-state prediction error,

$$\mathcal{L}_{\texttt{OE}}(\mathsf{K}) := \lim_{T\to\infty} \frac{1}{T}\left(\mathbb{E}\int_0^T \|\mathbf{z}(t) - \hat{\mathbf{z}}(t)\|^2 \mathrm{d}t\right) = \lim_{t\to\infty} \mathbb{E}\|\mathbf{z}(t) - \hat{\mathbf{z}}(t)\|^2. \tag{1.3}$$

In this paper, we study solving the `OE` problem via model-free methods, where the goal is to search for the optimal filter parameters $\mathsf{K} = (\mathbf{A}_\mathsf{K}, \mathbf{B}_\mathsf{K}, \mathbf{C}_\mathsf{K})$ using direct policy search without knowledge or estimation of the true system parameters $\mathbf{A}, \mathbf{C}, \mathbf{G}, \mathbf{W}_1, \mathbf{W}_2$; see Section 2 for further details. The `OE` problem is not a control problem per se, as decisions $\hat{\mathbf{z}}$ do not influence the state evolution of the true system (1.1). Nonetheless, it is an attractive stepping stone for studying model-free control. Like `LQG`, optimal policies for `OE` are *dynamic*, i.e., have memory. `OE` therefore serves as a valuable testbed to develop the theoretical foundations of direct policy search for this important class of policies which has thus far remained poorly understood.

Related work. Recent years have witnessed a resurgence of interest in direct policy search for control and reinforcement learning. Fazel et al. [2018] established global convergence of policy gradient methods on discrete-time `LQR`. Subsequent work has sharpened rates [Malik et al., 2019], analyzed convergence under more general frameworks [Bu et al., 2019], and extended the analysis to continuous-time [Mohammadi et al., 2021]. Beyond `LQR`, Zhang et al. [2020] analyzed policy search for mixed $\mathcal{H}_2/\mathcal{H}_\infty$ and risk-sensitive control. Sun and Fazel [2021] also considered analysis via convex reformulations, for state feedback problems. For discrete state-action `MDPs`, Agarwal et al. [2021] established convergence rates for a variety of policy gradient methods, cf. also Bhandari and Russo [2019]. All of these works considered *static* state-feedback policies, with perfect state information. For problems with partial observation - which necessitates the use of *dynamic* policies - the work most relevant to our own, is Tang et al. [2021], which studied policy search for `LQG`. They established that all stationary points corresponding to minimal controllers are globally optimal, and demonstrated that gradient descent may fail to converge to optimal policies.

1.1 Contributions

We propose a novel policy search method which provably converges to a globally optimal $\mathcal{L}_{\texttt{OE}}$ cost. Despite extensive prior work on *static* policy search (see above), our result constitutes the first rigorous guarantee for policy search over *dynamic* policies. While the `OE` problem admits a convex reformulation [Scherer et al., 1997], it is also well known that its landscape contains suboptimal critical points. This main seem like a contradiction at first, but we demonstrate such spurious critical points occur precisely when the policy loses what we term "informativity": that the steady covariance between the controller state and system state, $\mathbf{\Sigma}_{12} := \lim_{T\to\infty} \frac{1}{T}[\int_0^T \mathbf{x}(t)\hat{\mathbf{x}}(t)^\top \mathrm{d}t]$, becomes rank deficient. With this insight, we propose a regularizer $\mathcal{R}$ that ensures the internal state of the learned policy remains "uniformly informative" about the state of the true system, and prove that gradient descent on the *regularized* objective $\mathcal{L}_{\texttt{OE}}(\cdot) + \lambda \mathcal{R}(\cdot)$ converges at a $\mathcal{O}(1/T)$ rate to an optimal policy. The efficacy of the approach is illustrated via numerical experiments, cf. Appendix F.6. Code is provided in the supplementary material.

Our techniques. Searching over dynamic policies introduces two key challenges: (1) spurious critical points can arise when one or more factors become "degenerate" in a certain way; (2) changes of basis produce a continuum of "equivalent realizations" of the filter K, some of which are poorly conditioned. Similar challenges have been observed in problems with rotational symmetries, e.g. nonconvex matrix factorization. Neither challenge arises when searching over static policies [Fazel et al., 2018]. Our approach is centered around the idea of *convex reformulations* of control synthesis problems, and the following fact regarding functions which admit these reformulations, cf. Appendix H.1 for proof.

Fact 1.1. Let $f : \mathbb{R}^{n_x} \to \mathbb{R}$ be a differentiable, possibly nonconvex function such that $\min_{\boldsymbol{x}} f(\boldsymbol{x})$ is finite. There exists a differentiable function $\Psi : \mathbb{R}^{n_\nu} \to \mathbb{R}^{n_x}$ satisfying the following two properties: (i) the mapping Ψ is surjective, i.e. for all $\boldsymbol{x} \in \mathbb{R}^{n_x}$ there exists $\boldsymbol{\nu} \in \mathbb{R}^{n_\nu}$ such that $\boldsymbol{x} = \Psi(\boldsymbol{\nu})$, (ii) under the change of variables the function $f_{\mathrm{cvx}}(\boldsymbol{\nu}) := f(\Psi(\boldsymbol{\nu}))$ is differentiable and *convex*. Then all first-order stationary points, $\boldsymbol{x}$ s.t $\nabla f(\boldsymbol{x}) = 0$, are globally optimal.

The OE problem, LQG, and many other related control tasks admit convex reformulations [Scherer et al., 1997, Masubuchi et al., 1998]. Given that gradient descent (under mild regularity assumptions) converges to stationary points, we might hope that Fact 1.1 guarantees that direct policy search on the OE filter will succeed at finding an optimal policy, when applied to loss functions admitting such convex reformulations. Somewhat surprisingly, we find that this is emphatically not the case: gradient descent on the $\mathcal{L}_{\mathrm{OE}}$ objective fails to reliably converge to optimal solutions (see Section 3.1).[1]

To resolve this paradox, we show that the surjectivity condition of Fact 1.1 may fail for the convex reparametrization of OE: there are filters K with finite cost $\mathcal{L}_{\mathrm{OE}}(\mathsf{K})$, which are not in the image of the reformulation map $\Psi(\cdot)$. We find that degeneracy occurs precisely when *informativity*, defined in Section 1.1 as $\boldsymbol{\Sigma}_{12,\mathsf{K}}$ having full rank, fails to hold. Conversely, when $\boldsymbol{\Sigma}_{12,\mathsf{K}}$ is full-rank, the conditions of Fact 1.1 are met and the parametrization behaves as needed. Thus, we identify *non-informativity* - rank deficiency of $\boldsymbol{\Sigma}_{12,\mathsf{K}}$ - as the fundamental notion of degeneracy corresponding to challenge (1). Motivated by this observation, we introduce a novel "informativity regularizer" $\mathcal{R}_{\mathrm{info}}(\cdot)$ which enforces that $\boldsymbol{\Sigma}_{12,\mathsf{K}}$ is full rank. Our proposed algorithm, IR-PG alternates between gradient updates on the regularized loss $\mathcal{L}_\lambda(\cdot) := \mathcal{L}_{\mathrm{OE}}(\cdot) + \lambda \mathcal{R}_{\mathrm{info}}(\cdot)$, and "reconditioning" steps to ensure well-conditioned realizations of the filters K, thereby addressing challenge (2) above. We stress that our notion of informativity differs from the *minimality* criterion emphasized in Tang et al. [2021], whose limitations we discuss in Section 3.1.

In order to achieve our quantitative converge guarantees, we establish numerous results which may be of independent interest, including: (i) a *quantitative* analysis of the OE convex reformulation due to Scherer [1995], (ii) novel bounds on the magnitude of solution to Lyapunov equations under the closed-loop OE filter dynamics. Both arguments appeal to a (quantitative measure of) informativity. Finally, we develop a quantitative analogue of Fact 1.1 via a paradigm we call *differentiable convex liftings*, or DCLs, described at length in Section 4.1.

2 Preliminaries

Before presenting our main results in Section 3, we first introduce some of the relevant definitions, and provide the reader with some the relevant background on prediction in partially-observed dynamical systems. We adopt standard notation wherever possible, and for brevity, defer details to Appendix A. As outlined in the introduction, we consider the problem of predicting the outputs of a partially observed linear dynamical system. We refer to the dynamical system defined in Eq. (1.1) as the *true system*, with states $\mathbf{x}(t) \in \mathbb{R}^n$, observations $\mathbf{y}(t) \in \mathbb{R}^m$, and performance outputs $\mathbf{z}(t) \in \mathbb{R}^p$. To ensure the dynamics have a well-defined steady-state, we assume that $\mathbf{A}$ is stable.

Assumption 2.1. The matrix $\mathbf{A}$ is *Hurwitz stable*. That is, the real components of all its eigenvalues are strictly negative: $\Re[\lambda_i(\mathbf{A})] < 0$ for $i \in [n]$.

Because these policies only access the system outputs, and are only evaluated in relation to system outputs, we assume that the true system state is *observable*. Further, we assume that dynamics are subject to sufficiently rich noise excitations.

Assumption 2.2. The pair $(\mathbf{A}, \mathbf{C})$ in Eq. (1.1) is observable. That is, the observability Gramian defined as $\mathcal{G}_{\mathrm{obs}} := \int_0^\infty \exp(s\mathbf{A})^\top \mathbf{C}^\top \mathbf{C} \exp(s\mathbf{A}) \mathrm{d}s$ is strictly positive definite.

---

[1]Failure modes for the LQG problem were presented by Tang et al. [2021].

Assumption 2.3. We assume that the noise matrices $\mathbf{W}_1$ and $\mathbf{W}_2$ are strictly positive definite.[2]

As stated previously, we restrict our attention to finding the best dynamic filter within the parametric family described in Eq. (1.2). Note that this family contains the *Bayes optimal predictor* for the $\mathcal{L}_{\mathtt{OE}}$ objective, as we will later describe in more detail. We review a number of basic facts:

A. Steady state distributions. We define $\mathcal{K}_{\mathtt{stab}} := \{\mathsf{K} : \mathbf{A}_{\mathsf{K}} \text{ is Hurwitz-stable}\}$ to be the set of filters such that $\mathbf{A}_{\mathsf{K}}$ is stable. Under Assumption 2.1, Appendix E.1 shows this equivalent to stability of the closed-loop matrix $\mathbf{A}_{\mathrm{cl},\mathsf{K}}$.

$$\mathbf{A}_{\mathrm{cl},\mathsf{K}} := \begin{bmatrix} \mathbf{A} & 0 \\ \mathbf{B}_{\mathsf{K}}\mathbf{C} & \mathbf{A}_{\mathsf{K}} \end{bmatrix}, \quad \mathcal{K}_{\mathtt{stab}} = \{\mathsf{K} : \mathbf{A}_{\mathrm{cl},\mathsf{K}} \text{ is Hurwitz-stable}\}. \tag{2.1}$$

Stability of $\mathbf{A}_{\mathrm{cl},\mathsf{K}}$ is a sufficient condition for $\mathcal{L}_{\mathtt{OE}}(\mathsf{K})$ to be finite, and for the following limiting covariance to be well defined, $\boldsymbol{\Sigma}_{\mathsf{K}} = \lim_{t\to\infty} \mathbb{E}\begin{bmatrix} \mathbf{x}(t) \\ \hat{\mathbf{x}}_{\mathsf{K}}(t) \end{bmatrix}\begin{bmatrix} \mathbf{x}(t) \\ \hat{\mathbf{x}}_{\mathsf{K}}(t) \end{bmatrix}^{\top} \in \mathbb{S}_{+}^{2n}$. This steady-state covariance is given by the solution to the continuous-time Lyapunov equation,

$$\mathbf{A}_{\mathrm{cl},\mathsf{K}}\boldsymbol{\Sigma} + \boldsymbol{\Sigma}\mathbf{A}_{\mathrm{cl},\mathsf{K}}^{\top} + \mathbf{W}_{\mathrm{cl},\mathsf{K}} = 0, \quad \text{where } \mathbf{W}_{\mathrm{cl},\mathsf{K}} := \begin{bmatrix} \mathbf{W}_1 & 0 \\ 0 & \mathbf{B}_{\mathsf{K}}\mathbf{W}_2\mathbf{B}_{\mathsf{K}}^{\top} \end{bmatrix}. \tag{2.2}$$

Notice that $\boldsymbol{\Sigma}_{\mathsf{K}}$ depends only on $(\mathbf{A}_{\mathsf{K}}, \mathbf{B}_{\mathsf{K}})$, but not on $\mathbf{C}_{\mathsf{K}}$, and that the first $n \times n$ block of $\boldsymbol{\Sigma}_{\mathsf{K}}$ does not depend on K at all. To highlight these distinctions, we partition

$$\boldsymbol{\Sigma} = \begin{bmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} \\ \boldsymbol{\Sigma}_{12}^{\top} & \boldsymbol{\Sigma}_{22} \end{bmatrix}, \quad \boldsymbol{\Sigma}_{\mathsf{K}} = \begin{bmatrix} \boldsymbol{\Sigma}_{11,\mathrm{sys}} & \boldsymbol{\Sigma}_{12,\mathsf{K}} \\ \boldsymbol{\Sigma}_{12,\mathsf{K}}^{\top} & \boldsymbol{\Sigma}_{22,\mathsf{K}} \end{bmatrix}, \tag{2.3}$$

and define $\mathcal{K}_{\mathtt{ctrb}} := \{\mathsf{K} \in \mathcal{K}_{\mathtt{stab}} : \boldsymbol{\Sigma}_{22,\mathsf{K}} \succ 0\}$ as the set of filters whose internal state covariance is full rank. We refer to these as the *controllable* policies, as these are precise the policies for which the pair $(\mathbf{A}_{\mathsf{K}}, \mathbf{B}_{\mathsf{K}})$ is controllable.

B. Equivalent realizations. There are many different ways of parametrizing a given *dynamic* feedback policy, all of which have exactly the same input-output behavior. In particular, given an invertible matrix $\mathbf{S} \in \mathbb{GL}(n)$, the $\mathtt{OE}$ loss of a filter K is invariant under the following class of similarity transforms: $\mathsf{Sim}_{\mathbf{S}}(\mathsf{K}) : (\mathbf{A}_{\mathsf{K}}, \mathbf{B}_{\mathsf{K}}, \mathbf{C}_{\mathsf{K}}) \mapsto (\mathbf{S}\mathbf{A}_{\mathsf{K}}\mathbf{S}^{-1}, \mathbf{S}\mathbf{B}_{\mathsf{K}}, \mathbf{C}_{\mathsf{K}}\mathbf{S}^{-1})$. Formally, for any $\mathsf{K} \in \mathcal{K}_{\mathtt{stab}}$ and any $\mathbf{S} \in \mathbb{GL}(n)$, $\mathcal{L}_{\mathtt{OE}}(\mathsf{K}) = \mathcal{L}_{\mathtt{OE}}(\mathsf{Sim}_{\mathbf{S}}(\mathsf{K}))$. We say that K and K' are *equivalent realizations* if they are related by a similarity transformation $\mathsf{Sim}_{\mathbf{S}}(\mathsf{K}) = \mathsf{K}'$ for some $\mathbf{S} \in \mathbb{GL}(n)$. Note that the set $\mathcal{K}_{\mathtt{ctrb}}$ is also preserved under similarity transformation.

C. Optimal policies. The landmark result by Kalman shows that for the system defined by $(\mathbf{A}, \mathbf{C}, \mathbf{W}_1, \mathbf{W}_2)$ the Kalman filter $\mathsf{K}_{\star} = (\mathbf{A} - \mathbf{L}_{\star}\mathbf{C}, \mathbf{L}_{\star}, \mathbf{G})$ achieves minimal $\mathcal{L}_{\mathtt{OE}}$ loss. Here, $\mathbf{L}_{\star}$ is the Kalman gain which is defined in terms of the solution of the following Riccati equation:

$$\mathbf{A}\mathbf{P}_{\star} + \mathbf{P}_{\star}\mathbf{A}^{\top} - \mathbf{P}_{\star}\mathbf{C}^{\top}\mathbf{W}_2^{-1}\mathbf{C}\mathbf{P}_{\star} + \mathbf{W}_1 = 0, \qquad \mathbf{L}_{\star} = \mathbf{P}_{\star}\mathbf{C}^{\top}\mathbf{W}_2^{-1}. \tag{2.4}$$

We define the set of optimal filters $\mathcal{K}_{\mathtt{opt}}$ to be those which are equivalent to the Kalman filter:

$$\mathcal{K}_{\mathtt{opt}} := \bigcup_{\mathbf{S} \in \mathbb{GL}(n)} \{\mathsf{Sim}_{\mathbf{S}}(\mathbf{A} - \mathbf{L}_{\star}\mathbf{C}, \mathbf{L}_{\star}, \mathbf{G})\}. \tag{2.5}$$

D. Restricted problem setting. The problem description outlined above, including Assumptions 2.1 to 2.3, constitutes the standard $\mathtt{OE}$ problem, well-known in control theory, cf. [Doyle et al., 1989, §IV.D]. In this paper, we will make the following additional assumption that restricts the class of $\mathtt{OE}$ problems we consider. Further discussion on the utility and necessity of this assumption (for our analysis) is provided in Section 3.2 and Appendix D; the latter also shows that Assumption 2.4 holds for "generic" problem instances.

Assumption 2.4. The (cannonical) optimal policy $(\mathbf{A} - \mathbf{L}_{\star}\mathbf{C}, \mathbf{L}_{\star})$ is controllable, i.e. $\int_0^{\infty} \exp(s(\mathbf{A} - \mathbf{L}_{\star}\mathbf{C}))\mathbf{L}_{\star}\mathbf{L}_{\star}^{\top} \exp(s(\mathbf{A} - \mathbf{L}_{\star}\mathbf{C}))^{\top}\mathrm{d}s$ is strictly positive definite.

---

[2] We may relax this assumption to $(\mathbf{A}, \mathbf{W}_1)$ controllable.

E. Interaction protocol. In the spirit of model-free methods, we introduce algorithms which work only assuming access to cost and gradient evaluation oracles. We abstract away the particular implementation of these oracles to simplify our presentation and assume that they are exact, in order to focus on the overall optimization landscape of the $\mathtt{OE}$ problem. More formally, for any filter $\mathsf{K} \in \mathcal{K}_{\mathtt{stab}}$, $\mathsf{Eval}(\mathsf{K}, \mathcal{L}_{\mathtt{OE}})$ returns the $\mathtt{OE}$ cost, $\mathcal{L}_{\mathtt{OE}}(\mathsf{K})$ and $\mathsf{Grad}(\mathsf{K}, \mathcal{L}_{\mathtt{OE}})$ return the gradient of the $\mathtt{OE}$ cost, $\nabla \mathcal{L}_{\mathtt{OE}}(\mathsf{K})$. Despite this simplification, we would like to again emphasize that these can be efficiently approximated in finite samples, and purely on the basis of *observations* $\mathbf{y}_t$ subsampled in discrete intervals. For further discussion, please see Appendix C.

Lastly, in addition to standard cost and gradient evaluations of the $\mathcal{L}_{\mathtt{OE}}$ loss, as part of our algorithm, we further require access to gradient and cost evaluations of smooth functions of the stationary-state covariance. Specifically, if $f : \mathbb{S}_+^{2n} \to \mathbb{R}$ is a function of the covariance matrix K, we assume we can compute $\mathsf{Eval}_{\mathrm{cov}}(\mathsf{K}, f)$ which returns the $f(\mathbf{\Sigma}_\mathsf{K})$ and $\mathsf{Grad}_{\mathrm{cov}}(\mathsf{K}, f)$ which returns $\nabla_\mathsf{K} f(\mathbf{\Sigma}_\mathsf{K})$. In Appendix C.2, we show that these oracles can be implemented without direct state access by "subsampling" multiple observations at different time increments.

3 Main Results

In this section, we present the main contributions of our work. After demonstrating that the $\mathtt{OE}$ cost function contains stationary points that are not globally optimal, we present *informativity-regularized policy gradient* (IR-PG), a direct policy search algorithm based on a novel regularization strategy to preserve *informativity*, introduced in Section 1.1. We state a formal convergence result showing that IR-PG converges to a globally optimal filter at a $\mathcal{O}\left(1/T\right)$ rate. The efficacy of IR-PG is illustrated via numerical experiments in Appendix F.6, cf. Fig. 3 in particular to see how informativity regularization dramatically improves convergence of policy gradient.

3.1 Existence of suboptimal stationary points

Perhaps the simplest model-free approach to the $\mathtt{OE}$ problem is to run gradient descent on $\mathcal{L}_{\mathtt{OE}}(\cdot)$. Under mild assumptions on the loss function $\mathcal{L}_{\mathtt{OE}}$, gradient descent will converge to a first-order stationary point of $\mathcal{L}_{\mathtt{OE}}$. Unfortunately, despite the existence of a convex reformulation [Scherer et al., 1997] and Fact 1.1, the $\mathcal{L}_{\mathtt{OE}}$ loss function contains suboptimal stationary points:

Example 3.1. Consider the $\mathtt{OE}$ instance given by $\mathbf{A} = -\mathbf{I}_2$, $\mathbf{C} = \mathbf{I}_2$, $\mathbf{W}_1 = 3 \times \mathbf{I}_2$, $\mathbf{W}_2 = \mathbf{I}_2$, and the filter $\mathsf{K}_{\mathrm{bad}}$ given by $\mathbf{A}_{\mathrm{bad}} = -\varepsilon \times \mathbf{I}_2$, $\varepsilon > 0$, $\mathbf{B}_{\mathrm{bad}} = \mathbf{0}_2$, $\mathbf{C}_{\mathrm{bad}} = \mathbf{0}_2$. $\mathsf{K}_{\mathrm{bad}}$ constitutes a suboptimal stationary point of $\mathcal{L}_{\mathtt{OE}}$ for this $\mathtt{OE}$ instance.

The example is similar in spirit to Tang et al. [2021, Theorem 4.1]; details are given in Appendix F.1. Importantly, the cost is invariant under perturbations to any single parameter of the filter and $\mathsf{K}_{\mathrm{bad}}$, being equivalent to the zero-filter, is suboptimal. The same is true for any $\mathtt{OE}$ instance: every filter with $\mathbf{B}_{\mathrm{bad}} = \mathbf{0}$, $\mathbf{C}_{\mathrm{bad}} = \mathbf{0}$, and $\mathbf{A}_{\mathrm{bad}}$ being stable is a suboptimal stationary point of $\mathcal{L}_{\mathtt{OE}}$.

The perils of enforcing minimality. A filter K is *minimal* if $(\mathbf{A}_\mathsf{K}, \mathbf{B}_\mathsf{K})$ is controllable, and $(\mathbf{A}_\mathsf{K}, \mathbf{C}_\mathsf{K})$ is observable. Example 3.1 is the extreme case of a *non-minimal* filter, since $\mathbf{B}_{\mathrm{bad}} = \mathbf{C}_{\mathrm{bad}} = \mathbf{0}_2$. Conversely, as a special case of $\mathtt{LQG}$, the $\mathtt{OE}$ problem inherits the property that all stationary points corresponding to *minimal* filters are globally optimal [Tang et al., 2021, Theorem 4.3]. Therefore, it may be natural to ask: *can a local search algorithm enforce minimality to avoid suboptimal stationary points?* A classical result due to Brockett [1976] suggests not: the set of minimal n-th order single-input-single-output transfer functions (e.g. filters) is the disjoint union of $n + 1$ open sets. Thus it is impossible for a continuous path to pass from one of these open sets to another without entering a region corresponding to a non-minimal filter, suggesting that a local search algorithm regularized to ensure minimality at every iteration may never converge to the optimal solution. See Appendix F.2 for further discussion and supporting numerical experiments.

Futher challenges. Given the drawbacks of enforcing minimality, one may wonder whether it is sufficient to enforce controllability or observability alone. In Example 3.1 above - and indeed, for all the examples in Tang et al. [2021] of suboptimal stationary points in the $\mathtt{LQG}$ landscape - there is a loss of *both* observability ($\mathbf{C}_{\mathrm{bad}} = \mathbf{0}$) and controllability ($\mathbf{B}_{\mathrm{bad}} = \mathbf{0}$). Unfortunately, suboptimal stationary points can occur at controllable policies and observable policies, as shown in Proposition F.1, and detailed in Appendix F.3. There we also explain that approaches based on

221 saddle-point escape with appropriate random perturbations [Jin et al., 2017] or acceleration [Jin
222 et al., 2018] are also unlikely to be successful.

3.2 A provably convergent algorithm

224 The previous discussion puts us in a bind: we cannot regularize to preserve minimality because of
225 path-disconnectedness. Yet, neither controllability nor observability is not enough to rule out sub-
226 optimal stationary points. We identify a stronger condition called *informativity* which is sufficient.
227 We define the set of *informative* filters as $\mathcal{K}_{\texttt{info}} := \{\mathsf{K} \in \mathcal{K}_{\texttt{stab}} : \text{rank}(\boldsymbol{\Sigma}_{12,\mathsf{K}}) = n\}$. First, we
228 check that all optimal filters lie in this set, under Assumption 2.4 (see Appendix E.3 for proof)

229 **Lemma 3.1.** *Under Assumptions 2.1 to 2.4, then $\mathcal{K}_{\texttt{opt}} \subset \mathcal{K}_{\texttt{info}} \subset \mathcal{K}_{\texttt{ctrb}}$, and $\mathcal{K}_{\texttt{info}}$ is an open set.*

230 Our key insight is that informativity is also *sufficient* to ensure optimality of stationary points, but
231 does not cause path-connectedness issues as it did for minimality (see Appendix G.3 for the proof).

232 **Theorem 1.** *Let $\mathsf{K} \in \mathcal{K}_{\texttt{info}}$; then (i) there is a continuous path lying in $\mathcal{K}_{\texttt{info}}$ connecting K to some*
233 $\mathsf{K}_\star \in \mathcal{K}_{\texttt{opt}}$ *and (ii) if $\nabla\mathcal{L}_{\texttt{OE}}(\mathsf{K}) = \mathbf{0}$, then $\mathsf{K} \in \mathcal{K}_{\texttt{opt}}$.*

234 Theorem 1 suggests that gradient descent with enforced informativity should converge to optimal
235 filters. It is not, however, implied by the landscape analysis of Tang et al. [2021], which focuses
236 solely on *minimal* stationary points. But numerous challenges remain: (1) how can one enforce
237 informativity in a smooth fashion? (2) what quantitative measure of informativity provides quanti-
238 tative suboptimality guarantees on approximate first-order stationary points? (3) Given that $\mathcal{L}_{\texttt{OE}}(\mathsf{K})$
239 need not have compact level sets (see Appendix F.3), how does one ensure that the iterates of policy
240 search do not escape to infinity, or reach regions where the loss of smoothness is arbitrarily poor?

241 **The explained covariance matrix.** In light of Theorem 1, we design a policy search algorithm
242 which ensures that $\boldsymbol{\Sigma}_{12,\mathsf{K}}$ remains full-rank throughout the search, but does so in a quantitative
243 fashion. Our central object is the *explained covariance matrix*, which measures how much of the
244 covariance of the steady-state system $\mathbf{x}(t)$ is explained by the internal filter state $\hat{\mathbf{x}}_\mathsf{K}(t)$ in the large t
245 limit: $\mathbf{Z}_\mathsf{K} := \lim_{t\to\infty} (\text{Cov}[\mathbf{x}(t)] - \mathbb{E}[\text{Cov}[\mathbf{x}(t) \mid \hat{\mathbf{x}}_\mathsf{K}(t)]])$. When $\mathsf{K} \in \mathcal{K}_{\texttt{ctrb}}$, $\mathbf{Z}_\mathsf{K}$ admits an elegant
246 closed-form expression, which provides an alternative definition of $\mathcal{K}_{\texttt{info}}$:

$$\mathbf{Z}_\mathsf{K} = \boldsymbol{\Sigma}_{12,\mathsf{K}}\boldsymbol{\Sigma}_{22,\mathsf{K}}^{-1}\boldsymbol{\Sigma}_{12,\mathsf{K}}^\top, \text{ so that } \mathcal{K}_{\texttt{info}} = \{\mathsf{K} : \mathsf{K} \in \mathcal{K}_{\texttt{ctrb}} \text{ and } \mathbf{Z}_\mathsf{K} \succ 0\}.$$

247 Since $\mathbf{Z}_\mathsf{K}$ is invariant under similarity transformations, $\mathbf{Z}_\mathsf{K}$ can be interpreted as a normalized ana-
248 logue of $\boldsymbol{\Sigma}_{12,\mathsf{K}}$. Informally, the quadratic form $v^\top\mathbf{Z}_\mathsf{K}v$ is a sufficient statistic for how much infor-
249 mation $\hat{\mathbf{x}}(t)$ contains about the "v-direction" of $\mathbf{x}(t)$; see Appendix E.6 for a precise statement.

250 **Explained-covariance regularization.** We preserve informativity by ensuring our iterates satisfy
251 $\mathbf{Z}_\mathsf{K} \succ 0$. To this end, we run gradient descent on the regularized objective for some $\lambda > 0$:

$$\mathcal{L}_\lambda(\mathsf{K}) := \mathcal{L}_{\texttt{OE}}(\mathsf{K}) + \lambda \cdot \mathcal{R}_{\texttt{info}}(\mathsf{K}), \quad \text{where } \mathcal{R}_{\texttt{info}}(\mathsf{K}) := \begin{cases} \text{tr}[\mathbf{Z}_\mathsf{K}^{-1}] & \mathsf{K} \in \mathcal{K}_{\texttt{info}} \\ \infty & \text{otherwise.} \end{cases} \quad (3.1)$$

252 The proposed regularizer has several important properties. First, though $\mathcal{R}_{\texttt{info}}$ is nonconvex in
253 K, both $\mathcal{R}_{\texttt{info}}$ and $\mathcal{L}_{\texttt{OE}}$ can be made convex under the *same* change of variables, in the sense of
254 Fact 1.1, cf. Lemma I.2. In addition, $\mathcal{R}_{\texttt{info}}$ is non-negative, and tends to ∞ as $\mathbf{Z}_\mathsf{K}$ approaches
255 singularity. Furthermore, $\mathcal{R}_{\texttt{info}}$ is invariant under similarity transformations. Next, many of the
256 essential quantities arising in our analysis can be bounded in terms of $\mathbf{Z}_\mathsf{K}^{-1}$, justifying $\mathbf{Z}_\mathsf{K}$ is a natural
257 quantitative measure of informativity. Lastly, the set of global-minimizers of $\mathcal{R}_{\texttt{info}}(\cdot)$ are precisely
258 the optimal filters for the $\texttt{OE}$ problem, as per the following lemma (see Appendix E.4 for proof).

259 **Lemma 3.2** (Existence of maximal $\mathbf{Z}_\mathsf{K}$)**.** *Under Assumptions 2.1 to 2.3, there exists a unique $\mathbf{Z}_\star \succ 0$*
260 *such that $\mathbf{Z}_\star = \mathbf{Z}_\mathsf{K}$ if and only if $\mathsf{K} \in \mathcal{K}_{\texttt{opt}}$, and $\mathbf{Z}_\star \succeq \mathbf{Z}_\mathsf{K}$ for all $\mathsf{K} \in \mathcal{K}_{\texttt{ctrb}} \setminus \mathcal{K}_{\texttt{opt}}$. Consequently,*
261 $\mathcal{K}_{\texttt{opt}} = \arg\min_{\mathsf{K} \in \mathcal{K}_{\texttt{ctrb}}} \mathcal{R}_{\texttt{info}}(\mathsf{K})$.

262 Lemma 3.2 directly implies that the suboptimality of $\mathcal{L}_\lambda(\cdot)$ upper bounds the suboptimality in $\mathcal{L}_{\texttt{OE}}(\cdot)$,
263 so we can minimize $\mathcal{L}_\lambda$ as a proxy for minimizing $\mathcal{L}_{\texttt{OE}}$.

264 **Corollary 3.1.** *For any K, we have $\mathcal{L}_{\texttt{OE}}(\mathsf{K}) - \min_{\mathsf{K}'} \mathcal{L}_{\texttt{OE}}(\mathsf{K}') \leq \mathcal{L}_\lambda(\mathsf{K}) - \min_{\mathsf{K}'} \mathcal{L}_\lambda(\mathsf{K}')$.*

Algorithm 1 Informativity-regularized Policy Gradient (IR-PG)

1: **Input:** Initial $\mathsf{K}_0 \in \mathcal{K}_{\texttt{info}}$, step size $\eta > 0$, regularization parameter $\lambda > 0$
   `% Define` $\mathcal{L}_\lambda(\mathsf{K}) := \mathcal{L}_{\texttt{OE}}(\cdot) + \lambda \mathrm{tr}[\mathbf{Z}_\mathsf{K}^{-1}]$
2: **for** each iteration $s = 0, 1, 2, \ldots$ **do**
3:     **Recondition** $\widetilde{\mathsf{K}}_t = \mathrm{recond}(\mathsf{K}_t)$, where $\mathrm{recond}(\cdot)$ is defined in Eq. (3.2).
4:     **Compute** $\nabla_t = \nabla \mathcal{L}_\lambda(\widetilde{\mathsf{K}}_t)$.
5:     **Update** $\mathsf{K}_{t+1} \leftarrow \widetilde{\mathsf{K}}_t - \eta \nabla_t$.

Reconditioning. We introduce an additional normalization step between policy updates to ensure the iterates produced by our algorithm have well-conditioned covariance matrices; this in turn ensures the iterates produced by our algorithm remain in a compact set, and that the smoothness of $\mathcal{L}_\lambda$ is uniformly bounded. For any filter $\mathsf{K} \in \mathcal{K}_{\texttt{ctrb}}$ such that $\boldsymbol{\Sigma}_{22,\mathsf{K}} \succ 0$, the reconditioning operator $\mathrm{recond}(\mathsf{K})$ returns a filter K' which is equivalent to K, but for which $\boldsymbol{\Sigma}_{22,\mathsf{K}'} = \mathbf{I}_n$. Formally[3],

$$\mathrm{recond}(\mathsf{K}) := \mathsf{Sim}_\mathbf{S}(\mathbf{A}_\mathsf{K}, \mathbf{B}_\mathsf{K}, \mathbf{C}_\mathsf{K}), \quad \text{where } \mathbf{S} = \boldsymbol{\Sigma}_{22,\mathsf{K}}^{-1/2}. \tag{3.2}$$

Since K and $\mathsf{K}' = \mathrm{recond}(\mathsf{K})$ are equivalent realizations, we have $\mathcal{L}_{\texttt{OE}}(\mathsf{K}) = \mathcal{L}_{\texttt{OE}}(\mathsf{K}')$, $\mathcal{R}_{\texttt{info}}(\mathsf{K}) = \mathcal{R}_{\texttt{info}}(\mathsf{K}')$, and thus $\mathcal{L}_\lambda(\mathsf{K}) = \mathcal{L}_\lambda(\mathsf{K}')$.

Statement of IR-PG. We can now describe IR-PG, whose pseudocode is displayed in Algorithm 1. IR-PG applies gradient descent on the regularized $\mathcal{L}_{\texttt{OE}}$ objective, with an additional balancing step between gradient updates. Appendix B.1 provides a variant where the step size is chosen by backtracking line-search, which enjoys the same rigorous convergence guarantees. To guarantee convergence to an optimal filter (and finiteness of $\mathcal{L}_\lambda$), we need to initialize at a filter such that $\mathbf{Z}_{\mathsf{K}_0} \succ 0$, i.e. $\mathsf{K}_0 \in \mathcal{K}_{\texttt{info}}$. Fortunately, random initializations from a continuous distribution satisfy this condition with probability 1 (see Appendix E.7 for a formal statement and proof). Concerning the choice of λ, the formal guarantees in Section 3.3 below hold for any $\lambda > 0$. Nonetheless, the choice of λ can influence the numerical properties of the loss function, such as smoothness and the size of the gradients; cf. Appendix F.6 for further discussion.

3.3 Formal guarantees

We conclude this section by stating the formal convergence guarantee for IR-PG. Our results depend on natural problem quantities, among which is the minimum singular value of $\mathbf{P}_\star$ (as defined in Eq. (2.4)), which we show is always strictly positive.

Lemma 3.3. *Let $\mathbf{P}_\star$ be the solution to the Riccati equation in Eq. (2.4). Then under Assumptions 2.1 to 2.3, $\sigma_\star := \lambda_{\min}(\mathbf{P}_\star)$ is strictly positive. Moreover, $\mathbf{P}_\star = \boldsymbol{\Sigma}_{11,\mathrm{sys}} - \mathbf{Z}_\star$.*

In other words, $\mathbf{P}_\star$ is the limiting conditional covariance of the true state $\mathbf{x}(t)$ given the policy-state $\hat{\mathbf{x}}_\mathsf{K}(t)$ under (any) optimal policy. Lemma 3.3 states that this covariance is nonsingular, i.e. not even optimal policies contain perfect information about any mode of $\mathbf{x}(t)$. Given an initialization K_0, our convergence rate depends polynomially on the following problem parameters:

$$C_{\texttt{sys}} := \max\left\{\|\mathbf{A}\|, \|\mathbf{C}\|, \|\mathbf{G}\|, \|\mathbf{W}_2\|, \|\mathbf{W}_2^{-1}\|, \|\mathbf{W}_1^{-1}\|, \|\boldsymbol{\Sigma}_{11,\mathrm{sys}}\|, \sigma_\star^{-1}\right\}. \tag{3.3}$$

Theorem 2. *Fix $\lambda > 0$, $\mathsf{K}_0 \in \mathcal{K}_{\texttt{info}}$. There are terms $\mathcal{C}_1, \mathcal{C}_2 \geq 1$, which are at most polynomial in $n, m, C_{\texttt{sys}}, \lambda, \lambda^{-1}$ and $\mathcal{L}_\lambda(\mathsf{K}_0)$, such that the iterates of IR-PG with any stepsize $\eta \leq \frac{1}{\mathcal{C}_1}$ satisfy*

$$\mathcal{L}_{\texttt{OE}}(\mathsf{K}_s) - \min_\mathsf{K} \mathcal{L}_{\texttt{OE}}(\mathsf{K}) \leq \mathcal{L}_\lambda(\mathsf{K}_s) - \min_\mathsf{K} \mathcal{L}_\lambda(\mathsf{K}) \leq \frac{\mathcal{C}_2}{\eta} \cdot \frac{1}{s}, \quad \forall s \geq 1.$$

The formal guarantee for back-tracking stepsizes is nearly analogous, and given in Appendix B.1. The oracle complexity of IR-PG is described in Appendix B.2. We sketch the highlights of the proof in the following section, after introducing our `DCL` framework. A rigorous proof overview is deferred to Appendix G.2.

---

[3]Our proposed algorithm also works with an approximate balancing $\widetilde{\mathrm{recond}}(\cdot)$, where $\widetilde{\mathrm{recond}}(\mathsf{K})$ returns a K' which is equivalent to K, and $\|\boldsymbol{\Sigma}_{22,\mathsf{K}'} - \mathbf{I}_n\| \leq \varepsilon$ for some tolerance $\varepsilon > 0$ (e.g. $\varepsilon = 1/8$).

4 Analysis Framework

In this section, we sketch our analysis in two halves. In the first, we provide a general framework for analyzing gradient descent on objectives which admit convex formulations. This takes place in two parts: first, our technique of *differentiable convex liftings* (DCLs) which establish a weak form of gradient domination we term weak-PL. Second, we show that weak-PL, together with smoothness (i.e. upper bound on the magnitude of the Hessian) implies convergence of gradient descent at a $O(1/t)$ rate. The DCL formalizes the developments in the "our techniques" discussion of Section 1.1.

Next, we instatiate the analysis framework for regularized output estimation loss $\mathcal{L}_\lambda$ defined in the previous section. It suffices to establish both (a) smoothness of $\mathcal{L}_\lambda$ and (b) weak-PL by exhibiting a valid DCL. Interestingly, the upper bounds on both the Hessian magnitude and the weak-PL factor depend on the magnitude of $\mathrm{tr}[\mathbf{Z}_\mathsf{K}^{-1}]$, which is precisely what is controlled by the informativity regularizer in Eq. (3.1).

4.1 Differentiable convex liftings (DCLs)

This section introduces *differentiable convex liftings* (DCLs), a rigorous and flexible framework for operationalizing convex reformulations of nonconvex objectives.

Preliminaries. To neatly accommodate optimization over constrained domains, we express functions $f : \mathbb{R}^d \to \bar{\mathbb{R}}$ as taking values in the extended reals $\bar{\mathbb{R}} := \mathbb{R} \cup \{\infty\}$.[4] Given such an f, we denote its domain $\mathrm{dom}(f) : \{\boldsymbol{x} : f(\boldsymbol{x}) \neq \infty\}$ as the set on which f is finite; we say f is *proper* if $\mathrm{dom}(f) \neq \emptyset$; we define its minimal value $\inf(f) = \inf_{\boldsymbol{x} \in \mathbb{R}^d} f(\boldsymbol{x})$. We say $f \in \mathscr{C}^k(\mathcal{K})$ on a domain $\mathcal{K}$ if f is k-times continuously differentiable (and finite) on some open set containing $\mathcal{K} \subset \mathbb{R}^d$.

DCLs. The DCL is a generic template for convex reformulation that significantly generalizes the setup in Fact 1.1. Rather than relating f directly to a convex f_{cvx}, we "lift" f to a function f_{lft} by appending auxiliary variables. We then assume a reparametrization Φ mapping the domain of f_{lft} to that of f_{cvx} in this higher dimensional space; intuitively, Φ is the "inverse" of Ψ in Fact 1.1.

Definition 4.1. A triplet of functions $(f_{\mathrm{cvx}}, f_{\mathrm{lft}}, \Phi)$ is a DCL of a proper function $f : \mathbb{R}^d \to \bar{\mathbb{R}}$ if
(1) $f_{\mathrm{cvx}} : \mathbb{R}^{d_z} \to \bar{\mathbb{R}}$ is a a proper convex function whose minimum is attained by some $\boldsymbol{z}^\star$.
(2) For some additional number of parameters $d_\xi \geq 0$, $f_{\mathrm{lft}} : \mathbb{R}^{d+d_\xi} \to \bar{\mathbb{R}}$ is related to f; via partial minimization: $f(\boldsymbol{x}) = \min_{\boldsymbol{\xi} \in \mathbb{R}^{d_\xi}} f_{\mathrm{lft}}(\boldsymbol{x}, \boldsymbol{\xi})$.
(3) There is an open set $\mathcal{Y} \supseteq \mathrm{dom}(f_{\mathrm{lft}})$ for which $\Phi : \mathcal{Y} \to \mathrm{dom}(f_{\mathrm{cvx}})$ is $\mathscr{C}^1$ and satisfies $f_{\mathrm{lft}}(\cdot) = f_{\mathrm{cvx}}(\Phi(\cdot))$.

Above, d_z and d_ξ are arbitrary dimensions representing the arguments of $f_{\mathrm{cvx}}(\cdot)$ and $f_{\mathrm{lft}}(\boldsymbol{x}, \cdot)$, respectively. The mere existence of a DCL implies that approximate stationary points of f are also approximate minimizers, under conditions elaborated on below:

Theorem 3. *Let* $f : \mathbb{R}^d \to \bar{\mathbb{R}}$ *be a proper function with* DCL $(f_{\mathrm{cvx}}, f_{\mathrm{lft}}, \Phi)$. *Then, for any* $\boldsymbol{x} \in \mathrm{dom}(f)$ *at which* f *is differentiable,* f *satisfies the weak-PL condition:*

$$\|\nabla f(\boldsymbol{x})\| \geq \alpha_{\mathrm{DCL}}(\boldsymbol{x}) \cdot (f(\boldsymbol{x}) - \inf(f)), \quad \text{where } \alpha_{\mathrm{DCL}}(\boldsymbol{x}) := \max_{\substack{\boldsymbol{z}^\star \in \arg\min f_{\mathrm{cvx}}(\cdot) \\ \boldsymbol{\xi} \in \arg\min f_{\mathrm{lft}}(\boldsymbol{x}, \cdot)}} \frac{\sigma_{d_z}(\nabla \Phi(\boldsymbol{x}, \boldsymbol{\xi}))}{\|\Phi(\boldsymbol{x}, \boldsymbol{\xi}) - \boldsymbol{z}^\star\|}.$$

Theorem 3 strengthens Fact 1.1 in two respects. For one, it does not impose any smoothness restrictions on f_{lft} or f_{cvx}; in particular f_{cvx} can be highly non-smooth and, due to the extended-real function formulation, can also include constraints. And second, the lifting f_{lft} adds considerable flexibility, which we show is necessary to capture the convex reformulation of OE (Appendix I).

The factor $\alpha_{\mathrm{DCL}}(\boldsymbol{x})$ depends on two quantities. The numerator is the d_z-th singular value of $\nabla \Phi(\boldsymbol{x}, \boldsymbol{\xi})$ for any $\boldsymbol{\xi} \in \arg\min f_{\mathrm{lft}}(\boldsymbol{x}, \cdot)$. This captures how large perturbations of f_{lft}'s arguments must be in order to achieve a desired perturbation of the arguments of f_{cvx}, under the reparameterization Φ. The additional arguments in f_{lft} compared to f adds additional columns to $\nabla \Phi$ thereby making it easier to ensure $\sigma_{d_z}(\nabla \Phi(\cdot)) > 0$. On the other hand, the denominator measures the Euclidean distance between any minimizer of the convex function $f_{\mathrm{cvx}}(\cdot)$ and image of $(\boldsymbol{x}, \boldsymbol{\xi})$ under the reparameterization Φ, and can be bounded under quite benign conditions. *Proof Sketch.* The formal

---

[4]Because we solely consider minimizations, $\bar{\mathbb{R}}$ does not include $-\infty$

proof of Theorem 3 (given in Appendix H.2) takes special care to handle that allow f_{lft} and f_{cvx} to be finite only on restricted domains, and possible non-smoothness; still, the main ideas behind are intuitive. For f_{cvx} convex, $f_{\mathrm{cvx}}(\boldsymbol{z}) - \inf(f_{\mathrm{cvx}}) = \mathcal{O}(\|\nabla f_{\mathrm{cvx}}(\boldsymbol{z})\|)$. Using the DCL definition and analyzing the inverse image of a point under Φ, we can also establish a gradient domination result for f_{lft}. Weak-PL result for the original function f follows since f is related to f_{lft} by partial minimization, so its gradients must be larger than those of f_{lft}.

Gradient descent with DCLs. We now describe how DCLs yield quantitative convergence guarantees for gradient descent. A more general guarantee accommodating the reconditioning step in IR-PG is deferred to Appendix G.1, and encompasses the bound below as a special case. Given $\alpha > 0$, we say that proper $f : \mathbb{R}^d \to \bar{\mathbb{R}}$ satisfies α-*weak-PL* (named after the stronger Polyak-Łojasiewicz condition) on a domain $\mathcal{K} \subset \mathbb{R}^d$ if $f \in \mathscr{C}^1(\mathcal{K})$ and $\|\nabla f(\boldsymbol{x})\| \geq \alpha(f(\boldsymbol{x}) - \inf(f))$. From Theorem 3, we see f satisfies $\alpha_{\mathcal{K}}$-weak PL on $\mathcal{K}$ if f has a DCL and $\alpha_{\mathcal{K}} := \inf_{\boldsymbol{x} \in \mathcal{K}} \alpha_{\mathrm{DCL}}(\boldsymbol{x}) > 0$. To analyze gradient descent, we also require smoothness: we say f is β-*upper-smooth* on $\mathcal{K}$ if $f \in \mathscr{C}^2(\mathcal{K})$ and for all $\boldsymbol{x} \in \mathcal{K}$, $\nabla^2 f(\boldsymbol{x}) \preceq \beta \mathbf{I}$. The following follows from a standard descent lemma for smooth (though possibly nonconvex) functions.

Proposition 4.1. *Let $\boldsymbol{x}_0 \in \mathrm{dom}(f)$, and suppose that the level set $\mathcal{K}(\boldsymbol{x}_0) := \{\boldsymbol{x} : f(\boldsymbol{x}) \leq f(\boldsymbol{x}_0)\}$ is compact, and f satisfies $\alpha_{\boldsymbol{x}_0} > 0$-weak PL and $\beta_{\boldsymbol{x}_0} > 0$-upper smoothness on $\mathcal{K}(\boldsymbol{x})$. Then, $\boldsymbol{x}_0$ lies in the same path-connected component of some minimizer of f, and for any $\eta \leq 1/\beta_{\boldsymbol{x}_0}$ the updates $\boldsymbol{x}_{k+1} = \boldsymbol{x}_k - \eta \nabla f(\boldsymbol{x}_k)$ satisfy $f(\boldsymbol{x}_k) - \inf(f) \leq 2/(k \cdot \alpha_{\boldsymbol{x}_0}^2 \eta)$.*

4.2 Proof of Theorem 2.

In light of Proposition 4.1 (and its generalization to include reconditioning), it suffices to establish both the weak-PL and smoothness of the loss $\mathcal{L}_\lambda(\cdot)$ on the well-conditioned sublevel set $\mathcal{K}_0 := \{\mathsf{K} : \mathcal{L}_\lambda(\mathsf{K}) \leq \mathcal{L}_\lambda(\mathsf{K}_0), \frac{1}{2}\mathbf{I}_n \preceq \boldsymbol{\Sigma}_{22,\mathsf{K}} \preceq 2\mathbf{I}_n\}$. To establish weak-PL, we exhibit a DCL for which $\alpha_{\mathrm{DCL}}(\mathsf{K})$ depends only on the operator norms of $\boldsymbol{\Sigma}_\mathsf{K}, \boldsymbol{\Sigma}_\mathsf{K}^{-1}, \mathbf{Z}_\mathsf{K}$: as well as other system-quantities. Here, $\mathrm{poly}_{\mathrm{op}}(\dots)$ denotes a quantity at most polynomial in the operator norms of its arguments.

Proposition 4.2. *For any $\lambda \geq 0$ (non-strict), the objective $\mathcal{L}_\lambda(\mathsf{K})$ admits a DCL $(f_{\mathrm{cvx}}, f_{\mathrm{lft}}, \Phi)$ where the lifted parameter takes the form $(\mathsf{K}, \boldsymbol{\Sigma}_\mathsf{K}) \in \mathcal{K}_{\mathrm{info}} \times \mathbb{S}_{++}^{2n}$, $\mathcal{L}_\lambda(\mathsf{K}) = f_{\mathrm{lft}}(\mathsf{K}, \boldsymbol{\Sigma}_\mathsf{K}) = \min_{\boldsymbol{\Sigma} \in \mathbb{S}_+^{2n}} f_{\mathrm{lft}}(\mathsf{K}, \boldsymbol{\Sigma})$, and where*

$$\sigma_{d_z}(\nabla \Phi(\mathsf{K}, \boldsymbol{\Sigma}_\mathsf{K})) \geq 1/\mathrm{poly}_{\mathrm{op}}\left(\mathbf{A}, \mathbf{C}, \mathbf{W}_2^{-1}, \boldsymbol{\Sigma}_\mathsf{K}, \boldsymbol{\Sigma}_\mathsf{K}^{-1}, \mathbf{Z}_\mathsf{K}^{-1}, \mathcal{L}_{\mathrm{OE}}(\mathsf{K})\right)$$

$$\|\Phi(\mathsf{K}, \boldsymbol{\Sigma}_\mathsf{K})\|_{\ell_2} \leq (\max\{n, \sqrt{mn}\} + \sqrt{\mathcal{L}_{\mathrm{OE}}(\mathsf{K})}) \cdot \mathrm{poly}_{\mathrm{op}}\left(\mathbf{A}, \mathbf{C}, \mathbf{W}_2^{-1}, \boldsymbol{\Sigma}_\mathsf{K}, \boldsymbol{\Sigma}_\mathsf{K}^{-1}, \mathbf{Z}_\mathsf{K}^{-1}\right).$$

Furthermore, the norms of the parameters $\mathbf{A}_\mathsf{K}, \mathbf{B}_\mathsf{K}, \mathbf{C}_\mathsf{K}$ satisfy the following bounds:

$$\max\{\|\mathbf{A}_\mathsf{K}\|_{\mathrm{op}}, \|\mathbf{B}_\mathsf{K}\|_{\mathrm{op}}\} \leq \mathrm{poly}_{\mathrm{op}}\left(\mathbf{A}, \mathbf{C}, \mathbf{W}_2^{-1}, \mathbf{Z}_\mathsf{K}^{-1}, \boldsymbol{\Sigma}_\mathsf{K}, \boldsymbol{\Sigma}_\mathsf{K}^{-1}\right), \quad \|\mathbf{C}_\mathsf{K}\|_{\mathrm{F}} \leq \sqrt{\mathcal{L}_{\mathrm{OE}}(\mathsf{K})/\|\boldsymbol{\Sigma}_\mathsf{K}^{-1}\|}.$$
$$\tag{4.1}$$

Smoothness is established in (Proposition G.5), which relies on a novel (and rather challenging) bound on the solutions to Lyapunov equations involving $\mathbf{A}_{\mathrm{cl},\mathsf{K}}$:

Proposition 4.3 (Stability of $\mathbf{A}_{\mathrm{cl},\mathsf{K}}$). *Suppose that $\mathsf{K} \in \mathcal{K}_{\mathrm{info}}$. Then, for any matrix $\mathbf{Y} \in \mathbb{S}^{2n}$, the solution $\boldsymbol{\Sigma}_{\mathsf{K},\mathbf{Y}}$ to the Lyapunov equation $\mathbf{A}_{\mathrm{cl},\mathsf{K}} \boldsymbol{\Sigma}_{\mathsf{K},\mathbf{Y}} + \boldsymbol{\Sigma}_{\mathsf{K},\mathbf{Y}} \mathbf{A}_{\mathrm{cl},\mathsf{K}}^\top + \mathbf{Y} = 0$ satisfies*

$$\|\boldsymbol{\Sigma}_{\mathsf{K},\mathbf{Y}}\|_\circ \leq C_{\mathrm{lyap}}(\mathsf{K}) \cdot \|\mathbf{Y}\|_\circ, \quad \text{where } C_{\mathrm{lyap}}(\mathsf{K}) = \mathrm{poly}_{\mathrm{op}}\left(\boldsymbol{\Sigma}_\mathsf{K}, \boldsymbol{\Sigma}_\mathsf{K}^{-1}, \mathbf{Z}_\mathsf{K}^{-1}, \mathbf{W}_1^{-1}, \mathbf{W}_2^{-1}, \mathbf{C}\right),$$

and where $\|\cdot\|_\circ$ denotes either the operator, Frobenius, or nuclear norm.

The regularization $\mathcal{R}_{\mathrm{info}}(\mathsf{K})$ ensures $\|\mathbf{Z}_\mathsf{K}^{-1}\|$ remains bounded on the sublevel $\mathcal{K}_0$; we further show (Lemma G.7) that $\frac{1}{2}\mathbf{I}_n \preceq \boldsymbol{\Sigma}_{22,\mathsf{K}} \preceq \frac{3}{2}\mathbf{I}_n$ implies $\boldsymbol{\Sigma}_\mathsf{K}$ is invertible, and ensures $\|\boldsymbol{\Sigma}_\mathsf{K}\|, \|\boldsymbol{\Sigma}_\mathsf{K}^{-1}\|$ are uniformly bounded. Thus, the weak-PL constant and smoothness parameters are uniformly bounded on $\mathcal{K}_0$, concluding the proof. We stress that the proofs of Proposition 4.2 and Proposition 4.3 require several novel technical arguments, which may be of independent interest. A full proof roadmap, including formal statements of the aforementioned results, is given in Appendix G.2. The fact that $\|\boldsymbol{\Sigma}_\mathsf{K}\|, \|\boldsymbol{\Sigma}_\mathsf{K}^{-1}\|, \|\mathbf{Z}_\mathsf{K}^{-1}\|$ appear throughout the analysis suggests that (a) informativity as measured by $\mathbf{Z}_\mathsf{K}^{-1}$, and (b) the conditioning of $\boldsymbol{\Sigma}_{22,\mathsf{K}}$ may be fundamental to the OE landscape.

Proof of Theorem 1. The proof of Theorem 1 is simpler, as it relies mainly on the *existence* of a DCL for which α_{DCL} does not vanish. See Appendix G.3 for details.

References

Alekh Agarwal, Sham M Kakade, Jason D Lee, and Gaurav Mahajan. On the theory of policy gradient methods: Optimality, approximation, and distribution shift. *Journal of Machine Learning Research*, 22(98):1–76, 2021.

Naman Agarwal, Zeyuan Allen-Zhu, Brian Bullins, Elad Hazan, and Tengyu Ma. Finding approximate local minima faster than gradient descent. In *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing*, pages 1195–1199, 2017.

Michael Athans. The importance of Kalman filtering methods for economic systems. In *Annals of Economic and Social Measurement, Volume 3, number 1*, pages 49–64. NBER, 1974.

Jalaj Bhandari and Daniel Russo. Global optimality guarantees for policy gradient methods. *arXiv preprint arXiv:1906.01786*, 2019.

Stephen Boyd, Stephen P Boyd, and Lieven Vandenberghe. *Convex Optimization*. Cambridge University Press, 2004.

Roger Brockett. Some geometric questions in the theory of linear systems. *IEEE Transactions on Automatic Control*, 21(4):449–455, 1976. doi: 10.1109/TAC.1976.1101301.

Jingjing Bu, Afshin Mesbahi, Maryam Fazel, and Mehran Mesbahi. LQR through the lens of first order methods: Discrete-time case. *arXiv preprint arXiv:1907.08921*, 2019.

Sébastien Bubeck. Convex optimization: Algorithms and complexity. *arXiv preprint arXiv:1405.4980*, 2014.

Yair Carmon, John C Duchi, Oliver Hinder, and Aaron Sidford. Accelerated methods for nonconvex optimization. *SIAM Journal on Optimization*, 28(2):1751–1772, 2018.

J.C. Doyle, K. Glover, P.P. Khargonekar, and B.A. Francis. State-space solutions to standard h_2 and h_∞ control problems. *IEEE Transactions on Automatic Control*, 34(8):831–847, 1989. doi: 10.1109/9.29425.

Guang-Ren Duan and Ron J Patton. A note on hurwitz stability of matrices. *Automatica*, 34(4):509–511, 1998.

Maryam Fazel, Rong Ge, Sham Kakade, and Mehran Mesbahi. Global convergence of policy gradient methods for the linear quadratic regulator. In Jennifer Dy and Andreas Krause, editors, *Proceedings of the 35th International Conference on Machine Learning*, volume 80 of *Proceedings of Machine Learning Research*, pages 1467–1476. PMLR, 10–15 Jul 2018.

Abraham D. Flaxman, Adam Tauman Kalai, and H. Brendan McMahan. Online convex optimization in the bandit setting: gradient descent without a gradient. In *Proceedings of the Sixteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA '05, page 385–394, USA, 2005. Society for Industrial and Applied Mathematics.

Maxime Gautier and Ph Poignet. Extended kalman filtering and weighted least squares dynamic identification of robot. *Control Engineering Practice*, 9(12):1361–1372, 2001.

Moritz Hardt, Ben Recht, and Yoram Singer. Train faster, generalize better: Stability of stochastic gradient descent. In *International Conference on Machine Learning*, pages 1225–1234. PMLR, 2016.

Arieh Iserles. *Ordinary differential equations*, page 1–2. Cambridge Texts in Applied Mathematics. Cambridge University Press, 2 edition, 2008.

Chi Jin, Rong Ge, Praneeth Netrapalli, Sham M. Kakade, and Michael I. Jordan. How to escape saddle points efficiently. In Doina Precup and Yee Whye Teh, editors, *Proceedings of the 34th International Conference on Machine Learning*, volume 70 of *Proceedings of Machine Learning Research*, pages 1724–1732. PMLR, 06–11 Aug 2017.

Chi Jin, Praneeth Netrapalli, and Michael I. Jordan. Accelerated gradient descent escapes saddle points faster than gradient descent. In Sébastien Bubeck, Vianney Perchet, and Philippe Rigollet, editors, *Proceedings of the 31st Conference On Learning Theory*, volume 75 of *Proceedings of Machine Learning Research*, pages 1042–1085. PMLR, 06–09 Jul 2018.

Claude Lemaréchal. Cauchy and the gradient method. *Doc Math Extra*, 251(254):10, 2012.

Gabriele Lillacci and Mustafa Khammash. Parameter estimation and model selection in computational biology. *PLoS computational biology*, 6(3):e1000696, 2010.

Dhruv Malik, Ashwin Pananjady, Kush Bhatia, Koulik Khamaru, Peter Bartlett, and Martin Wainwright. Derivative-free methods for policy optimization: Guarantees for linear quadratic systems. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 2916–2925. PMLR, 2019.

Izumi Masubuchi, Atsumi Ohara, and Nobuhide Suda. LMI-based controller synthesis: A unified formulation and solution. *International Journal of Robust and Nonlinear Control*, 8(8):669–686, 1998. doi: https://doi.org/10.1002/(SICI)1099-1239(19980715)8:8<669::AID-RNC337>3.0.CO;2-W.

Hesameddin Mohammadi, Armin Zare, Mahdi Soltanolkotabi, and Mihailo R Jovanovic. Convergence and sample complexity of gradient methods for the model-free linear quadratic regulator problem. *IEEE Transactions on Automatic Control*, 2021.

Kevin Scaman and Cedric Malherbe. Robustness analysis of non-convex stochastic gradient descent using biased expectations. *Advances in Neural Information Processing Systems*, 33, 2020.

C. Scherer, P. Gahinet, and M. Chilali. Multiobjective output-feedback control via LMI optimization. *IEEE Transactions on Automatic Control*, 42(7):896–911, 1997. doi: 10.1109/9.599969.

Carsten Scherer. Mixed H2 /H∞ control. *Trends in control*, pages 173–216, 1995.

Yue Sun and Maryam Fazel. Learning optimal controllers by policy gradient: Global optimality via convex parameterization. In *2021 60th IEEE Conference on Decision and Control (CDC)*, pages 4576–4581. IEEE, 2021.

Yujie Tang, Yang Zheng, and Na Li. Analysis of the optimization landscape of linear quadratic Gaussian (LQG) control. In *Learning for Dynamics and Control*, pages 599–610. PMLR, 2021.

Joel A Tropp. An introduction to matrix concentration inequalities. *arXiv preprint arXiv:1501.01571*, 2015.

Stephen Wright, Jorge Nocedal, et al. Numerical optimization. *Springer Science*, 35(67-68):7, 1999.

Bin Yu. Rates of convergence for empirical processes of stationary mixing sequences. *The Annals of Probability*, pages 94–116, 1994.

Kaiqing Zhang, Bin Hu, and Tamer Basar. Policy optimization for $\mathcal{H}_2$ linear control with $\mathcal{H}_\infty$ robustness guarantee: Implicit regularization and global convergence. In Alexandre M. Bayen, Ali Jadbabaie, George Pappas, Pablo A. Parrilo, Benjamin Recht, Claire Tomlin, and Melanie Zeilinger, editors, *Proceedings of the 2nd Conference on Learning for Dynamics and Control*, volume 120 of *Proceedings of Machine Learning Research*, pages 179–190. PMLR, 10–11 Jun 2020.

Kemin Zhou, John Comstock Doyle, Keith Glover, et al. *Robust and Optimal Control*, volume 40. Prentice hall New Jersey, 1996.