Approximate Secular Equations for the Cubic Regularization Subproblem

Abstract

The cubic regularization method (CR) is a popular algorithm for unconstrained non-convex optimization. At each iteration, CR solves a cubically regularized quadratic problem, called the cubic regularization subproblem (CRS). One way to solve the CRS relies on solving the secular equation, whose computational bottleneck lies in the computation of all eigenvalues of the Hessian matrix. In this paper, we propose and analyze a novel CRS solver based on an approximate secular equation, which requires only some of the Hessian eigenvalues and is therefore much more efficient. Two approximate secular equations (ASEs) are developed. For both ASEs, we first study the existence and uniqueness of their roots and then establish an upper bound on the gap between the root and that of the standard secular equation. Such an upper bound can in turn be used to bound the distance from the approximate CRS solution based ASEs to the true CRS solution, thus offering a theoretical guarantee for our CRS solver. A desirable feature of our CRS solver is that it requires only matrix-vector multiplication but not matrix inversion, which makes it particularly suitable for high-dimensional applications of unconstrained non-convex optimization, such as low-rank recovery and deep learning. Numerical experiments with synthetic and real data-sets are conducted to investigate the practical performance of the proposed CRS solver. Experimental results show that the proposed solver outperforms two state-of-the-art methods.

1 Introduction

The cubic regularization method (CR) is a variant of Newton's method proposed by Griewank [8], and later independently by Nesterov and Polyak [10], and Weiser et al. [14]. It gained significant attention over the last decade due to its attractive theoretical properties, such as convergence to second-order critical points[10] and quadratic convergence rate under mild assumptions [15]. Each iteration of CR solves a problem of the following form, called the cubic regularization subproblem (CRS):

$$\min_{\mathbf{x} \in \mathbb{R}^n} f_{\mathbf{A},\mathbf{b},\rho}(\mathbf{x}) := \mathbf{b}^{\mathrm{T}}\mathbf{x} + \frac{1}{2}\mathbf{x}^{\mathrm{T}}\mathbf{A}\mathbf{x} + \frac{\rho}{3}\|\mathbf{x}\|^3, \tag{1}$$

where $\rho > 0$ is the regularization parameter, $\mathbf{b} \in \mathbb{R}^n$, and $\mathbf{A} \in \mathbb{R}^{n \times n}$ is a symmetric matrix, not necessarily positive semidefinite. Many variants and generalizations of CR are developed, including the Adaptive Regularization Using Cubics (ARC) which allows for a dynamic choice of ρ and inexact CRS solutions [3, 4], accelerated CR using momentum [13] and stochastic CR for solving stochastic optimization [12]. Despite the theoretical success, the practicality of CR and its variants relies critically on the efficiency of the CRS solver. The goal of this paper is to develop a novel, efficient CRS solver along with theoretical guarantees.

A popular approach for solving the CRS is via solving the so-called secular equation. We now review this approach. Towards that, we denote by $\lambda_1 \leq \cdots \leq \lambda_n$ the eigenvalues of $\mathbf{A}$ and

by $\mathbf{v}_1, \cdots, \mathbf{v}_n$ the corresponding eigenvectors. In other words, we have the eigendecomposition $\mathbf{A} = \sum_{i=1}^n \lambda_i \mathbf{v}_i \mathbf{v}_i^T = \mathbf{V}\mathbf{\Lambda}\mathbf{V}^T$, where $\mathbf{\Lambda} = \mathrm{diag}(\lambda_1, \ldots, \lambda_n)$ and $\mathbf{V} = [\mathbf{v}_1, \cdots, \mathbf{v}_n]$. Note that eigenvalues λ_i are not necessarily positive due to the indefiniteness of the matrix $\mathbf{A}$. Also, we denote the Euclidean norm by $\|\cdot\|$.

Proposition 1 ([10, 3]). *A vector $\mathbf{x}^*$ solves the CRS* (1) *if and only if it satisfies the system*

$$\begin{cases} (\mathbf{A} + \rho\|\mathbf{x}^*\|\mathbf{I})\mathbf{x}^* + \mathbf{b} = \mathbf{0}, & (2) \\ \mathbf{A} + \rho\|\mathbf{x}^*\|\mathbf{I} \succeq \mathbf{0}. & (3) \end{cases}$$

Moreover, if $\mathbf{A} + \rho\|\mathbf{x}^*\|\mathbf{I} \succ \mathbf{0}$, then $\mathbf{x}^*$ is the unique solution (and hence a critical point).

Proposition 2 ([1]). *Let $\mathbf{x}^*$ be a global solution of CRS* (1) *and the eigendecomposition for $\mathbf{A} = \sum_{i=1}^n \lambda_i \mathbf{v}_i \mathbf{v}_i^T = \mathbf{V}\mathbf{\Lambda}\mathbf{V}^T$, where $\mathbf{\Lambda} = diag(\lambda_1, \ldots, \lambda_n)$ and $\mathbf{V} = [\mathbf{v}_1, \cdots, \mathbf{v}_n]$. If $\mathbf{b}^T\mathbf{v}_1 \neq 0$, then $\mathbf{A} + \rho\|\mathbf{x}^*\|\mathbf{I} \succ \mathbf{0}$ and the solution $\mathbf{x}^*$ is the unique critical point (and hence the unique solution). Conversely, if $\mathbf{b}^T\mathbf{v}_1 = 0$, then the CRS* (1) *has multiple optimal solutions.*

From Proposition 2, if $\mathbf{b}^T\mathbf{v}_1 \neq 0$, then there is only one critical point, which is also the optimal solution, and hence the gradient norm $\|\nabla f_{\mathbf{A}, \mathbf{b}, \rho}(\mathbf{x})\|$ serves as an optimality measure. Throughout the paper, we assume $\mathbf{b}^T\mathbf{v}_1 \neq 0$, under which the CRS is said to be in the easy case. This is without much loss of generality as this holds generically true in practice. Moreover, we could easily avoid the hard case ($\mathbf{b}^T\mathbf{v}_1 = 0$) by slightly perturbing the vector $\mathbf{b}$, see [10, 2].

To introduce the secular equation, note that in the easy case, conditions (2) and (3) can be written as

$$\begin{cases} (\mathbf{\Lambda} + \sigma\mathbf{I}) \cdot \mathbf{y}^* = \mathbf{c}, \\ \lambda_1 + \sigma > 0. \end{cases}$$

where $\sigma =: \rho\|\mathbf{x}^*\|$, $[y_1^*, \cdots, y_n^*]^T := \mathbf{y}^* = \mathbf{V}^T\mathbf{x}^*$ and $[c_1, \cdots, c_n]^T := \mathbf{c} = -\mathbf{V}^T\mathbf{b}$. Therefore,

$$y_i^* = \frac{c_i}{\lambda_i + \sigma}, \qquad i = 1, \ldots, n.$$

Since the Euclidean norm is invariant to orthogonal transformation, we have

$$\frac{\sigma^2}{\rho^2} = \|\mathbf{x}^*\|^2 = \|\mathbf{y}^*\|^2 = \sum_{i=1}^n \frac{c_i^2}{(\lambda_i + \sigma)^2}.$$

Consequently, instead of solving the complicated nonlinear system (2)-(3), we could solve the CRS (1) by first finding the (unique) root $\sigma > \max\{-\lambda_1, 0\}$ of the equation

$$w(\sigma) = \sum_{i=1}^n \frac{c_i^2}{(\lambda_i + \sigma)^2} - \frac{\sigma^2}{\rho^2}, \tag{4}$$

called the secular equation, and then solves the linear system $(\mathbf{A} + \sigma\mathbf{I})\mathbf{x} = -\mathbf{b}$. The first step can be done efficiently by using existing root-finding algorithms (e.g., the bisection method and Newton's method etc.).

The disadvantage of the above CRS solver, based on the secular equation (4), is that it requires the full spectrum of $\mathbf{A}$, which costs $\mathcal{O}(n^3)$. This approach is viable only for low- to moderate-dimensional problems. However, when n is large, computing all eigenvalues of $\mathbf{A}$ is prohibitive. Worse still, after the root σ is solved, we still need to apply iterative methods (e.g., Lanczos method) to solve the large-scale linear system (2). We are thus motivated approximate the secular equation by using only some of the eigenvalues of $\mathbf{A}$, as opposed to all.

As our main contribution, we developed two different approximate secular equations (ASEs), both of which require computing $m < n$ eigenvalues of $\mathbf{A}$. The cost for forming the approximate secular equations is only $\mathcal{O}(mn^2)$, and hence the resulting CRS solver is much more efficient and scalable. On the theoretical side, for each of the proposed approximate secular equations, we first studied the existence and uniqueness of its root, and then derived an upper bound on the gap between the root and that of the standard secular equation (4). This upper bound can in turn be used to bound the distance from the approximate CRS solution based ASEs to the true CRS solution, thus offering a theoretical guarantee for the resulting CRS solver. A desirable feature of our CRS solver is that it requires only matrix-vector multiplication but not matrix inversion, which makes it particularly suitable for

73 high-dimensional applications of unconstrained non-convex optimization, such as low-rank recovery
74 and deep learning. On the empirical side, we conducted experiments with both synthetic and real
75 problem instances to investigate the practical performance of the proposed CRS solver and the
76 associated CR. Experimental results showed that the proposed solver outperforms two state-of-the-art
77 methods. The selection of m for the proposed ASEM is an interesting and crucial topic. We will
78 discuss related issues in section 4 and some numerical explorations are also conducted in section 5.

2 The First-Order Truncated Secular Equation

80 We define the first-order truncated secular equation by

$$w_1(\sigma; \mu) = \sum_{i=1}^{m} \frac{c_i^2}{(\lambda_i + \sigma)^2} + \sum_{i=m+1}^{n} \frac{c_i^2}{(\mu + \sigma)^2} - \frac{\sigma^2}{\rho^2}, \tag{5}$$

81 where $\mu \geq \lambda_m$ is an input parameter that approximates the unobserved eigenvalues $\lambda_{m+1}, \cdots, \lambda_n$,
82 $c_i = -\mathbf{b}^{\mathrm{T}} \mathbf{v}_i$ and $\sum_{i=m+1}^{n} c_i^2 = \|\mathbf{b}\|^2 - \sum_{i=1}^{m} c_i^2$. Note that only m eigenvalues $\lambda_1, \cdots, \lambda_m$ and
83 their corresponding eigenvectors $\mathbf{v}_1, \cdots, \mathbf{v}_m$ are needed to form (5), which is computationally
84 friendlier compared with (4). The name first-order truncated secular equation comes from the fact
85 that $w_1(\sigma, \mu)$ is the first-order Taylor approximation to the function $w(\sigma)$. Below we will first study
86 the existence and uniqueness for the root of (5). Then, we derive an error bound for the root.

2.1 Existence and Uniqueness for the Root

88 In the easy case that $\mathbf{b}^{\mathrm{T}} \mathbf{v}_1 \neq 0$ (equivalently, $c_1 \neq 0$), the solution $\mathbf{x}^*$ to the CRS (1) is unique, which
89 implies the existence and uniqueness for the root σ^* of (4). To show that our proposed CRS solver
90 is also well-defined, we prove the existence and uniqueness of the root of the first-order truncated
91 secular equation (5).

92 **Lemma 1.** *For any $\mu \geq \lambda_m$, the function $w_1(\cdot; \mu)$ as defined in (5) admits a unique root.*

93 *Proof.* **Existence.** We first consider the case when $\lambda_1 \leq 0$. Then, for any fixed $\mu \geq \lambda_m$,

$$\lim_{\sigma \to (-\lambda_1)^+} w_1(\sigma; \mu) = +\infty \quad \text{and} \quad \lim_{\sigma \to +\infty} w_1(\sigma; \mu) = -\infty,$$

94 By the intermediate value theorem, $w_1(\cdot; \mu)$ has a root in $(-\lambda_1, +\infty)$. For $\lambda_1 > 0$, we have

$$w_1(0; \mu) > 0 \quad \text{and} \quad \lim_{\sigma \to +\infty} w_1(\sigma; \mu) = -\infty.$$

95 Therefore, $w_1(\sigma; \mu)$ has a root in $(0, +\infty)$.

96 **Uniqueness.** Note that $w_1(\sigma; \mu)$ is monotonically decreasing for $\sigma \in (-\lambda_1, +\infty)$ and $\sigma \in (0, +\infty)$
97 when $\lambda_1 \leq 0$ and $\lambda_1 > 0$, respectively. Therefore, the uniqueness of the root for $w_1(\sigma; \mu)$ is
98 guaranteed. □

2.2 Error Analysis

100 In order to study the quality of the CRS solution based on our proposed solver using approximate
101 secular equations, we need the quality of the root to the first-order truncated secular equation, denoted
102 by σ_1^*. Towards that end, we provide an upper bound on the gap $|\sigma_1^* - \sigma^*|$ between σ_1^* and the root σ^*
103 of the exact secular equation (4).

104 **Theorem 1.** *Let σ_1^* and σ^* be the unique roots of $w_1(\sigma; \mu)$ and $w(\sigma)$, respectively. Then*

$$|\sigma_1^* - \sigma^*| \leq C_m \cdot \max_{m+1 \leq i \leq n} |\lambda_i - \mu|, \tag{6}$$

105 *where $C_m > 0$ is a constant, upper bounded by $\frac{2\|\mathbf{b}\|^2}{(\lambda_m - \lambda_1)^3} \cdot \min\left\{\frac{(\lambda_d + B_1)^3}{2\|\mathbf{b}\|^2}, \frac{\rho^2}{2B_1}\right\}$ with $B_1 =$*
106 *$\frac{-\lambda_1 + \sqrt{\lambda_1^2 + 4\rho \cdot \|\mathbf{b}\|}}{2}$ being an upper bound for $|\sigma_1^*|$.*

107 We clearly see that the right-hand side of inequality (1) is decreasing in m. This confirms that using
108 more eigen information (i.e., larger m) helps to reduce the error $|\sigma_1^* - \sigma^*|$. The proof of Theorem 1
109 is technical and quite long and hence relegated to Appendix A. The approximation quality of our
110 CRS solver is guaranteed by combining Theorem 1 with the following proposition.

Proposition 3. *Let $\mathbf{x}^*$ and $\tilde{\mathbf{x}}$ be solutions to the equations $(\mathbf{A} + \sigma^*\mathbf{I})\mathbf{x}^* = -\mathbf{b}$ and $(\mathbf{A} + \sigma_1^*\mathbf{I})\tilde{\mathbf{x}} = -\mathbf{b}$, respectively. Then, $\|\tilde{\mathbf{x}} - \mathbf{x}^*\| = \mathcal{O}(|\sigma_1^* - \sigma^*|)$.*

Proof. By definition, we have

$$\mathbf{x}^* = \sum_{i=1}^{n}(\lambda_i + \sigma^*)^{-1}\mathbf{v}_i\mathbf{v}_i^{\mathrm{T}} \cdot (-\mathbf{b}) = \sum_{i=1}^{n}(\lambda_i + \sigma^*)^{-1}c_i \cdot \mathbf{v}_i,$$

and

$$\tilde{\mathbf{x}} = \sum_{i=1}^{n}(\lambda_i + \sigma_1^*)^{-1}\mathbf{v}_i\mathbf{v}_i^{\mathrm{T}} \cdot (-\mathbf{b}) = \sum_{i=1}^{n}(\lambda_i + \sigma_1^*)^{-1}c_i \cdot \mathbf{v}_i,$$

then

$$
\begin{aligned}
\|\tilde{\mathbf{x}} - \mathbf{x}^*\| &= \left\|\sum_{i=1}^{n}\left((\lambda_i + \sigma_1^*)^{-1} - (\lambda_i + \sigma^*)^{-1}\right)\mathbf{v}_i\mathbf{v}_i^{\mathrm{T}} \cdot (-\mathbf{b})\right\| \\
&\leq \left(\max_{1 \leq i \leq n}\left\{(\lambda_i + \sigma_1^*)^{-1} - (\lambda_i + \sigma^*)^{-1}\right\}\right) \cdot \|\mathbf{b}\| \\
&= \mathcal{O}(|\sigma_1^* - \sigma^*|).
\end{aligned}
$$

This completes the proof. □

Before ending this section, some remarks are in order. First, the parameter μ acts as an approximation to $n - m$ unknown eigenvalues $\lambda_{m+1}, \cdots, \lambda_n$. An intuitive choice of μ that works well in practice and is computationally cheap is the average of unknown eigenvalues, i.e.,

$$\mu_1 = \frac{\sum_{i=m+1}^{n}\lambda_i}{n - m} = \frac{\mathrm{tr}(\mathbf{A}) - \sum_{i=1}^{m}\lambda_i}{n - m}. \tag{7}$$

Second, the error bound $C_m \cdot \max_{m+1 \leq i \leq n}|\lambda_i - \mu|$ in Theorem 1 depends on the distribution of eigenvalues of $\mathbf{A}$. If the unobserved eigenvalues $\lambda_{m+1}, \cdots, \lambda_n$ cluster around a small interval, then with a suitable choice of $\mu \in [\lambda_{m+1}, \lambda_n]$, $\max_{m+1 \leq i \leq n}|\lambda_i - \mu|$ is small. Conversely, if the unknown eigenvalues spread over a large interval, then it is hard to make the error $\max_{m+1 \leq i \leq n}|\lambda_i - \mu|$ small.

Third, it is instructive to study the error bound (6) under some random matrix model for $\mathbf{A}$. Suppose that $\mathbf{A} = \widetilde{\mathbf{A}}/\sqrt{2n}$, where $\widetilde{\mathbf{A}}$ is a symmetric random matrix with i.i.d. entries on and above the diagonal. By the Wigner semicircle law [6], as $n \to \infty$, the eigenvalues of $\mathbf{A}$ distribute according to a density of a semi-circle shape. In particular, we can deduce that with probability of $1 - o(1)$,

$$\max_{m+1 \leq i \leq n}|\lambda_i - \mu| \leq \mathcal{O}\left(\left(1 - \frac{m+1}{n}\right)^{2/3}\right) \approx \left(\frac{3\pi}{4\sqrt{2}}\right)^{2/3} \cdot \left(1 - \frac{m+1}{n}\right)^{2/3} \tag{8}$$

The detailed proof of (8) and further discussions under random $\mathbf{A}$ can be found in Appendix C.

3 The Second-Order Truncated Secular Equation

Similarly to the equation (5), but with the second-order Taylor approximation, we define the second-order truncated secular equation by

$$w_2(\sigma; \mu) = \sum_{i=1}^{m}\frac{c_i^2}{(\lambda_i + \sigma)^2} + \sum_{i=m+1}^{n}\frac{c_i^2}{(\mu + \sigma)^2} - 2\sum_{i=m+1}^{n}\frac{c_i^2 \cdot (\lambda_i - \mu)}{(\mu + \sigma)^3} - \frac{\sigma^2}{\rho^2}, \tag{9}$$

where $\mu \geq \lambda_m$ is an input parameter that approximates the unobserved eigenvalues $\lambda_{m+1}, \cdots, \lambda_n$.

3.1 Existence and Uniqueness for the Root

The lemma blow shows the existence and uniqueness of the root of $w_2(\cdot; \mu)$.

Lemma 2. *With*

$$\mu = \frac{\sum_{i=m+1}^{n} c_i^2 \cdot \lambda_i}{\sum_{i=m+1}^{n} c_i^2},$$ (10)

the function $w_2(\cdot; \mu)$ as defined in (9) admits a unique root.

Proof. When

$$\mu = \frac{\sum_{i=m+1}^{n} c_i^2 \cdot \lambda_i}{\sum_{i=m+1}^{n} c_i^2},$$

the third summation in the definition (9) vanishes, and hence $w_2(\sigma, \mu)$ becomes the same as $w_1(\sigma, \mu)$, except with a specific choice of μ. The desired conclusion then follows from Lemma 1. □

Unlike its first-order counterpart, we do not develop the existence and uniqueness of the root of the second-order truncated secular equation for arbitrary μ. The reason is that when $\lambda_1 > 0$, $w_2(0; \mu)$ can potentially be positive or negative.

3.2 Error Analysis

Similar to that for the first-order truncated secular equation, we can also derive an error bound for the root of the second-order truncated secular equation.

Theorem 2. *Let σ_2^* and σ^* be the unique root of $w_2(\sigma; \mu)$ and $w(\sigma)$, respectively, and*

$$\mu = \frac{\sum_{i=m+1}^{n} c_i^2 \cdot \lambda_i}{\sum_{i=m+1}^{n} c_i^2}.$$

Then,

$$|\sigma_2^* - \sigma^*| \le C_m \cdot \max_{m+1 \le i \le n} (\lambda_i - \mu)^2,$$ (11)

where $C_m > 0$ is a constant bounded by $\frac{3\|\mathbf{b}\|^2}{(\lambda_m - \lambda_1)^4} \cdot \min\left\{ \frac{(\lambda_n + B_1)^3}{2\|\mathbf{b}\|^2}, \frac{\rho^2}{2B_1} \right\}$ with $B_1 = \frac{-\lambda_1 + \sqrt{\lambda_1^2 + 4\rho \cdot \|\mathbf{b}\|}}{2}$ being an upper bound for $|\sigma_2^*|$.

The proof of Theorem 2 can be found in Appendix B. We can similarly estimate the approximation quality by combining Theorem 2 and Proposition 3. Again, the right-hand side of the error bound (11) is decreasing in m. We should also point out that the CRS solver based on the second-order secular equation outperforms the first-order counterpart only if $\max_{m+1 \le i \le n} |\lambda_i - \mu|/|\lambda_m - \lambda_1|$ is small enough. The computation of μ here requires $c_{m+1}, \cdots, c_n$, which seem to be inaccessible. We provide a tractable form for μ in (13) and will discuss it in the next part.

4 Implementation Details

We now discuss the implementation details for solving the proposed first-order secular equation for CRS. First, we obtain the partial eigen information $\{\lambda_1, \cdots, \lambda_m\}$ and $\{\mathbf{v}_1, \cdots, \mathbf{v}_m\}$ by Krylov subspace methods. Note that only Hessian-vector products are required. This is computationally friendlier than other methods that rely on matrix inversions and particularly suitable for modern, high-dimensional applications. Then, we solve the first-order secular equation (5) with μ defined in (7) or (10), using anyroot-finding algorithm, such as Newton's method. Finally, we solve the linear system $(\mathbf{A} + \sigma^* \mathbf{I})\mathbf{x} = -\mathbf{b}$ by iterative algorithms, e.g., the Lanczos method and the conjugate gradient method. The resulting CRS solver, namely the approximate secular equation method (ASEM), is summarized as follows:

Step 1: obtaining the partial eigen information $\{\lambda_1, \cdots, \lambda_m\}$ and $\{\mathbf{v}_1, \cdots, \mathbf{v}_m\}$ of $\mathbf{A}$.

Step 2: solving the secular equation (5) with μ defined in (7) or (10); we get σ^*.

Step 3: iteratively solving the linear system $(\mathbf{A} + \sigma^* \mathbf{I})\mathbf{x} + \mathbf{b} = \mathbf{0}$.

Output: the solution $\mathbf{x}$.

Details for step 1. The Krylov subspace is one of the most popular iterative methods in solving eigen problems with $\mathcal{O}(mn^2)$ computational cost [11]. The Lanczos decomposition for a real symmetric matrix $\mathbf{B}$ satisfies

$$\mathbf{B}\mathbf{U}_k = \mathbf{T}_k\mathbf{U}_k + \beta_k\mathbf{u}_{k+1}\mathbf{e}_k^{\mathrm{T}},$$

where $\mathbf{U}_k \in \mathbb{R}^{n\times k}$ is an orthonormal matrix (i.e., $\mathbf{U}_k^{\mathrm{T}}\mathbf{U}_k = \mathbf{I}_k$), $\mathbf{T}_k \in \mathbb{R}^{k\times k}$ is a symmetric tridiagonal matrix and $\mathbf{e}_k \in \mathbb{R}^k$ is the k-th standard basis vector in $\mathbb{R}^k$. Lanczos observed that even for comparatively small k, $\mathbf{T}_k$ has significant approximation to eigenvalues and eigenvectors of $\mathbf{B}$. Specifically, for a suitable eigenpair $(\gamma, \mathbf{w})$ of $\mathbf{T}_k$ with $\mathbf{T}_k\mathbf{w} = \gamma \cdot \mathbf{w}$, the pair $(\gamma, \mathbf{U}_k\mathbf{w})$ is an approximation to an eigenpair of $\mathbf{B}$, i.e., $\mathbf{B}\mathbf{z} \approx \gamma \cdot \mathbf{z}$ with $\mathbf{z} = \mathbf{U}_k\mathbf{w}$. Here, the Krylov subspace is constructed by $\mathbf{u}_1$, the first column of $\mathbf{U}_k = [\mathbf{u}_1, \cdots, \mathbf{u}_k]$, i.e., $\mathcal{K}_k(\mathbf{B}, \mathbf{u}_1)$. Note that $\mathbf{T}_k$ approximates $\mathbf{B}$ for eigenvalues with largest modulus (or absolute values) and the corresponding eigenvectors. Empirically, for calculating m eigenvalues of $\mathbf{B}$ with largest absolute values and the corresponding eigenvectors, we usually construct the Krylov subspace $\mathcal{K}_k(\mathbf{B}, \mathbf{u}_1)$ with dimension $k = \max\{2m, 20\}$. Moreover, the base vector $\mathbf{u}_1$ is also essential for the Krylov subspace method. Moreover, restarting is adopted to iteratively update the base vector $\mathbf{u}_1$. To the best of our knowledge, the mentioned iterative method for partial eigen informnation is supported in many softwares, e.g., Matlab (eigs function) and Python (Scipy package) etc. For more details, please refer to ARPACK [9]. Returning back to the proposed algorithm with $\mathbf{A}$, instead of calculating the largest (in terms of the absolute value) m eigenvalues of $\mathbf{A}$, we aim to get m (algebraically) smallest eigenvalues $\{\lambda_1, \cdots, \lambda_m\}$ and the corresponding eigenvectors $\{\mathbf{v}_1, \cdots, \mathbf{v}_m\}$. We first roughly calculate a shift value $\beta \geq \|\mathbf{A}\|$ by several steps of power iteration (Hessian-vector products). Then, let $\mathbf{B} = \beta\cdot\mathbf{I} - \mathbf{A}$, whose eigenvalues $\{\beta - \lambda_n, \cdots, \beta - \lambda_1\}$ are non-negative and the corresponding eigenvectors are $\{\mathbf{v}_n, \cdots, \mathbf{v}_1\}$. Applying the mentioned Krylov subspace algorithm, we first obtain an estimate of shifted eigenvalues $\{\lambda_1, \cdots, \lambda_m\}$ and the corresponding eigenvectors $\{\mathbf{v}_1, \cdots, \mathbf{v}_m\}$ for $\mathbf{A}$, since $\{\beta - \lambda_m, \cdots, \beta - \lambda_1\}$ are largest eigenvalues of $\mathbf{B}$. To further lower the computational cost, we may adopt $k = m$ dimensional Krylov subspace $\mathcal{K}_k(\mathbf{B}, \mathbf{u}_1)$ for m eigenvalues without restarting in implementation.

Details for step 2. Instead of directly solving (5), Cartis et al. [3] recommended to find the root for the equivalent equation:

$$\tilde{w}_1(\sigma; \mu) = \sqrt{\sum_{i=1}^{m} \frac{c_i^2}{(\lambda_i + \sigma)^2} + \sum_{i=m+1}^{n} \frac{c_i^2}{(\mu + \sigma)^2}} - \frac{\sigma}{\rho}, \tag{12}$$

which is convex on $(-\lambda_1, +\infty)$. Moreover, Newton method is proved to achieve (locally) quadratic convergence with perfect initialization. However, we numerically find that it depends much on the initialization and may converge to a point outside the feasible domain $(-\lambda_1, +\infty)$ if it has imperfect initialization. Here, we recommend to use the bisection method to find the root of (5) or (12) due to its linear convergence, stability (not beyond the feasible domain) and simple implementation. For the weighted average μ defined in (10), we can rewrite it as a more tractable but equivalent form that

$$\mu_2 = \frac{\sum_{i=m+1}^{n} c_i^2 \cdot \lambda_i}{\sum_{i=m+1}^{n} c_i^2} = \frac{\mathbf{b}^{\mathrm{T}}(\mathbf{A} - \mathbf{V}_m\mathbf{\Lambda}_m\mathbf{V}_m^{\mathrm{T}})\mathbf{b}}{\|\mathbf{b}\|^2 - \sum_{i=1}^{m} c_i^2} = \frac{\mathbf{b}^{\mathrm{T}}\mathbf{A}\mathbf{b} - \sum_{i=1}^{m} c_i^2 \cdot \lambda_i}{\|\mathbf{b}\|^2 - \sum_{i=1}^{m} c_i^2}, \tag{13}$$

where $\mathbf{V}_m = [\mathbf{v}_1, \cdots, \mathbf{v}_m]$ and $\mathbf{\Lambda_m} = \mathrm{diag}(\lambda_1, \cdots, \lambda_m)$.

Details for step 3. There are a bunch of and well-behaved iterative methods for (real symmetric) linear systems, e.g., Krylov subspace (Lanczos) methods and conjugate gradient methods etc. We do not discuss any details here since solving linear systems has been investigated for decades and achieved great success. We adopt the Lanczos method for solving the linear system $(\mathbf{A} + \sigma^*\mathbf{I})\mathbf{x} + \mathbf{b} = \mathbf{0}$, where multiple steps of Hessian-vector products are required.

In summary, the main computational cost comes from step 1 and step 3 for Hessian-vector products ($\mathcal{O}(mn^2)$), since $w_1(\sigma; \mu)$ is a one dimensional problem for σ in step 2 and is of $\mathcal{O}(n)$. Therefore, the total computational cost for the proposed algorithm is of $\mathcal{O}(mn^2)$, much lower than the method with full eigendecomposition ($\mathcal{O}(n^3)$).

The selection of m. We seek to set proper m in ASEM for solving CRS because overly large m would cause computational waste while too small m cannot achieve our expected accuracy. To achieve ε-accuracy (i.e., $|\sigma_1^* - \sigma^*| \leq \varepsilon$), with random Gaussian matrix $\mathbf{A}$, we may set $m \leq n$ such

that

$$\left(\frac{3\pi}{4\sqrt{2}}\right)^{2/3} \cdot \left(1 - \frac{m+1}{n}\right)^{2/3} \le \varepsilon,$$

according to (8). However, the error bound (8) may be overestimated for m. Moreover, for general problems without priors for eigen distribution of $\mathbf{A}$, it is hard to estimate m based on expected ε-accuracy, because the error bounds (6) and (11) are implicit for m. Therefore, adaptive methods (or heuristic methods) for selecting m are essential and urgent in practice. An intuitive way is to check the suboptimality (gradient norm) in each step, and enlarge m by m_0 (i.e., we set $m = \max\{m + m_0, m_{\max}\}$, where $m_{\max}$ is the maximal number of eigenvalues we adopt in ASEM), if the output does not satisfy the given condition for suboptimality. Moreover, numerical experiments on CUTEst problems (see Experiment 6 in section 5) shows that $m = 1$ is enough for most of the cases. We left the study for the selection of m as a future work. To the best of our knowledge, it is also unsolved for the Krylov subspace method [3, 2] in solving CRS.

5 Experimental Results

Without the loss of generality (see Appendix D), we adopt diagonal matrix $\mathbf{A}$ in the synthetic cubic regularized quadratic problem, for simplicity and fair comparison. Furthermore, we also test the proposed ASEM on CUTEst library [7]. All experiments were run on a Macbook Pro M1 laptop. For more experimental details, please refer to Appendix E.

Experiment 1. The distribution for eigenvalues of the matrix $\mathbf{A}$. In the error analysis (Theorem 1 and Theorem 2), the error is controlled by the approximation of μ to unobserved eigenvalues $\{\lambda_i\}_{i=m+1}^{n}$, i.e., $\max_{m+1 \le i \le n} |\lambda_i - \mu|$ and $\max_{m+1 \le i \le n} (\lambda_i - \mu)^2$. It further implies that the distribution of eigenvalues $\{\lambda_i\}_{i=m+1}^{n}$ is essential for the proposed method. Intuitively, if eigenvalues $\{\lambda_i\}_{i=m+1}^{n}$ cluster around a small interval, then the rough estimate (7) for μ is enough to approximate the unknown eigenvalues well. Conversely, if eigenvalues $\{\lambda_i\}_{i=m+1}^{n}$ are spread over a large interval, then we cannot expect a single μ to estimate all eigenvalues $\{\lambda_i\}_{i=m+1}^{n}$. Here, we have four specially designed cases for distributions of eigenvalues of the matrix $\mathbf{A}$ to illustrate our theoretical observations for the proposed method. Case 1 (evenly spaced): all eigenvalues $\{\lambda_i\}_{i=1}^{n}$ are evenly spaced in $[-1, 1]$; Case 2 (separated): half of eigenvalues are far away from the remaining, i.e., eigenvalues $\{\lambda_i\}_{i=1}^{n/2}$ are evenly spaced in $[-1, -4/5]$ and the remaining eigenvalues $\{\lambda_i\}_{i=n/2+1}^{n}$ are evenly spaced in $[4/5, 1]$; Case 3 (right centered): the minimal 2% of eigenvalues and the remaining 98% of eigenvalues gather together respectively, i.e., eigenvalues $\{\lambda_i\}_{i=1}^{n/50}$ and $\{\lambda_i\}_{i=n/50+1}^{n}$ are spaced evenly in $[-1, 4/5]$ and $[4/5, 1]$ respectively; Case 4 (left centered): the maximal 2% of eigenvalues and the remaining 98% of eigenvalues gather together respectively, i.e., eigenvalues $\{\lambda_i\}_{i=1}^{49/50n}$ and $\{\lambda_i\}_{i=49/50n+1}^{n}$ are evenly spaced in $[-1, 4/5]$ and $[4/5, 1]$ respectively. The vector $\mathbf{b}$ is proportional to $[1, \cdots, 1]^{\mathrm{T}}$ with $\|\mathbf{b}\| = 0.1$. The remaining parameters are $n = 5 \times 10^3$ and $\rho = 0.1$. We adopt the first-order method (i.e., μ is defined in (7)) here. Figure 1 validates our theories that the proposed algorithm converges fast if unknown eigenvalues $\{\lambda_i\}_{i=m+1}^{n}$ are close. Moreover, without the need to compute all eigenvalues, partial eigen information is enough to achieve satisfactory solutions in practice, except the hard case (e.g., case 4).

Figure 1: Trajectories of suboptimality (gradient norm $\|\nabla f_{\mathbf{A},\mathbf{b},\rho}(\mathbf{x})\|$) with different distributions for eigenvalues in Experiment 1.

Figure 2: Trajectories of suboptimality (gradient norm $\|\nabla f_{\mathbf{A},\mathbf{b},\rho}(\mathbf{x})\|$) with different μ in Experiment 2.

Figure 3: Trajectories of suboptimality (gradient norm $\|\nabla f_{\mathbf{A},\mathbf{b},\rho}(\mathbf{x})\|$) with exact and approximated eigenvalues and eigenvectors in Experiment 3.

Experiment 2. The effect of the parameter μ (first-order and second-order methods). For the first-order method, we select (7) for μ, while (10) for the second-order method based on our previous analysis. Here, we test the effect of μ for the proposed method in solving the cubic regularized quadratic problem, with other parameters fixed. Case 1 (first-order method): μ is adopted as the mean value of unknown eigenvalues, defined in (7); Case 2 (second-order method): μ is selected as the weighted average of unknown eigenvalues with respected to c_i^2 (please see (10)); Case 3 (first-order method): $\mu = \lambda_m$, the maximal eigenvalue we observe; Case 4: $\mu = 10^6$, much larger than eigenvalues of $\mathbf{A}$, as an approximation to $+\infty$. Different from the previous two experiments, the vector $\mathbf{b}$ is proportional to $[\lambda_1, \cdots, \lambda_n]^{\mathrm{T}}$ with length $\|\mathbf{b}\| = 0.1$. Eigenvalues of the matrix $\mathbf{A}$ are evenly spaced in $[-1, 1]$. The remaining parameters are $d = 5 \times 10^3$ and $\rho = 0.1$. Figure 2 shows the superiority of second-order method over the first-order method that it is more stable and converges faster when m is large, consistent with the theoretical analysis Theorem 1 and Theorem 2. Moreover, the results further imply the significance of the selection for μ. There are several observations from Figure 2. Firstly, we cannot discard the residual term with the unknown eigenvalues $\{\lambda_i\}_{i=m+1}^n$, where they still contain much information, as is shown in case 1 and case 4. Secondly, random selection of μ is not acceptable and may cause non-convergence (e.g., see case 3 and case 4). Furthermore, proper selection of μ contributes to the well-behaved algorithm (e.g., see case 1 and case 2).

Experiment 3. Approximation capabilities of the Krylov subspace method for ASEM. As is introduced in section 4, we adopt m dimensional Krylov subspace $\mathcal{K}_m(\mathbf{B}, \mathbf{u}_1)$ to approximately calculate m algebraically smallest eigenvalues $\{\lambda_1, \cdots, \lambda_m\}$ and the corresponding eigenvectors $\{\mathbf{v}_1, \cdots, \mathbf{v}_m\}$. Here, we investigate the performance of ASEM with estimated eigenvalues and eigenvectors. The vector $\mathbf{b}$ is proportional to the vector $[1, \cdots, 1]^{\mathrm{T}}$ with length $\|\mathbf{b}\| = 0.1$. Eigenvalues of the matrix $\mathbf{A}$ are evenly spaced in $[-1, 1]$. The remaining parameters are $n = 5 \times 10^3$ and $\rho = 0.1$. We adopt the first-order method (i.e., μ is defined in (7)) here. Trajectories for suboptimality with exact and approximated eigenvalues and eigenvectors are shown in Figure 3. The Krylov subspace $\mathcal{K}_m(\mathbf{B}, \mathbf{u}_1)$ with relatively low dimension m for ASEM matches well with ASEM with exact eigenvalues. It provides confidence that we can use the m dimensional Krylov subspace for m eigenvalues and eigenvectors in ASEM.

Experiment 4. Comparison of ASEM with the Krylov subspace method [3, 2] and the gradient descent method [1] on synthetic problems. For large scale problems, the Krylov subspace method and the gradient descent method are two state-of-the-art methods in solving the target problem (1). In this experiment, we compare the proposed ASEM (with partial eigen information) with the Krylov subspace method and the gradient descent method. The dominant computation steps for those three methods are Hessian-vector products ($\mathcal{O}(mn^2)$). The vector $\mathbf{b}$ is proportional to the vector $[1, \cdots, 1]^{\mathrm{T}}$ with length $\|\mathbf{b}\| = 0.1$. Eigenvalues of the matrix $\mathbf{A}$ are evenly spaced in $[-1, 1]$. Similar to the setting in [2], we define the condition number for (1) as $\kappa = \frac{\lambda_n + \rho \cdot \|\mathbf{x}^*\|}{\lambda_1 + \rho \cdot \|\mathbf{x}^*\|} = \frac{\lambda_n + \sigma^*}{\lambda_1 + \sigma^*}$. Then, we have $\sigma^* = \frac{\lambda_n - \kappa \cdot \lambda_1}{\kappa - 1}$ and $\rho = \frac{\sigma^*}{\|(\mathbf{A} + \sigma^* \mathbf{I})^{-1}\mathbf{b}\|}$. Case 1: easy case that $\kappa = 10^3$; Case 2: harder case that $\kappa = 10^6$. The remaining parameter is $n = 5 \times 10^3$. We adopt the first-order method (i.e., μ is defined in (7)) here. As are shown in Figure 4, ASEM outperforms both the gradient descent method and the Krylov subspace method when m is relatively large. It is reasonable that ASEM underperforms

(a) Case 1: $\kappa = 10^3$. (b) Case 2: $\kappa = 10^6$.

Figure 4: Trajectories of suboptimality (gradient norm $\|\nabla f_{\mathbf{A},\mathbf{b},\rho}(\mathbf{x})\|$) for ASEM, the Krylov subspace method and the gradient descent method in Experiment 4.

when m is small since the m dimensional Krylov subspace cannot well approximate eigenvalues and eigenvectors of $\mathbf{A}$. The results further display the performance of the proposed ASEM method.

Experiment 5. Comparison of ASEM with the Cauchy point method [5], the gradient descent method, and the Krylov subspace method on CUTEst problems [7]. CUTEst library collects various unconstrained and constrained optimization problems that arise in real applications. In this part, we compare the numerical performances of the ARC algorithm [3] on four unconstrained optimization problems from the CUTEst library, where subproblems are solved by the Cauchy point method (ARC-CP), the gradient descent method (ARC-GD), the Krylov subspace method (ARC-Krylov(k), where k is the number of Lanczos basis vectors) and the ASEM method (ARC-ASEM(m), where m is the number of eigenvalues for ASEM). The architecture of the ARC algorithm is provided in Appendix E.2. Four unconstrained optimization problems (e.g., TOINTGSS, BRYBAND, DIXMAANG and TQUARTIC) in the CUTEst library are adopted for testing, where their dimensions are 1000, 2000, 3000 and 5000 respectively. We use the first order ASEM (μ is defined in (7)) here, since we found that the performances of the first order method and the second order methods in ASEM does not differ so much for these problems. Numerical results are reported in Table 1, where $\mathbf{x}_{\text{out}}$, $\|\nabla f(\mathbf{x}_{\text{out}})\|$, $\lambda_1(\nabla^2 f(\mathbf{x}_{\text{out}}))$, iter and time represent the output of the ARC algorithm, the suboptimality (gradient norm) of the objective function, the minimal eigenvalue of the Hessian matrix, numbers of iteration for the ARC and CPU time, respectively. Here are several observations. Firstly, the proposed ASEM algorithm outperforms others in most of cases and is comparible with the Krylov subspace method sometimes, where the ASEM achieves lower suboptimality (gradient norms) or CPU time. Furthermore, only one eigenvalue is enough for the well-behaved ASEM (i.e., ARC-ASEM(m) with $m = 1$), which is a surprising result not observed from theoretical analysis. The main reason is that Hessian matrices in these problems are not left centered (i.e., case 4 in experiment 1), consistent with our analysis. For more experimental details, please refer to Appendix E.

6 Conclusion

We develop the first-order and the second-order truncated secular equations as surrogates to the secular equation with full eigendecomposition in solving the cubic regularized non-convex quadratic problem (1). The proposed ASEM algorithm is feasible since it reduces the computational cost from $\mathcal{O}(n^3)$ to $\mathcal{O}(mn^2)$. The theoretical analysis that more eigen information contributes to lower error shows the properties and reasonableness of the algorithm. We also introduce the implementation details for ASEM algorithm that multiple steps of Hessian-vector products are requisite for estimating eigenvalues and eigenvectors rather than dealing with a large matrix. Numerical experiments further investigate properties and performances of the proposed ASEM algorithm and validate the derived theoretical results. Moreover, the ASEM algorithm converges faster than the state-of-the-art methods on synthetic and CUTEst problems.

Table 1: Results on CUTEst problems in Experiment 5.

Problem	Method	$f(\mathbf{x}_{\text{out}})$	$\|\nabla f(\mathbf{x}_{\text{out}})\|$	$\lambda_1(\nabla^2 f(\mathbf{x}_{\text{out}}))$	iter	time(s)
TOINTGSS ($n = 1000$)	ARC-CP	3.60E+14	4.12E-05	1.40E-16	1000	6.08
	ARC-GD	3.60E+14	1.42E-06	3.89E-16	100	6.98
	ARC-Krylov(1)	3.60E+14	4.12E-05	1.20E-15	300	6.75
	ARC-Krylov(10)	3.60E+14	2.20E-08	1.29E-15	19	**1.87**
	ARC-ASEM(1)	3.60E+14	**8.01E-10**	-1.63E-15	19	2.17
	ARC-ASEM(10)	3.60E+14	8.17E-10	-7.67E-16	19	2.74
BRYBAND ($n = 2000$)	ARC-CP	7.49E+14	1.10E-03	5.40E+00	1000	8.27
	ARC-GD	**1.25E+05**	4.93E+03	4.40E+02	100	11.05
	ARC-Krylov(10)	7.49E+14	6.60E-06	5.40E+00	100	9.85
	ARC-Krylov(30)	7.49E+14	1.14E-07	5.40E+00	14	2.37
	ARC-ASEM(1)	7.49E+14	1.02E-07	5.40E+00	14	**2.24**
	ARC-ASEM(10)	7.49E+14	**1.01E-07**	5.40E+00	14	3.83
DIXMAANG ($n = 3000$)	ARC-CP	1.00E+00	3.13E-04	6.67E-04	2000	35.16
	ARC-GD	1.00E+00	9.24E-05	6.67E-04	200	33.83
	ARC-Krylov(10)	1.00E+00	3.44E-05	6.67E-04	500	32.18
	ARC-Krylov(30)	1.00E+00	9.06E-09	6.67E-04	46	**6.65**
	ARC-ASEM(1)	1.00E+00	5.53E-09	6.67E-04	**30**	7.51
	ARC-ASEM(10)	1.00E+00	**4.85E-09**	6.67E-04	42	18.74
TQUARTIC ($n = 5000$)	ARC-CP	8.04E-01	6.10E-02	-5.41E-05	500	71.92
	ARC-GD	8.05E-01	2.77E-02	-4.80E-05	100	98.14
	ARC-Krylov(1)	8.05E-01	2.76E-02	-4.71E-05	100	29.43
	ARC-Krylov(10)	**5.05E-14**	**8.48E-09**	4.00E-04	46	16.09
	ARC-ASEM(1)	7.43E-14	9.62E-09	4.00E-04	46	**15.47**
	ARC-ASEM(10)	7.43E-14	9.62E-09	4.00E-04	46	16.18

References

[1] Yair Carmon and John Duchi. Gradient descent finds the cubic-regularized nonconvex Newton step. *SIAM Journal on Optimization*, 29(3):2146–2178, 2019.

[2] Yair Carmon and John C Duchi. Analysis of Krylov subspace solutions of regularized nonconvex quadratic problems. *Advances in Neural Information Processing Systems*, 31, 2018.

[3] Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. Adaptive cubic regularisation methods for unconstrained optimization. Part I: motivation, convergence and numerical results. *Mathematical Programming*, 127(2):245–295, 2011.

[4] Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. Adaptive cubic regularisation methods for unconstrained optimization. Part II: worst-case function-and derivative-evaluation complexity. *Mathematical Programming*, 130(2):295–319, 2011.

[5] Andrew R Conn, Nicholas IM Gould, and Philippe L Toint. *Trust region methods*. SIAM, 2000.

[6] Peter J Forrester. Log-Gases and Random Matrices. Princeton University Press, 2010.

[7] Nicholas IM Gould, Dominique Orban, and Philippe L Toint. Cutest: a constrained and unconstrained testing environment with safe threads for mathematical optimization. *Computational Optimization and Applications*, 60(3):545–557, 2015.

[8] Andreas Griewank. The modification of newton's method for unconstrained optimization by bounding cubic terms. Technical report, Technical Report NA/12, 1981.

[9] Richard B Lehoucq, Danny C Sorensen, and Chao Yang. *ARPACK Users' Guide: Solution of Large-Scale Eigenvalue Problems with Implicitly Restarted Arnoldi Methods*. SIAM, 1998.

[10] Yurii Nesterov and Boris T Polyak. Cubic regularization of Newton method and its global performance. *Mathematical Programming*, 108(1):177–205, 2006.

[11] Gilbert W Stewart. A Krylov-Schur algorithm for large eigenproblems. *SIAM Journal on Matrix Analysis and Applications*, 23(3):601–614, 2002.

[12] Nilesh Tripuraneni, Mitchell Stern, Chi Jin, Jeffrey Regier, and Michael I Jordan. Stochastic cubic regularization for fast nonconvex optimization. *Advances in Neural Information Processing Systems*, 31, 2018.

[13] Zhe Wang, Yi Zhou, Yingbin Liang, and Guanghui Lan. Cubic regularization with momentum for nonconvex optimization. In *Uncertainty in Artificial Intelligence*, pages 313–322. PMLR, 2020.

[14] Martin Weiser, Peter Deuflhard, and Bodo Erdmann. Affine conjugate adaptive Newton methods for nonlinear elastomechanics. *Optimisation Methods and Software*, 22(3):413–431, 2007.

[15] Man-Chung Yue, Zirui Zhou, and Anthony Man-Cho So. On the quadratic convergence of the cubic regularization method under a local error bound condition. *SIAM Journal on Optimization*, 29(1):904–932, 2019.

A Proof of Theorem 1

The secular equation (4) can be decomposed into two parts:

$$w(\sigma) = \sum_{i=1}^{m} \frac{c_i^2}{(\lambda_i + \sigma)^2} + \sum_{i=m+1}^{n} \frac{c_i^2}{(\mu + \sigma)^2} - \frac{\sigma^2}{\rho^2} + \sum_{i=m+1}^{n} \frac{c_i^2}{(\lambda_i + \sigma)^2} - \sum_{i=m+1}^{n} \frac{c_i^2}{(\mu + \sigma)^2} \quad (14)$$
$$=: w_1(\sigma; \mu) + e_1(\sigma; \mu),$$

where $e_1(\sigma; \mu) = w(\sigma) - w_1(\sigma; \mu)$ represents the error between two secular equations.

By the Taylor's theorem, we have

$$0 = w(\sigma^*) = w(\sigma_1^*) + \frac{d}{d\sigma}w(\xi) \cdot (\sigma^* - \sigma_1^*) = w_1(\sigma_1^*; \mu) + e_1(\sigma_1^*; \mu) + \frac{d}{d\sigma}w(\xi) \cdot (\sigma^* - \sigma_1^*),$$

where $\xi \in (\sigma^*, \sigma_1^*)$. By the definition that $w_1(\sigma_1^*; \mu) = 0$ and $w(\sigma)$ is monotonically decreasing in the domain (i.e., $\frac{d}{d\sigma}w(\xi) \neq 0$), we have

$$|\sigma_1^* - \sigma^*| \leq \frac{|e_1(\sigma_1^*; \mu)|}{\left|\frac{d}{d\sigma}w(\xi)\right|}. \quad (15)$$

For the error term, we can further rewrite it as

$$e_1(\sigma; \mu) = \sum_{i=m+1}^{n} \frac{c_i^2}{(\lambda_i + \sigma)^2} - \sum_{i=m+1}^{n} \frac{c_i^2}{(\mu + \sigma)^2} = \sum_{i=m+1}^{n} \frac{-2c_i^2}{(\varsigma_i + \sigma)^3} \cdot (\lambda_i - \mu),$$

for some $\varsigma_i \in (\mu, \lambda_i)$, $i = m+1, \cdots, n$. Due to $\mu \geq \lambda_m$ and $\lambda_i \geq \lambda_m$, we have $\varsigma_i > \lambda_m$ and thus $\varsigma_i + \sigma > \lambda_m - \lambda_1 > 0$. Therefore, the error term can be bounded by

$$|e_1(\sigma_1^*; \mu)| \leq \frac{2\|\mathbf{b}\|^2}{(\lambda_m - \lambda_1)^3} \cdot \max_{m+1 \leq i \leq n} |\lambda_i - \mu|. \quad (16)$$

The derivative of the secular equation is

$$\frac{d}{d\sigma}w(\sigma) = \sum_{i=1}^{n} \frac{-2c_i^2}{(\lambda_i + \sigma)^3} - \frac{2}{\rho^2}\sigma < 0,$$

for $\sigma > \max\{-\lambda_1, 0\}$. Then, we have

$$\left|\frac{d}{d\sigma}w(\xi)\right| = \sum_{i=1}^{n} \frac{2c_i^2}{(\lambda_i + \xi)^3} + \frac{2}{\rho^2}\xi \geq \sum_{i=1}^{n} \frac{2c_i^2}{(\lambda_i + \xi)^3} \geq \frac{2\|\mathbf{b}\|^2}{\max_{1 \leq i \leq n}(\lambda_i + \xi)^3} = \frac{2\|\mathbf{b}\|^2}{(\lambda_n + \xi)^3},$$

and

$$\left|\frac{d}{d\sigma}w(\xi)\right| = \sum_{i=1}^{n} \frac{2c_i^2}{(\lambda_i + \xi)^3} + \frac{2}{\rho^2}\xi \geq \frac{2}{\rho^2}\xi.$$

Moreover, σ_1^* is a root of $w_1(\sigma; \mu) = 0$, which implies that

$$\sigma_1^{*2} = \rho^2 \cdot \left(\sum_{i=1}^{m} \frac{c_i^2}{(\lambda_i + \sigma_1^*)^2} + \sum_{i=m+1}^{n} \frac{c_i^2}{(\mu + \sigma_1^*)^2}\right) \leq \rho^2 \cdot \|\mathbf{b}\|^2 \cdot \frac{1}{(\lambda_1 + \sigma_1^*)^2},$$

and thus

$$0 \leq \sigma_1^* \leq \frac{-\lambda_1 + \sqrt{\lambda_1^2 + 4\rho \cdot \|\mathbf{b}\|}}{2} =: B_1.$$

Similarly, σ^* is a root of $w(\sigma) = 0$, which further implies that

$$\sigma^{*2} = \rho^2 \cdot \sum_{i=1}^{n} \frac{c_i^2}{(\lambda_i + \sigma^*)^2} \leq \rho^2 \cdot \|\mathbf{b}\|^2 \cdot \frac{1}{(\lambda_1 + \sigma^*)^2},$$

and thus $0 \leq \sigma^* \leq B_1$. Due to $\xi \in (\sigma^*, \sigma_1^*)$ (or $\xi \in (\sigma_1^*, \sigma^*)$), we have

$$0 \leq \xi \leq B_1. \quad (17)$$

Equipped with the above results, we can bound the derivative term by

$$\left|\frac{d}{d\sigma}w(\xi)\right| \geq \max\left\{\frac{2\|\mathbf{b}\|^2}{(\lambda_n + B_1)^3}, \frac{2B_1}{\rho^2}\right\}. \tag{18}$$

Note that $B_1 > -\lambda_1$ and thus $(\lambda_n + B_1)^3 > 0$, which implies that (18) makes sense. Combining with (15), (16) and (18), we have

$$|\sigma_1^* - \sigma^*| \leq \frac{2\|\mathbf{b}\|^2}{(\lambda_m - \lambda_1)^3} \cdot \min\left\{\frac{(\lambda_n + B_1)^3}{2\|\mathbf{b}\|^2}, \frac{\rho^2}{2B_1}\right\} \cdot \max_{m+1\leq i\leq n} |\lambda_i - \mu| =: C_m \cdot \max_{m+1\leq i\leq n} |\lambda_i - \mu|,$$

where the constant C_m is bounded by $\frac{2\|\mathbf{b}\|^2}{(\lambda_m - \lambda_1)^3} \cdot \min\left\{\frac{(\lambda_n + B_1)^3}{2\|\mathbf{b}\|^2}, \frac{\rho^2}{2B_1}\right\}$.

B Proof of Theorem 2

Similar steps as in the proof for Theorem 1. The secular equation (4) can be decomposed into two parts:

$$
\begin{aligned}
w(\sigma) &= \sum_{i=1}^{m} \frac{c_i^2}{(\lambda_i + \sigma)^2} + \sum_{i=m+1}^{n} \frac{c_i^2}{(\mu + \sigma)^2} - 2\sum_{i=m+1}^{n} \frac{c_i^2 \cdot (\lambda_i - \mu)}{(\mu + \sigma)^3} - \frac{\sigma^2}{\rho^2} \\
&\quad + \sum_{i=m+1}^{n} \frac{c_i^2}{(\lambda_i + \sigma)^2} - \sum_{i=m+1}^{n} \frac{c_i^2}{(\mu + \sigma)^2} + 2\sum_{i=m+1}^{n} \frac{c_i^2 \cdot (\lambda_i - \mu)}{(\mu + \sigma)^3} \\
&=: w_2(\sigma; \mu) + e_2(\sigma; \mu),
\end{aligned}
\tag{19}
$$

where $e_2(\sigma; \mu) = w(\sigma) - w_2(\sigma; \mu)$ represents the error between two secular equations. By Taylor's Theorem, we have

$$0 = w(\sigma^*) = w(\sigma_2^*) + \frac{d}{d\sigma}w(\xi) \cdot (\sigma^* - \sigma_2^*) = w_1(\sigma_2^*; \mu) + e_1(\sigma_2^*; \mu) + \frac{d}{d\sigma}w(\xi) \cdot (\sigma^* - \sigma_2^*),$$

where $\xi \in (\sigma^*, \sigma_2^*)$. By the definition that $w_2(\sigma_2^*; \mu) = 0$ and $w(\sigma)$ is monotonically decreasing in the domain (i.e., $\frac{d}{d\sigma}w(\varepsilon) \neq 0$), we have

$$|\sigma_2^* - \sigma^*| \leq \frac{|e_2(\sigma_2^*; \mu)|}{\left|\frac{d}{d\sigma}w(\xi)\right|}. \tag{20}$$

For the error term

$$e_2(\sigma; \mu) = \sum_{i=m+1}^{n} \frac{c_i^2}{(\lambda_i + \sigma)^2} - \sum_{i=m+1}^{n} \frac{c_i^2}{(\mu + \sigma)^2} + 2\sum_{i=m+1}^{n} \frac{c_i^2 \cdot (\lambda_i - \mu)}{(\mu + \sigma)^3} = \sum_{i=m+1}^{n} \frac{3c_i^2}{(\varsigma_i + \sigma)^4} \cdot (\lambda_i - \mu)^2$$

for some $\varsigma_i \in (\mu, \lambda_i)$, $i = m + 1, \cdots, n$. Due to $\mu > \lambda_m$, we have $\varsigma_i > \lambda_m$ and thus $\varsigma_i + \sigma > \lambda_m - \lambda_1 > 0$. Therefore, the error term can be bounded by

$$|e_2(\sigma_2^*; \mu)| \leq \frac{3\|\mathbf{b}\|^2}{(\lambda_m - \lambda_1)^4} \cdot \max_{m+1\leq i\leq n} (\lambda_i - \mu)^2. \tag{21}$$

Due to the special choice of $\mu = \mu_2$ defined in (10), same upper bound for the derivative term is achieved as that in Theorem 1, i.e.,

$$\left|\frac{d}{d\sigma}w(\xi)\right| \geq \max\left\{\frac{2\|\mathbf{b}\|^2}{(\lambda_n + B_1)^3}, \frac{2B_1}{\rho^2}\right\}. \tag{22}$$

Combining with (20), (21) and (22), we have

$$|\sigma_2^* - \sigma^*| \leq \frac{3\|\mathbf{b}\|^2}{(\lambda_m - \lambda_1)^4} \cdot \min\left\{\frac{(\lambda_d + B_1)^3}{2\|\mathbf{b}\|^2}, \frac{\rho^2}{2B_1}\right\} \cdot \max_{m+1\leq i\leq n} (\lambda_i - \mu)^2 =: C_m \cdot \max_{m+1\leq i\leq n} (\lambda_i - \mu)^2.$$

 # C   Random Gaussian Matrix

In this part, we further elaborate the error analysis in Theorem 1 for random Gaussian matrix, where we can access the eigen distribution. We first introduce the definition for a class of Gaussian random matrix.

Definition 1. *A $n \times n$ matrix $\mathbf{M}$ with entries m_{ij} is a Wigner matrix if $m_{ij} = m_{ji}$ and m_{ij} are randomly i.i.d. up to symmetry.*

Definition 2. *Let $\mathbf{M}$ be a real Wigner matrix with diagonal entries $m_{ii} \sim \mathcal{N}(0,1)$ and $m_{ij} \sim \mathcal{N}(0, 1/2)$, $i \neq j$. Then, $\mathbf{M}$ is said to belong to the Gaussian orthogonal ensemble (GOE).*

Remark. Consider a random $n \times n$ matrix $\mathbf{W}$ with all entries coming from $\mathcal{N}(0,1)$, then $\mathbf{M} = \frac{\mathbf{W}+\mathbf{W}^{\mathrm{T}}}{2}$ is a Gaussian orthogonal ensemble (GOE) matrix.

Theorem 3. *Suppose that the $n \times n$ matrix $\mathbf{A} = \frac{\widetilde{\mathbf{A}}}{\sqrt{2n}}$ with $\widetilde{\mathbf{A}}$ belonging to GOE, then with probability of $1 - o(1)$, we have*

$$\max_{m+1 \leq i \leq n} |\lambda_i - \mu| \leq \mathcal{O}\left(\left(1 - \frac{m+1}{n}\right)^{2/3}\right), \tag{23}$$

where $\mu \in [\lambda_{m+1}, \lambda_n]$ and n is large enough.

Proof. According to section 1.4 in [6], eigenvalues $\{\lambda_1, \cdots, \lambda_n\}$ of $\mathbf{A}$ follows the Wigner's semicircle law that p.d.f. of $\lambda \in \{\lambda_1, \cdots, \lambda_n\}$ satisfies

$$\lim_{n \to \infty} f_n(\lambda) = f(\lambda) = \frac{2}{\pi} \cdot \sqrt{1 - \lambda^2} \cdot \mathbb{I}_{[-1,1]}(\lambda),$$

where $\mathbb{I}_{[-1,1]}(\lambda)$ is the indicator function that values 1 if $\lambda \in [-1,1]$ and 0 if $\lambda \notin [-1,1]$. Therefore,

$$
\begin{aligned}
P(\lambda^* \leq \lambda \leq \lambda_{m+1}) &= P(\lambda \leq \lambda_{m+1}) - P(\lambda \geq \lambda^*) \\
&\approx \frac{m+1}{n} - \int_{-1}^{\lambda^*} f(s) \ ds \\
&= 1 - \frac{2}{\pi} \int_{-1}^{\lambda^*} \sqrt{1 - s^2} \ ds - \left(1 - \frac{m+1}{n}\right).
\end{aligned}
$$

Furthermore,

$$
\begin{aligned}
1 - \frac{2}{\pi} \int_{-1}^{\lambda^*} \sqrt{1 - s^2} \ ds &= \frac{1}{\pi} \cdot \left(\frac{\pi}{2} - \arcsin(\lambda^*) - \lambda^* \cos(\arcsin(\lambda^*))\right) \\
&\approx \frac{4\sqrt{2}}{3\pi} \cdot (1 - \lambda^*)^{3/2} \\
&= \mathcal{O}\left((1 - \lambda^*)^{3/2}\right),
\end{aligned}
$$

where the last equation comes from $\frac{\pi}{2} - \arcsin(\lambda^*) \approx \sqrt{2(1 - \lambda^*)} = \mathcal{O}\left(\sqrt{1 - \lambda^*}\right)$ when $\lambda^* \to 1^-$, and we may select λ^* with $1 - \lambda^* = \mathcal{O}\left(\left(1 - \frac{m+1}{n}\right)^{2/3}\right) \approx \left(\frac{3\pi}{4\sqrt{2}}\right)^{2/3} \cdot \left(1 - \frac{m+1}{n}\right)^{2/3}$ such that $P(\lambda^* \leq \lambda \leq \lambda_{m+1}) > 0$. Therefore, with probability of $1 - o(1)$, we have $\lambda^* \leq \lambda_{m+1} \leq \mu \leq \lambda_n \leq 1$ and thus

$$\max_{m+1 \leq i \leq n} |\lambda_i - \mu| \leq \lambda_n - \lambda_{m+1} \leq 1 - \lambda^* = \mathcal{O}\left(\left(1 - \frac{m+1}{n}\right)^{2/3}\right) \approx \left(\frac{3\pi}{4\sqrt{2}}\right)^{2/3} \cdot \left(1 - \frac{m+1}{n}\right)^{2/3}.$$

$\square$

Remark. The above theorem also holds for other kinds of symmetric random Gaussian matrix, e.g., the Gaussian unitary ensemble (GUE) matrix and the Gaussian symplectic ensemble (GSE) matrix etc., because eigen distribution for both GUE and GSE matrix follow the Wigner's semicircle law [6].

D Equivalence of Cubic Regularized Quadratic Problems

For the cubic regularized non-convex quadratic problem

$$\min_{\mathbf{x}} f_{\mathbf{A},\mathbf{b},\rho}(\mathbf{x}) := \mathbf{b}^{\mathrm{T}}\mathbf{x} + \frac{1}{2}\mathbf{x}^{\mathrm{T}}\mathbf{A}\mathbf{x} + \frac{\rho}{3}\|\mathbf{x}\|^3, \tag{24}$$

the Hessian matrix $\mathbf{A}$ is diagonalizable, i.e., there exist a diagonal matrix $\mathbf{\Lambda} \in \mathbb{R}^{n \times n} = \mathrm{diag}(\lambda_1, \cdots, \lambda_n)$ and a unitary matrix $\mathbf{V} \in \mathbf{R}^{n \times n}$, such that

$$\mathbf{A} = \mathbf{V}\mathbf{\Lambda}\mathbf{V}^T.$$

Then (24) is equivalent to the following problem (25) with diagonal Hessian matrix $\mathbf{\Lambda}$:

$$\min_{\mathbf{y}} f_{\mathbf{\Lambda},\mathbf{c},\rho}(\mathbf{x}) := \mathbf{c}^{\mathrm{T}}\mathbf{y} + \frac{1}{2}\mathbf{y}^{\mathrm{T}}\mathbf{\Lambda}\mathbf{y} + \frac{\rho}{3}\|\mathbf{y}\|^3, \tag{25}$$

where $\mathbf{c} = \mathbf{V}^{\mathrm{T}}\mathbf{b}$. Therefore, we adopt diagonal matrix $\mathbf{A}$ in numerical experiments since these two problems are equivalent.

E Experimental Details

E.1 The Cauchy Point

The Cauchy radius [5] for the cubic regularized quadratic problem

$$\min_{\mathbf{x}} f_{\mathbf{A},\mathbf{b},\rho}(\mathbf{x}) := \mathbf{b}^{\mathrm{T}}\mathbf{x} + \frac{1}{2}\mathbf{x}^{\mathrm{T}}\mathbf{A}\mathbf{x} + \frac{\rho}{3}\|\mathbf{x}\|^3$$

is the solution of

$$r = \arg\min_{\zeta \in \mathbb{R}} f_{\mathbf{A},\mathbf{b},\rho}(-\zeta \cdot \mathbf{b}/\|\mathbf{b}\|),$$

and the Cauthy point is defined as

$$\mathbf{x}_c = -r \cdot \mathbf{b}/\|\mathbf{b}\|.$$

Note that $f_{\mathbf{A},\mathbf{b},\rho}(\mathbf{x}_c) \leq f_{\mathbf{A},\mathbf{b},\rho}(\mathbf{0}) = 0$.

E.2 The ARC Algorithm

We present the ARC algorithm [3] (Algorithm 1) here for completeness and clarification.

E.3 Hyperparameters

In this subsection, we provide hyperparameters of the ARC algorithm in solving four optimization problems (e.g., TOINTGSS, BRYBAND, DIXMAANG and TQUARTIC) from the CUTEst library. In all experiments, we set $\gamma_1 = \gamma_2 = 2$, $\eta_1 = 0.1$, $\eta_2 = 0.9$ and $\rho_0 = 10^3$. The initial guess $\mathbf{x}_0$ is available for each problem (you can access it by *prob.x* command in MATLAB). For TOINTGSS, BRYBAND, DIXMAANG and TQUARTIC, we select the top 1000, 2000, 3000 and 5000 variables as unknown variables for optimization, respectively.

E.4 Additional Experiments

Experiment 6. The effect of the length $\|\mathbf{b}\|$. The derived upper bound in (6) and (11) are proportional to $\|\mathbf{b}\|^{3/2}$, which implies that larger $\|\mathbf{b}\|$ may cause worse approximation for σ^* and thus the unattainable optimality. Here, we tested the convergence of the proposed method for different length $\|\mathbf{b}\|$ with other parameters fixed. In case 1 to 6, we set $\|\mathbf{b}\| \in \{1, 0.5, 0.2, 0.1, 0.05, 0.01\}$. The vector $\mathbf{b}$ is proportional to the vector $[1, \cdots, 1]^{\mathrm{T}}$. Eigenvalues of the matrix $\mathbf{A}$ are evenly spaced in $[-1, 1]$. The remaining parameters are $n = 5 \times 10^3$ and $\rho = 0.1$. We adopt the first-order method (i.e., μ is defined in (7)) here. As is shown in Figure 5, the proposed method is stable for both large and small $\|\mathbf{b}\|$ while gradient descent [1] is ill-conditioned if $\|\mathbf{b}\|$ is tiny. Consistent with the error analysis, the proposed method converge faster for smaller length $\|\mathbf{b}\|$.

Algorithm 1 The ARC algorithm

Target: a local minimizer of the objective function $f(\mathbf{x})$, where $\mathbf{x} \in \mathbb{R}^n$.
Input: initial guess $\mathbf{x}_0$, $\gamma_2 \geq \gamma_1 > 1$, $1 > \eta_2 \geq \eta_1 > 0$, $\rho_0 > 0$ and the integer $T > 0$.
Output: the approximate solution $\mathbf{x}_{\text{out}} \in \mathbb{R}^n$.

1: **for** $t = 0, \cdots, T$ **do**
2: load $\mathbf{b}_t = \nabla f(\mathbf{x}_t)$, $\mathbf{A}_t = \nabla^2 f(\mathbf{x}_t)$
3: compute the Cauchy point $\mathbf{s}_c$ of $f_{\mathbf{A}_t, \mathbf{b}_t, \rho_t}(\mathbf{s}) := \mathbf{b}_t^{\mathrm{T}} \mathbf{s} + \frac{1}{2} \mathbf{s}^{\mathrm{T}} \mathbf{A}_t \mathbf{s} + \frac{\rho_t}{3} \|\mathbf{s}\|^3$
4: solve $f_{\mathbf{A}_t, \mathbf{b}_t, \rho_t}(\mathbf{s})$ by the gradient descent, the Krylov subspace or the ASEM method, the approximate solution is denoted as $\mathbf{s}_t$
5: **if** $f_{\mathbf{A}_t, \mathbf{b}_t, \rho_t}(\mathbf{s}_c) \leq f_{\mathbf{A}_t, \mathbf{b}_t, \rho_t}(\mathbf{s}_t)$ **then**
6:   $\mathbf{s}_t = \mathbf{s}_c$;
7: **end if**
8: compute

$$\kappa_t = \frac{f(\mathbf{x}_t) - f(\mathbf{x}_t + \mathbf{s}_t)}{-f_{\mathbf{A}_t, \mathbf{b}_t, \rho_t}(\mathbf{s}_t)};$$

9: set

$$\mathbf{x}_{t+1} = \begin{cases} \mathbf{x}_t + \mathbf{s}_t & \text{if } \kappa_t \geq \eta_1 \\ \mathbf{x}_t & \text{otherwise} \end{cases}$$

10: set

$$\rho_{t+1} \in \begin{cases} (0, \rho_t] & \text{if } \kappa_t > \eta_2 & \text{(very successful iteration)} \\ [\rho_t, \gamma_1 \rho_t] & \text{if } \eta_1 \leq \kappa_t \leq \eta_2 & \text{(successful iteration)} \\ [\gamma_1 \rho_t, \gamma_2 \rho_t] & \text{otherwise} & \text{(unsuccessful iteration)} \end{cases}$$

11: **end for**

Figure 5: Trajectories of suboptimality (gradient norm $\|\nabla f_{\mathbf{A}, \mathbf{b}, \rho}(\mathbf{x})\|$) with different length $\|\mathbf{b}\|$ in Experiment 6.