Scaled Neural Multiplicative Model for Tractable Optimization

Abstract

Challenging decision problems in retail and beyond are often solved using the predict-then-optimize paradigm. An initial effort to develop and parameterize a model of an uncertain environment is followed by a separate effort to identify the best possible solution of an optimization problem. Linear models are often used to ensure optimization problems are tractable. Remarkably accurate Deep Neural Network (DNN) models have recently been developed for various prediction tasks. Such models have been shown to scale to large datasets without loss of accuracy and with good computational performance. It can, however, be challenging to formulate tractable optimization problems based on DNN models. In this work we consider the problem of shelf space allocation for retail stores using DNN models. We highlight the trade-off between predictive performance and the tractability of optimization problems. We introduce a Scaled Neural Multiplicative Model (SNMM) with shape constraints for demand learning that leads to a tractable optimization formulation. Although, this work focuses on a specific application, the formulation of the models are general enough such that they can be extended to many real world applications.

1 Introduction

The predict-then-optimize framework is ubiquitous in applied research. A predictive model is first developed to approximate the true dynamics of a system under consideration to a given level of accuracy. A mathematical programming formulation based on the predictive model is then used to help researchers identify optimal policies for challenging real-world decision problems. Despite the fact that predictive models that are estimated based on the historical data need not be causal, there are recent works that shed some light on the principles behind this approach (Bertsimas & Kallus, 2020).

An alternative to the predict-then-optimize framework is integrating the two stages, predicting and optimizing at the same time, by utilizing a specific loss function in prediction models (Elmachtoub & Grigas, 2022). For example, model-free Reinforcement learning (RL) approaches explore their environment while also exploiting it to make optimal decisions.

Big-box retailers have thousands of stores located across the country. They use data-driven decision to optimize operations; e.g, assortment planning, pricing, and supply chain optimization. They work on problems related to shelf space allocation, making the best use of limited space in stores. Space planning for apparel is particularly challenging, due to the different shapes and sizes of the merchandise and the temporal shifts in brand importance.

The problem that motivated our work in this area involves deciding how much space in terms of *fixtures* to assign to each of several *brands* within a *category* of products for sale. These problems are typically solved at the *department* or *store* level. Products are arranged into departments and categories; for example, 'activewear tops' is a category within the women's apparel department. In this work, we introduce a novel approach based on the predict-then-optimize framework for solving a shelf space allocation problem.

In particular our contributions are as follows

- We begin by identifying certain key characteristics of relevant real-world data that make modeling challenging.

- We discuss predictive model selection considering the tractability of resulting optimization problem formulations and focus on the convexity of the formulations.

- We propose multiplicative models and establish conditions for tractability of the optimization. We also discuss why family of linear models are not suitable for the given application.

- We hypothesize that it is possible to convert an intractable optimization to a tractable one without loss of prediction accuracy. We achieve this via a Scaled Neural Multiplicative Model (SNMM) and demonstrate that our proposed model performs well relative to alternative models.

2 RELATED WORK

The relationship between the shelf space allocated to a product and that product's sales has been studied extensively (Bianchi-Aguiar et al., 2021; Hübner & Kuhn, 2012; Karampatsa et al., 2017). Analyses often focus on incremental returns, estimating the *space elasticity*. Under the assumption of diminishing returns to scale, the relationship between shelf space and sales can be modeled using a concave function (Curhan, 1972; Eisend, 2014). The assumption of diminishing returns makes intuitive sense and has been widely used in production functions (Gopalswamy & Uzsoy, 2019)(Aigner & Chu, 1968); the incremental gain in sales by adding more space to showcase a product decreases as the space allocated increases. This assumption can also have a side benefit of making shelf space allocation optimization problems easier to solve.

Optimization: There have been a number of scientific articles focused on the assortment optimization problem; a problem related to the one we consider in this work. Retailers solve this problem to determine which products to sell. Shelf space in stores is often the most prominent constraint. Kök & Fisher (2007), Yücel et al. (2009), Lo (2019), and numerous others focus on developing optimization frameworks that include complicated consumer choice models. This allows the authors to select the optimal product mix accounting for the fact that some consumers will substitute one product for another depending on what is and is not sold by a retailer. Our problem is somewhat different in that we are not selecting specific products to sell, but rather deciding which brands to carry and how much space to allocate to these brands. These brands will offer products that are relatively unique within a store.

Hübner & Kuhn (2011) points out the relationships between space allocated, consumer demand, and inventory costs. Assuming a fixed amount of space available, a retailer can choose to offer fewer facings of a more diverse set of products to increase consumer interest. This will, however, also increase inventory holding and replenishment costs. There will be increasing demands placed on store labor. Ryzin & Mahajan (1999) came to a similar conclusion earlier looking specifically at apparel. Our problem is, again, somewhat different. Brand managers are responsible for selecting product mix within brands sold in specific stores, managing inventory costs and the tradeoff between such costs and expected sales or profits.

Shape-Constrained Models: An integral part of our prediction model that makes the optimization tractable is shape constraints. These are models that are constrained to be monotonic, convex, concave or non-negative among others. Shape constraints provide effective regularization, reducing the chance that noisy training data or adversarial examples produce a model that does not behave as expected. These impose strong priors on the data and can be used effectively to produce well behaved structured prediction models. It is this property that we enforce in our prediction model to yield a tractable optimization. Shape constraints on neural networks have been studied in different applications (Gupta et al., 2018). Most of the prior works consider GAMs and neural networks separately. In our work, we combine the strength of neural networks and the simplicity of GAMs and propose a Scaled Neural Multiplicative model that can model concave, convex or monotone constraints effectively.

3 Preliminaries

Given a closed set $\mathbb{X}$ and function F_θ that is parameterized by θ, decision making problem is to find the solution to the optimization problem given below

$$\boldsymbol{x}^* = \arg\max_{\boldsymbol{x} \in \mathbb{X}} F_{\boldsymbol{\theta}}(\boldsymbol{x}) \tag{1}$$

The complexity of the underlying optimization model is based on the form of the function F_θ, the objective function, and the definition of the set $\mathbb{X}$, the constraints. Further, the problem is generally tractable only for concave functions with respect to $\boldsymbol{x}$ (including affine).

Although, more generic functional forms can be incorporated using Mixed Integer Programming (MIP) with piecewise linear approximations (Vielma et al., 2010)(Gopalswamy et al., 2019), the solution time for MIPs in general does not scale well for large problems. By tractability, we refer to the class of optimization problems that can be solved in polynomial time.

3.1 Linear Case

A linear functional form for the objective function leads to a relatively simple convex (linear) optimization problem if the set $\mathbb{X}$ is a convex polytope. In this case, the prediction model must be linear with respect to the optimization variable $\boldsymbol{x}$

$$F_{\boldsymbol{\theta}}(\boldsymbol{x}) = \boldsymbol{c}^t \boldsymbol{x} + \phi(\mathbb{P} \backslash \boldsymbol{x})$$

The $\boldsymbol{c}$ model parameters are coefficients in a linear function of $\boldsymbol{x}$ while the rest of the features $\mathbb{P} \backslash \boldsymbol{x}$ in the prediction model do not have any restrictions. The predictive model can be a generic neural network, for instance. While this approach leads to a tractable optimization problem with respect to $\boldsymbol{x}$, certain features cannot influence the optimal solution $\boldsymbol{x}^*$ and are thus irrelevant to the actual decision problem.

$$\boldsymbol{x}^* = \arg\max_{\boldsymbol{x} \in \mathbb{X}} \boldsymbol{c}^t \boldsymbol{x} + \phi(\mathbb{P} \backslash \boldsymbol{x}) = \arg\max_{\boldsymbol{x} \in \mathbb{X}} \boldsymbol{c}^t \boldsymbol{x}$$

3.2 Multiplicative Case

A multiplicative model form can be a better alternative to a linear model form for two reasons: *(i) it can handle heterogeneity in variance w.r.t features in $\mathbb{X}$ and (ii) other features in $\mathbb{P}$ can play a role in determining the optimal solution to equation 1*. A general multiplicative model is given by

$$F_{\boldsymbol{\theta}}(\boldsymbol{x}) = \prod_i x_i^{\beta_i} \phi(\mathbb{P} \backslash x_i) \tag{2}$$

If $\sum_i \beta_i = 1$ and $\phi(\mathbb{P} \backslash x_i) \in \mathbb{R}^+$, then our objective function is convex. Such conditions can be difficult to enforce on a general neural network model without explicit architectural design (Amos et al., 2017). Further, such models can be difficult to interpret and communicate to business stakeholders in general. To alleviate the above problems, we consider the following multiplicative form

$$F_{\boldsymbol{\theta}}(\boldsymbol{x}) = \sum_i x_i^{\beta_i} \prod_{p \in \mathbb{P} \backslash x_i} \phi_p(w_i^p) \tag{3}$$

where w_i^p is the feature p that depends on the index i of the optimization variable. For example, variable x_i could represent space for brand i whereas w_p^i could be description of brand i embedded using a language model. Details on how to constraint 3 to enforce convexity (concavity) will be discussed in the next section.

4 Problem Formulation

We discuss a specific instance of the optimization problem based on the predictive model F_θ that captures the relationship between sales and features. The variable of interest in the optimization, *fixture count*, defines the shelf space in a store. This work specifically considers the problem of finding optimal space allocation for each brand-category pair for a given department in each store to maximize revenue.

The tractability of the above formulation depends on the set $\mathbb{X}$ and the parametric function F_θ. In this work we will consider a family of functions from Generalized linear models (GLM) such as additive, multiplicative as well as DNNs that extend GAMs (Agarwal et al., 2021). We will analyze the models from the perspective of flexibility and convexity with respect to the fixture count feature

We approach this problem in two stages -

- Learn the functional form of $F_\theta(x)$ through demand learning.
- Use this relationship to optimize for space across all stores.

4.1 Data Sparsity

A major challenge in pushing research from academia to application rests with the data quality. It is non trivial to achieve the appropriate functional form or relationship that captures the expected model behavior without loss in performance. In the retail space in particular, we have found that the domains of the functions we would like to use are not fully observed in the historical data. This is due to the challenges in experimentation in physical retail stores at the granularity needed for making decisions. For example, in apparel there are a large number of items sold, while the sales traffic at each item level is very sparse. Ideally, one would be interested in estimating the space elasticity of item i for time period t. This estimation problem is challenging due to sparsity in data and non-dynamic space allocation in retail stores (i.e, space allocation for an item or category usually stays the same for a quarter to offset high labor costs involved). We consider models that have a single coefficient for fixture count. Further, the category and brand have multiplicative effects in the models we consider. This way, we are able to estimate the space elasticity by using data across different products and also estimate total effect for any given brand and category.

4.2 Generalized Linear Models

GLMs (Nelder & Wedderburn, 1972) are a general class of linear models that can capture structured non-linearity between the target and the covariates. Logistic regression (Wright, 1995) is one of the widely used GLMs in classification problems. Multinomial logit models have been used for discrete choice modeling (Ben-Akiva et al., 1985).

Linear Fixed Effects Model: We start by fitting a linear regression model where y, the \$ sales amount, is a linear combination of x and d_i, the features. Here, x denotes the fixture count variable and d_i denotes other features like brand, category and department.

$$y = \beta_0 + \beta_1 x + \sum_i \xi_i d_i \tag{4}$$

When used as an objective function with x as the decision variable, the linear model has constant space elasticity across all brand-category pairs and the effect of other features also becomes constant, and therefore irrelevant to the optimization.

Log-Log Model: A log-log model ensures that the effect of fixture count on sales is positive.

$$\log y = \beta_0 + \beta_1 \log x + \sum_i \xi_i d_i$$
$$y = e^{\beta_0} x^{\beta_1} e^{\sum_i \xi_i d_i}$$
$$y = F(\boldsymbol{d}) x^{\beta_1} \tag{5}$$

Moreover, unlike a simple linear model, the coefficients of log-log model have a multiplicative effect on the independent variable. The multiplicative effect of coefficients here ensures that the effect of different features like brand, category and department are directly related to the fixture count variable. These variations across brand, category and department learned from the historical data allows the optimization model to identify the right weights associated with each fixture count variable and assign space accordingly.

Convexity Analysis: The log-log model is convex when $\beta_1 > 1$ or $\beta_1 < 0$. In our motivating example, these would be the cases where allocating additional space to a brand would either reduce

sales or increase sales at a faster rate than before (increasing returns to scale). Neither of these cases seem realistic. Requiring $0 \leq \beta_1 \leq 1$ results in more intuitive models of retail operations as well as a function more amenable to optimization.

4.3 Deep Neural Networks

Neural Additive Models (NAM): Deep neural networks have been successful in prediction tasks. We consider a special form of neural networks that are additive in nature (Agarwal et al., 2021). NAMs learn a linear combination of networks, each of which attend to a single input feature: each f_i in 6 is parameterized by a neural network. NAMs are explainable, elegant and easy to understand models.

$$y = \beta + f_1(x_1) + f_2(x_2) + \cdots + f_K(x_K) \tag{6}$$

Neural Multiplicative Models (NMM): Similar to the multiplicative linear models, we consider multiplicative form for the neural additive model using the log transformation of the dependent variable y and fixture count x

$$y = e^\beta e^{f(\log x)} e^{\sum_i f_i^d(d_i)} \tag{7}$$

Equation 7 in general will not lead to a tractable optimization problem. The functional form considered in Agarwal et al. (2021) involves EXU layer $g(x) = e^w(x - b)$ and linear layers. We consider a linear functional form of f that can be modeled via linear layers. To improve learning and expressiveness we use hidden dimension of 100, thus forming a total of 2 linear layers.

$$y = e^\beta x^{\boldsymbol{w}_2' \boldsymbol{w}_1} e^{\boldsymbol{w}_2' \boldsymbol{b}_1 + b_2} e^{\sum_i f_i^d(d_i)} \tag{8}$$

The parameters of Equation 8 can be constrained to produce a concave function well suited for use as an objective function in a subsequent optimization problem. The constraint $0 \leq \boldsymbol{w}_2' \boldsymbol{w}_1 \leq 1$, can be modeled by considering $\sigma(\boldsymbol{w}_2), \sigma(\boldsymbol{w}_1)$ instead $\boldsymbol{w}_2, \boldsymbol{w}_1$, where $\sigma(\boldsymbol{x}) = \left\{ \frac{\exp x_i}{\sum_i \exp x_i} \right\}_{i=1}^n$. It can be shown that $0 \leq \sigma(\boldsymbol{w}_2)' \sigma(\boldsymbol{w}_1) \leq 1$.

Lemma 4.1. *For any $x, y \in \Delta$ where $\Delta = \{w \in \mathcal{R}_+^n \mid \sum_i w_i = 1\}$, we have $0 \leq \sum_i x_i y_i \leq 1$*

Proof. In appendix. □

Scaled Neural Multiplicative Models (SNMM): Generally, constraining neural network models leads to reduction in their capacity. The direct consequence being the inability to be a universal function approximator. We propose a NMM model where the scaling of the feature x before the log transformation can be learned in an end-to-end fashion.

$$y = e^\beta z^{\sigma(\boldsymbol{w}_2)' \sigma(\boldsymbol{w}_1)} e^{\sigma(\boldsymbol{w}_2)' \boldsymbol{b}_1 + b_2} e^{\sum_i f_i^d(d_i)} \tag{9a}$$

$$z = 1 + \max\left(0, \boldsymbol{s}_2' \boldsymbol{s}_1 x + \boldsymbol{s}_2' \boldsymbol{t}_1 + t_2\right) \tag{9b}$$

Equation 9 is tractable with respect to the optimization when $\boldsymbol{s}_2' \boldsymbol{s}_1 x + \boldsymbol{s}_2' \boldsymbol{t}_1 + t_2 \geq 0$. Although, $\max$ operator cannot be modeled without mixed integer program, we noticed that the above condition for tractability holds on the experiments we carried out with different number of stores. In our application, y represents sales in dollars and x represents fixture count. Equation 9 can be written for an explicit category i, brand j and department k as follows

$$y_{i,j,k} = a_{i,j,k} z_{i,j,k}^\gamma \tag{10a}$$

$$z_{i,j,k} = \boldsymbol{s}_2' \boldsymbol{s}_1 x_{i,j,k} + \boldsymbol{s}_2' \boldsymbol{t}_1 + t_2 \tag{10b}$$

$$a_{i,j,k} = e^\beta e^{\sigma(\boldsymbol{w}_2)' \boldsymbol{b}_1 + b_2} e^{f_i(d_i) + f_j(d_j) + f_k(d_k)} \tag{10c}$$

Note that $a_{i,j,k}$ is constant with respect to optimization variables.

4.4 Optimization

After learning the functional form of F_θ, we can formulate the problem for stage two as follows. Let the sets $\mathbb{I}$ and $\mathbb{K}$ denote the set of brand, category pairs (i, j) and set of departments, respectively. The parameter $a_{i,j,k}$ is the coefficient of category i, brand j and department k based on the multiplicative

model in 9. Let $x_{i,j,k}$ denote the variable fixture count for category i, brand j and department k, with $l_{i,j,k}$ and $u_{i,j,k}$ as lower and upper bounds on the variable, respectively. Further, lets define parameters s and s_k and γ as the total fixture count (across all departments), fixture count for a department k and space elasticity, respectively.

We define the optimization with explicit index (i,j) to indicate the dependency of the coefficients $a_{i,j,k}$ on brand, category and department encoding, even though it can be combined into a single index. The objective is to maximize sales revenue from equation 10. The explicit form of equation 10, $f(x_{i,j,k})$, depends on the model of choice (SNMM in our case).

$$\max \sum_{k \in \mathbb{K}} \sum_{i,j \in \mathbb{I}} y_{i,j,k} \tag{11a}$$

$$s.t.$$

$$y_{i,j,k} = f(x_{i,j,k}), \forall i,j,k \tag{11b}$$

$$\sum_{i,j \in \mathbb{I}} x_{i,j,k} \le s_k, \forall k \in \mathbb{K} \tag{11c}$$

$$\sum_{k \in \mathbb{K}} \sum_{i,j \in \mathbb{I}} x_{i,j,k} \le s \tag{11d}$$

$$l_{i,j,k} \le x_{i,j,k} \le u_{i,j,k}, \forall i,j,k \tag{11e}$$

$$x_{i,j,k} \ge 0, \forall i,j,k \tag{11f}$$

5 EXPERIMENTS

We demonstrate the ability of the proposed model to handle shape constraints and predictive power with two sets of experiments. In the first experiment, we compare SNMM with different models from (Gupta et al., 2018) using real world datasets on their ability to handle constraints such as concavity with respect to the input. In the second experiment, we provide results on a proprietary dataset from a retailer with respect to modeling space elasticity.

5.1 SHAPE-CONSTRAINED BENCHMARK

We consider three publically available datasets and test the ability of SNMM to handle convexity and concavity constraints based on established benchmarks.

Table 1: Synthetic experiments: Car Sales and Puzzle Review

Car Sales			Puzzle Sales		
Model	Val. MSE	Test MSE	Model	Val. MSE	Test MSE
SNMM	**1923**	**8486**	SNMM	3768	8457
DNN	2035	10931	**DNN**	**2189**	**5652**
SCNN conv.	2262	10613	SCNN conc.	2632	7931
SCNN conv. decr.	2442	10590	SCNN conc. incr.	2437	6927
Cal Lin. decr.	2271	10727	RTL incr.	4457	8838
Cal Lin. conv. decr.	2304	10593	RTL all	3543	8315
			Cal Lin. incr.	3589	8270
			Cal Lin. all	3617	8189

5.1.1 CAR SALES

For this tiny 1-d problem with 109 training, 14 validation, and 32 test examples (www.kaggle.com/hsinha53/car-sales/data), we predict monthly car sales (in thousands) from the price (in thousands). We enforce a convexity constraint on the price variable with

respect to the sales variable. The results can be seen in Table 1. SNMM performed relatively better compared to all models with the lowest Validation and Test MSE.

5.1.2 PUZZLE SALES FROM REVIEWS

For this small problem (3 features, 156 training, 169 validation, and 200 non-IID test examples, dataset courtesy of Artifact Puzzles and available at `www.kaggle.com/dbahri/puzzles`), we predict the 6-month sales of different wooden jigsaw puzzles from three features based on its Amazon reviews: its average star rating, the number of reviews, and the average word count of its reviews. Here we assume a convexity constraint on the star rating, and concavity constraints on number of reviews and word count. Table 1 shows that for SNMM, even though the Validation MSE is highest compared to the rest of the models, the Test MSE remains within the range of the highest Test MSE.

Table 2: Synthetic experiments: Wine Quality

| Model | Wine Quality | |
	Val. MSE	Test MSE
SNMM	6.95	6.98
DNN	**4.91**	**4.79**
SCNN conc.	5.96	7.22
SCNN conc. incr.	6.13	6.21
RTL incr.	4.96	4.85
RTL conc. incr.	4.96	4.83
Cal Lin. incr.	5.25	5.10
Cal Lin. conc. incr.	5.23	5.10

5.1.3 WINE ENTHUSIAST MAGAZINE REVIEWS

The goal is to predict a wine's quality measured in points [80, 100] based on price (the most important feature), country (21 Bools), and 39 Bool features based on the wine description from Wine Enthusiast Magazine (61 features, 84,642 training, 12,092 validation, and 24,185 test examples; `www.kaggle.com/dbahri/wine-ratings`). We constrain the price feature to be concave. From Table 2, it can be seen that while DNN has the lowest error, the Test MSE for SNMM is within the range of the highest Test MSE across all models.

Table 3: Benchmark results for 5 stores

| Model | log scale | | | linear scale | | |
	R^2	MSE	MAE	R^2	MSE	MAE
GLM-L	-	-	-	0.623 ± 0.01	61692 ± 2214	126 ± 0.67
GLM-M	0.759 ± 0.00	0.531 ± 0.00	0.568 ± 0.00	0.619 ± 0.01	62398 ± 1323	96 ± 0.64
GLM-CM	0.736 ± 0.00	0.582 ± 0.00	0.603 ± 0.02	0.618 ± 0.01	62525 ± 2202	97 ± 0.87
NAM	-	-	-	0.684 ± 0.01	55809 ± 2033	100 ± 1.90
NMM	0.769 ± 0.01	0.506 ± 0.01	0.564 ± 0.01	0.567 ± 0.01	69923 ± 1672	95 ± 1.46
NMM-L	0.824 ± 0.02	0.388 ± 0.05	0.483 ± 0.02	0.738 ± 0.06	43917 ± 1099	80 ± 5.44
NMM-CL	0.638 ± 0.01	0.798 ± 0.01	0.724 ± 0.01	0.407 ± 0.01	96364 ± 2771	120 ± 1.61
SNMM	$\mathbf{0.827 \pm 0.01}$	$\mathbf{0.382 \pm 0.02}$	$\mathbf{0.476 \pm 0.01}$	$\mathbf{0.775 \pm 0.02}$	$\mathbf{37830 \pm 2687}$	$\mathbf{75 \pm 2.32}$

5.2 CASE STUDY ON REAL DATA

In this section, we perform tests on a random selection of stores across the country from one retailer. We first describe the dataset and the features considered. We will then compare eight different

models for predicting sales with the metrics given below.

$$R^2 = 1 - \frac{\|\boldsymbol{y} - \hat{\boldsymbol{y}}\|^2}{\|\boldsymbol{y} - \bar{\boldsymbol{y}}\|^2} \quad (12)$$

$$MSE = N^{-1}\|\boldsymbol{y} - \hat{\boldsymbol{y}}\|^2 \quad (13)$$

$$MAE = N^{-1}|\boldsymbol{y} - \hat{\boldsymbol{y}}| \quad (14)$$

We consider linear and multiplicative models with variants. The variants are explored in lieu of search for computationally tractable optimization model. Further, certain models offer better modeling capacity with respect to the solution space. The functional relationship between the space feature (variable in optimization) and sales defines the objective function. We expect the models to capture the concave relationship from the data, but it seldom happens in reality. Real-word data is noisy and unobserved confounders can affect the data. Further, retail data in general depends on exogenous uncertainty such as market dynamics, competitors, economic factors, etc. Explicit shape constraint is placed on the model such as non-negativity of sales and concavity of sales with respect to space feature.

Table 4: Benchmark results for 10 stores

Model	log scale			linear scale		
	R^2	MSE	MAE	R^2	MSE	MAE
GLM-L	-	-	-	0.638 ± 0.00	56655 ± 1378	121 ± 0.43
GLM-M	0.753 ± 0.00	0.522 ± 0.00	0.564 ± 0.00	0.657 ± 0.01	53653 ± 1465	90 ± 0.39
GLM-CM	0.727 ± 0.00	0.577 ± 0.00	0.601 ± 0.00	0.616 ± 0.01	60084 ± 1404	92 ± 0.46
NAM	-	-	-	0.668 ± 0.00	53217 ± 827	98 ± 1.09
NMM	0.786 ± 0.01	0.452 ± 0.01	0.533 ± 0.01	0.622 ± 0.01	59894 ± 2255	86 ± 1.52
NMM-L	0.813 ± 0.01	0.396 ± 0.02	0.488 ± 0.01	0.707 ± 0.01	46438 ± 2315	78 ± 2.20
NMM-CL	0.648 ± 0.02	0.746 ± 0.04	0.7 ± 0.03	0.478 ± 0.04	82035 ± 7789	109 ± 5.24
SNMM	$\mathbf{0.817 \pm 0.01}$	$\mathbf{0.387 \pm 0.00}$	$\mathbf{0.479 \pm 0.00}$	$\mathbf{0.760 \pm 0.01}$	$\mathbf{37143 \pm 1996}$	$\mathbf{72 \pm 0.51}$

As discussed in sections 4.2 and 4.3, we enforce the constraints and report performance metrics. We present the metrics in both log scale and linear scale whenever applicable to offer complete comparison across all the models

The following models are used for the experiment:

- GLM-L: Classical linear regression model that assumes a linear relationship between the dependent and independent variables, Equation 4.

- GLM-M: Multiplicative model that takes a log transformation of dependent variable and space variable as in Equation 5.

- GLM-CM: Constrained GLM-M with space elasticity between 0 and 1.

- NAM: Neural Additive Model as described in Equation 6

- NMM: NAM that takes a log transformation of dependent variable and space variable as in Equation 7.

- NMM-L: NMM with linear layers for the space variable as in Equation 8

- NMM-CL: Constrained NMM-L with space elasticity between 0 and 1 using lemma 4.1

- SNMM: Proposed model in Equation 9

Dataset: The models consider historical space and sales data of Apparel departments from 2018 to 2022. We want to predict and optimize for space at the department-category-brand level, so we aggregate data to that level at a weekly granularity.

Model Evaluation: Numerical tests were carried out on a set of 5, 10 and 20 stores. The effectiveness of the models in sales prediction is evaluated using the metrics described above. The proposed model is compared with variety of different models and the results are shown in Tables 3,

Table 5: Data summary

Feature	Description
$ sales	Dependent variable representing the dollar sales amount
fixture count	Measure of space represented in a store
store no	Store number (categorical)
store area	Area of a store in sq. ft.
department no	Department number
category	Category description (vector embedding)
brand	Name of the brand (vector embedding)
demographic	Demographic data based on location of the store
income	Income data based on location of the store
time features	Time-based features based on the week of transaction

Table 6: Benchmark results for 20 stores

Model	log scale R^2	log scale MSE	log scale MAE	linear scale R^2	linear scale MSE	linear scale MAE
GLM-L	-	-	-	0.583 ± 0.00	71124 ± 1943	126 ± 0.29
GLM-M	0.742 ± 0.00	0.531 ± 0.00	0.568 ± 0.00	0.626 ± 0.01	63831 ± 3224	91 ± 0.56
GLM-CM	0.715 ± 0.00	0.587 ± 0.00	0.605 ± 0.00	0.575 ± 0.00	72437 ± 1849	92 ± 0.50
NAM	-	-	-	0.617 ± 0.00	65838 ± 1475	102 ± 0.57
NMM	0.793 ± 0.01	0.428 ± 0.01	0.516 ± 0.01	0.585 ± 0.01	72292 ± 841	84 ± 0.35
NMM-L	$\mathbf{0.807 \pm 0.01}$	$\mathbf{0.400 \pm 0.01}$	$\mathbf{0.487 \pm 0.01}$	0.682 ± 0.01	55934 ± 2198	79 ± 1.12
NMM-CL	0.623 ± 0.01	0.78 ± 0.03	0.713 ± 0.02	0.352 ± 0.02	113907 ± 4103	113 ± 2.49
SNMM	$\mathbf{0.806 \pm 0.02}$	$\mathbf{0.399 \pm 0.03}$	$\mathbf{0.486 \pm 0.02}$	$\mathbf{0.716 \pm 0.01}$	$\mathbf{48983 \pm 2228}$	$\mathbf{74 \pm 2.56}$

4 and 6. The results are reported on the `test-split`. Models are tuned for hyper-parameters - `batch-size {32,64,128,256,512}`, `hidden-size {10,50,100}`.

SNMM model performs the best across all the scenarios; MSE and MAE for SNMM are the lowest across all the models. Comparable performance can be seen in NMM-L model where we enforce linearity of the space feature in log space. We highlight the benefit of enforcing structured constraints by comparison to several other models that are unconstrained. Although constrained models usually have lower capacity, SNMM is able to recover the performance lost due to constraints when compared to the constrained models and even improve on them. This clearly shows that the generalization ability of the shape constrained model is due to regularization with respect to the structure.

6 CONCLUSION AND FUTURE WORK

In this work, we presented a shelf space allocation problem for retail use case. A general framework of predict and optimize is discussed in detail with regards to trade-off between tractability of the optimization and prediction model accuracy. We discuss the need for multiplicative models and the difficulty in estimating space elasticity on a sparse dataset (most often this is the case in real world problems). To that extent, we propose a Scaled Neural Multiplicative Model (SNMM) that satisfies the conditions: non-negativity of sales and concavity with respect to the space feature. The optimization problem is formulated as a convex problem which can solved using convex solvers. Specifically, we use the power cone formulation provided by `cvxpy`. The model proposed in this work is general enough to extend to many other applications such as advertisement optimization, revenue maximization, etc. In future, we plan to explore these areas and enhance the model to handle different sets of constraints jointly.

References

Rishabh Agarwal, Levi Melnick, Nicholas Frosst, Xuezhou Zhang, Ben Lengerich, Rich Caruana, and Geoffrey E Hinton. Neural additive models: Interpretable machine learning with neural nets. *Advances in Neural Information Processing Systems*, 34:4699–4711, 2021.

Dennis J Aigner and Shih-fan Chu. On estimating the industry production function. *The American Economic Review*, 58(4):826–839, 1968.

Brandon Amos, Lei Xu, and J Zico Kolter. Input convex neural networks. In *International Conference on Machine Learning*, pp. 146–155. PMLR, 2017.

Moshe E Ben-Akiva, Steven R Lerman, Steven R Lerman, et al. *Discrete choice analysis: theory and application to travel demand*, volume 9. MIT press, 1985.

Dimitris Bertsimas and Nathan Kallus. From predictive to prescriptive analytics. *Management Science*, 66(3):1025–1044, 2020.

Teresa Bianchi-Aguiar, Alexander Hübner, Maria Antónia Carravilla, and José Fernando Oliveira. Retail shelf space planning problems: A comprehensive review and classification framework. *European Journal of Operational Research*, 289(1):1–16, 2021.

Ronald C Curhan. The relationship between shelf space and unit sales in supermarkets. *Journal of Marketing Research*, 9(4):406–412, 1972.

Martin Eisend. Shelf space elasticity: A meta-analysis. *Journal of Retailing*, 90(2):168–181, 2014.

Adam N Elmachtoub and Paul Grigas. Smart "predict, then optimize". *Management Science*, 68(1): 9–26, 2022.

Karthick Gopalswamy and Reha Uzsoy. A data-driven iterative refinement approach for estimating clearing functions from simulation models of production systems. *International Journal of Production Research*, 57(19):6013–6030, 2019.

Karthick Gopalswamy, Yahya Fathi, and Reha Uzsoy. Valid inequalities for concave piecewise linear regression. *Operations Research Letters*, 47(1):52–58, 2019.

Maya Gupta, Dara Bahri, Andrew Cotter, and Kevin Canini. Diminishing returns shape constraints for interpretability and regularization. *Advances in neural information processing systems*, 31, 2018.

Alexander H Hübner and Heinrich Kuhn. Shelf and inventory management with space-elastic demand. In *Operations research proceedings 2010*, pp. 405–410. Springer, 2011.

Alexander H Hübner and Heinrich Kuhn. Retail category management: State-of-the-art review of quantitative research and software applications in assortment and shelf space management. *Omega*, 40(2):199–209, 2012.

Marina Karampatsa, Evangelos Grigoroudis, and Nikolaos F Matsatsinis. Retail category management: A review on assortment and shelf-space planning models. *Operational Research in Business and Economics*, pp. 35–67, 2017.

A Gürhan Kök and Marshall L Fisher. Demand estimation and assortment optimization under substitution: Methodology and application. *Operations Research*, 55(6):1001–1021, 2007.

Venus Hiu Ling Lo. *Capturing Product Complementarity in Assortment Optimization*. PhD thesis, Cornell University, 2019.

John Ashworth Nelder and Robert WM Wedderburn. Generalized linear models. *Journal of the Royal Statistical Society: Series A (General)*, 135(3):370–384, 1972.

Garrett van Ryzin and Siddharth Mahajan. On the relationship between inventory costs and variety benefits in retail assortments. *Management Science*, 45(11):1496–1509, 1999.

Juan Pablo Vielma, Shabbir Ahmed, and George Nemhauser. Mixed-integer models for nonseparable piecewise-linear optimization: Unifying framework and extensions. *Operations research*, 58 (2):303–315, 2010.

Raymond E Wright. Logistic regression. 1995.

Eda Yücel, Fikri Karaesmen, F Sibel Salman, and Metin Türkay. Optimizing product assortment under customer-driven demand substitution. *European Journal of Operational Research*, 199(3): 759–768, 2009.

A APPENDIX

Proof. **Lemma 4.1**

$$\left(\sum_i x_i y_i \right)^2 \le \left(\sum_i x_i^2 \right)\left(\sum_i y_i^2 \right) \quad \text{By Cauchy Shwartz inequality}$$

$$\le \left(\sum_i x_i \right)\left(\sum_i y_i \right) \quad \because x_i \le 1, y_i \le 1 \implies x_i^2 \le x_i, y_i^2 \le y_i$$

$$\sum_i x_i y_i \le 1 \quad \because \sum_i x_i = \sum_i y_i = 1$$

Trivially, $\sum_i x_i y_i \ge 0$ since $x_i, y_i \ge 0 \ \forall i$ $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad$ □