De-randomizing MCMC dynamics with the diffusion Stein operator

Abstract

Approximate Bayesian inference determines proper descriptors of an intractable target distribution – in essence, an optimization problem within a family of distributions. For example, Langevin dynamics (LD) extracts asymptotically exact samples from a diffusion process because the time evolution of its marginal distributions constitutes a curve that minimizes the KL-divergence via steepest descent in the Wasserstein space. Parallel to LD, Stein variational gradient descent (SVGD) similarly minimizes the KL, albeit endowed with a novel Stein-Wasserstein distance, by *deterministically* transporting a set of particle samples, thus de-randomizes the stochastic diffusion process. We propose de-randomized kernel-based particle samplers to all diffusion-based samplers known as MCMC dynamics. Following previous work in interpreting MCMC dynamics, we equip the Stein-Wasserstein space with a fiber-Riemannian Poisson structure, with the capacity of characterizing a fiber-gradient Hamiltonian flow that simulates MCMC dynamics. Such dynamics discretize into generalized SVGD (GSVGD), a Stein-type deterministic particle sampler, with particle updates coinciding with applying the *diffusion Stein operator* to a kernel function. We demonstrate empirically that GSVGD can de-randomize complicated MCMC dynamics, which combine the advantages of auxiliary momentum variables and Riemannian structure, while maintaining the high sample quality from an interacting particle system.

1 Introduction

Evaluating an un-normalized target distribution π is a centerpiece of Bayesian inference, due to its ubiquitous presence in posterior distributions. Markov chain Monte Carlo (MCMC) methods fulfill this objective by generating asymptotically exact random samples from the distribution, a significant subset of which involves discretization of continuous-time diffusion processes, most notably Langevin diffusion, stochastic gradient Hamiltonian Monte Carlo (HMC) (Chen et al., 2014) and their further generalizations, which we collectively call MCMC *dynamics* (Ma et al., 2015). Despite its simplicity and theoretical soundness, this diffusion-based sampling often suffers from slow convergence and small effective sample sizes, largely due to the auto-correlation of the samples.

As an alternative to the simulation of stochastic systems, particle-based variational inference (Liu and Wang, 2016; Chen et al., 2018; Liu et al., 2019a) partially addresses the shortcomings of MCMC by replacing the Langevin diffusion, the simplest MCMC dynamics, with a deterministic interacting particle system that transports a set of interacting particles towards the target distribution. Theoretically speaking, Langevin diffusion encodes an evolution of density that minimizes the KL-divergence through steepest descent in the 2-Wasserstein space (Jordan et al., 1998), and particle-based variational inference (PARVI) approximates such evolution using reproducing kernel Hilbert space (RKHS) (Liu, 2017; Liu et al., 2019a), thus de-randomizing the Langevin diffusion process.

Submitted to 35th Conference on Neural Information Processing Systems (NeurIPS 2021). Do not distribute.

Figure 1: A diagram showing the contribution of our paper: we extend and generalize previous work linking Langevin diffusion to gradient flow on the 2-Wasserstein metric W_2 through the Fokker-Planck equation, and the W_2 gradient flow linking to SVGD (Liu and Wang, 2016) through projection onto an RKHS; MCMC dynamics (Ma et al., 2015) are interpreted as a fiber-gradient Hamiltonian (fGH) flow on W_2 (Liu et al., 2019b), and a projection onto an RKHS yields GSVGD.

Given the elegant theoretical link between Langevin diffusion and PARVI, one crucial question arises: can we leverage the advantages of general MCMC dynamics onto de-randomized particle systems? While it is difficult to formulate them as a direct Wasserstein gradient flow, Liu et al. (2019b) propose an interpretation for "regular" MCMC dynamics that augments the 2-Wasserstein space with a fiber bundle, forming a fiber-Riemannian Poisson manifold, under which MCMC dynamics follows a Hamiltonian flow on the fiber bundle, and a gradient flow on each fiber.

We show that adapting the fiber-Riemannian Hamiltonian flow to the Stein-Wasserstein metric (Liu, 2017; Duncan et al., 2019) yields a vector field in the form of applying the *diffusion Stein operator* to the kernel function $k(\cdot, \boldsymbol{\theta})$. A discretization of this vector field gives generalized SVGD (GSVGD), an interacting particle system that generalizes SVGD, with particle updates balancing an attractive force maximizing the log-likelihood and a repulsive force preventing a "mode collapse" of particles. We further demonstrate that the connection drawn between LD and SVGD (Liu and Wang, 2016; Liu, 2017; Liu et al., 2019a) is retraced by GSVGD, reaffirming our claim that GSVGD mirrors SVGD in approximating a larger class of MCMC diffusion processes. Within the generality of our framework, we can develop PARVI algorithms that exploit two key types of possible acceleration in MCMC dynamics: auxiliary momentum variables and an adaptive Riemannian parametrization that allows for fast and efficient exploration of the probability space (Patterson and Teh, 2013), as shown in Table 1.

MCMC dynamics	$\mathbf{A}$	$\mathbf{C}$	auxiliary variable	Riemannian	PARVI variant
SGLD (Welling and Teh, 2011)	$\mathbf{I}$	$\mathbf{0}$	✗	✗	SVGD (Liu and Wang, 2016)
Blob (Chen et al., 2018)
SGRLD (Patterson and Teh, 2013)	$\mathbf{G}(\boldsymbol{\theta})^{-1}$	$\mathbf{0}$	✗	✓	Riemannian SVGD (Liu and Zhu, 2017)
SGHMC (Chen et al., 2014)	$\begin{pmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & A\mathbf{I} \end{pmatrix}$	$\begin{pmatrix} \mathbf{0} & -\mathbf{I} \\ \mathbf{I} & \mathbf{0} \end{pmatrix}$	✓	✗	HMC-blob (Liu et al., 2019b)
SGRHMC (Ma et al., 2015)	$\begin{pmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{G}(\boldsymbol{\theta})^{-1} \end{pmatrix}$	$\begin{pmatrix} \mathbf{0} & -\mathbf{G}(\boldsymbol{\theta})^{-1/2} \\ \mathbf{G}(\boldsymbol{\theta})^{-1/2} & \mathbf{0} \end{pmatrix}$	✓	✓	this work (SGRHMC-Stein)

Table 1: An overview of MCMC dynamics along with their PARVI approximations. Between **(a)**, **(c)** and **(b)**, **(d)**, MCMC dynamics are stochastic single-chain simulations of a diffusion process (shown by one instance of a particle trajectory); PARVI approximates the time evolution of MCMC densities with a set of particles with deterministic dynamics; Between **(a)**, **(b)** and **(c)**, **(d)**, the two systems follows different time evolutions, as **(a)**, **(b)** describe a gradient flow and **(c)**, **(d)** describe a fiber-gradient Hamiltonian flow (with momentum variable r).

2 MCMC dynamics – the relevant bits

In this paper, we consider the problem of extracting samples $\boldsymbol{\theta} \in \mathbb{R}^D = \Omega$ from an un-normalized target distribution $\pi(\boldsymbol{\theta}) \propto e^{-H(\boldsymbol{\theta})}$. We begin by reviewing the key characteristics of MCMC dynamics, with particular emphasis on Langevin dynamics and its Fokker-Planck equation (FPE), the partial differential equation (PDE) that depicts the time evolution of a diffusion process. As the FPE of LD conforms to a gradient flow structure in the 2-Wasserstein metric space of probability

measures $(\mathcal{P}(\Omega), W_2)$ (Jordan et al., 1998), we then briefly cover gradient flow on $\mathcal{P}(\Omega)$ as an infinite-dimensional Riemannian manifold. Through the lens of gradient flow, we can see SVGD as a deterministic interacting particle system that approximates the gradient flow of LD through (i) gradient flow on a novel Stein-Wasserstein metric (Liu, 2017) or (ii) projecting the gradient flow direction onto an RKHS (Liu et al., 2019a). With Wasserstein gradient flow (WGF) neatly linking to both LD and SVGD, we see that a diffusion process LD and an interacting particle system SVGD are, respectively, a stochastic instance and a deterministic approximation of the same Wasserstein gradient flow.

Notations: We use ∇f to denote the gradient of a scalar-valued function, and $\nabla \cdot \mathbf{f}$ the divergence of a vector-valued function, $\nabla \cdot \mathbf{A}$ applies the divergence operator to each row of a matrix-valued function, $\dot{\rho}_t$ the "partial derivative" with respect to t.

2.1 Langevin dynamics and its Fokker-Planck equation

In this paper, we consider MCMC dynamics in the form of Itō diffusion processes following the stochastic differential equation (SDE) formula

$$d\boldsymbol{\theta}_t = \mathbf{f}(\boldsymbol{\theta}_t)dt + \sqrt{2\boldsymbol{\Sigma}(\boldsymbol{\theta}_t)}d\mathbf{W}_t, \tag{1}$$

consisting of drift coefficient $\mathbf{f} : \mathbb{R}^D \mapsto \mathbb{R}^D$, diffusion coefficient $\sqrt{2\boldsymbol{\Sigma}}$ and a D-dimensional Brownian motion $\mathbf{W}_t$. Ma et al. (2015) provide a complete recipe of all Itō diffusion processes converging to the target measure π. The simplest MCMC dynamics takes the form of Langevin diffusion (Langevin, 1908), which moves towards higher densities with $\mathbf{f}(\boldsymbol{\theta}) = \nabla \log \pi(\boldsymbol{\theta})$ perturbed by white noise $\boldsymbol{\Sigma} = \mathbf{I}$. Given an initial distribution $\boldsymbol{\theta}_0 \sim \rho_0$, the Fokker-Planck equation describes the time evolution of the density of $\boldsymbol{\theta}_t$ (Risken, 1996)

$$\dot{\rho}_t + \nabla \cdot (\rho_t \nabla \log \pi) - (\nabla^\top \nabla) : (\rho_t \mathbf{I}) = 0, \tag{2}$$

Given the property of the differential operators, we can rewrite (2) as

$$\dot{\rho}_t = \nabla \cdot (\nabla \rho_t - \rho_t \nabla \log \pi) = \nabla \cdot \left(\rho_t \nabla \underbrace{\frac{\delta \mathrm{KL}\left[\rho_t \,\|\, \pi\right]}{\delta \rho_t}}_{\text{first variation of KL}\left[\rho_t \,\|\, \pi\right]} \right), \tag{3}$$

a crucial step in developing the Wasserstein gradient flow perspective of LD, as $\nabla \delta E / \delta \rho$ coincides with the differential of functionals induced by the Wasserstein metric.

2.2 Gradient flow on $(\mathcal{P}(\Omega), W_2)$

The conventional gradient flow (or the steepest descent curve) that minimizes a smooth function $F : \mathbb{R}^D \mapsto \mathbb{R}$ follows the PDE: $\dot{\mathbf{x}}_t + \nabla F(\mathbf{x}_t) = 0$. Defining gradient flow on $\mathcal{P}(\Omega)$, informally written as $\dot{\rho}_t + \nabla_W E(\rho_t) = 0$, requires a definition of the differentiation ∇_W, which requires an understanding of the Wasserstein metric, i.e., the inner product g_ρ defined on its tangent space.

To simplify the discussion, we restrict the discussion on measures with a density function and with finite second-order moments. For each ρ, the tangent space constitutes smooth functions integrating to zero, $T_\rho \mathcal{P}(\Omega) = \{f | f \in C^\infty(\Omega), \int f(\boldsymbol{\theta})d\boldsymbol{\theta} = 0\}$, in that a curve on $\mathcal{P}(\Omega)$ preserves volume. The cotangent space constitutes an equivalence class of smooth functions in differing constant, noted as the quotient space $T_\rho^* \mathcal{P}(\Omega) = C^\infty(\Omega)/\mathbb{R}$. We characterize the inner product space using the metric tensor $G(\rho) : T_\rho \mathcal{P} \mapsto T_\rho^* \mathcal{P}$, a one-to-one mapping between elements of the tangent space and those of the cotangent space. The inner product g_ρ is defined by the inverse of the metric tensor, often denoted as Onsager operators (Onsager, 1931)

$$g_\rho(f_1, f_2) = \int f_1 G(\rho) f_2 d\boldsymbol{\theta} = \int \phi_1 G(\rho)^{-1} \phi_2 d\mathbf{x}, \quad \phi_1 = G(\rho)f_1, \ \phi_2 = G(\rho)f_2. \tag{4}$$

The Wasserstein Onsager operator takes the form $G(\rho)^{-1} : \phi \mapsto -\nabla \cdot (\rho \nabla \phi)$. In the context of Wasserstein gradient flow, we define $\nabla_W E(\rho)$ as $G(\rho)^{-1} \frac{\delta E(\rho)}{\delta \rho}$, leading to the formulation:

$$0 = \dot{\rho}_t - G(\rho)^{-1} \frac{\delta E(\rho_t)}{\delta \rho_t} = \dot{\rho}_t + \nabla \cdot \left(\rho_t \nabla \frac{\delta E(\rho_t)}{\delta \rho_t} \right). \tag{5}$$

The first variation of $\mathrm{KL}\left[\rho \,\|\, \pi\right]$ takes the form of $\frac{\delta \mathrm{KL}[\rho \,\|\, \pi]}{\delta \rho} = \log \rho / \pi + 1$, leading to the conclusion by Jordan et al. (1998) that the time evolution of the Langevin diffusion follows a curve of steepest descent in the Wasserstein space, laying the foundation for approximation using particle interaction.

2.3 SVGD as gradient flow

PARVI transports a set N of interacting particles $\left\{ \boldsymbol{\theta}_t^{(i)} \right\}_{1 \leq i \leq N}$ towards the target distribution over time. Take SVGD for example, denoting the empirical measure at time t as $\hat{\rho}_t = \frac{1}{N} \sum_{i=1}^{N} \delta_{\boldsymbol{\theta}_t^{(i)}}$, SVGD (Liu and Wang, 2016) follows the update rule

$$\dot{\boldsymbol{\theta}}_t = \mathbb{E}_{\boldsymbol{\theta}' \sim \hat{\rho}_t} \left[k(\boldsymbol{\theta}_t, \boldsymbol{\theta}') \nabla \log \pi(\boldsymbol{\theta}') + \nabla_2 k(\boldsymbol{\theta}_t, \boldsymbol{\theta}') \right] = \mathbf{v}_{\mathcal{H}}(\boldsymbol{\theta}_t | \hat{\rho}_t), \tag{6}$$

where $k(\cdot, \cdot)$ defines a positive-definite kernel, and ∇_2 denotes the gradient of the second argument. The FPE of SVGD can be written as

$$\dot{\rho}_t - \nabla \cdot \left(\rho_t \mathcal{K}_\rho \nabla \frac{\delta \mathrm{KL} \left[\rho_t \parallel \pi \right]}{\delta \rho_t} \right) = 0, \tag{7}$$

where $\mathcal{K}_\rho$ is an integral operator: $\mathcal{K}_\rho \mathbf{f}(\boldsymbol{\theta}) = \int k(\boldsymbol{\theta}, \boldsymbol{\theta}') \mathbf{f}(\boldsymbol{\theta}') \mathrm{d}\rho(\boldsymbol{\theta})$. We shall denote the RKHS defined by k as $\mathcal{H}$. While significantly different from LD at first glance, SVGD approximates the WGF of LD by

- kernelizing the Wasserstein Onsager operator to give $G_{\mathcal{H}}(\rho)^{-1} : \phi \mapsto -\nabla \cdot (\rho \mathcal{K}_\rho \nabla \phi)$, thus forming the Stein-Wasserstein metric $W_{\mathcal{H}}$ (Liu, 2017; Duncan et al., 2019);

- projecting the gradient flow vector field $\mathbf{v}(\rho) = -\nabla \frac{\delta \mathrm{KL}[\rho \parallel \pi]}{\delta \rho} = \nabla \log \pi - \nabla \log \rho$ onto $\mathcal{H}$ (Liu et al., 2019a).

The interpretation of SVGD yields valuable insights. While gradient flow (7) on $(\mathcal{P}(\Omega), W_2)$ yields no closed-form energy functional (Chen et al., 2018), it behooves to absorb the operator $\mathcal{K}_\rho$ into the definition of the Stein-Wasserstein metric to guarantee a tractable energy functional. In the meantime, the kernelization trick transforms a gradient flow (2) simulated by a diffusion process into an approximate deterministic transportation of particles – in other words, de-randomizes it.

2.4 MCMC dynamics as fiber-gradient Hamiltonian flow

The concept of *flow* on $(\mathcal{P}(\Omega), W_2)$ is a generalization of the gradient flow: given a vector field $\mathbf{v} : \mathcal{P}(\Omega) \mapsto \bigcup_{\rho \in \mathcal{P}} T_\rho$, the flow of $\mathbf{v}$ is defined as $\dot{\rho}_t = \mathbf{v}(\rho_t)$. Liu et al. (2019b) interpret "regular" MCMC dynamics on $(\mathcal{P}(\Omega), W_2)$ as a flow, by combining gradient flow and Hamiltonian flow on the Wasserstein space, yielding a fiber-Riemannian manifold structure of $\mathcal{P}(\Omega)$, a fiber bundle consisting of Riemannian manifolds. In the context of a "regular" MCMC dynamics taking the form of underdamped Langevin dynamics (Chen et al., 2014), the diffusion matrix $\mathbf{A}$ determines the gradient flow on each fiber, and the curl matrix $\mathbf{C}$ determines the Hamiltonian flow on the fiber bundle. The Hamiltonian flow keeps $\mathrm{KL} \left[\rho \parallel \pi \right]$ constant while encouraging fast exploration of the probability space; the fiber gradient flow minimizes the KL. The fiber-gradient Hamiltonian flow determines a PARVI updated in the form of

$$\dot{\boldsymbol{\theta}}_t = \left(\mathbf{A}(\boldsymbol{\theta}_t) + \mathbf{C}(\boldsymbol{\theta}_t) \right) \left(\nabla \log \pi(\boldsymbol{\theta}_t) - \widehat{\nabla} \log \rho_t(\boldsymbol{\theta}_t) \right) = \mathbf{v}^{\mathbf{A}, \mathbf{C}}(\rho_t), \tag{8}$$

where the intractable $\widehat{\nabla} \log \rho_t$ is approximated via the Blob method (Carrillo et al., 2019).

2.5 Other relevant works

Myriad works (Chen et al., 2018; Liu et al., 2019a; Zhang et al., 2020) seek alternatives to approximating the gradient flow (2) through kernelization. Notably, Chen et al. (2018); Liu et al. (2019a) construct PARVI algorithms by approximating the term $\nabla \log \rho_t(\boldsymbol{\theta}_t)$ in the gradient flow, namely with Blob method (Carrillo et al., 2019), kernel density estimation (Liu et al., 2019a) and Stein gradient estimator (Li and Turner, 2017).

Our work connects to the "Nesterov acceleration" viewpoint for PARVI methods. Ma et al. (2019) demonstrate that underdamped LD (or the stochastic gradient HMC (Chen et al., 2014)) accelerates the steepest descent steps taken by the overdamped LD, forming an analog of Nesterov acceleration for MCMC methods. Wang and Li (2019) present a framework for Nesterov's accelerated gradient method in the Wasserstein space, which consists of augmenting the energy functional with the kinetic energy of an additional momentum variable.

Figure 2: Particle samples of an underlying 2-dimensional correlated mixture π extracted by SVGD (Liu and Wang, 2016), Riemannian SVGD (Liu and Zhu, 2017), SGHMC-Stein and SGRHMC-Stein. While all methods explore the mode nearest to initialization, only RHMC explores all three modes.

Figure 3: A visualisation of sampler trajectories on the tri-Gaussian density π. With the help of momentum variable and efficient Riemannian parametrization of $(\mathbf{A}, \mathbf{C})$, SGRHMC-Stein aggressively explores the space, while other methods cannot escape the local component.

3 PARVI for MCMC dynamics – a general recipe

In this section, we outline the main contribution of this work, which traces the roadmap delineated by previous works to explore the flow interpretation, as well as the approximation by interacting particle systems of general form MCMC dynamics (Ma et al., 2015) in the form of Itō diffusion:

$$\mathbf{f}(\boldsymbol{\theta}) = \frac{1}{\pi(\boldsymbol{\theta})} \nabla \cdot \left[\pi(\boldsymbol{\theta}) \left(\mathbf{A}(\boldsymbol{\theta}) + \mathbf{C}(\boldsymbol{\theta}) \right) \right], \tag{9}$$

$$\boldsymbol{\Sigma}(\boldsymbol{\theta}) = \mathbf{A}(\boldsymbol{\theta}), \tag{10}$$

where $\mathbf{A}$ and $\mathbf{C}$ are positive-semidefinite and skew-symmetric matrix-valued functions, respectively. This general framework covers *all* continuous Itō diffusion processes with stationary distribution π, most notably LD, stochastic gradient HMC (Chen et al., 2014), stochastic gradient Nosé-Hoover thermostat (SGNHT) (Ding et al., 2014) and stochastic gradient Riemannian Hamiltonian Monte Carlo (SGRHMC) (Ma et al., 2015).

We demonstrate in this section that applying the fiber-gradient Hamiltonian flow structure of MCMC dynamics to the Stein-Wasserstein metric yields a flow on $(\mathcal{P}(\Omega), W_{\mathcal{H}})$ that discretizes into GSVGD, a PARVI that updates particles with the diffusion Stein operator (Gorham et al., 2019), suggesting that the infinitesimal generator of MCMC diffusion processes offers a "kernel smoothing" derandomization. Apart from the Stein-Wasserstein metric, the analog of GSVGD generalizing SVGD extends to other interpretations of SVGD, namely that GSVGD takes steepest descent minimizing the KL-divergence in incremental transformation of particles, and that that GSVGD projects the fiber-gradient Hamiltonian flow onto an RKHS.

3.1 Fiber-gradient Hamiltonian flow on $(\mathcal{P}(\Omega), W_{\mathcal{H}})$

Similar to (2), we derive the FPE of MCMC dynamics (9)-(10)

$$\dot{\rho}_t = -\nabla \cdot (\rho_t \mathbf{f}) + (\nabla^{\top} \nabla) : (\rho_t \mathbf{A}) \tag{11}$$

$$= -\nabla \cdot (\rho_t \mathbf{f}) + (\nabla^{\top} \nabla) : (\rho_t (\mathbf{A} + \mathbf{C})) \tag{12}$$

$$= \nabla \cdot \left(\rho_t \left(\mathbf{A} + \mathbf{C} \right) \nabla \log \frac{\rho_t}{\pi} \right) = \nabla \cdot \left(\rho_t \left(\mathbf{A} + \mathbf{C} \right) \nabla \frac{\delta \mathrm{KL} \left[\rho_t \parallel \pi \right]}{\delta \rho_t} \right), \tag{13}$$

which is a curve in $\mathcal{P}(\Omega)$ following the continuity equation $\dot{\rho}_t + \nabla \cdot (\rho_t \mathbf{v}_t) = 0$.

We can derive GSVGD from the fiber-Riemannian manifold perspective of $(\mathcal{P}(\Omega), W_{\mathcal{H}})$ (Liu et al., 2019b). The Riemannian structure of the Stein-Wasserstein metric induces the tangent space $T_\rho \mathcal{P}(\Omega) = \overline{\{ \mathcal{K}_\rho \nabla f \mid f \in C^\infty(\Omega) \}}^{\mathcal{H}^D}$. An orthogonal projection $\mathfrak{p}_\rho$ from $\mathcal{L}^2_\rho(\Omega)$ to $T_\rho \mathcal{P}$ is uniquely defined as such vector in $T_\rho \mathcal{P}$ that satisfies $\nabla \cdot (\rho \mathcal{K}_\rho \phi) = \nabla \cdot (\rho \mathfrak{p}_\rho(\phi))$. Using Theorem 5 from Liu et al. (2019b), we arrive at a fiber-gradient Hamiltonian flow

$$\mathcal{W}_{\mathrm{KL}_\pi}(\rho) = \mathfrak{p}_\rho \left((\mathbf{A} + \mathbf{C}) \nabla \log \pi / \rho \right). \tag{14}$$

It is straightforward to verify that $\mathbf{v}_{\mathcal{H}}^{\mathbf{A}, \mathbf{C}}(\rho) = \mathcal{K}_\rho (\mathbf{A} + \mathbf{C}) \nabla \log \pi / \rho$ induces the same evolution of distribution as $\mathcal{W}_{\mathrm{KL}_\pi}(\rho)$.

3.2 fRH flow induces the diffusion Stein operator

The flow on $(\mathcal{P}(\Omega), W_{\mathcal{H}})$ following $\dot{\rho}_t = \nabla \cdot \left(\rho_t \mathbf{v}_{\mathcal{H}}^{\mathbf{A}, \mathbf{C}}(\rho_t) \right)$ requires spatial and temporal discretization suitable for a particle update. While $\mathbf{v}_{\mathcal{H}}^{\mathbf{A}, \mathbf{C}}$ constitutes a vector field on $\mathcal{P}(\Omega)$, its calculation does not involve an explicit term of $\nabla \log \rho$ inside the expectation.

$$\mathbf{v}_{\mathcal{H}}^{\mathbf{A}, \mathbf{C}}(\rho) = \mathcal{K}_\rho \left[(\mathbf{A} + \mathbf{C}) \nabla \log \pi / \rho \right] \tag{15}$$

$$= \mathbb{E}_{\boldsymbol{\theta}' \sim \rho} k(\cdot, \boldsymbol{\theta}') (\mathbf{A}(\boldsymbol{\theta}') + \mathbf{C}(\boldsymbol{\theta}')) \nabla \log \pi(\boldsymbol{\theta}') - \int k(\cdot, \boldsymbol{\theta}')(\mathbf{A}(\boldsymbol{\theta}') + \mathbf{C}(\boldsymbol{\theta}')) \nabla \rho(\boldsymbol{\theta}') \mathrm{d} \boldsymbol{\theta}'$$

$$= \mathbb{E}_{\boldsymbol{\theta}' \sim \rho} (\mathbf{A}(\boldsymbol{\theta}') + \mathbf{C}(\boldsymbol{\theta}')) \nabla \log \pi(\boldsymbol{\theta}') k(\cdot, \boldsymbol{\theta}') + \underbrace{\int \nabla \cdot \left((\mathbf{A}(\boldsymbol{\theta}') + \mathbf{C}(\boldsymbol{\theta}')) k(\cdot, \boldsymbol{\theta}') \right) \mathrm{d} \rho}_{\text{integration by parts}}$$

$$= \mathbb{E}_{\boldsymbol{\theta}' \sim \rho} \big[\underbrace{\mathbf{f}(\boldsymbol{\theta}') k(\cdot, \boldsymbol{\theta}')}_{\text{weighted drift coefficient (9)}} + \underbrace{(\mathbf{A} + \mathbf{C})(\boldsymbol{\theta}') \nabla_2 k(\cdot, \boldsymbol{\theta}')}_{\text{repulsive force}} \big]. \tag{16}$$

$$= \mathbb{E}_{\boldsymbol{\theta}' \sim \rho} \underbrace{\frac{1}{\pi} \nabla \cdot \left(\pi (\mathbf{A} + \mathbf{C}) k(\cdot, \boldsymbol{\theta}') \right)}_{\text{diffusion Stein operator } \mathcal{T}_\pi^{\mathbf{A}, \mathbf{C}} k(\cdot, \boldsymbol{\theta}')}. \tag{17}$$

Interestingly, the kernelized flow $\mathbf{v}_{\mathcal{H}}^{\mathbf{A}, \mathbf{C}}$ coincides with calculating the expectation of the diffusion Stein operator $\mathcal{T}_\pi^{\mathbf{A}, \mathbf{C}}$ (Gorham et al., 2019) applied to the kernel function $k(\boldsymbol{\theta}, \cdot)$, a family of operators mapping functions to zero-expectation functions under an (un-normalized) distribution π. When ρ_t is approximated by the empirical measure of its particles $\hat{\rho}_t = 1/N \sum_{i=1}^N \delta_{\boldsymbol{\theta}_t^{(i)}}$, we can derive the particle update as

$$\dot{\boldsymbol{\theta}}_t = \mathbf{v}_{\mathcal{H}}^{\mathbf{A}, \mathbf{C}}(\hat{\rho}_t) = \mathbb{E}_{\boldsymbol{\theta}' \sim \hat{\rho}_t} \mathcal{T}_\pi^{\mathbf{A}, \mathbf{C}} k(\boldsymbol{\theta}_t, \boldsymbol{\theta}') = \frac{1}{N} \sum_{i=1}^N \mathcal{T}_\pi^{\mathbf{A}, \mathbf{C}} k(\boldsymbol{\theta}_t, \boldsymbol{\theta}_t^{(i)}), \boldsymbol{\theta}_0^{(i)} \sim \rho_0. \tag{18}$$

Similar to SVGD, the particle update of GSVGD leverages between a weighted average of the diffusion vector and a repulsive force that prevents the particles from collapsing to the local mode. The fiber-gradient Hamiltonian flow on $(\mathcal{P}(\Omega), W_{\mathcal{H}})$ informs a duality between diffusion Stein operator and MCMC dynamics, extending the use of Stein operators beyond a generalization of kernel Stein discrepancy (Gorham et al., 2019).

3.3 How does GSVGD approximate MCMC dynamics?

While the derivation of GSVGD originates from the flow interpretation on $W_{\mathcal{H}}$, the analogy between SVGD and GSVGD expresses in equivalent alternative forms discussed by previous literature, further shedding light on the interpretation of GSVGD.

Originally, the SVGD update is expressed in the form of the functional gradient (Liu and Wang, 2016). For MCMC dynamics, we similarly derive the functional gradient with respect to the push-forward

measure $\rho_\epsilon = (\text{id} + (\mathbf{A} + \mathbf{C})\mathbf{v})_\# \, \rho, \mathbf{v} \in \mathcal{H}^D$,

$$\nabla_{\mathbf{v}} \, \text{KL} \left[\rho_\epsilon \parallel \pi \right] \big|_{\mathbf{v}=\mathbf{0}} = -\mathbb{E}_{\boldsymbol{\theta}' \sim \rho} \mathcal{T}_\pi^{\mathbf{A},\mathbf{C}} k(\cdot, \boldsymbol{\theta}') = -\mathbf{v}_{\mathcal{H}}(\rho). \tag{19}$$

This presents a generalization of the discussion in Liu and Wang (2016) that GSVGD takes incremental transformation of particles in the direction that minimizes KL-divergence, with such transformation warped by $(\mathbf{A}, \mathbf{C})$ and constrained by RKHS.

Alternatively, Liu et al. (2019a) explains SVGD as the projection of the original gradient flow direction $\mathbf{v}(\rho_t) = \nabla \log \pi / \rho_t$ onto an RKHS. We similarly generalize this explanation for SVGD, as it projects $\mathbf{v}^{\mathbf{A},\mathbf{C}}(\rho_t) = (\mathbf{A} + \mathbf{C})\nabla \log \pi / \rho_t \in \mathcal{L}_\rho^2$ onto $\mathcal{H}^D$, as

$$\mathbf{v}_{\mathcal{H}}^{\mathbf{A},\mathbf{C}}(\rho) = \max \cdot \arg\max_{\mathbf{v} \in \mathcal{H}^D, \|\mathbf{v}\|_{\mathcal{H}^D}=1} \langle \mathbf{v}^{\mathbf{A},\mathbf{C}}(\rho), \mathbf{v} \rangle_{\mathcal{L}_\rho^2}. \tag{20}$$

The analogy of GSVGD goes further when we consider the transportation of N particles as inferring a joint distribution $\pi^{\otimes N}$, which will be covered in the supplements.

3.4 GSVGD in practice

We can apply GSVGD updates (16) to de-randomize myriad MCMC dynamics (e.g., Table 1). Notably, GSVGD establishes the previously unexplored duality between Riemannian Langevin diffusion (Patterson and Teh, 2013) and Riemannian SVGD (Liu and Zhu, 2017).

To fully harness the capacity of our framework, we can introduce auxiliary (momentum) variables $\mathbf{r}$ to help with the exploration of probability space, namely augmenting the target distribution as $\pi(\boldsymbol{\theta}, \mathbf{r}) = \pi(\boldsymbol{\theta})\mathcal{N}(\mathbf{r}|\mathbf{0}, \boldsymbol{\Sigma})$, therefore achieving de-randomized PARVI variant of stochastic gradient Hamiltonian Monte Carlo (SGHMC) (Chen et al., 2014). Further leveraging a positive definite $\mathbf{G}(\boldsymbol{\theta})$ to efficiently explore the target distribution yields a PARVI variant of SGRHMC (Ma et al., 2015).

We can accelerate GSVGD with previous acceleration techniques applied to PARVI methods based on LD, most notably the Wasserstein Nesterov's method (Liu et al., 2019a) with the following updates on $\left(\boldsymbol{\theta}_k^{(i)} \right)$ and $\left(\boldsymbol{\xi}_k^{(i)} \right)$ initialized at the same location:

$$\boldsymbol{\xi}_{k+1}^{(i)} = \boldsymbol{\theta}_k^{(i)} + \epsilon \mathbf{v}_{\mathcal{H}}^{\mathbf{A},\mathbf{C}} \left(\boldsymbol{\theta}_k^{(i)} \Big| \hat{\rho}_k(\boldsymbol{\theta}_k) \right), \tag{21}$$

$$\boldsymbol{\theta}_{k+1}^{(i)} = c_1(c_2 - 1) \left(\boldsymbol{\xi}_{k+1}^{(i)} - \boldsymbol{\xi}_k^{(i)} \right). \tag{22}$$

When GSVGD is used in conjunction with auxiliary momentum variables, we can employ symmetric splitting for leapfrog-like steps for GSVGDs de-randomizing with momentum,

$$\mathbf{r}_{k+1/2}^{(i)} = \mathbf{r}_k^{(i)} + \frac{\epsilon}{2} \mathbf{v}_{\mathcal{H}}^{\mathbf{A},\mathbf{C}} \left(\mathbf{r}_k^{(i)} \Big| \hat{\rho}(\boldsymbol{\theta}_k, \mathbf{r}_k) \right), \tag{23}$$

$$\boldsymbol{\theta}_{k+1}^{(i)} = \boldsymbol{\theta}_k^{(i)} + \epsilon \mathbf{v}_{\mathcal{H}}^{\mathbf{A},\mathbf{C}} \left(\boldsymbol{\theta}_k^{(i)} \Big| \hat{\rho}(\boldsymbol{\theta}_k, \mathbf{r}_{k+1/2}) \right), \tag{24}$$

$$\mathbf{r}_{k+1}^{(i)} = \mathbf{r}_{k+1/2}^{(i)} + \frac{\epsilon}{2} \mathbf{v}_{\mathcal{H}}^{\mathbf{A},\mathbf{C}} \left(\mathbf{r}_{k+1/2}^{(i)} \Big| \hat{\rho}(\boldsymbol{\theta}_{k+1}, \mathbf{r}_{k+1/2}) \right). \tag{25}$$

4 Experiments

4.1 Stein PARVI on toy data

To demonstrate the efficacy of our methods, we use various PARVI to infer a 3-component mixture of crescent-shaped target measure inspired by Ma et al. (2015). The parametrizations of Riemannian SVGD and SGRHMC-Stein coincide with those in Ma et al. (2015). While all methods explore the nearest mode well (See figure 2), only the SGRHMC-Stein manage to effectively use the Riemannian formulation of $(\mathbf{A}, \mathbf{C})$ to explore the 2 other modes (see figure 3), showcasing the gain in efficiency with application-specific dynamics.

4.2 GSVGD on Bayesian neural networks

We apply GSVGD of advanced MCMC dynamics to the inference of Bayesian neural networks, taking a simple structure of one hidden layer, and an output with Gaussian likelihood. We

Figure 4: Evolution of RHMC dynamics (blue lines) over four highlighted time increments k (red points) on tri-Gaussian density. The RHMC explores the space by an aggressive outward expansion followed by convergence to the tri-Gaussian density π.

Figure 5: The trace plots of test log-likelihoods of BNN experiments: the x-axis determines the iterations. The figures show the effect of the momentum variables in SGHMC-Stein and SGNHT-Stein in exploring the posterior space, and quickly converging to better optima. The jaggedness of SVGD showcases that while SVGD empirically works with stochastic gradient, it is less robust with stochastic gradient noise.

opt for the fully Bayesian specification of BNN, where the precision parameter of the weight prior and the Gaussian likelihood follows $\text{Gamma}(1, 0.1)$. The GSVGD variant used in the experiments include SGHMC (Chen et al., 2014) and SGNHT (Ding et al., 2014), with target measure $\pi(\boldsymbol{\theta}, \mathbf{r}, \boldsymbol{\xi}) = \pi(\boldsymbol{\theta})\mathcal{N}(\mathbf{r}|\mathbf{0}, \sigma^2\mathbf{I})\mathcal{N}(\boldsymbol{\xi}|A\mathbf{1}, \mu^{-1}\mathbf{I})$, and $\mathbf{A} = \begin{pmatrix} \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & A\mathbf{I} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} \end{pmatrix}, \mathbf{C} = \begin{pmatrix} \mathbf{0} & -\mathbf{I} & \mathbf{0} \\ \mathbf{I} & \mathbf{0} & (\mu\sigma^2)^{-1}\text{diag}(\mathbf{r}) \\ \mathbf{0} & -(\mu\sigma^2)^{-1}\text{diag}(\mathbf{r}) & \mathbf{0} \end{pmatrix}$. As a MCMC form suitable when used in conjunction with stochastic gradients, SGNHT takes additional temperature parameter $\boldsymbol{\xi}$. The log-likelihood results in Table 2 demonstrates that the de-randomized MCMC dynamics achieve superior predictive performance than their LD counterparts, largely thanks to the exploration of probability space from the momentum variables (See figure 5), with a notable gain from symmetric splitting. SGNHT-Stein methods show robustness to hyperparameter selection and stochastic gradient noise.

Method	boston	concrete	energy	kin8nm	power	yacht	year	protein
LD	-2.52 (0.15)	-3.16 (0.04)	-2.36 (0.06)	0.10 (0.03)	-2.85 (0.03)	-1.60 (0.08)	-3.69 (0.01)	-3.05 (0.01)
SVGD	-2.60 (0.46)	-3.11 (0.10)	-1.95 (0.07)	1.03 (0.25)	-2.82 (0.03)	-1.66 (0.17)	-3.61 (0.00)	-2.91 (0.01)
Blob	-2.52 (0.32)	-3.19 (0.07)	-1.67 (0.03)	0.95 (0.03)	-2.81 (0.04)	-1.34 (0.07)	-3.60 (0.00)	-2.93 (0.01)
SGHMC-Blob	**-2.42 (0.12)**	**-2.95 (0.01)**	-1.36 (0.26)	1.23 (0.02)	-2.77 (0.04)	**-0.73 (0.46)**	-3.60 (0.02)	-3.73 (0.16)
SGNHT	-2.59 (0.13)	-3.41 (0.05)	-2.44 (0.04)	0.75 (0.02)	-2.86 (0.02)	-2.64 (0.04)	-3.69 (0.00)	-3.01 (0.01)
DE	-2.68 (0.63)	-2.96 (0.15)	**-0.45 (0.14)**	1.16 (0.02)	-2.80 (0.03)	-1.04 (0.40)	-3.68 (0.00)	-3.00 (0.01)
SGHMC-Stein	-2.81 (0.71)	-3.04 (0.25)	-1.40 (0.82)	**1.25 (0.02)**	**-2.76 (0.04)**	-0.86 (0.33)	**-3.59 (0.00)**	-2.90 (0.01)
SGNHT-Stein	-2.49 (0.30)	-2.97 (0.11)	**-0.44 (0.10)**	1.24 (0.03)	-2.78 (0.04)	-0.85 (0.21)	**-3.59 (0.00)**	**-2.85 (0.01)**

Table 2: The test log-likelihood (higher is better) results for selected datasets in UCI repository. The results are reported in mean (standard deviation) form averaged over 20 runs in the first 6 columns and 6 runs in the last 2, with the best performing model marked in boldface.

5 Discussion

In this section, we discuss open questions pertinent to the "optimality" of the diffusion Stein operator as a PARVI update, and a direction for future work regarding the fiber-gradient Hamiltonian flow of the chi-squared divergence, inspired by Chewi et al. (2020).

5.1 Alternatives to the diffusion Stein operator

The vector field $\mathbf{v}^{\mathbf{A},\mathbf{C}}(\rho) = (\mathbf{A} + \mathbf{C}) \nabla \log \frac{\pi}{\rho}$ is not the unique vector field that induces the time evolution of MCMC dynamics: indeed, two vector fields induce the same curve in $\mathcal{P}(\Omega)$ when they differ by a divergence-free field: $\nabla \cdot (\rho(\mathbf{v}_1(\rho) - \mathbf{v}_2(\rho))) = 0$. Notably, Liu et al. (2019b) propose an alternative vector field $\mathbf{v}_0^{\mathbf{A},\mathbf{C}}(\rho) = \mathbf{v}^{\mathbf{A},\mathbf{C}}(\rho) - \mathbf{C}\nabla \log \rho + \nabla \cdot \mathbf{C}$ [1]. A projection onto RKHS yields a Stein PARVI with update

$$\dot{\boldsymbol{\theta}}_t = \mathbb{E}_{\rho_t} \left[\frac{1}{\pi} \nabla \cdot (\pi (\mathbf{A} + \mathbf{C}) k(\boldsymbol{\theta}_t, \cdot)) + \mathbf{A}\nabla_2 k(\boldsymbol{\theta}_t, \cdot) \right], \tag{26}$$

consistent with the standard formulation infinitesimal generator. However, the expectation in (26) notably does not converge to zero when $\rho_t = \pi$ (Gorham et al., 2019), suggesting not all vector fields inducing the same curve translate into appropriate PARVI.

5.2 Generalizing LAWGD with the diffusion Stein operator

The application of the diffusion Stein operator generalizes Laplacian Adjusted Wasserstein Gradient Descent (LAWGD) (Chewi et al., 2020), which views SVGD as a kernelized Wasserstein gradient flow driven by the chi-squared divergence. Given the first variation of the chi-squared divergence $\frac{\delta\chi^2[\rho \, \| \, \pi]}{\delta\rho} = \frac{2\rho}{\pi}$, we can verify that the vector field $\mathbf{v}_{\mathcal{H}}^{\mathbf{A},\mathbf{C}} = \mathcal{K}_\rho \left[(\mathbf{A} + \mathbf{C}) \nabla \log \frac{\pi}{\rho} \right]$ is rewritten as

$$2\mathbf{v}_{\mathcal{H}}^{\mathbf{A},\mathbf{C}} = \mathcal{K}_\pi \left[(\mathbf{A} + \mathbf{C}) \nabla \frac{2\pi}{\rho} \right] = \mathcal{K}_\pi \left[(\mathbf{A} + \mathbf{C}) \nabla \frac{\delta\chi^2 [\rho_t \, \| \, \pi]}{\delta\rho_t} \right]. \tag{27}$$

The chi-squared divergence perspectives enriches GSVGD as a kernelized fiber-gradient Hamiltonian flow minimizing the chi-squared divergence. Replacing the vector field $\mathcal{K}_\pi \left[(\mathbf{A} + \mathbf{C}) \nabla \frac{\mathrm{d}\rho}{\mathrm{d}\pi} \right]$ with $\mathbf{u} = \nabla \cdot \left(\mathcal{K}_\pi \left[(\mathbf{A} + \mathbf{C}) \frac{\mathrm{d}\rho}{\mathrm{d}\pi} \right] \right)$, we get the dissipation formula for the KL-divergence

$$\frac{\mathrm{dKL}\,[\rho_t \, \| \, \pi]}{\mathrm{d}t} = -\mathbb{E}_\pi \left\langle \nabla \frac{\rho}{\pi}, \nabla \cdot \left(\mathcal{K}_\pi \left[(\mathbf{A} + \mathbf{C}) \frac{\rho}{\pi} \right] \right) \right\rangle \tag{28}$$

$$= -\mathbb{E}_\pi \left[\frac{\rho}{\pi} \mathcal{A}_\pi^{\mathbf{A},\mathbf{C}} \mathcal{K}_\pi \frac{\rho}{\pi} \right], \tag{29}$$

where $\mathcal{A}_\pi^{\mathbf{A},\mathbf{C}}$ represents the infinitesimal generator that induces the diffusion Stein operator: $\mathcal{A}_\pi^{\mathbf{A},\mathbf{C}} g = \frac{1}{2\pi} \nabla \cdot (\pi(\mathbf{A} + \mathbf{C})\nabla g)$. Following the construction of LAWGD, we choose the kernel so that $\mathcal{K}_\pi = \left(\mathcal{A}_\pi^{\mathbf{A},\mathbf{C}} \right)^{-1}$. While it remains to be seen what kernel induces this equality, the diffusion Stein operator yields a more flexible formulation, while retaining the exponential convergence of LAWGD.

6 Conclusion

We further the application of the Stein operator and its generalizations by proposing GSVGD, an interacting particle system transporting a set of particles towards a target distribution π, while mimicking MCMC dynamics with respect to its time evolution of marginal distributions. We showcase theoretically and empirically, that a generalization of SVGD helps bridge the well-researched field of efficient MCMC dynamics with a deterministic interacting particle system with high-quality samples.

---

[1]In fact, a divergence-free vector field $\mathbf{C}_0 \nabla \log \rho - \nabla \cdot \mathbf{C}_0$ can be constructed out of an arbitrary skew-symmetric matrix-valued function $\mathbf{C}_0$

References

Tianqi Chen, Emily Fox, and Carlos Guestrin. Stochastic gradient Hamiltonian Monte Carlo. In *International conference on machine learning*, pages 1683–1691, 2014.

Yi-An Ma, Tianqi Chen, and Emily B. Fox. A complete recipe for stochastic gradient MCMC. In *NIPS'15 Proceedings of the 28th International Conference on Neural Information Processing Systems - Volume 2*, volume 28, pages 2917–2925, 2015.

Qiang Liu and Dilin Wang. Stein variational gradient descent: A general purpose Bayesian inference algorithm. In *Advances in Neural Information Processing Systems*, volume 29, pages 2378–2386, 2016.

Changyou Chen, Ruiyi Zhang, Wenlin Wang, Bai Li, and Liqun Chen. A unified particle-optimization framework for scalable Bayesian sampling. In *UAI*, pages 746–755, 2018.

Chang Liu, Jingwei Zhuo, Pengyu Cheng, Ruiyi Zhang, and Jun Zhu. Understanding and accelerating particle-based variational inference. In *International Conference on Machine Learning*, pages 4082–4092, 2019a.

Richard Jordan, David Kinderlehrer, and Felix Otto. The variational formulation of the Fokker-Planck equation. *SIAM Journal on Mathematical Analysis*, 29(1):1–17, 1998.

Qiang Liu. Stein variational gradient descent as gradient flow. In *Advances in Neural Information Processing Systems*, volume 30, pages 3115–3123, 2017.

Chang Liu, Jingwei Zhuo, and Jun Zhu. Understanding MCMC dynamics as flows on the Wasserstein space. In *International Conference on Machine Learning*, pages 4093–4103, 2019b.

A. Duncan, N. Nuesken, and L. Szpruch. On the geometry of Stein variational gradient descent. *arXiv preprint arXiv:1912.00894*, 2019.

Sam Patterson and Yee Whye Teh. Stochastic gradient Riemannian Langevin dynamics on the probability simplex. In *Advances in Neural Information Processing Systems 26*, volume 26, pages 3102–3110, 2013.

Max Welling and Yee W. Teh. Bayesian learning via stochastic gradient Langevin dynamics. In *Proceedings of the 28th International Conference on Machine Learning*, pages 681–688, 2011.

Chang Liu and Jun Zhu. Riemannian Stein variational gradient descent for bayesian inference. In *AAAI*, pages 3627–3634, 2017.

P. Langevin. Sur la theorie du mouvement Brownien. *Compt. Rendus*, 146:530–533, 1908.

Hannes Risken. Fokker-Planck equation. pages 63–95, 1996.

Lars Onsager. Reciprocal relations in irreversible processes. ii. *Physical Review*, 37(12):2265–2279, 1931.

José Antonio Carrillo, Katy Craig, and Francesco S. Patacchini. A blob method for diffusion. *Calculus of Variations and Partial Differential Equations*, 58(2):1–53, 2019.

Jianyi Zhang, Ruiyi Zhang, Lawrence Carin, and Changyou Chen. Stochastic particle-optimization sampling and the non-asymptotic convergence theory. In *International Conference on Artificial Intelligence and Statistics*, pages 1877–1887, 2020.

Yingzhen Li and Richard E. Turner. Gradient estimators for implicit models. In *International Conference on Learning Representations*, 2017.

Yi-An Ma, Niladri S. Chatterji, Xiang Cheng, Nicolas Flammarion, Peter L. Bartlett, and Michael I. Jordan. Is there an analog of Nesterov acceleration for MCMC. *arXiv preprint arXiv:1902.00996*, 2019.

Yifei Wang and Wuchen Li. Accelerated information gradient flow. *arXiv preprint arXiv:1909.02102*, 2019.

Nan Ding, Youhan Fang, Ryan Babbush, Changyou Chen, Robert D Skeel, and Hartmut Neven. Bayesian sampling using stochastic gradient thermostats. In *Advances in Neural Information Processing Systems 27*, volume 27, pages 3203–3211, 2014.

Jackson Gorham, Andrew B. Duncan, Sebastian J. Vollmer, and Lester Mackey. Measuring sample quality with diffusions. *Annals of Applied Probability*, 29(5):2884–2928, 2019.

Sinho Chewi, Thibaut Le Gouic, Chen Lu, Tyler Maunu, and Philippe Rigollet. SVGD as a kernelized Wasserstein gradient flow of the chi-squared divergence. In *Advances in Neural Information Processing Systems*, volume 33, pages 2098–2109, 2020.