When is Momentum Extragradient Optimal?
A Polynomial-Based Analysis

Junhyung Lyle Kim[*]   *jlylekim@rice.edu*
Rice University, Department of Computer Science

Gauthier Gidel   *gauthier.gidel@umontreal.ca*
Université de Montréal, Department of Computer Science and Operations Research
Mila – Quebec AI Institute, Canada CIFAR AI Chair

Anastasios Kyrillidis   *anastasios@rice.edu*
Rice University, Department of Computer Science

Fabian Pedregosa   *pedregosa@google.com*
Google DeepMind

Reviewed on OpenReview: *https://openreview.net/forum?id=ZLVbQEu4Ab*

Abstract

The extragradient method has gained increasing attention due to its favorable convergence on differentiable games. The eigenvalues of the game vector field's Jacobian are distributed on the complex plane, making game dynamics more complicated than single objective minimization. Even for bilinear games, the simple gradient method diverges, while the extragradient method converges. Our work focuses on studying the momentum extragradient method, which was recently proven to converge at an accelerated rate for bilinear games (Azizian et al., 2020b). By using polynomial-based analysis, we identify three cases of game Jacobian eigenvalue distribution where the momentum extragradient method exhibits accelerated convergence rates. These cases include when the eigenvalues are distributed on the (positive) real line, both on the real line along with complex conjugates, and only as complex conjugates. We also derive the optimal hyperparameters for each scenario.

1 Introduction

While most machine learning problems are formulated as a minimization problem, a growing number of works rely instead on game formulations that involve multiple players and objectives. For example, generative adversarial networks (GANs) (Goodfellow et al., 2014), actor-critic models (Pfau & Vinyals, 2016), and sharpness aware minimization (Foret et al., 2021) can all be formulated as two-player games. Naturally, there is an increasing interest in the theoretical understanding of such differentiable games.

Due to the interaction of multiple players and objectives, there are issues in optimizing differentiable games that are not present in minimization problems. Most notably, the eigenvalues of the game Jacobian are distributed on the *complex plane*, exhibiting richer dynamics compared to single-objective minimization, where the Hessian eigenvalues lie on the *real line*. As a result, even for simple bilinear games, standard algorithms –like the gradient method– fail to converge (Mescheder et al., 2018; Balduzzi et al., 2018; Gidel et al., 2019; Berard et al., 2020).

On the contrary, the extragradient method (EG), originally introduced by Korpelevich (1976), converges for bilinear games (Tseng, 1995). As such, many recent works analyze EG from different perspectives, including

---

[*]Authors after JLK are listed in alphabetical order.
This paper extends Kim et al. (2022) presented at the NeurIPS 2022 Optimization for Machine Learning Workshop.

variational inequality (Gidel et al., 2018; Gorbunov et al., 2022), stochastic (Li et al., 2021), and distributed (Liu et al., 2020; Beznosikov et al., 2021) settings.

Most existing works, including those mentioned earlier, analyze EG and relevant algorithms by assuming some structure on the objectives, such as (strong) monotonicity or Lipschitzness (Solodov & Svaiter, 1999; Tseng, 1995; Daskalakis & Panageas, 2018; Ryu et al., 2019; Azizian et al., 2020a). Such assumptions, in the context of differentiable games, confine the distribution of the eigenvalues of the game Jacobian; for instance, strong monotonicity implies a lower bound on the real part of the eigenvalues, and the Lipschitz assumption implies an upper bound on the magnitude of the eigenvalues of the Jacobian.

Azizian et al. (2020b) showed that, however, such assumptions may be too crude to study the game dynamics, which is captured by the eigenvalue *distribution* on the complex plane. Through polynomial-based analysis, they show that some first-order methods can sometimes achieve faster rates using momentum, by replacing the smoothness and monotonicity assumptions with more precise assumptions on the distribution of the Jacobian eigenvalues, represented by simple shapes like ellipses or line segments. In particular, Azizian et al. (2020b) proved that the extragradient method can be accelerated with momentum for bilinear games.

In this work, we take a similar path, but ask the reverse question: for what shape of Jacobian spectrum does the momentum extragradient (MEG) method achieve optimal performance? Such "reverse" analysis enables us to study the behavior of MEG in specific settings depending on the hyperparameter setup, encompassing minimization (where the eigenvalues of the Jacobian are all on the positive real line), regularized bilinear games (where the eigenvalues are all complex conjugates), and the intermediate case (where the eigenvalues are both on the real line and as complex conjugates), as illustrated in Figure 1. Our contributions can be summarized as follows.

- We derive the residual polynomials of MEG when the game vector field is an affine function. We then identify three different modes of convergence for MEG depending on the hyperparameter setup. Such analysis can then be applied to different eigenvalue structures of the (game) Jacobian (c.f., Theorem 3).

- For each case of the eigenvalue structure, we derive the optimal hyperparameters of MEG and its (asymptotic) convergence rates. For minimization, MEG exhibits "super-acceleration," where a constant improvement upon classical lower bound rate is attained,[1] similarly to the gradient method with momentum (GDM) with cyclical step sizes (Goujaud et al., 2022). For the other two cases involving imaginary eigenvalues, MEG exhibits accelerated convergence rates with the derived optimal hyperparameters.

- Finally, we compare the convergence rates with other first-order methods, including the gradient (GD), the gradient method with momentum (GDM), and the extragradient (EG) methods. For the classes of games considered, none of the aforementioned methods (asymptotically) converge at an accelerated rate (c.f., Corollaries 1 and 2), in contrast to MEG. In Section 7, we validate our theoretical findings through numerical experiments, including a scenario that slightly deviates from our initial assumption.

2 Problem Setup and Related Work

Following Letcher et al. (2019); Balduzzi et al. (2018), we define the n-player differentiable game as a family of twice continuously differentiable losses $\ell_i : \mathbb{R}^d \to \mathbb{R}$, for $i = 1, \ldots, n$. The player i controls the parameter $w^{(i)} \in \mathbb{R}^{d_i}$. We denote the concatenated parameters by $w = [w^{(1)}, \ldots, w^{(n)}] \in \mathbb{R}^d$, where $d = \sum_{i=1}^n d_i$.

For this problem, a Nash equilibrium satisfies: $w^{(i)\star} \in \arg\min_{w^{(i)} \in \mathbb{R}^{d_i}} \ell_i \big(w^{(i)}, w^{(\neg i)\star}\big) \quad \forall i \in \{1, \ldots, n\}$, where the notation $\cdot^{(\neg i)}$ denotes all indices except for i. We also define the vector field v of the game as the concatenation of the individual gradients: $v(w) = [\nabla_{w^{(1)}} \ell_1(w) \cdots \nabla_{w^{(n)}} \ell_n(w)]^\top$, and denote its associated Jacobian with ∇v.

---

[1]Note that achieving this improvement is possible by having additional information beyond just the largest (smoothness) and smallest (strong convexity) eigenvalues of the Jacobian.

Figure 1: *Convergence rates of MEG in terms of the game Jacobian eigenvalues.* The step sizes for MEG, h and γ, and the momentum parameter m are set up according to each case of Theorem 3, illustrating three distinct convergence modes of MEG. For each case, the red line indicates the robust region (c.f., Definition 1) where MEG achieves the optimal convergence rate.

Unfortunately, finding Nash equilibria for general games is impractical to solve (Shoham & Leyton-Brown, 2008; Letcher et al., 2019).[2] Instead, we focus on finding a stationary point of the vector field v, since a Nash equilibrium is always a stationary point of the gradient dynamics. That is, we want to solve:

$$\text{Find} \quad w^\star \in \mathbb{R}^d \quad \text{such that} \quad v(w^\star) = 0. \tag{1}$$

Notation. $\mathfrak{R}(z)$ and $\mathfrak{I}(z)$ respectively denote the real and the imaginary part of a complex number z. The spectrum of a matrix M is denoted by $\text{Sp}(M)$, and its spectral radius by $\rho(M) := \max\{|\lambda| : \lambda \in \text{Sp}(M)\}$. $M \succ 0$ denotes that M is a positive-definite matrix. $\mathbb{C}_+$ denotes the complex plane with positive real part, and $\mathbb{R}_+$ denotes positive real numbers.

2.1 Related Work

The extragradient method, originally introduced in Korpelevich (1976), is a popular algorithm for solving (unconstrained) variational inequality problems in (1) (Gidel et al., 2018). There are several works that study the convergence rate of EG for (strongly) monotone problems (Tseng, 1995; Solodov & Svaiter, 1999; Nemirovski, 2004; Monteiro & Svaiter, 2010; Mokhtari et al., 2020; Gorbunov et al., 2022). Under similar settings, stochastic variants of EG are studied in Palaniappan & Bach (2016); Hsieh et al. (2019; 2020); Li et al. (2021). However, as mentioned earlier, assumptions like (strong) monotonicity or Lipchtizness may not accurately represent *how* Jacobian eigenvalues are distributed.

Instead, we make more fine-grained assumptions on these eigenvalues, to obtain the optimal hyperparameters and convergence rates for MEG via polynomial-based analysis. Such analysis dates back to the development of the conjugate gradient method (Hestenes & Stiefel, 1952), and is still actively used; for instance, to derive lower bounds (Arjevani & Shamir, 2016), to develop accelerated decentralized algorithms (Berthier et al., 2020), and to analyze average-case performance (Pedregosa & Scieur, 2020; Domingo-Enrich et al., 2021).

On that end, we use the following lemma (Chihara, 2011), which elucidates the connection between first-order methods and (residual) polynomials, when the vector field v is affine. First-order methods are the ones in which the sequence of iterates w_t is in the span of previous gradients: $w_t \in w_0 + \text{span}\{v(w_0), \ldots, v(w_{t-1})\}$.

Lemma 1 (Chihara (2011)). *Let w_t be the iterate generated by a first-order method after t iterations, with $v(w) = Aw + b$. Then, there exists a real polynomial P_t, of degree at most t, satisfying:*

$$w_t - w^\star = P_t(A)(w_0 - w^\star), \tag{2}$$

where $P_t(0) = 1$, and $v(w^\star) = Aw^\star + b = 0$.

---

[2]Finding Nash equilibria can be reformulated as a nonlinear complementarity problem, which is PPAD hard (Daskalakis et al., 2009; Letcher et al., 2019).

By taking ℓ_2-norms, (2) further implies the following worst-case convergence rate:

$$\|w_t - w^\star\| = \|P_t(A)(w_0 - w^\star)\| \leqslant \|P_t(Z\Lambda Z^{-1})\| \cdot \|w_0 - w^\star\| \leqslant \sup_{\lambda \in \mathcal{S}^\star} |P_t(\lambda)| \cdot \|Z\|\|Z^{-1}\| \cdot \|w_0 - w^\star\|, \quad (3)$$

where $A = Z\Lambda Z^{-1}$ is the diagonalization of A,[3] and the constant $\|Z\|\|Z^{-1}\|$ disappears if A is a normal matrix. Hence, the worst-case convergence rate of a first-order method can be analyzed by studying the associated residual polynomial P_t evaluated at the eigenvalues λ of the Jacobian $\nabla v = A$, distributed over the set $S^\star$.

Azizian et al. (2020b) took a similar approach, and proposed a geometric interpretation of the conditioning of a game via the notion of *spectral shape*, defined as the set containing all eigenvalues of the Jacobian. Specifically, let $\mathcal{A}_{\mathcal{S}^\star}$ be the set of matrices A whose spectrum, denoted by $\mathrm{Sp}(A)$, belong to a set $\mathcal{S}^\star$ on the complex plane with positive real part:

$$\mathcal{A}_{\mathcal{S}^\star} := \{A \in \mathbb{R}^{d \times d} : \mathrm{Sp}(A) \subset \mathcal{S}^\star \subset \mathbb{C}_+\}.$$

When $\mathcal{S}^\star$ is "simple" (e.g., a real line segment or a disc on the complex plane), Azizian et al. (2020b) characterized the lower bound and the optimality of some first-order methods (Nemirovskij & Yudin, 1983; Arjevani & Shamir, 2016); see also Section 5.

Yet, it may be possible that additional knowledge about the exact shape of $\mathcal{S}^\star$ could result in a more detailed analysis and faster rates. We follow this perspective: based on the (residual) polynomials of MEG, specific spectral shapes $\mathcal{S}^\star$ emerge, where MEG shows the optimal performance. Recently, similar directions have appeared in the optimization literature (Oymak, 2021; Goujaud et al., 2022), which focus on convex smooth minimization and demonstrate that knowing more than just the largest and smallest eigenvalues of the Jacobian (i.e., the smoothness and strong convexity) leads to better convergence rates.

3 Momentum Extragradient via Chebyshev Polynomials

In this section, we study the *momentum extragradient method* (MEG) and its performance, through the lens of (residual) polynomials. MEG iterates as follows:

$$\text{(MEG)} \quad w_{t+1} = w_t - hv(w_t - \gamma v(w_t)) + m(w_t - w_{t-1}). \quad (4)$$

Here, h is the step size, γ is the extrapolation step size, and m is the momentum parameter. The extragradient (EG) method (without momentum, i.e., $m = 0$) was first proposed by Korpelevich (1976) for saddle point problems. Recently, it has gained significant attention due to its convergence behavior for some differentiable games, such as bilinear games, where the standard gradient method fails to converge (Gidel et al., 2019; Azizian et al., 2020b;a). For completeness, we remind the gradient method with momentum (GDM):

$$\text{(GDM)} \quad w_{t+1} = w_t - hv(w_t) + m(w_t - w_{t-1}), \quad (5)$$

from which the gradient method (GD) can be obtained by setting $m = 0$.

As MEG in (4) is a first-order method (Arjevani & Shamir, 2016; Azizian et al., 2020b), we can apply Lemma 1 and study the residual polynomials of MEG, which are summarized in the next theorem.

Theorem 1 (Residual polynomials of MEG). *Consider the momentum extragradient method (MEG) in* (4). *If $v(w) = Aw + b$, the associated residual polynomials of MEG are:*

$$\tilde{P}_0(\lambda) = 1, \quad \tilde{P}_1(\lambda) = 1 - \frac{h\lambda(1-\gamma\lambda)}{1+m}, \quad and \quad \tilde{P}_{t+1}(\lambda) = (1 + m - h\lambda(1-\gamma\lambda))\tilde{P}_t(\lambda) - m\tilde{P}_{t-1}(\lambda).$$

Further, let $T_t(\cdot)$ and $U_t(\cdot)$ be the Chebyshev polynomials of the first and the second kind, respectively. Then, the above expressions simplify to:

$$P_t^{MEG}(\lambda) = m^{t/2}\left(\frac{2m}{1+m}T_t(\sigma(\lambda)) + \frac{1-m}{1+m}U_t(\sigma(\lambda))\right) \quad with \quad \sigma(\lambda) = \frac{1+m-h\lambda(1-\gamma\lambda)}{2\sqrt{m}}, \quad (6)$$

where we refer to the term $\sigma(\lambda) \equiv \sigma(\lambda; h, \gamma, m)$ as the link function.

---

[3]Note that almost all matrices are diagonalizable over $\mathbb{C}$, in the sense that the set of non-diagonalizable matrices has Lebesgue measure zero (Hetzel et al., 2007).

As can be seen in (6), MEG's link function is *quadratic* with respect to λ. Interestingly, GDM also admits a similar expression, but with a different link function. For completeness, we write below the residual polynomials of GDM, expressed in Chebyshev polynomials (Pedregosa, 2020):

$$P_t^{\text{GDM}}(\lambda) = m^{t/2}\left(\tfrac{2m}{1+m}T_t(\xi(\lambda)) + \tfrac{1-m}{1+m}U_t(\xi(\lambda))\right) \text{ with } \xi(\lambda) = \tfrac{1+m-h\lambda}{2\sqrt{m}}. \tag{7}$$

Notice that the residual polynomials of MEG in (6) and that of GDM in (7) are identical, except for the link functions $\sigma(\lambda)$ and $\xi(\lambda)$, which enter as arguments in $T_t(\cdot)$ and $U_t(\cdot)$. This difference is crucial, as the Chebyshev polynomials behave very differently based on the domain, per the following lemma:

Lemma 2 (Goujaud & Pedregosa (2022)). *Let z be a complex number, and let $T_t(\cdot)$ and $U_t(\cdot)$ be the Chebyshev polynomials of the first and the second kind, respectively. The sequence $\left\{\left|\tfrac{2m}{1+m}T_t(z) + \tfrac{1-m}{1+m}U_t(z)\right|\right\}_{t\geqslant 0}$ grows exponentially in t for $z \notin [-1,1]$, while for $z \in [-1,1]$, the following bounds hold:*

$$|T_t(z)| \leqslant 1 \quad and \quad |U_t(z)| \leqslant t+1. \tag{8}$$

Therefore, to study the optimal convergence behavior of MEG, we are interested in the case where the set of step sizes and the momentum parameters lead to $|\sigma(\lambda; h, \gamma, m)| \leqslant 1$ so that we can use the bounds in (8). We will refer to those sets of eigenvalues and hyperparameters as the *robust region*, as defined below.

Definition 1 (Robust region of MEG). *Consider the MEG method in (4) expressed via Chebyshev polynomials, as in (6). We define the set of eigenvalues and hyperparameters such that the image of the link function $\sigma(\lambda; h, \gamma, m)$ lies in the interval $[-1, 1]$ as the **robust region**, and denote it with $\sigma^{-1}([-1, 1])$.*

Although polynomial-based analysis requires the assumption that the vector field is affine, it captures intuitive insights into how various algorithms behave in different settings, as we remark below.

Remark 1. *From the definition of $\xi(\lambda)$ in (7), one can infer why negative momentum can help the convergence of GDM (Gidel et al., 2019) when $\lambda \in \mathbb{R}_+$: it forces GDM to stay within the robust region, $|\xi(\lambda)| \leqslant 1$. One can also infer the divergence of GDM in the presence of complex eigenvalues, unless, for instance, complex momentum is used (Lorraine et al., 2022). Similarly, the residual polynomial of GD is $P_t^{GD}(\lambda) = (1 - h\lambda)^t$ (Goujaud & Pedregosa, 2022, Example 4.2), and can easily diverge in the presence of complex eigenvalues, which can potentially be alleviated by using complex step sizes. On the contrary, thanks to the quadratic link function of MEG in (6), it can converge for much wider subsets of complex eigenvalues.*

By analyzing the residual polynomials of MEG, we can also characterize the asymptotic convergence rate of MEG for any combination of hyperparameters, as summarized in the next theorem.

Theorem 2 (Asymptotic convergence rate of MEG). *Suppose $v(w) = Aw + b$. The asymptotic convergence rate of MEG in (4) is:[4]*

$$\limsup_{t\to\infty} \sqrt[2t]{\frac{\|w_t - w^\star\|}{\|w_0 - w^\star\|}} = \begin{cases} \sqrt[4]{m}, & if \;\; \bar{\sigma} \leqslant 1 \;\;\; (robust\ region); \\ \sqrt[4]{m}(\bar{\sigma} + \sqrt{\bar{\sigma}^2 - 1})^{1/2}, & if \;\; \bar{\sigma} \in \left(1, \tfrac{1+m}{2\sqrt{m}}\right); \\ \geqslant 1 \;(no\ convergence), & otherwise, \end{cases} \tag{9}$$

where $\bar{\sigma} = \sup_{\lambda \in \mathcal{S}^\star} |\sigma(\lambda; h, \gamma, m)|$, and $\sigma(\lambda; h, \gamma, m) \equiv \sigma(\lambda)$ is the link function of MEG defined in (6).

Optimal hyperparameters for MEG that we obtain in Section 4 minimize the asymptotic convergence rate above. Note that the optimal hyperparameters vary based on the set $\mathcal{S}^\star$, which we detail in Section 3.2.

---

[4]The reason why we take the $2t$-th root is to normalize by the number of vector field computations; we compare in Section 4 the asymptotic rate of MEG in (9) with other gradient methods that use a single vector field computation in the recurrences, such as GD and GDM.

3.1 Three Modes of the Momentum Extragradient

Within the robust region of MEG, we can compute its worst-case convergence rate based on (3) as follows:

$$\sup_{\lambda \in \mathcal{S}^\star} |P_t^{\text{MEG}}(\lambda)| \overset{(6)}{\leqslant} m^{t/2}\Big(\tfrac{2m}{1+m} \sup_{\lambda \in \mathcal{S}^\star} |T_t(\sigma(\lambda))| + \tfrac{1-m}{1+m} \sup_{\lambda \in \mathcal{S}^\star} |U_t(\sigma(\lambda))| \Big)$$

$$\overset{(8)}{\leqslant} m^{t/2}\Big(\tfrac{2m}{1+m} + \tfrac{1-m}{1+m}(t+1) \Big) \leqslant m^{t/2}(t+1). \tag{10}$$

Since the Chebyshev polynomial expressions of MEG in (6) and that of GDM[5] are identical except for the link functions, the convergence rate in (10) applies to both MEG and GDM, as long as the link functions $|\sigma(\lambda)|$ and $|\xi(\lambda)|$ are bounded by 1. As a result, we see that the asymptotic convergence rate in (9) only depends on the momentum parameter m, when the hyperparameters are restricted to the robust region. This fact was utilized in tuning GDM for strongly convex quadratic minimization (Zhang & Mitliagkas, 2019).

The robust region of MEG can be described with the four extreme points below (derivation in the appendix):

$$\sigma^{-1}(-1) = \tfrac{1}{2\gamma} \pm \sqrt{\tfrac{1}{4\gamma^2} - \tfrac{(1+\sqrt{m})^2}{h\gamma}}, \quad \text{and} \quad \sigma^{-1}(1) = \tfrac{1}{2\gamma} \pm \sqrt{\tfrac{1}{4\gamma^2} - \tfrac{(1-\sqrt{m})^2}{h\gamma}}. \tag{11}$$

The above four points and their intermediate values characterize the set of Jacobian eigenvalues λ that can be mapped to $[-1, 1]$. The distribution of these eigenvalues can vary in three different modes depending on the selected hyperparameters of MEG, as stated in the following theorem.

Theorem 3. *Consider the momentum extragradient method in (4), expressed with the Chebyshev polynomials as in (6). Then, the robust region (c.f., Definition 1) have the following three modes:*

- **Case 1:** *If $\frac{h}{4\gamma} \geqslant (1 + \sqrt{m})^2$, then $\sigma^{-1}(-1)$ and $\sigma^{-1}(1)$ are all real numbers;*

- **Case 2:** *If $(1 - \sqrt{m})^2 \leqslant \frac{h}{4\gamma} < (1 + \sqrt{m})^2$, then $\sigma^{-1}(-1)$ are complex, and $\sigma^{-1}(1)$ are real;*

- **Case 3:** *If $(1 - \sqrt{m})^2 > \frac{h}{4\gamma}$, then $\sigma^{-1}(-1)$ and $\sigma^{-1}(1)$ are all complex numbers.*

Remark 2. *Theorem 3 offers basic guidance on how to set up the hyperparameters for MEG. This depends on the Jacobian spectra of the game problem being considered. For instance, if one observes only real eigenvalues (i.e., the problem is in fact minimization), the main step size h should be at least $4\times$ larger than the extrapolation step size γ, based on the condition $\frac{h}{4\gamma} \geqslant (1 + \sqrt{m})^2$.*

We illustrate Theorem 3 in Figure 1. We first set the hyperparameters according to each condition in Theorem 3. We then discretize the interval $[-1, 1]$, and plot $\sigma^{-1}([-1, 1])$ for each case, represented by red lines. We can see the quadratic link function induced by MEG allows interesting eigenvalue dynamics to be mapped onto the $[-1, 1]$ segment, such as the cross-shape observed in Case 2. Moreover, although MEG exhibits the best rates within the robust region, it does not necessarily diverge outside of it, as in the second case of Theorem 2. We illustrate the convergence region of MEG measured by $\sqrt[2t]{|P_t(\lambda)|} < 1$ from (6) for $t = 2000$, with different colors indicating varying convergence rates, which slow down as one moves away from the robust region. Interestingly, Figure 1 (right) shows that MEG can also converge in the absence of monotonicity (i.e., in the presence of Jacobian eigenvalues with negative real part) (Gorbunov et al., 2023).

3.2 Robust Region-Induced Problem Cases

According to Theorem 3, we can classify the problem classes into three distinct cases, induced by different modes of the robust region, as illustrated in Figure 2. We explain each case in detail below.

---

[5] Asymptotically, GDM enjoys $\sqrt{m}$ convergence rate instead of the $\sqrt[4]{m}$ of MEG, as it uses a single vector field computation per iteration instead of the two. However, these are not directly comparable, as the values of m that correspond to the robust region are not the same.

Figure 2: *Illustration of the three spectrum models where MEG achieves accelerated convergence rates.*

Case 1: The problem reduces to minimization, where the Jacobian eigenvalues are distributed on the (positive) real line, but as a *union* of two intervals. We can model such spectrum as:

$$\text{Sp}(\nabla v) \subset \mathcal{S}_1^\star = [\mu_1, L_1] \cup [\mu_2, L_2] \subset \mathbb{R}_+. \tag{12}$$

Above generalizes the Hessian spectrum that arise in minimizing μ-strongly convex and L-smooth functions, i.e. $\lambda \in [\mu, L]$. This spectrum can be obtained from (12) by setting $\mu_1 = \mu$, $L_2 = L$, and $L_1 = \mu_2$. It was empirically observed that, during DNN training, sometimes a few eigenvalues of the Hessian have significantly larger magnitudes (Papyan, 2020). In such cases, (12) can be more precise than a single interval $[\mu, L]$. In particular, Goujaud et al. (2022) utilized (12), and showed that the GDM with alternating step sizes can achieve a (constant factor) improvement over the traditional lower bound for strongly convex and smooth quadratic objectives.

In Section 4, we show that MEG enjoys similar improvement. To show that, we define the following quantities following Goujaud et al. (2022), which will be used to obtain the convergence rate of MEG in (18) for this problem class.

$$\rho := \frac{L_2 + \mu_1}{L_2 - \mu_1} = \frac{1 + \tau}{1 - \tau}, \quad \text{and} \quad R := \frac{\mu_2 - L_1}{L_2 - \mu_1} \in [0, 1). \tag{13}$$

Here, ρ is the ratio between the center of $\mathcal{S}_1^\star$ and its radius, and $\tau := L_2/\mu_1$ is the inverse condition number. R is the relative gap of $\mu_2 - L_1$ and $L_2 - \mu_1$, which becomes 0 if $\mu_2 = L_1$ (i.e., $\mathcal{S}_1^\star$ becomes $[\mu_1, L_2]$).

Case 2: In this case, the Jacobian eigenvalues are distributed both on the real line and as complex conjugates, exhibiting a *cross-shaped* spectrum. We model this spectrum as:

$$\text{Sp}(\nabla v) \subset \mathcal{S}_2^\star = [\mu, L] \cup \{z \in \mathbb{C} : \mathfrak{R}(z) = c' > 0, \ \mathfrak{I}(z) \in [-c, c]\}. \tag{14}$$

The first set $[\mu, L]$ denotes a segment on the real line, reminiscent of the Hessian spectrum for minimizing μ-strongly convex and L-smooth functions. The second set has a fixed real component ($c' > 0$), along with imaginary components symmetric across the real line (i.e., complex conjugates), as the Jacobian is real.

This is a strict generalization of the purely imaginary interval $\pm[ai, bi]$ commonly considered in the bilinear games literature (Liang & Stokes, 2019; Azizian et al., 2020b; Mokhtari et al., 2020). While many recent papers on bilinear games cite GANs (Goodfellow et al., 2014) as a motivation, the work in Berard et al. (2020, Figure 4) empirically shows that the spectrum of GANs is not contained in the imaginary axis; the cross-shaped spectrum model above might be closer to some of the observed GAN spectra.

Case 3: In this case, the Jacobain eigenvalues are distributed only as complex conjugates, with a fixed real component, exhibiting a *shifted imaginary* spectrum. We model this spectrum as:

$$\text{Sp}(\nabla v) \subset \mathcal{S}_3^\star = [c + ai, c + bi] \cup [c - ai, c - bi] \subset \mathbb{C}_+. \tag{15}$$

Again, (15) generalizes bilinear games, where the spectrum reduces to $\pm[ai, bi]$ with $c = 0$.

Examples of Cases 2 and 3 in quadratic games. To understand these spectra better, we provide examples using quadratic games. Consider the following two player quadratic game, where $x \in \mathbb{R}^{d_1}$ and $y \in \mathbb{R}^{d_2}$ are the parameters controlled by each player, whose loss functions respectively are:

$$\ell_1(x,y) = \frac{1}{2}x^\top S_1 x + x^\top M_{12} y + x^\top b_1 \quad \text{and} \quad \ell_2(x,y) = \frac{1}{2}y^\top S_2 y + y^\top M_{21} x + y^\top b_2, \tag{16}$$

where $S_1, S_2 \succ 0$. Then, the vector field can be written as:

$$v(x,y) = \begin{bmatrix} S_1 x + M_{12} y + b_1 \\ M_{21} x + S_2 y + b_2 \end{bmatrix} = Aw + b, \text{ where } A = \begin{bmatrix} S_1 & M_{12} \\ M_{21} & S_2 \end{bmatrix}, \ w = \begin{bmatrix} x \\ y \end{bmatrix}, \text{ and } b = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}. \tag{17}$$

If $S_1 = S_2 = 0$ and $M_{12} = -M_{21}^\top$, the game Jacobian $\nabla v = A$ has only purely imaginary eigenvalues (Azizian et al., 2020b, Lemma 7), recovering bilinear games.

As the second and the third spectrum models in (14) and (15) generalize bilinear games, we can consider more complex quadratic games, where S_1 and S_2 does not have to be 0. Specifically, when $M_{12} = -M_{21}^\top$, and they share common bases with S_1 and S_2 specified in the below proposition, then $\text{Sp}(A)$ has a cross-shaped spectrum in (14) of Case 2 and a shifted imaginary spectrum in (15) of Case 3.

Proposition 1. *Let A be a matrix of the form $\begin{bmatrix} S_1 & B \\ -B^\top & S_2 \end{bmatrix}$, where $S_1, S_2 \succ 0$. Without loss of generality, assume that $dim(S_1) > dim(S_2) = d$. Then,*

- **Case 2:** $\text{Sp}(A)$ *has a cross-shape if there exist orthonormal matrices U, V and diagonal matrices D_1, D_2 such that $S_1 = U\,diag(a, \ldots, a, D_1)U^\top$, $S_2 = V\,diag(a, \ldots, a)V^\top$, and $B = UD_2V^\top$.*

- **Case 3:** $\text{Sp}(A)$ *has a shifted imaginary shape if there exist orthonormal matrices U, V and diagonal matrix D_2 such that $S_1 = U\,diag(a, \ldots, a)U^\top$, $S_2 = V\,diag(a, \ldots, a)V^\top$, and $B = UD_2V^\top$.*

We can interpret Case 3 as a *regularized* bilinear game, where S_1 and S_2 are diagonal matrices with a constant eigenvalue. This implies that the players cannot control their parameter x and y arbitrarily, which can be seen in the loss functions in (16), where S_1 and S_2 appears in terms $x^\top S_1 x$ and $y^\top S_2 y$. Case 2 can be interpreted similarly, but player 1 (without loss of generality) has more flexibility in its parameter choice due to the additional diagonal matrix D_1 in the eigenvalue decomposition of S_1.

4 Optimal Parameters and Convergence Rates

In this section, we obtain the optimal hyperparameters of MEG, for each spectrum model discussed in the previous section. Intuitively, the optimal hyperparameters are the ones that minimize the asymptotic convergence rate of MEG in (9) of Theorem 2 while satisfying the constraint for each case in Theorem 3.

Case 1: minimization. If the condition in Case 1 of Theorem 3 is satisfied (i.e., $\frac{h}{4\gamma} \geqslant (1 + \sqrt{m})^2$), then $\sigma^{-1}(-1)$ and $\sigma^{-1}(1)$ (and their intermediate values) are all on real line, forming a union of two intervals (c.f., Figure 1, left). We can write the robust region $\sigma^{-1}_{\text{Case}_1}([-1,1])$ as:

$$\left[\frac{1}{2\gamma} - \sqrt{\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma}},\ \frac{1}{2\gamma} - \sqrt{\frac{1}{4\gamma^2} - \frac{(1+\sqrt{m})^2}{h\gamma}} \right] \bigcup \left[\frac{1}{2\gamma} + \sqrt{\frac{1}{4\gamma^2} - \frac{(1+\sqrt{m})^2}{h\gamma}},\ \frac{1}{2\gamma} + \sqrt{\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma}} \right] \subset \mathbb{R}_+.$$

For this case, the optimal hyperparameters of MEG in terms of the worst-case asymptotic convergence rate in (9) can be set as below.

Theorem 4 (Case 1). *Consider solving (1) for games where the Jacobian has the spectrum in (12). For this problem, the optimal hyperparameters for the momentum extragradient method in (4) are:*

$$h = \frac{4(\mu_1 + L_2)}{(\sqrt{\mu_2 + L_1} + \sqrt{\mu_1 + L_2})^2}, \quad \gamma = \frac{1}{\mu_1 + L_2} = \frac{1}{\mu_2 + L_1}, \quad m = \left(\frac{\sqrt{\mu_2 L_1} - \sqrt{\mu_1 L_2}}{\sqrt{\mu_2 L_1} + \sqrt{\mu_1 L_2}} \right)^2 = \left(\frac{\sqrt{\rho^2 - R^2} - \sqrt{\rho^2 - 1}}{\sqrt{\rho^2 - R^2} + \sqrt{\rho^2 - 1}} \right)^2.$$

Recalling (9), we immediately get the asymptotic convergence rate from Theorem 4. Further, this formula can be simplified in the ill-conditioned regime, where the inverse condition number $\tau := \mu_1/L_2 \to 0$:

$$\sqrt[4]{m} = \left(\frac{\sqrt{\rho^2 - R^2} - \sqrt{\rho^2 - 1}}{\sqrt{\rho^2 - R^2} + \sqrt{\rho^2 - 1}} \right)^{1/2} \underset{\tau \to 0}{=} 1 - \frac{2\sqrt{\tau}}{\sqrt{1 - R^2}} + o(\sqrt{\tau}). \tag{18}$$

From (18), we see that MEG achieves an accelerated convergence rate $1 - O(\sqrt{\tau})$, which is known to be "optimal" for this function class, and can be asymptotically achieved by GDM[6] (Polyak, 1987) (see also Theorem 8 with $\theta = 1$). Surprisingly, this rate can be further improved by the factor $\sqrt{1 - R^2}$, exhibiting "super-acceleration" phenomenon enjoyed by GDM with (optimal) cyclical step sizes (Goujaud et al., 2022). Note that achieving this improvement is possible by having additional information beyond just the largest (L_2) and smallest (μ_1) eigenvalues of the Hessian.

Case 2: cross-shaped spectrum. If the condition in Case 2 of Theorem 3 is satisfied (i.e., $(1 - \sqrt{m})^2 \leqslant \frac{h}{4\gamma} < (1 + \sqrt{m})^2$), then $\sigma^{-1}(-1)$ are complex, while $\sigma^{-1}(1)$ are real (c.f., Figure 1, middle). We can write the robust region $\sigma_{\text{Case}_2}^{-1}([-1,1])$ as:

$$\underbrace{\left[\frac{1}{2\gamma} - \sqrt{\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma}}, \frac{1}{2\gamma} + \sqrt{\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma}} \right]}_{\subset \mathbb{R}_+} \bigcup \underbrace{\left[\frac{1}{2\gamma} - \sqrt{\frac{1}{4\gamma^2} - \frac{(1+\sqrt{m})^2}{h\gamma}}, \frac{1}{2\gamma} + \sqrt{\frac{1}{4\gamma^2} - \frac{(1+\sqrt{m})^2}{h\gamma}} \right]}_{\subset \mathbb{C}_+}.$$

Here, the first interval lies on $\mathbb{R}_+$, as the square root term is real; conversely, in the second interval, the square root term is imaginary, with the fixed real component: $\frac{1}{2\gamma}$. We summarize the optimal hyperparameters for this case in the next theorem.

Theorem 5 (Case 2). *Consider solving* (1) *for games where the Jacobian has a cross-shaped spectrum as in* (14). *For this problem, the optimal hyperparameters for the momentum extragradient method in* (4) *are:*

$$h = \frac{16(\mu+L)}{(\sqrt{4c^2 + (\mu+L)^2} + \sqrt{4\mu L})^2}, \quad \gamma = \frac{1}{\mu+L}, \quad and \quad m = \left(\frac{\sqrt{4c^2 + (\mu+L)^2} - \sqrt{4\mu L}}{\sqrt{4c^2 + (\mu+L)^2} + \sqrt{4\mu L}} \right)^2.$$

We get the asymptotic rate from Theorem 5, which simplifies in the ill-conditioned regime $\tau := \mu/L \to 0$ as:

$$\sqrt[4]{m} = \left(\frac{\sqrt{4c^2 + (\mu+L)^2} - \sqrt{4\mu L}}{\sqrt{4c^2 + (\mu+L)^2} + \sqrt{4\mu L}} \right)^{1/2} \underset{\tau \to 0}{=} 1 - \frac{2\sqrt{\tau}}{\sqrt{(2c/L)^2 + 1}} + o(\sqrt{\tau}). \tag{19}$$

We see that MEG achieves accelerated convergence rate $1 - O(\sqrt{\mu/L})$, as long as $c = O(L)$. We remark that this rate is optimal in the following sense. The lower bound for the problems with cross-shaped spectrum in (14) must be slower than the existing ones for minimizing μ-strongly convex and L-smooth functions, as the former is strictly more general. Since we reach the same asymptotic optimal rate, this must be optimal.

Case 3: shifted imaginary spectrum. Lastly, if the condition in Case 3 of Theorem 3 is satisfied (i.e., $\frac{h}{4\gamma} < (1 - \sqrt{m})^2$), then $\sigma^{-1}(-1)$ and $\sigma^{-1}(1)$ (and the intermediate values) are all complex conjugates (c.f., Figure 1, right). We can write the robust region $\sigma_{\text{Case}_3}^{-1}([-1,1])$ as:

$$\left[\frac{1}{2\gamma} + \sqrt{\frac{1}{4\gamma^2} - \frac{(1+\sqrt{m})^2}{h\gamma}}, \frac{1}{2\gamma} + \sqrt{\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma}} \right] \bigcup \left[\frac{1}{2\gamma} - \sqrt{\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma}}, \frac{1}{2\gamma} - \sqrt{\frac{1}{4\gamma^2} - \frac{(1+\sqrt{m})^2}{h\gamma}} \right] \subset \mathbb{C}_+.$$

We modeled such spectrum in (15), which generalizes bilinear games, where the spectrum reduces to $\pm[ai, bi]$ (i.e., with $c = 0$). We summarize the optimal hyperparameters for this case below.

---

[6]Precisely, GDM with optimal step size and momentum asymptotically achieve $1 - 2\sqrt{\tau} + o(\sqrt{\tau})$ convergence rate, as $\tau \to 0$ (Goujaud & Pedregosa, 2022, Proposition 3.3).

Theorem 6 (Case 3)**.** *Consider solving* (1) *for games where the Jacobian has a shifted imaginary spectrum in* (15)*. For this problem, the optimal hyperparameters for the momentum extragradient method in* (4) *are:*

$$h = \frac{8c}{(\sqrt{c^2+a^2}+\sqrt{c^2+b^2})^2}, \quad \gamma = \frac{1}{2c}, \quad and \quad m = \left(\frac{\sqrt{c^2+b^2}-\sqrt{c^2+a^2}}{\sqrt{c^2+b^2}+\sqrt{c^2+a^2}}\right)^2.$$

Similarly to before, we compute the asymptotic convergence rate from Theorem 4 using (9).

$$\sqrt[4]{m} = \left(\frac{\sqrt{c^2+b^2}-\sqrt{c^2+a^2}}{\sqrt{c^2+b^2}+\sqrt{c^2+a^2}}\right)^{1/2} = \left(1 - \frac{2\sqrt{c^2+a^2}}{\sqrt{c^2+b^2}+\sqrt{c^2+a^2}}\right)^{1/2} \tag{20}$$

Note that by setting $c = 0$, the rate in (20) matches the lower bound of bilinear game: $\sqrt{\frac{b-a}{b+a}}$ (Azizian et al., 2020b, Proposition 5). Further, with $c > 0$, the convergence rate in (20) improves, highlighting the contrast between vanilla bilinear games and their regularized counterpart.

Remark 3. *Notice that the optimal momentum m in both Theorems 5 and 6 are positive. This is in contrast to Gidel et al. (2019), where the* **gradient** *method with negative momentum is studied. This difference elucidates distinct dynamics of how momentum interacts with the* **gradient** *and the* **extragradient** *methods.*

5 Comparison with Other Methods

In the previous section, we obtained the asymptotic convergence rates of MEG for each spectrum model. In this section, we compare them with other first-order methods, including the gradient (GD), gradient method with momentum (GDM), and extragradient (EG) methods.

Comparison with GD and EG. In Azizian et al. (2020a), GD and EG are interpreted as fixed-point iterations, following Polyak (1987). In this framework, iterates of a method are generated by:

$$w_{t+1} = F(w_t), \quad \forall t \geqslant 0, \tag{21}$$

where $F : \mathbb{R}^d \to \mathbb{R}^d$ is an operator representing the method. It can be difficult to analyze the scheme above in general settings since F can be any nonlinear operator. However, if F is twice differentiable and w is close to a stationary point $w^\star$, the analysis can be simplified by linearizing F at the stationary point: $F(w) \approx F(w^\star) + \nabla F(w^\star)(w - w^\star)$. Then, for w_0 in a neighborhood of $w^\star$, one can obtain an asymptotic convergence rate of (21) by studying the spectral radius of the Jacobian at the solution: $\rho(\nabla F(w^\star)) \leqslant \rho^\star < 1$. This implies that (21) locally converges linearly to $w^\star$ at the rate $O((\rho^\star + \varepsilon)^t)$ for $\varepsilon \geqslant 0$. Further, if F is linear, $\varepsilon = 0$ (Polyak, 1987).

The corresponding fixed point operators F_h^{GD} and F_h^{EG} of GD and EG[7] respectively are:

$$\text{(GD)} \quad w_{t+1} = w_t - hv(w_t) = F_h^{\mathrm{GD}}(w_t), \quad \text{and} \tag{22}$$

$$\text{(EG)} \quad w_{t+1} = w_t - hv(w_t - hv(w_t)) = F_h^{\mathrm{EG}}(w_t). \tag{23}$$

The local convergence rate can then be obtained by bounding the spectral radius of the Jacobian of the operators under certain assumptions. We summarize the relevant results below.

Theorem 7 (Azizian et al. (2020a); Gidel et al. (2019))**.** *Let $w^\star$ be a stationary point of v. Further, assume the eigenvalues of $\nabla v(w^\star)$ all have positive real parts. Then, denoting $\mathcal{S}^\star := Sp(\nabla v(w^\star))$,*

1. For the gradient method in (22) *with step size $h = \min_{\lambda \in \mathcal{S}^\star} \mathfrak{R}(1/\lambda)$, it satisfies:*[8]

$$\rho(\nabla F_h^{GD}(w^\star))^2 \leqslant 1 - \min_{\lambda \in \mathcal{S}^\star} \mathfrak{R}\left(\frac{1}{\lambda}\right) \min_{\lambda \in \mathcal{S}^\star} \mathfrak{R}(\lambda). \tag{24}$$

2. For the extragradient method in (23) *with step size $h = (4\sup_{\lambda \in \mathcal{S}^\star} |\lambda|)^{-1}$, it satisfies:*

$$\rho(\nabla F_h^{EG}(w^\star))^2 \leqslant 1 - \frac{1}{4}\left(\frac{\min_{\lambda \in \mathcal{S}^\star} \mathfrak{R}(\lambda)}{\sup_{\lambda \in \mathcal{S}^\star} |\lambda|} + \frac{1}{16}\frac{\min_{\lambda \in \mathcal{S}^\star} |\lambda|^2}{\sup_{\lambda \in \mathcal{S}^\star} |\lambda|^2}\right). \tag{25}$$

---

[7]Azizian et al. (2020a) assumes that EG uses the same step size h for both the main and the extrapolation steps.
[8]Note that the spectral radius ρ is squared, but asymptotically is almost the same as $\sqrt{1-x} \leqslant 1 - x/2$.

We can determine the convergence rate of GD and EG by using Theorem 7 since all three cases of our spectrum models in (12), (14), and (15) meet the condition that the eigenvalues of $\nabla v(w^\star)$ have positive real parts. The following corollary summarizes this result.

Corollary 1. *With the conditions in Theorem 7, for each case of the Jacobian spectrum $\mathcal{S}_1^\star$, $\mathcal{S}_2^\star$, and $\mathcal{S}_3^\star$, the gradient method in (22) and the extragradient method in (23) satisfy the following:*

- **Case 1:** $Sp(\nabla v) \subset \mathcal{S}_1^\star = [\mu_1, L_1] \cup [\mu_2, L_2] \in \mathbb{R}_+$:

$$\rho(\nabla F_h^{GD}(w^\star))^2 \leqslant 1 - \tfrac{\mu_1}{L_2}, \quad and \quad \rho(\nabla F_h^{EG}(w^\star))^2 \leqslant 1 - \tfrac{1}{4}\left(\tfrac{\mu_1}{L_2} + \tfrac{\mu_1^2}{16L_2^2}\right). \tag{26}$$

- **Case 2:** $Sp(\nabla v) \subset \mathcal{S}_2^\star = [\mu, L] \cup \{z \in \mathbb{C} : \mathfrak{R}(z) = c' > 0, \ \mathfrak{I}(z) \in [-c, c]\}$:

$$\rho(\nabla F_h^{GD}(w^\star))^2 \leqslant \begin{cases} 1 - \frac{2\mu}{4c^2/(L-\mu)+(L-\mu)} & if \ c \geqslant \sqrt{\frac{L^2-\mu^2}{4}}, \\ 1 - \frac{\mu}{L} & otherwise. \end{cases} \tag{27}$$

$$\rho(\nabla F_h^{EG}(w^\star))^2 \leqslant \begin{cases} 1 - \frac{1}{4}\left(\frac{\mu}{\sqrt{c^2+((L-\mu)/2)^2}} + \frac{\mu^2}{16(c^2+((L-\mu)/2)^2)}\right) & if \ c \geqslant \sqrt{\frac{3L^2+2L\mu-\mu^2}{4}}, \\ 1 - \frac{1}{4}\left(\frac{\mu}{L} + \frac{\mu^2}{16L^2}\right) & otherwise. \end{cases}$$

- **Case 3:** $Sp(\nabla v) \subset \mathcal{S}_3^\star = [c+ai, c+bi] \cup [c-ai, c-bi] \in \mathbb{C}_+$:

$$\rho(\nabla F_h^{GD}(w^\star))^2 \leqslant 1 - \tfrac{c^2}{c^2+b^2}, \quad and \quad \rho(\nabla F_h^{EG}(w^\star))^2 \leqslant 1 - \tfrac{1}{4}\left(\tfrac{c}{\sqrt{c^2+b^2}} + \tfrac{c^2+a^2}{16(c^2+b^2)}\right). \tag{28}$$

In Case 1, we see from (26) that both GD and EG have convergence rates $1 - O(\mu_1/L_2) = 1 - O(\tau)$. MEG, on the other hand, has an accelerated convergence rate of $1 - O(\sqrt{\tau})$, as well as an additional constant improvement by a factor of $\sqrt{1 - R^2}$, as we showed in (18). Moving on to Case 2, we showed in (19) that MEG enjoys an accelerated convergence rate of $1 - O(\sqrt{\mu/L})$ as long as $c = O(L)$. However, both GD and EG in (27) have non-accelerated convergence under the same condition. Lastly, for Case 3, we showed in (20) that MEG achieves an asymptotic rate that matches the known lower bound for bilinear games: $\sqrt{\frac{b-a}{b+a}}$, with $c = 0$; further, the rate of MEG improves if $c > 0$. On the contrary, GD and EG suffer from slower rates, as shown in (28).

Comparison with GDM. We now compare the convergence rate of MEG with that of GDM, which iterates as in (5). In Azizian et al. (2020b), it was shown that GD is the optimal method for games where the Jacobian eigenvalues are within a *disc* in the complex plane. This suggests that acceleration is not possible for this type of problem.[9] On the other hand, it is well-known that GDM achieves an accelerated convergence rate for strongly-convex (quadratic) minimization, where the eigenvalues of the Hessian lie on the (strictly positive) real line segment (Polyak, 1987). Hence, Azizian et al. (2020b) studies the intermediate case, where the Jacobian eigenvalues are within an ellipse, which can be thought of as the real segment $[\mu, L]$ perturbed with ϵ in an elliptic way. That is, they consider the spectral shape:[10]

$$K_\epsilon = \left\{ z \in \mathbb{C} : \left(\tfrac{\mathfrak{R}z - (\mu+L)/2}{(L-\mu)/2}\right)^2 + \left(\tfrac{\mathfrak{I}z}{\epsilon}\right)^2 \leqslant 1 \right\}.$$

Similarly to GD and EG above, in Azizian et al. (2020b), GDM is interpreted as a fixed point iteration:[11]

$$w_{t+1} = w_t - hv(w_t) + m(w_t - w_{t-1}) = F^{\text{GDM}}(w_t, w_{t-1}). \tag{29}$$

To study the convergence rate of GDM, we use the following theorem from Azizian et al. (2020b):

---

[9]Yet, one can consider the case where, e.g., a cross-shape is contained in a disc. Then, by knowing more fine-grained structure of the Jacobian spectrum, MEG can have faster convergence in (19).

[10]A visual illustration of this ellipse can be found in Azizian et al. (2020b, Figure 2).

[11]As GDM updates w_{t+1} using both w_t and w_{t-1}, Azizian et al. (2020b) uses an augmented fixed point operator; see Lemma 2 in that work for details.

Theorem 8 (Azizian et al. (2020b)). *Define $\epsilon(\mu, L)$ as $\epsilon(\mu, L)/L = (\mu/L)^\theta = \tau^\theta$ with $\theta > 0$ and $a \wedge b = \min(a, b)$. If $Sp(\nabla F^{GDM}(w^\star, w^\star)) \subset K_\epsilon$, and when $\tau \to 0$, it satisfies:*

$$\rho(\nabla F^{GDM}(w^\star, w^\star)) \leqslant \begin{cases} 1 - 2\sqrt{\tau} + O\left(\tau^{\theta \wedge 1}\right), & \text{if } \theta > \frac{1}{2} \\ 1 - 2(\sqrt{2} - 1)\sqrt{\tau} + O\left(\tau\right), & \text{if } \theta = \frac{1}{2} \\ 1 - \tau^{1-\theta} + O\left(\tau^{1 \wedge (2-3\theta)}\right), & \text{if } \theta < \frac{1}{2}, \end{cases} \tag{30}$$

where the hyperparametes h and m are functions of μ, L, and ϵ only.

For Case 1, GDM converges at the rate $1 - 2\sqrt{\tau} + O(\tau)$ (i.e., with $\theta = 1$ from the above), which is always slower than the rate of MEG in (18) by the factor of $\sqrt{1 - R^2}$. For Case 2, we see from Theorem 8 that GDM achieves an accelerated rate, i.e., $1 - O(\sqrt{\tau})$, until $\theta = \frac{1}{2}$. In other words, the biggest elliptic perturbation ϵ where GDM permits the accelerated rate is $\epsilon = \sqrt{\mu L}$.[12] We interpret Theorem 8 for games with cross-shaped Jacobian spectrum in (14) and shifted imaginary spectrum in (15) in the following corollary.

Corollary 2. *Consider the gradient method with momentum, interpreted as fixed point iteration as in (29). For games with cross-shaped Jacobian spectrum in (14) with $c = \frac{L-\mu}{2}$, GDM cannot achieve an accelerated rate when $\frac{L-\mu}{2} = c > \epsilon = \sqrt{\mu L}$. Since $L > \mu$, this further implies $\frac{L}{\mu} > \sqrt{5}$. That is, when the condition number exceeds $\sqrt{5} \approx 2.236$, GDM cannot achieve an accelerated convergence rate. On the contrary, as we showed in (19), MEG can converge at an accelerated rate in the ill-conditioned regime.*

The convergence rate of GDM for Case 3 cannot be determined from Theorem 8, as this theorem assumes the spectrum model of the real line segment $[\mu, L]$ with ϵ perturbation (along the imaginary axis), while $\mathcal{S}_3^\star$ in (15) has a fixed real component. Instead, we utilize the link function of GDM in (7) to show that it is unlikely for GDM to stay in the robust region: $\xi^{-1}([-1, 1])$.

Proposition 2. *Consider solving (1) for games where the Jacobian has a shifted imaginary spectrum in (15), using the gradient method with momentum in (5). For any complex number $z = p + qi \in \mathbb{C}_+$, if $\frac{2(1+m)}{h} < p$, then GDM cannot stay in the robust region, i.e., $|\xi(\lambda)| > 1$.*

Note that the condition $\frac{2(1+m)}{h} < p$ is hard to avoid even for small p, considering h is usually a small value.

6 Local Convergence for Non-affine Vector Fields

The optimal hyperparameters of MEG for each spectrum model and the associated convergence rate we obtained in Section 4 are attainable when the vector field is affine. A natural question is, then, what can we say about the convergence rate of MEG when the vector field is not affine? To that end, we provide the local convergence of MEG by restarting the momentum, as detailed below.

Let us consider the operator G representing the MEG in (4) such that:

$$[w_{t+1}, w_t] = G([w_t, w_{t-1}]) \quad \text{and} \quad G([w^\star, w^\star]) = [w^\star, w^\star].$$

In addition, we assume that $w_1 = w_0 - \frac{h}{1+m} v(w_0 - \gamma v(w_0))$, in order to induce the residual polynomials from Theorem 1; see also its proof and Algorithm 1 in the appendix. Now let us consider the following algorithm:

$$[w_{tk+i+1}, w_{tk+i}] = G([w_{tk+i}, w_{tk+i-1}]) \quad \text{for } 1 \leqslant i \leqslant k - 1, \quad \text{and then}$$
$$w_{(t+1)k+1} = w_{(t+1)k} - \frac{h}{1+m} v(w_{(t+1)k} - \gamma v(w_{(t+1)k})). \tag{31}$$

In other words, we repeat MEG for k steps, and then restart the momentum at $[w_{(t+1)k+1}, w_{(t+1)k}]$. The local convergence of the restarted MEG is established in the next theorem.

Theorem 9 (Local convergence). *Let $G : \mathbb{R}^{2d} \to \mathbb{R}^{2d}$ be the continuously differentiable operator representing the momentum extragradient method (MEG) in (4). Let $w^\star$ be a stationary point. Let w_t denote the output of MEG, which enjoys a convergence rate of the form $\|w_t - w^\star\| = C(1 - \rho)^t(t + 1)\|w_0 - w^\star\|$ for some*

---

[12]Observe that $(\mu/L)^{1/2} = \epsilon(\mu, L)/L \implies \epsilon(\mu, L) = \sqrt{\mu L}$.

$0 < \rho < 1$ *when the vector field is affine. Further, consider restarting the momentum of MEG after running* k *steps, as in* (31). *Then, for each* $\varepsilon > 0$, *there exists* $k > 0$ *and* $\delta > 0$ *such that, for all initializations* w_0 *satisfying* $\|w_0 - w^\star\| \leqslant \delta$, *the restarted MEG satisfies:*

$$\|w_t - w^\star\| = O((1 - \rho + \varepsilon)^t)\|w_0 - w^\star\|.$$

7 Experiments

Figure 3: *Illustration of the game Jacobian spectra and the performance of different algorithms considered.* Jacobian spectrum in the first plot matches $\mathcal{S}_2^\star$ in (14) precisely, while that in the third plot inexactly follows $\mathcal{S}_2^\star$. The second (fourth) plot shows the performance of different algorithms for solving quadratic games in (16) with the Jacobian spectrum following the first (third) plot.

In this section, we perform numerical experiments to optimize a game when the Jacobian has a cross-shaped spectrum in (14). We focus on this spectrum as it may be the most challenging case, involving both real and complex eigenvalues (c.f., Theorem 3). To test the robustness, we consider two cases where the Jacobian spectrum exactly follows $\mathcal{S}_2^\star$ in (14), as well as the inexact case. We illustrate them in Figure 3.

We focus on two-player quadratic games, where player 1 controls $x \in \mathbb{R}^{d_1}$ and player 2 controls $y \in \mathbb{R}^{d_2}$ with loss functions in (16). In our setting, the corresponding vector field in (17) satisfies $M_{12} = -M_{21}^\top$, but S_1 and S_2 can be nonzero symmetric matrices. Further, the Jacobian $\nabla v = A$ has the cross-shaped eigenvalue structure in (14), with $c = \frac{L-\mu}{2}$ (c.f., Proposition 1, Case 2). For the problem constants, we use $\mu = 1$, and $L = 200$. The optimum $[x^\star \; y^\star]^\top = w^\star \in \mathbb{R}^{200}$ is generated using the standard normal distribution. For simplicity, we assume $b = [b_1 \; b_2]^\top = [0 \; 0]^\top$. For the algorithms, we compare GD in (22), GDM in (5), EG in (23), and MEG in (4). All algorithms are initialized with 0. We plot the experimental results in Figure 3.

For MEG (optimal), we set the hyperparameters using Theorem 5. For GD (theory) and EG (theory), we set the hyperparameters using Theorem 7, both for the exact and the inexact settings. For GDM (grid search), we perform a grid search of h^{GDM} and m^{GDM}, and choose the best-performing ones, as Theorem 8 does not give a specific form for hyperparameter setup. Specifically, we consider $0.005 \leqslant h^{\text{GDM}} \leqslant 0.015$ with 10^{-3} increment, and $0.01 \leqslant m^{\text{GDM}} \leqslant 0.99$ with 10^{-2} increment. In addition, as Theorem 7 might be conservative, we conduct grid searches for GD and EG as well. For GD (grid search), we use the same setup as h^{GDM}. For EG (grid search), we use $0.001 \leqslant h^{\text{EG}} \leqslant 0.05$ with 10^{-4} increment.

There are several remarks to make. First, although the third plot in Figure 3 does not exactly follow the spectrum model in (14), MEG still works well with the optimal hyperparameters from Theorem 5. As expected, MEG (optimal) required more iterations in the inexact case compared to the exact case. Second, compared to other algorithms, MEG (optimal) indeed exhibits a significantly faster rate of convergence, even when compared to other methods that use grid-search hyperparameter tuning, supporting our theoretical findings in Section 4. Third, while EG (theory) is slower than GD (theory), which confirms Corrolary 1, EG (grid search) can be tuned to converge faster via grid search. Lastly, even though the best performance of GDM (grid search) is obtained through grid search, one can see the GD (grid search) obtains a slightly faster convergence rate than GDM (grid search), confirming Corollay 2.

8 Conclusion

In this work, we focused on the momentum extragradient method for finding a stationary point of differentiable games. Via polynomial-based analysis, we showed the momentum extragradient method converges in three different modes, depending on the Jacobian eigenvalue distribution. Then, for each case, we obtained the optimal hyperparameters for the momentum extragradient method, which enjoys accelerated asymptotic convergence rates. We compared the obtained rates with other first-order methods and showed that the considered methods do not achieve the accelerated convergence rate. Lastly, we validated our theoretical findings through numerical experiments on quadratic games.

Acknowledgments

The authors would like to thank Fangshuo Liao, Baptiste Goujaud, Damien Scieur, Miri Son, and Giorgio Young for their useful discussions and feedback.

This work is supported by NSF FET: Small No. 1907936, NSF MLWiNS CNS No. 2003137 (in collaboration with Intel), NSF CMMI No. 2037545, NSF CAREER award No. 2145629, NSF CIF No. 2008555, Rice InterDisciplinary Excellence Award (IDEA), and the Canada CIFAR AI Chairs program.

References

Yossi Arjevani and Ohad Shamir. On the iteration complexity of oblivious first-order optimization algorithms. In *International Conference on Machine Learning*, pp. 908–916. PMLR, 2016.

Waïss Azizian, Ioannis Mitliagkas, Simon Lacoste-Julien, and Gauthier Gidel. A tight and unified analysis of gradient-based methods for a whole spectrum of differentiable games. In *International Conference on Artificial Intelligence and Statistics*, pp. 2863–2873. PMLR, 2020a.

Waïss Azizian, Damien Scieur, Ioannis Mitliagkas, Simon Lacoste-Julien, and Gauthier Gidel. Accelerating smooth games by manipulating spectral shapes. In *International Conference on Artificial Intelligence and Statistics*, pp. 1705–1715. PMLR, 2020b.

David Balduzzi, Sebastien Racaniere, James Martens, Jakob Foerster, Karl Tuyls, and Thore Graepel. The mechanics of n-player differentiable games. In *International Conference on Machine Learning*, pp. 354–363. PMLR, 2018.

Hugo Berard, Gauthier Gidel, Amjad Almahairi, Pascal Vincent, and Simon Lacoste-Julien. A closer look at the optimization landscapes of generative adversarial networks. In *International Conference on Learning Representations*, 2020. URL https://openreview.net/forum?id=HJeVnCEKwH.

Raphaël Berthier, Francis Bach, and Pierre Gaillard. Accelerated gossip in networks of given dimension using jacobi polynomial iterations. *SIAM Journal on Mathematics of Data Science*, 2(1):24–47, 2020.

Aleksandr Beznosikov, Pavel Dvurechensky, Anastasia Koloskova, Valentin Samokhin, Sebastian U Stich, and Alexander Gasnikov. Decentralized local stochastic extra-gradient for variational inequalities. *arXiv preprint arXiv:2106.08315*, 2021.

Theodore S Chihara. *An introduction to orthogonal polynomials.* Courier Corporation, 2011.

Constantinos Daskalakis and Ioannis Panageas. The limit points of (optimistic) gradient descent in min-max optimization. *Advances in neural information processing systems*, 31, 2018.

Constantinos Daskalakis, Paul W Goldberg, and Christos H Papadimitriou. The complexity of computing a nash equilibrium. *SIAM Journal on Computing*, 39(1):195–259, 2009.

Carles Domingo-Enrich, Fabian Pedregosa, and Damien Scieur. Average-case acceleration for bilinear games and normal matrices. In *International Conference on Learning Representations*, 2021. URL https://openreview.net/forum?id=H0syOoy3Ash.

Pierre Foret, Ariel Kleiner, Hossein Mobahi, and Behnam Neyshabur. Sharpness-aware minimization for efficiently improving generalization. In *International Conference on Learning Representations*, 2021. URL https://openreview.net/forum?id=6Tm1mposlrM.

Gauthier Gidel, Hugo Berard, Gaëtan Vignoud, Pascal Vincent, and Simon Lacoste-Julien. A variational inequality perspective on generative adversarial networks. *arXiv preprint arXiv:1802.10551*, 2018.

Gauthier Gidel, Reyhane Askari Hemmat, Mohammad Pezeshki, Rémi Le Priol, Gabriel Huang, Simon Lacoste-Julien, and Ioannis Mitliagkas. Negative momentum for improved game dynamics. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pp. 1802–1811. PMLR, 2019.

Ian Goodfellow, Jean Pouget-Abadie, Mehdi Mirza, Bing Xu, David Warde-Farley, Sherjil Ozair, Aaron Courville, and Yoshua Bengio. Generative adversarial nets. In Z. Ghahramani, M. Welling, C. Cortes, N. Lawrence, and K.Q. Weinberger (eds.), *Advances in Neural Information Processing Systems*, volume 27. Curran Associates, Inc., 2014. URL https://proceedings.neurips.cc/paper/2014/file/5ca3e9b122f61f8f06494c97b1afccf3-Paper.pdf.

Eduard Gorbunov, Nicolas Loizou, and Gauthier Gidel. Extragradient method: O (1/k) last-iterate convergence for monotone variational inequalities and connections with cocoercivity. In *International Conference on Artificial Intelligence and Statistics*, pp. 366–402. PMLR, 2022.

Eduard Gorbunov, Adrien Taylor, Samuel Horváth, and Gauthier Gidel. Convergence of proximal point and extragradient-based methods beyond monotonicity: the case of negative comonotonicity. In *International Conference on Machine Learning*, pp. 11614–11641. PMLR, 2023.

Baptiste Goujaud and Fabian Pedregosa. Cyclical step-sizes, 2022. URL http://fa.bianp.net/blog/2022/cyclical/.

Baptiste Goujaud, Damien Scieur, Aymeric Dieuleveut, Adrien B Taylor, and Fabian Pedregosa. Super-acceleration with cyclical step-sizes. In *International Conference on Artificial Intelligence and Statistics*, pp. 3028–3065. PMLR, 2022.

Magnus R Hestenes and Eduard Stiefel. Methods of conjugate gradients for solving. *Journal of research of the National Bureau of Standards*, 49(6):409, 1952.

Andrew J Hetzel, Jay S Liew, and Kent E Morrison. The probability that a matrix of integers is diagonalizable. *The American Mathematical Monthly*, 114(6):491–499, 2007.

Yu-Guan Hsieh, Franck Iutzeler, Jérôme Malick, and Panayotis Mertikopoulos. On the convergence of single-call stochastic extra-gradient methods. *Advances in Neural Information Processing Systems*, 32, 2019.

Yu-Guan Hsieh, Franck Iutzeler, Jérôme Malick, and Panayotis Mertikopoulos. Explore aggressively, update conservatively: Stochastic extragradient methods with variable stepsize scaling. *Advances in Neural Information Processing Systems*, 33:16223–16234, 2020.

Junhyung Lyle Kim, Gauthier Gidel, Anastasios Kyrillidis, and Fabian Pedregosa. Momentum extragradient is optimal for games with cross-shaped spectrum. In *OPT 2022: Optimization for Machine Learning (NeurIPS 2022 Workshop)*, 2022.

Galina M Korpelevich. The extragradient method for finding saddle points and other problems. *Matecon*, 12:747–756, 1976.

Peter Lancaster and Hanafi K Farahat. Norms on direct sums and tensor products. *mathematics of computation*, 26(118):401–414, 1972.

Alistair Letcher, David Balduzzi, Sébastien Racaniere, James Martens, Jakob Foerster, Karl Tuyls, and Thore Graepel. Differentiable game mechanics. *The Journal of Machine Learning Research*, 20(1):3032–3071, 2019.

Chris Junchi Li, Yaodong Yu, Nicolas Loizou, Gauthier Gidel, Yi Ma, Nicolas Le Roux, and Michael I Jordan. On the convergence of stochastic extragradient for bilinear games with restarted iteration averaging. *arXiv preprint arXiv:2107.00464*, 2021.

Tengyuan Liang and James Stokes. Interaction matters: A note on non-asymptotic local convergence of generative adversarial networks. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pp. 907–915. PMLR, 2019.

Mingrui Liu, Wei Zhang, Youssef Mroueh, Xiaodong Cui, Jarret Ross, Tianbao Yang, and Payel Das. A decentralized parallel algorithm for training generative adversarial nets. *Advances in Neural Information Processing Systems*, 33:11056–11070, 2020.

Jonathan P. Lorraine, David Acuna, Paul Vicol, and David Duvenaud. Complex momentum for optimization in games. In Gustau Camps-Valls, Francisco J. R. Ruiz, and Isabel Valera (eds.), *Proceedings of The 25th International Conference on Artificial Intelligence and Statistics*, volume 151 of *Proceedings of Machine Learning Research*, pp. 7742–7765. PMLR, 28–30 Mar 2022.

Lars Mescheder, Andreas Geiger, and Sebastian Nowozin. Which training methods for gans do actually converge? In *International conference on machine learning*, pp. 3481–3490. PMLR, 2018.

Aryan Mokhtari, Asuman Ozdaglar, and Sarath Pattathil. A unified analysis of extra-gradient and optimistic gradient methods for saddle point problems: Proximal point approach. In *International Conference on Artificial Intelligence and Statistics*, pp. 1497–1507. PMLR, 2020.

Renato DC Monteiro and Benar Fux Svaiter. On the complexity of the hybrid proximal extragradient method for the iterates and the ergodic mean. *SIAM Journal on Optimization*, 20(6):2755–2787, 2010.

Arkadi Nemirovski. Prox-method with rate of convergence o (1/t) for variational inequalities with lipschitz continuous monotone operators and smooth convex-concave saddle point problems. *SIAM Journal on Optimization*, 15(1):229–251, 2004.

Arkadij Semenovič Nemirovskij and David Borisovich Yudin. Problem complexity and method efficiency in optimization. 1983.

Samet Oymak. Provable super-convergence with a large cyclical learning rate. *IEEE Signal Processing Letters*, 28:1645–1649, 2021.

Balamurugan Palaniappan and Francis Bach. Stochastic variance reduction methods for saddle-point problems. *Advances in Neural Information Processing Systems*, 29, 2016.

Vardan Papyan. Traces of class/cross-class structure pervade deep learning spectra. *The Journal of Machine Learning Research*, 21(1):10197–10260, 2020.

Fabian Pedregosa. Momentum: when chebyshev meets chebyshev, 2020. URL http://fa.bianp.net/blog/2020/momentum/.

Fabian Pedregosa and Damien Scieur. Acceleration through spectral density estimation. In *Proceedings of the 37th International Conference on Machine Learning*, pp. 7553–7562. PMLR, November 2020. URL https://proceedings.mlr.press/v119/pedregosa20a.html. ISSN: 2640-3498.

David Pfau and Oriol Vinyals. Connecting generative adversarial networks and actor-critic methods. *arXiv preprint arXiv:1610.01945*, 2016.

Boris T Polyak. Introduction to optimization. optimization software. *Inc., Publications Division, New York*, 1:32, 1987.

Ernest K Ryu, Kun Yuan, and Wotao Yin. Ode analysis of stochastic gradient methods with optimism and anchoring for minimax problems. *arXiv preprint arXiv:1905.10899*, 2019.

Yoav Shoham and Kevin Leyton-Brown. *Multiagent systems: Algorithmic, game-theoretic, and logical foundations.* Cambridge University Press, 2008.

Mikhail V Solodov and Benar F Svaiter. A hybrid approximate extragradient–proximal point algorithm using the enlargement of a maximal monotone operator. *Set-Valued Analysis*, 7(4):323–345, 1999.

Paul Tseng. On linear convergence of iterative methods for the variational inequality problem. *Journal of Computational and Applied Mathematics*, 60(1-2):237–252, 1995.

Jian Zhang and Ioannis Mitliagkas. Yellowfin and the art of momentum tuning. *Proceedings of Machine Learning and Systems*, 1:289–308, 2019.

A   Missing Proofs in Section 3

A.1   Proof of Lemma 1

Proof of Lemma 1 can be found for example in Azizian et al. (2020b, Section B).

A.2   Proof of Theorem 1

To obtain the residual polynomials of MEG, w_1 has to be set slightly differently from the rest of the iterates, as we write in the pseudocode below:

Algorithm 1: Momentum extragradient (MEG) method

Input: Initialization w_0, hyperparameters h, γ, m.
Set: $w_1 = w_0 - \frac{h}{1+m} v(w_0 - \gamma v(w_0))$
for $t = 1, 2, \ldots$ **do**
$\quad | \quad w_{t+1} = w_t - hv(w_t - \gamma v(w_t)) + m(w_t - w_{t-1})$
end

Derivation of the first part

Proof. We want to find the residual polynomial $\tilde{P}_t(A)$ of the extragradient with momentum (MEG) in (4). That is, we want to find

$$w_t - w^\star = \tilde{P}_t(A)(w_0 - w^\star), \tag{32}$$

where $\{w_t\}_{t=0}$ is the iterates generated by MEG, which is possible by Lemma 1, as MEG is a first-order method (Arjevani & Shamir, 2016; Azizian et al., 2020b). We now prove this is by induction. To do so, we will use the following properties. First, note that as we are looking for a stationary point, it holds that $v(w^\star) = 0$. Further, as v is linear by the assumption of Lemma 1, it holds that $v(w) = A(w - w^\star)$.

Base case. For $t = 0$, $\tilde{P}_0(A)$ is a degree-zero polynomial, and hence equals I_d, which denotes the identity matrix. Thus, $w_0 - w^\star = I_d(w_0 - w^\star)$ holds true.

For completeness, we also prove when $t = 1$. In that case, observe that MEG in proceeds as $w_1 = w_0 - \frac{h}{1+m} v(w_0 - \gamma v(w_0))$. Subtracting $w^\star$ on both sides, we have:

$$
\begin{aligned}
w_1 - w^\star &= w_0 - w^\star - \frac{h}{1+m} v(w_0 - \gamma v(w_0)) \\
&= w_0 - w^\star - \frac{h}{1+m} v(w_0 - \gamma A(w_0 - w^\star)) \\
&= w_0 - w^\star - \frac{h}{1+m} A(w_0 - \gamma A(w_0 - w^\star) - w^\star) \\
&= w_0 - w^\star - \frac{h}{1+m} A(w_0 - w^\star) + \frac{h\gamma}{1+m} A^2(w_0 - w^\star) \\
&= \left(I_d - \frac{h}{1+m} A + \frac{h\gamma}{1+m} A^2 \right)(w_0 - w^\star) \\
&= \left(I_d - \frac{h}{1+m} A(I_d - \gamma A) \right)(w_0 - w^\star) \\
&= \tilde{P}_1(A)(w_0 - w^\star).
\end{aligned}
$$

Induction step. As the induction hypothesis, assume $\tilde{P}_t$ satisfies (32). We want to prove this holds for $t + 1$. We have:

$$
\begin{aligned}
w_{t+1} &= w_t - hv(w_t - \gamma v(w_t)) + m(w_t - w_{t-1}) \\
&= w_t - hv(w_t - \gamma A(w_t - w^\star)) + m(w_t - w_{t-1}) \\
&= w_t - hA(w_t - \gamma A(w_t - w^\star) - w^\star) + m(w_t - w_{t-1}) \\
&= w_t - hA(w_t - w^\star) + h\gamma A^2(w_t - w^\star) + m(w_t - w_{t-1}) \\
&= w_t - hA(I_d - \gamma A)(w_t - w^\star) + m(w_t - w_{t-1}).
\end{aligned}
$$

Subtracting $w^\star$ on both sides, we have:

$$
\begin{aligned}
w_{t+1} - w^\star &= w_t - w^\star - hA(I_d - \gamma A)(w_t - w^\star) + m(w_t - w_{t-1}) \\
&= (I_d - hA(I_d - \gamma A))(w_t - w^\star) + m(w_t - w^\star - (w_{t-1} - w^\star)) \\
&\stackrel{(32)}{=} (I_d - hA(I_d - \gamma A))\tilde{P}_t(A)(w_0 - w^\star) + m(\tilde{P}_t(A)(w_0 - w^\star) - \tilde{P}_{t-1}(A)(w_0 - w^\star)) \\
&= (I_d + mI_d - hA(I_d - \gamma A))\tilde{P}_t(A)(w_0 - w^\star) - m\tilde{P}_{t-1}(A)(w_0 - w^\star) \\
&= \tilde{P}_{t+1}(A)(w_0 - w^\star),
\end{aligned}
$$

where in the third equality, we used the induction hypothesis in (32).

$\square$

Derivation of the second part in (6)

Proof. We show $P_t = \tilde{P}_t$ for all t via induction.

Base case. For $t = 0$, by the definition of Chebyshev polynomials of the first and the second kinds, we have $T_0(\lambda) = U_0(\lambda) = 1$. Thus,

$$
\begin{aligned}
P_0(\lambda) &= m^0 \left(\frac{2m}{1+m} T_0(\sigma(\lambda)) + \frac{1-m}{1+m} U_0(\sigma(\lambda)) \right) \\
&= \frac{2m}{1+m} + \frac{1-m}{1+m} = 1 = \tilde{P}_0(\lambda).
\end{aligned}
$$

Again, for completeness, we prove when $t = 1$ as well. In that case, by the definition of Chebyshev polynomials of the first and the second kinds, we have $T_1(\lambda) = \lambda$, and $U_1(\lambda) = 2\lambda$. Therefore,

$$
\begin{aligned}
P_1(\lambda) &= m^{t/2} \left(\frac{2m}{1+m} T_1(\sigma(\lambda)) + \frac{1-m}{1+m} U_1(\sigma(\lambda)) \right) \\
&= m^{t/2} \left(\frac{2m}{1+m} \sigma(\lambda) + \frac{1-m}{1+m} \cdot 2 \cdot \sigma(\lambda) \right) \\
&= m^{t/2} \left(\frac{2\sigma(\lambda)}{1+m} \right) \\
&= 1 - \frac{h\lambda(1 - \gamma\lambda)}{1+m} = \tilde{P}_1(\lambda).
\end{aligned}
$$

Induction step. As the induction hypothesis, assume that $P_t = \tilde{P}_t$ for t. In this step, we show that the same holds for $t+1$.

$$
\begin{aligned}
P_{t+1} &= m^{(t+1)/2}\left[\frac{2m}{1+m}T_{t+1}(\sigma(\lambda)) + \frac{1-m}{1+m}U_{t+1}(\sigma(\lambda))\right] \\
&= m^{(t+1)/2}\left[\frac{2m}{1+m}\Big(2\sigma(\lambda)T_t(\sigma(\lambda)) - T_{t-1}(\sigma(\lambda))\Big) \right.\\
&\qquad\qquad \left. + \frac{1-m}{1+m}\Big(2\sigma(\lambda)U_t(\sigma(\lambda) - U_{t-1}(\sigma(\lambda)))\Big)\right] \\
&= 2\sigma(\lambda)\cdot m^{1/2}\cdot \underbrace{m^{t/2}\left(\frac{2m}{1+m}T_t(\sigma(\lambda)) + \frac{1-m}{1+m}U_t(\sigma(\lambda))\right)}_{P_t(\lambda)} \\
&\qquad - m\cdot \underbrace{m^{(t-1)/2}\left(\frac{2m}{1+m}T_{t-1}(\sigma(\lambda)) + \frac{1-m}{1+m}U_{t-1}(\sigma(\lambda))\right)}_{P_{t-1}(\lambda)} \\
&= 2\sigma(\lambda)\cdot\sqrt{m}\cdot\tilde{P}_t(\lambda) - m\cdot\tilde{P}_{t-1}(\lambda) \\
&= (1 + m - h\lambda(1-\gamma\lambda))\tilde{P}_t(\lambda) - m\tilde{P}_{t-1}(\lambda),
\end{aligned}
$$

where in the second to last equality we use the induction hypothesis. $\qquad\square$

A.3 Proof of Lemma 2

Proof of Lemma 2 can be found in Goujaud & Pedregosa (2022).

A.4 Proof of Theorem 2

Proof. We first recall that using (3), we can upper bound the worst-case convergence rate as:

$$
\begin{aligned}
\sup_{\lambda\in\mathcal{S}^\star}|P_t(\lambda)| &\overset{(6)}{=} \sup_{\lambda\in\mathcal{S}^\star}\left|m^{t/2}\left(\tfrac{2m}{1+m}T_t(\sigma(\lambda)) + \tfrac{1-m}{1+m}U_t(\sigma(\lambda))\right)\right| \\
&\leqslant m^{t/2}\left(\tfrac{2m}{1+m}\sup_{\lambda\in\mathcal{S}^\star}|T_t(\sigma(\lambda))| + \tfrac{1-m}{1+m}\sup_{\lambda\in\mathcal{S}^\star}|U_t(\sigma(\lambda))|\right)
\end{aligned}
\tag{33}
$$

Now, denote $\bar{\sigma} := \sup_{\lambda\in\mathcal{S}^\star}|\sigma(\lambda;h,\gamma,m)|$. For the first case, if $\bar{\sigma} \leqslant 1$, both $T_t(x)$ and $U_t(x)$ behave nicely, per Lemma 2. Thus, we have

$$
(33) \overset{(8)}{\leqslant} m^{t/2}\left(\frac{2m}{1+m} + \frac{1-m}{1+m}(t+1)\right) \leqslant m^{t/2}(t+1) \implies \limsup_{t\to\infty}\left(m^{t/2}(t+1)\right)^{\frac{1}{2t}} = \sqrt[4]{m}.
$$

For the second case, we use the following expressions of Chebyshev polynomials:

$$
\begin{aligned}
T_n(x) &= \frac{\left(x - \sqrt{x^2-1}\right)^n + \left(x + \sqrt{x^2-1}\right)^n}{2}, \quad\text{and} \\
U_n(x) &= \frac{\left(x + \sqrt{x^2-1}\right)^{n+1} - \left(x - \sqrt{x^2-1}\right)^{n+1}}{2\sqrt{x^2-1}}.
\end{aligned}
$$

Therefore, in the second case where $\bar{\sigma} > 1$, we have both $T_n(x)$ and $U_n(x)$ growing at rate $(x + \sqrt{x^2-1})^n$. Hence, we have:

$$
(33) \leqslant O\left(m^{t/2}\left(\bar{\sigma} + \sqrt{\bar{\sigma}^2-1}\right)^t\right) \implies \limsup_{t\to\infty}\left(m^{t/2}\left(\bar{\sigma} + \sqrt{\bar{\sigma}^2-1}\right)^t\right)^{\frac{1}{2t}} = \sqrt[4]{m}\left(\bar{\sigma} + \sqrt{\bar{\sigma}^2-1}\right)^{1/2}.
$$

Finally, in order for MEG to converge in the second case, we need:

$$\sqrt[4]{m}\left(\bar{\sigma} + \sqrt{\bar{\sigma}^2 - 1}\right)^{1/2} < 1$$

which is equivalent to

$$\bar{\sigma} \leqslant \frac{\sqrt{m}(m+1)}{2m} = \frac{m+1}{2\sqrt{m}}.$$

$\square$

A.5 Derivation of extreme points of robust region in (11)

We first write a general formula for inverting a quadratic function. For $f(x) = ax^2 + bx + c$, its inverse is given by:

$$f(x) = ax^2 + bx + c := y$$
$$f^{-1}(y) = \frac{-b \pm \sqrt{b^2 - 4a(c - y)}}{2a},$$

with some abuse of notation (i.e., f^{-1} above is not a function).

Applying the above to the link function of MEG in (6), we get

$$\sigma^{-1}(y) = \frac{1}{2\gamma} \pm \sqrt{\frac{1}{4\gamma^2} - \frac{1+m}{h\gamma} + \frac{2\sqrt{m}}{h\gamma} \cdot y}.$$

With this formula, we can plug in 1 and -1 to get:

$$\sigma^{-1}(-1) = \frac{1}{2\gamma} \pm \sqrt{\frac{1}{4\gamma^2} - \frac{(1+\sqrt{m})^2}{h\gamma}} \quad \text{and} \quad \sigma^{-1}(1) = \frac{1}{2\gamma} \pm \sqrt{\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma}}.$$

A.6 Proof of Theorem 3

Proof. We analyze each case separately.

Case 1: There are two square roots: $\sqrt{\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma}}$ and $\sqrt{\frac{1}{4\gamma^2} - \frac{(1+\sqrt{m})^2}{h\gamma}}$. The second one is real if:

$$\frac{1}{4\gamma^2} \geqslant \frac{(1+\sqrt{m})^2}{h\gamma} \implies \frac{h\gamma}{4\gamma^2} = \frac{h}{4\gamma} \geqslant (1+\sqrt{m})^2,$$

which implies the first is real, as $(1+\sqrt{m})^2 \geqslant (1-\sqrt{m})^2$.

Case 3: There are two square roots: $\sqrt{\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma}}$ and $\sqrt{\frac{1}{4\gamma^2} - \frac{(1+\sqrt{m})^2}{h\gamma}}$. The first one is complex if:

$$\frac{1}{4\gamma^2} < \frac{(1-\sqrt{m})^2}{h\gamma} \implies \frac{h\gamma}{4\gamma^2} = \frac{h}{4\gamma} < (1-\sqrt{m})^2,$$

which implies the second is complex, as $(1+\sqrt{m})^2 \geqslant (1-\sqrt{m})^2$.

Case 2: This case follows automatically from the above two cases.

$\square$

A.7 Proof of Proposition 1

Proof. Define $D_3 = \text{diag}(a, \ldots, a)$ of dimensions $d \times d$. Let us prove that if there exists U, V orthonormal matrices and D_1, D_2 matrices with non-zeros coefficients only on the diagonal such that (with a slight abuse of notation)

$$S_1 = U\text{diag}(D_3, D_1)U^\top, S_2 = VD_3V^\top, \quad \text{and} \quad B = UD_2V^\top,$$

then the spectrum of A is crossed shaped. In that case, we have

$$A = \begin{bmatrix} U[D_3; D_1]U^\top & UD_2V^\top, \\ -VD_2^\top U^\top & VD_3V^\top \end{bmatrix}$$

$$= \begin{bmatrix} U & 0 \\ 0 & V \end{bmatrix} \begin{bmatrix} [D_3; D_1] & D_2 \\ -D_2^\top & D_3 \end{bmatrix} \begin{bmatrix} U & 0 \\ 0 & V \end{bmatrix}^\top.$$

Now by considering the basis $W = ((U_1, 0), (0, V_1), \ldots, (U_{d_v}, 0), (0, V_{d_v}), (U_{d_v+1}, 0), \ldots, (U_d, 0))$ we have that A can be block diagonalized in that basis as

$$A = W\text{diag}\left(\begin{bmatrix} a & [D_2]_{11} \\ -[D_2]_{11} & a \end{bmatrix}, \ldots, \begin{bmatrix} a & [D_2]_{d_v,d_v} \\ -[D_2]_{d_v d_v} & a \end{bmatrix}, [D_1]_1, \ldots, [D_1]_{d_u-d_v d_u-d_v}\right) W^\top. \tag{34}$$

Now, notice that

$$\text{Sp}\left(\begin{bmatrix} a & -b \\ b & a \end{bmatrix}\right) = \{a \pm bi\}, \tag{35}$$

since the associated characteristic polynomial of the above matrix is:

$$(a - \lambda)^2 + b^2 = 0 \implies a - \lambda = \pm bi \implies \lambda = a \pm bi.$$

Hence, using (35) in the formulation of A in (34), we have that the spectrum of A is cross-shaped.

$\square$

B Missing Proofs in Section 4

B.1 Proof of Theorem 4

Proof. We write the conditions required for Theorem 5 below:

$$\frac{1}{2\gamma} - \sqrt{\frac{1}{4\gamma^2} - \frac{(1 - \sqrt{m})^2}{h\gamma}} = \mu_1, \tag{36}$$

$$\frac{1}{2\gamma} - \sqrt{\frac{1}{4\gamma^2} - \frac{(1 + \sqrt{m})^2}{h\gamma}} = L_1, \tag{37}$$

$$\frac{1}{2\gamma} + \sqrt{\frac{1}{4\gamma^2} - \frac{(1 + \sqrt{m})^2}{h\gamma}} = \mu_2, \quad \text{and} \tag{38}$$

$$\frac{1}{2\gamma} + \sqrt{\frac{1}{4\gamma^2} - \frac{(1 - \sqrt{m})^2}{h\gamma}} = \mu_2. \tag{39}$$

By adding (37) and (38) (or equivalently by ading (36) and (39)), we get

$$\gamma = \frac{1}{\mu_1 + L_2} = \frac{1}{\mu_2 + L_1}. \tag{40}$$

From (36), we have:

$$\frac{1}{2\gamma} + \sqrt{\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma}} = \mu_1$$

$$\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma} = \left(\frac{1}{2\gamma} - \mu_1\right)^2$$

$$\frac{(1-\sqrt{m})^2}{h} = \mu_1(1-\gamma\mu_1)$$

$$h = \frac{(1-\sqrt{m})^2}{\mu_1(1-\gamma\mu_1)} = \frac{(1-\sqrt{m})^2(\mu_1+L_2)}{\mu_1 L_2} \tag{41}$$

Similarly, from (38), we have:

$$\frac{1}{2\gamma} + \sqrt{\frac{1}{4\gamma^2} - \frac{(1+\sqrt{m})^2}{h\gamma}} = \mu_2$$

$$\frac{1}{4\gamma^2} - \frac{(1+\sqrt{m})^2}{h\gamma} = \left(\frac{\mu_2-L_1}{2}\right)^2$$

$$\left(\frac{\mu_2+L_1}{2}\right)^2 - \left(\frac{\mu_2-L_1}{2}\right)^2 = \mu_2 L_1 = \frac{(1+\sqrt{m})^2}{h\gamma} \tag{42}$$

Combining (41) and (42), and solving for m, we get

$$\mu_2 L_1(1-\sqrt{m})^2 = \mu_1 L_2(1+\sqrt{m})^2$$

$$m = \left(\frac{\sqrt{\mu_2 L_1} - \sqrt{\mu_1 L_2}}{\sqrt{\mu_2 L_1} + \sqrt{\mu_1 L_2}}\right)^2 \overset{(13)}{=} \left(\frac{\sqrt{\rho^2-R^2} - \sqrt{\rho^2-1}}{\sqrt{\rho^2-R^2} + \sqrt{\rho^2-1}}\right)^2. \tag{43}$$

Finally, plugging (43) back to (41), we get:

$$h = \frac{(1-\sqrt{m})^2(\mu_1+L_2)}{\mu_1 L_2}$$

$$= \frac{4\mu_1 L_2}{(\sqrt{\mu_2+L_1} + \sqrt{\mu_1+L_2})^2} \cdot \frac{\mu_1+L_2}{\mu_1 L_2}$$

$$= \frac{4(\mu_1+L_2)}{(\sqrt{\mu_2+L_1} + \sqrt{\mu_1+L_2})^2}.$$

$\square$

B.2 Proof of Theorem 5

Proof. We write the conditions required for Theorem 5 below:

$$\frac{1}{2\gamma} - \sqrt{\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma}} = \mu, \tag{44}$$

$$\frac{1}{2\gamma} + \sqrt{\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma}} = L, \quad \text{and} \tag{45}$$

$$\sqrt{\frac{(1+\sqrt{m})^2}{h\gamma} - \frac{1}{4\gamma^2}} = c. \tag{46}$$

First, by adding (44) and (45), we get:

$$\frac{1}{\gamma} = \mu + L \implies \gamma = \frac{1}{\mu+L}. \tag{47}$$

Plugging (47) back into (44), we have:

$$\frac{1}{2\gamma} - \sqrt{\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma}} = \mu$$

$$\frac{\mu+L}{2} - \mu = \sqrt{\left(\frac{\mu+L}{2}\right)^2 - \frac{(1-\sqrt{m})^2(\mu+L)}{h}}$$

$$\left(\frac{L-\mu}{2}\right)^2 = \left(\frac{\mu+L}{2}\right)^2 - \frac{(1-\sqrt{m})^2(\mu+L)}{h}$$

$$\frac{(1-\sqrt{m})^2(\mu+L)}{h} = \left(\frac{\mu+L}{2}\right)^2 - \left(\frac{L-\mu}{2}\right)^2 = \mu L$$

$$h = \frac{(1-\sqrt{m})^2(\mu+L)}{\mu L}. \tag{48}$$

Plugging (47) and (48) into (46), we have:

$$\sqrt{\frac{(1+\sqrt{m})^2}{h\gamma} - \frac{1}{4\gamma^2}} = c$$

$$\sqrt{\frac{(1+\sqrt{m})^2 \cdot \mu L}{(1-\sqrt{m})^2} - \left(\frac{\mu+L}{2}\right)^2} = c$$

$$\frac{(1+\sqrt{m})^2 \cdot \mu L}{(1-\sqrt{m})^2} = c^2 + \left(\frac{\mu+L}{2}\right)^2 = \frac{4c^2 + (\mu+L)^2}{4}$$

$$\frac{(1+\sqrt{m})^2}{(1-\sqrt{m})^2} = \frac{4c^2 + (\mu+L)^2}{4\mu L}$$

$$(1+\sqrt{m})\sqrt{4\mu L} = (1-\sqrt{m})\sqrt{4c^2 + (\mu+L)^2}$$

$$\sqrt{m}(\sqrt{4c^2 + (\mu+L)^2} + \sqrt{4\mu L}) = \sqrt{4c^2 + (\mu+L)^2} - \sqrt{4\mu L}$$

$$\sqrt{m} = \frac{\sqrt{4c^2 + (\mu+L)^2} - \sqrt{4\mu L}}{\sqrt{4c^2 + (\mu+L)^2} + \sqrt{4\mu L}}. \tag{49}$$

Finally, to simplify (48) further, from (49), we have:

$$1 - \sqrt{m} = \frac{4\sqrt{\mu L}}{\sqrt{4c^2 + (\mu+L)^2} + \sqrt{4\mu L}}.$$

Hence, from (48),

$$h = \frac{(\mu+L)(1-\sqrt{m})^2}{\mu L} = \frac{\frac{16\mu L(\mu+L)}{(\sqrt{4c^2+(\mu+L)^2}+\sqrt{4\mu L})^2}}{\mu L} = \frac{16(\mu+L)}{(\sqrt{4c^2 + (\mu+L)^2} + \sqrt{4\mu L})^2}. \tag{50}$$

$\square$

B.3 Proof of Theorem 6

Proof. We write the conditions required for (6) below:

$$\frac{1}{2\gamma} + \sqrt{\frac{1}{4\gamma^2} - \frac{(1+\sqrt{m})^2}{h\gamma}} = c + bi, \tag{51}$$

$$\frac{1}{2\gamma} - \sqrt{\frac{1}{4\gamma^2} - \frac{(1+\sqrt{m})^2}{h\gamma}} = c + ai, \tag{52}$$

$$\frac{1}{2\gamma} + \sqrt{\frac{1}{4\gamma^2} - \frac{(1+\sqrt{m})^2}{h\gamma}} = c - ai, \quad \text{and} \tag{53}$$

$$\frac{1}{2\gamma} + \sqrt{\frac{1}{4\gamma^2} - \frac{(1-\sqrt{m})^2}{h\gamma}} = c - bi. \tag{54}$$

First, we can see from all cases that the optimal γ is

$$\gamma = \frac{1}{2c}. \tag{55}$$

(51) and (54) equivalently imply

$$\sqrt{\frac{(1+\sqrt{m})^2}{h\gamma} - \frac{1}{4\gamma^2}} = b$$

$$(1+\sqrt{m})^2 = h\gamma b^2 + \frac{h}{4\gamma} = \frac{h(c^2+b^2)}{2c}$$

$$h = \frac{2c(1+\sqrt{m})^2}{c^2+b^2}. \tag{56}$$

Similarly, (52) and (53) imply

$$\sqrt{\frac{(1-\sqrt{m})^2}{h\gamma} - \frac{1}{4\gamma^2}} = a$$

$$\frac{(1-\sqrt{m})^2}{h\gamma} = a^2 + \frac{1}{4\gamma^2} = a^2 + c^2$$

$$\frac{(1-\sqrt{m})^2(c^2+b^2)}{(1+\sqrt{m})^2} = a^2 + c^2$$

$$(1-\sqrt{m})\sqrt{c^2+b^2} = (1+\sqrt{m})\sqrt{c^2+a^2}$$

$$\sqrt{m} = \frac{\sqrt{c^2+b^2} - \sqrt{c^2+a^2}}{\sqrt{c^2+b^2} + \sqrt{c^2+a^2}} = 1 - \frac{2\sqrt{c^2+a^2}}{\sqrt{c^2+b^2} + \sqrt{c^2+a^2}} \tag{57}$$

Plugging (57) to (56), we get

$$h = \frac{2c(1+\sqrt{m})^2}{c^2+b^2} = \frac{8c}{(\sqrt{c^2+b^2} + \sqrt{c^2+a^2})^2}.$$

$\square$

C Missing Proofs in Sections 5

C.1 Proof of Corollary 1

Proof. To compute the convergence rates of GD and EG from Theorem 7 applied to each spectrum model in (12), (14), and (15), we need to compute $\min_{\lambda \in \Delta^\star} \Re(1/\lambda)$ and $\min_{\lambda \in \Delta^\star} \Re(\lambda)$ for GD. Similarly for EG, we need to compute additionally $\min_{\lambda \in \Delta^\star} |\lambda|$, $\min_{\lambda \in \Delta^\star} |\lambda|^2$, and $\sup_{\lambda \in \Delta^\star} |\lambda|^2$.

Case 1: It's straightforward to compute

$$\min_{\lambda \in \mathcal{S}_1^\star} \mathfrak{R}(1/\lambda) = 1/L_2, \quad \text{and} \quad \min_{\lambda \in \mathcal{S}_1^\star} \mathfrak{R}(\lambda) = \mu_1$$

Thus, GD for Case 1 has the rate

$$1 - \frac{\mu_1}{L_2} = 1 - \tau.$$

For EG, it's also simple to obtain

$$\min_{\lambda \in \mathcal{S}_1^\star} |\lambda| = L_2, \quad \min_{\lambda \in \mathcal{S}_1^\star} |\lambda|^2 = \mu_1^2, \quad \text{and} \quad \sup_{\lambda \in \mathcal{S}_1^\star} |\lambda|^2 = L_2^2.$$

Thus, EG for Case 1 has the rate

$$1 - \frac{1}{4}\left(\frac{\mu_1}{L_2} + \frac{1}{16}\left(\frac{\mu_1}{L_2}\right)^2\right).$$

Case 2: For a complex number $z = p + qi \in \mathbb{C}$, we can compute $\mathfrak{R}(1/z)$ as:

$$\frac{1}{z} = \frac{1}{p+qi} = \frac{p-qi}{p^2+q^2} = \frac{p}{p^2+q^2} - \frac{q}{p^2+q^2}i \implies \mathfrak{R}\left(\frac{1}{z}\right) = \frac{p}{p^2+q^2}.$$

The four extreme points of the cross-shaped spectrum model in (14) are:

$$\mu = \mu + 0i, \quad L = L + 0i, \quad \text{and} \quad \frac{L-\mu}{2} \pm ci.$$

Hence, $\mathfrak{R}(1/z)$ for each of the above points is:

$$\mathfrak{R}\left(\frac{1}{\mu}\right) = \frac{\mu}{\mu^2} = \frac{1}{\mu},$$

$$\mathfrak{R}\left(\frac{1}{L}\right) = \frac{L}{L^2} = \frac{1}{L}, \quad \text{and}$$

$$\mathfrak{R}\left(\frac{1}{\frac{L-\mu}{2} \pm ci}\right) = \frac{\frac{L-\mu}{2}}{\left(\frac{L-\mu}{2}\right)^2 + c^2}$$

$$= \frac{2(L-\mu)}{4c^2 + (L-\mu)^2}.$$

Therefore, $\min_{\lambda \in \mathcal{S}_2^\star} \mathfrak{R}\left(\frac{1}{\lambda}\right) = \frac{1}{L}$. As $\mu < L$, we only need to compare the last two values. Observe that:

$$c > \sqrt{\frac{L^2 - \mu^2}{4}}$$

$$4c^2 > (L-\mu)(L+\mu)$$

$$4c^2 > 2L(L-\mu) - (L-\mu)^2$$

$$\frac{1}{L} > \frac{2(L-\mu)}{4c^2 + (L-\mu)^2}.$$

Therefore,

$$\min_{\lambda \in \mathcal{S}_2^\star} \mathfrak{R}\left(\frac{1}{\lambda}\right) = \begin{cases} \frac{2(L-\mu)}{4c^2+(L-\mu)^2} & \text{if} \quad c > \sqrt{\frac{L^2-\mu^2}{4}} \\ \frac{1}{L} & \text{otherwise.} \end{cases}$$

For $\min_{\lambda \in \mathcal{S}_2^\star} \mathfrak{R}(\lambda)$, it's straightforward from the definition that

$$\min_{\lambda \in \mathcal{S}_2^\star} \mathfrak{R}(\lambda) = \mu.$$

Thus, GD for Case 2 has the rate

$$\begin{cases} 1 - \frac{2\mu(L-\mu)}{4c^2 + (L-\mu)^2} & \text{if} \quad c > \sqrt{\frac{L^2 - \mu^2}{4}} \\ 1 - \frac{\mu}{L} & \text{otherwise.} \end{cases}$$

Similarly for EG, we need to compute $\min_{\lambda \in \mathcal{S}_2^\star} \mathfrak{R}(\lambda)$, which was computed above; additionally, we need to compute $\min_{\lambda \in \mathcal{S}_2^\star} |\lambda|$, $\min_{\lambda \in \mathcal{S}_2^\star} |\lambda|^2$, and $\sup_{\lambda \in \mathcal{S}_2^\star} |\lambda|^2$. For $z = p + qi \in \mathbb{C}$, $|z| = \sqrt{p^2 + q^2}$. Hence, we have

$$|\mu + 0i| = \mu, \quad |L + 0i| = L, \quad \text{and} \quad \left| \frac{L-\mu}{2} \pm ci \right| = \sqrt{c^2 + \left(\frac{L-\mu}{2} \right)^2}.$$

Observe that:

$$c > \sqrt{\frac{3L^2 + 2L\mu - \mu^2}{4}}$$

$$c^2 > L^2 - \frac{L^2 - 2L\mu + \mu^2}{4}$$

$$c^2 + \left(\frac{L-\mu}{2} \right)^2 > L^2$$

Thus, for $\sup_{\lambda \in \mathcal{S}_2^\star} |\lambda|$, we have

$$\sup_{\lambda \in \mathcal{S}_2^\star} |\lambda| = \begin{cases} \sqrt{c^2 + \left(\frac{L-\mu}{2} \right)^2} & \text{if} \quad c > \sqrt{\frac{3L^2 + 2L\mu - \mu^2}{4}} \\ L & \text{otherwise,} \end{cases}$$

from which $\sup_{\lambda \in \mathcal{S}_2^\star} |\lambda|^2$ can also be obtained. Lastly, $\min_{\lambda \in \mathcal{S}_2^\star} |\lambda|^2 = \mu^2$, as we know $\mu < L$, and $(L-\mu)/2$ is the center of $[\mu, L]$.

Combining all three, we get the rate of EG for Case 3 is

$$\begin{cases} 1 - \frac{1}{4} \left(\frac{\mu}{\sqrt{c^2 + \left(\frac{L-\mu}{2} \right)^2}} + \frac{\mu^2}{16 \left(c^2 + \left(\frac{L-\mu}{2} \right)^2 \right)} \right) & \text{if } c \geqslant \sqrt{\frac{3L^2 + 2L\mu - \mu^2}{4}}, \\ 1 - \frac{1}{4} \left(\frac{\mu}{L} + \frac{\mu^2}{16L^2} \right) & \text{otherwise.} \end{cases}$$

Case 3: Since (15) has fixed real component, $\min_{\lambda \in \mathcal{S}_3^\star} \mathfrak{R}(\lambda) = c$.

For $\min_{\lambda \in \mathcal{S}_3^\star} \mathfrak{R}(1/\lambda)$ we can compute compare

$$\mathfrak{R}\left(\frac{1}{c + ai} \right) = \frac{c}{c^2 + a^2} > \frac{c}{c^2 + b^2} = \mathfrak{R}\left(\frac{1}{c + bi} \right),$$

since $a < b$ from (15). Thus, GD for Case 3 has the rate

$$1 - \frac{c^2}{c^2 + b^2}.$$

For EG, it's also simple to obtain

$$\min_{\lambda \in \mathcal{S}_3^\star} |\lambda| = \sqrt{c^2 + b^2}, \quad \min_{\lambda \in \mathcal{S}_3^\star} |\lambda|^2 = c^2 + a^2, \quad \text{and} \quad \sup_{\lambda \in \mathcal{S}_3^\star} |\lambda|^2 = c^2 + b^2.$$

Thus, EG has the rate

$$1 - \frac{1}{4} \left(\frac{c}{\sqrt{c^2 + b^2}} + \frac{1}{16} \frac{(c^2 + a^2)}{(c^2 + b^2)} \right).$$

$\square$

C.2 Proof of Corollary 2

Proof. Per Theorem 8, the largest ϵ that permits acceleration for GDM is $\epsilon = \sqrt{\mu L}$. Therefore, in the special case of (14) we consider, i.e., when $c = \frac{L-\mu}{2}$, GDM *cannot* achieve acceleration if $\frac{L-\mu}{2} > \sqrt{\mu L}$. Hence, we have:

$$\frac{L-\mu}{2} > \sqrt{\mu L}$$
$$L - \mu > 2\sqrt{\mu L}$$
$$L^2 + \mu^2 > 6\mu L > 6\mu^2 \quad (\because L > \mu)$$
$$L > \sqrt{5}\mu.$$

$\square$

C.3 Proof of Proposition 2

Proof. For an arbitrary complex number $p + qi$ with $p > 0$, and using the link function of GDM from (7), we have

$$|\xi(p+qi)| = \sqrt{\left(\frac{1+m-hp}{2\sqrt{m}}\right)^2 + \left(\frac{hq}{2\sqrt{m}}\right)^2} \leqslant 1$$
$$\frac{(1+m-hp)^2 + h^2q^2}{4m} \leqslant 1$$
$$(1+m-hp)^2 + h^2q^2 \leqslant 4m$$
$$(1-m)^2 + hp(hp - 2(1+m)) + h^2q^2 \leqslant 0$$
$$\frac{(1-m)^2 + h^2q^2}{hp} \leqslant 2(1+m) - hp$$

Notice that the LHS is positive. Therefore, if the RHS is negative, the above inequality cannot hold. In other words, if $\frac{2(1+m)}{h} < p$, GDM cannot stay in the robust region. This is very hard to satisfy, even with a small p.

$\square$

D Missing Proofs in Section 6

Let us consider an affine vector field $v(w) = Aw + b$ and its associated augmented MEG linear operator J:

$$\begin{bmatrix} w_{t+1} - w^\star \\ w_t - w^\star \end{bmatrix} = J \begin{bmatrix} w_t - w^\star \\ w_{t-1} - w^\star \end{bmatrix} \quad \text{with} \quad J = \begin{bmatrix} (1+\beta)I_d - hA(I_d - \gamma A) & -\beta I_d \\ I_d & 0_d \end{bmatrix}, \tag{58}$$

where I_d and 0_d respectively stand for the identity and the null matrices. To show the local convergence of (restarted) MEG for non-affine vector fields in Theorem 9, we first establish the following lemma, which connects the augmented state and the non-augmented one.

Lemma 3. *Let P_t^{MEG} be the residual polynomial associated with t updates of MEG (c.f., Theorem 1). Let J be defined as (58). If $w_1 = w_0 - \frac{h}{1+m}v(w_0 - \gamma v(w_0))$, we then have*

$$J^t \begin{bmatrix} w_1 - w^\star \\ w_0 - w^\star \end{bmatrix} = \begin{bmatrix} P_t^{MEG}(A)(w_1 - w^\star) \\ P_t^{MEG}(A)(w_0 - w^\star) \end{bmatrix}. \tag{59}$$

Consequently, if we denote $z_{t+1} := [w_{t+1}, w_t]$ and $z_* := [w^\star, w^\star]$, we have

$$\|z_{t+1} - z_*\| \leqslant C(t+1)(1-\rho)^t\|z_0 - z_*\|. \tag{60}$$

Proof. Let us express J^t such that

$$J^t = \begin{bmatrix} P_t^{11}(A) & P_t^{12}(A) \\ P_t^{21}(A) & P_t^{22}(A) \end{bmatrix}, \quad \text{and} \quad J^t \begin{bmatrix} w_1 - w^\star \\ w_0 - w^\star \end{bmatrix} = \begin{bmatrix} P_t^{11}(A)(w_0 - w^\star) + P_t^{12}(A)(w_0 - w^\star) \\ P_t^{21}(A)(w_0 - w^\star) + P_t^{22}(A)(w_0 - w^\star) \end{bmatrix}. \tag{61}$$

By writing $J^{t+1} = JJ^t$ and using the block-matrix form of J in (58), we get that for any $t \geqslant 0$,

$$P_{t+1}^{11}(A) = ((1+\beta)I_d - hA(I_d - \gamma A))P_t^{11}(A) - \beta P_t^{21}(A)$$
$$P_{t+1}^{21}(A) = P_t^{11}(A) \tag{62}$$
$$P_{t+1}^{12}(A) = ((1+\beta)I_d - hA(I_d - \gamma A))P_t^{12}(A) - \beta P_t^{22}(A)$$
$$P_{t+1}^{22}(A) = P_t^{12}(A). \tag{63}$$

Hence, we have that,

$$P_{t+1}^{11}(A) \overset{(62)}{=} ((1+\beta)I_d - hA(I_d - \gamma A))P_t^{11}(A) - \beta P_{t-1}^{11}(A) \tag{64}$$

$$P_{t+1}^{12}(A) \overset{(63)}{=} ((1+\beta)I_d - hA(I_d - \gamma A))P_t^{12}(A) - \beta P_{t-1}^{12}(A). \tag{65}$$

We claim that

$$P_t^{11}(A) + P_t^{12}(A) = P_t^{\text{MEG}}(A) \quad \text{for all} \quad t \geqslant 0. \tag{66}$$

We prove this via induction.

For the base case, using the fact that $w_1 = w_0 - \frac{h}{1+m}(w_0 - \gamma v(w_0))$, we have that

$$(P_1^{11}(A) + P_1^{12}(A))(w_0 - w^\star) = w_1 - w^\star = (I_d - \tfrac{h}{1+m}(I_d - \gamma A))(w_0 - w^\star) = P_1^{\text{MEG}}(A)(w_0 - w^\star),$$
$$(P_0^{11}(A) + P_0^{12}(A))(w_0 - w^\star) = (P_1^{21}(A) + P_1^{22}(A))(w_0 - w^\star) = I_d(w_0 - w^\star) = P_0^{\text{MEG}}(A)(w_0 - w^\star).$$

To show the induction step, by adding (64) and (65), we get

$$P_{t+1}^{11}(A) + P_{t+1}^{12}(A) = ((1+\beta)I_d - hA(I_d - \gamma A))(P_t^{11}(A) + P_t^{12}(A)) - \beta(P_{t-1}^{11}(A) + P_{t-1}^{12}(A))$$
$$\overset{(66)}{=} ((1+\beta)I_d - hA(I_d - \gamma A))P_t^{\text{MEG}}(A) - \beta P_{t-1}^{\text{MEG}}(A),$$

where in the last step we used the induction hypothesis. Also notice $P_{t+1}^{11}(A) + P_{t+1}^{12}(A) = P_{t+1}^{\text{MEG}}$ on the left-hand side.

Hence we have for any $t \geqslant 0$,

$$(P_t^{11}(A) + P_t^{12}(A))(w_0 - w^\star) = P_t^{\text{MEG}}(A)(w_0 - w^\star). \tag{67}$$

Therfore, going back to (61), we have:

$$\begin{bmatrix} w_{t+1} - w^\star \\ w_t - w^\star \end{bmatrix} = J^t \begin{bmatrix} w_1 - w^\star \\ w_0 - w^\star \end{bmatrix} = \begin{bmatrix} (P_t^{11}(A) + P_t^{12}(A))(w_0 - w^\star) \\ (P_t^{21}(A) + P_t^{22}(A))(w_0 - w^\star) \end{bmatrix}$$
$$\overset{(62),(63)}{=} \begin{bmatrix} (P_t^{11}(A) + P_t^{12}(A))(w_0 - w^\star) \\ (P_{t-1}^{11}(A) + P_{t-1}^{12}(A))(w_0 - w^\star) \end{bmatrix}$$
$$\overset{(66)}{=} \begin{bmatrix} P_t^{\text{MEG}}(A)(w_0 - w^\star) \\ P_{t-1}^{\text{MEG}}(A)(w_0 - w^\star) \end{bmatrix}$$
$$\overset{\text{Thm. } 1}{=} \begin{bmatrix} P_t^{\text{MEG}}(A)(w_0 - w^\star) \\ P_t^{\text{MEG}}(A)(w_{-1} - w^\star) \end{bmatrix}$$
$$= \begin{bmatrix} P_t^{\text{MEG}}(A) & 0 \\ 0 & P_t^{\text{MEG}}(A) \end{bmatrix} \cdot \begin{bmatrix} w_0 - w^\star \\ w_{-1} - w^\star \end{bmatrix}$$
$$= P_t^{\text{MEG}}(A) \otimes I_2 \cdot \begin{bmatrix} w_0 - w^\star \\ w_{-1} - w^\star \end{bmatrix},$$

where we use the convention that $w_0 = w_{-1}$. Finally, using the fact that $\|A \otimes B\| = \|A\|\|B\|$ for ℓ_2-operator norm (Lancaster & Farahat, 1972), we have

$$\|z_{t+1} - z_*\| \leqslant \|P_t^{\text{MEG}}(A)\|\|z_0 - z_*\| \overset{(10)}{\leqslant} C(t+1)(1-\rho)^t\|z_0 - z_*\|.$$

$\square$

D.1 Proof of Theorem 9

Proof. We first recall the restarted MEG algorithm we consider in (31):

$$[w_{tk+i+1}, w_{tk+i}] = G([w_{tk+i}, w_{tk+i-1}]) \quad \text{for} \quad 1 \leqslant i \leqslant k-1, \quad \text{and then}$$
$$w_{(t+1)k+1} = w_{(t+1)k} - \tfrac{h}{1+m}v(w_{(t+1)k} - \gamma v(w_{(t+1)k})).$$

In other words, we repeat MEG for k steps, and then re-start the momentum at $[w_{(t+1)k+1}, w_{(t+1)k}]$.

We can analyze this method as follows, where we denote $z_t := [w_t, w_{t-1}]$ and $z_* = [w^\star, w^\star]$:

$$\begin{aligned}
\|z_{(t+1)k} - z_*\| &= \|G^{(k)}(z_{tk}) - z_*\| \\
&\overset{(68)}{=} \|\nabla G^{(k)}(\tilde{z}_{tk})(z_{tk} - z_*)\| \\
&\leqslant \|\nabla G^{(k)}(z_*)(z_{tk} - z_*)\| + \|(\nabla G^{(k)}(\tilde{z}_{tk}) - \nabla G^{(k)}(z_*))(z_{tk} - z_*)\| \\
&\overset{(60)}{\leqslant} C(k+1)(1-\rho)^k\|z_{tk} - z_*\| + \|\nabla G^{(k)}(\tilde{z}_{tk}) - \nabla G^{(k)}(z_*)\|\|(z_{tk} - z_*)\|,
\end{aligned}$$

where in the second line we use the Mean Value Theorem:

$$\begin{aligned}
\exists \tilde{z}_{tk} \in [z_{tk}, z_*] \quad \text{such that} \quad G^{(k)}(z_{tk}) &= G^{(k)}(z_*) + \nabla G^{(k)}(\tilde{z}_{tk})(z_{tk} - z_*) \\
&= z_* + \nabla G^{(k)}(\tilde{z}_{tk})(z_{tk} - z_*) \quad \text{(since } z_* \text{ is the fixed point).} \quad (68)
\end{aligned}$$

In the fourth line we used the fact that $\nabla G^{(k)}(z_*)(z_{tk} - z_*)$ exactly correspond to k updates of MEG when the vector field is affine, as well as Lemma 3 to account for the augmented state.

Now let us consider $\rho > \varepsilon > 0$ and k large enough such that $C(k+1)(1-\rho)^k \leqslant (1-\rho+\frac{\varepsilon}{2})^k$. Since ∇G is assumed to be continuous, $\nabla G^{(k)}$ is continuous too. Therefore, there exists $\delta > 0$ such that $\|z_{tk} - z_*\| \leqslant \delta$ implies $\|\nabla G^{(k)}(\tilde{z}_{tk}) - \nabla G^{(k)}(z_*)\| \leqslant \varepsilon'$. In particular, choose $\varepsilon' = (1-\rho+\varepsilon)^k - (1-\rho+\frac{\varepsilon}{2})^k \sim \frac{k\varepsilon}{2(1-\rho)}$. Then, we have

$$\begin{aligned}
\|z_{(t+1)k} - z_*\| &\leqslant C(k+1)(1-\rho)^k\|z_{tk} - z_*\| + \|\nabla G^{(k)}(\tilde{z}_{tk}) - \nabla G^{(k)}(z_*)\|\|z_{tk} - z_*\| \\
&\leqslant (1-\rho+\tfrac{\varepsilon}{2})^k\|z_{tk} - z_*\| + \varepsilon'\|z_{tk} - z_*\| \\
&\leqslant (1-\rho+\varepsilon)^k\|z_{tk} - z_*\| < \|z_{tk} - z_*\| < \|z_0 - z_*\|.
\end{aligned}$$

From the above, we can conclude that for all $\varepsilon > 0$, there exists $k > 0$ and $\delta > 0$ such that, for all initialization satisfying $\|w_0 - w^\star\| \leqslant \delta$, the restarted MEG described above satisfies:

$$\|w_t - w^\star\| = O((1-\rho+\varepsilon)^t)\|w_0 - w^\star\|.$$

$\square$