ACCOUNTING FOR UNOBSERVED CONFOUNDING IN DOMAIN GENERALIZATION

ABSTRACT

The ability to extrapolate, or generalize, from observed to new related environments is central to any form of reliable machine learning, yet most methods fail when moving beyond $i.i.d$ data. In some cases, the reason lies in a misappreciation of the causal structure that governs the data, and in particular as a consequence of the influence of unobserved confounders that drive changes in observed distributions and distort correlations. In this paper, we argue for defining generalization with respect to a broader class of distribution shifts (defined as arising from interventions in the underlying causal model), including changes in observed, unobserved and target variable distributions. We propose a new robust learning principle that may be paired with any gradient-based learning algorithm. This learning principle has explicit generalization guarantees, and relates robustness with certain invariances in the causal model, clarifying why, in some cases, test performance lags training performance. We demonstrate the empirical performance of our approach on healthcare data from different modalities, including image and speech data.

1 INTRODUCTION

Prediction algorithms use data, necessarily sampled under specific conditions, to learn correlations that extrapolate to new or related data. If successful, the performance gap between these two domains is small, and we say that algorithms *generalize* beyond their training data. Doing so is difficult however, some form of uncertainty about the distribution of new data is unavoidable. The set of potential distributional changes that we may encounter is mostly unknown and in many cases may be large and varied. Some examples include covariate shifts (Bickel et al., 2009), interventions in the underlying causal system (Pearl, 2009), varying levels of noise (Fuller, 2009) and confounding (Pearl, 1998). All of these feature in modern applications, and while learning systems are increasingly deployed in practice, generalization of predictions and their reliability in a broad sense remains an open question.

A common approach to formalize learning with uncertain data is, instead of optimizing for correlations in a *fixed* distribution, to do so simultaneously for a *range* of different distributions in an uncertainty set $\mathcal{P}$ (Ben-Tal et al., 2009).

$$\underset{f}{\text{minimize}} \sup_{P \in \mathcal{P}} \mathbb{E}_{(x,y) \sim P}[\mathcal{L}(f(x), y)] \tag{1}$$

for some measure of error $\mathcal{L}$ of the function f that relates input and output examples $(x, y) \sim P$. Choosing different sets $\mathcal{P}$ leads to estimators with different properties. It includes as special cases, for instance, many approaches in domain adaptation, covariate shift, robust statistics and optimization (Kuhn et al., 2019; Bickel et al., 2009; Duchi et al., 2016; 2019; Sinha et al., 2017; Wozabal, 2012; Abadeh et al., 2015; Duchi & Namkoong, 2018). Robust solutions to problem (1) are said to generalize if potential shifted, test distributions are contained in $\mathcal{P}$, but also larger sets $\mathcal{P}$ result in *conservative* solutions (i.e. with sub-optimal performance) on data sampled from distribution away from worst-case scenarios, in general.

One formulation of causality is in fact also a version of this problem, for $\mathcal{P}$ defined as any distribution arising from *arbitrary* interventions on observed covariates x leading to shifts in their distribution P_x (see e.g. sections 3.2 and 3.3 in (Meinshausen, 2018)). The invariance to changes in covariate distributions of causal solutions is powerful for generalization, but implicitly assumes that all

Figure 1: **The challenges of generalization**. We consider the following learning paradigms: Ordinary Least Squares (OLS) learns the linear mapping that minimizes average training risk, Domain Robust Optimization (DRO) minimizes the maximum training risk among the two available datasets, and the causal solution, assumed known, has fixed coefficients $(0, 1)$ for (X_1, X_2). The proposed approach (Ours) is described in section 3.1. The data generating mechanism and a discussion of the results are given in the paragraph above our contributions.

covariates or other drivers of the outcome subject to change at test time are observed. Often shifts occur elsewhere, for example in the distribution of unobserved confounders, in which case also conditional distributions $P_{y|x}$ may shift. Perhaps surprisingly, in the presence of unobserved confounders, the goals of achieving robustness and learning a causal model can be *different* (and similar behaviour also occurs with varying measurement noise and missing data).

We introduce in Figure 1 a simple example designed to show empirically the challenges of out of distribution generalization. We assume access to observations of variables (X_1, X_2, Y) in two training datasets, each dataset sampled with differing variances ($\sigma^2 = 1$ and $\sigma^2 = 2$) from the following structural model $\mathbb{F}$,

$$X_2 := -H + E_{X_2}, \quad Y := X_2 + 3H + E_Y, \quad X_1 := Y + X_2 + E_{X_1},$$

where $E_{X_1}, E_{X_2} \sim \mathcal{N}(0, \sigma^2)$, $E_Y \sim \mathcal{N}(0, 1)$ are exogenous variables. In a first scenario (leftmost panel) we consider all data (training and testing) to be generated *without* unobserved confounders, $H := 0$; and, in a second scenario (remaining panels) all data *with* unobserved confounders, $H := E_H \sim \mathcal{N}(0, 1)$. Each panel shows performance on *new* data obtained after manipulating the underlying data generating system; the magnitude and type of intervention appears in the horizontal axis. We make two important observations that motivate this paper. First, observe that OLS and DRO absorb *spurious* correlations (due to H, and the fact that X_1 is caused by Y) with unstable performance under shifts in $p(X_1, X_2)$. Causal solutions, by contrast, are robust to shifts in $p(X_1, X_2)$, even on new data with large shifts, but underperform substantially under changes in the distribution of unobserved confounders $p(H)$. Second, the presence of unobserved confounding *hurts* generalization performance in general with higher errors for all methods, e.g. contrast the middle and leftmost panel. To the best of our knowledge, the influence of unobserved confounders has been minimally explored in the context of generalization of learning algorithms, even though, as Figure 1 shows, in this context different shifts in distribution may have important consequences for predictive performance.

Our Contributions. In this paper we provide a new choice of $\mathcal{P}$ and learning problem (1) that we show to be justified by certain statistical *invariances* across training and testing data, to be expected in the presence of unobserved confounders. This leads us to define a new differentiable, regularized objective for representation learning. Our proposal defines $\mathcal{P}$ as an *affine* combination of available training data distributions, and we show that solutions to this problem are robust to more general shifts in distribution than previously considered, spanning robustness to shifts in observed, unobserved, and target variables, depending on the properties of the available training data distributions. This approach has benefits for performance out-of-sample but also for tasks involving variable selection, where important features are consistently replicated across experiments with our objective.

2 INVARIANCES IN THE PRESENCE OF UNOBSERVED CONFOUNDERS

This section formally introduces the problem of out-of-distribution generalization. We describe in greater detail the reasons that popular learning principles, such as Empirical Risk Minimization (ERM), underperform in general, and define certain invariances to recover solutions that generalize.

We take the perspective that all potential distributions that may be observed over a system of variables arise from a causal model $\mathcal{M} = (\mathbb{F}, \mathbb{V}, \mathbb{U})$, characterized by endogenous variables, $\mathbb{V} \in \mathcal{V}$,

representing all variables determined by the system, either observed or not; exogenous variables, $\mathbb{U} \in \mathcal{U}$, in contrast imposed upon the model, and a sequence of structural equations $\mathbb{F} : \mathcal{U} \rightarrow \mathcal{V}$, describing how endogenous variables can be (deterministically) obtained from the exogenous variables (Pearl, 2009). An example is given in Figure 1, $\mathbb{V} = (X_1, X_2, H, Y)$ are endogenous and $\mathbb{U} = (E_{X_1}, E_{X_2}, E_H, E_Y)$ are exogenous variables.

Unseen data is generated from such a system $\mathcal{M}$ after manipulating the distribution of exogenous variables $\mathbb{U}$, which propagates across the system shifting the joint distribution of all variables $\mathbb{V}$, whether observed or unobserved, but keeping the causal mechanisms $\mathbb{F}$ unchanged. Representative examples include changes in data collection conditions, such as due to different measurement devices, or new data sources, such as patients in different hospitals or countries, among many others.

Our goal is to learn a representation $Z = \phi(X)$ acting on a set observed variables $X \subset \mathbb{V}$ with the ability to extrapolate to new unseen data, and doing so acknowledging that all relevant variables in $\mathbb{V}$ are likely not observed. Unobserved confounders (for the task at hand, say predicting $Y \in \mathbb{V}$) simultaneously cause X and Y, confounding or biasing the causal association between X and Y giving rise to spurious correlations that do not reproduce in general (Pearl, 1998; 2009). We present a brief argument below highlighting the systematic bias due to unobserved confounders in ERM.

2.1 THE BIASES OF UNOBSERVED CONFOUNDING

Consider the following structural equation for observed variables (X, Y),

$$Y := f \circ \phi(X) + E \tag{2}$$

where $f := f(\cdot; \beta_0)$ is a predictor acting on a representation $Z := \phi(X)$ and E stands for potential sources of mispecification and unexplained sources of variability. For a given sample of data (x, y) and $z = \phi(x)$, the optimal prediction rule $\hat{\beta}$ is often taken to minimize squared residuals, with $\hat{\beta}$ the solution to the normal equations: $\nabla_\beta f(z; \hat{\beta}) y = \nabla_\beta f(z; \hat{\beta}) f(z; \hat{\beta})$, where $\nabla_\beta f(z; \hat{\beta})$ denotes the column vector of gradients of f with respect to parameters β evaluated at $\hat{\beta}$. Consider the Taylor expansion of $f(z; \beta_0)$ around an estimate $\hat{\beta}$ sufficiently close to β_0, $f(z; \beta_0) \approx f(z; \hat{\beta}) + \nabla_\beta f(z; \hat{\beta})^T (\beta_0 - \hat{\beta})$. Using this approximation in our first order optimality condition we find,

$$\nabla_\beta f(z; \hat{\beta}) \nabla_\beta f(z; \hat{\beta})^T (\beta_0 - \hat{\beta}) + v = \nabla_\beta f(z; \hat{\beta}) \epsilon \tag{3}$$

where v is a scaled disturbance term that includes the rest of the linear approximation of f and is small asymptotically; $\epsilon := y - f(z; \hat{\beta})$ is the residual. $\hat{\beta}$ is consistent for the true β_0 if and only if $\nabla_\beta f(z; \hat{\beta}) \epsilon \rightarrow 0$ in probability. This assumption is satisfied if E (all sources of variation in Y not captured by X) are independent of X (i.e. exogenous) or in other words if all common causes or confounders to both X and Y have been observed. Conventional regression may assign significant associations to variables that are neither directly nor indirectly related to the outcome, and in this case, we have no performance guarantees on new data with changes in the distribution of these variables. Omitted variables are a common source of unobserved confounding but we note in Appendix A.1 that similar biases also arise from other prevalent model mispecifications, such as measurement error (Carroll et al., 2006), and missing data (Westreich et al., 2015).

2.2 INVARIANCES WITH MULTIPLE ENVIRONMENTS

The underlying structural mechanism $\mathbb{F}$, that also relates unobserved with observed variables, even if unknown, is stable irrespective of manipulations in exogenous variables that may give rise to heterogeneous data sources. Under certain conditions, statistical footprints emerge from this structural invariance across different data sources, properties testable from data that have been exploited recently, for example (Peters et al., 2016; Ghassami et al., 2017; Rothenhäusler et al., 2019).

We assume that such a heterogeneous data scenario applies, input and output pairs (X, Y) are observed across heterogeneous data sources or environments e, defined as a probability distribution P_e over an observation space $\mathcal{X} \times \mathcal{Y}$ that arises, just like new unseen data, from manipulations in the distribution of exogenous variables in an underlying model $\mathcal{M}$.

For the remainder of this section, consider restricting ourselves to data sources emerging from manipulations in exogenous E_X, appearing in the structural equations of X only in an underlying

additive model. It may be shown by considering the distributions of error terms $Y - f \circ \phi(X)$ and its correlation with any function of X, that the inner product $\nabla_\beta f(z; \beta_0)\epsilon$, even if *non-zero* due to unobserved confounding, converges to a fixed unknown value *equal* across training environments (see Appendix A.3.1 for the derivation). With a similar decomposition to the one given in equation (3), in the population case, it holds that up to disturbance terms,

$$\left(\mathop{\mathbb{E}}_{(x,y)\sim P_i} \nabla_\beta f(z; \beta^\star) \nabla_\beta f(z; \beta^\star)^T - \mathop{\mathbb{E}}_{(x,y)\sim P_j} \nabla_\beta f(z; \beta^\star) \nabla_\beta f(z; \beta^\star) \right)^T (\beta_0 - \beta^\star)$$

$$= \left(\mathop{\mathbb{E}}_{(x,y)\sim P_i} \nabla_\beta f(z; \beta^\star)\epsilon - \mathop{\mathbb{E}}_{(x,y)\sim P_j} \nabla_\beta f(z; \beta^\star)\epsilon \right) = 0 \qquad (4)$$

where $\beta^\star$ is a solution to,

$$\mathop{\mathbb{E}}_{(x,y)\sim P_i} \nabla_\beta f(z; \beta)(y - f(z; \beta)) - \mathop{\mathbb{E}}_{(x,y)\sim P_j} \nabla_\beta f(z; \beta)(y - f(z; \beta)) = 0. \qquad (5)$$

and is *consistent* for the causal parameters β_0 if unique. $i, j \in \mathcal{E}$ are the indeces of any two observed environments in an index set $\mathcal{E}$. This invariance across environments must hold for causal parameters (under certain conditions) *even* in the presence of unobserved confounders.

A few remarks are necessary concerning this relationship and its extrapolation properties.

- The first is based on the observation that, up to a constant, each inner product in (5) is the gradient of the squared error with respect to β. This reveals that the optimal predictor, in the presence of unobserved confounding, is not one that produces minimum loss but one that produces a *non-zero* loss gradient *equal* across environments. Seeking minimum error solutions, even in the population case, produces estimators with *necessarily* unstable correlations because the variability due to unobserved confounders is not explainable from observed data. Forcing gradients to be zero then *forces* models to utilize artifacts of the specific data collection process that are not related to the input-output relationship; and, for this reason, will not in general perform outside training data.

- From (5) we may pose a sequence of moment conditions for each pair of available environments. We may then seek solutions β that make all of them small simultaneously. Solutions are unique if the set of moments is sufficient to identify $\beta^\star$ exactly (and given our model assumptions may be interpreted as causal and robust to certain interventions). We revisit our introductory example to show in Appendix A.2 that, in contrast to ERM and Invariant Risk Minimization (IRM) (a related approach proposed in (Arjovsky et al., 2019) we discuss in more detail in later sections), this procedure does recover the underlying causal model correctly.

- However, in practice, only a set of solutions may be identified with no performance guarantees for any individual solutions. Moreover, even if accessible, we have seen that causal solutions, robust to certain distribution shifts, may not always be desirable under more general shifts.

3 GENERALIZATION FROM A ROBUST OPTIMIZATION PERSPECTIVE

While certain invariances may hold in the presence of unobserved confounding, we have in general no guarantees on performance under more general manipulations. In this section we motivate a relaxation of the ideas presented above by considering bounds on the worst-case performance on data arising from distribution shifts informed by the structure of available environments.

Optimizing for the worst case loss in a set of domains has been shown to ensure accurate prediction on any convex mixture of training environments (Ben-Tal et al., 2009). The space of convex mixtures, however, is restrictive. Systems of variables are in general high-dimensional and new manipulations likely occur at a new vertex not represented as a linear combination of training environments. By *extrapolation* we desire performance guarantees outside this convex hull. The extension we consider optimizes instead over an *affine* combination of training losses, similarly to (Krueger et al., 2020).

Let $\Delta_\alpha := \{\{\alpha_e\}_{e\in\mathcal{E}} : \alpha_e \geq -\alpha, \sum_{e\in\mathcal{E}} \alpha_e = 1\}$ be a collection of scalars and consider the set of distributions defined by $\mathcal{P} := \{\sum_{e\in\mathcal{E}} \alpha_e P_e : \{\alpha_e\} \in \Delta_\alpha\}$, all affine combinations of distributions defined by the available environments. $\alpha \in \mathbb{R}$ defines the strength of the extrapolation, $\alpha = 0$ corresponds to a convex hull of distributions but above that value the space of distributions is richer, going beyond what has been observed: affine combinations amplify the strength of manipulations

that generated the observed training environments. The following theorem presents an upperbound to the robust problem (1) with affine combinations of errors.

Theorem 1 *Let $\{P_e\}_{e\in\mathcal{E}}$, be a set of available environments. Further let the parameter space of β be open and bounded, such that the expected loss function $\mathcal{L}$ as a function of β belongs to a Sobolev space. Then, the following inequality holds,*

$$\sup_{\{\alpha_e\}\in\Delta_\alpha} \sum_{e\in\mathcal{E}} \alpha_e \mathop{\mathbb{E}}_{(x,y)\sim P_e} \mathcal{L}(f\circ\phi(x),y) \leq \mathop{\mathbb{E}}_{(x,y)\sim P_e, e\sim\mathcal{E}} \mathcal{L}(f\circ\phi(x),y)$$

$$+ (1+n\alpha)\cdot C \cdot \left\| \sup_{e\in\mathcal{E}} \mathop{\mathbb{E}}_{(x,y)\sim P_e} \nabla_\beta \mathcal{L}(f\circ\phi(x),y) - \mathop{\mathbb{E}}_{(x,y)\sim P_e, e\sim\mathcal{E}} \nabla_\beta \mathcal{L}(f\circ\phi(x),y) \right\|_{L_2}$$

where $\|\cdot\|_{L_2}$ denotes the L_2-norm, C depends on the domain of β, $n := |\mathcal{E}|$ is the number of available environments and $e\sim\mathcal{E}$ loosely denotes sampling indeces with equal probability from $\mathcal{E}$.

The proof is given in Appendix A.3.2. The tightness of the inequality is given by the constant C which depends on the geometry of the parameter space and loss function and is intractable in general. Interestingly however, this bound illustrates the trade-off between invariances (of the type explored under unobserved confounding) and prediction in-sample. A combination of them exhibits robustness to certain shifts in distributions, and depending how much we weight each objective (prediction versus invariance) we can expect solutions to be more or less robust. Specifically, for $\alpha = -1/n$ the objective reduces to empirical risk minimization, but otherwise the upperbound increasingly weights differences in loss derivatives (violations of the invariances of section 2.2), and in the limit ($\alpha \to \infty$) can be interpreted to be robust at least to *any* affine combination of training losses.

3.1 PROPOSED OBJECTIVE

Our proposal is to guide the optimization of ϕ and β towards solutions that minimize the upperbound in Theorem 1, satisfying approximately the moment conditions (5) and simultaneously optimizing for minimum average error. Using Lagrange multipliers we define the general objective,

$$\mathop{\text{minimize}}_{\beta,\phi} \mathop{\mathbb{E}}_{(x,y)\sim P_e, e\sim\mathcal{E}} \mathcal{L}(f\circ\phi(x),y) + \lambda\cdot \mathop{\text{Var}}_{e\sim\mathcal{E}} \left(\| \mathop{\mathbb{E}}_{(x,y)\sim P_e} \nabla_\beta \mathcal{L}(f\circ\phi(x),y) \|_{L_2} \right) \qquad (6)$$

where $\lambda \geq 0$. This objective shares similarities with the objective proposed in (Krueger et al., 2020). The authors considered enforcing equality in environment-specific losses, rather than derivatives, as regularization, which can also be related to a robust optimization problem over an affine combination of errors. We have seen in section 2.2 however that equality in losses is not expected to hold in the presence of unobserved confounders (e.g. due to changes in the irreducible error across environments, which may occur also after interventions on target variables).

3.2 ROBUSTNESS IN TERMS OF INTERVENTIONS

As is apparent in Theorem 1, performance guarantees on data from a new environment depend on the relationship of new distributions with those observed during training.

Let $f\circ\phi_{\lambda\to\infty}$ minimize $\mathcal{L}$ among all functions that satisfy all pairs of moment conditions defined in (5); that is, a solution to our proposed objective in (6) with $\lambda\to\infty$. At optimality, it holds that gradients evaluated at this solution are equal across environments. As a consequence of Theorem 1, the loss evaluated at this solution with respect to *any* affine combination of environments is bounded by the average loss computed in-sample (denoted L, say),

$$\sum_{e\in\mathcal{E}} \alpha_e \mathop{\mathbb{E}}_{(x,y)\sim P_e} \mathcal{L}(f\circ\phi(x),y) \leq L, \qquad \text{for any set of } \alpha_e \in \Delta_\alpha \qquad (7)$$

From the perspective of interventions in the underlying causal mechanism, this can be seen as a form of data-driven predictive stability across a range of distributions whose perturbations occur in the same direction as those observed during training. As a minimal example for intuition, consider a system of three variables (X, Y, H), with differing interventions on X only. Assume we have access to data sampled under two environments defined by joint distributions $p_1(X,Y,H) := p(Y,H|X)p_1(X)$ and $p_2(X,Y,H) := p(Y,H|X)p_2(X)$. Errors are bounded with respect to any (valid) distribution of

the form $p(Y, H|X)(\alpha_1 p_1(X) + \alpha_2 p_2(X))$. For instance, a particular shift in the mean or variance of X observed during training can be extrapolated in the extreme to *any* shift in the mean or variance of new data. With this reasoning, if environment-specific distributions arise from differing interventions on all observed covariates X, then solutions are robust to arbitrary shifts in these variables. If, in addition, solutions are unique we may interpret them as causal, irrespective of the presence or not of unobserved confounders. The generalization properties, however, go further. Interventions on unobserved variables and also target variables are accommodated for in $f \circ \phi_{\lambda\to\infty}$, if observed through shifted distributions in different available environments.

Using our simple example in Figure 1 to verify this intuition empirically, we consider 3 scenarios corresponding to interventions on exogenous variables of X, H and Y. In each, training data from two environments is generated with means in the distribution of the concerned variables set to a value of 0 and 1 respectively, everything else being equal ($\sigma^2 := 1, H := E_H \sim \mathcal{N}(0,1)$). Performance is evaluated on out-of-sample data generated by increasing the shift in the variable being studied up to a mean of 5. In all cases, we see in Figure 2 that performance is stable to increasing perturbations in the system as long as the heterogeneity in the data allows us to capture the direction of the unseen shift.

Figure 2: Stability to general shifts.

3.3 STABILITY OF CERTAIN OPTIMAL SOLUTIONS

A special case may also be considered when the underlying system of variables and the available environments allow for optimal solutions $f \circ \phi_{\lambda\to\infty}$ and $f \circ \phi_{\lambda=0}$ to coincide. In this case, the learned representation $\phi(x)$ results in a predictor f optimal on average *and* simultaneously with equal gradient in each environment, thus,

$$||\mathop{\mathbb{E}}_{(x,y)\sim P_e} \nabla_\beta \mathcal{L}(f \circ \phi(x), y)||_{L_2} = 0, \qquad \text{for all } e \in \mathcal{E} \tag{8}$$

For this representation ϕ, it follows that optimal solutions f learned on any new dataset sampled from an affine combination of training distributions coincides with this special solution. This gives us a sense of reproducibility of learning. In other words, if a specific feature is significant for predictions on the whole range of λ with the available data then it will likely be significant on new (related) data.

The above special case where all solutions in our hyperparameter range agree has important parallels with IRM (Arjovsky et al., 2019). The authors proposed a learning objective enforcing representations of data with minimum error on average and across environments, such that at optimum $\mathbb{E}_{P_i} Y|\phi^\star(X) = \mathbb{E}_{P_j} Y|\phi^\star(X)$ for any pair $(i, j) \in \mathcal{E}$. With unobserved confounding, both learning paradigms agree but, with unobserved confounding, minimum error solutions of IRM by design converge to spurious associations (see the discussion after equation (5)) and are not guaranteed to generalize to more general environments. For example, in the presence of additive unobserved confounding H, irrespective of ϕ, we may have $\mathbb{E}_{P_i} Y|\phi^\star(X) = \phi^\star(X) + \mathbb{E}_{P_i} H \neq \phi^\star(X) + \mathbb{E}_{P_j} H = \mathbb{E}_{P_j} Y|\phi^\star(X)$ if the means of H differ. The sought invariance then does not hold.

4 RELATED WORK

Causality. There has been a growing interest in interpreting shifts in distribution to fundamentally arise from interventions in the causal mechanisms of data. Peters et al. (Peters et al., 2016) exploited this link for causal inference: causal relationships by definition being invariant to the observational regime. Invariant solutions, as a result of this connection, may be interpreted also as robust to certain interventions (Meinshausen, 2018), and recent work has explored learning invariances in various problem settings (Arjovsky et al., 2019; Rothenhäusler et al., 2019; Krueger et al., 2020; Gimenez & Zou, 2020). Among those, we note the invariance proposed in (Rothenhäusler et al., 2019), the authors seek to recover causal solutions with unobserved confounding. Generalization properties of these solutions were rarely studied, with one exception being Anchor regression (Rothenhäusler et al., 2018). The authors proposed to interpolate between empirical risk minimization and causal solutions with explicit robustness to certain interventions in a linear model. The present work may be interpreted as a non-linear formulation of this principle with a more general study of generalization.

Domain generalization represent one direction of out-of-sample generalization by explicitly learning representations projecting out superficial environment-specific information. Recent work on domain generalization has included the use data augmentation (Volpi et al., 2018; Shankar et al., 2018), meta-learning to simulate domain shift (Li et al., 2018) and adversarially learning representations that are environment invariant (Ganin et al., 2016), even though explicitly aligning representations has important caveats when label distributions differ, articulated for instance in (Arjovsky et al., 2019).

Distributionally robust optimization explicitly solves a worst-case optimization problem (1). A popular approach is to define $\mathcal{P}$ as a ball around the empirical distribution $\hat{P}$, for example using f-divergences or Wasserstein balls of a defined radius (Kuhn et al., 2019; Duchi et al., 2016; 2019; Sinha et al., 2017; Wozabal, 2012; Abadeh et al., 2015; Duchi & Namkoong, 2018). These are general and multiple environments are not required, but this also means that sets are defined agnostic to the geometry of plausible shifted distributions, and may therefore lead to solutions, when tractable, that are overly conservative or do not satisfy generalization requirements (Duchi et al., 2019).

5 EXPERIMENTS

Data linkages, electronic health records, and bio-repositories, are increasingly being collected to inform medical practice. As a result, also prediction models derived from healthcare data are being put forward as potentially revolutionizing decision-making in hospitals. Recent studies (Cabitza et al., 2017; Venugopalan et al., 2019), however, suggest that their performance may reflect not only their ability to identify disease-specific features, but also their ability to exploit spurious correlations due to unobserved confounding (such as varying data collection practices): a major challenge for the reliability of decision support systems. In this section, we explore this pattern conducting a wide analysis of domain generalization on image, speech and tabular data from the medical domain.

Baseline algorithms. Comparisons are made with **ERM**, optimizing for minimum loss agnostic of data source. Domain Robust Optimization (**DRO**) (Sagawa et al., 2019), optimizing for minimum loss across the worst convex mixture of training environments. Domain adversarial training (**DANN**) (Ganin et al., 2016), using domain adversarial training to facilitate transfer by augmenting the neural network architecture with an additional domain classifier to enforce the distribution of $\phi(X)$ to be the same across training environments. Invariant Risk Minimization (**IRM**) (Arjovsky et al., 2019), that regularizes ERM ensuring representations $\phi(X)$ be optimal in every observed environment. All trained models use the same convolutional or fully-connected architecture, where appropriate. Performance results are given in Table 1. Further experimental details can be found in Appendix A.4.

5.1 DIAGNOSIS OF PNEUMONIA WITH CHEST X-RAY DATA

In this section, we attempt to replicate the study in (Zech et al., 2018). The authors observed a tendency of image models towards exploiting spurious correlations for the diagnosis on pneumonia from patient Chest X-rays that do not reproduce outside of training data. We use publicly available data from the National Institutes of Health (NIH) (Wang et al., 2017) and the Guangzhou Women and Children's Medical Center (GMC) (Kermany et al., 2018). Differences in distribution are apparent, for instance in the top edge of mean pneumonia-diagnosed X-rays shown in Figure 3.

In this experiment, we exploit this (spurious) pathology correlation to demonstrate the need for solutions robust to changes in site-specific features. We construct two training sets, in each case 90% and 80% of pneumonia-diagnosed patients were drawn from the NIH dataset and the remaining 10% and 20% of the pneumonia-diagnosed patients were drawn from the GMC dataset; the reverse logic (10%/90% split) was followed for the test set.

Figure 3: Average pneumonia X-ray.

5.2 DIAGNOSIS OF PARKINSON'S DISEASE WITH VOICE RECORDINGS

Parkinson's disease is a progressive nervous system disorder that affects movement. Symptoms start gradually, sometimes starting with a barely noticeable tremor in a patient's voice. This section investigates the performance of predictive models for the detection of Parskinson's disease, trained on voice recordings of a specific type (vowels and numbers) and tested on more involved forms of speech, such as words and short sentences, in new patients. We used the UCI Parkinson Speech Dataset with given training and testing splits (Sakar et al., 2013). Even though the distributions of

	Pneumonia Prediction		Parkinson Prediction		Survival Prediction	
	Training	**Testing**	**Training**	**Testing**	**Training**	**Testing**
ERM	91.4 (± .7)	52.4 (± 1)	95.7 (± .5)	62.9 (± 1)	93.2 (± .4)	75.3 (± .9)
DRO	91.2 (± .5)	53.1 (± .6)	94.0 (± .3)	69.8 (± 2)	90.5 (± .4)	75.5 (± .8)
DANN	92.3 (± 1)	57.1 (± 2)	91.6 (± 2)	51.4 (± 5)	89.3 (± .8)	73.9 (± .9)
IRM	89.5 (± 1)	58.6 (± 2)	93.6 (± 1)	71.4 (± 2)	91.9 (± .6)	75.7 (± .8)
Ours	84.3 (± 1)	63.7 (± 3)	93.1 (± 2)	72.8 (± 2)	91.4 (± .6)	77.9 (± 1)

Table 1: Accuracy of predictions in percentages (%). Uncertainty intervals are standard deviations. All datasets are approximately balanced, 50% performance is as good as random guessing.

features will differ in different types of recordings, we would expect the underlying patterns in speech to reproduce across different samples. However, this is not the case for correlations learned with baseline training paradigms (Table 1). This suggests that spurious correlations due to the specific type of recording, and in this case emphasized due to low sample sizes (120 examples in each training and testing sets here), may be responsible for poor generalization error. We found similar results even when tested on different recording of the same patients the model was trained on.

5.3 SURVIVAL PREDICTION WITH ELECTRONIC HEALTH RECORDS

This section investigates whether predictive models transfer across a collection of datasets from 30 different medical studies containing patients that experienced heart failure, the MAGGIC studies (MAGGIC, 2012). We do so with a twist however, explicitly introducing unobserved confounding by omitting predictive variables and selecting studies for testing where precisely the distribution of these variables happens to be different than those in training studies. The objective is to test performance on shifted distributions, where shifts occur predominantly due to variability in unobserved variables.

The problem is to predict survival within 5 years of experiencing heart failure from a total of 33 demographic variables. To construct confounded data, we start by running logistic regression, yielding significant influence of a patient's age on its risk of death, a feature also correlated with a number of other variables including blood pressure and body mass index. We chose 5 studies of over 500 examples with balanced death rates and different age distributions (age subsequently omitted to induce unobserved confounding). Performance results are averaged over 5 experiments (using every combination of 4 studies for training and the remaining study for testing).

Influential variables that reproduce across datasets. Recall that in the special case that the optimal predictor is invariant as a function $\lambda \in [0, \infty)$, optimal predictors estimated in new perturbed datasets in the span of observed shifted distributions, should be stable.

We consider here a form of diluted stability for feature selection. For a single layer network, we consider significant those covariates with estimated parameters bounded away from zero in all solutions in the range $\lambda \in [0, 1]$. Comparisons are made with ERM (conventional logistic regression), both methods trained separately on 100 random pairs of studies. Figure 4 shows how many features (in the top 10 of predictive features) from each model intersect across pairs of studies. In constrast to ERM, our objective recovers significant features more consistently.

Figure 4: Reproducible features.

6 CONCLUSIONS

We have studied the problem of out-of-sample generalization from a new perspective, grounded in the underlying causal mechanism generating new data that may arise from shifts in observed, unobserved or target variables. Our proposal is a new objective that is provably robust to certain shifts in distribution, and is informed by certain statistical invariances in the presence of unobserved confounders. Our experiments show that we may expect better generalization performance and also better reproducibility of influential features in problems of variable selection. A limitation of our approach is that robustness guarantees crucially depend on the (unobserved) properties of available data. Our hope nevertheless is to pave the way for exploring new connections between causality and machine learning that may lead to better behaved learning agents acting in the world.

REFERENCES

Soroosh Shafieezadeh Abadeh, Peyman Mohajerin Mohajerin Esfahani, and Daniel Kuhn. Distributionally robust logistic regression. In *Advances in Neural Information Processing Systems*, pp. 1576–1584, 2015.

Martin Arjovsky, Léon Bottou, Ishaan Gulrajani, and David Lopez-Paz. Invariant risk minimization. *arXiv preprint arXiv:1907.02893*, 2019.

Aharon Ben-Tal, Laurent El Ghaoui, and Arkadi Nemirovski. *Robust optimization*, volume 28. Princeton University Press, 2009.

Steffen Bickel, Michael Brückner, and Tobias Scheffer. Discriminative learning under covariate shift. *Journal of Machine Learning Research*, 10(Sep):2137–2155, 2009.

Federico Cabitza, Raffaele Rasoini, and Gian Franco Gensini. Unintended consequences of machine learning in medicine. *Jama*, 318(6):517–518, 2017.

Raymond J Carroll, David Ruppert, Leonard A Stefanski, and Ciprian M Crainiceanu. *Measurement error in nonlinear models: a modern perspective*. CRC press, 2006.

John Duchi and Hongseok Namkoong. Learning models with uniform performance via distributionally robust optimization. *arXiv preprint arXiv:1810.08750*, 2018.

John Duchi, Peter Glynn, and Hongseok Namkoong. Statistics of robust optimization: A generalized empirical likelihood approach. *arXiv preprint arXiv:1610.03425*, 2016.

John C Duchi, Tatsunori Hashimoto, and Hongseok Namkoong. Distributionally robust losses against mixture covariate shifts. *Under review*, 2019.

Wayne A Fuller. *Measurement error models*, volume 305. John Wiley & Sons, 2009.

Yaroslav Ganin, Evgeniya Ustinova, Hana Ajakan, Pascal Germain, Hugo Larochelle, François Laviolette, Mario Marchand, and Victor Lempitsky. Domain-adversarial training of neural networks. *The Journal of Machine Learning Research*, 17(1):2096–2030, 2016.

AmirEmad Ghassami, Saber Salehkaleybar, Negar Kiyavash, and Kun Zhang. Learning causal structures using regression invariance. In *Advances in Neural Information Processing Systems*, pp. 3011–3021, 2017.

Jaime Roquero Gimenez and James Zou. Identifying invariant factors across multiple environments with kl regression. *arXiv preprint arXiv:2002.08341*, 2020.

Daniel S Kermany, Michael Goldbaum, Wenjia Cai, Carolina CS Valentim, Huiying Liang, Sally L Baxter, Alex McKeown, Ge Yang, Xiaokang Wu, Fangbing Yan, et al. Identifying medical diagnoses and treatable diseases by image-based deep learning. *Cell*, 172(5):1122–1131, 2018.

David Krueger, Ethan Caballero, Joern-Henrik Jacobsen, Amy Zhang, Jonathan Binas, Remi Le Priol, and Aaron Courville. Out-of-distribution generalization via risk extrapolation (rex). *arXiv preprint arXiv:2003.00688*, 2020.

Daniel Kuhn, Peyman Mohajerin Esfahani, Viet Anh Nguyen, and Soroosh Shafieezadeh-Abadeh. Wasserstein distributionally robust optimization: Theory and applications in machine learning. In *Operations Research & Management Science in the Age of Analytics*, pp. 130–166. INFORMS, 2019.

Giovanni Leoni. *A first course in Sobolev spaces*. American Mathematical Soc., 2017.

Da Li, Yongxin Yang, Yi-Zhe Song, and Timothy M Hospedales. Learning to generalize: Meta-learning for domain generalization. In *Thirty-Second AAAI Conference on Artificial Intelligence*, 2018.

MAGGIC. The survival of patients with heart failure with preserved or reduced left ventricular ejection fraction: an individual patient data meta-analysis. *European heart journal*, 33(14): 1750–1757, 2012.

Nicolai Meinshausen. Causality from a distributional robustness point of view. In *2018 IEEE Data Science Workshop (DSW)*, pp. 6–10. IEEE, 2018.

Karthika Mohan, Judea Pearl, and Jin Tian. Graphical models for inference with missing data. In *Advances in neural information processing systems*, pp. 1277–1285, 2013.

Judea Pearl. Why there is no statistical test for confounding, why many think there is, and why they are almost right. 1998.

Judea Pearl. *Causality*. Cambridge university press, 2009.

Jonas Peters, Peter Bühlmann, and Nicolai Meinshausen. Causal inference by using invariant prediction: identification and confidence intervals. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 78(5):947–1012, 2016.

Dominik Rothenhäusler, Nicolai Meinshausen, Peter Bühlmann, and Jonas Peters. Anchor regression: heterogeneous data meets causality. *arXiv preprint arXiv:1801.06229*, 2018.

Dominik Rothenhäusler, Peter Bühlmann, Nicolai Meinshausen, et al. Causal dantzig: fast inference in linear structural equation models with hidden variables under additive interventions. *The Annals of Statistics*, 47(3):1688–1722, 2019.

Shiori Sagawa, Pang Wei Koh, Tatsunori B Hashimoto, and Percy Liang. Distributionally robust neural networks for group shifts: On the importance of regularization for worst-case generalization. *arXiv preprint arXiv:1911.08731*, 2019.

Betul Erdogdu Sakar, M Erdem Isenkul, C Okan Sakar, Ahmet Sertbas, Fikret Gurgen, Sakir Delil, Hulya Apaydin, and Olcay Kursun. Collection and analysis of a parkinson speech dataset with multiple types of sound recordings. *IEEE Journal of Biomedical and Health Informatics*, 17(4): 828–834, 2013.

Shiv Shankar, Vihari Piratla, Soumen Chakrabarti, Siddhartha Chaudhuri, Preethi Jyothi, and Sunita Sarawagi. Generalizing across domains via cross-gradient training. *arXiv preprint arXiv:1804.10745*, 2018.

Aman Sinha, Hongseok Namkoong, and John Duchi. Certifying some distributional robustness with principled adversarial training. *arXiv preprint arXiv:1710.10571*, 2017.

Subhashini Venugopalan, Arunachalam Narayanaswamy, Samuel Yang, Anton Gerashcenko, Scott Lipnick, Nina Makhortova, James Hawrot, Christine Marques, Joao Pereira, Michael Brenner, et al. It's easy to fool yourself: Case studies on identifying bias and confounding in bio-medical datasets. *arXiv preprint arXiv:1912.07661*, 2019.

Riccardo Volpi, Hongseok Namkoong, Ozan Sener, John C Duchi, Vittorio Murino, and Silvio Savarese. Generalizing to unseen domains via adversarial data augmentation. In *Advances in Neural Information Processing Systems*, pp. 5334–5344, 2018.

Xiaosong Wang, Yifan Peng, Le Lu, Zhiyong Lu, Mohammadhadi Bagheri, and Ronald M Summers. Chestx-ray8: Hospital-scale chest x-ray database and benchmarks on weakly-supervised classification and localization of common thorax diseases. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pp. 2097–2106, 2017.

Daniel Westreich, Jessie K Edwards, Stephen R Cole, Robert W Platt, Sunni L Mumford, and Enrique F Schisterman. Imputation approaches for potential outcomes in causal inference. *International journal of epidemiology*, 44(5):1731–1737, 2015.

David Wozabal. A framework for optimization under ambiguity. *Annals of Operations Research*, 193 (1):21–47, 2012.

John R Zech, Marcus A Badgeley, Manway Liu, Anthony B Costa, Joseph J Titano, and Eric K Oermann. Confounding variables can degrade generalization performance of radiological deep learning models. *arXiv preprint arXiv:1807.00431*, 2018.

A APPENDIX

A.1 EXAMPLES OF UNOBSERVED CONFOUNDING

Measurement error. The data generating processes described in the main body of this paper for instance, as well as most of machine learning, assume that all nuisance variability enters the causal mechanisms of the data; that is, observed data reflects *only* causal drivers. If this is not the case, for example because of independent measurement noise observed in data but that does not propagate to across causal children, regression is known to be inconsistent in general (Carroll et al., 2006) and its bias is analogous to a form of unobserved confounding.

Consider a simple model for illustration. Suppose (X, Y) are observed subject to measurement noise, $X^\star = X + E_x$ and $Y^\star = Y + E_y$, which are not causally related to one another but rather $Y = \beta X + E$. Let $E_x = \beta_x H$ and $E_y = \beta_y H$ be the structure of measurement error independent of X and Y. Then substituting our observed data $(X^\star, Y^\star)$ into the underlying (X, Y) relationship the observed model is,

$$Y^\star = \beta X^\star + (\beta_y - \beta_x \beta)H + E, \quad X^\star = \beta_x H + X \tag{9}$$

A special case of regression with unobserved confounders H.

Missing data in observed variables. Data observed with missing values in causal covariates, if missing not at random, will similarly introduce bias in the estimation of the relationship between variables that may also be related to a special case of unobserved confounders in the underlying data generating mechanism (see for instance section 2.1 in (Mohan et al., 2013) or (Westreich et al., 2015)).

A.2 FURTHER EXPERIMENTS TO ILLUSTRATE CONFOUNDING BIAS

Recall the data generating mechanism presented in the introductory example in Figure 1 of the main body of this paper. We have access to observations of variables (X_1, X_2, Y) in two training datasets, each dataset sampled with differing variances ($\sigma^2 = 1$ and $\sigma^2 = 2$) from the following structural model $\mathbb{F}$,

$$X_2 := -H + E_{X_2}, \quad Y := X_2 + 3H + E_Y, \quad X_1 := Y + X_2 + E_{X_1} \quad H := E_H$$

where $E_{X_1}, E_{X_2} \sim \mathcal{N}(0, \sigma^2)$, $E_Y \sim \mathcal{N}(0, 1)$, $E_H \sim \mathcal{N}(0, 1)$ are exogenous variables. H is an unobserved confounder, not observed during training but that does play an important role in the underlying data generation mechanism.

Our introductory example considered predictive performance under different data distributions. Here we inspect the weights learned in a simple one layer feed-forward neural network to infer whether unobserved confounding induces a given learning paradigm to exploit spurious correlations. We consider training with Empirical Risk Minimization (ERM), Invariant Risk Minimization (IRM) (Arjovsky et al., 2019) and training by equating moment conditions as described in Section 2.2, that is optimizing our objective for $\lambda \to \infty$. The results are given in Table 2 below. With regards to IRM, these results demonstrate that seeking invariant predictors to approximate causal solutions in the presence of unobserved confounding is not appropriate. Our invariance principle by contrast does approximately recover causal solutions asymptotically in λ.

	Truth	**ERM**	**IRM**	**Ours** ($\lambda \to \infty$)
Estimated parameters	[0, 1]	[0.91, -1.02]	[0.75, -0.76]	[0.01, 0.95]

Table 2: Bias in estimation due to unobserved confounders.

A.3 TECHNICAL RESULTS

A.3.1 INVARIANCES IN THE PRESENCE OF UNOBSERVED CONFOUNDING

Assume data sources emerge from manipulations in exogenous E_X, related to X only, in an underlying additive model $\mathbb{F}$ with also additive functions f_1, f_2, f_3, f_4, such as,

$$Y := f_1(X) + f_2(H) + E_Y, \qquad X := f_3(X) + f_4(H) + E_X, \qquad H := E_H \tag{10}$$

each exogenous variables (E_X, E_Y, E_H) with arbitrary distributions but only E_X varying across environments. Then if holds that $X = (I - f_3)^{-1} f_4(H) + (I - f_3)^{-1} E_X$ and that $\nabla_\beta f_1(X)(Y - f_1(X)) = (\nabla_\beta f_1(I - f_3)^{-1} f_4(H) + \nabla_\beta f_1(I - f_3)^{-1} E_X) \cdot (f_2(H) + E_Y)$. Since (E_X, E_Y, E_H) are mutually independent it follows that expectations of products of functions of these random variables are zero; that is the expectation of the inner product $\nabla_\beta f_1(X)(Y - f_1(X))$ does not depend on E_X and is thus stable across environments.

A.3.2 PROOF OF THEOREM 1

We restate the Theorem for convenience.

Theorem 1 *Let $\{P_e\}_{e \in \mathcal{E}}$, be a set of available environments. Further let the parameter space of β be open and bounded, such that the expected loss function $\mathcal{L}$ as a function of β belongs to a Sobolev space. Then, the following inequality holds,*

$$\sup_{\alpha_e \in \Delta_\alpha} \sum_{e \in \mathcal{E}} \alpha_e \mathop{\mathbb{E}}_{(x,y) \sim P_e} \mathcal{L}\left(f \circ \phi(x), y\right) \leq \mathop{\mathbb{E}}_{(x,y) \sim P_e, e \sim \mathcal{E}} \mathcal{L}\left(f \circ \phi(x), y\right)$$

$$+ (1 + n\alpha) \cdot C \cdot \left\| \sup_{e \in \mathcal{E}} \mathop{\mathbb{E}}_{(x,y) \sim P_e} \nabla_\beta \mathcal{L}\left(f \circ \phi(x), y\right) - \mathop{\mathbb{E}}_{(x,y) \sim P_e, e \sim \mathcal{E}} \nabla_\beta \mathcal{L}\left(f \circ \phi(x), y\right) \right\|_{L_2}$$

where C depends on the domain of β, $n := |\mathcal{E}|$ is the number of available environments and $e \sim \mathcal{E}$ loosely denotes sampling indeces with equal probability from $\mathcal{E}$.

Proof. Let Ω denote the parameter space of β. The following derivation shows the claim,

$$\sup_{\alpha_e \in \Delta_\alpha} \sum_{e \in \mathcal{E}} \alpha_e \mathop{\mathbb{E}}_{(x,y) \sim P_e} \mathcal{L}\left(f \circ \phi(x), y\right)$$

$$= (1 + n\alpha) \cdot \sup_{e \in \mathcal{E}} \mathop{\mathbb{E}}_{(x,y) \sim P_e} \mathcal{L}\left(f \circ \phi(x), y\right) - \alpha \sum_{e \sim \mathcal{E}} \mathbb{E}_{P_e} \mathcal{L}\left(f \circ \phi(x), y\right)$$

$$= \mathop{\mathbb{E}}_{(x,y) \sim P_e, e \sim \mathcal{E}} \mathcal{L}\left(f \circ \phi(x), y\right) + (1 + n\alpha) \cdot \sup_{e \in \mathcal{E}} \mathop{\mathbb{E}}_{(x,y) \sim P_e} \mathcal{L}\left(f \circ \phi(x), y\right)$$

$$- (\alpha + 1/n) \sum_{e \sim \mathcal{E}} \mathop{\mathbb{E}}_{(x,y) \sim P_e} \mathcal{L}\left(f \circ \phi(x), y\right)$$

$$= \mathop{\mathbb{E}}_{(x,y) \sim P_e, e \sim \mathcal{E}} \mathcal{L}\left(f \circ \phi(x), y\right)$$

$$+ (1 + n\alpha) \cdot \left(\sup_{e \in \mathcal{E}} \mathop{\mathbb{E}}_{(x,y) \sim P_e} \mathcal{L}\left(f \circ \phi(x), y\right) - \mathop{\mathbb{E}}_{(x,y) \sim P_e, e \sim \mathcal{E}} \mathcal{L}\left(f \circ \phi(x), y\right) \right)$$

$$\leq \mathop{\mathbb{E}}_{(x,y) \sim P_e, e \sim \mathcal{E}} \mathcal{L}\left(f \circ \phi(x), y\right)$$

$$+ (1 + n\alpha) \cdot M \cdot \left\| \sup_{e \in \mathcal{E}} \mathop{\mathbb{E}}_{(x,y) \sim P_e} \mathcal{L}\left(f \circ \phi(x), y\right) - \mathop{\mathbb{E}}_{(x,y) \sim P_e, e \sim \mathcal{E}} \mathcal{L}\left(f \circ \phi(x), y\right) \right\|_{L_2}$$

where the inequality is given by the property that the evaluation functional is a bounded linear operator in certain Sobolev spaces $\mathcal{W}$, for example with $\Omega = \mathbb{R}^d$ and L_2 norm. In particular this means that $|f(\beta)| \leq M||f||_{L_2}$ for all $f \in \mathcal{W}$. It follows then also that the above is,

$$\leq \mathop{\mathbb{E}}_{(x,y) \sim P_e, e \sim \mathcal{E}} \mathcal{L}\left(f \circ \phi(x), y\right) +$$

$$(1 + n\alpha) \cdot P \cdot M \cdot \left\| \sup_{e \in \mathcal{E}} \mathop{\mathbb{E}}_{(x,y) \sim P_e} \nabla_\beta \mathcal{L}\left(f \circ \phi(x), y\right) - \mathop{\mathbb{E}}_{(x,y) \sim P_e, e \sim \mathcal{E}} \nabla_\beta \mathcal{L}\left(f \circ \phi(x), y\right) \right\|_{L_2}$$

by Poincaré's inequality for Sobolev functions defined on an open, bounded parameter space, see e.g. (Leoni, 2017). The assumption we make here for this last inequality to hold is that the region where the difference in loss functions is near zero is large enough such that the integral of the gradient is also large enough to control the integral of the function. This holds however for functions defined on many "reasonable" parameter spaces (Lipschitz suffices).

A.4 EXPERIMENTAL DETAILS

X-ray Data

After resizing and creating training environments with different proportions of X-rays from our two

hospital sources such that that the positive and negative labels remained balanced, we were left with 2002 samples are available in each training environment and 1144 samples in the testing set.

All learning paradigms trained a CNN model, 2 layers deep, with each layer consisting of a convolution (kernel size of 3 and a stride of 1). All predictions were made off of the deepest layer of the network. The number of input channels was 64, doubled for each subsequent layer, and dropout was applied after each layer. We optimize the binary cross-entropy loss using Adam (learning rate 0.001) without further regularization on parameters and use Xavier initialization. While learning with IRM and the proposed approach, the respective penalty $\lambda = 1$ is added to the loss after 5 epochs of learning with $\lambda = 0$. Experiments are run for a maximum of 20 epochs with early stopping based on validation performance. All results are averaged over 10 trials with different random splits of the data.

Parkinson's Disease Speech Data
The data includes a total of 26 features recorded on each sample of speech and set training and testing splits which we use in our experiments. For each patient 26 different voice samples including sustained vowels, numbers, words and short sentences where recorded, which we considered as heterogeneous sources. We created three training environments by concatenating features from three number recordings, concatenating features from three word recording and concatenating features from three sentences; culminating in a total of 120 samples in each training environment. The available testing split contained recordings of vowels for a total of 168 samples. Positive and negative samples were balanced in both training and testing environments. We investigated other choices for our training and testing data splits, for which we didn't not observe a qualitative change in our conclusions.

On this data, for all learning paradigms we train a MLP with two hidden layers of size 64 with tanh activations and dropout ($p = 0.5$) after each layer. As in the image experiments, we optimize the binary cross-entropy loss using Adam (learning rate 0.001) with L_2 regularization on parameters and use Xavier initialization. While learning with IRM and the proposed approach, the respective penalty is added to the loss $\lambda = 1$ after 200 epochs of learning with $\lambda = 0$ to ensure stable optimization. Experiments are run for a maximum of 1000 epochs with early stopping based on the validation performance. All results are averaged over 10 trials with different random seeds.

MAGGIC Electronic Health Records
The same architecture and hyperparameters as in Parkinson's disease speech data experiments was used for MAGGIC data except that we increase the maximum training epochs to 5000. We preprocessed the data such that each feature is scaled to a $[0, 1]$ interval. After removing the confounder "age", selecting for balanced death rates and restricting ourselves to studies with more than 500 patients, 5 studies remained. Each was chosen in turn as a target environment with models trained on the other 4 training environments.

For the feature reproducibility experiments we give a bit more context for clarity and completeness. A natural objective for the consistency of health care and such that we may reproduce the experiments and their results in different scenarios is to find relevant features that are not specific to an individual medical study, but can also be found (replicated) on other studies with different patients. Heterogeneous patients and studies, along with different national guidelines and standards of care make this challenging. In our experiments we made comparisons of reproducibility in parameter estimates for models trained using Empirical Risk Minimization (ERM) and our proposed approach. We chose networks with a single layer with logistic activation and focused on the estimation of parameter to understand the variability in training among different data sources. Naturally, feature importance measured by parameter magnitudes makes sense only after normalization of the covariates to the same (empirical) variance (equal to 1) in each study separately. After this preprocessing step, for both ERM and the proposed approach we trained separate networks on 100 random pairs of studies (each pair concatenated for ERM) and returned the top 10 significant features (by the magnitude of parameters). Over all sets of significant parameters we then identified how many intersected across a fixed number of the 100 runs.