What's the Harm? Sharp Bounds on the Fraction Negatively Affected by Treatment

Abstract

The fundamental problem of causal inference – that we never observe counterfactuals – prevents us from identifying how many might be negatively affected by a proposed intervention. If, in an A/B test, half of users click (or buy, or watch, or renew, *etc.*), whether exposed to the standard experience A or a new one B, hypothetically it could be because the change affects no one, because the change positively affects half the user population to go from no-click to click while negatively affecting the other half, or something in between. While unknowable, this impact is clearly of material importance to the decision to implement a change or not, whether due to fairness, long-term, systemic, or operational considerations. We therefore derive the *tightest-possible* (*i.e.*, sharp) bounds on the fraction negatively affected (and other related estimands) given data with only factual observations, whether experimental or observational. Naturally, the more we can stratify individuals by observable covariates, the tighter the sharp bounds. Since these bounds involve unknown functions that must be learned from data, we develop a robust inference algorithm that is efficient almost regardless of how and how fast these functions are learned, remains consistent when some are mislearned, and still gives valid conservative bounds when most are mislearned. Our methodology altogether therefore strongly supports credible conclusions: it avoids spuriously point-identifying this unknowable impact, focusing on the best bounds instead, and it permits exceedingly robust inference on these. We demonstrate our method in simulation studies and in a case study of career counseling for the unemployed.

1 Introduction

Before making changes to an online platform, product managers regularly conduct experiments ("A/B tests") to make sure a change does not negatively impact users' experience. Similarly, extensive program evaluations, whether experimental or observational, are a corner stone of evidence-based policymaking and help avoid harmful policy changes. These tests and evaluations focus on assessing the causal effect of a change on key metrics, whether user engagement or retention on online platforms or outcomes like employment for social programs. Average treatment effects, heterogeneous treatment effects, and distributional treatment effects on these outcomes quantify the quality of the intervention in terms of aggregate impacts on the population or on subpopulations. This is in its essence a statistical solution to the *fundamental problem of causal inference*: assess aggregate effects since the impact on any one individual can never be observed. We can never know whether a user that churned under arm "B" would have been retained under arm "A," but we can characterize the distribution of retention under each arm, even segmenting the population by observable features.

While unknowable, the impact of changes on individuals is clearly of material importance to the decision to implement a change or not. This impact can have downstream effects, whether user behavioral, reputational, or operational. Moreover, quantifying how negative this impact can be is crucial to understanding fairness with respect to the welfare of the individuals, rather than just focusing

Submitted to 36th Conference on Neural Information Processing Systems (NeurIPS 2022). Do not distribute.

on social welfare, the welfare of the decision-making entity, or the welfare of observable groups. Specifically, even if the average treatment effect is zero, there can still be a sizable subpopulation that is negatively affected by the change. That is, a change with zero average effect cannot earnestly be called harmless. In this paper we consider measuring the fraction that are negatively affected. The aim is to be able to flag changes that may individually harm many. Unfortunately this fraction cannot be identified from data, even experimental data. However, we may still be able to bound it.

In this paper, we derive the *sharp* bounds on the fraction of individuals negatively affected when outcomes are binary and develop tools to conduct robust inferences on these bounds. That is, we characterize the *tightest-possible* interval that contains all values of this unknowable quantity consistent with observable information. We consider both the fraction negatively affected under a wholesale change from one treatment regime to another as well as under an optimal policy that assigns each observable subpopulation the treatment with best average outcomes for the subpopulation. Then, we tackle how to actually assess these theoretical bounds based on actual data, namely, how to estimate the interval endpoints and construct confidence intervals (CIs) that account for sampling error, on top of the inherent epistemic uncertainty characterized by the bounds. The bounds involve potentially complex functions such as the conditional average treatment effect (CATE) function, so we develop inference methodology that is exceedingly robust to learning these functions. Namely, our inference works even when these are learned nonparametrically at slow rates, and our estimated bounds remain valid even when these are learned inconsistently. We demonstrate our tools in a simulation study and a case study of hypothetical changes to French reemployment assistance programs.

2   Problem Set Up and the Fraction Negatively Affected

The Population   Each individual in the population is associated with baseline covariates (observable characteristics), $X \in \mathcal{X}$, and two *binary* potential outcomes, $Y^*(0), Y^*(1) \in \{0, 1\}$, corresponding to each of two treatment options, 0 and 1. Examples include: two versions of product on an online platform and whether user is retained in the next quarter, or two versions of a job training program for the unemployed and whether the participant is reemployed within the following year.

The individual treatment effect (ITE) is the difference in potential outcomes,

$$\text{ITE} = Y^*(1) - Y^*(0) \in \{-1, 0, 1\}.$$

We assume that an outcome value of 1 corresponds to a better outcome (*e.g.*, retained, reemployed) and 0 to a worse outcome (*e.g.*, churned, unemployed). Thus, if treatment 0 represents status quo and treatment 1 represents a proposed change, then individuals with ITE = 0 ($Y^*(0) = Y^*(1) = 0$ or $Y^*(0) = Y^*(1) = 1$) are unaffected by the change, individuals with ITE = 1 ($Y^*(0) = 0, Y^*(1) = 1$) benefit from the change, and individuals with ITE = -1 ($Y^*(0) = 1, Y^*(1) = 0$) are harmed by the change. And, this list of potential-outcome combinations is exhaustive.

The Data   We consider data from either a randomized experiment or an observational study, wherein we never observe both $Y^*(0)$ and $Y^*(1)$ simultaneously. Each individual is associated with a treatment $A \in \{0, 1\}$, and we observe the *factual outcome* $Y = Y^*(A)$ corresponding to A. We never observe the counterfactual $Y^*(1 - A)$. That is, the data consists of observations of (X, A, Y).

In particular, the distribution of the data only reflects part of the full population distribution involving both potential outcomes. Let $\mathbb{P}^*$ denote the distribution of $(X, A, Y^*(0), Y^*(1))$. Define the coarsening function $\mathcal{C} : (x, a, y_0, y_1) \mapsto (x, a, ay_1 + (1 - a)y_0)$, which has the pre-image $\mathcal{C}^{-1}(x, a, y) = \{(x, a, y_0, y_1) : ay_0 + (1 - a)y_1 = y\}$. Then the distribution induced on (X, A, Y) is given by $\mathbb{P} = \mathbb{P}^* \circ \mathcal{C}^{-1}$ (as measures). That is, the data distribution $\mathbb{P}$ is given by taking a draw from $\mathbb{P}^*$ and coarsening it by $\mathcal{C}$. The data then consists of n independent draws, $(X_i, A_i, Y_i) \sim \mathbb{P}$ for $1 \le i \le n$. We use $\mathbb{E}^*, \mathbb{E}$ to denote expectations with respect to $\mathbb{P}^*, \mathbb{P}$, respectively. Note that both have the same (X, A)-distribution, so anything involving only (X, A) will be the same. For any function f of X, A, Y we let $\|f\|_p$ be the L_p-norm with respect to $\mathbb{P}$.

We assume throughout that all endogeneity is explained by X, known as unconfoundedness: $\mathbb{P}^*(A = 1 \mid X) = \mathbb{P}^*(A = 1 \mid Y^*(0), X) = \mathbb{P}^*(A = 1 \mid Y^*(1), X)$. That is, controlling for X, treatment assignment is independent of an individual's idiosyncrasies as relevant to their potential outcomes. Randomized experiments ensure this by design, often with $X \perp\!\!\!\perp A$ (complete randomization). Our results extend to observational settings assuming unconfoundedness, which is a common assumption in this setting [26]. For our purposes, the only technical difference between the experimental and

observational settings is whether or not we exactly know the propensity score, defined as

$$e(X) = \mathbb{P}(A = 1 \mid X).$$

We assume throughout that $e(X) \in (0, 1)$ almost surely, known as overlap. Note that assuming $Y = Y^*(A)$ also encapsulates non-interference [48].

In the following it will also be useful to define the conditional mean of potential outcomes:

$$\mu(X, a) = \mathbb{E}^*[Y^*(a) \mid X] = \mathbb{E}[Y \mid X, A = a], \quad a = 0, 1, \tag{1}$$

where the last equality follows from unconfoundedness and overlap, showing μ only depends on $\mathbb{P}$. We also define the conditional average treatment effect (CATE) and sum (CATS) functions as

$$\tau_-(X) = \mathbb{E}^*[\text{ITE} \mid X] = \mu(X, 1) - \mu(X, 0), \quad \tau_+(X) = \mu(X, 1) + \mu(X, 0). \tag{2}$$

2.1 Parameters of Interest

The primary parameter of interest we consider is the fraction negatively affected (FNA) by a wholesale change from treatment 0 to treatment 1:

$$\text{FNA} = \mathbb{P}^*(\text{ITE} < 0) = \mathbb{P}^*(Y^*(0) = 1, Y^*(1) = 0).$$

In particular, this quantity stands in contrast to the average treatment effect, $\text{ATE} = \mathbb{E}^*[\text{ITE}] = \mathbb{E}^*[Y^*(1)] - \mathbb{E}^*[Y^*(0)]$. A zero or positive ATE is generally interpreted to indicate a neutral or favorable treatment. However, having $\text{ATE} \geq 0$ need not mean having $\text{FNA} = 0$. A simple example is $(Y(0), Y(1)) \sim \text{Unif}(\{0, 1\}^2)$, which has $\text{ATE} = 0$ but $\text{FNA} = 1/4$.

Given we observe baseline covariates, we can more generally consider personalized treatment policies that assign treatments based on the value of X. Let $\pi_0, \pi_1 : \mathcal{X} \to \{0, 1\}$ denote any two such policies and let us define the fraction harmed by a change from π_0 to π_1 as

$$\text{FNA}_{\pi_0 \to \pi_1} = \mathbb{P}^*((\pi_1(X) - \pi_0(X)) \cdot \text{ITE} < 0) = \mathbb{P}^*(Y^*(\pi_0(X)) = 1, Y^*(\pi_1(X)) = 0).$$

This more general construct gives rise to several parameters of interest:

1. First, we recover our primary parameter of interest, $\text{FNA} = \text{FNA}_{0 \to 1}$, where 0 (or, 1) stands for the constant policy taking value 0 (or, 1) everywhere.
2. Second, given a proposed personalized treatment policy $\pi : \mathcal{X} \to \{0, 1\}$, the quantity $\text{FNA}_{0 \to \pi}$ is the fraction negatively affected by changing from a status quo of 0 to deploying the proposed policy π, which intervenes to treat the subpopulation with covariates X such that $\pi(X) = 1$.
3. Third, we can recover the misclassification rate of the policy π:

$$\text{FNA}_{1-\pi \to \pi} = \mathbb{P}^*(\pi(X) \notin \text{argmax}_{a \in \{0,1\}} Y^*(a)).$$

   This represents the fraction of individuals negatively affected by being misclassified by π: they should be treated by one treatment but are being given the other.
4. Of particular interest is the misclassification rate of the *optimal* personalized treatment policy, which assigns the treatment with larger conditional average outcome:

$$\pi^*(X) \in \text{argmax}_{a \in \{0,1\}} \mathbb{E}^*[Y^*(a) \mid X].$$

   We are interested in $\text{FNA}_{1-\pi^* \to \pi^*}$. The policy π^* can equivalently be characterized as the policy *maximizing* social welfare, $\mathbb{E}^*[Y^*(\pi(X))]$, or as *minimizing* misclassification, $\text{FNA}_{1-\pi \to \pi}$, both over *all* policies π. Given this, one hopes but cannot rule out that π^* avoids negative impact.

3 Sharp Bounds

In this section, we derive sharp bounds on our parameters of interest. For the sake of generality we focus on the parameter $\text{FNA}_{\pi_0 \to \pi_1}$ as it captures all of our parameters of interest using different π_0, π_1. The sharp bounds describe the set of values that $\text{FNA}_{\pi_0 \to \pi_1}$ can take if all we know is the distribution of the data, $\mathbb{P}$. This characterizes the most we can hope to learn at the limit of infinite data since this distribution is the most we can learn from draws from it, that is, our data.

To define this formally, recall that $\mathbb{P} = \mathbb{P}^* \circ \mathcal{C}^{-1}$ is given by $\mathbb{P}^*$ via coarsening. However, this mapping from $\mathbb{P}^*$ to $\mathbb{P}$ need not be invertible. Given a parameter $\psi(\mathbb{P}^*)$ that depends on the full

distribution $\mathbb{P}^*$, we define its *identified set* as the set of all values consistent with $\mathbb{P}$, that is, that could be explained by some $\mathbb{P}^*$ that gives rise to the given data distribution and satisfies unconfoundedness:

$$\mathcal{S}(\psi;\mathbb{P}) = \left\{\psi(\mathbb{P}^*) \ : \ \mathbb{P}^* \circ \mathcal{C}^{-1} = \mathbb{P}, \ \mathbb{P}^*(A = 1 \mid X) = \mathbb{P}^*(A = 1 \mid X, Y^*(a)), \ a = 0, 1\right\}.$$

111 We are interested in the identified set of and sharp bounds on the parameter $\mathrm{FNA}_{\pi_0 \to \pi_1}$.

112 When $|\mathcal{S}(\psi;\mathbb{P})| = 1$, we say that ψ is *identifiable* because it is uniquely specified by the distribution
113 of the data. Otherwise, we say it is *unidentifiable*: even given infinite data (equivalently, the
114 distribution of the data), we cannot determine the value of $\psi(\mathbb{P}^*)$ based on the data alone since
115 many values are consistent with the observations. For example, ATE *is* identifiable because ATE $=$
116 $\mathbb{E}[\mu(X, 1)] - \mathbb{E}[\mu(X, 0)]$ and Eq. (1) shows that $\mu(X, a) = \mathbb{E}[Y \mid X, A = a]$ depends on $\mathbb{P}$ alone.

117 Is $\mathrm{FNA}_{\pi_0 \to \pi_1}$ identifiable? Our first result characterizes exactly when, showing that generally, *no*.

118 **Theorem 1** (Identifiability). $\mathrm{FNA}_{\pi_0 \to \pi_1}$ *is identifiable if and only if, for almost all X, either*
119 $\pi_0(X) = \pi_1(X)$ *or* $\mathrm{Var}(Y \mid X, A = 0) = 0$ *or* $\mathrm{Var}(Y \mid X, A = 1) = 0$.

120 Thm. 1 shows that only in degenerate cases is $\mathrm{FNA}_{\pi_0 \to \pi_1}$ identifiable. Generally, there will exist a
121 non-negligible (positive probability) set of X's where the policies π_0, π_1 differ and neither conditional
122 outcome distribution is constant. Then, by Thm. 1, $\mathrm{FNA}_{\pi_0 \to \pi_1}$ would be unidentifiable.

123 We are now prepared to state our main theorem characterizing the sharp bounds on $\mathrm{FNA}_{\pi_0 \to \pi_1}$.
124 Since $\mathrm{FNA}_{\pi_0 \to \pi_1}$ is generally unidentifiable, the best we can do using the data is measure
125 the set $\mathcal{S}(\mathrm{FNA}_{\pi_0 \to \pi_1};\mathbb{P})$. The *sharp lower and upper bounds* are $\inf(\mathcal{S}(\mathrm{FNA}_{\pi_0 \to \pi_1};\mathbb{P}))$ and
126 $\sup(\mathcal{S}(\mathrm{FNA}_{\pi_0 \to \pi_1};\mathbb{P}))$, respectively, as these are precisely the largest (resp., smallest) lower (resp.,
127 upper) bounds on the set of possible values. The following result gives the sharp bounds and moreover
128 shows the identified set is closed and convex so it is in fact an interval with the bounds as its endpoints.

129 **Theorem 2** (Sharp Bounds on FNA). *Fix any $\pi_0, \pi_1 : \mathcal{X} \to \{0, 1\}$. Set*

$$\mathrm{FNA}_{\pi_0 \to \pi_1}^- = \mathbb{E}[\max\{(\pi_0(X) - \pi_1(X))\tau_-(X), 0\}], \tag{3}$$

$$\mathrm{FNA}_{\pi_0 \to \pi_1}^+ = \mathbb{E}[\min\{\pi_1(X)(1 - \pi_0(X))\mu(X, 0) + \pi_0(X)(1 - \pi_1(X))(1 - \mu(X, 0)), \tag{4}$$
$$\pi_1(X)(1 - \pi_0(X))(1 - \mu(X, 1)) + \pi_0(X)(1 - \pi_1(X))\mu(X, 1)\}].$$

130 *Then, the identified set is the interval $\mathcal{S}(\mathrm{FNA}_{\pi_0 \to \pi_1};\mathbb{P}) = [\mathrm{FNA}_{\pi_0 \to \pi_1}^-, \mathrm{FNA}_{\pi_0 \to \pi_1}^+]$.*

131 The proof of Thm. 2 proceeds by applying the Fréchet-Hoeffding bounds [20, 49] for each level of X
132 and showing this remains sharp. Eqs. (3) and (4) simplify a lot for the change from all-0 to all-1:

$$\mathrm{FNA}^- = -\mathbb{E}[\min\{\tau_-(X), 0\}], \quad \mathrm{FNA}^+ = \mathbb{E}[\min\{\mu(X, 0), 1 - \mu(X, 1)\}]. \tag{5}$$

133 We can also simplify when considering the optimal policy. Plugging in the optimal policy we get:

$$\mathrm{FNA}_{1-\pi^* \to \pi^*}^- = 0, \ \mathrm{FNA}_{1-\pi^* \to \pi^*}^+ = \mathbb{E}[\min\{\mu(X, 0), 1 - \mu(X, 0), \mu(X, 1), 1 - \mu(X, 1)\}]. \tag{6}$$

134 Unlike most policies, it is always plausible that the optimal policy affects no one negatively: because
135 $\mathrm{FNA}_{1-\pi^* \to \pi^*}^- = 0$, we always have $0 \in \mathcal{S}(\mathrm{FNA}_{1-\pi^* \to \pi^*};\mathbb{P})$. On the other hand, it is generally also
136 plausible that it does, that is, $\mathrm{FNA}_{1-\pi^* \to \pi^*}^+$ is generally nonzero. In agreement with Thm. 1, it is in
137 fact zero *only* when $\mu(X, 0) \in \{0, 1\}$ or $\mu(X, 1) \in \{0, 1\}$ for almost all X.

Remark 1 (Tail Expectations of Individual Treatment Effects). Kallus [27] considers bounds on the conditional value at risk (CVaR) of individual treatment effects (ITEs): for $\alpha \in (0, 1]$,

$$\mathrm{CVaR}_\alpha(\mathrm{ITE}) = \sup_\beta \mathbb{E}^*\left[\beta + \alpha^{-1} \min\{\mathrm{ITE} - \beta, 0\}\right].$$

138 This is equal to the smallest subgroup-ATE among all α-sized fractions of the population, that is, the
139 average effect on the $(100\alpha)\%$-worst affected. It is generally unidentifiable from data. While Kallus
140 [27] considers general real-valued outcomes, in the special case of binary outcomes considered here,
141 $\mathrm{CVaR}_\alpha(\mathrm{ITE})$ is a function of just FNA and ATE:

142 **Lemma 1.** *For $\alpha \in (0, 1)$, $\mathrm{CVaR}_\alpha(\mathrm{ITE}) = \max\{-1, -\alpha^{-1}\mathrm{FNA}, 1 - \alpha^{-1}(\mathrm{ATE} - 1)\}$.*

143 As a consequence of Thm. 2 and Lemma 1, we immediately have that the identified set for
144 $\mathrm{CVaR}_\alpha(\mathrm{ITE})$ for $\alpha \in (0, 1)$ is the interval $\mathcal{S}(\mathrm{CVaR}_\alpha(\mathrm{ITE});\mathbb{P}) = [\max\{-1, -\alpha^{-1}\mathrm{FNA}^+, 1 -$
145 $\alpha^{-1}(\mathrm{ATE} - 1)\}, \max\{-1, -\alpha^{-1}\mathrm{FNA}^-, 1 - \alpha^{-1}(\mathrm{ATE} - 1)\}]$, with $\mathrm{FNA}^\pm$ as defined in Eq. (5).
146 Consequently, the endpoints are the sharp lower and upper bounds on $\mathrm{CVaR}_\alpha(\mathrm{ITE})$.

147 Kallus [27] gives the upper bound $\mathrm{CVaR}_\alpha(\mathrm{ITE}) \leq \mathrm{CVaR}_\alpha(\tau_-(X))$ and shows it is tight given only
148 $\tau_-(X)$ in the sense that is always realizable by some $\mathbb{P}^*$ with the same distribution of $\tau_-(X)$. How-
149 ever, this bound is generally not sharp, that is, it need not be realizable given $\mathbb{P}$, which characterizes

more than just $\tau_-(X)$. In contrast, plugging FNA$^-$ into Lemma 1 gives the sharp upper bound in the special case of binary outcomes, that is, this bound fully uses *all* the information in $\mathbb{P}$. Moreover, we have a finite sharp lower bound, whereas Kallus [27] shows that no lower bound exists when given only $\tau_-(X)$. Nonetheless, Kallus [27] crucially handles general real-valued outcomes. In particular, the CVaR curve as α varies is most interesting in settings with non-binary outcomes, since in binary settings, two numbers – the ATE and FNA – summarize all relevant information per Lemma 1.

4 Inference Methodology

In the previous section we characterized the sharp (meaning, tightest possible) bounds on FNA given the distribution of the data, that is, the best we could do if we had infinite data. We now address how to estimate these from actual data and characterize the uncertainty. That is, how to do inference on the parameters $\text{FNA}^\pm_{\pi_0 \to \pi_1}$. Recall our data is n independent draws $(X_i, A_i, Y_i) \sim \mathbb{P}$ for $i = 1, \ldots, n$.

In order to systematically handle all of parameters of interest, we will develop a generic, robust method for a large class of parameters that include all of our parameters of interest as special cases. We discuss implications for each parameter of interest along the way.

4.1 Average Hinge Effects

We consider a class parameters we call Average Hinge Effects (AHEs), parametrized by a positive integer $m \in \mathbb{N}$, a vector of signs $\rho \in \{-1, +1\}^m$, and a set of $3(m+1)$ functions $g_\ell(x) = (g_\ell^{(0)}(x), g_\ell^{(1)}(x), g_\ell^{(2)}(x)) \in [-1,1]^3$ for $\ell = 0, \ldots, m$:

$$\text{AHE}^\rho_{g_0,\ldots,g_m} = \mathbb{E}\big[g_0^{(0)}(X)\mu(X,0) + g_0^{(1)}(X)\mu(X,1) + g_0^{(2)}(X) \tag{7}$$
$$+ \textstyle\sum_{\ell=1}^m \rho_\ell \min\{0,\, g_\ell^{(0)}(X)\mu(X,0) + g_\ell^{(1)}(X)\mu(X,1) + g_\ell^{(2)}(X)\}\big].$$

The AHE is so called because $\min\{0, x\}$ is called the hinge function. AHEs cover all of our parameters of interest: $\text{FNA}^- = \text{AHE}^{(-)}_{(0,0,0),(-1,1,0)}$, $\text{FNA}^+ = \text{AHE}^{(+)}_{(1,0,0),(-1,-1,1)}$, $\text{FNA}^-_{\pi_0 \to \pi_1} = \text{AHE}^{(-)}_{(0,0,0),(\pi_0-\pi_1,\, \pi_1-\pi_0,\, 0)}$, $\text{FNA}^+_{\pi_0 \to \pi_1} = \text{AHE}^{(+)}_{(\pi_0(1-\pi_1),\, \pi_1-\pi_0,\, 0),\, (\pi_0-\pi_1,\, \pi_0-\pi_1,\, -1)}$, and $\text{FNA}^+_{1-\pi^* \to \pi^*} = \text{AHE}^{(+,+)}_{(1,0,0),(-1,1,0),(-1,-1,1)}$. For now, we focus on any given $m, \rho, g_0, \ldots, g_m$.

4.2 Re-formulating the AHE

The AHE is defined as an average of a function of X. The only unknown part of this function is $\mu(x,a)$. A naïve approach would be to estimate μ by $\hat\mu$, plug it into Eq. (7) and replace $\mathbb{E}$ by a sample average over $X_1, \ldots, X_n$. This estimate, however, will *not* behave like a sample average approximation of Eq. (7). In particular, $\hat\mu$ may only converge as fast as $O_p(1/\sqrt{n})$. Even if it converges that fast, this will introduce significant errors on the same order as the convergence of sample averages, which must be somehow accounted for in inference. If it converges more slowly, it can even affect the convergence rate of the final estimate. Generally, the behavior of the estimate will heavily depend on how exactly we fit $\hat\mu$, which is something we would like to avoid.

We will therefore develop a *robust* inference method, which can guarantee an estimate that behaves like a sample average (known as being asymptotically linear) almost *regardless* of how $\hat\mu$ is estimated, provided some lax conditions hold. Our approach would be to focus on another formulation of AHE as a sample average where plugging in unknowns does not affect the result too much.

Toward that end, let us first define $\eta_\ell : \mathcal{X} \to [-3, 3]$ by

$$\eta_\ell(x) = g_\ell^{(0)}(X)\mu(X,0) + g_\ell^{(1)}(X)\mu(X,1) + g_\ell^{(2)}(X). \tag{8}$$

Specifically, we have $\eta_1(X) = (\pi_1(X) - \pi_0(X))\tau_-(X)$ for $\text{FNA}^-_{\pi_0 \to \pi_1}$, $\eta_1(X) = (\pi_1(X) - \pi_0(X))(1 - \tau_+(X))$ for $\text{FNA}^+_{\pi_0 \to \pi_1}$, and $\eta_1(X) = \tau_-(X)$, $\eta_2(X) = 1 - \tau_+(X)$ for $\text{FNA}^+_{1-\pi^* \to \pi^*}$.

Given $\check{e}, \check{\mu}, \check{\eta}_1, \ldots, \check{\eta}_m$, understood as (possibly wrong) stand-ins for $e, \mu, \eta_1, \ldots, \eta_m$, define

$$\phi^\rho_{g_0,\ldots,g_m}(X, A, Y; \check{e}, \check{\mu}, \check{\eta}_1, \ldots, \check{\eta}_m) = g_0^{(2)}(X) + \textstyle\sum_{\ell=1}^m \rho_\ell \mathbb{I}[\check{\eta}_\ell(X) \le 0]g_\ell^{(2)}(X)$$
$$+ \big(g_0^{(0)}(X) + \textstyle\sum_{\ell=1}^m \rho_\ell \mathbb{I}[\check{\eta}_\ell(X) \le 0]g_\ell^{(0)}(X)\big)\frac{(A-\check{e}(X))\check{\mu}(X,0)+(1-A)Y}{1-\check{e}(X)}$$
$$+ \big(g_0^{(1)}(X) + \textstyle\sum_{\ell=1}^m \rho_\ell \mathbb{I}[\check{\eta}_\ell(X) \le 0]g_\ell^{(1)}(X)\big)\frac{(\check{e}(X)-A)\check{\mu}(X,1)+AY}{\check{e}(X)}.$$

Algorithm 1 Point estimate and CIs for $\text{AHE}^{\rho}_{g_0,\ldots,g_m}$

Input: Data $\{(X_i, A_i, Y_i) : i = 1, \ldots, n\}$, number of folds K, estimators for $e, \mu, \eta_1, \ldots, \eta_m$

1: **for** $k = 1, \ldots, K$ **do**
2:     Estimate $\hat{e}^{(k)}, \hat{\mu}^{(k)}, \hat{\eta}_1^{(k)}, \ldots, \hat{\eta}_m^{(k)}$ using data $\{(X_i, A_i, Y_i) : i \not\equiv k - 1 \pmod{K}\}$
3:     **for** $i \equiv k - 1 \pmod{K}$ **do** set $\phi_i = \phi^{\rho}_{g_0,\ldots,g_m}(X_i, A_i, Y_i; \hat{e}^{(k)}, \hat{\mu}^{(k)}, \hat{\eta}_1^{(k)}, \ldots, \hat{\eta}_m^{(k)})$
4: **end for**
5: Set $\widehat{\text{AHE}}^{\rho}_{g_0,\ldots,g_m} = \frac{1}{n}\sum_{i=1}^{n}\phi_i$, $\hat{\text{se}} = \sqrt{\frac{1}{n(n-1)}\sum_{i=1}^{n}(\phi_i - \widehat{\text{AHE}}^{\rho}_{g_0,\ldots,g_m})^2}$
6: Return point estimate $\widehat{\text{AHE}}^{\rho}_{g_0,\ldots,g_m}$ and β-CIs $[\widehat{\text{AHE}}^{\rho}_{g_0,\ldots,g_m} \pm \Phi^{-1}((1+\beta)/2)\hat{\text{se}}]$

If we plug in the correct values of the nuisance parameters $e, \mu, \eta_1, \ldots, \eta_m$, then we would have

$$\text{AHE}^{\rho}_{g_0,\ldots,g_m} = \mathbb{E}[\phi^{\rho}_{g_0,\ldots,g_m}(X, A, Y; e, \mu, \eta_1, \ldots, \eta_m)]. \tag{9}$$

Crucially, unlike the definition of AHE as an average in Eq. (7), we will show in Sec. 5 that this new representation as an average remains faithful *even* when we make small errors in the nuisances.

4.3 Inference Algorithm

Given our special construction of $\phi^{\rho}_{g_0,\ldots,g_m}$, our algorithm, presented in Alg. 1, follows a simple recipe: we estimate the nuisances in a cross-fold fashion and plug them into Eq. (9), approximating the expectation by a sample average. In Sec. 5 we will show that the special structure of $\phi^{\rho}_{g_0,\ldots,g_m}$ affords this procedure a lot of robustness. The use of cross-fitting ensures nuisance estimates are independent of samples applied thereto [12, 50, 61]. This allows our analysis to only require lax rate assumptions on the nuisance fitting without requiring any additional regularity on restricting *how* the fitting is done. (If we assume nuisance estimates belong to a Donsker class with probability tending to 1, all of our results will hold even without cross-fitting; we omit this option for brevity.)

4.4 Nuisance Fitting

Our algorithm requires methods for fitting the nuisances $e, \mu, \eta_1, \ldots, \eta_m$. Fitting e and μ amounts to binary regressions (probabilistic classification), which can be done with any of a variety of standard supervised learning methods, whether logistic regression, random forests, or neural nets.

There are different ways to fit η_ℓ. In particular, despite the fact that η_ℓ is determined by μ, we treat it as a separate nuisance to allow for specialized learners. Of course, the simplest way to fit it is to simply plugin an estimate for μ into the definition of η_ℓ in Eq. (8), and that is a legitimate option. For all of our parameters of interest, all η_ℓ nuisances are given by learning just two functions: τ_- and τ_+ as in Eq. (2). Therefore, we may alternatively learn these directly and plug them into η_ℓ. For example, there exists a wide literature specialized to learning τ_- motivated by the observation that effect signals are often more nuanced and can therefore be washed out by the noise in baselines if we simply difference μ-estimates [3, 25, 33, 36, 44, 57]. We can similarly apply of all these approaches to learning τ_+ by simply first mutating the outcome data as $Y \leftarrow (2A - 1)Y$. Notably, [33, 44] give rates of convergence for τ_--learning, which we can use to satisfy our assumptions in the next section.

5 Robustness Guarantees for Inference

We now show our inference method has some nice robustness guarantees. This will depend on showing that using our special ϕ renders Eq. (9) insensitive to errors in the nuisances. The specific level of errors allowed will depend on the *sharpness* of margin satisfied by η_ℓ, as defined below.

Definition 1. We say $f : \mathcal{X} \to \mathbb{R}$ satisfies a margin with sharpness $\alpha \in [0, \infty]$, or an α-margin, if there exists $t_0 > 0$ such that $\mathbb{P}(0 < |f(X)| \leq t) \leq (t/t_0)^{\alpha}$ for all $t > 0$. (We use the convention that $x^{\infty} = 0$ if $0 \leq x < 1$, $1^{\infty} = 1$, and $x^{\infty} = \infty$ if $x > 1$.)

Every f trivially satisfies a 0-margin, but usually a sharper margin holds. If $f(X)$ is almost surely either zero or bounded away from zero, then f satisfies an infinitely sharp margin:

Lemma 2. If $\mathbb{P}(|f(X)| \geq t_0) = \mathbb{P}(f(X) \neq 0)$ for some $t_0 > 0$ then f satisfies an ∞-margin.

In particular, if X has finite support, then we can always set $t_0 = \inf\{|f(x)| : x \in \mathcal{X}, f(x) \neq 0\}$. But, this also works more generally. Consider $\eta_1(X) = (\pi_1(X) - \pi_0(X))\tau_-(X)$ for $\text{FNA}^-_{\pi_0 \to \pi_1}$. If $\inf\{|\tau_-(x)| : \pi_1(x) \neq \pi_0(x), \tau_-(x) \neq 0\} > 0$, that is, the policies only differ either far from the boundary where τ_- is 0 or exactly on this boundary, then Lemma 2 ensures sharpness ∞.

More generally, we should expect a margin with sharpness 1 in continuous settings.

Lemma 3. *If the cumulative distribution function (CDF) of $f(X)$ is boundedly differentiable on $(-\epsilon, 0) \cup (0, \epsilon)$ for some $\epsilon > 0$, then f satisfies a 1-margin.*

Consider again the example of $\eta_1(X) = (\pi_1(X) - \pi_0(X))\tau_-(X)$. If $\tau_-(X)$ is a continuous random variable with a bounded density near 0, then Lemma 3 ensures a margin with sharpness 1. This holds for *any* two policies, including $\pi_1 = 1$, $\pi_0 = 0$, which differ everywhere. The same conclusion would also hold if the CDF of $\tau_-(X)$ was just continuously differentiable at 0.

Armed with the notion of margin, we can state in what sense the representation Eq. (9) is robust.

Theorem 3. *Fix any $\tilde{e}, \check{e} : \mathcal{X} \to [\bar{e}, 1 - \bar{e}]$ with $\bar{e} > 0$, $\tilde{\mu}, \check{\mu} : \mathcal{X} \times \{0, 1\} \to [0, 1]$, $\tilde{\eta}_\ell, \check{\eta}_\ell : \mathcal{X} \to [-3, 3]$ for $\ell = 1, \ldots, m$. Fix any $p_1, \ldots, p_\ell \in [1, \infty]$. Suppose that either $\tilde{e} = e$ or $\tilde{\mu} = \mu$ (or both). Set $\kappa_\ell = 1$ if $\tilde{\eta}_\ell = \eta_\ell$ and otherwise set $\kappa_\ell = 0$. Further assume that $\mathbb{P}(\tilde{\eta}_\ell(X) = 0, \check{\eta}_\ell(X) \neq 0) = 0$. Finally, suppose that $\tilde{\eta}_\ell$ satisfies a margin with sharpness α_ℓ. Then, for some constant $c > 0$, we have*

$$c\big|\mathbb{E}[\phi^\rho_{g_0,\ldots,g_m}(X, A, Y; \check{e}, \check{\mu}, \check{\eta}_1, \ldots, \check{\eta}_m)] - \mathbb{E}[\phi^\rho_{g_0,\ldots,g_m}(X, A, Y; \tilde{e}, \tilde{\mu}, \tilde{\eta}_1, \ldots, \tilde{\eta}_m)]\big|$$
$$\leq \|\check{e} - e\|_2 \|\check{\mu} - \mu\|_2 + \sum_{\ell=1}^m \|\check{\eta}_\ell - \tilde{\eta}_\ell\|_{p_\ell}^{\frac{p_\ell(\kappa_\ell + \alpha_\ell)}{p_\ell + \alpha_\ell}},$$

$$c\big\|\phi^\rho_{g_0,\ldots,g_m}(X, A, Y; \check{e}, \check{\mu}, \check{\eta}_1, \ldots, \check{\eta}_m) - \phi^\rho_{g_0,\ldots,g_m}(X, A, Y; \tilde{e}, \tilde{\mu}, \tilde{\eta}_1, \ldots, \tilde{\eta}_m)\big\|_2$$
$$\leq \|\check{e} - \tilde{e}\|_2 + \|\check{\mu} - \tilde{\mu}\|_2 + \sum_{\ell=1}^m \|\check{\eta}_\ell - \tilde{\eta}_\ell\|_{p_\ell}^{\frac{p\alpha_\ell}{2p + 2\alpha_\ell}}.$$

In the above, we use the convention that $\frac{\infty a + b}{\infty c + d} = \frac{a}{c}$.

5.1 Local Robustness, Confidence Intervals, and Efficiency

Thm. 3 has several important implications. Our first result shows that, if we estimate all nuisances correctly but potentially slowly at nonparametric rates, then we can ensure our estimator looks like a sample average approximation of Eq. (9) and our CIs are calibrated.

Theorem 4. *Suppose $e(X) \in [\bar{e}, 1 - \bar{e}]$ with $\bar{e} > 0$, η_ℓ satisfies an α_ℓ-margin, and for $k = 1, \ldots, K$*

$$\|\hat{e}^{(k)} - e\|_2 \|\hat{\mu}^{(k)} - \mu\|_2 = o_p(n^{-\frac{1}{2}}), \quad \|\hat{\eta}_\ell^{(k)} - \eta_\ell\|_{p_\ell} = o_p(n^{-\frac{p_\ell + \alpha_\ell}{2p_\ell(1 + \alpha_\ell)}}) \text{ with } p_\ell \in [1, \infty], \text{ and}$$

$$\mathbb{P}(\bar{e} \leq \hat{e}^{(k)}(x) \leq 1 - \bar{e}, 0 \leq \hat{\mu}^{(k)}(x) \leq 1, |\hat{\eta}_\ell^{(k)}(x)| \leq 3\mathbb{I}[\eta_\ell(x) \neq 0], \ell \leq m, \text{ a.e. } x) \to 1. \text{ Then,}$$

$$\widehat{\text{AHE}}^\rho_{g_0,\ldots,g_m} = \frac{1}{n} \sum_{i=1}^n \phi^\rho_{g_0,\ldots,g_m}(X_i, A_i, Y_i; e, \mu, \eta_1, \ldots, \eta_m) + o_p(n^{-\frac{1}{2}}),$$

$$\mathbb{P}(\text{AHE}^\rho_{g_0,\ldots,g_m} \in [\widehat{\text{AHE}}^\rho_{g_0,\ldots,g_m} \pm \Phi^{-1}((1 + \beta)/2)\hat{\text{se}}]) \to \beta \quad \forall \beta \in (0, 1).$$

The conditions of Thm. 4 allow that we learn $e, \mu, \eta_1, \ldots, \eta_m$ rather slowly. In particular, unlike naïvely plugging in a μ-estimate into Eq. (7) and taking sample averages, we do not depend at all on *how* μ is estimated and the uncertainty therein, provided some slow nonparametric rates hold. For example, learning e, μ in with $o_p(n^{-1/4})$-rates in L_2-error suffices. Or, if e is known, then consistently estimating μ *without* a rate suffices. For η_ℓ, if an ∞-margin holds, it suffices to learn η_ℓ consistently (no rate) in L_∞-error or at $o_p(n^{-1/4})$-rates in L_2-error. If a 1-margin holds, it suffices to learn at $o_p(n^{-1/4})$-rates in L_∞-error. While conditions like metric entropy imply L_2-error rates [58], for many classes like smooth functions the L_p error rates are even the same for all $p \in [1, \infty]$ [52].

In fact, under one more condition, our estimator achieves the semiparametric efficiency lower bound.

Theorem 5. *Suppose $\mathbb{P}(\eta_\ell(X) = 0, g_\ell^{(a)}(X) \neq 0) = 0$ for $\ell = 1, \ldots, m$, $a = 0, 1$. The semiparametric efficiency lower bound for $\text{AHE}^\rho_{g_0,\ldots,g_m}$ is $\text{Var}(\phi^\rho_{g_0,\ldots,g_m}(X, A, Y; e, \mu, \eta_1, \ldots, \eta_m))$, whether or not $e(X)$ is unknown (varies in the model) or known (fixed in the model).*

Corollary 1. *Under the assumptions of Thms. 4 and 5, $\widehat{\text{AHE}}^\rho_{g_0,\ldots,g_m}$ is efficient: it is regular and achieves the minimum asymptotic variance among all regular estimators.*

For all of our parameters of interest, the condition of Thm. 5 holds if $\mathbb{P}(\tau_-(X) = 0) = \mathbb{P}(\tau_+(X) = 0) = 0$ (*e.g.*, $\tau_-(X), \tau_+(X)$ are continuous). Similarly, the condition that $|\hat{\eta}_\ell^{(k)}(x)| \leq 3\mathbb{I}[\eta_\ell(x) \neq 0]$ in Thm. 4 is immediately satisfied if $\hat{\tau}_-^{(k)}(X), \hat{\tau}_+^{(k)}(X)$ are also almost never 0 (*e.g.*, continuous).

5.2 Double Robustness and Double Validity

Next we show that Alg. 1 is very robust to incorrectly learning some nuisances.

Theorem 6. *Fix any $\tilde{\mu}, \tilde{e}, \tilde{\eta}_1, \ldots, \tilde{\eta}_m$ with $\tilde{\mu}(X) \in [0, 1]$, $\tilde{e}(X) \in [\bar{e}, 1 - \bar{e}]$ with $\bar{e} > 0$. Set $\kappa_\ell = 1$ if $\tilde{\eta}_\ell = \eta_\ell$ and otherwise set $\kappa_\ell = 0$. Suppose $\tilde{\eta}_\ell$ satisfies a margin with sharpness α_ℓ, and for*

270   $k = 1, \ldots, K$ *that* $\|\hat{e}^{(k)} - \tilde{e}\|_2 = o_p(1)$, $\|\hat{\mu}^{(k)} - \tilde{\mu}\|_2 = o_p(1)$, $\|\hat{\eta}_\ell^{(k)} - \tilde{\eta}_\ell\|_{p_\ell} = O_p(n^{-\frac{p_\ell + \alpha_\ell}{2p_\ell(\kappa_\ell + \alpha_\ell)}})$

271   *for some* $p_\ell \in [1, \infty]$, *and* $\mathbb{P}(\bar{e} \leq \hat{e}^{(k)}(x) \leq 1 - \bar{e}, 0 \leq \hat{\mu}^{(k)}(x) \leq 1, |\hat{\eta}_\ell^{(k)}(x)| \leq 3\mathbb{I}[\tilde{\eta}_\ell(x) \neq$

272   $0], \ell \leq m, a.e.\ x) \to 1$. *If either* $\|\hat{e}^{(k)} - e\|_2 = O_p(n^{-1/2})$ *or* $\|\hat{\mu}^{(k)} - \mu\|_2 = O_p(n^{-1/2})$, *then*

$$\widehat{\mathrm{AHE}}_{g_0, \ldots, g_m}^{\rho} = \mathrm{AHE}_{g_0, \ldots, g_m}^{\rho} + O_p(n^{-\frac{1}{2}}) \ \ \textit{if } \kappa_1 = \cdots = \kappa_m = 1, \qquad \textit{(double robustness)}$$

$$\widehat{\mathrm{AHE}}_{g_0, \ldots, g_m}^{\rho} \geq \mathrm{AHE}_{g_0, \ldots, g_m}^{\rho} + O_p(n^{-\frac{1}{2}}) \ \ \textit{if } \rho_\ell = +\textit{ whenever } \kappa_\ell = 0, \quad \textit{(double validity, upper)}$$

$$\widehat{\mathrm{AHE}}_{g_0, \ldots, g_m}^{\rho} \leq \mathrm{AHE}_{g_0, \ldots, g_m}^{\rho} + O_p(n^{-\frac{1}{2}}) \ \ \textit{if } \rho_\ell = -\textit{ whenever } \kappa_\ell = 0. \quad \textit{(double validity, lower)}$$

273   Thm. 6 shows we remain consistent even if we incorrectly learn either e or μ. Moreover, even if we
274   learn η_ℓ in correctly, the direction of bias induced is pre-determined. Since all of our parameters of
275   interest had ρ all being positive for upper bounds and negative for lower bounds, the implication is that
276   even if we learn τ_-, τ_+ incorrectly, we will still obtain valid, albeit conservative, bounds. Therefore,
277   we can really trust the results of Alg. 1 are truly bounds on FNA, even if we make mistakes.

6   Empirical Investigation

279   We demonstrate our method in a simulation and a case study with data from a real experiment.
280   Replication code is given in the supplement. Experiments were run on an AWS c5.24xlarge instance.

Simulation Study   We consider 7-dimensional standard normal X and set $\mu(x, a) = (1 +$
282   $\exp(-\beta(2\mathbb{I}[\xi(x)x_2 > 0] - 1)(\mathbb{I}[\xi(x)x_1 \leq 0] + (2a - 1)\mathbb{I}[\xi(x)x_1 > 0])))^{-1}$, where $\beta \geq 0$ and
283   $\xi(x) = 2\,\mathrm{XOR}(\mathbb{I}[x_3 > 0], \ldots, \mathbb{I}[x_7 > 0]) - 1$. Here β controls how much X predicts $Y(a)$. The
284   dashed lines in Fig. 1 show the sharp bounds of Thm. 2 for FNA and $\mathrm{FNA}_{1-\pi^* \to \pi^*}$ as we vary β.
285   We can see that, as X becomes more predictive, the bounds go from $[0, 0.5]$ to a point at 0.25. Note
286   there is no *true* value for FNA as we are not specifying the joint distribution of potential outcomes;
287   instead, per Thm. 2, there are joint distributions with FNA equal to any point between the bounds.

288   Next, we estimate these true bounds. We fix $e(X) = (1+\exp(0.25-\mathbb{I}[x_3 > 0]+0.5\mathbb{I}[x_4 > 0])^{-1}$, and
289   draw n samples from $X \sim \mathcal{N}(0, I)$, $A \mid X \sim \mathrm{Bernoulli}(e(X))$, $Y \mid X, A \sim \mathrm{Bernoulli}(\mu(X, A))$.
290   We apply Alg. 1 to this data with $K = 5$, estimating e, μ using random regression forests and τ_-, τ_+
291   using causal forests (all using *R* package `grf` with default parameters). The solid lines in Fig. 1 show
292   the point estimate and 95%-CI output by Alg. 1 for a single draw of $n = 12800$ data for each β. In
293   Figs. 2 and 3 we plot the average root mean squared error of point estimates and coverage of CIs over
294   100 replications with $\beta = 3$ and varying n. We compare the performance to the (cross-fitted) "plugin"
295   estimator that just plugs in the (same) estimates for μ into Eq. (7) and approximate the expectation by
296   a sample average. Shaded intervals denote 95%-CIs for performance (*i.e.*, due to finite replications).

Case Study   Using data from [6] (BSD license), we compare three different assistance programs
298   offered to French unemployed individuals: the standard benefits (sta), access to public-run counseling
299   (pub), and to private-run counseling (pri). Our binary outcome is reemployment within six months.
300   For example, we can consider a hypothetical scenario where we change from a private to a public
301   counseling provider. As reported by [6], the ATE for this hypothetically change is slightly positive,
302   so a policymaker might therefore enact it, especially if, for example, it offered cost savings or other
303   operational benefits. We consider who may be harmed by this: how many who remain unemployed
304   under the new public program that would have been reemployed under the old private program.

305   For any pair of arms, we consider the sharp bounds on the FNA from one to other or the other way
306   and on the misclassification rate of the optimal policy choosing between the two. Following Kallus
307   [27], we set X to all pre-treatment covariates in table 2 of Behaghel et al. [6]., and we consider the
308   sharp bounds either with these X or given no X at all. We fit μ using linear regression and τ_-, τ_+
309   using doubly-robust pseudo-outcome linear regression. Since the propensity is known, in light of the
310   guarantees of Thm. 6, we do not worry about misspecifying μ, only using it for variance reduction by
311   accounting for main effects, and we do not worry about misspecifying τ_-, τ_+, only hoping to tighten
312   the bounds some from using no X. The results are shown in Fig. 4. We see that, while covariates are
313   somewhat uninformative, we are still able to tighten bounds compared to no X.

7   Connections, Limitations, Extensions, and Conclusions

Connections to partial identification   Partial identification of unknowable parameters has a long
316   tradition in econometrics [11, 34, 38, 39, 41, 43, 51, 53]. Some approaches focus on average treatment

Fig. 1: Sharp bounds, estimates, and 95%-CIs as X-predictiveness (β) varies

Fig. 2: Root mean squared error (RMSE) for estimating the sharp bounds as n varies

Fig. 3: Coverage of 95%-CIs using $\pm 1.96/\sqrt{n}$ standard deviations as n varies

(a) Lower bound FNA$^-$ (b) Upper bound FNA$^+$ (c) FNA$^+_{1-\pi^* \to \pi^*}$

Fig. 4: Estimated sharp bounds for FNA in the case study with covariates and without covariates

effects in the presence of confounding [8, 13, 15, 17, 30, 47, 54, 60], among which [8, 13, 17, 60] are notable for conducting semiparametric inference on bounds, but for confounding rather than for individual effects, as we do. Some others, like us, focus on understanding the joint distribution of potential outcomes [1, 16, 18, 24, 40]. Like us, these heavily leverage the Fréchet-Hoeffding bounds [9, 20, 46, 49, 59]. Unlike these, we use efficient and robust inference that can aggregate across covariates to tighten bounds by leveraging machine learning of conditional-mean outcome functions.

Connections to fairness   Understanding the distribution of impact is a central problem in algorithmic fairness [5, 42]. Here, we specifically focused on the *differential* impact of interventions, which brought up issues of identification. In contrast, in algorithmic fairness one usually measures disparities in observed outcomes, such as in the form of the loss function of a model on a labeled example [2, 7, 21, 32]. A line of work specifically focuses on identification issues in algorithmic fairness [10, 14, 19, 28, 29, 31, 35, 37]. Notably, [31] also conduct semiparametric inference on sharp bounds derived from the Fréchet-Hoeffding bounds, but in the context of algorithm evaluation with unobserved protected labels rather than interventions with unobserved counterfactuals. And, [29] do consider unobserved counterfactuals in assessing equality of opportunity [23] for intervention-prioritization policies, but (appropriately in their own context) they assume away our primary focus here by assuming an a priori known bound on FNA (possibly zero, *i.e.*, monotone treatment response).

Limitations and Extensions   Restricting outcomes to be binary is one limitation of our work, and extensions to continuous settings is an interesting avenue of future research. Another limitation is that, while we can bound the FNA, it can still be hard to asses *who* is negatively affected. This is, unfortunately, impossible for the same reason FNA is unidentifiable, but a place to start such analyses may be to consider who are individuals with $\tau_-(X) \leq 0$ and even characterize that group via summary statistics compared to the population. Indeed, per Eq. (5), the lower bound on FNA is exactly the (negative of the) ATE on this group. Another important consideration is whether the data is representative: FNA refers to the fraction of the *studied population*, which might differ from the population of interest. *E.g.*, if an experiment did not enroll a representative sample, we may be systematically excluding some groups from consideration. If such unrepresentativeness is explained by covariates X (*i.e.*, missing at random), the solution is simple: we reweight. If not explained by X (*i.e.*, missing *not* at random), then we need to also account for this additional source of unidentifiability. An avenue for future research is to combine such ambiguity with counterfactual ambiguity. Another concern is whether the outcome represents the impact we want to measure [45].

Conclusions   Our tools support drawing credible conclusions about the potential negative impact of interventions: they both account for ambiguity due to unobserved counterfactuals (while mitigating it using covariates) as well as strongly guard against slow or inconsistent learning of necessary nuisances functions. Robust inference on the lower bounds, in particular, crucially provides watertight demonstrations of negative impact, which can bolster efforts to mitigate harm and improve equity.

References

[1] Jaap H Abbring and James J Heckman. Econometric evaluation of social programs, part iii: Distributional treatment effects, dynamic treatment effects, dynamic discrete choice, and general equilibrium policy evaluation. *Handbook of econometrics*, 6:5145–5303, 2007.

[2] Alekh Agarwal, Alina Beygelzimer, Miroslav Dudík, John Langford, and Hanna Wallach. A reductions approach to fair classification. In *International Conference on Machine Learning*, pages 60–69. PMLR, 2018.

[3] Susan Athey and Guido Imbens. Recursive partitioning for heterogeneous causal effects. *Proceedings of the National Academy of Sciences*, 113(27):7353–7360, 2016.

[4] Jean-Yves Audibert and Alexandre B Tsybakov. Fast learning rates for plug-in classifiers. *The Annals of statistics*, 35(2):608–633, 2007.

[5] Solon Barocas, Moritz Hardt, and Arvind Narayanan. *Fairness and Machine Learning*. fairmlbook.org, 2019. http://www.fairmlbook.org.

[6] Luc Behaghel, Bruno Crépon, and Marc Gurgand. Private and public provision of counseling to job seekers: Evidence from a large controlled experiment. *American economic journal: applied economics*, 6(4):142–74, 2014.

[7] Sarah Bird, Miro Dudík, Richard Edgar, Brandon Horn, Roman Lutz, Vanessa Milan, Mehrnoosh Sameki, Hanna Wallach, and Kathleen Walker. Fairlearn: A toolkit for assessing and improving fairness in ai. *Microsoft, Tech. Rep. MSR-TR-2020-32*, 2020.

[8] Matteo Bonvini and Edward H Kennedy. Sensitivity analysis via the proportion of unmeasured confounding. *Journal of the American Statistical Association*, pages 1–11, 2021.

[9] Stamatis Cambanis, Gordon Simons, and William Stout. Inequalities for e k (x, y) when the marginals are fixed. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 36(4): 285–294, 1976.

[10] Jiahao Chen, Nathan Kallus, Xiaojie Mao, Geoffry Svacha, and Madeleine Udell. Fairness under unawareness: Assessing disparity when protected class is unobserved. In *Proceedings of the conference on fairness, accountability, and transparency*, pages 339–348, 2019.

[11] Victor Chernozhukov, Han Hong, and Elie Tamer. Estimation and confidence regions for parameter sets in econometric models 1. *Econometrica*, 75(5):1243–1284, 2007.

[12] Victor Chernozhukov, Denis Chetverikov, Mert Demirer, Esther Duflo, Christian Hansen, Whitney Newey, and James Robins. Double/debiased machine learning for treatment and structural parameters. *Econometrics Journal*, 21(1):C1–C68, 2018.

[13] Victor Chernozhukov, Carlos Cinelli, Whitney Newey, Amit Sharma, and Vasilis Syrgkanis. Omitted variable bias in machine learned causal models. *arXiv preprint arXiv:2112.13398*, 2021.

[14] Amanda Coston, Ashesh Rambachan, and Alexandra Chouldechova. Characterizing fairness over the set of good models under selective labels. In *International Conference on Machine Learning*, pages 2144–2155. PMLR, 2021.

[15] Philip J Cross and Charles F Manski. Regressions, short and long. *Econometrica*, 70(1): 357–368, 2002.

[16] Peng Ding, Avi Feller, and Luke Miratrix. Decomposing treatment effect variation. *Journal of the American Statistical Association*, 114(525):304–317, 2019.

[17] Jacob Dorn, Kevin Guo, and Nathan Kallus. Doubly-valid/doubly-sharp sensitivity analysis for causal inference with unmeasured confounding. *arXiv preprint arXiv:2112.11449*, 2021.

[18] Sergio Firpo and Geert Ridder. Bounds on functionals of the distribution of treatment effects. 2008.

[19] Riccardo Fogliato, Alexandra Chouldechova, and Max G'Sell. Fairness evaluation in presence of biased noisy labels. In *International Conference on Artificial Intelligence and Statistics*, pages 2325–2336. PMLR, 2020.

[20] Maurice Fréchet. Généralisation du théoreme des probabilités totales. *Fundamenta mathematicae*, 1(25):379–387, 1935.

[21] Avijit Ghosh, Lea Genuit, and Mary Reagan. Characterizing intersectional group fairness with worst-case comparisons. In *Artificial Intelligence Diversity, Belonging, Equity, and Inclusion*, pages 22–34. PMLR, 2021.

[22] Jinyong Hahn. On the role of the propensity score in efficient semiparametric estimation of average treatment effects. *Econometrica*, pages 315–331, 1998.

[23] Moritz Hardt, Eric Price, and Nati Srebro. Equality of opportunity in supervised learning. *Advances in neural information processing systems*, 29, 2016.

[24] James J Heckman, Jeffrey Smith, and Nancy Clements. Making the most out of programme evaluations and social experiments: Accounting for heterogeneity in programme impacts. *The Review of Economic Studies*, 64(4):487–535, 1997.

[25] Kosuke Imai and Marc Ratkovic. Estimating treatment effect heterogeneity in randomized program evaluation. *The Annals of Applied Statistics*, 7(1):443–470, 2013.

[26] Guido W Imbens and Donald B Rubin. *Causal inference in statistics, social, and biomedical sciences*. Cambridge University Press, 2015.

[27] Nathan Kallus. Treatment effect risk: Bounds and inference. *arXiv preprint arXiv:2201.05893*, 2022.

[28] Nathan Kallus and Angela Zhou. Residual unfairness in fair machine learning from prejudiced data. In *International Conference on Machine Learning*, pages 2439–2448. PMLR, 2018.

[29] Nathan Kallus and Angela Zhou. Assessing disparate impacts of personalized interventions: Identifiability and bounds. *arXiv preprint arXiv:1906.01552*, 2019.

[30] Nathan Kallus, Xiaojie Mao, and Angela Zhou. Interval estimation of individual-level causal effects under unobserved confounding. In *The 22nd international conference on artificial intelligence and statistics*, pages 2281–2290. PMLR, 2019.

[31] Nathan Kallus, Xiaojie Mao, and Angela Zhou. Assessing algorithmic fairness with unobserved protected class using data combination. *Management Science*, 2021.

[32] Michael Kearns, Seth Neel, Aaron Roth, and Zhiwei Steven Wu. Preventing fairness gerrymandering: Auditing and learning for subgroup fairness. In *International Conference on Machine Learning*, pages 2564–2572. PMLR, 2018.

[33] Edward H Kennedy. Optimal doubly robust estimation of heterogeneous causal effects. *arXiv preprint arXiv:2004.14497*, 2020.

[34] Gary King. A solution to the ecological inference problem. In *A Solution to the Ecological Inference Problem*. Princeton University Press, 2013.

[35] Jon Kleinberg, Himabindu Lakkaraju, Jure Leskovec, Jens Ludwig, and Sendhil Mullainathan. Human decisions and machine predictions. *The quarterly journal of economics*, 133(1):237–293, 2018.

[36] Sören R Künzel, Jasjeet S Sekhon, Peter J Bickel, and Bin Yu. Metalearners for estimating heterogeneous treatment effects using machine learning. *Proceedings of the national academy of sciences*, 116(10):4156–4165, 2019.

[37] Himabindu Lakkaraju, Jon Kleinberg, Jure Leskovec, Jens Ludwig, and Sendhil Mullainathan. The selective labels problem: Evaluating algorithmic predictions in the presence of unobservables. In *Proceedings of the 23rd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, pages 275–284, 2017.

[38] Charles F Manski. *Identification problems in the social sciences*. Harvard University Press, 1995.

[39] Charles F Manski. The mixing problem in programme evaluation. *The Review of Economic Studies*, 64(4):537–553, 1997.

[40] Charles F Manski. Monotone treatment response. *Econometrica*, pages 1311–1334, 1997.

[41] Charles F Manski. *Partial identification of probability distributions*, volume 5. Springer, 2003.

[42] Shira Mitchell, Eric Potash, Solon Barocas, Alexander D'Amour, and Kristian Lum. Algorithmic fairness: Choices, assumptions, and definitions. *Annual Review of Statistics and Its Application*, 8:141–163, 2021.

[43] Francesca Molinari. Microeconometrics with partial identification. *Handbook of econometrics*, 7:355–486, 2020.

[44] Xinkun Nie and Stefan Wager. Quasi-oracle estimation of heterogeneous treatment effects. *Biometrika*, 108(2):299–319, 2021.

[45] Samir Passi and Solon Barocas. Problem formulation and fairness. In *Proceedings of the Conference on Fairness, Accountability, and Transparency*, pages 39–48, 2019.

[46] Geert Ridder and Robert Moffitt. The econometrics of data combination. *Handbook of econometrics*, 6:5469–5547, 2007.

[47] Paul R. Rosenbaum. *Observtional Studies*. Springer, 2002.

[48] Donald B Rubin. Comment: Which ifs have causal answers. *Journal of the American statistical association*, 81(396):961–962, 1986.

[49] Ludger Rüschendorf. Sharpness of fréchet-bounds. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 57(2):293–302, 1981.

[50] Anton Schick. On asymptotically efficient estimation in semiparametric models. *Annals of Statistics*, 14(3):1139–1151, 09 1986.

[51] Vira Semenova. Debiased machine learning of set-identified linear models. *arXiv preprint arXiv:1712.10024*, 2017.

[52] Charles J Stone. Optimal global rates of convergence for nonparametric regression. *The annals of statistics*, pages 1040–1053, 1982.

[53] Elie Tamer. Partial identification in econometrics. *Annu. Rev. Econ.*, 2(1):167–195, 2010.

[54] Zhiqiang Tan. A distributional approach for causal inference using propensity scores. *Journal of the American Statistical Association*, 101(476):1619–1637, 2006.

[55] Anastasios A Tsiatis. Semiparametric theory and missing data. 2006.

[56] Aad W Van der Vaart. *Asymptotic statistics*. Cambridge University Press, 2000.

[57] Stefan Wager and Susan Athey. Estimation and inference of heterogeneous treatment effects using random forests. *Journal of the American Statistical Association*, 113(523):1228–1242, 2018.

[58] Martin J Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge University Press, 2019.

[59] Robert C Williamson and Tom Downs. Probabilistic arithmetic. i. numerical methods for calculating convolutions and dependency bounds. *International journal of approximate reasoning*, 4 (2):89–158, 1990.

[60] Steve Yadlowsky, Hongseok Namkoong, Sanjay Basu, John Duchi, and Lu Tian. Bounds on the conditional and average treatment effect with unobserved confounding factors. *arXiv preprint arXiv:1808.09521*, 2018.

[61] Wenjing Zheng and Mark J van der Laan. Cross-validated targeted minimum-loss-based estimation. In *Targeted Learning*, pages 459–474. Springer, 2011.