Online Learning and Pricing for Network Revenue Management with Reusable Resources

Abstract

We consider a price-based network revenue management problem with multiple products and multiple reusable resources. Each randomly arriving customer requests a product (service) that needs to occupy a sequence of reusable resources (servers). We adopt an incomplete information setting where the firm does not know the price-demand function for each product and the goal is to dynamically set prices of all products to maximize the total expected revenue of serving customers. We propose novel batched bandit learning algorithms for finding near-optimal pricing policies, and show that they admit a near-optimal cumulative regret bound of $\tilde{O}(J\sqrt{XT})$, where J, X, and T are the numbers of products, candidate prices, and service periods, respectively. As part of our regret analysis, we develop the first finite-time mixing time analysis of an open network queueing system (i.e., the Jackson Network), which could be of independent interest. Our numerical studies show very promising results of the proposed approaches.

1 Introduction

Network revenue management (NRM) (Gallego and Van Ryzin 1997) has attained remarkable results in a wide range of applications including airline management, electricity pricing, retailing, healthcare, and leisure (see, e.g., Williamson 1992, Talluri and Van Ryzin 2006, Özer et al. 2012). The bulk of existing literature focuses primarily on canonical settings with perishable resources in which the product once sold is permanently removed from inventory (Besbes and Zeevi 2012, Ferreira et al. 2018, Chen and Shi 2019). Little progress has been made for NRM with reusable resources, mainly due to the complexity of analyzing the underlying stochastic systems (typically manifested as stochastic queueing networks). In this paper, we develop easy-to-implement online learning and pricing policies for NRM with reusable resources shared by multiple products and analyze their theoretical and empirical performances.

1.1 Problem description

A service-providing firm is endowed with multiple types of reusable resources, each having a fixed capacity. The firm sells multiple types of products (services), each requiring the use of a sequence of different types of reusable resources (servers). (See, e.g., Figure 2.) Customers randomly arrive according to a price-dependent multivariate Poisson process to request products. Upon requesting one particular product, a customer enters the system by going through a sequence of product-specified reusable resources one by one. During the service process, if there is no available next type of resource upon finishing using the previous type in the sequence, she will join the queue in front of this next type of resource and wait until the next available unit. Both the mean demand rate of each product and the average usage time of each resource depend on the posted price vector. However, such dependence is unknown to the firm *a priori*. We note that under a fixed price vector, the service system can be reduced to the celebrated Jackson Network (Jackson 1963, 1957). The main objective of this paper is to design a "learning-while-doing" algorithm to dynamically post prices of different products, in order to maximize the expected total revenue collected over a finite time horizon.

Submitted to 36th Conference on Neural Information Processing Systems (NeurIPS 2022). Do not distribute.

It is evident that NRM with reusable resources finds many important real-world applications, including cloud computing, hospitality industry, project management, and leisure (Lei and Jasin 2020). For example, firms such as Amazon and Microsoft offer diverse cloud computing services, each requiring a different set of computational resources (virtual machines) for a certain duration of time to finish their computational jobs (see, e.g., Kaewpuang et al. 2013, Püschel et al. 2015).

1.2 Main results and contributions

We propose new batched bandit algorithms for NRM with reusable resources (RNRM). The performance measure is regret, which is the revenue difference between our learning algorithms and a clairvoyant optimal pricing policy under full information. We use J, X, T to denote the numbers of products, candidate prices, and service periods, respectively. We prove in Theorems 1 and 2 that the cumulative regret is $\tilde{O}(J\sqrt{XT})$, which matches the lower bound along the dimensions of X and T up to a logarithmic factor. It remains open whether one could improve the dependence on J since all prior studies (e.g., Miao and Wang 2021) show polynomial dependence on J at best (albeit for a non-reusable setting). We test diverse instances to show numerical performance of our approaches. The main contribution of our work is three-fold:

(Modeling) We are the first to consider the joint learning and optimization problem in NRM with reusable resources. Prior literature on reusable resources only focused on single-product settings (see Jia et al. (2020, 2022) and references therein). It is considerably more challenging to model and analyze the network setting where multiple products share multiple resources (with finite capacities), mainly due to the sequential use of resources (to finish up a service) and the evolving nature of available capacities in real-time. We establish a connection between our model and the celebrated *Jackson Network* (Jackson 1963, 1957).

(Algorithm) We first develop a linear programming (LP) upper bound on the optimal objective value given full information. The LP model provides a static pricing control policy. Therefore, we can cast our RNRM as a multi-armed bandits (MAB) problem where we select among candidate prices (analogs of arms) to minimize the cumulative regret. There are two key challenges. The first lies in the fact that whenever the system posts a new price, the arrival processes change immediately while the system still contains "old" customers from the previously posted price. Therefore, it is difficult to analyze the transient system performance and keep track of the cumulative revenue in a dynamic pricing scheme. We resolve this challenge by proposing batched algorithms. Our algorithms separate the time horizon into successive batches and select a price for each batch. We further divide each batch into two intervals. In the first interval, the system completes serving the previous customers and reaches the steady-state under the new price. In the second interval, the system maintains the steady-state (exploiting the new price). This batched framework not only allows for system stabilization (under new prices) but also allows for infrequent $O(\log T)$ price changes, which is more implementable in practical settings (see Cheung et al. 2017). The second challenge is more mathematical as we need to ensure that the regret does not scale exponentially in the number of products or resources (given the combinatorial nature of the underlying graph).

(Analysis) To cope with the first challenge mentioned above, we bound the *loss of nonstationarity* by bounding the length of the first interval of each batch. We prove that the system can reach the new steady-state reasonably fast (see mixing times of Jackson Network in Propositions 3 and 4). To the best of our knowledge, this is the first result giving a finite-time bound on mixing times of Jackson Networks. To cope with the second challenge, we decompose the resulting Jackson Network into topological layers and present a novel nested coupling argument (based on the layers). The coupling argument of Jia et al. (2020) in the single-resource single-product setting cannot be directly applied to our network setting. First, in the network of M/M/c queues, the arrival and waiting processes depend on previously visited nodes, which is significantly more complex to analyze upon price changes. To resolve this challenge, we have to carefully construct virtual and alternative queues to explicitly account for such dependency. On a high level, the difficulty lies in that the original queues have a time-varying rate from an initial rate to the stable rate upon any price change. One could construct virtual queues that could clear all previously arrived customers at a specific time τ, but it is far too strong for the coupling analysis. Instead, we construct alternative queues that start from zero initial rate and ramp up directly to the stable rate at a time stamp before τ so that the customer distribution matches the original queues in period τ. Second, even ignoring the subtleties of time-varying rates (upon price changes), if one applied the coupling argument of Jia et al. (2020) in a brute-force way to

every node in the network, the coupling probability would result in a undesirable exponential scaling in the number of products. We decompose the network into topological layers so that the resulting coupling probability only scales linearly in the number of products.

1.3 Literature review

NRM with Reusable Resources and Demand Learning. Network, demand learning, and reusable resources are three key elements of this study. We only review related literature that shares at least two of the three elements.

NRM with demand learning. Due to easy accessibility of real-time sales data, there have been recent studies on developing learning and pricing algorithms with probably near-optimal regret bound under NRM settings (see, e.g., Besbes and Zeevi 2012, Chen et al. 2019, Chen and Shi 2019, Badanidiyuru et al. 2013, Agrawal and Devanur 2016, Ferreira et al. 2018). All the aforementioned study considers the canonical settings (first defined by Gallego and Van Ryzin 1997), i.e., perishable (non-reusable) resources.

NRM with reusable resources. The only existing studies on RNRM are Owen and Simchi-Levi (2018), Rusmevichientong et al. (2020) and Lei and Jasin (2020). Lei and Jasin (2020) considered a deterministic service time assumption, and developed an asymptotically near-optimal real-time pricing control algorithm. Owen and Simchi-Levi (2018) considered a joint assortment and pricing problem under a multi-resource setting and provided constant factor performance guarantees. Rusmevichientong et al. (2020) considered a dynamic assortment and pricing problem with reusable products and proposed a provably near-optimal policy based on approximate dynamic programming.

Demand learning with reusable resources. The aforementioned literature on reusable resources all assumed known distributional information of the underlying model.There has been little work considering demand learning for service systems with reusable resources due to complex underlying system dynamics. The only studies that considered demand learning for reusable resources are Chen et al. (2020), Jia et al. (2020, 2022), and all considered only the single resource setting. Chen et al. (2020) proposed an online learning framework for a dynamic pricing and capacity sizing problem in a GI/GI/1 queue and Jia et al. (2020, 2022) developed MAB based algorithms for optimal dynamic pricing in an M/M/c queue for the single product single resource setting.

To the best of our knowledge, we are the first to thoroughly study all three elements.

Transient Analysis of Jackson Networks. Jackson Network has received much attention among the queueing networks because its steady-state distribution is particularly simple to compute as a product-form solution (Jackson 1963, 1957). The queue lengths across the nodes in Jackson Networks, i.e., a network of M/M/c queues, during the transient state can be bounded via techniques of stochastic dominance (see Massey 1986, 1984, 1987). However, how the network reacts to system changes remains an open question (see, e.g., Mohamed et al. 2003, 2005). Kelton and Law (1985) pointed out that a queueing system needs to run "long-enough" for removing the effects brought by the initial starting point and then reach an ensured "steady-state" portion of the run. The only work that explicitly developed a finite-time bound on mixing times of a queueing system is a recent paper by Jia et al. (2020). In contrast, our paper gives the first finite-time high probability bound on mixing times of Jackson Networks (consisting of a network of M/M/c queues) upon action changes.

Batched Bandit Approaches. UCB2 (Auer et al. 2002) and Improved-UCB (Auer and Ortner 2010) were implicitly M-batch policies with $M = \Theta(\log(T))$. Later, Perchet et al. (2016) studied batched MAB in the two-armed case under a static batch design and Gao et al. (2019) extended the algorithms to the general multiple-armed case and adaptively determined batch size. There is also rich literature focusing on batched MAB problems with switching cost. For example, Cesa-Bianchi et al. (2013) considered uniform switching cost in the objective and Simchi-Levi and Xu (2019) treated the finite switching cost budget as a hard constraint. In contrast, we construct the batched action framework to control the loss of the nonstationarity (resulted from the transient time after each action change).

2 Problem formulation

Consider a firm that manages $\mathcal{I}$ types of reusable resources to provide $\mathcal{J}$ types of products over a finite time horizon T. Denote the capacity of server type $i \in \mathcal{I}$ as c_i. At the beginning of each period,

the firm posts a price $\mathbf{x} = \{x_j : j \in \mathcal{J}\}$ for all products from a given price candidate set $\mathcal{X} \subset \mathbb{R}^{|\mathcal{J}|}$. Let $I = |\mathcal{I}|$, $J = |\mathcal{J}|$, and $X = |\mathcal{X}|$. Each arriving customer requests one product $j \in \mathcal{J}$, which is processed through an ordered sequence of resources. Customers arrive following a price-dependent multivariate Poisson process with a total rate $\alpha(\mathbf{x})$, which is unknown to the decision maker *a priori*. The service time of resource i follows an exponential distribution regardless of which product type it belongs to, which also depends on posted price $\mathbf{x}$. If there is not enough capacity for the current resource, the customer will join the queue of this resource until being served. The resources are released to process the next customers upon finishing serving the current ones. Customers pay a lump sum x_j to the firm for product j. This is referred to as the lump sum payment model. We also present an alternative service duration dependent payment model where the customer pays the amount that is equal to the product's per-unit-time price multiplied by its total service duration in Appendix F.

Consider a resource-consumption matrix $A_{ji} \in \mathbb{R}^{J \times I}$, where element $a_{ji} = 1$ indicates that service type j requires to be processed on server type i. We use an I-dimensional vector, $\mathbf{y} \in \{0, 1, \ldots, \infty\}^I$ to denote the number of customers across different types of resources. The firm aims at finding a pricing policy as $\pi : \{(\mathbf{y}, t) : \mathbf{y} \in \{0, 1, \ldots, \infty\}^I, t = 1, \ldots, T\} \to \mathcal{X}$, where the firm selects price $\pi(\mathbf{y}, t) \in \mathcal{X}$ for period t when there are $\mathbf{y}$ customers in the system at the beginning of period t, to maximize the expected cumulative revenue Y^π.

Assumption 1. *("Turn-off" price.) For each product $j \in \mathcal{J}$, there exists a "turn-off" price $\bar{p}_j < \infty$ such that, if the posted price is larger than this turn-off price, then the demand of this product is zero.*

Assumption 1 is widely assumed in NRM studies (see, e.g., Lei and Jasin 2020). This turn-off price can be used to effectively turn off demand of undesired products whenever needed. Therefore, for each candidate price $\mathbf{x} \in \mathcal{X}$, we can further decompose the products into turn-on product set $\mathcal{J}_{\mathrm{on}}(\mathbf{x})$ and a turn-off product set $\mathcal{J}_{\mathrm{off}}(\mathbf{x})$, where the latter can be empty.

Assumption 2. *There exists an order of server types $\mathcal{I}$ such that the processing sequence of servers $i_1, i_2, \ldots, i_n$ always has $i_1 < i_2 < \ldots < i_n$ for any product $j \in \mathcal{J}$.*

One can understand Assumption 2 via viewing a network. Consider a graph with resources as nodes and draw a directed arc from node i to i' if any product requires successive processing on resources i and i'. If the system satisfies Assumption 2, then this graph is acyclic, i.e., no cycles, and there exists at least one topological order such that all arcs are from smaller index to larger index. Without loss of generality, we consider one such topological order when referring to resource i in the rest part of this paper. To better utilize this network representation, we also add a source node s to represent customers entering the system and a sink node t to represent customers exiting the system. Thus, the consumption matrix can be extended to $A_{ji} \in \mathbb{R}^{J \times (I+2)}$ and $a_{js} = a_{jt} = 1$ for all products.

Assumption 3. *(Product-form demand.) Any given price vector $\mathbf{x}$ induces a (Markovian) routing matrix $P(\mathbf{x})$ where its (i, i')th entry $p_{i,i'}$ represents the product demand portion who will consume resource i' next out of those who consume resource i. Then the demand rate for any product $j \in \mathcal{J}$, represented by a sequence of resources $s, i_1, i_2, \ldots, i_n, t$, is given by*

$$\alpha_j = \alpha(\mathbf{x}) p_{s,i_1} p_{i_1,i_2} \ldots p_{i_{n-1},i_n} p_{i_n,t},$$

which means that the demand rate for product j is equal to the total demand rate $\alpha(\mathbf{x})$ times a sequence of routing probabilities of $P(\mathbf{x})$, under posted price vector $\mathbf{x}$.

We use Figure 2 in the Appendix to demonstrate a concrete example. With Assumptions 2 and 3, the service system under any posted price $\mathbf{x} \in \mathcal{X}$ can be reduced to a Jackson Network consisting with I number of M/M/c queues (when the system only contains customers arriving under this price, i.e., after finishing serving other customers arriving under the previously posted prices). The main idea is that there exists a mapping $\mathbf{x} \mapsto \alpha(\mathbf{x}), P(\mathbf{x})$ that gives rise to product demand pattern. Note that our assumptions are mild in the sense that this mapping can be nonparametric. The decision maker does not know this underlying mapping and needs to learn it while maximizing revenue on the fly.

Given any price vector $\mathbf{x}$, (i) on one hand, this price affects the total future demand $\alpha(\mathbf{x})$ entering the system. (ii) On the other hand, this price affects the internal competition between the products, which is reflected in the server routing matrix $P(\mathbf{x})$. As a result, $\alpha(\mathbf{x})$ and $P(\mathbf{x})$ define a Jackson Network well and the demand rate $\alpha_j(\mathbf{x})$ of each product type is also determined (use vector $\vec{\alpha}_{\mathbf{x}} \in \mathbb{R}^J$ to denote demand of products). We use $\lambda_i(\mathbf{x})$ to denote the demand rate of resource type i.

Assumption 4. *(Stability.) The queueing system of each resource is stable under any price, i.e., $c_i \mu_i(\mathbf{x}) > \lambda_i(\mathbf{x})$. Therefore, the Jackson Network is stable as well.*

194 Assumption 4 ensures that the underlying Jackson Network is stable under any candidate price (i.e.,
195 the queue length will not grow indefinitely). This assumption can be readily satisfied by putting
196 appropriate lower bounds on the price vector.

197 **Assumption 5.** *(Change of rates.) For any node, consider the arrival and service rates (λ_1, μ_1) and*
198 (λ_2, μ_2) *under* $\mathbf{x}_1,\ \mathbf{x}_2 \in \mathcal{X}$ *where* $\lambda_1 \geq \lambda_2$, *also denote the utilization factor by* $\rho_1 = \lambda_1/\mu_1$ *and*
199 $\rho_2 = \lambda_2/\mu_2$, *respectively, then 1)* $\lambda_1 - \lambda_2 \leq \frac{\rho_1}{-3e \log \rho_1}$ *and 2) if* $\mu_1 > \mu_2$, *then* $\rho_1 \geq \rho_2^2$.

200 This assumption is used to justify Lemmas 3 and 4 (used in the proof of Propositions 3 and 4) and the
201 original proofs are in Jia et al. (2020). These conditions are mild and easy to validate in practice in the
202 following sense. 1) The difference in arrival rates between two prices is bounded (by a large constant).
203 This can be easily satisfied by interpolating intermediate prices in the candidate price set $\mathcal{X}$. 2) If
204 both the service and arrival rate under a price is higher than or equal to another price, the utilization
205 factor in the former case must be greater than the latter squared. A simple sufficient condition to
206 guarantee $\rho_1 \geq \rho_2^2$ is $\rho_1 \geq \rho_2$, which reduces to $\lambda_1/\lambda_2 \geq \mu_1/\mu_2$, and thus the practitioners only
207 need to validate that the ratio of change in arrival rates exceeds the ratio of change in service rates.

208 **Assumption 6.** *We assume* $\log T \geq 4$ *(to ensure sufficient times for mixing).*

3 Regret, relaxed regret, and static price benchmark

210 Under the full information scenario, where the firm knows the underlying arrival rate $\alpha_j(\mathbf{x})$ of all
211 products $j \in \mathcal{J}$ under any candidate price $\mathbf{x} \in \mathcal{X}$. We denote a state-dependent optimal policy as π^*,
212 where $\pi^* = \arg\max_\pi Y^\pi$, and let $Y^* = Y^{\pi^*}$. Therefore, the regret of any pricing policy π can be
213 defined as $\text{Regret}(\pi, T) = Y^* - Y^\pi$. In short, we aim at finding heuristic pricing policies that lead
214 to a small regret. However, finding the state-dependent optimal pricing policy π^* is computationally
215 intractable even under the full information scenario. Firstly, it requires to solve a dynamic program
216 with an infinite number of potential states, in the form of $(\mathbf{y}, t)$, which suffers from the curse of
217 dimensionality. Moreover, the transient dynamics of a queueing network is complex and hard to
218 analyze especially when the offered price changes over periods. Thus, it is impossible to obtain π^*
219 and Y^*. To tackle this problem, we develop an upper bound on the expected revenue achievable by
220 the state-dependent optimal expected revenue (see a similar approach in Jia et al. 2020).

221 **An LP upper bound.** Consider continuous decision variables $\pi^{\mathbf{x}}_{\mathbf{y}t}$ for $\mathbf{y} \in \{0, 1, \ldots, \infty\}^I$, $t =$
222 $1, \ldots, T$, $\mathbf{x} \in \mathcal{X}$, $a_{\mathbf{y}t}$ for $\mathbf{y} \in \{0, 1, \ldots, \infty\}^I$, $t = 1, \ldots, T$, and $Y_{\mathbf{y}t}$ for $\mathbf{y} \in \{0, 1, \ldots, \infty\}^I$, $t =$
223 $1, \ldots, T$. We can formulate a linear program (LP) as follows.

$$Y^{\text{LP}} = \max_{\pi,a,J} \sum_{\mathbf{y} \in \{0,1,\ldots,\infty\}^I} \sum_{t=1}^{T} a_{\mathbf{y}t} Y_{\mathbf{y}t} \tag{1a}$$

$$\text{s.t.} \sum_{\mathbf{y} \in \{0,1,\ldots,\infty\}^I} a_{\mathbf{y}t} = 1, \quad \forall t = 1, \ldots, T \tag{1b}$$

$$\sum_{\mathbf{x} \in \mathcal{X}} \pi^{\mathbf{x}}_{\mathbf{y}t} = 1 \quad \forall \mathbf{y} \in \{0, 1, \ldots, \infty\}^I, \ \forall t = 1, \ldots, T \tag{1c}$$

$$Y_{\mathbf{y}t} \leq \sum_{\mathbf{x} \in \mathcal{X}} \pi^{\mathbf{x}}_{\mathbf{y}t} \cdot \vec{\alpha}_{\mathbf{x}}^T \mathbf{x} \quad \forall \mathbf{y} \in \{0, 1, \ldots, \infty\}^I, \ \forall t = 1, \ldots, T \tag{1d}$$

$$0 \leq a_{\mathbf{y}t} \leq 1 \quad \forall \mathbf{y} \in \{0, 1, \ldots, \infty\}^I, \ \forall t = 1, \ldots, T \tag{1e}$$

$$0 \leq \pi^{\mathbf{x}}_{\mathbf{y}t} \leq 1 \quad \forall \mathbf{y} \in \{0, 1, \ldots, \infty\}^I, \ \forall t = 1, \ldots, T, \ \forall \mathbf{x} \in \mathcal{X}. \tag{1f}$$

224 **Proposition 1.** *The LP solution provides an upper bound on the optimal revenue, i.e.,* $Y^* \leq Y^{LP}$,
225 *and* $Y^{LP} = T \cdot \vec{\alpha}_{\tilde{\mathbf{x}}}^T \tilde{\mathbf{x}}$ *where* $\tilde{\mathbf{x}} = \arg\max_{\mathbf{x} \in \mathcal{X}} \vec{\alpha}_{\mathbf{x}}^T \mathbf{x}$.

226 *Proof Sketch.* Consider an arbitrary policy $\hat{\pi}$. Let variable $\hat{a}_{\mathbf{y}t}$ take the value of probability that the
227 system has $\mathbf{y}$ customers at the beginning of period t for $\mathbf{y} \in \{0, 1, \ldots, \infty\}^I$ and $t = 1, \ldots, T$. Let
228 variable $\hat{\pi}^{\mathbf{x}}_{\mathbf{y}t} = 1$ if the policy $\hat{\pi}$ chooses price $\mathbf{x}$ at state $(\mathbf{y}, t)$, and 0 otherwise. Let variable $\hat{Y}_{\mathbf{y}t}$
229 take the value of expected revenue collected before the end of period T from the customers who
230 arrive during period t, when the system has $\mathbf{y}$ customers at the beginning of period t. Then we can
231 show that the decisions variables (π, a, Y) associated with any admissible policy must satisfy the

constraints of the linear program (1), and the expected revenue of this admissible policy is exactly the corresponding objective value in (1). We can readily check that $T \cdot \vec{\alpha}_{\tilde{\mathbf{x}}}^T \tilde{\mathbf{x}}$ is an upper bound of Y^{LP}. We can construct a solution which can obtain this upper bound and thus $Y^{\text{LP}} = T \cdot \vec{\alpha}_{\tilde{\mathbf{x}}}^T \tilde{\mathbf{x}}$. $\qquad\square$

Definition 1. *The relaxed regret is defined as* $\overline{Regret}(\pi, T) = Y^{LP} - Y^{\pi}$.

Based on Proposition 1, we have for any policy π, the corresponding relaxed regret is a valid upper bound of its regret, i.e., $\overline{\text{Regret}}(\pi, T) \geq \text{Regret}(\pi, T)$, and it suffices to bound the relaxed regret.

4 Batched bandit algorithms

The basis for implementing an Upper Confidence Bound (UCB) type of algorithm is identifying an upper confidence bound of the expected collected revenue of each candidate price. We start with the steady-state revenue rate. We can show that this steady-state revenue rate can be used to compute a lower bound of the expected revenue during consecutive periods (see proof of Theorem 1). Therefore, it is equivalent to deriving an upper confidence bound of the steady-state revenue rate based on historical sales information. Jia et al. (2020) established a concentration bound of the steady-state revenue rate under the single product setting. Here we develop a concentration bound for the multi-product multi-resource setting.

We use $n_j(\mathbf{x})$ to denote the number of arrived customers requesting product j under price $\mathbf{x}$. We use $\hat{d}_{jk}(\mathbf{x})$ to denote the observed time interval between the arrival time of customer k and that of customer $k-1$ who are requesting product j under the price $\mathbf{x}$. We use $\bar{d}_j(\mathbf{x})$ to denote the empirical mean of the inter-arrival time and thus $\bar{d}_j(\mathbf{x}) = \sum_{k=1}^{n_j(\mathbf{x})} \hat{d}_{jk}(\mathbf{x})/n_j(\mathbf{x})$, for $j \in \mathcal{J}$.

Proposition 2. *For any price* $\mathbf{x} \in \mathcal{X}$, *if the number of arrival time observations for each turn-on product* $n_j(\mathbf{x}) \geq 8 \log(J^{\frac{1}{4}}T)$ *for* $j \in \mathcal{J}_{on}(\mathbf{x})$, *then we have*

$$\mathbb{P}\left(\left|\vec{\alpha}_{\mathbf{x}}^T \mathbf{x} - \sum_{j \in \mathcal{J}_{on}(\mathbf{x})} \frac{x_j}{\bar{d}_j(\mathbf{x})}\right| \leq \sum_{j \in \mathcal{J}_{on}(\mathbf{x})} \frac{x_j \sqrt{32 \log(J^{\frac{1}{4}}T)}}{\bar{d}_j(\mathbf{x})\sqrt{n_j(\mathbf{x})}}\right) \geq 1 - \frac{2}{T^4}.$$

The proof of Proposition 2 is based on the concentration inequality for sub-exponential random variables (see, e.g., Boucheron et al. 2013, Jia et al. 2020) and is provided in Appendix C. Note that $\vec{\alpha}_{\mathbf{x}}^T \mathbf{x}$ is the steady-state revenue rate and denote it by $r(\vec{\alpha}_{\mathbf{x}}^T, \mathbf{x})$. Proposition 2 says that with a certain amount of arrival time observations of turn-on products under a candidate price, we can derive a valid upper bound of the steady-state revenue rate with a high probability guarantee. Hence, we define the corresponding upper and lower confidence bounds of the stationary revenue rate associated with price $\mathbf{x}$ by the end of period t as follows.

Definition 2. *Define* $\mathbf{Rad}_t(\mathbf{x}) = \sum_{j \in \mathcal{J}_{on}(\mathbf{x})} \frac{x_j \sqrt{32 \log(J^{\frac{1}{4}}T)}}{\bar{d}_j(\mathbf{x})\sqrt{n_j(\mathbf{x})}}$ *as the confidence radius by the end of period* t. *The upper and lower confidence bounds of the revenue rate associated with price* $\mathbf{x}$ *are:*

$$U_t(\mathbf{x}) = \sum_{j \in \mathcal{J}_{on}(\mathbf{x})} \frac{x_j}{\bar{d}_j(\mathbf{x})} + \mathbf{Rad}_t(\mathbf{x}), \qquad L_t(\mathbf{x}) = \sum_{j \in \mathcal{J}_{on}(\mathbf{x})} \frac{x_j}{\bar{d}_j(\mathbf{x})} - \mathbf{Rad}_t(\mathbf{x}). \qquad (2)$$

4.1 RNRM-UCB

With Definition 2, we can leverage the idea of UCB to solve RNRM. We separate T periods into two phases. The first phase is the *Warm-up* Phase, where we offer each candidate price one by one until we collect $8 \log(J^{1/4}T)$ number of arrival time intervals for each turn-on product $j \in \mathcal{J}_{on}(\mathbf{x})$ under each price. Therefore, the total length of this Warm-up Phase is $O(X \log(J^{1/4}T))$. After this Warm-up Phase, Proposition 2 is valid to use. The second phase is the *Learning* Phase. The UCB algorithm cannot be directly used because in order to estimate the expected revenue collected when offering a price, we have to wait until the system reaches the steady-state. The key idea is to reduce the number of price changes and further reduce the intractable transient state performance. We separate the Learning Phase into M number of batches. Define $I_m = 2^m$, $\forall m = 1, \ldots, M$. Each batch m contains $I_m L\tau$ periods, where $\tau = (\log T)^2$. Therefore, the batch size is exponentially increasing.

We can then add m in the subscript of notation $(n_{mj}(\mathbf{x}), \hat{d}_{jk}(\mathbf{x}), \bar{d}_{mj}(\mathbf{x}), \mathbf{Rad}_m(\mathbf{x}), U_m(\mathbf{x}), L_m(\mathbf{x}))$ to represent their values by the end of batch m. At the beginning of each batch $m + 1$, RNRM-UCB chooses price $\mathbf{x}$ with the highest $U_m(\mathbf{x})$ and offers this price during batch $m + 1$. We present the algorithmic details in Algorithm 1.

Algorithm 1 RNRM-UCB.

1: Input: T, $\mathcal{X}$, service network, turn-off price.
2: Initialize: L, τ, I_m, M, $m = 0$.
3: Warm-up Phase:
4: **for** $\mathbf{x} \in \mathcal{X}$ **do**
5:    Offer $\mathbf{x}$, record inter-arrival time $\hat{d}_{jk}(\mathbf{x})$ for arriving customers of product $j \in \mathcal{J}_{\text{on}}(\mathbf{x})$.
6:    **if** $n_{mj}(\mathbf{x}) \geq 8 \log(J^{1/4}T)$ **then**
7:       Continue.
8:    **end if**
9: **end for**
10: Compute and update UCB for all prices as defined in (2).
11: Learning Phase:
12: **for** $m = 1, \ldots, M$ **do**
13:    Choose $\mathbf{x}_m = \text{argmax}_{\mathbf{x} \in} U_{m-1}(\mathbf{x})$.
14:    Offer $\mathbf{x}_m$ in batch m, i.e., for $I_m L \tau$ periods.
15:    Record inter-arrival time $\hat{d}_{jk}(\mathbf{x}_m)$ for arriving customers of product $j \in \mathcal{J}_{\text{on}}(\mathbf{x}_m)$.
16:    Compute and update UCB of $\mathbf{x}_m$ as defined in (2).
17: **end for**

4.2 RNRM-TS

Considering the firm holds prior information about the rates, then we can add the Bayesian assumption to this RNRM problem. By the conjugate prior theory (see, e.g., Minton et al. 1962, Diaconis and Ylvisaker 1979), we assume that RNRM-TS holds Gamma priors on $\alpha_j(\mathbf{x})$ of product $j \in \mathcal{J}_{\text{on}}(\mathbf{x})$, $\forall \mathbf{x} \in \mathcal{X}$ and the posteriors are updated by the end of batch m, $\forall m = 1, \ldots, M$. Specifically, we set the prior distribution of $\alpha_j(\mathbf{x})$ as $\mathbf{Gamma}(\eta^{\alpha_j(\mathbf{x})}, \beta^{\alpha_j(\mathbf{x})})$. After observing one arrival time interval $\hat{d}_{jk}(\mathbf{x})$ of product j while offering price $\mathbf{x}$, the posterior distribution of $\alpha_j(\mathbf{x})$ is computed as $\mathbf{Gamma}(\eta^{\alpha_j(\mathbf{x})} + 1, \beta^{\alpha_j(\mathbf{x})} + \hat{d}_{jk}(\mathbf{x}))$. To implement RNRM-TS, at the beginning of batch m, we sample parameters $\hat{\alpha}_j(\mathbf{x})$ for product $j \in \mathcal{J}_{\text{on}}(\mathbf{x})$, $\forall \mathbf{x} \in \mathcal{X}$, following the updated posterior distributions. Then we offer the price with the maximum sampled stationary network revenue rate $r(\hat{\bar{\alpha}}_{\mathbf{x}}, \mathbf{x}) = \sum_{j \in \mathcal{J}_{\text{on}}(\mathbf{x})} \mathbf{x}_j \hat{\alpha}_j(\mathbf{x})$. We present the algorithmic details of RNRM-TS in Appendix.

5 Performance analysis

5.1 High probability bound for mixing time

When the firm starts service from an empty state, i.e., zero customers in the network service system, or switches to another price from the currently offered price, the service (queueing network) system enters a transient state and needs a certain amount of time to reach the steady-state again (referred to as the mixing time). Recall that the service system can be reduced to a Jackson Network under a specific offered price vector, upon completion of existing customers under the previously posted price. Therefore, we evaluate how quickly a Jackson Network can reach the steady-state. To the best of our knowledge, this is the first result giving a finite-time high probability bound on mixing times of Jackson Network system (upon action changes). Technically, we develop a new coupling argument between this target system and a virtual system starting from a state sampled from the steady-state distribution and maintaining the steady-state thereafter.

Poisson arrival and departure under steady-state. Under the steady-state of an M/M/c queue (if it exists, i.e., $\rho = \lambda/c\mu < 1$), then the process following which the customers leaving the system is also a Poisson process with rate λ. Therefore, one can easily derive that the arrival as well as the departure processes of each type of resource (each node in the network representation) are Poisson processes with the same rate λ_i under the steady-state (see the famous Burke's Theorem Burke 1956).

Jia et al. (2020) is the only work for studying finite-time bounds on the mixing time for an M/M/c queue. However, their results cannot be directly applied here because (i) the arrival process of each resource is not a well-defined Poisson process and (ii) the starting point of the later queues is not zero state nor a steady-state of old system parameters. Even with these two challenges tackled, this approach will induce a high-probability mixing time which is linear on the number of resources. This result is not desired for systems with multiple (especially large) number of resources. We develop a new mixing time analysis for Jackson Network. We divide resource of the service network into layers and derive a high-probability mixing time that is only linear on the number of layers.

Definition 3. *We divide the nodes (types of servers) into layers indexed by the maximum number of previously occupied nodes (including the virtual source node) along the service routes of all products. For example, Nodes 1 and 2 of the illustrative example (see Figure 2) is Layer-1 nodes and Node 3 is a Layer-2 node. We denote the number of layers by L and the set of Layer-l node as $\mathcal{I}_l$.*

Proposition 3. *(Coupling probability of a Jackson Network starting from empty state) Define $\mathcal{A}_t$ to be the event in which the Jackson Network reaches the steady-state by time t. For $t \geq L\tau$, where $\tau = (\log T)^2$, we have $\mathbb{P}(\mathcal{A}_t) \geq 1 - \frac{2I}{T^2}$.*

Proof Sketch. We use the aforementioned Poisson arrival and departure property of the nodes in a Jackson Network to bound the mixing time of the nodes layer by layer. We define events $\mathcal{A}_{t',l} = \{$all layer-l nodes reach the steady-state by time $t'\}$ and $\mathcal{A}_{t'}^i = \{$the node i reaches the steady-state by time $t'\}$. By treating the nodes in the same layer together,

$$\mathbb{P}(\mathcal{A}_t) = \mathbb{P}(\mathcal{A}_{L\tau,1}\mathcal{A}_{L\tau,2}\mathcal{A}_{L\tau,3}\ldots\mathcal{A}_{L\tau,L}) \geq \mathbb{P}(\mathcal{A}_{\tau,1})\prod_{l=2}^{L}\mathbb{P}(\mathcal{A}_{l\tau,l}|\mathcal{A}_{\tau,1}\ldots\mathcal{A}_{(l-1)\tau,l-1}).$$

We then prove that $\mathbb{P}(\mathcal{A}_{l\tau,l}|\mathcal{A}_{\tau,1}\ldots\mathcal{A}_{(l-1)\tau,l-1}) \geq 1 - (2I_l)/T^2$, where $I_l = |\mathcal{I}_l|$, by establishing a coupling argument between the target queue (with the help of an artificially constructed alternative queue) and a virtual queue starting from a state sampled from the steady-state distribution and maintaining the steady-state thereafter. The detailed proof is provided in Appendix E. $\square$

In addition, the transient state of the Jackson Network, as well as each resource queue (or each node), can be induced not only by starting from empty state but also by starting from a steady-state of a previously posted price. It is possible that when a new price is posted, the customers arrived under the previously posted price still remain in the system. Therefore we also need to ensure that the coupling also occurs with high probability when price changes.

Proposition 4. *(Coupling probability of Jackson Network starting from steady-state of previous price.) Define $\mathcal{B}_t$ to be the event in which the Jackson Network reaches the steady-state by time t. For $t \geq 2L\tau$, where $\tau = (\log T)^2$, we have $\mathbb{P}(\mathcal{B}_t) \geq 1 - \frac{4I}{T^2}$.*

Proof Sketch. This proposition is derived based on Proposition 3 and Assumption 5. We define $\mathcal{B}_{t',l}$ and $\mathcal{B}_{t'}^i$ similarly and bound $\mathbb{P}(\mathcal{B}_{2l\tau,l}|\mathcal{B}_{2\tau,1}\ldots\mathcal{B}_{2(l-1)\tau,l-1}) \geq 1 - \frac{4I_l}{T^2}$ by decomposing the scenarios by how the arrival and service rates are changing when considering each resource queue. $\square$

5.2 Regret bounds of RNRM-UCB and RNRM-TS

The high probability bounds for mixing time of Jackson Network developed in §5.1 help bound the loss due to the nonstationarity, and thus we are ready to carry out the regret analysis of RNRM-UCB.

Theorem 1. *The T-period cumulative regret of RNRM-UCB is bounded by $\tilde{O}\left(J\sqrt{XT}\right)$.*

Note that the regret lower bound for nominal MAB (where $J = 1$) is $\Omega(\sqrt{XT})$ (see, e.g., Theorem 2.1 in Slivkins (2019)), which suggests that our regret bound is tight, if the system size J is fixed.

Proof Sketch. The regret of the Warm-up phase is at most linear on the length of the Warm-up Phase, i.e., $O(XJ\log(J^{1/4}T))$. We decompose the relaxed regret in the Learning Phase into two parts. The first part is called *loss of nonstationarity* and it represents the revenue loss induced by the transient performance when changing the price. For batch m, we can derive that the expected revenue collected under the selected price $\mathbf{x}_m$ as $r(\vec{\alpha}_{\mathbf{x}_m}, \mathbf{x}_m)I_m L\tau - O(L\tau)$ with Proposition 4. The second part represents the *loss of suboptimality*, which is induced by not offering the optimal state-independent price in certain time periods. This part can be further rewritten as $\Delta(\mathbf{x}_m)I_m L\tau$ and

346 $\Delta(\mathbf{x}_m) = (r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) - U_m(\tilde{\mathbf{x}})) + (U_m(\mathbf{x}_m) - L_m(\mathbf{x}_m)) + (L_m(\mathbf{x}_m) - r(\vec{\alpha}_{\mathbf{x}_m}, \mathbf{x}_m))$. We bound
347 these three terms separately based on Proposition 2 and Lemma 2. Combine all results together, we
348 reach a $\tilde{O}(J\sqrt{XT})$ regret bound. We provided detailed proof in Appendix E. $\qquad\square$

349 We use *Bayesian regret* as the performance measure of RNRM-TS, which is typically used for the
350 Bayesian setting and is computed as the expected regret of the instance in expectation over all problem
351 instances. Similarly, we define a relaxed Bayesian regret:.

Definition 4. *Denoting all the events before period t by $\mathcal{F}_t$, $\forall t \geq 0$, we define the relaxed Bayesian regret of policy π by period T as*
$$\overline{BR}(\pi, T) = \mathbb{E}_{\{\alpha_j(\mathbf{x})\}_{j\in\mathcal{J}}\sim priori}\big[\mathbb{E}[Y^{LP} - Y^\pi | \mathcal{F}_T]\big].$$

352 The relaxed Bayesian regret bound of RNRM-TS is in the same order as RNRM-UCB (Appendix E).

353 **Theorem 2.** *The T-period cumulative Bayesian regret of RNRM-TS is bounded by $\tilde{O}(J\sqrt{XT})$.*

6 Numerical results

355 We consider $T = 2000$, $I = 2$ with capacity $c_1 = c_2 = 2$, $J = 3$ with product-resource orders
356 $\{1, 2, 1 \to 2\}$, and $\mathcal{X} = \{[1,2,3], [2,1,3], [2,2,3]\}$. We set the underlying arrival rates of candidate
357 prices as $\{[4,2,4], [3,4,3], [2.5,3,2.5]\}$ and use a uniform service rates $[5, 4.5]$ for all prices. We
358 adopt the blind network revenue management algorithm proposed in Besbes and Zeevi (2012) to
359 our settings as a benchmark. This benchmark algorithm tries each price for $\theta T/X$ periods and then
360 selects the price with the highest revenue in the rest of the periods. We set θ as 0.22, 0.1, and 0.05
361 in three Besbes and Zeevi (2012) benchmarks, respectively (the same as the settings in Table 1 in
362 the original paper Besbes and Zeevi 2012). To conclude, we implement five pricing algorithms:
363 RNRM-UCB, RNRM-TS, and three benchmark policies with different values of θ.

364 Figure 1a shows the offered price over periods of each algorithm and Figure 1b depicts the cumulative
365 time-average relaxed regret, i.e., $\sum_{t'=1}^{t}(Y_{t'}^{\text{LP}} - Y_{t'}^{\pi})/t$. For all algorithms (except "OPT"), the revenue
366 is collected from the simulation where the system starts with no customers at time 0. We can observe
367 that both RNRM-UCB and RNRM-TS identify the best price (P1) through the learning process and
368 the time average regret shrinks to zero very quickly.

(a) The offered prices.

(b) The time-average cumulative regret.

Figure 1: Offered price and time-average cumulative regret of policies. The price selections keep the same after Period 750 and we only plot for Periods 1 to 750 in subfigure (a).

7 Conclusion and social impacts

370 We proposed two new batch bandit algorithms for RNRM problems with multiple products and
371 multiple reusable resources. We established optimal regret upper bounds and demonstrated superior
372 computational results. This work solves an important practical problem and we believe that our
373 approach will open many doors to learning more complex stochastic systems (especially in the
374 network settings). There are two immediate future research questions: (1) how we can accommodate
375 continuous price space; and (2) how we can extend arrivals and services to general distributions. Both
376 directions would need significant methodological innovations to be developed.

References

Agrawal S, Devanur N (2016) Linear contextual bandits with knapsacks. *Advances in Neural Information Processing Systems* 29.

Auer P, Cesa-Bianchi N, Fischer P (2002) Finite-time analysis of the multiarmed bandit problem. *Machine learning* 47(2):235–256.

Auer P, Ortner R (2010) Ucb revisited: Improved regret bounds for the stochastic multi-armed bandit problem. *Periodica Mathematica Hungarica* 61(1-2):55–65.

Badanidiyuru A, Kleinberg R, Slivkins A (2013) Bandits with knapsacks. *2013 IEEE 54th Annual Symposium on Foundations of Computer Science*, 207–216 (IEEE).

Besbes O, Zeevi A (2012) Blind network revenue management. *Operations Research* 60(6):1537–1550.

Boucheron S, Lugosi G, Massart P (2013) *Concentration inequalities: A nonasymptotic theory of independence* (Oxford university press).

Burke PJ (1956) The output of a queuing system. *Operations Research* 4(6):699–704.

Cesa-Bianchi N, Dekel O, Shamir O (2013) Online learning with switching costs and other adaptive adversaries. *Advances in Neural Information Processing Systems*, 1160–1168.

Chen Q, Jasin S, Duenyas I (2019) Nonparametric self-adjusting control for joint learning and optimization of multiproduct pricing with finite resource capacity. *Mathematics of Operations Research* 44(2):601–631.

Chen X, Liu Y, Hong G (2020) An online learning approach to dynamic pricing and capacity sizing in service systems. Technical report, Chinese University of Hong Kong, Shenzhen, China. Available at arXiv:2009.02911.

Chen Y, Shi C (2019) Network revenue management with online inverse batch gradient descent method. Technical report, Forthcoming in Productions and Operations Management. Available at SSRN 3331939.

Cheung WC, Simchi-Levi D, Wang H (2017) Dynamic pricing and demand learning with limited price experimentation. *Operations Research* 65(6):1722–1731.

Diaconis P, Ylvisaker D (1979) Conjugate priors for exponential families. *The Annals of Statistics* 269–281.

Ferreira KJ, Simchi-Levi D, Wang H (2018) Online network revenue management using Thompson sampling. *Operations Research* 66(6):1586–1602.

Gallego G, Van Ryzin G (1997) A multiproduct dynamic pricing problem and its applications to network yield management. *Operations Research* 45(1):24–41.

Gao Z, Han Y, Ren Z, Zhou Z (2019) Batched multi-armed bandits problem. *Advances in Neural Information Processing Systems* 32.

Jackson JR (1957) Networks of waiting lines. *Operations Research* 5(4):518–521.

Jackson JR (1963) Jobshop-like queueing systems. *Management Science* 10(1):131–142.

Jia H, Shi C, Shen S (2020) Online learning and pricing for service systems with reusable resources. Technical report, University of Michigan at Ann Arbor, MI. Available at SSRN 3755902.

Jia H, Shi C, Shen S (2022) Online learning and pricing with reusable resources: Linear bandits with sub-exponential rewards. Technical report, University of Michigan at Ann Arbor, MI. Forthcoming in ICML 2022.

Kaewpuang R, Niyato D, Wang P, Hossain E (2013) A framework for cooperative resource management in mobile cloud computing. *IEEE Journal on Selected Areas in Communications* 31(12):2685–2700.

Kelton WD, Law AM (1985) The transient behavior of the m/m/s queue, with implications for steady-state simulation. *Operations Research* 33(2):378–396.

Lei Y, Jasin S (2020) Real-time dynamic pricing for revenue management with reusable resources, advance reservation, and deterministic service time requirements. *Operations Research* 68(3):676–685.

Massey WA (1984) An operator-analytic approach to the jackson network. *Journal of Applied Probability* 21(2):379–393.

Massey WA (1986) A family of bounds for the transient behavior of a jackson network. *Journal of Applied Probability* 23(2):543–549.

Massey WA (1987) Stochastic orderings for markov processes on partially ordered spaces. *Mathematics of Operations Research* 12(2):350–367.

Miao S, Wang Y (2021) Network revenue management with nonparametric demand learning: $\sqrt{T}$-regret and polynomial dimension dependency. Technical report, McGill University, Canada. Available at SSRN 3948140.

Minton P, Raiffa H, Schlaifer R (1962) Applied statistical decision theory. *American Mathematical Monthly* 69:72.

Mohamed AM, Lipsky L, Ammar R (2003) Transient model for jackson networks and its approximation. *International Conference On Principles Of Distributed Systems*, 198–209 (Springer).

Mohamed AM, Lipsky L, Ammar R (2005) Modeling parallel and distributed systems with finite workloads. *Performance Evaluation* 60(1-4):303–325.

Owen Z, Simchi-Levi D (2018) Price and assortment optimization for reusable resources. Technical report, Massachusetts Institute of Technology, Cambridge, MA. Available at SSRN 3070625.

Özer Ö, Ozer O, Phillips R (2012) *The Oxford Handbook of Pricing Management* (Oxford University Press).

Perchet V, Rigollet P, Chassang S, Snowberg E (2016) Batched bandit problems. *The Annals of Statistics* 44(2):660–681.

Püschel T, Schryen G, Hristova D, Neumann D (2015) Revenue management for cloud computing providers: Decision models for service admission control under non-probabilistic uncertainty. *European Journal of Operational Research* 244(2):637–647.

Rusmevichientong P, Sumida M, Topaloglu H (2020) Dynamic assortment optimization for reusable products with random usage durations. *Management Science* 66(7):2820–2844.

Simchi-Levi D, Xu Y (2019) Phase transitions and cyclic phenomena in bandits with switching constraints. *Available at SSRN 3380783* .

Slivkins A (2019) Introduction to multi-armed bandits. *Foundations and Trends in Machine Learning* 12(1-2):1–286.

Talluri KT, Van Ryzin GJ (2006) *The Theory and Practice of Revenue Management*, volume 68 (Springer Science & Business Media).

Williamson EL (1992) *Airline network seat inventory control: Methodologies and revenue impacts*. Ph.D. thesis, Massachusetts Institute of Technology.

A Summary of major notation

Table 1: Summary of Major Notation for Problem Formulation

$\mathcal{I}$	the set of reusable resource (server) types
$\mathcal{J}$	the set of product (service) types
c_i	the capacity of server type $i \in \mathcal{I}$
$\mu_i(\mathbf{x})$	the service rate of server type i under the price $\mathbf{x}$
$\mathbf{x}$	the price vector $\mathbf{x} = \{x_j : j \in \mathcal{J}\}$
$\mathcal{X}$	the set of candidate prices
T	the number of periods in consideration
$\alpha(\mathbf{x})$	the overall demand rate of the system under the price $\mathbf{x}$
$\alpha_j(\mathbf{x})$	the demand rate of product j under the price $\mathbf{x}$
$\vec{\alpha}_{\mathbf{x}}$	the vector to demand of products under the price $\mathbf{x}$, $\vec{\alpha}_{\mathbf{x}} \in \mathbb{R}^J$
$\lambda_i(\mathbf{x})$	the demand rate of server i under the price $\mathbf{x}$
A	the resource consumption matrix, $A_{ji} \in \mathbb{R}^{J \times I}$, where element $a_{ji} = 1$ indicates that service type j requires to be processed on server type i
$\mathcal{J}_{\text{on}}(\mathbf{x})$	the turn-on product set under price $\mathbf{x}$, containing products with non-zero demand rates
$\mathcal{J}_{\text{off}}(\mathbf{x})$	the turn-off product set under price $\mathbf{x}$, containing products with zero demand rates
P	the server routing matrix where $p_{ii'}$ denotes the demand portion that consumes resource i' straight after consuming resource i, $\sum_{i' \in \mathcal{I} \cup \{t\}} p_{ii'} = 1$.
$\mathbf{y}$	a J-dimensional vector, $\mathbf{y} \in \{0, 1, \ldots, \infty\}^I$, denote the number of current customers across different types of resources
Y_t^π	the expected revenue under pricing policy π during period t, $\forall t = 1, \ldots, T$,
Y^π	the cumulative expected revenue over periods $\{1, \ldots, T\}$, $Y^\pi = \sum_{t=1}^T Y_t^\pi$
$\tilde{\mathbf{x}}$	the (static) optimal state-independent price, $\tilde{\mathbf{x}} = \operatorname{argmax}_{\mathbf{x} \in \mathcal{X}} \vec{\alpha}_{\mathbf{x}}^T \mathbf{x}$
$r(\vec{\alpha}_{\mathbf{x}}, \mathbf{x})$	steady-state revenue rate of price $\mathbf{x}$, $r(\vec{\alpha}_{\mathbf{x}}, \mathbf{x}) = \sum_{j \in \mathcal{J}_{\text{on}}(\mathbf{x})} \mathbf{x}_j \alpha_j(\mathbf{x})$

Table 2: Summary of Major Notation for Algorithms and Regret Analysis

L	the number of layers
$\mathcal{I}_l$	resources of layer l
M	the number of batches
τ	the value for computing the length of each batch, $\tau = (\log(T))^2$
I_m	batch m, $m = 1, \ldots, M$ contains $I_m L\tau$ periods, where $I_m = 2^m$
$n_j(\mathbf{x})$	the number of arrived customers of product j under price $\mathbf{x}$
$\hat{d}_{jk}(\mathbf{x})$	as the observed time interval between the arrival time of customer k and that of customer $k - 1$ who are requesting product j under the price $\mathbf{x}$
$\bar{d}_j(\mathbf{x})$	an empirical estimate of arrival time, $\bar{d}_j(\mathbf{x}) = \sum_{k=1}^{n_j(\mathbf{x})} \hat{d}_{jk}(\mathbf{x})/n_j(\mathbf{x})$, for $j \in \mathcal{J}$
$\mathbf{Rad}_t(\mathbf{x})$	confidence radius of price $\mathbf{x}$ by the end of period t
$U_t(\mathbf{x})$	the upper confidence bound of expected stationary revenue rate of price $\mathbf{x}$
$L_t(\mathbf{x})$	the lower confidence bound of expected stationary revenue rate of price $\mathbf{x}$
$\mathbf{x}_m$	the offered price in batch m by the algorithm
$Y_m(\mathbf{x})$	the practical revenue collected in batch m with price $\mathbf{x}$
$\mathcal{M}_{\mathbf{x}} \subseteq \{1, \ldots, M\}$	the batches choosing price $\mathbf{x}$
$T_{\mathbf{x}} = \sum_{m \in \mathcal{M}_{\mathbf{x}}} I_m \tau$	the number of periods choosing price $\mathbf{x}$
$\mathbf{Gamma}(\eta^{\alpha_j \mathbf{x}}, \beta^{\alpha_j \mathbf{x}})$	the prior/posterior distribution of $\alpha_j(\mathbf{x})$, the demand of product j under price $\mathbf{x}$
$\mathcal{A}_t$	the Jackson Network reaches the steady-state by time t from empty state
$\mathcal{A}_{t,l}$	all layer-l nodes reach the steady-state by time t from empty state
$\mathcal{A}_t^i$	the node i reaches the steady-state by time t from empty state
$\mathcal{B}_t$	the Jackson Network reaches the steady-state by time t from steady-state of old price
$\mathcal{B}_{t,l}$	all layer-l nodes reach the steady-state by time t from steady-state of old price
$\mathcal{B}_t^i$	the node i reaches the steady-state by time t from steady-state of old price
$\mathcal{C}_{2L\tau}$	the event that the system reaches the steady-state under the new price within the first $2L\tau$ periods
$\bar{\mathcal{C}}_{2L\tau}$	the complementary event of $\mathcal{C}_{2L\tau}$ that the system does not reach the steady-state within $2L\tau$ periods

B Illustrative example

We refer to this simple illustrative example several times in the rest of the paper. Consider 4 types of resources and 4 types of products:

- Product No.1: $(s, 1, 3, t)$, i.e., SQL Database $\rightarrow$ EC2
- Product No.2: $(s, 1, 3, 4, t)$, i.e., SQL Database $\rightarrow$ EC2 $\rightarrow$ SAP
- Product No.3: $(s, 2, 3, t)$, i.e., NoSQL Database $\rightarrow$ EC2
- Product No.4: $(s, 2, 3, 4, t)$, i.e., NoSQL Database $\rightarrow$ EC2 $\rightarrow$ SAP

Typically, customers will choose between an SQL based or a NoSQL based database service to store and preprocess the data. After running algorithms on elastic cloud compute EC2, only a subset of customers may choose to further integrate SAP applications. The network representation is given by Figure 2, and it is straightforward to check that this acyclic graph satisfies Assumption 2.

Figure 2: Servers and products in a network system.

We assign a price vector $\mathbf{x} = (x_1, x_2, x_3, x_4)$ to these four products. Note that we do not assign prices to resources. We only sell individual product (as a bundle of resources). Each posted price vector induces two quantities, namely, $\alpha(\mathbf{x})$ and $P(\mathbf{x})$. Here $\alpha(\mathbf{x})$ is the total arrival rate of all products given price vector $\mathbf{x}$ while $P(\mathbf{x})$ is the so-called (Markovian) "routing matrix" that captures the demand flows for products. One can represent a product $j \in \mathcal{J}$ using an s-t path, i.e., $(s, i_1, \ldots, i_n, t)$. Assume that Assumption 3 holds. Then we can write the demand rate for this particular product j as

$$\alpha_j = \alpha(\mathbf{x}) p_{s,i_1} p_{i_1,i_2} \ldots p_{i_{n-1},i_n} p_{i_n,t},$$

where $p_{i,i'}$ are the (i, i')th entry of the routing probability matrix $P(\mathbf{x})$.

Note that in our setting, the mapping $\mathbf{x} \mapsto \alpha(\mathbf{x}), P(\mathbf{x})$ is unknown to the decision maker. Thus, we need to offer different price vector $\mathbf{x}$ to learn this mapping. And we also assume this mapping is nonparametric. For the sake of illustration, we give the following linear mapping as an example.

$$\alpha(\mathbf{x}) = 110 - \mathbf{1} \cdot \mathbf{x},$$
$$p_{ii'} = \frac{\sum_{j:a_{ji}=1, a_{ji'}=0} x_j}{\sum_{j:a_{ji}=1} x_j}.$$

Suppose the price vector $\mathbf{x} = (1, 2, 2, 5)$. In this case, the total arrival rate $\alpha(\mathbf{x}) = 100$ and the routing matrix $P(\mathbf{x})$ is given by

$$P(\mathbf{x}) = \begin{cases} p_{s1} = \dfrac{x_3 + x_4}{x_1 + x_2 + x_3 + x_4} = 0.7 \\[2mm] p_{s2} = \dfrac{x_1 + x_2}{x_1 + x_2 + x_3 + x_4} = 0.3 \\[2mm] p_{3t} = \dfrac{x_2 + x_4}{x_1 + x_2 + x_3 + x_4} = 0.7 \\[2mm] p_{34} = \dfrac{x_1 + x_3}{x_1 + x_2 + x_3 + x_4} = 0.3 \\[2mm] p_{13} = p_{23} = p_{4t} = 1 \end{cases},$$

giving rise to

$$\text{Product Demand:} \begin{cases} \alpha_1 = 100 \times 0.7 \times 0.7 = 49 \\ \alpha_2 = 100 \times 0.7 \times 0.3 = 21 \\ \alpha_3 = 100 \times 0.3 \times 0.7 = 21 \\ \alpha_4 = 100 \times 0.3 \times 0.3 = 9 \end{cases}, \quad \text{Server Demand:} \begin{cases} \lambda_1 = 100 \times 0.7 = 70 \\ \lambda_2 = 100 \times 0.3 = 30 \\ \lambda_3 = 100 \times 1 = 100 \\ \lambda_4 = 100 \times 0.3 = 30 \end{cases}.$$

Note that in this example, Assumption 3 also means that out of all the customers who have chosen EC2, the proportion of choosing SAP is 0.3 and the proportion of not choosing SAP is 0.7 (regardless of database options).

We also consider two payment models, the lump sum payment model (presented in the main manuscript) and the service duration dependent payment model (presented in Appendix F).

- **(Lump sum payment model.)** A customer of product $j \in \mathcal{J}$ pays a lump sum x_j to the firm. We provide justification for this model as follows: If a customers chooses product j, then the customer will pay x_j to the firm regardless of how long the services over resources are. In our illustrative example, each product can be quoted a fixed price (i.e., not service duration dependent), which could be seen a preset "cloud solution package". The customers may prefer or benefit from this fixed price (contract) from their own project/accounting/cash-flow management. Under this model, even though the customers are not sensitive to service durations, the cloud platform is nevertheless sensitive to service durations since they affect how many products/services can be completed within a finite horizon.

- **(Service duration dependent payment model.)** A customer of product $j \in \mathcal{J}$ pays the amount that is equal to the per-unit-time x_j multiplied by the total service duration. This payment scheme is duration dependent (e.g., how much time is spent using EC2 in our illustrative example). The model and analytical details are presented in Appendix F.

C Technical results in Section 3

Proof of Proposition 1.

We show that the decisions variables (π, a, Y) associated with any admissible policy must satisfy the constraints of the linear program (1), and the expected revenue of this admissible policy is exactly the corresponding objective value in (1). Consider an arbitrary policy $\hat{\pi}$. We obtain the values of decision variables by the following steps. Variable $\hat{a}_{\mathbf{y}t}$ takes the value of probability that the system has $\mathbf{y}$ customers at the beginning of period t under the policy $\hat{\pi}$ for $\mathbf{y} \in \{0, 1, \ldots, \infty\}^I$ and $t = 1, \ldots, T$. Therefore, constraints (1b) are immediately satisfied. Variable $\hat{\pi}^{\mathbf{x}}_{\mathbf{y}t} = 1$ if the policy $\hat{\pi}$ chooses price $\mathbf{x}$ at state $(\mathbf{y}, t)$, otherwise, $\hat{\pi}^{\mathbf{x}}_{\mathbf{y}t} = 0$, for $\mathbf{y} \in \{0, 1, \ldots, \infty\}^I$, $t = 1, \ldots, T$, $\mathbf{y} \in \mathcal{X}$. As a result, constraints (1c) hold because in the left-hand side we have one variable equal to 1 and others all zero. Variable $\hat{Y}_{\mathbf{y}t}$ takes the value of expected revenue collected before the end of period T from the customers who arrive during period t, when the system has $\mathbf{y}$ customers at the beginning of period t, for $\mathbf{y} \in \{0, 1, \ldots, \infty\}^I$. Consider arbitrary $\mathbf{y}$ and t, we denote the price chosen at state $(\mathbf{y}, t)$ by the policy $\hat{\pi}$ as $\hat{\mathbf{x}}$. Then, by plugging in $\hat{\pi}^{\hat{\mathbf{x}}}_{\mathbf{y}t} = 1$, in the right-hand side of constraints (1d), we have $\vec{\alpha}^T_{\mathbf{x}} \mathbf{x}$, where $\vec{\alpha}^T_{\mathbf{x}}$ denotes the average number of arrived customers for consuming products during period t and $\hat{\mathbf{x}}$ is the revenue vector of these products. Therefore, $\vec{\alpha}^T_{\mathbf{x}} \mathbf{x}$ denotes the expected revenue collected from customers who arrive during period t (where assuming all arrived customers being served successfully). Consequently, we can conclude that the right-hand side of (1d) is as large as the left-hand side, because the right-hand side may also compute the revenue collected after the end of period T. Hence, constraints (1d) are also satisfied. By the value assignment steps of $\hat{a}_{\mathbf{y}t}$ and $\hat{Y}_{\mathbf{y}t}$, for $\mathbf{y} \in \{0, 1, \ldots, \infty\}^I$, $t = 1, \ldots, T$, one has that the corresponding LP objective value at solution $(\hat{\pi}, \hat{a}, \hat{J})$ is $Y^{\hat{\pi}}$. Therefore, we have $Y^{LP} \geq Y^{\pi}$, for any admissible policy π, and thus also for the optimal state-dependent policy π^*.

By defining $\tilde{\mathbf{x}} = \operatorname{argmax}_{\mathbf{x} \in \mathcal{X}} \vec{\alpha}^T_{\mathbf{x}} \mathbf{x}$, we find an optimal solution $(\tilde{\pi}, \tilde{a}, \tilde{Y})$ to the LP in (1) as follows.

$$\tilde{\pi}^{\tilde{\mathbf{x}}}_t = 1, \quad \forall \mathbf{y} \in \{0, 1, \ldots, \infty\}^I, \ \forall t = 1, \ldots, T \tag{3a}$$

$$\tilde{\pi}^p_{\mathbf{y}t} = 0, \quad \forall \mathbf{y} \in \{0, 1, \ldots, \infty\}^I, \ \forall t = 1, \ldots, T, \ \forall \mathbf{x} \in \mathcal{X} \backslash \{\tilde{\mathbf{x}}\} \tag{3b}$$

$$\tilde{Y}_{\mathbf{y}_0 t} = \vec{\alpha}^T_{\tilde{\mathbf{x}}} \tilde{\mathbf{x}} \quad \forall t = 1, \ldots, T \tag{3c}$$

$$\tilde{Y}_{\mathbf{y}t} = 0 \quad \forall \mathbf{y} \in \{0, 1, \ldots, \infty\}^I \backslash \{\mathbf{y}_0\}, \ \forall t = 1, \ldots, T \tag{3d}$$

$$\tilde{a}_{\mathbf{y}_0 t} = 1 \quad \forall t = 1, \ldots, T \tag{3e}$$

$$\tilde{a}_{\mathbf{y}t} = 0 \quad \forall \mathbf{y} \in \{0, 1, \ldots, \infty\}^I \backslash \{\mathbf{y}_0\}, \ \forall t = 1, \ldots, T, \tag{3f}$$

where $\mathbf{y}_0 = [0]^J$. Then, we can compute $Y^{\mathrm{LP}} = T \cdot \vec{\alpha}^T_{\tilde{\mathbf{x}}} \tilde{\mathbf{x}}$. Note that $\vec{\alpha}^T_{\mathbf{x}} \mathbf{x}$ is the stationary revenue rate of price $\mathbf{x}$, denoted as $r(\vec{\alpha}_{\mathbf{x}}, \mathbf{x})$.

Q.E.D.

D Technical results in Section 4.

Algorithm 2 RNRM-TS.

1: Input: T, $\mathcal{X}$, service network, turn-off price, $\eta^{\alpha_j(\mathbf{x})}$, $\beta^{\alpha_j(\mathbf{x})}$, $\forall j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x})$, $\forall \mathbf{x} \in \mathcal{X}$.
2: Initialize: L, τ, I_m, M, $m = 0$.
3: Warm-up Phase:
4: **for** $\mathbf{x} \in \mathcal{X}$ **do**
5:     Offer $\mathbf{x}$, record inter-arrival time $\hat{d}_{jk}(\mathbf{x})$ for arriving customers of product $j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x})$.
6:     **if** $n_{mj}(\mathbf{x}) \geq 8 \log(J^{1/4}T)$ **then**
7:         Continue.
8:     **end if**
9: **end for**
10: Update $\eta^{\alpha_j(\mathbf{x})}$, $\beta^{\alpha_j(\mathbf{x})}$, $\forall j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x})$, $\forall \mathbf{x} \in \mathcal{X}$.
11: Learning Phase:
12: **for** $m = 1, \ldots, M$ **do**
13:     Sample $\hat{\alpha}_j(\mathbf{x}) \sim \mathbf{Gamma}(\eta^{\alpha_j(\mathbf{x})}, \beta^{\alpha_j(\mathbf{x})})$ for product $j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x})$, $\forall \mathbf{x} \in \mathcal{X}$
14:     Choose $\mathbf{x}_m = \mathrm{argmax}_{\mathbf{x}_m \in \mathcal{X}} r(\hat{\vec{\alpha}}_{\mathbf{x}}, \mathbf{x})$.
15:     Offer $\mathbf{x}_m$ in batch m, i.e., for $I_m L \tau$ periods.
16:     Record inter-arrival time $\hat{d}_{jk}(\mathbf{x}_m)$ for arriving customers of product $j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x}_m)$.
17:     Compute and update $\eta^{\alpha_j(\mathbf{x}_m)}$, $\beta^{\alpha_j(\mathbf{x}_m)}$, $\forall j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x}_m)$.
18: **end for**

Proof of Proposition 2.

The proof is based on the concentration bound for sub-exponential random variables. For product $j \in \mathcal{J}_{\mathrm{off}}$, their demand is off, i.e., $\alpha_j(\mathbf{x}) = 0$, $j \in \mathcal{J}_{\mathrm{off}}(\mathbf{x})$. In this proof, we only consider one price and thus we use α_j to represent $\alpha_j(\mathbf{x})$ for notation simplicity. We can rewrite the right-hand-side as:

$$
\begin{aligned}
&\mathbb{P}\left(\left| \vec{\alpha}_{\mathbf{x}}^T \mathbf{x} - \sum_{j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x})} \frac{x_j}{\bar{d}_j(\mathbf{x})} \right| \leq \sum_{j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x})} \frac{x_j \sqrt{32 \log(J^{\frac{1}{4}} T)}}{\bar{d}_j(\mathbf{x}) \sqrt{n_j(\mathbf{x})}} \right) \\
=&\mathbb{P}\left(\left| \sum_{j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x})} \left(\alpha_j x_j - \frac{x_j}{\bar{d}_j(\mathbf{x})} \right) \right| \leq \sum_{j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x})} \frac{x_j \sqrt{32 \log(J^{\frac{1}{4}} T)}}{\bar{d}_j(\mathbf{x}) \sqrt{n_j(\mathbf{x})}} \right) \\
\geq&\mathbb{P}\left(\sum_{j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x})} \left| \alpha_j x_j - \frac{x_j}{\bar{d}_j(\mathbf{x})} \right| \leq \sum_{j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x})} \frac{x_j \sqrt{32 \log(J^{\frac{1}{4}} T)}}{\bar{d}_j(\mathbf{x}) \sqrt{n_j(\mathbf{x})}} \right) \\
\geq& \prod_{j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x})} \mathbb{P}\left(\left| \alpha_j x_j - \frac{x_j}{\bar{d}_j(\mathbf{x})} \right| \leq \frac{x_j \sqrt{32 \log(J^{\frac{1}{4}} T)}}{\bar{d}_j(\mathbf{x}) \sqrt{n_j(\mathbf{x})}} \right).
\end{aligned}
\tag{4}
$$

We can derive that for one product $j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x})$:

$$
\mathbb{P}\left(\left| \alpha_j x_j - \frac{x_j}{\bar{d}_j(\mathbf{x})} \right| \leq \frac{x_j \sqrt{32 \log(J^{\frac{1}{4}} T)}}{\bar{d}_j(\mathbf{x}) \sqrt{n_j(\mathbf{x})}} \right) = \mathbb{P}\left(\left| \alpha_j - \frac{1}{\bar{d}_j(\mathbf{x})} \right| \leq \frac{\sqrt{32 \log(J^{\frac{1}{4}} T)}}{\bar{d}_j(\mathbf{x}) \sqrt{n_j(\mathbf{x})}} \right)
\tag{5a}
$$

$$
=\mathbb{P}\left(\left| \frac{\alpha_j - \frac{1}{\bar{d}_j(\mathbf{x})}}{\alpha_j \frac{1}{\bar{d}_j(\mathbf{x})}} \right| \leq \frac{\sqrt{32 \log(J^{\frac{1}{4}} T)}}{\alpha_j \frac{1}{\bar{d}_j(\mathbf{x})} \bar{d}_j(\mathbf{x}) \sqrt{n_j(\mathbf{x})}} \right) = \mathbb{P}\left(\left| \bar{d}_j(\mathbf{x}) - \frac{1}{\alpha_j} \right| \leq \frac{\sqrt{32 \log(J^{\frac{1}{4}} T)}}{\alpha_j \sqrt{n_j(\mathbf{x})}} \right).
\tag{5b}
$$

The exponential arrival time intervals are also sub-exponential random variables with parameters $(\frac{4}{\alpha_j^2}, \frac{2}{\alpha_j})$, denoted by $\mathrm{SE}(\frac{4}{\alpha_j^2}, \frac{2}{\alpha_j})$, see Lemma 1 in the appendix. Consider the concentration inequality

for i.i.d. exponential variables ξ_i, $i = 1, \ldots n$ with mean $1/\lambda$:

$$\mathbb{P}\left[\left|\frac{1}{\lambda} - \frac{1}{n}\sum_{i=1}^{n}\xi_i\right| \geq t\right] \leq \begin{cases} 2\exp\left(-\frac{nt^2}{2\frac{4}{\lambda^2}}\right) & \text{if } 0 \leq t \leq \frac{2}{\lambda} \\ 2\exp\left(-\frac{nt}{2\frac{2}{\lambda}}\right) & \text{if } t > \frac{2}{\lambda} \end{cases}. \tag{6}$$

By the assumption $n_j(\mathbf{x}) \geq 8\log(J^{\frac{1}{4}}T)$, we can compute

$$\left(\frac{\sqrt{32\log(J^{\frac{1}{4}}T)}}{\alpha_j\sqrt{n_m(\mathbf{x})}}\right)^2 = \frac{32\log(J^{\frac{1}{4}}T)}{\alpha_j^2 n_j(\mathbf{x})} \leq \frac{4}{\alpha_j^2},$$

and thus we can apply the first inequality in (6) to (5). Therefore, we can derive

$$\mathbb{P}\left(\left|\bar{d}_j(\mathbf{x}) - \frac{1}{\alpha_j}\right| \leq \frac{\sqrt{32\log(J^{\frac{1}{4}}T)}}{\alpha_j\sqrt{n_j(\mathbf{x})}}\right) \geq 1 - 2\exp\left(\frac{-n_j(\mathbf{x})\frac{32\log(J^{\frac{1}{4}}T)}{\alpha_j^2 n_j(\mathbf{x})}}{2\frac{4}{\alpha_j^2}}\right) \geq 1 - \frac{2}{J \cdot T^4} \tag{7}$$

Then, we can compute a lower bound of (4) as

$$\left(1 - \frac{2}{J \cdot T^4}\right)^{|\mathcal{J}_{\text{on}}(\mathbf{x})|} \geq \left(1 - \frac{2}{J \cdot T^4}\right)^J \geq 1 - \frac{2J}{J \cdot T^4} = 1 - \frac{2}{T^4}.$$

Q.E.D.

Lemma 1. *(Sub-exponential property, Lemma 3 in Jia et al. 2020) If a random variable X follows an exponential distribution with mean $1/\lambda$, then the random variable $X - \frac{1}{\lambda}$ is $(\frac{4}{\lambda^2}, \frac{2}{\lambda})$-sub-exponential.*

E Technical results in Section 5

Proof of Proposition 3.

In this proof, we analyze the coupling time of nodes layer by layer. We define events as follows (all starting from empty state).

$$\mathcal{A}_t = \{\text{the Jackson Network reaches the steady-state by time } t\},$$
$$\mathcal{A}_{t,l} = \{\text{all layer-l nodes reach the steady-state by time } t\},$$
$$\mathcal{A}_t^i = \{\text{the node } i \text{ reaches the steady-state by time } t\}.$$

First, we discuss the coupling analysis of an M/M/c queue, which is the basis for completing this proof. With the assumption $\log > 4$, the probability that an M/M/c queue reaches the steady-state from empty state after $(\log T)^2$ periods of time is bounded by $1 - 2/T^2$, see Lemma 3 (Proposition 4 in Jia et al. (2020)). This results are derived following the law of total probability, the first-order stochastic dominance between busy times of an M/M/c queue, and the concentration inequality for independent samples. We denote node i is a prerequisite node of i', $i < i'$ if there exists a product requires to use nodes i and i'. Thus, it is easy to show that the prerequisite nodes of a Layer-l node must have smaller layer indices.

In our Jackson Network, node i' reaches its steady-state indicates that the arrival process of node i' is a stable Poisson process, and therefore it requires that all the prerequisite nodes also reach the steady-state. In this proof, we analyze the nodes layer by layer. By treating the nodes in the same layer together, we can derive that

$$\begin{aligned}
\mathbb{P}(\mathcal{A}_{L\tau}) &= \mathbb{P}(\mathcal{A}_{L\tau,1}\mathcal{A}_{L\tau,2}\mathcal{A}_{L\tau,3}\ldots\mathcal{A}_{L\tau,L}) \\
&\geq \mathbb{P}(\mathcal{A}_{\tau,1}\mathcal{A}_{2\tau,2}\mathcal{A}_{3\tau,3}\ldots\mathcal{A}_{L\tau,L}) \\
&= \mathbb{P}(\mathcal{A}_{\tau,1})\mathbb{P}(\mathcal{A}_{2\tau,2}\mathcal{A}_{3\tau,3}\ldots\mathcal{A}_{L\tau,L}|\mathcal{A}_{\tau,1}) \\
&= \mathbb{P}(\mathcal{A}_{\tau,1})\mathbb{P}(\mathcal{A}_{2\tau,2}|\mathcal{A}_{\tau,1})\mathbb{P}(\mathcal{A}_{3\tau,3}\ldots\mathcal{A}_{L\tau,L}|\mathcal{A}_{\tau,1}\mathcal{A}_{2\tau,2}) \\
&= \mathbb{P}(\mathcal{A}_{\tau,1})\prod_{l=2}^{L}\mathbb{P}(\mathcal{A}_{l\tau,l}|\mathcal{A}_{\tau,1}\ldots\mathcal{A}_{(l-1)\tau,l-1}).
\end{aligned} \tag{8}$$

In short, we derive a lower bound of this mixing probability by requiring the Layer-l nodes reach the steady-state within the first $L \times \tau$ periods, conditional on that Layer-l' nodes also reach the steady-state within the first $l' \times \tau$ periods for $l' < l$, or equivalently, all prerequisite nodes of Layer-l nodes reach the steady-state within $(l-1) \times \tau$ periods.

Layer-1 Nodes. In the first τ periods, by applying the probability bound for an M/M/c queue, we can easily conclude that all Layer-1 nodes (the correspondent M/M/c queues) reach the steady-state with probability $1 - \frac{2}{T^3}$. Recall $\mathcal{I}_1$ is the collection of all Layer-1 nodes and $|\mathcal{I}_1|$ is the cardinality. Therefore we can compute that the probability that all Layer-1 nodes reach the steady-state by first τ periods is

$$\mathbb{P}(\mathcal{A}_{\tau,1}) = \left(1 - \frac{2}{T^3}\right)^{|\mathcal{I}_1|} \geq 1 - \frac{2|\mathcal{I}_1|}{T^2}, \tag{9}$$

because the queueing evolution of all Layer-1 nodes are independent with each other and thus we can simplify multiply the probability together, in the first equality in the above expression.

Layer-2 Nodes. We then analyze the Layer-2 nodes. Consider one arbitrary layer-2 node, denoted by i without loss of generality. We then focus on the corresponding queue of this node i conditional on that all Layer-1 nodes have reached steady-state. We firstly show a virtual queue can reach steady-state within the next τ periods with probability at least $1 - \frac{2}{T^2}$. Then we show that an alternative queue of the actual queue takes a shorter time to reach steady-state and thus the actual queue can also reach steady-state within the next τ periods with probability at least $1 - \frac{2}{T^2}$.

- *Virtual queue.* We consider a virtual queue which rejects all customers arrived before τ periods. Then by Lemma 3, this virtual queue reaches the steady-state with probability $1 - \frac{2}{T^2}$ after the next τ periods, i.e., after the first 2τ periods.

- *Alternative queue.* Denote the probability that the number of customers in Node i at the end of period τ is less than N by $\mathbb{P}_i(N)$, including the customers both using the server or in the waiting queue of node i. The actual queue accepts customers during the first τ periods with a changing rate from an initial rate to the stable arrival rate λ_i (conditional on all Layer-1 nodes reach steady-state and thus provides stable Poisson arrivals for nodes with high layer indices). Denote the demand of all products to the Jackson Network by α. The stable arrival rate $\lambda_i = \alpha p_{si} + \alpha \sum_{i':(i',i\in E)} p_{si'}p_{i'i}$, where αp_{si} is the initial rate, i.e., the demand rate of products that request node i as the first service, and $\alpha p_{si'}p_{i'i}$ is the stable demand flowing from prerequisite node i' to node i. Consider an alternative queue which accepts customers at the stable rate λ_i. If the queue starts from time 0, then the probability that this virtual queue has customers with a size less than N at the end of period τ is smaller than $\mathbb{P}_i(N)$. This is because this alternative queue accepts more customers while the service times are all following a same distribution. Then we postpone the starting point of this alternative queue to a time stamp between 0 and τ such that the probability that this alternative queue has customers with a size less than N at the end of period τ is the same as $\mathbb{P}_i(N)$. Note that this time stamps is exist, which can be easily proved by the memoryless property of the exponential service time assumptions. Then we have the following two results. First, this alternative queue has a same evolution path as the actual queue. Second, because this alternative queue starts accepting customers earlier than time τ, and thus it will reach steady-state earlier than the virtual queue.

We provide an illustration of the actual, alternative, and virtual queues in Figure 3.

To conclude, we have the actual queue can reach the steady-state with a higher probability after the first 2τ periods. Similarly, the queueing evolution of all Layer-2 nodes are independent with each other and thus we can simplify multiply the probability together. Therefore, we have that for all Layer-2 nodes

$$\mathbb{P}(\mathcal{A}_{2\tau,2}|\mathcal{A}_{\tau,1}) = \left(1 - \frac{2}{T^2}\right)^{|\mathcal{I}_2|} \geq 1 - \frac{2|\mathcal{I}_2|}{T^2}. \tag{10}$$

Layers with Higher Indices. Then we apply the same analysis steps of the Layer-2 nodes to nodes in layers with higher indices. Conditional on the events that all nodes in layers with lower indices have reached the steady-state by the end of $(l-1)\tau$ periods, we can compute that the probability the

Figure 3: The left part shows the arrival rates of the actual, alternative, and virtual queues (for a Layer-2 node i when the Jackson network starting from empty state). The right part shows the probability that the queue has different number of customers at time τ. This probability are the same for the alternative and the actual queues and thus they are equivalent systems after time τ.

all nodes of layer l can reach steady-state in the next τ period with a probability:

$$\mathbb{P}(\mathcal{A}_{l\tau,l}|\mathcal{A}_{\tau,1}\ldots\mathcal{A}_{(l-1)\tau,l-1}) \geq 1 - \frac{2I_l}{T^2}. \tag{11}$$

With the results in (9), (10), and (11), we are ready to bound the probability in (8):

$$\mathbb{P}(\mathcal{A}_{L\tau}) \geq \prod_{l=1}^{L}\left(1 - \frac{2|\mathcal{I}_l|}{T^2}\right) \geq 1 - \frac{2I}{T^2}.$$

We refer to Figure 4 for a direct representation of the above proof sketch, which shows the mixing time analysis of an illustrative example.

Figure 4: The mixing time analysis for an illustrative example starting from zero state.

Q.E.D.

Proof of Proposition 4.

Similar to the proof of Proposition 4, we analyze the mixing time of nodes layer by layer. Before conducting the proof, we introduce an existing work for an M/M/c queue mixing time. We similarly define the following events when the Jackson Network starts to operate under a new price starting from the steady-state of the old price:

$$\mathcal{B}_t = \{\text{the Jackson Network reaches the steady-state by time } t\},$$

$$\mathcal{B}_{t,l} = \{\text{all layer-l nodes reach the steady-state by time } t\},$$

$$\mathcal{B}_t^i = \{\text{the node } i \text{ reaches the steady-state by time } t\}.$$

Lemma 4 says that for one M/M/c queue where Assumption 5 holds, then the probability that the queue reaches a new steady-state from a old steady-state within 2τ time periods after the price change is bounded by $1 - 4/T^2$. To derive this result, the authors decompose possible price changes into the following two cases. The first is when the arrival rate changes from a higher value to a lower value and the second is vice versa. For each case, the authors further consider two sub-cases: the unit service rate increase or decrease.

Consider the Jackson Network system in our problem. We can bound the probability by:

$$\mathbb{P}(\mathcal{B}_{2L\tau}) \geq \mathbb{P}(\mathcal{B}_{2\tau,1}) \prod_{l=2}^{L} \mathbb{P}(\mathcal{B}_{2l\tau,l} | \mathcal{B}_{2\tau,1} \ldots \mathcal{B}_{2(l-1)\tau,l-1}). \tag{12}$$

The above inequality is derived following the same steps in (8) and we neglect the details here.

Assumption 5 allows us to apply Lemma 4 when we analyze one specific node, i.e., one specific M/M/c queue. The structure of the following analysis is similar to the structure of the proof of Proposition 3, we first analyze Layer-1 node and then the nodes with higher layer indices. For nodes with higher layer indices, we apply the aforementioned proposition to a virtual queue, and then show this virtual queue takes longer than an alternative system with the actual queue.

Layer-1 Nodes. In the first 2τ periods, by applying the probability bound for an M/M/c queue with price change (Lemma 4), we can easily conclude that all Layer-1 nodes (the correspondent M/M/c queues) reach the steady-state with probability $1 - \frac{4}{T^2}$. Consider the probabilistic independency of the coupling process among Layer-1 nodes, the probability that all Layer-1 nodes reach the steady-state by first 2τ periods is

$$\mathbb{P}(\mathcal{B}_{2\tau,1}) = \left(1 - \frac{4}{T^2}\right)^{|\mathcal{I}_1|} \geq 1 - \frac{4|\mathcal{I}_1|}{T^2}. \tag{13}$$

Layer-2 Nodes and Nodes with Higher Layer Indices. We then analyze nodes with a layer index 2 or higher. Consider one arbitrary Layer-l node, denoted by i without loss of generality. We then focus on the corresponding queue of this node i conditional on that all prerequisites nodes have reached steady-state. We firstly show a virtual queue can reach steady-state within the next 2τ periods with probability at least $1 - \frac{4}{T^2}$. Then we show that an alternative queue of the actual queue takes a shorter time to reach steady-state and thus the actual queue can also reach steady-state within the next 2τ periods with probability at least $1 - \frac{4}{T^2}$.

- *Virtual queue.* We consider a virtual queue which accepts customers under the old queueing parameters in the first $2(l-1)\tau$ periods and then accepts customers under the new queueing parameter in the next 2τ periods. With Assumption 5, we can directly apply the results for one M/M/c queue to reach steady-state starting from a steady-state and conclude that this virtual queue reaches the steady-state with probability $1 - \frac{4}{T^2}$ after the next 2τ periods, i.e., after the first $2l\tau$ periods.

- *Alternative queue.* Denote the probability that in Node i at the end of $2(l-1)\tau$ periods, the number of old customers is less than N_o and the number of new customers is less than N_n by $\mathbb{P}_i(N_o, N_n)$, including the customers both using the server or in the waiting queue of node i. Consider an alternative queue which starts accepting new customers under the new queueing system parameter from time 0 and denote the corresponding probability for having less than N_o number of old customers and less than N_n new customers at the end of $2(l-1)\tau$ periods by $\mathbb{P}'_i(N_o, N_n)$. By the analysis in Proposition 3, we know there is a mismatch between $\mathbb{P}_i(N_o, N_n)$. Then we postpone the starting point of this alternative queue to a time stamp between 0 and $2(L-1)\tau$ such that $\mathbb{P}'_i(N_o, N_n) = \mathbb{P}_i(N_o, N_n)$. Note that this

time stamps is always exist, which can be easily proved by the memoryless property of the exponential service time assumptions because we can always find this stamp backwards from any customer state at time $2(l-1)\tau$. Then we have the following two results. First, this alternative queue has a same evolution path as the actual queue. Second, because this alternative queue starts accepting new customers earlier than time $2(l-1)\tau$, and thus it will reach steady-state earlier than the virtual queue.

To conclude, we have the actual queue can reach the steady-state with a higher probability after the first 2τ periods. Therefore , conditional on the events that all nodes in layers with lower indices have reached the steady-state by the end of $2(l-1)\tau$ periods, we can compute that the probability the all nodes of layer l can reach steady-state in the next 2τ period with a probability:

$$\mathbb{P}(\mathcal{B}_{2l\tau,l}|\mathcal{B}_{2\tau,1}\ldots\mathcal{B}_{2(l-1)\tau,l-1}) \geq 1 - \frac{4I_l}{T^2}. \tag{14}$$

With the results in (13) and (14), we are ready to bound the probability in (12):

$$\mathbb{P}(\mathcal{B}_{2L\tau}) \geq \prod_{l=1}^{L}\left(1 - \frac{4|\mathcal{I}_l|}{T^2}\right) \geq 1 - \frac{4I}{T^2}.$$

Q.E.D.

Proof of Theorem 1.

We bound the relaxed regret during the Warm-up and Learning Phases separately. We further decompose the relaxed regret during the Learning Phase into two parts. The first part is the regret of suboptimality, which is induced by not offering the state-independent optimal price in certain time periods. We bound the loss due to suboptimality with the help of Proposition 2. The second part is the regret of nonstationarity, which represents the revenue loss induced by changing offered prices. We bound the loss due to nonstationarity with the help of Propositions 3 and 4. The structure of this proof is similar to that of Theorem 1 in Jia et al. (2020), where the authors consider a single reusable product setting.

Regret of the Warm-up Phase. The difference between $Y_{\text{Warm-up}}^{\text{LP}}$ and $Y_{\text{Warm-up}}^{\text{RNRM-UCB}}$ is at most linear on the length of the Warm-up Phase, which is $O(X\log(J^{1/4}T))$. Therefore we can directly have

$$\overline{\text{Regret}}_{\text{Warm-up}}(\text{RNRM-UCB}, T) = O(X\log(J^{1/4}T)). \tag{15}$$

Regret of nonstationarity during the Learning Phase. We denote the offered price of batch m in RNRM-UCB as $\mathbf{x}_m$. Therefore, the expected revenue collected in batch m is

$$Y_m^{\text{RNRM-UCB}}(\mathbf{x}_m) = Y_{m,0\sim2L\tau}^{\text{RNRM-UCB}}(\mathbf{x}_m) + Y_{m,2L\tau\sim I_mL\tau}^{\text{RNRM-UCB}}(\mathbf{x}_m),$$

where $Y_{m,0\sim2L\tau}^{\text{RNRM-UCB}}(\mathbf{x}_m)$ denotes the expected revenue collected in the first $2L\tau$ periods and $Y_{m,2L\tau\sim I_mL\tau}^{\text{RNRM-UCB}}(\mathbf{x}_m)$ denotes the expected revenue collected from $(2L\tau+1)$th period to the end of batch m. We define event $\mathcal{C}_{2L\tau}$ as that the network service system reaches the steady-state under the new price within the first $2L\tau$ periods and define $\bar{\mathcal{C}}_{2L\tau}$ as the complementary event of $\mathcal{C}_{2L\tau}$. By Propositions 3 and 4, we have $\mathbb{P}[\mathcal{C}_{2L\tau}] \geq 1 - 4I/T^2$. Consequently, we define $Y_{m,\mathcal{C}_{2L\tau}}^{\text{RNRM-UCB}}(\mathbf{x}_m)$ to be the expected value of $Y_{m,2L\tau\sim I_mL\tau}^{\text{RNRM-UCB}}(\mathbf{x}_m)$ conditional on $\mathcal{C}_{2L\tau}$ and $Y_{m,\bar{\mathcal{C}}_{2L\tau}}^{\text{RNRM-UCB}}(\mathbf{x}_m)$ similarly. As a result, we can express the expected revenue during batch m as:

$$
\begin{aligned}
Y_m^{\text{RNRM-UCB}}(\mathbf{x}_m) &= Y_{m,0\sim2L\tau}^{\text{RNRM-UCB}}(\mathbf{x}_m) + Y_{m,\mathcal{C}_{2L\tau}}^{\text{RNRM-UCB}}(\mathbf{x}_m)\mathbb{P}(\mathcal{C}_{2L\tau}) + Y_{m,\bar{\mathcal{C}}_{2L\tau}}^{\text{RNRM-UCB}}(\mathbf{x}_m)\left(1 - \mathbb{P}(\mathcal{C}_{2L\tau})\right) \\
&\geq O(L\tau) + \left(\vec{\alpha}_{\mathbf{x}_m}^T\mathbf{x}_m\right)(I_mL\tau - 2L\tau)\left(1 - \frac{4I}{T^2}\right) + O\left(\frac{(I_mL\tau - 2L\tau)I}{T^2}\right) \\
&= \left(\vec{\alpha}_{\mathbf{x}_m}^T\mathbf{x}_m\right)I_mL\tau - O(L\tau) \\
&= r(\vec{\alpha}_{\mathbf{x}_m}, \mathbf{x}_m)I_mL\tau - O(L\tau).
\end{aligned}
\tag{16}
$$

Regret of suboptimality during the Learning Phase. Recall the upper and lower confidence bounds of the revenue rate defined in Definition 2. It is clear that

$$U_m(\mathbf{x}) - L_m(\mathbf{x}) = 2\mathbf{Rad}_m(\mathbf{x}) = 2 \sum_{j \in \mathcal{J}_{\mathrm{on}}(\mathbf{x})} \frac{x_j \sqrt{32 \log(J^{\frac{1}{4}} T)}}{\bar{d}_j(\mathbf{x}) \sqrt{n_j(\mathbf{x})}}.$$

For batch m, we can compute the difference in the revenue rates between the optimal price and selected price $\mathbf{x}_m$ as

$$\Delta(\mathbf{x}_m) = r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) - r(\vec{\alpha}_{\mathbf{x}_m}, \mathbf{x}_m) \tag{17a}$$

$$= \big(r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) - U_m(\mathbf{x}_m)\big) + \big(U_m(\mathbf{x}_m) - L_m(\mathbf{x}_m)\big) + \big(L_m(\mathbf{x}_m) - r(\vec{\alpha}_{\mathbf{x}_m}, \mathbf{x}_m)\big) \tag{17b}$$

$$\le \big(r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) - U_m(\tilde{\mathbf{x}})\big) + \big(U_m(\mathbf{x}_m) - L_m(\mathbf{x}_m)\big) + \big(L_m(\mathbf{x}_m) - r(\vec{\alpha}_{\mathbf{x}_m}, \mathbf{x}_m)\big), \tag{17c}$$

where the last inequality holds because $U_m(\mathbf{x}_m) \ge U_m(\tilde{\mathbf{x}})$ by the algorithm construction.

Then the cumulative regret during the Learning Phase is

$$\overline{\mathrm{Regret}}_{\mathrm{Learning}}(\text{RNRM-UCB}, T) = Y_{\mathrm{Learning}}^{\mathrm{LP}} - Y_{\mathrm{Learning}}^{\text{RNRM-UCB}} \tag{18a}$$

$$= \sum_{m=1}^{M} \big(r(\alpha_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) I_m L \tau - Y_m^{\text{RNRM-UCB}}(\mathbf{x}_m)\big) \tag{18b}$$

$$= \sum_{m=1}^{M} \mathbb{E}\Big[\Delta(\mathbf{x}_m) I_m L \tau + O(L\tau)\Big]. \tag{18c}$$

By (17), we have

$$\sum_{m=1}^{M} \mathbb{E}\Big[\Delta(\mathbf{x}_m) I_m L \tau\Big] = \sum_{m=1}^{M} \mathbb{E}\Big[\big(r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) - U_m(\tilde{\mathbf{x}})\big)\Big] I_m L \tau + \sum_{m=1}^{M} \mathbb{E}\Big[\big(U_m(\mathbf{x}_m) - L_m(\mathbf{x}_m)\big)\Big] I_m L \tau$$

$$+ \sum_{m=1}^{M} \mathbb{E}\Big[\big(L_m(\mathbf{x}_m) - r(\vec{\alpha}_{\mathbf{x}_m}, \mathbf{x}_m)\big)\Big] I_m L \tau. \tag{19}$$

We first analyze the middle term of (19). Let $\mathcal{M}_{\mathbf{x}}$ denote the batches choosing price $\mathbf{x}$ and $T_{\mathbf{x}}$ the number of periods choosing price $\mathbf{x}$, i.e., $T_{\mathbf{x}} = \sum_{m \in \mathcal{M}_{\mathbf{x}}} I_m L \tau$. Then,

$$\sum_{m=1}^{M} \mathbb{E}\Big[\big(U_m(\mathbf{x}_m) - L_m(\mathbf{x}_m)\big) I_m L \tau\Big] = \sum_{\mathbf{x} \in \mathcal{X}} \sum_{m \in \mathcal{M}_{\mathbf{x}}} \mathbb{E}\big[U_m(\mathbf{x}) - L_m(\mathbf{x})\big] I_m L \tau \tag{20a}$$

$$= \sum_{\mathbf{x} \in \mathcal{X}} \sum_{m \in \mathcal{M}_{\mathbf{x}}} 2\mathbb{E}\big[\mathbf{Rad}_m(\mathbf{x})\big] I_m L \tau \tag{20b}$$

$$\le \sum_{\mathbf{x} \in \mathcal{X}} O\left(J \sqrt{|\mathcal{M}_{\mathbf{x}}| T_{\mathbf{x}} \log(J^{1/4} T)}\right) \tag{20c}$$

$$= O\left(J \sqrt{\log(J^{1/4} T)}\right) \sum_{\mathbf{x} \in \mathcal{X}} \sqrt{|\mathcal{M}_{\mathbf{x}}| T_{\mathbf{x}}} \tag{20d}$$

$$\le O\left(J \sqrt{\log(J^{1/4} T) \log T}\right) \sum_{\mathbf{x} \in \mathcal{X}} \sqrt{T_{\mathbf{x}}} \tag{20e}$$

$$\le O\left(J \sqrt{X T \log T \log(J^{1/4} T)}\right), \tag{20f}$$

where (20c) by Lemma 2, (20e) is derived by the number of the batches, and (20f) follows Jensen's inequality. The number of batches is $\log_2\left(1 + \frac{T - 8X \log(J^{1/4} T)}{2L \log(T)^2}\right) = O\big(\log(T)\big)$.

Below we analyze the first term of (19), and the same logic applies to the third term, of which we omit the details here.

$$\sum_{m=1}^{M} \mathbb{E}\Big[\big(r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) - U_m(\tilde{\mathbf{x}})\big)\Big] I_m L\tau \tag{21a}$$

$$\leq \sum_{m=1}^{M} \mathbb{E}\Big[\big(r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) - U_m(\tilde{\mathbf{x}})\big)\mathbb{1}\big(r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) > U_m(\tilde{\mathbf{x}})\big)\Big] I_m L\tau \tag{21b}$$

$$\leq \sum_{m=1}^{M} \mathbb{E}\Big[r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}})\mathbb{1}\big(r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) > U_m(\tilde{\mathbf{x}})\big) - U_m(\tilde{\mathbf{x}})\mathbb{1}\big(r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) > U_m(\tilde{\mathbf{x}})\big)\Big] I_m L\tau \tag{21c}$$

$$= \sum_{m=1}^{M} r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}})\mathbb{P}\big(r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) > U_m(\tilde{\mathbf{x}})\big) I_m L\tau - \sum_{m=1}^{M} \mathbb{E}\Big[U_m(\tilde{\mathbf{x}})\mathbb{1}\big(r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) > U_m(\tilde{\mathbf{x}})\big)\Big] I_m L\tau \tag{21d}$$

$$\leq \sum_{m=1}^{M} r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}})\mathbb{P}\big(r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) > U_m(\tilde{\mathbf{x}})\big) \cdot I_m L\tau \tag{21e}$$

$$\leq O\left(\frac{1}{T^3}\right), \tag{21f}$$

where (21e) is because $U_m(\tilde{\mathbf{x}}) > 0$ by Definition 2, and (21f) follows Proposition 2.

Given that $O\big(n_{Mj}(\mathbf{x})\big) = O(T_{\mathbf{x}})$, for any product $j \in \mathcal{J}_{\text{on}}(\mathbf{x})$, we have $\sum_{m=1}^{M} O(L\tau) = O\big(L(\log(T))^3\big)$. Combining the three terms together,

$$\overline{\text{Regret}}_{\text{Learning}}(\text{RNRM-UCB}, T) = \sum_{m=1}^{M} \mathbb{E}\Big[\Delta(\mathbf{x}_m)I_m L\tau\Big] + \sum_{m=1}^{M} O(L\tau)$$

$$= O\left(J\sqrt{XT\log T\log(J^{1/4}T)}\right) + O\left(\frac{1}{T^3}\right) + O\big(L(\log(T))^3\big) \tag{22}$$

$$= O\left(J\sqrt{XT\log T\log(J^{1/4}T)}\right).$$

Total Regret of RNRM-UCB algorithm. By adding up the relaxed regret during the Warm-up and Learning Phases, we have the regret of RNRM-UCB algorithm as:

$$\overline{\text{Regret}}(\text{RNRM-UCB}, T) = \overline{\text{Regret}}_{\text{Warm-up}}(\text{RNRM-UCB}, T) + \overline{\text{Regret}}_{\text{Learning}}(\text{RNRM-UCB}, T) \tag{23a}$$

$$= O(X\log(J^{1/4}T)) + O\left(J\sqrt{XT\log T\log(J^{1/4}T)}\right) \tag{23b}$$

$$= \tilde{O}(J\sqrt{XT}). \tag{23c}$$

Q.E.D.

Proof of Theorem 2.

The analysis here is similar to that in the proof of Theorem 1. We first bound the Bayesian regret of the Warm-up Phase and then for the Learning Phase.

Bayesian regret in the Warm-up Phase. Similarly, we can bound the Bayesian regret in the Warm-up Phase as

$$\overline{\text{BR}}_{\text{Warm-up}}(\text{RNRM-TS}, T) = O(X\log(J^{1/4}T)). \tag{24}$$

Bayesian regret in the Learning Phase. In batch m, the chosen price $\mathbf{x}_m$ and the state-independent optimal price $\tilde{\mathbf{x}}$ are identically distributed conditional on $\mathcal{F}_m$, which indicates that $\mathbb{E}\big[U_m(\mathbf{x}_m) - U_m(\tilde{\mathbf{x}})|\mathcal{F}_m\big] = 0$ (e.g., see the proof of Lemma 3.7 in Slivkins (2019)). Thus, the Bayesian regret in

batch m is:

$$\overline{\text{BR}}_m \tag{25a}$$

$$= \underset{\{\alpha_j(\mathbf{x})\}_{j\in\mathcal{J}} \sim \text{priori}}{\mathbb{E}} \left[\mathbb{E}\left[r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) I_m L\tau - Y_m^{\text{RNRM-TS}}(\mathbf{x}_m) | \mathcal{F}_m \right] \right] \tag{25b}$$

$$= \underset{\{\alpha_j(\mathbf{x})\}_{j\in\mathcal{J}} \sim \text{priori}}{\mathbb{E}} \left[\mathbb{E}\left[I_m L\tau U_m(\mathbf{x}_m) - Y_m^{\text{RNRM-TS}}(\mathbf{x}_m) + r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) I_m L\tau - I_m L\tau U_m(\tilde{\mathbf{x}}) | \mathcal{F}_m \right] \right] \tag{25c}$$

$$= \underset{\{\alpha_j(\mathbf{x})\}_{j\in\mathcal{J}} \sim \text{priori}}{\mathbb{E}} \left[\mathbb{E}\left[I_m L\tau U_m(\mathbf{x}_m) - Y_m^{\text{RNRM-TS}}(\mathbf{x}_m) | \mathcal{F}_m \right] \right] \tag{25d}$$

$$+ \underset{\{\alpha_j(\mathbf{x})\}_{j\in\mathcal{J}} \sim \text{priori}}{\mathbb{E}} \left[\mathbb{E}\left[r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) I_m L\tau - I_m L\tau U_m(\tilde{\mathbf{x}}) | \mathcal{F}_m \right] \right] \tag{25e}$$

$$= \mathbb{E}\left[I_m L\tau U_m(\mathbf{x}_m) - Y_m^{\text{RNRM-TS}}(\mathbf{x}_m) \right] + \mathbb{E}\left[r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) I_m L\tau - I_m L\tau U_m(\tilde{\mathbf{x}}) \right] \tag{25f}$$

$$\leq \mathbb{E}\left[I_m L\tau U_m(\mathbf{x}_m) - I_m L\tau \cdot r(\vec{\alpha}_{\mathbf{x}_m}, \mathbf{x}_m) + O(L\tau) \right] + \mathbb{E}\left[I_m L\tau \cdot r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) - I_m L\tau U_m(\tilde{\mathbf{x}}) \right] \tag{25g}$$

$$= I_m L\tau \left(\mathbb{E}\left[U_m(\mathbf{x}_m) - r(\vec{\alpha}_{\mathbf{x}_m}, \mathbf{x}_m) \right] + \mathbb{E}\left[r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) - U_m(\tilde{\mathbf{x}}) \right] \right) + O(L\tau), \tag{25h}$$

where (25g) is derived following the analysis of nonstationarity in (16).

Derived from Proposition 2, for some $\gamma > 0$, two properties hold (see, e.g., Slivkins 2019):

$$\mathbb{E}[(r(\vec{\alpha}_{\mathbf{x}}, \mathbf{x}) - U_m(\mathbf{x}))^+] \leq \frac{\gamma}{XT} \qquad \text{and} \qquad \mathbb{E}[(L_m(\mathbf{x}) - r(\vec{\alpha}_{\mathbf{x}}, \mathbf{x}))^+] \leq \frac{\gamma}{XT}. \tag{26}$$

Following these two properties of the stationary revenue rate in (26), we have

$$\mathbb{E}\left[r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) - U_m(\tilde{\mathbf{x}}) \right] \leq \mathbb{E}\left[(r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) - U_m(\tilde{\mathbf{x}}))^+ \right] \leq \sum_{\mathbf{x}\in\mathcal{X}} \mathbb{E}\left[(r(\vec{\alpha}_{\mathbf{x}}, \mathbf{x}) - U_m(\mathbf{x}))^+ \right] \leq \frac{\gamma}{T}, \tag{27}$$

and similarly,

$$\begin{aligned}
\mathbb{E}\left[U_m(\mathbf{x}_m) - r(\lambda_{\mathbf{x}_m}, \mu_{\mathbf{x}_m}, \mathbf{x}_m) \right] &= \mathbb{E}\left[2 \cdot \mathbf{Rad}_m(\mathbf{x}_m) + L_m(\mathbf{x}_m) - r(\vec{\alpha}_{\mathbf{x}_m}, \mathbf{x}_m) \right] \\
&\leq \mathbb{E}\left[2 \cdot \mathbf{Rad}_m(\mathbf{x}_m) \right] + \mathbb{E}\left[(L_m(\mathbf{x}) - r(\vec{\alpha}_{\mathbf{x}_m}, \mathbf{x}_m))^+ \right] \\
&\leq \mathbb{E}\left[2 \cdot \mathbf{Rad}_m(\mathbf{x}_m) \right] + \sum_{\mathbf{x}\in\mathcal{X}} \mathbb{E}\left[(L_m(\mathbf{x}) - r(\vec{\alpha}_{\mathbf{x}_m}, \mathbf{x}_m))^+ \right] \\
&\leq \mathbb{E}\left[2 \cdot \mathbf{Rad}_m(\mathbf{x}_m) \right] + \frac{\gamma}{T}.
\end{aligned} \tag{28}$$

Summing over all batches $m, \forall m = 1, \ldots, M$, we have the cumulative relaxed Bayesian regret by the end of the Learning Phase as:

$$\overline{\mathrm{BR}}_{\mathrm{Learning}}(\pi_{\mathrm{RNRM\text{-}TS}}, T) = \sum_{m=1}^{M} \overline{\mathrm{BR}}_m \tag{29a}$$

$$= \sum_{m=1}^{M} \left(I_m L \tau \Big(\mathbb{E}\big[U_m(\mathbf{x}_m) - r(\vec{\alpha}_{\mathbf{x}_m}, \mathbf{x}_m)\big] + \mathbb{E}\big[r(\vec{\alpha}_{\tilde{\mathbf{x}}}, \tilde{\mathbf{x}}) - U_m(\tilde{\mathbf{x}})\big] \Big) + O(L\tau) \right) \tag{29b}$$

$$\leq \sum_{m=1}^{M} \left(I_m L \tau \left(2\frac{\gamma}{T} + 2\mathbb{E}\big[\mathbf{Rad}_m(\mathbf{x}_m)\big] \right) \right) + O\big(\tau \log(T)\big) \tag{29c}$$

$$\leq 2 \sum_{m=1}^{M} I_m L \tau \frac{\gamma}{T} + 2 \sum_{\mathbf{x} \in \mathcal{X}} \sum_{m \in \mathcal{M}_{\mathbf{x}}} I_m L \tau \mathbb{E}\big[\mathbf{Rad}_m(\mathbf{x})\big] + O\big(\tau \log(T)\big) \tag{29d}$$

$$= 2\gamma + \leq \sum_{\mathbf{x} \in \mathcal{X}} O\left(J\sqrt{|\mathcal{M}_{\mathbf{x}}|T_{\mathbf{x}} \log(J^{1/4}T)} \right) + O\big(\tau \log(T)\big) \tag{29e}$$

$$\leq \sum_{\mathbf{x} \in \mathcal{X}} O\left(J\sqrt{|\mathcal{M}_{\mathbf{x}}|T_{\mathbf{x}} \log(J^{1/4}T)} \right) + O\big((\log(T))^3\big) \tag{29f}$$

$$= \sum_{\mathbf{x} \in \mathcal{X}} O\left(J\sqrt{|\mathcal{M}_{\mathbf{x}}|T_{\mathbf{x}} \log(J^{1/4}T)} \right) \tag{29g}$$

$$= O\left(J\sqrt{XT \log T \log(J^{1/4}T)} \right), \tag{29h}$$

where (29e) is by Lemma 2 and (29f) is by Jensen's inequality.

Regret of RNRM-TS algorithm. By adding up the relaxed regret during the Warm-up and Learning Phases, we have the regret of RNRM-TS algorithm as:

$$\overline{\mathrm{BR}}(\mathrm{RNRM\text{-}TS}, T) = \overline{\mathrm{BR}}_{\mathrm{Warm\text{-}up}}(\mathrm{RNRM\text{-}TS}, T) + \overline{\mathrm{BR}}_{\mathrm{Learning}}(\mathrm{RNRM\text{-}TS}, T) \tag{30a}$$

$$= O(X \log(J^{1/4}T)) + O\left(J\sqrt{XT \log T \log(J^{1/4}T)} \right) \tag{30b}$$

$$= \tilde{O}(J\sqrt{XT}). \tag{30c}$$

Q.E.D.

Lemma 2. *For any price* $\mathbf{x} \in \mathcal{X}$,

$$\sum_{m \in \mathcal{M}_{\mathbf{x}}} I_m L \tau \mathbb{E}\big[\mathbf{Rad}_m(p)\big] = O\left(J\sqrt{|\mathcal{M}_{\mathbf{x}}|T_{\mathbf{x}} \log(J^{1/4}T)} \right). \tag{31}$$

Proof of Lemma 2. Plugging the definition of $\mathbf{Rad}_m(\mathbf{x})$, $\forall m = 1, \ldots, M$, one has:

$$\sum_{m \in \mathcal{M}_\mathbf{x}} I_m L \tau \mathbb{E}\big[\mathbf{Rad}_m(\mathbf{x})\big] = O\left(\sqrt{\log(J^{1/4}T)}\right) \sum_{m \in \mathcal{M}_\mathbf{x}} I_m L \tau \cdot O\left(\sum_{j \in \mathcal{J}_{on}(\mathbf{x})} \frac{1}{\sqrt{n_{mj}(\mathbf{x})}}\right) \tag{32a}$$

$$\leq O\left(\sqrt{\log(J^{1/4}T)}\right) \sum_{m \in \mathcal{M}_\mathbf{x}} I_m L \tau \cdot O\left(\sum_{j \in \mathcal{J}_{on}(\mathbf{x})} \frac{1}{\sqrt{I_m L \tau}}\right) \tag{32b}$$

$$= O\left(\sqrt{\log(J^{1/4}T)}\right) \sum_{m \in \mathcal{M}_\mathbf{x}} I_m L \tau \cdot O\left(\frac{J}{\sqrt{I_m L \tau}}\right) \tag{32c}$$

$$= O\left(J\sqrt{\log(J^{1/4}T)}\right) \sum_{m \in \mathcal{M}_\mathbf{x}} \sqrt{I_m L \tau} \tag{32d}$$

$$\leq O\left(J\sqrt{\log(J^{1/4}T)}\right) \cdot \sqrt{|\mathcal{M}_\mathbf{x}| \sum_{m \in \mathcal{M}_\mathbf{x}} I_m L \tau} \tag{32e}$$

$$= O\left(J\sqrt{|\mathcal{M}_\mathbf{x}|T_\mathbf{x} \log(J^{1/4}T)}\right) \tag{32f}$$

$$\tag{32g}$$

where and (32b) is by $n_{mj}(\mathbf{x}) = O(I_m L \tau)$, $\forall j \in \mathcal{J}_{on}(\mathbf{x})$ and (32e) is by Jensen's inequality.

Lemma 3. *(Coupling probability of M/M/c queue starting from empty state, Proposition 4 in Jia et al. 2020) For $\tau = (\log T)^2$, if an M/M/c queue starts from empty state at time 0, then the probability that the system reaches the steady-state within τ is bounded by $1 - \frac{2}{T^2}$.*

Lemma 4. *(Coupling probability of M/M/c queue when price changes, Proposition 5 in Jia et al. 2020) For $\tau = (\log T)^2$, if an M/M/c queue starts from steady-state and changes system parameter at time 0, then the probability that the system reaches the steady-state within 2τ is bounded by $1 - \frac{4}{T^2}$.*

F Service Duration Dependent Payment Model

We consider a new variant payment model where the customer pays the product's per-unit-time price multiplied by its total service duration. The firm then needs to decide the per-unit-time price for each product $j \in \mathcal{J}$. Under this payment setting, we modify the definition of the steady-state revenue rate, derive a new concentration bound of historical observations and actual system parameters, and then correspondingly develop new UCB of the revenue rate of each price. With those modified definitions, we present a new RNRM-UCB variant and show that it achieves a same regret bound.

Data collection. Similar to n_j, $\hat{d}_{jk}(\mathbf{x})$, and $\bar{d}_j(\mathbf{x})$, we also collect service time observations over resources. We use $n_i^r(\mathbf{x})$ to denote the number of customers who successfully consume resource i under price $\mathbf{x}$. We use $\hat{g}_{ik}(\mathbf{x})$ to denote the service time of customer k over resource i under the price $\mathbf{x}$. We use $\bar{g}_i(\mathbf{x})$ to denote the empirical mean of the service time and $\bar{g}_i(\mathbf{x}) = \sum_{k=1}^{n_i^r(\mathbf{x})} \hat{g}_{ik}(\mathbf{x})/n_i^r(\mathbf{x})$, for $i \in \mathcal{I}$. We further assume that there is a known upper bound $r_{\max} = \max_{x,i,j} \alpha_j/\mu_i$, $n_j^{\min}(\mathbf{x}) = \min\{n_j(\mathbf{x}), \min_{i:a_{ji}=1} n_i^s(\mathbf{x})\}$.

Revenue rate. Under this duration-dependent payment model, we offer unit price $\mathbf{x} \in \mathbb{R}^J$ to products $\mathcal{J}$. Customer k who requests to use product j pays the per-unit-time price multiplied by the total service duration, i.e., $x_j \sum_{i:a_{ji}=1} \hat{g}_{ik}(\mathbf{x})$, to the firm. Different than the lump sum payment model, the steady-state revenue rate for this setting is $r^v(\mathbf{x}) = \sum_{j \in \mathcal{J}_{on}(\mathbf{x})} \alpha_j x_j \sum_{i:a_{ji}=1} 1/\mu_i(\mathbf{x})$.

Proposition 5. *For any price $\mathbf{x} \in \mathcal{X}$, if the number of arrival time observations for each turn-on product $n_j(\mathbf{x}) \geq 8\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)$ for $j \in \mathcal{J}_{on}(\mathbf{x})$ and the number of service time observations for each resource $n_i^s(\mathbf{x}) \geq 8\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)$ $\forall i \in \{i \in \mathcal{I} : \sum_{j \in \mathcal{J}_{on}(\mathbf{x})} a_{ji} \geq 1\}$, then we have*

$$\mathbb{P}\left(\left| r^v(\mathbf{x}) - \sum_{j \in \mathcal{J}_{on}(\mathbf{x})} \frac{x_j \sum_{i:a_{ji}=1} \bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})} \right| \leq \sum_{j \in \mathcal{J}_{on}(\mathbf{x})} \frac{\sum_{i:a_{ji}=1} (\frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})} + r_{max}) x_j \sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\sqrt{n_j^{min}(\mathbf{x})}} \right) \geq 1 - \frac{4}{T^4},$$

 where r_{max} is a known upper bound with r_{max} $=$ $\max_{x,i,j} \alpha_j/\mu_i$, $n_j^{min}(\mathbf{x})$ $=$
 $\min\{n_j(\mathbf{x}), \min_{i:a_{ji}=1} n_i^s(\mathbf{x})\}$, and L is the number of layers.

 Proof of Proposition 5.

The above probability can be bounded by the product of the following probabilities from below. Firstly we can bound that for i, j with $a_{ji} = 1$:

$$\mathbb{P}\left(\left|\frac{\alpha_j}{\mu_i(\mathbf{x})} - \frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})}\right| \leq \frac{(\frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})} + r_{\max})\sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\sqrt{\min\{n_j(\mathbf{x}), n_i^r(\mathbf{x})\}}}\right)$$

$$\geq \mathbb{P}\left(\left|\frac{\alpha_j}{\mu_i(\mathbf{x})} - \alpha_j\bar{g}_i(\mathbf{x}) + \alpha_j\bar{g}_i(\mathbf{x}) - \frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})}\right| \leq \frac{(\frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})} + r_{\max})\sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\sqrt{\min\{n_j(\mathbf{x}), n_i^r(\mathbf{x})\}}}\right)$$

$$\geq \mathbb{P}\left(\alpha_j\left|\frac{1}{\mu_i(\mathbf{x})} - \bar{g}_i(\mathbf{x})\right| + \bar{g}_i(\mathbf{x})\left|\alpha_j - \frac{1}{\bar{d}_j(\mathbf{x})}\right| \leq \frac{(\frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})} + r_{\max})\sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\sqrt{\min\{n_j(\mathbf{x}), n_i^r(\mathbf{x})\}}}\right)$$

$$\geq \mathbb{P}\left(\alpha_j\left|\frac{1}{\mu_i(\mathbf{x})} - \bar{g}_i(\mathbf{x})\right| \leq \frac{r_{\max}\sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\sqrt{n_i^r(\mathbf{x})}}\right) \cdot \mathbb{P}\left(\bar{g}_i(\mathbf{x})\left|\alpha_j - \frac{1}{\bar{d}_j(\mathbf{x})}\right| \leq \frac{\frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})}\sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\sqrt{n_j(\mathbf{x})}}\right)$$

$$\geq \mathbb{P}\left(\left|\frac{1}{\mu_i(\mathbf{x})} - \bar{g}_i(\mathbf{x})\right| \leq \frac{\sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\mu_i(\mathbf{x})\sqrt{n_i^r(\mathbf{x})}}\right) \cdot \mathbb{P}\left(\left|\alpha_j - \frac{1}{\bar{d}_j(\mathbf{x})}\right| \leq \frac{\sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\bar{d}_j(\mathbf{x})\sqrt{n_j(\mathbf{x})}}\right).$$

The first and second probability in the above equation can be bounded with the same technique as that we used in (7) with Lemma 1 and thus we have

$$\mathbb{P}\left(\left|\alpha_j - \frac{1}{\bar{d}_j(\mathbf{x})}\right| \leq \frac{\sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\bar{d}_j(\mathbf{x})\sqrt{n_j(\mathbf{x})}}\right) \geq 1 - \frac{2}{LJT^4}$$

and

$$\mathbb{P}\left(\left|\frac{1}{\mu_i(\mathbf{x})} - \bar{g}_i(\mathbf{x})\right| \leq \frac{\sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\mu_i(\mathbf{x})\sqrt{n_i^r(\mathbf{x})}}\right) \geq 1 - \frac{2}{LJT^4}.$$

Therefore, we have the above probability can be bounded by

$$\left(1 - \frac{4}{LJT^4}\right)^2 \geq 1 - \frac{4}{LJT^4}.$$

Then, for product $j \in \mathcal{J}_{\text{on}}(\mathbf{x})$, we can further bound

$$\mathbb{P}\left(\left|\sum_{i:a_{ji}=1}\frac{\alpha_j}{\mu_i(\mathbf{x})} - \sum_{i:a_{ji}=1}\frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})}\right| \leq \sum_{i:a_{ji}=1}\frac{(\frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})} + r_{\max})\sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\sqrt{n_j^{\min}(\mathbf{x})}}\right)$$

$$\geq \prod_{i:a_{ji}=1}\mathbb{P}\left(\left|\frac{\alpha_j}{\mu_i(\mathbf{x})} - \frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})}\right| \leq \frac{(\frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})} + r_{\max})\sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\sqrt{n_j^{\min}(\mathbf{x})}}\right)$$

$$\geq \left(1 - \frac{4}{LJT^4}\right)^{|\{i:a_{ji}=1\}|} \geq \left(1 - \frac{4}{LJT^4}\right)^L \geq 1 - \frac{4}{JT^4},$$

 because $\sum_{i\in\mathcal{I}} a_{ji} \leq L \; \forall j \in \mathcal{J}$.

Now we are ready to derive the concentration bound.

$$\mathbb{P}\left(\left|r^v(\mathbf{x}) - \sum_{j\in\mathcal{J}_{\text{on}}(\mathbf{x})} \frac{x_j \sum_{i:a_{ji}=1} \bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})}\right| \leq \sum_{j\in\mathcal{J}_{\text{on}}(\mathbf{x})} \frac{\sum_{i:a_{ji}=1}(\frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})}+r_{\max})x_j\sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\sqrt{n_j^{\min}(\mathbf{x})}}\right)$$

$$\geq \prod_{j\in\mathcal{J}_{\text{on}}(\mathbf{x})} \mathbb{P}\left(\left|\sum_{i:a_{ji}=1}\frac{\alpha_j}{\mu_i(\mathbf{x})} - \sum_{i:a_{ji}=1}\frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})}\right| \leq \sum_{i:a_{ji}=1}\frac{(\frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})}+r_{\max})\sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\sqrt{n_j^{\min}(\mathbf{x})}}\right)$$

$$\geq \left(1-\frac{4}{J\cdot T^4}\right)^{|\mathcal{J}_{\text{on}}(\mathbf{x})|} \geq \left(1-\frac{4}{J\cdot T^4}\right)^{J} \geq 1-\frac{4J}{J\cdot T^4} = 1-\frac{4}{T^4}.$$

Q.E.D.

According to the above proposition, we can define the confidence radius as follows. (We use the superscript v to denote this duration-dependent payment model.)

$$\mathbf{Rad}_t^v(\mathbf{x}) = \sum_{j\in\mathcal{J}_{\text{on}}(\mathbf{x})} \frac{\sum_{i:a_{ji}=1}(\frac{\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})}+r_{\max})x_j\sqrt{32\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)}}{\sqrt{n_j^{\min}(\mathbf{x})}}.$$

Therefore, the upper and lower confidence bound of the revenue rate associated with each price is

$$U_t^v(\mathbf{x}) = \sum_{j\in\mathcal{J}_{\text{on}}(\mathbf{x})} \frac{x_j\sum_{i:a_{ji}=1}\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})} + \mathbf{Rad}_t^v(\mathbf{x}), \qquad L_t^v(\mathbf{x}) = \sum_{j\in\mathcal{J}_{\text{on}}(\mathbf{x})} \frac{x_j\sum_{i:a_{ji}=1}\bar{g}_i(\mathbf{x})}{\bar{d}_j(\mathbf{x})} - \mathbf{Rad}_t^v(\mathbf{x}).$$

Algorithm design. The algorithmic framework remains unchanged except for two modifications. (i) In Step 6, the algorithm also requires the number of service time observations to be at least $8\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)$. (ii) In Steps 13 and 16, the algorithm uses the new UCB definition.

Algorithm 3 RNRM-UCB (Duration-Dependent Payment Model.)

1: Input: T, $\mathcal{X}$, service network, turn-off price.
2: Initialize: L, τ, I_m, M, $m=0$.
3: Warm-up Phase:
4: **for** $\mathbf{x} \in \mathcal{X}$ **do**
5:     Offer $\mathbf{x}$, record inter-arrival time $\hat{d}_{jk}(\mathbf{x})$ for arriving customers of product $j \in \mathcal{J}_{\text{on}}(\mathbf{x})$.
6:     **if** $n_{mj}^{\min}(\mathbf{x}) \geq 8\log(L^{\frac{1}{4}}J^{\frac{1}{4}}T)$ **then**
7:         Continue.
8:     **end if**
9: **end for**
10: Compute and update UCB for all prices as defined in (2).
11: Learning Phase:
12: **for** $m = 1, \ldots, M$ **do**
13:     Choose $\mathbf{x}_m = \arg\max_{\mathbf{x}\in} U_{m-1}^v(\mathbf{x})$.
14:     Offer $\mathbf{x}_m$ in batch m, i.e., for $I_m L\tau$ periods.
15:     Record inter-arrival time $\hat{d}_{jk}(\mathbf{x}_m)$ for arriving customers of product $j \in \mathcal{J}_{\text{on}}(\mathbf{x}_m)$.
16:     Compute and update $U_m^v(\mathbf{x_m})$.
17: **end for**

Regret analysis. We change the constraints (1d) in the LP benchmark to

$$Y_{\mathbf{y}t} \leq \sum_{\mathbf{x}\in\mathcal{X}} \pi_{\mathbf{y}t}^{\mathbf{x}} \cdot \sum_{j\in\mathcal{J}_{\text{on}}(\mathbf{x})} \alpha_j x_j \sum_{i:a_{ji}=1} \frac{1}{\mu_i(\mathbf{x})} \quad \forall\mathbf{y}\in\{0,1,\ldots,\infty\}^I, \; \forall t=1,\ldots,T.$$

Under this duration-dependent payment setting, we can follow the same analysis technique to show that this LP model also gives a valid upper bound of the optimal revenue. The results of Propositions 1, 3, and 4 and Theorem 1 remain the same for this duration-dependent payment model.