The Curse of Unrolling:
Rate of Differentiating Through Optimization

Abstract

Computing the Jacobian of the solution of an optimization problem is a central problem in machine learning, with applications in hyperparameter optimization, meta-learning, optimization as a layer, and dataset distillation, to name a few. Unrolled differentiation is a popular heuristic that approximates the solution using an iterative solver and differentiates it through the computational path. This work provides a non-asymptotic convergence-rate analysis of this approach on quadratic objectives for gradient descent and the Chebyshev method. We show that to ensure convergence of the Jacobian, we can either 1) choose a large learning rate leading to a fast asymptotic convergence but accept that the algorithm may have an arbitrarily long burn-in phase or 2) choose a smaller learning rate leading to an immediate but slower convergence. We refer to this phenomenon as the *curse of unrolling*. Finally, we discuss open problems relative to this approach, such as deriving a practical update rule for the optimal unrolling strategy and making novel connections with the field of Sobolev orthogonal polynomials.

Figure 1: **The Curse of Unrolling:** Better function suboptimality does not imply better Jacobian suboptimality. The speed of the function suboptimality (right) $f(\boldsymbol{x}_t(\boldsymbol{\theta}), \boldsymbol{\theta}) - f(\boldsymbol{x}_\star(\boldsymbol{\theta}), \boldsymbol{\theta})$ of gradient descent is not representative of the speed of convergence of the Jacobian $\|\partial_{\boldsymbol{\theta}} \boldsymbol{x}_t(\boldsymbol{\theta}) - \partial_{\boldsymbol{\theta}} \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F$ (left) at early iterations. Even worse, the optimal step size $2/(L + \ell)$ for function suboptimality yields a *burn-in phase* where the Jacobian suboptimality increases.

1   Introduction

Let $\boldsymbol{x}_\star(\boldsymbol{\theta})$ be a function defined implicitly as the solution to an optimization problem, $\boldsymbol{x}_\star(\boldsymbol{\theta}) = \arg\min_{\boldsymbol{x} \in \mathbb{R}^d} f(\boldsymbol{x}, \boldsymbol{\theta})$. Implicitly defined function of this form appear in different areas of machine learning, such as reinforcement learning (Du et al., 2017; Pfau et al., 2016), generative adversarial networks (Metz et al., 2016), hyper-parameter optimization (Bengio, 2000; Bertrand et al., 2020; Franceschi et al., 2017; Lorraine et al., 2020; Pedregosa, 2016), meta-learning (Franceschi et al., 2018; Rajeswaran et al., 2019), deep equilibrium models, (Bai et al., 2019) or optimization as a

layer (Amos et al., 2017; Kim et al., 2017; Wang et al., 2019), to name a few. The main computational burden of using implicit functions in a machine learning pipeline is that the Jacobian computation $\partial_{\boldsymbol{\theta}} \boldsymbol{x}_\star(\boldsymbol{\theta})$ is challenging: since the implicit function $\boldsymbol{x}_\star(\boldsymbol{\theta})$ does not usually admit an explicit formula, classical automatic differentiation techniques cannot be applied directly.

Two main approaches have emerged to compute the Jacobian of implicit functions: *implicit* differentiation and *unrolled* differentiation. This paper focuses on unrolled differentiationm while recent surveys on implicit differentiation are (Blondel et al., 2021; Duvenaud et al., 2020).

Unrolled differentiation, also known as iterative differentiation, starts by approximating the implicit function $\boldsymbol{x}_\star(\cdot)$ by the output of iterative algorithm, which we denote $\boldsymbol{x}_t(\cdot)$, and then differentiates through the algorithm's computational path (Deledalle et al., 2014; Domke, 2012; Franceschi et al., 2018; Shaban et al., 2019; Wengert, 1964) mode differentiation.

Contributions. We analyze the convergence of the unrolled Jacobian by establishing worst-case bounds on the Jacobian suboptimality $\|\partial \boldsymbol{x}_t(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|$ for different methods, more precisely:

1. We provide a general framework for analyzing unrolled differentiation on quadratic objectives for any gradient-based method (Theorem 1). For gradient descent and the Chebyshev iterative method, we derive closed-form worst-case convergence rates in Corollary 1 and Theorem 3.

2. We identify the "curse of unrolling" as a consequence of this analysis: A fast asymptotic rate inevitably leads to a condition number-long burn-in phase where the Jacobian suboptimality increases. While it is possible to reduce the length, or the peak, of this burn-in phase, this comes at the cost of a slower asymptotic rate (Figure 5).

3. Finally, we describe a novel approach to mitigate the curse of unrolling, motivated by the theory of Sobolev orthogonal polynomials (Theorem 4).

We consider Ablin et al. (2020) to be the closest related work, although it focuses on a more restrictive setting: the differentiation of functions defined as a minimum, instead of an *argmin* as in this work. While most of the problems mentioned in the introduction fit our setting, only a few of them have the specific structure studied by Ablin et al. (2020). They develop a complexity analysis for non-quadratic functions, the rate they obtain are valid only for monotone optimization algorithms, such as gradient descent with small step size. In our case, we obtain rates for non-monotone and accelerated algorithm such as GD with large step size, or the Chebyshev method.

2 Preliminaries and Notations

In this paper, we consider an objective function f parametrized by two variables $\boldsymbol{x} \in \mathbb{R}^d$ and $\boldsymbol{\theta} \in \mathbb{R}^n$. We are interested in the derivative of optimization problem solutions:

$$\text{\textbf{Goal}: approximate Jacobian } \partial \boldsymbol{x}_\star(\boldsymbol{\theta}), \text{ where } \boldsymbol{x}_\star(\boldsymbol{\theta}) = \underset{\boldsymbol{x} \in \mathbb{R}^d}{\arg \min} f(\boldsymbol{x}, \boldsymbol{\theta}). \tag{OPT}$$

We also assume $\boldsymbol{x}_\star(\boldsymbol{\theta}) \in \mathbb{R}^d$ is the unique minimizer of $f(\boldsymbol{x}, \boldsymbol{\theta})$, for some fixed value of $\boldsymbol{\theta}$. In particular, we will describe the rate of convergence of $\|\partial \boldsymbol{x}_t(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|$ in the specific case where f is a quadratic function in its first argument, and $\boldsymbol{x}_t$ is generated by a first-order method.

Notations. In this paper, we use upper-case letter for polynomials (P, Q), bold lower-case for vectors ($\boldsymbol{x}$, $\boldsymbol{b}$), and bold upper-case for matrices ($\boldsymbol{H}$). We write $\mathcal{P}_t$ the set of polynomials of degree at least t. We distinguish $\nabla f(\boldsymbol{x}, \boldsymbol{\theta})$, that refers to the gradient of the function f in its first argument, and $\partial_{\boldsymbol{\theta}} f(\boldsymbol{x}, \boldsymbol{\theta})$ is the partial derivative of f w.r.t. its second argument, evaluated at $(\boldsymbol{x}, \boldsymbol{\theta})$ (if there is no ambiguity, we write ∂ instead of $\partial_{\boldsymbol{\theta}}$). Similarly, $\partial \boldsymbol{x}(\boldsymbol{\theta})$ is the Jacobian of the vector-valued function $\boldsymbol{x}(\cdot)$ evaluated at $\boldsymbol{\theta}$ and $P'(\cdot)$ the derivative of the polynomial P. Finally, ℓ is the strong convexity constant, L the smoothness constant, and $\kappa = \ell/L$ is the inverse of the condition number.

2.1 Problem Setting and Main Assumptions

Throughout the paper, the following three assumptions hold: The first two assume the problem is quadratic and the Hessian commutes with its derivative respectively, while the third one restricts the class of algorithms to first-order methods.

Assumption 1 (Quadratic objective). *The function f is a quadratic function in its first argument,*

$$f(\boldsymbol{x}, \boldsymbol{\theta}) \stackrel{def}{=} \tfrac{1}{2}\boldsymbol{x}^\top \boldsymbol{H}(\boldsymbol{\theta})\boldsymbol{x} + \boldsymbol{b}(\boldsymbol{\theta})^\top \boldsymbol{x} \,, \tag{1}$$

where $\ell\boldsymbol{I} \preceq \boldsymbol{H}(\boldsymbol{\theta}) \preceq L\boldsymbol{I}$ for $0 < \ell < L$. We write $\boldsymbol{x}_\star(\boldsymbol{\theta})$ the minimizer of f w.r.t. the first argument.

Assumption 2 (Commutativity of Jacobian). *We assume $\partial\boldsymbol{H}(\boldsymbol{\theta})$ commutes with $\boldsymbol{H}(\boldsymbol{\theta})$.*

Remark. This assumption is critical for our polynomial-based analysis, as it allows to have a simpler expression for the Jacobian of powers of $\boldsymbol{H}^t$:

$$\partial\boldsymbol{H}^t(\boldsymbol{\theta}) = t\boldsymbol{H}^{t-1}(\boldsymbol{\theta})\partial\boldsymbol{H}(\boldsymbol{\theta}) \,, \text{ instead of } \partial\boldsymbol{H}^t(\boldsymbol{\theta}) = \sum_{i=0}^{t-1}\boldsymbol{H}^i(\boldsymbol{\theta})\partial\boldsymbol{H}(\boldsymbol{\theta})\boldsymbol{H}^{t-i-1}(\boldsymbol{\theta}) \,. \tag{2}$$

This assumption is verified for example for Ridge regression (see below). Although quite restrictive, we believe our it could potentially be removed in future work (see Appendix A).

Example 1 (Ridge regression). *Let us fix $\boldsymbol{A}, \bar{\boldsymbol{x}}, \boldsymbol{y}$, and let $\boldsymbol{H}(\theta) = \boldsymbol{A}^\top\boldsymbol{A} + \theta\boldsymbol{I}$, where θ is a scalar. The ridge regression problem*

$$f(\boldsymbol{x}, \theta) = \tfrac{1}{2}\left(\|\boldsymbol{A}\boldsymbol{x} - \boldsymbol{y}\|_2^2 + \theta\|\boldsymbol{x} - \bar{\boldsymbol{x}}\|_2^2\right) \,,$$

satisfies Assumptions 1, 2, as $f(\boldsymbol{x}, \theta)$ is quadratic in $\boldsymbol{x}$, and $\partial\boldsymbol{H}(\theta)\boldsymbol{H}(\theta) = \boldsymbol{H}(\theta)\partial\boldsymbol{H}(\theta) = \boldsymbol{H}(\theta)$.

Our last assumption concerns class of algorithms. We restrict ourselves to *first-order method*, widely used in large-scale optimization. This class includes methods like gradient descent or Polyak's heavy-ball method (Nesterov, 2004).

Assumption 3 (First-order method). *The iterates $\{\boldsymbol{x}_t\}_{t=0...}$ comes from first-order method:*

$$\boldsymbol{x}_t(\boldsymbol{\theta}) \in \boldsymbol{x}_0(\boldsymbol{\theta}) + \boldsymbol{Span}\{\nabla f(\boldsymbol{x}_0(\boldsymbol{\theta}), \boldsymbol{\theta}), \,\ldots, \, \nabla f(\boldsymbol{x}_{t-1}(\boldsymbol{\theta}), \boldsymbol{\theta})\} \,.$$

2.2 Polynomials and First-Order Methods on Quadratics

The polynomial formalism has seen a revival in recent years, thanks to its simple and constructive analysis (Agarwal et al., 2021; Pedregosa et al., 2020; Scieur et al., 2020a). It starts from a connection between optimization methods and polynomials that allows casting complexity analysis of optimization methods as a problem of bounding polynomials.

2.2.1 Connection with Residual Polynomials

There is a strong connection between first-order methods and polynomials: after t iterations, one can associate to any optimization method a polynomial $P_t(\lambda) = a_t\lambda^t + \cdots + 1$ of degree t such that $P_t(0) = 1$. The error at iteration t then can then be expressed as

$$\boldsymbol{x}_t(\boldsymbol{\theta}) - \boldsymbol{x}_\star(\boldsymbol{\theta}) = P_t(\boldsymbol{H}(\boldsymbol{\theta}))(\boldsymbol{x}_0(\boldsymbol{\theta}) - \boldsymbol{x}_\star(\boldsymbol{\theta})) \,. \tag{3}$$

This polynomial $P_t(\boldsymbol{H}(\boldsymbol{\theta}))$ is called the *residual polynomial* and represents the output of evaluating the originally real-valued polynomial $P_t(\cdot)$ at the matrix $\boldsymbol{H}$.

Example 2. *In the case of gradient descent, the update reads $\boldsymbol{x}_{t+1} - \boldsymbol{x}^\star = (I - \gamma\boldsymbol{H})(\boldsymbol{x}_t - \boldsymbol{x}^\star)$, which yields the residual polynomial $P_t(\boldsymbol{H}) = (I - \gamma\boldsymbol{H})^t$.*

2.2.2 Worst-Case Convergence Bound

Using the relationship described above between first-order method and its residual polynomial, one can quickly compute a worst-case on the convergence bound. Using the Cauchy-Schwartz inequality on (3), we obtain

$$\|\boldsymbol{x}_t(\boldsymbol{\theta}) - \boldsymbol{x}_\star(\boldsymbol{\theta})\| \le \|P_t(\boldsymbol{H}(\boldsymbol{\theta}))\| \, \|\boldsymbol{x}_0(\boldsymbol{\theta}) - \boldsymbol{x}_\star(\boldsymbol{\theta})\| \,.$$

We are interested in the performance of the first-order method on a class of quadratic functions (see Assumption 1), whose Hessian has bounded eigenvalues. Using the fact that the ℓ_2-norm of a matrix is equal to its largest singular value, the worst-case performance of the algorithm then reads

$$\|\boldsymbol{x}_t(\boldsymbol{\theta}) - \boldsymbol{x}_\star(\boldsymbol{\theta})\| \le \max_{\lambda\in[\ell,L]}|P_t(\lambda)| \, \|\boldsymbol{x}_0(\boldsymbol{\theta}) - \boldsymbol{x}_\star(\boldsymbol{\theta})\| \,. \tag{4}$$

Therefore, the worst-case convergence bound is a function of the polynomial associated with the first-order method, and depends on the bound over the eigenvalue of the Hessian ($\lambda \in [\ell, L]$).

2.2.3 Expected Spectral Density and Average-Case Complexity

We recall the average-case complexity framework (Cunha et al., 2022; Paquette et al., 2022; Pedregosa et al., 2020), which provides a finer-grained convergence analysis than the worst-case. This framework is crucial in developing an accelerated method for unrolled differentiation (Section 4).

Instead of considering the worst instance from a class of quadratic functions, average-case analysis considers that functions are drawn *at random* from the class. This means that, in Assumption 1, the matrix $\boldsymbol{H}(\boldsymbol{\theta})$, the vector $\boldsymbol{b}(\boldsymbol{\theta})$ and the initialization $\boldsymbol{x}_0(\theta)$ in Assumption 3 are sampled from some (potentially unknown) probability distributions. Surprisingly, we do not require the knowledge of these distributions, instead, the quantity of interest is the *expected spectral density* $\mu(\lambda)$, defined as

$$\mu(\lambda) \stackrel{\text{def}}{=} \mathbb{E}_{\boldsymbol{H}(\boldsymbol{\theta})}[\mu_{\boldsymbol{H}(\boldsymbol{\theta})}(\lambda)], \qquad \mu_{\boldsymbol{H}(\boldsymbol{\theta})}(\lambda) \stackrel{\text{def}}{=} \tfrac{1}{d} \sum_{i=1}^{d} \delta(\lambda - \lambda_i(\boldsymbol{H}(\boldsymbol{\theta}))). \tag{5}$$

In Equation 5, $\lambda_i(\boldsymbol{H}(\theta))$ is the i-th eigenvalue of $\boldsymbol{H}(\boldsymbol{\theta})$, $\delta(\cdot)$ is the Dirac's delta and $\mu_{\boldsymbol{H}(\boldsymbol{\theta})}(\lambda)$ is the *empirical spectral density* (i.e., $\mu_{\boldsymbol{H}(\boldsymbol{\theta})}(\lambda) = 1/d$ if λ is an eigenvalue of $\boldsymbol{H}(\boldsymbol{\theta})$ and 0 otherwise).

Assuming $\boldsymbol{x}_0(\boldsymbol{\theta}) - \boldsymbol{x}_\star(\boldsymbol{\theta})$ is independent of $\boldsymbol{H}(\boldsymbol{\theta})$,[1] the average-case complexity of the first-order method associated to the polynomial P_t as

$$\mathbb{E}[\|\boldsymbol{x}_t(\boldsymbol{\theta}) - \boldsymbol{x}_\star(\boldsymbol{\theta})\|^2] = \mathbb{E}[\|\boldsymbol{x}_0(\boldsymbol{\theta}) - \boldsymbol{x}_\star(\boldsymbol{\theta})\|^2] \int P_t^2(\lambda) \, \mathrm{d}\mu(\lambda). \tag{6}$$

Here, the term in P_t is algorithm-related, while the term in $\mathrm{d}\mu$ is related to the difficulty of the (distribution over the) problem class. As opposed to the worst-case analysis, where only the *worst* value of the polynomial impacts the convergence rate, the average-case rate depends on the expected value of the squared polynomial P_t over the *whole* distribution. Note that in the previous equation, the expectation is taken over problem instances and not over any stochasticity of the algorithm.

3 The Convergence Rate of differentiating through optimization

We now analyze the rate of convergence of gradient descent and Chebyshev algorithm (optimal on quadratics). We first introduce the *master identity* (Theorem 1), which draws a link between how well a first-order methods estimates $\partial \boldsymbol{x}_\star(\boldsymbol{\theta})$, its associated residual polynomial P_t, and its derivative P_t'.

Theorem 1 (Master identity). *Under Assumptions 1, 2, 3, let $\boldsymbol{x}_t(\boldsymbol{\theta})$ be the t^{th} iterate of a first-order method associated to the residual polynomial P_t. Then the Jabobian error can be written as*

$$\partial \boldsymbol{x}_t(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta}) = \big(P_t(\boldsymbol{H}(\boldsymbol{\theta})) - \boldsymbol{H}(\boldsymbol{\theta})P_t'(\boldsymbol{H}(\boldsymbol{\theta}))\big)(\partial \boldsymbol{x}_0(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta}))$$
$$+ P_t'(\boldsymbol{H}(\boldsymbol{\theta}))\partial_{\boldsymbol{\theta}} \nabla f(\boldsymbol{x}_0(\boldsymbol{\theta}), \boldsymbol{\theta}). \tag{7}$$

The above identity involves the **derivative** of the residual polynomials and not only the **residual** polynomial, as was the case for minimization (3). As we will see, this difference will generate different convergence rates for the Jacobian suboptimality than classical results for objective or iterate suboptimality.

For conciseness, our bounds make use of the following shorthand notation

$$G \stackrel{\text{def}}{=} \|\partial_{\boldsymbol{\theta}} \nabla f(\boldsymbol{x}_0(\boldsymbol{\theta}), \boldsymbol{\theta})\|_F.$$

3.1 Worst-case Rates for Gradient Descent

We consider the fixed-step gradient descent algorithm,

$$\boldsymbol{x}_t(\boldsymbol{\theta}) = \boldsymbol{x}_{t-1}(\boldsymbol{\theta}) - \nabla f(\boldsymbol{x}_{t-1}(\boldsymbol{\theta}), \boldsymbol{\theta}). \tag{8}$$

As mentioned in Example 2, the associated polynomial reads $P_t = (1 - h\lambda)^t$. We can deduce its convergence rate after injecting the polynomial in the master identity from Theorem 1.

Theorem 2. *Under Assumptions 1, 2, let $\boldsymbol{x}_t(\boldsymbol{\theta})$ be the t^{th} iterate of gradient descent scheme with step size $h > 0$. Then,*

$$\|\partial \boldsymbol{x}_t(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F \leq \max_{\lambda \in [\ell, L]} \left| (1 - h\lambda)^{t-1} \left\{ (1 + (t-1)h\lambda)\|\partial \boldsymbol{x}_0(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F + htG \right\} \right|.$$

---

[1]This assumption can be removed as in (Cunha et al., 2022) at the price of a more complicated expected spectral density.

Figure 2: **The Phases of Unrolling**. Gradient descent and the Chebyshev method exhibit two distinct phases during unrolled differentiation: an initial burn-in phase, where the Jacobian suboptimality increases, followed by a convergent phase with an asymptotic linear convergence. The maximum of the suboptimality is similar for both algorithms, but Chebyshev peaks sooner than gradient descent, after a number of iterations equal to (roughly) the square root of the one required by gradient descent. The distinct phases, as well as their relative duration, are predicted by the theoretical worst-case bounds of Corollary 1 and Theorem 3. Both plots are run on the same problem, a ridge regression loss on the `breast-cancer` dataset.

Discussion. The bound above is a product of two terms: the first term, $(1 - h\lambda)^{t-1}$, is the convergence rate of gradient descent and decreases exponentially for $h \leq \frac{2}{L+\ell}$, while the second term is increasing in t. This results in two distinct phases in training: an initial **burn-in** phase, where the second term dominates and the Jacobian suboptimality might be increasing, followed by an **linear convergence** phase where the exponential term dominates. This phenomenon can be seen empirically, see Figure 1. As an illustration, the next corollary exhibits explicit rates for gradient descent in the two special cases where $h = \frac{1}{L}$ (short steps) and $h = \frac{2}{L+\ell}$ (large steps, maximize asymptotic rate).

Corollary 1. *For the step size $h = 1/L$, the rate of Theorem 2 reads*

$$\|\partial \boldsymbol{x}_t(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F \leq \underbrace{(1 - \kappa)^{t-1}}_{\text{exponential decrease}} \left\{ \underbrace{(1 + \kappa(t-1))}_{\text{linear increase}} \|\partial \boldsymbol{x}_0(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F + \frac{t}{L} G \right\}.$$

Assuming $G = 0$, the above bound is monotonically decreasing.

If instead we take the worst-case optimal (for minimization) step size $h = 2/(L + \ell)$, then we have

$$\|\partial \boldsymbol{x}_t(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F \leq \underbrace{\left(\frac{1-\kappa}{1+\kappa}\right)^{t-1}}_{\text{exponential decrease}} \left\{ \underbrace{|2t - 1|}_{\text{linear increase}} \|\partial \boldsymbol{x}_0(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F + \frac{2t}{L + \ell} G \right\}.$$

Moreover, assuming $G = 0$, the maximum of the upper bound over t can go up to

$$\|\partial \boldsymbol{x}_t(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F \leq O_{\kappa \to 0}\left(\frac{1}{\kappa}\|\partial \boldsymbol{x}_0(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F\right) \quad at \quad t \approx \frac{1}{\kappa}.$$

In this Corollary, we see a trade-off between the linear convergence rate (exponential in t) and the linear growth in t. When the step size is small ($h = 1/L$), the linear rate is slightly slower than the rate of the larger step size ($h = 2/(\ell + L)$). However, the term in t is *way smaller* for the small step size. This makes a big difference in the convergence: for $h = 1/L$, there is *no* local increase, which is not the case for $h = \frac{2}{\ell + L}$. In the next Corollary we provide a bound on the step size h to guarantee a monotone convergence of the Jacobian.

Corollary 2. *Assuming $G = 0$, the bound of Theorem 2 is monotonically decreasing for $t \geq 1$ if the step size h from Theorem 2 satisfies $0 < h < \sqrt{2}/L$.*

This bound contrasts with the condition on the step size of gradient descent, which is $h \leq 2/L$, with an optimal value of $h = 2/(\ell + L)$ (Nesterov, 2004). This trade-off between asymptotic rate and length of the burn-in phase leads us to formulate:

The curse of unrolling

To ensure convergence of the Jacobian with gradient descent, we must either **1)** accept that the algorithm has a burn-in period proportional to the condition number $1/\kappa$, or **2)** choose a small step size that will *slow down* the algorithm's asymptotic convergence.

3.2 Worst-case Rates for the Chebyshev method

We now derive a convergence-rate analysis for the Chebyshev method, which achieves the best worst-case convergence rate for the minimization of a quadratic function with a bounded spectrum.

Chebyshev method and Chebyshev polynomials. We recall the properties of the Chebyshev method (see e.g. (d'Aspremont et al., 2021, Section 2) for a survey). As mentioned in §2.2.2, the rate of convergence of a first-order method associated with the residual polynomial P_t can be upper bounded by $\max_{\lambda \in [\ell, L]} |P_t(\lambda)|$. Let $\tilde{C}_t$ be the Chebyshev polynomial of the first kind of degree t, and define

$$C_t(\lambda) = \frac{\tilde{C}_t(m(\lambda))}{\tilde{C}_t(m(0))}, \qquad m : [\ell, L] \to [0, 1], \quad m(\lambda) = \frac{2\lambda - L - \ell}{L - \ell}. \tag{9}$$

A known property of Chebyshev polynomials is that the *shifted and normalized* Chebyshev polynomial C_t is the residual polynomial with smallest maximum value in the $[\ell, L]$ interval. This implies that the Chebyshev method, which is the method associated with this polynomial, enjoys the *best* worst-case convergence bound on quadratic functions. Algorithmically speaking, the Chebyshev method reads,

$$\boldsymbol{x}_t(\boldsymbol{\theta}) = \boldsymbol{x}_{t-1}(\boldsymbol{\theta}) - h_t \nabla f(\boldsymbol{x}_{t-1}(\boldsymbol{\theta}), \boldsymbol{\theta}) + m_t(\boldsymbol{x}_{t-1}(\boldsymbol{\theta}) - \boldsymbol{x}_{t-2}(\boldsymbol{\theta})),$$

where h_t is the step size and m_t the momentum. Those parameters are time-varying and depend only on ℓ and L. The following Proposition shows the rate of convergence of the Chebyshev method.

Theorem 3. *Under Assumptions 1,2, let $\xi \stackrel{def}{=} (1 - \sqrt{\kappa})/(1 + \sqrt{\kappa})$, and $\boldsymbol{x}_t(\boldsymbol{\theta})$ denote the t^{th} iterate of the Chebyshev method. Then, we have the following convergence rate*

$$\|\partial \boldsymbol{x}_t(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F \leq \underbrace{\left(\frac{2}{\xi^t + \xi^{-t}}\right)}_{exponential\ decrease} \left\{ \underbrace{\left|\frac{2t^2}{1 - \kappa} - 1\right|}_{quadratic\ increase} \|\partial \boldsymbol{x}_0(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F + \frac{2t^2}{L - \ell} G \right\}.$$

In short, the rate of the Chebyshev algorithm for unrolling in $O(t^2 \xi^t)$. Moreover, assuming $G = 0$, the maximum of the upper bound over t can go up to

$$\|\partial \boldsymbol{x}_t(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F \leq O_{\kappa \to 0} \left(\tfrac{2}{\kappa} \|\partial \boldsymbol{x}_0(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F\right) \quad at \quad t \approx 2\sqrt{\tfrac{1}{\kappa}}.$$

Discussion. Despite being optimal for minimization, the rate of the Chebyshev method has an additional $O(t^2)$ factor. Due to this term, the bound diverges at first, similarly to gradient descent with the optimal step size $h = \frac{2}{\ell + L}$, but sooner. This behavior is visible on Figure **??**.

4 Accelerated Unrolling: How fast can we differentiate through optimization?

We now show to accelerate unrolled differentiation. We first derive a lower bound on the Jacobian suboptimalit and then propose a method based on *Sobolev orthogonal polynomials* (Marcellán et al., 2015), which are extremal polynomials for a norm involving both the polynomial and its derivative.

4.1 Unrolling is at least as hard as optimization

The following Theorem produces a lower bound for the case where the initialization verifies $G = 0$.

Proposition 1. *Let $\boldsymbol{x}_t$ be the t-th iterate of a first-order method. Then, for all iteration t and for all $\boldsymbol{\theta}$, there exists a quadratic function f that verifies Assumption 1 such that $G = 0$, and*

$$\|\partial \boldsymbol{x}_t(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F \geq \frac{2}{\xi^t + \xi^{-t}} \|\partial \boldsymbol{x}_0(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F, \quad \xi = \frac{1 - \sqrt{\kappa}}{1 + \sqrt{\kappa}}. \tag{10}$$

This result tells us that unrolling is *at least* as difficult as optimization. Indeed, the rate in (10) is known to be the lower bound on the accuracy for minimizing smooth and strongly convex function (Nemirovski, 1995). Moreover, although we are not sure if the lower bound is tight for all t, we have that that when $t \to \infty$, the rate of Chebyshev method matches the above rate.

4.2 Average-Case Accelerated Unrolling with Sobolev Polynomials

We now describe an accelerated method for unrolling based on Sobolev polynomials. We first introduce the definition of the Sobolev scalar product for polynomials.

Definition 1. *The Sobolev scalar product (and its norm) for two polynomials P, Q and a density function μ is defined as*

$$\langle P, Q \rangle_\eta \stackrel{def}{=} \int_\mathbb{R} P(\lambda)Q(\lambda)\,\mathrm{d}\mu + \eta \int_\mathbb{R} P'(\lambda)Q'(\lambda)\,\mathrm{d}\mu, \quad \|P\|_\eta^2 \stackrel{def}{=} \langle P, P \rangle_\eta.$$

Although the above product is defined for any distribution $\mu > 0$, in the following we assume μ is the *expected spectral density* associated with the current problem class and discuss in the next section some practical choices. Using this scalar product, we can compute a (loose) upper-bound for $\|\partial \boldsymbol{x}_t(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F$.

Proposition 2. *Assume that $\|\partial \boldsymbol{H}(\boldsymbol{\theta})(\boldsymbol{x}_0(\boldsymbol{\theta}) - \boldsymbol{x}_\star(\boldsymbol{\theta}))\|_F \le \eta\|\partial \boldsymbol{x}_0(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F$. Then, under Assumption 1, 2 and 3, we have the following bound for the average-case rate*

$$\mathbb{E}_{\boldsymbol{H}(\boldsymbol{\theta})}\|\partial \boldsymbol{x}_t(\boldsymbol{\theta}) - \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F^2 \le 2\|P_t\|_\eta^2 \,\mathbb{E}_{\boldsymbol{H}(\boldsymbol{\theta})}\|\partial \boldsymbol{x}_0(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F^2.$$

Using tools from linear algebra, we can build a polynomial that minimizes the bound in Proposition 2.

Proposition 3. *Let $\{S_t\}$ be a sequence of orthogonal Sobolev polynomials, i.e., $\langle S_i,\ S_j \rangle > 0$ if $i = j$ and 0 otherwise, normalized such that $S_i(0) = 1$. Then, the residual polynomial that minimizes the Sobolev norm reads*

$$P_t^\star = \arg\min_{P \in \mathcal{P}_t : P(0)=1} \langle P,\ P \rangle_\eta = \frac{1}{A_t} \sum_{i=0}^{t} a_i S_i, \quad where \quad a_i = \frac{1}{\|S_t\|_\eta^2} \quad and \quad A_t = \sum_{i=0}^{t} a_i.$$

Moreover, we have that $\|P_t^\star\|_\eta^2 = 1/A_t$.

Limited burn-in phase Using the algorithm associated with $P^\star$ with parameters η and μ, we have

$$\mathbb{E}_{\boldsymbol{H}(\boldsymbol{\theta})}\|\partial \boldsymbol{x}_t(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F^2 \le 2 \cdot \mathbb{E}_{\boldsymbol{H}(\boldsymbol{\theta})}\|\partial \boldsymbol{x}_0(\boldsymbol{\theta}) - \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F^2.$$

This inequality follows directly from Proposition 2 and the optimality of P_t for $\|\cdot\|_\eta$: we have that $\|P_t\|_\eta \le \|P_{t-1}\|_\eta$ (because P_{t-1} is a feasible solution for P_t) and $\|P_t\|_\eta \le 1$ (because $P_t = 1$ is a feasible solution for any $t > 1$). That is much better than the maximum bump of $(O(1/\kappa)$ from gradient descend (Theorem 1) and from Chebyshev (Theorem 3).

4.3 Gegenbaueur-Sobolev Algorithm

In most practical scenarios don't have access to the expected spectral density μ. Furthermore, the rates of average-case accelerated algorithms have been shown to be robust with respect to distribution mismatch (Cunha et al., 2022). In these cases, we can approximate the expected spectral density by some distribution that has the same support. A classical choice is the following parametric family indexed by $\alpha \in \mathbb{R}$, which encompasses important distributions such as the density associated with Chebyshev's polynomials or the uniform distribution:

$$\mu(\lambda) = \tilde{\mu}(m(\lambda)), \quad \tilde{\mu}(x) = (1 - x^2)^{\alpha - \frac{1}{2}} \quad \text{and} \quad m : [\ell, L] \to [0, 1], m(\lambda) = \frac{2\lambda - L - \ell}{L - \ell}. \quad (11)$$

We'll call the sequence of Sobolev orthogonal polynomials for this distribution *Gegenbaueur-Sobolev* polynomials. Although in general Sobolev orthogonal polynomials don't enjoy a three-term recurrence as classical orthogonal polynomials do, in this particular case it is possible to build a recurrence for S_t involving only S_{t-2}, Q_t and Q_{t-2}, where $\{Q_t\}$ is sequence of Gegenbaueur polynomials (Marcellán et al., 1994). Unfortunately, existing work on Gegenbaueur and Gegenbaueur-Sobolev polynomials considers $\tilde{\mu}$, but not μ, and also doesn't consider the normalized polynomials $Q_t(0) = 1$ or $S_t(0) = 1$. After (painful) changes to shift and normalize the polynomials, we obtain a three-stages algorithm, summarized in the next Theorem.

Figure 3: (Theoretical) **Worst-case convergence rates** for different algorithms, with $\ell = 0.5$, $L = 10$, $\alpha = 1$ and $G = \|\partial_\theta \nabla f(\boldsymbol{x}(\boldsymbol{\theta}), \boldsymbol{\theta})\| = 0$. The upper bound of a first-order method associated to the sequence of polynomials $\{P_t\}$ is defined as $\max_{\lambda \in [\ell, L]} |P_t(\lambda)|$, where t is the iteration counter. The plot compares the following methods: Gradient descent with large ($\frac{2}{L+\mu}$) and small ($\frac{1}{L}$) step size, Chebyshev, Sobolev (with $\eta = L/\ell$), and their asymptotic variants. We recognize in those curves the peaks of gradient descent and Chebyshev from Figure **??**. We see that the Sobolev algorithm is the best performing method, because its has the shortest burn-in phase, does not locally diverge and has an accelerated asymptotic rate of convergence. Its asymptotic counterpart compromise the local divergence with a simpler implementation, along with depending on fewer hyperparameters.

Theorem 4 (Accelerated Unrolling). *Let $P_t^\star$ be defined in Proposition 3, in the case where the Sobolev product is defined with a Gegenbaueur density function μ (11). Then, the optimization algorithm associated to $P_t^\star$ reads*

$$\boldsymbol{y}_t = \boldsymbol{y}_{t-1} - h_t \nabla f(\boldsymbol{y}_{t-1}, \boldsymbol{\theta}) + m_t(\boldsymbol{y}_{t-1} - \boldsymbol{y}_{t-2}) \tag{12}$$

$$\boldsymbol{z}_t = c_t^{(1)} \boldsymbol{z}_{t-2} + c_t^{(2)} \boldsymbol{y}_t - c_t^{(3)} \boldsymbol{y}_{t-2} \tag{13}$$

$$\boldsymbol{x}_t = \frac{A_{t-1}}{A_t} \boldsymbol{x}_{t-1} + \frac{a_t}{A_t} \boldsymbol{z}_t\,, \tag{14}$$

for some step size h_t, momentum m_t, parameters $c_t^{(1)}$, $c_t^{(2)}$, $c_t^{(3)}$ that depend on α, ℓ, L and η, and A_t, a_t are defined in Proposition 3, whose recurrence are detailed in Appendix D. Moreover, when $t \to \infty$, the asymptotic dynamic simplifies into

$$\boldsymbol{y}_t = \boldsymbol{y}_{t-1} + h \nabla f(\boldsymbol{y}_{t-1}, \boldsymbol{\theta}) + m(\boldsymbol{y}_{t-1} - \boldsymbol{y}_{t-2}) \tag{15}$$

$$\boldsymbol{x}_t = \boldsymbol{y}_t + m(\boldsymbol{x}_{t-1} - \boldsymbol{y}_{t-2})\,, \tag{16}$$

where $m = \left(\frac{1-\sqrt{\kappa}}{1+\sqrt{\kappa}}\right)^2$ and $h = \left(\frac{2}{\sqrt{\ell}+\sqrt{L}}\right)^2$ are the momentum and step size of Polyak's Heavy Ball. Moreover, when $t \to \infty$, we have the same asymptotic linear convergence as the Chebyshev method:

$$\lim \sqrt[t]{\frac{\|\partial \boldsymbol{x}_t(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F}{\|\partial \boldsymbol{x}_0(\boldsymbol{\theta}) - \partial \boldsymbol{x}_\star(\boldsymbol{\theta})\|_F}} \le \sqrt{m}\,.$$

The accelerated algorithm for unrolling is divided into three parts. First, (12) corresponds to a Chebyshev-like algorithm, whose associated polynomials are Gegenbauer polynomials. This result is expected, as Pedregosa et al. (2020) have identified that all average-case accelerated algorithms for optimization take the form of gradient descent with momentum. Second, (13) builds the Sobolev polynomial that corresponds to a weighted average of $\boldsymbol{y}_t$. Finally, (14) is the weighted average of Sobolev polynomials that builds $P^\star$ in Proposition 2.

The non-asymptotic algorithm is rather complicated to implement; see Appendix D. Moreover, it requires a bound on the spectrum of $\boldsymbol{H}(\boldsymbol{\theta})$, namely $[\ell, L]$, and one also has to choose an associated expected spectral density $\mu(\lambda)$ (parametrized by α) and the parameter η. Nevertheless, this is the method that achieves the best performance for problems that satisfy our assumptions.

Surprisingly, the *asymptotic* version is extremely simple, as it corresponds to *a weighed average of Heavy-Ball iterates*: the only required parameters are ℓ and L. It means that asymptotically, the algorithm is *universally* optimal, i.e., it achieves the best performing rate as long as we can identify

Figure 4: **Empirical comparison** of the Sobolev method introduced in §4.3 (with $\alpha = 1$ and $\eta = 1$), the Chebyshev method and Gradient descent on 3 different datasets. The Sobolev method achieves the same asymptotic rate as the Chebyshev method without the initial loss of accuracy.

the bounds on the spectrum of $H(\theta)$. Such universal properties have been identified previously in (Scieur et al., 2020b), who showed that all average-case optimal algorithms converge to Polyak's momentum independently of the expected spectral density μ (up to mild assumptions). We have the same phenomenon here, but with the additional surprising (and counter-intuitive) feature that the asymptotic algorithm is also *independent of η*.

5 Experiments and Discussion

We compare multiple algorithms for estimating the Jacobian (OPT) of the solution of a ridge regression problem (Example (1)) for a fixed value of $\theta = 10^{-3}$. Figure 1 shows the objective and Jacobian suboptimality on a ridge regression problem with the breast-cancer[2] as underlying dataset. Figure 5 shows the Jacobian suboptimality as a function of the number of iterations, on both the breast-cancer and bodyfat[3] dataset, and for a synthetic dataset (where $H(\theta)$ is generated as $A^\top A$, where each entry in A is generated from a standard Gaussian distribution). We perform this comparison across 3 different algorithms (Gradient descent, Chebyshev, and Sobolev) solving Problem OPT, all tuned with their optimal worst-case parameter (see Appendix B for further details).

We observe the early suboptimality increase of Gradient descent and Chebyshev algorithm as predicted by Theorem 2 and Theorem 3. Compared to Figure 3, that showed the theoretical rates, we see that there's a remarkable agreement between theory and practice, as both the early increase, the asymptotic rate and the ordering of the methods matches the theoretical prediction. We also see that Sobolev is the best performing algorithm in practice, as it avoids the early increase while matching the accelerated asymptotic rate of Chebyshev.

Limitations. Our theoretical results are limited to first-order methods applied to quadratic functions. Many applications use first-order methods, the quadratic Assumption 1, as well as the commutativity Assumption 2, are somewhat restrictive. We believe that the observations made in Theorem 2 and Theorem 3 hold, i.e., that we can decompose the rate of convergence for unrolling by a burn-in phase, followed by an asymptotic linear convergence. The Sobolev algorithm developed in this paper, however, might not generalize well outside the scope of quadratics. Nevertheless, the development of this accelerated method for unrolling highlights that we can adapt the design of current optimization algorithms so that they might perform better for automatic differentiation. A potential extension to non-quadratics would be non-trivial, as the polynomial formalism does apply in this setting and would require an entirely different proof technique.

---

[2]https://archive.ics.uci.edu/ml/datasets/Breast+Cancer+Wisconsin+(Diagnostic)
[3]http://lib.stat.cmu.edu/datasets/

References

Ablin, Pierre, Gabriel Peyré, and Thomas Moreau (2020). "Super-efficiency of automatic differentiation for functions defined as a minimum". In: *Proceedings of the 37 th International Conference on Machine Learning*.

Agarwal, Naman, Surbhi Goel, and Cyril Zhang (2021). "Acceleration via Fractal Learning Rate Schedules". In: *Proceedings of the 38th International Conference on Machine Learning*. Ed. by Marina Meila and Tong Zhang. Proceedings of Machine Learning Research. PMLR.

Amos, Brandon and J Zico Kolter (2017). "Optnet: Differentiable optimization as a layer in neural networks". In: *International Conference on Machine Learning*.

Bai, Shaojie, J Zico Kolter, and Vladlen Koltun (2019). "Deep equilibrium models". In: *Advances in Neural Information Processing Systems*.

Bengio, Yoshua (2000). "Gradient-based optimization of hyperparameters". In: *Neural computation*.

Bertrand, Quentin et al. (2020). "Implicit differentiation of Lasso-type models for hyperparameter optimization". In: *International Conference on Machine Learning*. PMLR.

Blondel, Mathieu et al. (2021). "Efficient and Modular Implicit Differentiation". In: *arXiv preprint arXiv:2105.15183*.

Cunha, Leonardo et al. (2022). "Only tails matter: Average-Case Universality and Robustness in the Convex Regime". In.

d'Aspremont, Alexandre, Damien Scieur, and Adrien Taylor (2021). "Acceleration methods". In: *arXiv preprint arXiv:2101.09545*.

Deledalle, Charles-Alban et al. (2014). "Stein Unbiased GrAdient estimator of the Risk (SUGAR) for multiple parameter selection". In: *SIAM Journal on Imaging Sciences*.

Domke, Justin (2012). "Generic methods for optimization-based modeling". In: *Artificial Intelligence and Statistics*. PMLR.

Du, Simon S et al. (2017). "Stochastic variance reduction methods for valuation". In: *International Conference on Machine Learning*. PMLR.

Duvenaud, David, J. Zico Kolter, and Matthew Johnson (2020). "Deep Implicit Layers Tutorial - Neural ODEs, Deep Equilibirum Models, and Beyond". In: *Neural Information Processing Systems Tutorial*.

Franceschi, Luca et al. (2017). "Forward and reverse gradient-based hyperparameter optimization". In: *International Conference on Machine Learning*. PMLR.

Franceschi, Luca et al. (2018). "Forwardgramming for hyperparameter optimization and meta-learning". In: *International Conference on Machine Learning*. PMLR.

Kim, Yoon et al. (2017). "Structured attention networks". In: *arXiv preprint arXiv:1702.00887*.

Lorraine, Jonathan, Paul Vicol, and David Duvenaud (2020). "Optimizing millions of hyperparameters by implicit differentiation". In: *International Conference on Artificial Intelligence and Statistics*. PMLR.

Marcellán, Francisco, Teresa E Pérez, and Miguel A Piñar (1994). "Gegenbauer-Sobolev orthogonal polynomials". In: *Nonlinear Numerical Methods and Rational Approximation II*. Springer.

Marcellán, Francisco and Yuan Xu (2015). "On Sobolev orthogonal polynomials". In: *Expositiones Mathematicae*.

Metz, Luke et al. (2016). "Unrolled generative adversarial networks". In: *arXiv preprint arXiv:1611.02163*.

Nemirovski, Arkadi (1995). "Information-based complexity of convex programming". In: *Lecture Notes*.

Nesterov, Yurii (2004). *Introductory lectures on convex optimization*. Springer.

Paquette, Courtney et al. (2022). "Unro is predictable for large models: A universality property and average-case analysis". In: *Foundations of Computational Mathematics*.

Pedregosa, Fabian (2016). "Hyperparameter optimization with approximate gradient". In: *International Conference on Machine Learning*. PMLR.

Pedregosa, Fabian and Damien Scieur (2020). "Average-case Acceleration Through Spectral Density Estimation". In: *Proceedings of the 37th International Conference on Machine Learning (ICML)*.

Pfau, David and Oriol Vinyals (2016). "Connecting generative adversarial networks and actor-critic methods". In: *arXiv preprint arXiv:1610.01945*.

Rajeswaran, Aravind et al. (2019). "Meta-learning with implicit gradients". In: *Advances in Neural Information Processing Systems*.

Scieur, Damien, Alexandre d'Aspremont, and Francis Bach (2020a). "SATnlinear acceleration". In: *Mathematical Programming*.

Scieur, Damien and Fabian Pedregosa (2020b). "Rage-Case Optimality of Polyak Momentum". In: *Proceedings of the 37th International Conference on Machine Learning (ICML)*.

Shaban, Amirreza et al. (2019). "TrMiack-propagation for bilevel optimization". In: *The 22nd International Conference on Artificial Intelligence and Statistics*. PMLR.

Wang, Po-Wei et al. (2019). "Satnet: Bridging deep learning and logical reasoning using a differentiable satisfiability solver". In: *International Conference on Machine Learning*. PMLR.

Wengert, Robert Edwin (1964). "A simple automatic derivative evaluation program". In: *Communications of the ACM*.