FERMI: Fair Empirical Risk Minimization Via Exponential Rényi Mutual Information

Abstract

Several notions of fairness, such as demographic parity and equal opportunity, are defined based on statistical independence between a predicted target and a sensitive attribute. In machine learning applications, however, the data distribution is unknown to the learner and statistical independence is not verifiable. Hence, the learner could only resort to empirical evaluation of the degree of fairness violation. Many fairness violation notions are defined as a divergence/distance between the joint distribution of the target and sensitive attributes and the Kronecker product of their marginals, such as Rényi correlation, mutual information, L_1 distance, to name a few. In this paper, we propose another notion of fairness violation, called Exponential Rényi Mutual Information (ERMI) between sensitive attributes and the predicted target. We show that ERMI is a strong fairness violation notion in the sense that it provides an upper bound guarantee on all of the aforementioned notions of fairness violation. We also propose the Fair Empirical Risk Minimization via ERMI regularization framework, called FERMI. We provide scalable batch and stochastic optimization methods to solve FERMI with theoretical convergence guarantees. Our experiments show that FERMI achieves the most favorable tradeoffs between fairness violation and performance on test data across different problem setups, even when fairness violation is measured in notions other than ERMI.

1 Introduction

Ensuring that decisions made using machine learning algorithms are fair to different subgroups is of utmost importance. Without any mitigation strategy, machine learning algorithms may result in discrimination against certain subgroups based on sensitive attributes, such as gender or race, even if such discrimination is absent in the training data (Datta et al., 2015; Sweeney, 2013; Bolukbasi et al., 2016; Angwin et al., 2016; du Pin Calmon et al., 2017b; Feldman et al., 2015; Hardt et al., 2016; Fish et al., 2016; Woodworth et al., 2017; Zafar et al., 2017; Bechavod & Ligett, 2017; Kearns et al., 2018). To remedy such discrimination issues, several notions for imposing algorithmic fairness have been proposed in the literature.

A learning machine satisfies the *demographic parity* notion, if the predicted target is independent of the sensitive attributes (Dwork et al., 2012). Promoting demographic parity can lead to poor performance, especially if the true outcome is not independent of the sensitive attributes. To remedy this, (Hardt et al., 2016) proposed *equalized odds* to ensure that the predicted target is conditionally independent of the sensitive attributes given the true label. A further relaxed version of this notion is *equal opportunity* which is satisfied if predicted target is conditionally independent of sensitive attributes given that the true label is in an advantaged class (Hardt et al., 2016). *Note that the inherent assumption in such conditional notions is that the true labels are unbiased. These notions suffer from a potential amplification of the inherent biases that may exist in the targets/labels in the training data (e.g., data collection bias). Tackling such bias is beyond the scope of this work.*

In practice, the learner cannot empirically verify independence of random variables, and hence cannot verify demographic parity, equalized odds, or equal opportunity. This has led the machine learning community to define several notions of fairness violation that quantify the degree of independence between random variables, e.g., demographic parity/equalized odds, L_1 distance (Dwork et al., 2012; Hardt et al., 2016), mutual information (Kamishima et al., 2011; Rezaei et al., 2020; Steinberg et al., 2020), Pearson correlation (Zafar et al., 2017), false positive/negative rates (Bechavod & Ligett, 2017), Hilbert Schmidt independence criterion (HSIC) (Pérez-Suay et al.,

2017), and Rényi correlation (Baharlouei et al., 2020; Grari et al., 2020; 2019), to name a few. In this paper, we define yet another notion of fairness violation, called exponential Rényi mutual information (ERMI). We show that ERMI is easy to compute empirically and prove that ERMI provides an upper bound on the existing notions of fairness violation such as demographic parity, equalized odds, and equal opportunity.

Given a notion of fairness violation, it is still not straightforward to train an algorithm that satisfies a fairness violation constraint (Cotter et al., 2019). The fairness-promoting machine learning algorithms can be categorized in three main classes: *pre-processing*, *post-processing*, and *in-processing* methods. Pre-processing algorithms (Feldman et al., 2015; Zemel et al., 2013; du Pin Calmon et al., 2017b) transform the biased data features to a new space in which the labels and sensitive attributes are statistically independent. This transform is oblivious to the training procedure. Post-processing approaches (Hardt et al., 2016; Pleiss et al., 2017) mitigate the discrimination of the classifier by altering the the final decision, e.g., by changing the thresholds on soft labels, or reassigning the labels to impose notions of fairness. In-processing approaches focus on the training procedure and impose the notions of fairness as constraints or regularization terms in the optimization procedure. Several regularization-based methods are proposed in the literature to impose measures of fairness to decision-trees (Kamiran et al., 2010; Raff et al., 2018; Aghaei et al., 2019), support vector machines (Donini et al., 2018), neural networks (Grari et al., 2020), or (logistic) regression models (Zafar et al., 2017; Berk et al., 2017; Taskesen et al., 2020; Chzhen & Schreuder, 2020; Baharlouei et al., 2020; Jiang et al., 2020; Grari et al., 2019).

In this paper, we introduce a fair empirical risk minimization framework via ERMI regularization, and call it FERMI. We provide batch and stochastic gradient-based methods with guarantees for solving FERMI and demonstrate their effectiveness on multiple numerical experiments. We show that FERMI can be used to achieve more favorable tradeoffs between performance and fairness, even if fairness violation is measured in notions other than ERMI.

2 $(Z, \mathcal{Z})$-FAIRNESS: A GENERAL NOTION OF FAIRNESS

We consider a learner who trains a model to predict a target, $\widehat{Y}$, e.g., whether or not to extend a loan, supported on $\mathcal{Y}$ which can be discrete or continuous. The prediction is made using a set of features, $\mathbf{X}$, e.g., financial history features, length of credit, and amount of debt. We also assume that there is a set of discrete sensitive attributes, S, e.g., race and sex, supported on $\mathcal{S}$, associated with each sample. Further, let $\mathcal{A} \subseteq \mathcal{Y}$ denote an advantaged outcome class, e.g., the outcome where a loan is extended. Next, we will present the main fairness notion considered in this paper, which generalizes several existing ones.

Definition 1 (($Z, \mathcal{Z}$)-fairness). *Given a random variable Z, let $\mathcal{Z}$ be a subset of values that Z can take. We say that a learning machine satisfies $(Z, \mathcal{Z})$-fairness if for every $z \in \mathcal{Z}$, $\widehat{Y}$ is conditionally independent of S given $Z = z$. More precisely,*

$$p_{\widehat{Y}, S | Z}(\widehat{y}, s | z) = p_{\widehat{Y} | Z}(\widehat{y} | z) p_{S | Z}(s | z) \quad \forall \widehat{y} \in \mathcal{Y}, s \in \mathcal{S}, z \in \mathcal{Z}. \tag{1}$$

Lemma 1. *$(Z, \mathcal{Z})$-fairness recovers these important special cases:*

1. *$(Z, \mathcal{Z})$-fairness recovers demographic parity (Dwork et al., 2012) if $Z = 0$ and $\mathcal{Z} = \{0\}$.*

2. *$(Z, \mathcal{Z})$-fairness recovers equalized odds (Hardt et al., 2016) if $Z = Y$ and $\mathcal{Z} = \mathcal{Y}$.*

3. *$(Z, \mathcal{Z})$-fairness recovers equal opportunity (Hardt et al., 2016) if $Z = Y$ and $\mathcal{Z} = \mathcal{A}$.*

Refer to Appendix A for the proof and the definition of the existing notions of fairness. Demographic parity amounts to requiring equality of outcomes across sensitive groups. However, it can result in poor performance of a learned model, particularly if the true outcome Y is not independent of S. Equalized odds and equal opportunity aim to remedy this issue (Hardt et al., 2016). In the binary classification setting with binary sensitive attributes (e.g., male or female), equalized odds ensures equality of false negatives and false positives of a classifier across sensitive groups. Equal opportunity is a relaxation of the equalized odds criterion, which in the binary setting requires just equality of false negatives across sensitive groups. For example, this could be used to enforce that a face recognition software does not falsely classify people of one race as criminals more often than people of other races.

Note that verifying $(Z, \mathcal{Z})$-fairness requires having access to the joint distribution of random variables $(Z, \widehat{Y}, S)$. This joint distribution is unavailable to the learner in the context of machine learning, and hence the learner would resort to empirical estimation of the amount of violation of independence. In the next section, we propose exponential Rényi mutual information as a notion of the violation of $(Z, \mathcal{Z})$-fairness, and show that it is a stronger notion compared to several existing fairness violation notions, including demographic parity L_1 distance (Kearns et al., 2018), equalized odds L_1 distance (Hardt et al., 2016), and equal opportunity L_1 distance (Hardt et al., 2016).

3 EXPONENTIAL RÉNYI MUTUAL INFORMATION

In this section, we define ERMI and show that several existing fairness violation notions (which are mostly instances of f-divergences or distance metrics between the joint probability distribution and the Kronecker product of the marginals) are also upper bounded by ERMI, implying that ERMI is a stronger notion of fairness violation. In particular, this means that if ERMI is small, then we automatically obtain guarantees that all other notions of fairness violation must be small as well. We present all definitions and results for general $(Z, \mathcal{Z})$ fairness notion, which requires careful extension of several existing notions to the conditional case. These definitions and results will simplify significantly when $Z = 0$ and $\mathcal{Z} = \{0\}$, which will eliminate all conditional expectations.

Definition 2 (ERMI – exponential Rényi mutual information). *We define the exponential Rényi mutual information between $\widehat{Y}$ and S given $Z \in \mathcal{Z}$ as*

$$D_R(\widehat{Y}; S | Z \in \mathcal{Z}) := \mathbb{E}_{Z, \widehat{Y}, S} \left\{ \frac{p_{\widehat{Y}, S | Z}(\widehat{Y}, S | Z)}{p_{\widehat{Y} | Z}(\widehat{Y} | Z) p_{S | Z}(S | Z)} \middle| Z \in \mathcal{Z} \right\} - 1. \tag{2}$$

As we discuss in Appendix B, ERMI is an f-divergence between the joint distribution $p_{\widehat{Y}, S | Z}$ and the Kronecker product of marginals $p_{\widehat{Y} | Z} \otimes p_{S | Z}$. In particular, ERMI is non-negative, and zero if and only if $(Z, \mathcal{Z})$-fairness is satisfied. Hence, ERMI is a valid notion of fairness violation.

Definition 3 (Rényi mutual information (Rényi, 1961)). *Let the Rényi mutual information of order $\alpha > 1$ between random variables $\widehat{Y}$ and S given $Z \in \mathcal{Z}$ be defined as:*

$$I_\alpha(\widehat{Y}; S | Z \in \mathcal{Z}) := \frac{1}{\alpha - 1} \log \left(\mathbb{E}_{Z, \widehat{Y}, S} \left\{ \left(\frac{p_{\widehat{Y}, S | Z}(\widehat{Y}, S | Z)}{p_{\widehat{Y} | Z}(\widehat{Y} | Z) p_{S | Z}(S | Z)} \right)^{\alpha - 1} \middle| Z \in \mathcal{Z} \right\} \right), \tag{3}$$

which generalizes Shannon mutual information

$$I_1(\widehat{Y}; S | Z \in \mathcal{Z}) := \mathbb{E}_{Z, \widehat{Y}, S} \left\{ \log \left(\frac{p_{\widehat{Y}, S | Z}(\widehat{Y}, S | Z)}{p_{\widehat{Y} | Z}(\widehat{Y} | Z) p_{S | Z}(S | Z)} \right) \middle| Z \in \mathcal{Z} \right\}, \tag{4}$$

and recovers it as $\lim_{\alpha \to 1^+} I_\alpha(\widehat{Y}; S | Z \in \mathcal{Z}) = I_1(\widehat{Y}; S | Z \in \mathcal{Z})$.

Note that $I_\alpha(\widehat{Y}; S | Z \in \mathcal{Z}) \geq 0$ with equality if and only if $(Z, \mathcal{Z})$-fairness is satisfied.

Theorem 1 (ERMI is stronger than Shannon mutual information). *We have*

$$0 \leq I_1(\widehat{Y}; S | Z \in \mathcal{Z}) \leq I_2(\widehat{Y}; S | Z \in \mathcal{Z}) \leq e^{I_2(\widehat{Y}; S | Z \in \mathcal{Z})} - 1 = D_R(\widehat{Y}; S | Z \in \mathcal{Z}). \tag{5}$$

All proofs are relegated to the appendix. Theorem 1 establishes that ERMI is a stronger notion of fairness in the sense that driving it to zero would also bound the Shannon mutual information. It also shows that ERMI is exponentially related to the Rényi mutual information of order 2.

Definition 4 (Rényi correlation (Hirschfeld, 1935; Gebelein, 1941; Rényi, 1959)). *Let $\mathcal{F}$ and $\mathcal{G}$ be the set of measurable functions such that for random variables $\widehat{Y}$ and S, $\mathbb{E}_{\widehat{Y}} \{ f(\widehat{Y}; z) \} = \mathbb{E}_S \{ g(S; z) \} = 0$, $\mathbb{E}_{\widehat{Y}} \{ f(\widehat{Y}; z)^2 \} = \mathbb{E}_S \{ g(S; z)^2 \} = 1$, for all $z \in \mathcal{Z}$. Rényi correlation is:*

$$\rho_R(\widehat{Y}, S | Z \in \mathcal{Z}) := \sup_{f \in \mathcal{F}, g \in \mathcal{G}} \mathbb{E}_{Z, \widehat{Y}, S} \left\{ f(\widehat{Y}; Z) g(S; Z) \middle| Z \in \mathcal{Z} \right\}. \tag{6}$$

Rényi correlation generalizes Pearson correlation coefficient

$$\rho(\widehat{Y}, S | Z \in \mathcal{Z}) := \mathbb{E}_Z \left\{ \frac{\mathbb{E}_{\widehat{Y}, S} \{ \widehat{Y} S | Z \}}{\sqrt{\mathbb{E}_{\widehat{Y}} \{ \widehat{Y}^2 | Z \} \mathbb{E}_S \{ S^2 | Z \}}} \middle| Z \in \mathcal{Z} \right\} \tag{7}$$

to capture nonlinear dependencies between the random variables by finding functions of random variables that maximize the Pearson correlation coefficient between the random variables. In fact, it is true that $\rho_R(\widehat{Y}, S | Z \in \mathcal{Z}) \geq 0$ with equality if and only if $(Z, \mathcal{Z})$-fairness is satisfied. Due to these favorable properties, Rényi correlation has gained popularity as a measure of fairness violation (Baharlouei et al., 2020; Grari et al., 2020).

Theorem 2 (ERMI is stronger than Rényi correlation.). *We have*

$$0 \leq |\rho(\widehat{Y}, S | Z \in \mathcal{Z})| \leq \rho_R(\widehat{Y}, S | Z \in \mathcal{Z}) \leq D_R(\widehat{Y}; S | Z \in \mathcal{Z}), \tag{8}$$

and if $|\mathcal{S}| = 2$, $D_R(\widehat{Y}; S | Z \in \mathcal{Z}) = \rho_R(\widehat{Y}, S | Z \in \mathcal{Z})$.

Next, we turn to another popular notion of fairness violation and establish similar relationships.

Definition 5 (L_q fairness violation). *We define the L_q fairness violation for $q \geq 1$ by:*

$$L_q(\widehat{Y}, S | Z \in \mathcal{Z}) := \mathbb{E}_Z \left\{ \left(\int_{\widehat{y} \in \mathcal{Y}_0} \sum_{s \in \mathcal{S}_0} \left| p_{\widehat{Y}, S | Z}(\widehat{y}, s | Z) - p_{\widehat{Y} | Z}(\widehat{y} | Z) p_{S | Z}(s | Z) \right|^q dy \right)^{\frac{1}{q}} \middle| Z \in \mathcal{Z} \right\}. \tag{9}$$

Note that $L_q(\widehat{Y}, S | Z \in \mathcal{Z}) = 0$ if and only if $(Z, \mathcal{Z})$-fairness is satisfied. In particular, L_1 fairness violation recovers the demographic parity violation (Kearns et al., 2018, Definition 2.1) if we let $\mathcal{Z} = \{0\}$ and $Z = 0$. It also recovers equal opportunity violation (Hardt et al., 2016) if we let $\mathcal{Z} = \mathcal{A}$ and $Z = Y$. L_q fairness violation generalizes this notion by considering the L_q norm of the difference between the joint distribution $p_{\widehat{Y}, S}$ and the Kronecker product of the marginal distributions $p_{\widehat{Y}} \otimes p_S$.

Theorem 3 (ERMI is stronger than L_1 fairness violation). *Let $\widehat{Y}$ be a discrete or continuous random variable, and S be a discrete random variable supported on a finite set. Then,*[1]

$$0 \leq L_1(\widehat{Y}, S | Z \in \mathcal{Z}) \leq L_2(\widehat{Y}, S | Z \in \mathcal{Z}) \leq \sqrt{D_R(\widehat{Y}, S | Z \in \mathcal{Z})}. \tag{10}$$

The above theorem says that if a method controls ERMI value for imposing fairness, L_1 demographic parity violation (Kearns et al., 2018), L_1 equal opportunity violation (Hardt et al., 2016), or L_1 equalized odds violation (Hardt et al., 2016) is also guaranteed to be bounded.

4 FAIR RISK MINIMIZATION VIA ERMI

Our goal is to train a model that balances fairness and accuracy objectives. To this end, we introduce *fair risk minimization through exponential Rényi mutual information* framework defined below:[2]

Definition 6 (FRMI – fair risk minimization through exponential Rényi mutual information). *To balance fairness and accuracy, we consider the learning objective function:*

$$\min_{\boldsymbol{\theta}} \quad \mathbb{E}_{\mathbf{X}, Y, S} \left\{ \ell(\mathbf{X}, Y; \boldsymbol{\theta}) \right\} + \lambda D_R(\widehat{Y}_{\boldsymbol{\theta}}; S), \tag{11}$$

where ℓ denotes the loss function, such as L_2 loss or cross entropy loss; $\lambda > 0$ is a scalar balancing the accuracy versus fairness objectives; and $\widehat{Y}_{\boldsymbol{\theta}}$ is the output of the learned model (e.g., the output of a classification or a regression task, or the cluster number in a clustering task).

In the rest of this paper, we often leave the dependence of $\widehat{Y}$ on $\boldsymbol{\theta}$ implicit for brevity of notation. Notice that we have also left the dependence of the loss on the predicted outcome $\widehat{Y}$ implicit. FRMI is the objective we should solve if demographic parity is the desired fairness notion; if instead we are interested in equalized odds or equal opportunity, then $D_R(\widehat{Y}, S)$ should be replaced by $D_R(\widehat{Y}, S | Z \in \mathcal{Z})$ for an appropriate $(Z, \mathcal{Z})$ pair, per the discussion in Section 3. Since the theory is very similar in both cases, we stick with FRMI as defined in Definition 6. In practice, the true

---

[1]Note that a similar relationship with TV norm could be established as well.

[2]In this section, we present all results in the context of $\mathcal{Z} = \{0\}$ amd $Z = 0$, leaving off all conditional expectations for clarity of presentation. The results could be generalized for general $(Z, \mathcal{Z})$, as we have used the resulting algorithms for empirical experiments.

joint distribution of $(\mathbf{X}, S, Y, \widehat{Y})$ is unknown and we only have N samples at our disposal, making it impossible to solve FRMI. Hence, we turn into fair empirical risk minimization via exponential Rényi mutual information (FERMI) approach. While it is natural to estimate $\mathbb{E}_{\mathbf{X},Y,S} \left\{ \ell(\mathbf{X}, Y; \boldsymbol{\theta}) \right\}$ through the empirical risk, the estimation of $D_R(\widehat{Y}, S)$ in the objective function in Eq. (11) is not as straightforward. In what follows, we propose two approaches for estimating $D_R(\widehat{Y}, S)$. These two approaches result in two different algorithms for balancing fairness and accuracy, where we discuss the benefits and shortcomings of each.

4.1 FERMI VIA EMPIRICAL ESTIMATION OF THE PROBABILITY DISTRIBUTIONS

Let $\{\mathbf{x}_i, s_i, y_i, \widehat{y}_i\}_{i \in [N]}$ denote the features, sensitive attributes, targets, and the predictions of the model parameterized by $\boldsymbol{\theta}$ for samples $i \in [N]$. A natural approach to estimate the objective function in Eq. (11) and learning the parameter $\boldsymbol{\theta}$ is through solving the problem

$$\min_{\boldsymbol{\theta}} \left\{ \frac{1}{N} \sum_{i \in [N]} \ell(\mathbf{x}_i, y_i; \boldsymbol{\theta}) + \lambda \widehat{D}_R(\widehat{Y}_{\boldsymbol{\theta}}; S) \right\}, \quad (12)$$

Here $\widehat{D}_R(\widehat{Y}; S) := \sum_{s \in \mathcal{S}} \int_{\widehat{y} \in \mathcal{Y}} \frac{\widehat{p}_{\widehat{Y},S}(\widehat{y},s)^2}{\widehat{p}_{\widehat{Y}}(\widehat{y})\widehat{p}_S(s)} d\widehat{y} - 1$ is an empirical estimate of $D_R(\widehat{Y}, S)$ where $\widehat{p}_S(s)$, $\widehat{p}_{\widehat{Y}}(\widehat{y})$, and $\widehat{p}_{\widehat{Y},S}(\widehat{y}, s)$ are the empirical estimation of the corresponding probability functions. To make the objective function differentiable with respect to $\boldsymbol{\theta}$, we make the following assumption:

Assumption 1. *Assume the following soft conditional density to make $\widehat{D}_R$ differentiable w.r.t. $\boldsymbol{\theta}$:*

$$p_{\widehat{Y}|\mathbf{X}}(\widehat{y}|\mathbf{x}) = \frac{e^{-\tau \ell(\mathbf{X}, \widehat{y}; \boldsymbol{\theta})}}{\int_{y \in \mathcal{Y}} e^{-\tau \ell(\mathbf{X}, y; \boldsymbol{\theta})} dy}, \quad (13)$$

$$p_{\widehat{Y}}(\widehat{y}) = \mathbb{E}_{\mathbf{X}} \left\{ p_{\widehat{Y}|\mathbf{X}}(\widehat{y}|\mathbf{X}) \right\}, \quad (14)$$

$$p_{\widehat{Y}|S}(\widehat{y}|s) = \frac{\mathbb{E}_{\mathbf{X}} \left\{ p_{\widehat{Y}|\mathbf{X}}(\widehat{y}|\mathbf{x}) \mathbb{1}\{S = s\} \right\}}{\mathbb{E}_{\mathbf{X}} \left\{ \mathbb{1}\{S = s\} \right\}}, \quad (15)$$

where $\tau > 0$ controls the softness of the decision and $\tau \to \infty$ would recover the hard decision made by choosing y that minimizes the loss function.

Under this assumption, we will have the following empirical estimates of the distributions of the sensitive attribute and the predicted target:

$$\widehat{p}_S(s) := \frac{1}{N} \sum_{i \in [N]} \mathbb{1}\{s_i = s\}, \qquad \widehat{p}_{\widehat{Y}|S}(\widehat{y}|s) := \frac{\sum_{i \in [N]} \mathbb{1}\{s_i = s\} e^{-\tau \ell(\mathbf{x}_i, \widehat{y}; \boldsymbol{\theta})}}{\sum_{i \in [N]} \mathbb{1}\{s_i = s\} \int_{y \in \mathcal{Y}} e^{-\tau \ell(\mathbf{x}_i, y; \boldsymbol{\theta})} dy}, \quad (16)$$

which could be marginalized to define $\widehat{p}_{\widehat{Y}}(\cdot)$ as well. To solve Eq. (12), one can apply the gradient descent algorithm and use the dynamics

$$\boldsymbol{\theta}^{t+1} = \boldsymbol{\theta}^t - \eta \nabla_{\boldsymbol{\theta}} \left\{ \frac{1}{N} \sum_{i \in [N]} \ell(\mathbf{x}_i, y_i; \boldsymbol{\theta}^t) + \lambda \widehat{D}_R(\widehat{Y}_{\boldsymbol{\theta}^t}; S) \right\}, \quad (17)$$

where $\eta > 0$ is the learning rate/step-size. Under standard assumptions on the loss function and the learning rate, one can show that the dynamics in Eq. (17) find an $\epsilon-$stationary point, (i.e., a point with the norm of gradient being smaller than ϵ) in $O(\frac{1}{\epsilon^2})$ iterations (Nesterov, 2013).

Theorem 4. *(Informal statement) Gradient descent (i.e. Eq. (17)) converges to the set of ϵ-first order stationary points of the FERMI objective (Eq. (12)) in $O(\frac{1}{\epsilon^2})$ iterations (gradient evaluations).*

While this algorithm achieves the optimal rate of first-order methods for general smooth non-convex optimization problems, the empirical ERMI term in the objective in Eq. (12) is a biased estimator of ERMI in Eq. (11). This bias makes the optimization problem in Eq. (12) not suitable for using stochastic methods. For example, in the extreme case, where only one sample is available to the learner for updating the objective at each turn, $\widehat{D}_R(\cdot; \cdot)$ can be severely biased due to the nonlinearities in how it is defined. In the next subsection, we propose another approach for estimation of $D_R(\cdot; \cdot)$ that results in an unbiased estimator, which is amenable to stochastic optimization.

4.2 STOCHASTIC FERMI

In this subsection, we assume $\widehat{Y}$ is discrete and supported on a finite set $\mathcal{Y}$. The main challenge in solving Eq. (12) via stochastic optimization methods is obtaining an unbiased estimate of the gradient of $\widehat{D}_R(\widehat{Y}; S)$ through samples. To overcome this obstacle, we rely on the following key theorem:

Theorem 5. *In the discrete classification case, where $\widehat{Y} \in [m], S \in [k]$, we have*

$$D_R(\widehat{Y}; S) = \max_{W \in \mathbb{R}^{k \times m}} \left\{ -\operatorname{Tr}(W P_{\widehat{y}} W^T) + 2\operatorname{Tr}(W P_{\widehat{y}, s} P_s^{-1/2}) - 1 \right\}, \tag{18}$$

where $P_{\widehat{y}} = \begin{pmatrix} \widehat{p}_{\widehat{Y}}(1) & & 0 \\ & \ddots & \\ 0 & & \widehat{p}_{\widehat{Y}}(m) \end{pmatrix}$,

$$P_{\widehat{y}, s} = \begin{pmatrix} \widehat{p}_{\widehat{Y}, S}(1, 1) & \cdots & \widehat{p}_{\widehat{Y}, S}(1, k) \\ \vdots & \ddots & \vdots \\ \widehat{p}_{\widehat{Y}, S}(m, 1) & \cdots & \widehat{p}_{\widehat{Y}, S}(m, k) \end{pmatrix}, P_s = \begin{pmatrix} \widehat{p}_S(1) & & 0 \\ & \ddots & \\ 0 & & \widehat{p}_S(k) \end{pmatrix}.$$

The above theorem implies that Eq. (12) can be equivalently written as

$$\min_{\boldsymbol{\theta} \in \Theta} \max_{W \in \mathbb{R}^{k \times m}} \left\{ \frac{1}{N} \sum_{i \in [N]} \ell(\mathbf{x}_i, y_i; \boldsymbol{\theta}) + \lambda \left(-\operatorname{Tr}(W P_{\widehat{y}} W^T) + 2\operatorname{Tr}(W P_{\widehat{y}, s} P_s^{-1/2}) - 1 \right) \right\}. \tag{19}$$

This optimization problem can be solved through Algorithm 1. The convergence rate of this algorithm is analyzed in Theorem 6.

Algorithm 1 Two-Time Scale SGDA for FERMI

1: **Input**: $\boldsymbol{\theta}^0 \in \mathbb{R}^{d_\theta}$, $W^0 \in \mathcal{W} \subset \mathbb{R}^{k \times m}$, step-sizes (η_θ, η_w), mini-batch size M, fairness parameter $\lambda \geq 0$, iteration number T.
2: **for** $t = 0, 1, \dots, T$ **do**
3:     Draw M i.i.d. samples $\{\mathbf{z}_n\}_{n=1}^M$
4:     Set $\boldsymbol{\theta}^{t+1} \leftarrow \boldsymbol{\theta}^t - \eta_\theta \Big[\frac{1}{M} \sum_{n=1}^M \nabla_\theta \ell(\mathbf{x}_n, y_n, \boldsymbol{\theta}) - 2\lambda \nabla_\theta \operatorname{vec}(P_{\widehat{y}})^T \operatorname{vec}(W^T W) +$
    $2\lambda \nabla_\theta \operatorname{vec}(P_{\widehat{y}, s})^T \operatorname{vec}(W^T P_s^{-1/2}) \Big]$
5:     Set $W^{t+1} \leftarrow \Pi_{\mathcal{W}} \left(W^t + \eta_w \left[\frac{1}{M} \sum_{n=1}^M -2\lambda W P_{\widehat{y}} + 2\lambda P_s^{-1/2} P_{\widehat{y}, s}^T \right] \right)$
6: **end for**
7: Pick $\hat{t}$ uniformly at random from $\{1, \dots, T\}$
8: **Return:** $\boldsymbol{\theta}^{\hat{t}}$.

Theorem 6. *(Informal statement) Algorithm 1 converges to the set of ϵ-first order stationary points of the FERMI objective (c.f. Eq. 12) in $O(\frac{1}{\epsilon^4})$ iterations (stochastic gradient evaluations).*

The formal statement of this theorem can be found in Theorem 10 in Appendix D. Notice that while this algorithm has a slower rate of convergence than the one in the previous subsection, it is stochastic (each iteration is computationally cheap) and amenable to large-scale problems. Note also that a faster convergence rate of $O(\frac{1}{\epsilon^3})$ could be obtained by using the (more complicated) SREDA method of (Luo et al., 2020) instead of SGDA to solve Eq. (19). We omit the details here. In the next section, we numerically evaluate the performance of the algorithms described in this section.

5 NUMERICAL EXPERIMENTS

5.1 BINARY FAIR CLASSIFICATION WITH A BINARY SENSITIVE ATTRIBUTE

We start the experimental setup for binary classification problem with a binary sensitive attribute. This is a common setup among most existing baseline methods. Per Theorem 2, in this binary

Figure 1: Tradeoff of fairness violation vs test error for different baselines on German Credit and Adult datasets. The desired operation point is the lower left corner where both fairness violation and test error are small. FERMI achieves the best fairness vs performance tradeoff across all baselines.

classification case ERMI is equivalent to Rényi correlation (Baharlouei et al., 2020). While we solve FERMI to impose an ERMI regularizer, we still measure fairness violation via popular fairness violation notions, such as conditional demographic parity L_1 violation (Definition 10), conditional equal opportunity L_1 violation (Definition 11), and equalized odds violation. In Fig. 1, we report the fairness violation vs error for German Credit and Adult datasets. As can be seen, for all three popular notions of fairness, FERMI achieves the best trade-off between fairness and error probability on the test data. This could be partly due to smoothness of FERMI optimization problem, and partly due to the fact that ERMI upper bounds other fairness notions as discussed in Sec. 3.

5.2 NON-BINARY FAIR CLASSIFICATION WITH A NON-BINARY SENSITIVE ATTRIBUTE

Next, we consider a general classification problem with $|\mathcal{S}| > 2$. In this case, we consider the Communities and Crime dataset, which has 18 binary sensitive attributes in total, and we pick $\{7, 10, 14, 18\}$ sensitive attributes out of those for different experiments, which corresponds to $|\mathcal{S}| \in \{2^7, 2^{10}, 2^{14}, 2^{18}\}$. We discretize the target into three classes {high, medium, low} (ternary classification). The only baseline that we are aware of that can handle non-binary classification with non-binary is sensitive attributes is RFI (Baharlouei et al., 2020). The results are presented in Fig. 2, where we use conditional demographic parity L_1 violation (Definition 10) and conditional equal opportunity L_1 violation (Definition 11) as the fairness violation notion. As can be seen,

Figure 2: Comparison between FERMI and RFI (Baharlouei et al., 2020). FERMI achieves a better fairness vs performance tradeoff. Moreover, due to computationally expensive operations like performing singular value decomposition (SVD), RFI has poor scalability with the cardinality of sensitive features and target classes.

FERMI achieves a better tradeoff curve as compared with RFI. Furthermore, it is noteworthy that the per-iteration complexity of FERMI is far less than that of RFI, which requires solving a singular value decomposition at each iteration. Finally, the convergence rate for FERMI given in Theorem 4 guarantee a $O(\frac{1}{\epsilon^2})$ convergence vs $O(\frac{1}{\epsilon^4})$ for RFI (Baharlouei et al., 2020, Theorem 4.1). Empirically, we observed that FERMI converges $\sim$10x faster on this problem instance. Finally, we also show that conditional demographic parity L_1 violation (Definition 10) and square root of ERMI are approximately linearly related, which further justifies the use of ERMI regularizer in the FERMI framework.

5.3 LARGE-SCALE CLASSIFICATION WITH FERMI

In this experiment, we consider the color MNIST dataset (Li & Vasconcelos, 2019) where MNIST digits are colored with different colors drawn from a Gaussian distribution with variance σ around a certain average color. It is shown in (Li & Vasconcelos, 2019) that as $\sigma \to 0$, a convolutional network model overfits significantly to the color on this dataset, and hence will not be able to generalize on a regular black and white test set. We consider the extreme case of $\sigma \to 0$, where the performance on the test set is hit the most and compare FERMI with the existing baselines.

Figure 3: Color MNIST.

The result of the experiment is presented in Fig. 4. As expected, FERMI results in learning representations that are almost independent of the color, hence leading to better generalization. It is noteworthy that the test error achieved by FERMI when $\sigma = 0$ is 22.6%, as compared to 23.3% obtained using REPAIR (Li & Vasconcelos, 2019) for $\sigma = 0.1$. Further decreasing $\sigma \leq 0.05$ the test error using REPAIR sharply goes above 50%. We were unable to run REPAIR for $\sigma = 0$.

Figure 4: The application of FERMI to Color MNIST dataset (Li & Vasconcelos, 2019). As expected, FERMI achieves a tradeoff between demographic parity L_1 violation and train error on the colored training samples. On test data, however, reducing the demographic parity violation translates to less dependence on the color, which avoids overfitting and decreases the test error. Finally, as λ increases we see that the gap between train error and test error significantly decreases which shows that FERMI can avoid overfitting. We also observe that the stochastic variant (Algorithm 1) has better generalization performance (lower test error), while also providing better scalability (with around 100x speedup on per-iteration time complexity).

6 CONCLUSION

In this paper, we proposed a new notion of fairness, called exponential Rényi mutual information (ERMI). We showed that ERMI is a strong notion of fairness violation providing guarantees on several other popular notions, namely Pearson correlation, Rényi correlation, Shannon mutual information, Rényi mutual information, and L_1 distance violation. We proposed a Fair Empirical Risk Minimization framework with an ERMI regularizer to balance performance and fairness, and called it FERMI. We also proposed batch and stochastic algorithms for solving FERMI with convergence guarantees for smooth losses. From an experimental perspective, we showed that FERMI leads to better tradeoffs of fairness vs performance compared to the existing baselines. We also showed that FERMI could be efficiently solved for non-binary sensitive attributes and non-binary target variables.

REFERENCES

Sina Aghaei, Mohammad Javad Azizi, and Phebe Vayanos. Learning optimal and fair decision trees for non-discriminative decision-making. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 33, pp. 1418–1426, 2019.

Julia Angwin, Jeff Larson, Surya Mattu, and Lauren Kirchner. Machine bias. *ProPublica*, 2016.

Sina Baharlouei, Maher Nouiehed, Ahmad Beirami, and Meisam Razaviyayn. Rényi fair inference. In *ICLR*, 2020.

Yahav Bechavod and Katrina Ligett. Penalizing unfairness in binary classification. *arXiv preprint arXiv:1707.00044*, 2017.

Richard Berk, Hoda Heidari, Shahin Jabbari, Matthew Joseph, Michael Kearns, Jamie Morgenstern, Seth Neel, and Aaron Roth. A convex framework for fair regression. *arXiv preprint arXiv:1706.02409*, 2017.

Tolga Bolukbasi, Kai-Wei Chang, James Y Zou, Venkatesh Saligrama, and Adam T Kalai. Man is to computer programmer as woman is to homemaker? debiasing word embeddings. In *Advances in neural information processing systems*, pp. 4349–4357, 2016.

Evgenii Chzhen and Nicolas Schreuder. A minimax framework for quantifying risk-fairness trade-off in regression. *arXiv preprint arXiv:2007.14265*, 2020.

Andrew Cotter, Heinrich Jiang, Maya R Gupta, Serena Wang, Taman Narayan, Seungil You, and Karthik Sridharan. Optimization with non-differentiable constraints with applications to fairness, recall, churn, and other goals. *Journal of Machine Learning Research*, 20(172):1–59, 2019.

Thomas M Cover and Joy A Thomas. Information theory and statistics. *Elements of Information Theory*, 1(1):279–335, 1991.

Imre Csiszár and Paul C Shields. *Information theory and statistics: A tutorial*. Now Publishers Inc, 2004.

Amit Datta, Michael Carl Tschantz, and Anupam Datta. Automated experiments on ad privacy settings. *Proceedings on privacy enhancing technologies*, 2015(1):92–112, 2015.

A. Dembo and O. Zeitouni. *Large deviations techniques and applications*. Springer Science & Business Media, 2009.

Michele Donini, Luca Oneto, Shai Ben-David, John S Shawe-Taylor, and Massimiliano Pontil. Empirical risk minimization under fairness constraints. In *Advances in Neural Information Processing Systems*, pp. 2791–2801, 2018.

Flavio du Pin Calmon, Ali Makhdoumi, Muriel Médard, Mayank Varia, Mark Christiansen, and Ken R Duffy. Principal inertia components and applications. *IEEE Transactions on Information Theory*, 63(8):5011–5038, 2017a.

Flavio du Pin Calmon, Dennis Wei, Bhanukiran Vinzamuri, Karthikeyan Natesan Ramamurthy, and Kush R Varshney. Optimized pre-processing for discrimination prevention. In *Advances in Neural Information Processing Systems*, pp. 3992–4001, 2017b.

Cynthia Dwork, Moritz Hardt, Toniann Pitassi, Omer Reingold, and Richard Zemel. Fairness through awareness. In *Proceedings of the 3rd innovations in theoretical computer science conference*, pp. 214–226, 2012.

Michael Feldman, Sorelle A Friedler, John Moeller, Carlos Scheidegger, and Suresh Venkatasubramanian. Certifying and removing disparate impact. In *proceedings of the 21th ACM SIGKDD international conference on knowledge discovery and data mining*, pp. 259–268, 2015.

Benjamin Fish, Jeremy Kun, and Ádám D Lelkes. A confidence-based approach for balancing fairness and accuracy. In *Proceedings of the 2016 SIAM International Conference on Data Mining*, pp. 144–152. SIAM, 2016.

Hans Gebelein. Das statistische problem der korrelation als variations-und eigenwertproblem und sein zusammenhang mit der ausgleichsrechnung. *ZAMM-Journal of Applied Mathematics and Mechanics/Zeitschrift für Angewandte Mathematik und Mechanik*, 21(6):364–379, 1941.

Vincent Grari, Boris Ruf, Sylvain Lamprier, and Marcin Detyniecki. Fairness-aware neural Réyni minimization for continuous features. *arXiv preprint arXiv:1911.04929*, 2019.

Vincent Grari, Oualid El Hajouji, Sylvain Lamprier, and Marcin Detyniecki. Learning unbiased representations via Rényi minimization. *arXiv preprint arXiv:2009.03183*, 2020.

Moritz Hardt, Eric Price, and Nati Srebro. Equality of opportunity in supervised learning. In *Advances in neural information processing systems*, pp. 3315–3323, 2016.

Hermann O Hirschfeld. A connection between correlation and contingency. In *Proceedings of the Cambridge Philosophical Society*, volume 31, pp. 520–524, 1935.

Ray Jiang, Aldo Pacchiano, Tom Stepleton, Heinrich Jiang, and Silvia Chiappa. Wasserstein fair classification. In *Uncertainty in Artificial Intelligence*, pp. 862–872. PMLR, 2020.

Faisal Kamiran, Toon Calders, and Mykola Pechenizkiy. Discrimination aware decision tree learning. In *2010 IEEE International Conference on Data Mining*, pp. 869–874. IEEE, 2010.

Toshihiro Kamishima, Shotaro Akaho, and Jun Sakuma. Fairness-aware learning through regularization approach. In *2011 IEEE 11th International Conference on Data Mining Workshops*, pp. 643–650. IEEE, 2011.

Michael Kearns, Seth Neel, Aaron Roth, and Zhiwei Steven Wu. Preventing fairness gerrymandering: Auditing and learning for subgroup fairness. In *International Conference on Machine Learning*, pp. 2564–2572, 2018.

Yi Li and Nuno Vasconcelos. Repair: Removing representation bias by dataset resampling. In *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition*, pp. 9572–9581, 2019.

Tianyi Lin, Chi Jin, and Michael I. Jordan. On gradient descent ascent for nonconvex-concave minimax problems. *arXiv: 1906.00331v6*, 2020.

Luo Luo, Haishan Ye, and Tony Zhang. Stochastic recursive gradient descent ascent for stochastic nonconvex-strongly-concave minimax problems. *arXiv: 2001.03724*, 2020.

Yurii Nesterov. *Introductory lectures on convex optimization: A basic course*, volume 87. Springer Science & Business Media, 2013.

Adrián Pérez-Suay, Valero Laparra, Gonzalo Mateo-García, Jordi Muñoz-Marí, Luis Gómez-Chova, and Gustau Camps-Valls. Fair kernel learning. In *Joint European Conference on Machine Learning and Knowledge Discovery in Databases*, pp. 339–355. Springer, 2017.

Geoff Pleiss, Manish Raghavan, Felix Wu, Jon Kleinberg, and Kilian Q Weinberger. On fairness and calibration. In *Advances in Neural Information Processing Systems*, pp. 5680–5689, 2017.

Edward Raff, Jared Sylvester, and Steven Mills. Fair forests: Regularized tree induction to minimize model bias. In *Proceedings of the 2018 AAAI/ACM Conference on AI, Ethics, and Society*, pp. 243–250, 2018.

Alfréd Rényi. On measures of dependence. *Acta Mathematica Academiae Scientiarum Hungarica*, 10(3-4):441–451, 1959.

Alfréd Rényi. On measures of entropy and information. In *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability, Volume 1: Contributions to the Theory of Statistics*. The Regents of the University of California, 1961.

Ashkan Rezaei, Rizal Fathony, Omid Memarrast, and Brian D Ziebart. Fairness for robust log loss classification. In *AAAI*, pp. 5511–5518, 2020.

Daniel Steinberg, Alistair Reid, Simon O'Callaghan, Finnian Lattimore, Lachlan McCalman, and Tiberio Caetano. Fast fair regression via efficient approximations of mutual information. *arXiv preprint arXiv:2002.06200*, 2020.

Latanya Sweeney. Discrimination in online ad delivery. *arXiv preprint arXiv:1301.6822*, 2013.

Bahar Taskesen, Viet Anh Nguyen, Daniel Kuhn, and Jose Blanchet. A distributionally robust approach to fair classification. *arXiv preprint arXiv:2007.09530*, 2020.

Hans S Witsenhausen. On sequences of pairs of dependent random variables. *SIAM Journal on Applied Mathematics*, 28(1):100–113, 1975.

Blake Woodworth, Suriya Gunasekar, Mesrob I Ohannessian, and Nathan Srebro. Learning non-discriminatory predictors. *arXiv preprint arXiv:1702.06081*, 2017.

Muhammad Bilal Zafar, Isabel Valera, Manuel Gomez Rogriguez, and Krishna P Gummadi. Fairness constraints: Mechanisms for fair classification. In *Artificial Intelligence and Statistics*, pp. 962–970. PMLR, 2017.

Rich Zemel, Yu Wu, Kevin Swersky, Toni Pitassi, and Cynthia Dwork. Learning fair representations. In *International Conference on Machine Learning*, pp. 325–333, 2013.

A   EXISTING NOTIONS OF FAIRNESS

Let $(Y, \widehat{Y}, \mathcal{A}, S)$ denote the true target, predicted target, the advantaged outcome class, and the sensitive attribute, respectively. We review three major notions of fairness.

Definition 7 (demographic parity (Dwork et al., 2012))**.** *We say that a learning machine satisfies demographic parity if $\widehat{Y}$ is independent of S.*

Definition 8 (equalized odds (Hardt et al., 2016))**.** *We say that a learning machine satisfies equalized odds, if $\widehat{Y}$ is conditionally independent of S given Y.*

Definition 9 (equal opportunity (Hardt et al., 2016))**.** *We say that a learning machine satisfies equal opportunity with respect to $\mathcal{A}$, if $\widehat{Y}$ is conditionally independent of S given $Y = y$ for all $y \in \mathcal{A}$.*

Notice that the equal opportunity as defined here generalizes the definition in (Hardt et al., 2016). It recovers equalized odds if $\mathcal{A} = \mathcal{Y}$, and it recovers equal opportunity of (Hardt et al., 2016) for $\mathcal{A} = \{1\}$ in binary classification.

Proof of Lemma 1. The proof goes as follows:

- If $Z = 0$ and $\mathcal{Z} = \{0\}$, then conditioning on Z has no effect, and hence $(0, \{0\})$ fairness is equivalent to demographic parity.

- In the case, where $Z = Y$ and $\mathcal{Z} = \mathcal{Y}$, conditioning on Z is equivalent to conditioning on Y, which recovers the equalized odds notion of fairness.

- This is also similar to the previous case with $\mathcal{Y}$ replaced with $\mathcal{A}$.

$\square$

B   ERMI IS AN f-DIVERGENCE

Notice that ERMI is in fact the χ^2-divergence between the joint distribution, $p_{\widehat{Y}, S}$, and the Kronecker product of marginals, $p_{\widehat{Y}} \otimes p_S$. χ^2-divergence is an f-divergence with $f(t) = (t-1)^2$. See (Csiszár & Shields, 2004, Section 4) for a discussion. In the context of discrete random variables, ERMI is referred to as χ^2-information in (du Pin Calmon et al., 2017a). As an immediate result of this observation, we could state the following property of ERMI:

Lemma 2. $D_R(\widehat{Y}; S | Z \in \mathcal{Z}) \geq 0$ *with equality if and only if for all $z \in \mathcal{Z}$, $\widehat{Y}$ and S are conditionally independent given z.*

C   RELATIONS BETWEEN ERMI AND OTHER FAIRNESS VIOLATION NOTIONS

Proof of Theorem 1. We proceed to prove all the (in)equalities one by one:

- $0 \leq I_S(\widehat{Y}; S | Z \in \mathcal{Z})$. This is well known and the proof can be found in any information theory textbook (Cover & Thomas, 1991).

- $I_1(\widehat{Y}; S | Z \in \mathcal{Z}) \leq I_2(\widehat{Y}; S | Z \in \mathcal{Z})$. This is a known property of Rényi mutual information, but we provide a proof for completeness in Lemma 3.

- $I_2(\widehat{Y}; S | Z \in \mathcal{Z}) \leq e^{I_2(\widehat{Y}; S | Z \in \mathcal{Z})} - 1$. This follows from the fact that $x \leq e^x - 1$.

- $e^{I_2(\widehat{Y}; S) | Z \in \mathcal{Z}} - 1 = D_R(\widehat{Y}; S | Z \in \mathcal{Z})$. This follows from simple algebraic manipulation.

$\square$

Lemma 3. *Let $\widehat{Y}, S, Z$ be discrete or continuous random variables. Then:*

(a) For any $\alpha, \beta \in [1, \infty]$, $I_\beta(\widehat{Y}; S | Z \in \mathcal{Z}) \geq I_\alpha(\widehat{Y}; S | Z \in \mathcal{Z})$ if $\beta > \alpha$.

(b) $\lim_{\alpha \to 1^+} I_\alpha(\widehat{Y}; S | Z \in \mathcal{Z}) = I_1(\widehat{Y}; S) := \mathbb{E}_Z \left\{ D_{KL}(p_{\widehat{Y}, S | Z} || p_{\widehat{Y} | Z} \otimes p_{S | Z}) \middle| Z \in \mathcal{Z} \right\}$, where $I_1(\cdot; \cdot)$ denotes the Shannon mutual information and D_{KL} is Kullback–Leibler divergence (relative entropy).

(c) For all $\alpha \in [1, \infty]$, $I_\alpha(\widehat{Y}; S | Z \in \mathcal{Z}) \geq 0$ with equality if and only if for all $z \in \mathcal{Z}$, $\widehat{Y}$ and S are conditionally independent given z.

Proof. *(a)* First assume $0 < \alpha < \beta < \infty$ and that $\alpha, \beta \neq 1$. Define $a = \alpha - 1$, and $b = \beta - 1$. Then the function $\phi(t) = t^{b/a}$ is convex for all $t \geq 0$, so by Jensen's inequality we have:

$$\frac{1}{b} \log \left(\mathbb{E} \left\{ \left(\frac{p(\widehat{Y}, S | Z)}{p(\widehat{Y} | Z) p(S | Z)} \right)^b \middle| Z \in \mathcal{Z} \right\} \right) \geq \frac{1}{b} \log \left(\mathbb{E} \left\{ \left(\frac{p(\widehat{Y}, S | Z)}{p(\widehat{Y} | Z) p(S | Z)} \right)^a \middle| Z \in \mathcal{Z} \right\}^{b/a} \right)$$

$$= \frac{1}{a} \log \left(\mathbb{E} \left\{ \left(\frac{p(\widehat{Y}, S | Z)}{p(\widehat{Y} | Z) p(S | Z)} \right)^a \middle| Z \in \mathcal{Z} \right\} \right). \tag{20}$$

Now suppose $\alpha = 1$. Then by the monotonicity for $\alpha \neq 1$ proved above, we have $I_1(\widehat{Y}; S) = \lim_{\alpha \to 1^-} I_\alpha(\widehat{Y}; S) = \sup_{\alpha \in (0,1)} I_\alpha(\widehat{Y}; S) \leq \inf_{\alpha > 1} I_\alpha(\widehat{Y}; S)$. Also, $I_\infty(\widehat{Y}; S) = \lim_{\alpha \to \infty} I_\alpha(\widehat{Y}; S) = \sup_{\alpha > 0} I_\alpha(\widehat{Y}; S)$.

(b) This is a standard property of the cumulant generating function (see (Dembo & Zeitouni, 2009)).

(c) It is straightforward to observe that independence implies that Rényi mutual information vanishes. On the other hand, if Rényi mutual information vanishes, then part (a) implies that Shannon mutual information also vanishes, which implies the desired conditional independence. □

Proof of Theorem 2. The proof is completed using the following pieces.

- $0 \leq |\rho(\widehat{Y}, S | Z \in \mathcal{Z})| \leq \rho_R(\widehat{Y}, S | Z \in \mathcal{Z})$. This is obvious from the definition of $\rho_R(\widehat{Y}, S | Z \in \mathcal{Z})$.

- $\rho_R(\widehat{Y}, S | Z \in \mathcal{Z}) \leq D_R(\widehat{Y}; S | Z \in \mathcal{Z})$. This follows from Theorem 7.

- Notice that if $|\mathcal{S}| = 2$, Theorem 7 implies that $D_R(\widehat{Y}; S | Z \in \mathcal{Z}) = \rho_R(\widehat{Y}, S | Z \in \mathcal{Z})$.

□

Theorem 7. *Suppose that $\mathcal{S} = [k]$. Let the $k \times k$ matrix P be defined as $P = \{P_{ij}\}_{i,j \in [k] \times [k]}$, where*

$$P_{ij} := \frac{1}{\sqrt{p_S(i) p_S(j)}} \int_{y \in \mathcal{Y}} \left(\frac{p_{\widehat{Y}, S}(y, i) p_{\widehat{Y}, S}(y, j)}{p_{\widehat{Y}}(y)} \right) dy. \tag{21}$$

Let $1 = \sigma_1 \geq \sigma_2 \geq \ldots \geq \sigma_k \geq 0$ be the eigenvalues of P. Then,

$$\rho_R(\widehat{Y}, S) = \sigma_2, \tag{22}$$

$$D_R(\widehat{Y}; S) = \mathrm{Tr}(P) - 1 = \sum_{i=2}^{k} \sigma_i. \tag{23}$$

Proof. Eq. (22) is proved in (Witsenhausen, 1975, Section 3). To prove Eq. (23), notice that

$$\text{Tr}(P) = \sum_{i \in [k]} P_{ii}$$

$$= \sum_{i \in [k]} \frac{1}{p_S(i)} \int_{y \in \mathcal{Y}} \left(\frac{p_{\widehat{Y},S}(y,i)^2}{p_{\widehat{Y}}(y)} \right) dy$$

$$= E_{\widehat{Y},S} \left\{ \left(\frac{p_{\widehat{Y},S}(\widehat{Y},S)}{p_{\widehat{Y}}(\widehat{Y})p_S(S)} \right) \right\}$$

$$= 1 + D_R(\widehat{Y};S),$$

which completes the proof. $\square$

Proof of Theorem 3. The proof for the case where $Z = 0$ and $\mathcal{Z} = \{0\}$ follows from the following set of inequalities:

$$L_2^2(\widehat{Y},S|Z \in \mathcal{Z}) = \sum_{s \in \mathcal{S}} \int_{y \in \mathcal{Y}} \left(p_{\widehat{Y},S}(y,s) - p_{\widehat{Y}}(y)p_S(s) \right)^2 dy \tag{24}$$

$$= \sum_{s \in \mathcal{S}} \int_{y \in \mathcal{Y}} p_{\widehat{Y}}(y)p_S(s) \frac{\left(p_{\widehat{Y},S}(y,s) - p_{\widehat{Y}}(y)p_S(s) \right)^2}{p_{\widehat{Y}}(y)p_S(s)} dy \tag{25}$$

$$\leq \left(\sum_{s \in \mathcal{S}} \int_{y \in \mathcal{Y}} p_{\widehat{Y}}(y)p_S(s) dy \right) \left(\sup_{y \in \mathcal{Y}} \max_{s \in \mathcal{S}} \left\{ \frac{(p_{\widehat{Y},S}(y,s) - p_{\widehat{Y}}(y)p_S(s))^2}{p_{\widehat{Y}}(y)p_S(s)} \right\} \right) \tag{26}$$

$$\leq \sum_{s \in \mathcal{S}} \int_{y \in \mathcal{Y}} \left(\frac{(p_{\widehat{Y},S}(y,s) - p_{\widehat{Y}}(y)p_S(s))^2}{p_{\widehat{Y}}(y)p_S(s)} \right) dy \tag{27}$$

$$= D_R(\widehat{Y};S), \tag{28}$$

where Eq. (26) follows from Hölder's inequality, and Eq. (28) follows from Lemma 4. The extension to general Z and $\mathcal{Z}$ is immediate by observing that $\rho(\widehat{Y},S|Z \in \mathcal{Z}) = \mathbb{E}_Z \left[\rho(\widehat{Y},S|Z) \middle| Z \in \mathcal{Z} \right]$, $\rho_R(\widehat{Y},S|Z \in \mathcal{Z}) = \mathbb{E}_Z \left[\rho_R(\widehat{Y},S|Z) \middle| Z \in \mathcal{Z} \right]$, and $D_R(\widehat{Y},S|Z \in \mathcal{Z}) = \mathbb{E}_Z \left[D_R(\widehat{Y},S|Z) \middle| Z \in \mathcal{Z} \right]$. $\square$

Lemma 4. *We have*

$$D_R(\widehat{Y};S) = \sum_{s \in \mathcal{S}} \int_{y \in \mathcal{Y}} \left(\frac{(p_{\widehat{Y},S}(y,s) - p_{\widehat{Y}}(y)p_S(s))^2}{p_{\widehat{Y}}(y)p_S(s)} \right) dy. \tag{29}$$

Proof. The proof follows from the following set of identities:

$$\sum_{s \in \mathcal{S}} \int_{y \in \mathcal{Y}} \left(\frac{(p_{\widehat{Y},S}(y,s) - p_{\widehat{Y}}(y)p_S(s))^2}{p_{\widehat{Y}}(y)p_S(s)} \right) dy = \sum_{s \in \mathcal{S}} \int_{y \in \mathcal{Y}} \frac{(p_{\widehat{Y},S}(y,s))^2}{p_{\widehat{Y}}(y)p_S(s)} dy$$

$$- 2 \sum_{s \in \mathcal{S}} \int_{y \in \mathcal{Y}} p_{\widehat{Y},S}(y,s) dy$$

$$+ \sum_{s \in \mathcal{S}} \int_{y \in \mathcal{Y}} p_{\widehat{Y}}(y)p_S(s) dy \tag{30}$$

$$= E \left\{ \frac{p_{\widehat{Y},S}(\widehat{Y},S)}{p_{\widehat{Y}}(\widehat{Y})p_S(S)} \right\} - 1 \tag{31}$$

$$= D_R(\widehat{Y};S). \tag{32}$$

$\square$

Next, we present some alternative fairness definitions and show that they are also upper bounded by ERMI.

Definition 10 (conditional demographic parity L_1 violation). *Given a predictor $\widehat{Y}$ supported on $\mathcal{Y}$ and a discrete sensitive attribute S supported on a finite set $\mathcal{S}$, we define the conditional demographic parity violation by:*

$$\widetilde{dp}(\widehat{Y}|S) := \sup_{\widehat{y} \in \mathcal{Y}} \max_{s \in \mathcal{S}} \left| p_{\widehat{Y}|S}(\widehat{y}|s) - p_{\widehat{Y}}(\widehat{y}) \right|. \tag{33}$$

First, we show that $\widetilde{dp}(\widehat{Y}|S)$ is a reasonable notion of fairness violation.

Lemma 5. $\widetilde{dp}(\widehat{Y}|S) = 0$ *iff (if and only if) $\widehat{Y}$ and S are independent.*

Proof. By definition, $\widetilde{dp}(\widehat{Y}|S) = 0$ iff for all $\widehat{y} \in \mathcal{Y}, s \in \mathcal{S}$, $p_{\widehat{Y},S}(\widehat{y}|s) = p_{\widehat{Y}}(\widehat{y})$ iff $\widehat{Y}$ and S are independent (since we always assume $p(s) > 0$ for all $s \in \mathcal{S}$). □

Theorem 8 (ERMI is stronger than conditional demographic parity L_1 violation). *Let $\widehat{Y}$ be a discrete or continuous random variable supported on $\mathcal{Y}$, and S be a discrete random variable supported on a finite set $\mathcal{S}$. Denote $p_S^{\min} := \min_{s \in \mathcal{S}} p_S(s) > 0$. Then,*

$$0 \le \widetilde{dp}(\widehat{Y}|S) \le \frac{1}{p_S^{\min}} \sqrt{D_R(\widehat{Y}; S)}. \tag{34}$$

Proof. The proof follows from the following set of (in)equalities:

$$\left(\widetilde{dp}(\widehat{Y}|S) \right)^2 = \sup_{\widehat{y} \in \mathcal{Y}} \max_{s \in \mathcal{S}} \left(p_{\widehat{Y}|S}(\widehat{y}|s) - p_{\widehat{Y}}(\widehat{y}) \right)^2 \tag{35}$$

$$\le \frac{1}{(p_S^{\min})^2} \sup_{\widehat{y} \in \mathcal{Y}} \max_{s \in \mathcal{S}} \left(p_{\widehat{Y},S}(\widehat{y}, s) - p_{\widehat{Y}}(\widehat{y}) p_S(s) \right)^2 \tag{36}$$

$$\le \frac{1}{(p_S^{\min})^2} \int_{\widehat{y} \in \mathcal{Y}} \sum_{s \in \mathcal{S}} \left(p_{\widehat{Y},S}(\widehat{y}, s) - p_{\widehat{Y}}(\widehat{y}) p_S(s) \right)^2 \tag{37}$$

$$= \frac{1}{(p_S^{\min})^2} D_R(\widehat{Y}; S), \tag{38}$$

where Eq. (38) follows from Theorem 3. □

Definition 11 (conditional equal opportunity L_1 violation (Hardt et al., 2016)). *Let $Y, \widehat{Y}$ take values in $\mathcal{Y}$ and let $\mathcal{A} \subseteq \mathcal{Y}$ be a compact subset denoting the advantaged outcomes (For example, the decision "to interview" an individual or classify an individual as a "low risk" for financial purposes). We define the conditional equal opportunity L_1 violation of $\widehat{Y}$ with respect to the sensitive attribute S and the advantaged outcome $\mathcal{A}$ by*

$$\widetilde{eo}(\widehat{Y}|S, Y \in \mathcal{A}) := \mathbb{E}_Y \left\{ \sup_{\widehat{y} \in \mathcal{Y}} \max_{s \in \mathcal{S}} \left| p_{\widehat{Y},S|Y}(\widehat{y}|s, Y) - p_{\widehat{Y}|Y}(\widehat{y}|Y) \right| \Big| Y \in \mathcal{A} \right\}. \tag{39}$$

Theorem 9 (ERMI is stronger than generalized equal opportunity Kolmogorov violation - alternative definition). *Let $\widehat{Y}, Y$, be discrete or continuous random variables supported on $\mathcal{Y}$, and let S be a discrete random variable supported on a finite set $\mathcal{S}$. Let $\mathcal{A} \subseteq \mathcal{Y}$ be a compact subset of $\mathcal{Y}$. Denote $p_{S|\mathcal{A}}^{\min} = \min_{s \in \mathcal{S}, y \in \mathcal{A}} p_{S|Y}(s|y)$. Then,*

$$0 \le \widetilde{eo}(\widehat{Y}|S, Y \in \mathcal{A}) \le \frac{1}{p_{S|\mathcal{A}}^{\min}} \sqrt{D_R(\widehat{Y}; S|Y \in \mathcal{A})}. \tag{40}$$

Proof. Notice that the same proof for Theorem 8 would give that for all $y \in \mathcal{A}$:

$$0 \le \sup_{\widehat{y} \in \mathcal{Y}} \max_{s \in \mathcal{S}} \left| p_{\widehat{Y}, S|Y}(\widehat{y}|s, y) - p_{\widehat{Y}|Y}(\widehat{y}|y) \right| := \widetilde{\mathrm{eo}}(\widehat{Y}|S, Y = y)$$

$$\le \frac{1}{p_{S|y}^{\min}(y)} \sqrt{D_R(\widehat{Y}; S|Y = y)}$$

$$\le \frac{1}{p_{S|\mathcal{C}}^{\min}} \sqrt{D_R(\widehat{Y}; S|Y = y)}.$$

Hence,

$$\widetilde{\mathrm{eo}}(\widehat{Y}|S, Y \in \mathcal{A}) = \mathbb{E}_Y \left\{ \widetilde{\mathrm{eo}}(\widehat{Y}|S, Y) \,\middle|\, Y \in \mathcal{A} \right\}$$

$$\le \frac{1}{p_{S|\mathcal{A}}^{\min}} \mathbb{E}_Y \left\{ \sqrt{D_R(\widehat{Y}; S|Y)} \,\middle|\, Y \in \mathcal{A} \right\}$$

$$\le \frac{1}{p_{S|\mathcal{A}}^{\min}} \sqrt{\mathbb{E}_Y \left\{ D_R(\widehat{Y}; S|Y) \,\middle|\, Y \in \mathcal{A} \right\}}$$

$$= \frac{1}{p_{S|\mathcal{A}}^{\min}} \sqrt{D_R(\widehat{Y}; S|Y \in \mathcal{A})},$$

where the last inequality follows from Jensen's inequality. This completes the proof. $\qquad\square$

D   STOCHASTIC FERMI

Proof of Theorem 5. Denote $W^* \in \arg\max_{W \in \mathbb{R}^{k \times m}} - \mathrm{Tr}(W P_{\widehat{y}} W^T) + 2\,\mathrm{Tr}(W P_{\widehat{y}, s} P_s^{-1/2})$. Set the derivative of the expression on the RHS equal to zero:

$$-2W P_{\widehat{y}} + 2 P_s^{-1/2} P_{\widehat{y}, s}^T = 0 \implies W^* = P_{\widehat{y}}^{-1} P_{\widehat{y}, s}^T P_s^{-1/2}.$$

Plugging this expression for W^*, we have

$$\max_{W \in \mathbb{R}^{k \times m}} - \mathrm{Tr}(W P_{\widehat{y}} W^T) + 2\,\mathrm{Tr}(W P_{\widehat{y}, s} P_s^{-1/2})$$

$$= - \mathrm{Tr}(P_s^{-1/2} P_{\widehat{y}, s}^T P_{\widehat{y}}^{-1} P_{\widehat{y}} P_{\widehat{y}}^{-1} P_s^{-1/2}) + 2\,\mathrm{Tr}(P_s^{-1/2} P_{\widehat{y}, s}^T P_{\widehat{y}}^{-1} P_{\widehat{y}} P_{\widehat{y}}^{-1} P_s^{-1/2})$$

$$= \mathrm{Tr}(P_s^{-1/2} P_{\widehat{y}, s}^T P_{\widehat{y}}^{-1} P_{\widehat{y}, s} P_s^{-1/2})$$

$$= \mathrm{Tr}(P_s^{-1} P_{\widehat{y}, s}^T P_{\widehat{y}}^{-1} P_{\widehat{y}, s}).$$

Writing out the matrix multiplication explicitly in the last expression, we have

$$P_s^{-1} P_{\widehat{y}, s}^T P_{\widehat{y}}^{-1} P_{\widehat{y}, s} = U V^T,$$

where $U_{i,j} = \mathbb{P}(S = i)^{-1} \mathbb{P}(S = i, \widehat{Y} = j)$ and $V_{i,j} = \mathbb{P}(\widehat{Y} = j)^{-1} \mathbb{P}(\widehat{Y} = j, S = i)$, for $i \in [k], j \in [m]$. Hence

$$\max_{W \in \mathbb{R}^{k \times m}} - \mathrm{Tr}(W P_{\widehat{y}_\theta} W^T) + 2\,\mathrm{Tr}(W P_{\widehat{y}_\theta, s} P_s^{-1/2}) = \mathrm{Tr}(U V^T)$$

$$= \sum_{i \in [k]} \sum_{j \in [m]} \frac{\mathbb{P}(S = i, \widehat{Y} = j)^2}{\mathbb{P}(S = i) \mathbb{P}(\widehat{Y} = j)}$$

$$= \mathrm{Tr}(Q_\theta^T Q_\theta).$$

Hence the result follows from Theorem 7 (applied to the empirical probabilities). $\qquad\square$

Next, we move to the statement and proof of the precise version of Theorem 6. We first recall some basic definitions:

Definition 12. *A function f is β-smooth if for all $\mathbf{u}, \mathbf{u}'$, we have $\|\nabla f(\mathbf{u}) - \nabla f(\mathbf{u})\| \le \beta \|\mathbf{u} - \mathbf{u}'\|$.*

Definition 13. *A point $\boldsymbol{\theta}$ is an ϵ-stationary point of a differentiable function Φ if $\|\nabla\Phi(\boldsymbol{\theta})\| \leq \epsilon$.*

Assumption 2.
- ℓ *is twice differentiable, L_ℓ-Lipscthiz, and β_ℓ-smooth in $\boldsymbol{\theta}$.*

- $\|\nabla_\theta P_{\widehat{y}}\|_2 := \|\nabla_\theta \text{vec}(P_{\widehat{y}})\|_2 \leq L_y$ *and* $\max_{l\in[m]} \|\nabla_\theta ((P_{\widehat{y}})_{l,l})\|_2 \leq \widetilde{L}_y$

- $\max_{l\in[m]} \|\nabla_{\theta\theta}^2 (P_{\widehat{y}})_{l,l}\|_2 \leq \beta_y$.

- $\|\nabla_\theta P_{\widehat{y},s}^T\|_2 := \|\nabla_\theta \text{vec}(P_{\widehat{y},s}^T)\|_2 \leq L_{ys}$ *and* $\max_{l\in[m],j\in[k]} \|\nabla_\theta ((P_{\widehat{y},s})_{l,m})\|_2 \leq \widetilde{L}_{ys}$

- $\max_{l\in[m],j\in[k]} \|\nabla_{\theta\theta}^2 (P_{\widehat{y},s})_{l,j}\|_2 \leq \beta_{y,s}$.

Theorem 10 (Precise version of Theorem 6). *Denote*

$$f(\boldsymbol{\theta}, W) = \frac{1}{N}\sum_{i\in[N]} \ell(\mathbf{x}_i, y_i; \boldsymbol{\theta}) + \lambda\left(-\text{Tr}(WP_{\widehat{y}}W^T) + 2\text{Tr}(WP_{\widehat{y},s}P_s^{-1/2}) - 1\right).$$

Set $\mathcal{W} := B_F(0, 2D) \subset \mathbb{R}^{k\times m}$ *(Frobenius norm ball of radius $2D$),* $D := \frac{\sqrt{mk}}{\hat{p}_{\widehat{y}}^{\min}\sqrt{\hat{p}_s^{\min}}}$. *Denote* $\Delta_\Phi := \Phi(\theta_0) - \min_\theta \Phi(\theta)$, *where* $\Phi(\theta) := \max_{W\in\mathcal{W}} f(\boldsymbol{\theta}, W)$. *In Algorithm 1, choose the step-sizes as* $\eta_\theta = \Theta(1/\kappa^2\beta)$ *and* $\eta_W = \Theta(1/\beta)$ *and mini-batch size as* $M = \Theta\left(\max\{1, \kappa\sigma^2\epsilon^{-2}\}\right)$. *Then under Assumption 2, the iteration complexity of Algorithm 1 to return an ϵ-stationary point of f is bounded by*

$$\mathcal{O}\left(\frac{\kappa^2\beta\Delta_\Phi + \kappa\beta^2 D^2}{\epsilon^2}\right),$$

which gives the total stochastic gradient complexity of

$$\mathcal{O}\left(\frac{\kappa^2\beta\Delta_\Phi + \kappa\beta^2 D^2}{\epsilon^2} \max\{1, \kappa\sigma^2\epsilon^{-2}\}\right),$$

where

$$\beta = \beta_l + 8\lambda D^2\beta_y + 4\lambda\frac{1}{\sqrt{\hat{p}_s^{\min}}}\left(\sqrt{m}k^{3/2}D\beta_{ys}\right) + 2\lambda + 4\lambda\left(DL_y + \frac{L_{ys}}{\sqrt{\hat{p}_s^{\min}}}\right),$$

$$\kappa = \beta/\mu,$$

$$\sigma^2 = 2\left(L_\ell + 2\lambda\widetilde{L}_yD^2 + 4\lambda\frac{D}{\sqrt{\hat{p}_s^{\min}}}\sqrt{mk}\widetilde{L}_{ys}\right)^2 + 2\left(2\lambda D + 2(\hat{p}_s^{\min})^{-1/2}\sqrt{mk}\right)^2$$

The theorem follows from (Lin et al., 2020, Theorem 4.5) combined with the following technical lemmas. We assume Assumption 2 holds for the remainder of the proof of Theorem 10:

Lemma 6. *Let*

$$f(\boldsymbol{\theta}, W) = \frac{1}{N}\sum_{i\in[N]} \ell(\mathbf{x}_i, y_i; \boldsymbol{\theta}) + \lambda\left(-\text{Tr}(WP_{\widehat{y}}W^T) + 2\text{Tr}(WP_{\widehat{y},s}P_s^{-1/2}) - 1\right)$$

$$:= \frac{1}{N}\sum_{i\in[N]} g(\boldsymbol{\theta}, W, \mathbf{x}_i, y_i).$$

Then

1. *f is β-smooth, where* $\beta = \beta_l + 8\lambda D^2\beta_y + 4\lambda\frac{1}{\sqrt{\hat{p}_s^{\min}}}\left(\sqrt{m}k^{3/2}D\beta_{ys}\right) + 2\lambda + 4\lambda\left(DL_y + \frac{L_{ys}}{\sqrt{\hat{p}_s^{\min}}}\right)$.

2. *$f(\boldsymbol{\theta}, \cdot)$ is $2\lambda\hat{p}_{\widehat{y}}^{\min}$-strongly concave for all $\boldsymbol{\theta}$.*

3. *$\|W^*\|_F \leq D$, where D is as defined in Theorem 10 and W^* denotes any maximizer of $f(\boldsymbol{\theta}, W)$.*

Proof. By Assumption 2, g is twice continuously differentiable. Hence for part 1, it suffices to upper bound the spectral norm of the second derivative of $g(\cdot, \cdot, \mathbf{z})$ by β for all $\mathbf{z} = (\mathbf{x}, y)$, where we vectorize and then differentiate with respect to $w := \text{vec } W$ and/or $\boldsymbol{\theta}$, so that the resulting first and second derivatives are always vectors or a matrices (not tensors). Notice that $g(\boldsymbol{\theta}, w, \mathbf{z}) = \ell(\mathbf{z}, \boldsymbol{\theta}) - \lambda w^T (P_{\widehat{y}} \otimes \mathbf{I}) w + 2\lambda (\text{vec}(W))^T P_{\widehat{y},s} P_s^{-1/2} - \lambda$ and

$$\nabla^2 g(\boldsymbol{\theta}, w, \mathbf{z}) = \begin{pmatrix} \nabla^2_{\theta\theta} g(\boldsymbol{\theta}, w, \mathbf{z}) & \nabla^2_{\theta w} g(\boldsymbol{\theta}, w, \mathbf{z}) \\ \nabla^2_{w\theta} g(\boldsymbol{\theta}, w, \mathbf{z}) & \nabla^2_{ww} g(\boldsymbol{\theta}, w, \mathbf{z}) \end{pmatrix}.$$

Further, by the definition of operator norm, we have

$$\|\nabla^2 g(\boldsymbol{\theta}, w, \mathbf{z})\|_2 \le \|\nabla^2_{\theta\theta} g(\boldsymbol{\theta}, w, \mathbf{z})\|_2 + 2\|\nabla^2_{\theta w} g(\boldsymbol{\theta}, w, \mathbf{z})\|_2 + \|\nabla^2_{ww} g(\boldsymbol{\theta}, w, \mathbf{z})\|_2.$$

Now we vectorize all matrices and then compute derivatives of g with respect to θ and $\text{vec}(W)$:

$$\nabla_\theta g(\boldsymbol{\theta}, w, \mathbf{z}) = \nabla_\theta \ell(\mathbf{z}, \boldsymbol{\theta}) - 2\lambda \nabla_\theta \text{vec}(P_{\widehat{y}})^T \text{vec}(W^T W) + 2\lambda \nabla_\theta \text{vec}(P_{\widehat{y},s})^T \text{vec}(W^T P_s^{-1/2})$$

(41)

$$= \nabla_\theta \ell(\mathbf{z}, \boldsymbol{\theta}) - 2\lambda \left[\sum_{l \in [m], i \in [k]} W_{i,l}^2 \nabla_\theta ((P_{\widehat{y}})_{l,l}) \right]$$

$$+ 2\lambda \left[\sum_{j \in [m], i \in [k]} W_{i,j} (\nabla_\theta (P_{\widehat{y}s})_{j,i}) (P_s^{-1/2})_{i,i} \right];$$

(42)

$$\nabla_w g(\boldsymbol{\theta}, w, \mathbf{z}) = -2\lambda W P_{\widehat{y}} + 2\lambda P_s^{-1/2} P_{\widehat{y},s}^T.$$

(43)

Differentiating again yields:

$$\nabla^2_{ww} g(\boldsymbol{\theta}, w, \mathbf{z}) = -2\lambda P_{\widehat{y}} \otimes \mathbf{I}_k;$$

$$\nabla^2_{w\theta} g(\boldsymbol{\theta}, w, \mathbf{z}) = \frac{\partial}{\partial \theta} \frac{\partial g(\boldsymbol{\theta}, w, \mathbf{z})}{\partial w} = -2\lambda (\mathbf{I}_m \otimes W) \nabla_\theta P_{\widehat{y}} + 2\lambda (\mathbf{I}_m \otimes P_s^{-1/2}) \nabla_\theta \text{vec}(P_{\widehat{y},s}^T);$$

$$\nabla^2_{\theta\theta} g(\boldsymbol{\theta}, w, \mathbf{z}) = \nabla^2_\theta \ell(\mathbf{z}, \boldsymbol{\theta}) - 2\lambda \left[\sum_{l \in [m], i \in [k]} W_{i,l}^2 \nabla^2_{\theta\theta} ((P_{\widehat{y}})_{l,l}) \right]$$

$$+ 2\lambda \left[\sum_{j \in [m], i \in [k]} W_{i,j} (\nabla^2_{\theta\theta} (P_{\widehat{y}s})_{j,i}) (P_s^{-1/2})_{i,i} \right].$$

Then to establish part 1, use Assumption 2, Clairaut's theorem, the definitions of the matrices and fact that their entries are in $[0, 1]$, the relations $\|AB\|_2 \le \|A\|_2 \|B\|_2$ and $\|\text{vec } W\|_1 \le \sqrt{mk}\|\text{vec } W\|_2 = \sqrt{mk}\|W\|_F$, and the fact that $\|A \otimes B\|_2 = \|A\|_2 \|B\|_2$ to bound the spectral norm of each second derivative above.

The strong concavity statement follows by noticing $\nabla^2_{ww} g(\boldsymbol{\theta}, W) \preccurlyeq -\mu \mathbf{I}$ iff $P_{\widehat{y}} \succcurlyeq \frac{\mu}{2\lambda} \mathbf{I}$ iff $\min_{i \in [m]} p_{\widehat{y}}(i) \ge \frac{\mu}{2\lambda}$.

Part 3 follows from the expression for W^* in the proof of Theorem 5. $\qquad\square$

Lemma 7. *Consider f and g as defined above. Then we have*

$$\mathbb{E}_{\mathbf{z}} \nabla g(\boldsymbol{\theta}, W, \mathbf{z}) = \nabla f(\boldsymbol{\theta}, W),$$

$$\mathbb{E}_{\mathbf{z}} \|\nabla g(\boldsymbol{\theta}, W, \mathbf{z}) - \nabla f(\boldsymbol{\theta}, W)\|_2^2 \le 2 \left(L_\ell + 2\lambda \widetilde{L}_y D^2 + 4\lambda \frac{D}{\sqrt{\widehat{p}_s^{\min}}} \sqrt{mk} \widetilde{L}_{ys} \right)^2$$

$$+ 2 \left(2\lambda D + 2(\widehat{p}_s^{\min})^{-1/2} \sqrt{mk} \right)^2,$$

where both expectations are with respect to the empirical distribution on $\{\mathbf{z}_i\}_{i \in [N]}$.

Proof. The first statement is obvious. The second follows from Eq. (42) in the proof of Lemma 6, since

$$
\mathbb{E}_{\mathbf{z}} \| \nabla g(\boldsymbol{\theta}, W, \mathbf{z}) - \nabla f(\boldsymbol{\theta}, W) \|_2^2
$$

$$
= \frac{1}{N} \sum_{i=1}^{N} \| \nabla g(\boldsymbol{\theta}, W, z_i) \|_2^2 - \frac{1}{N^2} \sum_{i,j=1}^{N} \langle \nabla g(\boldsymbol{\theta}, W, \mathbf{z}_i), \nabla g(\boldsymbol{\theta}, W, \mathbf{z}_j) \rangle
$$

$$
\leq 2 \sup_{\mathbf{z}_i} \| \nabla g(\boldsymbol{\theta}, W, \mathbf{z}_i) \|_2^2
$$

$$
\leq 2 \sup_{\mathbf{z}} \left\{ \| \nabla_\theta g(\boldsymbol{\theta}, W, \mathbf{z}) \|^2 + \| \nabla_w g(\boldsymbol{\theta}, W, \mathbf{z}) \|^2 \right\}
$$

$$
\leq 2 \sup_{\mathbf{z}} \left\{ \left\| \nabla_\theta \ell(\mathbf{z}, \boldsymbol{\theta}) - 2\lambda \left[\sum_{l \in [m], i \in [k]} W_{i,l}^2 \nabla_\theta \left((P_{\widehat{y}})_{l,l} \right) \right] \right. \right.
$$

$$
\left. + 2\lambda \left[\sum_{j \in [m], i \in [k]} W_{i,j} (\nabla_\theta \left(P_{\widehat{y}s} \right)_{j,i}) (P_s^{-1/2})_{i,i} \right] \right\|_2^2 \right\}
$$

$$
+ 2 \left\| -2\lambda W P_{\widehat{y}} + 2\lambda P_s^{-1/2} P_{\widehat{y},s}^T \right\|_2^2 .
$$

Then use Assumption 2 and basic norm inequalities to bound the norm of each term. □

E   DATASETS DESCRIPTION

In this section we introduce the datasets used in numerical experiment discussed in Section Sec. 5. All of these datasets are publicly available at UCI repository.

- **German Credit Dataset.**[3] German Credit dataset consists of 20 features (13 categorical and 7 numerical) regarding to social, and economic status of 1000 customers. The assigned task is to classify customers as good or bad credit risks. Without imposing fairness, the DP violation of the trained model is larger than 20%. We chose first 800 customers as the training data, and last 200 customers as the test data. The sensitive attributes are gender, and marital-status.

- **Adult Dataset.**[4] Adult dataset contains the census information of individuals including education, gender, and capital gain. The assigned classification task is to predict whether a person earns over 50k annually. The train and test sets are two separated files consisting of 32,000 and 16,000 samples respectively. We consider gender and race as the sensitive attributes (For the experiments involving one sensitive attribute, we have chosen gender). Learning a logistic regression model on the training dataset (without imposing fairness) shows that only 3 features out of 14 have larger weights than the gender attribute. Note that removing the sensitive attribute (gender), and retraining the model does not eliminate the bias of the classifier. the optimal logistic regression classifier in this case is still highly biased. For the clustering task, we have chosen 5 continuous features (Capital-gain, age, fnlwgt, capital-loss, hours-per-week), and 10,000 samples to cluster. The sensitive attribute of each individual is gender.

- **Communities and Crime Dataset**. The dataset is cleaned and processed as described in (Kearns et al., 2018). Briefly, each record in this dataset summarizes aggregate socioeconomic information about both the citizens and police force in a particular U.S. community, and the problem is to predict whether the community has a high rate of violent crime. [5]

---

[3]https://archive.ics.uci.edu/ml/datasets/statlog+(german+credit+data)
[4]https://archive.ics.uci.edu/ml/datasets/adult.
[5]http://archive.ics.uci.edu/ml/datasets/communities+and+crime