# Learning to See Sharper: A Physics-Informed Artificial Intelligence Framework for Super-Resolving Galaxy Spectra

**Aryana Haghjoo**[1]     **Shoubaneh Hemmati**[2]

**Bahram Mobasher**[1]

[1]Dept. of Physics & Astronomy, UC Riverside     [2]IPAC, California Institute of Technology

ahagh010@ucr.edu

## Abstract

The scientific information recoverable from galaxy spectra is fundamentally set by the resolution, yet assembling large high-resolution samples remains observationally expensive. We present a three-stage physics-informed deep-learning framework for spectral super-resolution that enhances *James Webb Space Telescope* (JWST)/Near Infrared Spectrograph (NIRSpec) low-resolution prism spectra ($\mathcal{R} \sim 100$) by a factor of $\sim 10$ in resolving power ($\mathcal{R} \sim 1000$), recovering medium-resolution detail from prism-only observations. Trained on 1,187 paired *JWST Advanced Deep Extragalactic Survey* (JADES) observations, our model achieves noise-limited residuals across most of the 1–5 $\mu$m range and systematically improves the signal-to-noise (S/N) of key diagnostic emission lines, such as [O II], H$\beta$, [O III], and H$\alpha$, often by factors of several. The architecture embeds spectroscopic physics directly: a dedicated redshift-inference head provides global spectral context, while a line-token self-attention branch parameterizes individual emission features as Gaussian profiles constrained by physical inter-line relationships. The framework can be extended to forthcoming wide-field grism surveys (*Euclid*, *Roman Space Telescope*), enabling population-level diagnostics across millions of galaxies otherwise inaccessible at low resolution.

## 1   Introduction

The physical information encoded in galaxy spectra is gated by observed resolution. From redshifts and star-formation rates to gas-phase metallicities and active galactic nuclei (AGN) diagnostics, the richest spectroscopic diagnostics require well-resolved line features. Emission lines close in wavelength such as H$\alpha$/[N II], the [O III] $\lambda\lambda 4959, 5007$ doublet, blend irretrievably at low resolution, biasing every downstream physical quantity extraction (Faisst et al., 2018; Agostino et al., 2023).

Modern wide-area missions face an inherent tension between survey volume and spectral quality. Space observatories such as *Euclid* and the *Roman Space Telescope* will deliver grism spectroscopy for tens of millions of galaxies at $\mathcal{R} \sim 100\text{–}300$, far below what resolves fine diagnostic lines at many redshifts (Laureijs et al., 2010; Spergel et al., 2013). Targeted programs such as JADES with JWST/NIRSpec provide exquisite medium-resolution data but cover far fewer targets.

One way to circumvent this trade-off is to leverage the small but growing number of galaxies observed at both low and high spectral resolution, and use machine learning to learn the mapping between the two, enabling the resolution enhancement of the much larger low-resolution samples. Machine learning has found broad application in astronomical data analysis (Hemmati et al., 2022; Sanjaripour et al., 2024; Parker et al., 2024), yet spectral resolution enhancement remains largely unexplored. The model is trained on paired spectra from the *JWST Advanced Deep Extragalactic Survey* (JADES;

Conference on Physics and AI at Stanford University (PAI 2026).

Eisenstein et al. 2023), where low-resolution observations are matched with medium-resolution spectra of the same galaxies.

Figure 1: Example paired JWST/NIRSpec spectra for a representative JADES galaxy ($z = 3.235$). The prism (red) covers 0.6–5.3 $\mu$m at $\mathcal{R} \sim 100$; the three medium-resolution ($\mathcal{R} \sim 1000$) grating segments (G140M, G235M, G395M; green, blue, black) are overplotted. Vertical dashed lines mark key emission lines at the galaxy's redshift. The prism blends features that the gratings can resolve.

## 2  Data: Paired JWST/NIRSpec JADES Spectra

Our training dataset is drawn from the JADES dataset Eisenstein et al. 2023; Bunker et al. 2024; D'Eugenio et al. 2025; Curtis-Lake et al. 2026), conducted with the NIRSpec instrument. JADES observes the same galaxies in both the prism disperser ($\mathcal{R} \sim 100$, 0.6–5.3 $\mu$m) and three medium-resolution grating modes (G140M, G235M, G395M; combined 1–5.5 $\mu$m, $\mathcal{R} \sim 1000$), providing naturally co-observed paired spectra. An example of the prism and medium-resolution spectra for a representative JADES galaxy is shown in Figure 1.

**Pre-processing.** Sources observed in both modes are cross-matched by sky position (separation $< 0.2''$) and field name. The three grating spectra are merged into a single continuous reference by minimizing root mean square (RMS) flux differences in overlap regions and applying consistent flux scale factors. Both prism and stitched grating spectra are placed on a common wavelength grid defined by the native grating sampling; the prism is up-sampled via cubic interpolation, which preserves broadband shape without introducing artificial structure. Galaxies are retained only if their medium-resolution spectrum shows $S/N > 5$ in at least one of H$\alpha$, [O II], or [O III], yielding **1,187 paired spectra**.

**Augmentation.** Each pair generates 20 stochastic realizations via Gaussian redshift offsets ($\sigma_z = 0.3$) and 10% flux perturbations, yielding 24,927 augmented pairs that promote shift-robust, noise-tolerant learning.

## 3  Physics-Informed Super-Resolution Framework

Our pipeline comprises three sequentially trained components; Figure 2 shows the overall design.

**Stage 1 — Coarse Super-Resolution (SR1).** SR1 is a 1D residual convolutional neural network that maps the prism spectrum $x_{\mathrm{lr}}$ to an initial super-resolved estimate $x_{\mathrm{sr1}}$ and a predicted log-variance $\log \sigma_{\mathrm{sr1}}^2$. It is trained with a heteroscedastic negative log-likelihood loss that reflects the wavelength-dependent noise of the NIRSpec detector. The architecture uses $\sim 8$ residual blocks with small convolutional kernels and hidden dimension $\sim 128$, providing a physically plausible spectral backbone on which subsequent stages operate.

**Stage 2 — Redshift Inference (ZHead).** ZHead is a 1D CNN applied to the SR1 output that infers a redshift estimate $\hat{z}$ and its uncertainty $\log \sigma_{\hat{z}}^2$. Treating redshift as an explicit global latent parameter captures long-range spectral coherence that local convolutions cannot encode: line pairs relevant to one another (e.g., [O III] $\lambda\lambda 4959, 5007$ and H$\beta$) may be separated by hundreds of pixels in the observed frame. During SR2 training, a teacher-forcing schedule linearly decays the probability of substituting the catalog redshift for $\hat{z}$ from 0.8 to 0.2, transitioning the model from supervised to self-consistent redshift inference.

**Stage 3 — Physics-Informed Residual Refinement (SR2).** SR2 predicts a residual correction $\Delta x = \Delta x_{\mathrm{line}} + \Delta x_{\mathrm{CNN}}$ to the SR1 output via two parallel branches.

*Line-token self-attention branch.* For each of $K = 20$ known rest-frame emission lines, a local spectral window around the expected observed-frame wavelength at $\hat{z}$ is encoded by a small CNN into a compact feature vector; a learnable species embedding distinguishes morphologically similar lines. The $K$ tokens are processed by a Transformer encoder, enabling every line to exchange information

Figure 2: Three-stage super-resolution architecture. **SR1**: a 1D ResNet-CNN maps the prism spectrum to a coarse super-resolved estimate and predictive uncertainty. **ZHead**: a 1D CNN infers the galaxy redshift $\hat{z}$ from the SR1 output. **SR2**: a two-branch residual refiner conditioned on $\hat{z}$, combining a line-token self-attention branch (parametric Gaussian profiles per emission line) with a convolutional branch for smooth continuum correction.

globally. This explicitly encodes physical inter-line relations ($\approx 1{:}3$ [O III] doublet ratio, Balmer decrement) inaccessible to local convolutions. Each token decodes into a parametric Gaussian profile; a redshift-conditioned wavelength mask suppresses outputs where no feature is expected.

*Continuum correction branch.* A shallow ResNet-CNN with scaled skip connections ($\times 0.5$) predicts smooth continuum deviations $\Delta x_{\mathrm{CNN}}$, biased toward low-frequency adjustments. The two-branch design physically decomposes the galaxy spectrum into discrete emission features plus a slowly varying continuum.

**Hyperparameter Optimization.** We performed three sequential Bayesian sweeps using Weights & Biases (Biewald, 2020), with a total of $\sim 100{,}000$ trials, exploring learning rate, weight decay, dropout, hidden dimension, residual blocks, and activation per stage; SR2 additionally covered attention heads/layers, spectral window width, and L1 amplitude penalty. Optimal values across all stages consistently favor exponential linear unit (ELU) activations, learning rates $\sim 10^{-4}$, and $\sim 8$ residual blocks.

**Computational resources.** All stages were trained on a single **NVIDIA RTX 3090** (24 GB VRAM); final models require 2–4 hours per stage, with the full Bayesian sweep costing $\approx 400$ GPU-hours. The pipeline totals $\approx 2.5$M parameters; inference takes $< 1$ s per spectrum.

## 4 Results

We evaluate performance on the 20% held-out evaluation set.

**Spectral reconstruction.** Figure 3 shows example reconstructed spectra for two held-out galaxies. SR2 resolves the [O III] $\lambda\lambda 4959, 5007$ doublet into two peaks at the correct flux ratio ($\approx 1{:}3$) while the prism input shows only broad, blended features. Residual analysis confirms that SR2–high-resolution (HR) residuals are noise-limited: the robust scatter $\sigma_{\mathrm{resid}}(\lambda)$ converges to the median HR noise amplitude $\tilde{\sigma}_{\mathrm{HR}}(\lambda)$ above $\sim 2\,\mu$m.

**Emission-line S/N improvement.** For each galaxy and diagnostic line, we fit a Gaussian model $f(\lambda) = C_0 + C_1\lambda + A\exp[-(\lambda - \mu)^2/2\sigma^2]$ to the low-resolution (LR) and super-resolved (SR) spectra, computing S/N $= A/\delta A$ from the fit covariance. Figure 4 shows that for all four key diagnostics, the majority of galaxies lie above the one-to-one relation, confirming systematic improvement. These gains translate into higher-fidelity emission line detection and thus more precise estimates of the physical characteristics of the galaxy.

**Redshift consistency as a downstream diagnostic.** Identical redshift-inference networks trained on LR, SR, and HR spectra show SR reduces the normalized median absolute deviation [NMAD; $|\Delta z|/(1 + z)$] by $\approx 50\%$ relative to LR, cutting the outlier fraction from $\sim 16\%$ to $\sim 12\%$ (Figure 5).

Figure 3: Example super-resolution outputs for two held-out JADES galaxies. Low-resolution prism input (red dashed), super-resolved SR2 output (orange), and medium-resolution reference (purple) over the full 1–5 $\mu$m range. Insets zoom into the [O III] $\lambda\lambda 4959, 5007$ showing deblending of features entirely unresolved at prism resolution.

Figure 4: Emission-line S/N (LR vs. SR) for [O II], H$\beta$, [O III], and H$\alpha$. Points above the one-to-one line indicate improvement; color encodes $\log_{10}$ density.

Figure 5: Predicted vs. true redshifts for LR (left), SR2 (middle), and HR (right) networks. SR2 substantially reduces scatter and outlier fraction, approaching the HR oracle. The true redshifts are obtained from the medium-resolution JADES spectral catalog.

## 5 Discussion and Conclusion

We have presented a physics-informed deep-learning framework that achieves noise-limited spectral super-resolution of JWST/NIRSpec galaxy spectra. By embedding spectroscopic domain knowledge (e.g., redshift inference, line-token self-attention, and a two-branch refinement stage) the model recovers spectroscopic information consistently with underlying physics. The downstream scientific impact is substantial: improved S/N enables higher-fidelity Baldwin-Phillips-Terlevich (BPT) diagramsBaldwin et al. 1981, star-formation rate densities, and gas metallicity gradients. The broad 0.6–5.3 $\mu$m coverage makes the framework applicable to *Euclid* and *Roman Space Telescope* grism surveys delivering spectra for tens of millions of galaxies. Future extensions include incorporating broadband photometry, expanding the training set through upcoming JWST programs, and applying the framework to absorption-line features. Physics-informed spectral super-resolution can bridge the resolution gap between current deep spectroscopic programs and forthcoming wide-area surveys, enabling population-level diagnostics at a fraction of the observational cost.

**Limitations.** The training set of 1,187 spectra introduces mild selection bias toward emission-line galaxies. Performance degrades at $\lesssim 1.5\,\mu$m where both prism and HR data are observational noisy. Transfer to *Euclid* or *Roman* grism data will require domain adaptation for differences in resolution profile and detector noise.

# References

Agostino, C. J., Salim, S., Boquien, M., et al. 2023, Monthly Notices of the Royal Astronomical Society, 526, 4455

Baldwin, A., Phillips, M., & Terlevich, R. 1981, Publications of the Astronomical Society of the Pacific, 93, 817

Biewald, L. 2020, Experiment Tracking with Weights and Biases, `https://www.wandb.com/`

Bunker, A. J., Cameron, A. J., Curtis-Lake, E., et al. 2024, Astronomy & Astrophysics, 690, A288

Curtis-Lake, E., Cameron, A. J., Bunker, A. J., et al. 2026, Monthly Notices of the Royal Astronomical Society, stag836

D'Eugenio, F., Cameron, A. J., Scholtz, J., et al. 2025, The Astrophysical Journal Supplement Series, 277, 4

Eisenstein, D. J., Willott, C., Alberts, S., et al. 2023, arXiv preprint arXiv:2306.02465

Faisst, A. L., Masters, D., Wang, Y., et al. 2018, The Astrophysical Journal, 855, 132

Hemmati, S., Huff, E., Nayyeri, H., et al. 2022, The Astrophysical Journal, 941, 141

Laureijs, R. J., Duvet, L., Sanz, I. E., et al. 2010, in Space Telescopes and Instrumentation 2010: Optical, Infrared, and Millimeter Wave, Vol. 7731, SPIE, 453–458

Parker, L., Lanusse, F., Golkar, S., et al. 2024, Monthly Notices of the Royal Astronomical Society, 531, 4990

Sanjaripour, S., Hemmati, S., Mobasher, B., et al. 2024, The Astrophysical Journal, 977, 202

Spergel, D., Gehrels, N., Breckinridge, J., et al. 2013, arXiv preprint arXiv:1305.5425