TIGHT NON-ASYMPTOTIC INFERENCE VIA SUB-GAUSSIAN INTRINSIC MOMENT NORM

ABSTRACT

In non-asymptotic statistical inferences, variance-type parameters of sub-Gaussian distributions play a crucial role. However, direct estimation of these parameters based on the empirical moment generating function (MGF) is infeasible. To this end, we recommend using a sub-Gaussian intrinsic moment norm [Buldygin and Kozachenko (2000), Theorem 1.3] through maximizing a series of normalized moments. Importantly, the recommended norm can not only recover the exponential moment bounds for the corresponding MGFs, but also lead to tighter Hoeffding's sub-Gaussian concentration inequalities. In practice, intrinsic moment norm can be robustly and consistently estimated via a simple plug-in approach. Our theoretical results are applied to non-asymptotic analysis, including the multi-armed bandit.

1 INTRODUCTION

With the advancement of machine learning techniques, computer scientists have become more interested in establishing rigorous error bounds for desired learning procedures, especially those with finite sample validity (Wainwright, 2019; Zhang & Chen, 2021; Yang et al., 2020). In specific settings, statisticians, econometricians, engineers and physicist have developed non-asymptotic inferences to quantify uncertainty in data; see Romano & Wolf (2000); Chassang (2009); Arlot et al. (2010); Yang et al. (2020); Horowitz & Lee (2020); Armstrong & Kolesár (2021); Zheng & Cheng (2021); Lucas et al. (2008); Owhadi et al. (2013); Wang (2020). Therefore, the concentration-based statistical inference has received a considerable amount of attention, especially for bounded data (Romano & Wolf, 2000; Auer et al., 2002; Hao et al., 2019; Wang et al., 2021; Shiu, 2022) and Gaussian data (Arlot et al., 2010; Duy & Takeuchi, 2022; Bettache et al., 2021; Feng et al., 2021). For example, the Hoeffding's inequality can be applied to construct non-asymptotic confidence intervals based on bounded data[1].

However, in reality, it may be hard to know the support of data or its underlying distribution. In this case, misusing the Hoeffding's inequality (Hoeffding, 1963) for unbounded data will result in a notably loose confidence interval (CI); see Appendix A.1. Hence, it is a common practice to assume that data follow sub-Gaussian distribution (Kahane, 1960). By the Chernoff inequality[2], we have $P(X \geq t) \leq \inf_{s>0}\{\exp\{-st\}E\exp\{sX\}\}, \forall t \geq 0$. Hence, tightness of a confidence interval relies on how we upper bound the moment generating function (MGF) $E\exp\{sX\}$ for all $s > 0$. This can be further translated into the following optimal variance proxy of sub-Gaussian distribution, i.e., *sub-Gaussian parameter* $\sigma_{opt}(X)$.

Definition 1. *A r.v. X is sub-Gaussian (sub-G) with a variance proxy σ^2 [denoted as $X \sim \mathrm{subG}(\sigma^2)$] if its MGF satisfies $E\exp(tX) \leq \exp(\sigma^2 t^2/2)$ for all $t \in \mathbb{R}$. The sub-Gaussian parameter $\sigma_{opt}(X)$ is defined by the optimal variance proxy (Chow, 1966):*

$$\sigma_{opt}^2(X) := \inf\left\{\sigma^2 > 0 : E\exp(tX) \leq \exp\{\sigma^2 t^2/2\}, \quad \forall t \in \mathbb{R}\right\} = 2\sup_{t\in\mathbb{R}} t^{-2}\log[E\exp(tX)]. \quad (1)$$

Note that $\sigma_{opt}^2(X) \geq \mathrm{Var}\, X$; see (16) in Appendix A.2. When $\sigma_{opt}^2(X) = \mathrm{Var}\, X$, it is called strict sub-Gaussianity (Arbel et al., 2020). Based on Theorems 1.5 in Buldygin & Kozachenko (2000), we have

$$P(X \geq t) \leq \exp\left\{-\frac{t^2}{2\sigma_{opt}^2(X)}\right\}, \quad P\left(|\sum_{i=1}^n X_i| \geq t\right) \leq 2\exp\left\{-\frac{t^2}{2\sum_{i=1}^n \sigma_{opt}^2(X_i)}\right\}. \quad (2)$$

---

[1]Recently, Phan et al. (2021) obtained a sharper result than the Hoeffding's inequality for bounded data.
[2]For simplicity, we consider centered random variable (r.v.) with zero mean throughout the paper for all sub-Gaussian r.v..

for independent sub-G r.v.s X and $\{X_i\}_{i=1}^n$. The above inequality (2) provides the tightest upper bound over the form $\mathrm{P}(X > t) \leq \exp(-Ct^2)$ (or $\mathrm{P}(|\sum_{i=1}^n X_i| > t) \leq \exp(-Ct^2)$) for some positive constant C via Chernoff inequality.

Given $\{X_i\}_{i=1}^n \overset{\text{i.i.d.}}{\sim} \mathrm{subG}(\sigma_{opt}^2(X))$, a straightforward application of (2) gives the following non-asymptotic $100(1-\alpha)\%$ CI

$$\mathrm{E}X = 0 \in [\overline{X}_n \pm \sigma_{opt}(X)\sqrt{2n^{-1}\log(2/\alpha)}]. \tag{3}$$

A naive plug-in estimate[3] of $\sigma_{opt}^2(X) := 2\sup_{t\in\mathbb{R}} t^{-2}\log[\mathrm{E}\exp(tX)]$ (Arbel et al., 2020) is

$$\widehat{\sigma}_{opt}^2(X) := 2\sup_{t\in\mathbb{R}} t^{-2}\log[n^{-1}\Sigma_{i=1}^n \exp(tX_i)]. \tag{4}$$

However, two weaknesses of the naive estimator (4) substantially hinder its application: (i) the optimization result is unstable due to the possible non-convexity of the objective function; (ii) exponentially large n is required to ensure the variance term $\mathrm{Var}(n^{-1}\sum_{i=1}^n \exp(tX_i))$ not to explode when t is large. Figure 3 below present simulation evidences.

On the other hand, we are aware of other forms of variance-type parameter. For instance, van der Vaart & Wellner (1996) introduced the Orlicz norm as $\|X\|_{w_2} := \inf\{c > 0 : \mathrm{E}\exp\{|X|^2/c^2\} \leq 2\}$, frequently used in empirical process theory. Additionally, Vershynin (2010) suggested a norm based on the scale of moments as $\|X\|_{\psi_2} := \max_{k\geq 2} k^{-1/2}(\mathrm{E}|X|^k)^{1/k}$ in Page 6 of Buldygin & Kozachenko (2000). However, as shown in Table 1 and Appendix A.2.1, both types of norm fail to deliver sharp probability bounds even for strict sub-G distributions, such as the standard Gaussian distribution and symmetric beta distribution.

Table 1: Comparison of sub-Gaussian norms $\|\cdot\|_*$ for centralized and symmetric X.

$\|\cdot\|_*$-norm	variance bound	sharp tail for $\mathrm{P}(|X| \geq t)$	sharp MGF bound	easy to estimate
$\sigma_{opt}(X)$	$\sigma_{opt}^2(X) \geq \mathrm{Var}\,X$	Yes $[2\exp\{-\frac{t^2}{2}/\sigma_{opt}^2(X)\}]$	Yes $[\exp\{\sigma_{opt}^2(X)\frac{t^2}{2}\}]$	No
$2\|X\|_{w_2}$	$4\|X\|_{w_2}^2 \geq \mathrm{Var}\,X$	Yes $[2\exp\{-\frac{t^2}{2}/(2\|X\|_{w_2}^2)\}]$	No $[\exp\{(2\|X\|_{w_2})^2\frac{t^2}{2}\}]$	No
$\sqrt{2}\|X\|_{\psi_2}$	$2\|X\|_{\psi_2}^2 \geq \mathrm{Var}\,X$	No $[2\exp\{-\frac{t^2}{2}/(2e\|X\|_{\psi_2}^2)\}]$	No $[\exp\{(4\sqrt{e}\|X\|_{\psi_2})^2\frac{t^2}{2}\}]$	Yes
$\|X\|_G$ (Def. 2)	$\|X\|_G^2 \geq \mathrm{Var}\,X$	Yes $[2\exp\{-\frac{t^2}{2}/\|X\|_G^2\}]$	Yes $[\exp\{\|X\|_G^2\frac{t^2}{2}\}]$	Yes

1.1 CONTRIBUTIONS

In light of the above discussions, we advocate the use of the intrinsic moment norm in the Definition 2 in the construction of tight non-asymptotic CIs. There are two specific reasons: (i) it approximately recovers tight inequalities (2); (ii) it can be estimated friendly (with a closed form) and robustly.

The following definition 2 is from Page 6 and Theorem 1.3 in Buldygin & Kozachenko (2000).

Definition 2 (Intrinsic moment norm). $\|X\|_G := \max_{k\geq 1} \left[\frac{2^k k!}{(2k)!}\mathrm{E}X^{2k}\right]^{1/(2k)} = \max_{k\geq 1} \left[\frac{1}{(2k-1)!!}\mathrm{E}X^{2k}\right]^{1/(2k)}$.

From the sub-G characterization (see Theorem 2.6 in Wainwright (2019)), $\|X\|_G < \infty$ iff $\sigma_{opt}(X) < \infty$ *for any zero-mean r.v.* X. Hence, the finite intrinsic moment norm of a r.v. X ensures sub-Gaussianity (satisfying Definition 1).

Our contributions in this paper can be summarized as follows.

1. By the use of $\|X\|_G$, we achieve the Hoeffding-type inequalities with sharper constants under asymetric distribution; see Theorem 1(b).

2. Compared to the normal approximation based on Berry-Esseen (B-E) bounds, our results are more applicable to data of extremely small sample size. We illustrate Bernoulli observations with the comparison of two types of CIs based on the B-E-corrected CLT and Hoeffding's inequality in Figure 1; see Appendix A for details.

3. We propose a novel method called *sub-Gaussian plot* for judging whether the data are sub-Gaussian.

4. Finally, we employ the intrinsic moment norm in the non-asymptotic inference for a bandit problem: Bootstrapped UCB-algorithm for multi-armed bandits. This algorithm is shown to achieve feasible error bounds and competitive cumulative regret on unbounded sub-Gaussian data by the intrinsic moment norm estimation.

---

[3]We point out that a conservative and inconsistent estimator $2\inf_{t\in\mathbb{R}} \log(n^{-1}\sum_{i=1}^n \exp(tX_i))/t^2$ was proposed in statistical physics literature (Wang, 2020).

Figure 1: CIs via Hoeffding's inequality (red line) and B-E-corrected CLT (blue line). It describes a deficiency of B-E-corrected CLT under small sample, and it suggests that a simple Hoeffding's inequality can even perform better.

2 PROPERTIES OF INTRINSIC MOMENT NORM

In this section, we characterize two important properties of the intrinsic moment norm that are used in constructing non-asymptotic confidence intervals.

2.1 BASIC PROPERTIES

Lemma 1 below establishes that the intrinsic moment norm is estimable.

Lemma 1. *For sub-G X, we have* $\arg\max_{m \in 2\mathbb{N}} \left[\frac{\mathrm{E}X^m}{(m-1)!!} \right]^{1/m} < \infty$, *where* $2\mathbb{N} := \{2, 4, \cdots\}$ *is an even number set.*

Lemma 1 ensure that for any sub-Gaussian variable X, its intrinsic moment norm can be computed as

$$\|X\|_G := \max_{m \in 2\mathbb{N}} \left[\frac{\mathrm{E}X^m}{(m-1)!!} \right]^{1/m} = \max_{1 \le k \le k_X} \left[\frac{\mathrm{E}X^{2k}}{(2k-1)!!} \right]^{1/(2k)} \text{ with some finite } k_X < \infty.$$

This is an important property that other norms may not have. For example, the $\|X\|_{\psi_2} := \max_{k \ge 2} k^{-1/2}(\mathrm{E}|X|^k)^{1/k}$ for Gaussian X achieves its optimal point at $k = \infty$; see Example 3 in Appendix A.2.1. As for $\sigma^2_{opt}(X) := 2\sup_{t \in \mathbb{R}} \frac{\log[\mathrm{E}\exp(tX)]}{t^2}$, it is unclear that its value can be achieved at a finite t.

Note that if $k_X = 1$, one has $\|X\|_G^2 = \mathrm{Var}(X)$.

In the end, we present an example in calculating the values of k_X. Denote $\mathrm{Exp}(1)|_{[0,M]}$ as the truncated standard exponential distribution on $[0, M]$ with the density as $f(x) = \frac{e^{-x}}{\int_0^M e^{-x}\,dx} 1_{\{x \in [0,M]\}}$.

Example 1. *a.* $X \sim U[-a, a]$, $k_X = 1$ *for any* $a \in \mathbb{R}$; *b.* $X \sim \mathrm{Exp}(1)|_{[0,2.75]} - \mathrm{E}\,\mathrm{Exp}(1)|_{[0,2.75]}$, $k_X = 2$; *c.* $X \sim \mathrm{Exp}(1)|_{[0,3]} - \mathrm{E}\,\mathrm{Exp}(1)|_{[0,3]}$, $k_X = 3$. *Indeed, for any fixed* $k_0 \in \mathbb{N}$*, we can construct a truncated exponential r.v.* $X := \mathrm{Exp}(1)|_{[0,M]}$ *such that* $k_X = k_0$ *by properly adjusting the truncation level* M.

2.2 CONCENTRATION FOR SUMMATION

In what follows, we will show another property of $\|X\|_G$ that it recovers nearly tight MGF bounds in Definition 1. More powerfully, it enables us to derive the sub-G Hoeffding's inequality (2).

Theorem 1. *Suppose that* $\{X_i\}_{i=1}^n$ *are independent r.v.s with* $\max_{i \in [n]} \|X_i\|_G < \infty$. *We have*

(a). *If X_i is symmetric about zero, then* $\mathrm{E}\exp\{tX_i\} \le \exp\{t^2 \|X_i\|_G^2 /2\}$ *for any* $t \in \mathbb{R}$, *and*

$$\mathrm{P}\left(|\textstyle\sum_{i=1}^n X_i| \ge s\right) \le 2\exp\{-s^2/[2\textstyle\sum_{i=1}^n \|X_i\|_G^2]\}, \quad s \ge 0.$$

(b). *If X_i is not symmetric, then* $\mathrm{E}\exp\{tX_i\} \le \exp\{(17/12)t^2 \|X_i\|_G^2 /2\}$ *for any* $t \in \mathbb{R}$, *and*

$$\mathrm{P}\left(|\textstyle\sum_{i=1}^n X_i| \ge s\right) \le 2\exp\{-(12/17)s^2/[2\textstyle\sum_{i=1}^n \|X_i\|_G^2]\}, \quad s \ge 0.$$

Theorem 1(a) is an existing result in Theorem 2.6 of Wainwright (2019). For Theorem 1(b), we obtain $\sqrt{17/12} \approx 1.19$, while Lemma 1.5 in Buldygin & Kozachenko (2000) obtained $\mathrm{E}\exp\{tX_i\} \leq \exp\left\{\frac{t^2}{2}(\sqrt[4]{3.1}\|X_i\|_G)^2\right\}$ for $t \in \mathbb{R}$ with $\sqrt[4]{3.1} \approx 1.32$. A technical reason for this improvement is that $\|\cdot\|_G$ does not need Stirling's approximation for attaining the sharper MGF bounds when expanding the exponential function by Taylor's formula. In Appendix C, we gives some comparisons in Figure 6 with $\sigma_{opt}^2(X)$, $17\|X\|_G^2/12$, $2e\|X\|_{\psi_2}^2$ and $\mathrm{Var}\, X$ in the tail bound constants of Table 1, to show the tightness of Theorem 1(b), when X is Bernoulli or Beta distribution.

Before estimating $\|X\|_G$, the first step is to verify X is indeed sub-G given its i.i.d. copies $\{X_i\}_{i=1}^n$. Corollary 7.2 (b) in Zhang & Chen (2021) shows

$$\mathrm{P}(\max_{1\leq i\leq j} X_i \leq \sigma_{opt}(X)\sqrt{2(\log j + t)}) \geq 1 - \exp\{-t\}, \tag{5}$$

which implies $\max_{1\leq i\leq j} X_i = O_{\mathrm{P}}(\sqrt{\log j})$. Moreover, we will show the above rate is indeed sharp for a class of unbound sub-G r.v.s characterized by the *lower intrinsic moment norm* below.

Definition 3 (Lower intrinsic moment norm). *The lower intrinsic moment norm for a sub-G X is defined as*

$$\|X\|_{\tilde{G}} := \min_{k\geq 1}\{[(2k-1)!!]^{-1}\mathrm{E}X^{2k}\}^{1/(2k)}.$$

By the method in Theorem 1 of Zhang & Zhou (2020), we obtain the following the tight rate result with a lower bound.

Theorem 2. *(a). If $\|X\|_{\tilde{G}} > 0$ for i.i.d. symmetric sub-G r.v.s $\{X_i\}_{i=1}^n \sim X$, then with probability at least $1 - \delta$*

$$\frac{\|X\|_{\tilde{G}}/\|X\|_G}{2\sqrt{2\|X\|_G^2/\|X\|_{\tilde{G}}^2 - 1}}\sqrt{\log n - \log C^{-2}(X) - \log\log\left(\frac{2}{\delta}\right)} \leq \max_{1\leq i\leq n} \frac{X_i}{\|X\|_G} \leq \sqrt{2[\log n + \log\left(\frac{2}{\delta}\right)]},$$

where $C(X) < 1$ is constant defined in Lemma 2 below; (b) if X is bounded variable, then $\|X\|_{\tilde{G}} = 0$.

Figure 2: sub-Gaussian plot of standard Gaussian and standard Exponential distribution.

The upper bound follows from the proof of (5) similarly. The proof of lower bound relies on the sharp reverse Chernoff inequality from Paley–Zygmund inequality (see Paley & Zygmund (1932)).

Lemma 2 (A reverse Chernoff inequality). *Suppose $\|X\|_{\tilde{G}} > 0$ for a symmetric sub-G r.v. X. For $t > 0$, then*

$$\mathrm{P}(X \geq t) \geq C^2(X)\exp\{-4[2\|X\|_G^2/\|X\|_{\tilde{G}}^4 - \|X\|_{\tilde{G}}^{-2}]t^2\},$$

where $C(X) := \left(\frac{\|X\|_{\tilde{G}}^2}{4\|X\|_G^2 - \|X\|_{\tilde{G}}^2}\right)\left(\frac{4\|X\|_G^2 - 2\|X\|_{\tilde{G}}^2}{4\|X\|_G^2 - \|X\|_{\tilde{G}}^2}\right)^{2[2\|X\|_G^2/\|X\|_{\tilde{G}}^2 - 1]} \in (0, 1).$

Theorem 1 of Zhang & Zhou (2020) does not optimize the constant in Paley–Zygmund inequality. In contrast, our Lemma 2 has an optimal constant; see Appendix C for details.

By Theorem 2, we propose a novel *sub-Gaussian plot* check whether i.i.d data $\{X_i\}_{i=1}^n$ follow an unbound sub-G variable. Specifically, we plot the order statistics $\{\max_{1\leq i\leq j} X_i^*\}_{j=1}^n$ on the plane coordinate axis, where x axis represents $\sqrt{\log j}$ and y axis the value of $\max_{1\leq i\leq j} X_i^*$. Here for each j, $\{X_i^*\}_{i=1}^j$ are independently sampled from the empirical distribution $\mathbb{F}_n(x) = \frac{1}{n}\sum_{i=1}^n 1(X_i \leq x)$ of $\{X_i\}_{i=1}^n$. We check whether those points have a linear tendency: the more they are close to the tendency of a beeline, the more we can trust the data are sub-Gaussian.

The following graphs show *sub-Gaussian plot* of $N(0,1)$ and $\mathrm{Exp}(1)$. It can be seen that sub-Gaussian plot of $N(0,1)$ shows linear tendency, while $\mathrm{Exp}(1)$ shows quadratic tendency. For the quadratic tendency, we note that if $\{X_i\}_{i=1}^n$ have heavier tails such as sub-exponentiality, then $\max_{1\leq i\leq j} X_i = O_\mathrm{P}(\log j)$ instead of the order $O(\sqrt{\log j})$; see Corollary 7.3 in Zhang & Chen (2021).

3 ESTIMATION OF THE INTRINSIC MOMENT NORM

A first thought to estimate $\|X\|_G$ is by the plug-in approach. Although k_X is proven to be finite in Lemma 1, its (possibly large) exact value is still unknown in practice. Instead, we use a non-decreasing *index sequence* $\{\kappa_n\}$ to replace k_X in the estimation. Hence, we suggest a plug-in feasible estimator

$$\widehat{\|X\|}_G = \max_{1\leq k\leq \kappa_n} \left[\frac{1}{(2k-1)!!}\frac{1}{n}\sum_{i=1}^n X_i^{2k}\right]^{1/(2k)}. \tag{6}$$

Deriving the non-asymptotic property of the $\widehat{\|X\|}_G$ is not an easy task: the maximum point $\hat{k}(\kappa_n) := \arg\max_{1\leq k\leq \kappa_n}\left[\frac{1}{(2k-1)!!}\frac{1}{n}\sum_{i=1}^n X_i^{2k}\right]^{1/(2k)}$ will change with the sample size n even κ_n is fixed.

To resolve this, we first examine the oracle estimator defined as follows

$$\widetilde{\|X\|}_G = \left[\frac{1}{(2k_X-1)!!}\frac{1}{n}\sum_{i=1}^n X_i^{2k_X}\right]^{\frac{1}{2k_X}}. \tag{7}$$

It is trivial to show that $\widetilde{\|X\|}_G \xrightarrow{a.s.} \|X\|_G$. Hence, based on Orlicz norm $\|Y\|_{\psi_\theta} := \inf\{t > 0 : \mathrm{E}\exp\{|Y|^\theta/t^\theta\} \leq 2\}$ of sub-Weibull r.v. Y with $\theta > 0$ (Hao et al., 2019; Zhang & Wei, 2022), the following result gives the non-asymptotic concentration of $\widetilde{\|X\|}_G$.

Proposition 1. *Suppose* $\{X_i\}_{i=1}^n \overset{i.i.d.}{\sim} X$ *and X satisfies* $\|X\|_{\psi_{1/k_X}} < \infty$, *then for any $t > 0$,*

$$\mathrm{P}\left(\left|\widetilde{\|X\|}_G^{2k_X} - \|X\|_G^{2k_X}\right| \leq 2e\|X\|_{\psi_{1/k_X}} C(k_X^{-1})\left\{\sqrt{\frac{t}{n}} + \gamma^{2k_X} A(k_X^{-1})\frac{t^{k_X}}{n}\right\}\right) \geq 1 - 2e^{-t},$$

where the constant $\gamma \approx 1.78$, and the constant functions $C(\cdot)$ and $A(\cdot)$ are defined in Appendix C.

The exponential-moment condition $\|X\|_{\psi_{1/k_X}} < \infty$ is too strong for the error bound of $\widetilde{\|X\|}_G^{2k_X} - \|X\|_G^{2k_X}$ in Proposition 1, although it has exponential decay probability $1 - 2\exp(-t)$.

Except for the direct plug-in estimator, for estimators that are robust to outliers, here we resort to the median-of-means (MOM, Page244 in Nemirovskij & Yudin (1983)) plug-in estimator of intrinsic moment norm. Let m and b be a positive integer such that $n = mb$ and let $B_1, \ldots, B_b$ be a partition of $[n]$ into blocks of equal cardinality m. For any $s \in [b]$, let $\mathrm{P}_m^{B_s} X = m^{-1}\sum_{i\in B_s} X_i$ for independent data $\{X_i\}_{i=1}^n$. The MOM version intrinsic moment norm estimator is defined as

$$\widehat{\|X\|}_{b,G} := \max_{1\leq k\leq \kappa_n} \operatorname*{med}_{s\in[b]}\left\{\left[(2k-1)!!\right]^{-1}\mathrm{P}_m^{B_s} X^{2k}\right]^{1/(2k)}\right\}. \tag{8}$$

As stated in Proposition 1, the naive plug-in estimator $\widehat{\|X\|}_G = \widehat{\|X\|}_{1,G}$ is not robust. MOM estimators (8) with $b \gg 1$ have two merits: (a) it only needs finite moment conditions, but the exponential concentration bounds are still achieved; (b) it permits some outliers in the data. Non-asymptotic inferences require to bound for $\|X\|_G$ exactly by a feasible estimator $\widehat{\|X\|}_{b,G}$ up to a sharp constants. Next, we establish a high-probability upper bound for the estimated norm, if the data has finite moment assumptions as follows.

Figure 3: DE represents the naive plug-in estimator (6), MOM represents the MOM estimator (8), and OP is the estimator plug-in naive estimator (4) for the optimal variance proxy.

- (M.1) Suppose $\sqrt{\operatorname{Var} X^{2k}} \leq \sigma_k^k$, $1 \leq k \leq \kappa_n$ for a sequence $\{\sigma_k\}_{k \geq 1}$;

- (M.2) Define $\underline{g}_{k,m}(\sigma_k)$ and $\bar{g}_{k,m}(\sigma_k)$ as the sequences tending to zero as $m \to \infty$

$$\left[\mathbb{E} X^{2k}/(2k-1)!!\right]^{1/(2k)}(1 - \bar{g}_{k,m}(\sigma_k)) \leq \max_{1 \leq k \leq \kappa_n}\left[-2m^{-1/2}\sigma_k^k/(\mathbb{E} X^{2k}) + \mathbb{E} X^{2k}/(2k-1)!!\right]^{1/(2k)}; \quad (9)$$

$$[2m^{-1/2}\sigma_k^k/(\mathbb{E} X^{2k}) + 1]^{1/(2k)} = 1 + \underline{g}_{k,m}(\sigma_k) \text{ for any } m \in \mathbb{N} \text{ and } 1 \leq k \leq \kappa_n \text{ respectively.} \quad (10)$$

(M.1) is due to Chebyshev's inequality applying to the empirical mean of sub-sample. (M.2) is technical moment conditions in the proof, and the remainder terms in (9), (10) are from Taylor expansions. Below, the result concerns for estimated norm bound for finite sample size.

Theorem 3 (Guaranteed coverage). *Under (M.1), (M.2) with (9), we have for $\kappa_n \geq \kappa_X$*

$$\mathrm{P}\left\{\|X\|_G \leq [1 - \max_{1 \leq k \leq \kappa_n} \bar{g}_{k,m}(\sigma_k)]^{-1}\widehat{\|X\|}_{b,G}\right\} > 1 - \kappa_n \cdot \exp(-b/8);$$

and $\mathrm{P}\{\|X\|_G \geq [1 + \max_{1 \leq k \leq \kappa_n} \underline{g}_{k,m}(\sigma_k)]^{-1}\widehat{\|X\|}_{b,G}\} > 1 - \kappa_n \cdot \exp(-b/8)$ *by (M.1), (M.2) with (10).*

Theorem 3 ensures the consistency of the estimator $\widehat{\|X\|}_{b,G}$ when $\kappa_n \geq k_X$ under enough sample. When the data is i.i.d. Gaussian vector, Proposition 4.1 in Auer et al. (2002) also gave a high-probability estimated upper bound for ℓ_p-norm of the vector of Gaussian standard deviations, our result is for intrinsic moment norm. In practice, the block number b can be taken by the adaptation method based on the Lepski method (Depersin & Lecué, 2022). To guarantee high probability events in Theorem 3, it is required that the index sequence κ_n should not be very large for fixed b. The larger κ_n needs larger b in blocks $B_1, \ldots, B_b$. In the simulation, we will see that an increasing index sequence κ_n with slow rate will lead a good performance.

Here, we compare our two estimators (6) and (8), as well as the estimator (4). We consider the standard Gaussian and Rademacher variable distributed X, in the two case we have $\|X\|_G^2 = \sigma_{opt}^2(X) = \text{Var}(X) = 1$. The following figure shows the performance of three estimators under sample $n = 10$ to 1000 with κ_n just chosen as $\lceil \log n \rceil$. For the MOM method, we use five blocks in this simple setting. For a more complex case, one can use Lepski's method to choose b (see Page & Grünewälder (2021)), but some considerable computation cost may be introduced. From Figure 3, we know that the performance of the MOM estimator is best, while the naive estimator (4) is worst. For the high-quality data of extremely small sample size, we can apply the leave-one-out Hodges-Lehmann method (Rousseeuw & Verboven, 2002) in the estimation for further numerical improvement, and the details can be seen in Appendix B.

4   APPLICATION IN MULTI-ARMED BANDIT PROBLEM

In the multi-armed bandit problem (MAB), a player chooses between K different slot machines (an K-armed bandit), each with a different unknown random reward r.v.s $\{Y_k\}_{k=1}^K \subseteq \mathbb{R}$, while each realization of a fixed arm k is independent and shares the same distribution, i.e., the rewards from each arm are i.i.d. sample. We assume the rewards are sub-Gaussian, i.e.

$$\|Y_k - \mu_k\|_G \leq C < \infty, \qquad k \in [K]. \tag{11}$$

Our goal is to find the best arm with the largest expected reward, say Y_{t*}, by pulling arms. In each round $t \in [T]$, the player pulls an arm (an action) $A_t \in [K]$. Conditioning on $\{A_t = k\}$, we define the observed reward $\{Y_{k,t}\}_{t \in [T]} \overset{\text{i.i.d.}}{\sim} P_k$. The goal of the exploration in MAB is to minimize the cumulative regret after T steps: $\text{Reg}_T(Y, A) := \sum_{t=1}^T (\mu_{t^\star} - \mu_{A_t})$, i.e. the performance of any exploration strategy $\{A_t\}_{t \in [T]}$. The exploration performance is better, if we have smaller $\text{Reg}_T(Y, A)$. Without loss of generality, we assume $t^\star = 1$. Since $\text{Reg}_T(Y, A)$ is a random quantity, we seek to evaluate the expected bounds from the decomposition (see Lemma 4.5 in Lattimore & Szepesvári (2020)),

$$\text{Reg}_T := \mathbb{E}\,\text{Reg}_T(Y, A) = \sum_{k=1}^K \Delta_k \mathbb{E}\Big[\sum_{t=1}^T \mathbb{1}\{A_t = k\}\Big], \tag{12}$$

where $\mathbb{E}$ is taken on the randomness of the player's actions $\{A_t\}_{t \in [T]}$, and $\Delta_k = \mu_1 - \mu_k$ is the sub-optimality gap for arm $k \in [K]/\{1\}$. The upper bound of Reg_T is called problem-independent if the regret bound depends on the distribution of the data and does not rely on the gap Δ_k.

For each iteration t, let $T_k(t) := \text{card}\{1 \leq \tau \leq t : A_\tau = k\}$ be the number of pull for arm k until time t during the bandit process. Then if we define $\overline{Y}_{T_k(t)} := \frac{1}{T_k(t)} \sum_{\tau \leq t, A_\tau = k} Y_{k,\tau}$ as the running average of the rewards of arm k at time t. Suppose we obtain a $100(1 - \delta)\%$ CI $[\overline{Y}_{T_k(t)} - c_k(t), \overline{Y}_{T_k(t)} + c_k(t)]$ for μ_k from a tight concentration inequality, where $c_k(t)$ decreases with time t. Therefore, we confidently reckon that the reward of arm k is $\overline{Y}_{T_k(t)} + c_k(t)$, and play the arm $A_t = k$, hoping to maximize the reward with a high probability for finite t. This is upper confidence bound (UCB, Auer et al. (2002)) algorithms.

In details, suppose that $Y_k - \mu_k$ is symmetric around zero, by one-side version of Theorem 1, the (11) implies that for all k and all t, $\text{P}(\overline{Y}_{T_k(t)} > \mu_k + \|Y_k - \mu_k\|_G \sqrt{\frac{2}{T_k(t)} \log \frac{1}{\delta}}) \leq \delta$. Let sub-sample size m_k and block size b_k be positive integer such that $T_k(t) = m_k b_k$ for MOM estimators $\widehat{\|Y_k - \mu_k\|}_{b_k,G}$ in Section 3. Theorem 3 (a) guarantee that true norms can be replaced by MOM-estimated norms such that

$$\text{P}\left(\overline{Y}_{T_k(t)} \leq \mu_k + \frac{\widehat{\|Y_k - \mu_k\|}_{b_k,G}}{1 - o(1)} \sqrt{\frac{2}{T_k(t)} \log \frac{1}{\delta}}\right) \geq 1 - \delta - k_{Y_k} \cdot \exp(-b_k/8).$$

If the UCB algorithm is correctly applied, for a finite $T_k(t)$, with high probability, we will pull the best arm.

In practice, we nearly do not know any knowledge about the data. As a flexible way of uncertainty qualification, the multiplier bootstrap (Arlot et al., 2010) enables mimicking the non-asymptotic properties of the target statistic by reweighing its summands of the centralized empirical mean. The multiplier bootstrapped quantile for the i.i.d. observation $\mathbf{Y}_n := \{Y_i\}_{i=1}^n$ is the $(1 - \alpha)$-quantile of the distribution of $n^{-1}\sum_{i=1}^n w_i(Y_i - \overline{Y}_n)$, which is defined as

$$q_\alpha(\mathbf{Y}_n - \overline{Y}_n, \mathbf{w}) := \inf\{x \in \mathbb{R} \,|\, \mathrm{P}_w(n^{-1}\textstyle\sum_{i=1}^n w_i(Y_i - \overline{Y}_n) > x) \le \alpha\},$$

where $\mathbf{w} := \{w_i\}_{i=1}^n$ are bootstrap random weights independent of $\mathbf{Y}_n$. We denote the statistics $\widehat{\varphi}_G(\mathbf{Y}_n)$ as something satisfying $\mathrm{P}_{\mathbf{Y}_n}(|\overline{Y}_n - \mathrm{E}Y_1| \ge \widehat{\varphi}_G(\mathbf{Y}_n)) \le \alpha$.

Algorithm 1: Bootstrapped UCB

Input: $\widehat{\varphi}_G(\mathbf{Y}_{T_k(t)})$ is given by (13).

    for $t = 1, \ldots, K$ **do**
    | Pull each arm once to initialize the algorithm.
    end
    for $t = K + 1, \ldots, T$ **do**
      Set a confidence level $\alpha \in (0, 1)$.
      Calculate the boostrapped quantile $q_{\alpha/2}(\mathbf{Y}_{T_k(t)} - \overline{Y}_{T_k(t)}, \mathbf{w})$ with the Rademacher bootstrapped weights $\mathbf{w}$ independent with any Y.
      Pull the arm
$$A_t = \underset{k \in [K]}{\operatorname{argmax}} \, \mathrm{UCB}_k(t) := \underset{k \in [K]}{\operatorname{argmax}} \left(\overline{Y}_{T_k(t)} + q_{\alpha/2}(\mathbf{Y}_{T_k(t)} - \overline{Y}_{T_k(t)}, \mathbf{w}) + \widehat{\varphi}_G(\mathbf{Y}_{T_k(t)}) \right).$$
      Receive reward Y_{A_t}.
    end

Motivated by Hao et al. (2019), we design Algorithm 1 based on some estimators of the UCB. It guarantees a relatively small regret by bootstrapped threshold $q_{\alpha/2}(\mathbf{Y}_{T_k(t)} - \overline{Y}_{T_k(t)}, \mathbf{w})$ adding a concentration based second-order correction $\widehat{\varphi}_G(\mathbf{Y}_{T_k(t)})$ that is specified in Theorem 3. In the following regret bounds, we assume the mean reward from the k-th arm μ_k is known. In practice, it can be replaced by a robust estimator such as the MOM estimator, and we obtain the following results.

Theorem 4. *Consider a K-armed sub-G bandit under (11) and suppose that $Y_k - \mu_k$ is symmetric around zero. For any round T, according to moment conditions in Theorem 3, choosing $\widehat{\varphi}_G(\mathbf{Y}_{T_k(t)})$ as*

$$\widehat{\varphi}_G(\mathbf{Y}_{T_k(t)}) = \sqrt{\frac{2\log(4/\alpha)}{T_k(t)}} \frac{\|\widehat{Y_k - \mu_k}\|_{b_k, G}}{1 - T_k^{-1/2}(t)} \tag{13}$$

as a re-scaled version of MOM estimator $\|\widehat{Y_k - \mu_k}\|_{b_k, G}$ with block number b_k satisfying the moment assumptions C[UCB1] and C[UCB2] in Appendix C. Fix a confidence level $\alpha = 4/T^2$, if the player pull an arm $A_t \in [K]$ according to Algorithm 1, then we have the problem-dependent regret of Algorithm 1 is bounded by

$$\mathrm{Reg}_T \le 16(2 + \sqrt{2})^2 C^2 \log T \textstyle\sum_{k=2}^K \Delta_k^{-1} + (4T^{-1} + 2T^{-25-16\sqrt{2}} + 8) \textstyle\sum_{k=2}^K \Delta_k,$$

where Δ_k is the sub-optimality gap. Moreover, let $\mu_1^* := \max_{k_1 \in [K]} \mu_{k_1} - \min_{k_2 \in [K]} \mu_{k_2}$ be the range over the rewards, the problem-independent regret

$$\mathrm{Reg}_T \le 8(2 + \sqrt{2})C\sqrt{TK \log T} + (4T^{-1} + 2T^{-25-16\sqrt{2}} + 8)K\mu_1^*.$$

From Theorem 4, we know that the regret of our method achieve minimax rate $\log T$ for a problem-dependent problem and $\sqrt{KT}$ for a problem-independent case (see Tao et al. (2022)), so Algorithm 1 can be seen as an optimal algorithm.

Next, we give an simulation for Theorem 4 in two sub-G cases to verify the performance of estimated norms. Similar to Hao et al. (2019); Wang et al. (2020), we design the three methods as follows:

    1. Use our method $\widehat{\varphi}(\mathbf{Y}_{T_k(t)})$ with *Estimated Norm* in Theorem 4;

2. Use *Asymptotic Naive varphi* $\widetilde{\varphi}(\mathbf{Y}_{T_k(t)})$ satisfying $\mathrm{P}\big(|\overline{Y}_{T_k(t)} - \mu_k| \leq \widetilde{\varphi}(\mathbf{Y}_{T_k(t)})\big) \to \alpha$ by CLT, i.e. $\widetilde{\varphi}(\mathbf{Y}_{T_k(t)}) = \widehat{\sigma}_k \Phi^{-1}(1 - \alpha/2)/\sqrt{T_k(t)}$ with $\widehat{\sigma}_k = \sqrt{\frac{1}{T_k(t)} \sum_{\tau \leq t, A_\tau = k}(Y_{k,\tau} - \overline{Y}_{T_k(t)})^2}$ as the estimated standard deviation;

3. Regard all the unbounded rewards as bounded r.v. and use Hoeffding's inequality (*wrongly use Hoeffding's inequality*) to construct φ, i.e. $\check{\varphi}(\mathbf{Y}_{T_k(t)}) = \big[\max\{\mathbf{Y}_{T_k(t)}\} - \min\{\mathbf{Y}_{T_k(t)}\}\big]\sqrt{\frac{\log(2/\alpha)}{2T_k(t)}}$.

And for our detailed MAB simulation, we consider as follows, in each case, the number of arms is assigned as $K = 5$, and $(\mu_1, \ldots, \mu_5) = (0.1, 0.05, 0.02, 0.01, 0.01)^\top$,

$$\text{EG1. } Y_k \sim N(\mu_k, \mu_k^2); \qquad \text{EG2. } Y_k \sim 0.9N(\mu_k, \mu_k) + 0.1N(\mu_{k+1}, \mu_{k+1}) \text{ where } \mu_6 := \mu_5.$$

Figure 4: The regret of MAB with sub-G rewards under three methods. x-axis represents the round and y-axis is the cumulative regret.

As we can see, EG1 and EG2 are both sub-Gaussian rewards. In the simulation, μ_k is assigned sightly small for bounded $\max_k \Delta_k$, which is a standard-setting in the MAB problem (see the condition of Corollary 1 in Wang et al. (2020) for instance). The simulation results are shown in Figure 4, which illustrates that our method outperforms the other two methods under the unbounded sub-Gaussian rewards and small sample $T \in [1, 150]$. In the Gaussian case, Algorithm 1 can also give better results compared with the CLT-based UCB when the round is relatively small. For the Gaussian mixture case, the algorithm based on the intrinsic moment norm has smaller regret than the other two methods under $T \in [1, 800]$.

REFERENCES

Julyan Arbel, Olivier Marchal, and Hien D Nguyen. On strict sub-gaussianity, optimal proxy variance and symmetry for bounded random variables. *ESAIM: Probability and Statistics*, 24:39–55, 2020.

Sylvain Arlot, Gilles Blanchard, Etienne Roquain, et al. Some nonasymptotic results on resampling in high dimension, i: confidence regions. *The Annals of Statistics*, 38(1):51–82, 2010.

Timothy B. Armstrong and Michal Kolesár. Finite-sample optimal estimation and inference on average treatment effects under unconfoundedness. *Econometrica*, 89(3):1141–1177, 2021.

Peter Auer, Nicolo Cesa-Bianchi, and Paul Fischer. Finite-time analysis of the multiarmed bandit problem. *Machine learning*, 47(2):235–256, 2002.

Nayel Bettache, Cristina Butucea, and Marianne Sorba. Fast nonasymptotic testing and support recovery for large sparse toeplitz covariance matrices. *Journal of Multivariate Analysis*, pp. 104883, 2021.

Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities: A nonasymptotic theory of independence*. Oxford university press, 2013.

Valeriĭ Vladimirovich Buldygin and IU V Kozachenko. *Metric characterization of random variables and random processes*, volume 188. American Mathematical Soc., 2000.

Sylvain Chassang. Non-asymptotic tests of model performance. *Economic Theory*, 41(3):495–514, 2009.

YS Chow. Some convergence theorems for independent random variables. *The Annals of Mathematical Statistics*, 37 (6):1482–1493, 1966.

Jules Depersin and Guillaume Lecué. Robust sub-gaussian estimation of a mean vector in nearly linear time. *The Annals of Statistics*, 50(1):511–536, 2022.

Vo Nguyen Le Duy and Ichiro Takeuchi. Exact statistical inference for the wasserstein distance by selective inference. *Annals of the Institute of Statistical Mathematics*, 2022.

Yihao Feng, Ziyang Tang, na zhang, and qiang liu. Non-asymptotic confidence intervals of off-policy evaluation: Primal and dual bounds. In *International Conference on Learning Representations*, 2021.

Evarist Giné and Richard Nickl. *Mathematical foundations of infinite-dimensional statistical models*. Cambridge university press, 2016.

Botao Hao, Yasin Abbasi Yadkori, Zheng Wen, and Guang Cheng. Bootstrapping upper confidence bound. In *Advances in Neural Information Processing Systems*, volume 32, pp. 12123–12133, 2019.

Tim Hesterberg. Bootstrap. *Wiley Interdisciplinary Reviews: Computational Statistics*, 3(6):497–526, 2011.

Wassily Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association*, 58(301):13–30, 1963.

Joel Horowitz and Sokbae Lee. *Inference in a class of optimization problems: Confidence regions and finite sample bounds on errors in coverage probabilities*. Cemmap, Centre for Microdata Methuods and Practice, The Institute for Fiscal., 2020.

Jean-Pierre Kahane. Propriétés locales des fonctions à séries de fourier aléatoires. *Studia Mathematica*, 19(1):1–25, 1960.

Michael Kearns and Lawrence Saul. Large deviation methods for approximate probabilistic inference. In *Proceedings of the Fourteenth conference on Uncertainty in artificial intelligence*, pp. 311–319, 1998.

Aryeh Kontorovich. Concentration in unbounded metric spaces and algorithmic stability. In *International Conference on Machine Learning*, pp. 28–36. PMLR, 2014.

Tor Lattimore and Csaba Szepesvári. *Bandit algorithms*. Cambridge University Press, 2020.

Leonard J Lucas, H Owhadi, and M Ortiz. Rigorous verification, validation, uncertainty quantification and certification through concentration-of-measure inequalities. *Computer Methods in Applied Mechanics and Engineering*, 197(51-52):4591–4609, 2008.

Olivier Marchal, Julyan Arbel, et al. On the sub-gaussianity of the beta and dirichlet distributions. *Electronic Communications in Probability*, 22, 2017.

Arkadij Semenovič Nemirovskij and David Borisovich Yudin. Problem complexity and method efficiency in optimization. 1983.

Houman Owhadi, Clint Scovel, Timothy John Sullivan, Mike McKerns, and Michael Ortiz. Optimal uncertainty quantification. *Siam Review*, 55(2):271–345, 2013.

Stephen Page and Steffen Grünewälder. The goldenshluger–lepski method for constrained least-squares estimators over rkhss. *Bernoulli*, 27(4):2241–2266, 2021.

REAC Paley and Antoni Zygmund. A note on analytic functions in the unit circle. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 28, pp. 266–272. Cambridge University Press, 1932.

My Phan, Philip Thomas, and Erik Learned-Miller. Towards practical mean bounds for small samples. In *ICML 2021: 38th International Conference on Machine Learning*, pp. 8567–8576, 2021.

Phillippe Rigollet and Jan-Christian Hütter. High dimensional statistics. *Lecture notes for course 18S997*, 2019.

Joseph P Romano and Michael Wolf. Finite sample nonparametric inference and large sample efficiency. *Annals of Statistics*, pp. 756–778, 2000.

Peter J Rousseeuw and Sabine Verboven. Robust estimation in very small samples. *Computational Statistics & Data Analysis*, 40(4):741–758, 2002.

Irina Gennad'evna Shevtsova. On the absolute constants in the berry–esseen inequality and its structural and nonuniform improvements. *Informatika i Ee Primeneniya*, 7(1):124–125, 2013.

Daniel Shiu. Efficient computation of tight approximations to chernoff bounds. *Computational Statistics*, pp. 1–15, 2022.

Youming Tao, Yulian Wu, Peng Zhao, and Di Wang. Optimal rates of (locally) differentially private heavy-tailed multi-armed bandits. In *International Conference on Artificial Intelligence and Statistics*, pp. 1546–1574. PMLR, 2022.

A. W. van der Vaart and Jon A. Wellner. *Weak Convergence and Empirical Processes: With Applications to Statistics*. Springer, 1996.

Roman Vershynin. Introduction to the non-asymptotic analysis of random matrices. *arXiv preprint arXiv:1011.3027*, 2010.

Martin J Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge University Press, 2019.

Chi-Hua Wang, Yang Yu, Botao Hao, and Guang Cheng. Residual bootstrap exploration for bandit algorithms. *arXiv preprint arXiv:2002.08436*, 2020.

Jie Wang, Rui Gao, and Yao Xie. Two-sample test using projected wasserstein distance. In *Proc. ISIT*, volume 21, 2021.

Yan Wang. Sub-gaussian and subexponential fluctuation-response inequalities. *Physical Review E*, 102(5):052105, 2020.

Andreas Winkelbauer. Moments and absolute moments of the normal distribution. *arXiv preprint arXiv:1209.4340*, 2012.

Yun Yang, Zuofeng Shang, and Guang Cheng. Non-asymptotic analysis for nonparametric testing. In *Conference on Learning Theory*, pp. 3709–3755. PMLR, 2020.

Anru R Zhang and Yuchen Zhou. On the non-asymptotic and sharp lower tail bounds of random variables. *Stat*, 9(1): e314, 2020.

Huiming Zhang and Song Xi Chen. Concentration inequalities for statistical inference. *Communications in Mathematical Research*, 37(1):1–85, 2021.

Huiming Zhang and Haoyu Wei. Sharper sub-weibull concentrations. *Mathematics*, 10(13):2252, 2022.

Yao Zheng and Guang Cheng. Finite-time analysis of vector autoregressive models under linear restrictions. *Biometrika*, 108(2):469–489, 2021.

Anatolii Zolotukhin, Sergei Nagaev, and Vladimir Chebotarev. On a bound of the absolute constant in the berry–esseen inequality for iid bernoulli random variables. *Modern Stochastics: Theory and Applications*, 5(3):385–410, 2018.