Uniform Concentration Bounds toward a Unified Framework for Robust Clustering

Abstract

Recent advances in center-based clustering research continue to improve upon the drawbacks of Lloyd's effective k-means algorithm over 60 years since its introduction. Various methods seek to address poor local minima, sensitivity to outliers, and data that are not well-suited to Euclidean measures of fit, but many are supported largely empirically. Moreover, combining such approaches in a piecemeal manner can result in ad hoc methods, and the limited theoretical results supporting each individual contribution may no longer hold. To address these issues at once in a principled way, this paper proposes a cohesive robust framework for center-based clustering under a general class of dissimilarity measures. In particular, we present a rigorous theoretical treatment within a Median-of-Means (MoM) estimation framework, showing that it subsumes several popular k-means variants. In addition to unifying existing methods, we derive uniform concentration bounds that complete their analyses, and bridge these results to the MoM framework via Dudley's chaining arguments. Importantly, we neither require any assumptions on the distribution of the outlying observations nor on the relative number of observations n to features p. We establish strong consistency and an error rate of $O(n^{-1/2})$ under mild conditions, surpassing the best-known results in the literature. The methods are empirically validated thoroughly on real and synthetic datasets.

1 Introduction

Clustering is a fundamental task in unsupervised learning, which seeks to discover naturally occurring groups within a dataset. Among a plethora of algorithms in the literature, center-based methods remain widely popular, with k-means [29, 27] as the most prominent example. Given n data points $\mathcal{X} = \{\boldsymbol{X}_i : i = 1, \ldots, n\} \subset \mathbb{R}^p$, k-means seeks to partition the data into k mutually exclusive and exhaustive groups by minimizing the within cluster variance. Representing the cluster centroids $\boldsymbol{\Theta} = \{\boldsymbol{\theta}_1, \ldots, \boldsymbol{\theta}_k\} \subset \mathbb{R}^p$, the k-means problem is formulated as the minimization of the objective function $P(\boldsymbol{\Theta}) = \sum_{i=1}^n \min_{1 \leq j \leq k} d(\boldsymbol{X}_i, \boldsymbol{\theta}_j)$, where $d(\cdot, \cdot)$ is a dissimilarity measure. Taking the squared Euclidean distance $d(\boldsymbol{x}, \boldsymbol{y}) = \|\boldsymbol{x} - \boldsymbol{y}\|_2^2$ yields the classical k-means formulation.

Clustering via k-means is well documented to be sensitive to initialization [47], prone to getting stuck in poor local optima [50, 49, 13], poorly perform against linearly non-separable clusters [33], and is fragile in the presence of even a single outlying observation [21]. Researchers continue to tackle these well-documented drawbacks of k-means while preserving its simplicity and interpretability [4, 1, 41, 34, 51, 42]. Although often successful in practice, few of these approaches are grounded in rigorous theory or provide finite-sample statistical guarantees. The few consistency results that are established are mostly asymptotic in nature [13, 44, 45], lacking convergence rates or operating under restrictive assumptions on the relation between p and n [36]. For example, the recent large sample analysis [36] of the hard Bregman k-means algorithm [4] obtains an asymptotic error rate of

$O(\sqrt{\log n / n})$ under restrictive assumptions on the relation between n and p. The approach by [41] focuses on a unified framework from an optimization perspective, while [3, 42] focus on different divergence-based methods to better understand the underlying feature-space.

The presence of outliers only further complicates matters. Outlying data are common in real applications, but many of the aforementioned approaches are fragile to deviations from the assumed data generating mechanism. On the other hand, recent work on robust clustering methods [14, 17, 21] fail to integrate the practical advances surveyed above, while similarly lacking finite-sample guarantees. To bridge this methodological gap, the Median of Means (MoM) literature provides a promising and attractive framework to adapt center-based clustering methods to become outlier-robust. MoM estimators are not only insensitive to outliers, but are also equipped with exponential concentration results even under the mild condition of finite variance [28, 24, 6, 26, 22]. Recently, near-optimal results for mean estimation [31], classification [25], regression [30, 28], clustering [21, 11], bandits [12] and optimal transport [40] have been established from this perspective.

Under the MoM lens, we propose a unified framework for robust center-based clustering. Our treatment considers a family of Bregman loss functions not restricted to the classical squared Euclidean loss. We explore the exact sample error bounds for a general class of algorithms under mild regularity assumptions that apply to popular existing approaches, which we show to be special cases. In particular, we use metric entropy arguments and Rademacher complexities, which although extensively used in the empirical processes literature, remain largely unexplored in context to unsupervised learning and especially for clustering. Under the MoM philosophy, we also develop a general robust clustering framework for center-based clustering. The proposed framework allows for the data to be divided into two categories: the set of inliers ($\mathcal{I}$) and the set of outliers ($\mathcal{O}$). The inliers are assumed to be independent and identically distributed (i.i.d.), while absolutely no assumption is required on $\mathcal{O}$, allowing outliers to be unboundedly large, dependent on each other, sampled from a heavy-tailed distribution and so on. Our contributions within the MoM framework make use of Rademacher complexities [5, 7] and symmetrization arguments [46] which are powerful but so far have not been used to their full potential in the context of robust clustering.

The paper is organized as follows. In Section 2, we identify a general centroid-based clustering framework which encompasses k-means [29], k-harmonic means [50], and power k-means [49] as special cases, to name a few. We show how this framework is made robust via Median of Means estimation, yielding an array of center-based robust clustering methods. Within this framework, we derive uniform deviation bounds and concentration inequalities under standard regularity conditions through bounding Rademacher complexity by metric entropy via Dudley's chaining argument in Section 3. The analysis newly reveals the convergence rate for popularly used clustering methods such as k-harmonic means and power k-means, matching the known rate results for k-means, and elegantly carries over to their MoM counterparts. We empirically assess the resulting algorithms thorough simulated and real data experiments. In particular, we observe that power k-means [49] under the MoM paradigm outperforms the state-of-the-art in the presence of outliers.

2 Problem Setting and Proposed Method

We consider the problem of partitioning a set of n data points $\mathcal{X} = \{\boldsymbol{X}_i : i = 1, \ldots, n\} \subset \mathbb{R}^p$ into k mutually exclusive clusters. In a center-based clustering framework, we represent the j-th cluster by its centroid $\boldsymbol{\theta}_j \in \mathbb{R}^p$, $j \in \{1, \ldots, k\}$. Towards determining closeness, we allow the dissimilarity measure to be any Bregman divergence. Recall any differentiable, convex function $\phi : \mathbb{R}^p \to \mathbb{R}$ generates the Bregman divergence $d_\phi : \mathbb{R}^p \times \mathbb{R}^p \to \mathbb{R}_{\geq 0}$ ($\mathbb{R}_{\geq 0}$ is the set of all non-negative reals) defined as $d_\phi(\boldsymbol{x}, \boldsymbol{y}) = \phi(\boldsymbol{x}) - \phi(\boldsymbol{y}) - \langle \nabla \phi(\boldsymbol{y}), \boldsymbol{x} - \boldsymbol{y} \rangle$. For instance, $\phi(\boldsymbol{u}) = \|\boldsymbol{u}\|_2^2$ generates the Euclidean distance. Without loss of generality, one may assume $\phi(\boldsymbol{0}) = \nabla \phi(\boldsymbol{0}) = 0$. In this paradigm, clustering is achieved by minimizing the objective

$$\frac{1}{n} \sum_{i=1}^{n} \Psi_{\boldsymbol{\alpha}} \left(d_\phi(\boldsymbol{X}_i, \boldsymbol{\theta}_1), \ldots, d_\phi(\boldsymbol{X}_i, \boldsymbol{\theta}_k) \right). \tag{1}$$

Here $\Psi_{\boldsymbol{\alpha}} : \mathbb{R}_{\geq 0}^k \to \mathbb{R}_{\geq 0}$ is a component-wise non-decreasing function (such as a generalized mean) of the dissimilarities $\{d_\phi(\boldsymbol{X}, \boldsymbol{\Theta}_j)\}_{j=1}^k$ which satisfies $\Psi(\boldsymbol{0}) = 0$. The hyperparameter $\boldsymbol{\alpha} \in \mathcal{A} \subseteq \mathbb{R}^q$ is specified by the user, and we will additionally assume that Ψ is Lipschitz. For intuition, we begin by showing how this setup includes several popular clustering methods.

Examples: Suppose $\phi(\boldsymbol{u}) = \|\boldsymbol{u}\|_2^2$ and $\Psi(\boldsymbol{x}) = (\sum_{j=1}^k x_j^{-1})^{-1}$. Then the objective (1) becomes $\sum_{i=1}^n (\sum_{j=1}^k \|\boldsymbol{X}_i - \boldsymbol{\theta}_j\|_2^{-2})^{-1}$, which is the objective function of k-harmonic means clustering [50]. Now consider other generalized means: take $\Psi_s(\boldsymbol{x}) = M_s(\boldsymbol{x})$ where $M_s(\boldsymbol{x}) = (k^{-1} \sum_{i=1}^k x_i^s)^{1/s}$ for $s \in (-\infty, -1]$. Then the objective (1) coincides with the recent power k-means method [49], $\sum_{i=1}^n M_s\left(\|\boldsymbol{X}_i - \boldsymbol{\theta}_1\|_2^2, \ldots, \|\boldsymbol{X}_i - \boldsymbol{\theta}_k\|_2^2\right)$. If instead $\Psi(\boldsymbol{x}) = \min_{1 \leq j \leq k} x_j$, then (1) becomes equivalent to Bregman clustering $\sum_{i=1}^n \min_{1 \leq j \leq k} d_\phi(\boldsymbol{X}_i, \boldsymbol{\theta}_j)$ proposed in [4] for any valid ϕ, while the special case of $\phi(\boldsymbol{u}) = \|\boldsymbol{u}\|_2^2$ yields the familiar k-means problem under Euclidean distance [29].

In what follows, we derive concentration bounds that establish new theoretical guarantees such as consistency and convergence rates for clustering algorithms in this framework. These analyses lead us to a unified, robust framework by embedding this class of methods within the Median of Means (MoM) estimation paradigm. Via elegant connections between the properties of MoM estimators and Vapnik-Chervonenkis (VC) theory, our MoM estimators too will inherit uniform concentration inequalities from the preceding analysis, extending convergence guarantees to the robust setting.

Median of Means Instead of directly minimizing the empirical risk (1), MoM begins by partitioning the data into L sets $B_1, \ldots, B_L$ which each contain exactly b many elements (discarding a few observations when n is not divisible by L). The partitions can be assigned uniformly at random, or can be shuffled throughout the algorithm [25]. For notational simplicity, we abbreviate $f_{\boldsymbol{\Theta}}(\boldsymbol{X}) = \Psi_{\boldsymbol{\alpha}}\left(d_\phi(\boldsymbol{X}, \boldsymbol{\theta}_1), \ldots, d_\phi(\boldsymbol{X}, \boldsymbol{\theta}_k)\right)$. MoM then entails a robust version of the estimator under (1) by minimizing the following objective with respect to $\boldsymbol{\Theta}$:

$$\text{MoM}_L^n(\boldsymbol{\Theta}) = \text{Median}\left(\frac{1}{b}\sum_{i \in B_1} f_{\boldsymbol{\Theta}}(\boldsymbol{X}_i), \ldots, \frac{1}{b}\sum_{i \in B_L} f_{\boldsymbol{\Theta}}(\boldsymbol{X}_i)\right). \tag{2}$$

Intuitively, MoM estimators are more robust than their ERM counterparts since under mild conditions, only a subset of the partitions is contaminated by outliers while others are outlier-free. Taking the median over partitions negates the influence of these spurious partitions, thus reducing the effect of outliers. Formal analysis of robustness via breakdown points is also available for MoM estimators [23, 38]; we will make use of the nice concentration properties of MoM estimators in Section 3.4.

Algorithm 1 MoM Clustering via Adagrad

Input: $\boldsymbol{X} \in \mathbb{R}^{n \times p}$, k, L, $f_{\boldsymbol{\Theta}}(\cdot)$, η, ϵ.
Output: The cluster centroids $\boldsymbol{\Theta}$.
Initialization: Randomly partition $\{1, \ldots, n\}$ into L many partitions of equal length. Randomly pick k points without replacement from $\mathcal{X}$ to initialize $\boldsymbol{\Theta}_0$.
repeat
    Step 1: Find $\ell_t \in \{1, \ldots, L\}$, such that $\text{MoM}_L^n(\boldsymbol{\Theta}_t) = \frac{1}{b}\sum_{i \in B_{\ell_t}} f_{\boldsymbol{\Theta}_t}(\boldsymbol{X}_i)$.
    Step 2: $\boldsymbol{g}_t \leftarrow \frac{1}{b}\sum_{i \in B_{\ell_t}} \nabla_{\boldsymbol{\theta}_j} f_{\boldsymbol{\Theta}}(\boldsymbol{X}_i)$
    Step 3: Update $\boldsymbol{\Theta}$ by $\boldsymbol{\theta}_j^{(t+1)} \leftarrow \boldsymbol{\theta}_j^{(t)} - \frac{\eta}{\sqrt{\epsilon + \sum_{t'=1}^{(t)} \|\boldsymbol{g}_j^{(t)}\|_2^2}} \boldsymbol{g}_j^{(t)}$.
until objective (2) converges

Furthermore, optimizing (2) is tractable via gradient-based methods. We advocate the Adagrad algorithm, whose updates are given by $\boldsymbol{\theta}_j^{(t+1)} \leftarrow \boldsymbol{\theta}_j^{(t)} - \frac{\eta}{\sqrt{\epsilon + \sum_{t'=1}^{(t)} \|\boldsymbol{g}_j^{(t)}\|_2^2}} \boldsymbol{g}_j^{(t)}$ [15, 18] for hyperparameter $\epsilon > 0$ and learning rate $\eta > 0$. Here $\boldsymbol{g}_j^{(t)}$ is a subgradient of $\text{MoM}_L^n(\boldsymbol{\Theta})$ at $\boldsymbol{\Theta}_t$: if ℓ_t denotes the median partition step t, then $\nabla_{\boldsymbol{\Theta}_j} \text{MoM}_L^n(\boldsymbol{\Theta}_t) = \frac{1}{b}\sum_{i \in B_{\ell_t}} \nabla_{\boldsymbol{\theta}_j} f_{\boldsymbol{\Theta}}(\boldsymbol{X}_i)|_{\boldsymbol{\Theta}=\boldsymbol{\Theta}_t}$. For any $\Psi_{\boldsymbol{\alpha}}$ differentiable, by the chain rule, $\nabla_{\boldsymbol{\theta}_j} f_{\boldsymbol{\Theta}}(\boldsymbol{x}) = \partial_j \Psi_{\boldsymbol{\alpha}}\left(d_\phi(\boldsymbol{x}, \boldsymbol{\theta}_1^{(t)}), \ldots, d_\phi(\boldsymbol{x}, \boldsymbol{\theta}_k^{(t)})\right) \nabla_{\boldsymbol{\theta}} d_\phi(\boldsymbol{x}, \boldsymbol{\theta})$. More concretely, consider the power k-means objective $f_{\boldsymbol{\Theta}}(\boldsymbol{x}) = M_s(\|\boldsymbol{x} - \boldsymbol{\theta}_1\|_2^2, \ldots, \|\boldsymbol{x} - \boldsymbol{\theta}_k\|_2^2)$: differentiating immediately yields $\nabla_{\boldsymbol{\theta}_j} f_{\boldsymbol{\Theta}}(\boldsymbol{x}) = \frac{2}{k}\left(\frac{1}{k}\sum_{j'=1}^k \|\boldsymbol{x} - \boldsymbol{\theta}_{j'}\|_2^{2s}\right)^{1/s-1} \|\boldsymbol{x} - \boldsymbol{\theta}_j\|_2^{2(s-1)}(\boldsymbol{\theta}_j - \boldsymbol{x})$. As another example. the classic k-means objective $f_{\boldsymbol{\Theta}}(\boldsymbol{x}) = \min_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \|\boldsymbol{x} - \boldsymbol{\theta}\|_2^2$ requires the subgradient $\nabla_{\boldsymbol{\theta}_j} f_{\boldsymbol{\Theta}}(\boldsymbol{x}) = 2(\boldsymbol{\theta}_j - \boldsymbol{x})\mathbb{1}\{j \in \mathcal{J}(\boldsymbol{\Theta}, \boldsymbol{x})\}$, where $\mathcal{J}(\boldsymbol{\Theta}, \boldsymbol{x}) = \text{argmin}_{1 \leq j \leq k} \|\boldsymbol{x} - \boldsymbol{\theta}_j\|_2^2$. Here $\mathbb{1}\{\cdot\}$ denotes the indicator function.

A pseudocode summary appears in Algorithm 1. This method tends to find clusters efficiently: for instance, we incur $O(npk)$ per-iteration complexity applied to both the k-means and power k-means instances, which matches that of their original algorithms without robustifying via MoM [29, 27, 49].

3 Theoretical Analysis

We now analyze properties of the proposed framework (1), with complete proofs in the Supplement. Denoting $\mathcal{M}$ the set of all probability measures P with support on $[-M, M]^p$, i.e. $P([-M, M]^p) = 1$, we first make standard assumptions that data are i.i.d. with bounded components [9, 13, 36].

A1. $\boldsymbol{X}_1 \ldots, \boldsymbol{X}_n \overset{i.i.d.}{\sim} P$ such that $P \in \mathcal{M}$.

Let P_n be the empirical distribution based on $\boldsymbol{X}_1 \ldots, \boldsymbol{X}_n$, i.e. $P_n(A) = \sum_{i=1}^n \mathbb{1}\{\boldsymbol{X}_i \in A\}/n$, for any Borel set A. For notational simplicity, we write $\mu f := \int f d\mu$. Appealing to the Strong Law of Large Numbers (SLLN) [2], $P_n f_{\boldsymbol{\Theta}} \xrightarrow{a.s.} P f_{\boldsymbol{\Theta}}$. Suppose $\widehat{\boldsymbol{\Theta}}_n$ be a (global) minimizer of (1) and $\boldsymbol{\Theta}^*$ be the global minimizer of $P f_{\boldsymbol{\Theta}}$. Since the functions $P_n f_{\boldsymbol{\Theta}}$ and $P f_{\boldsymbol{\Theta}}$ are close to each other as n become large, we can expect that their respective minimizers, $\widehat{\boldsymbol{\Theta}}_n$ and $\boldsymbol{\Theta}^*$ are also close to one another. To show that $\widehat{\boldsymbol{\Theta}}_n$ converges to $\boldsymbol{\Theta}^*$ as $n \to \infty$, we consider bounding the uniform deviation $\sup_{\boldsymbol{\Theta}} |P_n f_{\boldsymbol{\Theta}} - P f_{\boldsymbol{\Theta}}|$. Towards establishing such bounds, we will posit two regularity assumptions on $\Psi_{\boldsymbol{\alpha}}(\cdot)$ and $\phi(\cdot)$, beginning with a $\tau_{\boldsymbol{\alpha},k}$-Lipschitz condition on $\Psi_{\boldsymbol{\alpha}}$.

A2. For all $\boldsymbol{\alpha} \in \mathcal{A}$ and any $\boldsymbol{x}, \boldsymbol{y} \in \mathbb{R}_{\geq 0}^k$, we have $|\Psi_{\boldsymbol{\alpha}}(\boldsymbol{x}) - \Psi_{\boldsymbol{\alpha}}(\boldsymbol{y})| \leq \tau_{\boldsymbol{\alpha},k} \|\boldsymbol{x} - \boldsymbol{y}\|_1$.

We also assume a weaker form of a standard condition that the gradient $\nabla \phi(\cdot)$ is H_p-Lipschitz [43]; unlike their work, note we do *not* additionally require strong convexity of ϕ.

A3. There exists $H_p \geq 0$ such that $\|\nabla \phi(\boldsymbol{x}) - \nabla \phi(\boldsymbol{y})\|_2 \leq H_p \|\boldsymbol{x} - \boldsymbol{y}\|_2$ for all $\boldsymbol{x}, \boldsymbol{y} \in [-M, M]^p$.

These conditions are mild, and can be seen to hold for all of the aforementioned popular special cases. For instance, taking $\phi(\boldsymbol{u}) = \|\boldsymbol{u}\|^2$ yields the squared Euclidean distances, and we see that $\nabla \phi(\boldsymbol{u}) = 2\boldsymbol{u}$ so that A3 is satisfied with constant $H_p = 2$. Now, let $\Psi(\boldsymbol{x}) = \min_{1 \leq j \leq k} x_j$ as in classical k-means, and denote $j^* \in \arg\min_{1 \leq j \leq k} y_j$. For any vectors $\boldsymbol{x}, \boldsymbol{y} \in \mathbb{R}_{\geq 0}^p$,

$$\Psi(\boldsymbol{x}) - \Psi(\boldsymbol{y}) = \min_{1 \leq j \leq k} x_j - \min_{1 \leq j \leq k} y_j = \min_{1 \leq j \leq k} x_j - y_{j^*} \leq x_{j^*} - y_{j^*} \leq \|\boldsymbol{x} - \boldsymbol{y}\|_1.$$

Thus, Ψ is clearly non-negative and componentwise non-decreasing, and satisfies A2. Similarly, if we take $\Psi_s(\boldsymbol{x}) = M_s(\boldsymbol{x})$, the conditions are met for power k-means: it is again non-negative and non-decreasing in its components, and satisfies A2 with $\tau_{\boldsymbol{\alpha},k} = k^{-1/s}$ due to [8].

3.1 Bounds on $\widehat{\boldsymbol{\Theta}}_n$ and $\boldsymbol{\Theta}^*$

Toward proving that $\widehat{\boldsymbol{\Theta}}_n$ converges to $\boldsymbol{\Theta}^*$, we will need to show that both $\widehat{\boldsymbol{\Theta}}_n$ and $\boldsymbol{\Theta}^*$ lie in $[-M, M]^{k \times p}$. To this end, we first establish the obtuse angle property for Bregman divergences.

Lemma 3.1. Let $\mathcal{C}$ be a convex set and suppose $P_{\mathcal{C}}(\boldsymbol{\theta})$ be the projection of $\boldsymbol{\theta}$ onto $\mathcal{C}$, with respect to the Bregman divergence $d_\phi(\cdot, \cdot)$, i.e. $P_{\mathcal{C}}(\boldsymbol{\theta}) = \arg\min_{\boldsymbol{x} \in \mathcal{C}} d_\phi(\boldsymbol{x}, \boldsymbol{\theta})$ (assuming it exists). Then,

$$d_\phi(\boldsymbol{x}, \boldsymbol{\theta}) \geq d_\phi(\boldsymbol{x}, P_{\mathcal{C}}(\boldsymbol{\theta})) + d_\phi(P_{\mathcal{C}}(\boldsymbol{\theta}), \boldsymbol{\theta}), \quad \text{for all } \boldsymbol{x} \in \mathcal{C}.$$

We next show that to minimize $P_n f_{\boldsymbol{\Theta}}$ or $P f_{\boldsymbol{\Theta}}$, it is enough to restrict the search to $[-M, M]^{k \times p}$.

Lemma 3.2. Let A2 hold, and $Q \in \mathcal{M}$. Let $d_\phi : \mathbb{R}^p \times \mathbb{R}^p \to \mathbb{R}_{\geq 0}$ be a Bregman divergence. Then for any $\boldsymbol{\Theta} \in \mathbb{R}^{k \times p}$, there exists $\boldsymbol{\Theta}' \in [-M, M]^{k \times p}$, such that $Q f_{\boldsymbol{\Theta}'} \leq Q f_{\boldsymbol{\Theta}}$.

Since we can restrict our attention to $[-M, M]^{k \times p}$ to minimize $Q f_{\boldsymbol{\Theta}}$, we have the following:

Corollary 3.1. Let $Q \in \mathcal{M}$ and $d_\phi : \mathbb{R}^p \times \mathbb{R}^p \to \mathbb{R}_{\geq 0}$ be a Bregman divergence. If $\boldsymbol{\Theta}_0 = \arg\min_{\boldsymbol{\Theta} \in \mathbb{R}^{k \times p}} \Psi(\boldsymbol{\Theta}, Q)$, then $\boldsymbol{\Theta}_0 \in [-M, M]^{k \times p}$.

Now note under A1 both P and P_n have support contained in $[-M, M]^{k \times p}$. The following corollary is thus implied by replacing Q by P and P_n.

162 **Corollary 3.2.** *Under A1—3, both* $\widehat{\boldsymbol{\Theta}}_n, \boldsymbol{\Theta}^* \in [-M, M]^{k \times p}$.

163 Now that we have bounded $\widehat{\boldsymbol{\Theta}}_n$ and $\boldsymbol{\Theta}^*$ in a compact set, the following section supplies probabilistic
164 bounds on the uniform deviation $2 \sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} |P_n f_{\boldsymbol{\Theta}} - P f_{\boldsymbol{\Theta}}|$ via metric entropy.

3.2 Concentration Inequality and Metric Entropy Bounds via Rademacher Complexity

166 We have proven that $\widehat{\boldsymbol{\Theta}}_n, \boldsymbol{\Theta}^* \in [-M, M]^{k \times p}$. To bound the difference $|P f_{\widehat{\boldsymbol{\Theta}}_n} - P f_{\boldsymbol{\Theta}^*}|$, we observe

$$
\begin{aligned}
|P f_{\widehat{\boldsymbol{\Theta}}_n} - P f_{\boldsymbol{\Theta}^*}| &= P f_{\widehat{\boldsymbol{\Theta}}_n} - P f_{\boldsymbol{\Theta}^*} = P f_{\widehat{\boldsymbol{\Theta}}_n} - P_n f_{\widehat{\boldsymbol{\Theta}}_n} + P_n f_{\widehat{\boldsymbol{\Theta}}_n} - P_n f_{\boldsymbol{\Theta}^*} + P_n f_{\boldsymbol{\Theta}^*} - P f_{\boldsymbol{\Theta}^*} \\
&\leq P f_{\widehat{\boldsymbol{\Theta}}_n} - P_n f_{\widehat{\boldsymbol{\Theta}}_n} + P_n f_{\boldsymbol{\Theta}^*} - P f_{\boldsymbol{\Theta}^*} \leq 2 \sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} |P_n f_{\boldsymbol{\Theta}} - P f_{\boldsymbol{\Theta}}|. \quad (3)
\end{aligned}
$$

167 Thus, to bound $|P f_{\widehat{\boldsymbol{\Theta}}_n} - P f_{\boldsymbol{\Theta}^*}|$, it is enough to prove bounds on $\sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} |P_n f_{\boldsymbol{\Theta}} - P f_{\boldsymbol{\Theta}}|$.
168 The main idea here is to bound this uniform deviation via Rademacher complexity, and in turn bound
169 the Rademacher complexity itself [16, 32]. Let $\mathcal{F} = \{f_{\boldsymbol{\Theta}} : \boldsymbol{\Theta} \in [-M, M]^{k \times p}\}$, and denote the
170 $\mathcal{F}$-norm [2] between two probability measures μ and ν as $\|\mu - \nu\|_{\mathcal{F}} = \sup_{f \in \mathcal{F}} |\int f d\mu - \int f d\nu|$.
171 We recall the definition of Rademacher complexity and covering number as follows.

172 **Definition 1.** *Let* ϵ_i*'s be i.i.d. Rademacher random variables independent of* $\mathcal{X} = \{\boldsymbol{X}_1, \dots, \boldsymbol{X}_n\}$,
173 *i.e.* $\mathbb{P}(\epsilon_i = 1) = \mathbb{P}(\epsilon_i = -1) = 0.5$, *The population Rademacher complexity of* $\mathcal{F}$ *is defined as*
174 $\mathcal{R}_n(\mathcal{F}) = \mathbb{E} \sup_{f \in \mathcal{F}} \frac{1}{n} \sum_{i=1}^{n} \epsilon_i f(\boldsymbol{X}_i)$, *where the expectation is over both* $\boldsymbol{\epsilon}$ *and* $\mathcal{X}$.

175 **Definition 2.** *(δ-cover and Covering Number) Let* (X, d) *be a metric space. The set* $X_\delta \subseteq X$ *is said*
176 *to be a δ-cover of* X *if for all* $x \in X$, $\exists\ x' \in X_\delta$, *such that* $d(x, x') \leq \delta$. *The δ-covering number of*
177 X *w.r.t.* d, *denoted by* $N(\delta; X, d)$, *is the size of the smallest δ-cover of* X *with respect to* d.

178 The following Lemma gives a bound for the δ-covering number of $\mathcal{F}$ under the supremum norm. The
179 main idea here is to use the Lipschitz property of $f_{\boldsymbol{\Theta}}$ and then to find a cover of the search space for
180 $\boldsymbol{\Theta}$, i.e. $[-M, M]^{k \times p}$. This automatically transcends to a cover of $\mathcal{F}$ under the sup norm.

181 **Lemma 3.3.** *Let* $N(\delta; \mathcal{F}, \| \cdot \|_\infty)$ *be the δ-covering number of* $\mathcal{F}$ *under* $\| \cdot \|_\infty$. *Then, under A1—3,*

$$
N(\delta; \mathcal{F}, \| \cdot \|_\infty) \leq \left(\max \left\{ \left\lfloor \frac{8M^2 \tau_{\boldsymbol{\alpha},k} H_p kp}{\delta} \right\rfloor, 1 \right\} \right)^{kp}.
$$

182 To bound the Rademacher complexity, we will also need to bound the diameter of $\mathcal{F}$ under $\| \cdot \|_\infty$:

183 **Lemma 3.4.** *Let* $diam(\mathcal{F}) = \sup_{f,g \in \mathcal{F}} \|f - g\|_\infty$. *Then, under A1—3,* $diam(\mathcal{F}) \leq 8\tau_{\boldsymbol{\alpha},k} H_p M^2 kp$.

184 We are now ready to state the bound on the Rademacher complexity $\mathcal{R}_n(\mathcal{F})$ in Theorem 3.1. We only
185 provide a proof sketch in this section. The complete proof details are available in the Supplement.

186 **Theorem 3.1.** *Under A1—3,* $\mathcal{R}_n(\mathcal{F}) \leq 48\sqrt{\pi} \tau_{\boldsymbol{\alpha},k} H_p M^2 (kp)^{3/2} n^{-1/2}$.

187 **Proof Sketch** The main idea of the proof is to use Dudley's chaining argument [16, 48] to bound
188 the Rademacher complexity in terms of an integral involving the metric entropy $\log N(\delta; \mathcal{F}, \| \cdot \|_\infty)$.
189 Let $\Delta = 8 H_p M^2 k^{1-1/s} p$. Using the chaining approach, one can show that

$$
\begin{aligned}
\mathcal{R}_n(\mathcal{F}) \leq \frac{12}{\sqrt{n}} \int_0^\Delta \sqrt{\log N(\epsilon; \mathcal{F}, \| \cdot \|_\infty)} d\epsilon &\leq \frac{12}{\sqrt{n}} \int_0^\Delta \sqrt{kp \log \left(\max \left\{ \frac{\Delta}{\epsilon}, 1 \right\} \right)} d\epsilon \quad (4) \\
&= \frac{12}{\sqrt{n}} \int_0^\Delta \sqrt{kp \log \left(\frac{\Delta}{\epsilon} \right)} d\epsilon \\
&= 48\sqrt{\pi} \tau_{\boldsymbol{\alpha},k} H_p M^2 (kp)^{3/2} n^{-1/2}.
\end{aligned}
$$

190 Here the second inequality (4) follows from Lemma 3.3. Before we proceed, we state the following
191 Lemma, which puts an uniform bound on $\|f\|_\infty, f \in \mathcal{F}$.

192 **Lemma 3.5.** *For all* $\boldsymbol{x} \in [-M, M]^p$ *and* $\boldsymbol{\Theta} \in [-M, M]^{k \times p}$,

$$
0 \leq \Psi_{\boldsymbol{\alpha}}(d_\phi(\boldsymbol{x}, \boldsymbol{\theta}_1), \dots, d_\phi(\boldsymbol{x}, \boldsymbol{\theta}_k)) \leq 4\tau_{\boldsymbol{\alpha},k} H_p M^2 pk.
$$

Now that we have established a bound on the Rademacher complexity $\mathcal{R}_n(\mathcal{F})$, we are now ready to prove a uniform concentration inequality on $\sup_{f \in \mathcal{F}} |P_n f - Pf|$ in the following Theorem.

Theorem 3.2. *Under A1—3, with probability at least $1 - \delta$, the following holds.*

$$\|P_n - P\|_{\mathcal{F}} \leq 96\sqrt{\pi}\tau_{\boldsymbol{\alpha},k}H_p M^2 (kp)^{3/2} n^{-1/2} + 4\tau_{\boldsymbol{\alpha},k}H_p M^2 pk \sqrt{\frac{\log(2/\delta)}{2n}}.$$

Using the bound derived in Theorem 3.2, we now provide a non-asymptotic bound on $|Pf_{\widehat{\boldsymbol{\Theta}}_n} - Pf_{\boldsymbol{\Theta}^*}|$:

Corollary 3.3. *Under A1—3, with probability at least $1 - \delta$, the following holds.*

$$|Pf_{\widehat{\boldsymbol{\Theta}}_n} - Pf_{\boldsymbol{\Theta}^*}| \leq 192\sqrt{\pi}\tau_{\boldsymbol{\alpha},k}H_p M^2 (kp)^{3/2} n^{-1/2} + 8\tau_{\boldsymbol{\alpha},k}H_p M^2 pk \sqrt{\frac{\log(2/\delta)}{2n}}.$$

Proof. From equation (3), we know that $|Pf_{\widehat{\boldsymbol{\Theta}}_n} - Pf_{\boldsymbol{\Theta}^*}| \leq 2\|P_n - P\|_{\mathcal{F}}$. The corollary follows immediately by application of Theorem 3.2. $\square$

3.3 Inference for Fixed p: Strong and $\sqrt{n}$ Consistency

We now discuss the classical domain where p is kept fixed [37, 13, 44] and show that the results above imply strong and $\sqrt{n}$-consistency. We first solidify the notion of convergence $\widehat{\boldsymbol{\Theta}}_n$ to $\boldsymbol{\Theta}^*$ following Pollard [37]. Since centroids are unique only up to label permutations, our notion of dissimilarity

$$\text{diss}(\boldsymbol{\Theta}_1, \boldsymbol{\Theta}_2) = \min_{M \in \mathcal{P}_k} \|\boldsymbol{\Theta}_1 - M\boldsymbol{\Theta}_2\|_F$$

considers $\mathcal{P}_k$ the set of all $k \times k$ real permutation matrices, where $\|\cdot\|_F$ denotes the Frobenius norm. Now, we say that the sequence $\boldsymbol{\Theta}_n$ converges to $\boldsymbol{\Theta}$ if $\lim_{n \to \infty} \text{diss}(\boldsymbol{\Theta}_n, \boldsymbol{\Theta}) = 0$. We begin by imposing the following standard identifiablity condition [37, 44, 13] on P for our analysis.

A4. *For all $\eta > 0$, there exists $\epsilon > 0$, such that $Pf_{\boldsymbol{\Theta}} > Pf_{\boldsymbol{\Theta}^*} + \epsilon$ whenever $\text{diss}(\boldsymbol{\Theta}, \boldsymbol{\Theta}^*) > \eta$.*

We now investigate the strong consistency properties of $\widehat{\boldsymbol{\Theta}}_n$, and also investigate the rate at which $|Pf_{\widehat{\boldsymbol{\Theta}}_n} - Pf_{\boldsymbol{\Theta}^*}|$ converges to 0. Theorem 3.3 states that indeed strong consistency holds, with convergence rate $O(n^{-1/2})$. Note that this rate is faster than that found previously in [36]. Before we proceed, recall we say that $X_n = O_P(a_n)$ if the sequence X_n/a_n is tight [2].

Theorem 3.3. *(Strong consistency and $\sqrt{n}$-consistency) If p is kept fixed then under A1—4, $\widehat{\boldsymbol{\Theta}}_n \xrightarrow{a.s.} \boldsymbol{\Theta}^*$, under P. Moreover, $|Pf_{\widehat{\boldsymbol{\Theta}}_n} - Pf_{\boldsymbol{\Theta}^*}| = O_P(n^{-1/2})$.*

3.4 Inference Under the MoM Framework

In this section, we extend results under (1) to the MoM setup under (2). Recall that in this setting, the data are partitioned into L equally sized blocks; without loss of generality, we take $n = L \cdot b$. We denote the set of all inliers by $\{\boldsymbol{X}_i\}_{i \in \mathcal{I}}$ and outliers by $\{\boldsymbol{X}_i\}_{i \in \mathcal{O}}$. Now let $\widehat{\boldsymbol{\Theta}}_n^{(\text{MoM})}$ denote the minimizer of (2): towards establishing the error rate at which $|Pf_{\widehat{\boldsymbol{\Theta}}_n^{(\text{MoM})}} - Pf_{\boldsymbol{\Theta}^*}|$ goes to 0, we assume the following:

A5. $\{\boldsymbol{X}_i\}_{i \in \mathcal{I}} \overset{i.i.d.}{\sim} P$ *with $P \in \mathcal{M}$.*

A6. *There exists $\eta > 0$ such that $L > (2 + \eta)|\mathcal{O}|$.*

We remark A5 is identical to A1, but imposed only on the inlying observations. A6 ensures at least half of the L partitions are free of outliers; note this is much weaker than requiring $L > 4|\mathcal{O}|$ as is done in recent work [25]. Importantly, we emphasize that *no distributional assumptions* regarding the outlying observations are made, allowing them to be unbounded, generated from heavy-tailed distributions, or dependent among each other. Proofs of the following results appear in the Supplement.

As a point of departure, we first establish that $\widehat{\boldsymbol{\Theta}}_n^{(\text{MoM})} \in [-M, M]^{k \times p}$.

Lemma 3.6. *Let A2—3 and A5—6 hold. Then for any $\boldsymbol{\Theta} \in \mathbb{R}^{k \times p}$, there exists $\boldsymbol{\Theta}' \in [-M, M]^{k \times p}$, such that $MoM_L^n(\boldsymbol{\Theta}') \leq MoM_L^n(\boldsymbol{\Theta})$.*

230   Again, we may restrict the search space for finding $\widehat{\boldsymbol{\Theta}}_n^{(\text{MoM})}$ in $[-M, M]^{k \times p}$ due to Lemma 3.6.

231   **Corollary 3.4.** *Under A2—3 and A5—6, $\widehat{\boldsymbol{\Theta}}_n^{(MoM)} \in [-M, M]^{k \times p}$.*

232   Similarly to Section 3, we derive a uniform bound on $\sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} |\text{MoM}_L^n(f_{\boldsymbol{\Theta}}) - Pf_{\boldsymbol{\Theta}}|$ and

233   then bound $|Pf_{\widehat{\boldsymbol{\Theta}}_n^{(\text{MoM})}} - Pf_{\boldsymbol{\Theta}^*}|$ in turn. For brevity we define $\delta := 2/(4 + \eta) - |\mathcal{O}|/L$; then the

234   uniform deviation bound is as follows. Here the notation "$\lesssim$" supresses the absolute constants.

235   **Theorem 3.4.** *Under A2—3 and A5—6, with probability at least $1 - 2e^{-2L\delta^2}$, the following holds.*

$$\sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} |MoM_L^n(f_{\boldsymbol{\Theta}}) - Pf_{\boldsymbol{\Theta}}| \lesssim \tau_{\boldsymbol{\alpha},k} H_p \max\left\{ kp\sqrt{\frac{L}{n}}, (kp)^{3/2} \frac{\sqrt{|\mathcal{I}|}}{n} \right\}.$$

236   We give an outline of our proof for Theorem 3.4 below, with details appearing in the Supplement.

237   **Proof sketch**   Let P_{B_ℓ} be the empirical distribution of $\{\boldsymbol{X}_i\}_{i \in B_\ell}$. We will bound the probability

238   that $\sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} |\text{MoM}_L^n(f_{\boldsymbol{\Theta}}) - Pf_{\boldsymbol{\Theta}}| > \epsilon$, and later detail choice of $\epsilon > 0$. We first note that

239   if $\sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} \sum_{\ell=1}^{L} \mathbb{1}\{(P - P_{B_\ell})f_{\boldsymbol{\Theta}} > \epsilon\} > L/2$, we have $\sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} (Pf_{\boldsymbol{\Theta}} -$

240    $\text{MoM}_L^n(f_{\boldsymbol{\Theta}})) > \epsilon$. Now let $\varphi(t) = (t - 1)\mathbb{1}\{1 \le t \le 2\} + \mathbb{1}\{t > 2\}$.

241   We observe that $\sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} \sum_{\ell=1}^{L} \mathbb{1}\{(P - P_{B_\ell})f_{\boldsymbol{\Theta}} > \epsilon\} \le \xi_1 + \xi_2 + |\mathcal{O}|$, where

242    $\xi_1 == \sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} \sum_{\ell \in \mathcal{L}} [\varphi(2(P - P_{B_\ell})f_{\boldsymbol{\Theta}}/\epsilon) - \mathbb{E}\varphi(2(P - P_{B_\ell})f_{\boldsymbol{\Theta}}/\epsilon)]$ and $\xi_2 =$

243    $\sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} \sum_{\ell \in \mathcal{L}} \mathbb{E}\varphi(2(P - P_{B_\ell})f_{\boldsymbol{\Theta}}/\epsilon)$. We can bound ξ_2 by appealing to Hoeffding's

244   inequality since $\mathbb{E}\varphi\left(\frac{2(P-P_{B_\ell})f_{\boldsymbol{\Theta}}}{\epsilon}\right) \le \exp\left\{-b\epsilon^2/(32\tau_{\boldsymbol{\alpha},k}^2 H_p^2 M^4 k^2 p^2)\right\}$. ξ_1 can be tackled us-

245   ing McDiarmid's inequality, so that with probability at least $1 - e^{-2L\delta^2}$, it holds that $\xi_1 \le$

246    $\frac{8}{n\epsilon}\mathbb{E}\left[\sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} \sum_{i \in \mathcal{J}} \gamma_i f_{\boldsymbol{\Theta}}(\boldsymbol{X}_i)\right] + \delta \le \frac{384}{n\epsilon}\sqrt{\pi}\tau_{\boldsymbol{\alpha},k} H_p M^2 (kp)^{3/2}\sqrt{|\mathcal{I}|} + \delta$. The last in-

247   equality holds due to Theorem 3.1.

248   Thus, combining equations the bounds on ξ_1 and ξ_2, we conclude that, with probability of at least

249    $1 - e^{-2L\delta^2}$, $\sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} \sum_{\ell=1}^{L} \mathbb{1}\{(P - P_{B_\ell})f_{\boldsymbol{\Theta}} > \epsilon\} \le L\left(\exp\left\{-\frac{b\epsilon^2}{32\tau_{\boldsymbol{\alpha},k}^2 H_p^2 M^4 k^2 p^2}\right\} +\right.$

250    $\left.\frac{|\mathcal{O}|}{L} + \frac{384}{n\epsilon}\sqrt{\pi}\tau_{\boldsymbol{\alpha},k} H_p M^2 (kp)^{3/2}\sqrt{|\mathcal{I}|} + \delta\right)$. We choose $\delta = \frac{2}{4+\eta} - \frac{|\mathcal{O}|}{L}$ and $\epsilon =$

251    $2\max\left\{\sqrt{32\tau_{\boldsymbol{\alpha},k}^2 H_p^2 M^4 \log(4(\eta+4)/\eta)}kp\sqrt{L/n}, 1536(\eta+4)\eta^{-1}\tau_{\boldsymbol{\alpha},k} H_p M^2\sqrt{\pi}(kp)^{3/2}\sqrt{|\mathcal{I}|}/n\right\}$.

252   This makes the right hand side of the above inequality strictly smaller than $L/2$, implying that

253    $\sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} (Pf_{\boldsymbol{\Theta}} - \text{MoM}_L^n(f_{\boldsymbol{\Theta}})) \le \epsilon$, with probability at least $1 - e^{-2L\delta^2}$. Similarly, one

254   can show that $\mathbb{P}(\sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} (\text{MoM}_L^n(f_{\boldsymbol{\Theta}}) - Pf_{\boldsymbol{\Theta}}) > \epsilon) \le e^{-2L\delta^2}$. Combining the above

255   two inequalities, we get, $\mathbb{P}\left(\sup_{\boldsymbol{\Theta} \in [-M,M]^{k \times p}} |\text{MoM}_L^n(f_{\boldsymbol{\Theta}}) - Pf_{\boldsymbol{\Theta}}| > \epsilon\right) \le 2e^{-2L\delta^2}$. Finally,

256   plugging in the values of ϵ and δ yields the desired result.

257   The following corollary gives a non-asymptotic control over the difference $|Pf_{\widehat{\boldsymbol{\Theta}}_n^{(\text{MoM})}} - Pf_{\boldsymbol{\Theta}^*}|$ in

258   terms of the model parameters.

259   **Corollary 3.5.** *Under A2—3 and A5—6, with probability at least $1 - 2e^{-2L\delta^2}$, the following holds.*

$$|Pf_{\widehat{\boldsymbol{\Theta}}_n^{(MoM)}} - Pf_{\boldsymbol{\Theta}^*}| \lesssim \tau_{\boldsymbol{\alpha},k} H_p \max\left\{ kp\sqrt{\frac{L}{n}}, (kp)^{3/2} \frac{\sqrt{|\mathcal{I}|}}{n} \right\}.$$

260   ## 3.5   Inference for Fixed k and p Under the MoM Framework

261   We now focus on the classical domain where the numbers of clusters and features remain fixed. To

262   show $\widehat{\boldsymbol{\Theta}}_n^{(\text{MoM})}$ is consistent for $\boldsymbol{\Theta}^*$, we need to impose conditions such that the RHS of the bounds

263   presented in Corollary 3.5 decrease to 0 as $n \to \infty$. We state the required conditions as follows.

264   **A7.** *The number of partitions $L = o(n)$, and $L \to \infty$ as $n \to \infty$.*

These conditions are natural: as n grows, so too must L in order to maintain a proportion of outlier-free partitions. On the other hand, L must grow slowly relative to n to ensure each partition can be assigned sufficient numbers of datapoints. We note that A7 implies $|\mathcal{O}| = o(n)$, an intuitive and standard condition [25, 40, 35] as outliers should be few by definition.

In the following corollary, we focus on the squared Euclidean distance for center-based clustering under the MoM framework. We show that the (global) estimates obtained from MoM k-means, MoM power k-means etc. are consistent. We stress that the obtained convergence rate, as a function of n, L and $|\mathcal{I}|$, does *not* depend on the choice of $\Psi_{\boldsymbol{\alpha}}(\cdot)$, as long as it satisfies A2, i.e. is Lipschitz. In particular, replacing $\Psi_{\boldsymbol{\alpha}}(\cdot)$ with $\min_{1 \leq j \leq k} x_j$, $M_s(\boldsymbol{x})$ and $(\sum_{j=1}^{k} x_j^{-1})^{-1}$, the rate in Corollary 3.6 applies to robustified MoM versions of k-means, power k-means and k-harmonic means alike.

Corollary 3.6. *Suppose* $\phi(\boldsymbol{u}) = \|\boldsymbol{u}\|_2^2$ *and* $\Psi_{\boldsymbol{\alpha}}(\cdot)$ *satisfy A2. Then under, A2—3 and A5—7,* $|Pf_{\widehat{\boldsymbol{\Theta}}_n^{(MoM)}} - Pf_{\boldsymbol{\Theta}^*}| = O_P\left(\max\left\{L^{1/2}n^{-1/2}, n^{-1}\sqrt{|\mathcal{I}|}\right\}\right)$ *and* $Pf_{\widehat{\boldsymbol{\Theta}}_n^{(MoM)}} \xrightarrow{P} Pf_{\boldsymbol{\Theta}^*}$. *Further, under A4,* $\widehat{\boldsymbol{\Theta}}_n^{(MoM)} \xrightarrow{P} \boldsymbol{\Theta}^*$.

Remark   These results imply that any MoM center-based algorithm under our paradigm admits a convergence rate of $O\left(\max\left\{L^{1/2}n^{-1/2}, n^{-1}\sqrt{|\mathcal{I}|}\right\}\right)$, when equipped with squared Euclidean distance. Note that as $L \geq 1$, $\max\left\{L^{1/2}n^{-1/2}, n^{-1}\sqrt{|\mathcal{I}|}\right\} = \Omega\left(n^{-1/2}\right)$. Thus, the convergence rates for MoM variants in our framework are generally slower than their ERM counterparts, for which the rate is $O(n^{-1/2})$. This is unsurprising as MoM operates on outlier-contaminated data; there is "no free lunch" in trading off robustness for rate of convergence. However, if the number of partitions L grows slowly relative to n (say, $L = O(\log n)$ so that $|\mathcal{O}| = O(\log n)$), the convergence rates for MoM estimates become comparable to the ERM counterparts at $\widetilde{O}(n^{-1/2})$.

4   Empirical Studies and Results

We now turn to an empirical comparison of the proposed and peer clustering methods as evaluated under the Adjusted Rand Index (ARI) [19]. ARI values range between 0 and 1, where 1 denotes a perfect match with the ground truth. We consider a broad range of competing methods, comparing with k-means [29], Partition Around Medoids (PAM) [20], Robust kmeans++ (RKMpp) [14], Robust Continuous Clustering (RCC) [39], Bregman Trimmed k-means (BTKM) [17], MoM k-means (MOMKM) [21] and a novel MOM power k-means (MOMPKM) implied by our framework.

We consider two thorough simulated experiments below, with additional simulations and large-scale real data results, including a KDD Cup benchmark with roughly 5 million observations. These are available in the Supplement due to space restrictions, and convey the same trends as below. In all settings, we generate data in $p = 5$ dimensions, varying the number of clusters and the outlier percentages. True centers are spaced uniformly on a grid with $\theta_k = \frac{k-1}{10}$ and observations are drawn from Gaussians around their ground truth centers with variance 0.1. In the experiments, we take the number of partitions L to be roughly double the number of outliers, and η and α to be 1.02 and 1 as default values. Results are averaged over 20 random restarts, and all competitors are initialized and tuned according to the standard implementations described in their original papers.

Experiment 1: increasing number of clusters   The first experiment assesses performance as the number of true clusters grows, while keeping the proportion of outliers fixed at 25%. Datasets are generated with the number of clusters k varying between 3 and 100. For each setting, we create balanced clusters drawing 30 points from each true center. The 25% outliers are generated from a uniform distribution with support on the range of the inlying observations. We repeat this data generation process 30 times for each parameter setting. The average ARI values, plotted against the number of clusters along with error bars ($\pm$ sd), are shown in Fig. 1. We see that the robustified version of power k-means implied by our framework (labeled MOMPKM) achieves the best performance. This may be unsurprising as the recent power k-means method was shown to significantly reduce sensitivity to local minima (which tend to increase with k), while MoM further protects the algorithm from outlier influence.

Figure 1: Average ARI values, along with error bars (±sd), comparing the peer algorithms in Experiments 1 (left) and 2 (right). MOMPKM remains relatively stableeven when k or outlier percentages increase, maintaining the best performance among peer methods.

Experiment 2: increasing outlier percentage Following the same data generation process in Experiment 1, we now fix $k = 20$ while varying the outlier percentage from 0% to 50%. For each parameter setting, we again replicate the experiment 30 times. Average ARI values comparing the inlying observations to their ground truths are plotted in the right panel of Figure 1. Not only does this study convey similar trends as Experiment 1, but we see that competing methods continue to deteriorate with increasing outliers, while MOMPKM remarkably remains stable.

It is likely that the ERM-based methods like BTKM, MOMKM, PAM fail due to the presence of high number of clusters, whereas RKMpp often stops short at poor local optima, made worse by large numbers of outliers despite initialization via clever seeding techniques. This phenomenon is also reported in literature for their non-robust counterparts [49, 13]. Overall, the empirical study suggests that a novel MoM-based power k-means method shows promise to handle several data challenges at once, in line with our theoretical analysis.

5 Discussion

In this paper, we proposed a paradigm for center-based clustering that gives a closure or unification of a suite of center-based clustering algorithms. Under this unifying view, developed a simple yet efficient framework for robust versions of such algorithms by appealing to the Median of Means (MoM) philosophy. Using gradient-based methods, the MoM objectives can be solved with the same per-iteration complexity of Lloyd's k-means algorithm, retaining its simplicity. Importantly, we derive a thorough analysis of the statistical properties and convergence rates by establishing uniform concentration bounds under i.i.d. sampling of the data. Our novel theoretical contributions demonstrate how arguments utilizing Rademacher complexities and Dudley's chaining arguments can be leveraged in the robust clustering context. As a result, we are able to obtain error rates that do not require asymptotic assumptions, nor restrictions on the relation between n and p. These imply asymptotic results such as strong consistency and $\sqrt{n}$-consistency under classical assumptions.

As shown in the paper, the robustness of MoM estimators comes at a cost of slower convergence rates than their ERM counterparts. We emphasize that there is *no median of means magic*, and the efficacy of MoM depends on the interplay between the partitions and the outliers. If the number of partitions circumvents the impact of the outliers, the performance of MoM clustering estimates under our framework scales with the block size b as $1/\sqrt{b} = \sqrt{L/n}$. Since L can be chosen to be approximately $2|\mathcal{O}|$, the obtained error rate is roughly $O(\sqrt{|\mathcal{O}|/n})$. If $|\mathcal{O}|$ scales proportionately with n, however, the error bound of $O(\sqrt{|\mathcal{O}|/n})$ becomes meaningless. For our consistency results to hold, it is crucial that $|\mathcal{O}| = o(n)$, which in turn allows us to choose L that satisfies A7. If $|\mathcal{O}| = O(n^\beta)$, for some $0 < \beta < 1$, the error rate is $O(n^{(\beta-1)/2})$. Future research in this direction might render fruitful avenues in improving the "slow" ERM rates via finding so called fast rates under additional assumptions [10, 48]. One can also focus on finding lower bounds on the approximation error or can explore high-dimensional robust center-based clustering under the proposed paradigm.

References

[1] D. Aloise, A. Deshpande, P. Hansen, and P. Popat. Np-hardness of euclidean sum-of-squares clustering. *Machine Learning*, 75(2):245–248, 2009.

[2] K. B. Athreya and S. N. Lahiri. *Measure theory and probability theory*. Springer Science & Business Media, 2006.

[3] M.-F. Balcan, A. Blum, and S. Vempala. A discriminative framework for clustering via similarity functions. In *Proceedings of the fortieth annual ACM symposium on Theory of computing*, pages 671–680, 2008.

[4] A. Banerjee, S. Merugu, I. S. Dhillon, J. Ghosh, and J. Lafferty. Clustering with bregman divergences. *Journal of Machine Learning Research*, 6(10), 2005.

[5] P. L. Bartlett and S. Mendelson. Rademacher and gaussian complexities: Risk bounds and structural results. *Journal of Machine Learning Research*, 3(Nov):463–482, 2002.

[6] P. L. Bartlett, S. Boucheron, and G. Lugosi. Model selection and error estimation. *Machine Learning*, 48(1):85–113, 2002.

[7] P. L. Bartlett, O. Bousquet, and S. Mendelson. Local rademacher complexities. *The Annals of Statistics*, 33(4):1497–1537, 2005.

[8] G. Beliakov, T. Calvo, and S. James. On lipschitz properties of generated aggregation functions. *Fuzzy Sets and Systems*, 161(10):1437–1447, 2010.

[9] S. Ben-David. A framework for statistical clustering with constant time approximation algorithms for k-median and k-means clustering. *Machine Learning*, 66(2):243–257, 2007.

[10] S. Boucheron, O. Bousquet, and G. Lugosi. Theory of classification: A survey of some recent advances. *ESAIM: Probability and Statistics*, 9:323–375, 2005.

[11] C. Brunet-Saumard, E. Genetay, and A. Saumard. K-bmom: a robust lloyd-type clustering algorithm based on bootstrap median-of-means. *arXiv preprint arXiv:2002.03899*, 2020.

[12] S. Bubeck, N. Cesa-Bianchi, and G. Lugosi. Bandits with heavy tail. *IEEE Transactions on Information Theory*, 59(11):7711–7717, 2013.

[13] S. Chakraborty, D. Paul, S. Das, and J. Xu. Entropy weighted power k-means clustering. In *International Conference on Artificial Intelligence and Statistics*, pages 691–701. PMLR, 2020.

[14] A. Deshpande, P. Kacham, and R. Pratap. Robust k-means++. In *Conference on Uncertainty in Artificial Intelligence*, pages 799–808. PMLR, 2020.

[15] J. Duchi, E. Hazan, and Y. Singer. Adaptive subgradient methods for online learning and stochastic optimization. *Journal of Machine Learning Research*, 12(61):2121–2159, 2011. URL http://jmlr.org/papers/v12/duchi11a.html.

[16] R. M. Dudley. The sizes of compact subsets of hilbert space and continuity of gaussian processes. *Journal of Functional Analysis*, 1(3):290–330, 1967.

[17] A. Fischer, C. Levrard, and C. Brécheteau. Robust bregman clustering. *Annals of Statistics*, 2020.

[18] I. Goodfellow, Y. Bengio, A. Courville, and Y. Bengio. *Deep learning*, volume 1. MIT press Cambridge, 2016.

[19] L. Hubert and P. Arabie. Comparing partitions. *Journal of Classification*, 2(1):193–218, 1985.

[20] L. Kaufman and P. J. Rousseeuw. *Finding groups in data: an introduction to cluster analysis*, volume 344. John Wiley & Sons, 2009.

[21] Y. Klochkov, A. Kroshnin, and N. Zhivotovskiy. Robust k-means clustering for distributions with two moments. *arXiv preprint arXiv:2002.02339*, 2020.

[22] P. Laforgue, S. Clémençon, and P. Bertail. On medians of (randomized) pairwise means. In *International Conference on Machine Learning*, pages 1272–1281. PMLR, 2019.

[23] G. Lecué and M. Lerasle. Learning from mom's principles: Le cam's approach. *Stochastic Processes and Their Applications*, 129(11):4385–4410, 2019.

[24] G. Lecué and M. Lerasle. Robust machine learning by median-of-means: theory and practice. *Annals of Statistics*, 48(2):906–931, 2020.

[25] G. Lecué, M. Lerasle, and T. Mathieu. Robust classification via mom minimization. *Machine Learning*, 109(8):1635–1665, 2020.

[26] M. Lerasle. Lecture notes: Selected topics on robust statistical learning theory. *arXiv preprint arXiv:1908.10761*, 2019.

[27] S. Lloyd. Least squares quantization in PCM. *IEEE Transactions on Information Theory*, 28(2): 129–137, 1982.

[28] G. Lugosi and S. Mendelson. Regularization, sparse recovery, and median-of-means tournaments. *Bernoulli*, 25(3):2075–2106, 2019.

[29] J. MacQueen. Some methods for classification and analysis of multivariate observations. In *Proceedings of the fifth Berkeley symposium on mathematical statistics and probability*, volume 1, pages 281–297. Oakland, CA, USA, 1967.

[30] T. Mathieu and S. Minsker. Excess risk bounds in robust empirical risk minimization. *Information and Inference: A Journal of the IMA*, 2021.

[31] S. Minsker. Uniform bounds for robust mean estimators. *arXiv preprint arXiv:1812.03523*, 2018.

[32] M. Mohri, A. Rostamizadeh, and A. Talwalkar. *Foundations of machine learning*. MIT press, 2018.

[33] A. Ng, M. Jordan, and Y. Weiss. On spectral clustering: Analysis and an algorithm. *Advances in Neural Information Processing Systems*, 14:849–856, 2001.

[34] R. Ostrovsky, Y. Rabani, L. J. Schulman, and C. Swamy. The effectiveness of lloyd-type methods for the k-means problem. *Journal of the ACM (JACM)*, 59(6):1–22, 2013.

[35] D. Paul, S. Chakraborty, and S. Das. Robust principal component analysis: A median of means approach. *arXiv preprint arXiv:2102.03403*, 2021.

[36] D. Paul, S. Chakraborty, and S. Das. On the uniform concentration bounds and large sample properties of clustering with bregman divergences. *Stat*, page e360, 2021.

[37] D. Pollard. Strong consistency of k-means clustering. *The Annals of Statistics*, 9(1):135–140, 1981.

[38] D. Rodriguez and M. Valdora. The breakdown point of the median of means tournament. *Statistics & Probability Letters*, 153:108–112, 2019.

[39] S. A. Shah and V. Koltun. Robust continuous clustering. *Proceedings of the National Academy of Sciences*, 114(37):9814–9819, 2017.

[40] G. Staerman, P. Laforgue, P. Mozharovskyi, and F. d'Alché Buc. When ot meets mom: Robust estimation of wasserstein distance. In A. Banerjee and K. Fukumizu, editors, *Proceedings of The 24th International Conference on Artificial Intelligence and Statistics*, volume 130 of *Proceedings of Machine Learning Research*, pages 136–144. PMLR, 13–15 Apr 2021. URL `http://proceedings.mlr.press/v130/staerman21a.html`.

[41] M. Teboulle. A unified continuous optimization framework for center-based clustering methods. *Journal of Machine Learning Research*, 8(1), 2007.

[42] M. J. Telgarsky and S. Dasgupta. Agglomerative bregman clustering. In *29th International Conference on Machine Learning, ICML 2012*, pages 1527–1534, 2012.

[43] M. J. Telgarsky and S. Dasgupta. Moment-based uniform deviation bounds for κ-means and friends. *Advances in Neural Information Processing Systems*, 2013.

[44] Y. Terada. Strong consistency of reduced k-means clustering. *Scandinavian Journal of Statistics*, 41(4):913–931, 2014.

[45] Y. Terada. Strong consistency of factorial k-means clustering. *Annals of the Institute of Statistical Mathematics*, 67(2):335–357, 2015.

[46] V. Vapnik. *The nature of statistical learning theory*. Springer science & business media, 2013.

[47] S. Vassilvitskii and D. Arthur. k-means++: The advantages of careful seeding. In *Proceedings of the Eighteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1027–1035, 2006.

[48] M. J. Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge University Press, 2019.

[49] J. Xu and K. Lange. Power k-means clustering. In *International Conference on Machine Learning*, pages 6921–6931. PMLR, 2019.

[50] B. Zhang, M. Hsu, and U. Dayal. K-harmonic means-a data clustering algorithm. *Hewlett-Packard Labs Technical Report HPL-1999-124*, 55, 1999.

[51] Z. Zhang, K. Lange, and J. Xu. Simple and scalable sparse k-means clustering via feature ranking. *Advances in Neural Information Processing Systems*, 33, 2020.