LAST ITERATE CONVERGENCE IN MONOTONE MEAN FIELD GAMES

ABSTRACT

Mean Field Game (MFG) is a framework utilized to model and approximate the behavior of a large number of agents, and the computation of equilibria in MFG has been a subject of interest. Despite the proposal of methods to approximate the equilibria, algorithms where the sequence of updated policy converges to equilibrium, specifically those exhibiting last-iterate convergence, have been limited. We propose the use of a simple, proximal-point-type algorithm to compute equilibiria for MFGs. Subsequently, we provide the first last-iterate convergence guarantee under the Lasry–Lions-type monotonicity condition. We further employ the Mirror Descent algorithm for the regularized MFG to efficiently approximate the update rules of the proximal point method for MFGs. We demonstrate that the algorithm can approximate with an accuracy of ε after $\mathcal{O}(\log(1/\varepsilon))$ iterations. This research offers a tractable approach for large-scale and large-population games.

1 INTRODUCTION

Mean Field Games (MFGs) provide a simple and powerful framework for approximating the behavior of large populations of interacting agents. Originally formulated by Lasry & Lions (2007); Huang et al. (2006), MFGs model the collective behavior of homogeneous agents in continuous time and state settings using Partial Differential Equations (PDEs) (Cardaliaguet & Hadikhanloo, 2017; Lavigne & Pfeiffer, 2023; Inoue et al., 2023). Subsequently, the formulation of MFGs using Markov Decision Processes (Bertsekas & Shreve, 1978; Puterman, 1994) has enabled the study of discrete-time and discrete-state models (Gomes et al., 2010), broadening the applicability of MFGs to Multi-Agent Reinforcement Learning (MARL) (Yang et al., 2018). Moreover, it has become possible to capture interactions among heterogeneous agents (Gao & Caines, 2017; Caines & Huang, 2019).

The applicability of MFGs to MARL drives research into their computational aspects. Under fairly general assumptions, the problem of finding an equilibrium in MFGs is known to be PPAD-complete (Yardim et al., 2024). Consequently, it would be essential to impose assumptions that allow for the existence of algorithms capable of efficiently computing an equilibrium. One of the assumptions is contractivity (Xie et al., 2021; Anahtarci et al., 2023; Yardim et al., 2023). However, it is known that many problems are not contractive in practice (Cui & Koeppl, 2021). One of the more realistic assumptions is monotonicity (Pérolat et al., 2022; Zhang et al., 2023; Yardim & He, 2024), which intuitively implies that as more

Figure 1: Overview of Algorithms

agents converge to a single state, the reward monotonically decreases. Under the monotonicity assumption, Online Mirror Descent (OMD) has been proposed and widely adopted (Pérolat et al., 2022; Cui & Koeppl, 2022; Lauriere et al., 2022; Fabian et al., 2023). OMD, especially when combined with function approximation via deep learning, has enabled the application of MFGs to MARL (Yang & Wang, 2021; Zhang et al., 2021; Cui et al., 2022).

Theoretically, *last-iterate convergence (LIC)*, which ensures that the policy obtained in the final iteration converges, is particularly important in deep learning settings due to the constraints imposed by neural networks (NN). In NNs, calculating the time-averaged policy like in the celebrated Fictitious

Play method (Brown, 1951; Perrin et al., 2020) may be less meaningful due to nonlinearity in the parameter space. These considerations have spurred significant research into developing algorithms that achieve LIC in finite N-player games, as seen in, e.g., Mertikopoulos et al. (2018); Piliouras et al. (2022); Abe et al. (2023; 2024).

Despite its importance, the literature on LIC results in MFG is quite limited. The only exception is Pérolat et al. (2022), who proved the LIC result for the continuous-time version of OMD without the quantified rates under the strict monotonicity condition. The aim of this research is to establish an online learning algorithm that can achieve LIC in MFGs under *non-strict* monotonicity conditions.

In this paper, we propose a novel proximal-point (PP) type algorithm and prove that it achieves LIC under the non-strict monotonicity assumption. Furthermore, we demonstrate that the update rule of the PP can be approximated efficiently by sequentially using the Regularized Mirror Descent (RMD). We further show that RMD achieves the approximation with the accuracy of ε within $\mathcal{O}(\log(1/\varepsilon))$ iterations. Figure 1 summarizes the overview of the algorithms in this paper.

In summary, the contributions of this paper are as follows:

> **Contribution**
>
> (i) We construct the first algorithm based on the celebrated PP method that achieves LIC for general monotone MFGs (Theorem 4.3).
>
> (ii) We prove for the first time that regularized Mirror Descent achieves exponential convergence for monotone MFGs (Theorem 4.4).
>
> (iii) We combine these two algorithms as shown in Figure 1 to develop a tractable algorithm that approximates the PP-based method (Algorithm 2).

The organization of this paper is as follows: In Section 2, we review the fundamental concepts of MFGs. In Section 3, we introduce the PP method and its convergence results. In Section 4, we present the RMD algorithm and its convergence properties. Finally, in Section 5, we propose a combined approximation method, demonstrating its convergence through experimental validation.

2 SETTING AND PRELIMINARY FACT

2.1 NOTATION

For a positive integer $N \in \mathbb{N}$, $[N] := \{1, \ldots, N\}$. For a finite set X, $\Delta(X) := \{p \in \mathbb{R}_{\geq 0}^{|X|} \mid \sum_{x \in X} p(x) = 1\}$. For a function $f: X \to \mathbb{R}$ and a probability $\pi \in \Delta(X)$, $\langle f, \pi \rangle := \langle f(\bullet), \pi(\bullet) \rangle := \sum_{x \in X} f(x)\pi(x)$. For $p^0, p^1 \in \Delta(X)$, define the Kullback–Leibler (KL) divergence $D_{\mathrm{KL}}(p^0, p^1) := \sum_{x \in X} p^0(x) \log(p^0(x)/p^1(x))$, and the total variation (TV) distance as $\|p^0 - p^1\| := \sum_{x \in X} |p^0(x) - p^1(x)|$.

2.2 MEAN-FIELD GAMES

Consider a *Mean-Field Game (MFG)* that is defined through a tuple $(\mathcal{S}, \mathcal{A}, H, P, r, \mu_1)$. Here, $\mathcal{S}$ is a finite discrete space of states, $\mathcal{A}$ is a finite discrete space of actions, $H \in \mathbb{N}_{\geq 2}$ is a time horizon, and $P = (P_h)_{h=1}^H$ is a family of transition kernels $P_h: \mathcal{S} \times \mathcal{A} \to \Delta(\mathcal{S})$, that is, if a player with state $s_h \in \mathcal{S}$ takes action $a_h \in \mathcal{A}$ at time $h \in [H]$, the next state $s_{h+1} \in \mathcal{S}$ will transition according to $s_{h+1} \sim P_h(\cdot \mid s_h, a_h)$. In addition, $r = (r_h)_{h=1}^H$ is a family of reward functions $r_h: \mathcal{S} \times \mathcal{A} \times \Delta(\mathcal{S}) \to [0, 1]$, and $\mu_1 \in \Delta(\mathcal{S})$ is an initial probability of state. Note that, in the context of theoretical analysis of the online learning method for MFG (Pérolat et al., 2022; Zhang et al., 2023), P is assumed to be independent of the state distribution. It is reasonable to assume that at any time h, every state $s' \in \mathcal{S}$ is reachable:

Assumption 2.1. For each $(h, s') \in [H] \times \mathcal{S}$, there exists $(s, a) \in \mathcal{S} \times \mathcal{A}$ such that $P_h(s' \mid s, a) > 0$.

In this paper, we focus on rewards r that satisfy the following two typical conditions, which are also assumed in Perrin et al. (2020; 2022); Pérolat et al. (2022); Fabian et al. (2023); Zhang et al. (2023). The first one is *monotonicity* of the type introduced by Lasry & Lions (2007), which means,

under a state distribution $\mu = (\mu_h)_{h=1}^H \in \Delta(\mathcal{S})^H$, if players choose a strategy—called a policy $\pi = (\pi_h)_{h=1}^H \in (\Delta(\mathcal{A})^{\mathcal{S}})^H$ to be planned—that concentrates on a state or action, they will receive a small reward.

Assumption 2.2 (weak monotonicity of r). For all $\mu, \widetilde{\mu} \in \Delta(\mathcal{S})^H$, $\pi, \widetilde{\pi} \in (\Delta(\mathcal{A})^{\mathcal{S}})^H$, it holds that

$$\sum_{h=1}^H \sum_{(s,a) \in \mathcal{S} \times \mathcal{A}} (r_h(s, a, \mu_h) - r_h(s, a, \widetilde{\mu}_h))(\pi_h(a \mid s)\, \mu_h(s) - \widetilde{\pi}_h(a \mid s)\, \widetilde{\mu}_h(s)) \le 0.$$

For example, a reward r that satisfies these assumptions includes a model of a crowd that avoids overcrowding.

The second is the Lipschitz continuity of the reward r with respect to $\mu \in (\Delta(\mathcal{S}))^H$, which is a standard assumption in the field of MFGs (Cui & Koeppl, 2021; Fabian et al., 2023; Zhang et al., 2023).

Assumption 2.3 (Lipschitz continuity of r). There exists a constant L such that for every $h \in [H]$, $s \in \mathcal{S}$, $a \in \mathcal{A}$, and $\mu, \mu' \in \Delta(\mathcal{S})$:

$$|r_h(s, a, \mu) - r_h(s, a, \mu')| \le L\|\mu - \mu'\|.$$

Given a policy π, the probabilities $m[\pi] = (m[\pi]_h)_{h=1}^H \in \Delta(\mathcal{S})^H$ of the state is recursively defined as follows: $m[\pi]_1 = \mu_1$ and

$$m[\pi]_h(s_h) = \sum_{(s_{h-1}, a_{h-1}) \in \mathcal{S} \times \mathcal{A}} \pi_{h-1}(a_{h-1} \mid s_{h-1})\, P_{h-1}(s_h \mid s_{h-1}, a_{h-1})\, m[\pi]_{h-1}(s_{h-1}),$$

(2.1)

if $h = 2, \ldots, H$. We plan to maximize the following cumulative reward

$$J(\mu, \pi) \coloneqq \sum_{h=1}^H \sum_{(s_h, a_h) \in \mathcal{S} \times \mathcal{A}} \pi_h(a_h \mid s_h)\, m[\pi]_h(s_h) r_h(s_h, a_h, \mu_h), \tag{2.2}$$

under a probability $\mu \in \Delta(\mathcal{S})^H$ of states. The *mean-field equilibrium* defined below means the pair of probabilities μ and policies π that achieves the maximum under the constraints (2.1).

Definition 2.4. A pair $(\mu^\star, \pi^\star) \in \Delta(\mathcal{S})^H \times (\Delta(\mathcal{A})^{\mathcal{S}})^H$ is a *mean-field equilibrium* if it satisfies (i) $J(\mu^\star, \pi^\star) = \max_{\pi \in \Delta(\mathcal{S})^H} J(\mu^\star, \pi)$, and (ii) $\mu^\star = m[\pi^\star]$. In addition, set $\Pi^\star \subset (\Delta(\mathcal{A})^{\mathcal{S}})^H$ as the set of all policies that are in equilibrium.

Under Assumptions 2.2 and 2.3, there exists a mean-field equilibrium, see the proof of (Saldi et al., 2018, Theorem 3.3.) and (Pérolat et al., 2022, Proposition 1.). Note that the equilibrium may not be unique if the inequality in Assumption 2.2 is non-strict. In other words, the set $\Pi^\star \subset (\Delta(\mathcal{A})^{\mathcal{S}})^H$ is not singleton in general. As an illustrative example, one might consider the trivial case where $r \equiv 0$. Our goal is to construct an algorithm that generates policies that converge to $\Pi^\star$.

3 PROXIMAL POINT-TYPE METHOD FOR MFG

3.1 ALGORITHM

This section presents an algorithm motivated by the proximal point (PP) method. Let $\lambda > 0$ be a sufficiently small positive number, roughly "the inverse of learning rate." In the algorithm proposed in this paper, we generate a sequence $((\sigma^k, \mu^k))_{k=0}^\infty \subset (\Delta(\mathcal{A})^{\mathcal{S}})^H \times \Delta(\mathcal{S})^H$ as

$$\sigma^{k+1} = \arg\max_{\pi \in (\Delta(\mathcal{A})^{\mathcal{S}})^H} \left\{ J(\mu^{k+1}, \pi) - \lambda D_{m[\pi]}(\pi, \sigma^k) \right\}, \quad \mu^{k+1} = m[\sigma^{k+1}], \tag{3.1}$$

where m is defined in (2.1) and $D_\mu(\pi, \sigma^k) \coloneqq \sum_h \mathbb{E}_{s \sim \mu_h} \left[D_{\mathrm{KL}}(\pi_h(s), \sigma_h^k(s)) \right]$ with a probability $\mu \in \Delta(\mathcal{S})^H$. If the initial policy π^0 has full support, i.e., $\min_{(h,s,a) \in H \times \mathcal{S} \times \mathcal{A}} \pi_h^0(a \mid s) > 0$, the rule (3.1) is well-defined, see Proposition C.1.

Algorithm 1: Proximal point (PP) method with KL divergence for MFG

Input: MFG $(\mathcal{S}, \mathcal{A}, H, P, r, \mu_1)$, initial policy π^0, number of iterations N, parameter $\lambda > 0$

1 Initialization: Set $k \leftarrow 0$ and $\sigma^k \leftarrow \pi^0$;

2 while $k < N$ **do**

3   Compute $(\mu^{k+1}, \sigma^{k+1})$ by solving the regularized MFG;

4

$$
\begin{cases}
\sigma^{k+1} = \underset{\pi \in (\Delta(\mathcal{A})^\mathcal{S})^H}{\arg\max} \ \{ J(\mu^{k+1}, \pi) - \lambda D_{m[\pi]}(\pi, \sigma^k) \}, \\
\mu^{k+1} = m[\sigma^{k+1}]
\end{cases}
$$

Hard to solve explicitly

    Update $k \leftarrow k + 1$;

Output: $\sigma^k (\approx \pi^\star)$

Interestingly, the rule (3.1) is similar to the traditional proximal point (PP) method with KL divergence in mathematical optimization and Optimal Transport, see (Censor & Zenios, 1992; Xie et al., 2019) and the pseudocode in Algorithm 1. Therefore, we also refer to this update rule as the PP method. On the other hand, unlike the traditional PP method, our method changes the objective function $J(\mu^k, \bullet) \colon (\Delta(\mathcal{A})^\mathcal{S})^H \to \mathbb{R}$ with each iteration $k \in \mathbb{N}$. Therefore, the convergence of our traditional method is not directly derived from traditional theory.

3.2 LAST-ITERATE CONVERGENCE RESULT

The following theorem implies the last-iterate convergence of the policies generated by (3.1). Specifically, it shows that under the assumptions above, the sequence of policies converges to the equilibrium set. This result is crucial for the effectiveness of the algorithm in reaching an optimal policy.

Theorem 3.1. *Let $(\sigma^k)_{k=0}^\infty$ be the sequence defined by Algorithm 1. In addition to Assumptions 2.1 to 2.3, assume that the initial policy π^0 has full support, i.e., $\min_{(h,s,a) \in H \times \mathcal{S} \times \mathcal{A}} \pi_h^0(a \mid s) > 0$. Then, the sequence $(\sigma^k)_{k=0}^\infty$ converges to the set $\Pi^\star$ of equilibrium, i.e.,*

$$\lim_{k \to \infty} \operatorname{dist}(\sigma^k, \Pi^\star) = 0,$$

where $\operatorname{dist}(\pi, \Pi^\star) \coloneqq \inf_{\pi^\star \in \Pi^\star} \sum_{(h,s) \in [H] \times \mathcal{S}} \|\pi_h(s) - \pi_h^\star(s)\|$.

Proof sketch of Theorem 3.1. If we accept the next two lemmas, we can easily prove Theorem 3.1: The first implies that the KL divergence from an equilibrium to the generated policy becomes smaller as the cumulative reward J increases.

Lemma 3.2. *Suppose Assumption 2.2. Then, for any equilibrium $(\mu^\star, \pi^\star)$ it holds that*

$$D_{\mu^\star}(\pi^\star, \sigma^{k+1}) - D_{\mu^\star}(\pi^\star, \sigma^k) \leq J(\mu^\star, \sigma^{k+1}) - J(\mu^\star, \pi^\star).$$

Furthermore, we can control the right-hand side of the inequality in Lemma 3.2 by the distance:

Lemma 3.3. *There exist positive constants α and C such that, for any $\pi \in (\Delta(\mathcal{A})^\mathcal{S})^H$,*

$$J(\mu^\star, \pi) - J(\mu^\star, \pi^\star) \leq -C(\operatorname{dist}(\pi, \Pi^\star))^\alpha.$$

Combining these lemmas yields that $D_{\mu^\star}(\pi^\star, \sigma^{k+1}) - D_{\mu^\star}(\pi^\star, \sigma^k) \leq -C\big(\text{dist}(\sigma^{k+1}, \Pi^\star)\big)^\alpha$. Thus, the telescoping sum of this inequality yields

$$\sum_{k=1}^{\infty} \big(\text{dist}(\sigma^k, \Pi^\star)\big)^\alpha \leq \frac{1}{C} D_{\mu^\star}(\pi^\star, \pi^0) < +\infty.$$

Therefore, $\lim_{k\to\infty} \text{dist}(\sigma^k, \Pi^\star) = 0$. $\square$

Thus, the non-trivial aspects of the last-iterate convergence lie in the proof of Lemmas 3.2 and 3.3; see Appendix B.

4 APPROXIMATING PROXIMAL POINT WITH MIRROR DESCENT IN REGULARIZED MFG

As in the previous section, in the PP method in Algorithm 1, it is necessary to solve the regularized MFG (3.1) at each iteration. Therefore, this section introduces Regularized Mirror Descent (RMD), which approximates the solution $(\mu^{k+1}, \sigma^{k+1})$ of (3.1) for each policy σ^k. The novel result in this section is that the divergence between the sequence of RMD and the equilibrium exponentially decays.

4.1 APPROXIMATION OF THE UPDATE RULE OF PP WITH REGULARIZED MFG

Fortunately, solving (3.1) corresponds to finding an equilibrium for *KL-regularized MFG* introduced in Cui & Koeppl (2021); Zhang et al. (2023). Let us review the settings for the regularized MFG. For each parameter $\lambda > 0$ and policy $\sigma \in (\Delta(\mathcal{A})^{\mathcal{S}})^H$, which plays the role of σ^k in Algorithm 1, we define the *regularized cumulative reward* $J^{\lambda,\sigma} : \Delta(\mathcal{S})^H \times (\Delta(\mathcal{A})^{\mathcal{S}})^H \ni (\mu, \pi) \mapsto J^{\lambda,\sigma}(\mu, \pi) \in \mathbb{R}$ to be

$$J^{\lambda,\sigma}(\mu, \pi) := J(\mu, \pi) - \lambda D_{m[\pi]}(\pi, \sigma). \tag{4.1}$$

Since σ is a representative of $(\sigma^k)_k$, the assumption of full support is also imposed on σ:

Assumption 4.1. The base σ has full support, i.e., $\sigma_{\min} := \min_{(s,a,h)\mathcal{S}\times\mathcal{A}\times[H]} \sigma_h(a \mid s) > 0$.

For the reward $J^{\lambda,\sigma}$, we introduce a *regularized equilibrium*:

Definition 4.2. A pair $(\mu^*, \varpi^*) \in \Delta(\mathcal{S})^H \times (\Delta(\mathcal{A})^{\mathcal{S}})^H$ is *regularized equilibrium* of $J^{\lambda,\sigma}$ if it satisfies (i) $J^{\lambda,\sigma}(\mu^*, \varpi^*) = \max_{\pi \in \Delta(\mathcal{S})^H} J^{\lambda,\sigma}(\mu^*, \pi)$, and (ii) $\mu^* = m[\varpi^*]$.

Specifically, $(\mu^{k+1}, \sigma^{k+1})$ can be characterized as the regularized equilibrium of J^{λ,σ^k} for $k \in \mathbb{N}$. Note that the regularized equilibrium is unique under Assumption 4.1, see Appendix C.

In the next subsection, we will introduce RMD using *value functions*, which are defined as follows: for each $h \in [H]$, $s \in \mathcal{S}$, $a \in \mathcal{A}$, $\mu \in \Delta(\mathcal{S})^H$ and $\pi \in \Delta(\mathcal{A})^{\mathcal{S}}$, define the *state value function* $V_h^{\lambda,\sigma} : \mathcal{S} \times \Delta(\mathcal{S})^H \times (\Delta(\mathcal{A})^{\mathcal{S}})^H \to \mathbb{R}_{\geq 0}$ and the *state-action value function* $Q_h^{\lambda,\sigma} : \mathcal{S} \times \mathcal{A} \times \Delta(\mathcal{S})^H \times (\Delta(\mathcal{A})^{\mathcal{S}})^H \to \mathbb{R}_{\geq 0}$ as

$$V_h^{\lambda,\sigma}(s, \mu, \pi) := \mathbb{E}_{((s_l,a_l))_{l=h}^H} \left[\sum_{l=h}^H \left(r_l(s_l, a_l, \mu_l) - \lambda D_{\mathrm{KL}}(\pi_l(s_l), \sigma_l(s_l)) \right) \,\Bigg|\, s_h = s \right], \tag{4.2}$$

$$V_{H+1}^{\lambda,\sigma}(s, \mu, \pi) := 0,$$

$$Q_h^{\lambda,\sigma}(s, a, \mu, \pi) = r_h(s, a, \mu_h) + \mathbb{E}_{s_{h+1}\sim P(s,a,\mu_h)} \left[V_{h+1}^{\lambda,\sigma}(s_{h+1}, \mu, \pi) \right]. \tag{4.3}$$

Here, the discrete time stochastic process $((s_l, a_l))_{l=h}^H$ is induced recursively as $s_{l+1} \sim P_l(s_l, a_l), a_l \sim \pi_l(s_l)$ for each $l \in \{h, \ldots, H-1\}$ and $a_H \sim \pi_H(s_H)$. Note that the the objective function $J^{\lambda,\sigma}$ in Definition 4.2 can be expressed as $J^{\lambda,\sigma}(\mu, \pi) = \mathbb{E}_{s\sim\mu_1}[V_1^{\lambda,\sigma}(s, \mu, \pi)]$.

4.2 AN EXPONENTIAL CONVERGENCE RESULT OF REGULARIZED MIRROR DESCENT

In this subsection, we introduce the iterative method for finding the regularized equilibrium proposed by Zhang et al. (2023) as RMD. The method constructs a sequence $((\pi^t, \mu^t))_{t=0}^{\infty} \subset (\Delta(\mathcal{A})^{\mathcal{S}})^H \times$

Algorithm 2: Practical version of Algorithm 1 for MFG

Input: $\mathtt{MFG}(\mathcal{S}, \mathcal{A}, H, P, r, \mu_1)$, initial policy π^0, number of iterations N, parameter $\lambda > 0$

1 Initialization: Set $k \leftarrow 0$ and $\sigma^k \leftarrow \pi^0$;

2 while $k < N$ **do**

3   Compute $(\mu^{k+1}, \sigma^{k+1})$ by solving the regularized MFG;

4

$$
\begin{cases}
\sigma^{k+1} = \mathtt{RMD}(\mathtt{MFG}, \sigma^k, \lambda, \eta, \sigma^k, \tau), \\
\mu^{k+1} = m[\sigma^{k+1}]
\end{cases}
$$

Tractable approximation of Algorithm 1 via RMD

    Update $k \leftarrow k + 1$;

Output: $\sigma^k (\approx \pi^\star)$

5

---

6 Function $\mathtt{RMD}(\mathtt{MFG}, \pi^0, \lambda, \eta, \sigma^0, \tau)$**:**

7   **Initialization:** Set $t \leftarrow 0$, $\pi^t \leftarrow \pi^0$ and $\sigma \leftarrow \sigma^0$;

8   **while** $t < \tau$ **do**

9   Compute $\mu^t = m[\pi^t]$;

10   Compute $Q_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t)$ $((h, s, a) \in [H] \times \mathcal{S} \times \mathcal{A})$ by (4.3);

11   Compute π^{t+1} as, for $(h, s, a) \in [H] \times \mathcal{S} \times \mathcal{A}$,

$$
\pi_h^{t+1}(a \mid s) = \frac{(\sigma_h(a \mid s))^{\lambda\eta} (\pi_h^t(a \mid s))^{1-\lambda\eta} \exp\left(\eta Q_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t)\right)}{\sum_{a' \in \mathcal{A}} (\sigma_h(a' \mid s))^{\lambda\eta} (\pi_h^t(a' \mid s))^{1-\lambda\eta} \exp\left(\eta Q_h^{\lambda,\sigma}(s, a', \pi^t, \mu^t)\right)}
$$

       Update $t \leftarrow t + 1$;

12   **return** π^t;

$\Delta(\mathcal{S})^H$ approximating the regularized equilibrium of $J^{\lambda,\sigma}$ using the following rule:

$$
\pi_h^{t+1}(s) = \arg\max_{p \in \Delta(\mathcal{A})} \left\{ \frac{\eta}{1 - \lambda\eta} \left(\left\langle Q_h^{\lambda,\sigma}(s, \bullet, \pi^t, \mu^t), p \right\rangle - \lambda D_{\mathrm{KL}}(p, \sigma_h(s)) \right) - D_{\mathrm{KL}}(p, \pi_h^t(s)) \right\},
$$

$$
\mu^{t+1} = m[\pi^{t+1}], \tag{4.4}
$$

where $\eta > 0$ is another learning rate, and $Q_h^{\lambda,\sigma}$ is the state-action value function defined in (4.3). We give the pseudo-code of RMD in Algorithm 2.

For the sequence of policies in RMD, we can establish the convergence result as follows:

> **Theorem 4.3.** *Let* $((\mu^t, \pi^t))_{t=0}^\infty \subset \Delta(\mathcal{S})^H \times (\Delta(\mathcal{A})^{\mathcal{S}})^H$ *be the sequence generated by* (4.4), *and* $(\mu^*, \varpi^*) \in \Delta(\mathcal{S})^H \times (\Delta(\mathcal{A})^{\mathcal{S}})^H$ *be the regularized equilibrium given in Definition 4.2. In addition to Assumptions 2.2, 2.3, and 4.1, suppose that* $\eta \leq \eta^*$, *where* $\eta^* > 0$ *is the upper bound of the learning rate defined in* (D.5), *which only depends on* λ, σ, H *and* $|\mathcal{A}|$. *Then, the sequence* $(\pi^t)_{t=0}^\infty$ *satisfies*
>
> $$
> D_{\mu^*}(\varpi^*, \pi^{t+1}) \leq \left(1 - \frac{\lambda\eta}{2}\right) D_{\mu^*}(\varpi^*, \pi^t) \quad (t = 0, 1, \dots).
> $$
>
> *Accordingly,* $D_{\mu^*}(\varpi^*, \pi^t) \leq D_{\mu^*}(\varpi^*, \pi^0) \exp(-\lambda\eta t/2)$. *Clearly, the inequality states that an approximate policy* π^t *satisfying* $D_{\mu^*}(\varpi^*, \pi^t) < \varepsilon$ *can be obtained in* $\mathcal{O}(\log(1/\varepsilon))$ *iterations.*

4.3 Intuition for exponential convergence: Continuous-time version of Regularized Mirror Descent

The convergence of $(\pi^t)_{t=0}^{\infty}$ can be intuitively explained by considering a continuous limit $(\pi^t)_{t\geq 0}$ with respect to the time t of RMD. In this paragraph, we will use the idea of mirror flow (Krichene et al., 2015; Tzen et al., 2023; Deb et al., 2023) and continuous dynamics in games (Taylor & Jonker, 1978; Mertikopoulos et al., 2018; Pérolat et al., 2021; 2022) to observe the exponential convergence of the flow to equilibrium. According to Deb et al. (2023, (2.1)), the continuous curve of π should satisfy that

$$\frac{\mathrm{d}}{\mathrm{d}t}\pi_h^t\left(a \mid s\right) = \pi_h^t\left(a \mid s\right)\left(Q_h^{\lambda,\sigma}(s,a,\pi^t,\mu^t) - \lambda\log\frac{\pi_h^t\left(a \mid s\right)}{\sigma_h\left(a \mid s\right)}\right). \tag{4.5}$$

The flow induced by the dynamical system (4.5) converges to equilibrium *exponentially* as time t goes to infinity.

> **Theorem 4.4.** *Let π^t be a solution of* (4.5) *and ϖ^* be a regularized equilibrium defined in Definition 4.2. Suppose that Assumption 2.2. Then, it holds that*
>
> $$\frac{\mathrm{d}}{\mathrm{d}t}D_{\mu^*}(\varpi^*,\pi^t) \leq -\lambda D_{\mu^*}(\varpi^*,\pi^t),$$
>
> *for all $t \geq 0$. Moreover, the inequality implies $D_{\mu^*}(\varpi^*,\pi^t) \leq D_{\mu^*}(\varpi^*,\pi^0)\exp\left(-\lambda t\right)$.*

Technically, the non-Lipschitz continuity of the value function $Q_h^{\lambda,\sigma}(s,a,\bullet,\mu^t)$ in the right-hand side of (4.5) is non-trivial for the existence of the solution $\pi:[0,+\infty) \to (\Delta(\mathcal{A})^{\mathcal{S}})^H$ of the differential equation (4.5), see, e.g., (Coddington & Levinson, 1984). The proof of this existence and Theorem 4.4 are given in Appendix C.

4.4 Proof sketch of the convergence result for Regularized Mirror Descent

Let us return from continuous-time dynamics (4.5) to the discrete-time algorithm (4.4). The technical difficulty in the proof of Theorem 4.3 is the non-Lipschitz continuity of the value function $Q_h^{\lambda,\sigma}$ in (4.4), that is, the derivative of $Q_h^{\lambda,\sigma}(s,a,\pi,\mu)$ with respect to the policy π can blow up as π approaches the boundary of the space $(\Delta(\mathcal{A})^{\mathcal{S}})^H$ of probability simplices.

We can overcome this difficulty as shown in the following sketch of proof:

> *Proof sketch of Theorem 4.3.* In a similar way to Theorem 4.4, we can obtain the following inequality with a discretization error:
>
> $$D_{\mu^*}(\varpi^*,\pi^{t+1}) - D_{\mu^*}(\varpi^*,\pi^t) \leq -\lambda\eta D_{\mu^*}(\varpi^*,\pi^t) + \boxed{D_{\mu^*}(\pi^t,\pi^{t+1}).} \tag{4.6}$$
>
> discretization error

The remainder of the proof is almost entirely dedicated to showing that the above error term is sufficiently small and bounded compared to the other terms in the inequality. As a result, we obtain the following claim:

> **Claim 4.5.** *Suppose that the learning rate η is less than the upper bound η^* in (D.5). Then, we have*
>
> $$\boxed{D_{\mu^*}(\pi^t,\pi^{t+1})} \leq C\eta^2 D_{\mu^*}(\varpi^*,\pi^t),$$
>
> *where $C > 0$ is the constant defined in (D.4), which satisfies $C\eta^* \leq \lambda/2$.*

The key to proving Claim 4.5 is leveraging another claim that, over the sequence $(\pi^t)_t$, the value function $Q_h^{\lambda,\sigma}$ behaves well, almost as if it were a Lipschitz continuous function, see Lemma D.3 for details. Therefore, applying Claim 4.5 to (4.6) completes the proof. $\qquad\square$

The complete proof of Theorem 4.3 is given in Appendix D.

4.5 APPROXIMATED PROXIMAL POINT METHOD

Let us consider an approximation of Algorithm 1 using RMD of (4.4). We can simply replace the intractable computation in line 4 of Algorithm 1 with RMD. In the end, this means that after repeating (4.4) a sufficient number of times, we also update σ to the most recently obtained policy σ^{k+1} using RMD. The pseudo-code that summarizes this idea is presented in Algorithm 2.

5 NUMERICAL EXPERIMENT

We numerically demonstrate that the proposed algorithm (Algorithm 2), which is the approximated version of Algorithm 1, can achieve convergence to the mean-field equilibrium.

Algorithms. In this experiment, we implement Algorithm 2. For comparison, we also implement RMD (i.e., Algorithm 2 without the update of σ_k) in (4.4). For both algorithms, the learning rate is fixed at $\eta = 0.1$, and we vary the regularization parameter λ and update time T to run the experiments.

Evaluations. We evaluate the convergence of our proposed method using the Beach Bar Process introduced by Perrin et al. (2020), a standard benchmark for MFGs. In particular, the transition kernel P in this benchmark gives a random walk on a one-dimensional discretized torus $\mathcal{S} = \{0, \ldots, |\mathcal{S}| - 1\}$, and the reward is set to be $r_h(s, a, \mu) = -|a|/|\mathcal{S}| - |s - |\mathcal{S}|/2|/|\mathcal{S}| - \log \mu_h(s)$ with $a \in \mathcal{A} := \{-1, \pm 0, +1\}$. See Appendix F for further details. Since the mean-field equilibrium in this benchmark cannot be computed exactly, we follow Pérolat et al. (2022); Zhang et al. (2023) and employ the exploitability of a policy $\pi \in (\Delta(\mathcal{A})^{\mathcal{S}})^H$ defined by

$$\text{Exploit}(\pi) := \max_{\pi' \in (\Delta(\mathcal{A})^{\mathcal{S}})^H} \{J(m[\pi], \pi')\} - J(m[\pi], \pi) \geq 0,$$

as our convergence criterion. Note that from Definition 2.4, $\text{Exploit}(\pi) = 0$ if and only if $(m[\pi], \pi)$ is mean-field equilibrium.

Discussion. Figure 2 is a summary of the results of the experiment. The most noteworthy aspect is the convergence of the exploitability, as shown in Figure 2b. Our proposed method decreases the exploitability with each iteration when we update σ.

Figures 2a and 2c illustrate the qualitative validity of the approximation achieved by our proposed method. In this benchmark, the equilibrium is expected to lie at the vertices of the probability simplex. Therefore, RMD, which can shift the equilibrium to the interior of the probability simplex, seems unable to find the mean-field equilibrium accurately. On the other hand, the sequence $(\pi^t)_t$ of policies generated by our proposed method shows a behavior that converges to the vertices.

In summary, Algorithm 2 experimentally shows the last-iterate convergence to the mean-field equilibrium. This is evidenced by the decreasing exploitability and the qualitative behavior in our proposed method, which align with the theoretical guarantees.

6 COMPARISON OF THE RESULTS

Last-iterate convergence (LIC) results for MFG. Pérolat et al. (2022) showed that Mirror Descent achieves LIC only under *strictly* monotone conditions, i.e., if the equality in the Lemma E.2 is satisfied only if $\pi = \widetilde{\pi}$. In contrast, our work establishes LIC even in *non-strictly* monotone scenarios. While the distinction regarding strictness might seem subtle, it is profoundly significant. Indeed, non-strictly monotone MFGs encompass the fundamental examples of finite-horizon Markov Decision Processes. Moreover, in strictly monotone cases, mean-field equilibria become unique. Consequently, as Zeng et al. (2024) also noted, strictly monotone rewards fail to represent MFGs with diverse equilibria.

(a) The difference between RMD and the proposed method in the evolution of μ_h.

(b) Convergence of the exploitability

(c) Trajectory of $(\pi_h^t(s))_{t=1}^N$ with $h = 1$, $s = 1$.

Figure 2: Experimental results for Algorithm 2 for Beach Bar Process

Regularized MFG. Theorem 4.3, which supports the efficient execution of RMD, is novel in two respects: RMD achieves LIC, and the divergence to the equilibrium decays exponentially. Indeed, one of the few works that analyze the convergence rate of RMD states that the time-averaged policy $\frac{1}{T}\sum_{t=0}^T \pi^t$ up to time T converges to the equilibrium in $\mathcal{O}(1/\varepsilon^2)$ iterations (Zhang et al., 2023). Additionally, although it is a different approach from MD, it is known that applying fixed-point iteration to regularized MFG achieves an exponential convergence rate under the assumption that the regularization parameter λ is sufficiently large (Cui & Koeppl, 2021). In contrast, our work derives the convergence rate for cases where λ is sufficiently small.

Other type of learning method of MFG. Recently, in addition to Mirror Descent and Fictitious Play, a new type of learning method using the characterization of MFGs as optimization problems has been proposed (Guo et al., 2024; Hu & Zhang, 2024). In this work, the authors establish local convergence of the algorithms without the assumption of monotonicity. Specifically, it is proved that an optimization method can achieve LIC if the initial guess of the algorithm is sufficiently close to the Nash equilibrium. In contrast, our convergence results state "global" convergence under the assumption of monotonicity, complementing their results.

See Table 1 in the Appendix for a comparison of our results with the more comprehensive previous studies.

7 CONCLUSION

This paper proposes noble algorithms that can achieve last-iterate convergence under the monotonicity condition. The main idea behind the derivation of the main algorithm (Algorithm 2) is to approximate the proximal-point type algorithm (Algorithm 1) using RMD. Theorem 3.1 guarantees that the proximal-point-type algorithm achieves LIC, and Theorem 4.3 guarantees the exponential convergence of RMD. An important future task of this study is to prove the convergence rates of Algorithm 2. Specifically, we aim to make the convergence result of Theorem 3.1 quantitative. As

the experimental results suggest in Figure 2b, we conjecture that the algorithm converges with a rate of $\mathcal{O}(1/t^\alpha)$ for some $\alpha > 0$.

REFERENCES

Kenshi Abe, Kaito Ariu, Mitsuki Sakamoto, Kentaro Toyoshima, and Atsushi Iwasaki. Last-iterate convergence with full and noisy feedback in two-player zero-sum games. In *AISTATS*, volume 206 of *Proceedings of Machine Learning Research*, pp. 7999–8028. PMLR, 2023.

Kenshi Abe, Kaito Ariu, Mitsuki Sakamoto, and Atsushi Iwasaki. Adaptively perturbed mirror descent for learning in games. In *ICML*, 2024.

Alekh Agarwal, Nan Jiang, Sham M. Kakade, and Wen Sun. Reinforcement learning: Theory and algorithms, 2022. URL https://rltheorybook.github.io/rltheorybook_AJKS.pdf. online.

Berkay Anahtarci, Can Deha Kariksiz, and Naci Saldi. Q-learning in regularized mean-field games. *Dynamic Games and Applications*, 13(1):89–117, Mar 2023. ISSN 2153-0793. doi: 10.1007/s13235-022-00450-2. URL https://doi.org/10.1007/s13235-022-00450-2.

Andrea Angiuli, Jean-Pierre Fouque, and Mathieu Laurière. Unified reinforcement q-learning for mean field game and control problems. *Math. Control. Signals Syst.*, 34(2):217–271, 2022.

Andrea Angiuli, Jean-Pierre Fouque, Mathieu Laurière, and Mengrui Zhang. Analysis of multi-scale reinforcement q-learning algorithms for mean field control games, 2024. URL https://arxiv.org/abs/2405.17017v3.

Dimitri P. Bertsekas and Steven E. Shreve. *Stochastic optimal control*, volume 139 of *Mathematics in Science and Engineering*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1978. ISBN 0-12-093260-1. The discrete time case.

George W. Brown. Iterative solution of games by fictitious play. In T. C. Koopmans (ed.), *Activity Analysis of Production and Allocation*. Wiley, New York, 1951.

Peter E. Caines and Minyi Huang. Graphon mean field games and the GMFG equations: ϵ-nash equilibria. In *CDC*, pp. 286–292. IEEE, 2019.

Pierre Cardaliaguet and Saeed Hadikhanloo. Learning in mean field games: The fictitious play. *ESAIM: COCV*, 23(2):569–591, 2017. doi: 10.1051/cocv/2016004. URL https://doi.org/10.1051/cocv/2016004.

Y. Censor and S. A. Zenios. Proximal minimization algorithm withd-functions. *Journal of Optimization Theory and Applications*, 73(3):451–464, Jun 1992. ISSN 1573-2878. doi: 10.1007/BF00940051. URL https://doi.org/10.1007/BF00940051.

Ralph Chill, Eva Fasangova, and Univerzita Fakulta. Gradient systems. *13th International Internet Seminar*, 6 2010.

A. Coddington and N. Levinson. *Theory of Ordinary Differential Equations*. International series in pure and applied mathematics. R.E. Krieger, 1984. ISBN 9780898747553.

Kai Cui and Heinz Koeppl. Approximately solving mean field games via entropy-regularized deep reinforcement learning. In *AISTATS*, volume 130 of *Proceedings of Machine Learning Research*, pp. 1909–1917. PMLR, 2021.

Kai Cui and Heinz Koeppl. Learning graphon mean field games and approximate nash equilibria. In *ICLR*. OpenReview.net, 2022.

Kai Cui, Anam Tahir, Gizem Ekinci, Ahmed Elshamanhory, Yannick Eich, Mengguang Li, and Heinz Koeppl. A survey on large-population systems and scalable multi-agent reinforcement learning, 2022. URL https://arxiv.org/abs/2209.03859.

Nabarun Deb, Young-Heon Kim, Soumik Pal, and Geoffrey Schiebinger. Wasserstein mirror gradient flow as the limit of the sinkhorn algorithm, 2023. URL https://arxiv.org/abs/2307.16421.

Christian Fabian, Kai Cui, and Heinz Koeppl. Learning sparse graphon mean field games. In *AISTATS*, volume 206 of *Proceedings of Machine Learning Research*, pp. 4486–4514. PMLR, 2023.

Shuang Gao and Peter E. Caines. The control of arbitrary size networks of linear systems via graphon limits: An initial investigation. In *CDC*, pp. 1052–1057. IEEE, 2017.

Diogo A. Gomes, Joana Mohr, and Rafael Rigão Souza. Discrete time, finite state space mean field games. *Journal de Mathématiques Pures et Appliquées*, 93(3):308–328, 2010. ISSN 0021-7824. doi: https://doi.org/10.1016/j.matpur.2009.10.010. URL https://www.sciencedirect.com/science/article/pii/S002178240900138X.

Xin Guo, Anran Hu, and Junzi Zhang. Mf-omo: An optimization formulation of mean-field games. *SIAM Journal on Control and Optimization*, 62(1):243–270, 2024. doi: 10.1137/22M1524084. URL https://doi.org/10.1137/22M1524084.

Anran Hu and Junzi Zhang. Mf-oml: Online mean-field reinforcement learning with occupation measures for large population games, 2024. URL https://arxiv.org/abs/2405.00282.

Jiawei Huang, Batuhan Yardim, and Niao He. On the statistical efficiency of mean-field reinforcement learning with general function approximation. In *AISTATS*, volume 238 of *Proceedings of Machine Learning Research*, pp. 289–297. PMLR, 2024.

Minyi Huang, Roland P. Malhamé, and Peter E. Caines. Large population stochastic dynamic games: closed-loop McKean-Vlasov systems and the Nash certainty equivalence principle. *Communications in Information & Systems*, 6(3):221 – 252, 2006.

Daisuke Inoue, Yuji Ito, Takahito Kashiwabara, Norikazu Saito, and Hiroaki Yoshida. A fictitious-play finite-difference method for linearly solvable mean field games. *ESAIM: M2AN*, 57(4):1863–1892, 2023. doi: 10.1051/m2an/2023026. URL https://doi.org/10.1051/m2an/2023026.

Walid Krichene, Alexandre M. Bayen, and Peter L. Bartlett. Accelerated mirror descent in continuous and discrete time. In *NIPS*, pp. 2845–2853, 2015.

Jean-Michel Lasry and Pierre-Louis Lions. Mean field games. *Jpn. J. Math.*, 2(1):229–260, 2007. ISSN 0289-2316,1861-3624. doi: 10.1007/s11537-007-0657-8. URL https://doi.org/10.1007/s11537-007-0657-8.

Mathieu Lauriere, Sarah Perrin, Sertan Girgin, Paul Muller, Ayush Jain, Theophile Cabannes, Georgios Piliouras, Julien Perolat, Romuald Elie, Olivier Pietquin, and Matthieu Geist. Scalable deep reinforcement learning algorithms for mean field games. In Kamalika Chaudhuri, Stefanie Jegelka, Le Song, Csaba Szepesvari, Gang Niu, and Sivan Sabato (eds.), *Proceedings of the 39th International Conference on Machine Learning*, volume 162 of *Proceedings of Machine Learning Research*, pp. 12078–12095. PMLR, 17–23 Jul 2022. URL https://proceedings.mlr.press/v162/lauriere22a.html.

Pierre Lavigne and Laurent Pfeiffer. Generalized conditional gradient and learning in potential mean field games. *Applied Mathematics & Optimization*, 88(3):89, Oct 2023. ISSN 1432-0606. doi: 10.1007/s00245-023-10056-8. URL https://doi.org/10.1007/s00245-023-10056-8.

Stanisław Łojasiewicz. Sur les ensembles semi-analytiques. In *Actes du Congrès International des Mathématiciens (Nice, 1970), Tome 2,*, pp. 237–241. 1971.

Weichao Mao, Haoran Qiu, Chen Wang, Hubertus Franke, Zbigniew Kalbarczyk, Ravishankar K. Iyer, and Tamer Basar. A mean-field game approach to cloud resource management with function approximation. In *NeurIPS*, 2022.

Panayotis Mertikopoulos, Christos Papadimitriou, and Georgios Piliouras. Cycles in adversarial regularized learning. In *SODA*, pp. 2703–2717, 2018. doi: 10.1137/1.9781611975031.172. URL https://epubs.siam.org/doi/abs/10.1137/1.9781611975031.172.

Julien Pérolat, Rémi Munos, Jean-Baptiste Lespiau, Shayegan Omidshafiei, Mark Rowland, Pedro A. Ortega, Neil Burch, Thomas W. Anthony, David Balduzzi, Bart De Vylder, Georgios Piliouras, Marc Lanctot, and Karl Tuyls. From Poincaré recurrence to convergence in imperfect information games: Finding equilibrium via regularization. In *ICML*, volume 139 of *Proceedings of Machine Learning Research*, pp. 8525–8535. PMLR, 2021.

Julien Pérolat, Sarah Perrin, Romuald Elie, Mathieu Laurière, Georgios Piliouras, Matthieu Geist, Karl Tuyls, and Olivier Pietquin. Scaling mean field games by online mirror descent. In *AAMAS*, pp. 1028–1037. International Foundation for Autonomous Agents and Multiagent Systems (IFAAMAS), 2022.

Sarah Perrin, Julien Pérolat, Mathieu Laurière, Matthieu Geist, Romuald Elie, and Olivier Pietquin. Fictitious play for mean field games: Continuous time analysis and applications. In *NeurIPS*, 2020.

Sarah Perrin, Mathieu Laurière, Julien Pérolat, Romuald Élie, Matthieu Geist, and Olivier Pietquin. Generalization in mean field games by learning master policies. In *AAAI*, pp. 9413–9421. AAAI Press, 2022.

Georgios Piliouras, Ryann Sim, and Stratis Skoulakis. Beyond time-average convergence: Near-optimal uncoupled online learning via clairvoyant multiplicative weights update. In *NeurIPS*, 2022.

Martin L. Puterman. *Markov decision processes: discrete stochastic dynamic programming*. Wiley Series in Probability and Mathematical Statistics: Applied Probability and Statistics. John Wiley & Sons, Inc., New York, 1994. ISBN 0-471-61977-9. A Wiley-Interscience Publication.

Naci Saldi, Tamer Başar, and Maxim Raginsky. Markov–nash equilibria in mean-field games with discounted cost. *SIAM Journal on Control and Optimization*, 56(6):4256–4287, 2018. doi: 10.1137/17M1112583. URL https://doi.org/10.1137/17M1112583.

Peter D. Taylor and Leo B. Jonker. Evolutionary stable strategies and game dynamics. *Mathematical Biosciences*, 40(1):145–156, 1978. ISSN 0025-5564. doi: https://doi.org/10.1016/0025-5564(78)90077-9. URL https://www.sciencedirect.com/science/article/pii/0025556478900779.

Belinda Tzen, Anant Raj, Maxim Raginsky, and Francis R. Bach. Variational principles for mirror descent and mirror langevin dynamics. *IEEE Control. Syst. Lett.*, 7:1542–1547, 2023.

Qiaomin Xie, Zhuoran Yang, Zhaoran Wang, and Andreea Minca. Learning while playing in mean-field games: Convergence and optimality. In *ICML*, volume 139 of *Proceedings of Machine Learning Research*, pp. 11436–11447. PMLR, 2021.

Yujia Xie, Xiangfeng Wang, Ruijia Wang, and Hongyuan Zha. A fast proximal point method for computing exact Wasserstein distance. In *UAI*, volume 115 of *Proceedings of Machine Learning Research*, pp. 433–453. AUAI Press, 2019.

Yaodong Yang and Jun Wang. An overview of multi-agent reinforcement learning from game theoretical perspective, 2021. URL https://arxiv.org/abs/2011.00583.

Yaodong Yang, Rui Luo, Minne Li, Ming Zhou, Weinan Zhang, and Jun Wang. Mean field multi-agent reinforcement learning. In *ICML*, volume 80 of *Proceedings of Machine Learning Research*, pp. 5567–5576. PMLR, 2018.

Batuhan Yardim and Niao He. Exploiting approximate symmetry for efficient multi-agent reinforcement learning, 2024. URL https://arxiv.org/abs/2408.15173.

Batuhan Yardim, Semih Cayci, Matthieu Geist, and Niao He. Policy mirror ascent for efficient and independent learning in mean field games. In *ICML*, volume 202 of *Proceedings of Machine Learning Research*, pp. 39722–39754. PMLR, 2023.

Batuhan Yardim, Artur Goldman, and Niao He. When is mean-field reinforcement learning tractable and relevant? In *AAMAS*, pp. 2038–2046. International Foundation for Autonomous Agents and Multiagent Systems / ACM, 2024.

Muhammad Aneeq uz Zeman, Alec Koppel, Sujay Bhatt, and Tamer Basar. Oracle-free reinforcement learning in mean-field games along a single sample path. In *AISTATS*, volume 206 of *Proceedings of Machine Learning Research*, pp. 10178–10206. PMLR, 2023.

Sihan Zeng, Sujay Bhatt, Alec Koppel, and Sumitra Ganesh. Learning in herding mean field games: Single-loop algorithm with finite-time convergence analysis, 2024. URL https://arxiv.org/abs/2408.04780v4.

Fengzhuo Zhang, Vincent Y. F. Tan, Zhaoran Wang, and Zhuoran Yang. Learning regularized monotone graphon mean-field games. In *NeurIPS*, 2023.

Kaiqing Zhang, Zhuoran Yang, and Tamer Başar. *Multi-Agent Reinforcement Learning: A Selective Overview of Theories and Algorithms*, pp. 321–384. Springer International Publishing, Cham, 2021. ISBN 978-3-030-60990-0. doi: 10.1007/978-3-030-60990-0_12. URL https://doi.org/10.1007/978-3-030-60990-0_12.

A RELATED WORKS

Several previous studies have derived convergence results for MFG for various algorithms. Note that the meaning of convergence is different in the previous studies. Table 1 shows which type of convergence is obtained for which type of algorithm in the previous studies and our results.

Table 1: Related work of convergence in MFGs

	Learning algorithm	**Summary of convergence results**
Xie et al. (2021)	Fictious play	time-averaging convergence
Zhang et al. (2023)	RMD	time-averaging convergence (to regularized equilibrium) under monotonicity
Mao et al. (2022)	Actor-critic	time-averaging convergence (to regularized equilibrium)
Zeman et al. (2023)	Q-learning	time-averaging convergence
Yardim et al. (2023)	Mirror Descent	LIC under contraction
Zeng et al. (2024)	Actor-critic	best-iterate convergence under Herding
Huang et al. (2024)	Maximum Likelihood Estimation	N/A
Angiuli et al. (2022)	(two-time scale) Q-Learning	N/A
Angiuli et al. (2024)	(three-time scale) Q-learning	LIC under contraction
Pérolat et al. (2021)	RMD	LIC under strict monotonicity in continuous time
Our work (Theorem 3.1)	PP	LIC (to regularized equilibrium) under monotonicity
Our work (Theorem 4.4)	RMD	LIC under monotonicity

B PROOF OF THEOREM 3.1

Proof of Lemma 3.2. Let $(\mu^\star, \pi^\star)$ be a mean-field equilibrium defined in Definition 2.4. By the update rule (3.1) and Lemma E.1, we have

$$\left\langle Q_h^{\lambda,\sigma^k}(s, \bullet, \sigma^{k+1}, \mu^{k+1}) - \lambda \log \frac{\sigma_h^{k+1}(s)}{\sigma_h^k(s)}, (\pi_h^\star - \sigma_h^{k+1})(s) \right\rangle \le 0,$$

for each $h \in [H]$, $s \in \mathcal{S}$ and $k \in \mathbb{N}$, i.e.,

$$D_{\mathrm{KL}}(\pi_h^\star(s), \sigma_h^{k+1}(s)) - D_{\mathrm{KL}}(\pi_h^\star(s), \sigma_h^k(s)) - D_{\mathrm{KL}}(\sigma_h^{k+1}(s), \sigma_h^k(s))$$

$$\le \frac{1}{\lambda} \left\langle Q_h^{\lambda,\sigma^k}(s, \bullet, \sigma^{k+1}, \mu^{k+1}), (\sigma_h^{k+1} - \pi_h^\star)(s) \right\rangle. \tag{B.1}$$

Taking the expectation with respect to $s \sim \mu_h^\star$ and summing (B.1) over $h \in [H]$ yields

$$D_{\mu^\star}(\pi^\star, \sigma^{k+1}) - D_{\mu^\star}(\pi^\star, \sigma^k) + D_{\mu^\star}(\sigma^{k+1}, \sigma^k)$$

$$\le \frac{1}{\lambda} \sum_{h=1}^H \mathbb{E}_{s \sim \mu_h^\star} \left[\left\langle Q_h^{\lambda,\sigma^k}(s, \bullet, \sigma^{k+1}, \mu^{k+1}), (\sigma_h^{k+1} - \pi_h^\star)(s) \right\rangle \right].$$

By virtue of Lemmas E.2 and E.4, we further have

$$\sum_{h=1}^H \mathbb{E}_{s \sim \mu_h^\star} \left[\left\langle Q_h^{\lambda,\sigma^k}(s, \bullet, \sigma^{k+1}, \mu^{k+1}), (\sigma_h^{k+1} - \pi_h^\star)(s) \right\rangle \right]$$

$$\le J^{\lambda,\sigma^k}(\mu^{k+1}, \sigma^{k+1}) - J^{\lambda,\sigma^k}(\mu^{k+1}, \pi^\star) - \lambda D_{\mu^\star}(\pi^\star, \sigma^k) + \lambda D_{\mu^\star}(\sigma^{k+1}, \sigma^k)$$

$$\le J^{\lambda,\sigma^k}(\mu^\star, \sigma^{k+1}) - J^{\lambda,\sigma^k}(\mu^\star, \pi^\star) - \lambda D_{\mu^\star}(\pi^\star, \sigma^k) + \lambda D_{\mu^\star}(\sigma^{k+1}, \sigma^k)$$

$$\le J(\mu^\star, \sigma^{k+1}) - J(\mu^\star, \pi^\star) - \lambda D_{\mu^{k+1}}(\sigma^{k+1}, \sigma^k) + \lambda D_{\mu^\star}(\sigma^{k+1}, \sigma^k),$$

where we use the identity $J^{\lambda,\sigma^k}(\mu^\star, \pi) = J(\mu^\star, \pi) - \lambda D_{m[\pi]}(\pi, \sigma^k)$ for $\pi \in (\Delta(\mathcal{A})^\mathcal{S})^H$, and Definition 2.4. ∎

Proof of Lemma 3.3. Note that the function $J(\mu^\star, \bullet): (\Delta(\mathcal{A})^\mathcal{S})^H \ni \pi \mapsto J(\mu^\star, \pi) \in \mathbb{R}$ is real-analytic. Therefore, we can apply (Łojasiewicz, 1971, §18, Théorème 2.). ∎

C PROOF OF THEOREM 4.4

Proof of Theorem 4.4. Let $h^\star: \mathbb{R}^{|\mathcal{A}|} \to \mathbb{R}$ be the convex conjugate of h, i.e., $h^\star(y) = \sum_{a \in \mathcal{A}} \exp(y(a))$ for $y \in \mathbb{R}^{|\mathcal{A}|}$. From direct computations, we have

$$\frac{\mathrm{d}}{\mathrm{d}t} D_{\mu^*}(\varpi^*, \pi^t)$$

$$= \sum_{h=1}^H \mathbb{E}_{s \sim \mu_h^*} \left[\frac{\mathrm{d}}{\mathrm{d}t} D_{\mathrm{KL}}(\varpi_h^*(s), \pi^t(s)) \right]$$

$$= \sum_{h=1}^H \mathbb{E}_{s \sim \mu_h^*} \left[\left\langle 1 - \frac{\varpi_h^*(s)}{\pi_h^t(s)}, \frac{\mathrm{d}}{\mathrm{d}t} \pi_h^t(s) \right\rangle \right]$$

$$= \sum_{h=1}^H \mathbb{E}_{s \sim \mu_h^*} \left[\left\langle 1 - \frac{\varpi_h^*(s)}{\pi_h^t(s)}, \pi_h^t(a \mid s) \left(Q_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t) - \lambda \log \frac{\pi_h^t(a \mid s)}{\sigma_h(a \mid s)} \right) \right\rangle \right]$$

$$= \sum_{h=1}^H \mathbb{E}_{s \sim \mu_h^*} \left[\left\langle (\pi_h^t - \varpi_h^*)(s), Q_h^{\lambda,\sigma}(s, \bullet, \pi^t, \mu^t) - \lambda \log \frac{\pi_h^t(a \mid s)}{\sigma_h(a \mid s)} \right\rangle \right]$$

$$= \sum_{h=1}^H \mathbb{E}_{s \sim \mu_h^*} \left[\left\langle (\pi_h^t - \varpi_h^*)(s), Q_h^{\lambda,\sigma}(s, \bullet, \pi^t, \mu^t) \right\rangle \right] - \lambda \sum_{h=1}^H \mathbb{E}_{s \sim \mu_h^*} \left[\left\langle (\pi_h^t - \varpi_h^*)(s), \log \frac{\pi_h^t(s)}{\sigma_h(s)} \right\rangle \right].$$

We apply Lemma E.4 for the first term and get

$$
\begin{aligned}
&\sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\left\langle (\pi_h^t - \varpi_h^*)(s), Q_h^{\lambda,\sigma}(s, \bullet, \pi^t, \mu^t) \right\rangle \right] \\
&= J^{\lambda,\sigma}(\mu^t, \pi^t) - J^{\lambda,\sigma}(\mu^t, \varpi^*) - \lambda D_{\mu^*}(\varpi^*, \sigma) + \lambda D_{\mu^*}(\pi^t, \sigma).
\end{aligned} \tag{C.1}
$$

Similarly, we apply Lemma E.5 for the second term and get

$$
\sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\left\langle (\pi_h^t - \varpi_h^*)(s), \log \frac{\pi_h^t(s)}{\sigma_h(s)} \right\rangle \right] = D_{\mu^*}(\pi^t, \sigma) - D_{\mu^*}(\varpi^*, \sigma) + D_{\mu^*}(\varpi^*, \pi^t). \tag{C.2}
$$

Combining (C.1) and (C.2) yields

$$
\frac{\mathrm{d}}{\mathrm{d}t} D_{\mu^*}(\varpi^*, \pi^t) = J^{\lambda,\sigma}(\mu^t, \pi^t) - J^{\lambda,\sigma}(\mu^t, \varpi^*) - \lambda D_{\mu^*}(\varpi^*, \pi^t).
$$

By virtue of the definition of mean-field equilibrium and Lemma E.2, we find

$$
J^{\lambda,\sigma}(\mu^t, \pi^t) - J^{\lambda,\sigma}(\mu^t, \varpi^*) \leq J^{\lambda,\sigma}(\mu^*, \pi^t) - J^{\lambda,\sigma}(\mu^*, \varpi^*) \leq 0.
$$

Therefore, we obtain

$$
\frac{\mathrm{d}}{\mathrm{d}t} D_{\mu^*}(\varpi^*, \pi^t) \leq -\lambda D_{\mu^*}(\varpi^*, \pi^t).
$$

∎

Proposition C.1. *Assume the same assumption as in Theorem 3.1. Then, there exists a unique maximizer of $J^{\lambda,\sigma^k}(\mu^k, \bullet) \colon (\Delta(\mathcal{A})^{\mathcal{S}})^H \to \mathbb{R}$ for each $k \in \mathbb{N}$.*

The uniqueness of Proposition C.1 is a new result. The proof uses a continuous-time dynamics shown in Theorem 4.4, see Appendix C. In the following proof, we employ the same proof strategy as in (Chill et al., 2010, Theorem 2.10). Before the proof, set $v_{s,h}^{\lambda,\sigma}(\pi) \coloneqq \pi_h(a \mid s) \left(Q_h^{\lambda,\sigma}(s, a, \pi, m[\pi]) - \lambda \log \frac{\pi_h(a \mid s)}{\sigma_h(a \mid s)} \right)$ for $\pi \in (\Delta(\mathcal{A})^{\mathcal{S}})^H$.

Proof of Proposition C.1. The existence is shown by a slightly modified version of (Zhang et al., 2023, Theorem 2). It remains to prove the uniqueness. Fix the regularized equilibrium $\varpi^* \in (\Delta(\mathcal{A})^{\mathcal{S}})^H$.

First of all, we prove the global existence of (4.5). By the local Lipschitz continuity of the right-hand side of the dynamics (4.5) and Picard–Lindelöf theorem, there exists a unique maximal solution π of (4.5) with the initial condition $\pi|_{t=0} = \pi^0$. Namely, there exist $T \in (0, +\infty]$ and $\pi \colon [0, T) \to \mathbb{R}^{|\mathcal{A}|}$ such that π is differentiable on $(0, T)$ and it holds that (4.5) for all $t \in (0, T)$. Thus, Theorem 4.4 ensures that

$$
D_{\mu^*}(\varpi^*, \pi^t) + \lambda \int_0^t D_{\mu^*}(\varpi^*, \pi^\tau) \, \mathrm{d}\tau \leq D_{\mu^*}(\varpi^*, \pi^0) =: c < +\infty,
$$

for every $t \in [0, T)$. As a result, the trajectory $\{\pi^t \in (\Delta(\mathcal{A})^{\mathcal{S}})^H \mid t \in [0, T)\}$ is included in $K_c \coloneqq \{\pi \in (\Delta(\mathcal{A})^{\mathcal{S}})^H \mid D_{\mu^*}(\varpi^*, \pi) \leq c\}$. Note that K_c is compact from Pinsker inequality.

Since the right-hand side of (4.5) is continuous on K_c, we obtain $\sup_{t \in [0,+\infty)} \left\| v_{s,h}^{\lambda,\sigma}(\pi^t) \right\| < +\infty$. Thus, the equation (4.5) implies $\left\| \frac{\mathrm{d}\pi^t}{\mathrm{d}t} \right\|$ is uniformly bounded on $[0, T)$. Hence, π extends to a continuous function on $[0, T]$.

To obtain a contradiction, we assume $T < +\infty$. Then, there exists the solution π' of (4.5) on a larger interval than π with a new initial condition $\pi'|_{t'=T} = \pi^T$, which contradicts the maximality of the solution π.

Therefore, the limit $\lim_{t \to \infty} \pi^t$ exists and is equal to ϖ^*. Here, ϖ^* is arbitrary, so the regularized equilibrium is unique. ∎

D Proof of Theorem 4.3

Lemma D.1. *It holds that*

$$\left\langle \eta\left(Q_h^{\lambda,\sigma}(s,\bullet,\pi^t,\mu^t) - \lambda\log\frac{\pi_h^{t+1}(s)}{\sigma_h(s)}\right) - (1-\lambda\eta)\log\frac{\pi_h^{t+1}(s)}{\pi_h^t(s)}, \delta\right\rangle = 0,$$

for all $\delta \in \mathbb{R}^{|\mathcal{A}|}$ such that $\sum_a \delta(a) = 0$.

We introduce the following lemma:

Lemma D.2. *Let $(\pi^t)_t$ be the sequence defined by (4.4) and ϖ^* be the policy satisfies Definition 4.2. Assume that there exist vectors w_h^σ and $w_h^0(s) \in \mathbb{R}^{|\mathcal{A}|}$ satisfying*

$$\lambda H \log\sigma_{\min} \leq w_h^\sigma(a \mid s) \leq -\lambda H \log\sigma_{\min}, \qquad \sigma_h(a \mid s) \propto \exp\left(\frac{w_h^\sigma(a \mid s)}{\lambda}\right),$$

$$2\lambda H \log\sigma_{\min} \leq w_h^0(a \mid s) \leq H, \qquad \pi_h^0(a \mid s) \propto \exp\left(\frac{w_h^0(a \mid s)}{\lambda}\right).$$

for all $a \in \mathcal{A}.\pi^0 \in (\Delta(\mathcal{A})^{\mathcal{S}})^H$, $h \in [H]$ and $s \in \mathcal{S}$. Then, for any $h \in [H], s \in \mathcal{S}$, and $t \geq 0$, it holds that

$$\max\left\{\left\|\log\pi_h^t(s)\right\|_\infty, \left\|\log\pi_h^*(s)\right\|_\infty\right\} \leq \frac{H(1-\lambda\log\sigma_{\min})}{\lambda} + \log|\mathcal{A}|.$$

Proof. We first show that π_h^t can be written as

$$\pi_h^t(a \mid s) \propto \exp\left(\frac{w_h^t(a \mid s)}{\lambda}\right), \tag{D.1}$$

for a vector $w_h^t(s) \in \mathbb{R}^{|\mathcal{A}|}$ satisfying $2\lambda H \log\sigma_{\min} \leq w_h^t(a \mid s) \leq H$. We prove it by induction on t. Suppose that there exist $t \in \mathbb{N}$ and w_h^t satisfying (D.1). By the update rule (4.4), we have

$$\pi_h^{t+1}(a \mid s) \propto (\sigma_h(a \mid s))^{\lambda\eta}\left(\pi_h^t(a \mid s)\right)^{1-\lambda\eta}\exp\left(\eta Q_h^{\lambda,\sigma}(s,a,\pi^t,\mu^t)\right)$$

$$\propto \exp\left(\frac{\lambda\eta w_h^\sigma(a \mid s) + (1-\eta\lambda)w_h^t(a \mid s) + \lambda\eta Q_h^{\lambda,\sigma}(s,a,\pi^t,\mu^t)}{\lambda}\right).$$

Set $w_h^{t+1}(a \mid s) := \lambda\eta w_h^\sigma(a \mid s) + (1-\eta\lambda)w_h^t(a|s) + \lambda\eta Q_h^{\lambda,\sigma}(s,a,\pi^t,\mu^t)$, we get $\pi_h^{t+1}(a|s) \propto e^{\frac{w_h^{t+1}(a \mid s)}{\lambda}}$. From Lemma E.3 and the hypothesis of the induction, we get $2\lambda H \log\sigma_{\min} \leq w_h^{t+1}(a \mid s) \leq H$.

Then we have for any $a_1, a_2 \in \mathcal{A}$:

$$\frac{\pi_h^t(a_1 \mid s)}{\pi_h^t(a_2 \mid s)} = \exp\left(\frac{w_h^t(a_1 \mid s) - w_h^t(a_2 \mid s)}{\lambda}\right) \leq \exp\left(\frac{H(1-\lambda\log\sigma_{\min})}{\lambda}\right).$$

It follows that:

$$\min_{a \in \mathcal{A}}\pi^t(a|s) \geq \exp\left(\frac{-H(1-\lambda\log\sigma_{\min})}{\lambda}\right)\max_{a' \in \mathcal{A}}\pi_h^t(a \mid s) \geq |\mathcal{A}|^{-1}\exp\left(\frac{-H(1-\lambda\log\sigma_{\min})}{\lambda}\right).$$

Therefore, we have:

$$\left\|\log\pi_h^t(s)\right\|_\infty \leq \frac{H(1-\lambda\log\sigma_{\min})}{\lambda} + \log|\mathcal{A}|.$$

From Lemmas E.1 and E.3, we have for π_h^* and $a_1, a_2 \in \mathcal{A}$:

$$\frac{\pi_h^*(a_1 \mid s)}{\pi_h^*(a_2 \mid s)} = \exp\left(\frac{Q_h^{\lambda,\sigma}(s,a_1,\pi^t,\mu^t) + w_h^\sigma(a_1 \mid s) - Q_h^{\lambda,\sigma}(s,a_2,\pi^t,\mu^t) - w_h^\sigma(a_2 \mid s)}{\lambda}\right)$$

$$\leq \exp\left(\frac{H(1-\lambda\log\sigma_{\min})}{\lambda}\right),$$

and, we get $\left\|\log\pi_h^*(s)\right\|_\infty \leq \frac{H(1-\lambda\log\sigma_{\min})}{\lambda} + \log|\mathcal{A}|$. ∎

Lemma D.3. *Let* $G_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t) \coloneqq Q_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t) - \lambda \log \dfrac{\pi_h^t(a \mid s)}{\sigma_h(a \mid s)}.$

$$\left| G_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t) - G_h^{\lambda,\sigma}(s, a', \pi^t, \mu^t) \right|$$

$$\leq 2L \sum_{l=h}^{H} \left\| \mu_l^t - \mu_l^* \right\|_1 + C^{\lambda,\sigma,H,|\mathcal{A}|} \left(E_h(a, \pi^t, \varpi^*) + E_h(a', \pi^t, \varpi^*) \right),$$

for $a, a' \in \mathcal{A}$. *Here,*

$$C^{\lambda,\sigma,H,|\mathcal{A}|} \coloneqq 2\lambda|\mathcal{A}|\mathrm{e}^{\frac{H(1 - \lambda \log \sigma_{\min})}{\lambda}} + 2(1 + H) - \lambda(1 + 2H) \log \sigma_{\min} + 2\lambda \log |\mathcal{A}|,$$

and

$$E_h(a, \pi^t, \varpi^*) \coloneqq \mathbb{E}\left[\sum_{l=h}^{H} \left\| \pi_l^*(s_l) - \pi_l^t(s_l) \right\|_1 \,\middle|\, \begin{array}{c} s_h = s, a_h = a, \\ s_{l+1} \sim P_l(s_l, a_l), \\ a_l \sim \varpi_l^*(s_l) \\ \text{for each } l \in \{h, \dots, H\} \end{array} \right].$$

Proof of [Lemma D.3](#). We first compute the absolute value as follows:

$$\left| G_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t) - G_h^{\lambda,\sigma}(s, a', \pi^t, \mu^t) \right|$$

$$= \left| \left(Q_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t) - \lambda \log \frac{\pi_h^t(a \mid s)}{\sigma_h(a \mid s)} \right) - \left(Q_h^{\lambda,\sigma}(s, a', \pi^t, \mu^t) - \lambda \log \frac{\pi_h^t(a' \mid s)}{\sigma_h(a' \mid s)} \right) \right|$$

$$\leq \left| \left(Q_h^{\lambda,\sigma}(s, a, \varpi^*, \mu^*) - \lambda \log \frac{\pi_h^t(a \mid s)}{\sigma_h(a \mid s)} \right) - \left(Q_h^{\lambda,\sigma}(s, a', \varpi^*, \mu^*) - \lambda \log \frac{\pi_h^t(a' \mid s)}{\sigma_h(a' \mid s)} \right) \right|$$

$$+ \left| \left(Q_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t) - Q_h^{\lambda,\sigma}(s, a, \varpi^*, \mu^*) \right) - \left(Q_h^{\lambda,\sigma}(s, a', \pi^t, \mu^t) - Q_h^{\lambda,\sigma}(s, a', \varpi^*, \mu^*) \right) \right|.$$

$$\text{(D.2)}$$

By [Lemmas D.2](#) and [E.1](#), the first term of right-hand side in [(D.3)](#) can be computed as

$$\left| \left(Q_h^{\lambda,\sigma}(s, a, \varpi^*, \mu^*) - \lambda \log \frac{\pi_h^t(a \mid s)}{\sigma_h(a \mid s)} \right) - \left(Q_h^{\lambda,\sigma}(s, a', \varpi^*, \mu^*) - \lambda \log \frac{\pi_h^t(a' \mid s)}{\sigma_h(a' \mid s)} \right) \right|$$

$$= \left| \left(\lambda \log \frac{\varpi_h^*(a \mid s)}{\sigma_h(a \mid s)} - \lambda \log \frac{\pi_h^t(a \mid s)}{\sigma_h(a \mid s)} \right) - \left(\lambda \log \frac{\varpi_h^*(a' \mid s)}{\sigma_h(a' \mid s)} - \lambda \log \frac{\pi_h^t(a' \mid s)}{\sigma_h(a' \mid s)} \right) \right|$$

$$\leq \lambda \left(\left| \log \frac{\varpi_h^*(a \mid s)}{\pi_h^t(a \mid s)} \right| + \left| \log \frac{\varpi_h^*(a' \mid s)}{\pi_h^t(a' \mid s)} \right| \right)$$

$$\leq \lambda \left(\frac{1}{\varpi_{\min}^*} + \frac{1}{\min_{a \in \mathcal{A}} \pi_h^t(a \mid s)} \right) \left(\left| \varpi_h^*(a \mid s) - \pi_h^t(a \mid s) \right| + \left| \varpi_h^*(a' \mid s) - \pi_h^t(a' \mid s) \right| \right)$$

$$\leq 2\lambda|\mathcal{A}| \exp\left(\frac{H(1 - \lambda \log \sigma_{\min})}{\lambda} \right) \left(\left| \varpi_h^*(a \mid s) - \pi_h^t(a \mid s) \right| + \left| \varpi_h^*(a' \mid s) - \pi_h^t(a' \mid s) \right| \right).$$

$$\text{(D.3)}$$

By [Proposition E.8](#) and [Lemma E.6](#), the second term is bounded as

$$\left| \left(Q_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t) - Q_h^{\lambda,\sigma}(s, a, \varpi^*, \mu^*) \right) - \left(Q_h^{\lambda,\sigma}(s, a', \pi^t, \mu^t) - Q_h^{\lambda,\sigma}(s, a', \varpi^*, \mu^*) \right) \right|$$

$$\leq 2L \sum_{l=h}^{H} \left\| \mu_l^t - \mu_l^* \right\|_1$$

$$+ C^{\lambda,\sigma}(\pi^t, \varpi^*) \mathbb{E}\left[\sum_{l=h+1}^{H} \left\| \pi_l^*(s_l) - \pi_l^t(s_l) \right\|_1 \,\middle|\, \begin{array}{c} s_{h+1} \sim P_h(\bullet \mid s, a), \\ s_{l+1} \sim P_l(s_l, a_l), \\ a_l \sim \varpi_l^*(s_l) \\ \text{for each } l \in \{h+1, \dots, H\} \end{array} \right]$$

$$+ C^{\lambda,\sigma}(\pi^t, \varpi^*) \mathbb{E}\left[\sum_{l=h+1}^{H} \left\| \pi_l^*(s_l) - \pi_l^t(s_l) \right\|_1 \,\middle|\, \begin{array}{c} s_{h+1} \sim P_h(\bullet \mid s, a'), \\ s_{l+1} \sim P_l(s_l, a_l), \\ a_l \sim \varpi_l^*(s_l) \\ \text{for each } l \in \{h+1, \dots, H\} \end{array} \right].$$

Furthermore, $C^{\lambda,\sigma}(\pi^t, \varpi^*)$ can be bounded as

$$
\begin{aligned}
C^{\lambda,\sigma}(\pi^t, \varpi^*) \leq & 2 - \lambda \log \sigma_{\min} + 2\lambda \left(\frac{H(1 - \lambda \log \sigma_{\min})}{\lambda} + \log |\mathcal{A}| \right) \\
= & 2(1 + H) - \lambda(1 + 2H) \log \sigma_{\min} + 2\lambda \log |\mathcal{A}|.
\end{aligned}
$$

∎

Proof of Theorem 4.3. Set

$$
C := 4H^2 \left(L^2 H^2 + \frac{\left(C^{\lambda,\sigma,H,|\mathcal{A}|} \right)^2}{|\mathcal{A}| \exp \left(\frac{H(1 - \lambda \log \sigma_{\min})}{\lambda} \right)} \right) \tag{D.4}
$$

$$
= 4H^2 \left(L^2 H^2 + \frac{\left(2\lambda |\mathcal{A}| e^{\frac{H(1 - \lambda \log \sigma_{\min})}{\lambda}} + 2(1 + H) - \lambda(1 + 2H) \log \sigma_{\min} + 2\lambda \log |\mathcal{A}| \right)^2}{|\mathcal{A}| e^{\frac{H(1 - \lambda \log \sigma_{\min})}{\lambda}}} \right)
$$

$$
\eta^* = \min \left\{ \frac{1}{2H(L + C^{\lambda,\sigma,H,|\mathcal{A}|})}, \frac{\lambda}{2C} \right\}, \tag{D.5}
$$

where $C^{\lambda,\sigma,H,|\mathcal{A}|}$ is the constant defined in Lemma D.3. We prove the inequality by induction on t.

(I) Base step $t = 0$: It is obvious.

(II) Inductive step: Suppose that there exists $t \in \mathbb{N}$ such that $\pi^t \in \Omega$. Lemma D.1 yields that

$$
\begin{aligned}
& D_{\mu^*}(\varpi^*, \pi^{t+1}) - D_{\mu^*}(\varpi^*, \pi^t) - D_{\mu^*}(\pi^t, \pi^{t+1}) \\
= & \sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\left\langle \log \frac{\pi_h^t(s)}{\pi_h^{t+1}(s)}, (\varpi_h^* - \pi_h^t)(s) \right\rangle \right] \\
= & -\sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\left\langle \frac{\eta}{1 - \lambda\eta} \left(Q_h^{\lambda,\sigma}(s, \bullet, \pi^t, \mu^t) - \lambda \log \frac{\pi_h^{t+1}(s)}{\sigma_h(s)} \right), (\varpi_h^* - \pi_h^t)(s) \right\rangle \right] \\
= & -\frac{\eta}{1 - \lambda\eta} \underbrace{\sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\left\langle Q_h^{\lambda,\sigma}(s, \bullet, \pi^t, \mu^t), (\pi_h^* - \pi_h^t)(s) \right\rangle \right]}_{=: \mathrm{I}} \\
& + \frac{\lambda\eta}{1 - \lambda\eta} \sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\left\langle \log \frac{\pi_h^{t+1}(s)}{\sigma_h(s)}, (\pi_h^* - \pi_h^{t+1})(s) \right\rangle \right] \\
\leq & -\frac{\eta}{1 - \lambda\eta} \left(\lambda D_{\mu^*}(\varpi^*, \sigma) - \lambda D_{\mu^*}(\pi^{t+1}, \sigma) \right) \\
& + \frac{\lambda\eta}{1 - \lambda\eta} \left(D_{\mu^*}(\varpi^*, \sigma) - D_{\mu^*}(\varpi^*, \pi^{t+1}) - D_{\mu^*}(\pi^{t+1}, \sigma) \right) \\
\leq & -\frac{\lambda\eta}{1 - \lambda\eta} D_{\mu^*}(\varpi^*, \pi^{t+1}),
\end{aligned} \tag{D.6}
$$

where I is bounded from below as follows: By Lemma E.4, we get

$$
\mathrm{I} = J^{\lambda,\sigma}(\mu^{t+1}, \varpi^*) - J^{\lambda,\sigma}(\mu^{t+1}, \pi^{t+1}) + \lambda D_{\mu^*}(\varpi^*, \sigma) - \lambda D_{\mu^*}(\pi^{t+1}, \sigma). \tag{D.7}
$$

By virtue of the definition of mean-field equilibrium and Lemma E.2, we find

$$
J^{\lambda,\sigma}(\mu^{t+1}, \varpi^*) - J^{\lambda,\sigma}(\mu^{t+1}, \pi^{t+1}) \geq J^{\lambda,\sigma}(\mu^*, \varpi^*) - J^{\lambda,\sigma}(\mu^*, \pi^{t+1}) \geq 0.
$$

Then, we obtain

$$
\mathrm{I} \geq \lambda D_{\mu^*}(\varpi^*, \sigma) - \lambda D_{\mu^*}(\pi^{t+1}, \sigma).
$$

For the last term $D_{\mu^*}(\pi^t, \pi^{t+1})$ of the leftmost hand of (D.6), we can employ a similar argument to (Abe et al., 2023, Lemma 5.4), that is, we can estimate $D_{\mu^*}(\pi^t, \pi^{t+1})$ as follows: Set $G(a) :=$
$G_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t) = Q_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t) - \lambda \log \dfrac{\pi_h^t(a \mid s)}{\sigma_h(a \mid s)}$. Note that $\max_{a,a' \in \mathcal{A}} |G(a') - G(a)| \leq$
η^{*-1} by Lemma D.3. By the update rule (4.4) and concavity of the logarithmic function $\log$, we have

$$D_{\mu^*}(\pi^t, \pi^{t+1})$$

$$= \sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\sum_{a \in \mathcal{A}} \pi_h^t(a \mid s) \log \frac{\pi_h^t(a \mid s)}{\pi_h^{t+1}(a \mid s)} \right]$$

$$= \sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\sum_{a \in \mathcal{A}} \pi_h^t(a \mid s) \log \frac{\sum_{a' \in \mathcal{A}} (\sigma_h(a' \mid s))^{\lambda\eta}(\pi_h^t(a' \mid s))^{1-\lambda\eta} \exp\left(\eta Q_h^{\lambda,\sigma}(s, a', \pi^t, \mu^t)\right)}{(\sigma_h(a \mid s))^{\lambda\eta}(\pi_h^t(a \mid s))^{-\lambda\eta} \exp\left(\eta Q_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t)\right)} \right]$$

$$= \sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\sum_{a \in \mathcal{A}} \pi_h^t(a \mid s) \log \frac{\sum_{a' \in \mathcal{A}} \pi_h^t(a' \mid s) \exp\left(\eta Q_h^{\lambda,\sigma}(s, a', \pi^t, \mu^t) - \lambda\eta \log \frac{\pi_h^t(a' \mid s)}{\sigma_h(a' \mid s)}\right)}{\exp\left(\eta Q_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t) - \lambda\eta \log \frac{\pi_h^t(a \mid s)}{\sigma_h(a \mid s)}\right)} \right]$$

$$\leq \sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\log \sum_{a \in \mathcal{A}} \pi_h^t(a \mid s) \frac{\sum_{a' \in \mathcal{A}} \pi_h^t(a' \mid s) \exp\left(\eta Q_h^{\lambda,\sigma}(s, a', \pi^t, \mu^t) - \lambda\eta \log \frac{\pi_h^t(a' \mid s)}{\sigma_h(a' \mid s)}\right)}{\exp\left(\eta Q_h^{\lambda,\sigma}(s, a, \pi^t, \mu^t) - \lambda\eta \log \frac{\pi_h^t(a \mid s)}{\sigma_h(a \mid s)}\right)} \right].$$

$$(D.8)$$

If we take η to be $\eta \leq \eta^*$, it follows that

$$\eta(G(a') - G(a)) \leq 1,$$

for $a, a' \in \mathcal{A}$. Thus, we can use the inequality $e^x \leq 1 + x + x^2$ for $x \leq 1$ and obtain

$$D_{\mu^*}(\pi^t, \pi^{t+1})$$

$$\leq \sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\log \sum_{a,a' \in \mathcal{A}} \pi_h^t(a \mid s) \pi_h^t(a' \mid s) e^{\eta(G(a')-G(a))} \right]$$

$$=$$

$$\leq \sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\log \sum_{a,a' \in \mathcal{A}} \pi_h^t(a \mid s) \pi_h^t(a' \mid s) \left(1 + \eta(G(a') - G(a)) + \eta^2(G(a') - G(a))^2\right) \right]$$

$$= \sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\log \sum_{a,a' \in \mathcal{A}} \pi_h^t(a \mid s) \pi_h^t(a' \mid s) \left(1 + (G(a') - G(a))^2\right) \right]$$

$$= \sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\log \left(1 + \eta^2 \sum_{a,a' \in \mathcal{A}} \pi_h^t(a \mid s) \pi_h^t(a' \mid s) (G(a') - G(a))^2\right) \right]$$

$$\leq \eta^2 \sum_{h=1}^{H} \mathbb{E}_{s \sim \mu_h^*} \left[\sum_{a,a' \in \mathcal{A}} \pi_h^t(a \mid s) \pi_h^t(a' \mid s) (G(a') - G(a))^2 \right].$$

By Lemma D.3, we can see that

$$\sum_{a,a' \in \mathcal{A}} \pi_h^t(a \mid s) \pi_h^t(a' \mid s) (G(a') - G(a))^2$$

$$\leq \sum_{a,a' \in \mathcal{A}} \pi_h^t(a \mid s) \pi_h^t(a' \mid s) \left(2L \sum_{l=h}^{H} \left\|\mu_l^t - \mu_l^*\right\|_1 + C^{\lambda,\sigma,H,|\mathcal{A}|}\left(E_h(a, \pi^t, \varpi^*) + E_h(a', \pi^t, \varpi^*)\right)\right)^2$$

$$\leq \sum_{a,a'\in\mathcal{A}} \pi_h^t(a\mid s)\,\pi_h^t(a'\mid s)\left(8L^2\left(\sum_{l=h}^{H}\left\|\mu_l^t-\mu_l^*\right\|_1\right)^2 + 4\left(C^{\lambda,\sigma,H,|\mathcal{A}|}\right)^2\left(E_h^2(a,\pi^t,\varpi^*)+E_h^2(a',\pi^t,\varpi^*)\right)\right)$$

$$\leq 8L^2H\sum_{l=h}^{H}\left\|\mu_l^t-\mu_l^*\right\|_1^2 + 8\left(C^{\lambda,\sigma,H,|\mathcal{A}|}\right)^2\sum_{a\in\mathcal{A}}\pi_h^t(a\mid s)\,E_h^2(a,\pi^t,\varpi^*)$$

$$= 8L^2H\sum_{l=h}^{H}\left\|\mu_l^t-\mu_l^*\right\|_1^2 + 8\left(C^{\lambda,\sigma,H,|\mathcal{A}|}\right)^2\sum_{a\in\mathcal{A}}\frac{\pi_h^t(a\mid s)}{\varpi_h^*(a\mid s)}\varpi_h^*(a\mid s)\,E_h^2(a,\pi^t,\varpi^*)$$

$$\leq 8L^2H\sum_{l=h}^{H}\left\|\mu_l^t-\mu_l^*\right\|_1^2 + \frac{8\left(C^{\lambda,\sigma,H,|\mathcal{A}|}\right)^2}{|\mathcal{A}|\exp\left(\frac{H(1-\lambda\log\sigma_{\min})}{\lambda}\right)}\sum_{a\in\mathcal{A}}\varpi_h^*(a\mid s)\,E_h^2(a,\pi^t,\varpi^*)$$

$$\leq 8L^2H\sum_{l=h}^{H}\left\|\mu_l^t-\mu_l^*\right\|_1^2 + \frac{8H\left(C^{\lambda,\sigma,H,|\mathcal{A}|}\right)^2}{|\mathcal{A}|\exp\left(\frac{H(1-\lambda\log\sigma_{\min})}{\lambda}\right)}\sum_{l=h}^{H}\mathbb{E}_{s_l\sim\mu_l^*}\left[\left\|\pi_l^*(s_l)-\pi_l^t(s_l)\right\|_1^2\right]$$

$$\leq 8L^2H\sum_{l=h}^{H}\left\|\mu_l^t-\mu_l^*\right\|_1^2 + \frac{4H\left(C^{\lambda,\sigma,H,|\mathcal{A}|}\right)^2}{|\mathcal{A}|\exp\left(\frac{H(1-\lambda\log\sigma_{\min})}{\lambda}\right)}D_{\mu^*}(\varpi^*,\pi^t).$$

Moreover, Lemma E.6 bounds $\sum_{l=h}^{H}\left\|\mu_l^t-\mu_l^*\right\|_1^2$ as

$$\sum_{l=h}^{H}\left\|\mu_l^t-\mu_l^*\right\|_1^2$$

$$\leq \sum_{l=h}^{H}\left(\sum_{k=0}^{l-1}\mathbb{E}_{s_k\sim\mu_k^*}\left[\left\|\pi_k^*(s_k)-\pi_k^t(s_k)\right\|\right]\right)^2$$

$$\leq H\sum_{l=h}^{H}\sum_{k=0}^{l-1}\mathbb{E}_{s_k\sim\mu_k^*}\left[\left\|\pi_k^*(s_k)-\pi_k^t(s_k)\right\|^2\right]$$

$$\leq \frac{1}{2}H^2 D_{\mu^*}(\varpi^*,\pi^t).$$

Therefore, we finally obtain

$$D_{\mu^*}(\varpi^*,\pi^{t+1}) \leq \left(1-\lambda\eta+C\eta^2\right)D_{\mu^*}(\varpi^*,\pi^t) \leq \left(1-\frac{1}{2}\lambda\eta\right)D_{\mu^*}(\varpi^*,\pi^t), \tag{D.9}$$

where we use $C\eta \leq C\eta^* \leq 1/2$. $\blacksquare$

E   USEFUL LEMMAS

For Mean-field games, one can write down the *Bellman optimality equation* as follows: for a function $Q':\mathcal{S}\to\Delta(\mathcal{A})$, a policy $\pi':\mathcal{S}\to\Delta(\mathcal{A})$, $\sigma':\mathcal{S}\to\Delta(\mathcal{A})$ and $s\in\mathcal{S}$ set

$$f_s^{\sigma'}(Q',\pi') = \langle Q'(s),\pi'(s)\rangle - \lambda D_{\mathrm{KL}}(\pi'(s),\sigma'(s)). \tag{E.1}$$

Lemma E.1. *Let (μ^*,ϖ^*) be equilibrium in the sense of Definition 4.2. Then, it holds that*

$$\varpi_h^*(s) = \arg\max_{p\in\Delta(\mathcal{A})}f_s^{\sigma_h}\left(Q_h^{\lambda,\sigma}(s,\bullet,\varpi^*,\mu^*),p\right) \propto \sigma_h(\bullet\mid s)\exp\left(\frac{Q_h^{\lambda,\sigma}(s,\bullet,\varpi^*,\mu^*)}{\lambda}\right),$$

for each $s\in\mathcal{S}$ and $h\in[H]$. Moreover,

$$\left\langle Q_h^{\lambda,\sigma}(s,\bullet,\varpi^*,\mu^*) - \lambda\log\frac{\pi_h^*(s)}{\sigma_h(s)},\delta\right\rangle = 0,$$

for all $\delta\in\mathbb{R}^{|\mathcal{A}|}$ such that $\sum_a\delta(a) = 0$.

Proof. See the Bellman optimality equation (e.g., (Agarwal et al., 2022, Theorem 1.9)). ∎

Lemma E.2. *Under Assumption 2.2, it holds that, for all $\pi, \widetilde{\pi} \in (\Delta(\mathcal{A})^{\mathcal{S}})^H$,*

$$J^{\lambda,\sigma}(m[\pi], \pi) + J^{\lambda,\sigma}(m[\widetilde{\pi}], \widetilde{\pi}) - J^{\lambda,\sigma}(m[\pi], \widetilde{\pi}) - J^{\lambda,\sigma}(m[\widetilde{\pi}], \pi) \leq 0,$$

where m is defined in (2.1).

Proof of Lemma E.2. The proof is similar to (Zhang et al., 2023, §H). Set $\mu = m[\pi]$ and $\widetilde{\mu} = m[\widetilde{\pi}]$. One can obtain that

$$J^{\lambda,\sigma}(m[\pi], \pi) + J^{\lambda,\sigma}(m[\widetilde{\pi}], \widetilde{\pi}) - J^{\lambda,\sigma}(m[\pi], \widetilde{\pi}) - J^{\lambda,\sigma}(m[\widetilde{\pi}], \pi)$$

$$= (J^{\lambda,\sigma}(\mu, \pi) - J^{\lambda,\sigma}(\widetilde{\mu}, \pi)) + (J^{\lambda,\sigma}(\widetilde{\mu}, \widetilde{\pi}) - J^{\lambda,\sigma}(\mu, \widetilde{\pi}))$$

$$= \sum_{h=1}^{H} \sum_{s_h \in \mathcal{S}} m[\pi]_h(s_h) \sum_{a_h \in \mathcal{A}} \pi_h(a_h \mid s_h)(r_h(s_h, a_h, \mu_h) - r_h(s_h, a_h, \widetilde{\mu}_h))$$

$$+ \sum_{h=1}^{H} \sum_{s_h \in \mathcal{S}} m[\widetilde{\pi}]_h(s_h) \sum_{a_h \in \mathcal{A}} \widetilde{\pi}_h(a_h \mid s_h)(r_h(s_h, a_h, \widetilde{\mu}_h) - r_h(s_h, a_h, \mu_h))$$

$$= \sum_{h,s,a} (\pi_h(a \mid s)\mu_h(s) - \widetilde{\pi}_h(a \mid s)\widetilde{\mu}_h(s))(r_h(s_h, a_h, \mu_h) - r_h(s_h, a_h, \widetilde{\mu}_h)),$$

and the right-hand side of the above inequality is less than 0 by Assumption 2.2. ∎

Lemma E.3. *Let $V_h^{\lambda,\sigma}$ be the state value function defined in (4.2) and $Q_h^{\lambda,\sigma}$ be the state action value function defined in (4.3). For any $s \in \mathcal{A}$, $a \in \mathcal{A}$, and $h \in [H]$, it holds that*

$$\lambda(H - h + 1) \log \sigma_{\min} \leq V_h^{\lambda,\sigma}(s, \mu, \pi) \leq H - h + 1,$$

$$\lambda(H - h + 1) \log \sigma_{\min} \leq Q_h^{\lambda,\sigma}(s, a, \mu, \pi) \leq H - h + 2.$$

Proof. We prove the inequalities by backward induction on h. By definition, we have

$$\max_{s \in \mathcal{S}} V_h^{\lambda,\sigma}(s, \mu, \pi) = \mathbb{E}\left[\sum_{l=h}^{H} (r_l(s_l, a_l, \mu_l) - \lambda D_{\mathrm{KL}}(\pi_l(s_l), \sigma_l(s_l))) \,\Bigg|\, s_h = s\right]$$

$$= \langle r_h(s, \bullet, \mu_h), \pi_h(s) \rangle - \lambda D_{\mathrm{KL}}(\pi_h(s_h), \sigma_h(s_h))$$

$$+ \sum_{s_{h+1} \in \mathcal{S}} V_{h+1}^{\lambda,\sigma}(s_{h+1}, \mu, \pi) \sum_{a_h \in \mathcal{A}} P_h(s_{h+1} \mid s, a_h)\pi_h(a_h \mid s)$$

$$\leq 1 + \max_{s_{h+1} \in \mathcal{S}} V_{h+1}^{\lambda,\sigma}(s_{h+1}, \mu, \pi),$$

and

$$\min_{s \in \mathcal{S}} V_h^{\lambda,\sigma}(s, \mu, \pi) = \langle r_h(s, \bullet, \mu_h), \pi_h(s) \rangle - \lambda D_{\mathrm{KL}}(\pi_h(s_h), \sigma_h(s_h))$$

$$+ \sum_{s_{h+1} \in \mathcal{S}} V_{h+1}^{\lambda,\sigma}(s_{h+1}, \mu, \pi) \sum_{a_h \in \mathcal{A}} P_h(s_{h+1} \mid s, a_h)\pi_h(a_h \mid s)$$

$$\geq \lambda \log \sigma_{\min} + \max_{s_{h+1} \in \mathcal{S}} V_{h+1}^{\lambda,\sigma}(s_{h+1}, \mu, \pi).$$

Then, we have

$$V_h^{\lambda,\sigma}(s, \mu, \pi) \in [\lambda(H - h + 1) \log \sigma_{\min}, H - h + 1],$$

by the induction. The definition of $Q_h^{\lambda,\sigma}$ in (4.3) immediately yields the bound. ∎

Lemma E.4. *For all $\pi, \widetilde{\pi} \in (\Delta(\mathcal{A})^{\mathcal{S}})^H$, it holds that*

$$\sum_{h=1}^{H} \mathbb{E}_{s \sim m[\widetilde{\pi}]_h}\left[\left\langle (\pi_h - \widetilde{\pi}_h)(s), Q_h^{\lambda,\sigma}(s, \bullet, \pi, \mu) \right\rangle\right] = J^{\lambda,\sigma}(\mu, \pi) - J^{\lambda,\sigma}(\mu, \widetilde{\pi}) - \lambda D_{m[\widetilde{\pi}]}(\widetilde{\pi}, \sigma) + \lambda D_{m[\widetilde{\pi}]}(\pi, \sigma).$$

Proof. From the definition of $V^{\lambda,\sigma}$ and $Q^{\lambda,\sigma}$ in (4.2) and (4.3), we have

$$\sum_{h=1}^{H} \mathbb{E}_{s\sim m[\widetilde{\pi}]_h} \left[\left\langle \pi_h(s), Q_h^{\lambda,\sigma}(s, \bullet, \pi, \mu) \right\rangle \right]$$

$$= \sum_{h=1}^{H} \mathbb{E}_{s\sim m[\widetilde{\pi}]_h} \left[\left\langle \pi_h(s), r_h(s, \bullet, \mu_h) + \mathbb{E}\left[V_{h+1}^{\lambda,\sigma}(s_{h+1}, \mu, \pi) \mid s_{h+1} \sim P(s, \bullet, \mu_h) \right] \right\rangle \right]$$

$$= \sum_{h=1}^{H} \mathbb{E}_{s_h\sim m[\widetilde{\pi}]_h} \left[\mathbb{E}_{a_h\sim \pi_h(s)} \left[r_h(s_h, a_h, \mu_h) - \lambda D_{\mathrm{KL}}(\pi(s_h), \sigma(s_h)) \right] \right] + \lambda D_{m[\widetilde{\pi}]}(\pi, \sigma)$$

$$+ \sum_{h=1}^{H} \mathbb{E}_{s\sim m[\widetilde{\pi}]_h} \left[\mathbb{E}\left[V_{h+1}^{\lambda,\sigma}(s_{h+1}, \mu, \pi) \mid s_{h+1} \sim P(s, a_h, \mu_h), a_h \sim \pi_h(s) \right] \right] \tag{E.2}$$

$$= \sum_{h=1}^{H} \mathbb{E}_{s_h\sim m[\widetilde{\pi}]_h} \left[V_h^{\lambda,\sigma}(s_h, \mu, \pi) - \mathbb{E}\left[V_{h+1}^{\lambda,\sigma}(s_{h+1}, \mu, \pi) \;\middle|\; \begin{matrix} s_{h+1} \sim P(s, a_h, \mu_h), \\ a_h \sim \pi_h(s) \end{matrix} \right] \right]$$

$$+ \lambda D_{m[\widetilde{\pi}]}(\pi, \sigma)$$

$$+ \sum_{h=1}^{H} \mathbb{E}_{s\sim m[\widetilde{\pi}]_h} \left[\mathbb{E}\left[V_{h+1}^{\lambda,\sigma}(s_{h+1}, \mu, \pi) \;\middle|\; \begin{matrix} s_{h+1} \sim P(s, a_h, \mu_h), \\ a_h \sim \pi_h(s) \end{matrix} \right] \right]$$

$$= \sum_{h=1}^{H} \mathbb{E}_{s\sim m[\widetilde{\pi}]_h} \left[V_h^{\lambda,\sigma}(s, \mu, \pi) \right] + \lambda D_{m[\widetilde{\pi}]}(\pi, \sigma).$$

Similarly, (4.1) and (2.1) gives us

$$\sum_{h=1}^{H} \mathbb{E}_{s\sim m[\widetilde{\pi}]_h} \left[\left\langle \widetilde{\pi}_h(s), Q_h^{\lambda,\sigma}(s, \bullet, \pi, \mu) \right\rangle \right]$$

$$= \sum_{h=1}^{H} \mathbb{E}_{s_h\sim m[\widetilde{\pi}]_h} \left[\mathbb{E}_{a_h\sim \widetilde{\pi}_h(s)} \left[r_h(s_h, a_h, \mu_h) - \lambda D_{\mathrm{KL}}(\widetilde{\pi}(s_h), \sigma(s_h)) \right] \right] + \lambda D_{m[\widetilde{\pi}]}(\widetilde{\pi}, \sigma)$$

$$+ \sum_{h=1}^{H} \mathbb{E}_{s\sim m[\widetilde{\pi}]_h} \left[\mathbb{E}\left[V_{h+1}^{\lambda,\sigma}(s_{h+1}, \mu, \pi) \mid s_{h+1} \sim P(s, a_h, \mu_h), a_h \sim \widetilde{\pi}_h(s) \right] \right] \tag{E.3}$$

$$= J^{\lambda,\sigma}(\mu, \widetilde{\pi}) + \lambda D_{m[\widetilde{\pi}]}(\widetilde{\pi}, \sigma) + \sum_{h=1}^{H} \mathbb{E}_{s\sim m[\widetilde{\pi}]_{h+1}} \left[V_{h+1}^{\lambda,\sigma}(s, \mu, \pi) \right].$$

Combining (E.2) and (E.3) yields

$$\sum_{h=1}^{H} \mathbb{E}_{s\sim m[\widetilde{\mu}]_h} \left[\left\langle (\pi_h - \widetilde{\pi}_h)(s), Q_h^{\lambda,\sigma}(s, \bullet, \pi, \mu) \right\rangle \right]$$

$$= \left(\sum_{h=1}^{H} \mathbb{E}_{s\sim m[\widetilde{\pi}]_h} \left[V_h^{\lambda,\sigma}(s, \mu, \pi) \right] + \lambda D_{m[\widetilde{\pi}]}(\pi, \sigma) \right)$$

$$- \left(J^{\lambda,\sigma}(\mu, \widetilde{\pi}) + \lambda D_{m[\widetilde{\pi}]}(\widetilde{\pi}, \sigma) + \sum_{h=1}^{H} \mathbb{E}_{s\sim m[\widetilde{\pi}]_{h+1}} \left[V_{h+1}^{\lambda,\sigma}(s, \mu, \pi) \right] \right)$$

$$= \left(\mathbb{E}_{s\sim m[\widetilde{\pi}]_1} \left[V_1^{\lambda,\sigma}(s, \mu, \pi) \right] + \lambda D_{m[\widetilde{\pi}]}(\pi, \sigma) \right) - \left(J^{\lambda,\sigma}(\mu, \widetilde{\pi}) + \lambda D_{m[\widetilde{\pi}]}(\widetilde{\pi}, \sigma) \right)$$

$$= \mathbb{E}_{s\sim \mu_1} \left[V_1^{\lambda,\sigma}(s, \mu, \pi) \right] - J^{\lambda,\sigma}(\mu, \widetilde{\pi}) + \lambda D_{m[\widetilde{\pi}]}(\pi, \sigma) - \lambda D_{m[\widetilde{\pi}]}(\widetilde{\pi}, \sigma),$$

which concludes the proof. ∎

Lemma E.5. *For all* $\pi, \widetilde{\pi} \in (\Delta(\mathcal{A})^{\mathcal{S}})^H$, *it holds that*

$$\sum_{h=1}^{H} \mathbb{E}_{s\sim m[\widetilde{\pi}]_h} \left[\left\langle (\pi_h - \widetilde{\pi}_h)(s), \log \frac{\pi_h(s)}{\sigma_h(s)} \right\rangle \right] = D_{m[\widetilde{\pi}]}(\pi, \sigma) - D_{m[\widetilde{\pi}]}(\widetilde{\pi}, \sigma) + D_{\widetilde{\pi}}(\widetilde{\pi}, \pi).$$

Proof. A direct computation yields

$$\sum_{h=1}^{H} \mathbb{E}_{s\sim m[\widetilde{\pi}]_h}\left[\left\langle (\pi_h - \widetilde{\pi}_h)(s), \log\frac{\pi_h(s)}{\sigma_h(s)}\right\rangle\right]$$

$$=D_{m[\widetilde{\pi}]}(\pi,\sigma) - \sum_{h=1}^{H}\mathbb{E}_{s\sim m[\widetilde{\pi}]_h}\left[\left\langle \widetilde{\pi}_h(s), \log\frac{\widetilde{\pi}_h(s)}{\sigma_h(s)} - \log\frac{\widetilde{\pi}(s)}{\pi(s)}\right\rangle\right]$$

$$=D_{m[\widetilde{\pi}]}(\pi,\sigma) - D_{m[\widetilde{\pi}]}(\widetilde{\pi},\sigma) + D_{m[\widetilde{\pi}]}(\widetilde{\pi},\pi).$$

∎

Lemma E.6. *The operator m defined in (2.1) is 1-Lipschitz, namely, it holds that*

$$\|m[\pi]_{h+1} - m[\pi']_{h+1}\| \leq \sum_{l=0}^{h} \mathbb{E}_{s_l\sim m[\pi]_l}\left[\|\pi_l(s_l) - \pi'_l(s_l)\|\right], \tag{E.4}$$

for $\pi, \pi' \in (\Delta(\mathcal{A})^{\mathcal{S}})^H$ and all $h \in \{0,\ldots,H\}$. Here, we set $\pi_0(s) = \pi'_0(s) = \mathrm{U}_{\mathcal{A}}$ for all $s \in \mathcal{S}$.

Proof. Fix $\pi, \pi' \in (\Delta(\mathcal{A})^{\mathcal{S}})^H$. We prove the inequality by induction on h.

(I) Base step $h = 0$: It is obvious because $\|m[\pi]_1 - m[\pi']_1\| = \|\mu_1 - \mu_1\| = 0$.

(II) Inductive step: Suppose that there exists $h \in [H]$ satisfying the inequality (E.4). By (2.1), we obtain

$\|m[\pi]_{h+2} - m[\pi']_{h+2}\|$

$$\leq \sum_{\substack{s_{h+2}\in\mathcal{S},\\ (s_{h+1},a_{h+1})\in\mathcal{S}\times\mathcal{A}}} P_{h+1}(s_{h+2} \mid s_{h+1}, a_{h+1}) m[\pi]_{h+1}(s_{h+1})\big|\pi_{h+1}(a_{h+1} \mid s_{h+1}) - \pi'_{h+1}(a_{h+1} \mid s_{h+1})\big|$$

$$+ \sum_{\substack{s_{h+2}\in\mathcal{S},\\ (s_{h+1},a_{h+1})\in\mathcal{S}\times\mathcal{A}}} P_{h+1}(s_{h+2} \mid s_{h+1}, a_{h+1}) \pi'_{h+1}(a_{h+1} \mid s_{h+1})\big|m[\pi]_{h+1}(s_{h+1}) - m[\pi']_{h+1}(s_{h+1})\big|$$

$$\leq \sum_{(s_{h+1},a_{h+1})\in\mathcal{S}\times\mathcal{A}} m[\pi]_{h+1}(s_{h+1})\big|\pi_{h+1}(a_{h+1} \mid s_{h+1}) - \pi'_{h+1}(a_{h+1} \mid s_{h+1})\big|$$

$$+ \sum_{s_{h+1}\in\mathcal{S}} \big|m[\pi]_{h+1}(s_{h+1}) - m[\pi']_{h+1}(s_{h+1})\big|$$

$$= \mathbb{E}_{s_{h+1}\sim m[\pi]_{h+1}}\left[\big\|\pi_{h+1}(s_{h+1}) - \pi'_{h+1}(s_{h+1})\big\|\right] + \|m[\pi]_{h+1} - m[\pi']_{h+1}\|.$$

By the hypothesis of the induction, we finally obtain

$\|m[\pi]_{h+2} - m[\pi']_{h+2}\|$

$$\leq \mathbb{E}_{s\sim m[\pi]_{h+1}}\left[\big\|\pi_{h+1}(s) - \pi'_{h+1}(s)\big\|\right] + \sum_{l=1}^{h} \mathbb{E}_{s\sim m[\pi]_l} \|\pi_l(s) - \pi'_l(s)\|$$

$$\leq \sum_{l=1}^{h+1} \mathbb{E}_{s\sim m[\pi]_l} \|\pi_l(s) - \pi'_l(s)\|.$$

∎

Lemma E.7. *Let $\pi, \pi' \in (\Delta(\mathcal{A})^{\mathcal{S}})^H$, $\mu, \mu' \in \Delta(\mathcal{S})^H$, $s \in \mathcal{S}$, and $h \in \{1,\ldots,H+1\}$. Assume*

$$\min_{(h,a,s)\in[H]\times\mathcal{A}\times\mathcal{S}} \min\{\pi_h(a \mid s), \pi'_h(a \mid s)\} > 0,$$

and set $\mu_{H+1} = \mu'_{H+1} = \mathrm{U}_{\mathcal{S}}$, $\pi_{H+1}(s) = \pi'_{H+1}(s) = \mathrm{U}_{\mathcal{A}}$ for all $s \in \mathcal{S}$.

$$\left|V_h^{\lambda,\sigma}(s,\pi,\mu) - V_h^{\lambda,\sigma}(s,\pi',\mu')\right|$$

$$\leq \mathbb{E}\left[\sum_{l=h}^{H+1}\left(C^{\lambda,\sigma}(\pi,\pi')\|\pi_l(s_l) - \pi'_l(s_l)\|_1 + L\|\mu_l - \mu'_l\|_1\right)\ \middle|\ \begin{array}{c} s_h = s, \\ s_{l+1}\sim P_l(s_l,a_l), \\ a_l \sim \pi_l(s_l) \\ \text{for each } l \in \{h,\ldots,H+1\} \end{array}\right]$$

for Here, $C^{\lambda,\sigma}(\pi,\pi') > 0$ is defined in Proposition E.8, and the discrete time stochastic process $(s_l)_{l=h}^H$ is induced recursively as $s_{l+1} \sim P_l(s_l, a_l), a_l \sim \pi_l(s_l)$ for each $l \in \{h, \ldots, H-1\}$.

Proof. Fix π, π', μ and μ'. We prove the inequality by backward induction on h.

(I) Base step $h = H+1$: It is obvious because $\left|V_{H+1}^{\lambda,\sigma}(s,\pi,\mu) - V_{H+1}^{\lambda,\sigma}(s,\pi',\mu')\right| = |0-0| = 0$.

(II) Inductive step: Suppose that there exists $h \in [H]$ satisfying

$$
\left|V_{h+1}^{\lambda,\sigma}(s,\pi,\mu) - V_{h+1}^{\lambda,\sigma}(s,\pi',\mu')\right|
$$

$$
\leq \mathbb{E}\left[\sum_{l=h+1}^{H+1}\left(C^{\lambda,\sigma}(\pi,\pi')\|\pi_l(s_l) - \pi'_l(s_l)\|_1 + L\|\mu_h - \mu'_h\|_1\right) \left|\begin{array}{c} s_{h+1} = s, \\ s_{l+1} \sim P_l(s_l,a_l), \\ a_l \sim \pi_l(s_l) \\ \text{for each } l \in \{h+1,\ldots,H+1\} \end{array}\right.\right],
$$

$$(E.5)$$

for all $s \in \mathcal{S}$. By the definition of the value function in (4.2) and Assumption 2.3, we have

$$
\left|V_h^{\lambda,\sigma}(s,\pi,\mu) - V_h^{\lambda,\sigma}(s,\pi',\mu')\right|
$$

$$
\leq \left|\sum_{a_h \in \mathcal{A}}\left(\pi_h(a_h \mid s)r_h(s,a_h,\mu_h) - \pi'_h(a_h \mid s)r_h(s,a_h,\mu'_h)\right)\right|
$$

$$
+ \lambda|D_{\mathrm{KL}}(\pi_h(s),\sigma_h(s)) - D_{\mathrm{KL}}(\pi'_h(s),\sigma_h(s))|
$$

$$
+ \left|\sum_{\substack{a_h \in \mathcal{A}, \\ s_{h+1} \in \mathcal{S}}} P_h(s_{h+1} \mid s,a_h)\left(\pi_h(a_h \mid s)V_{h+1}^{\lambda,\sigma}(s_{h+1},\pi,\mu) - \pi'(a_h \mid s)V_{h+1}^{\lambda,\sigma}(s_{h+1},\pi',\mu')\right)\right|
$$

$$
\leq \|\pi_h(s) - \pi'_h(s)\|_1 + \sum_{a_h \in \mathcal{A}}\pi_h(a_h \mid s)|r_h(s,a_h,\mu_h) - r_h(s,a_h,\mu'_h)|
$$

$$
+ \lambda\left|\sum_{a_h \in \mathcal{A}}\left(\pi_h(a_h \mid s)\left(\log\frac{\pi_h(a_h \mid s)}{\sigma_h(a_h \mid s)} - 1\right) - \pi'_h(a_h \mid s)\left(\log\frac{\pi'_h(a_h \mid s)}{\sigma_h(a_h \mid s)} - 1\right)\right)\right|
$$

$$
+ \|\pi_h(s) - \pi'_h(s)\|_1
$$

$$
+ \sum_{\substack{a_h \in \mathcal{A}, \\ s_{h+1} \in \mathcal{S}}} P_h(s_{h+1} \mid s,a_h)\pi_h(a_h \mid s)\left|V_{h+1}^{\lambda,\sigma}(s_{h+1},\pi,\mu) - V_{h+1}^{\lambda,\sigma}(s_{h+1},\pi',\mu')\right|
$$

$$
\leq 2\|\pi_h(s) - \pi'_h(s)\|_1 + L\|\mu_h - \mu'_h\|_1
$$

$$
+ \lambda \max_{(h,a,s)} \log\frac{1}{(\sigma\pi\pi')_h(a \mid s)}\|\pi_h(s) - \pi'_h(s)\|_1
$$

$$
+ \sum_{\substack{a_h \in \mathcal{A}, \\ s_{h+1} \in \mathcal{S}}} P_h(s_{h+1} \mid s,a_h)\pi_h(a_h \mid s)\left|V_{h+1}^{\lambda,\sigma}(s_{h+1},\pi,\mu) - V_{h+1}^{\lambda,\sigma}(s_{h+1},\pi',\mu')\right|
$$

$$
\leq C^{\lambda,\sigma}(\pi,\pi')\|\pi_h(s) - \pi'_h(s)\|_1 + L\|\mu_h - \mu'_h\|_1
$$

$$
+ \mathbb{E}\left[\left|V_{h+1}^{\lambda,\sigma}(s_{h+1},\pi,\mu) - V_{h+1}^{\lambda,\sigma}(s_{h+1},\pi',\mu')\right| \left|\begin{array}{c} s_h = s, \\ s_{h+1} \sim P_h(s_h,a_h), \\ a_h \sim \pi_h(s_h) \end{array}\right.\right].
$$

Combining the above inequality and the hypothesis of the induction completes the proof. $\blacksquare$

Proposition E.8. *Let $Q^{\lambda,\sigma}$ be the function defined by (4.3), and $(\pi, \pi') \in \left((\Delta(\mathcal{A})^{\mathcal{S}})^H\right)^2$ be policies with full supports. Under Assumptions 2.3 and 4.1, it holds that*

$$\left| Q_h^{\lambda,\sigma}(s, a, \pi, \mu) - Q_h^{\lambda,\sigma}(s, a, \pi', \mu') \right|$$

$$\leq L \sum_{l=h}^H \|\mu_l - \mu_l'\| + C^{\lambda,\sigma}(\pi, \pi') \, \mathbb{E}_{(s_l)_{l=h+1}^H} \left[\left. \sum_{l=h+1}^H \|\pi_l(s_l) - \pi_l'(s_l)\| \, \right| \, s_h = s \right],$$

for $(h, s, a) \in [H] \times \mathcal{S} \times \mathcal{A}$ and $\mu, \mu' \in \Delta(\mathcal{S})^H$. Here, the random variables $(s_l)_{l=h+1}^H$ follows the stochastic process starting from state s at time h, induced from P and π, and the function $C^{\lambda,\sigma}: \left((\Delta(\mathcal{A})^{\mathcal{S}})^H\right)^2 \to \mathbb{R}$ is given by $C^{\lambda,\sigma}(\pi, \pi') = 2 - \lambda \inf_{(h,s,a) \in [H] \times \mathcal{S} \times \mathcal{A}} \log (\sigma \pi \pi')_h (a \mid s)$.

Proof of Proposition E.8. Let h be larger than 2. By the definition of $Q_h^{\lambda,\sigma}$ given in (4.3) and Lemma E.7, we have

$$\left| Q_{h-1}^{\lambda,\sigma}(s, a, \pi, \mu) - Q_{h-1}^{\lambda,\sigma}(s, a, \pi', \mu') \right|$$

$$\leq \left| r_{h-1}(s, a, \mu_{h-1}) - r_{h-1}(s, a, \mu_{h-1}') \right| + \mathbb{E}_{s_h \sim P_{h-1}(s,a)} \left[\left| V_h^{\lambda,\sigma}(s_h, \pi, \mu) - V_h^{\lambda,\sigma}(s_h, \pi', \mu') \right| \right]$$

$$\leq L \left\| \mu_{h-1} - \mu_{h-1}' \right\| + \mathbb{E}_{s_h \sim P_{h-1}(s,a)} \left[\left| V_h^{\lambda,\sigma}(s_h, \pi, \mu) - V_h^{\lambda,\sigma}(s_h, \pi', \mu') \right| \right].$$

Combining the above inequality and Lemma E.7 completes the proof. ∎

F EXPERIMENT DETAILS

We ran experiments on a laptop with an 11th Gen Intel Core i7-1165G7 8-core CPU, 16GB RAM, running Windows 11 Pro with WSL. As is clear from Algorithm 2, our proposed method is deterministic. Thus, we ran the algorithm only once for each experimental setting. We implemented our proposed method using Python. The computation of $Q^{\lambda,\sigma}$ and μ in Algorithm 2 was based on the implementation provided by Fabian et al. (2023).

We show further details for Beach Bar Process. We set $H = 10, |\mathcal{S}| = 10, \mathcal{A} = \{-1, \pm 0, +1\}, \lambda = 0.1, \eta = 0.1$, and

$$P_h\left(s' \mid s, a\right) = \begin{cases} 1 - \varepsilon & \text{if } a = \pm 0 \ \& \ s' = s, \\ \dfrac{\varepsilon}{2} & \text{if } a = \pm 1 \ \& \ s' = s \pm 1, \\ 0 & \text{otherwise}, \end{cases}$$

where we choose $\epsilon = 0.1$. In addition, we initialize σ^0 and π^0 in Algorithm 2 as the uniform distributions on $\mathcal{A}$.