Branch-Price-and-Cut for Causal Discovery

Abstract

We show how to extend the integer programming (IP) approach to score-based causal discovery by including *pricing*. Pricing allows the addition of new IP variables during solving, rather than requiring them all to be present initially. The dual values of acyclicity constraints allow this addition to be done in a principled way. We have an implemented system PRICEILP which uses the SCIP library to effect a branch-price-and-cut method for DAG learning. Empirical results are presented showing how PRICEILP learns optimal DAGs and that implementing a *delayed pricing* approach can be beneficial. The current pricing algorithm in PRICEILP is slow, so further work on fast pricing is required.

Keywords: Causal discovery, integer linear programming, pricing

1. Introduction

We consider score-based approaches to causal discovery. In the score-based approach each causal model has a score—determined by the observed data, and perhaps prior knowledge—and the goal is to find whichever causal model has the best score. Typical scores include posterior probability and log-likelihood with some suitable complexity penalty. The problem of causal discovery thus reduces to an optimisation problem.

The principal problem with this approach is that the resulting optimisation problem can be NP-hard. For example, Chickering et al. (2004) showed that Bayesian network learning (i.e. causal discovery assuming causal sufficiency) is NP-hard even with 'large' samples. Assuming $P \neq NP$ this means that solving the causal discovery optimisation problem exactly (i.e. finding a provably globally optimal solution) is impractical in the general case. As a result most approaches to causal discovery are inexact (sometimes called *heuristic*): a search for high-scoring causal models is conducted and the best-scoring model is returned, but there is no guarantee that a better-scoring model does not exist. Many inexact methods use local search.

The great advantage of resorting to inexact methods is that it is often possible in practice to find high-scoring models quickly even when learning from large datasets. However, when exact learning is possible it is clearly preferable; the goal of this paper is to explore whether a particular method—*pricing*—can extend the applicability of exact learning.

2. Bayesian network structure learning using integer programming

We will focus on learning DAGs under the assumption of causal sufficiency, postponing discussion of other casual discovery tasks until Section 7. This problem is often called *Bayesian network structure learning* and we will abbreviate it to BNSL throughout. We will assume that the score is *decomposable*. Let P be the random variables observed in the data which correspond to the vertices of candidate DAGs. A decomposable score is one which is a sum of *local scores*, one for each BN variable, where the local score for each BN variable $i \in P$ is determined by the choice of *parents*

$$\text{MAXIMISE} \quad \sum_{\substack{i \in P \\ J \subseteq P \setminus \{i\}}} c'_{i \leftarrow J} x_{i \leftarrow J} \tag{1}$$

SUBJECT TO:

$$\sum_{J \subseteq P \setminus \{i\}} x_{i \leftarrow J} = 1 \qquad\qquad i \in P \tag{2}$$

$$\sum_{i \in C} \sum_{\substack{J \subseteq P \setminus \{i\} \\ J \cap C \neq \emptyset}} x_{i \leftarrow J} \leq |C| - 1 \qquad\qquad C \subseteq P, |C| \geq 2 \tag{3}$$

$$x_{i \leftarrow J} \in \{0, 1\}, i \in P, J \subseteq P \setminus \{i\} \tag{4}$$

Figure 1: Integer program for score-based BNSL with a decomposable score

$J \subseteq P \setminus \{i\}$ for that variable. BN variable j is a parent of BN variable i iff there is a directed edge $i \leftarrow j$ in the DAG in which case i is called a *child* of j.

Let $x_{i \leftarrow J}$ be the binary variable indicating that BN variable i has parent set J. The $x_{i \leftarrow J}$ will be called *family variables* since they indicate the choice of parents for a particular BN (child) variable. Clearly any directed graph has a unique encoding as a zero-one vector indexed by the family variables. Denote the local score for i having parents J by $c'_{i \leftarrow J}$. Each $c'_{i \leftarrow J}$ can be computed from the data we are learning from. If a score is decomposable it follows that the score for any DAG is $\sum c'_{i \leftarrow J} x_{i \leftarrow J}$ where the $x_{i \leftarrow J}$ encode the DAG.

Let x denote both the vector of all $x_{i \leftarrow J}$ family variables and also particular values of x—where the context will make clear which is meant. Finding an x which maximises $\sum c'_{i \leftarrow J} x_{i \leftarrow J}$ does not solve the BNSL problem since typically x will not represent a DAG. It is necessary to add *constraints* to ensure that the only feasible values of x are those which represent DAGs. One suitable set of constraints is given in Fig 1 where score-based BNSL with a decomposable score is represented as an *integer linear program*, or *integer program (IP)* for short. Fig 1 represents an IP since the objective function is linear and constraints are either linear (2,3) or are integrality constraints (4).

In the IP in Fig 1 the equations (2) ensure that exactly one parent set J is selected for each BN variable i, so that each feasible x at least represents a directed graph. Next note that for $C \subseteq P, |C| \geq 2$ if there is a cycle with vertices C then each $i \in C$ has a parent in C and we have $\sum_{i \in C} \sum_{J \subseteq P \setminus \{i\}, J \cap C \neq \emptyset} x_{i \leftarrow J} = |C|$, so the inequalities (3) are sufficient to ensure that only *acyclic* directed graphs are feasible. These inequalities (in a different formulation) were introduced by Jaakkola et al. (2010) under the name *cluster constraints*.

3. Solving the BNSL linear relaxation

Exact optimisation has two tasks. Just like inexact optimisation it aims to find solutions with high objective values (high scoring DAGs in this paper). But unlike inexact optimisation it also computes an upper bound on the objective of an optimal solution. When we have a solution whose objective value equals this upper bound then we know that it is a globally optimal solution.

The best solution found at any point in the solving process is known as the *incumbent* and in modern IP solvers incumbents are found by all sorts of heuristic algorithms. The current version of the IP solver SCIP (Bestuzheva et al., 2021) has no fewer than 58 heuristic algorithms available! Note that, in principle, *any* DAG learning algorithm could be used by an IP solver to provide an initial incumbent DAG, the task of the IP solver being then to search for yet better incumbents and to compute upper bounds in the hope of establishing optimality.

Upper bounds on optimal objective values are provided by solving *relaxations* of the original problem. A relaxation is produced by removing some constraints and solving the resulting problem to optimality. The objective value of an optimal solution of the relaxed problem provides the desired upper bound: it will always be no worse (i.e. not lower than) the objective value of an optimal solution to the original problem, precisely because the former is a less constrained version of the latter. It is, of course, crucial that this optimal solving can be performed quickly. In addition, it is preferable if the relaxation is *tight*, so that the upper bound it provides is close to the true optimal objective value.

For an IP problem the standard approach is to solve its *linear relaxation* where integrality constraints are removed, thus producing a *linear program (LP)*. LPs can be solved to optimality in polynomial time, although in practice they are quickly solved by the simplex algorithm which has worst-case exponential time. The linear relaxation of our BNSL IP (Fig 1) just replaces $x_{i \leftarrow J} \in \{0, 1\}$ with $x_{i \leftarrow J} \in [0, 1]$.

However, solving this linear relaxation is non-trivial since there are $2^p - p - 1$ cluster constraints (where $p = |P|$) and $p2^{p-1}$ family variables. So unless p is small, constructing and solving an LP with all variables and constraints present is impractical. Cussens (2011) addressed the too-many-cluster-constraints problem by adding these constraints as *cutting planes*. An initial LP is created with only a small set C of cluster constraints. This LP is solved and then a search is conducted for cutting planes—cluster constraints not satisfied by the solution to this LP. The set C of cluster constraints is then updated to include these cutting planes and the resulting new LP solved, and a new search for cutting planes is conducted. This iterative approach is continued until no cutting planes can be found or the linear relaxation provides a sufficiently good bound so that there is no point spending time searching for new cutting planes. Note that each cluster constraint $C \in C$ is a *facet*, a maximally tight inequality of the convex hull of zero-one vectors encoding DAGs (Cussens et al., 2017a,b). This is why they provide tight linear relaxations and why it is worth the effort to find them even though cycles can be ruled out with merely a quadratic number of constraints (Cussens, 2010).

Although the problem of finding these cutting planes is NP-hard (Cussens et al., 2017b), this has proved a reasonable approach to producing tight linear relaxations *if there are not too many family variables*. The key contribution of this paper is to investigate the use of *pricing* to iteratively introduce family variables, when adding all of them initially is impractical.

3.1. Pricing-in variables

To understand how pricing can be inserted into the cutting plane approach for BNSL one needs to consider the *dual* of the linear program that is solved at each stage. Let C be the current set of cluster constraints. Consider multiplying the equations (2) by coefficients λ_i and the inequalities (3) by λ_C where $C \in C, \lambda_C \geq 0$, then the sum of resulting inequalities is a valid inequality on the $x_{i \leftarrow J}$ since it was implied by the original constraints. (Each equation (2) can be viewed as a pair of

inequalities, the sign of λ_i indicating which one was selected.) Let $d_{i \leftarrow J}$ be the coefficient for $x_{i \leftarrow J}$ in this new inequality, then, as long as $c'_{i \leftarrow J} \leq d_{i \leftarrow J}$, the new inequality is an upper bound on the objective, equal to $\sum_{i \in P} \lambda_i + \sum_{C \in \mathcal{C}} (|C| - 1)\lambda_C$. The dual of the linear program is just the problem of finding values for the λ_i and the λ_C which lead to a lowest such upper bound. By the strong duality of linear programming the objective values of the solutions to the dual and the original (*primal*) linear program are equal (assuming both have a solution).

Let λ^* be the solution to the dual linear program. Note that any reasonable LP solver will compute λ^* in the course of solving the primal LP, a separate optimisation process is not required. Suppose that instead of including all family variables, only a subset $\mathcal{V}$ are in the LP, thus most probably not allowing an optimal solution to the (primal) LP. Let $x_{i \leftarrow J}$ be a family variable not in $\mathcal{V}$ whose local score $c'_{i \leftarrow J}$ is high enough so that $\lambda_i^* + \sum_{C \in \mathcal{C}, i \in C, C \cap J \neq \emptyset} \lambda_C^* < c'_{i \leftarrow J}$. Then if one were to add $x_{i \leftarrow J}$ to $\mathcal{V}$, the new dual linear program thus produced would have an additional constraint $\lambda_i + \sum_{C \in \mathcal{C}, i \in C, C \cap J \neq \emptyset} \lambda_C \geq c'_{i \leftarrow J}$ which the earlier solution λ^* does not satisfy. The addition of the variable $x_{i \leftarrow J}$ to the primal LP thus adds a cutting plane to the dual LP which renders the dual solution λ^* no longer feasible.

Pricing is the process of finding and adding new variables $x_{i \leftarrow J}$ which render the current dual solution infeasible. Typically adding such variables to the primal LP will result in an LP solution with a higher objective value. If no such variables can be found then it follows that the LP has been solved to optimality *even though only a subset of variables have been used*. All potential variables which are not included in the LP are implicitly set to 0 in this optimal solution.

The condition $\lambda_i^* + \sum_{C \in \mathcal{C}, i \in C, C \cap J \neq \emptyset} \lambda_C^* < c'_{i \leftarrow J}$ is equivalent to $c'_{i \leftarrow J} - \lambda_i^* - \sum_{C \in \mathcal{C}, i \in C, C \cap J \neq \emptyset} \lambda_C^* > 0$. We will call $c'_{i \leftarrow J} - \lambda_i^* - \sum_{C \in \mathcal{C}, i \in C, C \cap J \neq \emptyset} \lambda_C^*$ the *reduced local score* for $x_{i \leftarrow J}$. The goal of pricing is to find variables which, given the current dual values λ_i^*, λ_C^*, have positive reduced local score. If such variables are found they are added to $\mathcal{V}$ producing a new LP whose dual is solved leading to new dual values and a new pricing problem. Note that a new variable must be added to all constraints where it has non-zero coefficient. So $x_{i \leftarrow J}$ must be added to the constraint $\sum_{J \subseteq P \setminus \{i\}} x_{i \leftarrow J} = 1$ as well as to any cluster constraint whose cluster C has $i \in C$, $J \cap C \neq \emptyset$. In this way new variables are added to the problem in *pricing rounds*.

Although BNSL is most naturally represented as a maximisation problem, we will, from now on, recast it as a minimisation problem. This is because the solver we will actually use to do branch-price-and-cut, namely SCIP, always (internally) formulates optimisation as minimisation, and it is advantageous for our presentation to exactly match what is implemented. In this minimisation reformulation denote the objective coefficient for each family variable $x_{i \leftarrow J}$ to be $c_{i \leftarrow J}$ and set $c_{i \leftarrow J} = -c'_{i \leftarrow J}$. Call $c_{i \leftarrow J}$ the *cost* for $x_{i \leftarrow J}$. The dual LP will now be finding a lower bound and so for each $C \in \mathcal{C}$ we will have $\lambda_C \leq 0$. Now rather than seeking new variables with positive reduced local score we seek variables with *negative reduced cost* where the reduced cost for $x_{i \leftarrow J}$ is $c_{i \leftarrow J} - \lambda_i^* - \sum_{C \in \mathcal{C}, i \in C, C \cap J \neq \emptyset} \lambda_C^*$.

The minimal any pricing method must deliver is, on each pricing round, to find at least one variable with negative reduced cost or establish that no such variable exists. However, it is typically more efficient to add in several variables with negative reduced cost on each pricing round with a view to reducing the number of rounds. Also variables with more negative reduced costs are preferred in the hope that they correspond to 'deeper' cutting planes in the dual. One simple pricing strategy is, for each $i \in P$, to find a parent set J with minimal reduced cost (so that at most p variables are added in any pricing round). The pricing problem using this approach is show in Fig 2.

$$\text{MINIMISE}_J \qquad z = c_{i \leftarrow J} - \lambda_i^* - \sum_{\substack{C \in \mathcal{C}, i \in C \\ C \cap J \neq \emptyset}} \lambda_C^*$$

$$\text{SUBJECT TO} \qquad z < 0$$

Figure 2: Pricing problem for BN variable i where $\mathcal{C}$ is the current set of clusters and λ_i^* and λ_C^* are the current dual values for the constraints. This is for BNSL cast as a minimisation problem.

$$\text{MINIMISE}_J \qquad z = n \log \sigma_{i \leftarrow J}^2 + \Lambda^2 |J| - \sum_{\substack{C \in \mathcal{C}, i \in C \\ C \cap J \neq \emptyset}} \lambda_C^*$$

$$\text{SUBJECT TO} \qquad z < \lambda_i^*$$

Figure 3: Gaussian ℓ_0 pricing problem for BN variable i where $\mathcal{C}$ is the current set of clusters and λ_i^* and λ_C^* are the current dual values for the constraints.

3.2. Pricing for BNSL: a worked example

We will illustrate pricing when the BN score is the Gaussian ℓ_0-penalised maximum likelihood score for scoring Gaussian DAGs (van de Geer and Bühlmann, 2013). The ℓ_0-penalised maximum likelihood score is a decomposable score and the local score for variable i having parents J is:

$$c'_{i \leftarrow J} = L(i, J) - \Lambda^2 |J| \tag{5}$$

where

$$L(i, J) = -(n/2)[\log(2\pi) + 1 + \log \sigma_{i \leftarrow J}^2] \tag{6}$$

and where n is the size of the data, $n\sigma_{i \leftarrow J}^2$ is the sum of squared residuals resulting from OLS regression using variables J to predict variable i and Λ^2 is the tuning parameter for the ℓ_0 penalty $\Lambda^2 |J|$.

The cost is $c_{i \leftarrow J} = (n/2)[\log(2\pi) + 1 + \log \sigma_{i \leftarrow J}^2] + \Lambda^2 |J|$. After removing the constant terms $\log(2\pi) + 1$ and rescaling Λ^2 we end up with a simplified cost which when plugged into the pricing problem shown in Fig 2 and then rearranging leads to the pricing problem specific to Gaussian ℓ_0-penalised maximum likelihood shown in Fig 3. The objective in Fig 3 shows that this is a *doubly penalised regression problem*. There is $\Lambda^2 |J|$, the normal ℓ_0 penalty against overfitting, but also $-\sum_{C \in \mathcal{C}, i \in C C \cap J \neq \emptyset} \lambda_C^*$ which is a penalty for causing cycles (recall that $\lambda_C^* \leq 0, \forall C \in \mathcal{C}$). Note that λ_i^* is independent of J and so has been removed from the objective in Fig 3.

We now illustrate the price-and-cut method to solve a linear relaxation using the `gaussian.test` dataset from the BNLEARN R package (Scutari, 2010). This dataset consists of $n = 5000$ samples generated from a Gaussian distribution defined by choosing parameters for the 7-node DAG shown in Fig 4. The parameters can be found in the script `gaussian.test.R` (available from BN-LEARN) which was used to generate the sample. The ℓ_0 tuning parameter Λ^2 was set to 1. (In this

Figure 4: DAG used to generate `gaussian.test` dataset

example, the costs were not simplified by removing the constant terms $\log(2\pi) + 1$ and rescaling Λ^2.)

We start with an initial LP (LP 1) where we only have family variables for empty parent sets: $\mathcal{V} = \{x_{i\leftarrow\emptyset} : i = 1, \ldots, 7\}$ and $C = \emptyset$, so the only constraints are the $p = 7$ linear constraints (2). There is only one feasible solution to this LP, which is thus also optimal: all 7 of the variables are set to 1.

Solving the 7 Gaussian ℓ_0 pricing problems (Fig 3) for $i = 1, \ldots, 7$ using the dual values associated with this optimal solution finds the following family variables with maximal reduced local score: $x_{1\leftarrow\{2,3,4,5,6,7\}}$, $x_{2\leftarrow\{1,3,4\}}$, $x_{3\leftarrow\{1,2\}}$, $x_{4\leftarrow\{1,2,5,6,7\}}$, $x_{5\leftarrow\{1,4,6,7\}}$, $x_{6\leftarrow\{1,3,4,5,7\}}$ and $x_{7\leftarrow\{1,3,4,5,6\}}$. These variables have costs -466.13, -1921.01, 3718.43, 1238.48, 6507.84, 7076.24 and 4800.51 respectively and *reduced costs* of -7581.44, -14564.78, -12624.79, -13440.29, -4026.65, -9132.33 and -5718.30, respectively. Since at this point there is no penalty for cycles these are, inevitably, just the parent sets with minimal cost (=maximal local score) for each $i \in P$, with the reduced cost quantifying how much better these parent sets are than the empty parent set. As expected, each minimal cost parent set approximates the Markov blanket of the child variable in the true data-generating BN (Fig 4). For $i = 1, 2, 3, 4, 5$ exactly the Markov blanket is returned, for $i = 6, 7$ the parent 3 is added to the Markov blanket, which is probably a result of having a low value (i.e. 1) for Λ^2. Adding these 7 variables (so that now there is a total of 14 variables), resolving the dual LP and pricing again produces no new variables. So we have an optimal solution to the LP with no acyclicity constraints.

At this point a cutting plane algorithm is run. The particular cutting plane method chosen for this illustrative example limits the number of cutting planes to 20 and 20 are indeed found. Once these 20 cuts are added we have a new LP (LP 2) which must be solved to optimality using pricing. It turns out that 7 pricing rounds are required to solve this LP introducing 4,4,5,4,3,1 and 0 family variables respectively. This interleaved process of pricing and cutting is continued until no new cuts can be found. The process is summarised in Table 1. Eventually we obtain a value of 45301 for the solution to the linear relaxation of IP shown in Fig 1. This is achieved using 52 family variables. Running the cutting plane algorithm again but this time with all $7 \times 2^6 = 448$ family variables included and thus no need for pricing produces exactly the same objective value for the linear relaxation.

4. Branch-price-and-cut for BNSL

Pricing allows us to solve LPs without explicitly introducing all possible variables. But linear relaxations merely provide bounds on IP solutions. Almost always the solution to the linear relaxation of the BNSL IP (Fig 1) will contain non-integer values and thus violate the integrality constraints (4). To make further progress it is necessary to *branch*: pick a variable (typically one that is non-integer

LP	$|C|$	$|\mathcal{V}|$	Rounds	Obj
1	0	14	7,0	20954
2	20	35	4,4,5,4,3,1,0	38006
3	40	50	4,5,3,2,1,0	43158
4	60	52	2,0	45301

Table 1: Cut-and-price for the `gaussian.test` dataset. Learning cast as a minimisation problem. Each of the 4 progressively tighter linear relaxations solved to optimality using pricing.

in the linear relaxation solution) and create two subproblems: one with that variable set to 0 and one with that variable set to 1. (All IP variables will be binary.)

Each subproblem thus created is an IP and it is useful to solve its linear relaxation—and we do this using pricing. So pricing is used *at every node of the branch-and-bound tree* used to solve the IP. Cuts may or may not be added: doing so tightens the linear relaxation thus providing better (local) bounds but also takes time.

Instead of branching on family variables $x_{i \leftarrow J}$, we create new binary *arrow variables* $x_{i \leftarrow j}$ to branch on since this produces a more balanced search tree. $x_{i \leftarrow j} = 1$ iff there is an arrow from parent j to child i. Arrow and family variables satisfy the constraints $x_{i \leftarrow j} = \sum_{J:j \in J} x_{i \leftarrow J}$. In any node of the branch-and-bound tree there will be forbidden ($x_{i \leftarrow j} = 0$) and obligatory ($x_{i \leftarrow j} = 1$) arrows. The pricing algorithm for each node is thus (easily) altered to only look for new family variables which are consistent with the set of forbidden and obligatory arrows at that node.

Suppose we are at a node where $x_{i \leftarrow j} = 1$ and $x_{j \leftarrow k} = 1$ then, due to acyclicity we cannot have $x_{k \leftarrow i} = 1$ so $x_{k \leftarrow i}$ should immediately be fixed to 0. To facilitate such propagations a third kind of binary variable is introduced: *partial order variables* $x_{i \looparrowleft j}$. These variables satisfy the following constraints: $x_{i \leftarrow j} \leq x_{i \looparrowleft j}$, $x_{i \looparrowleft j} + x_{j \looparrowleft i} \leq 1$ and $x_{i \looparrowleft j} + x_{j \looparrowleft k} - x_{i \looparrowleft k} \leq 1$. In this way the partial order variables represent a partial order consistent with the ordering induced by the arrow variables and thus permit the desired propagation.

We now have the required ingredients for solving the BNSL IP: *pricing* to solve large LPs, *cutting* to produce LPs that provide good bounds and *branching* since linear relaxations alone are not enough to solve an IP. However to effectively solve large BNSL IP problems some additional steps are required.

Firstly, we need to ensure that unnecessary lower and upper bounds on variables do not appear as constraints in any LP, since if they do they will have associated dual values which will complicate the pricing algorithm. We will be using the SCIP system (Bestuzheva et al., 2021) since it provides support for pricing, cutting and branching. SCIP allows one to declare lower and upper variable bounds to be *lazy* which prevents the bound appearing in the LP. Here we set the upper bound (i.e. 1) for family variables to be lazy (since they are implied by the constraints (2) in Fig 1), and both the lower and upper bounds of arrow variables to be lazy.

Secondly, we need to represent the constraints between family, arrow and partial order variables appropriately. In any node where $x_{i \leftarrow j} = 1$ we clearly want all family variables where child i lacks j as a parent to be ruled out and similarly when $x_{i \leftarrow j} = 0$ all parent sets for i containing j should go. If we represent these constraints as general linear constraints we will not get these propagations. Since new variables can be created by pricing and then added to existing constraints

$$\text{MINIMISE}_J \qquad z = c_{i \leftarrow J} - \sum_{j \in J} \lambda^*_{i \leftarrow j} - \sum_{\substack{C \in \mathcal{C}, i \in C \\ C \cap J \neq \emptyset}} \lambda^*_C$$

$$\text{SUBJECT TO} \qquad z < \lambda^*_i$$

Figure 5: Pricing problem for BN variable i where $\mathcal{C}$ is the current set of clusters and λ^*_i, $\lambda^*_{i \leftarrow j}$ and λ^*_C are the current dual values for the constraints.

only propagations that remain valid no matter how the constraint is thus altered can be permitted. In our actual implementation the constraint $x_{i \leftarrow j} = \sum_{J : j \in J} x_{i \leftarrow J}$ is replaced by two *set partitioning constraints* $\sum_{J : j \in J} x_{i \leftarrow J} + \neg x_{i \leftarrow j} = 1$ and $\sum_{J : j \notin J} x_{i \leftarrow J} + x_{i \leftarrow j} = 1$. A set partitioning constraint states that exactly one of a set of binary variables has value 1. Since SCIP is explicitly informed that these are set partitioning constraints it knows that only binary variables with coefficient 1 will be added. From this it follows that SCIP can legitimately (and does!) rule out all family variables for child i lacking j as soon as $x_{i \leftarrow j} = 1$ and similarly those containing j as soon as $\neg x_{i \leftarrow j} = 1$ (i.e. $x_{i \leftarrow j} = 0$). Such propagations are valid irrespective of whether additional variables are later added to the set partitioning constraints. For similar reasons the constraint $x_{i \leftarrow j} \leq x_{i \leftsquigarrow j}$ is actually posted as a *set packing constraint* $x_{i \leftarrow j} + \neg x_{i \leftsquigarrow j} \leq 1$.

Thirdly we do not want constraints that only exist to allow propagations to enter the LP. Doing so creates extra dual values for the pricer to deal with and moreover makes the LP unnecessarily bigger (and thus slower to solve). For this reason all constraints involving partial order variables and the constraint $\sum_{J : j \notin J} x_{i \leftarrow J} + x_{i \leftarrow j} = 1$ are not included in the linear relaxation (LP). This is done in SCIP by setting the constraints' *initial* and *separate* flag appropriately when the constraint is added to the problem. Note that this means partial order variables are not in the LP.

Fourthly, even though the constraints $x_{i \leftsquigarrow j} + x_{j \leftsquigarrow i} \leq 1$ and $x_{i \leftsquigarrow j} + x_{j \leftsquigarrow k} - x_{i \leftsquigarrow k} \leq 1$ are not in the LP there are many of them. So instead of adding these constraints we create a single *partial order constraint* containing all partial order variables. This is done by using a slightly altered version of the 'LOP' linear ordering constraint handler (written by March Pfetsch) that is supplied as an example with the SCIP system. The end result is that we get the propagations we want.

Fifthly, for each arrow $i \leftarrow j$, the constraints $\sum_{J : j \in J} x_{i \leftarrow J} + \neg x_{i \leftarrow j} = 1$ are present in the LP and thus have associated dual values which must be taken in account when pricing. Let $\lambda^*_{i \leftarrow j}$ be the dual value for $\sum_{J : j \in J} x_{i \leftarrow J} + \neg x_{i \leftarrow j} = 1$. The pricing problems in Figs 2 and 3 can accommodate arrow dual values by adding $- \sum_{j \in J} \lambda^*_{i \leftarrow j}$ to the objective. Fig 5 is the pricing problem with dual values for the arrow constraints included.

5. Optimisations

Let us refer to our implementation of the above-described pricing approach as PRICEILP. A number of simple optimisations have been added to PRICEILP. The first of these is to simply do less pricing by precomputing the local scores for all potentially optimal parent sets (i.e. family variables) up to some predetermined cardinality limit k. This can speed up later pricing since we can add in

the constraint $\sum_j y_j \geq k + 1$ into (6) to indicate that all necessary parent sets of size up to k have already been found.

A second optimisation is to find, for each child variable i, the highest scoring parent set and to add the corresponding family variables initially to PRICEILP. At least one of these parent sets will be present in any optimal DAG (since each DAG has at least one sink), but one can go further and compute a lower bound (possibly greater than 1) on the number of these 'best parent sets' that must be in any optimal DAG.

In a third optimisation we add constraints that rule out, for each child variable i, any parent set that is a proper superset of 'best parent set' for that child. These constraints can be expressed using arrow indicator variables $x_{i \leftarrow j}$ alone and thus are not involved in the pricing process (which only introduces new family variables). A fourth optimisation adds cutting planes additional to cluster constraints in the form of k-cluster constraints (Cussens, 2011). These tighten the linear relaxations and only slightly complicate the pricing algorithm.

Our last optimisation concerns how pricing and cutting are interleaved. In the standard approach to price-and-cut, pricing is used to solve each of a series of linear programs to optimality. Once each LP is solved a cutting plane algorithm is run to produce a new LP which is a tighter linear relaxation. This is the method used in the example outlined in Section 3.2 and described in detail in Appendix A. It is also how the SCIP solver expects price-and-cut to be implemented. But why spend time solving an LP to optimality if it will soon be replaced by a tighter relaxation due to the addition of cutting planes? Based on this argument PRICEILP allows the user to *delay pricing*. In this approach pricing is only run once the cutting plane algorithm can find no new cutting planes.

6. Does it work?

The preceding sections have outlined an integer programming approach to learning DAGs under causal sufficiency where potential parent sets for individual random variables are introduced in the course of the learning (=optimisation). The issue is whether this an effective approach for this sort of causal discovery.

The short answer is that it is possible to learn optimal DAGs using pricing (together with cutting and branching), but the time it takes to do this depends crucially on how quickly the pricing problem (Fig 5) can be solved. This is because several pricing rounds will be run *for every node in the IP search tree* to ensure the linear relaxation for that node is solved to optimality. Consider, for example, the Gaussian ℓ_0 pricing problem (shown, with dual values for 'arrow constraints' omitted, in Fig 3). One strategy for solving this problem is to encode it in such a way that an existing solver can solve it. This is the approach we take: the encoding as a non-convex constrained optimisation problem is given in Fig 6, the solver we use is SCIP. In Fig 6 each binary variable y_j indicates that j is in the new parent set for i. The β_j are linear regression coefficients, X_i is the vector of observed values for random variable i in the data, X_{-i} is the data matrix of all variables other than i and S is the covariance matrix $X_{-i}^\top X_{-i}$. The variable x_{σ^2} represents squared error (when predicting random variable i in a linear regression model with coefficients β_j) and $x_{\log \sigma^2}$ is the log of that squared error. The 'SOS-1' constraints state that $y_j = 0 \rightarrow \beta_j = 0$. Intermediate variables γ_j have been introduced so that the constraint between linear regression coefficients and squared error can be expressed as the *second-order cone* constraint (8). This has been shown (empirically) to lead to faster solving than expressing this relationship directly.

$$\text{MINIMISE} \qquad z = nx_{\log \sigma^2} + \sum_{j \in P \setminus \{i\}} (\Lambda^2 - \lambda^*_{i \leftarrow j}) y_j - \sum_{C \in \mathcal{C}} \lambda^*_C y_C \tag{7}$$

$$\text{SUBJECT TO} \qquad \sum_{j \in P \setminus \{i\}} \gamma_j^2 + c \leq nx_{\sigma^2} \tag{8}$$

$$x_{\log \sigma^2} = \log x_{\sigma^2} \tag{9}$$

$$\gamma = S^{1/2} \boldsymbol{\beta} - S^{-1/2} \boldsymbol{X}^\top_{-i} \boldsymbol{X}_i \tag{10}$$

$$(\beta_j, 1 - y_j) : \text{SOS-1} \qquad\qquad j \in P \setminus \{i\} \tag{11}$$

$$y_C = \bigvee_{j \in C} y_j \qquad\qquad C \in \mathcal{C} \tag{12}$$

$$z < \lambda^*_i$$

$$x_{\sigma^2} \in \mathbb{R}_+ \ \ x_{\log \sigma^2}, \gamma_j, \beta_j \in \mathbb{R} \ \ y_j, y_C \in \{0, 1\}$$

Figure 6: Gaussian ℓ_0 pricing for BN variable i as nonlinear constrained optimisation.

k	NVars	Solving Time	Pricing time
1	321	478	475
2	316	463	459
3	283	151	150
4	282	80	77
5	-	-	-
6	276	2	0

Table 2: Solving times for PRICEILP on the small `gaussian.test` dataset using BIC ℓ_0 penalised log squared error. NVars indicates the number of IP variables in the problem at the point it is solved.

Table 2 shows solving times (in seconds) for different values of k (upper bound on initial parent set size) using PRICEILP on the `gaussian.test` dataset (see Section 3.2). The first thing to note that this is a very easy BNSL problem: it can be solved to optimality in 2 seconds if k is set to 6 (recall there are 7 random variables) so that no pricing is ever done. The optimal DAG when BIC is used to choose the level of ℓ_0 regularisation is the true data-generating DAG shown in Fig 4. The second point is that pricing is extremely slow, taking up almost all of the solving time for all values of $k < 6$. The third point is that PRICEILP is pricing in too many IP variables; the fewer IP variables we start with the more we eventually end up with! (For $k = 5$, SCIP's LP solver encounters numerical problems and aborts because, for unknown reasons, an infeasible LP occurs.)

The results in Table 2 show that pricing is a bad idea when using BIC scoring for a small dataset like `gaussian.test`. However, if we alter the score, results are more encouraging. The problem with the pricing problem in Fig 6 is the log in the objective. In linear regression, squared error is a convex quadratic function of the linear regression coefficients, but its log is merely a pseudoconvex

k	NVars	Solving Time	Pricing time
1	151	12.7	12.6
2	194	11.8	11.7
3	243	9.2	9.1
4	261	3.1	3.0
5	262	0.1	0
6	262	0.1	0

Table 3: Solving times for PRICEILP on the small `gaussian.test` dataset using BIC ℓ_0 penalised squared error. NVars indicates the number of IP variables in the problem at the point it is solved.

k	NVars	Solving Time	Pricing time
3	≥ 3022	> 2457	> 2457
6	19386	2206	2192

Table 4: Solving times for PRICEILP on `notears20.dat` dataset using BIC ℓ_0 penalised squared error. Pricing **not** delayed. NVars indicates the number of IP variables in the problem at the point it is solved.

function. If we use ℓ_0 penalised squared error (rather than its log) with ℓ_0 again chosen using BIC as the objective then we get the results in Table 3 (time still in seconds).

The problem is even easier than before, and although pricing still takes up most of the time, at least it is done more quickly. Note also that the more we rely on pricing to introduce IP variables the fewer we end up with, so the 'acyclicity penalty' that is there when pricing is helping us avoid creating unnecessary family variables. The downside is that the BIC choice of ℓ_0 penalty turns out to be too small: the optimal DAG produced is not the true one (Fig 4), since we get one extra incorrect edge from vertex 1 to vertex 2.

With this easier objective, larger problems can be addressed. Tables 4 and 5 show results for some experiments on a dataset `notears20.dat` with 20 random variables and 100 datapoints generated using the NOTEARS (Zheng et al., 2018) software from a known true Gaussian DAG. The latter Table uses delayed pricing, the former does not. Note that the run for $k = 3$ in Table 4 was aborted after 2457 seconds. Note also that in this table, as in those above, 'Solving Time' does not include time taken to construct the IP problem instance. For example, in the $k = 6$ run in Table 4 an additional 51 seconds was needed to compute local scores for initial family variables.

It is clear that for this DAG learning problem, delayed pricing brings big improvements. However, this is still only a 20 variable example and also with BIC scoring the ℓ_0 penalty is too small. The optimal DAG has 18 edges which are not present in the true data-generating DAG, although all 9 true edges are recovered. In contrast NOTEARS under default parameters has a stronger complexity penalty and is able to recover the true DAG exactly. Unfortunately, increasing the ℓ_0 penalty (which should allow faster and more accurate learning) led to aborted runs due to numerical problems during LP solving.

k	NVars	Solving Time	Pricing time
3	3035	1326	1324
6	19386	351	335

Table 5: Solving times for PRICEILP on `notears20.dat` dataset using BIC ℓ_0 penalised squared error and delayed pricing. NVars indicates the number of IP variables in the problem at the point it is solved.

7. Discussion

We have a working implementation—PRICEILP—that extends the IP approach to DAG learning to allow new candidate parent sets to be created during the solving/learning process via *pricing*.[1] This is done in a principled way by considering *reduced local scores* for these candidate parent sets which take into account the dual values associated with constraints ensuring acyclicity. We have shown, unsurprisingly, that fast pricing is required for fast learning and shown that, perhaps less obviously, delaying pricing can bring benefits.

Although we view this contribution as a useful first step in extending score-based causal discovery, it is clear that there is some way to go before this approach is a state-of-the-art method. Some of the necessary work concerns the details of implementation (tracking down the cause of LPs with numerical problems) and some is more algorithmic: designing fast pricing algorithms for a range of popular scores for causal model learning.

In this paper we have focused exclusively on DAG learning with the assumption of causal sufficiency. However, it is actually score-based learning of more complex causal models, e.g. those with latent variables where pricing can be of most benefit. For example, Chen et al. (2021) showed how to encode BIC score-based learning of ancestral acyclic directed mixed graphs (ADMGs) as an IP problem. But ADMG learning requires even more IP variables than DAG learning: one for every candidate district. It would be interesting to investigate the feasibility of pricing in candidate districts while learning the ADMG.

---

1. This software will be made available once anonymity is no longer a concern.

References

Ksenia Bestuzheva, Mathieu Besançon, Wei-Kun Chen, Antonia Chmiela, Tim Donkiewicz, Jasper van Doornmalen, Leon Eifler, Oliver Gaul, Gerald Gamrath, Ambros Gleixner, Leona Gottwald, Christoph Graczyk, Katrin Halbig, Alexander Hoen, Christopher Hojny, Rolf van der Hulst, Thorsten Koch, Marco Lübbecke, Stephen J. Maher, Frederic Matter, Erik Mühmer, Benjamin Müller, Marc E. Pfetsch, Daniel Rehfeldt, Steffan Schlein, Franziska Schlösser, Felipe Serrano, Yuji Shinano, Boro Sofranac, Mark Turner, Stefan Vigerske, Fabian Wegscheider, Philipp Wellner, Dieter Weninger, and Jakob Witzig. The SCIP Optimization Suite 8.0. Technical report, Optimization Online, December 2021. URL http://www.optimization-online.org/DB_HTML/2021/12/8728.html.

Rui Chen, Sanjeeb Dash, and Tian Gao. Integer programming for causal structure learning in the presence of latent variables. In Marina Meila and Tong Zhang, editors, *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pages 1550–1560. PMLR, 18–24 Jul 2021. URL https://proceedings.mlr.press/v139/chen21c.html.

David M. Chickering, David Heckerman, and Christopher Meek. Large-sample learning of Bayesian networks is NP-hard. *Journal of Machine Learning Research*, 20:1287–1330, October 2004.

James Cussens. Maximum likelihood pedigree reconstruction using integer programming. In *Proceedings of the Workshop on Constraint Based Methods for Bioinformatics (WCB-10)*, Edinburgh, July 2010.

James Cussens. Bayesian network learning with cutting planes. In Fabio G. Cozman and Avi Pfeffer, editors, *Proceedings of the 27th Conference on Uncertainty in Artificial Intelligence (UAI 2011)*, pages 153–160, Barcelona, 2011. AUAI Press.

James Cussens, David Haws, and Milan Studený. Polyhedral aspects of score equivalence in Bayesian network structure learning. *Mathematical Programming*, 164:285–324, 2017a. doi:10.1007/s10107-016-1087-2.

James Cussens, Matti Järvisalo, Janne H. Korhonen, and Mark Bartlett. Bayesian network structure learning with integer programming: Polytopes, facets, and complexity. *Journal of Artificial Intelligence Research*, 58:185–229, 2017b.

Tommi Jaakkola, David Sontag, Amir Globerson, and Marina Meila. Learning Bayesian network structure using LP relaxations. In *Proceedings of 13th International Conference on Artificial Intelligence and Statistics (AISTATS 2010)*, volume 9, pages 358–365, 2010. Journal of Machine Learning Research Workshop and Conference Proceedings.

Marco Scutari. Learning Bayesian networks with the bnlearn R package. *Journal of Statistical Software*, 35(3):1–22, 2010. doi: 10.18637/jss.v035.i03.

Sara van de Geer and Peter Bühlmann. ℓ_0-penalized maximum likelihood for sparse directed acyclic graphs. *The Annals of Statistics*, 41(2):536–567, 2013.

Xun Zheng, Bryon Aragam, Pradeep K Ravikumar, and Eric P Xing. Dags with no tears: Continuous optimization for structure learning. In S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 31. Curran Associates, Inc., 2018. URL https://proceedings.neurips.cc/paper/2018/file/e347c51419ffb23ca3fd5050202f9c3d-Paper.pdf.

Appendix A. Pricing for BNSL: a worked example in detail

This section contains full details of the Gaussian ℓ_0-penalised maximum likelihood pricing example summarised in Section 3.2. The objectives here are given in terms of *scores* rather than *costs* and are not rounded to the nearest integer. So, for example, the objective for the first LP (which has no cluster constraints) is shown here to have an optimal solution with objective value of -20954.37 rather than 20954 as given in Section 3.2. Similarly, the two values given for a priced-in family variable are its local score and its reduced local score. Note that for priced-in variables the reduced local score is always positive (i.e. the reduced cost is always negative).

$\mathcal{V}_1 = x_{1\leftarrow\{\}}, x_{2\leftarrow\{\}}, x_{C\leftarrow\{\}}, x_{4\leftarrow\{\}}, x_{5\leftarrow\{\}}, x_{6\leftarrow\{\}}, x_{7\leftarrow\{\}}$

LP. $C = \emptyset$. $\mathcal{V} = \mathcal{V}_1$. Solution is $x_{1\leftarrow\{\}} = 1$, $x_{2\leftarrow\{\}} = 1$, $x_{3\leftarrow\{\}} = 1$, $x_{4\leftarrow\{\}} = 1$, $x_{5\leftarrow\{\}} = 1$, $x_{6\leftarrow\{\}} = 1$, $x_{7\leftarrow\{\}} = 1$. Objective is -88043.06. Dual values: $\lambda_i^* = -7115.31, -12643.78, -16343.24, -14678.79, -10534.51, -16208.60, -10518.84$.

Pricing. New variables: $\mathcal{V}_2 = (x_{1\leftarrow\{2,3,4,5,6,7\}}, 466.13, 7581.44)$, $(x_{2\leftarrow\{1,3,4\}}, 1921.01, 14564.78)$, $(x_{3\leftarrow\{1,2\}}, -3718.43, 12624.79)$, $(x_{4\leftarrow\{1,2,5,6,7\}}, -1238.48, 13440.29)$, $(x_{5\leftarrow\{1,4,6,7\}}, -6507.84, 4026.65)$, $(x_{6\leftarrow\{1,3,4,5,7\}}, -7076.24, 9132.33)$, $(x_{7\leftarrow\{1,3,4,5,6\}}, -4800.51, 5718.30)$.

LP. $C = \emptyset$. $\mathcal{V} = \dot{\bigcup}_{i=1}^{2}\mathcal{V}_i$. Solution is $x_{1\leftarrow\{2,3,5,6,4,7\}} = 1, x_{2\leftarrow\{1,3,4\}} = 1, x_{3\leftarrow\{1,2\}} = 1, x_{4\leftarrow\{1,2,5,6,7\}} = 1, x_{5\leftarrow\{1,4,6,7\}} = 1, x_{6\leftarrow\{1,3,4,5,7\}} = 1, x_{7\leftarrow\{1,3,4,5,6\}} = 1$. Objective is -20954.37. Dual values: $\lambda_i^* = 466.13, 1921.01, -3718.43, -1238.48, -6507.84, -7076.24, -4800.51$.

Pricing. New variables: None

Separation. New clusters: $C_1 = \{1,2\}, \{2,4\}, \{4,5\}, \{1,3\}, \{1,4\}, \{5,6\}, \{1,5\}, \{2,3\}, \{1,2,3\}, \{1,2,4\}, \{1,3,4\}, \{2,3,4\}, \{1,2,3,4\}, \{1,3,5\}, \{1,4,5\}, \{1,5,6\}, \{2,3,4,5\}, \{1,2,4,5\}, \{2,4,7\}, \{1,3,4,5\}$.

LP. $C = C_1$. $\mathcal{V} = \dot{\bigcup}_{i=1}^{2}\mathcal{V}_i$. Solution is $x_{1\leftarrow\emptyset} = 1, x_{2\leftarrow\emptyset} = 1, x_{3\leftarrow\{1,2\}} = 1, x_{4\leftarrow\{1,2,5,6,7\}} = 1, x_{5\leftarrow\emptyset} = 1, x_{6\leftarrow\{1,3,4,5,7\}} = 1, x_{7\leftarrow\{1,3,4,5,6\}} = 1$. Objective is -47127.29. Dual values: $\lambda_i^* = -7115.31, -12643.78, -12424.37, -14678.78, -10534.51, -11102.91, -4800.51$. $\lambda_{\{2,4\}}^* = 13440.30$, $\lambda_{\{1,3\}}^* = 7581.45$, $\lambda_{\{5,6\}}^* = 4026.66$, $\lambda_{\{2,3\}}^* = 1124.49$.

Pricing. New variables: $\mathcal{V}_3 = (x_{1\leftarrow\{4,5,6,7\}}, -3071.68, 4043.62)$, $(x_{2\leftarrow\{1,3,5,6,7\}}, 97.25, 11616.53)$, $(x_{3\leftarrow\{4,5,6,7\}}, -7713.50, 4710.86)$, $(x_{4\leftarrow\{1,3,5,6,7\}}, -3060.54, 11618.23)$.

LP. $C = C_1$. $\mathcal{V} = \dot{\bigcup}_{i=1}^{3}\mathcal{V}_i$. Solution is $x_{1\leftarrow\emptyset} = 1/2, x_{2\leftarrow\emptyset} = 1/2, x_{4\leftarrow\emptyset} = 1/2, x_{5\leftarrow\emptyset} = 1, x_{1\leftarrow\{2,3,4,5,6,7\}} = 1/2, x_{2\leftarrow\{1,3,4\}} = 1/2 \; x_{3\leftarrow\{1,2\}} = 1/2, x_{4\leftarrow\{1,2,5,6,7\}} = 1/2, x_{6\leftarrow\{1,3,4,5,7\}} = 1, x_{7\leftarrow\{1,3,4,5,6\}} = 1, x_{3\leftarrow\{4,5,6,7\}} = 1/2$. Objective is -44771.85. Dual values: $\lambda_i^* = -7115.31, -12643.78, -14105.73, -14678.78, -10534.51, -11102.91, -4800.51, \lambda_{\{1,2\}}^* = 2355.43, \lambda_{\{2,4\}}^* = 1822.05, \lambda_{\{1,4\}}^* = 5226.01, \lambda_{\{5,6\}}^* = 4026.66, \lambda_{\{2,3\}}^* = 3995.07, \lambda_{\{2,3,4\}}^* = 6392.23$.

Pricing. New variables: $\mathcal{V}_4 = (x_{1\leftarrow\{2,3,5,6,7\}}, 418.10, 5177.97)$, $(x_{2\leftarrow\{1,5,6,7\}}, -5192.93, 5095.39)$, $(x_{3\leftarrow\{1,5,6,7\}}, -9010.62, 5095.10)$, $(x_{4\leftarrow\{2,3,5,6,7\}}, -1287.31, 5177.17)$.

LP. $C = C_1$. $\mathcal{V} = \dot{\bigcup}_{i=1}^{4}\mathcal{V}_i$. Solution is $x_{1\leftarrow\emptyset} = 1/2, x_{2\leftarrow\emptyset} = 1/2, x_{4\leftarrow\emptyset} = 1/2, x_{5\leftarrow\emptyset} = 1, x_{1\leftarrow\{2,3,4,5,6,7\}} = 1/2, x_{2\leftarrow\{1,3,4\}} = 1/2 \; x_{3\leftarrow\{1,2\}} = 1/2, x_{4\leftarrow\{1,2,5,6,7\}} = 1/2, x_{6\leftarrow\{1,3,4,5,7\}} = 1$,

$x_{7\leftarrow\{1,3,4,5,6\}} = 1$, $x_{3\leftarrow\{4,5,6,7\}} = 1/2$. Objective is -44771.85. Dual values: $\lambda_i^* = -7115.31$, -12643.784, -14188.30, -14678.78, -10534.51, -11102.91, -9895.92, $\lambda_{\{1,2\}}^* = 2355.43$, $\lambda_{\{2,4\}}^* = 1822.05$ $\lambda_{\{1,4\}}^* = 48.03$, $\lambda_{\{5,6\}}^* = 4026.66$, $\lambda_{\{2,3\}}^* = 3995.07$, $\lambda_{\{1,3,4\}}^* = 5177.98$, $\lambda_{\{2,3,4\}}^* = 1296.82$, $\lambda_{\{2,4,7\}}^* = 5095.40$.

Pricing. New variables: $\mathcal{V}_5 = (x_{1\leftarrow\{2,5,6,7\}}, -3304.52, 1455.33)$, $(x_{2\leftarrow\{4\}}, 503.85, 4933.34)$, $(x_{3\leftarrow\{5,6,7\}}, -9273.27, 4915.02)$, $(x_{4\leftarrow\{2,5,6,7\}}, -1471.23, 4993.26)$, $(x_{7\leftarrow\{1,3,5,6\}}, -5361.48, 4534.43)$.

LP. $C = C_1$. $\mathcal{V} = \dot{\bigcup}_{i=1}^5 \mathcal{V}_i$. Solution is $x_{1\leftarrow\emptyset} = 1/2$, $x_{2\leftarrow\emptyset} = 1/2$, $x_{5\leftarrow\emptyset} = 1$, $x_{2\leftarrow\{1,3,4\}} = 1/2$, $x_{3\leftarrow\{1,2\}} = 1/2$, $x_{4\leftarrow\{1,2,5,6,7\}} = 1/2$, $x_{6\leftarrow\{1,3,4,5,7\}} = 1$, $x_{7\leftarrow\{1,3,4,5,6\}} = 1/2$, $x_{4\leftarrow\{1,3,5,6,7\}} = 1/2$, $x_{1\leftarrow\{2,3,5,6,7\}} = 1/2$, $x_{3\leftarrow\{5,6,7\}} = 1/2$, $x_{7\leftarrow\{1,3,5,6\}} = 1/2$. Objective value is -40047.12. Dual values: $\lambda_i^* = -7115.31$, -12643.78, -9273.27, -14678.78, -10534.51, -10858.29, -5361.48, $\lambda_{\{1,2\}}^* = 1417.15$, $\lambda_{\{2,4\}}^* = 1822.05$, $\lambda_{\{4,5\}}^* = 244.62$, $\lambda_{\{1,3\}}^* = 453.24$, $\lambda_{\{1,4\}}^* = 48.03$, $\lambda_{\{5,6\}}^* = 3782.04$, $\lambda_{\{1,2,4\}}^* = 5663.02$, $\lambda_{\{2,3,4\}}^* = 5101.60$, $\lambda_{\{2,4,7\}}^* = 560.96$.

Pricing. New variables: $\mathcal{V}_6 = (x_{1\leftarrow\{3,4,5,6,7\}}, -495.90, 455.10)$, $(x_{2\leftarrow\{5,6,7\}}, -8632.31, 3450.49)$, $(x_{4\leftarrow\{5,6,7\}}, -10605.66, 3267.53)$, $(x_{5\leftarrow\{1,2,6,7\}}, -6729.11, 23.33)$.

LP. $C = C_1$. $\mathcal{V} = \dot{\bigcup}_{i=1}^6 \mathcal{V}_i$. Solution is $x_{1\leftarrow\emptyset} = 1/2$, $x_{1\leftarrow 2,3,5,6,7} = 1/2$, $x_{2\leftarrow 5,6,7} = 1/2$, $x_{2\leftarrow 1,3,4} = 1/2$, $x_{3\leftarrow 1,2} = 1/2$, $x_{3\leftarrow 5,6,7} = 1/2$, $x_{4\leftarrow 1,2,5,6,7} = 1/2$, $x_{4\leftarrow 1,3,5,6,7} = 1/2$, $x_{5\leftarrow\emptyset} = 1$, $x_{6\leftarrow 1,3,4,5,7} = 1$, $x_{7\leftarrow 1,3,5,6} = 1$. Objective is -38321.86. Dual values: $\lambda_i^* = -7115.31$, -9193.27, -9273.27, -11204.93, -10534.51, -10881.64, -5361.48, $\lambda_{\{1,2\}}^* = 962.04$, $\lambda_{\{2,4\}}^* = 1822.05$, $\lambda_{\{4,5\}}^* = 221.27$, $\lambda_{\{1,3\}}^* = 2178.49$, $\lambda_{\{1,4\}}^* = 48.03$, $\lambda_{\{5,6\}}^* = 3805.39$, $\lambda_{\{2,3\}}^* = 455.11$, $\lambda_{\{1,2,4\}}^* = 4392.88$, $\lambda_{\{2,3,4\}}^* = 2921.23$, $\lambda_{\{2,4,7\}}^* = 560.96$.

Pricing. New variables: $\mathcal{V}_7 = (x_{1\leftarrow\{5,6,7\}}, -6743.90, 371.39)$, $(x_{2\leftarrow\{1,3,5,6\}}, -202.07, 259.92)$, $(x_{4\leftarrow\{1,3,6\}}, -3634.97, 207.79)$.

LP. $C = C_1$. $\mathcal{V} = \dot{\bigcup}_{i=1}^7 \mathcal{V}_i$. Solution is $x_{1\leftarrow\{2,3,5,6,7\}} = 1/2$, $x_{1\leftarrow\{5,6,7\}} = 1/2$, $x_{2\leftarrow\{5,6,7\}} = 1/2$, $x_{2\leftarrow\{1,3,5,6\}} = 1/2$. $x_{3\leftarrow\{1,2\}} = 1/2$, $x_{3\leftarrow\{5,6,7\}} = 1/2$, $x_{4\leftarrow\{1,2,5,6,7\}} = 1$, $x_{5\leftarrow\emptyset} = 1$, $x_{6\leftarrow\{1,3,4,5,7\}} = 1$, $x_{7\leftarrow\{1,3,4,5,6\}} = 1/2$, $x_{7\leftarrow\{1,3,5,6\}} = 1/2$, Objective is -38006.19. Dual values: $\lambda_i^* = -6743.90$, -9193.27, -9273.27, -11204.93, -10534.51, -10881.64, -5361.48, $\lambda_{\{1,2\}}^* = 962.04$, $\lambda_{\{2,4\}}^* = 1614.25$, $\lambda_{\{4,5\}}^* = 221.27$, $\lambda_{\{1,3\}}^* = 1862.82$, $\lambda_{\{1,4\}}^* = 48.03$, $\lambda_{\{5,6\}}^* = 3805.39$, $\lambda_{\{2,3\}}^* = 507.23$, $\lambda_{\{1,2,4\}}^* = 4337.14$, $\lambda_{\{2,3,4\}}^* = 3184.78$, $\lambda_{\{2,4,7\}}^* = 560.96$.

Pricing. New variables: $\mathcal{V}_8 = (x_{4\leftarrow\{2,3,6,7\}}, -1460.31, 47.46)$,

LP. $C = C_1$. $\mathcal{V} = \dot{\bigcup}_{i=1}^8 \mathcal{V}_i$. Solution is $x_{1\leftarrow\{5,6,7\}} = 1/2$, $x_{1\leftarrow\{2,3,5,6,7\}} = 1/2$, $x_{2\leftarrow\{5,6,7\}} = 1/2$, $x_{2\leftarrow\{1,3,5,6\}} = 1/2$. $x_{3\leftarrow\{1,2\}} = 1/2$, $x_{3\leftarrow\{5,6,7\}} = 1/2$, $x_{4\leftarrow\{1,2,5,6,7\}} = 1$, $x_{5\leftarrow\emptyset} = 1$, $x_{6\leftarrow\{1,3,4,5,7\}} = 1$, $x_{7\leftarrow\{1,3,4,5,6\}} = 1/2$, $x_{7\leftarrow\{1,3,5,6\}} = 1/2$, Objective is -38006.19 Dual values $\lambda_i^* = -6743.90$, -9193.27, -9273.27, -11157.45, -10534.51, -10929.12, -5361.48, $\lambda_{\{1,2\}}^* = 962.04$, $\lambda_{\{2,4\}}^* = 1661.73$, $\lambda_{\{4,5,\}}^* = 173.79$, $\lambda_{\{1,3\}}^* = 1862.8$, $\lambda_{\{1,4\}}^* = 48.03$, $\lambda_{\{5,6\}}^* = 3852.87$, $\lambda_{\{2,3\}}^* = 554.71$, $\lambda_{\{1,2,4\}}^* = 4337.14$, $\lambda_{\{2,3,4\}}^* = 3137.30$, $\lambda_{\{2,4,7\}}^* = 560.96$.

Pricing. New variables: None.

Separation. New clusters: $C_2 = \{1, 7\}$, $\{2, 3, 7\}$, $\{6, 7\}$, $\{1, 2, 7\}$, $\{2, 6, 7\}$, $\{1, 3, 7\}$, $\{1, 2, 3, 7\}$, $\{4, 6, 7\}$, $\{1, 4, 7\}$, $\{2, 4, 6, 7\}$, $\{1, 2, 4, 7\}$, $\{1, 3, 4, 7\}$, $\{2, 3, 4, 7\}$, $\{1, 2, 3, 4, 7\}$, $\{1, 4, 6, 7\}$, $\{4, 6\}$, $\{1, 3, 4, 6, 7\}$, $\{1, 4, 6\}$, $\{1, 2, 4, 6, 7\}$, $\{2, 4, 6\}$.

LP. $C = \dot{\bigcup}_{i=1}^2 C_i$. $\mathcal{V} = \dot{\bigcup}_{i=1}^8 \mathcal{V}_i$. Solution is $x_{1\leftarrow\emptyset} = 1/2$, $x_{1\leftarrow\{2,3,4,5,6,7\}} = 1/2$, $x_{2\leftarrow\{1,3,4\}} = 1/2$, $x_{2\leftarrow\{4\}} = 1/2$, $x_{3\leftarrow\{1,2\}} = 1/2$ $x_{3\leftarrow\{5,6,7\}} = 1/2$, $x_{4\leftarrow\emptyset} = 1/2$, $x_{4\leftarrow\{1,3,5,6,7\}} = 1/2$ $x_{5\leftarrow\emptyset} = 1/2$, $x_{5\leftarrow\{1,4,6,7\}} = 1/2$ $x_{6\leftarrow\emptyset} = 1/2$, $x_{6\leftarrow\{1,3,4,5,7\}} = 1/2$ $x_{7\leftarrow\emptyset} = 1/2$, $x_{7\leftarrow\{1,3,4,5,6\}} = 1/2$. Objective value -45300.94 Dual values: $\lambda_i^* = -7115.31$, -10683.44, -11023.37, -14678.78, -10534.51, -16208.59, -10518.83, $\lambda_{\{1,2\}}^* = 962.04$, $\lambda_{\{4,2\}}^* = 1822.05$, $\lambda_{\{5,4\}}^* = 221.27$, $\lambda_{\{1,3\}}^* = 2016.53$, $\lambda_{\{4,1\}}^* =$

0.55, $\lambda^*_{\{5,6\}} = 3805.39$, $\lambda^*_{\{2,3\}} = 455.11$, $\lambda^*_{\{4,1,2\}} = 4230.91$, $\lambda^*_{\{4,2,3\}} = 3083.20$, $\lambda^*_{\{4,7,2\}} = 301.03$, $\lambda^*_{\{1,7\}} = 371.40$, $\lambda^*_{\{6,7\}} = 3295.78$, $\lambda^*_{\{4,7,2,3\}} = 1750.09$, $\lambda^*_{\{4,6\}} = 2031.167$.

Pricing. New variables: $\mathcal{V}_9 = (x_{1\leftarrow\{5,6\}}, -6845.99, 269.30)$, $(x_{2\leftarrow\{5,6\}}, -10383.55, 299.87)$, $(x_{4\leftarrow\{2\}}, -1531.14, 1960.32)$, $(x_{6\leftarrow\{1,2\}}, -13663.57, 2545.00)$.

LP. $C = \dot{\bigcup}_{i=1}^{2} C_i$. $\mathcal{V} = \dot{\bigcup}_{i=1}^{9} \mathcal{V}_i$. Solution is $x_{1\leftarrow\emptyset} = 1/2$, $x_{1\leftarrow\{2,3,4,5,6,7\}} = 1/2$, $x_{2\leftarrow\emptyset} = 1/2$, $x_{2\leftarrow\{1,3,4\}} = 1/2$, $x_{3\leftarrow\{1,2\}} = 1/2$, $x_{3\leftarrow\{5,6,7\}} = 1/2$, $x_{4\leftarrow\{1,3,5,6,7\}} = 1/2$, $x_{4\leftarrow\{2\}} = 1/2$, $x_{5\leftarrow\emptyset} = 1/2$, $x_{5\leftarrow\{1,4,6,7\}} = 1/2$, $x_{6\leftarrow\emptyset} = 1/2$, $x_{6\leftarrow\{1,3,4,5,7\}} = 1/2$, $x_{7\leftarrow\{1,3,4,5,6\}} = 1/2$, $x_{7\leftarrow\emptyset} = 1/2$. Objective value is -45300.94. Dual values: $\lambda^*_i = -7115.31, -12643.78, -10438.69, -14678.78, -10534.51, -16208.59, -10518.83$, $\lambda^*_{\{1,2\}} = 962.04$, $\lambda^*_{\{4,2\}} = 1822.05$, $\lambda^*_{\{4,5,\}} = 221.27$, $\lambda^*_{\{1,3\}} = 1431.85$, $\lambda^*_{\{1,4\}} = 0.55$, $\lambda^*_{\{5,6\}} = 3536.07$, $\lambda^*_{\{1,5\}} = 269.31$, $\lambda^*_{\{2,3\}} = 455.11$, $\lambda^*_{\{1,2,4\}} = 3646.24$, $\lambda^*_{\{2,3,4\}} = 3667.87$, $\lambda^*_{\{2,4,7\}} = 301.03$, $\lambda^*_{\{1,7\}} = 1271.44$, $\lambda^*_{\{6,7\}} = 2980.42$, $\lambda^*_{\{2,3,4,7\}} = 1165.42$, $\lambda^*_{\{4,6\}} = 70.82$, $\lambda^*_{\{2,4,6\}} = 2545.01$.

Pricing. New variables: $\mathcal{V}_{10} = (x_{1\leftarrow\{6,2,3\}}, -61.03, 1014.12)$, $(x_{2\leftarrow\{1,3\}}, -237.38, 2509.69)$, $(x_{4\leftarrow\{1,3\}}, -3690.99, 2507.68)$, $(x_{6\leftarrow\{1,3\}}, -13691.81, 2516.76)$, $x_{7\leftarrow\{2,3,5,6,4\}}, -5316.08, 755.86)$.

LP. $C = \dot{\bigcup}_{i=1}^{2} C_i$. $\mathcal{V} = \dot{\bigcup}_{i=1}^{10} \mathcal{V}_i$. Solution is $x_{1\leftarrow\emptyset} = 1/4$, $x_{1\leftarrow\{2,3,4,5,6,7\}} = 1/2$, $x_{1\leftarrow\{5,6,7\}} = 1/4$, $x_{2\leftarrow\emptyset} = 1/4$, $x_{2\leftarrow\{1,3\}} = 1/2$, $x_{2\leftarrow\{5,6,7\}} = 1/4$, $x_{3\leftarrow\{1,2\}} = 1/2$, $x_{3\leftarrow\{5,6,7\}} = 1/2$, $x_{4\leftarrow\{1,2,5,6,7\}} = 1/4$, $x_{4\leftarrow\{2\}} = 3/4$, $x_{5\leftarrow\emptyset} = 3/4$, $x_{5\leftarrow\{1,4,6,7\}} = 1/4$, $x_{6\leftarrow\{1,3,4,5,7\}} = 3/4$, $x_{6\leftarrow\{1,3\}} = 1/4$ $x_{7\leftarrow\emptyset} = 3/4$, $x_{7\leftarrow\{1,3,4,5,6\}} = 1/4$, Objective is -43970.52. Dual values: $\lambda^*_i = -7115.31, -12643.78, -12804.34, -14678.78, -10534.51, -13961.12, -10518.83$, $\lambda^*_{\{1,2\}} = 962.04$, $\lambda^*_{\{4,2\}} = 1822.05$, $\lambda^*_{\{1,3\}} = 1944.77$, $\lambda^*_{\{5,6\}} = 2246.83$, $\lambda^*_{\{2,3\}} = 599.17$, $\lambda^*_{\{4,1,2\}} = 4303.22$, $\lambda^*_{\{4,2,3\}} = 3010.89$, $\lambda^*_{\{5,4,2,3\}} = 1779.82$, $\lambda^*_{\{1,7\}} = 102.09$, $\lambda^*_{\{6,7\}} = 3864.98$, $\lambda^*_{\{4,7,2,3\}} = 1751.24$, $\lambda^*_{\{4,6\}} = 23.34$, $\lambda^*_{\{4,1,6\}} = 269.31$, $\lambda^*_{\{4,6,2\}} = 480.39$.

Pricing. New variables: $\mathcal{V}_{11} = (x_{2\leftarrow\{6\}}, -10741.98, 1421.39)$, $(x_{4\leftarrow\{6\}}, -12755.05, 1150.66)$, $(x_{6\leftarrow\{3\}}, -13734.11, 227.00)$.

LP. $C = \dot{\bigcup}_{i=1}^{2} C_i$. $\mathcal{V} = \dot{\bigcup}_{i=1}^{11} \mathcal{V}_i$. Solution is $x_{1\leftarrow\emptyset} = 1/2$, $x_{5\leftarrow\emptyset} = 1/2$, $x_{7\leftarrow\emptyset} = 1/2$, $x_{1\leftarrow\{2,3,4,5,6,7\}} = 1/2$, $x_{3\leftarrow\{1,2\}} = 1/2$, $x_{4\leftarrow\{1,2,5,6,7\}} = 1/2$, $x_{5\leftarrow\{1,4,6,7\}} = 1/2$, $x_{6\leftarrow\{1,3,4,5,7\}} = 1/2$, $x_{7\leftarrow\{1,3,4,5,6\}} = 1/2$, $x_{3\leftarrow\{5,6,7\}} = 1/2$, $x_{4\leftarrow\{2\}} = 1/2$, $x_{2\leftarrow\{1,3\}} = 1/2$, $x_{6\leftarrow\{1,3\}} = 1/2$, $x_{2\leftarrow\{6\}} = 1/2$, Objective is -43259.82. $\lambda^*_i = -7115.31, -11078.32, -12030.71, -13113.32, -10534.51, -13734.11, -10518.83$, $\lambda^*_{\{1,2\}} = 962.04$, $\lambda^*_{\{4,2\}} = 1822.05$, $\lambda^*_{\{5,4\}} = 221.27$, $\lambda^*_{\{1,3\}} = 2020.17$, $\lambda^*_{\{4,1\}} = 0.55$, $\lambda^*_{\{5,6\}} = 3219.95$, $\lambda^*_{\{2,3\}} = 455.11$, $\lambda^*_{\{4,1,2\}} = 3813.80$, $\lambda^*_{\{4,2,3\}} = 3079.55$, $\lambda^*_{\{5,1,3\}} = 227.01$, $\lambda^*_{\{5,4,2,3\}} = 358.42$, $\lambda^*_{\{1,7\}} = 515.56$, $\lambda^*_{\{6,7\}} = 3030.75$, $\lambda^*_{\{4,7,2,3\}} = 2172.00$, $\lambda^*_{\{4,6\}} = 28.52$, $\lambda^*_{\{4,1,6\}} = 42.30$, $\lambda^*_{\{4,6,2\}} = 336.33$.

Pricing. New variables: $\mathcal{V}_{12} = (x_{1\leftarrow\{6\}}, -6869.54, 203.45)$, $(x_{4\leftarrow\{1,2,6,7\}}, -1459.12, 0.62)$.

LP. $C = \dot{\bigcup}_{i=1}^{2} C_i$. $\mathcal{V} = \dot{\bigcup}_{i=1}^{12} \mathcal{V}_i$. Solution is $x_{5\leftarrow\emptyset} = 1/2$, $x_{7\leftarrow\emptyset} = 1/2$, $x_{1\leftarrow\{2,3,4,5,6,7\}} = 1/2$, $x_{3\leftarrow\{1,2\}} = 1/2$, $x_{4\leftarrow\{1,2,5,6,7\}} = 1/2$, $x_{5\leftarrow\{1,4,6,7\}} = 1/2$, $x_{6\leftarrow\{1,3,4,5,7\}} = 1/2$, $x_{7\leftarrow\{1,3,4,5,6\}} = 1/2$, $x_{3\leftarrow\{5,6,7\}} = 1/2$, $x_{4\leftarrow\{2\}} = 1/2$, $x_{2\leftarrow\{1,3\}} = 1/2$, $x_{2\leftarrow\{6\}} = 1/2$, $x_{6\leftarrow\{3\}} = 1/2$, $x_{1\leftarrow\{6\}} = 1/2$, Objective is -43158.09. Dual values $\lambda^*_i = -6940.37, -11078.32, -12002.83, -13113.32, -10534.51, -13762.64, -10518.83$, $\lambda^*_{\{1,2\}} = 962.04$, $\lambda^*_{\{4,2\}} = 1822.05$, $\lambda^*_{\{5,4\}} = 220.63$, $\lambda^*_{\{1,3\}} = 2093.38$, $\lambda^*_{\{4,1\}} = 1.19$, $\lambda^*_{\{5,6\}} = 3424.05$, $\lambda^*_{\{2,3\}} = 455.11$, $\lambda^*_{\{4,1,2\}} = 3739.95$, $\lambda^*_{\{4,2,3\}} = 3006.34$, $\lambda^*_{\{5,1,3\}} = 23.54$, $\lambda^*_{\{5,4,2,3\}} = 358.42$, $\lambda^*_{\{1,7\}} = 515.56$, $\lambda^*_{\{6,7\}} = 2855.17$, $\lambda^*_{\{4,7,2,3\}} = 2319.05$, $\lambda^*_{\{3,6,4,1,7\}} = 28.52$, $\lambda^*_{\{4,1,6\}} = 42.30$, $\lambda^*_{\{4,6,2\}} = 336.33$.

Pricing. New variables: $\mathcal{V}_{13} = (x_{1\leftarrow\{2,3,5,6\}}, -47.52, 3.08)$.

LP. $C = \dot{\bigcup}_{i=1}^{2} C_i$. $\mathcal{V} = \dot{\bigcup}_{i=1}^{13} \mathcal{V}_i$. Solution is $x_{5\leftarrow\emptyset} = 1/2$, $x_{7\leftarrow\emptyset} = 1/2$, $x_{1\leftarrow\{2,3,4,5,6,7\}} = 1/2$, $x_{3\leftarrow\{1,2\}} = 1/2$, $x_{4\leftarrow\{1,2,5,6,7\}} = 1/2$, $x_{5\leftarrow\{1,4,6,7\}} = 1/2$, $x_{6\leftarrow\{1,3,4,5,7\}} = 1/2$, $x_{7\leftarrow\{1,3,4,5,6\}} = 1/2$, $x_{3\leftarrow\{5,6,7\}} = 1/2$, $x_{4\leftarrow\{2\}} = 1/2$, $x_{2\leftarrow\{1,3\}} = 1/2$, $x_{2\leftarrow\{6\}} = 1/2$, $x_{6\leftarrow\{3\}} = 1/2$, $x_{1\leftarrow\{6\}} = 1/2$, Objective is -43158.09. Dual values: $-6940.37, -11078.32, -12005.92, -13113.32, -10534.51, -13762.64, -10518.83$, $\lambda^*_{\{1,2\}} = 962.04$, $\lambda^*_{\{4,2\}} = 1822.05$, $\lambda^*_{\{5,4\}} = 220.63$, $\lambda^*_{\{1,3\}} = 2096.47$, $\lambda^*_{\{4,1\}} = 1.19$,

$\lambda^*_{\{5,6\}}$ = 3424.05, $\lambda^*_{\{2,3\}}$ = 455.11, $\lambda^*_{\{4,1,2\}}$ = 3739.95, $\lambda^*_{\{4,2,3\}}$ = 3003.25, $\lambda^*_{\{5,1,3\}}$ = 23.54, $\lambda^*_{\{5,4,2,3\}}$ = 358.42, $\lambda^*_{\{1,7\}}$ = 512.47, $\lambda^*_{\{6,7\}}$ = 2855.17, $\lambda^*_{\{4,7,2,3\}}$ = 2322.14, $\lambda^*_{\{3,6,4,1,7\}}$ = 28.52, $\lambda^*_{\{4,1,6\}}$ = 42.30, $\lambda^*_{\{4,6,2\}}$ = 336.33.

Pricing. New variables: None

Separation. New clusters: C_3 = {2, 3, 6}, {1, 3, 6}, {2, 3, 4, 6}, {1, 2, 3, 6}, {1, 2, 3, 4, 6}, {1, 6}, {1, 2, 6}, {3, 6}, {1, 2, 4, 6}, {2, 3, 4, 5, 6}, {2, 3, 4, 6, 7}, {1, 3, 4, 6}, {1, 3, 5, 6}, {1, 3, 6, 7}, {1, 2, 3, 5, 6}, {2, 3, 6, 7}, {2, 3, 5, 6}, {1, 2, 3, 4, 5, 6}, {1, 2, 3, 4, 6, 7}, {1, 2, 3, 6, 7}.

LP. $C = \bigcup_{i=1}^{3} C_i$. $\mathcal{V} = \bigcup_{i=1}^{13} \mathcal{V}_i$. Solution is $x_{1\leftarrow\emptyset}$ = 1/2, $x_{4\leftarrow\emptyset}$ = 1/2, $x_{5\leftarrow\emptyset}$ = 1/2, $x_{6\leftarrow\emptyset}$ = 1/2, $x_{7\leftarrow\emptyset}$ = 1/2, $x_{1\leftarrow\{2,3,4,5,6,7\}}$ = 1/2, $x_{2\leftarrow\{1,3,4\}}$ = 1/2, $x_{3\leftarrow\{1,2\}}$ = 1/2, $x_{5\leftarrow\{1,4,6,7\}}$ = 1/2, $x_{6\leftarrow\{1,3,4,5,7\}}$ = 1/2, $x_{7\leftarrow\{1,3,4,5,6\}}$ = 1/2, $x_{4\leftarrow\{1,3,5,6,7\}}$ = 1/2, $x_{2\leftarrow\{4\}}$ = 1/2, $x_{3\leftarrow\{5,6,7\}}$ = 1/2. Objective is -45300.94. Dual values: -7115.31, -12643.78, -13942.09, -14678.78, -10534.51, -16208.59, -10518.83, $\lambda^*_{\{1,2\}}$ = 962.04, $\lambda^*_{\{4,2\}}$ = 1822.05, $\lambda^*_{\{5,4\}}$ = 220.63, $\lambda^*_{\{1,3\}}$ = 2190.82, $\lambda^*_{\{4,1\}}$ = 1.19, $\lambda^*_{\{5,6\}}$ = 3806.03, $\lambda^*_{\{2,3\}}$ = 455.11, $\lambda^*_{\{4,1,2\}}$ = 3645.60, $\lambda^*_{\{4,2,3\}}$ = 2908.90, $\lambda^*_{\{4,7,2\}}$ = 301.03, $\lambda^*_{\{1,7\}}$ = 512.47, $\lambda^*_{\{6,7\}}$ = 2710.47, $\lambda^*_{\{4,7,2,3\}}$ = 2194.34, $\lambda^*_{\{4,6\}}$ = 70.82, $\lambda^*_{\{6,1,3\}}$ = 198.77, $\lambda^*_{\{4,6,2,3\}}$ = 2275.69, $\lambda^*_{\{1,6\}}$ = 70.54.

Pricing. New variables: $\mathcal{V}_{14} = (x_{1\leftarrow\{2,3\}}, -106.68, 11.36). (x_{6\leftarrow\{2,3\}}, -13710.19, 23.91).$

LP. $C = \bigcup_{i=1}^{3} C_i$. $\mathcal{V} = \bigcup_{i=1}^{14} \mathcal{V}_i$. Solution is $x_{1\leftarrow\emptyset}$ = 1/2, $x_{5\leftarrow\emptyset}$ = 1/2, $x_{6\leftarrow\emptyset}$ = 1/2, $x_{7\leftarrow\emptyset}$ = 1/2, $x_{1\leftarrow\{2,3,4,5,6,7\}}$ = 1/2, $x_{2\leftarrow\{1,3,4\}}$ = 1/2, $x_{3\leftarrow\{1,2\}}$ = 1/2, $x_{5\leftarrow\{1,4,6,7\}}$ = 1/2, $x_{6\leftarrow\{1,3,4,5,7\}}$ = 1/2, $x_{7\leftarrow\{1,3,4,5,6\}}$ = 1/2, $x_{4\leftarrow\{1,3,5,6,7\}}$ = 1/2, $x_{3\leftarrow\{5,6,7\}}$ = 1/2, $x_{4\leftarrow\{2\}}$ = 1/2, $x_{2\leftarrow\emptyset}$ = 1/2, Objective value is -45300.93. -7115.31, -12643.76, -13942.09, -14678.76, -10534.51, -16208.59, -10518.83, $\lambda^*_{\{1,2\}}$ = 962.04, $\lambda^*_{\{4,2\}}$ = 1822.05, $\lambda^*_{\{5,4\}}$ = 220.63, $\lambda^*_{\{1,3\}}$ = 2190.83, $\lambda^*_{\{4,1\}}$ = 1.19, $\lambda^*_{\{5,6\}}$ = 3806.03, $\lambda^*_{\{2,3\}}$ = 455.11, $\lambda^*_{\{4,1,2\}}$ = 3645.59, $\lambda^*_{\{4,2,3\}}$ = 2908.89, $\lambda^*_{\{4,7,2\}}$ = 301.03, $\lambda^*_{\{1,7\}}$ = 512.47, $\lambda^*_{\{6,7\}}$ = 2710.47, $\lambda^*_{\{4,7,2,3\}}$ = 2194.34, $\lambda^*_{\{4,6\}}$ = 70.82, $\lambda^*_{\{4,6,2\}}$ = 23.92, $\lambda^*_{\{6,1,3\}}$ = 222.69, $\lambda^*_{\{4,6,2,3\}}$ = 2251.77, $\lambda^*_{\{1,6\}}$ = 46.61.

Pricing. New variables: None

Separation. New clusters: None