Optimal prediction of Markov chains with and without spectral gap

Abstract

We study the following learning problem with dependent data: Observing a trajectory of length n from a stationary Markov chain with k states, the goal is to predict the next state. For $3 \leq k \leq O(\sqrt{n})$, using techniques from universal compression, the optimal prediction risk in Kullback-Leibler divergence is shown to be $\Theta(\frac{k^2}{n} \log \frac{n}{k^2})$, in contrast to the optimal rate of $\Theta(\frac{\log \log n}{n})$ for $k = 2$ previously shown in [FOPS16]. These rates, slower than the parametric rate of $O(\frac{k^2}{n})$, can be attributed to the memory in the data, as the spectral gap of the Markov chain can be arbitrarily small. To quantify the memory effect, we study irreducible reversible chains with a prescribed spectral gap. In addition to characterizing the optimal prediction risk for two states, we show that, as long as the spectral gap is not excessively small, the prediction risk in the Markov model is $O(\frac{k^2}{n})$, which coincides with that of an iid model with the same number of parameters.

1 Introduction

Learning distributions from samples is a central question in statistics and machine learning. While significant progress has been achieved in property testing and estimation based on independent and identically distributed (iid) data, for many applications, most notably natural language processing, two new challenges arise: (a) Modeling data as independent observations fails to capture their temporal dependency; (b) Distributions are commonly supported on a large domain whose cardinality is comparable to or even exceeds the sample size. Continuing the progress made in [FOPS16, HOP18], in this paper we study the following prediction problem with dependent data modeled as Markov chains.

Suppose $X_1, X_2, \ldots$ is a stationary first-order Markov chain on state space $[k] \triangleq \{1, \ldots, k\}$ with unknown statistics. Observing a trajectory $X^n \triangleq (X_1, \ldots, X_n)$, the goal is to predict the next state X_{n+1} by estimating its distribution conditioned on the present data. We use the Kullback-Leibler (KL) divergence as the loss function: For distributions P, Q on $[k]$, $D(P\|Q) = \sum_{i=1}^{k} p_i \log \frac{p_i}{q_i}$ if $p_i = 0$ whenever $q_i = 0$ and $D(P\|Q) = \infty$ otherwise. The minimax prediction risk is given by

$$\mathsf{Risk}_{k,n} \triangleq \inf_{\widehat{M}} \sup_{\pi, M} \mathbb{E}[D(M(\cdot|X_n)\|\widehat{M}(\cdot|X_n))] = \inf_{\widehat{M}} \sup_{\pi, M} \sum_{i=1}^{k} \mathbb{E}[D(M(\cdot|i)\|\widehat{M}(\cdot|i))\mathbf{1}_{\{X_n=i\}}] \quad (1)$$

where the supremum is taken over all stationary distributions π and transition matrices M (row-stochastic) such that $\pi M = \pi$, the infimum is taken over all estimators $\widehat{M} = \widehat{M}(X_1, \ldots, X_n)$, and $M(\cdot|i)$ denotes the ith row of M. Our main objective is to characterize this minimax risk within universal constant factors as a function of n and k.

Submitted to 35th Conference on Neural Information Processing Systems (NeurIPS 2021). Do not distribute.

The prediction problem (1) is distinct from the parameter estimation problem such as estimating the transition matrix [Bar51, AG57, Bil61, WK19] or its properties [CS00, KV16, HJL$^+$18, HKL$^+$19], in that the quantity to be estimated (conditional distribution of the next state) depends on the sample path itself. This is precisely what renders the prediction problem closely relevant to natural applications such as autocomplete and text generation. In addition, this formulation allows more flexibility with far less assumptions compared to the estimation framework. For example, if certain state has very small probability under the stationary distribution, consistent estimation of the transition matrix may not be possible, whereas the prediction problem is unencumbered by such rare states.

In the special case of iid data, the prediction problem reduces to estimating the distribution in KL divergence. In this setting the optimal risk is well understood, which is known to be $\frac{k-1}{2n}(1 + o(1))$ when k is fixed and $n \to \infty$ [BFSS02] and $\Theta(\frac{k}{n})$ for $k = O(n)$ [Pan04, KOPS15].[1] Typical in parametric models, this rate $\frac{k}{n}$ is commonly referred to the "parametric rate", which leads to a sample complexity that scales proportionally to the number of parameters and inverse proportionally to the desired accuracy.

In the setting of Markov chains, however, the prediction problem is much less understood especially for large state space. Recently the seminal work [FOPS16] showed the surprising result that for stationary Markov chains on two states, the optimal prediction risk satisfies

$$\text{Risk}_{2,n} = \Theta\left(\frac{\log\log n}{n}\right), \tag{2}$$

which has a nonparametric rate even when the problem has only two parameters. The follow-up work [HOP18] studied general k-state chains and showed a lower bound of $\Omega(\frac{k \log\log n}{n})$ for uniform (not necessarily stationary) initial distribution; however, the upper bound $O(\frac{k^2 \log\log n}{n})$ in [HOP18] relies on implicit assumptions on mixing time such as spectral gap conditions. The following theorem, which is the main result of this paper, resolves the optimal risk for k-state Markov chains:

Theorem 1 (Optimal rates without spectral gap). *There exists a universal constant $C > 0$ such that for all $3 \le k \le \sqrt{n}/C$,*

$$\frac{k^2}{Cn}\log\left(\frac{n}{k^2}\right) \le \text{Risk}_{k,n} \le \frac{Ck^2}{n}\log\left(\frac{n}{k^2}\right). \tag{3}$$

Furthermore, the lower bound continues to hold even if the Markov chain is restricted to be irreducible and reversible.

Remark 1. The optimal prediction risk of $O(\frac{k^2}{n}\log\frac{n}{k^2})$ can be achieved by an average version of the *add-one estimator* (i.e. Laplace's rule of succession). Given a trajectory $x^n = (x_1, \ldots, x_n)$ of length n, denote the transition counts (with the convention $N_i \equiv N_{ij} \equiv 0$ if $n = 0, 1$)

$$N_i = \sum_{\ell=1}^{n-1} \mathbf{1}_{\{x_\ell=i\}}, \quad N_{ij} = \sum_{\ell=1}^{n-1} \mathbf{1}_{\{x_\ell=i, x_{\ell+1}=j\}}. \tag{4}$$

The add-one estimator for the transition probability $M(j|i)$ is given by

$$\widehat{M}_{x^n}^{+1}(j|i) \triangleq \frac{N_{ij} + 1}{N_i + k}, \tag{5}$$

which is an additively smoothed version of the empirical frequency. Finally, the optimal rate in (3) can be achieved by the following estimator $\widehat{M}$ defined as an average of add-one estimators over different sample sizes:

$$\widehat{M}_{x^n}(x_{n+1}|x_n) \triangleq \frac{1}{n}\sum_{t=1}^{n} \widehat{M}_{x_{n-t+1}^n}^{+1}(x_{n+1}|x_n). \tag{6}$$

In other words, we apply the add-one estimator to the most recent t observations $(X_{n-t+1}, \ldots, X_n)$ to predict the next X_{n+1}, then average over $t = 1, \ldots, n$. Such Cesàro-mean-type estimators have been introduced before in the density estimation literature (see, e.g., [YB99]). It remains open

---

[1]Here and below $\asymp, \lesssim, \gtrsim$ or $\Theta(\cdot), O(\cdot), \Omega(\cdot)$ denote equality and inequalities up to universal multiplicative constants.

whether the usual add-one estimator (namely, the last term in (6) which uses all the data) or any add-c estimator for constant c achieves the optimal rate. In contrast, for two-state chains the optimal risk (2) is attained by a hybrid strategy [FOPS16], applying add-c estimator for $c = \frac{1}{\log n}$ for trajectories with at most one transition and $c = 1$ otherwise.

Theorem 1 shows that the departure from the parametric rate of $\frac{k^2}{n}$, first discovered in [FOPS16, HOP18] for binary chains, is even more pronounced for larger state space. As will become clear in the proof, there is some fundamental difference between two-state and three-state chains, resulting in $\mathsf{Risk}_{3,n} = \Theta(\frac{\log n}{n}) \gg \mathsf{Risk}_{2,n} = \Theta(\frac{\log \log n}{n})$. It is instructive to compare the sample complexity for prediction in the iid and Markov model. Denote by d the number of parameters, which is $k-1$ for the iid case and $k(k-1)$ for Markov chains. Define the sample complexity $n^*(d, \epsilon)$ as the smallest sample size n in order to achieve a prescribed prediction risk ϵ. For $\epsilon = O(1)$, we have

$$n^*(d, \epsilon) \asymp \begin{cases} \frac{d}{\epsilon} & \text{iid} \\ \frac{d}{\epsilon} \log \log \frac{1}{\epsilon} & \text{Markov with 2 states} \\ \frac{d}{\epsilon} \log \frac{1}{\epsilon} & \text{Markov with } k \geq 3 \text{ states.} \end{cases} \tag{7}$$

At a high level, the nonparametric rates in the Markov model can be attributed to the memory in the data. On the one hand, Theorem 1 as well as (2) affirm that one can obtain meaningful prediction without imposing any mixing conditions;[2] such decoupling between learning and mixing has also been observed in other problems such as learning linear dynamics [SMT$^+$18, DMM$^+$19]. On the other hand, the dependency in the data does lead to a strictly higher sample complexity than that of the iid case; in fact, the lower bound in Theorem 1 is proved by constructing chains with spectral gap as small as $O(\frac{1}{n})$ (see Section 3). Thus, it is conceivable that with sufficiently favorable mixing conditions, the prediction risk improves over that of the worst case and, at some point, reaches the parametric rate. To make this precise, we focus on Markov chains with a prescribed spectral gap.

It is well-known that for an irreducible and reversible chain, the transition matrix M has k real eigenvalues satisfying $1 = \lambda_1 \geq \lambda_2 \geq \ldots \lambda_k \geq -1$. The *absolute spectral gap* of M, defined as

$$\gamma_* \triangleq 1 - \max \left\{ |\lambda_i| : i \neq 1 \right\}, \tag{8}$$

quantifies the memory of the Markov chain. For example, the mixing time is determined by $1/\gamma^*$ (relaxation time) up to logarithmic factors. As extreme cases, the chain which does not move (M is identity) and which is iid (M is rank-one) have spectral gap equal to 0 and 1, respectively. We refer the reader to [LP17] for more background. Given $\gamma_0 \in (0, 1)$, define $\mathcal{M}_k(\gamma_0)$ as the set of transition matrices corresponding to irreducible and reversible chains whose absolute spectral gap exceeds γ_0. Restricting (1) to this subcollection and noticing the stationary distribution here is uniquely determined by M, we define the corresponding minimax risk:

$$\mathsf{Risk}_{k,n}(\gamma_0) \triangleq \inf_{\widehat{M}} \sup_{M \in \mathcal{M}_k(\gamma_0)} \mathbb{E} \left[D(M(\cdot|X_n) \| \widehat{M}(\cdot|X_n)) \right] \tag{9}$$

Extending the result (2) of [FOPS16], the following theorem characterizes the optimal prediction risk for two-state chains with prescribed spectral gaps:

Theorem 2 (Spectral gap dependent rates for binary chain). *For any $\gamma_0 \in (0, 1)$*

$$\mathsf{Risk}_{2,n}(\gamma_0) \asymp \frac{1}{n} \max \left\{ 1, \log \log \left(\min \left\{ n, \frac{1}{\gamma_0} \right\} \right) \right\}.$$

Theorem 2 shows that for binary chains, parametric rate $O(\frac{1}{n})$ is achievable if and only if the spectral gap is nonvanishing. While this holds for bounded state space (see Corollary 4 below), for large state space, it turns out that much weaker conditions on the absolute spectral gap suffice to guarantee the parametric rate $O(\frac{k^2}{n})$, achieved by the add-one estimator applied to the entire trajectory. In other words, as long as the spectral gap is not excessively small, the prediction risk in the Markov model behaves in the same way as that of an iid model with equal number of parameters. Similar conclusion has been established previously for the sample complexity of estimating the entropy rate of Markov chains in [HJL$^+$18, Theorem 1].

---

[2]To see this, it is helpful to consider the extreme case where the chain does not move at all or is periodic, in which case predicting the next state is in fact easy.

Theorem 3. *The add-one estimator in* (5) *achieves the following risk bound.*

    (i) For any $k \geq 2$, $\mathsf{Risk}_{k,n}(\gamma_0) \lesssim \frac{k^2}{n}$ provided that $\gamma_0 \gtrsim (\frac{\log k}{k})^{1/4}$.

    (ii) In addition, for $k \gtrsim (\log n)^6$, $\mathsf{Risk}_{k,n}(\gamma_0) \lesssim \frac{k^2}{n}$ provided that $\gamma_0 \gtrsim \frac{(\log(n+k))^2}{k}$.

Corollary 4. *For any fixed $k \geq 2$, $\mathsf{Risk}_{k,n}(\gamma_0) = O(\frac{1}{n})$ if and only if $\gamma_0 = \Omega(1)$.*

1.1 Proof techniques

The proof of Theorem 1 deviates from existing approaches based on concentration inequalities for Markov chains. For instance, the standard program for analyzing the add-one estimator (5) involves proving concentration of the empirical counts on their population version, namely, $N_i \approx n\pi_i$ and $N_{ij} \approx n\pi_i M(j|i)$, and bounding the risk in the atypical case by concentration inequalities, such as the Chernoff-type bounds in [Lez98, Pau15], which have been widely used in recent work on statistical inference with Markov chains [KV16, HJL$^+$18, HOP18, HKL$^+$19, WK19]. However, these concentration inequalities inevitably depends on the spectral gap of the Markov chain, leading to results which deteriorate as the spectral gap becomes smaller. For two-state chains, results free of the spectral gap are obtained in [FOPS16] using explicit joint distribution of the transition counts; this refined analysis, however, is difficult to extend to larger state space as the probability mass function of (N_{ij}) is given by Whittle's formula [Whi55] which takes an unwieldy determinantal form.

Eschewing concentration-based arguments, the crux of our proof of Theorem 1, for both the upper and lower bound, revolves around the following quantity known as *redundancy*:

$$\mathsf{Red}_{k,n} \triangleq \inf_{Q_{X^n}} \sup_{P_{X^n}} D(P_{X^n} \| Q_{X^n}) = \inf_{Q_{X^n}} \sup_{P_{X^n}} \sum_{x^n} P_{X^n}(x^n) \log \frac{P_{X^n}(x^n)}{Q_{X^n}(x^n)}. \tag{10}$$

Here the supremum is taken over all joint distributions of stationary Markov chains X^n on k states, and the infimum is over all joint distributions Q_{X^n}. A central quantity which measures the minimax regret in universal compression, the redundancy (10) corresponds to minimax cumulative risk (namely, the total prediction risk when the sample size ranges from 1 to n), while (1) is the individual minimax risk at sample size n – see Section 2 for a detailed discussion. We prove the following reduction between prediction risk and redundancy:

$$\frac{1}{n}\mathsf{Red}^{\mathsf{sym}}_{k-1,n} - \frac{\log k}{n} \lesssim \mathsf{Risk}_{k,n} \leq \frac{1}{n-1}\mathsf{Red}_{k,n} \tag{11}$$

where $\mathsf{Red}^{\mathsf{sym}}$ denotes the redundancy for *symmetric* Markov chains. The upper bound is standard: thanks to the convexity of the loss function and stationarity of the Markov chain, the risk of the Cesàro-mean estimator (6) can be related to the cumulative risk and, in turn, the redundancy. The proof of the lower bound is more involved. Given a $(k-1)$-state chain, we embed it into a larger state space by introducing a new state, such that with constant probability, the chain starts from and gets stuck at this state for a period time that is approximately uniform in $[n]$, then enters the original chain. Effectively, this scenario is equivalent to a prediction problem on $k-1$ states with a *random* (approximately uniform) sample size, whose prediction risk can then be related to the cumulative risk and redundancy. This intuition can be made precise by considering a Bayesian setting, in which the $(k-1)$-state chain is randomized according to the least favorable prior for (10), and representing the Bayes risk as conditional mutual information and applying the chain rule.

Given the above reduction, it suffices to show both redundancies in (10) are on the order of $\frac{k^2}{n} \log \frac{n}{k^2}$. The redundancy is upper bounded by *pointwise redundancy*, which replaces the average in (10) by the maximum over all trajectories. Following [DMPW81, CS04], we consider an explicit probability assignment defined by add-one smoothing and using combinatorial arguments to bound the pointwise redundancy, shown optimal by information-theoretic arguments.

The optimal spectral gap-dependent rate in Theorem 2 relies on the key observation in [FOPS16] that, for binary chains, the dominating contribution to the prediction risk comes from trajectories with a single transition, for which we may apply an add-c estimator with c depending appropriately on the spectral gap. The lower bound is shown using a Bayesian argument similar to that of [HOP18, Theorem 1]. The proof of Theorem 3 relies on more delicate concentration arguments as the spectral gap is allowed to be vanishingly small. Notably, for small k, direct application of existing Bernstein

inequalities for Markov chains in [Lez98, Pau15] falls short of establishing the parametric rate of $O(\frac{k^2}{n})$ (see Remark 5 in Section 7.2 for details); instead, we use a fourth moment bound which turns out to be well suited for analyzing concentration of empirical counts conditional on the terminal state.

For large k, we further improve the spectral gap condition using a simulation argument for Markov chains using independent samples [Bil61, HJL$^+$18]. A key step is a new concentration inequality for $D(P\|\widehat{P}_{n,k}^{+1})$, where $\widehat{P}_{n,k}^{+1}$ is the add-one estimator based on n iid observations of P supported on $[k]$:

$$\mathbb{P}\left(D(P\|\widehat{P}_{n,k}^{+1}) \geq c \cdot \frac{k}{n} + \frac{\mathsf{polylog}(n) \cdot \sqrt{k}}{n}\right) \leq \frac{1}{\mathsf{poly}(n)}, \qquad (12)$$

for some absolute constant $c > 0$. Note that an application of the classical concentration inequality of McDiarmid would result in the second term being $\mathsf{polylog}(n)/\sqrt{n}$, and (12) crucially improves this to $\mathsf{polylog}(n) \cdot \sqrt{k}/n$. Such an improvement has been recently observed by [MJT$^+$20, Agr20, GR20] in studying the similar quantity $D(\widehat{P}_n\|P)$ for the (unsmoothed) empirical distribution $\widehat{P}_n$; however, these results, based on either the method of types or an explicit upper bound of the moment generating function, are not directly applicable to (12) in which the true distribution P appears as the first argument in the KL divergence.

1.2 Related work

While the exact prediction problem studied in this paper has recently been in focus since [FOPS16, HOP18], there exists a large body of literature on relate works. As mentioned before some of our proof strategies draws inspiration and results from the study of redundancy in universal compression, its connection to mutual information, as well as the perspective of sequential probability assignment as prediction, dating back to [Dav73, DMPW81, Ris84, Sht87, Rya88]. Asymptotic characterization of the minimax redundancy for Markov sources, both average and pointwise, were obtained in [Dav83, Att99, JS02], in the regime of fixed alphabet size k and large sample size n. Non-asymptotic characterization was obtained in [Dav83] for $n \gg k \log^2 k$ and recently extended to $n \asymp k^2$ in [TJW18], which further showed that the behavior of the redundancy remains unchanged even if the Markov chain is very close to being iid in terms of spectral gap $\gamma^* = 1 - o(1)$.

The current paper adds to a growing body of literature devoted to statistical learning with dependent data, in particular those dealing with Markov chains. Estimation of the transition matrix [Bar51, AG57, Bil61, Sin64] and testing the order of Markov chains [CS00] have been well studied in the large-sample regime. More recently attention has been shifted towards large state space and nonasymptotics. For example, [WK19] studied the estimation of transition matrix in $\ell_\infty \to \ell_\infty$ induced norm for Markov chains with prescribed pseudo spectral gap and minimum probability mass of the stationary distribution, and determined sample complexity bounds up to logarithmic factors. Similar results have been obtained for estimating properties of Markov chains, including mixing time and spectral gap [HKL$^+$19], entropy rate [KV16, HJL$^+$18, OS20], graph statistics based on random walk [BHOP18], as well as identity testing [DDG18, CB19, WK20, FW21]. Most of these results rely on assumptions on the Markov chains such as lower bounds on the spectral gap and the stationary distribution, which afford concentration for sample statistics of Markov chains. In contrast, one of the main contributions in this paper, in particular Theorem 1, is that optimal prediction can be achieved without these assumptions, thereby providing a novel way of tackling these seemingly unavoidable issues. This is ultimately accomplished by information-theoretic and combinatorial techniques from universal compression.

1.3 Notations and preliminaries

For $n \in \mathbb{N}$, let $[n] \triangleq \{1, \ldots, n\}$. Denote $x^n = (x_1, \ldots, x_n)$ and $x_t^n = (x_t, \ldots, x_n)$. The distribution of a random variable X is denoted by P_X. In a Bayesian setting, the distribution of a parameter θ is referred to as a prior, denoted by P_θ. We recall the following definitions from information theory [CK82, CT06]. The conditional KL divergence is defined as as an average of KL divergence between conditional distributions:

$$D(P_{A|B}\|Q_{A|B}|P_B) \triangleq \mathbb{E}_{B \sim P_B}[D(P_{A|B}\|Q_{A|B})] = \int P_B(db)D(P_{A|B=b}\|Q_{A|B=b}). \qquad (13)$$

The mutual information between random variables A and B with joint distribution P_{AB} is $I(A;B) \triangleq D(P_{B|A}\|P_B|P_A)$; similarly, the conditional mutual information is defined as $I(A;B|C) \triangleq D(P_{B|A,C}\|P_{B|C}|P_{A,C})$. The following variational representation of (conditional) mutual information is well-known

$$I(A;B) = \min_{Q_B} D(P_{B|A}\|Q_B|P_A), \quad I(A;B|C) = \min_{Q_{B|C}} D(P_{B|A,C}\|Q_{B|C}|P_{AC}). \quad (14)$$

The entropy of a discrete random variables X is $H(X) \triangleq \sum_x P_X(x) \log \frac{1}{P_X(x)}$.

The rest of the paper is organized as follows. In Section 2 we describe the general paradigms of minimax redundancy and prediction risk and their dual representation in terms of mutual information. We give a general redundancy-based bound on the prediction risk, which leads to the upper bound in Theorem 1. Section 3 presents the main idea of the lower bound construction. In the supplement, Section 4 discusses the assumptions and implications of our results and several open problems. Sections 5 and 6 contain the full proof for the upper and lower bound in Theorem 1. Due to space limitations, proofs for spectral-gap dependent risk bounds in Theorems 2 and 3 are given in Section 7.

2 Two general paradigms

2.1 Redundancy, prediction risk, and mutual information representation

For $n \in \mathbb{N}$, let $\mathcal{P} = \{P_{X^{n+1}|\theta} : \theta \in \Theta\}$ be a collection of joint distributions parameterized by θ.

"Compression". Consider a sample $X^n \triangleq (X_1, \ldots, X_n)$ of size n drawn from $P_{X^n|\theta}$ for some unknown $\theta \in \Theta$. The *redundancy* of a probability assignment (joint distribution) Q_{X^n} is defined as the worst-case KL risk of fitting the joint distribution of X^n, namely

$$\mathsf{Red}(Q_{X^n}) \triangleq \sup_{\theta \in \Theta} D(P_{X^n|\theta}\|Q_{X^n}). \quad (15)$$

Optimizing over Q_{X^n}, the minimax redundancy is defined as

$$\mathsf{Red}_n \triangleq \inf_{Q_{X^n}} \mathsf{Red}_n(Q_{X^n}), \quad (16)$$

where the infimum is over all joint distribution Q_{X^n}. This quantity can be operationalized as the redundancy (i.e. regret) in the setting of universal data compression, that is, the excess number of bits compared to the optimal compressor of X^n that knows θ [CT06, Chapter 13].

The capacity-redundancy theorem (see [Kem74] for a very general result) provides the following mutual information characterization of (16):

$$\mathsf{Red}_n = \sup_{P_\theta} I(\theta; X^n), \quad (17)$$

where the supremum is over all distributions (priors) P_θ on Θ. In view of the variational representation (14), this result can be interpreted as a minimax theorem:

$$\mathsf{Red}_n = \inf_{Q_{X^n}} \sup_{P_\theta} D(P_{X^n|\theta}\|Q_{X^n}|P_\theta) = \sup_{P_\theta} \inf_{Q_{X^n}} D(P_{X^n|\theta}\|Q_{X^n}|P_\theta).$$

Typically, for fixed model size and $n \to \infty$, one expects that $\mathsf{Red}_n = \frac{d}{2} \log n(1 + o(1))$, where d is the number of parameters; see [Ris84] for a general theory of this type. Indeed, on a fixed alphabet of size k, we have $\mathsf{Red}_n = \frac{k-1}{2} \log n(1 + o(1))$ for iid model [Dav73] and $\mathsf{Red}_n = \frac{k^m(k-1)}{2} \log n(1 + o(1))$ for m-order Markov models [Tro74], with more refined asymptotics shown in [XB97, SW12]. For large alphabets, nonasymptotic results have also been obtained. For example, for first-order Markov model, $\mathsf{Red}_n \asymp k^2 \log \frac{n}{k^2}$ provided that $n \gtrsim k^2$ [Dav83, TJW18].

"Prediction". Consider the problem of predicting the next unseen data point X_{n+1} based on the observations $X_1, \ldots, X_n$, where $(X_1, \ldots, X_{n+1})$ are jointly distributed as $P_{X^{n+1}|\theta}$ for some unknown $\theta \in \Theta$. Here, an estimator is a distribution (for X_{n+1}) as a function of X^n, which, in turn, can be written as a conditional distribution $Q_{X_{n+1}|X^n}$. As such, its worst-case average risk is

$$\mathsf{Risk}(Q_{X_{n+1}|X^n}) \triangleq \sup_{\theta \in \Theta} D(P_{X_{n+1}|X^n,\theta}\|Q_{X_{n+1}|X^n}|P_{X^n|\theta}), \quad (18)$$

where the conditional KL divergence is defined in (13). The minimax prediction risk is then defined as

$$\mathsf{Risk}_n \triangleq \inf_{Q_{X_{n+1}|X^n}} \mathsf{Risk}_n(Q_{X_{n+1}|X^n}), \tag{19}$$

While (16) does not directly correspond to a statistical estimation problem, (19) is exactly the familiar setting of "density estimation", where $Q_{X_{n+1}|X^n}$ is understood as an estimator for the distribution of the unseen X_{n+1} based on the available data $X_1, \ldots, X_n$.

In the Bayesian setting where θ is drawn from a prior P_θ, the Bayes prediction risk coincides with the conditional mutual information as a consequence of the variational representation (14):

$$\inf_{Q_{X_{n+1}|X^n}} \mathbb{E}_\theta[D(P_{X_{n+1}|X^n,\theta}\|Q_{X_{n+1}|X^n}|P_{X^n|\theta})] = I(\theta; X_{n+1}|X^n). \tag{20}$$

Furthermore, the Bayes estimator that achieves this infimum takes the following form:

$$Q^{\mathsf{Bayes}}_{X_{n+1}|X^n} = P_{X^{n+1}|X^n} = \frac{\int_\Theta P_{X^{n+1}|\theta}\, dP_\theta}{\int_\Theta P_{X^n|\theta}\, dP_\theta}, \tag{21}$$

known as the Bayes predictive density [Dav73, LB04]. These representations play a crucial role in the lower bound proof of Theorem 1. Under appropriate conditions which hold for Markov models (see Lemma 21 in Appendix A), the minimax prediction risk (19) also admits a dual representation analogous to (17):

$$\mathsf{Risk}_n = \sup_{\theta \sim \pi} I(\theta; X_{n+1}|X^n), \tag{22}$$

which, in view of (20), show that the principle of "minimax=worst-case Bayes" continues to hold for prediction problem in Markov models.

The following result (proved in Section 5) relates the redundancy and the prediction risk.

Lemma 5. *For any model $\mathcal{P}$,*

$$\mathsf{Red}_n \leq \sum_{t=0}^{n-1} \mathsf{Risk}_t. \tag{23}$$

In addition, suppose that each $P_{X^n|\theta} \in \mathcal{P}$ is stationary and m^{th}-order Markov. Then for all $n \geq m+1$,

$$\mathsf{Risk}_n \leq \mathsf{Risk}_{n-1} \leq \frac{\mathsf{Red}_n}{n-m}. \tag{24}$$

Furthermore, for any joint distribution Q_{X^n} factorizing as $Q_{X^n} = \prod_{t=1}^n Q_{X_t|X^{t-1}}$, the prediction risk of the estimator

$$\widetilde{Q}_{X_n|X^{n-1}}(x_n|x^{n-1}) \triangleq \frac{1}{n-m} \sum_{t=m+1}^{n} Q_{X_t|X^{t-1}}(x_n|x_{n-t+1}^{n-1}) \tag{25}$$

is bounded by the redundancy of Q_{X^n} as

$$\mathsf{Risk}(\widetilde{Q}_{X_n|X^{n-1}}) \leq \frac{1}{n-m}\mathsf{Red}(Q_{X^n}). \tag{26}$$

Remark 2. Note that the upper bound (23) on redundancy, known as the "estimation-compression inequality" [KOPS15, FOPS16], holds without conditions, while the lower bound (24) relies on stationarity and Markovity. For iid data, the estimation-compression inequality is almost an equality; however, this is not the case for Markov chains, as both sides of (23) differ by an unbounded factor of $\Theta(\log\log n)$ for $k = 2$ and $\Theta(\log n)$ for fixed $k \geq 3$ – see (2) and Theorem 1. On the other hand, Markov chains with at least three states offers a rare instance where (24) is tight, namely, $\mathsf{Risk}_n \asymp \frac{\mathsf{Red}_n}{n}$ (cf. Lemma 6).

2.2 Proof of the upper bound part of Theorem 1

Specializing to first-order stationary Markov chains with k states, we denote the redundancy and prediction risk in (16) and (19) by $\mathsf{Red}_{k,n}$ and $\mathsf{Risk}_{k,n}$, the latter of which is precisely the quantity

previously defined in (1). Applying Lemma 5 yields $\text{Risk}_{k,n} \leq \frac{1}{n-1}\text{Red}_{k,n}$. To upper bound $\text{Red}_{k,n}$, consider the following probability assignment:

$$Q(x_1, \cdots, x_n) = \frac{1}{k} \prod_{t=1}^{n-1} \widehat{M}_{x^t}^{+1}(x_{t+1}|x_t) \tag{27}$$

where $\widehat{M}^{+1}$ is the add-one estimator defined in (5). This Q factorizes as $Q(x_1) = \frac{1}{k}$ and $Q(x_{t+1}|x^t) = \widehat{M}_{x^t}^{+1}(x_{t+1}|x_t)$. The following lemma (proved in Section 5.2) bounds the redundancy of Q:

Lemma 6. $\text{Red}(Q) \leq k(k-1)\left[\log\left(1 + \frac{n-1}{k(k-1)}\right) + 1\right] + \log k$.

Combined with Lemma 5, Lemma 6 shows that $\text{Risk}_{k,n} \leq C\frac{k^2}{n}\log\frac{n}{k^2}$ for all $k \leq \sqrt{n/C}$ and some universal constant C, achieved by the estimator (6), which is obtained by applying the rule (25) to (27).

3   Optimal rates without spectral gap

In this section, we sketch the proof of the lower bound part of Theorem 1, which shows the optimality of the average version of the add-one estimator (25). We first describe the lower bound construction for three-state chains, which is subsequently extended to k states.

3.1   Warmup: an $\Omega(\frac{\log n}{n})$ lower bound for three-state chains

Theorem 7. $\text{Risk}_{3,n} = \Omega\left(\frac{\log n}{n}\right)$.

To show Theorem 7, consider the following one-parameter family of transition matrices:

$$\mathcal{M} = \left\{M_p = \begin{bmatrix} 1 - \frac{2}{n} & \frac{1}{n} & \frac{1}{n} \\ \frac{1}{n} & 1 - \frac{1}{n} - p & p \\ \frac{1}{n} & p & 1 - \frac{1}{n} - p \end{bmatrix} : 0 \leq p \leq 1 - \frac{1}{n}\right\}. \tag{28}$$

Note that each transition matrix in $\mathcal{M}$ is symmetric (hence doubly stochastic), whose corresponding chain is reversible with a uniform stationary distribution and spectral gap $\Theta(\frac{1}{n})$; see Fig. 1.

Figure 1: Lower bound construction for three-state chains.

The proof of Theorem 7 is given in Section 6.1 using a Bayesian argument. The main idea is as follows. Notice that by design, with constant probability, the trajectory is of the following form: The chain starts and stays at state 1 for t steps, and then transitions into state 2 or 3 and never returns to state 1, where $t = 1, \ldots, n-1$. Since p is the single unknown parameter, the only useful observations are the visits to state 2 and 3 and each visit entails one observation about p by flipping a coin with bias roughly p. Thus the effective sample size for estimating p is $n - t - 1$ and we expect the best

estimation error is of the order of $\frac{1}{n-t}$. However, t is not fixed. In fact, conditioned on the trajectory is of this form, t is roughly uniformly distributed between 1 and $n-1$. As such, we anticipate the estimation error of p is approximately

$$\frac{1}{n-1}\sum_{i=1}^{n-1}\frac{1}{n-t} = \Theta\left(\frac{\log n}{n}\right).$$

Intuitively speaking, the construction in Fig. 1 "embeds" a symmetric two-state chain (with states 2 and 3) with unknown parameter p into a space of three states, by adding a "nuisance" state 1, which effectively slows down the exploration of the useful part of the state space, so that in a trajectory of length n, the effective number of observations we get to make about p is roughly uniformly distributed between 1 and n. This explains the extra log factor in Theorem 7, which actually stems from the harmonic sum in $\mathbb{E}[\frac{1}{\text{Uniform}([n])}]$. We will fully explore this embedding idea in Section 3.2 to deal with larger state space.

3.2 k-state chains

The above lower bound construction for 3-state chains can be generalized to k-state chains with $k \geq 3$. The high-level argument is again to augment a $(k-1)$-state chain into a k-state chain. Specifically, we partition the state space $[k]$ into two sets $\mathcal{S}_1 = \{1\}$ and $\mathcal{S}_2 = \{2, 3, \cdots, k\}$. Consider a k-state Markov chain such that the transition probabilities from $\mathcal{S}_1$ to $\mathcal{S}_2$, and from $\mathcal{S}_2$ to $\mathcal{S}_1$, are both very small (on the order of $\Theta(1/n)$). At state 1, the chain either stays at 1 with probability $1 - 1/n$ or moves to one of the states in $\mathcal{S}_2$ with equal probability $\frac{1}{n(k-1)}$; at each state in $\mathcal{S}_2$, the chain moves to 1 with probability $\frac{1}{n}$; otherwise, within the state subspace $\mathcal{S}_2$, the chain evolves according to some symmetric transition matrix T. See Fig. 2 in Section 6.2 for the precise transition diagram.

The key feature of such a chain is as follows. Let $\mathcal{X}_t$ be the event that $X_1, X_2, \cdots, X_t \in \mathcal{S}_1$ and $X_{t+1}, \cdots, X_n \in \mathcal{S}_2$. For each $t \in [n-1]$, one can show that $\mathbb{P}(\mathcal{X}_t) \geq c/n$ for some absolute constant $c > 0$. Moreover, conditioned on the event $\mathcal{X}_t$, $(X_{t+1}, \ldots, X_n)$ is equal in law to a stationary Markov chain $(Y_1, \cdots, Y_{n-t})$ on state space $\mathcal{S}_2$ with symmetric transition matrix T. It is not hard to show that estimating M and T are nearly equivalent. Consider the Bayesian setting where T is drawn from some prior. We have

$$\inf_{\widehat{M}} \mathbb{E}_T\left[\mathbb{E}[D(M(\cdot|X_n)\|\widehat{M}(\cdot|X_n))|\mathcal{X}_t]\right] \approx \inf_{\widehat{T}} \mathbb{E}_T\left[\mathbb{E}[D(T(\cdot|Y_{n-t})\|\widehat{T}(\cdot|Y_{n-t}))]\right] = I(T; Y_{n-t+1}|Y^{n-t}),$$

where the last equality follows from the representation (20) of Bayes prediction risk as conditional mutual information. Lower bounding the minimax risk by the Bayes risk, we have

$$\begin{aligned}
\text{Risk}_{k,n} &\geq \inf_{\widehat{M}} \mathbb{E}_T\left[\mathbb{E}[D(M(\cdot|X_n)\|\widehat{M}(\cdot|X_n))]\right] \\
&\geq \inf_{\widehat{M}} \sum_{t=1}^{n-1} \mathbb{E}_M\left[\mathbb{E}[D(M(\cdot|X_n)\|\widehat{M}(\cdot|X_n))|\mathcal{X}_t] \cdot \mathbb{P}(\mathcal{X}_t)\right] \\
&\geq \frac{c}{n} \cdot \sum_{t=1}^{n-1} \inf_{\widehat{M}} \mathbb{E}_M\left[\mathbb{E}[D(M(\cdot|X_n)\|\widehat{M}(\cdot|X_n))|\mathcal{X}_t]\right] \\
&\approx \frac{c}{n} \cdot \sum_{t=1}^{n-1} I(T; Y_{n-t+1}|Y^{n-t}) = \frac{c}{n} \cdot (I(T; Y^n) - I(T; Y_1)).
\end{aligned} \tag{29}$$

Note that $I(T; Y_1) \leq H(Y_1) \leq \log(k-1)$ since Y_1 takes values in $\mathcal{S}_2$. Maximizing the right hand side over the prior P_T and recalling the dual representation for redundancy in (17), the above inequality (29) leads to a risk lower bound of $\text{Risk}_{k,n} \gtrsim \frac{1}{n}(\text{Red}^{\text{sym}}_{k-1,n} - \log k)$, where $\text{Red}^{\text{sym}}_{k-1,n} = \sup I(T; Y_1)$ is the redundancy for *symmetric* Markov chains with $k-1$ states and sample size n. Since symmetric transition matrices still have $\Theta(k^2)$ degrees of freedom, it is expected that $\text{Red}^{\text{sym}}_{k,n} \asymp k^2 \log \frac{n}{k^2}$ for $n \gtrsim k^2$, so that (29) yields the desired lower bound $\text{Risk}_{k,n} = \Omega(\frac{k^2}{n} \log \frac{n}{k^2})$ in Theorem 1. The rigorous proof is carried out in Section 6.2.

References

[AG57] Theodore W Anderson and Leo A Goodman. Statistical inference about Markov chains. *The Annals of Mathematical Statistics*, pages 89–110, 1957.

[Agr20] Rohit Agrawal. Finite-sample concentration of the multinomial in relative entropy. *IEEE Transactions on Information Theory*, 66(10):6297–6302, 2020.

[Att99] K. Atteson. The asymptotic redundancy of bayes rules for Markov chains. *IEEE Transactions on Information Theory*, 45(6):2104–2109, 1999.

[Bar51] Maurice S Bartlett. The frequency goodness of fit test for probability chains. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 47, pages 86–95. Cambridge University Press, 1951.

[BFSS02] Dietrich Braess, Jürgen Forster, Tomas Sauer, and Hans U Simon. How to achieve minimax expected Kullback-Leibler distance from an unknown finite distribution. In *Algorithmic Learning Theory*, pages 380–394. Springer, 2002.

[BHOP18] Anna Ben-Hamou, Roberto I Oliveira, and Yuval Peres. Estimating graph parameters via random walks with restarts. In *Proceedings of the Twenty-Ninth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1702–1714. SIAM, 2018.

[Bil61] P. Billingsley. Statistical methods in Markov chains. *The Annals of Mathematical Statistics*, pages 12–40, 1961.

[CB19] Yeshwanth Cherapanamjeri and Peter L Bartlett. Testing symmetric Markov chains without hitting. In *Conference on Learning Theory*, pages 758–785. PMLR, 2019.

[CK82] Imre Csiszár and János Körner. *Information Theory: Coding Theorems for Discrete Memoryless Systems*. Academic Press, Inc., 1982.

[CS00] Imre Csiszár and Paul C Shields. The consistency of the BIC Markov order estimator. *The Annals of Statistics*, 28(6):1601–1619, 2000.

[CS04] I Csiszár and PC Shields. Information theory and statistics: a tutorial. *Foundations and Trends in Communications and Information Theory*, 1(4):417–527, 2004.

[CT06] Thomas M. Cover and Joy A. Thomas. *Elements of information theory, 2nd Ed.* Wiley-Interscience, New York, NY, USA, 2006.

[Dav73] L. Davisson. Universal noiseless coding. *IEEE Transactions on Information Theory*, 19(6):783–795, 1973.

[Dav83] L. Davisson. Minimax noiseless universal coding for Markov sources. *IEEE Transactions on Information Theory*, 29(2):211–215, 1983.

[DDG18] Constantinos Daskalakis, Nishanth Dikkala, and Nick Gravin. Testing symmetric Markov chains from a single trajectory. In *Conference On Learning Theory*, pages 385–409. PMLR, 2018.

[DMM+19] Sarah Dean, Horia Mania, Nikolai Matni, Benjamin Recht, and Stephen Tu. On the sample complexity of the linear quadratic regulator. *Foundations of Computational Mathematics*, pages 1–47, 2019.

[DMPW81] L Davisson, R McEliece, M Pursley, and Mark Wallace. Efficient universal noiseless source codes. *IEEE Transactions on Information Theory*, 27(3):269–279, 1981.

[Fan53] Ky Fan. Minimax theorems. *Proceedings of the National Academy of Sciences*, 39(1):42–47, 1953.

[FOPS16] M. Falahatgar, A. Orlitsky, V. Pichapati, and A.T. Suresh. Learning Markov distributions: Does estimation trump compression? In *2016 IEEE International Symposium on Information Theory (ISIT)*, pages 2689–2693. IEEE, July 2016.

[FW21] Sela Fried and Geoffrey Wolfer. Identity testing of reversible Markov chains. *arXiv preprint arXiv:2105.06347*, 2021.

[GR20] F Richard Guo and Thomas S Richardson. Chernoff-type concentration of empirical probabilities in relative entropy. *IEEE Transactions on Information Theory*, 67(1):549–558, 2020.

[HJL+18] Y. Han, J. Jiao, C.Z. Lee, T. Weissman, Y. Wu, and T. Yu. Entropy rate estimation for Markov chains with large state space. In *Advances in Neural Information Processing Systems*, pages 9781–9792, 2018.

[HKL+19] Daniel Hsu, Aryeh Kontorovich, David A Levin, Yuval Peres, Csaba Szepesvári, and Geoffrey Wolfer. Mixing time estimation in reversible Markov chains from a single sample path. *Annals of Applied Probability*, 29(4):2439–2480, 2019.

[HOP18] Yi Hao, A. Orlitsky, and V. Pichapati. On learning Markov chains. *In Advances in Neural Information Processing Systems*, pages 648–657, 2018.

[JS02] Philippe Jacquet and Wojciech Szpankowski. A combinatorial problem arising in information theory: Precise minimax redundancy for Markov sources. In *Mathematics and Computer Science II*, pages 311–328. Springer, 2002.

[Kem74] JHB Kemperman. On the Shannon capacity of an arbitrary channel. In *Indagationes Mathematicae (Proceedings)*, volume 77, pages 101–115. North-Holland, 1974.

[KOPS15] S. Kamath, A. Orlitsky, D. Pichapati, and A.T. Suresh. On learning distributions from their samples. In *Conference on Learning Theory*, pages 1066–1100, June 2015.

[KV16] Sudeep Kamath and Sergio Verdú. Estimation of entropy rate and Rényi entropy rate for Markov chains. In *Information Theory (ISIT), 2016 IEEE International Symposium on*, pages 685–689. IEEE, 2016.

[LB04] Feng Liang and Andrew Barron. Exact minimax strategies for predictive density estimation, data compression, and model selection. *IEEE Transactions on Information Theory*, 50(11):2708–2726, 2004.

[Lez98] P. Lezaud. Chernoff-type bound for finite Markov chains. *Annals of Applied Probability*, 8(3):849–867, 1998.

[LP17] D.A. Levin and Y. Peres. *Markov chains and mixing times*, volume 107. American Mathematical Soc., 2017.

[MJT+20] Jay Mardia, Jiantao Jiao, Ervin Tánczos, Robert D Nowak, and Tsachy Weissman. Concentration inequalities for the empirical distribution of discrete distributions: beyond the method of types. *Information and Inference: A Journal of the IMA*, 9(4):813–850, 2020.

[OS20] Maciej Obremski and Maciej Skorski. Complexity of estimating Rényi entropy of Markov chains. In *2020 IEEE International Symposium on Information Theory (ISIT)*, pages 2264–2269. IEEE, 2020.

[Pan04] Liam Paninski. Variational minimax estimation of discrete distributions under KL loss. *Advances in Neural Information Processing Systems*, 17:1033–1040, 2004.

[Pau15] D. Paulin. Concentration inequalities for Markov chains by Marton couplings and spectral methods. *Electronic Journal of Probability*, pages 1–20, 2015.

[Ris84] Jorma Rissanen. Universal coding, information, prediction, and estimation. *IEEE Transactions on Information theory*, 30(4):629–636, 1984.

[Rya88] B.Y. Ryabko. Prediction of random sequences and universal coding. *Prob. Pered. Inf.*, 24(2):87–96, 1988.

[Sht87] Yuri M Shtarkov. Universal sequential coding of single messages. *Prob. Pered. Inf.*, 23:175–186, 1987.

[Sin64] Richard Sinkhorn. A relationship between arbitrary positive matrices and doubly stochastic matrices. *The Annals of Mathematical Statistics*, 35(2):876–879, 1964.

[SMT$^+$18] Max Simchowitz, Horia Mania, Stephen Tu, Michael I Jordan, and Benjamin Recht. Learning without mixing: Towards a sharp analysis of linear system identification. In *Conference On Learning Theory*, pages 439–473. PMLR, 2018.

[SW12] Wojciech Szpankowski and Marcelo J Weinberger. Minimax pointwise redundancy for memoryless models over large alphabets. *IEEE transactions on information theory*, 58(7):4094–4104, 2012.

[TJW18] Kedar Tatwawadi, Jiantao Jiao, and Tsachy Weissman. Minimax redundancy for Markov chains with large state space. In *2018 IEEE International Symposium on Information Theory (ISIT)*, pages 216–220. IEEE, 2018.

[Tro74] Viktor Kupriyanovich Trofimov. Redundancy of universal coding of arbitrary Markov sources. *Prob. Pered. Inf.*, 10(4):16–24, 1974.

[Whi55] P. Whittle. Some distribution and moment formulae for the Markov chain. *Journal of the Royal Statistical Society: Series B (Methodological)*, 17(2):235–242, 1955.

[WK19] Geoffrey Wolfer and Aryeh Kontorovich. Minimax learning of ergodic Markov chains. In *Algorithmic Learning Theory*, pages 904–930. PMLR, 2019.

[WK20] Geoffrey Wolfer and Aryeh Kontorovich. Minimax testing of identity to a reference ergodic Markov chain. In *International Conference on Artificial Intelligence and Statistics*, pages 191–201. PMLR, 2020.

[XB97] Qun Xie and Andrew R Barron. Minimax redundancy for the class of memoryless sources. *IEEE Transactions on Information Theory*, 43(2):646–657, 1997.

[YB99] Y. Yang and A. R. Barron. Information-theoretic determination of minimax rates of convergence. *The Annals of Statistics*, 27(5):1564–1599, 1999.