Mean Estimation with User-level Privacy under Data Heterogeneity

Rachel Cummings[*]
Department of Industrial Engineering and Operations Research
Columbia University
New York, NY 10027
`rac2239@columbia.edu`

Vitaly Feldman
Apple
Cupertino, CA 95014

Audra McMillan
Apple
Cupertino, CA 95014
`audra.mcmillan@apple.com`

Kunal Talwar
Apple
Cupertino, CA 95014
`ktalwar@apple.com`

Abstract

A key challenge in many modern data analysis tasks is that user data is heterogeneous. Different users may possess vastly different numbers of data points. More importantly, it cannot be assumed that all users sample from the same underlying distribution. This is true, for example in language data, where different speech styles result in data heterogeneity. In this work we propose a simple model of heterogeneous user data that differs in both distribution and quantity of data, and we provide a method for estimating the population-level mean while preserving user-level differential privacy. We demonstrate asymptotic optimality of our estimator and also prove general lower bounds on the error achievable in our problem.

1 Introduction

Many practical problems in statistical data analysis and machine learning deal with the setting in which each user generates multiple data points. In such settings the distribution of each user's data may be (somewhat) different and, furthermore, users may possess vastly different numbers of samples. This issue is one the key challenges in federated learning [20] leading to considerable interest in models and algorithms that address the issue.

As an example, consider the task of next-word prediction for a keyboard. Different users typing on a keyboard may have different styles of writing, leading to different distributions. There are aspects of the language that are common to all users, and likely additional aspects of style that are common to large groups of users. Thus while each user has their own data distribution, there are commonalities to all the distributions, and additional commonalities amongst distributions corresponding to subsets of users. Modeling and learning such relationships between users' distributions is crucial for building a better global model, as well as for personalizing models for users.

The focus of this work is on differentially private algorithms for such settings. We assume that there is an unknown global meta-distribution $\mathcal{D}$. For each user i, a personal data distribution $\mathcal{D}_i$ is chosen randomly from $\mathcal{D}$ (for example, by sampling a set of parameters that define $\mathcal{D}_i$). Each user then receives some number k_i of i.i.d. samples from $\mathcal{D}_i$. The goal is to solve an analysis task relative to $\mathcal{D}$, with an eye towards better modeling of each $\mathcal{D}_i$ even when k_i is small. This abstract setting

---

[*]Part of this work was completed while the author was at Apple.

36th Conference on Neural Information Processing Systems (NeurIPS 2022).

can model many practical settings where the relationships between the $\mathcal{D}_i$'s take different forms. Indeed the standard loss in federated learning is the unweighted average over users of a per-user loss function [20, Sec. 3.3.2], which corresponds to learning $\mathcal{D}$. Little theoretical work has been done in this setting and even the most basic statistical tasks are poorly understood. Thus we start by focusing on the fundamental problem of mean estimation in this setting. Specifically, in our model, $\mathcal{D}$ is a distribution on the interval $[0, 1]$ with unknown mean p and unknown variance σ_p^2. Further, we assume that $\mathcal{D}_i$ is simply a Bernoulli distribution with mean $p_i \sim \mathcal{D}$.

While the general $\mathcal{D}_i$ setting is of interest, there are many settings where users generate Boolean signals. For example, each sample from the Bernoulli distribution could represent whether or not the user has clicked on an ad. Another common example is model evaluation, where the user produces a Bernoulli sample by engaging or not engaging with a feature (e.g., phone keyboard next word suggestion, crisis helpline link, search engine knowledge panels, sponsored link in search results, etc.). As a concrete example, a language model is used to make the next word suggestions on a phone keyboard. A new version of this model would be first tested to measure the average suggestion acceptance rate over users. Each user would thus generate a set of independent Bernoulli r.v.'s with each individual mean p_i corresponding to the model accuracy for the user. Heterogeneity comes from different users typing differently (and hence model accuracy varying across users) and using the keyboard with different frequency. Note that the distribution of model accuracies among users is the meta distribution $\mathcal{D}$ in our work. More generally, measuring the average accuracy of a classification model among a large group of users is an important task in itself. Such models are deployed in privacy-sensitive applications such as health and finance. The resulting statistics may need to be shared with third parties or other teams within a company, raising potential user privacy concerns.

Our main contribution is a differentially private algorithm that estimates p and σ_p in this setting. We first study this question in an idealized setting with known σ_p and no privacy constraints. Here the optimal non-private estimator for p_i is simple and linear: it is a weighted linear combination of the individual user means with weights that depend on the k_i's and on σ_p. The variance of this estimate is $\sigma_{ideal}^2 \approx (\sum_i \min(k_i, \sigma_p^{-2}))^{-1}$. This expression has a natural interpretation: this is the variance from using $\min(k_i, \sigma_p^{-2})$ samples from user i and averaging all the Bernoulli samples thus obtained. The restriction on using at most σ_p^{-2} samples from each user ensures that the estimator is not too affected by their individual mean p_i.

Even in the case where it is known that $\sigma_p^2 = 0$, the solution is non-trivial and, to the best of our knowledge, no optimal private algorithm was previously known. In this case, each user samples from the same distribution, but there may be deviations in the number of samples that each user holds. In the absence of privacy constraints, this setting poses no additional complexity over the case where each user has a single data point, since the data points all come from the same distribution. However, with the requirement of user-level differential privacy, additional care needs to be taken to hide *all* samples from any individual user. In this case, we already need to employ many of the technical tools developed in this work, as we show in Appendix C.

We show that under mild assumptions, there is no asymptotic price to privacy (and to not knowing σ_p). We provide a differentially private estimator for p with variance $O(\sigma_{ideal}^2)$. Interestingly, the estimator achieving this bound in the private setting is non-linear. Further, we show that σ_{ideal}^2 is near-optimal, under some mild technical conditions.

Our technical results highlight several of the challenges associated with ensuring user-level privacy when data is heterogeneous. For example, the optimal choice of weights for each user contribution itself depends on p and σ_p that we are trying to estimate. Further, we show a novel approach to proving lower bounds for private statistical estimation in the heterogenous setting. Our approach builds on the proof of the Cramér-Rao lower bound in statistics, and we show how privacy terms can be incorporated in this approach to show near optimality of our algorithms for nearly every setting of k_i's. These tools and insights should be useful for modeling and designing algorithms for more involved data analysis tasks.

Our work lays the foundation for similar model-driven exploration in other settings. There have been attempts to handle heterogeneity by phrasing the problem as meta-learning or multi-task learning [20, Sec 3.3.3], which implicitly makes some assumptions about the different distributions. Our goal is to start with a more principled approach that makes explicit the assumptions on the relationship between different distributions and use that to derive the algorithm. For example, if were to model the D_i's

as having means coming from a mixture of Gaussians, the estimation of cluster means would be a necessary step in an EM-type algorithm. Our choice of $\mathcal{D}_i$'s being Bernoulli is meant to capture discrete distribution learning problems that have been extensively studied in private federated settings. Our techniques are general and would extend naturally to real-valued settings where, e.g., $\mathcal{D}_i$ is a Gaussian with mean p_i and known variance. While we make minimal assumptions on $\mathcal{D}$, our results asymptotically match the lower bounds for the case of $\mathcal{D}$ being Gaussian with known variance. Our techniques also extend in natural ways to higher dimensions.

Our main results involve two estimators; a non-realisable estimator $\widehat{p}_\epsilon^{\text{ideal}}$ that assumes that the mean and variance of $\mathcal{D}$ are known to the estimator, and a realisable estimator $\widehat{p}_\epsilon^{\text{realistic}}$ that is private with respect to the user's samples, but not with respect to each user's number of samples k_i. Let $\widehat{p}_i$ be the mean of the k_i samples from user i. The estimator $\widehat{p}_\epsilon^{\text{realistic}}$ requires as input initial, less accurate (ϵ, δ)-DP mean and variance estimators $\texttt{mean}_{\epsilon,\delta}$ and $\texttt{variance}_{\epsilon,\delta}$. The main results of this paper can be (informally) summarised as follows:

- **Near optimality of $\widehat{p}_\epsilon^{\text{ideal}}$ [Theorem 5.1].** For any parameterized family of distributions $p \mapsto \mathcal{D}_p$, if the Fisher information of $\widehat{p}_i$ is inversely proportional to the variance of $\widehat{p}_i$ for all i, and each $\widehat{p}_i$ is sufficiently well concentrated (sub-Gaussian is sufficient) then $\widehat{p}_\epsilon^{\text{ideal}}$ is minimax optimal, up to logarithmic factors, among all unbiased estimators of p in the range $p \in [1/3, 2/3]$. The proof of this result involves a Cramér-Rao style proof which may be of independent interest.

- **Near optimality of $\widehat{p}_\epsilon^{\text{realistic}}$ [Theorem 4.1].** Under mild conditions on the accuracy of $\texttt{mean}_{\epsilon,\delta}$ and $\texttt{variance}_{\epsilon,\delta}$, and assuming the $\max$ and and median k_i are within a factor of $(n\epsilon/\log n) - 1$, then $\text{Var}(\widehat{p}_\epsilon^{\text{realistic}}) = O(\text{Var}(\widehat{p}_\epsilon^{\text{ideal}}))$.

- **Lower bound in terms of k_i [Corollary 5.5].** We give an explicit formula for the minimax optimal error in terms of the sequence $k_1, \cdots, k_n$ and variance σ_p^2.

Our main algorithmic results require concentration of the meta-distribution $\mathcal{D}$. We note that in practice, this is not an unreasonable assumption. For example, in the case of model evaluation, it may be be reasonable to assume that a general model has similar accuracy for the vast majority of users, or formally, that the model accuracy is well-concentrated.

1.1 Related Work

Frequency estimation in the example-level privacy model has been well-studied in the central [12, 11] and local models [18, 14, 7, 1, 2]. Similarly, private mean estimation has been well studied in both central [12, 16] and local models [9, 8, 5] of privacy. These works have focused on providing example-level privacy (rather than user-level) in settings with homogeneous data, i.e., i.i.d. samples.

[23] recently studied the problem of learning discrete distributions in the homogeneous cases (same distribution and same number of samples per user) with user-level differential privacy, and [22] extended such results to other statistical tasks. These works also consider the setting with different number of samples per user although only via a reduction to same number of samples by discarding the data of users that have less than the median number of samples and effectively only using the median number of samples from all the other users. This approach can be asymptotically suboptimal for many natural distributions of k_i's and is also likely to be worse in practice. Previously, [27] showed how to build a (user-level) differentially private recommendation system, and [25] showed how to train a language model with user-level differential privacy.

User-level differential privacy in the context of heterogeneous data distributions has been studied in the constant k_i setting [31]. Much of the complexity in our setting arises from variation in the k_i values, which makes it challenging to maintain user-level privacy while leveraging the additional data points from users with a large number of data points. The challenges to optimization due to data heterogeneity have also been studied; [34, 15], and Eichner et al. [13] study the approach of using different models for different groups from a convex optimization point-of-view. Mathematically, similar issues are addressed in meta-analysis [6, 33], where the heterogeneity comes from different studies instead of different users. The non-private approach of inverse variance weighting that we recap in Section 3 is standard in that context.

2 Model and Preliminaries

Let $\mathcal{D}$ be a distribution on $[0,1]$ with (unknown) mean p and variance σ_p^2. We assume a population of $n \in \mathbb{N}$ users, where each user $i \in [n]$ has a hidden variable $p_i \sim \mathcal{D}$ and $k_i \in \mathbb{N}$ samples $x_i^1, \ldots, x_i^{k_i} \sim_{i.i.d.} Ber(p_i)$. That is, the samples of user i are i.i.d. from a Bernoulli distribution with parameter p_i, which we will denote $\mathcal{D}_i = Ber(p_i)$. Assume without loss of generality that individuals are sorted by their k_i, so that $k_1 \geq \cdots \geq k_n$. The samples x_i^j and hidden variables p_i of each user are unknown to the analyst, and we start with assuming that the k_i's are publicly known. In Appendix D, we extend our results to the general case where the k_i's are also private.

The analyst's goal is to estimate the population mean p with an estimator of minimum variance in a manner that is differentially private with respect to user data (p_i and $\{x_i^j\}$). Each user provides their own estimate of their p_i to the analyst based on their data x_i: $\widehat{p}_i = \frac{1}{k_i} \sum_{j=1}^{k_i} x_i^j$. The analyst can then aggregate these (possibly along with other information) into her estimate of p.

Let us first give some intuition for the distribution of these $\widehat{p}_i$. Let $\mathcal{D}(k)$ be the distribution that first samples $p_i \sim \mathcal{D}$, then samples $x_1, \cdots, x_k \sim Ber(p_i)$ and finally outputs $\widehat{p}_i = \frac{1}{k} \sum_{i=1}^k x_i$. The following lemma (proven in Appendix A) shows that the variance of $\widehat{p}_i$ is larger than σ_p^2 and transitions from $p(1-p)$ to σ_p^2 as k increases (equivalently as $\widehat{p}_i$ concentrates around p_i).

Lemma 2.1. *For all distributions $\mathcal{D}$ supported on $[0,1]$ with mean p and variance σ_p^2, $\sigma_p^2 \leq p(1-p)$. Further, $\mathbb{E}[\mathcal{D}(k)] = p$ and $\mathrm{Var}(\mathcal{D}(k)) = \frac{1}{k}p(1-p) + \left(1 - \frac{1}{k}\right)\sigma_p^2$.*

We assume that k_i and p_i are independent, so the amount of data an individual has is independent of her data distribution. This is crucial for the problem setup: in order for learning from the heterogeneous population to be advantageous, there must a common meta-distribution is shared across all individuals in the population, rather than a meta-distribution only for each fixed k_i.

2.1 Differential Privacy

Differential privacy (DP) [12] informally limits the inferences that can be made about an individual as a result of computations on a large dataset containing their data. The definition of DP requires a pairwise *neighbouring relation* between datasets, and DP algorithms ensure that differences between all pairs of neighboring datasets should be hidden by the private algorithm.

In our setting where users have multiple data points, we must distinguish between *user-level* and *event-level* DP. The former considers D and D' neighbours if they differ on all data points associated with a single user, whereas the latter considers D and D' neighbours only if they differ on a *single* data point, regardless of the number of data points contributed by that user. Naturally, user-level DP provides substantially stronger privacy guarantees, and is often more challenging to achieve from a technical perspective. In this work, we will provide user-level DP guarantees.

Definition 2.2 (User-level (ϵ, δ)-Differential Privacy [12]). *Given $\epsilon \geq 0$, $\delta \in [0,1]$ and a neighbouring relation $\sim$, a randomized mechanism $\mathcal{M} : \mathcal{X}^{k_1} \times \cdots \times \mathcal{X}^{k_n} \to \mathcal{Y}$ is (ϵ, δ)-differentially private if for all neighboring datasets $D \sim D' \in \mathcal{X}^{k_1} \times \cdots \times \mathcal{X}^{k_n}$, and all events $E \subseteq \mathcal{Y}$, $\Pr[\mathcal{M}(D) \in E] \leq e^\epsilon \cdot \Pr[\mathcal{M}(D') \in E] + \delta$, where the probabilities are taken over the random coins of $\mathcal{M}$. When $\delta = 0$, we refer to this as ϵ-differential privacy.*

One standard tool for achieving ϵ-differential privacy is the *Laplace Mechanism*. For a given function f to be evaluated on a dataset D, the Laplace Mechanism first computes $f(D)$ and then adds Laplace noise which depends on the *sensitivity* of f, defined for real-valued functions as $\Delta f = \max_{D, D' \text{ neighbors}} |f(D) - f(D')|$. The Laplace Mechanism outputs $\mathcal{M}_L(D, f, \epsilon) = f(D) + Lap(\Delta f / \epsilon)$ and is $(\epsilon, 0)$-DP.

3 A Non-Private Estimator

We begin by illustrating the procedure for computing an optimal estimator $\widehat{p}$ in the non-private setting. The general structure of the estimator will be the same in both settings. The analyst will compute the

Algorithm 1 Non-private Heterogeneous Mean Estimation

Input: number of users n, number of samples held by each user $(k_1, \ldots, k_n)$ $(k_i \geq k_{i+1})$, user-level estimates $(\widehat{p}_1, \cdots, \widehat{p}_n)$.

1: **Initial Estimates**
2: $\widehat{p}^{\text{initial}} = \sum_{i=9n/10}^{n} x_i^1$
3: $\widehat{\sigma}_p^2 = \frac{1}{\log n (\log n - 1)} \sum_{i,j \in [\log n]} (\widehat{p}_i - \widehat{p}_j)^2$

4: **Defining weights**
5: **for** $i = \log n$ to $9n/10$ **do**
6:     Compute $\widehat{\sigma}_i^2 = \frac{1}{k_i}(\widehat{p}^{\text{initial}} - (\widehat{p}^{\text{initial}})^2) + (1 - \frac{1}{k_i})\widehat{\sigma}_p^2$.
7:     $\widehat{w}_i = \frac{1/\widehat{\sigma}_i^2}{\sum_{j=\log n}^{9n/10} 1/\widehat{\sigma}_j^2}$

8: **Final Estimate**
9: **return** $\widehat{p}^{\text{realistic}} = \sum_{i=\log n}^{n} \widehat{w}_i \widehat{p}_i$

---

population-level mean estimate $\widehat{p}$ as a weighted linear combination of the user-level estimates $\widehat{p}_i$.[2] The key question is how to derive the weights so that individuals with more reliable estimates (i.e., larger k_i) have more influence over the final result.

Let σ_i^2 be the variance of $\widehat{p}_i$. In an idealized setting where the σ_i^2 are all known, the analyst can minimize the variance of the estimator by weighting each user's estimate $\widehat{p}_i$ proportionally to the inverse variance of their estimate. The weights are normalised to ensure the estimate is unbiased. This approach yields the following estimator, which is optimal in the non-private setting [17]:

$$\widehat{p}^{\text{ideal}} = \sum_{i=1}^{n} w_i^* \widehat{p}_i \text{ where } w_i^* = \frac{1/\sigma_i^2}{\sum_{j=1}^{n} 1/\sigma_j^2}. \tag{1}$$

In practice, the σ_i^2s are unknown, so the analyst must rely on estimates to assign weights. Fortunately, the user-level variance σ_i^2 can be expressed as a function of k_i and the population statistics p and σ_p^2, as shown in Lemma 2.1:

$$\sigma_i^2 = \frac{1}{k_i}(p - p^2) + (1 - \frac{1}{k_i})\sigma_p^2. \tag{2}$$

Now, p and σ_p^2 are also unknown but since they are population statistics, we can use simple estimators to obtain initial estimates. These initial statistics can then be used to define the weights, resulting in a refined estimate of the mean p. Specifically, as outlined in Algorithm 1, we split users into three groups. The $\log n$ individuals with the most data are used to produce an estimate of $\text{Var}(\mathcal{D}(k_{\log n}))$, which serves as a proxy for σ_p^2. The $1/10$th of individuals with the least data are used to produce an initial estimate of the mean p. The remaining $9n/10 - \log n$ individuals are used to produce the final estimate. We split the individuals into separate groups to ensure the initial estimates and the final estimate are independent so we can easily obtain variance bounds on the final estimate. The specific sizes of the three groups are heuristic, the exact fraction $1/10$ is not necessary. Under some mild conditions on $\mathcal{D}$, and if n is large enough, the error incurred by $\widehat{p}^{\text{realistic}}$ is within a constant factor of the error incurred by the ideal estimator $\widehat{p}^{\text{ideal}}$.[3]

4   A Framework for Private Estimators

We now turn to our main result, which is a framework for designing differentially private estimators for the mean p of the meta-distribution $\mathcal{D}$. We discussed in Section 3 the need for initial estimates of p and σ_p^2 to weight the contributions of the users. In the non-private setting, there are canonical, optimal choices of these estimators; the empirical mean and empirical variance. In the private setting, these choices are not canonical, and different estimators may perform better in different settings. There is a considerable literature exploring various mean and variance estimators for the homogeneous,

---

[2]In the non-private setting, this restriction is without loss of generality since the optimal estimator takes this form. In the private setting this is still near-optimal; see Section 5 for more details.

[3]This can be observed by viewing the non-private setting as a simplified version of the setting studied in Section 5, which proves near-optimality of (truncated) linear estimators for this problem.

single-data-point-per-user setting. As such, we leave the choice of the specific initial mean and variance estimators as parameters of the framework. This allows us to focus on the nuances of the heterogeneous setting, not addressed in prior work. In Appendix F, we give a specific pair of private mean and variance estimators that provably perform well in our framework.

As in the previous section, we will define two estimators: a ideal estimator $\widehat{p}_\epsilon^{\text{ideal}}$ (only implementable if all the σ_i^2 are known), and a realisable estimator $\widehat{p}_\epsilon^{\text{realistic}}$. The main result in this section (Theorem 4.1) is that under some mild conditions and assuming n is sufficiently large, there exists an (ϵ, δ)-DP estimator $\widehat{p}_\epsilon^{\text{realistic}}$ (Algorithm 2) such that for some constant C, $\text{Var}(\widehat{p}_\epsilon^{\text{realistic}}) \leq C \cdot \text{Var}(\widehat{p}_\epsilon^{\text{ideal}})$.

4.1 The Complete Information Private Estimator

As in Section 3, we begin with a discussion of the ideal estimator if the σ_i were known. This ideal estimator $\widehat{p}_\epsilon^{\text{ideal}}$ has a similar form to $\widehat{p}^{\text{ideal}}$ with some crucial differences. The first main distinction is that Laplace noise is added to achieve DP, where the standard deviation of the noise must be scaled to the sensitivity of the statistic. A natural solution would be to add noise directly to the non-private estimator $\widehat{p}^{\text{ideal}}$, but the sensitivity of this statistic is too high. In fact, the worst case sensitivity of $\widehat{p}^{\text{ideal}}$ is 1, which would result in the noise that completely masks the signal. Thus, the first change we make is to limit the weight of any individual's contribution by setting $w_i = \frac{\min\{1/\sigma_i^2, T/\sigma_i\}}{\sum_{j=1}^n \min\{1/\sigma_j^2, T/\sigma_j\}}$ for some truncation parameter T. Intuitively, the parameter T controls the trade-off between variance of the weighted sum of individual estimates (which is minimized by assigning high weight to low variance estimators) and variance of the noise added for privacy (which is minimized by assigning roughly equal weight to all users).

We make one final modification to lower the sensitivity of the statistic. Inspired by the Gaussian mean estimator of [21], we truncate the individual contributions $\widehat{p}_i$ into a sub-interval of $[0, 1]$. The truncation intervals $[a_i, b_i]$ are chosen to be as small as possible (to reduce the sensitivity and hence the noise added for privacy), while simultaneously ensuring that $\widehat{p}_i \in [a_i, b_i]$ with high probability (to avoid truncating relevant information for the estimation). In order to achieve this, we need a tail bound on the distribution $\mathcal{D}$. To maintain generality for now, we assume there exists a known function $f_{\mathcal{D}}^k(n, \sigma_p^2, \beta)$ that gives high-probability concentration guarantees of $\widehat{p}_i$ around p, and is defined such that $\Pr\left(\forall i, |\widehat{p}_i - p| \leq f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)\right) \geq 1 - \beta$. Appendix G presents a more detailed discussion of the structure of these concentration functions and how they may be estimated if they are unknown to the analyst.

We can now describe the full information, or *ideal*, estimator $\widehat{p}_\epsilon^{\text{ideal}}$:

$$\widehat{p}_\epsilon^{\text{ideal}} = \sum_{i=1}^n w_i^* [\widehat{p}_i]_{a_i}^{b_i} + \text{Lap}(\frac{\max_i w_i^* |b_i - a_i|}{\epsilon}), \tag{3}$$

where $[\widehat{p}_i]_{a_i}^{b_i}$ denotes the projection of $\widehat{p}_i$ onto the interval $[a_i, b_i]$ and

$$a_i = p - f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta), \quad b_i = p + f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta), \text{ and } w_i^* = \frac{\min\{1/\sigma_i^2, T^*/\sigma_i\}}{\sum_{j=1}^n \min\{1/\sigma_j^2, T^*/\sigma_j\}}. \tag{4}$$

We would like to choose the truncation parameter T^* to minimise the variance of the resulting estimator:

$$\text{Var}(\widehat{p}_\epsilon^{\text{ideal}}) = \sum_{i=1}^n (w_i^*)^2 \text{Var}([\widehat{p}_i]_{a_i}^{b_i}) + \max_i \frac{(w_i^*)^2 |b_i - a_i|^2}{\epsilon^2}. \tag{5}$$

Although we do not know $\text{Var}([\widehat{p}_i]_{a_i}^{b_i})$ exactly, we do know that $[\widehat{p}_i]_{a_i}^{b_i} = \widehat{p}_i$ with high probability, and thus we can approximate $\text{Var}([\widehat{p}_i]_{a_i}^{b_i})$ with σ_i. Throughout the remainder of the paper, we will assume that β is chosen such that $\frac{1}{2}\sigma_i^2 \leq \text{Var}([\widehat{p}_i]_{a_i}^{b_i})$. Thus, we will approximate the optimal truncation parameter by

$$T^* = \arg\min_T \sum_{i=1}^n (w_i^*)^2 \sigma_i^2 + \max_i \frac{(w_i^*)^2 |b_i - a_i|^2}{\epsilon^2}$$

$$= \arg\min_T \frac{1}{(\sum_{j=1}^n \min\{1/\sigma_j^2, T/\sigma_i\})^2} (\sum_{i=1}^n \min\{1/\sigma_i^2, T^2\} + \max_i \frac{\min\{1/\sigma_i^4, T^2/\sigma_i^2\} |b_i - a_i|^2}{\epsilon^2}). \tag{6}$$

We'll show in Section 5 that under some conditions on the Fisher information of $\mathcal{D}(k)$, $\widehat{p}_\epsilon^{\text{ideal}}$ is optimal up to logarithmic factors among all private unbiased estimators for heterogeneous mean estimation.

Example 1. *As a simple example, suppose that $p \in (\frac{1}{3}, \frac{2}{3})$, $\sigma_p = 1/\sqrt{n}$, and $k_i = \lceil \frac{n}{i} \rceil$. In this case, an asymptotically optimal non-private estimator averages all the $\sum k_i = O(n \log n)$ available samples. It can be shown that this gives us an unbiased estimator with standard deviation $\Theta(\frac{1}{\sqrt{n \log n}})$. A naive sensitivity-based noise addition method will give us privacy error $O(\frac{1}{\varepsilon \log n})$, since the weight of the first user in this average is $\Theta(1/\log n)$. Our truncation-based algorithm will truncate the ith user's contribution to a range of width $\sqrt{\frac{\log n}{k_i}} \approx \sqrt{\frac{i \log n}{n}}$. Applying our algorithm would then give us privacy error $\Theta(\frac{1}{\varepsilon \sqrt{n \log n}})$. In other words, for constant ε, privacy does not have an asymptotic cost. We remark that in this case, any uniform weighted average will incur asymptotically larger standard deviation $\Omega(\frac{1}{\sqrt{n}})$.*

4.2 Realizable Private Heterogeneous Mean Estimation

Our goal in this section is to design a realizable estimator $\widehat{p}_\epsilon^{\text{realistic}}$ that is competitive with the ideal estimator $\widehat{p}_\epsilon^{\text{ideal}}$. As in the non-private setting, we divide the individuals into three groups. The first group, consisting of the $n/10$ individuals with the lowest k_i will be used to compute the initial mean estimate $\widehat{p}_\epsilon^{\text{initial}}$. The $\log n$ individuals with the largest k_i will be used to compute the initial variance estimate $\widehat{\sigma}_p^2$. These initial estimates will be plugged into expressions to compute $\widehat{\sigma}_i^2$, $\widehat{a}_i$, and $\widehat{b}_i$ for the remaining individuals $\log n + 1 \leq i \leq 9n/10$. As in the non-private setting, the specific sizes of these groups are heuristic. The important thing is that the size of the first two groups are large enough that the resulting mean and variance estimates are sufficiently accurate, and the last group contains $\Theta(n)$-users whose k_i is above the median.

Since the estimate $\widehat{p}_\epsilon^{\text{initial}}$ used in $\widehat{a}_i$ and $\widehat{b}_i$ may had additional error up to α, we shift these estimates by an additive α to account for this error. Next, all of these intermediate estimates and the user-level mean estimates $\widehat{p}_i$ from users $\log n + 1 \leq i \leq 9n/10$ will be used to compute the optimal weight cutoff $\widehat{T}^*$, the optimal weights $\widehat{w}_i^*$ for each user $\log n + 1 \leq i \leq 9n/10$, and finally the estimator $\widehat{p}_\epsilon^{\text{realistic}}$ as a weighted sum of the truncated user-level estimates $[\widehat{p}_i]_{\widehat{a}_i}^{\widehat{b}_i}$ plus Laplace noise. This procedure is presented in full detail in Algorithm 2.

For the remainder of this section, we turn to establishing the accuracy requirements of $\texttt{mean}_{\epsilon,\delta}$ and $\texttt{variance}_{\epsilon,\delta}$ that ensure that the error of $\widehat{p}_\epsilon^{\text{realistic}}$ is within a constant factor of the error of $\widehat{p}_\epsilon^{\text{ideal}}$.

Theorem 4.1. *For any $\epsilon > 0$, $\delta \in [0,1]$, Algorithm 2 is (ϵ, δ)-DP. If,*

- *$\texttt{mean}_{\epsilon,\delta}$ is such that given $n/10$ samples from $\mathcal{D}$, with probability $1 - \beta$ $|p - \widehat{p}_\epsilon^{\text{initial}}| \leq f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$ and $\widehat{p}_\epsilon^{\text{initial}}(1 - \widehat{p}_\epsilon^{\text{initial}}) \in \left[\frac{1}{2}p(1-p), \frac{3}{2}p(1-p)\right]$,*

- *$\texttt{variance}_{\epsilon,\delta}$ is such that given $\log n$ samples from $\mathcal{D}(k)$, with probability $1 - \beta$, $\widehat{\sigma}_p^2 \in [\text{Var}(\mathcal{D}(k)), 8\text{Var}(\mathcal{D}(k))]$,*

- *the k_is are such that $\frac{k_1}{k_{n/2}} \leq \frac{n/2 - \log n}{\log n}$,*

then with probability $1 - 2\beta$, $\text{Var}(\widehat{p}_\epsilon^{\text{realistic}}) \leq C \cdot \text{Var}(\widehat{p}_\epsilon^{\text{ideal}})$ for some absolute constant C.

A full proof of Theorem 4.1 is given in Appendix B; we present intuition and a proof sketch here. The first two conditions of Theorem 4.1 ensure that the mean and variance estimates are sufficiently accurate to use in the remainder of the algorithm. Notice that the initial estimates do not need to be especially accurate. In fact, provided p is not too close to 0 or 1, the DP mean estimator that simply adds noise to the sample mean achieves the right accuracy (see Lemma F.1 for details). In Appendix F, we also give a DP variance estimator that achieves the desired accuracy guarantee using only $\log n$ samples, under some mild conditions (Lemma F.4). Thus the set of mean and variance estimators that satisfy the accuracy requirements of Theorem 4.1 are non-empty. We note that the constants $1/2$, $3/2$ and 8 in Theorem 4.1 are not intrinsic; any constant multiplicative factors will suffice. We also note that the specific sizes of the three groups outlined in Algorithm 2 are heuristic and can be varied to ensure that the initial estimator achieves the required accuracy.

Algorithm 2 Private Heterogeneous Mean Estimation

Input: (ϵ,δ)-DP mean estimator $\texttt{mean}_{\epsilon,\delta}$, (ϵ,δ)-DP variance estimator $\texttt{variance}_{\epsilon,\delta}$, number of users n, number of samples held by each user $(k_1,\ldots,k_n \ s.t. \ k_i \geq k_{i+1})$, user-level estimates $(\widehat{p}_1,\cdots,\widehat{p}_n)$, error guarantee on $\texttt{mean}_{\epsilon,\delta}$ $\alpha > 0$, and desired high probability bound $\beta \in [0,1]$.

1: **Initial Estimates**
2: $\widehat{p}_\epsilon^{\text{initial}} = \texttt{mean}_{\epsilon,\delta}(\mathrm{x}_{9n/10+1}^1,\cdots,\mathrm{x}_n^1)$
3: $\widehat{\sigma}_p^2 = \texttt{variance}_{\epsilon,\delta}(\widehat{p}_1,\cdots,\widehat{p}_{\log n})$

4: **Defining weights and truncation**
5: **for** $i = \log n + 1$ to $9n/10$ **do**
6: $\quad$ Compute $\widehat{\sigma}_i^2 = \frac{1}{k_i}(\widehat{p}_\epsilon^{\text{initial}} - (\widehat{p}_\epsilon^{\text{initial}})^2) + (1-\frac{1}{k_i})\widehat{\sigma}_p^2$.
7: $\quad$ $\widehat{a}_i = \widehat{p}_\epsilon^{\text{initial}} - \alpha - f_{\mathcal{D}}^{k_i}(n,\widehat{\sigma_p^2},\beta)$
8: $\quad$ $\widehat{b}_i = \widehat{p}_\epsilon^{\text{initial}} + \alpha + f_{\mathcal{D}}^{k_i}(n,\widehat{\sigma_p^2},\beta)$
9:
10: $\widehat{T}^* = \arg\min_T \dfrac{(\sum_{i=\log n+1}^{9n/10} \min\{\frac{1}{\widehat{\sigma}_i^2},T^2\}+\max_{\log n+1 \leq i \leq 9n/10} \frac{\min\{1/\widehat{\sigma}_i^4,T^2/\widehat{\sigma}_i^2\}|\widehat{b}_i-\widehat{a}_i|^2}{\epsilon^2})}{(\sum_{i=\log n+1}^{9n/10} \min\{1/\widehat{\sigma}_j^2,T/\widehat{\sigma}_i\})^2}$
11: **for** $i = \log n + 1$ to $9n/10$ **do**
12: $\quad$ $\widehat{w_i^*} = \dfrac{\min\{1/\widehat{\sigma}_i^2,\widehat{T}^*/\widehat{\sigma}_i\}}{\sum_{j=\log n+1}^{9n/10} \min\{1/\widehat{\sigma}_j^2,\widehat{T}^*/\widehat{\sigma}_i\}}$

13: **Final Estimate**
14: $\Lambda = \max_{i \in [\log n+1,9n/10]} \dfrac{\min\{1/\widehat{\sigma}_i^2,\widehat{T}^*/\widehat{\sigma}_i\}|\widehat{b}_i-\widehat{a}_i|}{\sum_{j=\log n+1}^{9n/10} \min\{1/\widehat{\sigma}_j^2,\widehat{T}^*/\widehat{\sigma}_i\}}$
15: Sample $Y \sim \text{Lap}\left(\frac{\Lambda}{\epsilon}\right)$
16: **return** $\widehat{p}_\epsilon^{\text{realistic}} = \sum_{i=\log n+1}^{9n/10} \widehat{w_i^*}[\widehat{p}_i]_{\widehat{a}_i}^{\widehat{b}_i} + Y$

The final assumption ensures that the $\log n$ users with the most data can not estimate the mean of meta-distribution alone. Note that up to logarithmic factors, this condition simply requires that the number of data points held by the user with the most data is at most n times the number of data points of the median user. If n is large, then this is unlikely to be a limiting factor.

The main distinction between $\widehat{p}_\epsilon^{\text{ideal}}$ and $\widehat{p}_\epsilon^{\text{realistic}}$ is the use of the output of the estimators $\texttt{mean}_{\epsilon,\delta}$ and $\texttt{variance}_{\epsilon,\delta}$ to estimate σ_i^2, a_i and b_i. Thus, the main component of the proof of Theorem 4.1 is to show that the conditions stated in the theorem are enough to ensure that $\widehat{\sigma}_i^2$, $\widehat{a}_i$ and $\widehat{b}_i$ are sufficiently accurate.

Lemma 4.2. *Given* $\widehat{p}_\epsilon^{\text{initial}}$, $\widehat{\sigma}_p^2$, *and* k_i, *define* $\widehat{\sigma}_i^2 = \frac{1}{k_i}\widehat{p}_\epsilon^{\text{initial}}(1 - \widehat{p}_\epsilon^{\text{initial}}) + \frac{k_i-1}{k_i}\widehat{\sigma}_p^2$. *Under the conditions of Theorem 4.1, for all* $i > \log n$, *we have* $\widehat{\sigma}_i^2 \in \left[\frac{1}{2}\sigma_i^2, 9.5\sigma_i^2\right]$ *and* $|\widehat{b}_i - \widehat{a}_i| \leq 4|b_i - a_i|$.

A detailed proof of Lemma 4.2 is presented in Appendix B. Lemma 4.2 implies that the individual variance estimates used in the weights, and the truncation parameters are accurate up to constant multiplicative factors. The main ingredient left then is to show that using only a subset of the population in the final estimate only affects the performance up to a multiplicative factor. Under the assumption that $\frac{k_{\max}}{k_{\text{med}}} \leq \frac{n/2-\log n}{\log n}$, where $\sigma_{k_{\max}}^2 = \text{Var}(\widehat{p}_1)$ and $\sigma_{k_{\text{med}}}^2 = \text{Var}(\widehat{p}_{n/2})$ then

$$\sigma_{k_{\text{med}}}^2 = \frac{1}{k_{\text{med}}}p(1-p) + (1 - \frac{1}{k_{\text{med}}})\sigma_p^2 \leq \frac{n/2-\log n}{\log n}\sigma_{k_{\max}}^2. \tag{7}$$

We use this to show that for any truncation parameter T, $\sum_{i=1}^n \min\{\frac{1}{\sigma_i^2},\frac{T}{\sigma_i}\} \leq 4\sum_{i=\log n+1}^{9n/10} \min\{\frac{1}{\sigma_i^2},\frac{T}{\sigma_i}\}$. Using this, along with the bounds on estimated quantities from Lemma 4.2, we show that with high probability, the variance of the our estimator $\widehat{p}_\epsilon^{\text{realistic}}$ is within a constant factor of $\text{Var}(\widehat{p}_\epsilon^{\text{ideal}})$, as given in Equation (5).

We remark that this framework is amenable to being performed in a federated manner if one has private federated mean and variance estimators. Steps (6) - (8) and Step (12) can be performed locally. Steps (10) and the final sum in Step (16) would need to be altered to fit the federated framework.

We'll see in Appendix D that it is sufficient to replace Step (10) with an estimate of $\frac{1}{\sigma_{\log n}}$ (the inverse standard deviation of the user with the $\log n$-th most data). The final step is then a simple addition with output perturbation, which can be performed in a federated manner (e.g., [24, 20]).

In Appendix D, we extend this result to the case where k_is are private and unknown to the analyst (Algorithm 3, Theorem D.1). We'll need considerably more machinery in this setting where both the sensitivity of the final estimator and the truncation parameter T are data dependent.

5 Near Optimality and Lower Bounds

In Section 4, we showed that the variance of our realisable private estimator $\widehat{p}_\epsilon^{\text{realistic}}$ is within a constant of that of the complete information estimator $\widehat{p}_\epsilon^{\text{ideal}}$. In this section, we will show that in fact, $\widehat{p}_\epsilon^{\text{realistic}}$ performs as well (up to logarithmic factors) as the true optimal private estimator. We'll also give a lower bound on the performance of the optimal estimator in terms of the k_i. This will give us some intuition into the types of distributions of k_i's that benefit from this refined analysis.

5.1 Minimax Optimality of $\widehat{p}_\epsilon^{\text{realistic}}$

The goal of this section is to show that the estimator $\widehat{p}_\epsilon^{\text{realistic}}$ discussed in Section 4.2 is minimax optimal up to logarithmic factors. In light of Theorem 4.1, it suffices to show that the estimator $\widehat{p}_\epsilon^{\text{ideal}}$ is minimax optimal up to logarithmic factors. Let $\mathcal{P}$ be a parameterized family of distributions $p \mapsto \mathcal{D}_p$, where $\mathbb{E}[\mathcal{D}_p] = p$ and $\mathcal{D}_p$ is supported on $[0, 1]$. For $p \in [0, 1]$ and $k \in \mathbb{N}$, let $\phi_{p,k}$ be the probability density function of $\mathcal{D}_p(k)$.

Our lower bound will show that the estimation error must consist of a statistical term and a privacy term. Such a lower bound thus must generalize a statistical lower bound. We will rely on the Cramer-Rao approach to proving statistical lower bounds; as we show, it is particularly amenable to incorporating a privacy term. This approach relates the variance of any unbiased estimator of the mean of a distribution to the inverse of the Fischer information; the proof naturally extends to the case where we are given samples from a set of distributions with the same mean but different variances, as is the case in our setting. For many distributions of interest, e.g., Gaussian and Bernoulli, the Fischer information of a single sample is the inverse of the variance, and we make that assumption for $\mathcal{D}_p$. We also assume that the $\mathcal{D}_p$ has sub-Gaussian tails. Thus, as long as the set of permissible meta-distributions includes distributions with this property, e.g., included truncated Gaussians, our lower bound applies.

Theorem 5.1. *Let $\mathcal{P}$ be a parameterized family of distributions $p \mapsto \mathcal{D}_p$ and suppose that for all $p \in [0, 1]$ and $k \in \mathbb{N}$, the Fisher information of $\phi_{p,k}$ is inversely proportional to the variance, $\text{Var}(\mathcal{D}_p(k))$:*

$$\int (\tfrac{\partial}{\partial p} \log \phi_{p,k}(x))^2 \phi_{p,k}(x) dx = O(\tfrac{1}{\text{Var}(\mathcal{D}_p(k))}), \tag{8}$$

and for all p, $n > 0$, $k \in \mathbb{N}$ and $\beta \in [1/3, 2/3]$, $f_{\mathcal{D}_p}^k(n, \sigma_p^2, \beta) = \tilde{O}(\text{Var}(\mathcal{D}_p(k)))$, then

$$\max_{p \in [1/3, 2/3]} [\text{Var}_{\forall i \in [n], x_i \sim \mathcal{D}(k_i), M}(\widehat{p}_\epsilon^{\text{ideal}})] = \tilde{O}(\min_{M, \text{ unbiased}} \max_{p \in [1/3, 2/3]} [\text{Var}_{\forall i \in [n], x_i \sim \mathcal{D}(k_i), M}(M)]).$$

Further, under the conditions of Theorem 4.1,

$$\max_{p \in [1/3, 2/3]} [\text{Var}_{\forall i \in [n], x_i \sim \mathcal{D}(k_i), M}(\widehat{p}_\epsilon^{\text{realistic}})] = \tilde{O}(\min_{M, \text{ unbiased}} \max_{p \in [1/3, 2/3]} [\text{Var}_{\forall i \in [n], x_i \sim \mathcal{D}(k_i), M}(M)]).$$

We will prove Theorem 5.1 in three steps. The following class of noisy linear estimators, NLE, will act as an intermediary in our proof. The notation σ_i denotes $\text{Var}(x_i)$, which accounts for the randomness in generating x_i.

$$\text{NLE} = \{M_{\text{NL}}(\mathbf{x}; \mathbf{w}) = \textstyle\sum_{i=1}^n w_i x_i + \text{Lap}(\tfrac{\max_i w_i \sigma_i}{\epsilon}) \mid w_i \in [0, 1], \textstyle\sum_{i=1}^n w_i = 1\}.$$

Similar to $\widehat{p}_\epsilon^{\text{ideal}}$, this class of estimators is not realizable since we only have access to an estimate of $\sigma_i = \text{Var}(\mathcal{D}_p(k_i))$. Additionally, the estimators in NLE are not necessarily ϵ-DP.

The proof Theorem 5.1 has three main steps outlined below. The proof of each Lemma is contained in Appendix E. The first step is shown in Lemma 5.2, which shows that the weights used in $\widehat{p}_\epsilon^{\text{ideal}}$ are optimal (i.e., variance-minimizing) among all estimators in the set NLE.

Lemma 5.2. *Given* $\widehat{p}_i \sim \mathcal{D}_p(k_i)$ *with variance* σ_i^2 *for all* $i \in [n]$ *and* $w \in [0,1]^n$ *such that* $\sum_{i=1}^n w_i = 1$, *let* $\widehat{p} = \sum_{i=1}^n w_i \widehat{p}_i + \mathrm{Lap}(\frac{\max_i w_i \sigma_i}{\epsilon})$. *The variance of* $\widehat{p}$ *is minimized by the following weights:* $\tilde{w}_i^* = \frac{\min\{1/\sigma_i^2, T/\sigma_i\}}{\sum_{j=1}^n \min\{1/\sigma_j^2, T/\sigma_j\}}$ *for some* T.

Since the threshold T^* in $\widehat{p}_\epsilon^{\mathrm{ideal}}$ was chosen to minimize $\mathrm{Var}(\widehat{p}_\epsilon^{\mathrm{ideal}})$, then we know that the weights w_i^* in $\widehat{p}_\epsilon^{\mathrm{ideal}}$ are optimal.

Now, let us turn to the second – and main – component of the proof of Theorem 5.1. Lemma 5.3 formalises the statement that an estimator inside the class NLE is minimax optimal among unbiased estimators. That is, for any unbiased estimator M, there exists an estimator $M_{\mathrm{NL}} \in \mathrm{NLE}$ with lower worst-case variance.

Lemma 5.3. *Let* $\mathcal{P}$ *be a parameterized family of distributions* $p \mapsto \mathcal{D}_p$ *and suppose that* M : $[0,1]^n \to [0,1]$ *is an* ϵ-*DP estimator such that for all* $p \in [1/3, 2/3]$, *(1)* M *is unbiased,* $\mu_M(p) = p$, *and (2) the Fisher information of* ϕ_{p,k_i} *is inversely proportional to the variance* $\mathrm{Var}(\mathcal{D}_p(k_i))$, $\int (\frac{\partial}{\partial p} \log \phi_{p,k_i}(x_i))^2 \phi_{p,k_i}(x_i) dx_i = O(\frac{1}{\mathrm{Var}(\mathcal{D}_p(k_i))})$, *then there exists an estimator* $M_{\mathrm{NL}} \in \mathrm{NLE}$ *such that*
$\max_{p \in [1/3, 2/3]}[\mathrm{Var}_{\forall i \in [n], x_i \sim \mathcal{D}(k_i), M_{\mathrm{NL}}}(M_{\mathrm{NL}})] \leq O(\max_{p \in [1/3, 2/3]}[\mathrm{Var}_{\forall i \in [n], x_i \sim \mathcal{D}(k_i), M}(M)])$.

The final component needed for the proof of Theorem 5.1 is a translation from the estimators in NLE, which are not ϵ-DP to the corresponding ϵ-DP estimator. For any weight vector $\mathbf{w}$, we can define an ϵ-DP estimator by truncating the data point x_i and calibrating the noise appropriately:

$$M_{\mathrm{TNL}}(x_1, \cdots, x_n; \mathbf{w}) = \sum_{i=1}^n w_i [x_i]_{p - f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)}^{p + f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)} + \mathrm{Lap}(\frac{\max_i 2 w_i f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)}{\epsilon}). \qquad (9)$$

Provided $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) \approx \mathrm{Var}(\mathcal{D}(k_i))$, the estimators M_{TNL} have approximately the same variance as the corresponding element of NLE, but are slightly biased. This is formalized in the following lemma.

Lemma 5.4. *For any distribution* $\mathcal{D}$, $n > 0$ *and* $\beta \in [0,1]$, *if for all* k_i, $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) = \tilde{O}(\mathrm{Var}(\mathcal{D}(k_i))$ *then for any* $\mathbf{w} \in [0,1]^n$ *such that* $\sum_{i=1}^n w_i = 1$, *we have* $\mathrm{Var}(M_{\mathrm{TNL}}(\cdot \; ; \mathbf{w})) = \tilde{O}(\mathrm{Var}(M_{\mathrm{NL}}(\cdot \; ; \mathbf{w})))$. *Further, the bias of* M_{TNL} *is at most* β.

Finally, we have the tools to prove the main theorem in this section, Theorem 5.1:

$$\min_{M \text{ unbiased}} \max_{p \in [1/3, 2/3]} [\mathrm{Var}_{\mathcal{D}_p}(M)] = \Omega(\min_{M \in \mathrm{NLE}} \max_{p \in [1/3, 2/3]} [\mathrm{Var}_{\mathcal{D}_p}(M)]) = \Omega(\max_{p \in [1/3, 2/3]} [\mathrm{Var}_{\mathcal{D}_p}(p_\epsilon^{\mathrm{NLE}})])$$

$$= \tilde{\Omega}(\max_{p \in [1/3, 2/3]} [\mathrm{Var}_{\mathcal{D}_p}(\widehat{p}_\epsilon^{\mathrm{ideal}})])$$

$$= \tilde{\Omega}(\max_{p \in [1/3, 2/3]} [\mathrm{Var}_{\mathcal{D}_p}(\widehat{p}_\epsilon^{\mathrm{realistic}})])$$

where $p_\epsilon^{\mathrm{NLE}} \in \mathrm{NLE}$ has the same weights as $\widehat{p}_\epsilon^{\mathrm{ideal}}$. The equalities follow from Lemmas 5.3, 5.2, 5.4, and Theorem 4.1, respectively.

5.2 Minimax Lower Bound on Estimation Rate

In addition to establishing the near optimality of $\widehat{p}_\epsilon^{\mathrm{realistic}}$, we will also give a lower bound on minimax rate of estimation in terms of the parameters $k_1, \cdots, k_n$ and σ_p^2. Note that we can view the truncation of the weights w_i as establishing an effective upper bound on k_i. Given $k_1, \cdots, k_n \in \mathbb{N}$, and $\epsilon > 0$, let $k^* = \arg\min_k \frac{\frac{k}{\epsilon^2} + \sum_{i=1}^n \min\{k_i, k\}}{(\sum_{i=1}^n \min\{k_i, k\})^2}$. Intuitively, in the case that $\sigma_p = 0$, we want to use as many samples as possible, but one user contributing many samples leads to larger sensitivity and thus privacy cost. Limiting to k_{max} the number of samples per user allows us to limit the sensitivity to be about $w_{max}(1/\sqrt{k_{max}})$. Since w_i is proportional to the number of samples used, the variance when using at most k samples per user is the above expression being minimized. Our lower bound below is close to this value for reasonable k_i's.

Corollary 5.5. *Given* $k_1, \cdots, k_n \in \mathbb{N}$, *and* σ_p, *there exists a family of distributions* $\mathcal{D}_p$ *such that*

$$\min_{M, \text{ unbiased}} \max_{p \in [1/3, 2/3]} \mathrm{Var}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i)}[M(x_1, \cdots, x_n)] \geq \tilde{\Omega}(\min\{\frac{\frac{k^*}{\epsilon^2} + \sum_{i=1}^n \min\{k_i, k^*\}}{(\sum_{i=1}^n \min\{k_i, \sqrt{k_i k^*}\})^2}, \frac{\sigma_p^2}{n}\}).$$

References

[1] Jayadev Acharya and Ziteng Sun. Communication complexity in locally private distribution estimation and heavy hitters. *arXiv preprint arXiv:1905.11888*, 2019.

[2] Jayadev Acharya, Ziteng Sun, and Huanyu Zhang. Hadamard response: Estimating distributions privately, efficiently, and with little communication. volume 89 of *Proceedings of Machine Learning Research*, pages 1120–1129. PMLR, 16–18 Apr 2019.

[3] Daniel Alabi, Audra McMillan, Jayshree Sarathy, Adam Smith, and Salil Vadhan. Differentially private simple linear regression, 2020.

[4] Hilal Asi and John C Duchi. Instance-optimality in differential privacy via approximate inverse sensitivity mechanisms. In H. Larochelle, M. Ranzato, R. Hadsell, M. F. Balcan, and H. Lin, editors, *Advances in Neural Information Processing Systems*, volume 33, pages 14106–14117. Curran Associates, Inc., 2020. URL `https://proceedings.neurips.cc/paper/2020/file/a267f936e54d7c10a2bb70dbe6ad7a89-Paper.pdf`.

[5] Abhishek Bhowmick, John Duchi, Julien Freudiger, Gaurav Kapoor, and Ryan Rogers. Protection against reconstruction and its applications in private federated learning, 2019.

[6] Michael Borenstein, Larry V Hedges, Julian PT Higgins, and Hannah R Rothstein. *Introduction to meta-analysis*. John Wiley & Sons, 2021.

[7] Wei-Ning Chen, Peter Kairouz, and Ayfer Özgür. Breaking the communication-privacy-accuracy trilemma. *arXiv preprint arXiv:2007.11707*, 2020.

[8] John Duchi and Ryan Rogers. Lower bounds for locally private estimation via communication complexity. In Alina Beygelzimer and Daniel Hsu, editors, *Proceedings of the Thirty-Second Conference on Learning Theory*, volume 99 of *Proceedings of Machine Learning Research*, pages 1161–1191, Phoenix, USA, 25–28 Jun 2019. PMLR. URL `http://proceedings.mlr.press/v99/duchi19a.html`.

[9] John C Duchi, Michael I Jordan, and Martin J Wainwright. Minimax optimal procedures for locally private estimation. *Journal of the American Statistical Association*, 113(521):182–201, 2018.

[10] Cynthia Dwork and Jing Lei. Differential privacy and robust statistics. In *Proceedings of the Forty-First Annual ACM Symposium on Theory of Computing*, STOC '09, page 371–380, New York, NY, USA, 2009. Association for Computing Machinery. ISBN 9781605585062.

[11] Cynthia Dwork and Aaron Roth. *The Algorithmic Foundations of Differential Privacy*, volume 9. 2014. URL `http://dx.doi.org/10.1561/0400000042`.

[12] Cynthia Dwork, Frank McSherry, Kobbi Nissim, and Adam Smith. Calibrating noise to sensitivity in private data analysis. volume Vol. 3876, pages 265–284, 01 2006. doi: 10.1007/11681878_14.

[13] Hubert Eichner, Tomer Koren, Brendan Mcmahan, Nathan Srebro, and Kunal Talwar. Semicyclic stochastic gradient descent. In Kamalika Chaudhuri and Ruslan Salakhutdinov, editors, *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pages 1764–1773. PMLR, 09–15 Jun 2019.

[14] Úlfar Erlingsson, Vasyl Pihur, and Aleksandra Korolova. RAPPOR: randomized aggregatable privacy-preserving ordinal response. In *ACM SIGSAC Conference on Computer and Communications Security*, pages 1054–1067, 2014.

[15] Filip Hanzely and Peter Richtárik. Federated learning of a mixture of global and local models. *arXiv preprint arXiv:2002.05516*, 2020.

[16] Moritz Hardt and Kunal Talwar. On the geometry of differential privacy. In *Proceedings of the Forty-Second ACM Symposium on Theory of Computing*, STOC '10, page 705–714, New York, NY, USA, 2010. Association for Computing Machinery. ISBN 9781450300506.

[17] Joachim Hartung, Guido Knapp, and Bimal Sinha. Statistical meta-analysis with applications. 08 2008. doi: 10.1002/9780470386347.

[18] Justin Hsu, Sanjeev Khanna, and Aaron Roth. Distributed private heavy hitters. In *International Colloquium on Automata, Languages, and Programming*, pages 461–472. Springer, 2012.

[19] Aaron Johnson and Vitaly Shmatikov. Privacy-preserving data exploration in genome-wide association studies. In *Proceedings of the 19th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, KDD '13, page 1079–1087, New York, NY, USA, 2013. Association for Computing Machinery. ISBN 9781450321747.

[20] Peter Kairouz, H. Brendan McMahan, Brendan Avent, Aurélien Bellet, Mehdi Bennis, Arjun Nitin Bhagoji, Kallista Bonawitz, Zachary Charles, Graham Cormode, Rachel Cummings, Rafael G. L. D'Oliveira, Hubert Eichner, Salim El Rouayheb, David Evans, Josh Gardner, Zachary Garrett, Adrià Gascón, Badih Ghazi, Phillip B. Gibbons, Marco Gruteser, Zaid Harchaoui, Chaoyang He, Lie He, Zhouyuan Huo, Ben Hutchinson, Justin Hsu, Martin Jaggi, Tara Javidi, Gauri Joshi, Mikhail Khodak, Jakub Konecný, Aleksandra Korolova, Farinaz Koushanfar, Sanmi Koyejo, Tancrède Lepoint, Yang Liu, Prateek Mittal, Mehryar Mohri, Richard Nock, Ayfer Özgür, Rasmus Pagh, Hang Qi, Daniel Ramage, Ramesh Raskar, Mariana Raykova, Dawn Song, Weikang Song, Sebastian U. Stich, Ziteng Sun, Ananda Theertha Suresh, Florian Tramèr, Praneeth Vepakomma, Jianyu Wang, Li Xiong, Zheng Xu, Qiang Yang, Felix X. Yu, Han Yu, and Sen Zhao. Advances and open problems in federated learning. *Foundations and Trends® in Machine Learning*, 14(1–2):1–210, 2021. ISSN 1935-8237. doi: 10.1561/2200000083. URL http://dx.doi.org/10.1561/2200000083.

[21] Vishesh Karwa and Salil P. Vadhan. Finite sample differentially private confidence intervals. volume abs/1711.03908 of *Innovations in Theoretical Computer Science '18*, 2018.

[22] Daniel Levy, Ziteng Sun, Kareem Amin, Satyen Kale, Alex Kulesza, Mehryar Mohri, and Ananda Theertha Suresh. Learning with user-level privacy. In *Neural Information Processing Systems (NeurIPS 2021)*, 2021.

[23] Yuhan Liu, Ananda Theertha Suresh, Felix Xinnan X Yu, Sanjiv Kumar, and Michael Riley. Learning discrete distributions: user vs item-level privacy. In H. Larochelle, M. Ranzato, R. Hadsell, M. F. Balcan, and H. Lin, editors, *Advances in Neural Information Processing Systems*, volume 33, pages 20965–20976. Curran Associates, Inc., 2020. URL https://proceedings.neurips.cc/paper/2020/file/f06edc8ab534b2c7ecbd4c2051d9cb1e-Paper.pdf.

[24] Brendan McMahan, Eider Moore, Daniel Ramage, Seth Hampson, and Blaise Aguera y Arcas. Communication-Efficient Learning of Deep Networks from Decentralized Data. In Aarti Singh and Jerry Zhu, editors, *Proceedings of the 20th International Conference on Artificial Intelligence and Statistics*, volume 54 of *Proceedings of Machine Learning Research*, pages 1273–1282, Fort Lauderdale, FL, USA, 20–22 Apr 2017. PMLR.

[25] H. Brendan McMahan, Daniel Ramage, Kunal Talwar, and Li Zhang. Learning differentially private recurrent language models. In *6th International Conference on Learning Representations, ICLR 2018, Vancouver, BC, Canada, April 30 - May 3, 2018, Conference Track Proceedings*. OpenReview.net, 2018. URL https://openreview.net/forum?id=BJ0hF1Z0b.

[26] Audra McMillan, Adam Smith, and Jon Ullman. Instance Optimal Differentially Private Estimation. In preparation, 2022.

[27] Frank McSherry and Ilya Mironov. Differentially private recommender systems: Building privacy into the netflix prize contenders. In *Proceedings of the 15th ACM SIGKDD international conference on Knowledge discovery and data mining*, pages 627–636, 2009.

[28] Frank McSherry and Kunal Talwar. Mechanism design via differential privacy. In *48th Annual IEEE Symposium on Foundations of Computer Science (FOCS'07)*, pages 94–103, 2007. doi: 10.1109/FOCS.2007.66.

[29] Ion Mihoc and Cristina Fătu. Fisher's information measure and truncated normal distributions (ii). *Revue d'Analyse Numérique et de Théorie de l'Approximation*, 32, 01 2003.

[30] Frank Nielsen. *Cramér-Rao Lower Bound and Information Geometry*, pages 18–37. Hindustan Book Agency, Gurgaon, 2013. URL `https://doi.org/10.1007/978-93-86279-56-9_2`.

[31] Kaan Ozkara, Antonious Girgis, Deepesh Data, and Suhas Diggavi. A generative framework for personalized learning and estimation: Theory, algorithms, and privacy, 07 2022.

[32] Abhradeep Guha Thakurta and Adam Smith. Differentially private feature selection via stability arguments, and the robustness of the lasso. In Shai Shalev-Shwartz and Ingo Steinwart, editors, *Proceedings of the 26th Annual Conference on Learning Theory*, volume 30 of *Proceedings of Machine Learning Research*, pages 819–850, Princeton, NJ, USA, 12–14 Jun 2013. PMLR.

[33] Wikipedia contributors. Meta-analysis — Wikipedia, the free encyclopedia, 2021. URL `https://en.wikipedia.org/w/index.php?title=Meta-analysis&oldid=1023577278`. [Online; accessed May 2021].

[34] Fan Zhou and Guojing Cong. On the convergence properties of a k-step averaging stochastic gradient descent algorithm for nonconvex optimization. In *Proceedings of the Twenty-Seventh International Joint Conference on Artificial Intelligence, IJCAI-18*, pages 3219–3227. International Joint Conferences on Artificial Intelligence Organization, 7 2018. doi: 10.24963/ijcai.2018/447. URL `https://doi.org/10.24963/ijcai.2018/447`.

A Proofs from Section 2

Lemma 2.1. *For all distributions $\mathcal{D}$ supported on $[0,1]$ with mean p and variance σ_p^2, $\sigma_p^2 \leq p(1-p)$. Further, $\mathbb{E}[\mathcal{D}(k)] = p$ and $\mathrm{Var}(\mathcal{D}(k)) = \frac{1}{k}p(1-p) + \left(1 - \frac{1}{k}\right)\sigma_p^2$.*

Proof of Lemma 2.1. Firstly, note that,

$$\sigma_p^2 = \mathbb{E}_{x \sim \mathcal{D}}[x^2] - p^2 \leq \mathbb{E}_{x \sim \mathcal{D}}[x] - p^2 = p(1-p),$$

where the inequality follows from the fact that $\mathcal{D}$ is supported on $[0,1]$.

Next,

$$\mathbb{E}[x_i] = \int_{x=0}^{1} \Pr(p_i = x)\Pr(\mathrm{Ber}(x) = 1)dx = \int_{x=0}^{1} \Pr(p_i = x)xdx = p,$$

which by linearity of expectation implies that $\mathbb{E}[\mathcal{D}(k)] = p$.

By the Law of Total Variation, the variance of $\widehat{p}_i$ is:

$$\begin{aligned}
\mathrm{Var}(\widehat{p}_i) &= \mathbb{E}_{p_i}[\mathrm{Var}_{x_i}(\widehat{p}_i|p_i)] + \mathrm{Var}_{p_i}(\mathbb{E}_{x_i}[\widehat{p}_i|p_i]) \\
&= \mathbb{E}_{p_i}[\frac{1}{k_i}p_i(1-p_i)] + \mathrm{Var}_{p_i}(p_i) \\
&= \frac{1}{k_i}(p - \sigma_p^2 - p^2) + \sigma_p^2 \\
&= \frac{1}{k_i}(p - p^2) + (1 - \frac{1}{k_i})\sigma_p^2. \\
&= \frac{1}{k_i}\mathrm{Var}(\mathrm{Ber}(p)) + (1 - \frac{1}{k_i})\sigma_p^2.
\end{aligned}$$

$\square$

B Proofs from Section 4.2

Theorem 4.1. *For any $\epsilon > 0$, $\delta \in [0,1]$, Algorithm 2 is (ϵ, δ)-DP. If,*

- *$\mathtt{mean}_{\epsilon,\delta}$ is such that given $n/10$ samples from $\mathcal{D}$, with probability $1 - \beta$ $|p - \widehat{p}_\epsilon^{\mathrm{initial}}| \leq f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$ and $\widehat{p}_\epsilon^{\mathrm{initial}}(1 - \widehat{p}_\epsilon^{\mathrm{initial}}) \in [\frac{1}{2}p(1-p), \frac{3}{2}p(1-p)]$,*

- *$\mathtt{variance}_{\epsilon,\delta}$ is such that given $\log n$ samples from $\mathcal{D}(k)$, with probability $1 - \beta$, $\widehat{\sigma}_p^2 \in [\mathrm{Var}(\mathcal{D}(k)), \ 8\mathrm{Var}(\mathcal{D}(k))]$,*

- *the k_is are such that $\frac{k_1}{k_{n/2}} \leq \frac{n/2 - \log n}{\log n}$,*

then with probability $1 - 2\beta$, $\mathrm{Var}(\widehat{p}_\epsilon^{\mathrm{realistic}}) \leq C \cdot \mathrm{Var}(\widehat{p}_\epsilon^{\mathrm{ideal}})$ for some absolute constant C.

First, let us show that the conditions of Theorem 4.1 imply that the variance and truncation parameter estimates of each individual data subject are correct up to constant factors.

Lemma 4.2. *Given $\widehat{p}_\epsilon^{\mathrm{initial}}$, $\widehat{\sigma}_p^2$, and k_i, define $\widehat{\sigma}_i^2 = \frac{1}{k_i}\widehat{p}_\epsilon^{\mathrm{initial}}(1 - \widehat{p}_\epsilon^{\mathrm{initial}}) + \frac{k_i - 1}{k_i}\widehat{\sigma}_p^2$. Under the conditions of Theorem 4.1, for all $i > \log n$, we have $\widehat{\sigma}_i^2 \in \left[\frac{1}{2}\sigma_i^2, 9.5\sigma_i^2\right]$ and $|\widehat{b}_i - \widehat{a}_i| \leq 4|b_i - a_i|$.*

Proof of Lemma 4.2. Note that $\widehat{\sigma}_p^2$ is actually an estimate of the variance of $\mathcal{D}(k_{\log n})$ since it has access to samples from this distribution rather than $\mathcal{D}$ itself. Therefore, $\widehat{\sigma}_p^2 \in [\mathrm{Var}(\mathcal{D}(k_{\log n})), 8 \cdot \mathrm{Var}(\mathcal{D}(k_{\log n}))]$ implies $\widehat{\sigma}_p^2 \in \left[\sigma_p^2, 8\left(\frac{1}{k_{\log n}}p(1-p) + \sigma_p^2\right)\right]$. Then for every

$i \geq \log n$ (i.e., with $k_i \leq k_{\log n}$),

$$\widehat{\sigma}_i^2 = \frac{1}{k_i}\widehat{p}_\epsilon^{\text{initial}}(1 - \widehat{p}_\epsilon^{\text{initial}}) + \frac{k_i - 1}{k_i}\widehat{\sigma}_p^2$$

$$\geq \frac{1}{k_i}\frac{1}{2}p(1-p) + \frac{k_i - 1}{k_i}\sigma_p^2$$

$$\geq \frac{1}{2}\left(\frac{1}{k_i}p(1-p) + \frac{k_i - 1}{k_i}\sigma_p^2\right)$$

$$= \frac{1}{2}\sigma_i^2,$$

where the first inequality follows from the accuracy conditions on $\texttt{mean}_{\epsilon,\delta}$ and $\texttt{variance}_{\epsilon,\delta}$ in Theorem 4.1, and the last equality follows from the definition of σ_i^2 in Lemma 2.1. Also,

$$\widehat{\sigma}_i^2 = \frac{1}{k_i}\widehat{p}_\epsilon^{\text{initial}}(1 - \widehat{p}_\epsilon^{\text{initial}}) + \frac{k_i - 1}{k_i}\widehat{\sigma}_p^2$$

$$\leq \frac{1}{k_i}\frac{3}{2}p(1-p) + 8\frac{k_i - 1}{k_i}\left(\frac{1}{k_{\log n}}p(1-p) + \sigma_p^2\right)$$

$$= \left(\frac{3}{2} + 8\frac{k_i - 1}{k_{\log n}}\right)\frac{1}{k_i}p(1-p) + 8\frac{k_i - 1}{k_i}\sigma_p^2$$

$$\leq 9.5\left(\frac{1}{k_i}p(1-p) + \frac{k_i - 1}{k_i}\sigma_p^2\right)$$

$$= 9.5\sigma_i^2,$$

where again, the first inequality follows from the accuracy conditions on $\texttt{mean}_{\epsilon,\delta}$ and $\texttt{variance}_{\epsilon,\delta}$ in Theorem 4.1, and the last equality follows from the definition of σ_i^2 in Lemma 2.1. The intermediate steps are simply algebraic manipulations. These two facts give us the desired bounds on $\widehat{\sigma}_i^2$.

Next we turn to the truncation parameters $\widehat{a}_i$ and $\widehat{b}_i$. Using the definition of $\widehat{a}_i$ in Algorithm 2, we have,

$$\widehat{a}_i = \widehat{p}_\epsilon^{\text{initial}} - \alpha - f_{\mathcal{D}}^{k_i}(n, \widehat{\sigma}_p^2, \beta/2)$$

$$\leq p - f_{\mathcal{D}}^{k_i}(n, \widehat{\sigma}_p^2, \beta/2))$$

$$\leq p - f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$$

$$= a_i,$$

where the two inequalities respectively follow from the accuracy conditions on $\texttt{mean}_{\epsilon,\delta}$ and $\texttt{variance}_{\epsilon,\delta}$ in Theorem 4.1. A symmetric result that $\widehat{b}_i \geq b_i$ follows similarly.

Finally,

$$|\widehat{b}_i - \widehat{a}_i| = 2\alpha + 2f_{\mathcal{D}}^{k_i}(n, \widehat{\sigma}_p^2, \beta)$$

$$\leq 2f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) + 2f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$$

$$= 4f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$$

$$= 4|b_i - a_i|.$$

The inequalities again follows from the accuracy conditions on $\texttt{mean}_{\epsilon,\delta}$ and $\texttt{variance}_{\epsilon,\delta}$ in Theorem 4.1. □

Proof of Theorem 4.1. To see that Algorithm 2 is differentially private, consider the three cohorts into which users are placed. The first cohort, containing the $n/10$ users with the smallest k_i will have their data used in $\texttt{mean}_{\epsilon,\delta}$, which is (ϵ, δ)-DP. Similarly, the second cohort containing the $\log n$ users with the largest k_i will have their data used in $\texttt{variance}_{\epsilon,\delta}$, which is also (ϵ, δ)-DP. The intermediate estimators of $\widehat{\sigma}_i^2$, $\widehat{T}^*$, $\widehat{a}_i$, $\widehat{b}_i$, and sensitivity Λ are all computed as post-processing on the private outputs of these initial estimation subroutines and on the public k_is, and thus do not incur

any additional privacy cost. The third cohort contains the middle users $i \in [\log n + 1, 9n/10]$. These users' data are only used in the final estimate, which is an $(\epsilon, 0)$-DP instantiation of the Laplace Mechanism [12].

Since these cohorts are disjoint and private algorithms are applied to each cohort's data separately, parallel composition applies, and the overall privacy parameters are the maximum of those experienced by any cohort, so the overall algorithm is (ϵ, δ)-DP.

For accuracy of the $\widehat{p}_\epsilon^{\text{realistic}}$ estimator produced by Algorithm 2, first notice that under the assumption that $\frac{k_{\max}}{k_{\text{med}}} \leq \frac{n/2 - \log n}{\log n}$, if $\sigma_{k_{\max}}^2 = \text{Var}(\widehat{p}_1)$ and $\sigma_{k_{\text{med}}}^2 = \text{Var}(\widehat{p}_{n/2})$ then

$$\sigma_{k_{\text{med}}}^2 = \frac{1}{k_{\text{med}}}p(1-p) + \left(1 - \frac{1}{k_{\text{med}}}\right)\sigma_p^2 \leq \frac{n/2 - \log n}{\log n}\frac{1}{k_{\max}}p(1-p) + \left(1 - \frac{1}{k_{\max}}\right)\sigma_p^2 \leq \frac{n/2 - \log n}{\log n}\sigma_{k_{\max}}^2.$$

Therefore, for any truncation parameter T,

$$\frac{1}{2}\sum_{i=1}^{n}\min\left\{\frac{1}{\sigma_i^2}, \frac{T}{\sigma_i}\right\} \leq \sum_{i=1}^{n/2}\min\left\{\frac{1}{\sigma_i^2}, \frac{T}{\sigma_i}\right\}$$

$$= \sum_{i=1}^{\log n}\min\left\{\frac{1}{\sigma_i^2}, \frac{T}{\sigma_i}\right\} + \sum_{i=\log n+1}^{n/2}\min\left\{\frac{1}{\sigma_i^2}, \frac{T}{\sigma_i}\right\}$$

$$\leq \log n \cdot \min\left\{\frac{1}{\sigma_{k_{\max}}^2}, \frac{T}{\sigma_{k_{\max}}}\right\} + \sum_{i=\log n+1}^{n/2}\min\left\{\frac{1}{\sigma_i^2}, \frac{T}{\sigma_i}\right\}$$

$$\leq (n/2 - \log n) \cdot \min\left\{\frac{1}{\sigma_{k_{\text{med}}}^2}, \frac{T}{\sigma_{k_{\text{med}}}}\right\} + \sum_{i=\log n+1}^{n/2}\min\left\{\frac{1}{\sigma_i^2}, \frac{T}{\sigma_i}\right\}$$

$$\leq 2\sum_{i=\log n+1}^{n/2}\min\left\{\frac{1}{\sigma_i^2}, \frac{T}{\sigma_i}\right\}$$

$$\leq 2\sum_{i=\log n+1}^{9n/10}\min\left\{\frac{1}{\sigma_i^2}, \frac{T}{\sigma_i}\right\}, \tag{10}$$

where the first, second, and fourth inequalities follow from our assumed ordering on the k_is. The third inequality comes from our assumption on $k_{\max}$ and k_{med}, and the final inequality follows from the fact that the summands $\min\{\frac{1}{\sigma_i^2}, \frac{T}{\sigma_i}\}$ are positive so adding more terms only increases the sum.

Therefore,

$$\operatorname{Var}(\widehat{p}_\epsilon^{\text{realistic}}) = \frac{1}{(\sum_{j=\log n+1}^{9n/10} \min\{1/\widehat{\sigma}_j^2, \frac{\widehat{T}^*}{\widehat{\sigma}_i}\})^2} \left(\sum_{i=\log n+1}^{9n/10} \min\{\frac{1}{\widehat{\sigma}_i^4}, \frac{\widehat{T}^{*2}}{\widehat{\sigma}_i^2}\}\sigma_i^2 + \max_i \frac{\min\{\frac{1}{\widehat{\sigma}_i^4}, \frac{\widehat{T}^{*2}}{\widehat{\sigma}_i^2}\}|\widehat{b}_i - \widehat{a}_i|^2}{\epsilon^2} \right)$$

$$\leq \frac{1}{(\sum_{j=\log n+1}^{9n/10} \min\{1/\widehat{\sigma}_j^2, \frac{\widehat{T}^*}{\widehat{\sigma}_i}\})^2} \left(\sum_{i=\log n+1}^{9n/10} \min\{\frac{1}{\widehat{\sigma}_i^4}, \frac{\widehat{T}^{*2}}{\widehat{\sigma}_i^2}\}2\widehat{\sigma}_i^2 + \max_i \frac{\min\{\frac{1}{\widehat{\sigma}_i^4}, \frac{\widehat{T}^{*2}}{\widehat{\sigma}_i^2}\}|\widehat{b}_i - \widehat{a}_i|^2}{\epsilon^2} \right)$$

$$\leq 2\frac{1}{(\sum_{j=\log n+1}^{9n/10} \min\{1/\widehat{\sigma}_j^2, \frac{\widehat{T}^*}{\widehat{\sigma}_i}\})^2} \left(\sum_{i=\log n+1}^{9n/10} \min\{\frac{1}{\widehat{\sigma}_i^2}, \widehat{T}^{*2}\} + \max_i \frac{\min\{\frac{1}{\widehat{\sigma}_i^4}, \frac{\widehat{T}^{*2}}{\widehat{\sigma}_i^2}\}|\widehat{b}_i - \widehat{a}_i|^2}{\epsilon^2} \right)$$

$$\leq 2\frac{1}{(\sum_{j=\log n+1}^{9n/10} \min\{1/\widehat{\sigma}_j^2, \frac{T^*}{\widehat{\sigma}_i}\})^2} \left(\sum_{i=\log n+1}^{9n/10} \min\{\frac{1}{\widehat{\sigma}_i^2}, T^{*2}\} + \max_i \frac{\min\{\frac{1}{\widehat{\sigma}_i^4}, \frac{T^{*2}}{\widehat{\sigma}_i^2}\}|\widehat{b}_i - \widehat{a}_i|^2}{\epsilon^2} \right)$$

$$\leq 2\frac{1}{(\sum_{j=\log n+1}^{9n/10} \min\{1/10\sigma_j^2, \frac{\sqrt{2}T^*}{\sigma_i}\})^2} \left(\sum_{i=\log n+1}^{9n/10} \min\{\frac{2}{\sigma_i^2}, T^{*2}\} + \max_i \frac{\min\{\frac{4}{\sigma_i^4}, \frac{2T^{*2}}{\sigma_i^2}\}6|b_i - a_i|^2}{\epsilon^2} \right)$$

$$\leq 480\frac{1}{(\sum_{j=\log n+1}^{9n/10} \min\{1/\sigma_j^2, \frac{T^*}{\sigma_i}\})^2} \left(\sum_{i=\log n+1}^{9n/10} \min\{\frac{1}{\sigma_i^2}, T^{*2}\} + \max_i \frac{\min\{\frac{1}{\sigma_i^4}, \frac{T^{*2}}{\sigma_i^2}\}|b_i - a_i|^2}{\epsilon^2} \right)$$

$$\leq 480\frac{1}{\frac{1}{16}(\sum_{j=1}^{n} \min\{1/\sigma_j^2, \frac{T^*}{\sigma_i}\})^2} \left(\sum_{i=1}^{n} \min\{\frac{1}{\sigma_i^2}, T^{*2}\} + \max_i \frac{\min\{\frac{1}{\sigma_i^4}, \frac{T^{*2}}{\sigma_i^2}\}|b_i - a_i|^2}{\epsilon^2} \right)$$

$$= 7680 \cdot \operatorname{Var}(\widehat{p}_\epsilon^{\text{ideal}})$$

The first equality simply follows from the definition of the estimator and basic properties of the variance, as well as the fact that $\operatorname{Var}([\widehat{p}_i]_{a_i}^{b_i}) \leq \sigma_i$. The first inequality follows from the fact that $\sigma_i^2 \leq 2\widehat{\sigma}_i^2$, which was shown in Lemma 4.2. The second inequality is simply pulling out the constant to the front. The third inequality follows from the definition of $\widehat{T}$ as the optimiser of the variance using the approximations $\widehat{\sigma}_i^2$, $\widehat{b}_i$ and $\widehat{a}_i$. The fourth inequality follows from the fact that $\widehat{\sigma}_i^2 \in \left[\frac{1}{2}\sigma_i^2, 10\sigma_i^2\right]$ and $|\widehat{b}_i - \widehat{a}_i| \leq 4|b_i - a_i|$, as shown in Lemma 4.2, and will hold with probability $1 - 2\beta$, by taking a union bound over the β failure probabilities from each of the $\text{mean}_{\epsilon,\delta}$ and $\text{variance}_{\epsilon,\delta}$ subroutines. The fifth inequality simply pulls out the constants (480=2*10*4*6). The final inequality follows from Equation (10) above. The final equality follows from definition of $\widehat{p}_\epsilon^{\text{ideal}}$ and the assumption that $\frac{1}{2}\sigma_i^2 \leq \operatorname{Var}([\widehat{p}_i]_{a_i}^{b_i})$.

$\square$

C   Special Case: The constant p_i case.

In the previous section, we considered the setting where there was heterogeneity in both the users' distributions (i.e., the p_is were not constant), as well as the number of data points that they each held (i.e., the k_is were not constant). In the absence of variation in the p_i, each user is sampling from the same distribution $\text{Ber}(p)$. When privacy is not a concern, this setting reduces to the single-data-point-per-user setting where the sample size is increased to $\sum_{i=1}^n k_i$. However, under the constraint of user-level differential privacy, this setting is distinct from the single-data-point-per-user setting, since we need to protect the entirety of each users data set. In fact, much of the complexity of Algorithm 2 is required even in this simpler case. In particular, the truncated inverse variance weighting is still required in this case when there is variation in the k_i. In fact, the only step of Algorithm 2 that is not required is Step **??**, since already know that $\sigma_p^2 = 0$. Since there is no variance in $\mathcal{D}$, the high probability bound $f_{\mathcal{D}}^{k_i}(n, \widehat{\sigma_p^2}, \beta)$ is just due to the randomness in the binomial distribution $\text{Bin}(k_i, p)$, which comes from averaging k_i samples drawn from $\text{Ber}(p)$.

When $\sigma_p^2 = 0$, σ_i has the simple formula $\sigma_i = \frac{\sqrt{p(1-p)}}{k_i}$ and we can directly translate from the truncation threshold T on σ_i to a truncation threshold k on k_i, $T = \frac{\sqrt{p(1-p)}}{k}$. Further, if we assume that all the k_i are large enough ($\min k_i \geq 2\ln(1/\delta)/p$) then we also have the simple formula $f_{\mathcal{D}}^{k_i}(n, \widehat{\sigma_p^2}, \beta) = \sqrt{\frac{3p\ln(2/\beta)}{k_i}}$. We can plug these into Equation (6) (recall that T^* is defined as the truncation threshold that minimizes the variance of $\widehat{p}_\epsilon^{\text{ideal}}$) to obtain the following formula for the variance of $\widehat{p}_\epsilon^{\text{ideal}}$, and hence $\widehat{p}_\epsilon^{\text{realistic}}$:

$$\min_k \frac{p(1-p)\sum_{i=1}^n \min\{k_i, k\} + 6p\ln(2/\beta)\frac{k}{\epsilon^2}}{(\sum_{j=1}^n \min\{k_i, \sqrt{k_i k}\})^2}. \tag{11}$$

Even in the private setting, one can reduce to the single-data-point-per-user setting by reducing the sample size by a factor of 2, and forcing the $n/2$ users with the most data points to produce their estimate $\hat{p}_i$ using only k_{med} (the median k_i) data points. Then each estimate $\hat{p}_i$ is a sample from the same distribution and we can compute their mean. To the best of our knowledge, all the prior work in the private literature that handles variations in k_i follows this formula. However, not only does this algorithm reduce the sample size by a factor of 2, it also unnecessarily hinders the contribution of users with many data points. As a simple example, suppose that all the users have a single data point, except for $\sqrt{n}$ users, which have n data points. Then the algorithm which forces $n/2$ of the users to use the median number of data points has an error rate of $\Theta(\frac{1}{n} + \frac{1}{n^2\epsilon^2})$ assuming that p is bounded away from 0 or 1. Letting $k = n$ in Equation 11 implies that that the truncated inverse variance weighted algorithm in the previous section is better able to utilise the data of the users with high k_is, resulting in an error rate of $O(\frac{1}{n^{3/2}} + \frac{1}{n^2\epsilon^2})$.

D Extension: private-size user-level differential privacy setting

When defining user-level DP where users have heterogeneous quantities of data, we also need to distinguish between settings where the number of data points held by each user is protected information, and settings where it is publicly known. We'll refer to the former as *private-size user-level differential privacy*, where the entry that differs between neighboring databases can have arbitrarily different number of data points, and the latter as *public-size user-level differential privacy*, where the amount of data held by each user is the same in neighboring databases. Formally, let $D_i = \{x_i^1, \cdots, x_i^{k_i}\}$ be the data of user i for each $i \in [n]$. For private-size user-level differential privacy, we say D and D' are neighbours if there exists an index i such that for all $j \in [n] \backslash \{i\}$, $D_j = D_j'$. For public-size user-level differential privacy, we say D and D' are neighbours if they are neighbours under private-size user-level differential privacy and additionally $|D_i| = |D_i'|$ for all $i \in [n]$.

Let us now turn to our problem in the private-size user-level differential privacy setting, where the k_is are private information and require formal privacy protections. We will need to make several changes to Algorithm 2 to make it private under this stronger notion of privacy. Under public-size user-level privacy, the quantities $\hat{T}^*$ (the weight truncation parameter) and Λ (the sensitivity of the final estimate) in Algorithm 2 do not pose privacy concerns since they only depend on the data points $\hat{p}_i$ through the $\widehat{p}_\epsilon^{\text{initial}}$ and $\widehat{\sigma}_i^2$, which are both produced differentially privately. However, both these quantities depend on the k_i directly, and hence care needs to be taken when using them under private-size user-level DP.

In Algorithm 3, we outline the extension of Algorithm 2 to satisfy private-size user-level differential privacy. The core component of the final estimate is described in Algorithm 4. The function $\mathcal{M}$ as described in Algorithm 4 incorporates the truncation in a slightly different (but equivalent) manner to Algorithm 2, but is otherwise the same, without the addition of noise.

The first key difference is how the weights are truncated. Observe that choosing a truncation parameter T is equivalent to choosing an integer k such that $T = 1/\text{Var}(\mathcal{D}(k))$. So $\widehat{k_{\log n}}$ plays the role in Algorithm 3 that T^* plays in Algorithm 2. The statistic $\widehat{k_{\log n}}$ is a private estimate of the $\log n$-th order statistic of the set $\{k_1, \cdots, k_n\}$. Since the only users that participate in the final estimate (and hence have their data truncated) all have $k_i < k_{\log n}$, this algorithm attempts to find the smallest truncation parameter such that no data is actually truncated. We will show that provided ϵ is not too small, this level of truncation is sufficient.

Algorithm 3 Private Heterogeneous Mean Estimation $\hat{p}_\epsilon^{\text{realistic, unknown size}}$

Input: (ϵ, δ)-DP mean estimator $\texttt{mean}_{\epsilon,\delta}$, (ϵ, δ)-DP variance estimator $\texttt{variance}_{\epsilon,\delta}$, ϵ-DP estimator of the ℓth order statistic $\texttt{EM}_\epsilon(\cdot; \ell)$, number of users n, number of samples held by each user $(k_1, \ldots, k_n \ s.t. \ k_i \geq k_{i+1})$, an upper bound on the total number of data points held by a single user $k_{\max}$, user-level estimates $(\widehat{p}_1, \cdots, \widehat{p}_n)$, a constant $C > 0$, error guarantee on $\texttt{mean}_{\epsilon,\delta}$ $\alpha > 0$, and desired high probability bound $\beta \in [0, 1]$.

1: **Initial Estimates**
2: $\widehat{p}^{\text{initial}} = \texttt{mean}_{\epsilon,\delta}(\text{x}_{9n/10+1}^1, \cdots, \text{x}_n^1)$ $\hspace{2cm}$ Initial mean estimate
3: $\widehat{\sigma}_p^2 = \texttt{variance}_{\epsilon,\delta}(\widehat{p}_1, \cdots, \widehat{p}_{\log n})$ $\hspace{2cm}$ Initial variance estimate

4: **Compute Sensitivity Proposal**
5: $\widehat{k_T} = \texttt{EM}_\epsilon(k_1, \cdots, k_n; \frac{2\log n}{\epsilon}, k_{\max})$ $\hspace{2cm}$ Compute weight truncation
6: **for** $i \in [\log n + 1, n/2]$ **do**
7: $\quad k_i' = \min\{k_i, \widehat{k_T}\}$
8: $\quad \widehat{\sigma_i'}^2 = \frac{1}{k_i'}(\widehat{p}_\epsilon^{\text{initial}} - (\widehat{p}_\epsilon^{\text{initial}})^2) + (1 - \frac{1}{k_i'})\widehat{\sigma}_p^2.$
9: $\quad v_i = \frac{1}{\widehat{\sigma_i'}^2}$ $\hspace{2cm}$ Compute truncated, unnormalised weights
10: $\widehat{\sigma_{\min}}^2 = \frac{1}{\widehat{k_T}}(\widehat{p}_\epsilon^{\text{initial}} - (\widehat{p}_\epsilon^{\text{initial}})^2) + (1 - \frac{1}{\widehat{k_T}})\widehat{\sigma}_p^2.$
11: $\widehat{N} = \sum_{j=\log n+1}^{9n/10} v_i + \text{Lap}\left(\frac{1}{\epsilon\widehat{\sigma_{\min}}^2}\right) - \frac{1}{\epsilon\widehat{\sigma_{\min}}^2}\ln(2\delta)$ $\hspace{1cm}$ Compute noisy normalisation term
12: $\Upsilon = \frac{\log(1/\delta)}{\epsilon} + \frac{\ln(1/\delta)\ln(1/\gamma)}{\epsilon}$
13: $\Lambda = 8\left(\sqrt{\Upsilon + 1}\log n + \frac{\log(1/\beta)}{\sqrt{\Upsilon+1}}\right)\frac{f_{\mathcal{D}}^{k_{\max}}(n, \widehat{\sigma}_p^2, \beta)}{\widehat{\sigma_{\max}}^2 \widehat{N}}$ $\hspace{1cm}$ Compute local sensitivity proposal

14: **Propose-Test-Release on** $\mathcal{M}(\cdot; \widehat{k_{\log n}}, n, \widehat{p}_\epsilon^{\text{initial}}, \widehat{\sigma}_p^2, \alpha)$
15: $D_T = \{(\widehat{p}_i, k_i)\}_{i \in [\log n+1:9n/10]}$
16: $\kappa^* = \arg\max\{\kappa \in \mathbb{N} \mid \forall D' \text{ s.t. } D' \text{ is a } \kappa\text{-neighbor of } D_T, \Delta(\mathcal{M}(\cdot; \widehat{k_T}, n, \widehat{p}_\epsilon^{\text{initial}}, \widehat{\sigma}_p^2, \alpha); D') \leq \Lambda\}$ $\hspace{2cm}$ Compute distance to high sensitivity dataset
17: $\tilde{\kappa} = \kappa^* + \text{Lap}(1/\epsilon)$
18: **if** $\tilde{\kappa} < \frac{\log(1/\delta)}{\epsilon}$ **then**
19: $\quad$ **return** $\hat{p}_\epsilon^{\text{realistic, unknown size}} = \widehat{p}_\epsilon^{\text{initial}}$ $\hspace{0.5cm}$ Return initial estimate if proposed local sens. too small
20: **else**
21: $\quad$ Sample $Y \sim \text{Lap}\left(\frac{\Lambda}{\epsilon}\right)$ $\hspace{2cm}$ Sample noise added for privacy
22: $\quad$ **return** $\hat{p}_\epsilon^{\text{realistic, unknown size}} = \mathcal{M}(D_T; \widehat{k_T}, n, \widehat{p}_\epsilon^{\text{initial}}, \widehat{\sigma}_p^2, \alpha) + Y$ $\hspace{1cm}$ Final estimate

---

The second significant change in Algorithm 3 is how the sensitivity parameter Λ is chosen. The final statistic is more sensitive under the view of private-size user level privacy; the weight of every user can change as a result of a single user changing the amount of data they hold (due to the normalisation constant changing). Thus, the formula for the scale of the noise given in Algorithm 3 is higher than the noise added in Algorithm 2. Further, Λ as defined in Algorithm 3 is not an upper bound on the local sensitivity for *all* databases, although with high probability it is an upper bound on the local sensitivity of all databases that lie in a neighbourhood of D. So, we use a standard framework from the differential privacy literature called propose-test-release (PTR) to privately verify that Λ is indeed an upper bound on the local sensitivity of all databases in a neighbourhood of D, which allows us to safely add noise proportional to Λ to privatise the final statistic.

There are several existing algorithms in the literature that can be used to privately estimate the $\frac{\log n}{\epsilon}$-th order statistic $\widehat{k_T}$. A simple algorithm [10, 32, 19, 3, 4] that estimates the order statistic using the common differential privacy framework called the exponential mechanism [28] is sufficient up to a constant factor. For a full description of this algorithm, as well as its accuracy guarantees see [4]. In order for this algorithm to produce accurate results, we need an upper bound on the maximum number of data points a single user can have; we will call this number $k_{\max}$.

Algorithm 4 Truncated weighted mean, $\mathcal{M}(\cdot; k_{\max}, n, \hat{p}, \hat{\sigma_p}^2, \alpha)$

Input: number of users n, number of samples held by each user $(k_1, \ldots, k_n)$, user-level estimates $(\widehat{p}_1, \cdots, \widehat{p}_n)$, upper bound $k_{\max}$, lower bound $k_{\min}$, mean estimate $\hat{p}$, variance estimate $\hat{\sigma}_p^2$, accuracy on mean estimate α

1: **for** $i \in [n]$ **do**
2: $\quad \widehat{a}_i = \hat{p} - \alpha - f_{\mathcal{D}}^{k_i}(n, \hat{\sigma}_p^2, \beta)$
3:
4: $\quad \widehat{b}_i = \hat{p} + \alpha + f_{\mathcal{D}}^{k_i}(n, \hat{\sigma}_p^2, \beta)$
5:
6: $\quad k_i' = \min\{k_i, k_{\max}\}$
7: $\quad \widehat{\sigma}_i^2 = \frac{1}{k_i'}(\hat{p} - (\hat{p})^2) + (1 - \frac{1}{k_i'})\hat{\sigma}_p^2.$
8: $\quad v_i = \frac{1}{\widehat{\sigma}_i^2}$
9: Return $\dfrac{\sum_{i \in [n]} v_i [\widehat{p}_i]_{\widehat{a}_i}^{\widehat{b}_i}}{\sum_{i \in [n]} v_i}$

A κ-neighbour of D is a database D' of size n that differs from D on the data of at most κ data subjects. Given a function f from the set of databases to $\mathbb{R}$, and a database D, the local sensitivity of f at D is defined by $\Delta(f; D) = \max_{D' \text{neighbour of} D} |f(D) - f(D')|$.

Theorem D.1. *For any $\epsilon > 0$ and $\delta \in [0, 1]$, Algorithm 3 is $(3\epsilon, 2\delta)$-DP. If the conditions of Theorem 4.1 hold and*

- $\epsilon \geq \frac{2 \log n}{n}$,

- $k_{\max} \leq \sqrt{n}$ *and* $\beta \leq 1/\sqrt{n}$,

- *if k_{med} is the median k_i value then* $\frac{k_{\max}}{k_{med}} \leq$
  $\min\left\{\frac{1}{2}(n - \Upsilon - 1), \frac{n-1}{\Upsilon+1}, \frac{\epsilon^3(n/2 - \log n - 1)}{\log n \log(1/\delta) \log(1/\beta)}, \frac{(n/4-1)\epsilon}{3\ln(2/\delta)}, \frac{n/2 - \frac{\log n}{\epsilon}}{\frac{\log n}{\epsilon}}\right\}$,

- $\max\{\alpha, \sigma_{k_{\max}}\} \leq f_{\mathcal{D}}^{k_{\max}}(n, \hat{\hat{\sigma}}_p^2, \beta)$, *and*

- *for any set $I \subset \{1, m\}$ then with probability $1 - \beta$,* $\left|\frac{\sum_{i \in I} v_i \widehat{p}_i}{\sum_{i \in I} v_i} - p\right| \leq$
  $2\text{Var}\left(\frac{\sum_{i \in I} v_i \widehat{p}_i}{\sum_{i \in I} v_i}\right) \log(1/\beta)$,

then with probability $1 - 4\beta$, $\text{Var}(\hat{p}_\epsilon^{\text{realistic, unknown size}}) \leq \tilde{O}\left(\text{Var}(\widehat{p}^{\text{realistic}}) + \frac{\text{Var}(\hat{p}_\epsilon^{\text{realistic}})}{\sqrt{\epsilon}}\right)$

Theorem D.1 implies that when ϵ is sufficiently large ($\epsilon \geq \frac{\log n}{n}$) then the variance of $\hat{p}_\epsilon^{\text{realistic, unknown size}}$ is within a constant of $\hat{p}_\epsilon^{\text{realistic}}$, except that the final noise added for privacy is $\tilde{O}(1/\sqrt{\epsilon})$ larger. While the conditions of this theorem may seem intimidating, note that none of them are particularly stringent. The first condition $\epsilon \geq 2 \log n/n$ is a reasonable assumption on ϵ (we note that essentially no learning is possible with $\epsilon < 1/n$). The second condition on $k_{\max}$ is a reasonable assumption if the number of users n is large, as may be the case in practice. The third condition, up to logarithmic factors, can be viewed as $k_{\max}/k_{\text{med}} \leq \epsilon^3 n$. This is a stronger assumption than was required for Theorem 4.1 but provided reasonable values of ϵs, it is not prohibitive in practice. The final condition is simply a concentration bound on the weighted sums; this condition holds for Gaussians.

The proof that Algorithm 3 is $(3\epsilon, 2\delta)$-DP is fairly routine so we will present it first. The population is broken into three cohorts. Let us consider each cohort individually. First, consider the $\log n$ individuals with the most data. They participate in private releases in lines (3) ((ϵ, δ)-DP), and (5) (ϵ-DP). Using the simple composition rule of differential privacy [12], Algorithm 3 is $(2\epsilon, \delta)$-DP with respect to these users.

Next, consider the $1/10$th of users with the least data. These users participate in lines (2) ((ϵ, δ)-DP) and (5) (ϵ-DP). Again using the simple composition rule of differential privacy, Algorithm 3 is $(2\epsilon, \delta)$-DP with respect to these users.

Finally, let us consider the the group consisting of users $i \in [\log n + 1, 9n/10]$. These users first participate in line (5) (ϵ-DP). The post-processing inequality of differential privacy states that we can now use these statistics in the subsequent computations without paying additionally for their privacy. Lines (7) - (9) are pre-processing for the computation of $\tilde{N}$. The algorithm releasing $\tilde{N}$ is a simple application of the Laplace mechanism since each $v_i \in [0, \frac{1}{\widehat{\sigma_{\min}}^2}]$, and hence is ϵ-differentially private. The computation of Λ in line (13) does not additionally touch the users data. The final estimate $\widehat{p}_\epsilon^{\text{realistic}}$ is an application of the propose-test-release framework on the function $\mathcal{M}(\cdot; \widehat{k_T}, n, \widehat{p}_\epsilon^{\text{initial}}, \widehat{\sigma}_p^2)$ with proposed sensitivity Λ. This is a generic application of the propose-test-release framework, so we refer the reader to [10] for a proof that this final step of the algorithm is (ϵ, δ)-differentially private. Therefore, again using the composition theorem, Algorithm 3 is $(3\epsilon, 2\delta)$-DP with respect to this final set of users.

For the remainder of this section, we will focus on outlining the proof of the utility claims. Let us focus first on the impact of the use of propose-test-release (PTR). The two relevant components for the how the PTR component of Algorithm 3 affects the utility are the scale of Λ/ϵ and the probability that the proposed sensitivity is too small resulting in the algorithm ending in line (19), rather than line (22). The impact of the former is easy to analyse since the noise added is simply output perturbation. In order to show that the PTR ends in line (22) with high probability, we need to show that with high probability (over the randomness in the samples), κ^* as defined in line (16) is large enough. Since this claim is really about $\mathcal{M}(\cdot; k_{\max}, n, \hat{p}, \hat{\sigma}_p^2)$, we will state and prove this claim in the notation of Algorithm 4.

Lemma D.2. *Given $k_{\min} < k_{\max}$, $n \in \mathbb{N}$, $\hat{p} \in [0, 1]$, $\hat{\sigma}_p^2 \in [0, 1]$, $\Upsilon \in \mathbb{N}$, and a database $D = \{(\widehat{p}_i, k_i)\}_{i=1}^n$ such that $\widehat{p}_i \sim \mathcal{D}(k_i)$. If the conditions of Theorem D.1 hold then with probability $1 - \beta$, D is such that for any κ-neighbour of D, D' where $0 \le \kappa \le \Upsilon$ then*

$$\Delta(\mathcal{M}(\cdot; k_{\max}, m, \hat{p}, \hat{\sigma}_p^2, \alpha); D') \le 8B \frac{v_{k_{\max}} f_{\mathcal{D}}^{k_{\max}}(n, \hat{\sigma}_p^2, \beta)}{\sum_{i=1}^n v_i},$$

where $B = \sqrt{\Upsilon + 1} \log n + \frac{\log(1/\beta)}{\sqrt{\Upsilon + 1}}$.

Up to the factor B, the local sensitivity is exactly what we expect from the Algorithm 2. In order for our application of propose-test-release in Algorithm 3 to end in the desired result, we need to set $\Upsilon = \frac{\log(1/\delta)}{\epsilon} + \frac{\ln(1/\delta)\ln(1/\gamma)}{\epsilon}$. This means that up to logarithmic terms, the additional factor B is $O(1/\sqrt{\epsilon})$. If ϵ is constant then this factor is also constant, although it may be a notable factor when ϵ is small. This inclusion of this extra factor is actually a result of a union bound in the proof of Lemma D.2, and we leave as an open question whether this factor can be improved or perhaps removed entirely.

Proof of Lemma D.2. Let $\sigma_{\max}^2 = \frac{1}{k_{\min}}(\hat{p} - (\hat{p})^2) + (1 - \frac{1}{k_{\min}})\hat{\sigma}_p^2$, $\sigma_{\min}^2 = \frac{1}{k_{\max}}(\hat{p} - (\hat{p})^2) + (1 - \frac{1}{k_{\max}})\hat{\sigma}_p^2$, $v_{\max} = 1/\sigma_{\min}^2$ and $v_{\min} = 1/\sigma_{\max}^2$, and $A = \frac{1}{2}\frac{n - \Upsilon - 1}{\Upsilon + 1}$. Note that as in Equation (7), the condition that $k_{\max}/k_{\min} \le A$ implies that $\sigma_{\max}^2 \le A\sigma_{\min}^2$ and, equivalently, $v_{\max} \le Av_{\min}$.

Let $D = \{(\widehat{p}_i, k_i)\}_{i=1}^n$ be a dataset of size n where each $\widehat{p}_i \sim \mathcal{D}(k_i)$ where $\mathcal{D}$ has mean p and variance σ_p^2. It suffices to show that for any database D', which is a κ-neighbour of D where $0 \le \kappa \le \Upsilon + 1$, and any $j \in [n]$, if D'_{-j} is D' where the data of the jth data subject has been removed, then,

$$\left| \mathcal{M}(D'; k_{\max}, n, \hat{p}, \hat{\sigma}_p^2, \alpha) - \mathcal{M}(D'_{-j}; k_{\max}, n, \hat{p}, \hat{\sigma}_p^2, \alpha) \right|$$

$$\le 4\left(\sqrt{\Upsilon + 1}\log n + \frac{\log(1/\beta)}{\sqrt{\Upsilon + 1}}\right) \frac{v_{k_{\max}} f_{\mathcal{D}}^{k_{\max}}(n, \hat{\sigma}_p^2, \beta)}{\sum_{i=1}^n v_i}. \tag{12}$$

The final result is then a simple application of the triangle inequality.

Our proof that Equation (12) holds with high probability for all κ-neighbours of D relies on the fact that the data points in D are samples from $\mathcal{D}(k_i)$. In particular, it will rely on the fact that with

probability $1 - \beta$, D is such that all subsets S of D of size at least $n - \Upsilon - 1$, $\mathcal{M}(S; k_{\max}, n, \hat{p}, \hat{\sigma_p}^2)$ is concentrated around p. Let I be a subset of $[n]$ of size $n - \kappa$ where $\kappa \leq \Upsilon + 1$. Then

$$\text{Var}(\mathcal{M}(S; k_{\max}, n, \hat{p}, \hat{\sigma_p}^2, \alpha)) = \text{Var}\left(\frac{\sum_{i \in I} v_i \widehat{p}_i}{\sum_{i \in I} v_i}\right)$$

$$= \frac{\sum_{i \in I} \frac{1}{(\hat{\sigma}_i^2)^2} \sigma_i^2}{\left(\sum_{i \in I} \frac{1}{\hat{\sigma}_i^2}\right)^2}$$

$$\leq 2 \frac{1}{\sum_{i \in I} \frac{1}{\hat{\sigma}_i^2}}$$

$$\leq 2 \frac{1}{\frac{n-\kappa}{A} \frac{1}{\sigma_{\min}^2}}$$

$$= \frac{2A}{n - \kappa} \sigma_{\min}^2$$

$$\leq \frac{\sigma_{\min}^2}{\Upsilon + 1}$$

where the first inequality follows from $\sigma_i^2 \leq 2\hat{\sigma}_i^2$ (by Lemma 4.2), the second from the fact that $\hat{\sigma}_i^2 \leq A\sigma_{\min}^2$ for all $i \in [n]$ and the final from the definition of A. Let $\Gamma = \sum_{\kappa=0}^{\Upsilon+1} \binom{n}{\kappa}$ be the number of subsets of D of size greater than $n - \upsilon - 1$. Therefore, by the concentration on the assumption of $\mathcal{M}(S; k_{\max}, n, \hat{p}, \hat{\sigma_p}^2)$, with probability $1 - \frac{\beta}{\Gamma}$,

$$\left|\mathcal{M}(S; k_{\max}, n, \hat{p}, \hat{\sigma_p}^2, \alpha)) - p\right| \leq \frac{2\sigma_{\min} \log \frac{\Gamma}{\beta}}{\sqrt{\Upsilon + 1}} \tag{13}$$

Applying a union bound, with probability $1 - \beta$, eqn (13) holds simultaneously for all subsets of D of sufficiently large size. For the remainder of the proof, let us assume that this holds.

Let D' be a κ-neighbour of D where $0 \leq \kappa \leq \Upsilon + 1$. Without loss of generality, assume that $D' = \{(\widehat{p}'_i, k'_i)\}_{i=1}^n$ where $(\widehat{p}'_i, k'_i) = (\widehat{p}_i, k_i)$ for $i \in [n - k]$. In order to use this simplification, we will not assume that the k'_i are in descending order. Let the v_i be the un-normalised weights corresponding to D', as defined in line (8) of Algorithm 4. Note that the v_i depends only on the data of user i, not the data of any other individual in the data set. Then

$$\left|\mathcal{M}(D'; k_{\max}, n, \hat{p}, \hat{\sigma_p}^2, \alpha) - \mathcal{M}(D'_{-j}; k_{\max}, n, \hat{p}, \hat{\sigma_p}^2, \alpha)\right| = \left|\frac{\sum_{i=1}^n v_i \widehat{p}'_i}{\sum_{i=1}^n v_i} - \frac{\sum_{i=1, i \neq j}^n v_i \widehat{p}'_i}{\sum_{i=1, i \neq j}^n v_i}\right|$$

$$= \frac{v_j}{\sum_{i=1}^n v_i} \left|\widehat{p}'_j - \frac{\sum_{i=1, i \neq j}^n v_i \widehat{p}'_i}{\sum_{i=1, i \neq j}^n v_i}\right|$$

$$\leq \frac{v_j}{\sum_{i=1}^n v_i} \left(\left|\widehat{p}'_j - \frac{\sum_{i=1}^{n-\kappa} v_i \widehat{p}_i}{\sum_{i=1}^{n-\kappa} v_i}\right| + \left|\frac{\sum_{i=1}^{n-\kappa} v_i \widehat{p}_i}{\sum_{i=1}^{n-\kappa} v_i} - \frac{\sum_{i=1, i \neq j}^n v_i \widehat{p}'_i}{\sum_{i=1, i \neq j}^n v_i}\right|\right).$$
$$\tag{14}$$

We will bound the two terms separately. For the first term in Equation (14), we will use the fact that $\frac{\sum_{i=1}^{n-\kappa} v_i \widehat{p}'_i}{\sum_{i=1}^{n-\kappa} v_i}$ is concentrated around p, and $\hat{p}'_j$ is truncated to within $\alpha + f_{\mathcal{D}}^{k'_j}(n, \sigma_p^2, \beta)$ of p. So

$$\left|\widehat{p}'_j - \frac{\sum_{i=1}^{n-\kappa} v_i \widehat{p}_i}{\sum_{i=1}^{n-\kappa} v_i}\right| \leq \max\left\{2(\alpha + f_{\mathcal{D}}^{k'_j}(n, \sigma_p^2, \beta)), \frac{2\sigma_{\min} \log \frac{\Gamma}{\beta}}{\sqrt{\Upsilon + 1}}\right\} \leq 4 f_{\mathcal{D}}^{k'_j}(n, \sigma_p^2, \beta) \frac{\log \frac{\Gamma}{\beta}}{\sqrt{\Upsilon + 1}},$$

where the second inequality follows since $\max\{\alpha, \sigma_{\min}\} \leq f_{\mathcal{D}}^{k'_j}(n, \sigma_p^2, \beta)$.

Next, let us handle the second term in Equation (14):

$$\left| \frac{\sum_{i=1}^{n-1} v_i \widehat{p}_i'}{\sum_{i=1}^{n-1} v_i} - \frac{\sum_{i=1}^{n-\kappa} v_i \widehat{p}_i'}{\sum_{i=1}^{n-\kappa} v_i} \right| = \left| \frac{\sum_{i=n-\kappa+1}^{n-1} v_i}{\sum_{i=1}^{n-1} v_i} \left(\frac{\sum_{i=n-\kappa+1}^{n-1} v_i \widehat{p}_i'}{\sum_{i=n-\kappa+1}^{n-1} v_i} - \frac{\sum_{i=1}^{n-\kappa} v_i \widehat{p}_i'}{\sum_{i=1}^{n-\kappa} v_i} \right) \right|$$

$$\leq \left(\frac{\sum_{i=n-\kappa+1}^{n-1} v_i}{\sum_{i=1}^{n-1} v_i} \right) \left(\frac{\sum_{i=n-\kappa+1}^{n-1} v_i \left| \widehat{p}_i' - \frac{\sum_{i=1}^{n-\kappa} v_i \widehat{p}_i'}{\sum_{i=1}^{n-\kappa} v_i} \right|}{\sum_{i=n-\kappa+1}^{n-1} v_i} \right)$$

$$\leq \left(\frac{\sum_{i=n-\kappa+1}^{n-1} v_i}{\sum_{i=1}^{n-1} v_i} \right) \left(\frac{\sum_{i=n-\kappa+1}^{n-1} v_i \max\{(2\alpha + 2f_{\mathcal{D}}^{k_i'}(n, \widehat{\sigma}_p^2, \beta)), 2\sigma_{\min} \log \frac{\Gamma}{\beta}\}}{\sum_{i=n-\kappa+1}^{n-1} v_i} \right)$$

$$\leq \left(\frac{\sum_{i=n-\kappa+1}^{n-1} v_i}{\sum_{i=1}^{n-1} v_i} \right) \left(\frac{\sum_{i=n-\kappa+1}^{n-1} v_i 4 f_{\mathcal{D}}^{k_i'}(n, \widehat{\sigma}_p^2, \beta) \frac{\log \frac{\Gamma}{\beta}}{\sqrt{\Upsilon+1}}}{\sum_{i=n-\kappa+1}^{n-1} v_i} \right)$$

$$\leq \left(\frac{4 \sum_{i=n-\kappa+1}^{n-1} v_i f_{\mathcal{D}}^{k_i'}(n, \widehat{\sigma}_p^2, \beta) \frac{\log \frac{\Gamma}{\beta}}{\sqrt{\Upsilon+1}}}{\sum_{i=1}^{n-1} v_i} \right)$$

$$\leq 4\kappa \frac{\log \frac{\Gamma}{\beta}}{\sqrt{\Upsilon+1}} \left(\frac{\max_i v_i f_{\mathcal{D}}^{k_i'}(n, \widehat{\sigma}_p^2, \beta)}{\sum_{i=1}^{n-1} v_i} \right)$$

Since $v_i = \frac{1}{\sigma_{k_i'}^2}$ and $f_{\mathcal{D}}^{k_i'}(n, \sigma_p^2, \beta) \leq 2\sigma_{k_i'}^2 \log n$, $\max_i v_i f_{\mathcal{D}}^{k_i'}(n, \sigma_p^2, \beta) \leq v_{k_{\max}} f_{\mathcal{D}}^{k_{\max}}(n, \sigma_p^2, \beta)$.
Also, $\sigma_{k_i'}^2 \leq A\sigma_{k_{\max}}^2$ so we have $\sum_{i=1}^{n-1} v_i \geq \frac{n-1}{A} v_{k_{\max}}$. Therefore,

$$\left| \frac{\sum_{i=1}^{n-1} v_i \widehat{p}_i'}{\sum_{i=1}^{n-1} v_i} - \frac{\sum_{i=1}^{n-\kappa} v_i \widehat{p}_i'}{\sum_{i=1}^{n-\kappa} v_i} \right| \leq 4\kappa \frac{\log \frac{\Gamma}{\beta}}{\sqrt{\Upsilon+1}} \frac{v_{k_{\max}} f_{\mathcal{D}}^{k_{\max}}(n, \sigma_p^2, \beta)}{\frac{n-1}{A} v_{k_{\max}}}$$

$$\leq \frac{4\sqrt{\Upsilon+1} A \log \frac{\Gamma}{\beta}}{n-1} f_{\mathcal{D}}^{k_{\max}}(n, \sigma_p^2, \beta)$$

$$\leq 2 \frac{\log \frac{\Gamma}{\beta}}{\sqrt{\Upsilon+1}} f_{\mathcal{D}}^{k_{\max}}(n, \sigma_p^2, \beta)$$

where the second inequality follows from $\kappa \leq \Upsilon+1$ and the third inequality follows from $A \leq \frac{n-1}{2(\Upsilon+1)}$ by assumption. Therefore,

$$\left| \mathcal{M}(D'; k_{\max}, n, \hat{p}, \hat{\sigma_p}^2, \alpha) - \mathcal{M}(D'_{-j}; k_{\max}, n, \hat{p}, \hat{\sigma_p}^2, \alpha) \right|$$

$$\leq \frac{v_j}{\sum_{i=1}^n v_i} \cdot 4 \frac{\log \frac{\Gamma}{\beta}}{\sqrt{\Upsilon+1}} (f_{\mathcal{D}}^{k_j'}(n, \hat{\sigma_p}^2, \beta) + f_{\mathcal{D}}^{k_{\max}}(n, \hat{\sigma_p}^2, \beta))$$

$$\leq 4 \frac{\log \frac{\Gamma}{\beta}}{\sqrt{\Upsilon+1}} \frac{v_j}{\sum_{i=1}^n v_i} \cdot f_{\mathcal{D}}^{k_j'}(n, \hat{\sigma_p}^2, \beta)$$

$$\leq 4 \frac{\log \frac{\Gamma}{\beta}}{\sqrt{\Upsilon+1}} \frac{v_j}{\sum_{i=1}^n v_i} \cdot f_{\mathcal{D}}^{k_j'}(n, \hat{\sigma_p}^2, \beta)$$

Note that $\Gamma \leq n^{\Upsilon+1}$. Taking the max over j, we again have that $\max_j v_j f_{\mathcal{D}}^{k_j'}(n, \hat{\sigma}_p{}^2, \beta) \leq v_{k_{\max}} f_{\mathcal{D}}^{k_{\max}}(n, \hat{\sigma}_p{}^2, \beta)$ so for all j,

$$\left| \mathcal{M}(D'; k_{\max}, n, \hat{p}, \hat{\sigma}_p{}^2, \alpha) - \mathcal{M}(D'_{-j}; k_{\max}, n, \hat{p}, \hat{\sigma}_p{}^2, \alpha) \right|$$

$$\leq 4 \frac{\log \frac{\Gamma}{\beta}}{\sqrt{\Upsilon+1}} \frac{v_{k_{\max}} f_{\mathcal{D}}^{k_{\max}}(n, \hat{\sigma}_p{}^2, \beta)}{\sum_{i=1}^n v_i}$$

$$\leq 4 \left(\sqrt{\Upsilon+1} \log n + \frac{\log(1/\beta)}{\sqrt{\Upsilon+1}} \right) \frac{v_{k_{\max}} f_{\mathcal{D}}^{k_{\max}}(n, \hat{\sigma}_p{}^2, \beta)}{\sum_{i=1}^n v_i}$$

$\square$

Proof of Theorem D.1. The main component remaining to prove is that truncating at $\frac{1}{\widehat{\sigma_{\min}}^2}$ rather than the optimal truncation does not affect the utility by more than a constant factor, provided ϵ is not too small. Recall $k_1 \geq k_2 \geq \cdots \geq k_n$. Firstly, we need to show that $\widehat{k_T}$ is a sufficiently good estimate of $k_{\frac{2 \log n}{\epsilon}}$. A careful instantiation of the exponential mechanism provides us with a ϵ-DP estimator of the $\frac{2 \log n}{\epsilon}$-th order statistic that has the guarantee that with probability $1 - \beta$, $k_{\frac{2 \log n}{\epsilon} + \frac{1}{\epsilon}(\ln k_{\max} + \ln(1/\beta))} \leq \widehat{k_T} \leq k_{\frac{2 \log n}{\epsilon} - \frac{1}{\epsilon}(\ln k_{\max} + \ln(1/\beta))}$. Since by assumption $k \leq \sqrt{n}$ and $\beta \leq 1/\sqrt{n}$, this implies that with probability $1 - \beta$, $k_{\frac{3 \log n}{\epsilon}} \leq \widehat{k_T} \leq k_{\frac{\log n}{\epsilon}}$. That is, only $\frac{\log n}{\epsilon}$ more data points than desired will be truncated.

Next, we need to show that truncating at any point within this range provides an estimator with accuracy competitive with the optimal truncation. The variance of $\hat{p}_\epsilon^{\text{realistic, unknown size}}$ can be written as two terms, the variance that exists in the non-private setting, and the additional noise due to privacy;

$$\text{Var}(\hat{p}_\epsilon^{\text{realistic}}) = \underbrace{\frac{\sum_{i=\log n+1}^{9n/10} \min\left\{ \frac{T^2}{\hat{\sigma}_i^2}, \frac{1}{\hat{\sigma}_i^4} \right\} \text{Var}([\hat{p}_i]_{\hat{a}_i}^{\hat{b}_i})}{\left(\sum_{i=\log n+1}^{9n/10} \min\left\{ \frac{T}{\hat{\sigma}_i}, \frac{1}{\hat{\sigma}_i^2} \right\} \right)^2}}_{\text{non-private term}} + \underbrace{\frac{\left(8 \left(\sqrt{\Upsilon+1} \log n + \frac{\log(1/\beta)}{\sqrt{\Upsilon+1}} \right) \frac{f_{\mathcal{D}}^{\widehat{k_T}}(n, \hat{\sigma}_p{}^2, \beta)}{\widehat{\sigma_{\min}}^2 \widehat{N}} \right)^2}{\epsilon^2}}_{\text{private term}},$$

where $\Upsilon = \frac{\log(1/\delta)}{\epsilon} + \frac{\ln(1/\delta) \ln(1/\gamma)}{\epsilon}$. The truncation has opposite effects on each of these terms. As T increases, the private term decreases while the non-private term increases. When we set $T = 1/\text{Var}(\mathcal{D}(k_{\frac{2 \log n}{\epsilon} + K}))$, where $K \in [-\frac{\log n}{\epsilon}, \frac{\log n}{\epsilon}]$ then if K is negative, no truncation occurs and the non-private term is optimal. If K is positive then only a small number of data points are truncated so the non-private term is close to it's optimal value:

$$\frac{\sum_{i=\log n+1}^{9n/10} \min\left\{ \frac{T^{*2}}{\hat{\sigma}_i^2}, \frac{1}{\hat{\sigma}_i^4} \right\} \text{Var}([\hat{p}_i]_{\hat{a}_i}^{\hat{b}_i})}{\left(\sum_{i=\log n+1}^{9n/10} \min\left\{ \frac{T^*}{\hat{\sigma}_i}, \frac{1}{\hat{\sigma}_i^2} \right\} \right)^2} \leq O\left(\frac{\sum_{i=\log n+K}^{9n/10} \frac{1}{\hat{\sigma}_i^4} \text{Var}([\hat{p}_i]_{\hat{a}_i}^{\hat{b}_i})}{\left(\sum_{i=\log n+K}^{9n/10} \frac{1}{\hat{\sigma}_i^2} \right)^2} \right)$$

$$\leq O\left(\frac{\sum_{i=\log n+1}^{9n/10} \frac{1}{\hat{\sigma}_i^4} \text{Var}([\hat{p}_i]_{\hat{a}_i}^{\hat{b}_i})}{\left(\sum_{i=\log n+1}^{9n/10} \frac{1}{\hat{\sigma}_i^2} \right)^2} \right)$$

where the first inequality follows from the fact that if $k_{\max}/k_{\text{med}} \leq \frac{n/2 - \frac{\log n}{\epsilon}}{\frac{\log n}{\epsilon}} \leq \frac{n/2 - \frac{2 \log n}{\epsilon} - K}{\frac{2 \log n}{\epsilon} - K}$ and $\frac{\log n}{\epsilon} \leq n/2$ then deleting K points has only a constant factor impact on the variance (this argument is identical to that made in eqn (10) in the proof of Theorem 4.1). The second inequality follows from the fact that adding more high quality data points only improves the variance of the estimator. Therefore, the non-private term in the variance is within a constant factor of optimal.

Next, we will show under the conditions outlines in the theorem, the non-private term is dominates the variance. The normalisation term also appears in the private term but as an approximation:

$$\widehat{N} = \sum_{j=\log n+1}^{n/2} \min\left\{ \frac{T}{\hat{\sigma}_i}, \frac{1}{\hat{\sigma}_i^2} \right\} + \text{Lap}\left(\frac{1}{\epsilon \widehat{\sigma_{\min}}^2} \right) - \frac{1}{\epsilon \widehat{\sigma_{\min}}^2} \ln(2\delta).$$

With probability $1 - \beta$,

$$\tilde{N} \geq \sum_{j=\log n+1}^{9n/10} \min\left\{\frac{T}{\widehat{\sigma}_i}, \frac{1}{\widehat{\sigma}_i^2}\right\} - 2\frac{1}{\epsilon\widehat{\sigma_{\min}}^2}\ln(2\delta)$$

$$\geq \sum_{j=\log n+1}^{n/4} \min\left\{\frac{T}{\widehat{\sigma}_i}, \frac{1}{\widehat{\sigma}_i^2}\right\} + \sum_{j=n/4+1}^{n/2} \min\left\{\frac{T}{\widehat{\sigma}_i}, \frac{1}{\widehat{\sigma}_i^2}\right\} - 2\frac{1}{\epsilon\widehat{\sigma_{\min}}^2}\ln(2\delta)$$

$$\geq \sum_{j=\log n+1}^{n/4} \min\left\{\frac{T}{\widehat{\sigma}_i}, \frac{1}{\widehat{\sigma}_i^2}\right\} + (n/4-1)\frac{1}{\widehat{\sigma}_{k_{\mathrm{med}}}^2} - 2\frac{1}{\epsilon\widehat{\sigma_{\min}}^2}\ln(2\delta)$$

$$\geq \sum_{j=\log n+1}^{n/4} \min\left\{\frac{T}{\widehat{\sigma}_i}, \frac{1}{\widehat{\sigma}_i^2}\right\}$$

$$\geq \frac{1}{2} \sum_{j=\log n+1}^{9n/10} \min\left\{\frac{T}{\widehat{\sigma}_i}, \frac{1}{\widehat{\sigma}_i^2}\right\}$$

where the final inequality comes from high probability bounds on the Laplacian distribution, the second inequality is simply separating the sum into two pieces and removing the contribution of users $i \in [n/2 + 1, 9n/10]$, the third inequality comes from the fact that any user with more than k_{med} data points has weight larger than $1/\widehat{\sigma}_{k_{\mathrm{med}}}^2$. The fourth inequality follows from $\frac{\widehat{\sigma}_{k_{\mathrm{med}}}}{\widehat{\sigma_{\min}}^2} \leq \frac{(n/4-1)\epsilon}{3\ln(2/\delta)}$. Now, let us turn to the proof that the non-private noise is dominant when ϵ is not too large. To see this note that the non-private term satisfies

$$\frac{\sum_{i=\log n+1}^{9n/10} \min\left\{\frac{T^2}{\widehat{\sigma}_i^2}, \frac{1}{\widehat{\sigma}_i^4}\right\} \mathrm{Var}([\widehat{p}_i]_{\widehat{a}_i}^{\widehat{b}_i})}{\left(\sum_{i=\log n+1}^{9n/10} \min\left\{\frac{T}{\widehat{\sigma}_i}, \frac{1}{\widehat{\sigma}_i^2}\right\}\right)^2} \geq \Omega\left(\frac{\sum_{i=\log n+1}^{9n/10} \min\{T^2, \frac{1}{\sigma_i^2}\}}{\left(\sum_{i=\log n+1}^{9n/10} \min\{\frac{T}{\sigma_i}, \frac{1}{\sigma_i^2}\}\right)^2}\right)$$

where the inequality comes from $\mathrm{Var}([\widehat{p}_i]_{\widehat{a}_i}^{\widehat{b}_i}) \geq \frac{1}{2}\sigma_i^2$ and $\widehat{\sigma}_i^2$ is within a constant multiplicative factor of σ_i^2. Further, the private term satisfies

$$\frac{\left(8\left((\Upsilon+1)\log n + \log(1/\beta)\right)\frac{f_{\mathcal{D}}^{\widehat{k_T}}(n,\widehat{\sigma_p}^2,\beta)}{\widehat{\sigma_{\min}}^2\widehat{N}}\right)^2}{\epsilon^2} = O\left(\frac{(\log n\log(1/\delta)\log(1/\beta))^2}{\sigma_{\min}^2 N^2\epsilon^3}\right)$$

$$= O\left(\frac{(\log n\log(1/\delta)\log(1/\beta))^2}{\sigma_T^2\epsilon^3(\sum_{i=\log n+1}^{9n/10}\min\{\frac{T}{\sigma_i}, \frac{1}{\sigma_i^2}\})^2}\right)$$

Now, comparing these two terms we can see that the non-private term dominates when:

$$\frac{\sum_{i=\log n+1}^{9n/10}\min\{T, \frac{1}{\sigma_i^2}\}}{\left(\sum_{i=\log n+1}^{9n/10}\min\{\frac{T}{\sigma_i}, \frac{1}{\sigma_i^2}\}\right)^2} = \Omega\left(\frac{(\log n\log(1/\delta)\log(1/\beta))^2}{\sigma_T^2\epsilon^3(\sum_{i=\log n+1}^{9n/10}\min\{\frac{T}{\sigma_i}, \frac{1}{\sigma_i^2}\})^2}\right).$$

That is, when:

$$\sum_{i=\log n+1}^{9n/10} \min\left\{T, \frac{1}{\sigma_i^2}\right\} \geq \left(\frac{\log n\log(1/\delta)\log(1/\beta)}{\sigma_{\min}\epsilon^{1.5}}\right)^2.$$

Noting that,

$$\sum_{i=\log n+1}^{9n/10} \min\left\{T^*, \frac{1}{\sigma_i^2}\right\} \geq (n/2 - \log n - 1)\frac{1}{\sigma_{\mathrm{med}}^2} \geq \left(\frac{\log n\log(1/\delta)\log(1/\beta)}{\sigma_{\min}\epsilon^{1.5}}\right)^2$$

where the first inequality is simply because more than $(n/2 - \log n)$ of the user have weight larger than the median weight, and the second inequality follows from the assumption that $\frac{k_{\max}}{k_{\mathrm{med}}} \leq \frac{\epsilon^3(n/2 - \log n - 1)}{\log n \log(1/\delta) \log(1/\beta)}$. Therefore, with high probability (based on the accuracy of $\widehat{k_T}$), truncating at $1/\sigma_{\min}^2$ rather than the optimal truncation T does not affect the variance of the estimator by more than a constant factor.

Now that we have established that the noise added for privacy is not too large, the only remaining potential point of failure for the algorithm is that the PTR component fails and the algorithm outputs $\widehat{p}_\epsilon^{\mathrm{initial}}$ rather than the more accurate weighted estimate. The fact that this does not happen with high probability is a direct corollary of Lemma D.2. $\qquad\square$

E Proofs from Section 5

Lemma 5.2. *Given $\widehat{p}_i \sim \mathcal{D}_p(k_i)$ with variance σ_i^2 for all $i \in [n]$ and $w \in [0,1]^n$ such that $\sum_{i=1}^n w_i = 1$, let $\widehat{p} = \sum_{i=1}^n w_i \widehat{p}_i + \mathrm{Lap}(\frac{\max_i w_i \sigma_i}{\epsilon})$. The variance of $\widehat{p}$ is minimized by the following weights: $\tilde{w}_i^* = \frac{\min\{1/\sigma_i^2, T/\sigma_i\}}{\sum_{j=1}^n \min\{1/\sigma_j^2, T/\sigma_j\}}$ for some T.*

Proof of Lemma 5.2. Let

$$w^* = \arg\min_{\substack{w \in [0,1]^n \\ \sum_{i=1}^n w_i = 1}} \mathrm{Var}(\widehat{p}) = \arg\min_{\substack{w \in [0,1]^n \\ \sum_{i=1}^n w_i = 1}} \sum_{i=1}^n w_i^2 \sigma_i^2 + \frac{\max_k w_k^2 \sigma_k^2}{\epsilon^2}$$

be an optimal weight vector that minimizes variance of $\widehat{p}$. We start with a few observations on structural properties of the optimal weights. Let $M = \{\arg\max_k w_k^* \sigma_k\}$ be the set of all users with maximum weighted-variance contribution to the estimate $\widehat{p}$.

First, notice that for all $i, j \in [n]$, if $w_i^* > w_j^*$ then $\sigma_i^2 \leq \sigma_j^2$. This follows since if $\sigma_i^2 > \sigma_j^2$ then $w_i^* \sigma_j^2 + w_j^* \sigma_i^2 < w_i^* \sigma_i^2 + w_j^* \sigma_j^2$ and $\max\{w_i^* \sigma_j^2, w_j^* \sigma_i^2\} \leq w_i^* \sigma_i$ which implies that swapping the weights of i and j would result in an estimator with lower variance. This is a contradiction given the definition of w^*.

Next, we show that if $i, j \notin M$ then $w_i^* \sigma_i^2 = w_j^* \sigma_j^2$. Suppose towards a contradiction that $w_i^* \sigma_i^2 < w_j^* \sigma_j^2$. Let $\alpha = \min\{\frac{w_j^* \sigma_j^2 - w_i^* \sigma_i^2}{\sigma_i^2 + \sigma_j^2}, \frac{\max_k w_k^* \sigma_k - w_i^* \sigma_i}{\sigma_i}, w_j^*\}$. Then $\alpha > 0$, and $(w_j^* - \alpha)\sigma_j, (w_i^* + \alpha)\sigma_i \in [0, \max_k w_k^* \sigma_k]$. Also,

$$(w_j^* - \alpha)^2 \sigma_j^2 + (w_i^* + \alpha)^2 \sigma_i^2 = w_j^{*2} \sigma_j^2 + w_i^{*2} \sigma_i^2 + \alpha^2(\sigma_i^2 + \sigma_j^2) - 2\alpha(w_j^* \sigma_j^2 - w_i^* \sigma_i^2)$$
$$= w_j^{*2} \sigma_j^2 + w_i^{*2} \sigma_i^2 + \alpha(\alpha(\sigma_i^2 + \sigma_j^2) - 2(w_j^* \sigma_j^2 - w_i^* \sigma_i^2))$$
$$< w_j^{*2} \sigma_j^2 + w_i^{*2} \sigma_i^2.$$

This implies that shifting α weight from w_i^* to w_j^* would reduce the variance of the estimator $\widehat{p}$ without changing the maximum weighted-variance, which is a contradiction of the optimality of w^*.

Define $H = \max_k w_k^* \sigma_k$ and note that there exists $R > 0$ such that $w_i^* = R/\sigma_i^2$ for all $i \notin M$. From these observations, there must exist some threshold T such that if $\sigma_i \geq 1/T$, then $w_i^* = R/\sigma_i^2$, and if $\sigma_i < 1/T$, then $w_i^* = H/\sigma_i$. By continuity, $H = RT$, and we can write the optimal weights as: $w_i^* = \min\{1/\sigma_i^2, T/\sigma_i\}R$. Since the weights w_i^* must sum to 1, we know that $R = \frac{1}{\sum_{j=1}^n \min\{1/\sigma_j^2, T/\sigma_j\}}$.

Thus the optimal weights are:

$$w_i^* = \frac{\min\{1/\sigma_i^2, T/\sigma_i\}}{\sum_{j=1}^n \min\{1/\sigma_j^2, T/\sigma_j\}},$$

for some appropriate threshold T.

$\qquad\square$

Let us recall some notation. Let $\mathcal{P}$ be a parameterized family of distributions $p \mapsto \mathcal{D}_p$, so $\mathbb{E}[\mathcal{D}_p]$. Given an estimator M, vector $\boldsymbol{q} \in [0,1]^n$ and set $I \subset [n]$, let

$$\mu_M(x_{[n] \setminus I}; \boldsymbol{q}) = \mathbb{E}_{\forall i \in I, x_i \sim \mathcal{D}_{q_i}(k_i), M}[M(x_1, \cdots, x_n)]$$

be the expectation taken only over the randomness of I and M. Note that in this notation, user i is sampling from a meta-distribution with mean q_i, which may be different for each user. We will abuse notation slightly and for $p \in [0, 1]$, we will let $\mu_M(x_{[n] \setminus I}; p) = \mu_M(x_{[n] \setminus I}; (p, \cdots, p))$. Let $\mu_M(\boldsymbol{q}) = \mu_M(\emptyset; \boldsymbol{q})$. When the estimator M is clear from context, we will simply use the notation $\mu(x_{[n] \setminus I}; \boldsymbol{p})$. Recall that for $p \in [0, 1]$ and $k \in \mathbb{N}$, $\phi_{p,k}$ is the probability density function of $\mathcal{D}_p(k)$.

Lemma 5.3. *Let $\mathcal{P}$ be a parameterized family of distributions $p \mapsto \mathcal{D}_p$ and suppose that M : $[0, 1]^n \to [0, 1]$ is an ϵ-DP estimator such that for all $p \in [1/3, 2/3]$, (1) M is unbiased, $\mu_M(p) = p$, and (2) the Fisher information of ϕ_{p,k_i} is inversely proportional to the variance $\mathrm{Var}(\mathcal{D}_p(k_i))$, $\int(\frac{\partial}{\partial p} \log \phi_{p,k_i}(x_i))^2 \phi_{p,k_i}(x_i) dx_i = O(\frac{1}{\mathrm{Var}(\mathcal{D}_p(k_i))})$, then there exists an estimator $M_{NL} \in \mathrm{NLE}$ such that*

$\max_{p \in [1/3, 2/3]}[\mathrm{Var}_{\forall i \in [n], x_i \sim \mathcal{D}(k_i), M_{NL}}(M_{NL})] \leq O(\max_{p \in [1/3, 2/3]}[\mathrm{Var}_{\forall i \in [n], x_i \sim \mathcal{D}(k_i), M}(M)])$.

Before we formally prove Lemma 5.3, let us start with some intuition for the proof. Given an estimator $M_{NL} \in \mathrm{NLE}$, the variance of M_{NL} can be written as

$$\mathrm{Var}(M_{NL}) \leq \sum_{i=1}^{n} w_i^2 \mathrm{Var}(\mathcal{D}(k_i)) + O(\frac{\max w_i \sigma_i}{\epsilon})^2. \tag{15}$$

That is, it can be decomposed as the variance contribution of each individual coordinate, and the variance contribution of the additional noise due to privacy. Lemma E.1 (proved in Appendix E) shows that the variance of any estimator M can be lower bounded by a similar decomposition. Since this involves considering the impact of each coordinate individually, the following notation will be useful. Given an estimator M, vector $\boldsymbol{q} \in [0, 1]^n$ and set $I \subset [n]$, let $\mu_M(x_{[n] \setminus I}; \boldsymbol{q}) = \mathbb{E}_{\forall i \in I, x_i \sim \mathcal{D}_{q_i}(k_i), M}[M(x_1, \cdots, x_n)]$ be the expectation over only randomness in I and M. Note that in this notation, user i is sampling from a meta-distribution with mean q_i, which may be different for each user. We will abuse notation slightly to let $\mu_M(\boldsymbol{q}) = \mu_M(\emptyset; \boldsymbol{q})$, and for $p \in [0, 1]$, we will let $\mu_M(x_{[n] \setminus I}; p) = \mu_M(x_{[n] \setminus I}; (p, \cdots, p))$. When the estimator M is clear from context, we will omit it.

The following lemma is proved later, outside the proof of Lemma 5.3.

Lemma E.1. *For any randomised mechanism $M : [0, 1]^n \to [0, 1]$,*

$$\mathrm{Var}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i), M}(M) \tag{16}$$
$$= \mathbb{E}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i), M}[(M(x_1, ..., x_n) - \mu(p))^2]$$
$$\geq \sum_{i=1}^{n} \mathbb{E}_{x_i \sim \mathcal{D}_p(k_i)}[(\mu(x_i; p) - \mu(p))^2] + \mathbb{E}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i), M}[(M(x_1, ..., x_n) - \mu(x_1, ..., x_n; p))^2]$$

In Equation (16), the first term is the sum of contributions to the variance of the individual terms x_i, and the second term is the contribution to the variance of the noise added for privacy. Now we want to define a weight vector $\mathbf{w}$ such that the terms in Equation (16) are lower bounded by the corresponding terms in Equation (15). The key component of the proof is the observation that if we let

$$w_i(p) = \frac{\partial}{\partial q_i} \mu(\boldsymbol{q}) \big|_{\boldsymbol{q} = (p, \cdots, p)} \tag{17}$$

then we can show that there exists a constant c such that

$$\mathbb{E}_{x_i \sim \mathcal{D}_p(k_i)}[(\mu(x_i; p) - \mu(p))^2] \geq c \cdot w_i(p)^2 \mathrm{Var}(\mathcal{D}_p(k_i)). \tag{18}$$

This controls the contribution of each individual coordinate to the variance of M. It remains only to control the contribution of the noise due to privacy. We show that there exists x_i, x_i' such that $|\mu(x_i; p) - \mu(x_i'; p)| \geq \Omega(w_i(p) \cdot \sqrt{\mathrm{Var}(\mathcal{D}_p(k_i))})$, which we show implies that,

$$\mathbb{E}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i), M}[(M(x_1, \cdots, x_n) - \mu(x_1, \cdots, x_n; p))^2] \geq \Omega(\frac{w_i(p)^2 \mathrm{Var}(\mathcal{D}_p(k_i))}{\epsilon^2}). \tag{19}$$

Intuitively, the worst-case $|\mu(x_i; p) - \mu(x_i'; p)|$ plays an analogous role to the sensitivity, since it captures the impact of changing one user's data. Since M is an ϵ-DP mechanism and $|\mu(x_i; p) - \mu(x_i'; p)|$ is at least $\Omega(w_i(p) \cdot \sqrt{\mathrm{Var}(\mathcal{D}_p(k_i))})$, we show that it must include noise with standard deviation of at least this magnitude over ϵ. This is consistent with, e.g., the Laplace Mechanism that adds noise with standard deviation $\Theta(\Delta f/\epsilon)$.

Combining Lemma E.1 with Equations (18) and (19) gives that the variance of M is at least,

$\mathrm{Var}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i), M}(M) \geq \sum_{i=1}^{n} c \cdot w_i(p)^2 \mathrm{Var}(\mathcal{D}_p(k_i)) + \Omega(\frac{w_i(p)^2 \mathrm{Var}(\mathcal{D}_p(k_i))}{\epsilon^2})$.

Finally, we must create a corresponding $M_{\texttt{NL}} \in \texttt{NLE}$ for comparison, using the same weights. Since $\sum_{i=1}^n w_i(p)$ as defined in Equation (17) need not equal 1, these weights will need to be normalized to sum to 1 to create an estimator in $\texttt{NLE}$. We need to show this normalisation does not substantially increase the variance of the resulting estimator. In order to show this, we show that there exists a $p^* \in [1/3, 2/3]$ such that $\sum_{i=1}^n w_i(p^*) \geq 1$, since normalizing the estimator by a factor of $\frac{1}{\sum_{i=1}^n w_i(p^*)}$ will affect the variance by a factor of $\frac{1}{(\sum_{i=1}^n w_i(p^*))^2}$, and thus if $\sum_{i=1}^n w_i(p^*) \geq 1$, then this will decrease variance. This desired fact follows from the definition of w_i, and the fact that M is unbiased. Now, if we define

$$M_{\texttt{NL}}(\mathbf{x}) = \frac{\sum_{i=1}^n w_i(p^*)x_i + \text{Lap}(\frac{\max_i w_i(p^*)\sqrt{\text{Var}(\mathcal{D}_p(k_i))}}{\epsilon})}{\sum_{i=1}^n w_i(p^*)},$$

then $M_{\texttt{NL}} \in \texttt{NLE}$ and $\text{Var}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i), M_{\texttt{TNL}}}(M_{\texttt{NL}}) = \Theta\left(\text{Var}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i), M}(M)\right).$

Proof of Lemma 5.3. We first apply Lemma E.1 to decompose the variance of the estimate computed by M as:

$\text{Var}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i), M}(M)$

$$\geq \sum_{i=1}^n \mathbb{E}_{x_i \sim \mathcal{D}_p(k_i)}[(\mu(x_i; p) - \mu(p))^2] + \mathbb{E}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i), M}[(M(x_1, \cdots, x_n) - \mu(x_1, \cdots, x_n; p))^2]$$

The first term is the sum of contributions to the variance of the individual terms x_i, and the second term is the contribution to the variance of the noise added for privacy. We will proceed by bounding these terms separately, starting with the first term.

First note that by definition,

$$\int (\mu(x_i; \boldsymbol{q}) - \mu(\boldsymbol{q}))\phi_{q_i, k_i}(x_i)dx_i = \mathbb{E}_{x_i \sim \mathcal{D}_{q_i}(k_i)}[\mu(x_i; \boldsymbol{q})] - \mu(\boldsymbol{q}) = 0.$$

Therefore, by taking the partial derivative with respect to q_i we have

$$\int \left[\left(\frac{\partial}{\partial q_i}(\mu(x_i; \boldsymbol{q}) - \mu(\boldsymbol{q})) \right) \phi_{q_i, k_i}(x_i) + (\mu(x_i; \boldsymbol{q}) - \mu(\boldsymbol{q}))\frac{\partial}{\partial q_i}\phi_{q_i, k_i}(x_i) \right] dx_i = 0.$$

Note that $\mu(x_i; \boldsymbol{q})$ is constant in q_i so rearranging, and noting that $\frac{\partial}{\partial q_i}\phi_{q_i, k_i}(x_i) = \phi_{q_i, k_i}(x_i)\left(\frac{\partial}{\partial q_i}\log\phi_{q_i, k_i}(x_i)\right)$ we have,

$$\int \left(\frac{\partial}{\partial q_i}\mu(\boldsymbol{q})\right)\phi_{q_i, k_i}(x_i)dx_i$$

$$= \int (\mu(x_i; \boldsymbol{q}) - \mu(\boldsymbol{q}))\phi_{q_i, k_i}(x_i)\left(\frac{\partial}{\partial q_i}\log\phi_{q_i, k_i}(x_i)\right)dx_i \qquad (20)$$

$$\leq \sqrt{\left(\int (\mu(x_i; \boldsymbol{q}) - \mu(\boldsymbol{q}))^2\phi_{q_i, k_i}(x_i)dx_i\right)\left(\int \left(\frac{\partial}{\partial q_i}\log\phi_{q_i, k_i}(x_i)\right)^2\phi_{q_i, k_i}(x_i)dx_i\right)}.$$
$$(21)$$

Let

$$w_i(p) = \int \left(\frac{\partial}{\partial q_i}\mu(\boldsymbol{q})\right)\phi_{q_i, k_i}(x_i)dx_i \Big|_{\boldsymbol{q}=(p,\cdots,p)} = \frac{\partial}{\partial q_i}\mu(\boldsymbol{q})\Big|_{\boldsymbol{q}=(p,\cdots,p)}$$

and note that by assumption there exists a constant c such that for all $i \in [n]$ and $q_i \in [1/3, 2/3]$,

$$\int \left(\frac{\partial}{\partial q_i}\log\phi_{q_i, k_i}(x_i)\right)^2\phi_{q_i, k_i}(x_i)dx_i \leq \frac{1}{c \cdot \text{Var}(\mathcal{D}_{q_i}(k_i))}.$$

Then evaluating both sides of Equation (21) at the constant vector $\boldsymbol{q} = (p, \cdots, p)$, we have

$$\left(\int (\mu(x_i; p) - \mu(p))^2\phi_{p, k_i}(x_i)dx_i\right) \geq \frac{w_i(p)^2}{\int \left(\frac{\partial}{\partial p}\log\phi_{p, k_i}(x_i)\right)^2\phi_{p, k_i}(x_i)dx_i} \geq c \cdot w_i(p)^2\text{Var}(\mathcal{D}_p(k_i)).$$

Now we have controlled the contribution of each individual coordinate to the variance of M, and it remains to control the contribution of the noise due to privacy.

We will show that for two independent samples x_i, x_i' drawn from $\mathcal{D}_p(k_i)$,

$$\mathbb{E}[(\mu(x_i; p) - \mu(x_i'; p))^2] \geq \Omega\Big(w_i(p)^2 \cdot \text{Var}(\mathcal{D}_p(k_i))\Big). \tag{22}$$

Letting

$$\alpha = \sqrt{\mathbb{E}[(\mu(x_i; p) - \mu(x_i'; p))^2]},$$

we can write

$$
\begin{aligned}
w_i(p) &= \frac{\partial \mu(\boldsymbol{q})}{\partial q_i}\bigg|_{\boldsymbol{q}=(p,\cdots,p)} \\
&= \frac{\partial(\mu(\boldsymbol{q}) - \mu(x_i'; \boldsymbol{q}))}{\partial q_i}\bigg|_{\boldsymbol{q}=(p,\cdots,p)} \\
&= \frac{\partial}{\partial q_i} \int_{x_i} (\mu(x_i; \boldsymbol{q}) - \mu(x_i'; \boldsymbol{q})) \phi_{q_i,k_i}(x_i) dx_i \bigg|_{\boldsymbol{q}=(p,\cdots,p)} \\
&= \int_{x_i} (\mu(x_i; p) - \mu(x_i'; p)) \left(\frac{\partial \phi_{q_i,k_i}(x_i)}{\partial q_i}\bigg|_{\boldsymbol{q}=(p,\cdots,p)} \right) dx_i \\
&= \int_{x_i} (\mu(x_i; p) - \mu(x_i'; p)) \left(\frac{\partial \log \phi_{q_i,k_i}(x_i)}{\partial q_i}\bigg|_{\boldsymbol{q}=(p,\cdots,p)} \right) \phi_{p,k_i}(x_i) dx_i \\
&\leq \sqrt{\left(\int_{x_i} (\mu(x_i; p) - \mu(x_i'; p))^2 \phi_{p,k_i}(x_i) dx_i \right) \left(\int_{x_i} \left(\frac{\partial \log \phi_{q_i,k_i}(x)}{\partial q_i}\bigg|_{\boldsymbol{q}=(p,\cdots,p)} \right)^2 \phi_{p,k_i}(x) dx_i \right)} \\
&\leq \alpha \cdot \sqrt{\int_{x_i} \left(\frac{\partial \log \phi_{p_i,k_i}(x_i)}{\partial p_i}\bigg|_{\boldsymbol{p}=(p,\cdots,p)} \right)^2 \phi_{p_i,k_i}(x_i) dx_i} \\
&\leq \alpha \cdot \sqrt{\frac{1}{c \cdot \text{Var}(\mathcal{D}_p(k_i))}}
\end{aligned}
$$

The first equality is by definition. The second equality follows from the fact that $\mu(x_i'; \mathbf{q})$ is constant with respect to q_i, so its derivative is 0. The third inequality simply expands out the definition of $\mu(\mathbf{q})$. The fourth equality follows from the linearity of derivatives, the fact that $\mu(x_i; \mathbf{q}) - \mu(x_i', \mathbf{q})$ is constant with respect to q_i, and the fact that $(\mu(x_i; \mathbf{q}) - \mu(x_i', \mathbf{q}))|_{\mathbf{q}=(p,\cdots,p)} = (\mu(x_i; p) - \mu(x_i', p))$. The fifth equality follows from the formula $\frac{\partial}{\partial x} \ln f(x) = \frac{\frac{\partial}{\partial x} f(x)}{f(x)}$, which holds for any differentiable function f. The first inequality is a result of the Cauchy-Schwarz inequality. The second inequality follows from the definition of α, and the final inequality follows from Assumption 2 of Lemma 5.3.

Therefore,

$$\alpha \geq w_i(p) \cdot \sqrt{c \cdot \text{Var}(\mathcal{D}_p(k_i))}.$$

We now argue that any $(\epsilon, \epsilon^2/100)$-differentially private mechanism should have variance $\Omega(\alpha^2 \log \frac{1}{\epsilon}/10\epsilon^2)$. Suppose that we had a mechanism that violated this property. Then by running this mechanism $\frac{1}{\epsilon^2 \log \frac{1}{\epsilon}}$ times and averaging, the advanced composition theorem implies that this average is $(1, 1/100)$-DP. This averaged output however has variance $O(\alpha^2/10)$. Thus given samples

x_i, and x_i' such that $|\mu(x_i;p) - \mu(x_i';p)| \geq \alpha/2$, if the noise had variance $O(\alpha^2/10)$ on (x_i, x_{-i}) as well as on (x_i', x_{-i}) (when x_{-i} is drawn randomly), then these two inputs would be distinguishable with probability at least $9/10$. This however violates the $(1, 1/100)$-DP of the averaged algorithm. This implies that for random x_i, the noise added by the DP algorithm is at least $\Omega(\alpha^2 \log \frac{1}{\epsilon}/20\epsilon^2)$

Thus the variance of M is,

$$\text{Var}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i), M}(M) \geq \sum_{i=1}^n c \cdot w_i(p)^2 \text{Var}(\mathcal{D}_p(k_i)) + \Omega\left(\frac{w_i(p)^2 \text{Var}(\mathcal{D}_p(k_i))}{\epsilon^2}\right) \qquad (23)$$

Finally, since the weights $w_i(p)$ that we defined need not sum to 1, they will need to be normalized to sum to 1 to satisfy the conditions of NLE. We need to show this normalisation does not substantially increase the variance of the estimator in NLE defined by these weights. This is equivalent to showing that the normalisation term, $\sum_{i=1}^n w_i(p)$ is large for some p. For $p \in [1/3, 2/3]$, let $\gamma : [1/3, 2/3] \to [0,1]^n$, defined by $\gamma(p) = (p, \cdots, p)$, be a path in $[0,1]^n$ then by the fundamental theorem of line integrals,

$$3\int_{1/3}^{2/3} \sum_{i=1}^n w_i(p)dp = 3\int_{1/3}^{2/3} \left(\sum_{i=1}^n \left(\frac{\partial}{\partial q_i}\mu(\boldsymbol{q})\right)\bigg|_{\boldsymbol{q}=(p,\cdots,p)}\right)dp$$

$$= 3\int_\gamma \nabla\mu(\boldsymbol{q}) \cdot \mathbf{1}d\boldsymbol{q}$$

$$= 3(\mu(2/3, \cdots, 2/3) - \mu(1/3, \cdots, 1/3))$$

$$= 1$$

This implies that there exists $p^* \in [1/3, 2/3]$ such that $\sum_{i=1}^n w_i(p^*) \geq 1$. Define

$$M_{\text{NL}}(x_1, \cdots, x_n) = \sum_{i=1}^n \frac{w_i(p^*)}{\sum_{j=1}^n w_j(p^*)x_j} + \text{Lap}\left(\frac{\max_i \frac{w_i(p^*)}{\sum_{j=1}^n w_j(p^*)}\sqrt{\text{Var}(\mathcal{D}_p(k_i))}}{\epsilon}\right)$$

$$= \frac{1}{\sum_{i=1}^n w_i(p^*)}\left(\sum_{i=1}^n w_i(p^*)x_i + \text{Lap}\left(\frac{\max_i w_i(p^*)\sqrt{\text{Var}(\mathcal{D}_p(k_i))}}{\epsilon}\right)\right),$$

where the second equality follows from properties of the Laplace distribution. Now,

$$\text{Var}_{\mathcal{D}_p}(M_{\text{NL}}) \leq \frac{1}{(\sum_{i=1}^n w_i(p^*))^2}\left(\sum_{i=1}^n w_i(p^*)^2\text{Var}(\mathcal{D}_p(k_i)) + O\left(\frac{\max_i w_i(p^*)^2\text{Var}(\mathcal{D}_p(k_i))}{\epsilon^2}\right)\right)$$

$$\leq \sum_{i=1}^n w_i(p^*)^2\text{Var}(\mathcal{D}_p(k_i)) + O\left(\frac{\max_i w_i(p^*)^2\text{Var}(\mathcal{D}_p(k_i))}{\epsilon^2}\right),$$

where the second inequality comes from the fact that $\sum_{i=1}^n w_i(p^*) \geq 1$. Comparing this with Equation 23, we see that specifically, at $p = p^*$,

$$\text{Var}_{\mathcal{D}_{p^*}}(M_{\text{NL}}) \leq O\left(\text{Var}_{\mathcal{D}_{p^*}}(M)\right).$$

Now, if $p, p^* \in [1/3, 2/3]$ then $\text{Var}(\mathcal{D}_p(k_i)) = \Theta(\text{Var}(\mathcal{D}_{p^*}(k_i)))$ so $\text{Var}_{D_p}(M_{\text{NL}}) = \Theta(\text{Var}_{D_{p^*}}(M_{\text{NL}}))$. Therefore, the worst case variance of M_{NL} is less than the worst case variance of M over all $p \in [1/3, 2/3]$, as required. $\qquad \square$

Lemma E.1. *For any randomised mechanism* $M : [0,1]^n \to [0,1]$,

$$\text{Var}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i), M}(M) \qquad (16)$$

$$= \mathbb{E}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i), M}[(M(x_1, ..., x_n) - \mu(p))^2]$$

$$\geq \sum_{i=1}^n \mathbb{E}_{x_i \sim \mathcal{D}_p(k_i)}[(\mu(x_i;p) - \mu(p))^2] + \mathbb{E}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i), M}[(M(x_1, ..., x_n) - \mu(x_1, ..., x_n; p))^2]$$

Proof of Lemma E.1. Let $M : [0,1]^n \to [0,1]$ be a randomised mechanism and suppose that each $x_i \sim \mathcal{D}(p_i, k_i)$ where $p_i \sim \mathcal{D}$. Now, our goal is to decompose the variance of M into

the variance conditioned on each coordinate, and the variance inherent in the mechanism itself. Let $\mu = \mathbb{E}_{x_1 \sim \mathcal{D}(k_1), \cdots, x_n \sim \mathcal{D}(k_n), M}[M(x_1, \cdots, x_n)]$ be the expectation and for any $I \subset [n]$, let $\mu(x_{[n] \setminus I}) = \mathbb{E}_{\forall i \in I, x_i \sim \mathcal{D}(k_i), M}[M(x)]$ be the expectation conditioned only on the randomness in I. So,

$$\mathrm{Var}(M) = \mathbb{E}_{x_1 \sim \mathcal{D}(k_1), \cdots, x_n \sim \mathcal{D}(k_n), M}[(M(x_1, \cdots, x_n) - \mu)^2]$$

$$= \mathbb{E}_{x_1 \sim \mathcal{D}(k_1)} \mathbb{E}_{x_2 \sim \mathcal{D}(k_2), \cdots, x_n \sim \mathcal{D}(k_n), M}[(M(x_1, \cdots, x_n) - \mu_1(x_1) + \mu_1(x_1) - \mu)^2]$$

$$= \mathbb{E}_{x_1 \sim \mathcal{D}(k_1)} \mathbb{E}_{x_2 \sim \mathcal{D}(k_2), \cdots, x_n \sim \mathcal{D}(k_n), M}[(M(x_1, \cdots, x_n) - \mu_1(x_1))^2$$

$$+ 2(M(x_1, \cdots, x_n) - \mu_1(x_1))(\mu_1(x_1) - \mu) + (\mu_1(x_1) - \mu)^2]$$

$$= \mathbb{E}_{x_1 \sim \mathcal{D}(k_1)}[(\mu_1(x_1) - \mu)^2] + \mathbb{E}_{x_1 \sim \mathcal{D}(k_1)} \mathbb{E}_{x_2 \sim \mathcal{D}(k_2), \cdots, x_n \sim \mathcal{D}(k_n), M}[(M(x_1, \cdots, x_n) - \mu_1(x_1))^2].$$

Now, by induction we obtain the following decomposition of the variance of M,

$$\mathrm{Var}(M) = \sum_{i=1}^{n} \mathbb{E}_{x_1 \sim \mathcal{D}(k_1), \cdots, x_i \sim \mathcal{D}(k_i)}[(\mu(x_{j \leq i}) - \mu(x_{j < i}))^2]$$

$$+ \mathbb{E}_{x_1 \sim \mathcal{D}(k_1), \cdots, x_n \sim \mathcal{D}(k_n), M}[(M(x_1, \cdots, x_n) - \mu(x_1, \cdots, x_n))^2]$$

$$\geq \sum_{i=1}^{n} \mathbb{E}_{x_i \sim \mathcal{D}(k_i)}[(\mu(x_i) - \mu)^2] + \mathbb{E}_{x_1 \sim \mathcal{D}(k_1), \cdots, x_n \sim \mathcal{D}(k_n), M}[(M(x_1, \cdots, x_n) - \mu(x_1, \cdots, x_n))^2]$$

where the second inequality follows from Jensen's inequality:

$$\mathbb{E}_{x_1 \sim \mathcal{D}(k_1), \cdots, x_i \sim \mathcal{D}(k_i)}[(\mu(x_{j \leq i}) - \mu(x_{j < i}))^2] \geq \mathbb{E}_{x_i \sim \mathcal{D}(k_i)}[(\mathbb{E}_{x_1 \sim \mathcal{D}(k_1), \cdots, x_{i-1} \sim \mathcal{D}(k_i)}[\mu(x_{j \leq i}) - \mu(x_{j < i})])^2]$$

$$= \mathbb{E}_{x_i \sim \mathcal{D}(k_i)}[(\mu(x_i) - \mu)^2].$$

$\square$

Lemma 5.4. *For any distribution $\mathcal{D}$, $n > 0$ and $\beta \in [0,1]$, if for all k_i, $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) = \tilde{O}(\mathrm{Var}(\mathcal{D}(k_i)))$ then for any $\mathbf{w} \in [0,1]^n$ such that $\sum_{i=1}^{n} w_i = 1$, we have $\mathrm{Var}(M_{\mathit{TNL}}(\cdot\;;\mathbf{w})) = \tilde{O}(\mathrm{Var}(M_{\mathit{NL}}(\cdot\;;\mathbf{w})))$. Further, the bias of M_{TNL} is at most β.*

Proof of Lemma 5.4. The variance claim follows immediately from noting that $\mathrm{Var}\left([x_i]_{p - f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)}^{p + f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)}\right) \leq \mathrm{Var}(x_i)$, and the assumption that $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) = \tilde{O}(\mathrm{Var}(\mathcal{D}(k_i)))$. The bias claim follows from noting that with probability $1 - \beta$, $[x_i]_{p - f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)}^{p + f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)} = x_i$. This implies that M_{TNL} is within β in total variation distance to an unbiased estimator. Since M_{TNL} takes values in $[0,1]$, this implies the mean is in $[p - \beta, p + \beta]$. $\square$

Corollary 5.5. *Given $k_1, \cdots, k_n \in \mathbb{N}$, and σ_p, there exists a family of distributions $\mathcal{D}_p$ such that*

$$\min_{M, \text{ unbiased}} \max_{p \in [1/3, 2/3]} \mathrm{Var}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i)}[M(x_1, \cdots, x_n)] \geq \tilde{\Omega}\left(\min\left\{\frac{\frac{k^*}{\epsilon^2} + \sum_{i=1}^{n} \min\{k_i, k^*\}}{(\sum_{i=1}^{n} \min\{k_i, \sqrt{k_i k^*}\})^2}, \frac{\sigma_p^2}{n}\right\}\right).$$

Proof of Corollary 5.5. Firstly, suppose that $\sigma_p = 0$, so the meta-distribution is constant, and $\mathcal{D}_p(k_i) = \mathrm{Bin}(k_i, p)$. Then the Fisher information of ϕ_{p, k_i} is $\int \left(\frac{\partial}{\partial p} \log \phi_{p, k_i}(x_i)\right)^2 \phi_{p, k_i}(x_i) dx_i = \frac{k_i}{p(1-p)}$ and $\mathrm{Var}(\mathcal{D}_p(k_i)) = \frac{p(1-p)}{k_i}$, so $\mathcal{D}_p(k_i)$ satisfies Condition 8 of Theorem 5.1. Additionally,

$$\min_{M, \text{ unbiased}} \max_{p \in [1/3, 2/3]} \mathrm{Var}_{\mathcal{D}_p}[M] = \tilde{\Omega}\left(\max_{p \in [1/3, 2/3]} \mathrm{Var}_{\mathcal{D}_p}[\widehat{p}_{\epsilon}^{\text{ideal}}]\right) \quad \text{(under conditions of Thm 4.1)}$$

We can view the truncation as simply choosing a maximum k^* so that $T = \sqrt{\frac{k^*}{p(1-p)}}$. Now, the un-normalised weights of $\widehat{p}_{\epsilon}^{\text{ideal}}$ are

$$\min\left\{\frac{1}{\mathrm{Var}(\mathcal{D}_p(k_i))}, \frac{T}{\sqrt{\mathrm{Var}(\mathcal{D}_p(k_i))}}\right\} = \min\left\{\frac{k_i}{p(1-p)}, \frac{\sqrt{k_i k^*}}{p(1-p)}\right\}.$$

Further, $\mathrm{Var}([\widehat{p}_i]_{a_i}^{b_i}) \leq \mathrm{Var}(\mathcal{D}(k_i))$ and we assume thoughout this paper that $\mathrm{Var}([\widehat{p}_i]_{a_i}^{b_i}) \geq [(1/2)\mathrm{Var}(\mathcal{D}(k_i))]$. So, $\mathrm{Var}([\widehat{p}_i]_{a_i}^{b_i}) = \Theta(\mathrm{Var}(\mathcal{D}(k_i))) = \Theta(\frac{p(1-p)}{k_i})$. Finally, since binomials are highly concentrated, $|b_i - a_i| = \Omega(\sigma_i)$, which implies that $\frac{\max_i w_i^* |b_i - a_i|}{\epsilon}$ as defined in eqn (3) is achieved at $k_i = k^*$. Thus,

$$
\min_{M,\text{ unbiased}} \max_{p \in [1/3, 2/3]} \mathrm{Var}_{\mathcal{D}_p}[M] = \max_{p \in [1/3, 2/3]} \frac{\Omega\left(\frac{k^*}{p(1-p)\epsilon^2}\right) + \sum_{i=1}^n \left(\min\left\{\frac{k_i}{p(1-p)}, \frac{\sqrt{k_i k^*}}{p(1-p)}\right\}\right)^2 \frac{1}{2}\frac{p(1-p)}{k_i}}{\left(\sum_{i=1}^n \min\left\{\frac{k_i}{p(1-p)}, \frac{\sqrt{k_i k^*}}{p(1-p)}\right\}\right)^2}
$$

$$
= \tilde{\Omega}\left(\max_{p \in [1/3, 2/3]} p(1-p)\frac{\frac{k^*}{\epsilon^2} + \sum_{i=1}^n \min\{k_i, k^*\}}{(\sum_{i=1}^n \min\{k_i, \sqrt{k_i k^*}\})^2}\right)
$$

$$
= \tilde{\Omega}\left(\frac{\frac{k^*}{\epsilon^2} + \sum_{i=1}^n \min\{k_i, k^*\}}{(\sum_{i=1}^n \min\{k_i, \sqrt{k_i k^*}\})^2}\right),
$$

where the first equality comes from Theorem 5.1, the second equality pulls out common factors, and the third equality is because p is bounded away from 0 and 1.

For the other component of the bound we will let $\mathcal{D}_p$ be a truncated Gaussian distribution. Let ϕ and Φ respectively be the probability density function and cumulative density function of the standard Gaussian $\mathcal{N}(0, 1)$. Let W be such that $\gamma := \Phi(W) - \Phi(-W) \geq 9/10$ and $\lambda := \frac{2W\phi(W)}{\Phi(W)-\Phi(-W)} \leq 1/2$. Define the truncated Gaussian $\mathcal{D}_p$ with mean p on $[p - \frac{\sigma_p}{\sqrt{1-\lambda}}W, p + \frac{\sigma_p}{\sqrt{1-\lambda}}W]$ by the probability density function:

$$
\phi_p(q) = \begin{cases} \frac{1}{\gamma}\phi\left((q-p)\frac{\sqrt{1-\lambda}}{\sigma_p}\right) & q \in [p - \frac{\sigma_p}{\sqrt{1-\lambda}}W, p + \frac{\sigma_p}{\sqrt{1-\lambda}}W] \\ 0 & \text{otherwise.} \end{cases}
$$

Now, the variance of $\mathcal{D}_p$ is σ_p^2 and the Fisher information of $\mathcal{D}_p$ is given by [29]

$$
\frac{1}{\sigma_p^2}(1-\lambda)^2 \in \left[\frac{1}{4\sigma_p^2}, \frac{1}{\sigma_p^2}\right]. \tag{24}
$$

Since any sample from $\mathcal{D}$ can be post-processed into a sampling from $\mathcal{D}(k)$ for any $k \in \mathbb{N}$, we have

$$
\min_{M,\text{ unbiased}} \max_{p \in [1/3, 2/3]} \mathrm{Var}_{\forall i \in [n], x_i \sim \mathcal{D}_p(k_i)}[M(x_1, \cdots, x_n)] \geq \min_{M,\text{ unbiased}} \max_{p \in [1/3, 2/3]} \mathrm{Var}_{p_1, \cdots, p_n \sim \mathcal{D}_p}[M(p_1, \cdots, p_n)]
$$

$$
\geq \max_{p \in [1/3, 2/3]} O\left(\frac{\sigma_p^2}{n}\right)
$$

$$
= O(\frac{\sigma_p^2}{n}),
$$

where the second inequality follows from the Cramer-Rao bound [30] and Equation (24). $\qquad\square$

F Example Initial Estimators

In this section we give example initial mean and variance estimation procedures that can be used in the framework described in Section 4. For both estimators, we show that they satisfy the conditions of Theorem 4.1, and thus can be used as initial estimators in Algorithm 2, assuming all other technical conditions are satisfied. This also immediately implies that the set of initial mean and variance estimators which satisfy the conditions of Theorem 4.1 is non-empty.

We note again that the estimators described in this section are examples of estimators that achieve the conditions of Theorem 4.1, and that any private mean and variance estimators that satisfy these conditions could be used instead. As discussed in Section 4.2, one may choose to use different estimators of these initial quantities in different settings (for example, if local differential privacy is required or if different distributional assumptions are known).

F.1 Initial Mean Estimation

We will begin with the initial mean estimate $\widehat{p}_\epsilon^{\text{initial}}$. We consider the simplest mean estimation subroutine, where the analyst collects a single data point from the $n/10$ users with the smallest k_i, then privately computes the empirical mean of these points using the Laplace Mechanism. The following lemma shows that this process is differentially private and satisfies the accuracy conditions of Theorem 4.1, i.e., that with high probability, $\widehat{p}_\epsilon^{\text{initial}}$ is close to p and $\widehat{p}_\epsilon^{\text{initial}}(1 - \widehat{p}_\epsilon^{\text{initial}})$ is close to $p(1-p)$.

Lemma F.1. *Fix any $\epsilon > 0$ and let $\widehat{p}_\epsilon^{\text{initial}}(x^1_{(9n/10)+1}, \cdots, x^1_n) = \frac{1}{n/10} \sum_{i=(9n/10)+1}^n x_i +$ Lap $\left(\frac{10}{\epsilon n} \right)$. Then $\widehat{p}_\epsilon^{\text{initial}}$ is ϵ-differentially private, $\mathbb{E}[\widehat{p}_\epsilon^{\text{initial}}(x^1_{(9n/10)+1}, \cdots, x^1_n)] = p$ and if $p \geq \frac{20 \log(1/\beta)}{n}$, then for n sufficiently large,*

$$\Pr[|\widehat{p}_\epsilon^{\text{initial}} - p| \leq \alpha] \leq \beta \text{ for } \alpha = 2 \max \left\{ \sqrt{\frac{12 \widehat{p}_\epsilon^{\text{initial}} \log(4/\beta)}{n/10} + \frac{36 \log^2(4/\beta)}{n^2/100}} + \frac{6 \log(4/\beta)}{n/10}, \frac{\log(2/\beta)}{\epsilon n/10} \right\} \leq f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta).$$

Further, if $\min\{p, 1-p\} \geq 12 \max \left\{ \frac{3 \log(4/\beta)}{n/10}, \frac{\log(2/\beta)}{\epsilon n/10} \right\}$ then with probability $1 - \beta$, $\widehat{p}_\epsilon^{\text{initial}} \in [\frac{1}{2}p, \frac{3}{2}p]$ and $\widehat{p}_\epsilon^{\text{initial}}(1 - \widehat{p}_\epsilon^{\text{initial}}) \in [\frac{p(1-p)}{2}, \frac{3p(1-p)}{2}].$

The concentration bound follows from noticing that $\mathcal{D} = \text{Ber}(p)$ and using the concentration of binomial random variables.

Note that the expression of α depends only on quantities known to the analyst – including $\widehat{p}_\epsilon^{\text{initial}}$, which will be observed as output – so that α can be computed directly for use in Algorithm 2. Although our presentation of Algorithm 2 requires α to be specified up front as input to the algorithm, it could equivalently be computed internally by the algorithm as a function of $\widehat{p}_\epsilon^{\text{initial}}$ and other input parameters.

Proof. Firstly, the privacy guarantees follows immediately from the Laplace Mechanism in differential privacy [12] noting that $\frac{10}{n} \sum_{i=(9n/10)+1}^n x_i^1$ has sensitivity $\frac{10}{n}$.

Now, let us turn to the two accuracy guarantees. We will start with the guarantee that $\widehat{p}_\epsilon^{\text{initial}}$ is close to p with high-probability. Note that $\mathcal{D}$ is simply a Bernoulli random variable with mean p so since each sample is independent, $\frac{10}{n} \sum_{i=(9n/10)+1}^n x_i^1 = \text{Bin}(n/10, p)$. Thus, if $n \geq \frac{20 \log(1/\beta)}{p}$, a Chernoff bound gives

$$\Pr \left[\left| \frac{10}{n} \sum_{i=(9n/10)+1}^n x_i^1 - p \right| \geq \sqrt{\frac{3 \min\{p, 1-p\} \log(4/\beta)}{n/10}} \right] \leq \beta/2.$$

Therefore, combining with a high probability bound on the Laplace distribution,

$$\Pr \left[|\widehat{p}_\epsilon^{\text{initial}} - p| \geq \sqrt{\frac{3 \min\{p, 1-p\} \log(4/\beta)}{n/10}} + \frac{\log(2/\beta)}{\epsilon n/10} \right] \leq \beta.$$

We will condition on the following event for the remainder of the proof, which will occur with probability $1 - \beta$:

$$|\widehat{p}_\epsilon^{\text{initial}} - p| \leq 2 \max \left\{ \sqrt{\frac{3 \min\{p, 1-p\} \log(4/\beta)}{n/10}}, \frac{\log(2/\beta)}{\epsilon n/10} \right\}.$$

Now if $|\widehat{p}_\epsilon^{\text{initial}} - p| \leq 2 \sqrt{\frac{3 \min\{p, 1-p\} \log(4/\beta)}{n/10}}$. Since we need α in terms of $\widehat{p}_\epsilon^{\text{initial}}$ rather than p (since $\widehat{p}_\epsilon^{\text{initial}}$ is known to the algorithm), we need to rework this formula. Squaring both sides and bringing all the terms to the same side, we obtain

$$p^2 - 2 \left(\widehat{p}_\epsilon^{\text{initial}} + \frac{6 \log(4/\beta)}{n/10} \right) p + (\widehat{p}_\epsilon^{\text{initial}})^2 \leq 0.$$

Completing the square we obtain

$$\left(p - \widehat{p}_\epsilon^{\text{initial}} - \frac{6 \log(4/\beta)}{n/10} \right)^2 + (\widehat{p}_\epsilon^{\text{initial}})^2 - \left(\widehat{p}_\epsilon^{\text{initial}} + \frac{6 \log(4/\beta)}{n/10} \right)^2 \leq 0.$$

Now, rearranging and taking the square root, we obtain

$$\left| p - \widehat{p}_\epsilon^{\text{initial}} - \frac{6\log(4/\beta)}{n/10} \right| \leq \sqrt{\left(\widehat{p}_\epsilon^{\text{initial}} + \frac{6\log(4/\beta)}{n/10} \right)^2 - (\widehat{p}_\epsilon^{\text{initial}})^2}$$

then by squaring both sides, using the fact that $\min\{p, 1-p\} \leq p$, and rearranging we have

$$|\widehat{p}_\epsilon^{\text{initial}} - p| \leq \sqrt{\frac{12\widehat{p}_\epsilon^{\text{initial}}\log(4/\beta)}{n/10} + \frac{36\log^2(4/\beta)}{n^2/100} + \frac{6\log(4/\beta)}{n/10}}$$

which implies that,

$$|\widehat{p}_\epsilon^{\text{initial}} - p| \leq 2\max\left\{ \sqrt{\frac{12\widehat{p}_\epsilon^{\text{initial}}\log(4/\beta)}{n/10} + \frac{36\log^2(4/\beta)}{n^2/100} + \frac{6\log(4/\beta)}{n/10}}, \frac{\log(2/\beta)}{\epsilon n/10} \right\}.$$

We need to show that this expression is less than or equal to $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$ because $\alpha = O(1/\sqrt{n})$. To see this, note that $\alpha = O(1/\sqrt{n})$ and $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$ is increasing towards 1 as n grows large. Thus for n sufficiently large, $\alpha \leq f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$ will be satisfied.

Next we turn to proving the second accuracy claim, that $\widehat{p}_\epsilon^{\text{initial}}(1 - \widehat{p}_\epsilon^{\text{initial}})$ is concentrated around $p(1-p)$. Let $\mathcal{E} = \widehat{p}_\epsilon^{\text{initial}} - p$ so

$$\widehat{p}_\epsilon^{\text{initial}}(1 - \widehat{p}_\epsilon^{\text{initial}}) = (p + \mathcal{E})(1 - p - \mathcal{E}) = p(1-p) + (1 - 2p)\mathcal{E} - \mathcal{E}^2$$

Now, if $\min\{p, 1-p\} \geq K\max\left\{ \frac{3\log(4/\beta)}{n/10}, \frac{\log(2/\beta)}{\epsilon n/10} \right\}$ for some constant K, then

$$|\mathcal{E}| \leq \sqrt{\frac{3\min\{p, 1-p\}\log(4/\beta)}{n/10}} + \frac{\log(2/\beta)}{\epsilon n/10}$$

$$\leq \sqrt{\frac{\min\{p, 1-p\}\min\{p, (1-p)\}}{K}} + \frac{\min\{p, (1-p)\}}{K}$$

$$\leq \frac{2\min\{p, 1-p\}}{K}.$$

Thus, combining this with the fact that $1 - 2p \leq \max\{p, 1-p\}$ for $p \in [0, 1]$,

$$|(1 - 2p)\mathcal{E} - \mathcal{E}^2| \leq \max\{p, 1-p\}\frac{2\min\{p, 1-p\}}{K} + \left(\frac{2\min\{p, 1-p\}}{K} \right)^2$$

$$\leq \frac{6p(1-p)}{K}$$

Finally, choosing $K = 12$ gives,

$$\widehat{p}_\epsilon^{\text{initial}}(1 - \widehat{p}_\epsilon^{\text{initial}}) \in \left[\frac{p(1-p)}{2}, \frac{3p(1-p)}{2} \right].$$

$\square$

F.2 Initial Variance Estimation

We now turn to estimating σ_p^2. Let us first provide some background on privately estimating the standard deviation of well-behaved distributions. Lemma F.2 guarantees the existence of a differentially private algorithm for estimating standard deviation within a small constant factor with high probability, as long as the sample size is sufficiently large. The following is a slight generalisation of the estimation of the standard deviation of a Gaussian given in [21] due to [26].

Lemma F.2 (DP standard deviation estimation). *For all $n \in \mathbb{N}$, $\sigma_{min} < \sigma_{max} \in [0, \infty], \epsilon > 0, \delta \in (0, \frac{1}{n}], \beta \in (0, 1/2), \zeta > 0$, there exists an (ϵ, δ)-differentially private algorithm $\mathcal{M}$ that satisfies: if $x_1, \ldots, x_n$ are i.i.d. draws from a distribution P which has standard deviation $\sigma \in [\sigma_{min}, \sigma_{max}]$ and absolute central third moment $\rho = \mathbb{E}[|x - \mu(P)|^3]$ such that $\frac{\rho}{\sigma^3} \leq \zeta$, then if $n \geq c\zeta^2\min\{\frac{1}{\epsilon}\ln(\frac{\ln\frac{\sigma_{max}}{\sigma_{min}}}{\beta}), \frac{1}{\epsilon}\ln(\frac{1}{\delta\beta})\}$, (where c is a universal constant), then $\mathcal{M}$ produces an estimate $\widehat{\sigma}$ of the standard deviation such that $\Pr_{x_1, \ldots, x_n \sim P, \mathcal{M}}(\sigma^2 \leq \widehat{\sigma}^2 \leq 8\sigma^2) \geq 1 - \beta$.*

In order to estimate σ_p^2, we will use the estimator promised by Lemma F.2 on the data of the $\log n$ users with the largest k_i. Let $k = k_{\log n}$, so the top $\log n$ individuals all have at least k data points. We will have these individuals report $\widehat{p}_i^k := \frac{1}{k}\sum_{j=1}^k x_j^i$, which is the empirical mean of their first k data points. Thus, we are running the estimator promised in Lemma F.2 on $\mathcal{D}(k)$ with $\log n$ data points. In order to utilise Lemma F.2, we first need to ensure that $\mathcal{D}(k)$ satisfies the moment condition that ρ/σ^3 is bounded, which is shown in Lemma F.3.

Lemma F.3. *For $k \in \mathbb{N}$, suppose $p \in [\frac{1}{k}, 1 - \frac{1}{k}]$, $\sigma_p \geq \frac{1}{k}$, $k \geq 2$, and there exists $\gamma > 0$ such that $\frac{\rho_\mathcal{D}}{\sigma_p^3} \leq \gamma$ where $\rho_\mathcal{D}$ denotes the absolute central third moment of $\mathcal{D}$. Then $\frac{\rho_{\mathcal{D}(k)}}{\mathrm{Var}(\mathcal{D}(k))^{3/2}} \leq 8(3\sqrt{3} + \gamma)$.*

Proof. Note that $\mathbb{E}[\mathcal{D}(k)] = p$. Then we can bound the absolute third central moment as follows,

$$\mathbb{E}_{x \sim \mathcal{D}(k)}[|x - p|^3] = \mathbb{E}_{p_i \sim \mathcal{D}}\mathbb{E}_{y \sim \mathrm{Bin}(k, p_i)}[|(\frac{1}{k}y - p_i) - (p - p_i)|^3]$$

$$\leq 4\left(\mathbb{E}_{p_i \sim \mathcal{D}}\mathbb{E}_{y \sim \mathrm{Bin}(k, p_i)}[|\frac{1}{k}y - p_i|^3] + \mathbb{E}_{p_i \sim \mathcal{D}}[|p - p_i|^3]\right)$$

$$\leq 4\left(\frac{1}{k^3}\mathbb{E}_{p_i \sim \mathcal{D}}\left[\sqrt{\mathbb{E}_{y \sim \mathrm{Bin}(k, p_i)}[|y - k \cdot p_i|^2]\mathbb{E}_{y \sim \mathrm{Bin}(k, p_i)}[|y - p_i|^4]}\right] + \gamma\sigma_p^3\right)$$
(by Cauchy-Schwarz inequality)

$$\leq 4\left(\frac{1}{k^3}\mathbb{E}_{p_i \sim \mathcal{D}}\left[\sqrt{k^2(p_i(1 - p_i))^2(1 + 3kp_i(1 - p_i))}\right] + \gamma\sigma_p^3\right)$$

$$\leq 4\left(\frac{1}{k^3}\mathbb{E}_{p_i \sim \mathcal{D}}[k(p_i(1 - p_i))] + \frac{1}{k^3}\mathbb{E}_{p_i \sim \mathcal{D}}[\sqrt{3k^3(p_i(1 - p_i))^3}] + \gamma\sigma_p^3\right)$$

$$\leq 4\left(\frac{1}{k^2}p(1 - p) + \frac{\sqrt{3}}{k^{3/2}}\mathbb{E}_{p_i \sim \mathcal{D}}[\sqrt{(p_i(1 - p_i))^3}] + \gamma\sigma_p^3\right)$$
(by Jensen's inequality)

$$\leq 4\left(\frac{1}{k^{3/2}}\sqrt{(p(1 - p))^3} + \frac{\sqrt{3}}{k^{3/2}}\mathbb{E}_{p_i \sim \mathcal{D}}[\sqrt{(p_i(1 - p_i))^3}] + \gamma\sigma_p^3\right),$$

where the first inequality follows from the following inequality that holds for all real valued a and b: $|a - b|^3 \leq 4(|a|^3 + |b|^3)$. The second to last inequality follows from Jensen's inequality since $h(x) = x(1 - x)$ is concave, and the last inequality follows since $\frac{1}{\sqrt{k}} \leq \sqrt{p(1 - p)}$. Now, we will use a generalised form of Jensen's inequality to bound $\mathbb{E}_{p_i \sim \mathcal{D}}[\sqrt{(p_i(1 - p_i))^3}]$. Let $h(x) = (x(1 - x))^{3/2}$ and

$$\phi(x) = \frac{h(x) - h(p)}{(x - p)^2} - \frac{h'(p)}{x - p}.$$

Since $p \in [\frac{1}{k}, 1 - \frac{1}{k}]$,

$$\max_{x \in [\frac{1}{2k}, 1 - \frac{1}{2k}]} \phi(x) \leq (1/2)\max_{x \in [\frac{1}{2k}, 1 - \frac{1}{2k}]} h''(x) \leq h''\left(\frac{1}{2k}\right) = \frac{3(8(\frac{1}{2k})^2 - 8(\frac{1}{2k}) + 1)}{4\sqrt{(1 - \frac{1}{2k})\frac{1}{2k}}} = \frac{3(8 - 16k + 4k^2)}{8k\sqrt{(2k - 1)}} \leq \frac{3}{2}\sqrt{k}.$$

If $x \notin [\frac{1}{2k}, 1 - \frac{1}{2k}]$ then $|x - p| \geq \frac{1}{2k}$ and $h(x) < h(p)$, so

$$\phi(x) \leq \frac{|h'(p)|}{|x - p|} = \frac{3|1 - 2p|\sqrt{p(1 - p)}}{2|p - x|} \leq \frac{3}{2}\frac{\sqrt{p(1 - p)}}{|p - x|} \leq \max\left\{\frac{3}{2}\frac{\sqrt{\frac{1}{k}(1 - \frac{1}{k})}}{|\frac{1}{k} - x|}, \frac{3}{2}\frac{\sqrt{\frac{1}{k}(1 - \frac{1}{k})}}{|1 - \frac{1}{k} - x|}\right\} \leq 3\sqrt{k - 1} \leq 3\sqrt{k}.$$

Therefore, by the generalised Jensen's inequality,

$$\mathbb{E}_{p_i \sim \mathcal{D}}[\sqrt{(p_i(1 - p_i))^3}] \leq \sqrt{(p(1 - p))^3} + \sigma_p^2 \cdot 3\sqrt{k} \leq \sqrt{(p(1 - p))^3} + \sigma_p^2 \cdot 3\sqrt{k}.$$

Continuing to bound the absolute central third moment as above,

$$\mathbb{E}_{x \sim \mathcal{D}(k)}[|x - p|^3] \leq 4 \left(\frac{1}{k^{3/2}} \sqrt{(p(1-p))^3} + \frac{\sqrt{3}}{k^{3/2}} \mathbb{E}_{p_i \sim \mathcal{D}}[\sqrt{(p_i(1-p_i))^3}] + \gamma \sigma_p^3 \right)$$

$$\leq 4 \left(\frac{1}{k^{3/2}} \sqrt{(p(1-p))^3} + \frac{\sqrt{3}}{k^{3/2}} \sqrt{(p(1-p))^3} + 3\sqrt{3} \frac{\sigma_p^2}{k} + \gamma \sigma_p^3 \right)$$

$$\leq 4 \left(\frac{1}{k^{3/2}} \sqrt{(p(1-p))^3} + \frac{\sqrt{3}}{k^{3/2}} \sqrt{(p(1-p))^3} + 3\sqrt{3} \sigma_p^3 + \gamma \sigma_p^3 \right)$$

$$\leq 4(3\sqrt{3} + \gamma) \left(\frac{1}{k^{3/2}} \sqrt{(p(1-p))^3} + \sigma_p^3 \right)$$

$$\leq 4(3\sqrt{3} + \gamma) \left(\frac{1}{k} p(1-p) + \sigma_p^2 \right)^{3/2}$$

$$\leq 8(3\sqrt{3} + \gamma) \left(\frac{1}{k} p(1-p) + \frac{k-1}{k} \sigma_p^2 \right)^{3/2},$$

where the first and second inequalities follow from above, the third inequality follows because $k \geq 1$, the fourth is simply rearranging the terms, the fifth follows from the fact that for all positive, real numbers a and b: $a^{3/2} + b^{3/2} < (a + b)^{3/2}$, and the last inequality follows since if $k \geq 2$ then $(k-1)/k > 1/2$. $\qquad \square$

With this result, we can apply Lemma F.2 to our setting to privately achieve an estimate $\widehat{\sigma}_{p,k}^2$ that is close to the true population-level variance σ_p^2, as shown in Lemma F.4. Note that as k grows large, the allowable range for p approaches the full support $[0, 1]$ and the allowable standard deviation σ_p approaches any non-negative number.

Lemma F.4 combines these two results to show that Lemma F.2 can be applied to the individual reports $\widehat{p}_i^k$ from the top $\log n$ users, and the resulting variance estimate will satisfy the accuracy conditions of Theorem 4.1.

Lemma F.4. *Given $\sigma_{min} < \sigma_{max} \in [0, \infty], \epsilon > 0, \delta \in (0, \frac{1}{n}], \beta \in (0, 1/2)$, and $\zeta > 0$, let $\mathcal{M}$ be the (ϵ, δ)-differentially private mechanism given by Lemma F.2, and let $\widehat{\sigma}_{p,k}^2 = \mathcal{M}(\widehat{p}_1^k, \cdots, \widehat{p}_{\log n}^k)$, where $\widehat{p}_1^k, \cdots, \widehat{p}_{\log n}^k \sim \mathcal{D}(k)$. If there exists $\zeta > 0$ such that $\frac{\rho_{\mathcal{D}}}{\sigma_p^3} \leq \zeta$ where $\rho_{\mathcal{D}} = \mathbb{E}_{x \sim \mathcal{D}}[|x - p|^3], \sqrt{\frac{1}{k}p(1-p) + \frac{k-1}{k}\sigma_p^2} \in [\sigma_{min}, \sigma_{max}], \sigma_p > \frac{1}{k}, p \in \left[\frac{1}{k}, 1 - \frac{1}{k}\right]$, and $\log n \geq c(8(3\sqrt{3} + \zeta))^2 \min\{\frac{1}{\epsilon} \ln(\frac{\ln(\frac{\sigma_{max}}{\sigma_{min}})}{\beta}), \frac{1}{\epsilon} \ln(\frac{1}{\delta \beta})\}$, then with probability $1 - \beta$, $\widehat{\sigma}_{p,k}^2 \in [\mathrm{Var}(\mathcal{D}(k)), 8\mathrm{Var}(\mathcal{D}(k))]$.*

Proof. Note that the conditions are sufficient to ensure from Lemma F.3 that $\frac{\rho_{\mathcal{D}(k)}}{\mathrm{Var}(\mathcal{D}(k))^{3/2}} \leq 8(3\sqrt{3} + \gamma)$. Then Lemma F.2 and Lemma 2.1 imply that

$$\mathrm{Var}(\mathcal{D}(k)) = \frac{1}{k}p(1-p) + \frac{k-1}{k}\sigma_p^2 \leq \widehat{\sigma}_{p,k}^2 \leq 8 \left(\frac{1}{k}p(1-p) + \frac{k-1}{k}\sigma_p^2 \right) = 8\mathrm{Var}(\mathcal{D}(k)).$$

$\qquad \square$

G   Interpretation and Estimation of Concentration Functions

Recall that $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$ describes the concentration of $\widehat{p}_i \sim \mathcal{D}(k_i)$ and is defined as $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) = \arg\inf\{\alpha \mid \Pr_{\widehat{p}_1, \cdots, \widehat{p}_n \sim \mathcal{D}(k_i)}(\max_i |\widehat{p}_i - p| \geq \alpha) \leq \beta\}$. In the main body of the paper, we assumed that this function was known to the analyst, even if the input value σ_p^2 was unknown and had to be estimated. In this appendix, we interpret the structure of this concentration function and show that even when this informational assumption is relaxed, our Algorithm 2 can still be implemented with some minor modifications.

We start by introducing two additional functions: $f_{\mathcal{D}}(n, \sigma_p^2, \beta)$, which describes the concentration of $p_i \sim \mathcal{D}$, and $f_{\text{Bin}}(k_i, p_i, \beta)$, which describes the high probability tail bound on the binomial $\text{Bin}(k_i, p_i)$:

$$f_{\mathcal{D}}(n, \sigma_p^2, \beta) = \arg\inf\{\alpha \mid \Pr_{p_1, \cdots, p_n \sim \mathcal{D}}(\max_i |p - p_i| \geq \alpha) \leq \beta\}.$$

$$f_{\text{Bin}}(k_i, p_i, \beta) = \arg\inf\{\alpha \mid \Pr_{x \sim \text{Bin}(k_i, p_i)}(|\frac{1}{k_i}x - p_i| \geq \alpha) \leq \beta\}$$

In this appendix, we will assume that only the function $f_{\mathcal{D}}(n, \cdot, \beta)$ is known to the analyst, but the input variance parameter σ_p^2 of the distribution are not known. For example, the analyst may know that $\mathcal{D}$ is Gaussian with unknown mean and variance, and thus she can express the concentration of p_i as a function of the variance. Also note that for any values k_i, p_i and β, we can empirically compute $f_{\text{Bin}}(k_i, p_i, \beta)$.

The following lemma shows how we can translate high probability bounds on $\mathcal{D}$ to high probability bounds on $\mathcal{D}(k)$, using this binomial tail bound of $\text{Bin}(k_i, p_i)$. Specifically, it shows that our quantity of interest $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$ of the $\hat{p}_i$s can be upper and lower bounded by concentration of the p_is (as described by $f_{\mathcal{D}}(n, \sigma_p^2, \beta)$) plus a binomial tail bound.

Lemma G.1. *Suppose that $\mathcal{D}$ is supported on $[0, 1/2]$. Given $k_i, n \in \mathbb{N}$, σ_p^2, and $\beta \in [0, 1]$, define $\beta' = 2\sqrt{1 - \sqrt[n]{1 - \beta}} = \Theta(\sqrt{\beta/n})$ and assume that for all p_i in the support of $\mathcal{D}$,*

$$\Pr_{\hat{p}_i \sim \text{Bin}(k_i, p_i)}(p_i - \hat{p}_i \geq f_{\text{Bin}}(k_i, p_i, \beta')) \geq \frac{1}{2}\beta' \quad and \quad \Pr_{\hat{p}_i \sim \text{Bin}(k_i, p_i)}(\hat{p}_i - p_i \geq f_{\text{Bin}}(k_i, p_i, \beta')) \geq \frac{1}{4}\beta'.$$

Then for all $\beta \in [0, 1]$, for all $i \in [n]$,

$$f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) \leq f_{\mathcal{D}}(n, \sigma_p^2, \beta/2) + f_{\text{Bin}}(k_i, p_{\max}, \beta/n),$$

where $p_{\max} = \min\{1/2, p + f_{\mathcal{D}}(n, \sigma_p, \beta/2)\}$. Further, for all $i \in [n]$,

$$f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) \geq f_{\mathcal{D}}(1, \sigma_p^2, \beta') + f_{\text{Bin}}(k_i, p_{\max}, \beta').$$

We note that the conditions on $\mathcal{D}$ and $\text{Bin}(k_i, p_i)$ are mild. The condition on the tails of $\text{Bin}(k_i, p_i)$ is intuitively claiming that $\text{Bin}(k_i, p_i)$ is symmetric. This occurs whenever k_i is large enough, and p_i is bounded away from 0 or 1. We conjecture that the condition that $\mathcal{D}$ is supported on $[0, 1/2]$ can be relaxed but leave the relaxation to future work.

Proof of Lemma G.1. Notice that if $p < q < 1/2$ then $f_{\text{Bin}}(k_i, p, \beta) \leq f_{\text{Bin}}(k_i, q, \beta)$. Let us first consider the upper bound first. With probability $1 - \frac{\beta}{2}$,

$$\text{for all } i, |p - p_i| \leq f_{\mathcal{D}}(n, \sigma_p, \beta/2). \tag{25}$$

Further if Equation 25 holds then we have that with probability $1 - \frac{\beta}{2n}$,

$$|\hat{p}_i - p_i| \leq f_{\text{Bin}}(k_i, p_i, \frac{\beta}{2n}) \leq f_{\text{Bin}}(k_i, p_{\max}, \frac{\beta}{2n}).$$

Thus, for all i,

$$|p - p_i| \leq f_{\mathcal{D}}(n, \sigma_p, \beta/2) + f_{\text{Bin}}(k_i, p_{\max}, \frac{\beta}{2n}).$$

Now, for the lower bound, let $\beta' = \sqrt{8}\sqrt{1 - \sqrt[n]{1 - \beta}}$ and $\alpha = f_{\mathcal{D}}(1, \sigma_p^2, \beta')$. Note that either

$$\Pr_{p_i \sim \mathcal{D}}(p_i - p \geq f_{\mathcal{D}}(1, \sigma_p^2, \beta')) \geq \frac{1}{2}\beta' \quad \text{or} \quad \Pr_{p_i \sim \mathcal{D}}(p - p_i \geq f_{\mathcal{D}}(1, \sigma_p^2, \beta')) \geq \frac{1}{2}\beta'.$$

Assume without loss of generality that $\Pr_{p_i \sim \mathcal{D}}(p_i - p \geq f_{\mathcal{D}}(1, \sigma_p^2, \beta')) \geq \frac{1}{2}\beta'$. Then by assumption,

$$\Pr_{\hat{p}_i \sim \text{Bin}(k_i, p_i)}(\hat{p}_i - p_i \geq f_{\text{Bin}}(k_i, p_i, \beta')) \geq \frac{1}{4}\beta'$$

Then

$$\Pr\left(\max_i |\widehat{p}_i - p| \geq f_{\mathcal{D}}(1, \sigma_p^2, \beta') + f_{\mathrm{Bin}}(k_i, p + \alpha, \beta')\right)$$
$$\geq \Pr\left(\exists i \text{ s.t. } p_i - p \geq f_{\mathcal{D}}(1, \sigma_p^2, \beta') \text{ and } \widehat{p}_i - p_i \geq f_{\mathrm{Bin}}(k_i, p_i, \beta')\right)$$
$$= 1 - \Pr\left(\forall i, p_i - p \leq f_{\mathcal{D}}(1, \sigma_p^2, \beta') \text{ or } \widehat{p}_i - p_i \leq f_{\mathrm{Bin}}(k_i, p + \alpha, \beta')\right)$$
$$= 1 - \left(\Pr\left(p_i - p \leq f_{\mathcal{D}}(1, \sigma_p^2, \beta') \text{ or } \widehat{p}_i - p_i \leq f_{\mathrm{Bin}}(k_i, p + \alpha, \beta')\right)\right)^n.$$

Now,

$$\Pr\left(p_i - p \leq f_{\mathcal{D}}(1, \sigma_p^2, \beta') \text{ or } \widehat{p}_i - p_i \leq f_{\mathrm{Bin}}(k_i, p + \alpha, \beta')\right)$$
$$= 1 - \Pr\left(p_i - p \geq f_{\mathcal{D}}(1, \sigma_p^2, \beta') \text{ and } \widehat{p}_i - p_i \geq f_{\mathrm{Bin}}(k_i, p + \alpha, \beta')\right)$$
$$= 1 - \Pr\left(p_i - p \geq f_{\mathcal{D}}(1, \sigma_p^2, \beta')\right) \Pr\left(\widehat{p}_i - p_i \geq f_{\mathrm{Bin}}(k_i, p + \alpha, \beta') \mid p_i - p \geq f_{\mathcal{D}}(1, \sigma_p^2, \beta')\right)$$
$$\leq 1 - \Pr\left(p_i - p \geq f_{\mathcal{D}}(1, \sigma_p^2, \beta')\right) \Pr\left(\widehat{p}_i - p_i \geq f_{\mathrm{Bin}}(k_i, p_i, \beta') \mid p_i - p \geq f_{\mathcal{D}}(1, \sigma_p^2, \beta')\right)$$
$$\leq 1 - \Pr\left(p_i - p \geq f_{\mathcal{D}}(1, \sigma_p^2, \beta')\right) \Pr\left(\widehat{p}_i - p_i \geq f_{\mathrm{Bin}}(k_i, p_i, \beta')\right)$$
$$\leq 1 - \frac{1}{8}(\beta')^2$$

where the first inequality comes from $p_i \geq p + \alpha$, so $f_{\mathrm{Bin}}(k_i, p + \alpha, \beta') \leq f_{\mathrm{Bin}}(k_i, p_i, \beta')$ So, finally,

$$\Pr\left(\max_i |\widehat{p}_i - p| \geq f_{\mathcal{D}}(1, \sigma_p^2, \beta') + f_{\mathrm{Bin}}(k_i, p + \alpha, \beta')\right) \geq 1 - (1 - (\beta'/\sqrt{8})^2)^n = \beta,$$

which implies the result. $\qquad\square$

G.1 Extending Our Results to Unknown $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$ settings

Lemma G.1 gives both upper bound and lower bounds on $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$, which can be used to modify Algorithm 2 and extend Theorem 4.1 to apply in the setting where $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$ is unknown, but $f_{\mathcal{D}}(n, \sigma_p^2, \beta)$ is known instead.

Recall that the concentration bound $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$ is used in Algorithm 2 to define the truncation parameters $\widehat{a}_i$ and $\widehat{b}_i$, and that we would like to define a truncation window $[\widehat{a}_i, \widehat{b}_i]$ that both contains $[a_i, b_i]$ (so that with high probability none of the $\widehat{p}_i$ are truncated), and is not too wide, so $|\widehat{b}_i - \widehat{a}_i| \leq 6|b_i - a_i|$ (in order to invoke Lemma 4.2).

The following lemma proposes new values for $\hat{a}_i$ and $\hat{b}_i$ for the setting where only $f_{\mathcal{D}}(n, \sigma_p^2, \beta)$ is known, but not $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$. It combines the bounds on $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$ from Lemma G.1, with the bounds on $\widehat{p}^{initial}$ from Lemma F.1 to show that $|\widehat{b}_i - \widehat{a}_i| \leq 6|b_i - a_i|$, as desired.

Lemma G.2. *For $\alpha > 0$, let*

$$\widehat{a}_i = \max\left\{0, \widehat{p} - \alpha - f_{\mathcal{D}}(n, \widehat{\sigma_p^2}, \beta/2) - f_{\mathrm{Bin}}(k_i, \widehat{p} + \alpha + f_{\mathcal{D}}(n, \widehat{\sigma_p}, \beta/2), \beta/n)\right\}$$

and

$$\widehat{b}_i = \min\left\{1, \widehat{p} + \alpha + f_{\mathcal{D}}(n, \widehat{\sigma_p^2}, \beta/2) + f_{\mathrm{Bin}}(k_i, \widehat{p} + \alpha + f_{\mathcal{D}}(n, \widehat{\sigma_p^2}, \beta/2), \beta/n)\right\}.$$

If $\widehat{\sigma_p^2} \geq \sigma_p^2$, and $|p - \widehat{p}| \leq \alpha$, then for all $i \in [n]$,

$$[a_i, b_i] \subset [\widehat{a}_i, \widehat{b}_i].$$

Further, if $\alpha \leq f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$ and $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) \geq \Omega(f_{\mathcal{D}}(n, \sigma_p^2, \beta) + f_{\mathrm{Bin}}(k_i, \min\{1/2, p + f_{\mathcal{D}}(n, \sigma_p, \beta/2)\}, \beta/n))$ then

$$|\widehat{b}_i - \widehat{a}_i| \leq 6|b_i - a_i|.$$

Proof of Lemma G.2. Let us first show that $[\widehat{a}_i, \widehat{b}_i] \subset [a_i, b_i]$. Using our modified definition of $\widehat{a}_i$ given above, we have,

$$
\begin{aligned}
\widehat{a}_i &= \widehat{p}_\epsilon^{\text{initial}} - \alpha - f_{\mathcal{D}}(n, \widehat{\sigma_p^2}, \beta/2) - f_{\text{Bin}}(k_i, \widehat{p}_\epsilon^{\text{initial}} + \alpha + f_{\mathcal{D}}(n, \widehat{\sigma_p^2}, \beta/2), \beta/n) \\
&\leq p - f_{\mathcal{D}}(n, \widehat{\sigma_p^2}, \beta/2) - f_{\text{Bin}}(k_i, p + f_{\mathcal{D}}(n, \widehat{\sigma_p^2}, \beta/2), \beta/n) \\
&\leq p - f_{\mathcal{D}}(n, \sigma_p^2, \beta/2) - f_{\text{Bin}}(k_i, p + f_{\mathcal{D}}(n, \sigma_p, \beta/2), \beta/n) \\
&\leq p - f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) \\
&= a_i.
\end{aligned}
$$

The first two inequalities respectively follow from the accuracy conditions on $\texttt{mean}_{\epsilon,\delta}$ and $\texttt{variance}_{\epsilon,\delta}$ in Theorem 4.1; the third inequality comes from Lemma G.1; and the final equality is by the definition of a_i. A symmetric result that $\widehat{b}_i \geq b_i$ follows similarly.

The second statement of this lemma ensures that the width of the truncation parameter is not more than a constant factor larger than the ideal. Specifically,

$$
\begin{aligned}
|\widehat{b}_i - \widehat{a}_i| &\leq 2\alpha + 2\left(f_{\mathcal{D}}(n, \widehat{\sigma_p^2}, \beta/2) + f_{\text{Bin}}(k_i, \widehat{p} + \alpha + f_{\mathcal{D}}(n, \widehat{\sigma_p^2}, \beta/2), \beta/n) \right) \\
&\leq 2 f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) + O(f_{\mathcal{D}}(1, \sigma_p^2, \beta') - f_{\text{Bin}}(k_i, p + \alpha, \beta')) \\
&\leq 2 f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) + 2\left(2 f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) \right) \\
&\leq 6 f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) \\
&= 6|b_i - a_i|
\end{aligned}
$$

$\square$

We note that Lemma 4.2 as stated requires $|\widehat{b}_i - \widehat{a}_i| \leq 4|b_i - a_i|$, rather than $6|b_i - a_i|$, this difference of constants will only affect the constant C in Theorem 4.1, and the main claim of a constant approximation in variance will still hold with these new $\hat{a}_i$ and $\hat{b}_i$ values.

We will, however, have to add an additional assumption to Theorem 4.1 in this setting. We will need to assume that $\mathcal{D}$ is s.t. $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta) \geq \Omega(f_{\mathcal{D}}(n, \sigma_p^2, \beta) + f_{\text{Bin}}(k_i, \min\{1/2, p + f_{\mathcal{D}}(n, \sigma_p, \beta/2)\}, \beta/n))$, to satisfy the condition of Lemma G.2. This condition is related to the high probability bound on $\mathcal{D}(k)$. The right hand side of this condition is the high probability bound on $\mathcal{D}(k)$ that is inherited directly from the high probability bounds on $\mathcal{D}$ and $\text{Bin}(k, p)$. Without further assumptions on $\mathcal{D}$, this is the best upper bound on $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$ that we can obtain, and hence is the bound used in the truncation in $\widehat{p}_\epsilon^{\text{realistic}}$. The condition states that this upper bound is within a constant multiplicative factor of the true value $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$. We note that this condition is guaranteed by the lower bound on $f_{\mathcal{D}}^{k_i}(n, \sigma_p^2, \beta)$ in Lemma G.1 for $\mathcal{D}$ with support on $[0, 1/2]$, and we conjecture that it holds more broadly.