On counterfactual inference with unobserved confounding

Abstract

Given an observational study with n independent but heterogeneous units and one p-dimensional sample per unit containing covariates, interventions, and outcomes, our goal is to learn counterfactual distribution for each unit. We consider studies with unobserved confounding which introduces statistical biases between interventions and outcomes as well as exacerbates the heterogeneity across units. Modeling the underlying joint distribution as an exponential family and under suitable conditions, we reduce learning the n unit-level counterfactual distributions to learning n exponential family distributions with heterogeneous parameters and only one sample per distribution. We introduce a convex objective that pools all n samples to jointly learn all n parameters and provide a unit-wise mean squared error bound that scales linearly with the metric entropy of the parameter space. For example, when the parameters are s-sparse linear combination of k known vectors, the error is $O(s \log k/p)$. En route, we derive sufficient conditions for compactly supported distributions to satisfy the logarithmic Sobolev inequality.

1 Introduction

We are interested in the problem of unit-level counterfactual inference owing to the increasing importance of personalized decision-making in many domains. As a motivating example, consider a recommender system interacting with a user over time. At each time, the user is exposed to a product based on observed demographic factors as well as certain unobserved factors, and the user's engagement level is recorded. The engagement level at any time can depend sequentially on the prior interaction in addition to the ongoing interaction (see Fig. 1(a)). The system can also sequentially adapt its recommendation. Given historical data of many heterogeneous users, the system wants to infer each user's average engagement level if it were exposed to a different sequence of products while the observed and the unobserved factors remain unchanged. This task is challenging since: (a) the *unobserved* factors could give rise to spurious associations, (b) the users could be *heterogeneous* in that they may have different responses to same sequence of products, and (c) each user only provides a *single* interaction trajectory.

In a general problem, we consider an observational setting where a unit undergoes interventions denoted by **a**. We denote the outcomes of interest by **y**, and allow the interventions **a** and the outcomes **y** to be confounded by observed covariates **v** as well as unobserved covariates **z**. The graphical structure shown in Fig. 1(b) captures these interactions and is at the heart of our problem. We consider n heterogeneous and independent units indexed by $i \in [n] \triangleq \{1, \cdots, n\}$, and assume access to one observation per unit with $\boldsymbol{v}^{(i)}$, $\boldsymbol{a}^{(i)}$, and $\boldsymbol{y}^{(i)}$ denoting the realizations of **v**, **a**, and **y** for unit i respectively.

We operate with the Neyman-Rubin potential outcomes framework [21, 24] and denote the potential outcome of unit $i \in [n]$ under interventions $\boldsymbol{a}$ by $\boldsymbol{y}^{(i)}(\boldsymbol{a})$. Given the realizations

(a) A graphical model for sequential recommender system   (b) A generic model for our setting

Figure 1: Two graphical models illustrating aspects of our work. The directed and the bi-directed arrows indicate causation and association respectively. Panel **(a)** visualizes a sequential recommender system interacting with a user for 3 time points where v_t, a_t, and y_t denote the user's observed demographic factors, the product exposed to the user, and the user's engagement level respectively at time t, and $\mathbf{z}$ denotes unobserved factors. The left plot illustrates the dependency of the observed variables (v_t, a_t, y_t) at time t, on the observed variables at time $t-1$ via $\dashrightarrow$, and on the unobserved covariates $\mathbf{z}$ via $\leftdashedarrow$; these dependencies for time 1 and 2 are expanded in the right plot. Panel **(b)** exhibits a generic graphical model depicting the relationship between $\mathbf{z}$, $\mathbf{v}$, $\mathbf{a}$, and $\mathbf{y}$ for every unit.

37 $\left\{ (\boldsymbol{v}^{(i)}, \boldsymbol{a}^{(i)}, \boldsymbol{y}^{(i)}) \right\}_{i=1}^{n}$, our goal is to answer counterfactual questions for these n units, e.g., what
38 would the potential outcomes $\boldsymbol{y}^{(i)}(\widetilde{\boldsymbol{a}}^{(i)})$ for interventions $\widetilde{\boldsymbol{a}}^{(i)} \neq \boldsymbol{a}^{(i)}$ be, while the observed and
39 unobserved covariates remain unchanged? Under the stable unit treatment value assumption (SUTVA),
40 i.e., the potential outcomes of unit i are not affected by the interventions at other units, learning
41 unit-level counterfactual distributions is equivalent to learning unit-level conditional distributions

$$\left\{ f_{\mathbf{y}|\mathbf{a},\mathbf{z},\mathbf{v}}(\mathbf{y} = \cdot | \mathbf{a} = \cdot, \boldsymbol{z}^{(i)}, \boldsymbol{v}^{(i)}) \right\}_{i=1}^{n}. \tag{1}$$

42 Here, the i-th distribution represents the conditional distribution for the outcomes $\mathbf{y}$ as a function of
43 the interventions $\mathbf{a}$, while keeping the observed covariates $\mathbf{v}$ and the unobserved covariates $\mathbf{z}$ fixed at
44 the corresponding realizations for unit i, i.e., $\boldsymbol{v}^{(i)}$ and $\boldsymbol{z}^{(i)}$ respectively.

45 It is infeasible to answer such questions without any structural assumptions due to two key challenges:
46 (a) unobserved confounding and (b) heterogeneity in unit-level conditional distributions. First,
47 the unobserved covariates $\mathbf{z}$ introduces statistical dependence between interventions and outcomes,
48 known as *unobserved confounding*, which results in biased estimates. Second, the realizations
49 $\left\{ (\boldsymbol{z}^{(i)}, \boldsymbol{v}^{(i)}) \right\}_{i=1}^{n}$ could be different for different units leading to *heterogeneity* in conditional
50 distributions across units. The heterogeneity is crucial since we only observe one realization, namely
51 the outcomes $\boldsymbol{y}^{(i)}(\boldsymbol{a}^{(i)})$ under the interventions $\boldsymbol{a}^{(i)}$, that is consistent with the unit-level conditional
52 distribution $f_{\mathbf{y}|\mathbf{a},\mathbf{z},\mathbf{v}}(\boldsymbol{y}|\boldsymbol{a}, \boldsymbol{z}^{(i)}, \boldsymbol{v}^{(i)})$. As a result, one needs to learn n heterogeneous conditional
53 distributions while having access to only one sample from each of them.

54 In this work, we model the joint distribution of the unobserved covariates, the observed covariates,
55 the intervention, and the outcomes of interest as an exponential family distribution in accordance
56 with the principle of maximum entropy.[1] With this modeling assumption, we show that both the
57 aforementioned challenges can be tackled. In particular, we show that the n unit-level conditional
58 distributions in (1) lead to n distributions from the same exponential family albeit with parameters
59 that vary across units. The parameter corresponding to the i^{th} unit, say $\phi^{\star(i)}$, captures the effect of
60 $\boldsymbol{z}^{(i)}$, and helps tackle the challenge of unobserved confounding. However, the challenge still remains
61 to learn n heterogeneous exponential family distributions with one sample per distribution. This
62 has been addressed in two specific scenarios in the literature: (a) if the unobserved confounding
63 is identical across units, i.e., the parameters $\{\phi^{\star(i)}\}_{i=1}^{n}$ were all equal, then the challenge boils
64 down to learning parameters of a single exponential family distribution from n samples which has
65 been well-studied (see [26] for an overview); (b) if $\mathbf{v}$, $\mathbf{a}$, and $\mathbf{y}$ take binary values and have pairwise
66 interactions, and the dependencies between them are known, then the challenge boils down to learning
67 certain parameters of an Ising model with one sample which has been studied in [12, 18]. In this

---

[1]Exponential family distributions are the maximum entropy distributions given linear constraints on distributions such as bounded moments (see [8, 10]). The exponential family considered in this work arise when the first and second moments of the joint vector $(\mathbf{z}, \mathbf{v}, \mathbf{a}, \mathbf{y})$ are bounded.

work, we consider a generalized setting, where **v**, **a**, and **y** can be either discrete, continuous, or both, and assume no knowledge of the underlying dependencies.

Our contributions. As the main contribution, this work introduces a method to learn unit-level counterfactual distributions from observational studies in the presence of unobserved confounding with one sample per unit. Specifically, for every unit $i \in [n]$, we reduce learning its counterfactual distribution to learning $\phi^{\star(i)}$ from one sample $(\boldsymbol{v}^{(i)}, \boldsymbol{a}^{(i)}, \boldsymbol{y}^{(i)})$ where $\phi^{\star(i)}$ is the parameter of the exponential family distribution corresponding to unit i. Our technical contributions are as follows.

1. We introduce a convex loss function (Def. 1) that pools the data $\left\{(\boldsymbol{v}^{(i)}, \boldsymbol{a}^{(i)}, \boldsymbol{y}^{(i)})\right\}_{i=1}^{n}$ across all n samples to jointly learn all n parameters $\{\phi^{\star(i)}\}_{i=1}^{n}$.

2. For every unit i, we establish that the mean squared errors of our estimates of (a) $\phi^{\star(i)}$ (Thm. 1), and (b) the expected potential outcomes under alternate interventions (Thm. 2) scale linearly with the metric entropy of the parameter space, i.e., the set $\phi^{\star(i)}$ comes from. In particular, when $\phi^{\star(i)}$ is s-sparse linear combination of k known vectors (see Sec. 4), the error—just with one sample—decays as $O(s \log k/p)$, where p is the dimension of the tuple (**v**, **a**, **y**).

3. As part of our analysis, we (a) derive sufficient conditions for a continuous random vector supported on a compact set to satisfy the logarithmic Sobolev inequality (Prop. 1) and (b) provide new concentration bounds for arbitrary functions of a continuous random vector that satisfies the logarithmic Sobolev inequality (Prop. 2). These results could be of independent interest.

Notation. For a vector $\boldsymbol{u} \in \mathbb{R}^p$, we use u_t to denote its t^{th} coordinate and $v_{-t} \in \mathbb{R}^{p-1}$ to denote the vector after deleting the t^{th} coordinate. We denote the ℓ_0, ℓ_p ($p \geq 1$), and ℓ_∞ norms of a vector $\boldsymbol{v}$ by $\|\boldsymbol{v}\|_0$, $\|\boldsymbol{v}\|_p$, and $\|\boldsymbol{v}\|_\infty$ respectively. For a matrix $\mathbf{M} \in \mathbb{R}^{p \times p}$, we denote the element in t^{th} row and u^{th} column by $\mathbf{M}_{tu}$, the t^{th} row by $\mathbf{M}_t$, and the matrix maximum norm by $\|\mathbf{M}\|_{\max}$.

Related work. We provide an overview of related work that focus either on exponential family learning or on unit-level counterfactual inference with unobserved confounding. We refer the reader to [17, 15] for counterfactual inference with no unobserved confounding (as well as closely related concepts of ignorability in statistics and selection on the observables in economics). Likewise, we refer the reader to [22, 23] for counterfactual inference when the underlying causal mechanism (i.e., the directed acyclic graph) is known unlike our work.

Exponential family learning. There has been extensive work on learning parameters of a single exponential family distribution from multiple samples (see [26] for an overview). Of particular interest are the works that focus on learning sparse Markov random fields with (a) discrete variables [28] and (b) continuous variables [27] which inspire our loss function. Recently, a few works [18, 12] have focused on learning Ising model, i.e., sparse Markov random fields with binary variables, with one sample. However, these works focus on special cases where either the dependencies between the variables or a specific subset of the parameters are already known.

Unit-level counterfactual inference. For unit-level inference with unobserved confounding, prior work has largely focused on latent factor models, where the interventions and potential outcomes are assumed to be independent conditional on latent factors. These include popular frameworks of difference-in-differences [9, 5], synthetic controls [2, 1], synthetic difference-in-differences [6], and synthetic interventions [3]. A recent work [13] provides a latent factor model based approach for counterfactual inference in sequential experiments where the treatment mechanism is designed and known, and there is no confounding by definition. Notably these works allow only for finitely many interventions, and need multiple units to be simultaneously treated with the same interventions for a period of time (for their estimation strategies to work). Another key difference is that these works directly learn the outcomes, and not the distributions like we do.

2 Problem formulation

We consider a counterfactual inference task where units go through $p_a \geq 1$ interventions. For every unit, we observe $p_y \geq 1$ outcomes of interest. The interventions and the outcomes could be confounded by $p_v \geq 1$ observed covariates as well as $p_z \geq 1$ unobserved covariates. Additionally, the observed covariates and the unobserved covariates could be arbitrarily associated. We denote the random vector associated with the interventions, the outcomes, the observed covariates, and the unobserved covariates by $\mathbf{a} \triangleq (a_1, \cdots, a_{p_a}) \in \mathcal{A}^{p_a}$, $\mathbf{y} = (y_1, \cdots, y_{p_y}) \in \mathcal{Y}^{p_y}$, $\mathbf{v} \triangleq (v_1, \cdots, v_{p_v}) \in \mathcal{V}^{p_v}$, and $\mathbf{z} \triangleq (z_1, \cdots, z_{p_z}) \in \mathcal{Z}^{p_z}$ respectively where $\mathcal{A}, \mathcal{Y}, \mathcal{V}$, and $\mathcal{Z}$ denote the support of

interventions, outcomes, observed covariates, and unobserved covariates respectively. We allow these sets to contain discrete, continuous, or mixed values. We summarize the causal relationship between the random vectors $\mathbf{z}$, $\mathbf{v}$, $\mathbf{a}$, and $\mathbf{y}$ in Fig. 1(b) where we denote the arbitrary association between $\mathbf{z}$ and $\mathbf{v}$ by a bi-directed arrow, and the causal association between (i) $(\mathbf{z}, \mathbf{v})$ and $\mathbf{a}$, (ii) $(\mathbf{z}, \mathbf{v})$ and $\mathbf{y}$, and (iii) $\mathbf{a}$ and $\mathbf{y}$ by directed arrows. Fig. 1(a) provides an example for sequential recommender systems covered by our work where (i) a_{t+1} depends on a_t in addition to v_{t+1} and $\mathbf{z}$, and (ii) y_{t+1} depends on a_t and y_t in addition to a_{t+1}, v_{t+1} and $\mathbf{z}$. We do not assume any knowledge of such low-level causal links between elements of $\mathbf{z}$, $\mathbf{v}$, $\mathbf{a}$, and $\mathbf{y}$.

Data. We are interested in answering counterfactual questions regarding n heterogeneous but independent units in a population. To do so, we assume access to one observation of the observed covariates, the interventions, and the outcomes per unit, and index it by $i \in [n]$, i.e., $\boldsymbol{v}^{(i)}$, $\boldsymbol{a}^{(i)}$, and $\boldsymbol{y}^{(i)}$ denote the realizations of $\mathbf{v}$, $\mathbf{a}$, and $\mathbf{y}$ for unit i respectively. For every realized tuple $(\boldsymbol{v}^{(i)}, \boldsymbol{a}^{(i)}, \boldsymbol{y}^{(i)})$, there is a corresponding realization $\boldsymbol{z}^{(i)}$ of the unobserved covariates $\mathbf{z}$ that is unobserved.

Potential outcomes framework. We adopt the potential outcomes framework and denote the potential outcomes of unit $i \in [n]$ under interventions $\boldsymbol{a} \in \mathcal{A}^{p_a}$ by $\boldsymbol{y}^{(i)}(\boldsymbol{a})$. We work under the stable unit treatment value assumption (SUTVA) where the potential outcomes of any unit i are unaffected by the interventions at other units. In fact, we assume independence across units implying the potential outcomes of any unit i are also unaffected by the covariates and the potential outcomes at other units. Then, the observations are related to the potential outcomes as $\boldsymbol{y}^{(i)} = \boldsymbol{y}^{(i)}(\boldsymbol{a}^{(i)})$ for all $i \in [n]$. To establish equivalence between unit-level counterfactual distribution and unit-level conditional distribution, consider unit $i \in [n]$ and fix the observed covariates and the unobserved covariates at $\boldsymbol{v}^{(i)}$ and $\boldsymbol{z}^{(i)}$ respectively. Suppose $\widetilde{\boldsymbol{y}}^{(i)}$ is the realized value of $\mathbf{y}$ when $\mathbf{a} = \widetilde{\boldsymbol{a}}^{(i)}$. We are interested in the distribution of the potential outcomes of unit i for interventions $\widetilde{\boldsymbol{a}}^{(i)}$, i.e., the distribution of $\boldsymbol{y}^{(i)}(\widetilde{\boldsymbol{a}}^{(i)})|\mathbf{z} = \boldsymbol{z}^{(i)}, \mathbf{v} = \boldsymbol{v}^{(i)}$. Under the causal framework considered here (see Fig. 1(b)), it is equivalent to the distribution of $\boldsymbol{y}^{(i)}(\widetilde{\boldsymbol{a}}^{(i)})|\mathbf{a} = \widetilde{\boldsymbol{a}}^{(i)}, \mathbf{z} = \boldsymbol{z}^{(i)}, \mathbf{v} = \boldsymbol{v}^{(i)}$ since $(\mathbf{z}, \mathbf{v})$ satisfy ignorability [22, 17], i.e., the potential outcomes are independent of the interventions given $(\mathbf{z}, \mathbf{v})$. Further, under SUTVA, it is equivalent to distribution of $\widetilde{\boldsymbol{y}}^{(i)}|\mathbf{a} = \widetilde{\boldsymbol{a}}^{(i)}, \mathbf{z} = \boldsymbol{z}^{(i)}, \mathbf{v} = \boldsymbol{v}^{(i)}$, i.e., $f_{\mathbf{y}|\mathbf{a},\mathbf{z},\mathbf{v}}(\mathbf{y} = \cdot|\mathbf{a} = \widetilde{\boldsymbol{a}}^{(i)}, \boldsymbol{z}^{(i)}, \boldsymbol{v}^{(i)})$. Therefore, our goal is to learn the n unit-level conditional distributions in (1). To that end, we let the joint distribution of the $\mathbf{z}$, $\mathbf{v}$, $\mathbf{a}$, and $\mathbf{y}$ belong to an exponential family.

Modeling data as exponential family. Let $\widetilde{p} \triangleq p_z + p_v + p_a + p_y$. We parameterize the joint probability distribution $f_{\mathbf{w}}$ of the $\widetilde{p}$-dimensional random vector $\mathbf{w} \triangleq (\mathbf{z}, \mathbf{v}, \mathbf{a}, \mathbf{y})$ by natural statistics $\mathbf{w}$ and $\mathbf{w}\mathbf{w}^\top$, and by natural parameters $\phi \in \mathbb{R}^{\widetilde{p} \times 1}$ and $\Phi \in \mathbb{R}^{\widetilde{p} \times \widetilde{p}}$ as follows

$$f_{\mathbf{w}}(\boldsymbol{w}; \phi, \Phi) \propto \exp\left(\phi^\top \boldsymbol{w} + \boldsymbol{w}^\top \Phi \boldsymbol{w}\right), \quad \text{where} \quad \boldsymbol{w} \triangleq (\boldsymbol{z}, \boldsymbol{x}, \boldsymbol{a}, \boldsymbol{y}),$$

and $\boldsymbol{z} \triangleq (z_1, \cdots, z_{p_z})$, $\boldsymbol{v} \triangleq (v_1, \cdots, v_{p_v})$, $\boldsymbol{a} \triangleq (a_1, \cdots, a_{p_a})$, and $\boldsymbol{y} \triangleq (y_1, \cdots, y_{p_y})$ denote realizations of $\mathbf{z}$, $\mathbf{v}$, $\mathbf{a}$, and $\mathbf{y}$ respectively. Without loss of generality, we can assume Φ^* to be a symmetric matrix. Then, recognizing that the (unit-level) conditional distribution of $\mathbf{y}$ conditioned on $\mathbf{a} = a$, $\mathbf{z} = z$, and $\mathbf{v} = v$ belongs to an exponential family with natural statistics $\mathbf{y}$ and $\mathbf{y}\mathbf{y}^\top$, we can write

$$f_{\mathbf{y}|\mathbf{a},\mathbf{z},\mathbf{v}}(\boldsymbol{y}|\boldsymbol{a}, \boldsymbol{z}, \boldsymbol{v}) \propto \exp\left(\left[\phi^{(y)^\top} + 2\boldsymbol{z}^\top \Phi^{(z,y)} + 2\boldsymbol{v}^\top \Phi^{(v,y)} + 2\boldsymbol{a}^\top \Phi^{(a,y)}\right]\boldsymbol{y} + \boldsymbol{y}^\top \Phi^{(y,y)}\boldsymbol{y}\right), \quad (2)$$

where $\phi^{(y)} \in \mathbb{R}^{p \times 1}$ is the component of ϕ corresponding to $\mathbf{y}$ and $\Phi^{(u,y)} \in \mathbb{R}^{p_u \times p_y}$ is the component of Φ corresponding to $\mathbf{u}$ and $\mathbf{y}$ for all $\mathbf{u} \in \{\mathbf{z}, \mathbf{v}, \mathbf{a}, \mathbf{y}\}$. We make two key observations: (a) The term $\Phi^{(z,y)^\top}z$ captures the effect of unobserved covariates z on $f_{\mathbf{y}|\mathbf{a},\mathbf{z},\mathbf{v}}(\mathbf{y} = \cdot|\mathbf{a} = \cdot, \boldsymbol{z}, \boldsymbol{v})$ in (2). (b) The task of learning $f_{\mathbf{y}|\mathbf{a},\mathbf{z},\mathbf{v}}(\mathbf{y} = \cdot|\mathbf{a} = \cdot, \boldsymbol{z}, \boldsymbol{v})$ in (2) as a function of $\mathbf{a}$ reduces to learning

$$\text{(i) } \phi^{(y)} + 2\Phi^{(z,y)\top}\boldsymbol{z} + 2\Phi^{(v,y)\top}\boldsymbol{v}, \qquad \text{(ii) } \Phi^{(a,y)}, \quad \text{and} \quad \text{(iii) } \Phi^{(y,y)}. \qquad (3)$$

Now, we argue that learning (i), (ii), and (iii) in (3) is subsumed in learning the parameters of the (unit-level) conditional distribution $f_{\mathbf{x}|\mathbf{z}}$ of the random vector $\mathbf{x} \triangleq (\mathbf{v}, \mathbf{a}, \mathbf{y})$ conditioned on $\mathbf{z} = z$. To that end, we recognize that $f_{\mathbf{x}|\mathbf{z}}$ belongs to an exponential family with natural statistics $\mathbf{x}$ and $\mathbf{x}\mathbf{x}^\top$. For all $\mathbf{u} \in \{\mathbf{v}, \mathbf{a}, \mathbf{y}\}$, let $\phi^{(u)} \in \mathbb{R}^{p_u \times 1}$ be the component of ϕ corresponding to $\mathbf{u}$, and $\Phi^{(z,u)} \in \mathbb{R}^{p_z \times p_u}$

be the component of Φ corresponding to $\mathbf{z}$ and $\mathbf{u}$. We parameterize $f_{\mathbf{x}|\mathbf{z}}$ as

$$f_{\mathbf{x}|\mathbf{z}}\big(\boldsymbol{x}|\boldsymbol{z};\theta(\boldsymbol{z}),\Theta\big) \propto \exp\left(\big[\theta(\boldsymbol{z})\big]^{\top}\boldsymbol{x} + \boldsymbol{x}^{\top}\Theta\boldsymbol{x}\right) \text{ where } \theta(\boldsymbol{z}) \triangleq \begin{bmatrix} \phi^{(v)}+2\Phi^{(z,v)\top}\boldsymbol{z} \\ \phi^{(a)}+2\Phi^{(z,a)\top}\boldsymbol{z} \\ \phi^{(y)}+2\Phi^{(z,y)\top}\boldsymbol{z} \end{bmatrix} \in \mathbb{R}^{p\times 1}, \quad (4)$$

$p \triangleq p_v + p_a + p_y$, $\Theta \in \mathbb{R}^{p\times p}$ denotes the component of Φ corresponding to $\mathbf{x}$, $\boldsymbol{x} \triangleq (\boldsymbol{v}, \boldsymbol{a}, \boldsymbol{y})$, and $\boldsymbol{v}$, $\boldsymbol{a}$, and $\boldsymbol{y}$ denote realizations of $\mathbf{v}$, $\mathbf{a}$, and $\mathbf{y}$ respectively. Now, if we learn $\theta(\boldsymbol{z})$ and Θ, we can obtain (i), (ii), and (iii) in (3) for any $\mathbf{v} = \boldsymbol{v}$ by using the appropriate components of $\theta(\boldsymbol{z})$ and Θ.

Inference tasks. Let $f_{\mathbf{w}}(\cdot; \phi^*, \Phi^*)$ denote the true data generating distribution (Sec. 2) of $\mathbf{w}$, and let $f_{\mathbf{x}|\mathbf{z}}\big(\cdot\,|\boldsymbol{z};\theta^\star(\boldsymbol{z}),\Theta^\star\big)$ denote the true distribution of $\mathbf{x}$ conditioned on $\mathbf{z} = \boldsymbol{z}$. Then, for all $i \in [n]$, we note that the realization $\boldsymbol{x}^{(i)} \triangleq (\boldsymbol{v}^{(i)}, \boldsymbol{a}^{(i)}, \boldsymbol{y}^{(i)})$ is consistent with the conditional distribution $f_{\mathbf{x}|\mathbf{z}}\big(\cdot\,|\boldsymbol{z}^{(i)};\theta^\star(\boldsymbol{z}^{(i)}),\Theta^\star\big)$ where we do not observe $\boldsymbol{z}^{(i)}$. Our primary goal is to learn the n unit-level counterfactual distributions, which, as per our earlier discussion simplifies to estimating

(i) Unit-level parameters $\theta^{\star(i)} \triangleq \theta^\star(\boldsymbol{z}^{(i)})$ for $i \in [n]$, and (ii) Population-level parameter $\Theta^\star$. (5)

Our secondary goal is to estimate the expected potential outcomes for any given unit i (with $\mathbf{z} = \boldsymbol{z}^{(i)}, \mathbf{v} = \boldsymbol{v}^{(i)}$) and an alternate intervention $\widetilde{\boldsymbol{a}}^{(i)}$. In particular, our estimand of interest is

$$\mu^{(i)}(\widetilde{\boldsymbol{a}}^{(i)}) \triangleq \mathbb{E}[\boldsymbol{y}^{(i)}(\widetilde{\boldsymbol{a}}^{(i)})|\mathbf{z} = \boldsymbol{z}^{(i)}, \mathbf{v} = \boldsymbol{v}^{(i)}]. \quad (6)$$

Assumptions. We now state the assumptions we require to learn the parameters in (5). For the ease of exposition, we let the sets $\mathcal{V}$, $\mathcal{A}$, and $\mathcal{Y}$ be continuous valued. However, our results apply equally to discrete and mixed cases. Without loss of generality, we let these sets be equal, and let $\mathcal{X} \triangleq \mathcal{V} = \mathcal{A} = \mathcal{Y}$ be bounded and symmetric around 0, i.e., $\mathcal{X} = \{-x_{\max}, x_{\max}\}$ where $x_{\max} < \infty$.

Assumption 1 (Bounded and sparse parameters). *The true model parameters satisfy the following:*

(a) $\theta^{\star(i)}$ and $\Theta^\star$ are bounded for all $i \in [n]$,[2] i.e., $\max\left\{\max_{i\in[n]} \left\|\theta^{\star(i)}\right\|_{\infty}, \|\Theta^\star\|_{\max}\right\} \leq \alpha$.

(b) $\Theta^\star$ is row-wise sparse and has zeros on the diagonal, i.e., $\|\Theta_t^\star\|_0 \leq \beta$ and $\Theta_{tt}^\star = 0$ for all $t \in [p]$. These imply that each $\mathsf{x}_t \in \mathbf{x}$ interacts with only a few other $\mathsf{x}_u \in \mathbf{x}$ in (4).

Below, we define the set Λ_θ such that it contains all $p \times 1$ vectors θ satisfying Assum. 1(a) and the set Λ_Θ such that it contains all $p \times p$ symmetric matrices Θ satisfying Assum. 1(a) and (b).

$$\Lambda_\theta \triangleq \left\{\theta \in \mathbb{R}^{p\times 1} : \|\theta\|_{\infty} \leq \alpha\right\} \text{ and}$$
$$\Lambda_\Theta \triangleq \left\{\Theta \in \mathbb{R}^{p\times p} : \Theta = \Theta^{\top}, \Theta_{tt} = 0 \text{ for all } t \in [p], \|\Theta\|_{\max} \leq \alpha, \max_{t\in[p]} \|\Theta_t\|_0 \leq \beta\right\}. \quad (7)$$

3 Algorithm

We propose a computationally tractable loss function to estimate the unit-level and the population-level parameters in (5). Our algorithm jointly learns all the parameters of interest by pooling the observations across all n units and exploiting the exponential family structure of $\mathbf{v}$, $\mathbf{a}$, and $\mathbf{y}$ conditioned on $\mathbf{z} = \boldsymbol{z}$ in (4). In particular, our loss explicitly utilizes the fact that the population-level parameter $\Theta^\star$ is shared across units. We defer our estimate of the expected potential outcomes in (6) to App. A.1.

Loss function and parameter estimate. Consider any $t \in [p]$. Our loss function is inspired by the conditional distribution $f_{\mathsf{x}_t|\mathbf{x}_{-t},\mathbf{z}}$ of the random variable x_t conditioned on $\mathbf{x}_{-t} = \boldsymbol{x}_{-t}$ and $\mathbf{z} = \boldsymbol{z}$ which is given by

$$f_{\mathsf{x}_t|\mathbf{x}_{-t},\mathbf{z}}\big(x_t|\boldsymbol{x}_{-t},\boldsymbol{z};\theta_t^\star(\boldsymbol{z}),\Theta_t^\star\big) \propto \exp\left(\big[\theta_t^\star(\boldsymbol{z}) + \Theta_t^{\star\top}\boldsymbol{x}\big]x_t\right), \quad (8)$$

where $\theta_t^\star(\boldsymbol{z})$ is the t^{th} element of $\theta^\star(\boldsymbol{z})$, $\Theta_t^\star$ is the t^{th} row of $\Theta^\star$, and $\boldsymbol{x}$ denotes a realization of $\mathbf{x}$.

---

[2]This bound is necessary for model identifiability [25].

Definition 1 (**Loss function**). *Given a sample $\boldsymbol{x}^{(i)}$ for every unit $i \in [n]$, our loss function maps $\underline{\Theta} \in \mathbb{R}^{p \times (n+p)}$ to $\mathcal{L}(\underline{\Theta}) \in \mathbb{R}$ defined as*

$$\mathcal{L}(\underline{\Theta}) = \frac{1}{n} \sum_{t \in [p]} \sum_{i \in [n]} \exp\left(-[\theta_t^{(i)} + \Theta_t^\top \boldsymbol{x}^{(i)}]x_t^{(i)}\right) \text{ where } \underline{\Theta} \triangleq \begin{bmatrix} \underline{\Theta}_1^\top \\ \vdots \\ \underline{\Theta}_p^\top \end{bmatrix} \text{ with } \underline{\Theta}_t \triangleq \left\{\theta_t^{(1)}, \cdots, \theta_t^{(n)}, \Theta_t\right\}. \tag{9}$$

Our estimate of $\underline{\Theta}^\star$ (defined analogous to $\underline{\Theta}$) is obtained by minimizing the convex function $\mathcal{L}(\underline{\Theta})$ over all $\theta^{(i)} \in \Lambda_\theta$ for all $i \in [n]$ and $\Theta \in \Lambda_\Theta$, i.e.,

$$\widehat{\underline{\Theta}} \in \underset{\underline{\Theta} \in \Lambda_\theta^n \times \Lambda_\Theta}{\arg\min} \, \mathcal{L}(\underline{\Theta}). \tag{10}$$

We note (10) is a convex optimization problem, and a projected gradient descent algorithm returns an ϵ-optimal estimate with $\tau = O(p/\epsilon)$ iterations[3] where $\widehat{\underline{\Theta}}_\epsilon$ is said to be an ϵ-optimal estimate if $\mathcal{L}(\widehat{\underline{\Theta}}_\epsilon) \leq \mathcal{L}(\widehat{\underline{\Theta}}) + \epsilon$ for any $\epsilon > 0$.

4 Main results

In this section, we provide our guarantee on estimating the unit-level and the population-level parameters in (5). We defer our guarantee on estimating the expected potential outcomes in (6) to App. A.2. Our guarantees depend on the metric entropy of the set Λ_θ which provides some notion of the richness of Λ_θ. To that end, we define the notions of ε-covering number and metric entropy.

Definition 2 (ε-covering number and metric entropy). *Given a set $\mathcal{V} \subset \mathbb{R}^p$ and a scalar $\varepsilon > 0$, we use $\mathcal{C}(\mathcal{V}, \varepsilon)$ to denote the ε-covering number of $\mathcal{V}$ with respect to $\|\cdot\|_2$, i.e., $\mathcal{C}(\mathcal{V}, \varepsilon)$ denotes the minimum cardinality over all possible covers $\mathcal{U} \subset \mathcal{V}$ that satisfy $\mathcal{V} \subset \cup_{u \in \mathcal{U}} \mathcal{B}(u; \varepsilon)$, where $\mathcal{B}(u; \varepsilon) \triangleq \{v \in \mathbb{R}^p : \|u - v\|_2 \leq \varepsilon\}$ denotes a ball of radius ε in $\mathbb{R}^p$ centered at u. Further, we use $\mathcal{M}(\mathcal{V}, \varepsilon)$ to denote the metric entropy of $\mathcal{V}$, i.e., $\mathcal{M}(\mathcal{V}, \varepsilon) \triangleq \log \mathcal{C}(\mathcal{V}, \varepsilon)$. Lastly, we use the shorthand notation $\mathcal{M}_\theta(\varepsilon) = \mathcal{M}(\Lambda_\theta, \varepsilon)$ to denote the metric entropy of Λ_θ.*

The following result provides the estimation error for the estimate $\widehat{\underline{\Theta}}$ obtained in (10). We divide the proof across App. B and App. C.[4]

Theorem 1 (**Guarantee on parameter estimate**). *Fix an $\varepsilon > 0$ and $\delta \in (0, 1)$, and define*

$$R(\varepsilon, \delta) \triangleq \max\{ce^{c'\beta}\sqrt{\log\log(2p/\delta)) + \mathcal{M}_\theta(r)}, \varepsilon\gamma\}, \quad r \triangleq \frac{ce^{-c'\beta}}{\gamma}, \text{ and } \gamma \triangleq \max_{\theta, \overline{\theta} \in \Lambda_\theta} \frac{\|\theta - \overline{\theta}\|_1}{\|\theta - \overline{\theta}\|_2}. \tag{11}$$

Then, with probability at least $1 - \delta$, the estimates $\widehat{\Theta}, \widehat{\theta}^{(1)}, \cdots, \widehat{\theta}^{(n)}$ defined in (10) satisfy

$$\max_{t \in [p]} \|\widehat{\Theta}_t - \Theta_t^\star\|_2 \leq \varepsilon \qquad \text{whenever} \quad n \geq \frac{ce^{c'\beta}\log\frac{p}{\delta}}{\varepsilon^4} \quad \text{and}$$

$$\max_{i \in [n]} \|\widehat{\theta}^{(i)} - \theta^{\star(i)}\|_2 \leq R(\varepsilon, \delta/n) \quad \text{whenever} \quad n \geq \frac{ce^{c'\beta}(\log\frac{np}{\delta} + \mathcal{M}_\theta(r))}{\varepsilon^4}.$$

Example. Consider the case where $\theta^{\star(i)}$ is s-sparse linear combination of k known vectors for all $i \in [n]$, i.e., $\theta^{\star(i)} = \mathbf{B}\mathbf{a}^{(i)}$ for a known matrix $\mathbf{B} \in \mathbb{R}^{p \times k}$ and $\mathbf{a}^{(i)} \in \mathbb{R}^{k \times 1}$ with $\|\mathbf{a}^{(i)}\|_0 \leq s$ such that $\|\theta^{\star(i)}\|_\infty \leq \alpha$. Then, the parameter set Λ_θ can be re-parameterized as the set $\Lambda_\mathbf{a} = \{\mathbf{a} \in \mathbb{R}^{k \times 1} : \|\mathbf{a}\|_0 \leq s \text{ and } \|\mathbf{B}\mathbf{a}\|_\infty \leq \alpha\}$ whose metric entropy is given by $\mathcal{M}_\mathbf{a}(r) = \left(1 + \frac{c}{r}\right)^s \binom{k}{s}$ for some constant c [12, Corollary 4]. Using this bound, and substituting the worst case $\gamma = \sqrt{p}$, the guarantee in Thm. 1 for $\{\widehat{\theta}^{(i)}\}_{i=1}^n$ simplifies as follows: For any $\varepsilon > 0$ and $\delta \in (0, 1)$, with probability at least $1 - \delta$,

$$\max_{i \in [n]} \|\widehat{\theta}^{(i)} - \theta^{\star(i)}\|_2 \leq \max\{\varepsilon, ce^{c'\beta}\sqrt{s\log pk + \log\log(1/\delta)}\} \quad \text{whenever} \quad n \geq \frac{ce^{c'\beta}sp^2\log\frac{npk}{\delta}}{\varepsilon^4}.$$

---

[3]This follows from [7, Theorem 10.6] by noting that $\mathcal{L}(\underline{\Theta})$ is $O(p)$ smooth function of $\underline{\Theta}$.

[4]To simplify presentation, we use c and c' to denote universal constants or constants that depend on the model-parameters α and $x_{\max}$, and can take a different value in each appearance.

References

[1] A. Abadie, A. Diamond, and J. Hainmueller. Synthetic control methods for comparative case studies: Estimating the effect of california's tobacco control program. *Journal of the American Statistical Association*, 2010.

[2] A. Abadie and J. Gardeazabal. The economic costs of conflict: A case study of the basque country. *American Economic Review*, 2003.

[3] A. Agarwal, D. Shah, and D. Shen. Synthetic interventions. *arXiv preprint arXiv:2006.07691*, 2021.

[4] S. Aida and D. Stroock. Moment estimates derived from poincaré and logarithmic sobolev inequalities. *Mathematical Research Letters*, 1(1):75–86, 1994.

[5] J. D. Angrist and J.-S. Pischke. *Mostly harmless econometrics: An empiricist's companion*. Princeton university press, 2009.

[6] D. Arkhangelsky, S. Athey, D. A. Hirshberg, G. W. Imbens, and S. Wager. Synthetic difference-in-differences. *American Economic Review*, 111(12):4088–4118, December 2021.

[7] M. Bande and N. He. Lower bounds & projected gradient descent. 2016.

[8] O. Barndorff-Nielsen. *Information and exponential families: in statistical theory*. John Wiley & Sons, 2014.

[9] M. Bertrand, E. Duflo, and S. Mullainathan. How much should we trust differences-in-differences estimates? *The Quarterly journal of economics*, 119(1):249–275, 2004.

[10] L. D. Brown. Fundamentals of statistical exponential families: with applications in statistical decision theory. Ims, 1986.

[11] R. Busa-Fekete, D. Fotakis, B. Szörényi, and M. Zampetakis. Optimal learning of mallows block model. In *Conference on Learning Theory*, pages 529–532. PMLR, 2019.

[12] Y. Dagan, C. Daskalakis, N. Dikkala, and A. V. Kandiros. Learning ising models from one or multiple samples. In *Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing*, pages 161–168, 2021.

[13] R. Dwivedi, S. Murphy, and D. Shah. Counterfactual inference for sequential experimental design. *arXiv preprint arXiv:2202.06891*, 2022.

[14] W. Ghang, Z. Martin, and S. Waruhiu. The sharp log-sobolev inequality on a compact interval. 2014.

[15] M. Hernán and J. Robins. *Causal Inference: What If*. Boca Raton: Chapman & Hall/CRC, 2020.

[16] R. Holley and D. W. Stroock. Logarithmic sobolev inequalities and stochastic ising models. 1986.

[17] G. W. Imbens and D. B. Rubin. *Causal inference in statistics, social, and biomedical sciences*. Cambridge University Press, 2015.

[18] V. Kandiros, Y. Dagan, N. Dikkala, S. Goel, and C. Daskalakis. Statistical estimation from dependent data. In *International Conference on Machine Learning*, pages 5269–5278. PMLR, 2021.

[19] M. Ledoux. Logarithmic sobolev inequalities for unbounded spin systems revisited. In *Séminaire de Probabilités XXXV*, pages 167–194. Springer, 2001.

[20] K. Marton. Logarithmic sobolev inequalities in discrete product spaces: a proof by a transportation cost distance. *arXiv preprint arXiv:1507.02803*, 2015.

[21] J. Neyman. Sur les applications de la théorie des probabilités aux experiences agricoles: Essai des principes. *Roczniki Nauk Rolniczych*, 10(1):1–51, 1923.

[22] J. Pearl. *Causality*. Cambridge university press, 2009.

[23] J. Pearl, M. Glymour, and N. P. Jewell. *Causal inference in statistics: A primer*. John Wiley & Sons, 2016.

[24] D. B. Rubin. Estimating causal effects of treatments in randomized and nonrandomized studies. *Journal of educational Psychology*, 66(5):688, 1974.

[25] N. P. Santhanam and M. J. Wainwright. Information-theoretic limits of selecting binary graphical models in high dimensions. *IEEE Transactions on Information Theory*, 58(7):4117–4134, 2012.

[26] A. Shah, D. Shah, and G. Wornell. A computationally efficient method for learning exponential family distributions. *Advances in Neural Information Processing Systems*, 34:15841–15854, 2021.

[27] A. Shah, D. Shah, and G. Wornell. On learning continuous pairwise markov random fields. In *International Conference on Artificial Intelligence and Statistics*, pages 1153–1161. PMLR, 2021.

[28] M. Vuffray, S. Misra, and A. Y. Lokhov. Efficient learning of discrete graphical models. *Journal of Statistical Mechanics: Theory and Experiment*, 2021(12):124017, 2022.

Appendix

A   Causal estimate and associated guarantees

In this section, we provide our estimate of the expected potential outcomes under alternate intervention and the corresponding estimation error.

A.1   Causal estimate

Assuming the estimate $\widehat{\Theta}$ of $\Theta^\star$ is given from Sec. 3, we define our estimate of the expected potential outcome $\mu^{(i)}(\widetilde{\boldsymbol{a}}^{(i)})$ (see (6)) for any given unit $i \in [n]$ under an alternate intervention $\widetilde{\boldsymbol{a}}^{(i)} \in \mathcal{A}^{p_a}$. First, we identify $\widehat{\Phi}^{(u,y)} \in \mathbb{R}^{p_u \times p_y}$ to be the component of $\widehat{\Theta}$ corresponding to u and y for all $\mathsf{u} \in \{\mathsf{v}, \mathsf{a}, \mathsf{y}\}$ and $\widehat{\theta}^{(i,y)} \in \mathbb{R}^{p_y}$ to be the component of $\widehat{\theta}^{(i)}$ corresponding to y. Then, our estimate of the conditional distribution of y as a function of the interventions a, while keeping v and z fixed at the corresponding realizations for unit i, i.e., $\boldsymbol{v}^{(i)}$ and $\boldsymbol{z}^{(i)}$ respectively, is as follows:

$$\widehat{f}^{(i)}_{\mathsf{y}|\mathsf{a}}(\boldsymbol{y}|\boldsymbol{a}) \propto \exp\left(\left[\widehat{\theta}^{(i,y)} + 2\boldsymbol{v}^{(i)\top}\widehat{\Phi}^{(v,y)} + 2\boldsymbol{a}^\top \widehat{\Phi}^{(a,y)}\right]\boldsymbol{y} + \boldsymbol{y}^\top \widehat{\Phi}^{(y,y)}\boldsymbol{y}\right). \tag{12}$$

Finally, our estimate of $\mu^{(i)}(\widetilde{\boldsymbol{a}}^{(i)})$ is given by

$$\widehat{\mu}^{(i)}(\widetilde{\boldsymbol{a}}^{(i)}) \triangleq \mathbb{E}_{\widehat{f}^{(i)}_{\mathsf{y}|\mathsf{a}}}[\mathsf{y}|\mathsf{a} = \widetilde{\boldsymbol{a}}^{(i)}]. \tag{13}$$

A.2   Guarantee on outcome estimate

The following result provides the estimation error for the estimate $\widehat{\mu}^{(i)}(\widetilde{\boldsymbol{a}}^{(i)})$ (see (13)) of the expected potential outcomes for any unit $i \in [n]$ under an alternate intervention $\widetilde{\boldsymbol{a}}^{(i)} \in \mathcal{A}^{p_a}$. The result requires the operator norm of the covariance matrix of y conditioned on a, z, and v to remain bounded for minor perturbation in the parameters. We denote this covariance matrix by $\mathbb{E}_{\theta,\Theta}[\mathsf{y}\mathsf{y}^\top|\boldsymbol{a},\boldsymbol{z},\boldsymbol{v}]$ where the expectation is with respect to the conditional distribution of y conditioned on $\mathsf{a} = \boldsymbol{a}$, $\mathsf{z} = \boldsymbol{z}$, and $\mathsf{v} = \boldsymbol{v}$ which is fully parameterized by θ and Θ, i.e., replace $\theta(\boldsymbol{z})$ by θ in (4). For simplicity, we assume $p_v = p_a = p_y$. See the proof below for the general case.

Theorem 2 (**Guarantee on outcome estimate**). *Fix an $\varepsilon > 0$ and $\delta \in (0, 1)$. Then, with probability at least $1 - \delta$, the estimates $\{\widehat{\mu}^{(i)}(\widetilde{\boldsymbol{a}}^{(i)})\}_{i=1}^n$ (13) for any $\{\widetilde{\boldsymbol{a}}^{(i)} \in \mathcal{A}^{p_a}\}_{i=1}^n$ satisfy*

$$\max_{i\in[n]} \frac{\|\mu^{(i)}(\widetilde{\boldsymbol{a}}^{(i)}) - \widehat{\mu}^{(i)}(\widetilde{\boldsymbol{a}}^{(i)})\|_2}{\sup_{\theta,\Theta\in\mathbb{B}_i}\|\mathbb{E}_{\theta,\Theta}[\mathsf{y}\mathsf{y}^\top|\widetilde{\boldsymbol{a}}^{(i)},\boldsymbol{z}^{(i)},\boldsymbol{v}^{(i)}]\|_{\mathrm{op}}} \leq R(\varepsilon,\delta/n) + p\varepsilon,$$

whenever

$$n \geq \frac{ce^{c'\beta}(\log\frac{np}{\delta} + \mathcal{M}_\theta(r))}{\varepsilon^4},$$

where $R(\varepsilon,\delta)$ and r were defined in (11), and

$$\mathbb{B}_i \triangleq \left\{\theta \in \Lambda_\theta : \|\theta - \theta^{\star(i)}\|_2 \leq R(\varepsilon,\delta/n)\right\} \times \left\{\Theta \in \Lambda_\Theta : \max_{t\in[p]}\|\Theta_t - \Theta_t^\star\|_2 \leq \varepsilon\right\}.$$

Proof. Fix any unit $i \in [n]$ and an alternate intervention $\widetilde{\boldsymbol{a}}^{(i)} \in \mathcal{A}^{p_a}$. Then, we have

$$\mu^{(i)}(\widetilde{\boldsymbol{a}}^{(i)}) \overset{(6)}{=} \mathbb{E}[\boldsymbol{y}^{(i)}(\widetilde{\boldsymbol{a}}^{(i)})|\mathsf{z} = \boldsymbol{z}^{(i)}, \mathsf{v} = \boldsymbol{v}^{(i)}] \overset{(a)}{=} \mathbb{E}[\mathsf{y}|\mathsf{a} = \widetilde{\boldsymbol{a}}^{(i)}, \mathsf{z} = \boldsymbol{z}^{(i)}, \mathsf{v} = \boldsymbol{v}^{(i)}],$$

where (a) follows because the unit-level counterfactual distribution is equivalent to unit-level conditional distribution under the causal framework considered as described in Sec. 2. To obtain a convenient expression for $\mathbb{E}[\mathbf{y}|\mathbf{a} = \widetilde{\boldsymbol{a}}^{(i)}, \mathbf{z} = \boldsymbol{z}^{(i)}, \mathbf{v} = \boldsymbol{v}^{(i)}]$, we identify $\Phi^{\star(u,y)} \in \mathbb{R}^{p_u \times p_y}$ to be the component of $\Theta^\star$ corresponding to $\mathbf{u}$ and $\mathbf{y}$ for all $\mathbf{u} \in \{\mathbf{v}, \mathbf{a}, \mathbf{y}\}$ and $\theta^{\star(i,y)} \in \mathbb{R}^{p_y}$ to be the component of $\theta^{\star(i)}$ corresponding to $\mathbf{y}$. Then, the conditional distribution of $\mathbf{y}$ as a function of the interventions $\mathbf{a}$, while keeping $\mathbf{v}$ and $\mathbf{z}$ fixed at the corresponding realizations for unit i, i.e., $\boldsymbol{v}^{(i)}$ and $\boldsymbol{z}^{(i)}$ respectively, can be written as

$$f^{(i)}{}_{\mathbf{y}|\mathbf{a}}(\boldsymbol{y}|\boldsymbol{a}) \propto \exp\left(\left[\theta^{\star(i,y)} + 2\boldsymbol{v}^{(i)\top}\Phi^{\star(v,y)} + 2\boldsymbol{a}^\top\Phi^{\star(a,y)}\right]\boldsymbol{y} + \boldsymbol{y}^\top\Phi^{\star(y,y)}\boldsymbol{y}\right). \quad (14)$$

Therefore, we have

$$\mathbb{E}[\mathbf{y}|\mathbf{a} = \widetilde{\boldsymbol{a}}^{(i)}, \mathbf{z} = \boldsymbol{z}^{(i)}, \mathbf{v} = \boldsymbol{v}^{(i)}] = \mathbb{E}_{f^{(i)}{}_{\mathbf{y}|\mathbf{a}}}[\mathbf{y}|\mathbf{a} = \widetilde{\boldsymbol{a}}^{(i)}].$$

Now, consider the p_u dimensional random vector $\mathbf{u}$ supported on $\mathcal{X}^{p_u}$ with distribution $f_\mathbf{u}$ parameterized by $\psi \in \mathbb{R}^{p_y}$ and $\Psi \in \mathbb{R}^{p_y \times p_y}$ as follows

$$f_\mathbf{u}(\boldsymbol{u}|\psi, \Psi) \propto \exp(\psi^\top\boldsymbol{u} + \boldsymbol{u}^\top\Psi\boldsymbol{u}). \quad (15)$$

Then, note that $\widehat{f}^{(i)}_{\mathbf{y}|\mathbf{a}}(\boldsymbol{y}|\boldsymbol{a})$ in (12) and $f^{(i)}{}_{\mathbf{y}|\mathbf{a}}(\boldsymbol{y}|\boldsymbol{a})$ in (14) belong to the set $\{f_\mathbf{u}(\cdot|\psi, \Psi) : \psi \in \mathbb{R}^{p_y}, \Psi \in \mathbb{R}^{p_y \times p_y}\}$ for some ψ and Ψ. Now, we consider any two distributions in this set, namely $f_\mathbf{u}(\boldsymbol{u}|\widehat{\psi}, \widehat{\Psi})$ and $f_\mathbf{u}(\boldsymbol{u}|\psi^\star, \Psi^\star)$. Then, we claim that the two norm of the difference of the mean vectors of these distributions is bounded as below. We provide a proof at the end.

Lemma 1 (**Perturbation in the mean vector**). *For any $\psi \in \mathbb{R}^{p_y}$ and $\Psi \in \mathbb{R}^{p_y \times p_y}$, let $\mu(\psi, \Psi) \in \mathbb{R}^{p_u}$ and $\mathbb{C}ov(\psi, \Psi) \in \mathbb{R}^{p_u \times p_u}$ denote the mean vector and the covariance matrix of $\mathbf{u}$ respectively with respect to $f_\mathbf{u}$ in (15). Then, for any $\widehat{\psi}, \psi^\star \in \mathbb{R}^{p_y}$ and $\widehat{\Psi}, \Psi^\star \in \mathbb{R}^{p_y \times p_y}$, there exists $t \in [0, 1]$ such that*

$$\|\mu(\widehat{\psi}, \widehat{\Psi}) - \mu(\psi^\star, \Psi^\star)\|_2 \leq \|\mathbb{C}ov(t\widehat{\psi} + (1-t)\psi^\star, t\widehat{\Psi} + (1-t)\Psi^\star)\|_{\mathrm{op}}\|\widehat{\psi} - \psi^\star\|_2.$$

Given this lemma, we proceed with the proof. By applying this lemma to $\widehat{f}^{(i)}_{\mathbf{y}|\mathbf{a}}(\boldsymbol{y}|\boldsymbol{a})$ in (12) and $f^{(i)}{}_{\mathbf{y}|\mathbf{a}}(\boldsymbol{y}|\boldsymbol{a})$ in (14), we see that it is sufficient to show the following bound

$$\|(\theta^{\star(i,y)} - \widehat{\theta}^{(i,y)}) + 2\boldsymbol{v}^{(i)\top}(\Phi^{\star(v,y)} - \widehat{\Phi}^{(v,y)}) + 2\widetilde{\boldsymbol{a}}^{(i)\top}(\Phi^{\star(a,y)} - \widehat{\Phi}^{(a,y)})\|_2 \leq R(\varepsilon, \delta/n) + p\varepsilon.$$

To that end, we have

$$\|(\theta^{\star(i,y)} - \widehat{\theta}^{(i,y)}) + 2\boldsymbol{v}^{(i)\top}(\Phi^{\star(v,y)} - \widehat{\Phi}^{(v,y)}) + 2\widetilde{\boldsymbol{a}}^{(i)\top}(\Phi^{\star(a,y)} - \widehat{\Phi}^{(a,y)})\|_2$$

$$\overset{(a)}{\leq} \|\theta^{\star(i,y)} - \widehat{\theta}^{(i,y)}\|_2 + 2\|\boldsymbol{v}^{(i)\top}(\Phi^{\star(v,y)} - \widehat{\Phi}^{(v,y)})\|_2 + 2\|\widetilde{\boldsymbol{a}}^{(i)\top}(\Phi^{\star(a,y)} - \widehat{\Phi}^{(a,y)})\|_2$$

$$\overset{(b)}{\leq} \|\theta^{\star(i,y)} - \widehat{\theta}^{(i,y)}\|_2 + 2\|\boldsymbol{v}^{(i)}\|_2\|\Phi^{\star(v,y)} - \widehat{\Phi}^{(v,y)}\|_{\mathrm{op}} + 2\|\widetilde{\boldsymbol{a}}^{(i)}\|_2\|(\Phi^{\star(a,y)} - \widehat{\Phi}^{(a,y)})\|_{\mathrm{op}}$$

$$\overset{(c)}{\leq} \|\theta^{\star(i)} - \widehat{\theta}^{(i)}\|_2 + 2\left(\|\boldsymbol{v}^{(i)}\|_2 + \|\widetilde{\boldsymbol{a}}^{(i)}\|_2\right)\|\Theta^\star - \widehat{\Theta}\|_{\mathrm{op}}$$

$$\overset{(d)}{\leq} \|\theta^{\star(i)} - \widehat{\theta}^{(i)}\|_2 + 2\left(\|\boldsymbol{v}^{(i)}\|_2 + \|\widetilde{\boldsymbol{a}}^{(i)}\|_2\right)\|\Theta^\star - \widehat{\Theta}\|_1 \overset{(e)}{\leq} R(\varepsilon, \delta/n) + 4x_{\max}\sqrt{p_y}\sqrt{p}\varepsilon,$$

where (a) follows from triangle inequality, (b) follows because induced matrix norms are submultiplicative, (c) follows because operator norm of any sub-matrix is no more than operator norm of the matrix and ℓ_2 norm of any sub-vector is no more than ℓ_2 norm of the vector, (d) follows because $\Theta^\star - \widehat{\Theta}$ is symmetric and because matrix operator norm is bounded by square root of the product of matrix one norm and matrix infinity norm, and (e) follows from Thm. 1 by using the relationship between vector norms. The proof is complete by rescaling ε and absorbing the constants in c.

Proof of Lem. 1: Perturbation in the mean vector. Let $Z(\psi, \Psi) \in \mathbb{R}_+$ denote the log-partition function of $f_{\mathbf{u}}(\cdot | \psi, \Psi)$ in (15). Then, from [11, Theorem 1], we have

$$\|\mu(\widehat{\psi}, \widehat{\Psi}) - \mu(\psi^\star, \Psi^\star)\|_2 = \|\nabla_{\widehat{\psi}} Z(\widehat{\psi}, \widehat{\Psi}) - \nabla_{\psi^\star} Z(\psi^\star, \Psi^\star)\|_2. \tag{16}$$

For some $t \in (0,1)$, $\widetilde{\psi} \triangleq t\widehat{\psi} + (1-t)\psi^\star$ and $\widetilde{\Psi} \triangleq t\widehat{\psi} + (1-t)\widetilde{\psi}$, we have the following from the mean value theorem

$$\|\nabla_{\widehat{\psi}} Z(\widehat{\psi}, \widehat{\Psi}) - \nabla_{\psi^\star} Z(\psi^\star, \Psi^\star)\|_2 = \|\nabla^2_{\widetilde{\psi}} Z(\widetilde{\psi}, \widetilde{\Psi}) \cdot (\widehat{\psi} - \psi^\star)\|_2 \overset{(a)}{\leq} \|\nabla^2_{\widetilde{\psi}} Z(\widetilde{\psi}, \widetilde{\Psi})\|_{\mathrm{op}} \|(\widehat{\psi} - \psi^\star)\|_2$$
$$\overset{(b)}{=} \|\mathbb{C}\mathrm{ov}(\widetilde{\psi}, \widetilde{\Psi})\|_{\mathrm{op}} \|(\widehat{\psi} - \psi^\star)\|_2, \tag{17}$$

where (a) follows because induced matrix norms are submultiplicative and (b) follows because for any regular exponential family a sub-matrix of the Hessian of the log partition function is the corresponding sub-matrix of the covariance matrix of the associated sufficient statistic. Combining (16) and (17) completes the proof. $\qquad\square$

B Proof of Thm. 1 Part I: Recovering population-level parameter

To analyze our estimate of the population-level parameter, we note that the Λ_Θ in (7) places independent constraints on the rows of Θ[5]. Therefore, we look at p independent convex optimization problems by decomposing the loss function $\mathcal{L}$ in (9) and the estimate $\widehat{\Theta}$ in (10) as follows:

$$\mathcal{L}_t(\underline{\Theta}_t) \triangleq \frac{1}{n} \sum_{i \in [n]} \exp\left(-[\theta_t^{(i)} + \Theta_t^\top x^{(i)}] x_t^{(i)}\right) \quad \text{and} \quad \widehat{\underline{\Theta}}_t \in \operatorname*{arg\,min}_{\underline{\Theta} \in \Lambda_\theta^n \times \Lambda_\Theta} \mathcal{L}_t(\underline{\Theta}_t) \quad \text{for all} \quad t \in [p], \tag{18}$$

where $\underline{\Theta}_t = \{\theta_t^{(1)}, \cdots, \theta_t^{(n)}, \Theta_t\}$ as defined in (9). Now, fix any $t \in [p]$, and consider any direction $\underline{\Omega}_t \triangleq \{\omega_t^{(1)}, \cdots, \omega_t^{(n)}, \Omega_t\} \in \mathbb{R}^{n+p}$ along the parameter $\underline{\Theta}_t$, i.e.,

$$\underline{\Omega}_t = \underline{\Theta}_t - \underline{\Theta}_t^\star, \quad \text{and} \quad \Omega_t = \Theta_t - \Theta_t^\star. \tag{19}$$

Without loss of generality, we let $\Omega_{tt} = 0$ since $\Theta_{tt}^\star = 0$. We denote the first-order and the second-order directional derivatives of the loss function $\mathcal{L}_t$ in (18) along the direction $\underline{\Omega}_t$ evaluated at $\underline{\Theta}_t$ by $\partial_{\underline{\Omega}_t} \mathcal{L}_t(\underline{\Theta}_t)$ and $\partial^2_{\underline{\Omega}_t^2} \mathcal{L}_t(\underline{\Theta}_t)$ respectively. Below, we state a lemma (with proof divided across App. B.1 and App. B.2) that provides us a control on $\partial_{\underline{\Omega}_t} \mathcal{L}_t(\underline{\Theta}_t)$ and $\partial^2_{\underline{\Omega}_t^2} \mathcal{L}_t(\underline{\Theta}_t)$. The assumptions of Thm. 1 remain in force. The lemma statement makes use of the following constants that depend on model-parameters $\tau \triangleq (\alpha, \beta, x_{\max})$:

$$C_{1,\tau} \triangleq \alpha(1 + 2\beta x_{\max}) \quad \text{and} \quad C_{3,\tau} \triangleq \exp\left(\alpha x_{\max}(1 + \beta x_{\max})\right). \tag{20}$$

Lemma 2 (Control on first and second directional derivatives). *For any fixed $\varepsilon_1, \varepsilon_2 > 0$, $\delta_1, \delta_2 \in (0,1)$, $\underline{\Theta} \in \Lambda_\theta^n \times \Lambda_\Theta$ defined in (9) with $\underline{\Omega}_t$ and Ω_t defined in (19), we have the following:*

(a) **Concentration of first directional derivative**: *with probability at least $1 - \delta_1$,*

$$\left|\partial_{\underline{\Omega}_t} \mathcal{L}_t(\underline{\Theta}_t^\star)\right| \leq \varepsilon_1 \quad \text{for} \quad n \geq \frac{8 C_{1,\tau}^2 C_{3,\tau}^2 x_{\max}^2 \cdot \log(2p/\delta_1)}{\varepsilon_1^2},$$

uniformly for all $t \in [p]$.

(b) **Anti-concentration of second directional derivative**: *with probability at least $1 - \delta_2$,*

$$\partial^2_{\underline{\Omega}_t^2} \mathcal{L}_t(\underline{\Theta}_t) \geq \frac{x_{\max}^2 \|\Omega_t\|_2^2}{\pi^2 e^2 (\beta + 1) C_{3,\tau}^5} - \varepsilon_2 \quad \text{for} \quad n \geq \frac{32 C_{1,\tau}^4 x_{\max}^4 \cdot \log(2p/\delta_2)}{\varepsilon_2^2 C_{3,\tau}^2},$$

uniformly for all $t \in [p]$.

---

[5]To ensure that the final estimate is symmetric, we can take the average of Θ and $\Theta^\top$.

Given this lemma, we now proceed with the proof. Consider the direction $\underline{\Omega}_t = \widehat{\underline{\Theta}}_t - \underline{\Theta}_t^\star$, and define a function $g : [0, 1] \to \mathbb{R}^{n+p}$

$$g(a) \triangleq \underline{\Theta}_t^\star + a(\widehat{\underline{\Theta}}_t - \underline{\Theta}_t^\star).$$

Notice that $g(0) = \underline{\Theta}_t^\star$ and $g(1) = \widehat{\underline{\Theta}}_t$ as well as

$$\frac{d\mathcal{L}_t(g(a))}{da} = \partial_{\underline{\Omega}_t} \mathcal{L}_t(\underline{\Theta}_t)\big|_{\underline{\Theta}_t = g(a)} \quad \text{and} \quad \frac{d^2\mathcal{L}_t(g(a))}{da^2} = \partial_{\underline{\Omega}_t^2}^2 \mathcal{L}_t(\underline{\Theta}_t)\big|_{\underline{\Theta}_t = g(a)}. \tag{21}$$

By the fundamental theorem of calculus, we have

$$\frac{d\mathcal{L}_t(g(a))}{da} \geq \frac{d\mathcal{L}_t(g(a))}{da}\Big|_{a=0} + a \min_{a \in (0,1)} \frac{d^2\mathcal{L}_t(g(a))}{da^2}. \tag{22}$$

Integrating both sides of (22) with respect to a, we obtain

$$\mathcal{L}_t(g(a)) - \mathcal{L}_t(g(0)) \geq a \frac{d\mathcal{L}_t(g(a))}{da}\Big|_{a=0} + \frac{a^2}{2} \min_{a \in (0,1)} \frac{d^2\mathcal{L}_t(g(a))}{da^2}$$

$$\stackrel{(21)}{=} a\partial_{\underline{\Omega}_t} \mathcal{L}_t(\underline{\Theta}_t)\big|_{\underline{\Theta}_t = g(0)} + \frac{a^2}{2} \min_{a \in (0,1)} \partial_{\underline{\Omega}_t^2}^2 \mathcal{L}_t(\underline{\Theta}_t)\big|_{\underline{\Theta}_t = g(a)}$$

$$\stackrel{(a)}{=} a\partial_{\underline{\Omega}_t} \mathcal{L}_t(\underline{\Theta}_t^\star) + \frac{a^2}{2} \min_{a \in (0,1)} \partial_{\underline{\Omega}_t^2}^2 \mathcal{L}_t(\underline{\Theta}_t)\big|_{\underline{\Theta}_t = g(a)}$$

$$\stackrel{(b)}{\geq} -a\big|\partial_{\underline{\Omega}_t} \mathcal{L}_t(\underline{\Theta}_t^\star)\big| + \frac{a^2}{2} \min_{a \in (0,1)} \partial_{\underline{\Omega}_t^2}^2 \mathcal{L}_t(\underline{\Theta}_t)\big|_{\underline{\Theta}_t = g(a)}, \tag{23}$$

where (a) follows because $g(0) = \underline{\Theta}_t^\star$ and (b) follows by the triangle inequality. Plugging in $a = 1$ in (23), and noting that by the optimality of $\widehat{\underline{\Theta}}_t$ in (18), we have $\mathcal{L}_t(\underline{\Theta}_t^\star) \geq \mathcal{L}_t(\widehat{\underline{\Theta}}_t)$, we find that

$$0 \geq \mathcal{L}_t(\widehat{\underline{\Theta}}_t) - \mathcal{L}_t(\underline{\Theta}_t^\star) \geq -\big|\partial_{\underline{\Omega}_t} \mathcal{L}_t(\underline{\Theta}_t^\star)\big| + \frac{1}{2} \min_{a \in (0,1)} \partial_{\underline{\Omega}_t^2}^2 \mathcal{L}_t(\underline{\Theta}_t)\big|_{\underline{\Theta}_t = g(a)}.$$

Now, using Lem. 2 with $\delta_1 = \delta/2$ and $\delta_2 = \delta/2$, and because $\Omega_t = \widehat{\Theta}_t - \Theta_t^\star$, we have

$$0 \geq -\varepsilon_1 + \frac{1}{2}\left(\frac{x_{\max}^2 \|\widehat{\Theta}_t - \Theta_t^\star\|_2^2}{\pi^2 e^2 (\beta + 1) C_{3,\tau}^5} - \varepsilon_2 \right), \tag{24}$$

uniformly for all $t \in [p]$ with probability at least $1 - \delta$ as long as

$$n \geq \max\left(\frac{8C_{1,\tau}^2 C_{3,\tau}^2 x_{\max}^2 \cdot \log(4p/\delta)}{\varepsilon_1^2}, \frac{32C_{1,\tau}^4 x_{\max}^4 \cdot \log(4p/\delta)}{\varepsilon_2^2 C_{3,\tau}^2} \right).$$

Choosing

$$\varepsilon_1 = \frac{\varepsilon^2 x_{\max}^2}{4\pi^2 e^2 (\beta + 1) C_{3,\tau}^5} \quad \text{and} \quad \varepsilon_2 = \frac{\varepsilon^2 x_{\max}^2}{2\pi^2 e^2 (\beta + 1) C_{3,\tau}^5},$$

and rearranging (24), we have $\|\widehat{\Theta}_t - \Theta_t^\star\|_2 \leq \varepsilon$ with probability at least $1 - \delta$ as long as

$$n \geq \frac{128\pi^4 e^4 (\beta + 1)^2 C_{1,\tau}^2 C_{3,\tau}^{12} \cdot \log(4p/\delta)}{\varepsilon^4 x_{\max}^2}.$$

B.1 Proof of Lem. 2 (a): Concentration of first directional derivative

For every $t \in [p]$ with $\underline{\Omega}_t$ defined in (19), we claim that the first-order directional derivative of the loss function defined in (18) is given by

$$\partial_{\underline{\Omega}_t} \mathcal{L}_t(\underline{\Theta}_t) = -\frac{1}{n} \sum_{i \in [n]} \left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)} \right) \exp\left(-[\theta_t^{(i)} + \Theta_t^\top \boldsymbol{x}^{(i)}] x_t^{(i)} \right), \tag{25}$$

where $\Delta_t^{(i)} \triangleq \begin{bmatrix} \omega_t^{(i)} \\ \Omega_t \end{bmatrix} \in \mathbb{R}^{p+1}$ and $\widetilde{\boldsymbol{x}}^{(i)} \triangleq \begin{bmatrix} 1 \\ \boldsymbol{x}^{(i)} \end{bmatrix} \in \mathbb{R}^{p+1}$ for all $i \in [n]$. We provide a proof in App. B.1.1.

Next, we claim that the mean of the first-order directional derivative evaluated at the true parameter is zero. We provide a proof in App. B.1.2

Lemma 3 (**Zero-meanness of first directional derivative**). *For every $t \in [p]$ with $\underline{\Omega}_t$ defined in (19), we have $\mathbb{E}\big[\partial_{\underline{\Omega}_t}\mathcal{L}_t(\underline{\Theta}_t^\star)\big] = 0$.*

Given these, we proceed to show the concentration of the first-order directional derivative evaluated at the true parameter. Fix any $t \in [p]$. From (25), we have

$$\partial_{\underline{\Omega}_t}\mathcal{L}_t(\underline{\Theta}_t^\star) \stackrel{(25)}{=} -\frac{1}{n}\sum_{i\in[n]}\left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)}\right)\exp\left(-[\theta_t^{\star(i)} + \Theta_t^{\star\top}\boldsymbol{x}^{(i)}]x_t^{(i)}\right).$$

Each term in the above summation is an independent random variable and is bounded as follows:

$$\left|\left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)}\right)\times\exp\left(-[\theta_t^{\star(i)} + \Theta_t^{\star\top}\boldsymbol{x}^{(i)}]x_t^{(i)}\right)\right|$$

$$\stackrel{(a)}{=}\left|\left(\omega_t^{(i)}x_t^{(i)} + \Omega_t^\top\boldsymbol{x}^{(i)}x_t^{(i)}\right)\times\exp\left(-[\theta_t^{\star(i)} + \Theta_t^{\star\top}\boldsymbol{x}^{(i)}]x_t^{(i)}\right)\right|$$

$$\stackrel{(b)}{\leq}\left|\omega_t^{(i)} + \Omega_t^\top\boldsymbol{x}^{(i)}\right|\times x_{\max}\times\exp\left(|\theta_t^{\star(i)} + \Theta_t^{\star\top}\boldsymbol{x}^{(i)}|x_{\max}\right)$$

$$\stackrel{(c)}{\leq}\left||\omega_t^{(i)}| + \|\Omega_t\|_1\|\boldsymbol{x}^{(i)}\|_\infty\right|\times x_{\max}\times\exp\left(\left(|\theta_t^{\star(i)}| + \|\Theta_t^\star\|_1\|\boldsymbol{x}^{(i)}\|_\infty\right)x_{\max}\right)$$

$$\stackrel{(d)}{\leq}\left(2\alpha + 4\alpha\beta x_{\max}\right)\times x_{\max}\times\exp\left((\alpha + \alpha\beta x_{\max})x_{\max}\right)\stackrel{(20)}{=}2C_{1,\tau}C_{3,\tau}x_{\max},$$

where (a) follows by plugging in $\Delta_t^{(i)}$ and $\widetilde{\boldsymbol{x}}^{(i)}$, (b) follows because $\|\boldsymbol{x}^{(i)}\|_\infty \leq x_{\max}$ for all $i \in [n]$, (c) follows by triangle inequality and Cauchy–Schwarz inequality, and (d) follows because $\theta^{\star(i)} \in \Lambda_\theta$ for all $i \in [n]$, $\Theta^\star \in \Lambda_\Theta$, $\omega^{(i)} \in 2\Lambda_\theta$ for all $i \in [n]$, $\Omega \in 2\Lambda_\Theta$, and $\|\boldsymbol{x}^{(i)}\|_\infty \leq x_{\max}$ for all $i \in [n]$.

Further, from Lem. 3, we have $\mathbb{E}\big[\partial_{\underline{\Omega}_t}\mathcal{L}_t(\underline{\Theta}_t^\star)\big] = 0$. Therefore, using the Hoeffding's inequality results in

$$\mathbb{P}\left(\left|\partial_{\underline{\Omega}_t}\mathcal{L}_t(\underline{\Theta}_t^\star)\right| > \varepsilon_1\right) < 2\exp\left(-\frac{n\varepsilon_1^2}{8C_{1,\tau}^2C_{3,\tau}^2x_{\max}^2}\right).$$

The proof follows by using the union bound over all $t \in [p]$.

B.1.1 Proof of (25): Expression for first directional derivative

Fix any $t \in [p]$. The first-order partial derivatives of $\mathcal{L}_t$ with respect to entries of $\underline{\Theta}_t$ defined in (9) are given by

$$\frac{\partial\mathcal{L}_t(\underline{\Theta}_t)}{\partial\theta_t^{(i)}} = -\frac{1}{n}x_t^{(i)}\exp\left(-[\theta_t^{(i)} + \Theta_t^\top\boldsymbol{x}^{(i)}]x_t^{(i)}\right)\quad\text{for all}\quad i\in[n],\quad\text{and}$$

$$\frac{\partial\mathcal{L}_t(\underline{\Theta}_t)}{\partial\Theta_{tu}} = -\frac{1}{n}\sum_{i\in[n]}x_t^{(i)}x_u^{(i)}\exp\left(-[\theta_t^{(i)} + \Theta_t^\top\boldsymbol{x}^{(i)}]x_t^{(i)}\right)\quad\text{for all}\quad u\in[p]\setminus\{t\}\,.$$

Now, we can write the first-order directional derivative of $\mathcal{L}_t$ as

$$\partial_{\underline{\Omega}_t}\mathcal{L}_t(\underline{\Theta}_t) \triangleq \lim_{h\to 0}\frac{\mathcal{L}_t(\underline{\Theta}_t + h\underline{\Omega}_t) - \mathcal{L}_t(\underline{\Theta}_t)}{h} = \sum_{i\in[n]}\omega_t^{(i)}\frac{\partial\mathcal{L}_t(\underline{\Theta}_t)}{\partial\theta_t^{(i)}} + \sum_{u\in[p]\setminus\{t\}}\Omega_{tu}\frac{\partial\mathcal{L}_t(\underline{\Theta}_t)}{\partial\Theta_{tu}}$$

$$= -\frac{1}{n}\sum_{i\in[n]}\left(\omega_t^{(i)}x_t^{(i)} + \sum_{u\in[p]\setminus\{t\}}\Omega_{tu}x_t^{(i)}x_u^{(i)}\right)\exp\left(-[\theta_t^{(i)} + \Theta_t^\top\boldsymbol{x}^{(i)}]x_t^{(i)}\right)$$

$$= -\frac{1}{n}\sum_{i\in[n]}\left(\omega_t^{(i)}x_t^{(i)} + \Omega_t^\top\boldsymbol{x}^{(i)}x_t^{(i)}\right)\exp\left(-[\theta_t^{(i)} + \Theta_t^\top\boldsymbol{x}^{(i)}]x_t^{(i)}\right)$$

$$\stackrel{(a)}{=} -\frac{1}{n}\sum_{i\in[n]}\left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)}\right)\exp\left(-[\theta_t^{(i)} + \Theta_t^\top\boldsymbol{x}^{(i)}]x_t^{(i)}\right),$$

where (a) follows from the definitions of $\Delta_t^{(i)}$ and $\widetilde{\boldsymbol{x}}^{(i)}$.

B.1.2 Proof of Lem. 3: Zero-meanness of first directional derivative.

Fix any $t \in [p]$. From (25), we have

$$
\mathbb{E}\big[\partial_{\underline{\Omega}_t} \mathcal{L}_t(\Theta_t^\star)\big] \overset{(25)}{=} -\frac{1}{n} \sum_{i \in [n]} \mathbb{E}\bigg[\Big([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)}\Big) \exp\Big(-[\theta_t^{\star(i)} + \Theta_t^{\star\top} \boldsymbol{x}^{(i)}]x_t^{(i)}\Big)\bigg]
$$

$$
\overset{(a)}{=} -\frac{1}{n} \sum_{i \in [n]} \sum_{u \in [p+1]} \Delta_{tu}^{(i)} \mathbb{E}_{\boldsymbol{x}^{(i)}, \boldsymbol{z}^{(i)}}\bigg[\widetilde{\boldsymbol{x}}_u^{(i)} x_t^{(i)} \exp\Big(-[\theta_t^\star(\boldsymbol{z}^{(i)}) + \Theta_t^{\star\top} \boldsymbol{x}^{(i)}]x_t^{(i)}\Big)\bigg]
$$

$$
\overset{(b)}{=} -\frac{1}{n} \sum_{i \in [n]} \sum_{u \in [p+1] \setminus \{i+1\}} \Delta_{tu}^{(i)} \mathbb{E}_{\boldsymbol{x}^{(i)}, \boldsymbol{z}^{(i)}}\bigg[\widetilde{\boldsymbol{x}}_u^{(i)} x_t^{(i)} \exp\Big(-[\theta_t^\star(\boldsymbol{z}^{(i)}) + \Theta_t^{\star\top} \boldsymbol{x}^{(i)}]x_t^{(i)}\Big)\bigg],
$$

where (a) follows by linearity of expectation and by plugging in $\theta_t^{\star(i)} = \theta_t^\star(\boldsymbol{z}^{(i)})$ and (b) follows because $\Delta_{tu}^{(i)} = \Omega_{tt} = 0$ for $u = t+1$ for every $i \in [n]$. To complete the proof, we show

$$
\mathbb{E}_{\boldsymbol{x}^{(i)}, \boldsymbol{z}^{(i)}}\bigg[\widetilde{\boldsymbol{x}}_u^{(i)} x_t^{(i)} \exp\Big(-[\theta_t^\star(\boldsymbol{z}^{(i)}) + \Theta_t^{\star\top} \boldsymbol{x}^{(i)}]x_t^{(i)}\Big)\bigg] = 0.
$$

for all $t \in [p], i \in [n], u \in [p+1] \setminus \{t+1\}$. To that end, fix any $t \in [p], i \in [n], u \in [p+1] \setminus \{t+1\}$. We have

$$
\mathbb{E}_{\boldsymbol{x}^{(i)}, \boldsymbol{z}^{(i)}}\bigg[\widetilde{\boldsymbol{x}}_u^{(i)} x_t^{(i)} \exp\Big(-[\theta_t^\star(\boldsymbol{z}^{(i)}) + \Theta_t^{\star\top} \boldsymbol{x}^{(i)}]x_t^{(i)}\Big)\bigg]
$$

$$
= \int_{\mathcal{X}^p \times \mathcal{Z}^{p_z}} \widetilde{x}_u^{(i)} x_t^{(i)} \exp\Big(-[\theta_t^\star(\boldsymbol{z}^{(i)}) + \Theta_t^{\star\top} \boldsymbol{x}^{(i)}]x_t^{(i)}\Big) f_{\mathsf{x},\mathsf{z}}\big(\boldsymbol{x}^{(i)}, \boldsymbol{z}^{(i)}\big) d\boldsymbol{x}^{(i)} d\boldsymbol{z}^{(i)}
$$

$$
= \int_{\mathcal{X}^p \times \mathcal{Z}^{p_z}} \widetilde{x}_u^{(i)} x_t^{(i)} \exp\Big(-[\theta_t^\star(\boldsymbol{z}^{(i)}) + \Theta_t^{\star\top} \boldsymbol{x}^{(i)}]x_t^{(i)}\Big) f_{\mathsf{x}_{-t},\mathsf{z}}\big(\boldsymbol{x}_{-t}^{(i)}, \boldsymbol{z}^{(i)}\big) \times
$$

$$
f_{\mathsf{x}_t | \mathsf{x}_{-t}, \mathsf{z}}\big(x_t^{(i)} | \boldsymbol{x}_{-t}^{(i)}, \boldsymbol{z}^{(i)}; \theta_t^\star(\boldsymbol{z}^{(i)}), \Theta_t^\star\big) d\boldsymbol{x}^{(i)} d\boldsymbol{z}^{(i)}
$$

$$
\overset{(a)}{=} \int_{\mathcal{X}^p \times \mathcal{Z}^{p_z}} \frac{\widetilde{x}_u^{(i)} x_t^{(i)} f_{\mathsf{x}_{-t},\mathsf{z}}\big(\boldsymbol{x}_{-t}^{(i)}, \boldsymbol{z}^{(i)}\big) d\boldsymbol{x}^{(i)} d\boldsymbol{z}^{(i)}}{\int_{\mathcal{X}} \exp\Big([\theta_t^\star(\boldsymbol{z}^{(i)}) + \Theta_t^{\star\top} \boldsymbol{x}^{(i)}]x_t^{(i)}\Big) dx_t^{(i)}}
$$

$$
\overset{(b)}{=} \int_{\mathcal{X}^{p-1} \times \mathcal{Z}^{p_z}} \bigg[\int_{\mathcal{X}} x_t^{(i)} dx_t^{(i)}\bigg] \frac{\widetilde{x}_u^{(i)} f_{\mathsf{x}_{-t},\mathsf{z}}\big(\boldsymbol{x}_{-t}^{(i)}, \boldsymbol{z}^{(i)}\big) d\boldsymbol{x}_{-t}^{(i)} d\boldsymbol{z}^{(i)}}{\int_{\mathcal{X}} \exp\Big([\theta_t^\star(\boldsymbol{z}^{(i)}) + \Theta_t^{\star\top} \boldsymbol{x}^{(i)}]x_t^{(i)}\Big) dx_t^{(i)}} \overset{(c)}{=} 0,
$$

where (a) follows by plugging in $f_{\mathsf{x}_t | \mathsf{x}_{-t}, \mathsf{z}}\big(x_t^{(i)} | \boldsymbol{x}_{-t}^{(i)}, \boldsymbol{z}^{(i)}; \theta_t^\star(\boldsymbol{z}^{(i)}), \Theta_t^\star\big)$ from (8), (b) follows by re-arranging and observing that $\widetilde{x}_u^{(i)} \neq x_t^{(i)}$ for any $u \in [p+1] \setminus \{t+1\}$, and (c) follows because $\int_{\mathcal{X}} x_t^{(i)} dx_t^{(i)} = 0$ when $\mathcal{X}$ is symmetric around 0.

B.2 Proof of Lem. 2 (b): Anti-concentration of second directional derivative

We start by claiming that the second-order directional derivative can be lower bounded by a quadratic form. We provide a proof in App. B.2.1.

Lemma 4 (**Lower bound on the second directional derivative**). *For every $t \in [p]$ with $\underline{\Omega}_t$ defined in (19), we have*

$$
\partial_{\underline{\Omega}_t^2}^2 \mathcal{L}_t(\Theta_t) \geq \frac{1}{nC_{3,\tau}} \sum_{i \in [n]} \Big([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)}\Big)^2,
$$

where $\Delta_t^{(i)} \triangleq \begin{bmatrix} \omega_t^{(i)} \\ \Omega_t \end{bmatrix} \in \mathbb{R}^{p+1}$ and $\widetilde{\boldsymbol{x}}^{(i)} \triangleq \begin{bmatrix} 1 \\ \boldsymbol{x}^{(i)} \end{bmatrix} \in \mathbb{R}^{p+1}$ for all $i \in [n]$, and the constant $C_{3,\tau}$ was defined in (20).

Next, we claim that the conditional variance of $x_t^{(i)}$ conditioned on $\mathsf{x}_{-t} = \boldsymbol{x}_{-t}^{(i)}$ and $\mathsf{z} = \boldsymbol{z}^{(i)}$ is lower bounded by a constant for every $t \in [p]$ and $i \in [n]$. We provide a proof in App. B.2.2.

Lemma 5 (Lower bound on the conditional variance). *For every $t \in [p]$ and $i \in [n]$, we have*

$$\mathbb{V}ar\big(x_t^{(i)} \big| \boldsymbol{x}_{-t}^{(i)}, \boldsymbol{z}^{(i)}\big) \geq \frac{x_{\max}}{\pi e C_{3,\tau}^2},$$

where the constant $C_{3,\tau}$ was defined in (20).

Given these, we proceed to show the anti-concentration of the second-order directional derivative. Fix any $t \in [p]$ and any $\underline{\Theta} \in \Lambda_\theta^n \times \Lambda_\Theta$. From Lem. 4, we have

$$\partial_{\underline{\Omega}_t^2}^2 \mathcal{L}_t(\underline{\Theta}_t) \geq \frac{1}{nC_{3,\tau}} \sum_{i \in [n]} \left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)} \right)^2. \tag{26}$$

First, using the Hoeffding's inequality, let us show concentration of $\frac{1}{n}\sum_{i \in [n]} \left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)}\right)^2$ around its mean. We observe that each term in the summation is an independent random variable and is bounded as follows

$$\begin{aligned}
\left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)}\right)^2 &\stackrel{(a)}{=} \left(\omega_t^{(i)} x_t^{(i)} + \Omega_t^\top \boldsymbol{x}^{(i)} x_t^{(i)}\right)^2 \stackrel{(b)}{\leq} \left(\omega_t^{(i)} + \Omega_t^\top \boldsymbol{x}^{(i)}\right)^2 x_{\max}^2 \\
&\stackrel{(c)}{\leq} \left(|\omega_t^{(i)}| + \|\Omega_t\|_1 \|\boldsymbol{x}^{(i)}\|_\infty\right)^2 x_{\max}^2 \\
&\stackrel{(d)}{\leq} \left(2\alpha + 4\alpha\beta x_{\max}\right)^2 x_{\max}^2 \stackrel{(20)}{=} 4C_{1,\tau}^2 x_{\max}^2,
\end{aligned}$$

where (a) follows by plugging in $\Delta_t^{(i)}$ and $\widetilde{\boldsymbol{x}}^{(i)}$, (b) follows because $\|\boldsymbol{x}^{(i)}\|_\infty \leq x_{\max}$ for all $i \in [n]$, (c) follows by triangle inequality and Cauchy–Schwarz inequality, and (d) follows because $\Omega \in 2\Lambda_\Theta$, $\omega^{(i)} \in 2\Lambda_\theta$, and $\|\boldsymbol{x}^{(i)}\|_\infty \leq x_{\max}$ for all $i \in [n]$. Then, from the Hoeffding's inequality, we have

$$\mathbb{P}\left(\left|\frac{1}{n}\sum_{i \in [n]}\left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)}\right)^2 - \frac{1}{n}\sum_{i \in [n]}\mathbb{E}\left[\left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)}\right)^2\right]\right| > \varepsilon\right) < 2\exp\left(-\frac{n\varepsilon^2}{32C_{1,\tau}^4 x_{\max}^4}\right).$$

Applying the union bound over all $t \in [p]$, for any $\delta \in (0,1)$ and uniformly for all $t \in [p]$, we have

$$\frac{1}{n}\sum_{i \in [n]}\left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)}\right)^2 \geq \frac{1}{n}\sum_{i \in [n]}\mathbb{E}\left[\left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)}\right)^2\right] - \varepsilon, \tag{27}$$

with probability at least $1 - \delta$ as long as

$$n \geq \frac{32C_{1,\tau}^4 x_{\max}^4}{\varepsilon^2}\log\left(\frac{2p}{\delta}\right).$$

Now, we lower bound $\mathbb{E}\left[\left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)}\right)^2\right]$ for every $t \in [p]$ and every $i \in [n]$. Fix any $t \in [p]$ and $i \in [n]$. We have

$$\mathbb{E}\left[\left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)}\right)^2\right] \stackrel{(a)}{\geq} \mathbb{V}ar\left[[\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)}\right] = \mathbb{V}ar\left[\omega_t^{(i)} x_t^{(i)} + \Omega_t^\top \boldsymbol{x}^{(i)} x_t^{(i)}\right], \tag{28}$$

where (a) follows from the fact that for any random variable a, $\mathbb{E}[a^2] \geq \mathbb{V}ar[a]$. We define the following set to lower bound $\mathbb{V}ar\left[\omega_t^{(i)} x_t^{(i)} + \Omega_t^\top \boldsymbol{x}^{(i)} x_t^{(i)}\right]$:

$$\mathcal{E}(\Theta^\star) \triangleq \left\{(t,u) \in [p]^2 : t < u, \Theta_{tu}^\star \neq 0\right\}, \tag{29}$$

and consider the graph $\mathcal{G}(\Theta^\star) = ([p], \mathcal{E}(\Theta^\star))$ with $[p]$ as nodes and $\mathcal{E}(\Theta^\star)$ as edges such that $f_{\boldsymbol{x}|\boldsymbol{z}}\big(\boldsymbol{x}|\boldsymbol{z}; \theta^\star(\boldsymbol{z}), \Theta^\star\big)$ is Markov with respect to $\mathcal{G}(\Theta^\star)$. We claim that there exists a non-empty set $\mathcal{R}_t \subset [p] \setminus \{t\}$ such that

    (i) $\mathcal{R}_t$ is an independent set of $\mathcal{G}(\Theta^\star)$, i.e., there are no edges between any pair of nodes in $\mathcal{R}_t$, and

    (ii) the row vector Ω_t satisfies $\sum_{u \in \mathcal{R}_t} |\Omega_{tu}|^2 \geq \frac{1}{\beta+1}\|\Omega_t\|_2^2$.

Taking this claim as given at the moment, we continue our proof. Denoting $\mathcal{R}_t^c \triangleq [p] \setminus \mathcal{R}_t$, and using the law of total variance, the variance term in (28) can be lower bounded as

$$
\begin{aligned}
\mathbb{V}\mathrm{ar}\Big[\omega_t^{(i)} x_t^{(i)} + \Omega_t^\top \boldsymbol{x}^{(i)} x_t^{(i)}\Big] &\geq \mathbb{E}\Big[\mathbb{V}\mathrm{ar}\Big[\omega_t^{(i)} x_t^{(i)} + \Omega_t^\top \boldsymbol{x}^{(i)} x_t^{(i)} \Big| \boldsymbol{x}_{\mathcal{R}_t^c}^{(i)}, \boldsymbol{z}^{(i)}\Big]\Big] \\
&\overset{(a)}{=} \mathbb{E}\Big[(x_t^{(i)})^2 \mathbb{V}\mathrm{ar}\Big(\sum_{u \in \mathcal{R}_t} \Omega_{tu} x_u^{(i)} \Big| \boldsymbol{x}_{\mathcal{R}_t^c}^{(i)}, \boldsymbol{z}^{(i)}\Big)\Big] \\
&\overset{(b)}{=} \mathbb{E}\Big[(x_t^{(i)})^2 \sum_{u \in \mathcal{R}_t} \Omega_{tu}^2 \mathbb{V}\mathrm{ar}\Big(x_u^{(i)} \Big| \boldsymbol{x}_{\mathcal{R}_t^c}^{(i)}, \boldsymbol{z}^{(i)}\Big)\Big] \\
&\overset{(c)}{=} \mathbb{E}\Big[(x_t^{(i)})^2 \sum_{u \in \mathcal{R}_t} \Omega_{tu}^2 \mathbb{V}\mathrm{ar}\Big(x_u^{(i)} \Big| \boldsymbol{x}_{-u}^{(i)}, \boldsymbol{z}^{(i)}\Big)\Big] \\
&\overset{(d)}{\geq} \frac{x_{\max}}{\pi e C_{3,\tau}^2} \sum_{u \in \mathcal{R}_t} \Omega_{tu}^2 \mathbb{E}\Big[(x_t^{(i)})^2\Big] \\
&\overset{(e)}{\geq} \frac{x_{\max}}{\pi e C_{3,\tau}^2} \sum_{u \in \mathcal{R}_t} \Omega_{tu}^2 \mathbb{V}\mathrm{ar}\Big(x_t^{(i)} \Big| \boldsymbol{x}_{-t}^{(i)}, \boldsymbol{z}^{(i)}\Big) \\
&\overset{(f)}{\geq} \frac{x_{\max}^2}{\pi^2 e^2 C_{3,\tau}^4} \sum_{u \in \mathcal{R}_t} \Omega_{tu}^2 \overset{(ii)}{\geq} \frac{x_{\max}^2 \|\Omega_t\|_2^2}{\pi^2 e^2 (\beta+1) C_{3,\tau}^4},
\end{aligned} \tag{30}
$$

where (a) follows because $(x_u^{(i)})_{u \in \mathcal{R}_t^c}$ are deterministic when conditioned on themselves, and $t \in \mathcal{R}_t^c$, (b) follows because $(x_u^{(i)})_{u \in \mathcal{R}_t}$ are conditionally independent given $\boldsymbol{x}_{\mathcal{R}_t^c}^{(i)}$ and $\boldsymbol{z}^{(i)}$ which is a direct consequence of (i), (c) follows because of the local Markov property (as the conditioning set includes all the neighbors in $\mathcal{G}(\Theta^\star)$ of each node in $\mathcal{R}_t$), (d) and (f) follow from Lem. 5, and (e) follows because $\mathbb{E}\Big[(x_t^{(i)})^2\Big] = \mathbb{E}\Big[\mathbb{E}\Big[(x_t^{(i)})^2 \Big| \boldsymbol{x}_{-t}^{(i)}, \boldsymbol{z}^{(i)}\Big]\Big] \geq \mathbb{V}\mathrm{ar}\Big(x_t^{(i)} \Big| \boldsymbol{x}_{-t}^{(i)}, \boldsymbol{z}^{(i)}\Big)$.

Combining (26) to (28) and (30), for any $\delta \in (0,1)$ and uniformly for all $t \in [p]$, we have

$$
\partial_{\underline{\Omega}_t^2}^2 \mathcal{L}_t(\underline{\Theta}_t) \geq \frac{1}{C_{3,\tau}} \left(\frac{x_{\max}^2 \|\Omega_t\|_2^2}{\pi^2 e^2 (\beta+1) C_{3,\tau}^4} - \varepsilon \right),
$$

with probability at least $1 - \delta$ as long as

$$
n \geq \frac{32 C_{1,\tau}^4 x_{\max}^4}{\varepsilon^2} \log\left(\frac{2p}{\delta}\right).
$$

Choosing $\varepsilon_2 = \varepsilon/C_{3,\tau}$ and $\delta_2 = \delta$ yields the claim.

It remains to construct the set $\mathcal{R}_t$ that is an independent set of $\mathcal{G}(\Theta^\star)$ and satisifies (ii).

Construction of the set $\mathcal{R}_t$: We select $r_1 \in [p] \setminus \{t\}$ such that

$$
|\Omega_{tr_1}| \geq |\Omega_{tu}| \quad \text{for all} \quad u \in [p] \setminus \{t, r_1\}.
$$

Next, we identify $r_2 \in [p] \setminus \{t, r_1, \mathcal{N}(r_1)\}$ such that

$$
|\Omega_{tr_2}| \geq |\Omega_{tu}| \quad \text{for all} \quad u \in [p] \setminus \{t, r_1, \mathcal{N}(r_1), r_2\}.
$$

We continue identifying $r_3, \ldots, r_s$ in such a manner till no more nodes are left, where s denotes the total number of nodes selected. Now we define $\mathcal{R}_t \triangleq \{r_1, \cdots, r_s\}$. Further, for any $u \in [p]$, let $\mathcal{N}(u)$ denote the set of neighbors of u in $\mathcal{G}(\Theta^\star)$. We have $|\mathcal{N}(u)| \leq \|\Theta_u^\star\|_0 \leq \beta$ from (29) and Assum. 1(b). We can now see that $\mathcal{R}_t$ is an independent set of $\mathcal{G}(\Theta^\star)$ as claimed in (i) such that it satisfies (ii) by construction.

 B.2.1   Proof of Lem. 4: Lower bound on the second directional derivative

503 For every $t \in [p]$ with $\underline{\Omega}_t$ defined in (19), we claim that the second-order directional derivative of the
504 loss function defined in (18) is given by

$$\partial_{\underline{\Omega}_t^2}^2 \mathcal{L}_t(\underline{\Theta}_t) = \frac{1}{n} \sum_{i \in [n]} \left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)} \right)^2 \exp\left(- [\theta_t^{(i)} + \Theta_t^\top \boldsymbol{x}^{(i)}] x_t^{(i)} \right), \qquad (31)$$

505 where $\Delta_t^{(i)} \triangleq \begin{bmatrix} \omega_t^{(i)} \\ \Omega_t \end{bmatrix} \in \mathbb{R}^{p+1}$ and $\widetilde{\boldsymbol{x}}^{(i)} \triangleq \begin{bmatrix} 1 \\ \boldsymbol{x}^{(i)} \end{bmatrix} \in \mathbb{R}^{p+1}$ for all $i \in [n]$. We provide a proof at the
506 end.

507 Given this, we proceed to prove the lower bound on the second directional derivative. Fix any $t \in [p]$.
508 From (31), we have

$$
\begin{aligned}
\partial_{\underline{\Omega}_t^2}^2 \mathcal{L}_t(\underline{\Theta}_t) &= \frac{1}{n} \sum_{i \in [n]} \left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)} \right)^2 \times \exp\left(- [\theta_t^{(i)} + \Theta_t^\top \boldsymbol{x}^{(i)}] x_t^{(i)} \right) \\
&\overset{(a)}{\geq} \frac{1}{n} \sum_{i \in [n]} \left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)} \right)^2 \times \exp\left(- |\theta_t^{(i)} + \Theta_t^\top \boldsymbol{x}^{(i)}| x_{\max} \right) \\
&\overset{(b)}{\geq} \frac{1}{n} \sum_{i \in [n]} \left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)} \right)^2 \times \exp\left(- \left(|\theta_t^{(i)}| + \|\Theta_t\|_1 \|\boldsymbol{x}^{(i)}\|_\infty \right) x_{\max} \right) \\
&\overset{(c)}{\geq} \frac{1}{n} \sum_{i \in [n]} \left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)} \right)^2 \times \exp\left(- (\alpha + \alpha\beta x_{\max}) x_{\max} \right) \\
&\overset{(20)}{=} \frac{1}{C_{3,\tau} n} \sum_{i \in [n]} \left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)} \right)^2,
\end{aligned}
$$

509 where (a) follows because $\|\boldsymbol{x}^{(i)}\|_\infty \leq x_{\max}$ for all $i \in [n]$, (b) follows by triangle inequality and
510 Cauchy–Schwarz inequality, and (c) follows because $\theta^{(i)} \in \Lambda_\theta$ for all $i \in [n]$, $\Theta \in \Lambda_\Theta$, and
511 $\|\boldsymbol{x}^{(i)}\|_\infty \leq x_{\max}$ for all $i \in [n]$.

512 **Proof of (31): Expression for second directional derivative.**   Fix any $t \in [p]$. The second-order
513 partial derivatives of $\mathcal{L}_t$ with respect to entries of $\underline{\Theta}_t$ defined in (9) are given by

$$
\frac{\partial^2 \mathcal{L}_t(\underline{\Theta}_t)}{\partial [\theta_t^{(i)}]^2} = \frac{1}{n} [x_t^{(i)}]^2 \exp\left(- [\theta_t^{(i)} + \Theta_t^\top \boldsymbol{x}^{(i)}] x_t^{(i)} \right) \text{ for all } i \in [n],
$$

$$
\frac{\partial^2 \mathcal{L}_t(\underline{\Theta}_t)}{\partial \Theta_{tu} \Theta_{tv}} = \frac{1}{n} \sum_{i \in [n]} [x_t^{(i)}]^2 x_u^{(i)} x_v^{(i)} \exp\left(- [\theta_t^{(i)} + \Theta_t^\top \boldsymbol{x}^{(i)}] x_t^{(i)} \right) \text{ for all } u, v \in [p] \setminus \{i\}, \text{ and}
$$

$$
\frac{\partial^2 \mathcal{L}_t(\underline{\Theta}_t)}{\partial \Theta_{tu} \theta_t^{(i)}} = \frac{\partial^2 \mathcal{L}_t(\underline{\Theta}_t)}{\partial \theta_t^{(i)} \Theta_{tu}} = \frac{1}{n} [x_t^{(i)}]^2 x_u^{(i)} \exp\left(- [\theta_t^{(i)} + \Theta_t^\top \boldsymbol{x}^{(i)}] x_t^{(i)} \right) \text{ for all } i \in [n], u \in [p] \setminus \{t\}.
$$

514 Now, we can write the second-order directional derivative of $\mathcal{L}_t$ as

$$
\begin{aligned}
&\partial_{\underline{\Omega}_t^2}^2 \mathcal{L}_t(\underline{\Theta}_t) \\
&\triangleq \lim_{h \to 0} \frac{\partial_{\underline{\Omega}_t} \mathcal{L}_t(\underline{\Theta}_t + h\underline{\Omega}_t) - \partial_{\underline{\Omega}_t} \mathcal{L}_t(\underline{\Theta}_t)}{h} \\
&= \sum_{i \in [n]} [\omega_t^{(i)}]^2 \frac{\partial^2 \mathcal{L}_t(\underline{\Theta}_t)}{\partial [\theta_t^{(i)}]^2} + \sum_{u \in [p] \setminus \{t\}} \sum_{v \in [p] \setminus \{t\}} \Omega_{tu} \Omega_{tv} \frac{\partial^2 \mathcal{L}_t(\underline{\Theta}_t)}{\partial \Theta_{tu} \Theta_{tv}} + 2 \sum_{i \in [n]} \sum_{u \in [p] \setminus \{t\}} \omega_t^{(i)} \Omega_{tu} \frac{\partial^2 \mathcal{L}_t(\underline{\Theta}_t)}{\partial \Theta_{tu} \theta_t^{(i)}} \\
&= \frac{1}{n} \sum_{i \in [n]} \left([\omega_t^{(i)} x_t^{(i)}]^2 + \sum_{u \in [p] \setminus \{t\}} \Omega_{tu} x_t^{(i)} x_u^{(i)} \sum_{v \in [p] \setminus \{t\}} \Omega_{tv} x_t^{(i)} x_v^{(i)} + 2 \omega_t^{(i)} x_t^{(i)} \sum_{u \in [p] \setminus \{t\}} \Omega_{tu} x_t^{(i)} x_u^{(i)} \right) \\
&\qquad\qquad \times \exp\left(- [\theta_t^{(i)} + \Theta_t^\top \boldsymbol{x}^{(i)}] x_t^{(i)} \right)
\end{aligned}
$$

$$= \frac{1}{n} \sum_{i \in [n]} \left(\omega_t^{(i)} x_t^{(i)} + \Omega_t^\top \boldsymbol{x}^{(i)} x_t^{(i)} \right)^2 \exp \left(- [\theta_t^{(i)} + \Theta_t^\top \boldsymbol{x}^{(i)}] x_t^{(i)} \right)$$

$$\stackrel{(a)}{=} \frac{1}{n} \sum_{i \in [n]} \left([\Delta_t^{(i)}]^\top \widetilde{\boldsymbol{x}}^{(i)} x_t^{(i)} \right)^2 \exp \left(- [\theta_t^{(i)} + \Theta_t^\top \boldsymbol{x}^{(i)}] x_t^{(i)} \right),$$

where (a) follows from the definitions of $\Delta_t^{(i)}$ and $\widetilde{\boldsymbol{x}}^{(i)}$.

B.2.2 Proof of Lem. 5: Lower bound on the conditional variance

For any random variable x, let $h(x)$ denote the differential entropy of x. Fix any $t \in [p]$ and $i \in [n]$. Then, we have

$$2\pi e \mathbb{V}\mathrm{ar}\big(x_t^{(i)} | \boldsymbol{x}_{-t}^{(i)}, \boldsymbol{z}^{(i)}\big) \stackrel{(a)}{\geq} \exp\left(h\big[x_t^{(i)} | \boldsymbol{x}_{-t}^{(i)}, \boldsymbol{z}^{(i)}\big] \right)$$

$$= \exp\left(-\int_{\mathcal{X}^p \times \mathcal{Z}^{p_z}} f_{\mathsf{x},\mathsf{z}}(\boldsymbol{x}^{(i)}, \boldsymbol{z}^{(i)}) \log \left(f_{\mathsf{x}_t | \mathsf{x}_{-t}, \mathsf{z}}\big(x_t^{(i)} | \boldsymbol{x}_{-t}^{(i)}, \boldsymbol{z}^{(i)}; \theta_t^\star(\boldsymbol{z}^{(i)}), \Theta_t^\star\big) \right) d\boldsymbol{x}^{(i)} d\boldsymbol{z}^{(i)} \right)$$

$$= \exp\left(-\int_{\mathcal{X}^p \times \mathcal{Z}^{p_z}} f_{\mathsf{x},\mathsf{z}}(\boldsymbol{x}^{(i)}, \boldsymbol{z}^{(i)}) \log \left(\frac{\exp\left([\theta_t^\star(\boldsymbol{z}^{(i)}) + \Theta_t^{\star\top} \boldsymbol{x}^{(i)}] x_t^{(i)} \right)}{\int_{\mathcal{X}} \exp\left([\theta_t^\star(\boldsymbol{z}^{(i)}) + \Theta_t^{\star\top} \boldsymbol{x}^{(i)}] x_t^{(i)} \right) dx_t^{(i)}} \right) d\boldsymbol{x}^{(i)} d\boldsymbol{z}^{(i)} \right)$$

$$\stackrel{(b)}{\geq} \exp\left(-\int_{\mathcal{X}^p \times \mathcal{Z}^{p_z}} f_{\mathsf{x},\mathsf{z}}(\boldsymbol{x}^{(i)}, \boldsymbol{z}^{(i)}) \log \left(\frac{\exp\left(|\theta_t^\star(\boldsymbol{z}^{(i)}) + \Theta_t^{\star\top} \boldsymbol{x}^{(i)}| x_{\max} \right)}{\int_{\mathcal{X}} \exp\left(-|\theta_t^\star(\boldsymbol{z}^{(i)}) + \Theta_t^{\star\top} \boldsymbol{x}^{(i)}| x_{\max} \right) dx_t^{(i)}} \right) d\boldsymbol{x}^{(i)} d\boldsymbol{z}^{(i)} \right)$$

$$\stackrel{(c)}{\geq} \exp\left(-\int_{\mathcal{X}^p \times \mathcal{Z}^{p_z}} f_{\mathsf{x},\mathsf{z}}(\boldsymbol{x}^{(i)}, \boldsymbol{z}^{(i)}) \log \left(\frac{\exp\left((|\theta_t^\star(\boldsymbol{z}^{(i)})| + \|\Theta_t^\star\|_1 \|\boldsymbol{x}^{(i)}\|_\infty) x_{\max} \right)}{\int_{\mathcal{X}} \exp\left(-(|\theta_t^\star(\boldsymbol{z}^{(i)})| + \|\Theta_t^\star\|_1 \|\boldsymbol{x}^{(i)}\|_\infty) x_{\max} \right) dx_t^{(i)}} \right) d\boldsymbol{x}^{(i)} d\boldsymbol{z}^{(i)} \right)$$

$$\stackrel{(d)}{\geq} \exp\left(-\int_{\mathcal{X}^p \times \mathcal{Z}^{p_z}} f_{\mathsf{x},\mathsf{z}}(\boldsymbol{x}^{(i)}, \boldsymbol{z}^{(i)}) \log \left(\frac{\exp\left((\alpha + \alpha \beta x_{\max}) x_{\max} \right)}{\int_{\mathcal{X}} \exp\left(-(\alpha + \alpha \beta x_{\max}) x_{\max} \right) dx_t^{(i)}} \right) d\boldsymbol{x}^{(i)} d\boldsymbol{z}^{(i)} \right)$$

$$\stackrel{(e)}{=} \exp\left(-\int_{\mathcal{X}^p \times \mathcal{Z}^{p_z}} f_{\mathsf{x},\mathsf{z}}(\boldsymbol{x}^{(i)}, \boldsymbol{z}^{(i)}) \log \left(\frac{C_{3,\tau}^2}{2 x_{\max}} \right) d\boldsymbol{x}^{(i)} d\boldsymbol{z}^{(i)} \right) = \frac{2 x_{\max}}{C_{3,\tau}^2},$$

where (a) follows from Shannon's entropy inequality $(h(\cdot) \leq \log \sqrt{2\pi \mathbb{V}\mathrm{ar}(\cdot)})$, (b) follows because $\|\boldsymbol{x}^{(i)}\|_\infty \leq x_{\max}$ for all $i \in [n]$, (c) follows by triangle inequality and Cauchy–Schwarz inequality, and (d) follows because $\theta^\star(\boldsymbol{z}^{(i)}) \in \Lambda_\theta$ for all $i \in [n]$, $\Theta^\star \in \Lambda_\Theta$, $\|\boldsymbol{x}^{(i)}\|_\infty \leq x_{\max}$ for all $i \in [n]$, and (e) follows because $\int_{\mathcal{X}} dx_t^{(i)} = 2 x_{\max}$.

C Proof of Thm. 1 Part II: Recovering unit-level parameters

Throughout the proof, c denotes either a universal constant or a constant depending on the model-parameters α and $x_{\max}$, and may change appearance line by line.

To analyze our estimate of the unit-level parameters, we use the estimate $\widehat{\Theta}$ of the population-level parameter $\Theta^\star$ along with the associated guarantee provided in Thm. 1. We note that the constraints on the unit-level parameters in (10) are independent across units, i.e., $\theta^{(i)} \in \Lambda_\theta$ independently for all $i \in [n]$. Therefore, we look at n independent convex optimization problems by decomposing the loss function $\mathcal{L}$ in (9) and the estimate $\widehat{\Theta}$ in (10) as follows:

$$\mathcal{L}^{(i)}\big(\theta^{(i)}\big) \triangleq \sum_{t \in [p]} \exp\left(- [\theta_t^{(i)} + \widehat{\Theta}_t^\top \boldsymbol{x}^{(i)}] x_t^{(i)} \right) \quad \text{and} \quad \widehat{\theta}^{(i)} = \underset{\theta^{(i)} \in \Lambda_\theta}{\arg\min} \mathcal{L}^{(i)}\big(\theta^{(i)}\big) \quad \text{for all} \quad i \in [n],$$

$$\tag{32}$$

Now, fix any $i \in [n]$. From (32), we have $\mathcal{L}^{(i)}(\widehat{\theta}^{(i)}) \leq \mathcal{L}^{(i)}(\theta^{\star(i)})$. Therefore, to prove this part, it is sufficient to show that all points $\theta^{(i)} \in \Lambda_\theta$ that satisfy $\|\theta^{(i)} - \theta^{\star(i)}\|_2 \geq R(\varepsilon, \delta, p)$ also uniformly satisfy

$$\mathcal{L}^{(i)}(\theta^{(i)}) \geq \mathcal{L}^{(i)}(\theta^{\star(i)}) + R^2(\varepsilon, \delta, p) \text{ for } n \geq \frac{c \exp(c\beta)}{\varepsilon^4} \cdot \left(\log \frac{p}{\delta} + \mathcal{M}(\Lambda_\theta, \eta) \right), \qquad (33)$$

with probability at least $1 - \delta$ where $R(\varepsilon, \delta, p)$ and η were defined in (11), and the metric entropy $\mathcal{M}$ was defined in Def. 2.

To that end, first, we claim that for any fixed $\theta^{(i)} \in \Lambda_\theta$, if $\theta^{(i)}$ is far from $\theta^{\star(i)}$, then with high probability $\mathcal{L}^{(i)}(\theta^{(i)})$ will be significantly larger than $\mathcal{L}^{(i)}(\theta^{\star(i)})$. We provide a proof in App. C.1.

Lemma 6 (**Gap between the loss function for a fixed parameter**). *Fix any $\varepsilon_1 > 0$, $\delta_1 \in (0, 1)$, and $i \in [n]$. Then, for any $\theta^{(i)} \in \Lambda_\theta$ such that $\|\theta^{(i)} - \theta^{\star(i)}\|_2 \geq \varepsilon_1 \gamma$ (see (11)), we have*

$$\mathcal{L}^{(i)}(\theta^{(i)}) \geq \mathcal{L}^{(i)}(\theta^{\star(i)}) + \frac{\sqrt{2}\alpha\beta x_{\max}^3}{\pi e C_{3,\tau}^4} \|\theta^{(i)} - \theta^{\star(i)}\|_2^2 \quad for \quad n \geq \frac{c \exp(c\beta) \log(p/\delta_1)}{\varepsilon_1^4},$$

with probability at least $1 - \delta_1 - c\beta^2 \log p \cdot \exp(- \exp(-c\beta)\|\theta^{(i)} - \theta^{\star(i)}\|_2^2)$ where $C_{3,\tau}$ was defined in (20).

Next, we claim that the loss function $\mathcal{L}^{(i)}$ is Lipschitz and capture this property via the following lemma. We provide a proof in App. C.2.

Lemma 7 (**Lipschitzness of the loss function**). *Consider any $i \in [n]$. Then, the loss function $\mathcal{L}^{(i)}$ is Lipschitz with respect to the ℓ_1 norm $\|\cdot\|_1$ and with Lipschitz constant $x_{\max} C_{3,\tau}$, i.e.,*

$$\left| \mathcal{L}^{(i)}(\widetilde{\theta}^{(i)}) - \mathcal{L}^{(i)}(\theta^{(i)}) \right| \leq x_{\max} C_{3,\tau} \|\widetilde{\theta}^{(i)} - \theta^{(i)}\|_1 \quad for\ all \quad \theta^{(i)}, \widetilde{\theta}^{(i)} \in \Lambda_\theta, \qquad (34)$$

where the constant $C_{3,\tau}$ was defined in (20).

Given these lemmas, we now proceed with the proof.

Proof strategy. As mentioned earlier, the idea is to show that all points $\theta^{(i)} \in \Lambda_\theta$ that satisfy $\|\theta^{(i)} - \theta^{\star(i)}\|_2 \geq R(\varepsilon, \delta, p)$ also uniformly satisfy (33) with probability at least $1 - \delta$. To do so, we consider the set of points $\Lambda_\theta^{r,i} \subset \Lambda_\theta$ whose distance from $\theta^{\star(i)}$ is at least $r > 0$ in ℓ_2 norm. Then, using an appropriate covering set of $\Lambda_\theta^{r,i}$ and the Lipschitzness of $\mathcal{L}^{(i)}$, we show that the value of $\mathcal{L}^{(i)}$ at all points in $\Lambda_\theta^{r,i}$ is uniformly $O(r^2)$ larger than the value of $\mathcal{L}^{(i)}$ at $\theta^{\star(i)}$ with high probability. Finally, we choose an r small enough to make the failure probability smaller than δ.

Gap between the loss function for all parameters in the covering set. Consider any $r \geq \varepsilon\gamma$ (where γ is defined in (11)) and the set of elements $\Lambda_\theta^{r,i} \triangleq \{ \theta^{(i)} \in \Lambda_\theta : \|\theta^{\star(i)} - \theta^{(i)}\|_2 \geq r \}$. Let $\mathcal{C}(\Lambda_\theta^{r,i}, \varepsilon')$ be the ε'-covering number of the set $\Lambda_\theta^{r,i}$ and let $\mathcal{U}(\Lambda_\theta^{r,i}, \varepsilon')$ be the associated ε'-cover (see Def. 2) where

$$\varepsilon' \triangleq \frac{\alpha\beta x_{\max}^2 r^2}{\sqrt{2}\pi e C_{3,\tau}^5 \gamma}. \qquad (35)$$

Now, we apply Lem. 6 to each element in $\mathcal{U}(\Lambda_\theta^{r,i}, \varepsilon')$ and argue by a union bound that the value of $\mathcal{L}^{(i)}$ at all points in $\mathcal{U}(\Lambda_\theta^{r,i}, \varepsilon')$ is uniformly $O(r^2)$ larger than the value of $\mathcal{L}^{(i)}$ at $\theta^{\star(i)}$ with high probability. We start by considering any $\theta^{(i)} \in \mathcal{U}(\Lambda_\theta^{r,i}, \varepsilon')$. We have

$$\|\theta^{\star(i)} - \theta^{(i)}\|_2 \overset{(a)}{\geq} r, \qquad (36)$$

where (a) follows because $\mathcal{U}(\Lambda_\theta^{r,i}, \varepsilon') \subseteq \Lambda_\theta^{r,i}$. Now, applying Lem. 6 with $\varepsilon_1 = \varepsilon$ and $\delta_1 = \delta/2\mathcal{C}(\Lambda_\theta^{r,i}, \varepsilon')$, we have

$$\mathcal{L}^{(i)}(\theta^{(i)}) \geq \mathcal{L}^{(i)}(\theta^{\star(i)}) + \frac{\sqrt{2}\alpha\beta x_{\max}^3}{\pi e C_{3,\tau}^4} \|\theta^{\star(i)} - \theta^{(i)}\|_2^2 \overset{(36)}{\geq} \mathcal{L}^{(i)}(\theta^{\star(i)}) + \frac{\sqrt{2}\alpha\beta x_{\max}^3 r^2}{\pi e C_{3,\tau}^4},$$

with probability at least $1 - \delta/2\mathcal{C}(\Lambda_\theta^{r,i}, \varepsilon') - c\beta^2 \log p \cdot \exp\left(-\exp(-c\beta)\|\theta^{(i)} - \theta^{\star(i)}\|_2^2\right)$ whenever

$$n \geq \frac{c\exp(c\beta)\log\left(2\mathcal{C}(\Lambda_\theta^{r,i}, \varepsilon') \cdot p/\delta\right)}{\varepsilon^4}. \tag{37}$$

By applying the union bound over $\mathcal{U}(\Lambda_\theta^{r,i}, \varepsilon')$, as long as n satisfies (37), we have

$$\mathcal{L}^{(i)}\left(\theta^{(i)}\right) \geq \mathcal{L}^{(i)}\left(\theta^{\star(i)}\right) + \frac{\sqrt{2}\alpha\beta x_{\max}^3 r^2}{\pi e C_{3,\tau}^4} \text{ uniformly for every } \theta^{(i)} \in \mathcal{U}(\Lambda_\theta^{r,i}, \varepsilon'), \tag{38}$$

with probability at least $1 - \delta/2 - c\beta^2\mathcal{C}(\Lambda_\theta^{r,i}, \varepsilon')\log p \cdot \exp\left(-\exp(-c\beta)\|\theta^{(i)} - \theta^{\star(i)}\|_2^2\right)$ which can lower bounded by $1 - \delta/2 - c\beta^2\mathcal{C}(\Lambda_\theta^{r,i}, \varepsilon')\log p \cdot \exp\left(-\exp(-c\beta)r^2\right)$ using (36).

Generalize beyond the covering set. Next, we assume that (38) holds, and generalize from $\mathcal{U}(\Lambda_\theta^{r,i}, \varepsilon')$ to all of $\Lambda_\theta^{r,i}$. Consider any $\widetilde{\theta}^{(i)} \in \Lambda_\theta^{r,i}$ and let $\theta^{(i)}$ be any point in $\mathcal{U}(\Lambda_\theta^{r,i}, \varepsilon')$ that satisfies $\|\theta^{(i)} - \widetilde{\theta}^{(i)}\|_2 \leq \varepsilon'$ (see Def. 2). Then, from Lem. 7, we have

$$
\begin{aligned}
\mathcal{L}^{(i)}\left(\widetilde{\theta}^{(i)}\right) \geq \mathcal{L}^{(i)}\left(\theta^{(i)}\right) - x_{\max}C_{3,\tau}\|\theta^{(i)} - \widetilde{\theta}^{(i)}\|_1 &\overset{(11)}{\geq} \mathcal{L}^{(i)}\left(\theta^{(i)}\right) - x_{\max}C_{3,\tau}\gamma\|\theta^{(i)} - \widetilde{\theta}^{(i)}\|_2 \\
&\overset{(a)}{\geq} \mathcal{L}^{(i)}\left(\theta^{(i)}\right) - x_{\max}C_{3,\tau}\gamma\varepsilon' \\
&\overset{(35)}{\geq} \mathcal{L}^{(i)}\left(\theta^{(i)}\right) - \frac{\alpha\beta x_{\max}^3 r^2}{\sqrt{2}\pi e C_{3,\tau}^4} \\
&\overset{(38)}{\geq} \mathcal{L}^{(i)}\left(\theta^{\star(i)}\right) + \frac{\alpha\beta x_{\max}^3 r^2}{\sqrt{2}\pi e C_{3,\tau}^4},
\end{aligned}
$$

where (a) follows because $\|\theta^{(i)} - \widetilde{\theta}^{(i)}\|_2 \leq \varepsilon'$.

Bounding n. To bound n in (37), we bound the covering number $\mathcal{C}(\Lambda_\theta^{r,i}, \varepsilon')$ as follows

$$\mathcal{C}(\Lambda_\theta^{r,i}, \varepsilon') \overset{(a)}{\leq} \mathcal{C}(\Lambda_\theta, \varepsilon'/2), \tag{39}$$

where (a) follows from the fact that for any sets $\mathcal{U} \subseteq \mathcal{V}$ and any ε, it holds that $\mathcal{C}(\mathcal{U}, \varepsilon) \leq \mathcal{C}(\mathcal{V}, \varepsilon/2)$. Then using (39) in (37) and observing that $\varepsilon' = \frac{r^2}{c\exp(c\beta)\gamma}$, it is sufficient for

$$n \geq \frac{c\exp(c\beta)}{\varepsilon^4} \cdot \left(\log\frac{p}{\delta} + \mathcal{M}\left(\Lambda_\theta, r^2\eta\right)\right).$$

Bounding the failure probability. To bound the failure probability by δ, it is sufficient to chose r such that

$$
\begin{aligned}
\delta &\geq \delta/2 + c\beta^2\mathcal{C}(\Lambda_\theta^{r,i}, \varepsilon')\log p \cdot \exp\left(-\exp(-c\beta)r^2\right) \\
&\overset{(39)}{\geq} \delta/2 + c\beta^2\mathcal{C}(\Lambda_\theta, \varepsilon'/2)\log p \cdot \exp\left(-\exp(-c\beta)r^2\right).
\end{aligned} \tag{40}
$$

Re-arranging and taking logarithm on both sides of (40), and observing that $\varepsilon' = \frac{r^2}{c\exp(c\beta)\gamma}$, we have

$$\log\delta \geq c\left[\log\left(\beta^2\log p\right) + \mathcal{M}\left(\Lambda_\theta, r^2\eta\right) - \exp(-c\beta)r^2\right]. \tag{41}$$

Finally, (41) holds whenever

$$r \geq c\exp\left(c\beta\right)\sqrt{\log\frac{\beta^2\log p}{\delta} + \mathcal{M}\left(\Lambda_\theta, r^2\eta\right)}.$$

Recalling that the choice of r was such that $r \geq \varepsilon\gamma$ completes the proof.

 C.1   Proof of Lem. 6: Gap between the loss function for a fixed parameter

582 Fix any $\varepsilon_1 > 0$, any $\delta_1 \in (0,1)$, and any $i \in [n]$. Consider any direction $\omega^{(i)} \in \mathbb{R}^p$ along the
583 parameter $\theta^{(i)}$, i.e.,

$$\omega^{(i)} = \theta^{(i)} - \theta^{\star(i)}. \tag{42}$$

584 We denote the first-order and the second-order directional derivatives of the loss function $\mathcal{L}^{(i)}$ in
585 (32) along the direction $\omega^{(i)}$ evaluated at $\theta^{(i)}$ by $\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{(i)}))$ and $\partial^2_{[\omega^{(i)}]^2}\mathcal{L}^{(i)}(\theta^{(i)})$ respectively.
586 Below, we state a lemma (with proof divided across App. C.1.1 and App. C.1.2) that provides us a
587 control on $\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{\star(i)}))$ and $\partial^2_{[\omega^{(i)}]^2}\mathcal{L}^{(i)}(\theta^{(i)})$. The assumptions of Lem. 6 remain in force.

588 **Lemma 8** (**Control on first and second directional derivatives**). *For any fixed $\varepsilon_2, \varepsilon_3 > 0$, $\delta_2 \in$*
589 *$(0,1)$, $i \in [n]$, $\theta^{(i)} \in \Lambda_\theta$ with $\omega^{(i)}$ defined in (42), we have the following:*

590 *(a)* **Concentration of first directional derivative***:*

$$\left|\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{\star(i)}))\right| \leq \varepsilon_2\|\omega^{(i)}\|_1 + \varepsilon_3\|\omega^{(i)}\|_2^2 \quad for \quad n \geq O\left(\frac{\exp\left(O(\beta)\right)\log(p/\delta_2)}{\varepsilon_2^4}\right),$$

591 *with probability at least $1 - \delta_2 - O\left(\beta^2\log p \exp\left(\dfrac{-\varepsilon_3^2\|\omega^{(i)}\|_2^2}{\exp\left(O(\beta)\right)}\right)\right)$.*

592 *(b)* **Anti-concentration of second directional derivative***:*

$$\partial^2_{[\omega^{(i)}]^2}\mathcal{L}^{(i)}(\theta^{(i)}) \geq \frac{8\sqrt{2}\alpha\beta x_{\max}^3}{\pi e C_{3,\tau}^3}\|\omega^{(i)}\|_2^2,$$

593 *with probability at least $1 - O\left(\beta^2\log p \exp\left(\dfrac{-\|\omega^{(i)}\|_2^2}{\exp\left(O(\beta)\right)}\right)\right)$ where $C_{3,\tau}$ was defined in*
594 *(20).*

595 Given this lemma, we now proceed to prove Lem. 6. Fix any $i \in [n]$. Define a function $g : [0,1] \to \mathbb{R}^p$
596 as follows:

$$g(a) = \theta^{\star(i)} + a(\theta^{(i)} - \theta^{\star(i)}).$$

597 Notice that $g(0) = \theta^{\star(i)}$ and $g(1) = \theta^{(i)}$ as well as

$$\frac{d\mathcal{L}^{(i)}(g(a))}{da} = \partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{(i)}))\big|_{\theta^{(i)}=g(a)} \quad \text{and} \quad \frac{d^2\mathcal{L}^{(i)}(g(a))}{da^2} = \partial^2_{[\omega^{(i)}]^2}\mathcal{L}^{(i)}(\theta^{(i)})\big|_{\theta^{(i)}=g(a)}. \tag{43}$$

598 By the fundamental theorem of calculus, we have

$$\frac{d\mathcal{L}^{(i)}(g(a))}{da} \geq \frac{d\mathcal{L}^{(i)}(g(a))}{da}\Big|_{a=0} + a\min_{a\in(0,1)}\frac{d^2\mathcal{L}^{(i)}(g(a))}{da^2}. \tag{44}$$

599 Integrating both sides of (44) with respect to a, we obtain

$$
\begin{aligned}
\mathcal{L}^{(i)}(g(a)) - \mathcal{L}^{(i)}(g(0)) &\geq a\frac{d\mathcal{L}^{(i)}(g(a))}{da}\Big|_{a=0} + \frac{a^2}{2}\min_{a\in(0,1)}\frac{d^2\mathcal{L}^{(i)}(g(a))}{da^2} \\
&\overset{(43)}{=} a\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{(i)}))\big|_{\theta^{(i)}=g(0)} + \frac{a^2}{2}\min_{a\in(0,1)}\partial^2_{[\omega^{(i)}]^2}\mathcal{L}^{(i)}(\theta^{(i)})\big|_{\theta^{(i)}=g(a)} \\
&\overset{(a)}{=} a\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{\star(i)})) + \frac{a^2}{2}\min_{a\in(0,1)}\partial^2_{[\omega^{(i)}]^2}\mathcal{L}^{(i)}(\theta^{(i)})\big|_{\theta^{(i)}=g(a)} \\
&\overset{(b)}{\geq} -a\left|\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{\star(i)}))\right| + \frac{a^2}{2}\min_{a\in(0,1)}\partial^2_{[\omega^{(i)}]^2}\mathcal{L}^{(i)}(\theta^{(i)})\big|_{\theta^{(i)}=g(a)}, \tag{45}
\end{aligned}
$$

600 where (a) follows because $g(0) = \theta^{\star(i)}$, and (b) follows by the triangle inequality. Plugging in $a = 1$
601 in (45) as well as using $g(0) = \theta^{\star(i)}$ and $g(1) = \theta^{(i)}$, we find that

$$\mathcal{L}^{(i)}(\theta^{(i)}) - \mathcal{L}^{(i)}(\theta^{\star(i)}) \geq -\left|\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{\star(i)}))\right| + \frac{1}{2}\min_{a\in(0,1)}\partial^2_{[\omega^{(i)}]^2}\mathcal{L}^{(i)}(\theta^{(i)})\big|_{\theta^{(i)}=g(a)}.$$

602 Now, we use Lem. 8 with $\varepsilon_2 = \sqrt{2}\alpha\beta x_{\max}^3 \varepsilon_1 / \pi e C_{3,\tau}^4$, $\varepsilon_3 = 2\sqrt{2}\alpha\beta x_{\max}^3 / \pi e C_{3,\tau}^4$, and $\delta_2 = \delta_1$.

603 Therefore, with probability at least $1 - \delta_1 - O\left(\beta^2 \log p \exp\left(\frac{-\|\omega^{(i)}\|_2^2}{\exp\left(O(\beta)\right)}\right)\right)$ and as long as

604 $n \geq O\left(\frac{\exp\left(O(\beta)\right)\log(p/\delta_1)}{\varepsilon_1^4}\right)$, we have

$$
\begin{aligned}
\mathcal{L}^{(i)}(\theta^{(i)}) - \mathcal{L}^{(i)}(\theta^{\star(i)}) &\geq -\frac{\sqrt{2}\alpha\beta x_{\max}^3 \varepsilon_1}{\pi e C_{3,\tau}^4}\|\omega^{(i)}\|_1 - \frac{2\sqrt{2}\alpha\beta x_{\max}^3}{\pi e C_{3,\tau}^4}\|\omega^{(i)}\|_2^2 + \frac{4\sqrt{2}\alpha\beta x_{\max}^3}{\pi e C_{3,\tau}^4}\|\omega^{(i)}\|_2^2 \\
&= -\frac{\sqrt{2}\alpha\beta x_{\max}^3 \varepsilon_1}{\pi e C_{3,\tau}^4}\|\omega^{(i)}\|_1 + \frac{2\sqrt{2}\alpha\beta x_{\max}^3}{\pi e C_{3,\tau}^4}\|\omega^{(i)}\|_2^2 \\
&\overset{(11)}{\geq} -\frac{\sqrt{2}\alpha\beta x_{\max}^3 \varepsilon_1 \gamma}{\pi e C_{3,\tau}^4}\|\omega^{(i)}\|_2 + \frac{2\sqrt{2}\alpha\beta x_{\max}^3}{\pi e C_{3,\tau}^4}\|\omega^{(i)}\|_2^2 \\
&\overset{(a)}{\geq} -\frac{\sqrt{2}\alpha\beta x_{\max}^3}{\pi e C_{3,\tau}^4}\|\omega^{(i)}\|_2^2 + \frac{2\sqrt{2}\alpha\beta x_{\max}^3}{\pi e C_{3,\tau}^4}\|\omega^{(i)}\|_2^2 = \frac{\sqrt{2}\alpha\beta x_{\max}^3}{\pi e C_{3,\tau}^4}\|\omega^{(i)}\|_2^2,
\end{aligned}
$$

605 where (a) follows because $\|\omega^{(i)}\|_2 = \|\theta^{(i)} - \theta^{\star(i)}\|_2 \geq \varepsilon_1 \gamma$ according to the lemma statement.

606 C.1.1 Proof of Lem. 8 (a): Concentration of first directional derivative

607 For every $i \in [n]$ with $\omega^{(i)}$ defined in (42), we claim that the first-order directional derivative of $\mathcal{L}^{(i)}$
608 defined in (32) is given by

$$
\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{(i)})) = -\sum_{t \in [p]} \omega_t^{(i)} x_t^{(i)} \exp\left(-[\theta_t^{(i)} + \widehat{\Theta}_t^\top \boldsymbol{x}^{(i)}]x_t^{(i)}\right). \tag{46}
$$

609 We provide a proof at the end. Assuming the claim at the moment, we proceed by fixing some
610 $i \in [n]$ and some $\theta^{(i)} \in \Lambda_\theta$. We note that the pair $\{\mathbf{x}, \mathbf{z}\}$ corresponds to a τ-SGM (Def. 8) with
611 $\tau \triangleq (\alpha, \alpha\beta, x_{\max}, \Theta)$. To show the concentration, we decompose $\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{\star(i)}))$ as a sum of
612 $L = 256\alpha^2\beta^2 x_{\max}^4 \log 4p$ terms using the conditioning trick (see App. E) with $\lambda = \frac{1}{2\sqrt{2}x_{\max}^2}$ and
613 focus on these L terms. Consider the L subsets $S_1, \cdots, S_L \in [p]$ obtained from Lem. 18 with
614 $\lambda = \frac{1}{2\sqrt{2}x_{\max}^2}$ and define

$$
\psi_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) \triangleq \sum_{t \in S_u} \omega_t^{(i)} x_t^{(i)} \exp\left(-[\theta_t^{\star(i)} + \widehat{\Theta}_t^\top \boldsymbol{x}^{(i)}]x_t^{(i)}\right) \quad \text{for every} \quad u \in L. \tag{47}
$$

615 Now, we decompose $\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{\star(i)}))$ as a sum of the L terms defined above. More precisely, we
616 have

$$
\begin{aligned}
\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{\star(i)})) &\overset{(46)}{=} -\sum_{t \in [p]} \omega_t^{(i)} x_t^{(i)} \exp\left(-[\theta_t^{\star(i)} + \widehat{\Theta}_t^\top \boldsymbol{x}^{(i)}]x_t^{(i)}\right) \\
&\overset{(a)}{=} -\frac{1}{L'}\sum_{u \in [L]}\sum_{t \in S_u} \omega_t^{(i)} x_t^{(i)} \exp\left(-[\theta_t^{\star(i)} + \widehat{\Theta}_t^\top \boldsymbol{x}^{(i)}]x_t^{(i)}\right) \\
&\overset{(47)}{=} -\frac{1}{L'}\sum_{u \in [L]} \psi_u(\boldsymbol{x}^{(i)}; \omega^{(i)}),
\end{aligned} \tag{48}
$$

617 where (a) follows because each $t \in [p]$ appears in exactly $L' = \lceil L/16\sqrt{2}\alpha\beta x_{\max}^2 \rceil$ of the sets
618 $S_1, \cdots, S_L$ according to Lem. 18(a) (with $\lambda = \frac{1}{2\sqrt{2}x_{\max}^2}$). Now, we focus on the L terms in (48).
619

620 First, we claim that for every $i \in [n]$, and $\theta^{(i)} \in \Lambda_\theta$, given $\boldsymbol{x}_{-S_u}^{(i)}$ and $\boldsymbol{z}^{(i)}$, the expected value of
621 $\psi_u(\boldsymbol{x}^{(i)}; \omega^{(i)})$ can be upper bounded uniformly across all $u \in [L]$. We provide a proof at the end of
622 this section.

623 **Lemma 9 (Upper bound on expected ψ_u).** *For any fixed $\varepsilon_5 > 0$, $\delta_5 \in (0,1)$, $i \in [n]$ and $\theta^{(i)} \in \Lambda_\theta$*
624 *with $\omega^{(i)}$ defined in (42), given $\boldsymbol{x}_{-S_u}^{(i)}$ and $\boldsymbol{z}^{(i)}$, we have*

$$
\max_{u \in [L]} \mathbb{E}\left[\psi_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) \mid \boldsymbol{x}_{-S_u}^{(i)}, \boldsymbol{z}^{(i)}\right] \leq \varepsilon_5 \|\omega^{(i)}\|_1 \quad \text{for} \quad n \geq \frac{O(\exp(\beta))\log(4p/\delta_5)}{\varepsilon_5^4},
$$

625 *with probability at least $1 - \delta_5$, where the constants $C_{1,\tau}, C_{2,\tau}$, and $C_{3,\tau}$ were defined in (20).*

626 Next, we claim that for every $i \in [n]$, $u \in [L]$, and $\theta^{(i)} \in \Lambda_\theta$, given $\boldsymbol{x}^{(i)}_{-S_u}$ and $\boldsymbol{z}^{(i)}$, $\psi_u(\boldsymbol{x}^{(i)}; \omega^{(i)})$

627 concentrates around its conditional expected value.

628 **Lemma 10 (Concentration of ψ_u).** *For any fixed $\varepsilon_6 > 0$, $i \in [n]$, $u \in [L]$, $\theta^{(i)} \in \Lambda_\theta$ with $\omega^{(i)}$*

629 *defined in (42), given $\boldsymbol{x}^{(i)}_{-S_u}$ and $\boldsymbol{z}^{(i)}$, we have*

$$\left| \psi_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) - \mathbb{E}\big[\psi_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) \mid \boldsymbol{x}^{(i)}_{-S_u}, \boldsymbol{z}^{(i)}\big]\right| \leq \varepsilon_6,$$

630 *with probability at least $1 - \exp\left(\dfrac{-\varepsilon_6^2}{\exp\left(O(\beta)\right)\|\omega^{(i)}\|_2^2}\right)$.*

631 Given these lemmas, we proceed to show the concentration of $\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{\star(i)}))$. To that end, for any

632 $u \in [L]$, given $\boldsymbol{x}^{(i)}_{-S_u}$ and $\boldsymbol{z}^{(i)}$, let E_u denote the event that

$$\psi_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) \leq \mathbb{E}\big[\psi_u(\boldsymbol{x}^{(i)}; \omega^{(i)})|\boldsymbol{x}^{(i)}_{-S_u}, \boldsymbol{z}^{(i)}\big] + \frac{1}{16\sqrt{2}\alpha\beta x_{\max}^2}\varepsilon_3\|\omega^{(i)}\|_2^2. \qquad (49)$$

633 Since E_u in an indicator event, using the law of total expectation results in

$$\mathbb{P}(E_u) = \mathbb{E}\Big[\mathbb{P}(E_u|\boldsymbol{x}^{(i)}_{-S_u}, \boldsymbol{z}^{(i)})\Big] \overset{(a)}{\geq} 1 - \exp\left(\frac{-\varepsilon_3^2\|\omega^{(i)}\|_2^2}{\exp\left(O(\beta)\right)}\right).$$

634 where (a) follows from Lem. 10 with $\varepsilon_6 = \dfrac{\varepsilon_3\|\omega^{(i)}\|_2^2}{16\sqrt{2}\alpha\beta x_{\max}^2}$. Now, by applying the union bound over

635 all $u \in [L]$ where $L = 256\alpha^2\beta^2 x_{\max}^4 \log 4p$, we have

$$\mathbb{P}\Big(\bigcap_{u \in L} E_u\Big) \geq 1 - O\left(\beta^2 \log p \exp\left(\frac{-\varepsilon_3^2\|\omega^{(i)}\|_2^2}{\exp\left(O(\beta)\right)}\right)\right).$$

636 Now, assume the event $\cap_{u \in L} E_u$ holds. Whenever this holds, we also have

$$\left|\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{\star(i)}))\right| \overset{(48)}{\leq} \frac{1}{L'}\sum_{u \in [L]}\left|\psi_u(\boldsymbol{x}^{(i)}; \omega^{(i)})\right|$$

$$\overset{(49)}{\leq} \frac{1}{L'}\sum_{u \in [L]}\left|\mathbb{E}\big[\psi_u(\boldsymbol{x}^{(i)}; \omega^{(i)})|\boldsymbol{x}^{(i)}_{-S_u}, \boldsymbol{z}^{(i)}\big] + \frac{1}{16\sqrt{2}\alpha\beta x_{\max}^2}\varepsilon_3\|\omega^{(i)}\|_2^2\right|, \quad (50)$$

637 where $L' = \lceil L/16\sqrt{2}\alpha\beta x_{\max}^2\rceil$. Further, using Lem. 9 in (50) with $\varepsilon_5 = \dfrac{\varepsilon_2}{16\sqrt{2}\alpha\beta x_{\max}^2}$ and $\delta_5 = \delta_2$,

638 whenever

$$n \geq O\left(\frac{\exp\left(O(\beta)\right)\log(p/\delta_2)}{\varepsilon_2^4}\right),$$

639 with probability at least $1 - \delta_2$, we have,

$$\left|\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{\star(i)}))\right| \leq \frac{1}{L'}\sum_{u \in [L]}\left(\frac{1}{16\sqrt{2}\alpha\beta x_{\max}^2}\varepsilon_2\|\omega^{(i)}\|_1 + \frac{1}{16\sqrt{2}\alpha\beta x_{\max}^2}\varepsilon_3\|\omega^{(i)}\|_2^2\right)$$

$$= \frac{L}{16\sqrt{2}\alpha\beta x_{\max}^2 L'}\left(\varepsilon_2\|\omega^{(i)}\|_1 + \varepsilon_3\|\omega^{(i)}\|_2^2\right) \overset{(a)}{\leq} \varepsilon_2\|\omega^{(i)}\|_1 + \varepsilon_3\|\omega^{(i)}\|_2^2,$$

640 where (a) follows because $L' = \lceil L/16\sqrt{2}\alpha\beta x_{\max}^2\rceil$.

641 **Proof of (46): Expression for first directional derivative.**   Fix any $i \in [n]$. The first-order partial

642 derivatives of $\mathcal{L}^{(i)}$ (defined in (32)) with respect to the entries of the parameter vector $\theta^{(i)}$ are given

643 by

$$\frac{\partial \mathcal{L}^{(i)}(\theta^{(i)})}{\partial \theta_t^{(i)}} = -x_t^{(i)}\exp\left(-[\theta_t^{(i)} + \widehat{\Theta}_t^\top \boldsymbol{x}^{(i)}]x_t^{(i)}\right) \quad \text{for all} \quad t \in [p].$$

Now, we can write the first-order directional derivative of $\mathcal{L}^{(i)}$ as

$$\partial_{\omega^{(i)}}(\mathcal{L}^{(i)}(\theta^{(i)})) \triangleq \lim_{h \to 0} \frac{\mathcal{L}^{(i)}(\theta^{(i)} + h\omega^{(i)}) - \mathcal{L}^{(i)}(\theta^{(i)})}{h} = \sum_{t \in [p]} \omega_t^{(i)} \frac{\partial \mathcal{L}^{(i)}(\theta^{(i)})}{\partial \theta_t^{(i)}}$$

$$= - \sum_{t \in [p]} \omega_t^{(i)} x_t^{(i)} \exp\left(- [\theta_t^{(i)} + \widehat{\Theta}_t^\top x^{(i)}] x_t^{(i)} \right).$$

Proof of Lem. 9: Upper bound on expected ψ_u. Fix any $i \in [n]$, $u \in [L]$, and $\theta^{(i)} \in \Lambda_\theta$. Then, given $x_{-S_u}^{(i)}$ and $z^{(i)}$, we have

$$\mathbb{E}\left[\psi_u(x^{(i)}; \omega^{(i)}) \mid x_{-S_u}^{(i)}, z^{(i)} \right]$$

$$\overset{(a)}{=} \mathbb{E}\left[\sum_{t \in S_u} \omega_t^{(i)} x_t^{(i)} \exp\left(- [\theta_t^{\star(i)} + \widehat{\Theta}_t^\top x^{(i)}] x_t^{(i)} \right) \mid x_{-S_u}^{(i)}, z^{(i)} \right]$$

$$\overset{(b)}{=} \sum_{t \in S_u} \omega_t^{(i)} \mathbb{E}\left[x_t^{(i)} \exp\left(- [\theta_t^{\star(i)} + \widehat{\Theta}_t^\top x^{(i)}] x_t^{(i)} \right) \mid x_{-S_u}^{(i)}, z^{(i)} \right]$$

$$\overset{(c)}{=} \sum_{t \in S_u} \omega_t^{(i)} \mathbb{E}\left[\mathbb{E}\left[x_t^{(i)} \exp\left(- [\theta_t^{\star(i)} + \widehat{\Theta}_t^\top x^{(i)}] x_t^{(i)} \right) \mid x_{-t}^{(i)}, z^{(i)} \right] \mid x_{-S_u}^{(i)}, z^{(i)} \right], \qquad (51)$$

where (a) follows from the definition of $\psi_u(x^{(i)}; \omega^{(i)})$ in (47), (b) follows from linearity of expectation, and (c) follows from the law of total expectation, i.e., $\mathbb{E}[\mathbb{E}[Y|X,Z]|Z] = \mathbb{E}[Y|Z]$ since $x_{-S_u}^{(i)} \subseteq x_{-t}^{(i)}$.

Now, for every $t \in S_u$, we will bound $\mathbb{E}\left[x_t^{(i)} \exp\left(- [\theta_t^{\star(i)} + \widehat{\Theta}_t^\top x^{(i)}] x_t^{(i)}, z^{(i)} \right) \mid x_{-t}^{(i)}, z^{(i)} \right]$. We have

$$\mathbb{E}\left[x_t^{(i)} \exp\left(- [\theta_t^{\star(i)} + \widehat{\Theta}_t^\top x^{(i)}] x_t^{(i)} \right) \mid x_{-t}^{(i)}, z^{(i)} \right]$$

$$= \int_{\mathcal{X}} x_t^{(i)} \exp\left(- [\theta_t^{\star(i)} + \widehat{\Theta}_t^\top x^{(i)}] x_t^{(i)} \right) f_{\mathsf{x}_t | \mathsf{x}_{-t}, \mathsf{z}}(x_t^{(i)} | x_{-t}^{(i)}, z^{(i)}; \theta_t^\star(z^{(i)}), \Theta_t^\star) dx_t^{(i)}$$

$$= \frac{\int_{\mathcal{X}} x_t^{(i)} \exp\left(- [\theta_t^{\star(i)} + \widehat{\Theta}_t^\top x^{(i)}] x_t^{(i)} \right) \exp\left([\theta_t^\star(z^{(i)}) + \Theta_t^{\star\top} x^{(i)}] x_t^{(i)} \right) dx_t^{(i)}}{\int_{\mathcal{X}} \exp\left([\theta_t^\star(z^{(i)}) + \Theta_t^{\star\top} x^{(i)}] x_t^{(i)} \right) dx_t^{(i)}}$$

$$\overset{(a)}{=} \frac{\int_{\mathcal{X}} x_t^{(i)} \exp\left([\Theta_t^\star - \widehat{\Theta}_t]^\top x^{(i)} x_t^{(i)} \right) dx_t^{(i)}}{\int_{\mathcal{X}} \exp\left([\theta_t^\star(z^{(i)}) + \Theta_t^{\star\top} x^{(i)}] x_t^{(i)} \right) dx_t^{(i)}}$$

$$\overset{(b)}{=} \frac{\int_{\mathcal{X}} x_t^{(i)} \left[1 + ([\Theta_t^\star - \widehat{\Theta}_t]^\top x^{(i)} x_t^{(i)}) + ([\Theta_t^\star - \widehat{\Theta}_t]^\top x^{(i)} x_t^{(i)})^2 + o([\Theta_t^\star - \widehat{\Theta}_t]^\top x^{(i)} x_t^{(i)})^3 \right] dx_t^{(i)}}{\int_{\mathcal{X}} \exp\left([\theta_t^\star(z^{(i)}) + \Theta_t^{\star\top} x^{(i)}] x_t^{(i)} \right) dx_t^{(i)}}$$

$$\overset{(c)}{=} \frac{2 x_{\max}^3 [\Theta_t^\star - \widehat{\Theta}_t]^\top x^{(i)}}{3 \int_{\mathcal{X}} \exp\left([\theta_t^\star(z^{(i)}) + \Theta_t^{\star\top} x^{(i)}] x_t^{(i)} \right) dx_t^{(i)}} + \frac{x_{\max}^5 ([\Theta_t^\star - \widehat{\Theta}_t]^\top x^{(i)})^3 o(1)}{\int_{\mathcal{X}} \exp\left([\theta_t^\star(z^{(i)}) + \Theta_t^{\star\top} x^{(i)}] x_t^{(i)} \right) dx_t^{(i)}}, \quad (52)$$

where (a) follows because $\theta^{\star(i)} = \theta^\star(z^{(i)}) \, \forall i \in [n]$, (b) follows by using the Taylor series expansion $\exp(y) = 1 + y + y^2 + o(y^3)$ around zero, (c) follows because $\int_{\mathcal{X}} x_t^{(i)} dx_t^{(i)} = 0$, $\int_{\mathcal{X}} (x_t^{(i)})^2 dx_t^{(i)} = 2x_{\max}^3/3$, $\int_{\mathcal{X}} (x_t^{(i)})^3 dx_t^{(i)} = 0$, and $\int_{\mathcal{X}} (x_t^{(i)})^4 dx_t^{(i)} = 2x_{\max}^5/5$.

Now, we bound the numerators in (52) by using $\|\Theta_t^\star - \widehat{\Theta}_t\|_1 \le 2\beta \|\Theta_t^\star - \widehat{\Theta}_t\|_\infty \le 2\beta \|\Theta_t^\star - \widehat{\Theta}_t\|_2$ which follows from the order of norms on Euclidean space as well as $\Theta^\star \in \Lambda_\Theta$ and $\widehat{\Theta} \in \Lambda_\Theta$. Then, we invoke Thm. 1 to bound $\|\Theta_t^\star - \widehat{\Theta}_t\|_2$ by $\varepsilon = \frac{3\varepsilon_5}{2\beta C_{3,\tau} x_{\max}^3}$. Therefore, we subsume the second term by the first term resulting in the following bound:

$$\mathbb{E}\left[x_t^{(i)} \exp\left(- [\theta_t^{\star(i)} + \widehat{\Theta}_t^\top x^{(i)}] x_t^{(i)} \right) \mid x_{-t}^{(i)}, z^{(i)} \right] \le \frac{2\beta C_{3,\tau} x_{\max}^3 \|\Theta_t^\star - \widehat{\Theta}_t\|_2}{3}, \qquad (53)$$

where we have used $\int_{\mathcal{X}} dx_t^{(i)} = 2x_{\max}$, $\|\boldsymbol{x}^{(i)}\|_\infty \leq x_{\max}$ for all $i \in [n]$, and the arguments used in the proof of Lem. 5 to lower bound the denominator.

Using Thm. 1 in (53) with $\varepsilon = \dfrac{3\varepsilon_5}{2\beta C_{3,\tau} x_{\max}^3}$ and $\delta = \delta_5$, we have

$$\mathbb{E}\Big[x_t^{(i)} \exp\Big(-[\theta_t^{\star(i)} + \widehat{\Theta}_t^\top \boldsymbol{x}^{(i)}]x_t^{(i)}\Big) \ \Big| \ \boldsymbol{x}_{-t}^{(i)}, \boldsymbol{z}^{(i)}\Big] \leq \varepsilon_5, \tag{54}$$

with probability at least $1 - \delta_5$ as long as

$$n \geq \frac{(130\beta^2 x_{\max}^6 C_{1,\tau} C_{2,\tau} C_{3,\tau}^8)^2 \cdot \log(4p/\delta_5)}{\varepsilon_5^4}, \quad \text{i.e.,} \quad n \geq \frac{O(\exp(\beta))\log(4p/\delta_5)}{\varepsilon_5^4}. \tag{55}$$

Using (54) and triangle inequality in (51), we have

$$\mathbb{E}\Big[\psi_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) \ \big| \ \boldsymbol{x}_{-S_u}^{(i)}, \boldsymbol{z}^{(i)}\Big] \leq \varepsilon_5 \sum_{t \in S_u} |\omega_t^{(i)}| \leq \varepsilon_5 \|\omega^{(i)}\|_1,$$

with probability at least $1 - \delta_5$ as long as n satisfies (55).

Proof of Lem. 10: Concentration of ψ_u. To show this concentration result, we use Cor. 1 (120) for the function q_1. To that end, we note that the pair $\{\mathbf{x}, \mathbf{z}\}$ corresponds to a τ-SGM (Def. 8) with $\tau \triangleq (\alpha, \alpha\beta, x_{\max}, \Theta)$. However, the random vector $\mathbf{x}$ conditioned on $\mathbf{z}$ need not satisfy the Dobrushin's uniqueness condition (Def. 4). Therefore, we cannot apply Cor. 1 (120) as is. To resolve this, we resort to the conditioning trick (Lem. 18) with $\lambda = \frac{1}{2\sqrt{2}x_{\max}^2}$.

Fix any $u \in [L]$. Then, from Lem. 18(b), (a) the pair of random vectors $\{\mathbf{x}_{S_u}, (\mathbf{x}_{-S_u}, \mathbf{z})\}$ corresponds to a τ_1-SGM with $\tau_1 \triangleq (\alpha + \alpha\beta x_{\max}, \frac{1}{2\sqrt{2}x_{\max}^2}, x_{\max}, \Theta_{S_u})$, and (b) the random vector $\mathbf{x}_{S_u}$ conditioned on $(\mathbf{x}_{-S_u}, \mathbf{z})$ satisfies the Dobrushin's uniqueness condition (Def. 4) with coupling matrix $\sqrt{2}x_{\max}^2 \Theta_{S_u}$. Now, for any fixed $i \in [n]$, we apply Cor. 1 (120) for the function q_1 with $\varepsilon = \varepsilon_6$ for a given $\boldsymbol{x}_{-S_u}^{(i)}$ and $\boldsymbol{z}^{(i)}$, to obtain

$$\mathbb{P}\Bigg(\Big|\psi_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) - \mathbb{E}\Big[\psi_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) \ \Big| \ \boldsymbol{x}_{-S_u}^{(i)}, \mathbf{z}\Big]\Big| \geq \varepsilon_6 \ \Big| \ \boldsymbol{x}_{-S_u}^{(i)}, \mathbf{z}\Bigg) \leq \exp\Bigg(\frac{-\varepsilon_6^2}{\exp\big(O(\beta)\big)\|\omega^{(i)}\|_2^2}\Bigg).$$

C.1.2 Proof of Lem. 8 (b): Anti-concentration of second directional derivative

For every $i \in [n]$ with $\omega^{(i)}$ defined in (42), we claim that the second-order directional derivative of $\mathcal{L}^{(i)}$ defined in (32) is given by

$$\partial_{[\omega^{(i)}]^2}^2 \mathcal{L}^{(i)}(\theta^{(i)}) = \sum_{t \in [p]} \big(\omega_t^{(i)} x_t^{(i)}\big)^2 \exp\Big(-[\theta_t^{(i)} + \widehat{\Theta}_t^\top \boldsymbol{x}^{(i)}]x_t^{(i)}\Big). \tag{56}$$

We provide a proof at the end. Assuming the claim at the moment, we proceed by fixing some $i \in [n]$ and some $\theta^{(i)} \in \Lambda_\theta$, Then, we lower bound $\partial_{[\omega^{(i)}]^2}^2 \mathcal{L}^{(i)}(\theta^{(i)})$ by a quadratic form as follows:

$$\begin{aligned}
\partial_{[\omega^{(i)}]^2}^2 \mathcal{L}^{(i)}(\theta^{(i)}) &\overset{(a)}{\geq} \sum_{t \in [p]} \big(\omega_t^{(i)} x_t^{(i)}\big)^2 \times \exp\Big(-|\theta_t^{(i)} + \widehat{\Theta}_t^\top \boldsymbol{x}^{(i)}|x_{\max}\Big) \\
&\overset{(b)}{\geq} \sum_{t \in [p]} \big(\omega_t^{(i)} x_t^{(i)}\big)^2 \times \exp\Big(-\big(|\theta_t^{(i)}| + \|\widehat{\Theta}_t\|_1\|\boldsymbol{x}^{(i)}\|_\infty\big)x_{\max}\Big) \\
&\overset{(c)}{\geq} \sum_{t \in [p]} \big(\omega_t^{(i)} x_t^{(i)}\big)^2 \times \exp\Big(-(\alpha + \alpha\beta x_{\max})x_{\max}\Big) \overset{(20)}{=} \frac{1}{C_{3,\tau}} \sum_{t \in [p]} \big(\omega_t^{(i)} x_t^{(i)}\big)^2,
\end{aligned} \tag{57}$$

where (a) follows from (56) because $\|\boldsymbol{x}^{(i)}\|_\infty \leq x_{\max}$ for all $i \in [n]$, (b) follows by triangle inequality and Cauchy–Schwarz inequality, and (c) follows because $\widehat{\Theta} \in \Lambda_\Theta$, $\theta^{(i)} \in \Lambda_\theta$, and

$\|\boldsymbol{x}^{(i)}\|_\infty \leq x_{\max}$ for all $i \in [n]$.

Now, to show the anti-concentration of $\partial^2_{[\omega^{(i)}]^2}\mathcal{L}^{(i)}(\theta^{(i)})$, we show the anti-concentration of the quadratic form in (57). To that end, we note that the pair $\{\mathbf{x}, \mathbf{z}\}$ corresponds to a τ-SGM (Def. 8) with $\tau \triangleq (\alpha, \alpha\beta, x_{\max}, \Theta)$. Then, we decompose the quadratic form in (57) as a sum of $L = 256\alpha^2\beta^2 x_{\max}^4 \log 4p$ terms using the conditioning trick (see App. E) with $\lambda = \frac{1}{2\sqrt{2}x_{\max}^2}$ and focus on these L terms. Consider the L subsets $S_1, \cdots, S_L \in [p]$ obtained from Lem. 18 and define

$$\overline{\psi}_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) \triangleq \sum_{t \in S_u} \left(\omega_t^{(i)} x_t^{(i)}\right)^2 \quad \text{for every} \quad u \in L. \tag{58}$$

Then, we have

$$\sum_{t \in [p]} \left(\omega_t^{(i)} x_t^{(i)}\right)^2 \stackrel{(a)}{=} \frac{1}{L'} \sum_{u \in [L]} \sum_{t \in S_u} \left(\omega_t^{(i)} x_t^{(i)}\right)^2 \stackrel{(47)}{=} \frac{1}{L'} \sum_{u \in [L]} \overline{\psi}_u(\boldsymbol{x}^{(i)}; \omega^{(i)}), \tag{59}$$

where (a) follows because each $t \in [p]$ appears in exactly $L' = \lceil L/16\sqrt{2}\alpha\beta x_{\max}^2 \rceil$ of the sets $S_1, \cdots, S_L$ according to Lem. 18(a) (with $\lambda = \frac{1}{2\sqrt{2}x_{\max}^2}$). Now, we focus on the L terms in (59).

First, we claim that for every $i \in [n]$ and $\theta^{(i)} \in \Lambda_\theta$, given $\boldsymbol{x}^{(i)}_{-S_u}$ and $\boldsymbol{z}^{(i)}$, the expected value of $\overline{\psi}_u(\boldsymbol{x}^{(i)}; \omega^{(i)})$ can be lower bounded uniformly across all $u \in [L]$. We provide a proof at the end of this section.

Lemma 11 (Lower bound on expected $\overline{\psi}_u$). *For any fixed $i \in [n]$ and $\theta^{(i)} \in \Lambda_\theta$ with $\omega^{(i)}$ defined in (42), given $\boldsymbol{x}^{(i)}_{-S_u}$ and $\boldsymbol{z}^{(i)}$, we have*

$$\min_{u \in [L]} \mathbb{E}\left[\overline{\psi}_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) \mid \boldsymbol{x}^{(i)}_{-S_u}, \boldsymbol{z}^{(i)}\right] \geq \frac{x_{\max}}{\pi e C_{3,\tau}^2}\|\omega^{(i)}\|_2^2,$$

where the constant $C_{3,\tau}$ was defined in (20).

Next, we claim that for every $i \in [n]$, $u \in [L]$, and $\theta^{(i)} \in \Lambda_\theta$, given $\boldsymbol{x}^{(i)}_{-S_u}$ and $\boldsymbol{z}^{(i)}$, $\overline{\psi}_u(\boldsymbol{x}^{(i)}; \omega^{(i)})$ concentrates around its conditional expected value.

Lemma 12 (Concentration of $\overline{\psi}_u$). *For any fixed $\varepsilon_7 > 0$, $i \in [n]$, $u \in [L]$, $\theta^{(i)} \in \Lambda_\theta$ with $\omega^{(i)}$ defined in (42), given $\boldsymbol{x}^{(i)}_{-S_u}$ and $\boldsymbol{z}^{(i)}$, we have*

$$\left|\overline{\psi}_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) - \mathbb{E}\left[\overline{\psi}_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) \mid \boldsymbol{x}^{(i)}_{-S_u}, \boldsymbol{z}^{(i)}\right]\right| \leq \varepsilon_7,$$

with probability at least $1 - \exp\left(\dfrac{-\varepsilon_7^2}{\exp\left(O(\beta)\right)\|\omega^{(i)}\|_2^2}\right)$.

Given these lemmas, we proceed to show the anti-concentration of the quadratic form in (57) implying the anti-concentration of $\partial^2_{[\omega^{(i)}]^2}\mathcal{L}^{(i)}(\theta^{(i)})$. To that end, for any $u \in [L]$, given $\boldsymbol{x}^{(i)}_{-S_u}$ and $\boldsymbol{z}^{(i)}$, let E_u denote the event that

$$\overline{\psi}_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) \geq \mathbb{E}\left[\overline{\psi}_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) | \boldsymbol{x}^{(i)}_{-S_u}, \boldsymbol{z}^{(i)}\right] - \frac{x_{\max}}{2\pi e C_{3,\tau}^2}\|\omega^{(i)}\|_2^2. \tag{60}$$

Since E_u in an indicator event, using the law of total expectation results in

$$\mathbb{P}(E_u) = \mathbb{E}\left[\mathbb{P}(E_u | \boldsymbol{x}^{(i)}_{-S_u}, \boldsymbol{z}^{(i)})\right] \stackrel{(a)}{\geq} 1 - \exp\left(\frac{\|\omega^{(i)}\|_2^2}{\exp\left(O(\beta)\right)}\right),$$

where (a) follows from Lem. 12 with $\varepsilon_7 = \frac{x_{\max}}{2\pi e C_{3,\tau}^2}\|\omega^{(i)}\|_2^2$. Now, by applying the union bound over all $u \in [L]$ where $L = 256\alpha^2\beta^2 x_{\max}^4 \log 4p$, we have

$$\mathbb{P}\left(\bigcap_{u \in L} E_u\right) \geq 1 - O\left(\beta^2 \log p \exp\left(\frac{\|\omega^{(i)}\|_2^2}{\exp\left(O(\beta)\right)}\right)\right).$$

Now, assume the event $\cap_{u\in L}E_u$ holds. Whenever this holds, we also have

$$
\sum_{t\in[p]}\left(\omega_t^{(i)}x_t^{(i)}\right)^2 \stackrel{(59)}{=} \frac{1}{L'}\sum_{u\in[L]}\overline{\psi}_u(\boldsymbol{x}^{(i)};\omega^{(i)})
$$

$$
\stackrel{(60)}{\geq} \frac{1}{L'}\sum_{u\in[L]}\left(\mathbb{E}\big[\overline{\psi}_u(\boldsymbol{x}^{(i)};\omega^{(i)})|\boldsymbol{x}_{-S_u}^{(i)},\boldsymbol{z}^{(i)}\big] - \frac{x_{\max}}{2\pi e C_{3,\tau}^2}\|\omega^{(i)}\|_2^2\right)
$$

$$
\stackrel{(a)}{\geq} \frac{1}{L'}\sum_{u\in[L]}\frac{x_{\max}}{2\pi e C_{3,\tau}^2}\|\omega^{(i)}\|_2^2 = \frac{x_{\max}L}{2\pi e L'C_{3,\tau}^2}\|\omega^{(i)}\|_2^2, \tag{61}
$$

where $L' = \lceil L/16\sqrt{2}\alpha\beta x_{\max}^2\rceil$ and (a) follows from Lem. 11. Finally, approximating $L' = L/16\sqrt{2}\alpha\beta x_{\max}^2$ and using (57), we have

$$
\partial^2_{[\omega^{(i)}]^2}\mathcal{L}^{(i)}(\theta^{(i)}) \geq \frac{1}{C_{3,\tau}}\sum_{t\in[p]}\left(\omega_t^{(i)}x_t^{(i)}\right)^2 \stackrel{(61)}{\geq} \frac{8\sqrt{2}\alpha\beta x_{\max}^3}{\pi e C_{3,\tau}^3}\|\omega^{(i)}\|_2^2,
$$

which completes the proof.

Proof of (56): Expression for second directional derivative. Fix any $i\in[n]$. The second-order partial derivatives of $\mathcal{L}^{(i)}$ (defined in (32)) with respect to the entries of the parameter vector $\theta^{(i)}$ are given by

$$
\frac{\partial^2\mathcal{L}^{(i)}(\theta^{(i)})}{\partial\big[\theta_t^{(i)}\big]^2} = \big[x_t^{(i)}\big]^2\exp\Big(-[\theta_t^{(i)}+\widehat{\Theta}_t^\top\boldsymbol{x}^{(i)}]x_t^{(i)}\Big) \quad\text{for all}\quad t\in[p].
$$

Now, we can write the second-order directional derivative of $\mathcal{L}^{(i)}$ as

$$
\partial^2_{[\omega^{(i)}]^2}\mathcal{L}^{(i)}(\theta^{(i)}) \triangleq \lim_{h\to 0}\frac{\partial_{\omega^{(i)}}\mathcal{L}^{(i)}(\theta^{(i)}+h\omega^{(i)})-\partial_{\omega^{(i)}}\mathcal{L}^{(i)}(\theta^{(i)})}{h} = \sum_{t\in[p]}\big[\omega_t^{(i)}\big]^2\frac{\partial^2\mathcal{L}^{(i)}(\theta^{(i)})}{\partial\big[\theta_t^{(i)}\big]^2}
$$

$$
= \sum_{t\in[p]}\left(\omega_t^{(i)}x_t^{(i)}\right)^2\exp\Big(-[\theta_t^{(i)}+\widehat{\Theta}_t^\top\boldsymbol{x}^{(i)}]x_t^{(i)}\Big).
$$

Proof of Lem. 11: Lower bound on expected $\overline{\psi}_u$. Fix any $i\in[n]$, $u\in[L]$, and $\theta^{(i)}\in\Lambda_\theta$. Then, given $\boldsymbol{x}_{-S_u}^{(i)}$ and $\boldsymbol{z}^{(i)}$, we have

$$
\mathbb{E}\Big[\overline{\psi}_u(\boldsymbol{x}^{(i)};\omega^{(i)})\mid\boldsymbol{x}_{-S_u}^{(i)},\boldsymbol{z}^{(i)}\Big] \stackrel{(58)}{=} \mathbb{E}\Big[\sum_{t\in S_u}\left(\omega_t^{(i)}x_t^{(i)}\right)^2\mid\boldsymbol{x}_{-S_u}^{(i)},\boldsymbol{z}^{(i)}\Big]
$$

$$
\stackrel{(a)}{=} \sum_{t\in S_u}\mathbb{E}\Big[\left(\omega_t^{(i)}x_t^{(i)}\right)^2\mid\boldsymbol{x}_{-S_u}^{(i)},\boldsymbol{z}^{(i)}\Big]
$$

$$
\stackrel{(b)}{=} \sum_{t\in S_u}\mathbb{E}\Big[\mathbb{E}\Big[\left(\omega_t^{(i)}x_t^{(i)}\right)^2\Big|\boldsymbol{x}_{-t}^{(i)},\boldsymbol{z}^{(i)}\Big]\mid\boldsymbol{x}_{-S_u}^{(i)},\boldsymbol{z}^{(i)}\Big]
$$

$$
\stackrel{(c)}{\geq} \sum_{t\in S_u}\mathbb{E}\Big[\mathbb{V}\text{ar}\Big(\omega_t^{(i)}x_t^{(i)}\Big|\boldsymbol{x}_{-t}^{(i)},\boldsymbol{z}^{(i)}\Big)\mid\boldsymbol{x}_{-S_u}^{(i)},\boldsymbol{z}^{(i)}\Big]
$$

$$
\stackrel{(d)}{\geq} \frac{x_{\max}}{\pi e C_{3,\tau}^2}\|\omega^{(i)}\|_2^2,
$$

where (a) follows from linearity of expectation, (b) follows from the law of total expectation i.e., $\mathbb{E}[\mathbb{E}[Y|X,Z]|Z] = \mathbb{E}[Y|Z]$ since $\boldsymbol{x}_{-S_u}^{(i)}\subseteq\boldsymbol{x}_{-t}^{(i)}$, (c) follows follows from the fact that for any random variable a, $\mathbb{E}[a^2]\geq\mathbb{V}\text{ar}[a]$, and (d) follows from Lem. 5.

Proof of Lem. 12: Concentration of $\overline{\psi}_u$. To show this concentration result, we use Cor. 1 (120) for the function q_2. To that end, we note that the pair $\{\mathsf{x},\mathsf{z}\}$ corresponds to a τ-SGM (Def. 8)

with $\tau \triangleq (\alpha, \alpha\beta, x_{\max}, \Theta)$. However, the random vector **x** conditioned on **z** need not satisfy the Dobrushin's uniqueness condition (Def. 4). Therefore, we cannot apply Cor. 1 (120) as is. To resolve this, we resort to the conditioning trick (Lem. 18) with $\lambda = \frac{1}{2\sqrt{2}x_{\max}^2}$.

Fix any $u \in [L]$. Then, from Lem. 18(b), (a) the pair of random vectors $\{\mathbf{x}_{S_u}, (\mathbf{x}_{-S_u}, \mathbf{z})\}$ corresponds to a τ_1-SGM with $\tau_1 \triangleq (\alpha + \alpha\beta x_{\max}, \frac{1}{2\sqrt{2}x_{\max}^2}, x_{\max}, \Theta_{S_u})$, and (b) the random vector $\mathbf{x}_{S_u}$ conditioned on $(\mathbf{x}_{-S_u}, \mathbf{z})$ satisfies the Dobrushin's uniqueness condition (Def. 4) with coupling matrix $\sqrt{2}x_{\max}^2 \Theta_{S_u}$. Now, for any fixed $i \in [n]$, we apply Cor. 1 (120) for the function q_2 with $\varepsilon = \varepsilon_7$ for a given $\boldsymbol{x}_{-S_u}^{(i)}$ and $\boldsymbol{z}^{(i)}$, to obtain

$$\mathbb{P}\left(\left|\overline{\psi}_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) - \mathbb{E}\left[\overline{\psi}_u(\boldsymbol{x}^{(i)}; \omega^{(i)}) \,\Big|\, \boldsymbol{x}_{-S_u}^{(i)}, \boldsymbol{z}^{(i)}\right]\right| \geq \varepsilon_7 \,\Big|\, \boldsymbol{x}_{-S_u}^{(i)}, \boldsymbol{z}^{(i)}\right) \leq \exp\left(\frac{-\varepsilon_7^2}{\exp\left(O(\beta)\right)\|\omega^{(i)}\|_2^2}\right).$$

C.2 Proof of Lem. 7: Lipschitzness of the loss function

Fix any $i \in [n]$, any $\theta^{(i)} \in \Lambda_\theta$, and any $\widetilde{\theta}^{(i)} \in \Lambda_\theta$. Consider the direction $\omega^{(i)} = \widetilde{\theta}^{(i)} - \theta^{(i)}$, and define the function $q : [0, 1] \to \mathbb{R}$ as follows:

$$q(a) = \mathcal{L}^{(i)}\big(\theta^{(i)} + a(\widetilde{\theta}^{(i)} - \theta^{(i)})\big). \tag{62}$$

Then, the desired inequality in (34) is equivalent to $|q(1) - q(0)| \leq x_{\max} C_{3,\tau}\|\omega^{(i)}\|_1$. From the mean value theorem, there exists $a' \in (0, 1)$ such that

$$|q(1) - q(0)| = \left|\frac{dq(a')}{da}\right|. \tag{63}$$

Therefore, we have

$$
\begin{aligned}
\big|q(1) - q(0)\big| &\overset{(63)}{=} \left|\frac{dq(a')}{da}\right| \\
&\overset{(62)}{=} \left|\frac{d\mathcal{L}^{(i)}\big(\theta^{(i)} + a'(\widetilde{\theta}^{(i)} - \theta^{(i)})\big)}{da}\right| \\
&\overset{(43)}{=} \left|\partial_{\omega^{(i)}}\big(\mathcal{L}^{(i)}(\theta^{(i)})\big)\big|_{\theta^{(i)} = \theta^{(i)} + a'(\widetilde{\theta}^{(i)} - \theta^{(i)})}\right| \\
&\overset{(46)}{=} \left|\sum_{t \in [p]} \omega_t^{(i)} x_t^{(i)} \exp\left(-[\theta_t^{(i)} + a'(\widetilde{\theta}_t^{(i)} - \theta_t^{(i)}) + \widehat{\Theta}_t^\top \boldsymbol{x}^{(i)}]x_t^{(i)}\right)\right| \\
&\overset{(a)}{\leq} x_{\max} \sum_{t \in [p]} \big|\omega_t^{(i)}\big| \exp\left(\left[\big|(1 - a')\theta_t^{(i)}\big| + \big|a'\widetilde{\theta}_t^{(i)}\big| + \|\widehat{\Theta}_t\|_1\|\boldsymbol{x}^{(i)}\|_\infty\right]x_{\max}\right) \\
&\overset{(b)}{\leq} x_{\max} \exp\left(\big((1 - a')\alpha + a'\alpha + \alpha\beta x_{\max}\big)x_{\max}\right) \sum_{t \in [p]} \big|\omega_t^{(i)}\big| \overset{(20)}{=} x_{\max} C_{3,\tau}\|\omega^{(i)}\|_1,
\end{aligned}
$$

where (a) follows from triangle inequality, Cauchy–Schwarz inequality, and because $\|\boldsymbol{x}^{(i)}\|_\infty \leq x_{\max}$ for all $i \in [n]$ and (b) follows because $\theta^{(i)}, \widetilde{\theta}^{(i)} \in \Lambda_\theta$, $\widehat{\Theta} \in \Lambda_\Theta$, and $\|\boldsymbol{x}^{(i)}\|_\infty \leq x_{\max}$ for all $i \in [n]$.

D Logarithmic Sobolev inequality and tail bounds

In this section, we present two results which may be of independent interest. First, we show that a random vector supported on a compact set satisfies the logarithmic Sobolev inequality (to be defined) if it satisfies the Dobrushin's uniqueness condition (to be defined). This result is a generalization of the result in [20] to continuous random vectors supported on a compact set. Next, we show that if a random vector satisfies the logarithmic Sobolev inequality, then any arbitrary function of the random vector concentrates around its mean. This result is a generalization of the result in [12] to continuous random vectors.

Throughout this section, we consider the random vector $\mathbf{x}$ to have support $\mathcal{X}^p$ with distribution $f_{\mathbf{x}}$. We start by defining the logarithmic Sobolev inequality (LSI). We use the convention $0 \log 0 = 0$.

Definition 3 (**Logarithmic Sobolev inequality**). *A random vector $\mathbf{x}$ satisfies the logarithmic Sobolev inequality with constant $\sigma^2 > 0$ (abbreviated as $\mathrm{LSI}_{\mathbf{x}}(\sigma^2)$) if*

$$\mathsf{Ent}_{\mathbf{x}}\left(q^2\right) \leq \sigma^2 \mathbb{E}_{\mathbf{x}}\left[\|\nabla_{\mathbf{x}} q(\mathbf{x})\|_2^2\right] \quad \text{for all} \quad q : \mathcal{X}^p \to \mathbb{R}, \tag{64}$$

where $\mathsf{Ent}_{\mathbf{x}}\left(g\right) \triangleq \mathbb{E}_{\mathbf{x}}[g(\mathbf{x}) \log g(\mathbf{x})] - \mathbb{E}_{\mathbf{x}}[g(\mathbf{x})] \log \mathbb{E}_{\mathbf{x}}[g(\mathbf{x})]$ denotes the entropy of the function $g : \mathcal{X}^p \to \mathbb{R}_+$.

Next, we restate the Dobrushin's uniqueness condition [20].

Definition 4. *[20, Dobrushin's uniqueness condition] A random vector $\mathbf{x}$ satisfies the Dobrushin's uniqueness condition with coupling matrix $\Theta \in \mathbb{R}^{p \times p}$ with $\Theta_{tt} = 0$ for all $t \in [p]$, if $\|\Theta\|_{\mathrm{op}} < 1$, and for every $t \in [p], u \in [p] \setminus \{t\}$, and $\boldsymbol{x}_{-t}, \widetilde{\boldsymbol{x}}_{-t} \in \mathcal{X}^{p-1}$ differing only in the u^{th} coordinate,*

$$\|f_{\mathsf{x}_t | \mathbf{x}_{-t} = \boldsymbol{x}_{-t}} - f_{\mathsf{x}_t | \mathbf{x}_{-t} = \widetilde{\boldsymbol{x}}_{-t}}\|_{\mathsf{TV}} \leq \Theta_{tu}. \tag{65}$$

From hereon, we let $\mathcal{X}^p$ be compact unless otherwise specified. Moreover, we define

$$f_{\min} \triangleq \min_{t \in [p], \boldsymbol{x} \in \mathcal{X}^p} f_{\mathsf{x}_t | \mathbf{x}_{-t}}(x_t | \boldsymbol{x}_{-t}). \tag{66}$$

Now, we provide the first main result of this section with a proof in App. D.1.

Proposition 1 (**Logarithmic Sobolev inequality**). *If a random vector $\mathbf{x}$ with $f_{\min} > 0$ (66) satisfies (a) the Dobrushin's uniqueness condition (Def. 4) with coupling matrix $\Theta \in \mathbb{R}^{p \times p}$, and (b) $\mathsf{x}_t | \mathbf{x}_{-t}$ satisfies $\mathrm{LSI}_{\mathsf{x}_t | \mathbf{x}_{-t} = \boldsymbol{x}_{-t}}(\sigma^2)$ for all $t \in [p]$ and $\boldsymbol{x}_{-t} \in \mathcal{X}^{p-1}$, then it satisfies $\mathrm{LSI}_{\mathbf{x}}(2\sigma^2 / (f_{\min}(1 - \|\Theta\|_{\mathrm{op}})^2))$.*

Finally, we define the notion of pseudo derivative and pseudo Hessian that come in handy in our proofs for providing upper bounds on the norm of the derivative and the Hessian.

Definition 5 (**Pseudo derivative and Hessian**). *For a function $q : \mathcal{X}^p \to \mathbb{R}$, the functions $\widetilde{\nabla} q : \mathcal{X}^p \to \mathbb{R}^{p_1}$ and $\widetilde{\nabla}^2 q : \mathcal{X}^p \to \mathbb{R}^{p_1 \times p_2}$ $(p_1, p_2 \geq 1)$ are respectively called a pseudo derivative and a pseudo Hessian for q if for all $\boldsymbol{y} \in \mathcal{X}^p$ and $\rho \in \mathbb{R}^{p_1 \times 1}$, we have*

$$\|\widetilde{\nabla} q(\boldsymbol{y})\|_2 \geq \|\nabla q(\boldsymbol{y})\|_2 \quad \text{and} \quad \|\rho^\top \widetilde{\nabla}^2 q(\boldsymbol{y})\|_2 \geq \|\nabla\left[\rho^\top \widetilde{\nabla} q(\boldsymbol{y})\right]\|_2. \tag{67}$$

Next, we provide the second main result of this section with a proof in App. D.2.

Proposition 2 (**Tail bounds for arbitrary functions under LSI**). *Given a random vector $\mathbf{x}$ satisfying $\mathrm{LSI}_{\mathbf{x}}(\sigma^2)$, any function $q : \mathcal{X}^p \to \mathbb{R}$ with a pseudo derivative $\widetilde{\nabla} q$, and pseudo Hessian $\widetilde{\nabla}^2 q$ (Def. 5) satisfies a tail bound, namely for any fixed $\varepsilon > 0$, we have*

$$\mathbb{P}\left[\left|q(\mathbf{x}) - \mathbb{E}\left[q(\mathbf{x})\right]\right| \geq \varepsilon\right] \leq \exp\left(\frac{-c}{\sigma^2} \min\left(\frac{\varepsilon^2}{\mathbb{E}\left[\|\widetilde{\nabla} q(\mathbf{x})\|_2\right]^2 + \max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{F}}^2}, \frac{\varepsilon}{\max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{op}}}\right)\right),$$

where c is a universal constant.

D.1 Proof of Prop. 1: Logarithmic Sobolev inequality

We start by defining the notion of W_2 distance [20] which is useful in the proof. We note that W_2 distance is a metric on the space of probability measures and satisfies triangle inequality.

Definition 6. *[20, W_2 distance] For random vectors $\mathbf{x}$ and $\mathbf{y}$ supported on $\mathcal{X}^p$ with distributions f and g respectively, the W_2 distance is given by $W_2^2(g_{\mathbf{y}}, f_{\mathbf{x}}) \triangleq \inf_\pi \sum_{t \in [p]}\left[\mathbb{P}_\pi(\mathsf{x}_t \neq \mathsf{y}_t)\right]^2$, where the infimum is taken over all couplings $\pi(\mathbf{x}, \mathbf{y})$ such that $\pi(\mathbf{x}) = f(\mathbf{x})$ and $\pi(\mathbf{y}) = g(\mathbf{y})$.*

Given Def. 6, our next lemma states that if appropriate W_2 distances are bounded, then the KL divergence and the entropy approximately tensorize. We provide a proof in App. D.1.1.

Lemma 13 (Approximate tensorization of KL divergence and entropy). *Given random vectors* **x** *and* **y** *supported on* $\mathcal{X}^p$ *with distributions* f *and* g *respectively such that* $f_{\min} > 0$ (66)*, if*

$$W_2^2\big(g_{\mathbf{y}_S|\mathbf{y}_{S^C}=\boldsymbol{y}_{S^C}}, f_{\mathbf{x}_S|\mathbf{x}_{S^C}=\boldsymbol{y}_{S^C}}\big) \leq C \sum_{t \in S} \mathbb{E}\Big[\|g_{y_t|\mathbf{y}_{-t}=\boldsymbol{y}_{-t}} - f_{x_t|\mathbf{x}_{-t}=\boldsymbol{y}_{-t}}\|_{\mathsf{TV}}^2 \Big| \mathbf{y}_{S^C} = \boldsymbol{y}_{S^C}\Big], \quad (68)$$

almost surely, for all subsets $S \subseteq [p]$ *with some constant* $C \geq 1$*, then*

$$\mathsf{KL}\left(g_{\mathbf{y}} \| f_{\mathbf{x}}\right) \leq \frac{2C}{f_{\min}} \sum_{t \in [p]} \mathbb{E}\big[\mathsf{KL}\left(g_{y_t|\mathbf{y}_{-t}=\boldsymbol{y}_{-t}} \| f_{x_t|\mathbf{x}_{-t}=\boldsymbol{y}_{-t}}\right)\big], \quad and \quad (69)$$

$$\mathsf{Ent}_{\mathbf{x}}\left(q\right) \leq \frac{2C}{f_{\min}} \sum_{t \in [p]} \mathbb{E}_{\mathbf{x}_{-t}}\big[\mathsf{Ent}_{x_t|\mathbf{x}_{-t}}\left(q\right)\big] \quad for\ any\ function\ q : \mathcal{X}^p \to \mathbb{R}_+. \quad (70)$$

Next, we claim that if the random vector **x** satisfies Dobrushin's uniqueness condition, then the condition (68) of Lem. 13 is naturally satisfied. We provide a proof in App. D.1.2.

Lemma 14 (Dobrushin's uniqueness implies approximate tensorization). *Given random vectors* **x** *and* **y** *supported on* $\mathcal{X}^p$ *with distributions* f *and* g *respectively, if* **x** *satisfies Dobrushin's uniqueness condition (Def. 4) with coupling matrix* $\Theta \in \mathbb{R}^{p \times p}$*, then*

$$W_2^2\big(g_{\mathbf{y}_S|\mathbf{y}_{S^C}=\boldsymbol{y}_{S^C}}, f_{\mathbf{x}_S|\mathbf{x}_{S^C}=\boldsymbol{y}_{S^C}}\big) \leq \frac{1}{\left(1-\|\Theta\|_{\mathrm{op}}\right)^2} \sum_{t \in S} \mathbb{E}\Big[\|g_{y_t|\mathbf{y}_{-t}=\boldsymbol{y}_{-t}} - f_{x_t|\mathbf{x}_{-t}=\boldsymbol{y}_{-t}}\|_{\mathsf{TV}}^2 \Big| \mathbf{y}_{S^C} = \boldsymbol{y}_{S^C}\Big], \tag{71}$$

almost surely for all subsets $S \subseteq [p]$*.*

Now to prove Prop. 1, applying Lem. 13 and 14 for an arbitrary function $f : \mathcal{X}^p \to \mathbb{R}$, we find that

$$
\begin{aligned}
\mathsf{Ent}_{\mathbf{x}}\left(q^2\right) &\leq \frac{2}{f_{\min}\left(1-\|\Theta\|_{\mathrm{op}}\right)^2} \sum_{t \in [p]} \mathbb{E}_{\mathbf{x}_{-t}}\Big[\mathsf{Ent}_{x_t|\mathbf{x}_{-t}}\left(q^2\right)\Big] \\
&\overset{(a)}{\leq} \frac{2\sigma^2}{f_{\min}\left(1-\|\Theta\|_{\mathrm{op}}\right)^2} \sum_{t \in [p]} \mathbb{E}_{\mathbf{x}_{-t}}\Big[\mathbb{E}_{x_t|\mathbf{x}_{-t}}\Big[\|\nabla_{x_t} q(x_t; \mathbf{x}_{-t})\|_2^2\Big]\Big] \\
&\overset{(b)}{=} \frac{2\sigma^2}{f_{\min}\left(1-\|\Theta\|_{\mathrm{op}}\right)^2} \mathbb{E}_{\mathbf{x}_{-t}}\Big[\mathbb{E}_{x_t|\mathbf{x}_{-t}}\Big[\sum_{t \in [p]} \|\nabla_{x_t} q(x_t; \mathbf{x}_{-t})\|_2^2\Big]\Big] \\
&\overset{(c)}{=} \frac{2\sigma^2}{f_{\min}\left(1-\|\Theta\|_{\mathrm{op}}\right)^2} \mathbb{E}_{\mathbf{x}}\Big[\|\nabla_{\mathbf{x}} q(\mathbf{x})\|_2^2\Big],
\end{aligned}
$$

where (a) follows because $x_t|\mathbf{x}_{-t}$ satisfies $\mathsf{LSI}_{x_t|\mathbf{x}_{-t}=\boldsymbol{x}_{-t}}(\sigma^2)$ for all $t \in [p]$ and $\boldsymbol{x}_{-t} \in \mathcal{X}^{p-1}$, (b) follows by the linearity of expectation and (b) follows by the law of total expectation. The claim follows.

D.1.1 Proof of Lem. 13: Approximate tensorization of KL divergence and entropy

We start by establishing a reverse-Pinsker style inequality for distributions with compact support to bound their KL divergence by their total variation distance. We provide a proof at the end.

Lemma 15 (Reverse-Pinsker inequality). *For any distributions* f *and* g *supported on* $\mathcal{X} \subset \mathbb{R}$ *such that* $f_{\min} \triangleq \min_{x \in \mathcal{X}} f(x) > 0$*, we have* $\mathsf{KL}\left(g \| f\right) \leq \frac{4}{f_{\min}}\|g-f\|_{\mathsf{TV}}^2$*.*

Given Lem. 15, we proceed to prove Lem. 13. To prove (69), we show that the following inequality holds using the technique of mathematical induction on p:

$$\mathsf{KL}\left(g_{\mathbf{y}} \| f_{\mathbf{x}}\right) \leq \frac{4C}{f_{\min}} \sum_{t \in [p]} \mathbb{E}\Big[\|g_{y_t|\mathbf{y}_{-t}=\boldsymbol{y}_{-t}} - f_{x_t|\mathbf{x}_{-t}=\boldsymbol{y}_{-t}}\|_{\mathsf{TV}}^2\Big]. \tag{72}$$

Then, (69) follows by using Pinsker's inequality to bound the right hand side of (72).

Base case: $p = 1$. For the base case, we need to establish that the claim holds for all distributions supported on $\mathcal{X}$ that satisfy the required conditions. In other words, we need to show that

$$\mathsf{KL}\left(g_y \,\|\, f_x\right) \le \frac{4C}{f_{\min}} \|g_y - f_x\|_{\mathsf{TV}}^2 \quad \text{for every} \quad t \in [p].$$

for all random variables x and y supported on $\mathcal{X}$ such that $f_{\min} = \min_{x \in \mathcal{X}} f_x(x) > 0$. This follows from Lem. 15 by observing that $C \ge 1$.

Inductive step. Now, we assume that the claim holds for all distributions supported on $\mathcal{X}^{p-1}$ that satisfy the required conditions, and establish it for distributions supported on $\mathcal{X}^p$. From the chain rule of KL divergence, we have

$$\mathsf{KL}\left(g_{\mathbf{y}} \,\|\, f_{\mathbf{x}}\right) = \mathsf{KL}\left(g_{y_t} \,\|\, f_{x_t}\right) + \mathbb{E}\left[\mathsf{KL}\left(g_{\mathbf{y}_{-t}|y_t} \,\|\, f_{\mathbf{x}_{-t}|x_t}\right)\right] \quad \text{for every} \quad t \in [p].$$

Taking an average over all $t \in [p]$, we have

$$\mathsf{KL}\left(g_{\mathbf{y}} \,\|\, f_{\mathbf{x}}\right) = \frac{1}{p}\sum_{t\in[p]} \mathsf{KL}\left(g_{y_t} \,\|\, f_{x_t}\right) + \frac{1}{p}\sum_{t\in[p]} \mathbb{E}\left[\mathsf{KL}\left(g_{\mathbf{y}_{-t}|y_t} \,\|\, f_{\mathbf{x}_{-t}|x_t}\right)\right]. \tag{73}$$

Now, we bound the first term in (73). Let π^* be such that

$$\pi^* = \operatorname*{arg\,min}_{\pi:\pi(\mathbf{x})=f(\mathbf{x}),\pi(\mathbf{y})=g(\mathbf{y})} \sum_{t\in[p]} \left[\mathbb{P}_\pi(x_t \ne y_t)\right]^2. \tag{74}$$

Then, we have

$$\frac{1}{p}\sum_{t\in[p]} \mathsf{KL}\left(g_{y_t} \,\|\, f_{x_t}\right) \overset{(a)}{\le} \frac{1}{p}\sum_{t\in[p]} \frac{4}{f_{\min}} \|g_{y_t} - f_{x_t}\|_{\mathsf{TV}}^2 \overset{(b)}{\le} \frac{4}{pf_{\min}}\sum_{t\in[p]}\left[\mathbb{P}_{\pi^*}(x_t \ne y_t)\right]^2$$

$$\overset{(c)}{=} \frac{4}{pf_{\min}} W_2^2(g_{\mathbf{y}}, f_{\mathbf{x}})$$

$$\overset{(68)}{\le} \frac{4C}{pf_{\min}}\sum_{t\in[p]}\mathbb{E}\left[\|g_{y_t|\mathbf{y}_{-t}=\boldsymbol{y}_{-t}} - f_{x_t|\mathbf{x}_{-t}=\boldsymbol{y}_{-t}}\|_{\mathsf{TV}}^2\right], \tag{75}$$

where (a) follows from Lem. 15 by observing that the marginals are lower bounded if the conditional are lower bounded, i.e., $\min_{t\in[p],x_t\in\mathcal{X}} f_{x_t}(x_t) = \min_{t\in[p],x_t\in\mathcal{X}} \int_{\boldsymbol{x}_{-t}\in\mathcal{X}^{p-1}} f_{x_t|\mathbf{x}_{-t}}(x_t|\boldsymbol{x}_{-t})f_{\mathbf{x}_{-t}}(\boldsymbol{x}_{-t})d\boldsymbol{x}_{-t} > f_{\min}$, (b) follows from the connections of total variation distance to optimal transportation cost, i.e., $\|g_y - f_x\|_{\mathsf{TV}} = \inf_{\pi:\pi(x)=f(x),\pi(y)=g(y)} \mathbb{P}_\pi(x \ne y)$, and (c) follows from Def. 6 and (74).

Next, we bound the second term in (73). We have

$$\sum_{t\in[p]} \mathbb{E}\left[\mathsf{KL}\left(g_{\mathbf{y}_{-t}|y_t} \,\|\, f_{\mathbf{x}_{-t}|x_t}\right)\right] \overset{(a)}{\le} \sum_{t\in[p]}\mathbb{E}\left[\frac{4C}{f_{\min}}\sum_{u\in[p]\setminus\{t\}}\mathbb{E}\left[\|g_{y_u|\mathbf{y}_{-u}=\boldsymbol{y}_{-u}} - f_{x_u|\mathbf{x}_{-u}=\boldsymbol{y}_{-u}}\|_{\mathsf{TV}}^2\Big|y_t = y_t\right]\right]$$

$$\overset{(b)}{=} \frac{4C}{f_{\min}}\sum_{t\in[p]}\sum_{u\in[p]\setminus\{t\}}\mathbb{E}\left[\|g_{y_u|\mathbf{y}_{-u}=\boldsymbol{y}_{-u}} - f_{x_u|\mathbf{x}_{-u}=\boldsymbol{y}_{-u}}\|_{\mathsf{TV}}^2\right]$$

$$= \frac{4C(p-1)}{f_{\min}}\sum_{u\in[p]}\mathbb{E}\left[\|g_{y_u|\mathbf{y}_{-u}=\boldsymbol{y}_{-u}} - f_{x_u|\mathbf{x}_{-u}=\boldsymbol{y}_{-u}}\|_{\mathsf{TV}}^2\right], \tag{76}$$

where (a) follows from the inductive hypothesis and (b) follows from the law of total expectation. Then, (72) follows by putting (73), (75), and (76) together.

To prove (70), we note that (69) holds for any random vector $\mathbf{y}$ supported on $\mathcal{X}^p$. Consider $\mathbf{y}$ be such that $q(\mathbf{x})/\mathbb{E}_{\mathbf{x}}[q(\mathbf{x})]$ is the Radon-Nikodym derivative of $g_{\mathbf{y}}$ with respect to $f_{\mathbf{x}}$. Then, we have

$$\frac{dg_{\mathbf{y}_{-t}}}{df_{\mathbf{x}_{-t}}} = \frac{\mathbb{E}_{x_t|\mathbf{x}_{-t}}\left[q(\mathbf{x})\right]}{\mathbb{E}_{\mathbf{x}}\left[q(\mathbf{x})\right]} \quad \text{and} \quad \frac{dg_{y_t|\mathbf{y}_{-t}}}{df_{x_t|\mathbf{x}_{-t}}} = \frac{q(\mathbf{x})}{\mathbb{E}_{x_t|\mathbf{x}_{-t}}\left[q(\mathbf{x})\right]} \quad \text{for all} \quad t \in [p]. \tag{77}$$

We have

$$\mathsf{KL}\left(g_\mathbf{y}\,\|\,f_\mathbf{x}\right) \overset{(a)}{=} \mathbb{E}_\mathbf{x}\left[\frac{dg_\mathbf{y}}{df_\mathbf{x}}\log\frac{dg_\mathbf{y}}{df_\mathbf{x}}\right]$$

$$\overset{(b)}{=} \mathbb{E}_\mathbf{x}\left[\frac{q(\mathbf{x})}{\mathbb{E}_\mathbf{x}\big[q(\mathbf{x})\big]}\log\frac{q(\mathbf{x})}{\mathbb{E}_\mathbf{x}\big[q(\mathbf{x})\big]}\right]$$

$$= \frac{1}{\mathbb{E}_\mathbf{x}\big[q(\mathbf{x})\big]}\Big(\mathbb{E}_\mathbf{x}\big[q(\mathbf{x})\log q(\mathbf{x})\big] - \mathbb{E}_\mathbf{x}\big[q(\mathbf{x})\big]\log\mathbb{E}_\mathbf{x}\big[q(\mathbf{x})\big]\Big) = \frac{\mathsf{Ent}_\mathbf{x}\left(q\right)}{\mathbb{E}_\mathbf{x}\big[q(\mathbf{x})\big]}, \qquad (78)$$

where (a) follows from the definition of KL divergence and (b) follows from the choice of **y**. Similarly, for every $t \in [p]$, we have

$$\mathbb{E}_{\mathbf{y}_{-t}}\left[\mathsf{KL}\left(g_{y_t|\mathbf{y}_{-t}=\boldsymbol{y}_{-t}}\,\|\,f_{x_t|\mathbf{x}_{-t}=\boldsymbol{y}_{-t}}\right)\right]$$

$$\overset{(a)}{=} \mathbb{E}_{\mathbf{y}_{-t}}\left[\mathbb{E}_{y_t|\mathbf{y}_{-t}}\left[\log\frac{dg_{y_t|\mathbf{y}_{-t}}}{df_{x_t|\mathbf{x}_{-t}}}\right]\right]$$

$$\overset{(b)}{=} \mathbb{E}_\mathbf{y}\left[\log\frac{dg_{y_t|\mathbf{y}_{-t}}}{df_{x_t|\mathbf{x}_{-t}}}\right]$$

$$\overset{(c)}{=} \mathbb{E}_\mathbf{x}\left[\frac{dg_\mathbf{y}}{df_\mathbf{x}}\log\frac{dg_{y_t|\mathbf{y}_{-t}}}{df_{x_t|\mathbf{x}_{-t}}}\right]$$

$$\overset{(d)}{=} \mathbb{E}_\mathbf{x}\left[\frac{q(\mathbf{x})}{\mathbb{E}_\mathbf{x}\big[q(\mathbf{x})\big]}\log\frac{q(\mathbf{x})}{\mathbb{E}_{x_t|\mathbf{x}_{-t}}\big[q(\mathbf{x})\big]}\right]$$

$$\overset{(e)}{=} \frac{\mathbb{E}_{\mathbf{x}_{-t}}\big[\mathbb{E}_{x_t|\mathbf{x}_{-t}}\big[q(\mathbf{x})\log q(\mathbf{x})\big] - \mathbb{E}_{x_t|\mathbf{x}_{-t}}\big[q(\mathbf{x})\log\mathbb{E}_{x_t|\mathbf{x}_{-t}}\big[q(\mathbf{x})\big]\big]\big]}{\mathbb{E}_\mathbf{x}\big[q(\mathbf{x})\big]} \overset{(f)}{=} \frac{\mathbb{E}_{\mathbf{x}_{-t}}\big[\mathsf{Ent}_{x_t|\mathbf{x}_{-t}}\left(q\right)\big]}{\mathbb{E}\big[q(\mathbf{x})\big]},$$

$$\tag{79}$$

where (a) follows from the definition of KL divergence, (b) follows from the law of total expectation, (c) follows from the definition of Radon-Nikodym derivative, (d) follows from the choice of **y** and (77), (e) follows from the law of total expectation, (f) follows from the definition of entropy. Then, (70) follows by putting (69), (78), and (79) together.

Proof of Lem. 15: Reverse-Pinsker inequality. Using the facts (a) $\log a \geq 1 - \frac{1}{a}$ for all $a > 0$, and (b) $f_{\min} > 0$, we find that

$$\log\frac{f(x)}{g(x)} \geq 1 - \frac{g(x)}{f(x)} \quad \text{for every} \quad x \in \mathcal{X}. \tag{80}$$

Multiplying both sides of (80) by $g(x) \geq 0$ and rearranging terms yields that

$$g(x)\log\frac{g(x)}{f(x)} \leq \frac{g^2(x)}{f(x)} - g(x) \quad \text{for every} \quad x \in \mathcal{X}. \tag{81}$$

Now, we have

$$\mathsf{KL}\left(g\,\|\,f\right) = \int_{x\in\mathcal{X}}g(x)\log\frac{g(x)}{f(x)}dx \overset{(81)}{\leq} \int_{x\in\mathcal{X}}\left(\frac{g^2(x)}{f(x)} - g(x)\right)dx$$

$$= \int_{x\in\mathcal{X}}\frac{\left(g(x) - f(x)\right)^2}{f(x)}dx$$

$$\overset{(a)}{\leq} \frac{1}{f_{\min}}\int_{x\in\mathcal{X}}\left(g(x) - f(x)\right)^2 dx$$

$$\overset{(b)}{\leq} \frac{1}{f_{\min}}\left(\int_{x\in\mathcal{X}}\big|g(x) - f(x)\big|dx\right)^2$$

$$\overset{(c)}{=} \frac{1}{f_{\min}}\left(2\|g-f\|_{\mathsf{TV}}\right)^2 = \frac{4}{f_{\min}}\|g-f\|_{\mathsf{TV}}^2,$$

where (a) follows by definition of $f_{\min}$, (b) follows by using the order of norms on Euclidean space, and (c) follows by the definition of the total variation distance.

 D.1.2   Proof of Lem. 14: Dobrushin's uniqueness implies approximate tensorization

842 We start by defining the notion of Gibbs sampler which is useful in the proof.

843 **Definition 7.** *[20, **Gibbs Sampler**] For a random vector $\mathbf{x}$ with distribution f, define the Markov*
844 *kernels and the Gibbs sampler as follows:*

$$\Gamma_t(\boldsymbol{x}|\boldsymbol{x}') \triangleq \mathbb{1}(\boldsymbol{x}_{-t} = \boldsymbol{x}'_{-t})f_{\mathsf{x}_t|\mathbf{x}_{-t}}(x_t|\boldsymbol{x}'_{-t}) \quad and \quad \Gamma(\boldsymbol{x}|\boldsymbol{x}') \triangleq p^{-1}\sum_{t\in[p]}\Gamma_t(\boldsymbol{x}|\boldsymbol{x}'), \quad (82)$$

845 *for $t \in [p]$ and $x, x' \in \mathcal{X}^p$. That is, the kernel Γ_t leaves all but the t^{th} coordinate unchanged, and*
846 *updates the t^{th} coordinate according to $f_{\mathsf{x}_t|\mathbf{x}_{-t}}$, and the sampler Γ selects an index $t \in [p]$ at random,*
847 *and applies Γ_t. Further, for a random vector $\mathbf{y}$ with distribution g supported on $\mathcal{X}^p$, we also define*

$$g_\mathbf{y}\Gamma_t(\boldsymbol{y}) \triangleq \int g_\mathbf{y}(\boldsymbol{y}')\Gamma_t(\boldsymbol{y}|\boldsymbol{y}')d\boldsymbol{y}' \text{ for } t \in [p], \quad and \quad g_\mathbf{y}\Gamma(\boldsymbol{y}) \triangleq \int g_\mathbf{y}(\boldsymbol{y}')\Gamma(\boldsymbol{y}|\boldsymbol{y}')d\boldsymbol{y}' \text{ for } \boldsymbol{y} \in \mathcal{X}^p.$$
$$(83)$$

848 We now proceed to prove Lem. 14 and split it in two cases: (i) $S = [p]$, and (ii) $S \subset [p]$.

849 **Case (i)** $(S = [p])$:   Let Γ be the Gibbs sampler associated with the distribution f. Then,

$$W_2\big(g_{\mathbf{y}_S|\mathbf{y}_{S^C}}, f_{\mathbf{x}_S|\mathbf{x}_{S^C}}\big) = W_2(g_\mathbf{y}, f_\mathbf{x}) \stackrel{(a)}{\leq} W_2(g_\mathbf{y}, g_\mathbf{y}\Gamma) + W_2(g_\mathbf{y}\Gamma, f_\mathbf{x}), \quad (84)$$

850 where (a) follows from the triangle inequality. We claim that

$$W_2(g_\mathbf{y}, g_\mathbf{y}\Gamma) \leq \frac{1}{p}\sqrt{\sum_{t\in[p]}\mathbb{E}_{\mathbf{y}_{-t}}\left[\|g_{y_t|\mathbf{y}_{-t}=\boldsymbol{y}_{-t}} - f_{\mathsf{x}_t|\mathbf{x}_{-t}=\boldsymbol{y}_{-t}}\|_{\mathsf{TV}}^2\right]}, \quad and \quad (85)$$

$$W_2(g_\mathbf{y}\Gamma, f_\mathbf{x}) \leq \left(1 - \frac{(1 - \|\Theta\|_{\mathrm{op}})}{p}\right)W_2(g_\mathbf{y}, f_\mathbf{x}). \quad (86)$$

851 Putting (84) to (86) together, we have

$$W_2(g_\mathbf{y}, f_\mathbf{x}) \leq \frac{1}{p}\sqrt{\sum_{t\in[p]}\mathbb{E}_{\mathbf{y}_{-t}}\left[\|g_{y_t|\mathbf{y}_{-t}=\boldsymbol{y}_{-t}} - f_{\mathsf{x}_t|\mathbf{x}_{-t}=\boldsymbol{y}_{-t}}\|_{\mathsf{TV}}^2\right]} + \left(1 - \frac{(1 - \|\Theta\|_{\mathrm{op}})}{p}\right)W_2(g_\mathbf{y}, f_\mathbf{x}).$$
$$(87)$$

852 Rearranging (87) results in (71) for $S = [p]$ as desired. It remains to prove our earlier claims (85)
853 and (86) which we now do one-by-one.

854 **Proof of bound (85) on $W_2(g_\mathbf{y}, g_\mathbf{y}\Gamma)$.**   To bound $W_2(g_\mathbf{y}, g_\mathbf{y}\Gamma)$, we construct a random vector $\mathbf{y}^\Gamma$
855 such that it is coupled with the random vector $\mathbf{y}$. We select an index $b \in [p]$ at random, and define

$$y_v^\Gamma \triangleq y_v \quad \text{for all} \quad v \in [p] \setminus \{b\}.$$

856 Then, given b and $\mathbf{y}_{-b} = \boldsymbol{y}_{-b}$, we define the joint distribution of (y_b, y_b^Γ) to be the maximal coupling
857 of $g_{y_b|\mathbf{y}_{-b}=\boldsymbol{y}_{-b}}$ and $f_{\mathsf{x}_b|\mathbf{x}_{-b}=\boldsymbol{y}_{-b}}$ that achieves $\|g_{y_b|\mathbf{y}_{-b}=\boldsymbol{y}_{-b}} - f_{\mathsf{x}_b|\mathbf{x}_{-b}=\boldsymbol{y}_{-b}}\|_{\mathsf{TV}}$. It is easy to see that the
858 marginal distribution of $\mathbf{y}$ is $g_\mathbf{y}$ and the marginal distribution of $\mathbf{y}^\Gamma$ is $g_\mathbf{y}\Gamma$ (see Def. 7). Then, we have

$$\begin{aligned} W_2^2(g_\mathbf{y}, g_\mathbf{y}\Gamma) &\stackrel{(a)}{\leq} \sum_{t\in[p]}\left[\mathbb{P}(b=t)\mathbb{P}(y_t \neq y_t^\Gamma|b=t) + \mathbb{P}(b \neq t)\mathbb{P}(y_t \neq y_t^\Gamma|b \neq t)\right]^2 \\ &\stackrel{(b)}{=} \sum_{t\in[p]}\left[\frac{1}{p}\mathbb{P}(y_t \neq y_t^\Gamma|b=t)\right]^2 \\ &\stackrel{(c)}{=} \frac{1}{p^2}\sum_{t\in[p]}\left[\int_{\boldsymbol{y}_{-t}\in\mathcal{X}^{p-1}}\mathbb{P}(y_t \neq y_t^\Gamma|b=t, \mathbf{y}_{-t}=\boldsymbol{y}_{-t})g_{\mathbf{y}_{-t}|b=t}(\boldsymbol{y}_{-t}|b=t)d\boldsymbol{y}_{-t}\right]^2 \\ &\stackrel{(d)}{=} \frac{1}{p^2}\sum_{t\in[p]}\left[\int_{\boldsymbol{y}_{-t}\in\mathcal{X}^{p-1}}\|g_{y_t|\mathbf{y}_{-t}=\boldsymbol{y}_{-t}} - f_{\mathsf{x}_t|\mathbf{x}_{-t}=\boldsymbol{y}_{-t}}\|_{\mathsf{TV}}g_{\mathbf{y}_{-t}}(\boldsymbol{y}_{-t})d\boldsymbol{y}_{-t}\right]^2 \end{aligned}$$

$$= \frac{1}{p^2} \sum_{t \in [p]} \left[\mathbb{E}_{\mathbf{y}_{-t}} \left[\| g_{y_t | \mathbf{y}_{-t} = \mathbf{y}_{-t}} - f_{x_t | \mathbf{x}_{-t} = \mathbf{y}_{-t}} \|_{\mathsf{TV}} \right] \right]^2, \tag{88}$$

where (a) follows from Def. 6 and the Bayes rule, (b) follows because $\mathbb{P}(b = t) = \frac{1}{p}$ and $\mathbb{P}(y_t \neq y_t^\Gamma | b \neq t) = 0$, (c) follows by the law of total probability, and (d) follows because $g_{\mathbf{y}_{-t} | b=t}(\mathbf{y}_{-t} | b = t) = g_{\mathbf{y}_{-t}}(\mathbf{y}_{-t})$ and by the construction of the coupling between $\mathbf{y}$ and $\mathbf{y}^\Gamma$. Then, (85) follows by using Jensen's inequality in (88).

Proof of bound (86) on $W_2(g_{\mathbf{y}}\Gamma, f_{\mathbf{x}})$. We first show that $f_{\mathbf{x}}$ is an invariant measure for Γ, i.e., $f_{\mathbf{x}} = f_{\mathbf{x}}\Gamma$, implying $W_2(g_{\mathbf{y}}\Gamma, f_{\mathbf{x}}) = W_2(g_{\mathbf{y}}\Gamma, f_{\mathbf{x}}\Gamma)$, and then Γ is a contraction with respect to the W_2 distance with rate $1 - \frac{(1 - \|\Theta\|_{\mathrm{op}})}{p}$, i.e., $W_2(g_{\mathbf{y}}\Gamma, f_{\mathbf{x}}\Gamma) \leq \left(1 - \frac{(1 - \|\Theta\|_{\mathrm{op}})}{p} \right) W_2(g_{\mathbf{y}}, f_{\mathbf{x}})$, implying (86).

Proof of $f_{\mathbf{x}}$ being an invariant measure for Γ. We have

$$f_{\mathbf{x}}\Gamma(\boldsymbol{x}) \overset{(83)}{=} \int_{\boldsymbol{x}' \in \mathcal{X}^p} f_{\mathbf{x}}(\boldsymbol{x}')\Gamma(\boldsymbol{x}|\boldsymbol{x}')d\boldsymbol{x}' \overset{(82)}{=} \int_{\boldsymbol{x}' \in \mathcal{X}^p} f_{\mathbf{x}}(\boldsymbol{x}')\left(\frac{1}{p} \sum_{t \in [p]} \Gamma_t(\boldsymbol{x}|\boldsymbol{x}') \right) d\boldsymbol{x}'$$

$$\overset{(82)}{=} \frac{1}{p} \sum_{t \in [p]} \int_{\boldsymbol{x}' \in \mathcal{X}^p} f_{\mathbf{x}}(\boldsymbol{x}')\mathbb{1}(\boldsymbol{x}_{-t} = \boldsymbol{x}'_{-t})f_{x_t|\mathbf{x}_{-t}}(x_t|\boldsymbol{x}'_{-t})d\boldsymbol{x}'$$

$$= \frac{1}{p} \sum_{t \in [p]} f_{x_t|\mathbf{x}_{-t}}(x_t|\boldsymbol{x}_{-t}) \int_{x'_t \in \mathcal{X}} f_{\mathbf{x}}(\boldsymbol{x}_{-t}, x'_t)dx'_t$$

$$= \frac{1}{p} \sum_{t \in [p]} f_{x_t|\mathbf{x}_{-t}}(x_t|\boldsymbol{x}_{-t})f_{\mathbf{x}_{-t}}(\boldsymbol{x}_{-t}) = f_{\mathbf{x}}(\boldsymbol{x}).$$

Proof of Γ being a contraction w.r.t the W_2 distance. Let π^* be the coupling between $\mathbf{x}$ and $\mathbf{y}$ that achieves $W_2(g_{\mathbf{y}}, f_{\mathbf{x}})$ i.e.,

$$\pi^* = \operatorname*{arg\,min}_{\pi : \pi(\mathbf{x}) = f(\mathbf{x}), \pi(\mathbf{y}) = g(\mathbf{y})} \sqrt{\sum_{t \in [p]} \left[\mathbb{P}_\pi(x_t \neq y_t) \right]^2}. \tag{89}$$

We construct random variables $\mathbf{x}'$ and $\mathbf{y}'$ as well as a coupling π' between them such that the marginal distribution of $\mathbf{x}'$ is $f_{\mathbf{x}}\Gamma$ and the marginal distribution of $\mathbf{y}'$ is $g_{\mathbf{y}}\Gamma$. We start by selecting an index $b \in [p]$ at random, and defining

$$y'_v \triangleq y_v \quad \text{and} \quad x'_v \triangleq x_v \quad \text{for all} \quad v \neq b. \tag{90}$$

Then, given b, $\mathbf{y}'_{-b} = \boldsymbol{y}_{-b}$, and $\mathbf{x}'_{-b} = \boldsymbol{x}_{-b}$, we define the joint distribution of (y'_b, x'_b) to be the maximal coupling of $f_{x_b|\mathbf{x}_{-b}}(\cdot|\boldsymbol{y}_{-b})$ and $f_{x_b|\mathbf{x}_{-b}}(\cdot|\boldsymbol{x}_{-b})$ that achieves $\|f_{x_b|\mathbf{x}_{-b} = \boldsymbol{y}_{-b}} - f_{x_b|\mathbf{x}_{-b} = \boldsymbol{x}_{-b}}\|_{\mathsf{TV}}$.

Now, for every $t \in [p]$, we bound $\mathbb{P}_{\pi'}(y'_t \neq x'_t)$ in terms of $\mathbb{P}_{\pi^*}(y_t \neq x_t)$. To that end, we have

$$\mathbb{P}_{\pi'}(y'_t \neq x'_t) \overset{(a)}{=} \mathbb{P}(b = t)\mathbb{P}_{\pi'}(y'_t \neq x'_t | b = t) + \mathbb{P}(b \neq t)\mathbb{P}_{\pi'}(y'_t \neq x'_t | b \neq t)$$

$$\overset{(b)}{=} \frac{1}{p}\mathbb{P}_{\pi'}(y'_t \neq x'_t | b = t) + \left(1 - \frac{1}{p} \right)\mathbb{P}_{\pi^*}(y_t \neq x_t), \tag{91}$$

where (a) follows from the Bayes rule and (b) follows because $\mathbb{P}(b = t) = \frac{1}{p}$ and (90). Focusing on $\mathbb{P}_{\pi'}(y'_t \neq x'_t | b = t)$ and using the law of total probability, we have

$$\mathbb{P}_{\pi'}(y'_t \neq x'_t | b = t)$$

$$= \int_{\boldsymbol{y}_{-t}, \boldsymbol{x}_{-t} \in \mathcal{X}^{p-1}} \mathbb{P}_{\pi'}(y'_t \neq x'_t | b = t, \mathbf{y}'_{-t} = \boldsymbol{y}_{-t}, \mathbf{x}'_{-t} = \boldsymbol{x}_{-t})\pi'_{\mathbf{y}'_{-t}, \mathbf{x}'_{-t}|b=t}(\boldsymbol{y}_{-t}, \boldsymbol{x}_{-t}|b = t)d\boldsymbol{y}_{-t}d\boldsymbol{x}_{-t}$$

$$\overset{(a)}{=} \int_{\boldsymbol{y}_{-t}, \boldsymbol{x}_{-t} \in \mathcal{X}^{p-1}} \|f_{x_t|\mathbf{x}_{-t} = \boldsymbol{y}_{-t}} - f_{x_t|\mathbf{x}_{-t} = \boldsymbol{x}_{-t}}\|_{\mathsf{TV}}\pi^*_{\mathbf{y}_{-t}, \mathbf{x}_{-t}}(\boldsymbol{y}_{-t}, \boldsymbol{x}_{-t})d\boldsymbol{y}_{-t}d\boldsymbol{x}_{-t}$$

$$= \mathbb{E}_{\pi^*_{\mathbf{y}_{-t},\mathbf{x}_{-t}}} \Big[\| f_{\mathsf{x}_t | \mathbf{x}_{-t} = \mathbf{y}_{-t}} - f_{\mathsf{x}_t | \mathbf{x}_{-t} = \mathbf{x}_{-t}} \|_{\mathsf{TV}} \Big]$$

$$\overset{(b)}{\leq} \mathbb{E}_{\pi^*_{\mathbf{y}_{-t},\mathbf{x}_{-t}}} \Big[\sum_{u \in [p] \setminus \{t\}} \mathbb{1}(y_u \neq \mathsf{x}_u) \mathbb{1}(\mathbf{y}_{-\{t,u\}} = \mathbf{x}_{-\{t,u\}}) \| f_{\mathsf{x}_t | \mathbf{x}_{-t}} - f_{\mathsf{x}_t | \mathbf{x}_{-t}}(\cdot | \mathbf{x}_{-t}) \|_{\mathsf{TV}} \Big]$$

$$\overset{(65)}{\leq} \mathbb{E}_{\pi^*_{\mathbf{y}_{-t},\mathbf{x}_{-t}}} \Big[\sum_{u \in [p] \setminus \{t\}} \Theta_{tu} \mathbb{1}(y_u \neq \mathsf{x}_u) \Big] = \sum_{u \in [p] \setminus \{t\}} \Theta_{tu} \mathbb{P}_{\pi^*}(y_u \neq \mathsf{x}_u), \tag{92}$$

where (a) follows by the construction of the coupling between $\mathbf{y}'$ and $\mathbf{x}'$, and (b) follows by triangle inequality. Putting together (91) and (92), we have

$$\mathbb{P}_{\pi'}(y'_t \neq \mathsf{x}'_t) \leq \frac{1}{p} \sum_{u \in [p] \setminus \{t\}} \Theta_{tu} \mathbb{P}_{\pi^*}(y_u \neq \mathsf{x}_u) + \Big(1 - \frac{1}{p} \Big) \mathbb{P}_{\pi^*}(y_t \neq \mathsf{x}_t). \tag{93}$$

Next, we use (93) to show contraction of Γ. We have

$$W_2^2(g_{\mathbf{y}} \Gamma, f_{\mathbf{x}} \Gamma) \overset{(a)}{\leq} \sum_{t \in [p]} \Big[\mathbb{P}_{\pi'}(y'_t \neq \mathsf{x}'_t) \Big]^2$$

$$\overset{(93)}{\leq} \sum_{t \in [p]} \Big[\frac{1}{p} \sum_{j \in [p] \setminus \{i\}} \Theta_{tu} \mathbb{P}_{\pi^*}(y_j \neq \mathsf{x}_j) + \Big(1 - \frac{1}{p} \Big) \mathbb{P}_{\pi^*}(y_t \neq \mathsf{x}_t) \Big]^2$$

$$\overset{(b)}{\leq} \Big\| \Big(1 - \frac{1}{p} \Big) I + \frac{1}{p} \Theta \Big\|_{\mathrm{op}}^2 \sum_{t \in [p]} \Big[\mathbb{P}_{\pi^*}(y_t \neq \mathsf{x}_t) \Big]^2$$

$$\overset{(c)}{=} \Big\| \Big(1 - \frac{1}{p} \Big) I + \frac{1}{p} \Theta \Big\|_{\mathrm{op}}^2 W_2^2(g_{\mathbf{y}}, f_{\mathbf{x}})$$

$$\overset{(d)}{\leq} \Big(\Big(1 - \frac{1}{p} \Big) + \frac{1}{p} \| \Theta \|_{\mathrm{op}} \Big)^2 W_2^2(g_{\mathbf{y}}, f_{\mathbf{x}}), \tag{94}$$

where (a) follows from Def. 6, (b) follows by some linear algebraic manipulations, (c) follows from Def. 6 and (89), and (d) follows from the triangle inequality. Then, contraction of Γ follows by taking square root on both sides of (94).

Case (ii) $(S \subset [p])$**:**   We can directly verify that the matrix $\Theta_S \triangleq \{\Theta_{tu}\}_{t,u \in S}$ is such that $\| \Theta_S \|_{\mathrm{op}} \leq \| \Theta \|_{\mathrm{op}}$. Further, we note that for any $\mathbf{y}_{S^C} \in \mathcal{X}^{p - |S|}$, the random vector $\mathbf{x}_S | \mathbf{x}_{S^C} = \mathbf{y}_{S^C}$ with distribution $f_{\mathbf{x}_S | \mathbf{x}_{S^C} = \mathbf{y}_{S^C}}$ satisfies the Dobrushin's uniqueness condition (Def. 4) with coupling matrix Θ_S. Then, by performing an analysis similar to the one above, we have

$$W_2 \big(g_{\mathbf{y}_S | \mathbf{y}_{S^C}}, f_{\mathbf{x}_S | \mathbf{x}_{S^C}} \big) \leq \frac{1}{(1 - \| \Theta_S \|_{\mathrm{op}})} \sqrt{\sum_{t \in S} \mathbb{E} \Big[\| g_{y_t | \mathbf{y}_{-t} = \mathbf{y}_{-t}} - f_{\mathsf{x}_t | \mathbf{x}_{-t} = \mathbf{y}_{-t}} \|_{\mathsf{TV}}^2 \Big| \mathbf{y}_{S^C} = \mathbf{y}_{S^C} \Big]}$$

$$\overset{(a)}{\leq} \frac{1}{(1 - \| \Theta \|_{\mathrm{op}})} \sqrt{\sum_{t \in S} \mathbb{E} \Big[\| g_{y_t | \mathbf{y}_{-t} = \mathbf{y}_{-t}} - f_{\mathsf{x}_t | \mathbf{x}_{-t} = \mathbf{y}_{-t}} \|_{\mathsf{TV}}^2 \Big| \mathbf{y}_{S^C} = \mathbf{y}_{S^C} \Big]},$$

where (a) follows because $\frac{1}{(1 - \| \Theta_S \|_{\mathrm{op}})} \leq \frac{1}{(1 - \| \Theta \|_{\mathrm{op}})}$. This completes the proof.

D.2   Proof of Prop. 2: Tail bounds for arbitrary functions under LSI

Fix any pseudo derivative $\widetilde{\nabla} q$ and any pseudo Hessian $\widetilde{\nabla}^2 q$. To prove Prop. 2, we bound the p-th moment of $q(\mathbf{x}) - \mathbb{E}\big[q(\mathbf{x})\big]$. To that end, first, we claim that in order to control the p-th moment of $q(\mathbf{x}) - \mathbb{E}\big[q(\mathbf{x})\big]$, it is sufficient to control the p-th moment of $\| \nabla q(\mathbf{x}) \|_2$. We provide a proof in App. D.2.1.

Lemma 16 (**Bounded p-th moment of $q(\mathbf{x}) - \mathbb{E}\big[q(\mathbf{x})\big]$**)**.** *If a random vector $\mathbf{x}$ satisfies* $\mathrm{LSI}_{\mathbf{x}}(\sigma^2)$, *then for any arbitrary function* $q : \mathcal{X}^p \to \mathbb{R}$, *and any* $p \geq 2$,

$$\big\| q(\mathbf{x}) - \mathbb{E}\big[q(\mathbf{x})\big] \big\|_{L_p} \leq \sigma \sqrt{2p} \, \big\| \| \nabla q(\mathbf{x}) \|_2 \big\|_{L_p}.$$

We note that the p-th moment of $\|\nabla q(\mathbf{x})\|_2$ is bounded by the p-th moment of $\|\widetilde{\nabla} q(\mathbf{x})\|_2$. Next, we claim that the p-th moment of $\|\widetilde{\nabla} q(\mathbf{x})\|_2$ is appropriately bounded by a linear combination of certain norms of $\widetilde{\nabla}^2 q$ and $\mathbb{E}_{\mathbf{x}}\big[\widetilde{\nabla} q(\mathbf{x})\big]$. We provide a proof in App. D.2.2.

Lemma 17 (**Bounded p-th moment of** $\|\widetilde{\nabla} q(\mathbf{x})\|_2$)**.** *If a random vector $\mathbf{x}$ satisfies $\mathrm{LSI}_{\mathbf{x}}(\sigma^2)$, then for any arbitrary function $q : \mathcal{X}^p \to \mathbb{R}$, any pseudo derivative $\widetilde{\nabla} q(\boldsymbol{x})$, any pseudo Hessian $\widetilde{\nabla}^2 q(\boldsymbol{x})$, and any even $p \geq 2$,*

$$\big\|\|\widetilde{\nabla} q(\mathbf{x})\|_2\big\|_{L_p} \leq 2c\sigma\Big(\max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{F}} + \sqrt{p} \max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{op}} \Big) + 4\|\mathbb{E}_{\mathbf{x}}\big[\widetilde{\nabla} q(\mathbf{x})\big]\|_2,$$

where $c \geq 0$ is a universal constant.

Given these lemmas, we proceed to prove Prop. 2. Combining Lem. 16 and 17 for any even $p \geq 2$, there exists a universal constant c' such that

$$\big\|q(\mathbf{x}) - \mathbb{E}\big[q(\mathbf{x})\big]\big\|_{L_p} \leq c'\sigma^2\Big(\sqrt{p} \max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{F}} + p \max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{op}} + \sqrt{p}\|\mathbb{E}_{\mathbf{x}}\big[\widetilde{\nabla} q(\mathbf{x})\big]\|_2\Big). \tag{95}$$

Now, we complete the proof by using (95) along with Markov's inequality for a specific choice of p. For any even $p \geq 2$, we have

$$\mathbb{P}\Big[\big|q(\mathbf{x}) - \mathbb{E}\big[q(\mathbf{x})\big]\big| > ec'\sigma^2\Big(\sqrt{p} \max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{F}} + p \max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{op}} + \sqrt{p}\|\mathbb{E}_{\mathbf{x}}\big[\widetilde{\nabla} q(\mathbf{x})\big]\|_2\Big)\Big]$$

$$= \mathbb{P}\Big[\big|q(\mathbf{x}) - \mathbb{E}\big[q(\mathbf{x})\big]\big|^p > \big(ec'\sigma^2\big)^p\Big(\sqrt{p} \max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{F}} + p \max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{op}} + \sqrt{p}\|\mathbb{E}_{\mathbf{x}}\big[\widetilde{\nabla} q(\mathbf{x})\big]\|_2\Big)^p\Big]$$

$$\overset{(a)}{\leq} \frac{\mathbb{E}\Big[\big|q(\mathbf{x}) - \mathbb{E}\big[q(\mathbf{x})\big]\big|^p\Big]}{\big(ec'\sigma^2\big)^p\Big(\sqrt{p}\max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{F}} + p\max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{op}} + \sqrt{p}\|\mathbb{E}_{\mathbf{x}}\big[\widetilde{\nabla} q(\mathbf{x})\big]\|_2\Big)^p}$$

$$\overset{(95)}{\leq} e^{-p},$$

where (a) follows from Markov's inequality. The proof is complete by choosing

$$p = \frac{1}{4ec'\sigma^2} \min\Big(\frac{1}{ec'\sigma^2} \cdot \frac{\varepsilon^2}{\mathbb{E}\big[\|\widetilde{\nabla} q(\mathbf{x})\|_2\big]^2 + \max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{F}}^2}, \frac{\varepsilon}{\max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{op}}} \Big),$$

and performing basic algebraic manipulations.

D.2.1 Proof of Lem. 16: Bounded p-th moment of $q(\mathbf{x}) - \mathbb{E}\big[q(\mathbf{x})\big]$

We start by using the following result from [4, Theorem 3.4] since $\mathbf{x}$ satisfies $\mathrm{LSI}_{\mathbf{x}}(\sigma^2)$:

$$\big\|q(\mathbf{x}) - \mathbb{E}\big[q(\mathbf{x})\big]\big\|_{L_p}^2 \leq \big\|q(\mathbf{x}) - \mathbb{E}\big[q(\mathbf{x})\big]\big\|_{L_2}^2 + 2\sigma^2(p - 2)\big\|\|\nabla q(\mathbf{x})\|_2\big\|_{L_p}^2. \tag{96}$$

Then, we bound the first term in (96) by using the fact that logarithmic Sobolev inequality implies Poincare inequality with the same constant:

$$\big\|q(\mathbf{x}) - \mathbb{E}\big[q(\mathbf{x})\big]\big\|_{L_2}^2 = \mathbb{V}\mathrm{ar}(q(\mathbf{x})) \leq \sigma^2\mathbb{E}_{\mathbf{x}}\Big[\|\nabla q(\mathbf{x})\|_2^2\Big]. \tag{97}$$

Putting together (96) and (97), we have

$$\big\|q(\mathbf{x}) - \mathbb{E}\big[q(\mathbf{x})\big]\big\|_{L_p}^2 \leq \sigma^2\mathbb{E}_{\mathbf{x}}\Big[\|\nabla q(\mathbf{x})\|_2^2\Big] + 2\sigma^2(p - 2)\big\|\|\nabla q(\mathbf{x})\|_2\big\|_{L_p}^2$$

$$\overset{(a)}{\leq} \sigma^2\Big(\mathbb{E}_{\mathbf{x}}\Big[\|\nabla q(\mathbf{x})\|_2^p\Big]\Big)^{2/p} + 2\sigma^2(p - 2)\big\|\|\nabla q(\mathbf{x})\|_2\big\|_{L_p}^2$$

$$\overset{(b)}{=} \sigma^2\big\|\|\nabla q(\mathbf{x})\|_2\big\|_{L_p}^2 + 2\sigma^2(p - 2)\big\|\|\nabla q(\mathbf{x})\|_2\big\|_{L_p}^2 \leq 2\sigma^2 p\big\|\|\nabla q(\mathbf{x})\|_2\big\|_{L_p}^2, \tag{98}$$

where (a) follows by Jensen's inequality and (b) follows by the definition of p-th moment. Taking square root on both sides of (98) completes the proof.

D.2.2 Proof of Lem. 17: Bounded p-th moment of $\|\widetilde{\nabla}q(\mathbf{x})\|_2$

Fix any pseudo derivative $\widetilde{\nabla}q$ and any pseudo Hessian $\widetilde{\nabla}^2 q$. We start by obtaining a convenient bound on $\|\widetilde{\nabla}q(\boldsymbol{x})\|_2$ for every $\boldsymbol{x} \in \mathcal{X}^p$ and then proceed to bound the p-th moment of $\|\widetilde{\nabla}q(\mathbf{x})\|_2$.

Consider a p-dimensional standard normal random vector $\mathbf{g}$ independent of $\mathbf{x}$. For a given $\mathbf{x} = \boldsymbol{x} \in \mathcal{X}^p$, the random variable $\dfrac{\widetilde{\nabla}q(\boldsymbol{x})^\top \mathbf{g}}{\|\widetilde{\nabla}q(\boldsymbol{x})\|_2}$ is a standard normal random variable. Then, for every $\boldsymbol{x} \in \mathcal{X}^p$, we have

$$\left\| \frac{\widetilde{\nabla}q(\boldsymbol{x})^\top \mathbf{g}}{\|\widetilde{\nabla}q(\boldsymbol{x})\|_2} \right\|_{L_p} \overset{(a)}{=} \left(\mathbb{E}_{\mathbf{g}|\mathbf{x}=\boldsymbol{x}} \left[\left(\frac{\widetilde{\nabla}q(\boldsymbol{x})^\top \mathbf{g}}{\|\widetilde{\nabla}q(\boldsymbol{x})\|_2} \right)^p \right] \right)^{1/p} \overset{(b)}{\geq} \frac{\sqrt{p}}{2}. \tag{99}$$

where (a) follows from the definition of p-th moment, and (b) follows since $\|g\|_{L_p} \geq \frac{\sqrt{p}}{2}$ for any standard normal random variable g and even p. Rearranging (99), we have

$$\|\widetilde{\nabla}q(\boldsymbol{x})\|_2 \leq \frac{2}{\sqrt{p}} \left(\mathbb{E}_{\mathbf{g}|\mathbf{x}=\boldsymbol{x}} \left[(\widetilde{\nabla}q(\boldsymbol{x})^\top \mathbf{g})^p \right] \right)^{1/p}. \tag{100}$$

Now, we proceed to bound the p-th moment of $\|\widetilde{\nabla}q(\mathbf{x})\|_2$ as follows:

$$\begin{aligned}
\left\| \|\widetilde{\nabla}q(\mathbf{x})\|_2 \right\|_{L_p} &\overset{(a)}{=} \left(\mathbb{E}_{\mathbf{x}} \left[\|\widetilde{\nabla}q(\mathbf{x})\|_2^p \right] \right)^{1/p} \\
&\overset{(100)}{\leq} \frac{2}{\sqrt{p}} \left(\mathbb{E}_{\mathbf{x},\mathbf{g}} \left[(\widetilde{\nabla}q(\mathbf{x})^\top \mathbf{g})^p \right] \right)^{1/p} \\
&\overset{(b)}{=} \frac{2}{\sqrt{p}} \left\| \widetilde{\nabla}q(\mathbf{x})^\top \mathbf{g} \right\|_{L_p} \\
&\overset{(c)}{\leq} \frac{2}{\sqrt{p}} \left(\left\| \widetilde{\nabla}q(\mathbf{x})^\top \mathbf{g} - \mathbb{E}_{\mathbf{x}} \left[\widetilde{\nabla}q(\mathbf{x})^\top \mathbf{g} \right] \right\|_{L_p} + \left\| \mathbb{E}_{\mathbf{x}} \left[\widetilde{\nabla}q(\mathbf{x})^\top \mathbf{g} \right] \right\|_{L_p} \right),
\end{aligned} \tag{101}$$

where (a) and (b) follow from the definition of p-th moment and (c) follows by Minkowski's inequality. We claim that

$$\left\| \widetilde{\nabla}q(\mathbf{x})^\top \mathbf{g} - \mathbb{E}_{\mathbf{x}} \left[\widetilde{\nabla}q(\mathbf{x})^\top \mathbf{g} \right] \right\|_{L_p} \leq c\sigma \left(\sqrt{p} \max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{F}} + p \max_{\boldsymbol{x} \in \mathcal{X}^p} \|\widetilde{\nabla}^2 q(\boldsymbol{x})\|_{\mathrm{op}} \right), \text{ and} \tag{102}$$

$$\left\| \mathbb{E}_{\mathbf{x}} \left[\widetilde{\nabla}q(\mathbf{x})^\top \mathbf{g} \right] \right\|_{L_p} \leq 2\sqrt{p} \left\| \mathbb{E}_{\mathbf{x}} \left[\widetilde{\nabla}q(\mathbf{x}) \right] \right\|_2, \tag{103}$$

where $c \geq 0$ is a universal constant. Putting together (101) to (103) completes the proof. It remains to prove our claims (102) and (103) which we now do one-by-one.

Proof of bound (102). We start by bounding $\left(\mathbb{E}_{\mathbf{x}|\mathbf{g}=\boldsymbol{g}} \left[(\widetilde{\nabla}q(\mathbf{x})^\top \boldsymbol{g} - \mathbb{E}_{\mathbf{x}|\mathbf{g}=\boldsymbol{g}} [\widetilde{\nabla}q(\mathbf{x})^\top \boldsymbol{g}])^p \right] \right)^{1/p}$ for every $\mathbf{g} = \boldsymbol{g}$. Then, we proceed to bound $\|\widetilde{\nabla}q(\mathbf{x})^\top \mathbf{g} - \mathbb{E}_{\mathbf{x}} [\widetilde{\nabla}q(\mathbf{x})^\top \mathbf{g}]\|_{L_p}$.

To that end, we define $h_{\boldsymbol{g}}(\mathbf{x}) \triangleq \widetilde{\nabla}q(\mathbf{x})^\top \boldsymbol{g} - \mathbb{E}_{\mathbf{x}|\mathbf{g}=\boldsymbol{g}} [\widetilde{\nabla}q(\mathbf{x})^\top \boldsymbol{g}]$ and observe that $\mathbb{E}_{\mathbf{x}|\mathbf{g}=\boldsymbol{g}} [h_{\boldsymbol{g}}(\mathbf{x})] = 0$. Now, applying Lem. 16 to $h_{\boldsymbol{g}}(\cdot)$, we have

$$\begin{aligned}
\|h_{\boldsymbol{g}}(\mathbf{x})\|_{L_p} \leq \sigma\sqrt{2p} \left(\mathbb{E}_{\mathbf{x}|\mathbf{g}=\boldsymbol{g}} \left[\|\nabla h_{\boldsymbol{g}}(\mathbf{x})\|_2^p \right] \right)^{1/p} &\overset{(a)}{\leq} \sigma\sqrt{2p} \left(\mathbb{E}_{\mathbf{x}|\mathbf{g}=\boldsymbol{g}} \left[\left\| \nabla \left[\boldsymbol{g}^\top \widetilde{\nabla}q(\mathbf{x}) \right] \right\|_2^p \right] \right)^{1/p} \\
&\overset{(67)}{\leq} \sigma\sqrt{2p} \left(\mathbb{E}_{\mathbf{x}|\mathbf{g}=\boldsymbol{g}} \left[\left\| \boldsymbol{g}^\top \widetilde{\nabla}^2 q(\mathbf{x}) \right\|_2^p \right] \right)^{1/p},
\end{aligned} \tag{104}$$

where (a) follows from the definition of $h_{\boldsymbol{g}}(\mathbf{x})$. Now, to obtain a bound on the RHS of (104), we further fix $\mathbf{x} = \boldsymbol{x}$. Then, we let $\mathbf{g}'$ be another p-dimensional standard normal vector and apply an inequality similar to (100) to $\boldsymbol{g}^\top \widetilde{\nabla}^2 q(\boldsymbol{x})$ obtaining

$$\left\| \boldsymbol{g}^\top \widetilde{\nabla}^2 q(\boldsymbol{x}) \right\|_2 \leq \frac{2}{\sqrt{p}} \left(\mathbb{E}_{\mathbf{g}'|\mathbf{x}=\boldsymbol{x},\mathbf{g}=\boldsymbol{g}} \left[\left(\boldsymbol{g}^\top \widetilde{\nabla}^2 q(\boldsymbol{x})\mathbf{g}' \right)^p \right] \right)^{1/p},$$

which implies

$$\left(\mathbb{E}_{\mathbf{x}|\mathbf{g}=g}\left[\left\|g^{\top}\widetilde{\nabla}^2 q(\mathbf{x})\right\|_2^p\right]\right)^{1/p} \leq \frac{2}{\sqrt{p}}\left(\mathbb{E}_{\mathbf{x},\mathbf{g}'|\mathbf{g}=g}\left[\left(\nabla g^{\top}\widetilde{\nabla}^2 q(\mathbf{x})\mathbf{g}'\right)^p\right]\right)^{1/p}. \tag{105}$$

Putting together (104) and (105), and using the definition of $h_g(\mathbf{x})$, we have

$$\left(\mathbb{E}_{\mathbf{x}|\mathbf{g}=g}\left[\left(\widetilde{\nabla}q(\mathbf{x})^{\top}g - \mathbb{E}_{\mathbf{x}|\mathbf{g}=g}\left[\widetilde{\nabla}q(\mathbf{x})^{\top}g\right]\right)^p\right]\right)^{1/p} \leq 2\sqrt{2}\sigma\left(\mathbb{E}_{\mathbf{x},\mathbf{g}'|\mathbf{g}=g}\left[\left(g^{\top}\widetilde{\nabla}^2 q(\mathbf{x})\mathbf{g}'\right)^p\right]\right)^{1/p}. \tag{106}$$

Now, we proceed to bound $\|\widetilde{\nabla}q(\mathbf{x})^{\top}\mathbf{g} - \mathbb{E}_{\mathbf{x}}[\widetilde{\nabla}q(\mathbf{x})^{\top}\mathbf{g}]\|_{L_p}$ as follows:

$$\left\|\widetilde{\nabla}q(\mathbf{x})^{\top}\mathbf{g} - \mathbb{E}_{\mathbf{x}}\left[\widetilde{\nabla}q(\mathbf{x})^{\top}\mathbf{g}\right]\right\|_{L_p} \overset{(a)}{=} \left(\mathbb{E}_{\mathbf{x},\mathbf{g}}\left[\left(\widetilde{\nabla}q(\mathbf{x})^{\top}\mathbf{g} - \mathbb{E}_{\mathbf{x}}\left[\widetilde{\nabla}q(\mathbf{x})^{\top}\mathbf{g}\right]\right)^p\right]\right)^{1/p}$$
$$\overset{(106)}{\leq} 2\sqrt{2}\sigma\left(\mathbb{E}_{\mathbf{g},\mathbf{x},\mathbf{g}'}\left[\left(\mathbf{g}^{\top}\widetilde{\nabla}^2 q(\mathbf{x})\mathbf{g}'\right)^p\right]\right)^{1/p}, \tag{107}$$

where (a) follows from the definition of p-th moment. Finally, to bound the RHS of (107), we fix $\mathbf{x} = x$ and bound the p-th norm of the quadratic form $\mathbf{g}^{\top}\widetilde{\nabla}^2 q(x)\mathbf{g}'$ by the Hanson-Wright inequality resulting in

$$\left(\mathbb{E}_{\mathbf{g},\mathbf{g}'|\mathbf{x}=x}\left[\left(\mathbf{g}^{\top}\widetilde{\nabla}^2 q(x)\mathbf{g}'\right)^p\right]\right)^{1/p} \leq c\left(\sqrt{p}\|\widetilde{\nabla}^2 q(x)\|_{\mathrm{F}} + p\|\widetilde{\nabla}^2 q(x)\|_{\mathrm{op}}\right)$$
$$\leq c\left(\sqrt{p}\max_{x\in\mathcal{X}^p}\|\widetilde{\nabla}^2 q(x)\|_{\mathrm{F}} + p\max_{x\in\mathcal{X}^p}\|\widetilde{\nabla}^2 q(x)\|_{\mathrm{op}}\right), \tag{108}$$

where $c \geq 0$ is a universal constant. Then, (102) follows by putting together (107) and (108).

Proof of bound (103). By linearity of expectation, we have

$$\|\mathbb{E}_{\mathbf{x}}[\widetilde{\nabla}q(\mathbf{x})^{\top}\mathbf{g}]\|_{L_p} = \|(\mathbb{E}_{\mathbf{x}}[\widetilde{\nabla}q(\mathbf{x})])^{\top}\mathbf{g}\|_{L_p}. \tag{109}$$

We note that the random variable $\dfrac{(\mathbb{E}_{\mathbf{x}}[\widetilde{\nabla}q(\mathbf{x})])^{\top}\mathbf{g}}{\|\mathbb{E}_{\mathbf{x}}[\widetilde{\nabla}q(\mathbf{x})]\|_2}$ is a standard normal random variable. Therefore,

$$\left\|\frac{(\mathbb{E}_{\mathbf{x}}[\widetilde{\nabla}q(\mathbf{x})])^{\top}\mathbf{g}}{\|\mathbb{E}_{\mathbf{x}}[\widetilde{\nabla}q(\mathbf{x})]\|_2}\right\|_{L_p} \overset{(a)}{=} \left(\mathbb{E}_{\mathbf{g}}\left[\left(\frac{(\mathbb{E}_{\mathbf{x}}[\widetilde{\nabla}q(\mathbf{x})])^{\top}\mathbf{g}}{\|\mathbb{E}_{\mathbf{x}}[\widetilde{\nabla}q(\mathbf{x})]\|_2}\right)^p\right]\right)^{1/p} \overset{(b)}{\leq} 2\sqrt{p}, \tag{110}$$

where (a) follows from the definition of p-th moment, and (b) follows since $\|g\|_{L_p} \leq 2\sqrt{p}$ for any standard normal variable g. Then, (103) follows by using (110) in (109).

E The conditioning trick

In App. D, we derived that a random vector (supported on a compact set) satisfies the logarithmic Sobolev inequality if it satisfies the Dobrushin's uniqueness condition. As a result, to use the tail bound in App. D, we need the random vector to satisfy the Dobrushin's uniqueness condition. In this section, we show that any random vector (outside Dobrushin's regime) that is a τ-Sparse Graphical Model can be reduced to satisfy the Dobrushin's uniqueness condition. In particular, we show that by conditioning on a subset of the random vector, the unconditioned subset of the random vector (in the conditional distribution) satisfies the Dobrushin's uniqueness condition. We exploit this conditioning trick throughout Lem. 6 to enable application of the tail bound in App. D. The result below is a generalization of the result in [12] to continuous random vectors.

We start by defining the notion of τ-Sparse Graphical Model.

Definition 8 (τ-**Sparse Graphical Model**). *A pair of random vectors $\{\mathbf{x}, \mathbf{z}\}$ supported on $\mathcal{X}^p \times \mathcal{Z}^{p_z}$ is a τ-Sparse Graphical Model for model-parameters $\tau \triangleq (\alpha, \zeta, x_{\max}, \Theta)$ and denoted by τ-SGM if $\mathcal{X} = \{-x_{\max}, x_{\max}\}$, and*

965   *1. for any realization $z \in \mathcal{Z}^{p_z}$, the conditional probability distribution of $\mathbf{x}$ given $\mathbf{z} = z$ is*

966   *given by $f_{\mathbf{x}|\mathbf{z}}(\,\cdot\,|z; \theta(z), \Theta)$ in (4) for a vector $\theta(z) \in \mathbb{R}^p$ depending on z and a symmetric*

967   *matrix $\Theta \in \mathbb{R}^{p \times p}$ (independent of z) with $\Theta_{tt}^\star = 0$ for all $t \in [p]$,*

968   *2. $\max\left\{\max_{z \in \mathcal{Z}^{p_z}} \|\theta(z)\|_\infty, \|\Theta\|_{\max}\right\} \leq \alpha$, and*

969   *3. $\|\Theta\|_\infty \leq \zeta$.*

970 Now, we provide the main result of this section.

971 **Lemma 18** (**The conditioning trick**). *Given a pair of random vectors $\{\mathbf{x}, \mathbf{z}\}$ supported on $\mathcal{X}^p \times \mathcal{Z}^{p_z}$*

972 *that is a τ-SGM (Def. 8) with $\tau \triangleq (\alpha, \zeta, x_{\max}, \Theta)$, and a scalar $\lambda \in (0, \zeta]$, there exists $L \triangleq$*

973 *$32\zeta^2 \log 4p/\lambda^2$ subsets $S_1, \cdots, S_L \subseteq [p]$ that satisfy the following properties:*

974   *(a) For any $t \in [p]$, we have $\sum_{u=1}^L \mathbb{1}(t \in S_u) = \lceil \lambda L/(8\zeta) \rceil$.*

975   *(b) For any $u \in [L]$,*

976   *(i) the pair of random vectors $\{\mathbf{x}_{S_u}, (\mathbf{x}_{-S_u}, \mathbf{z})\}$ correspond to a τ_1-SIM with $\tau_1 \triangleq (\alpha +$*

977   *$x_{\max}\zeta, \lambda, x_{\max}, \Theta_{S_u})$ where $\Theta_{S_u} \triangleq \{\Theta_{tv}\}_{t,v \in S_u}$, and*

978   *(ii) the random vector $\mathbf{x}_{S_u}$ conditioned on $(\mathbf{x}_{-S_u}, \mathbf{z})$ satisfies the Dobrushin's uniqueness*

979   *condition (Def. 4) with coupling matrix $\sqrt{2}x_{\max}^2 \Theta_{S_u}$ whenever $\lambda \in \left(0, \frac{1}{\sqrt{2}x_{\max}^2}\right]$.*

980 *Proof of Lem. 18: The conditioning trick.* We prove each part one-by-one.

981 **Proof of part (a)** From Def. 8, for any realization $z \in \mathcal{Z}^{p_z}$, the conditional probability distribution

982 of $\mathbf{x}$ given $\mathbf{z} = z$ is given by $f_{\mathbf{x}|\mathbf{z}}(\,\cdot\,|z; \theta(z), \Theta)$ in (4) where $\theta(z) \in \mathbb{R}^p$ is a vector and $\Theta \in \mathbb{R}^{p \times p}$

983 is a symmetric matrix with $\Theta_{tt} = 0$ for all $t \in [p]$ and $\|\Theta\|_\infty \leq \zeta$. Consider the matrix $A \triangleq \frac{1}{\zeta}\Theta$.

984 Since A has zeros on the diagonal and $\|A\|_\infty \leq 1$, we can apply [12, Lem. 12] on A with $\eta = \frac{\lambda}{\zeta}$.

985 Then part (a) follows directly from [12, Lem. 12.1].

986 **Proof of part (b) (i)** To prove this part, consider the conditional distribution of $\mathbf{x}_{S_u}$ conditioned on

987 $\mathbf{x}_{-S_u} = \boldsymbol{x}_{-S_u}$ and $\mathbf{z} = \boldsymbol{z}$ for any $u \in [L]$. We have

$$f_{\mathbf{x}_{S_u}|\mathbf{x}_{-S_u},\mathbf{z}}(\boldsymbol{x}_{S_u}|\boldsymbol{x}_{-S_u}, \boldsymbol{z}; \theta(\boldsymbol{z}), \Theta) \propto \exp\left(\sum_{t \in S_u}\left(\theta_t(\boldsymbol{z}) + \sum_{v \notin S_u}\Theta_{tv}x_v\right)x_t + \sum_{t \in S_u}\sum_{v \in S_u}\Theta_{tv}x_t x_v\right).$$

(111)

988 We can re-parameterize $f_{\mathbf{x}_{S_u}|\mathbf{x}_{-S_u},\mathbf{z}}(\boldsymbol{x}_{S_u}|\boldsymbol{x}_{-S_u}, \boldsymbol{z}; \theta(\boldsymbol{z}), \Theta)$ in (111) as follows

$$f_{\mathbf{x}_{S_u}|\mathbf{x}_{-S_u},\mathbf{z}}(\boldsymbol{x}_{S_u}|\boldsymbol{x}_{-S_u}, \boldsymbol{z}; \upsilon(\boldsymbol{z}, \boldsymbol{x}_{-S_u}), \Upsilon) \propto \exp\left([\upsilon(\boldsymbol{z}, \boldsymbol{x}_{-S_u})]^\top \boldsymbol{x}_{S_u} + \boldsymbol{x}_{S_u}^\top \Upsilon \boldsymbol{x}_{S_u}\right), \quad \text{where}$$

$$\upsilon(\boldsymbol{z}, \boldsymbol{x}_{-S_u}) \in \mathbb{R}^{|S_u| \times 1}, \quad \text{with} \quad \upsilon_t(\boldsymbol{z}, \boldsymbol{x}_{-S_u}) \triangleq \theta_t(\boldsymbol{z}) + \sum_{k \notin S_u}\Theta_{tv}x_k \quad \text{for} \quad t \in S_u, \quad \text{and} \quad (112)$$

$$\Upsilon = \Upsilon^\top \in \mathbb{R}^{|S_u| \times |S_u|} \quad \text{with} \quad \Upsilon_{tv} \triangleq \Theta_{tv}, \text{ and } \Upsilon_{tt} = 0 \text{ for all } t, v \in S_u. \tag{113}$$

989 Thus, to show that the random vector $\mathbf{x}_{S_u}$ conditioned on $\mathbf{x}_{-S_u}$ and $\mathbf{z}$ corresponds to an τ_1-SGM with

990 $\tau_1 \triangleq (\alpha + x_{\max}\zeta, \lambda, x_{\max}, \Theta_{S_u})$, it suffices to establish that

$$\max\left\{\max_{\boldsymbol{z} \in \mathcal{Z}^{p_z}}\|\upsilon(\boldsymbol{z}, \boldsymbol{x}_{-S_u})\|_\infty, \|\Upsilon\|_{\max}\right\} \overset{(i)}{\leq} \alpha + x_{\max}\zeta \quad \text{and} \quad \|\Upsilon\|_\infty \overset{(ii)}{\leq} \lambda. \tag{114}$$

991 To establish (i) in (114), we note that

$$\|\Upsilon\|_{\max} \overset{(113)}{\leq} \|\Theta\|_{\max} \overset{(a)}{\leq} \alpha \quad \text{and} \tag{115}$$

$$\|\upsilon(\boldsymbol{z}, \boldsymbol{x}_{-S_u})\|_\infty \overset{(b)}{\leq} \|\theta(\boldsymbol{z})\|_\infty + \max_{t \in S_u}\|\Theta_t\|_1 \|\boldsymbol{x}\|_\infty \overset{(c)}{\leq} \|\theta(\boldsymbol{z})\|_\infty + x_{\max}\|\Theta\|_\infty \overset{(d)}{\leq} \alpha + x_{\max}\zeta,$$

(116)

where (a) and (d) follow from Def. 8, (b) follows from (112) and the triangle inequality, and (c) follows from the definition of $\|\cdot\|_\infty$ and Def. 8. Then, from (115) and (116), we have

$$\max\left\{ \max_{\boldsymbol{z}\in\mathcal{Z}^{p_z}} \|\upsilon(\boldsymbol{z},\boldsymbol{x}_{-S_u})\|_\infty, \|\!|\Upsilon|\!\|_{\max} \right\} \le \alpha + x_{\max}\zeta,$$

as claimed. Next, to establish (ii) in (114), we again apply [12, Lem. 12] on the matrix $A = \frac{1}{\zeta}\Theta$ with $\eta = \frac{\lambda}{\zeta}$. Then, [12, Lem. 12.2] implies that

$$\sum_{v\in S_u} \left| \frac{\Theta_{tv}}{\zeta} \right| \le \frac{\lambda}{\zeta} \quad \text{for all } t\in S_u,\, u\in[L]. \tag{117}$$

Therefore, we have

$$\|\!|\Upsilon|\!\|_\infty = \max_{t\in S_u}\Big(\sum_{v\in S_u} |\Upsilon_{tv}| \Big) \stackrel{(113)}{=} \max_{t\in S_u}\Big(\sum_{v\in S_u} |\Theta_{tv}| \Big) \stackrel{(117)}{\le} \lambda, \tag{118}$$

as desired. The proof for this part is now complete.

Proof of part (b) (ii)   From the analysis in part (b) (i), for any $u\in S_u$, we have

$$\|\!|\Theta_{S_u}|\!\|_\infty \stackrel{(113)}{=} \|\!|\Upsilon|\!\|_\infty \stackrel{(118)}{\le} \lambda.$$

Therefore, $\|\!|\sqrt{2}x_{\max}^2\Theta_{S_u}|\!\|_\infty \le 1$ whenever $\lambda \le 1/\sqrt{2}x_{\max}^2$. It remains to show that for every $u\in[L]$, $t\in S_u$, $v\in S_u\setminus\{t\}$, $\boldsymbol{z}=\boldsymbol{z}$, and $\boldsymbol{x}_{-t},\widetilde{\boldsymbol{x}}_{-t}\in\mathcal{X}^{p-1}$ differing only in the v^{th} coordinate,

$$\|f_{\mathsf{x}_t|\mathbf{x}_{-t}=\boldsymbol{x}_{-t},\mathbf{z}=\boldsymbol{z}} - f_{\mathsf{x}_t|\mathbf{x}_{-t}=\widetilde{\boldsymbol{x}}_{-t},\mathbf{z}=\boldsymbol{z}}\|_{\mathsf{TV}} \le \sqrt{2}x_{\max}^2\Theta_{tv}.$$

To that end, fix any $u\in[L]$, any $t\in S_u$, any $v\in S_u\setminus\{t\}$, any $\mathbf{z}=\boldsymbol{z}$, and any $\boldsymbol{x}_{-t},\widetilde{\boldsymbol{x}}_{-t}\in\mathcal{X}^{p-1}$ differing only in the v^{th} coordinate. We have

$$
\begin{aligned}
\|f_{\mathsf{x}_t|\mathbf{x}_{-t}=\boldsymbol{x}_{-t},\mathbf{z}=\boldsymbol{z}} - f_{\mathsf{x}_t|\mathbf{x}_{-t}=\widetilde{\boldsymbol{x}}_{-t},\mathbf{z}=\boldsymbol{z}}\|_{\mathsf{TV}}^2 &\stackrel{(a)}{\le} \frac{1}{2}\mathsf{KL}\left(f_{\mathsf{x}_t|\mathbf{x}_{-t}=\boldsymbol{x}_{-t},\mathbf{z}=\boldsymbol{z}} \,\big\|\, f_{\mathsf{x}_t|\mathbf{x}_{-t}=\widetilde{\boldsymbol{x}}_{-t},\mathbf{z}=\boldsymbol{z}} \right) \\
&\stackrel{(b)}{=} \frac{1}{2}\Theta_{tv}^2(x_v-\widetilde{x}_v)^2 \mathbb{C}\mathrm{ov}(\mathsf{x}_v,\mathsf{x}_v) \\
&\stackrel{(c)}{\le} 2x_{\max}^4\Theta_{tv}^2,
\end{aligned}
$$

where (a) follows from Pinsker's inequality, (b) follows by applying [11, Theorem 1] to the conditional distribution $f_{\mathsf{x}_t|\mathbf{x}_{-t},\mathbf{z}}$ (see (8)) and because for any regular exponential family the Hessian of the log partition function is the covariance matrix of the associated sufficient statistic, and (c) follows from Def. 8. This completes the proof. $\square$

F   Supporting concentration results

In this section, we provide a corollary of Prop. 2 that is used to prove the concentration results in Lem. 10 and Lem. 12. To show any concentration result for the random vector $\mathbf{x}$ conditioned on $\mathbf{z}$ via Prop. 2, we need $\mathbf{x}|\mathbf{z}$ to satisfy the logarithmic Sobolev inequality (see (64)). From Prop. 1, for this to be true, we need the random vector x_t conditioned on $(\mathbf{x}_{-t},\mathbf{z})$ to satisfy the logarithmic Sobolev inequality for all $t\in[p]$. In the result below, we show this holds with a proof in App. F.1.

Lemma 19 (**Logarithmic Sobolev inequality for $\mathsf{x}_t|\mathbf{x}_{-t},\mathbf{z}$**). *Given a pair of random vectors $\{\mathbf{x},\mathbf{z}\}$ supported on $\mathcal{X}^p\times\mathcal{Z}^{p_z}$ that is a τ-SGM (Def. 8) with $\tau\triangleq(\alpha,\zeta,x_{\max},\Theta)$, $\mathsf{x}_t|\mathbf{x}_{-t},\mathbf{z}$ satisfies $\mathrm{LSI}_{\mathsf{x}_t|\mathbf{x}_{-t}=\boldsymbol{x}_{-t},\mathbf{z}=\boldsymbol{z}}\left(\frac{8x_{\max}^2}{\pi^2}\exp\left(2x_{\max}(\alpha+\zeta x_{\max})\right) \right)$ for all $t\in[p]$, $\boldsymbol{x}_{-t}\in\mathcal{X}^{p-1}$, and $\boldsymbol{z}\in\mathcal{Z}^{p_z}$.*

Now, we state the desired corollary of Prop. 2 with a proof in App. F.2. The corollary makes use of some $\tau\triangleq(\alpha,\zeta,x_{\max},\Theta)$-dependent constants:

$$C_{6,\tau}\triangleq 1+\alpha x_{\max}+2x_{\max}^2\zeta,\;\; C_{7,\tau}\triangleq\exp\left(x_{\max}(\alpha+\zeta x_{\max})\right),\text{ and }C_{8,\tau}\triangleq\frac{32x_{\max}^3 C_{7,\tau}^4}{\pi^2}. \tag{119}$$

Corollary 1. *Suppose a pair of random vectors $\{\mathbf{x}, \mathbf{z}\}$ supported on $\mathcal{X}^p \times \mathcal{Z}^{p_z}$ corresponds to a τ-SGM (Def. 8) with $\tau \triangleq (\alpha, \zeta, x_{\max}, \Theta)$, and the random vector $\mathbf{x}$ conditioned on $\mathbf{z}$ satisfy the Dobrushin's uniqueness condition (Def. 4) with coupling matrix $\overline{\Theta}$. Define the functions q_1 and q_2 as*

$$q_1(\mathbf{x}) \triangleq \sum_{t \in [p]} (\omega_t x_t)^2 \quad and \quad q_2(\mathbf{x}) \triangleq \sum_{t \in [p]} \omega_t x_t \exp\Big(-[\theta_t + \Theta_t^\top \mathbf{x}] x_t \Big).$$

Then for any $\theta, \overline{\theta} \in \Lambda_\theta$, $\Theta \in \Lambda_\Theta$, and $\varepsilon > 0$

$$\mathbb{P}\Big[\big| q_i(\mathbf{x}) - \mathbb{E}\big[q_i(\mathbf{x})\big|\mathbf{z}\big] \big| \geq \varepsilon \Big| \mathbf{z} \Big] \leq \exp\left(\frac{-c\big(1 - \|\overline{\Theta}\|_{\mathrm{op}}\big)^2 \varepsilon^2}{c_i \|\omega\|_2^2} \right) \quad for \quad i = 1, 2, \tag{120}$$

where $\omega = \overline{\theta} - \theta$, c is a universal constant, $c_1 \triangleq 16\alpha^2 x_{\max}^2 C_{8,\tau}$, and $c_2 \triangleq C_{6,\tau}^2 C_{7,\tau}^2 C_{8,\tau}$.

F.1 Proof of Lem. 19: Logarithmic Sobolev inequality for $x_t | \mathbf{x}_{-t}, \mathbf{z}$

Let u be the uniform distribution on $\mathcal{X}$. Then, u satisfies $\mathrm{LSI}_u\left(\frac{8x_{\max}^2}{\pi^2}\right)$ (see [14, Corollary 2.4]). Then, using the Holley-Stroock perturbation principle (see [16, Page 31], [19, Lemma 1.2]), for every $t \in [p]$, $\boldsymbol{x}_{-t} \in \mathcal{X}^{p-1}$, and $\boldsymbol{z} \in \mathcal{Z}^{p_z}$, $x_t | \mathbf{x}_{-t} = \boldsymbol{x}_{-t}, \mathbf{z} = \boldsymbol{z}$ satisfies the logarithmic Sobolev inequality with a constant bounded by

$$\frac{8x_{\max}^2 \exp(\sup_{x_t \in \mathcal{X}} \psi(x_t; \boldsymbol{x}_{-t}, \boldsymbol{z}) - \inf_{x_t \in \mathcal{X}} \psi(x_t; \boldsymbol{x}_{-t}, \boldsymbol{z}))}{\pi^2},$$

where $\psi(x_t; \boldsymbol{x}_{-t}, \boldsymbol{z}) \triangleq -[\theta_t(\boldsymbol{z}) + \Theta_t^\top \boldsymbol{x}] x_t$. We have

$$\frac{8x_{\max}^2 \exp(\sup_{x_t \in \mathcal{X}} \psi(x_t; \boldsymbol{x}_{-t}, \boldsymbol{z}) - \inf_{x_t \in \mathcal{X}} \psi(x_t; \boldsymbol{x}_{-t}, \boldsymbol{z}))}{\pi^2}$$

$$\overset{(a)}{=} \frac{8x_{\max}^2}{\pi^2} \exp\big(2|\theta_t(\boldsymbol{z}) + \Theta_t^\top \boldsymbol{x}| x_{\max}\big) \overset{(b)}{\leq} \frac{8x_{\max}^2}{\pi^2} \exp(2\alpha x_{\max} + 2\zeta x_{\max}^2) \overset{(119)}{=} \frac{8x_{\max}^2 C_{7,\tau}^2}{\pi^2},$$

where (a) and (b) follow from Def. 8.

F.2 Proof of Cor. 1

To apply Prop. 2 to the random vector $\mathbf{x}$ conditioned on $\mathbf{z}$, we need $\mathbf{x}|\mathbf{z}$ to satisfy the logarithmic Sobolev inequality. From Prop. 1, this is true if (i) $f_{\min} = \min_{t \in [p], \boldsymbol{x} \in \mathcal{X}^p, \boldsymbol{z} \in \mathcal{X}^{p_z}} f_{x_t | \mathbf{x}_{-t}, \mathbf{z}}(x_t | \boldsymbol{x}_{-t}, \boldsymbol{z}) > 0$ (see (66)), (ii) $\mathbf{x}|\mathbf{z}$ satisfies the Dobrushin's uniqueness condition, and (iii) $x_t | \mathbf{x}_{-t}, \mathbf{z}$ satisfies the logarithmic Sobolev inequality for all $t \in [p]$. By assumption, $\mathbf{x}|\mathbf{z}$ satisfies the Dobrushin's uniqueness condition with coupling matrix $\overline{\Theta}$. From Lem. 19, $x_t | \mathbf{x}_{-t}, \mathbf{z}$ satisfies $\mathrm{LSI}_{x_t | \mathbf{x}_{-t} = \boldsymbol{x}_{-t}, \mathbf{z} = \boldsymbol{z}}\left(\frac{8x_{\max}^2 C_{7,\tau}^2}{\pi^2}\right)$. It remains to show (i). Consider any $t \in [p]$, any $\boldsymbol{x} \in \mathcal{X}^p$, and any $\boldsymbol{z} \in \mathcal{X}^{p_z}$. We have

$$f_{x_t | \mathbf{x}_{-t}, \mathbf{z}}(x_t | \boldsymbol{x}_{-t}, \boldsymbol{z}) \overset{(a)}{=} \frac{\exp\Big([\theta_t(\boldsymbol{z}) + \Theta_t^\top \boldsymbol{x}] x_t\Big)}{\int_{\mathcal{X}} \exp\Big([\theta_t(\boldsymbol{z}) + \Theta_t^\top \boldsymbol{x}] x_t\Big) dx_t}$$

$$\overset{(b)}{\geq} \frac{\exp\Big(-|\theta_t(\boldsymbol{z}) + \Theta_t^\top \boldsymbol{x}| x_{\max}\Big)}{\int_{\mathcal{X}} \exp\Big(|\theta_t(\boldsymbol{z}) + \Theta_t^\top \boldsymbol{x}| x_{\max}\Big) dx_t}$$

$$\overset{(c)}{\geq} \frac{\exp\Big(-\big(|\theta(\boldsymbol{z})| + \|\Theta_t\|_1 \|\boldsymbol{x}\|_\infty\big) x_{\max}\Big)}{\int_{\mathcal{X}} \exp\Big(\big(|\theta(\boldsymbol{z})| + \|\Theta_t\|_1 \|\boldsymbol{x}\|_\infty\big) x_{\max}\Big) dx_t}$$

$$\overset{(d)}{\geq} \frac{\exp\Big(-(\alpha + \zeta x_{\max}) x_{\max}\Big)}{\int_{\mathcal{X}} \exp\Big((\alpha + \zeta x_{\max}) x_{\max}\Big) dx_t} \overset{(e)}{=} \frac{1}{2x_{\max} C_{7,\tau}^2},$$

where (a) follows from (8) and Def. 8, (b) and (d) follow from Def. 8, (c) follows by triangle inequality and Cauchy–Schwarz inequality, and (e) follows because $\int_{\mathcal{X}} dx_t = 2x_{\max}$. Therefore, $f_{\min} = \frac{1}{2x_{\max} C_{7,\tau}^2}$. Putting (i), (ii), and (iii) together, $\mathsf{x}|\mathsf{z}$ satisfies $\mathrm{LSI}_{\mathsf{x}}\left(\frac{C_{8,\tau}}{(1-\|\overline{\Theta}\|_{\mathrm{op}})^2}\right)$ from Prop. 1. Now, we apply Prop. 2 to q_1 and q_2 one-by-one. The general strategy is to choose appropriate pseudo derivatives and pseudo Hessians for both q_1 and q_2, and evaluate the corresponding terms appearing in Prop. 2.

Concentration for q_1.   Fix any $\boldsymbol{x} \in \mathcal{X}^p$. We start by decomposing $q_1(\boldsymbol{x})$ as follows:

$$q_1(\boldsymbol{x}) = \overline{\omega}^\top r(\boldsymbol{x}), \tag{121}$$

where $\overline{\omega} \triangleq (\omega_1^2, \cdots, \omega_p^2)$ and $r(\boldsymbol{x}) \triangleq (r_1(\boldsymbol{x}), \cdots, r_p(\boldsymbol{x}))$ with $r_t(\boldsymbol{x}) = x_t^2$ for every $t \in [p]$. Next, we define $H : \mathcal{X}^p \to \mathbb{R}^{p \times p}$ such that

$$H_{tu}(\boldsymbol{x}) = \frac{dq_u(\boldsymbol{x})}{dx_t} \quad \text{for every } t, u \in [p]. \tag{122}$$

Pseudo derivative.   We bound the ℓ_2 norm of the gradient of $q_1(\boldsymbol{x})$ as follows:

$$\|\nabla q_1(\boldsymbol{x})\|_2^2 = \sum_{t \in [p]} \left(\frac{dq_1(\boldsymbol{x})}{dx_t}\right)^2 \overset{(121)}{=} \sum_{t \in [p]} \left(\frac{\overline{\omega}^\top dr(\boldsymbol{x})}{dx_t}\right)^2 \overset{(122)}{=} \|H(\boldsymbol{x})\overline{\omega}\|_2^2 \overset{(a)}{\leq} \|H(\boldsymbol{x})\|_{\mathrm{op}}^2 \|\overline{\omega}\|_2^2$$

$$\overset{(b)}{\leq} \|H(\boldsymbol{x})\|_1 \|H(\boldsymbol{x})\|_\infty \|\overline{\omega}\|_2^2, \tag{123}$$

where (a) follows because induced matrix norms are submultiplicative and (b) follows because the matrix operator norm is bounded by square root of the product of matrix one norm and matrix infinity norm. Now, we claim that the one norm and the infinity norm of $H(\boldsymbol{x})$ are bounded as follows:

$$\max\left\{\max_{\boldsymbol{x} \in \mathcal{X}^p} \|H(\boldsymbol{x})\|_1, \max_{\boldsymbol{x} \in \mathcal{X}^p} \|H(\boldsymbol{x})\|_\infty\right\} \leq 2x_{\max}. \tag{124}$$

Taking this claim as given at the moment, we continue with our proof. Combining (123) and (124), we have

$$\max_{\boldsymbol{x} \in \mathcal{X}^p} \|\nabla q_1(\boldsymbol{x})\|_2^2 \leq 4x_{\max}^2 \|\overline{\omega}\|_2^2 = 4x_{\max}^2 \sum_{t \in [p]} \omega_t^4 \leq 4x_{\max}^2 \max_{u \in [p]} \omega_u^2 \sum_{t \in [p]} \omega_t^2 \overset{(a)}{\leq} 16x_{\max}^2 \alpha^2 \|\omega\|_2^2,$$

where (a) follows because $\omega \in 2\Lambda_\theta$. Therefore, we choose the pseudo derivative (see Def. 5) as follows:

$$\widetilde{\nabla} q_1(\boldsymbol{x}) = 4x_{\max}\alpha \|\omega\|_2. \tag{125}$$

Pseudo Hessian.   Fix any $\rho \in \mathbb{R}$. We bound $\|\nabla(\rho^\top \widetilde{\nabla} q_1(\boldsymbol{x}))\|_2^2$ (see Def. 5) as follows:

$$\|\nabla(\rho^\top \widetilde{\nabla} q_1(\boldsymbol{x}))\|_2^2 = \sum_{u \in [p]} \left(\frac{d\rho^\top \widetilde{\nabla} q_1(\boldsymbol{x})}{dx_u}\right)^2 \overset{(125)}{=} 0.$$

Therefore, we choose the pseudo Hessian (see Def. 5) as follows:

$$\widetilde{\nabla}^2 q_1(\boldsymbol{x}) = 0. \tag{126}$$

The claimed result now follows by applying Prop. 2 with the pseudo discrete derivative defined in (125), and the pseudo discrete Hessian defined in (126).

It remains to show that the one-norm and the infinity-norm of $H(\boldsymbol{x})$ are bounded as in (124).

Bounds on the one-norm and the infinity-norm of $H(\boldsymbol{x})$. We have

$$H_{tu}(\boldsymbol{x}) = \begin{cases} 2x_t & \text{if} \quad t = u, \\ 0 & \text{otherwise.} \end{cases} \tag{127}$$

Therefore,

$$\|H(\boldsymbol{x})\|_1 = \max_{u \in [p]} \sum_{t \in [p]} |H_{tu}(\boldsymbol{x})| \overset{(127)}{\leq} \max_{u \in [p]} 2|x_u| \overset{(a)}{\leq} 2x_{\max} \text{ and}$$

$$\|H(\boldsymbol{x})\|_\infty = \max_{t \in [p]} \sum_{u \in [p]} |H_{tu}(\boldsymbol{x})| \overset{(127)}{\leq} \max_{t \in [p]} 2|x_t| \overset{(a)}{\leq} 2x_{\max},$$

where (a) follows from Def. 8.

Concentration for q_2. Fix any $\boldsymbol{x} \in \mathcal{X}^p$. We start by decomposing $q_2(\boldsymbol{x})$ as follows:

$$q_2(\boldsymbol{x}) = \omega^\top r(\boldsymbol{x}), \tag{128}$$

where $r(\boldsymbol{x}) \triangleq (r_1(\boldsymbol{x}), \cdots, r_p(\boldsymbol{x}))$ with $r_t(\boldsymbol{x}) = x_t \exp\left(-[\theta_t + \Theta_t^\top \boldsymbol{x}]x_t\right)$ for every $t \in [p]$. Next, we define $H : \mathcal{X}^p \to \mathbb{R}^{p \times p}$ such that

$$H_{tu}(\boldsymbol{x}) = \frac{dr_u(\boldsymbol{x})}{dx_t} \quad \text{for every } t, u \in [p]. \tag{129}$$

Pseudo derivative. We bound the ℓ_2 norm of the gradient of $q_2(\boldsymbol{x})$ as follows:

$$\|\nabla q_2(\boldsymbol{x})\|_2^2 = \sum_{t \in [p]} \left(\frac{dq_2(\boldsymbol{x})}{dx_t}\right)^2 \overset{(128)}{=} \sum_{t \in [p]} \left(\frac{\omega^\top dr(\boldsymbol{x})}{dx_t}\right)^2 \overset{(129)}{=} \|H(\boldsymbol{x})\omega\|_2^2 \overset{(a)}{\leq} \|H(\boldsymbol{x})\|_{\text{op}}^2 \|\omega\|_2^2$$

$$\overset{(b)}{\leq} \|H(\boldsymbol{x})\|_1 \|H(\boldsymbol{x})\|_\infty \|\overline{\omega}\|_2^2, \tag{130}$$

where (a) follows because induced matrix norms are submultiplicative and (b) follows because the matrix operator norm is bounded by square root of the product of matrix one norm and matrix infinity norm. Now, we claim that the one norm and the infinity norm of $H(\boldsymbol{x})$ are bounded as follows:

$$\max\left\{\|H(\boldsymbol{x})\|_1, \|H(\boldsymbol{x})\|_\infty\right\} \leq C_{6,\tau}C_{7,\tau}. \tag{131}$$

where $C_{6,\tau}$ and $C_{7,\tau}$ were defined in (119). Taking this claim as given at the moment, we continue with our proof. Combining (130) and (131), we have

$$\|\nabla q_2(\boldsymbol{x})\|_2^2 \leq C_{6,\tau}^2 C_{7,\tau}^2 \|\omega\|_2^2.$$

Therefore, we choose the pseudo derivative (see Def. 5) as follows:

$$\widetilde{\nabla} q_2(\boldsymbol{x}) = C_{6,\tau}C_{7,\tau} \|\omega\|_2. \tag{132}$$

Pseudo Hessian. Fix any $\rho \in \mathbb{R}$. We bound $\|\nabla(\rho^\top \widetilde{\nabla} q_2(\boldsymbol{x}))\|_2^2$ (see Def. 5) as follows:

$$\|\nabla(\rho^\top \widetilde{\nabla} q_2(\boldsymbol{x}))\|_2^2 = \sum_{u \in [p]} \left(\frac{d\rho^\top \widetilde{\nabla} q_2(\boldsymbol{x})}{dx_u}\right)^2 \overset{(132)}{=} 0.$$

Therefore, we choose the pseudo Hessian (see Def. 5) as follows:

$$\widetilde{\nabla}^2 q_2(\boldsymbol{x}) = 0. \tag{133}$$

The claimed result now follows by applying Prop. 2 with the pseudo discrete derivative defined in (132), and the pseudo discrete Hessian defined in (133).

It remains to show that the one-norm and the infinity-norm of $H(\boldsymbol{x})$ are bounded as in (131).

Bounds on the one-norm and the infinity-norm of H. We have

$$H_{tu}(\boldsymbol{x}) = \begin{cases} \left[1 - [\theta_u + \Theta_u^\top \boldsymbol{x}]x_u\right] \exp\left(-[\theta_u + \Theta_u^\top \boldsymbol{x}]x_u\right) & \text{if} \quad t = u, \\ -\Theta_{tu}x_u^2 \exp\left(-[\theta_u + \Theta_u^\top \boldsymbol{x}]x_u\right) & \text{otherwise.} \end{cases} \tag{134}$$

Therefore,

$$\interleave H(\boldsymbol{x})\interleave_1$$
$$= \max_{u \in [p]} \sum_{t \in [p]} |H_{tu}(\boldsymbol{x})|$$
$$\overset{(134)}{=} \max_{u \in [p]} \left|1 - [\theta_u + \Theta_u^\top \boldsymbol{x}]x_u\right| \exp\left(-[\theta_u + \Theta_u^\top \boldsymbol{x}]x_u\right) + \max_{u \in [p]} x_u^2 \exp\left(-[\theta_u + \Theta_u^\top \boldsymbol{x}]x_u\right) \sum_{t \neq u} |\Theta_{tu}|$$
$$\overset{(a)}{\leq} \left(1 + \alpha x_{\max} + 2x_{\max}^2 \zeta\right) \exp\left(x_{\max}(\alpha + \zeta x_{\max})\right) \overset{(119)}{=} C_{6,\tau}C_{7,\tau},$$

where (a) follows from Def. 8. Similarly, we have

$$\interleave H(\boldsymbol{x})\interleave_\infty$$
$$= \max_{t \in [p]} \sum_{u \in [p]} |H_{tu}(\boldsymbol{x})|$$
$$\overset{(134)}{=} \max_{t \in [p]} \left|1 - [\theta_t + \Theta_t^\top \boldsymbol{x}]x_t\right| \exp\left(-[\theta_t + \Theta_t^\top \boldsymbol{x}]x_t\right) + \max_{t \in [p]} \sum_{u \neq t} |\Theta_{tu}|x_u^2 \exp\left(-[\theta_u + \Theta_u^\top \boldsymbol{x}]x_u\right)$$
$$\overset{(a)}{\leq} \left(1 + \alpha x_{\max} + 2x_{\max}^2 \zeta\right) \exp\left(x_{\max}(\alpha + \zeta x_{\max})\right) \overset{(119)}{=} C_{6,\tau}C_{7,\tau},$$

where (a) follows from Def. 8.