A sharp NMF result with applications in network modeling

1 • Text marked in red: added new text.

2 • Obvious changes (e.g., typo corrections) are not highlighted in red.

Abstract

3   Given an $n \times n$ non-negative rank-K matrix Ω where m eigenvalues are nega-
4   tive, when can we write $\Omega = ZPZ'$ for non-negative matrices $Z \in \mathbb{R}^{n,K}$ and
5     $P \in \mathbb{R}^{K,K}$? While most existing works focused on the case of $m = 0$, our primary
6   interest is on the case of general m. With new proof ideas we develop, we present
7   sharp results on when the NMF problem is solvable, which significantly extend ex-
8   isting results on this topic. The NMF problem is partially motivated by applications
9   in network modeling. For a network with K communities, rank-K models are pop-
10   ular, with many proposals. The Degree-Corrected Mixed-Membership (DCMM)
11   model is a recent rank-K model which is especially useful and interpretable in
12   practice. To enjoy such properties, it is of interest to study when a rank-K model
13   can be rewritten as a DCMM model. Using our NMF results, we show that for a
14   rank-K model with parameters in the most interesting range, we can always rewrite
15   it as a DCMM model.

1 Introduction

17 Fix (n, K, m) where $n \geq K \geq 2$ and $0 \leq m \leq K - 1$. We are interested in the following
18 *Non-negative Matrix Factorization (NMF)* problem.

> The NMF problem: given an $n \times n$ symmetric non-negative irreducible matrix Ω with
>
> rank K where exactly m of the K nonzero eigenvalues are negative, when can we find   (1.1)
>
> non-negative matrices $Z \in \mathbb{R}^{n,K}$ and $P \in \mathbb{R}^{K,K}$ such that $\Omega = ZPZ'$?

19
20 **Definition 1.1** *We say a matrix Ω non-negative if all of its entries are non-negative, and we say it*
21 *positive if all of its entries are (strictly) positive. We say the NMF problem is solvable for Ω is we can*
22 *find non-negative matrices (Z, P) as above such that $\Omega = ZPZ'$.*

23 We assume $K \geq 2$ for the case of $K = 1$ is trivial, and we assume $m \leq K - 1$ for an irreducible
24 non-negative matrix has at least one positive eigenvalue (e.g., by Perron's theorem [10]).

25 NMF is a fundamental problem and has applications in areas such as image processing [5, 22], text
26 learning [20], hyper-spectral unmixing, and social network analysis [13]. Our setting is a special
27 case of NMF where both Ω and P are symmetric, so we may call it *symmetric NMF*. In the literature,
28 symmetric NMF was widely used in clustering of nonlinearly separable data from a similarity matrix
29 [8], where for a non-negative symmetric matrix Ω, it aims to find a non-negative matrix Z such that

$$\Omega = ZZ', \qquad \text{where } Z \in \mathbb{R}^{n,N} \text{ and } N \geq K. \qquad (1.2)$$

Note that, first, this implicitly requires that Ω is positive semi-definite. Second, it is understood that for many non-negative and positive semi-definite matrices Ω, the smallest N we can find in the factorization of (1.2) is strictly larger than K (the rank of Ω). See the 2021 book by Shaked-Monderer and Berman [25]. The book is 551 pages and summarizes nicely most existing results on NMF.

Unfortunately, our setting in (1.1) is significantly different from that in (1.2), so existing results on NMF do not directly apply. Especially, our NMF setting is motivated applications of social network modeling, where we must

- allow Ω to have negative eigenvalues,
- require that Z has exactly K columns ($K = \mathrm{rank}(\Omega)$),
- have a factorization of $\Omega = ZPZ'$ instead of $\Omega = ZZ'$ (we will soon see that both (P, Z) have practical meanings in our setting).

Below, in Section 1.1, we introduce several recent network models. In Section 1.2, we explain why the NMF problem (1.1) is important and relevant in social network modeling.

1.1 Several recent rank-K network models, and especially the DCMM model

Consider a symmetric connected network with n nodes and let A be the adjacency matrix:

$$A(i,j) = \begin{cases} 1, & \text{if there is an edge connecting nodes } i \text{ and } j, \\ 0, & \text{otherwise,} \end{cases} \qquad 1 \le i, j \le n.$$

As a convention, we do not allow self edges, so all diagonal entries of A are 0. We assume the network has K perceivable communities (communities are scientifically meaningful but mathematically hard to define; intuitively, they are clusters of nodes that have more edges "within" than "across" [12, 29]):

$$\mathcal{C}_1, \mathcal{C}_2, \ldots, \mathcal{C}_K.$$

In many network models, we assume that the upper triangular entries of A are independent Bernoulli random variables, and that there is an $n \times n$ non-negative matrix Ω such that $\Omega(i,j) = \mathbb{P}(A(i,j) = 1)$ for all $1 \le i \ne j \le n$. Let $\mathrm{diag}(\Omega) \in \mathbb{R}^{n,n}$ be the diagonal matrix where the i-th diagonal entry is $\Omega(i,i)$ and let $W \in \mathbb{R}^{n,n}$ be the matrix where $W(i,j) = A(i,j) - \Omega(i,j)$ if $i \ne j$ and $W(i,j) = 0$ otherwise. The matrix W is known as the generalized Wigner matrix. With these notations,

$$A = \Omega - \mathrm{diag}(\Omega) + W. \tag{1.3}$$

We call Ω the *Bernoulli probability matrix*. Frequently, we assume a *rank-K* model for Ω:

$$\Omega \text{ is an irreducible non-negative matrix where the rank is } K. \tag{1.4}$$

Note that K is the number of communities and has important practical meanings. Also, irreducibility is a natural assumption as we assume the network is connected (otherwise, we can study each connected component of the network separately). Below are some examples of rank-K models.

Example 1 (RDPG Model). In a Random Dot Product Graph (RDPG) model [27], we fix a K-dimensional distribution F, generate $y_i \overset{iid}{\sim} F$, and let $\Omega(i,j) = (y_i, y_j)$ (inner product), $1 \le i, j \le n$. If we write $Y = [y_1, y_2, \ldots, y_n]'$ (which is an $n \times K$ matrix), then $\Omega = YY'$. The model is well-known in network and graph modeling. However, a noteworthy issue is that, the matrix Ω defined in this way is always positive semi-definite. This makes the model relatively restrictive (e.g., [24]).

Example 2 (GRDPG Model). To address the issue above, Rubin-Delanchy *et al* [24] proposed the generalized RDPG (GRDPG). Fix K and $0 \le m < K$. Let $J_{K,m} = \mathrm{diag}(1, 1, \ldots, -1, \ldots, -1)$ be the $K \times K$ diagonal matrix where the first $(K - m)$ diagonal entries are 1 and the remaining diagonal entries are -1. With a similar Y matrix as in RDPG, GRDPG assumes $\Omega = Y J_{K,m} Y'$. An Ω defined in this way has negative eigenvalues, but we have to choose $(Y, J_{K,m})$ carefully to make sure that Ω is non-negative; this problem is not immediately clear.

Example 3. It was argued (e.g., [4]) that the Bernoulli probability matrix Ω in a graphon model can be well-approximated by a low-rank matrix provided with some regularity conditions.

In all these examples above, the parameters do not have explicit practical meanings (at least not directly or not sufficiently), so in a real application example, it remains unclear how to interpret the estimates of these parameters. Therefore, it is desirable to have models where the parameters have more explicit meanings in practice and so are easier to interpret.

The Degree-Corrected Mixed-Membership (DCMM) model is one of such models. Proposed by [15] (see also [28]), the model is motivated by the observation that natural networks usually have severe degree heterogeneity and mixed-memberships. To accommodate both features, for each node i, $1 \leq i \leq n$, we use a (strictly positive) parameter θ_i to model the degree heterogeneity and a weight vector $\pi_i \in \mathbb{R}^K$ to model the memberships, where $\pi_i(k)$ = weight node i puts in $\mathcal{C}_k$, $1 \leq k \leq K$. We call node i pure if π_i is degenerate (i.e., only one entry is nonzero) and mixed otherwise. We also model the community structure by a symmetric and non-negative matrix $P \in \mathbb{R}^{K,K}$:

$$P(k, \ell) = \text{baseline probability where a node in } \mathcal{C}_k \text{ and a node in } \mathcal{C}_\ell \text{ have an edge}, \ 1 \leq k, \ell \leq K.$$

DCMM assumes that for all $1 \leq i, j \leq n$, $\Omega(i,j) = \theta_i \theta_j \pi_i' P \pi_j$. If we let $\theta = (\theta_1, \ldots, \theta_n)'$, $\Pi = [\pi_1, \ldots, \pi_n]'$, and Θ be the $n \times n$ diagonal matrix where $\Theta(i,i) = \theta_i$, $1 \leq i \leq n$, then we have

$$\Omega = \Theta \Pi P \Pi' \Theta, \tag{1.5}$$

Conventionally, we assume $\text{rank}(\Pi) = \text{rank}(P) = K$, so DCMM is also a rank-K model.

Remark 1. The DCMM model can be viewed as the extension of several models, including the classical block model. In fact, (a) DCMM reduces to *Degree-Corrected Block Model (DCBM)* [19] if all nodes are pure, (b) DCMM reduces to the *Mixed-Membership Stochastic Block Model (MMSBM)* [1, 2, 23] if all θ_i are equal, and (c) DCMM reduces to the classical *Stochastic Block Model (SBM)* [9] if all nodes are pure and all θ_i are equal (as above, node i is pure if π_i is degenerate).

1.2 When is a rank-K network model also a DCMM model?

A DCMM model is a rank-K model, but compared to other rank-K models, all parameter matrices (Θ, Π, P) in the DCMM model have practical meanings and are easy to interpret. These make the DCMM model especially appealing in practice, and motivate the following problem:

$$\text{When is a rank-}K\text{ network model also a DCMM model?} \tag{1.6}$$

To explain why this is important, we use the dynamic co-citation networks in [11] as an example. The paper presented 21 co-citation networks for the same set of nodes (i.e., authors) in statistics, each for a different time window. We are interested in (a) how many research areas in statistics, (b) what are baseline citation exchanges between different research areas, and (c) how the research interests of individual authors evolve over time. Here, a co-citation network is a symmetrized citation network where each node is an author, and two nodes have an edge if they have been co-cited for at least N times (for an N they picked) in the corresponding time window. The paper suggested that there are 3 primary research areas in statistics (which was interpreted as "Bayes", "Biostatistics", and "Non-parametric") and a handful of sub-areas, and that it is convenient to model each co-citation network by a DCMM model with $K = 3$. In detail, for each author i and time window t, $1 \leq i \leq n$, $1 \leq t \leq T$, they used a $K \times K$ matrix $P^{(t)}$ to model the baseline citation exchanges between the primary research areas, a positive number θ_{it} to model the relative influence (in citations) of author i, and a weight vector π_{it} to model the research interest of author i. If we similarly let $\Theta^{(t)} = \text{diag}(\theta_{1t}, \theta_{2t}, \ldots, \theta_{nt})$ and $\Pi^{(t)} = [\pi_{1t}, \pi_{2t}, \ldots, \pi_{nt}]'$, then the Bernoulli probability matrix of the DCMM model at time t is $\Omega^{(t)} = \Theta^{(t)} \Pi^{(t)} P^{(t)} (\Pi^{(t)})' \Theta^{(t)}$. Using the DCMM model, they discovered a research triangle of statisticians (reminiscent of Efron's triangle for statistical philosophy [6]), and used it to visualize the trajectories of research interests of a handful of individual authors.

Imagine that, if we use a different rank-K model (e.g., GRDPG) to model these networks, say, with $\Omega^{(t)} = Y^{(t)} J^{(t)} (Y^{(t)})'$ for some matrices $(Y^{(t)}, J^{(t)})$. It is unclear how to relate $Y^{(t)}$ to baseline citation exchanges, research interests and relative influence of individual authors. This explains why (1.6) is of interest: given a rank-K network model, we wish to know when we can rewrite it as DCMM model, and so we can enjoy the properties and interpretability of the DCMM model.

We now come back to (1.6). Seemingly, NMF is to key to answer this question. Consider a positive matrix Ω with rank K and suppose that it has an NMF as in (1.1) for two non-negative matrices $Z \in \mathbb{R}^{n,K}$ and $P \in \mathbb{R}^{K,K}$: $\Omega = ZPZ'$. Write $Z = [z_1, z_2, \ldots, z_n]'$ so z_i' is the i-th row. Without loss of generality, assume all z_i are nonzero vectors. Let $\Theta(i,i) = \|z_i\|_1$ and $\pi_i = z_i / \|z_i\|_1$, $1 \leq i \leq n$. It is seen that $\Theta(i,i) > 0$, that each π_i is a weight vector, and that $\Omega = ZPZ' = \Theta \Pi P \Pi' \Theta$. Therefore, we can always rewrite a rank-K model as a DCMM model if Ω has an NMF as in (1.1). This explains our motivation underline the NMF problem (1.1).

Note that to answer the question in (1.1), a study on the NMF problem in (1.2) would be not be relevant. For example, in a DCMM model, K is the number of communities, so an NMF in (1.2) with an $N > K$ would not be useful. For this reason, we have to focus on the NMF problem in (1.1).

1.3 Results and contributions

Write $\Omega = Y J_{K,m} Y'$ as in Example 2, where $J_{K,m} = \mathrm{diag}(1, \ldots, 1, -1, \ldots, -1)$ is a $K \times K$ diagonal matrix and $Y = [y_1, y_2, \ldots, y_n]' \in \mathbb{R}^{n,K}$. Let λ_k be the k-th eigenvalue of Ω and let ξ_k be the corresponding eigenvector. For $1 \leq i \leq n$, define $r_i \in \mathbb{R}^{K-1}$ by $r_i(k) = \xi_{k+1}(i)/\xi_1(i)$, $1 \leq k \leq K - 1$. For any unit-norm vector $y_0 \in \mathbb{R}^K$, let $c(y_0) = \max_{\{1 \leq i \leq n\}} \{|(y_i, y_0)|/\|y_i\|\}$. In Section 2, we show that the NMF problem for Ω is solvable if $m \leq K/2$ and $c(y_0) \geq \sqrt{1 - 1/K}$ for some y_0; let us call this the main condition. We show that, in order for the NMF problem to be solvable, the constant $\sqrt{1 - 1/K}$ can not be further reduced. Therefore, in this sense, our results are sharp. Using this, we deduce several other results. Especially, we show that the NMF problem is solvable for Ω if $\sum_{k=1}^{K-1}(|\lambda_{k+1}| \cdot r_i^2(k)) \leq |\lambda_1|/(K-1)$ for all $1 \leq i \leq n$. We also extend our results to the case of $m > K/2$, and explain why we need a different proof in this case.

In Section 3, we apply our results on NMF to network modeling. We argue that for parameters in the most interesting range, we have (A) all $\|r_i\|$ are bounded, and (B) $\max_{2 \leq k \leq K} \{|\lambda_k/\lambda_1|\} \to 0$, and so the condition just mentioned holds. This implies that we can alway rewrite a rank-K network model as a DCMM model if the parameters are in the most interesting range. We also discuss how to check the main condition in practice where Ω is unknown. We tackle this by proposing an approach to estimating Ω, and support our results by some real networks.

Our contributions are two fold. First, we develop several new results on symmetric NMF (a problem of interest in many applications [5]). Existing works on symmetric NMF have been focused on the case of $m = 0$ (so Ω is positive semi-definite; m is the number of negative eigenvalues of Ω). In this case, the best result is seen to be [25, Theorem 3.137], which can be viewed as a special case of our results; see Remark 2. This suggests that our results are sharp, for they are hard to improve even in the special case of $m = 0$. Note that our case allows m to take any possible values, so it is clearly harder to study. For example, to show the results for the case of $m = 0$, it suffices if we can find a $K \times K$ orthogonal matrix Q such that YQ' is non-negative, since $J_{K,m}$ is the identity matrix in this case. For our case, we must find a Q such that YQ' and $QJ_{K,m}Q'$ are simultaneously non-negative. Clearly, this requires new ideas. We tackle this by constructing a special class of matrices Q; see our proofs for details. Our approach is quite different from that of [25, Theorem 3.137] and is new.

Second, we shed interesting new light on different rank-K network models. In the literature, it is not unusual that many similar models are proposed for the same type of data sets. But in the end, we need to understand the advantages and disadvantages of different models, and pick the most suitable one. Our study recommends DCMM model, for it offers desired practical interpretability which other rank-K models do not have, and points out that a general rank-K model is also a DCMM model if the parameters are in the most interesting range. Such findings are valuable for they can help us identify the most suitable models in real applications.

1.4 Notations

In this paper, $e_0 = K^{-1/2} \mathbf{1}_K = K^{-1/2}(1, 1, \ldots, 1)'$ (where "$'$" is the transpose), and $e_1, e_2, \ldots, e_K$ denote the standard basis vectors of K-dimensional Euclidean space. For any $q > 0$ and vector x, $\|x\|_q$ denotes the ℓ^q-norm (when $q = 2$, we drop the subscript and write $\|x\|$). For any two vectors x and y of the same dimension, (x, y) denotes the inner product. For a vector $a \in \mathbb{R}^n$, $\mathrm{diag}(a)$ denotes the $n \times n$ diagonal matrix where the i-th diagonal entry is a_i, $1 \leq i \leq n$. When Ω is an $n \times n$ matrix, $\mathrm{diag}(\Omega)$ denotes the $n \times n$ diagonal matrix where the i-th entry is $\Omega(i, i)$, $1 \leq i \leq n$.

2 Main results on NMF

This section presents our results on NMF. Results on network modeling are in Section 3. Consider an $n \times n$ irreducible non-negative matrix Ω with rank K, where n is usually much larger than K. By Perron's theorem [10], at least one eigenvalue of Ω is positive. Fix $0 \leq m \leq K - 1$ and suppose Ω has m negative eigenvalues. Let $J_{K,m} = \mathrm{diag}(1, \ldots, 1, -1, \ldots, -1)$ be the $K \times K$ diagonal matrix

172   as in Example 2. By basic algebra, we can always write

$$\Omega = Y J_{K,m} Y', \qquad \text{for a full rank matrix } Y \in \mathbb{R}^{n,K}. \tag{2.7}$$

173   We can also show (e.g., an exercise with the Weyl's theorem [10]) that for any matrix as in (2.7), the
174   numbers of positive and negative eigenvalues are $(K - m)$ and m, respectively. Write

$$Y = [y_1, y_2, \ldots, y_n]', \qquad \text{so that } y_i' \text{ is row } i \text{ of } Y, 1 \le i \le n. \tag{2.8}$$

175   Define the subset of K-dimensional vectors that live on the unit-sphere where the last m entries are 0:

$$\mathcal{S}_{K,m} = \{x = (x_1, \ldots, x_K)' \in \mathbb{R}^K, \|x\| = 1, x_{K-m+1} = \ldots = x_K = 0\}.$$

176   When $m = 0$, $\mathcal{S}_m$ is the unit sphere of $\mathbb{R}^K$. The following theorem is proved in the supplement.

177   **Theorem 2.1** *Fix $K \ge 2$, $n \ge K$, and $0 \le m \le K/2$. Consider the NMF problem (1.1) where*
178   *$\Omega = Y J_{K,m} Y'$ and Y are as in (2.7). Suppose there is a vector $y_0 \in \mathcal{S}_{K,m}$ such that*

$$|(y_0, y_i)|/\|y_i\| \ge \sqrt{1 - 1/K}, \qquad \text{for all } 1 \le i \le n. \tag{2.9}$$

179   *There exists a $K \times K$ orthogonal matrix Q such that both YQ' and $QJ_{K,m}Q'$ are non-negative. As*
180   *a result, the NMF problem for Ω is solvable: $\Omega = ZPZ'$ with $Z = YQ'$ and $P = QJ_{K,m}Q'$.*

181   We have several comments. First, Theorem 2.1 assumes two conditions: $m \le K/2$ and (2.9). When
182   $K \le 2$, both conditions hold automatically, so the NMF problem is always solvable in this case; see
183   Section 2.1. As far as we know, our proof is different from existing approaches. Second, in Theorem
184   2.1, we require $y_0 \in \mathcal{S}_m$. This may seem restrictive, but is not. This is because y_0 is a vector we
185   choose for our own convenience. In fact, one of the most interesting settings for NMF seems to be
186   that in Section 2.3, where we choose $y_0 = (1, 0, \ldots, 0)'$, so the requirement is satisfied automatically.
187   Also, when the last m entries of y_0 are nonzero but sufficiently small, Theorem 2.1 continues to
188   hold if we modify the term $\sqrt{1 - 1/K}$ slightly. Third, from a practical view point, the condition
189   of $m \le K/2$ is mild: we rarely see a rank-K network model with $m > K/2$ (note here m can be
190   estimated using the eigenvalues of the adjacency matrix A). For theoretical completeness, the case of
191   $m > K/2$ is also interesting, but there does not exist an orthogonal matrix Q such that $QJ_{K,m}Q'$
192   is non-negative. This is because for any such Q, $\text{trace}(QJ_{K,m}Q') = K - 2m < 0$. Therefore, we
193   must find a different way to solve the NMF problem in this case. We discuss this in Section 2.4. Last,
194   an interesting question is whether our idea is extendable to asymmetric NMF or complex NMF [7].
195   As a simple extension to asymmetric NMF, consider an $n \times p$ positive matrix Ω of rank-K. By SVD,
196   $\Omega = YZ'$ for an $n \times K$ matrix Y and a $p \times K$ matrix Z. Let y_i' be i-th row of Y and z_j' be the j-th
197   row of Z, respectively. If there is a $y_0 \in \mathcal{S}_{K,m}$ such that for all i and j, $|(y_i, y_0)|/\|y_i\| \ge \sqrt{1 - 1/K}$
198   and $|(z_j, y_0)|/\|z_j\| \ge \sqrt{1 - 1/K}$, then we can find a $K \times K$ orthogonal matrix Q which rotates all
199   rows of Y and Z to the first orthant simultaneously. In this case, the asymmetric NMF problem is
200   solvable for Ω. For reasons of space, we leave further study along this line to the future.

201   Our result is sharp for the constant $\sqrt{1 - 1/K}$ in (2.9) can not be further reduced. While we can
202   show this for general K, we illustrate with the case of $K = 2$ for instruction purpose. In this case,
203   we can rotate n unit-norm vectors $y_1, y_2, \ldots y_n$ in $\mathbb{R}^2$ simultaneously to the first orthant if and only
204   if there is a unit-norm vector y_0 such that $|(y_0, y_i)| \ge \sqrt{1 - 1/2}$ (i.e., the angle between them is
205   $\le \pi/4$) for all $1 \le i \le n$. See Section 2.1 and Remark 3 for more discussion. Another way to see
206   the sharpness is to consider the case of $m = 0$ (so Ω is positive semi-definite). In this case, condition
207   (2.9) is hard to improve and is the weakest we have so far in the literature; see Remark 2.

2.1   The case of $K = 2$

209   In this case, the NMF problem is always solvable, as the two conditions of Theorem 2.1, $m \le K/2$
210   and (2.9), hold automatically. In fact, first, since Ω has at least one positive eigenvalues and $K = 2$,
211   we have either $m = 0$ or $m = 1$, and so $m \le K/2$. Second, we can always find a $y_0 \in \mathcal{S}_m$ such that
212   (2.9) is satisfied. In detail, let $0 \le \theta_i < 2\pi$ be the angle from e_1 ($e_1 = (1, 0)$) to y_i counterclockwise,
213   and let θ_{min} and θ_{max} be the smallest and largest values of all θ_i. Now, when $m = 0$, let y_0 be the
214   unit vector where the angle from e_1 to y_0 is $(\theta_{max} + \theta_{min})/2$, counterclockwise. When $m = 1$, take
215   $y_0 = (1, 0)$. The following theorem is proved in the supplement.

216   **Theorem 2.2** *Fix $K = 2$, $0 \le m \le K - 1$, $n \ge K$, and let y_0 be as above. In this case, $m \le K/2$*
217   *and (2.9) holds for the y_0 above, so the NMF problem is always solvable for Ω.*

2.2 When y_0 is a scaled weighted average of y_i's

For the y_0 in (2.9), an interesting choice is to let it be proportional to a weighted average of y_i's. Call $w \in \mathbb{R}^n$ a weight vector if all of its entries are non-negative with a sum of 1. Recall that $\Omega = Y J_{K,m} Y'$. Define a proxy of Ω by $\widetilde{\Omega} = YY'$. Note that $\widetilde{\Omega} = \Omega$ if $m = 0$. Introduce $y^{(w)} \in \mathbb{R}^K$ and $\beta^{(w)} \in \mathbb{R}^n$ by $y^{(w)} = \sum_{i=1}^n w_i y_i = Y'w$ and $\beta^{(w)} = \widetilde{\Omega}w$. Since Y is full rank, $y^{(w)} \neq 0$. Take $y_0 = y^{(w)}/\|y^{(w)}\|$. Condition (2.9) reduces to

$$|\beta_i^{(w)}|/\sqrt{\widetilde{\Omega}(i,i)(w'\widetilde{\Omega}w)} \geq \sqrt{1 - 1/K}, \qquad \text{for all } 1 \leq i \leq n. \tag{2.10}$$

Theorem 2.3 *Fix $K \geq 3$, $0 \leq m \leq K/2$, and $n \geq K$. The NMF problem (1.1) is solvable for Ω if the last m entries of $y^{(w)}$ are 0 and (2.10) holds.*

Theorem 2.3 follows from Theorem 2.1 by direct calculations, so the proof is omitted. We require that the last m entries of $y^{(w)}$ are 0, for we need $y_0 \in \mathcal{S}_m$ in Theorem 2.1. As explained before, this may seem restrictive, but it is not, as in the most interesting case to be discussed in Section 2.3, we take $y^{(w)} = (1, 0, \ldots, 0)$, so the requirement is satisfied automatically. See details therein.

When $m = 0$, $\widetilde{\Omega} = \Omega$, and $\beta^{(w)} = \Omega w$. In this case, condition (2.10) reduces to

$$|\beta_i^{(w)}|/\sqrt{\Omega(i,i)(w'\Omega w)} \geq \sqrt{1 - 1/K}. \tag{2.11}$$

We have the following corollary, the proof of which is straightforwards so is omitted.

Corollary 2.1 *Fix $K \geq 3$ and $n \geq K$. The NMF problem (1.1) is solvable for Ω if $m = 0$ and (2.11) holds.*

Remark 2. If we take $w = n^{-1}\mathbf{1}_n$, then (2.10) reduces to $|\beta_i|/\sqrt{\Omega(i,i)(\mathbf{1}_n'\Omega\mathbf{1}_n)} \geq \sqrt{1 - 1/K}$ with $\beta = \Omega\mathbf{1}_n$, and Corollary 2.1 reduces to [25, Theorem 3.137], where $m = 0$ and Ω is positive semi-definite. Our setting is more general as Ω may have m negative eigenvalues for any $m \leq K/2$. For the case of $m = 0$, [25, Theorem 3.137] (see also [26]) is by far the best results we can have. The book [25] presents several other results on this topic, but they need some conditions which are less intuitive or harder to check. Recall that the constant $\sqrt{1 - 1/K}$ in (2.9) can not be further reduced. These suggest that Theorem 2.1 is hard to improve and our results are sharp.

Remark 3. (*When can we rotate n vectors to the first orthant?*) As a stylized application, consider the following problem. Let $x_1, x_2, \ldots, x_n$ be n unit-norm vectors in $\mathbb{R}^K$, $n \geq K$, and let $\alpha_K(x_1, x_2, \ldots, x_n) = \min_{1 \leq i,j \leq n}\{(x_i, x_j)\}$. For what values of $\alpha_K(x_1, x_2, \ldots, x_n)$ can we rotate all n points simultaneously to the first orthant? Let $X = [x_1, x_2, \ldots, x_n]'$ and assume X is full rank without loss of generality. The matrix $\Omega = XX'$ is symmetric and positive semi-definite. Let $\alpha_K^* = 0$ if $K = 2$ and $\alpha_K^* = \sqrt{1 - 1/K}$ if $K \geq 3$. Applying Theorem 2.1 with $m = 0$, it follows that when $\alpha_K(x_1, x_2, \ldots, x_n) \geq \alpha_K^*$, we can rotate all n points to the first orthant. Note that we can not do so if $\alpha_K(x_1, x_2, \ldots, x_n) < 0$.

2.3 When Y is constructed by the spectral decomposition of Ω

So far, we have tried to keep our results as general as we can, and Y can be any matrix satisfying $\Omega = Y J_{K,m} Y'$. An interesting special case is when Y is constructed using the spectral decomposition of Ω, which we now discuss. For $1 \leq k \leq K$, let λ_k be the k-th largest eigenvalue of Ω, and let ξ_k be the corresponding (unit-norm) eigenvector. In the literature λ_1 and ξ_1 are called the Perron root and Perron vector, respectively, where we can always assume all entries of ξ_1 are positive since Ω is irreducible and non-negative (e.g., [25]). Write $\Xi = [\xi_1, \xi_2, \ldots, \xi_K]$ and define the $n \times (K-1)$ so-called *matrix of entry-wise ratio* R by $R(i,k) = \xi_{k+1}(i)/\xi_1(k)$, $1 \leq k \leq K-1, 1 \leq i \leq n$ [12, 16]. Introduce

$$D = \text{diag}(|\lambda_1|, |\lambda_2|, \ldots, |\lambda_K|), \qquad D_0 = \text{diag}(|\lambda_2|, \ldots, |\lambda_K|), \tag{2.12}$$

and write

$$R = [r_1, r_2, \ldots, r_n]', \qquad Y = \Xi D^{1/2} = [y_1, y_2, \ldots, y_n]'. \tag{2.13}$$

By spectral decomposition, $\Omega = \Xi D^{1/2} J_{K,m} D^{1/2} \Xi' = Y J_{K,m} Y'$. Now, in Section 2.2, if we take $w = c\xi_1$ where $c = 1/\|\xi_1\|_1$, then by basic algebra and definitions, it is seen $y^{(w)} = c\sqrt{\lambda_1}e_1$ and so

262 $y_0 = e_1$ and especially $y_0 \in S_m$. Moreover, $\beta_i^{(w)} = c\lambda_1\xi_1(i)$, $w'\widetilde{\Omega}w = c^2\lambda_1$, and $\widetilde{\Omega}(i,i) = y_i'Dy_i$.
263 Combining these, condition (2.10) reduces to

$$r_i'D_0r_i \equiv \sum_{k=1}^{K-1} (|\lambda_{k+1}| \cdot r_i^2(k)) \leq |\lambda_1|/(K-1), \qquad \text{for all } 1 \leq i \leq n. \qquad (2.14)$$

264 The following theorem is proved in the supplement.

265 **Theorem 2.4** *Fix* $K \geq 3$, $m \leq \frac{K}{2}$, *and* $n \geq K$. *The NMF problem (1.1) is solvable if (2.14) holds.*

266 Note that as in most works on NMF (e.g., [25]), the main goal is to find *easy-to-check* conditions
267 under which the NMF is solvable. Such conditions are sufficient but are not necessary.

268 **2.4 The case of** $m > K/2$

269 So far, we have been focused on the case of $m \leq K/2$, which is the case that is most frequently
270 found in real networks. For completeness, we now consider the case where $m > K/2$. Since
271 $0 \leq m \leq K-1$, such a case only exists when $K \geq 3$. In Theorem 2.1, we show that when $m \leq K/2$,
272 we can find an orthogonal matrix Q such that $QJ_{K,m}Q'$ is non-negative. When $m > K/2$, we can
273 not do this, as for any such Q, $\text{trace}(QJ_{K,m}Q') = (K - 2m) < 0$. Therefore, we need a new
274 approach.

275 A convenient approach is to redefine $J_{K,m}$ where we select a subset of the positive diagonal entries of
276 $J_{K,m}$ and add a positive number for each of them. Success has been shown in a related setting (e.g.,
277 [3]). Using such a trick, we can extend all our main results to the case of $m > K/2$. For reasons of
278 space, we only consider an extension of Theorem 2.4, as the claim of the theorem is probably the
279 most explicit. Also for reasons of space, we only consider the case where we add a number to the
280 first diagonal entry of $J_{K,m}$. Note that the idea is readily extendable to more general cases.

281 Let $\mathcal{Q}$ be the set of all orthogonal matrices where the first column is $K^{-1/2}(1, 1, \ldots, 1)'$. Fix
282 $1 \leq m \leq K - 1$. For any $Q \in \mathcal{Q}$, write $Q = [Q^{(K-m)}, Q^{(m)}]$, where $Q^{(K-m)}$ and $Q^{(m)}$ are
283 the sub-matrix of Q consisting the first $(K - m)$ columns and the other m columns, respectively.
284 Introduce a constant by $a_m = 1 + K \inf_{Q \in \mathcal{Q}} \max_{1 \leq i,j \leq K}\{H(i,j) : H = 2Q^{(m)}(Q^{(m)})' - I_K\}$ (I_K:
285 $K \times K$ identity matrix). Theorem 2.5 extends Theorem 2.4 and is proved in the supplement.

286 **Theorem 2.5** *Fix* $K \geq 3$, $0 \leq m \leq (K-1)$, *and* $n \geq K$. *We have* $a_m = 1$ *if* $m \leq K/2$
287 *and* $a_m = (K - 1)$ *if* $m = K - 1$. *Also, the NMF problem is solvable for* Ω *if* $r_i'D_0r_i \equiv$
288 $\sum_{k=1}^{K-1} |\lambda_{k+1}|r_i^2(k) \leq 1/[a_m(K-1)]$ *for all* $1 \leq i \leq n$.

289 When $m \leq K/2$, $a_m = 1$. In this case, the claim here is the same as that in Theorem 2.4.

290 **Remark 4**. When the NMF problem for Ω is solvable, the solution is usually not unique without a
291 proper regularity condition (e.g., [5]). In our setting, once we can write $\Omega = \Theta\Pi P\Pi'\Theta$ for some
292 non-negative matrices (Θ, Π, P) as in (1.5), the factorization is unique if (a) for each $1 \leq k \leq K$,
293 there is at least one i such that $\pi_j = e_k$, where e_k is the k-th standard Euclidean basis vector of $\mathbb{R}^K$,
294 and (b) all diagonal entries of P are 1 (see [15, 16] for a proof).

295 **Remark 5**. When condition (2.9) of Theorem 2.1 holds for some vectors y_0, how to find such a y_0
296 and the orthgonal matrix Q in Theorem 2.1 numerically? This is an interesting question and we
297 discuss it in Section F of the supplement.

298 # 3 When is a rank-K network model also a DCMM model

299 So far, we focus on general NMF settings where we show that the NMF problem (1.1) is solvable
300 when, for example, (2.14) holds. We now apply the results to networks and study when we can
301 rewrite a rank-K network model as a DCMM model. Network analysis (e.g., community detection,
302 membership estimation, link prediction) is a well-studied area, where we have a lot of knowledge
303 on what is the regime of major interest and what conditions are reasonable [16, 15, 18, 28]. In fact,
304 in network analysis, we usually use an asymptotic framework where $n \to \infty$, K is fixed, and other
305 parameters may vary with n, where it is quite acceptable to assume

306 • (A) all $\|r_i\|$ are bounded,

- (B) $\max_{2 \le k \le K}\{|\lambda_k/\lambda_1|\} \to 0$;

the notations are the same as those in Theorem 2.4. In fact, (A)-(B) model the *most interesting regime* in network analysis. In Theorem 2.4, the main condition (e.g., (2.14)) is $r_i' D_0 r_i \le |\lambda_1|/(K-1)$ for all $1 \le i \le n$. It is seen that once (A)-(B) hold, $(1/|\lambda_1|)D_0 \to 0$ and (2.14) holds. Therefore, we can always rewrite a rank-K network model as a DCMM model when (A)-(B) hold.

The remaining question is then, why (A)-(B) are reasonable assumptions in network analysis, and why they model the most interesting regime in network analysis. We now explain these in details.

Let Ω be the Bernoulli probability matrix as in (1.3). Suppose $\Omega = YPY'$, where $Y \in \mathbb{R}^{n,K}$ and is full rank, $P \in \mathbb{R}^{K,K}$, and (Y, P) are not necessarily non-negative. Denote $G = Y'Y$. Note that G is a $K \times K$ symmetric and positive definite matrix. Let $G^{1/2}$ be the (unique) square root of $Y'Y$. We usually assume Y is balanced in that (a) the ℓ^2-norm of all K columns are in the same order, and (b) no severe linearity between the K columns [15, 18]. As a result, all eigenvalues of G are at the same order. By basic algebra, there is a $K \times K$ orthogonal matrix Q such that $\Xi = [\xi_1, \xi_2, \ldots, \xi_K] = YB$, where $B = G^{-1/2}Q$. Write $B = [b_1, b_2, \ldots, b_K]$ and let $0 \le \alpha_i < 2\pi$ be the angle between b_1 and y_i (counterclockwise). Let $M(\Omega) = \max_{\{1 \le i \le n\}}\{1/|\cos(\alpha_i)|\}$ and define matrix $V \in \mathbb{R}^{K,K-1}$ by

$$V(i,k) = b_{k+1}(i)/b_1(i), \qquad 1 \le i \le K, \ 1 \le k \le K-1. \tag{3.15}$$

Write $V = [v_1, v_2, \ldots, v_K]'$ so v_k' is row-k of V, $1 \le k \le K$. For any symmetrical matrix P, $\lambda_k(P)$ denotes the k-th largest eigenvalue; to be consistent with earlier notations, we simply write $\lambda_k(\Omega)$ as λ_k. Lemma 3.1 is proved in the supplement.

Lemma 3.1 *We have $B = \mathrm{diag}(b_1)[\mathbf{1}_K, V]$, $P = B\mathrm{diag}(\lambda_1, \ldots, \lambda_K)B'$, b_1 is an eigenvector of PG, and $P(k,k) = b_1^2(k)[\lambda_1 + v_k'\mathrm{diag}(\lambda_2, \ldots, \lambda_K)v_k]$, $1 \le k \le K$. Moreover, if as $n \to \infty$, $\lambda_1(G) \le c_0\lambda_K(G)$ for a constant $c_0 > 0$, then condition (B) holds if and only if $\max_{\{2 \le k \le K\}}\{|\lambda_k(P)/\lambda_1(P)|\} \to 0$, and $\max_{\{1 \le i \le n\}}\{\|r_i\|\} \le CM(\Omega)$.*

It is seen that conditions (A)-(B) hold if $M(\Omega) \le C$ and $\max_{2 \le k \le K}\{|\lambda_k(P)/\lambda_1(P)|\} \to 0$. The first one is mild: it only requires that no y_i is nearly orthogonal to b_1. To boil these conditions down to a more explicit and vivid form, we consider the DCMM model. It is fine to consider the DCMM model here for (a) we only use the model to explain why conditions (A)-(B) are reasonable, and (b) the argument below is extendable beyond the DCMM model. In the DCMM model, $\Omega = \Theta\Pi P\Pi'\Theta$. Therefore, we can write $\Omega = YPY'$ if we let $Y = \Theta\Pi$, where we note that (Y, P) are non-negative. Recall that $G = Y'Y$ (a positive definite $K \times K$ matrix). Lemma 3.2 is proved in the supplement.

Lemma 3.2 *If (Y, P) are non-negative, then first, PG is an irreducible non-negative matrix and b_1 is the Perron vector, so all entries of b_1 are strictly positive. Second, all rows of r_i lives with a simplex with $v_1, v_2, \ldots, v_K$ being the vertices, so $\max_{\{1 \le i \le n\}}\{\|r_i\|\} \le \max_{\{1 \le k \le K\}}\{\|v_k\|\}$. Last, if $\lambda_1(G) \le c_0\lambda_K(G)$, then $\max_{\{1 \le i \le n\}}\{\|r_i\|\} \le CM(\Omega) \le C\max_{1 \le k \le K}\{\|b_1\|/b_1(k)\}$.*

Now, first, in a DCMM model, the matrix $P(k, \ell)$ measures the baseline probability where there is an edge between a node in community k and a node in community ℓ. Therefore, the *most difficult or most interesting case* is where all $P(k, \ell)$ have similar values. In this case, P is close to rank-1, or in other words, $\max_{2 \le k \le K}\{|\lambda_k(P)/\lambda_1(P)|\} \to 0$, and so $\max_{2 \le k \le K}\{|\lambda_k/\lambda_1|\} \to 0$. See for example [15, 18], where it was further pointed out that the most difficult case for network analysis is when $\max_{\{2 \le k \le K\}}\{|\lambda_k|\} \le L_n \cdot \sqrt{\lambda_1}$ for a multi-log(n) factor L_n. Therefore, condition (A) models the most difficult case of network analysis and so is of major interest. Moreover, by Lemma 3.2, $\max_{\{1 \le i \le n\}}\{\|r_i\|\} \le C$ if all entries of b_1 are at the same order. This is only a mild condition for b_1 is the Perron vector of PG. Last, by Lemma 3.2, we also have $\max_{\{1 \le i \le n\}}\{\|r_i\|\} \le C$ if we alternatively assume $\max_{\{1 \le k \le K\}}\{\|v_k\|\} \le C$. Recall that $B = G^{-1/2}Q = [b_1, b_2, \ldots, b_K]$ and $v_1', v_2', \ldots, v_K'$ are rows of V formed by dividing $b_2, b_3, \ldots, b_K$ by b_1 entry-wise, where b_1 is the Perron vector. Since G is positive definite where all eigenvalues are at the same order, Q is orthogonal, and V is properly scaled (and all of them have small-sizes), it is only a mild condition to assume $\max_{\{1 \le k \le K\}}\{\|v_k\|\} \le C$. These explain why conditions (A)-(B) are mild condition and they model the most challenging regime for network analysis.

4 Real data examples, and especially how to check condition (2.14)

Let $a_i = (1/|\lambda_1|)r_i' D_0 r_i$, $1 \le i \le n$. Condition (2.14) can be rewritten as $a_i \le 1/(K-1)$, for all $1 \le i \le n$. In applications, Ω is unknown, so it is unclear how to obtain a_i. A straightforward approach is to estimate a_i with the eigenvalues and eigenvectors of the adjacency matrix A, but the estimates may be too noisy. We propose the following approach, which is inspired by Lemmas 3.1-3.2 and the recent Mixed-SCORE approach [16]. Let (Y, V) be as above. Mixed-SCORE suggests an interesting idea for estimating V and (a normalized version of) Y, denoted by Π; see details therein. Let $\hat{\lambda}_k$ be the k-th eigenvalue of A and let $\hat{\xi}_k$ be the corresponding eigenvector. Write $\widehat{\Xi} = [\hat{\xi}_1, \hat{\xi}_2, \ldots, \hat{\xi}_K] = [\hat{z}_1, \hat{z}_2, \ldots, \hat{z}_n]'$, so $\hat{z}_i'$ is row-i of $\widehat{\Xi}$. Our approach runs as follows.

- Apply Mixed-SCORE and obtain an estimate $(\widehat{V}, \widehat{\Pi})$ for (V, Π). Let $\hat{v}_k'$ be row k or $\widehat{V}$ and let $\hat{\pi}_i'$ be row i of $\widehat{\Pi}$, $1 \le k \le K$, $1 \le i \le n$.

- Estimate b_1 by $\hat{b}_1$ where $\hat{b}_1(k) = [\hat{\lambda}_1 + \sum_{k=2}^{K} \hat{\lambda}_k \hat{v}_k' \mathrm{diag}(\hat{\lambda}_2, \ldots, \hat{\lambda}_K)\hat{v}_k]^{-1/2}$. Let $\widehat{B} = \mathrm{diag}(\hat{b}_1)[\mathbf{1}_K, \widehat{V}]$, and estimate P by $\widehat{P} = \widehat{B}\mathrm{diag}(\hat{\lambda}_1, \hat{\lambda}_2, \ldots, \hat{\lambda}_K)\widehat{B}'$. Let $\hat{y}_i = (\|z_i\|_1/\|\widehat{B}'\hat{\pi}_i\|_1)\hat{\pi}_i$, $1 \le i \le n$, and let $\widehat{Y} = [\hat{y}_1, \hat{y}_2, \ldots, \hat{y}_n]'$.

- Let $\hat{\mu}_k$ be the k-th eigenvalue of the matrix $\widehat{\Omega} = \widehat{Y}\widehat{P}\widehat{Y}'$, and let $\hat{\eta}_k$ be the corresponding eigenvector. In the definition of a_i (see above and (2.14)), replace (λ_k, ξ_k) by $(\hat{\mu}_k, \hat{\eta}_k)$ and denote the resultant quantity $\hat{a}_i$, $1 \le i \le n$. These are our estimates for a_i.

The approach can be shown to be consistent for Ω under some regularity conditions. We skip the study for it is beyond the scope of this paper. In this algorithm, $(\widehat{Y}, \widehat{P})$ are not automatically non-negative, and to check whether NMF is solvable for $\widehat{\Omega}$, we can check if

$$\hat{a}_i \le 1/(K-1), \qquad \text{for all } 1 \le i \le n. \tag{4.16}$$

Remark 4. Condition (2.14) of Theorem 2.4 is only a sufficient condition for NMF; they are not necessary conditions. It could happen that an NMF is solvable for an Ω but (2.14) does not hold.

We now consider some real examples. The weblog is a well-known data set [21], where with some light preprocessing, the network has $1,222$ node (each is a blog) and $16,714$ edges (each is a two-way hyperlink). The network has two communities: democratic and republican. For this data set, a rank-2 model is appropriate, so we have $(n, K) = (1, 222, 2)$ (e.g., [29, 12, 18]). Let Ω be the Bernoulli probability matrix as in (1.3). By Theorem 2.2, when $K = 2$, we can always decompose Ω as $\Omega = YPY'$ for a non-negative $n \times 2$ matrix Y and a 2×2 non-negative matrix P. Now, by the paragraph right above Remark 1, we can rewrite $\Omega = \Theta\Pi P\Pi\Theta$ as in (1.3), so Ω satisfies a DCMM model. Same claim can be drawn for the karate data set [29, 12], where we similarly have $K = 2$.

As another example, we consider the UKFaculty network (e.g., see [17, Table 1]). It is reasonable to model the network with a rank-K model with $(n, K) = (81, 3)$ and $m \le K/2$. By Theorem 2.4, the model can be rewritten as a DCMM model if (4.16) holds. Following the discussion above, we first obtain an estimate $\widehat{\Omega}$ for Ω. We then use $\widehat{\Omega}$ to obtain $\hat{a}_i$ and check if (4.16) holds. The results are in the figure below (left), where the maximum of $\hat{a}_1, \hat{a}_2, \ldots, \hat{a}_n$ is slightly smaller than 0.5 ($1/(K-1) = 0.5$ as $K = 3$), suggesting that (4.16) holds. Moreover, let $\hat{\mu}_k$ be the k-th eigenvector of $\widehat{\Omega}$ and let $\hat{\eta}_k$ be the corresponding eigenvector. Let $\widehat{D} = \mathrm{diag}(\hat{\mu}_1, \ldots, \hat{\mu}_K)$ and $\widehat{Y} = [\hat{\eta}_1, \ldots, \hat{\eta}_K]\widehat{D}^{1/2}$. We have $\widehat{\Omega} = \widehat{Y}J_{K,m}\widehat{Y}'$. Let Q be the 3×3 matrix where the three rows are $(1/\sqrt{3}, 1/\sqrt{6}, 1/\sqrt{2})$, $(1/\sqrt{3}, 1/\sqrt{6}, -1/\sqrt{2})$, and $(1/\sqrt{3}, -2/\sqrt{6}, 0)$, respectively. Define $\widehat{Z} = \widehat{Y}Q'$. It is seen $\widehat{\Omega} = \widehat{Y}J_{K,m}\widehat{Y}' = \widehat{Z}[QJ_{K,m}Q']\hat{Z}'$, where $QJ_{K,m}Q'$ is seen to be non-negative. Moreover, for $1 \le i \le n$, let $\hat{z}_i$ be the smallest entry in row-i of $\widehat{Z}$. The figure below (right) plots the histogram for $\{\hat{z}_i\}_{i=1}^n$. The results suggest that all $\hat{z}_i$ are non-negative, so the matrix $\widehat{Y}Q'$ is non-negative. Therefore, $\widehat{\Omega}$ has an NFM by $\widehat{\Omega} = \widehat{Z}[QJ_{K,m}Q']\hat{Z}'$. These suggest that for the UKFaculty data set, (4.16) holds and it is reasonable to model the UKFaculty with a DCMM model. In summary, in many recent works on network analysis, we frequently assume that a DCMM model holds for the settings at hand, but we rarely checked if such an assumption is valid. Our NMF results provide an approach to checking whether the network satisfies DCMM model.

5 Discussion

We derive a sharp NMF result and apply it to network modeling. Both NMF and network analysis are important areas in machine learning, with applications in image processing, social media, NLP, and cancer study [5, 22, 20]. In comparison, NMF is more theoretically oriented and network analysis is more application oriented. Our paper makes an interesting connection of the two areas. On one hand, we find a new application of NMF theory. This may open the door for a line of research where we find new applications of NMF in areas such as text learning [20] and tensor analysis [14]. On the other hand, we gain valuable insight on what are the most suitable network models in applications. This is crucial, for a suitable model is the starting point for methods and theory. Our study may help researchers identify the right network models and so can channel their strengths to the right direction. Our work may also help develop new methods. For example, compared to the general rank-K model, the DCMM model has more structures which we can exploit (see [16, 18] where they discovered a simplex structure in the spectral domain, using some specific features which the DCMM model has but a general rank-K model does not). Our approach is useful for it ensures us that in certain settings, we can use a more specific model and exploit the structures the model provides.

Another point is that, existing NMF theory usually requires some crucial conditions. However, whether such conditions are reasonable in real applications remains unclear, especially when the conditions are on matrices that are not directly observable. In Section 3-4, we tackle this problem by providing (a) a detailed explanation for why our NMF assumptions are reasonable in network analysis and (b) new ideas for checking the NMF conditions in real applications when the NMF conditions are on matrices that are not directly observable. We hope our efforts many spark some new research along this line.

Figure 1: Histograms of $\{\hat{a}_i\}_{i=1}^n$ (left) and $\{\hat{z}_i\}_{i=1}^n$ (right). The results suggest that all $\hat{a}_i$ are smaller than 0.5 (note that $1/(K-1) = .5$ as $K = 3$) so condition (4.16) is satisfied, and that all $\hat{z}_i$ are non-negative so the matrix $\widehat{Z}$ is non-negative. See above for definitions of $\hat{a}_i$ and $\hat{z}_i$.

References

[1] Edoardo M. Airoldi, David M. Blei, Stephen E. Fienberg, and Eric P. Xing. Mixed membership stochastic blockmodels. *J. Mach. Learn. Res.*, 9:1981–2014, 2008.

[2] Animashree Anandkuma, Rong Ge, Daniel Hsu, and Sham Kakade. A tensor approach to learning mixed membership community labels. *J. Mach. Learn. Res.*, 15(1), 2014.

[3] Damjana Kokol Bukovsek and Smigoc Helena. Semi symmetric non-negative matrix factorization. *arXiv.2106.14437*, 2021.

[4] Sourav Chatterjee. Matrix estimation by universal singular value thresholding. *Ann. Statist.*, 43(1):177–214, 2015.

[5] David L. Donoho and Victoria Stodden. When does non-negative matrix factorization give a correct decomposition into parts? *Advances in Neural Information Processing Systems 16*, 2003.

[6] Bradley Efron. Fisher in the 21st century. *Statist. Sci.*, 13(2):95–114, 1998.

[7] K. Kashino H. Kameoka, N. Ono and S. Sagayama. Complex nmf: a new sparse representation for acoustic signals. *Proc. of IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, 2019.

[8] Zhaoshui He, Shengli Xie, Rafal Zdunek, Guoxu Zhou, and Andrzej Cichocki. Symmetric nonnegative matrix factorization: Algorithms and applications to probabilistic clustering. *IEEE Transactions on Neural Networks*, 22(12):2117–2131, 2011.

[9] Paul W. Holland, Kathryn Blackmond Laskey, and Samuel Leinhardt. Stochastic blockmodels: First steps. *Soc. Netw.*, 5(2):109 – 137, 1983.

[10] Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge University Press, 2nd edition, 2013.

[11] Pengsheng Ji, Jiashun Jin, Zheng Tracy Ke, and Wanshan Li. Co-citation and co-authorship networks for statisticians (with discussions). *J. Bus. Econ. Statist.*, 40(2):469–485, 2022.

[12] Jiashun Jin. Fast community detection by SCORE. *Ann. Statist.*, 43(1):57–89, 2015.

[13] Jiashun Jin and Zheng Tracy Ke. The SCORE normalization, especially for heterogeneous network and text data. *Manuscript*, 2021.

[14] Jiashun Jin, Zheng Tracy Ke, and Jiajun Liang. Sharp impossibility results for hypergraph testing. *Manuscript*, 2021.

[15] Jiashun Jin, Zheng Tracy Ke, and S. Luo. Optimal adaptivity of signed-polygon statistics for network testing. *Ann. Statist.*, 49(6):3408–3433, 2021.

[16] Jiashun Jin, Zheng Tracy Ke, and Shengming Luo. Estimating network memberships by simplex vertex hunting. *arXiv:1708.07852*, 2017.

[17] Jiashun Jin, Zheng Tracy Ke, and Shengming Luo. Improvements on score: especially for weak signals. *Sankhya A*, pages 1–36, 2021.

[18] Jiashun Jin, Zheng Tracy Ke, Shengming Luo, and Minzhe Wang. Optimal estimation of the number of network communities. *J. Amer. Statist. Assoc.*, pages 1–16, 2022.

[19] Brian Karrer and Mark E. J. Newman. Stochastic blockmodels and community structure in networks. *Phys. Rev. E*, 83:016107, 2011.

[20] Zheng Tracy Ke and Minzhe Wang. A new SVD approach to optimal topic estimation. *arXiv:1704.07016*, 2017.

[21] Adamic L. and Glance N. The political blogosphere and the 2004 u.s. election: Divided htey blog. *Proceeding of the 3rd International Workshop on Link Discovery*, pages 36–43, 2005.

[22] Daniel D Lee and H Sebastian Seung. Learning the parts of objects by non-negative matrix factorization. *Nature*, 401(6755):788–791, 1999.

[23] X Mao, Sarkar P, and D Chakrabarti. On mixed memberships and symmetric nonnegative matrix factorizations. *International conference on machine learning*, pages 2324–2333, 2017.

[24] Patrick Rubin-Delanchy, Joshua Cape, Minh Tang, and Carey E Priebe. A statistical interpretation of spectral embedding: the generalised random dot product graph. *arXiv.1709.05506*, 2021.

[25] Naomi Shaked-Monderer and Abraham Berman. *Composite and completely positive matrices*. World Scientific, 2021.

[26] Wasin So and Changqing Xu. When does cp-rank equal rank? *arXiv.1308.3193*, 2013.

[27] S. Young and E. Scheinerman. Random dot product graph models for social networks. *Procceddings of the 5th international conference on algorithms and models for web-graph*, pages 138–149, 2007.

[28] Yuan Zhang, Elizaveta Levina, and Ji Zhu. Detecting overlapping communities in networks using spectral methods. *SIAM J. Math. Data Sci.*, 2(2):265–283, 2020.

[29] Yunpeng Zhao, Elizaveta Levina, and Ji Zhu. Consistency of community detection in networks under degree-corrected stochastic block models. *Ann. Statist.*, 40(4):2266–2292, 2012.