On the detrimental effect of invariances in the likelihood for variational inference

Abstract

Variational Bayesian posterior inference often requires simplifying approximations such as mean-field parametrisation to ensure tractability. However, prior work has associated the variational mean-field approximation for Bayesian neural networks with underfitting in the case of small datasets or large model sizes. In this work, we show that invariances in the likelihood function of over-parametrised models contribute to this phenomenon because these invariances complicate the structure of the posterior by introducing discrete and/or continuous modes which cannot be well approximated by Gaussian mean-field distributions. In particular, we show that the mean-field approximation has a gap in the evidence lower bound objective compared to a purpose-built posterior that takes into account the known invariances. Importantly, the invariance gap is not constant; it vanishes as the approximation reverts to the prior. We proceed by first considering translation invariances in a linear model with a single data point in detail. We show that, while the true posterior can be constructed from a mean-field parametrisation, this is achieved only if the objective function takes into account the invariance gap. Then, we transfer our analysis of the linear model to neural networks. Our analysis provides a framework for future work to explore solutions to the invariance problem.

1 Introduction

Bayesian neural networks (BNNs) have several appealing advantages compared to deterministic neural networks (NN) such as improving generalization [35], capturing epistemic uncertainty [16], and providing a framework for continual learning methods [17, 21]. Unfortunately, reaping these theoretical benefits has thus far been impeded [34]. In particular, several practical issues with variational Bayesian inference methods—which are the de-facto standard technique for scaling inference in BNNs to large datasets [14, 1]—have been identified to be partially responsible for this performance gap [15]. The most common variational approximation of the posterior is a product of independent Normal distributions, commonly referred to as the *mean-field* approximation. It has been observed that mean-field variational BNNs (VBNNs) suffer from severe underfitting for large models and small dataset sizes [12]. Recent work [4] showed that under certain assumptions such as an odd Lipschitz activation function and a finite dataset with bounded likelihood, the optimal mean-field approximation *collapses* to the prior as the NN width increases, *ignoring the data*.

In this work, we study the invariances of the likelihood function and the resulting non-identifiability in overparametrised models. We identify it as the primary reason why mean-field variational distributions cannot approximate the posterior, ultimately leading to the reversion to the prior analysed in [4]. It is well known that NNs exhibit several invariances *with respect to their parameters*, including node permutation invariance and sign-flip invariance (in the case of odd activation functions) [26, 2], and scaling invariance in the case of piece-wise linear activations [23]. Recent work showed that the permutation invariance introduces an additional gap in the evidence lower bound (ELBO) for

a mean-field approximation compared to an (intractable) mixture that includes all modes [18]. We show that the parameter space of BNNs additionally has subspaces with *translation invariance*.

Invariance in the likelihood function does not necessarily pose a challenge if maximum likelihood point estimation through stochastic gradient descent is used, because convergence to any of the equivalent optima (resulting from the invariance) suffices for prediction. However, we show that the variational Bayesian approach is particularly impaired by invariances in the likelihood, since the resulting posterior has a complicated structure that can not be well approximated by the mean-field approximation. In order to isolate the impact of the invariances, we define a posterior approximation in Sec. 3 that explicitly models the invariances by integrating over all transformations that leave the likelihood invariant. We *construct* this *invariance-abiding* distribution from a *mean-field likelihood* approximation such that it induces an identical posterior predictive distribution compared to an alternative *mean-field posterior* approximation that is agnostic to the invariances. Both approximations are parametrised with the same variational parameters and result in the same expected log-likelihood, but differ in their modelling of the invariances, and, hence, the corresponding ELBO objective differs only in the KL regularisation term. We refer to the difference between these two KL terms as the *invariance-gap*. While our constructed approximation is intractable in general, it serves as a useful tool for studying the invariance in simple models where it is indeed tractable, and it provides a framework for future work to devise practical remedies.

The main contribution of this work is a detailed analysis (Sec. 4) of a linear model that implements *translation invariance* and explains the reversion of the posterior to the prior as the result of optimizing the ELBO that exhibits a non-trivial invariance-gap. This gap vanishes for approximations that collapse to the prior and grows linearly in the number of dimensions with translation invariance for mean-field approximations that have the same posterior predictive distribution as the true posterior. We transfer our analysis of the linear model to VBNNs in Sec. 5, showing that subspaces in BNNs exist that exhibit translation invariance. We start by introducing the basic concepts and recent results on which we build our contribution in Sec. 2.

2 Background

2.1 Variational Bayesian Neural Networks

Neural network functional model. Deep NNs are layered models that progressively transform their inputs in each layer. More formally, an L–layer NN computes algebraically

$$f(\mathbf{x}) = h_L \left(\mathbf{w}_{L,1}^{\mathrm{T}} \mathbf{z}_{L-1} \right) , \quad z_{1,i} = h_1 \left(\mathbf{w}_{1,i}^{\mathrm{T}} \mathbf{x} \right) , \quad z_{l,i} = h_l \left(\mathbf{w}_{l,i}^{\mathrm{T}} \mathbf{z}_{l-1} \right) ,$$

where the $h_l : \mathbb{R} \to \mathbb{R}$ are monotonic transfer functions that introduce non-linearities. Such a network has $n_1 + n_2 + \cdots + n_L$ many nodes $z_{l,i}$ which are each a weighted linear combination of either the input vector $\mathbf{x}$ (first hidden layer) or the value of the hidden units $\mathbf{z}_l$ (all other hidden layers and the output $f(\mathbf{x})$). We use $\mathbf{w} = [\mathbf{w}_1^{\mathrm{T}}, \ldots, \mathbf{w}_{n_1+\cdots+n_L}^{\mathrm{T}}]^{\mathrm{T}}$ to denote all learnable parameters of the model.

Variational Bayesian treatment. If the weights and biases $\mathbf{w}$ are treated as random variables with a prior distribution $p(\mathbf{w})$, then the posterior $p(\mathbf{w} \,|\, \mathcal{D})$ (induced by the dataset $\mathcal{D}$ through the likelihood function $\ell(\mathbf{w}; \mathcal{D}) := p(\mathcal{D} \,|\, \mathbf{w})$ defined via $f(\mathbf{x})$) is referred to as a Bayesian neural network (BNN). The variational Bayesian method approximates the posterior by a distribution $q_{\boldsymbol{\theta}}(\mathbf{w}) \approx p(\mathbf{w} \,|\, \mathcal{D})$ with variational parameters $\boldsymbol{\theta}$, casting inference as an optimization problem

$$q_{\boldsymbol{\theta}}^*(\mathbf{w}) = \underset{q_{\boldsymbol{\theta}} \,\in\, \mathcal{Q}}{\operatorname{argmin}} \, \mathrm{KL} \left[q_{\boldsymbol{\theta}} \,||\, p(\cdot \,|\, \mathcal{D}) \right] , \tag{1}$$

where $\mathrm{KL} \left[q_{\boldsymbol{\theta}} \,||\, p(\cdot \,|\, \mathcal{D}) \right] := \mathbb{E}_{\mathbf{w} \sim q_{\boldsymbol{\theta}}} \left[\ln q_{\boldsymbol{\theta}}(\mathbf{w}) - \ln p(\mathbf{w}|\mathcal{D}) \right]$ and $\mathcal{Q}$ is a family of distributions over $\mathbf{w}$. The optimization of (1) is achieved by maximising a lower bound to the (log) model evidence $\ln p(\mathcal{D})$,

$$\mathcal{L}_{\mathrm{ELBO}} \left(q_{\boldsymbol{\theta}}, \mathcal{D} \right) = \ln p(\mathcal{D}) - \mathrm{KL} \left[q_{\boldsymbol{\theta}} \,||\, p(\cdot \,|\, \mathcal{D}) \right] = \mathbb{E}_{\mathbf{w} \sim q_{\boldsymbol{\theta}}} \left[\ln \frac{\ell(\mathbf{w}; \mathcal{D}) \, p(\mathbf{w})}{q_{\boldsymbol{\theta}}(\mathbf{w})} \right]$$
$$= \mathrm{ELL} \left(q_{\boldsymbol{\theta}}, \, \mathcal{D} \right) - \mathrm{KL} \left[q_{\boldsymbol{\theta}} \,||\, p \right] , \tag{2}$$

where $\mathrm{ELL} \left(q_{\boldsymbol{\theta}}, \, \mathcal{D} \right) := \mathbb{E}_{\mathbf{w} \sim q_{\boldsymbol{\theta}}} \left[\ln \ell(\mathbf{w}; \mathcal{D}) \right]$ is the expected log-likelihood.

Definition 1 (Mean-field variational BNN). A *mean-field* variational BNN (VBNN) is the minimizer of (1) where $\mathcal{Q}$ is the family of Gaussians with diagonal covariance matrix.

2.2 Data-related bound on the KL divergence

The term $\mathrm{KL}\left[q_{\boldsymbol{\theta}} \,\|\, p\right]$ in (2) admits a data-dependent upper bound that naturally occurs as a consequence of the finite information provided by a finite dataset in the presence of noise. This result has been shown in [4] and we recall the result for a Gaussian likelihood with homogeneous noise (for other likelihood functions, we refer to App. F in [4]).

Assume a VBNN with Gaussian observation noise defined through $y = f_{\mathbf{w}}(\mathbf{x}) + \epsilon$, with $\epsilon \sim \mathcal{N}(0, \sigma_y)$, an isotropic Gaussian prior $p(\mathbf{w}) = \mathcal{N}(\mathbf{w}; \mathbf{0}, \mathbf{I})$, and a mean-field variational approximation $q_{\boldsymbol{\theta}}(\mathbf{w}) = \mathcal{N}(\mathbf{w}; \mathbf{m}, \mathrm{Diag}(\mathbf{v}))$. Define the NN output variance $\sigma_L\left(\mathbf{x}^{(n)}\right) := \mathbb{V}_{\mathbf{w} \sim p}\left[f(\mathbf{x}^{(n)}; \mathbf{w})\right]$ for some fixed input $\mathbf{x}^{(n)}$, where L indicates the last layer. Then,

$$\mathrm{KL}\left[q_{\boldsymbol{\theta}} \,\|\, p\right] \leq \mathrm{ELL}\left(q^*, \mathcal{D}\right) - \mathrm{ELL}\left(p, \mathcal{D}\right) = \sum_{n=1}^{N} \frac{\sigma_L\left(\mathbf{x}^{(n)}\right) + \left(y^{(n)}\right)^2}{2\sigma_y}, \tag{3}$$

where q^* is a hypothetical approximation that predicts the data perfectly up to the noise variance σ_y (see App. F for details). Prior work [4] has used the data-related bound to prove *that* the predictive distribution of mean-field BNNs converges to the prior predictive distribution if the network width is large and the activation function is odd, ultimately resulting in posterior collapse to the prior. In contrast, we address the question *why* the mean-field approximation cannot approximate the posterior by relating this data-related bound to invariances in neural network functions in the following section.

3 Invariance-abiding variational approximation

We develop a framework that enables modelling the invariances in the likelihood and understanding their impact on the ELBO objective. To achieve this, we approximate the likelihood by a Gaussian function with variational parameters and marginalise over all transformations to which the true likelihood is invariant. By taking the product with the prior, we construct an *invariance-abiding* posterior approximation q_{mix} which can be related to a *mean-field* approximation q_0 that does not model invariances. We then describe conditions under which both approximations yield an identical posterior predictive, while the KL regularisation term of q_0 is lower bounded by the KL of q_{mix}.

Variational likelihood and posterior approximations. Non-identifiability implies that the posterior does not concentrate on a single set of parameters irrespective of the dataset size, because the same likelihood is assigned to different parameter values [11]. We model this invariance through transformations $t(\cdot, \mathbf{r})$ to which the likelihood $\ell(\mathbf{w}; \mathcal{D}) = p(\mathcal{D} \,|\, \mathbf{w})$ is invariant via variables $\mathbf{r}$:

$$\forall \mathbf{r} \sim p(\mathbf{r}): \quad \ell(t(\mathbf{w}, \mathbf{r}); \mathcal{D})) = \ell(\mathbf{w}; \mathcal{D}). \tag{4}$$

From (4) it follows that the likelihood is also invariant wrt. the marginalisation

$$\ell\left(\mathbf{w}; \mathcal{D}\right) = \mathbb{E}_{\mathbf{r} \sim p}\left[\ell\left(t\left(\mathbf{w}, \mathbf{r}\right); \mathcal{D}\right)\right].$$

We assume that each of the equivalent parametrisations $\mathbf{w}' = t(\mathbf{w}, \mathbf{r})$ of the likelihood has the same probability a priori. For discontinuous transformations such as node permutations (see App. E), $p(\mathbf{r})$ is a uniform distribution over discrete variables index these transformations. For the continuous translation invariance (see Sec. 4), we model the uniform distribution as a Gaussian with infinite variance. It is conceivable that the likelihood can be constructed by marginalising over these transformations of a simpler function $\ell_0(\mathbf{w}; \mathcal{D})$ such that $\ell(\mathbf{w}; \mathcal{D}) = \mathbb{E}_{\mathbf{r} \sim p}\left[\ell_0(t(\mathbf{w}, \mathbf{r}); \mathcal{D})\right]$. For instance, permutation invariance induces a factorial number of discontinuous modes that are each equivalent (cf. [18], App. E); and as we show in Sec. 4, the full covariance Gaussian likelihood and posterior of an over-parametrised Bayesian linear regression model can be constructed from a product of independent Gaussians. Curiously, it may be sufficient to approximate ℓ_0 while taking into account the known invariances. To this end, we define a Gaussian *variational likelihood approximation* $g_0(\mathbf{w}; \boldsymbol{\theta}) \approx \ell_0(\mathbf{w}, \mathcal{D})$, with variational parameters $\boldsymbol{\theta}$. From this single mode approximation, we model an *invariance-abiding likelihood* approximation from the same parametrisation through

$$g_{\mathrm{mix}}(\mathbf{w}; \boldsymbol{\theta}) := \mathbb{E}_{\mathbf{r} \sim p}\left[g_0(t(\mathbf{w}, \mathbf{r}); \boldsymbol{\theta})\right] \approx \mathbb{E}_{\mathbf{r} \sim p}\left[\ell_0(t(\mathbf{w}, \mathbf{r}), \mathcal{D})\right] = \ell(\mathbf{w}; \mathcal{D}). \tag{5}$$

We consider mean-field Gaussians for the prior $p(\mathbf{w})$ and likelihood approximation $g_0(\mathbf{w}; \boldsymbol{\theta})$, where $\boldsymbol{\theta} = \{\mathbf{m}, \boldsymbol{\lambda}\}$ are means and variances of g_0. We then define a *mean-field posterior* as the product

$$q_0(\mathbf{w}; \boldsymbol{\theta}) := Z_0^{-1} \, p(\mathbf{w}) \cdot g_0(\mathbf{w}; \boldsymbol{\theta}), \quad Z_0 = \int p(\mathbf{w}) \cdot g_0(\mathbf{w}; \boldsymbol{\theta}) d\mathbf{w}. \tag{6a}$$

Similarly, we define an *invariance-abiding posterior* as the product of prior and invariant likelihood:

$$q_{\mathrm{mix}}(\mathbf{w};\boldsymbol{\theta}) := Z_{\mathrm{mix}}^{-1}\, p(\mathbf{w}) \cdot g_{\mathrm{mix}}(\mathbf{w};\boldsymbol{\theta}), \quad Z_{\mathrm{mix}} = \int p(\mathbf{w}) \cdot g_{\mathrm{mix}}(\mathbf{w};\boldsymbol{\theta})d\mathbf{w}, \quad (6b)$$

where Z_0 and Z_{mix} are normalisation constants. While $q_0(\mathbf{w};\boldsymbol{\theta})$ is a mean-field approximation, the invariance-abiding approximation $q_{\mathrm{mix}}(\mathbf{w};\boldsymbol{\theta})$ is a mixture with infinite continuous or finite discrete modes depending on the type of invariance. We will describe continuous translation invariance in Sec. 4; for the discrete node permutation invariance, see App. E of the supplementary material.

Posterior predictive equivalence. Next, we discuss conditions under which we can construct approximations $q_{\mathrm{mix}}(\mathbf{w};\boldsymbol{\theta})$ and $q_0(\mathbf{w};\boldsymbol{\theta})$ that yield the same predictive distribution. We first write the density q_{mix} as an expectation of product densities,

$$q_{\mathrm{mix}}(\mathbf{w};\boldsymbol{\theta}) = Z_{\mathrm{mix}}^{-1}\, p(\mathbf{w}) \cdot \int p(\mathbf{r})\, g_0(t(\mathbf{w},\mathbf{r});\boldsymbol{\theta})d\mathbf{r} = \int p(\mathbf{r})\frac{p(\mathbf{w})\cdot g_0(t(\mathbf{w},\mathbf{r});\boldsymbol{\theta})}{Z_{\mathrm{mix}}}d\mathbf{r}.$$

Then, we assume that there exists a mapping $\mathbf{r}' = \varphi(\mathbf{r})$ such that

$$\forall \mathbf{r} \sim p(\mathbf{r}) : p(\mathbf{w}) \cdot g_0(t(\mathbf{w},\mathbf{r});\boldsymbol{\theta}) = p(t(\mathbf{w},\varphi(\mathbf{r}))) \cdot g_0(t(\mathbf{w},\varphi(\mathbf{r}));\boldsymbol{\theta}). \quad (7)$$

This assumption allows us to exploit the invariance property of the likelihood (approximation) also for each product density $q_0(t(\mathbf{w},\varphi(\mathbf{r}));\boldsymbol{\theta}) = p(t(\mathbf{w},\varphi(\mathbf{r}))) \cdot g_0(t(\mathbf{w},\varphi(\mathbf{r}));\boldsymbol{\theta})$, because then

$$q_{\mathrm{mix}}(\mathbf{w};\boldsymbol{\theta}) = \int \frac{Z_0(\mathbf{r})}{Z_{\mathrm{mix}}}p(\mathbf{r}) \cdot q_0(t(\mathbf{w},\varphi(\mathbf{r}));\boldsymbol{\theta})d\mathbf{r},$$

where the normalisation constants are $Z_0(\mathbf{r}) = \int p(\mathbf{w}) \cdot g_0(t(\mathbf{w},\varphi(\mathbf{r}));\boldsymbol{\theta})d\mathbf{w}$, and Z_{mix} is defined in (6b). We show that the condition defined in (7) holds for translation and permutation invariance in App. C of the supplementary material.

The second condition is that all invariance transformations $t(\cdot,\mathbf{r})$ must be volume-preserving, i.e.

$$\forall \mathbf{r} \sim p(\mathbf{r}) : \left| \det \frac{\partial t(\mathbf{w},\mathbf{r})}{\partial \mathbf{w}} \right|^{-1} = 1. \quad (8)$$

Although (7) and (8) are fairly restricting, common invariances such as translation and permutation invariance fulfil this condition for Gaussian priors and likelihood approximations (see App. C). With (7) and (8), we can show the posterior predictive equivalence (see Lemma 1 in App. C)

$$\mathbb{E}_{\mathbf{w}\sim q_{\mathrm{mix}}}\left[\ln p(\mathcal{D}\,|\,\mathbf{w})\right] = \mathbb{E}_{\mathbf{w}\sim q_0}\left[\ln p(\mathcal{D}\,|\,\mathbf{w})\right]. \quad (9)$$

Invariance gap. If the conditions in (7) and (8) are met, we also show (see Lemma 2 in App. C)

$$\mathcal{L}_{\mathrm{ELBO}}\left(q_0,\mathcal{D}\right) - \mathcal{L}_{\mathrm{ELBO}}\left(q_{\mathrm{mix}},\mathcal{D}\right) = \mathrm{KL}\left[q_0\,||\,p\right] - \mathrm{KL}\left[q_{\mathrm{mix}}\,||\,p\right] = \mathrm{KL}\left[q_0\,||\,q_{\mathrm{mix}}\right]. \quad (10)$$

Interestingly, the gap between the two respective ELBO objectives is given exactly by the relative entropy between the mean-field and invariance-abiding approximation; we therefore refer to it as the *invariance gap*. We make the following observation about the detrimental effect of the invariances: when maximising the standard ELBO $\mathcal{L}_{\mathrm{ELBO}}\left(q_0,\mathcal{D}\right)$ instead of the tighter objective $\mathcal{L}_{\mathrm{ELBO}}\left(q_{\mathrm{mix}},\mathcal{D}\right)$ wrt. the parameters of the variational *likelihood* approximation $g_0(\mathbf{w};\boldsymbol{\theta})$ (which is used to construct both q_0 and q_{mix}), we see that the former objective favors solutions where q_0 and q_{mix} coincide. One such suboptimal solution is when the likelihood is uninformative (uniform), and, thus, both q_0 and q_{mix} revert to the prior. In Sec. 4, we study a simple model that exhibits translation invariance and show that this additional gap is significant.

Implication of data-related bound. Since $\mathrm{KL}\left[q_0\,||\,p\right]$ is upper bounded by the best- and worst-case ELL as described in Sec. 2.2, the invariance gap is also bounded (using (10) and (3)):

$$\mathrm{KL}\left[q_0\,||\,q_{\mathrm{mix}}\right] \le \mathrm{KL}\left[q_0\,||\,p\right] \le \mathrm{ELL}\left(q^*,\mathcal{D}\right) - \mathrm{ELL}\left(p,\mathcal{D}\right). \quad (11)$$

Consequently, this bound puts a constraint on the achievable solutions to the maximisation of the ELBO objective wrt. the variational parameters of the mean-field approximation $q_0(\mathbf{w};\boldsymbol{\theta})$ One particular parametrization that guarantees this behaviour is the mean-field posterior collapse, $q_0(\mathbf{w};\boldsymbol{\theta}) \approx p(\mathbf{w})$ introduced above. However, in order to show that the invariance-gap not only

159 admits posterior collapse as a potential parametrization but indeed incentivises posterior collapse, we
160 next tackle the question how the invariance-gap grows for non-collapsed approximations.

161 We observe in the subsequent sections that the invariance-gap increases with the model size
162 for non-collapsed approximations and the invariances we consider. Examples of the invariance
163 include permutation invariance (interchangeability of neurons in the same layer of a BNN) in the
164 supplementary material. We show that $\mathrm{KL}\left[q_0 \,\|\, q_{\mathrm{mix}}\right]$ is between zero and the logarithm of the
165 number of discrete permutation-induced modes, which grows factorial with the number of neurons
166 per layer. Another example and the focus of the main body of this work is continuous translation
167 invariance. In Sec. 4, we analyse translation invariance in over-parametrised linear models, show that
168 the gap grows linear with the number of dimensions (weights) for parameter values $\boldsymbol{\theta}_{\mathrm{mix}}^*$ that are
169 optimal in terms of the posterior predictive and the ELBO optimized wrt. $q_{\mathrm{mix}}(\mathbf{w}, \boldsymbol{\theta})$.

4 Translation invariance in linear models

171 We study an over-parametrised Bayesian linear regression model as the canonical model that exhibits
172 *translation invariance.* The model serves as a useful tool to understand the detrimental effect of
173 translation invariance, since all interesting quantities can be computed analytically, incl. the mean-
174 field and invariance-abiding distribution defined in (6), and the invariance gap from (10). We show
175 how to lift results from this canonical model to the more general case of NNs in Sec. 5. We start by
176 specifiying the canonical model in the following.

177 **Likelihood model.** Consider a linear model with K latent variables $\mathbf{w} = [w_1, \ldots, w_K]^{\mathrm{T}}$ and
178 assume that we have N observations of variables y given dependent inputs $\mathbf{x}$. To simplify the setting,
179 we will assume that $\mathbf{x} = \mathbf{1}$ (see App. D in the supplementary material for the more general case). We
180 further assume that the observation depends only on the inner product with the inputs and additive
181 Gaussian noise. That is, $y = \frac{1}{K}\mathbf{1}^{\mathrm{T}}\mathbf{w} + \epsilon, \quad \epsilon \sim \mathcal{N}(0, \sigma_y)$. Note that any change in $\mathbf{w}$ which leaves
182 the sum of the elements unaffected does not change the likelihood. In general, we can model this
183 translation invariance using a $K - 1$ dimensional vector $\boldsymbol{\Delta} \in \mathbb{R}^{K-1}$ and observing that

$$\mathbf{1}^{\mathrm{T}}\mathbf{w} = \mathbf{1}^{\mathrm{T}}(\mathbf{w}+\mathbf{B}\boldsymbol{\Delta}), \quad \mathbf{B} := \left[\begin{array}{c} \mathbf{I} \\ -\mathbf{1}^{\mathrm{T}} \end{array} \right], \tag{12}$$

184 since $\mathbf{1}^{\mathrm{T}}\mathbf{B}\boldsymbol{\Delta} = \mathbf{0}$. This over-parametrised model exhibits translation invariance $t(\mathbf{w}, \Delta) = \mathbf{w} - \mathbf{B}\Delta$.

185 **Prior.** We assume a Gaussian prior $p(\mathbf{w}) = \mathcal{N}(\mathbf{w}; \boldsymbol{\mu}, \boldsymbol{\Sigma})$ with diagonal covariance $\boldsymbol{\Sigma} = \mathrm{Diag}(\boldsymbol{\sigma})$,
186 where we consider $\boldsymbol{\sigma} = K \cdot \sigma_0 \cdot \mathbf{1}$ proportional to the number of dimension K such that the predictive
187 variance $\mathbb{V}\left[\frac{1}{K}\sum_k w_k + \epsilon\right]$ is constant for different K. Another way to view this is to model a prior
188 over parameters that induces the same prior over functions for different K.

189 **Posterior.** The posterior of this Gaussian linear model can be computed as (see App. D.3)

$$p(\mathbf{w} \,|\, \mathbf{y}) = \mathcal{N}\left(\mathbf{w}; \mathbf{m}_p^*, \mathbf{V}_p^*\right), \quad \mathbf{V}_p^* = \left(\frac{N}{K^2\sigma_y}\mathbf{1}\mathbf{1}^{\mathrm{T}} + \boldsymbol{\Sigma}^{-1}\right)^{-1}, \quad \mathbf{m}_p^* = \mathbf{V}_p^*\frac{\sum_i y_i}{K\sigma_y}\mathbf{1}. \tag{13}$$

190 As can be seen, the resulting posterior has full covariance with a diagonal plus rank-1 matrix. We will
191 now show that we can construct this posterior from the prior and a mean-field Gaussian approximation
192 of the likelihood by marginalising over all translations $\Delta \sim p(\Delta)$ to which the likelihood is invariant.

4.1 Mean-field parametrisation of the likelihood function

194 Next, we construct the mean-field and invariance-abiding posterior approximations defined in (6)
195 from the prior defined above and the mean-field likelihood approximation with locations $\mathbf{m}$ and
196 variances $\boldsymbol{\lambda}$. We model that the likelihood is translation invariant by computing the marginal from (5)
197 and consider $p(\Delta) = \mathcal{N}\left(\boldsymbol{\Delta}; \mathbf{0}, \beta^2\mathbf{I}\right)$ with $\beta \to \infty$ as the uniform distribution over all translations:

$$g_{\mathrm{mix}}(\mathbf{w}; \boldsymbol{\theta}) = \int g_0(t(\mathbf{w}, \Delta); \boldsymbol{\theta}) \cdot p(\boldsymbol{\Delta}) \, d\boldsymbol{\Delta}$$

$$= \lim_{\beta \to \infty} \int \mathcal{N}(\mathbf{w}; \mathbf{m} + \mathbf{B}\boldsymbol{\Delta}, \mathrm{Diag}(\boldsymbol{\lambda})) \cdot \mathcal{N}\left(\boldsymbol{\Delta}; \mathbf{0}, \beta^2\mathbf{I}\right) \, d\boldsymbol{\Delta} =: \mathcal{N}(\mathbf{w}; \mathbf{m}_{\mathrm{mix}}, \mathbf{V}_{\mathrm{mix}}).$$

Writing the uniform distribution as the limiting case of a Gaussian with infinite variance allows us compute the integral analytically. As shown in App. D, the resulting function is a multivariate Gaussian with a degenerate rank-1 covariance matrix and the same location parameter as $g_0(\mathbf{w})$:

$$\mathbf{m}_{\mathrm{mix}} = \mathbf{m}, \qquad \mathbf{V}_{\mathrm{mix}}^{-1} = \frac{1}{\mathbf{1}^{\mathrm{T}}\boldsymbol{\lambda}}\mathbf{1}\mathbf{1}^{\mathrm{T}}. \tag{14}$$

However, the invariance-abiding posterior is non-degenerate; it can be computed efficiently as

$$q_{\mathrm{mix}}\left(\mathbf{w};\boldsymbol{\theta}\right) = \mathcal{N}\left(\mathbf{w};\boldsymbol{\mu} + \frac{\mathbf{1}^{\mathrm{T}}\left(\mathbf{m}-\boldsymbol{\mu}\right)}{\mathbf{1}^{\mathrm{T}}\left(\boldsymbol{\lambda}+\boldsymbol{\sigma}\right)}\boldsymbol{\sigma}, \mathrm{Diag}\left(\boldsymbol{\sigma}\right) - \frac{1}{\mathbf{1}^{\mathrm{T}}\left(\boldsymbol{\lambda}+\boldsymbol{\sigma}\right)}\cdot\boldsymbol{\sigma}\boldsymbol{\sigma}^{\mathrm{T}}\right). \tag{15}$$

As can be seen, this covariance matrix is not diagonal, but has an additive rank-1 term that vanishes as $\lambda \to \infty$. Interestingly, the location parameter of q_{mix} is translated from the *prior* location $\boldsymbol{\mu}$ along the direction of the *prior* variance vector $\boldsymbol{\sigma}$. This is because g_{mix} is uniform along the $K-1$ dimensions hyper-plane determined by its normal vector $\mathbf{1}$; taking the Gaussian product then translates the location in the direction $\mathrm{Diag}(\boldsymbol{\sigma})\mathbf{1} = \boldsymbol{\sigma}$. Note also that the invariance-abiding posterior can be written in the same form as the true posterior in (13) (see App. D.3). The optimal invariance-abiding posterior is thus the true posterior. We visualise the two respective posterior approximations and the corresponding likelihood with two different parametrisations (cf. Sec. 4.3) in Fig. 3, App. D.

4.2 Invariance-gap

In order to quantify the detrimental effect of the translation invariance in the considered linear model we will now compute the invariance-gap in (11). Note again that we assume a scaled standard normal prior with variance $\boldsymbol{\sigma} = K \cdot \sigma_0 \cdot \mathbf{1}$. In order to simplify the form of the KL divergence, we then assume that all variances and means take the same value for in the likelihood (and thus also in the posterior) approximation, i.e.

$$\forall k : \lambda_k = \hat{\lambda},\ m_k = \hat{m},\ \sigma_k = \hat{\sigma},\ \mu_k = \hat{\mu}.$$

This choice is motivated by the fact that the posterior approximation is a function of the sum $\mathbf{1}^{\mathrm{T}}\boldsymbol{\lambda}$ only (cf. (15)), and, hence, it makes no difference for q_{mix}; however, since the prior variance is proportional to $\mathbf{1}$, the Gaussian likelihood resulting in the highest ELBO for the approximation q_0 also has a variance vector proportional to $\mathbf{1}$. Using this assumption, the invariance gap is

$$\mathrm{KL}\left[q_0 \,||\, q_{\mathrm{mix}}\right] = \frac{K-1}{2}\left[\ln\left(\frac{\hat{\sigma}+\hat{\lambda}}{\hat{\lambda}}\right) + \frac{\hat{\lambda}}{\hat{\sigma}+\hat{\lambda}} - 1\right]. \tag{16}$$

This is a convex function in $\phi = \frac{\hat{\sigma}+\hat{\lambda}}{\hat{\lambda}}$ with a minimum at $\phi = 1$; it is minimised as $\hat{\lambda}\to\infty$. It is however not evident from the above whether the gap has a large magnitude compared to the rest of the regularisation term and over-regularises in practice. In the next section, we therefore analyse this term at the optima for the mean-field and the invariance-abiding approximation defined in (17), respectively. The corresponding invariance gaps are visualised in Fig. 1.

4.3 Optimal mean-field and invariance-abiding parametrisations

To better understand the detrimental effect of the invariance gap, we now analyse the ELL and KL terms of the ELBO objective for the mean-field and invariance-abiding variational approximations, respectively. We compare these terms for both distributions with the parameters optimized for both posterior distributions, giving 2×2 combinations. We denote the optimal parameters wrt. the invariance-abiding approximation and the mean-field approximation, respectively, as

$$\boldsymbol{\theta}_{\mathrm{mix}}^{*} = \operatorname*{argmax}_{\boldsymbol{\theta}} \mathcal{L}_{\mathrm{ELBO}}\left(q_{\mathrm{mix}}\left(\cdot\,;\boldsymbol{\theta}\right),\mathcal{D}\right), \tag{17a}$$

$$\boldsymbol{\theta}_{0}^{*} = \operatorname*{argmax}_{\boldsymbol{\theta}} \mathcal{L}_{\mathrm{ELBO}}\left(q_{0}\left(\cdot\,;\boldsymbol{\theta}\right),\mathcal{D}\right). \tag{17b}$$

Perhaps the simplest way compute these optimal parameters is to notice that q_{mix} in (15) can be written in the same form as the true posterior and then simply read out the optimal parameters. This is shown in App. D.3; the resulting optimal variational parameters are

$$g_0(\mathbf{w}\,;\boldsymbol{\theta}_{\mathrm{mix}}^{*}) = \mathcal{N}\left(\mathbf{w};\mathbf{m}_{\mathrm{mix}}^{*},\mathrm{Diag}\left(\boldsymbol{\lambda}_{\mathrm{mix}}^{*}\right)\right), \quad \mathbf{m}_{\mathrm{mix}}^{*} = \frac{1}{N}\sum_{n=1}^{N} y^{(n)}\cdot\mathbf{1}, \quad \boldsymbol{\lambda}_{\mathrm{mix}}^{*} = \frac{K\sigma_y}{N}\cdot\mathbf{1}. \tag{18}$$

(a) $N = 10$ observations.   (b) $N = 100$ observations.

Figure 1: Invariance-gap evaluated at different optima (cf. (17)), i.e. $\mathrm{KL}\left[q_0(\cdot;\boldsymbol{\theta}_{\mathrm{mix}}^*) \,\|\, q_{\mathrm{mix}}(\cdot;\boldsymbol{\theta}_{\mathrm{mix}}^*)\right]$ and $\mathrm{KL}\left[q_0(\cdot;\boldsymbol{\theta}_0^*)\,\|\,q_{\mathrm{mix}}(\cdot;\boldsymbol{\theta}_0^*)\right]$. Prior variances are $\sigma = K\cdot\mathbf{1}$, where K is the number of dimensions; the noise variance is $\sigma_y = (2\pi\mathrm{e})^{-1}$; all observations $y = 1$ and $\mathbf{x} = \mathbf{1}$ are identical. As K increases, the invariance-gap vanishes in case of the optimal parameters $\boldsymbol{\theta}_0^*$. In contrast, the gap for $\boldsymbol{\theta}_{\mathrm{mix}}^*$, which induces the true posterior predictive, grows linearly. As the data-related bound (cf. Sec. 2.2) can not be exceeded by the optimal parameters, $\boldsymbol{\theta}_0^*$ can not coincide with the optimal $\boldsymbol{\theta}_{\mathrm{mix}}^*$.

234 The optimal parameters for the mean-field *posterior* can also be computed analytically (see [31]),
235 since the Gaussian mean-field distribution that minimises the KL divergence to the multivariate
236 Gaussian true posterior is known. The resulting optimal parameters of the mean-field *likelihood* are

$$g_0(\mathbf{w}\,;\,\boldsymbol{\theta}_0^*) = \mathcal{N}\left(\mathbf{w};\mathbf{m}_0^*,\mathrm{Diag}\left(\boldsymbol{\lambda}_0^*\right)\right), \quad \mathbf{m}_0^* = \mathbf{m}_{\mathrm{mix}}^*, \quad \boldsymbol{\lambda}_0^* = \frac{K^2\sigma_y}{N}\,\mathbf{1}. \tag{19}$$

237 As can be seen, the two respective optimal parameters differ by the factor K in the likelihood variance.
238 We then construct the two respective posterior approximations q_0 and q_{mix} with these two optimal
239 parameters and compare the 2×2 combinations in Fig. 2 where we visualise the ELBO terms.

240 Note again that we model the prior variance $\boldsymbol{\sigma} = K\sigma_0 \cdot \mathbf{1}$ such that the prior and posterior over
241 functions are identical for any K. Due to this choice and since $q_{\mathrm{mix}}(\mathbf{w};\boldsymbol{\theta}_{\mathrm{mix}}^*)$ approximates the true
242 posterior exactly, it does not suffer from over-parametrisation. This can be seen by the constant (in K)
243 loss terms in Fig. 2. In contrast, since $\boldsymbol{\lambda}_0^*$ depends quadratically on K and $\boldsymbol{\sigma}$ only linearly, $q_{\mathrm{mix}}(\mathbf{w};\boldsymbol{\theta}_0^*)$
244 *collapses* to the prior as $K \to \infty$. This can be seen e.g. by the shrinking KL regularisation (Fig. 2a)
245 and ELL (in 2b) term. As a consequence, and in line with Coker et al. [4], the posterior predictive
246 variance of the optimal mean-field approximation reverts to the prior predictive variance (Fig. 2d).

247 We have shown that—in the simplified setting of a linear model with dependent observations—the
248 consequence of not handling translation invariance is indeed posterior collapse as $K \to \infty$, while
249 modelling the invariance coincides with the true posterior. An important observation and key takeaway
250 is that while we need to optimise for (17a), the mean-field posterior approximation $q_0(\mathbf{w};\boldsymbol{\theta}_{\mathrm{mix}}^*)$ is
251 sufficient for prediction. Indeed, for the linear model, we could compute the invariance-gap exactly
252 and thereby correct the ELBO objective for the invariances of this model class. For more complex non-
253 linear models such as BNNs, this section may serve as a direction for approximating the invariance
254 gap. To this end, the next section describes the layer-wise translation invariance exhibited by BNNs.

5   Translation invariance in Bayesian neural networks

256 Let us now go back to the general NN model in Sec. 2.1. In order to describe the set of all invariances,
257 note that for each node $z_{l,j}$ in layer l (or the output node y), there is a subspace that keeps the value
258 $z_{l,j}$ (or y) invariant when using the translation-invariance model in (12) because the value itself only
259 depends on the *actual* activation $\mathbf{z}_{l-1}$ (or the input $\mathbf{x}$).

260 More formally, the following equations model all translation invariances of a NN at a data point $\mathbf{x}$:

$$\forall\boldsymbol{\Delta}_{L,1}: \quad f(\mathbf{x}) = h_L\left(\mathbf{w}_{L,1}^{\mathsf{T}}\mathbf{z}_{L-1}\right) = h_L\left(\left(\mathbf{w}_{L,1} + \mathbf{B}_{\mathbf{z}_{L-1}}\boldsymbol{\Delta}_{L,1}\right)^{\mathsf{T}}\mathbf{z}_{L-1}\right), \tag{20}$$

$$\forall j \in \{1,\dots,n_l\},\boldsymbol{\Delta}_{l,j}: \quad z_{l,j} = h_l\left(\mathbf{w}_{l,j}^{\mathsf{T}}\mathbf{z}_{l-1}\right) h_l\left(\left(\mathbf{w}_{l,j} + \mathbf{B}_{\mathbf{z}_{l-1}}\boldsymbol{\Delta}_{l,j}\right)^{\mathsf{T}}\mathbf{z}_{l-1}\right), \tag{21}$$

$$\forall j \in \{1,\dots,n_1\},\boldsymbol{\Delta}_{1,j}: \quad z_{1,j} = h_1\left(\mathbf{w}_{1,j}^{\mathsf{T}}\mathbf{x}\right) = h_1\left(\left(\mathbf{w}_{1,j} + \mathbf{B}_{\mathbf{x}}\boldsymbol{\Delta}_{1,j}\right)^{\mathsf{T}}\mathbf{x}\right), \tag{22}$$

(a) $\mathrm{KL}\,[q\,||\,p]$.

(b) $\mathbb{E}_q\,[\ln p(\mathcal{D}\,|\,\mathbf{w})]$.

(c) $\mathcal{L}_{\mathrm{ELBO}}(q,\mathcal{D}) = \mathbb{E}_q\,[\ln p(\mathcal{D}\,|\,\mathbf{w})] - \mathrm{KL}\,[q\,||\,p]$.

(d) $\mathbb{V}_{\mathbf{w}\sim q}\left[\frac{1}{K}\mathbf{1}^{\mathsf{T}}\mathbf{w}\right] + \sigma_y$.

Figure 2: ELBO loss terms and predictive variance for different posterior approximations. Lines/dots indicate the invariance-abiding/mean-field posterior, respectively. Red/blue indicates the optimal invariance-abiding/mean-field parameters (cf. (17)). Prior variance is $\boldsymbol{\sigma} = K \cdot \mathbf{1}$, and $\sigma_y = 1/\left(2\pi\mathrm{e}\right)$; we used $N = 10$ identical observations with value $y = 1$ and inputs $\mathbf{x} = \mathbf{1}$. Notably, both posterior approximations yield the same predictive distribution as can be seen by the ELL (b) and predictive variance (d). The variance of the mean-field approximation reverts to the prior variance as $K \to \infty$.

where the layer index l ranges from $2,\dots,L-1$ and $\mathbf{B_z}$ is given by

$$\mathbf{B_z} := \left[\begin{array}{ccc} & \mathbf{I} & \\ -\frac{z_1}{z_k} & \cdots & -\frac{z_{k-1}}{z_k} \end{array} \right].$$

In order to efficiently compute the invariance-abiding likelihood q_{mix}, note that it also decomposes over the NN layers and the associated parameters as follows: For the n_1 nodes in the first layer we have

$$q_{\mathrm{mix}}(\mathbf{w}_{1,j}) = \mathcal{N}\left(\mathbf{w}_{1,j}; \boldsymbol{\mu} + \frac{\mathbf{x}^{\mathsf{T}}(\mathbf{m}-\boldsymbol{\mu})}{\mathbf{x}^{\mathsf{T}}(\mathbf{V}+\boldsymbol{\Sigma})\mathbf{x}}(\boldsymbol{\Sigma}\mathbf{x}), \boldsymbol{\Sigma} - \frac{1}{\mathbf{x}^{\mathsf{T}}(\mathbf{V}+\boldsymbol{\Sigma})\mathbf{x}}(\boldsymbol{\Sigma}\mathbf{x})(\boldsymbol{\Sigma}\mathbf{x})^{\mathsf{T}} \right), \quad (23)$$

$$P(z_{1,j}) = \mathbb{E}_{\mathbf{w}\sim q_{\mathrm{mix}}(\mathbf{w}_{1,j})}\left[h_1\left(\mathbf{w}^{\mathsf{T}}\mathbf{x}\right) \right], \quad (24)$$

where $\mathbf{m}$, $\mathbf{V}$, $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$ are constrained to the parts of the overall weight vector that correspond to $\mathbf{w}_{1,j}$. Now, since the invariance-abiding mechanism for nodes in layer l only depends on the *value* $\mathbf{z}_{l-1}$ of the hidden units in layer $n-1$ (see (20) and (21)), we have

$$q_{\mathrm{mix}}(\mathbf{w}_{l,j}) = \mathbb{E}_{\mathbf{z}}\left[\mathcal{N}\left(\mathbf{w}_{l,j}; \boldsymbol{\mu} + \frac{\mathbf{z}^{\mathsf{T}}(\mathbf{m}-\boldsymbol{\mu})}{\mathbf{z}^{\mathsf{T}}(\mathbf{V}+\boldsymbol{\Sigma})\mathbf{z}}(\boldsymbol{\Sigma}\mathbf{z}), \boldsymbol{\Sigma} - \frac{1}{\mathbf{z}^{\mathsf{T}}(\mathbf{V}+\boldsymbol{\Sigma})\mathbf{z}}(\boldsymbol{\Sigma}\mathbf{z})(\boldsymbol{\Sigma}\mathbf{z})^{\mathsf{T}} \right) \right], \quad (25)$$

$$P(z_{l,j}) = \mathbb{E}_{\mathbf{z}}\left[\mathbb{E}_{\mathbf{w}\sim q_{\mathrm{mix}}(\mathbf{w}_{l,j})}\left[h_l\left(\mathbf{w}^{\mathsf{T}}\mathbf{z}_{l-1}\right) \right] \right], \quad (26)$$

where again $\mathbf{m}$, $\mathbf{V}$, $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$ are constrained to the parts of the overall weight vector that correspond to $\mathbf{w}_{l,j}$ and the expectation over $\mathbf{z}$ is taken with respect to $\mathbf{z} \sim P(\mathbf{z}_{l-1})$.

Thus, in a practical application, the invariance-abiding mean-field approximation could e.g. be computed with a layer-by-layer iterative optimization: First, given the data $\mathbf{x}$ and a prior $p(\mathbf{w})$, we use (23) and point estimates for all weight vectors in later layers to optimize the mean $\mathbf{m}$ and diagonal covariance $\mathbf{V}$ of the weights associated with the first layer, i.e., $\{\mathbf{w}_{1,j}|j = 1,\dots,n_1\}$. Then, we can use (24) to generate P samples $\mathbf{z}_{1,n}$ of the activations of the first layer under the (now fitted) optimal $q_{\mathrm{mix}}(\{\mathbf{w}_{1,j}\})$. Next, for each of these samples $\mathbf{z}_{1,n}$ we use (25) and point estimates for all weight vectors in later layers to optimize the mean $\mathbf{m}$ and diagonal covariance $\mathbf{V}$ of the weights associated with the second layer and average them for the optimal q_{mix} of the weights associated with

277 the second layer. Finally, we can use (26) to generate samples $\mathbf{z}_{2,i}$ for the second layer. This process
278 can be repeated until the last layer and the whole procedure is iterated until the whole distribution
279 $q_{\mathrm{mix}}(\mathbf{w})$ does not change in one full optimization loop. The complexity of this procedure is no larger
280 than the optimization of the weight matrices of a single layer because all previous layers are fully
281 characterized by the distribution of the latent activations $\mathbf{z}_l$ of the hidden layers.

6 Related work

283 Poor empirical performance of VBNNs has been reported in several previous works, e.g., [34, 15, 30].
284 For instance, it has been shown that single-layer mean-field VBNNs with ReLU activations can
285 not have large predictive uncertainty between regions of low uncertainty [9]; In contrast, Farquhar
286 et al. [7] argue that mean-field approximations are expressive enough for deep BNNs; they prove
287 a universality result that the predictive distribution of mean-field approximations of BNNs with at
288 least 2 layers of hidden units *can* approximate any true posterior distribution over function values
289 arbitrarily closely. This result seems in conflict with our work and [4] who show that the predictive
290 distribution of mean-field VBNNs reverts to the prior predictive distribution as the network width
291 increases. The discrepancy between these two results is resolved by noting that Farquhar et al. [7]
292 only shows that the expressive power of the model is large enough but not that the approximate
293 inference algorithms will converge to this solution. Our work sheds light on why the mean-field
294 approximation fails to approximate expressive posteriors via the invariance-gap in the ELBO.

295 Other attempts have proposed to approximate the posterior predictive directly, thereby circumventing
296 the non-identifiability issue [25, 19, 33]. While these approaches have shown promising empirical
297 performance, estimating the KL regularisation term in function space is more complicated and can
298 even be ill-defined due to regions of zero prior probability mass [3]. In a similar vein, previous work
299 also proposed to map the BNN prior to a Gaussian process (functional) prior [8, 29].

300 Surprisingly, restricting the parametrisation of the variational posterior results in competitive or
301 even better performance [27, 20, 28, 6]. For example, Dusenberry et al. [6] proposes a variational
302 approximation only for rank-1 factors, resulting in inference of a lower-dimensional subspace; the
303 outer product of these low-rank factors perturb a weight matrix that is treated deterministically
304 through maximum a posteriori (MAP) estimation. This approach avoids the invariance problem
305 associated with variational Bayesian inference since the MAP estimate is not impeded by this problem
306 and the subspace of the variational weights do not possess the same invariances.

307 More broadly, the (non-)identifiability of parameters and latent variables in probabilistic models is a
308 widely-studied topic both from frequentist [24] and Bayesian perspectives [5, 22, 10]. In a recent
309 example, Wang et al. [32] attribute posterior collapse in the context of variational auto-encoders to
310 non-identifiability of latent variables.

7 Conclusion

312 We have associated the posterior collapse phenomenon of mean-field VBNNs with invariances in
313 the likelihood function, in particular translation invariance. While the invariance does not affect
314 the predictive distribution, the approximations of the posterior which abide and do not abide the
315 invariance differ in the KL regularisation term and consequently in the tightness of the ELBO
316 objective. We proved that the objectives of the two approximations differ by the relative entropy (KL)
317 between the standard mean-field approximation and the invariance-abiding distribution. We related
318 this to a data-related bound on the KL regularisation which prevents fits for which the gap is large.

319 We studied over-parametrised Bayesian linear regression as the canonical model that exhibits
320 *translation invariance* and for which the relevant terms can be computed exactly. A detailed analysis
321 of this model confirms our hypothesis that the invariance leads to a significant additional gap in the
322 ELBO objective compared to approximations that model the invariance. It is this very gap which
323 prevents mean-field approximation to achieve the same fit as an invariance-abiding approximation
324 and instead leads to a collapse of the posterior variances to the prior variance.

325 While we can compute the invariance gap for the over-parametrised linear model, we have not yet
326 identified a computationally efficient procedure for layered models or for other invariances. However,
327 our work provides the mathematical tools to address over-regularisation due to invariances. Future

work will focus on approximations of the invariance gap in order to correct the ELBO objective for translation and permutation invariances in general neural network functions.

References

[1] C. Blundell, J. Cornebise, K. Kavukcuoglu, and D. Wierstra. Weight uncertainty in neural networks. In *Proceedings of the 32nd International Conference on International Conference on Machine Learning - Volume 37*, ICML'15, page 1613–1622. JMLR.org, 2015.

[2] J. Brea, B. Simsek, B. Illing, and W. Gerstner. Weight-space symmetry in deep networks gives rise to permutation saddles, connected by equal-loss valleys across the loss landscape. *ArXiv*, abs/1907.02911, 2019.

[3] D. R. Burt, S. W. Ober, A. Garriga-Alonso, and M. van der Wilk. Understanding variational inference in function-space. In *Third Symposium on Advances in Approximate Bayesian Inference*, 2021.

[4] B. Coker, W. P. Bruinsma, D. R. Burt, W. Pan, and F. Doshi-Velez. Wide mean-field variational Bayesian neural networks ignore the data. In *Proceedings of the 25th International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2022.

[5] A. P. Dawid. Conditional independence in statistical theory. *Journal of the Royal Statistical Society. Series B (Methodological)*, 41(1):1–31, 1979. ISSN 00359246.

[6] M. W. Dusenberry, G. Jerfel, Y. Wen, Y.-A. Ma, J. Snoek, K. Heller, B. Lakshminarayanan, and D. Tran. Efficient and scalable Bayesian neural nets with rank-1 factors. In *Proceedings of the 37th International Conference on Machine Learning*, ICML'20. JMLR.org, 2020.

[7] S. Farquhar, L. Smith, and Y. Gal. Liberty or depth: Deep Bayesian neural nets do not need complex weight posterior approximations. In H. Larochelle, M. Ranzato, R. Hadsell, M. Balcan, and H. Lin, editors, *Advances in Neural Information Processing Systems*, volume 33, pages 4346–4357. Curran Associates, Inc., 2020.

[8] D. Flam-Shepherd. Mapping gaussian process priors to Bayesian neural networks. In *Bayesian Deep Learning NeurIPS workshop*, 2017.

[9] A. Foong, D. Burt, Y. Li, and R. Turner. On the expressiveness of approximate inference in Bayesian neural networks. In H. Larochelle, M. Ranzato, R. Hadsell, M. Balcan, and H. Lin, editors, *Advances in Neural Information Processing Systems*, volume 33, pages 15897–15908. Curran Associates, Inc., 2020.

[10] A. E. Gelfand and S. K. Sahu. Identifiability, improper priors, and gibbs sampling for generalized linear models. *Journal of the American Statistical Association*, 94(445):247–253, 1999.

[11] A. Gelman, D. Simpson, and M. Betancourt. The prior can often only be understood in the context of the likelihood. *Entropy*, 19(10), 2017. ISSN 1099-4300. doi: 10.3390/e19100555.

[12] S. Ghosh, J. Yao, and F. Doshi-Velez. Structured variational learning of Bayesian neural networks with horseshoe priors. In J. Dy and A. Krause, editors, *Proceedings of the 35th International Conference on Machine Learning*, volume 80 of *Proceedings of Machine Learning Research*, pages 1744–1753. PMLR, 10–15 Jul 2018.

[13] R. Herbrich. *Learning Kernel Classifiers: Theory and Algorithms*. The MIT Press, 2nd edition edition, 2002.

[14] G. E. Hinton and D. van Camp. Keeping the neural networks simple by minimizing the description length of the weights. In *Proceedings of the Sixth Annual Conference on Computational Learning Theory*, COLT '93, page 5–13. Association for Computing Machinery, 1993. ISBN 0897916115. doi: 10.1145/168304.168306.

[15] P. Izmailov, S. Vikram, M. D. Hoffman, and A. G. G. Wilson. What are Bayesian neural network posteriors really like? In M. Meila and T. Zhang, editors, *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pages 4629–4640. PMLR, 18–24 Jul 2021.

[16] A. Kendall and Y. Gal. What uncertainties do we need in Bayesian deep learning for computer vision? In I. Guyon, U. V. Luxburg, S. Bengio, H. Wallach, R. Fergus, S. Vishwanathan, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 30. Curran Associates, Inc., 2017.

[17] R. Kurle, B. Cseke, A. Klushyn, P. van der Smagt, and S. Günnemann. Continual learning with Bayesian neural networks for non-stationary data. In *8th International Conference on Learning Representations, ICLR 2020, Addis Ababa, Ethiopia, April 26-30, 2020*. OpenReview.net, 2020.

[18] R. Kurle, T. Januschowski, J. Gasthaus, and B. Wang. On symmetries in variational Bayesian neural nets. In *Bayesian Deep Learning NeurIPS workshop*, 2021.

[19] C. Ma, Y. Li, and J. M. Hernandez-Lobato. Variational implicit processes. In K. Chaudhuri and R. Salakhutdinov, editors, *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pages 4222–4233. PMLR, 09–15 Jun 2019.

[20] A. Mishkin, F. Kunstner, D. Nielsen, M. W. Schmidt, and M. E. Khan. Slang: Fast structured covariance approximations for Bayesian deep learning with natural gradient. In *NeurIPS*, pages 6248–6258, 2018.

[21] C. V. Nguyen, Y. Li, T. D. Bui, and R. E. Turner. Variational continual learning. In *International Conference on Learning Representations*, 2018.

[22] D. J. Poirier. Revising beliefs in nonidentified models. *Econometric Theory*, 14(4):483–509, 1998.

[23] A. Pourzanjani, R. M. Jiang, and L. Petzold. Improving the identifiability of neural networks for Bayesian inference. In *Bayesian Deep Learning NeurIPS workshop*, 2017.

[24] B. L. S. Prakasa Rao. *Identifiability in Stochastic Models*. Academic Press, Oxford, 1992.

[25] S. Sun, G. Zhang, J. Shi, and R. B. Grosse. Functional variational Bayesian neural networks. *ArXiv*, abs/1903.05779, 2019.

[26] H. J. Sussmann. Uniqueness of the weights for minimal feedforward nets with a given input-output map. *Neural Networks*, 5(4):589–593, 1992. ISSN 0893-6080. doi: https://doi.org/10.1016/S0893-6080(05)80037-1.

[27] J. Swiatkowski, K. Roth, B. S. Veeling, L. Tran, J. V. Dillon, J. Snoek, S. Mandt, T. Salimans, R. Jenatton, and S. Nowozin. The k-tied normal distribution: A compact parameterization of gaussian mean field posteriors in Bayesian neural networks. In *Proceedings of the 37th International Conference on Machine Learning*, ICML'20. JMLR.org, 2020.

[28] M. Tomczak, S. Swaroop, and R. Turner. Efficient low rank gaussian variational inference for neural networks. In H. Larochelle, M. Ranzato, R. Hadsell, M. Balcan, and H. Lin, editors, *Advances in Neural Information Processing Systems*, volume 33, pages 4610–4622. Curran Associates, Inc., 2020.

[29] B.-H. Tran, S. Rossi, D. Milios, and M. Filippone. All you need is a good functional prior for Bayesian deep learning. *Journal of Machine Learning Research*, 23(74):1–56, 2022.

[30] B. Trippe and R. Turner. Overpruning in variational Bayesian neural networks. *arXiv preprint arXiv:1801.06230*, 2018.

[31] M. J. Wainwright, M. I. Jordan, et al. Graphical models, exponential families, and variational inference. *Foundations and Trends® in Machine Learning*, 1(1–2):1–305, 2008.

[32] Y. Wang, D. Blei, and J. P. Cunningham. Posterior collapse and latent variable non-identifiability. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P. Liang, and J. W. Vaughan, editors, *Advances in Neural Information Processing Systems*, volume 34, pages 5443–5455. Curran Associates, Inc., 2021.

[33] Z. Wang, T. Ren, J. Zhu, and B. Zhang. Function space particle optimization for Bayesian neural networks. In *International Conference on Learning Representations*, 2019.

[34] F. Wenzel, K. Roth, B. Veeling, J. Swiatkowski, L. Tran, S. Mandt, J. Snoek, T. Salimans, R. Jenatton, and S. Nowozin. How good is the Bayes posterior in deep neural networks really? In H. D. III and A. Singh, editors, *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pages 10248–10259. PMLR, 13–18 Jul 2020.

[35] A. G. Wilson and P. Izmailov. Bayesian deep learning and a probabilistic perspective of generalization. *CoRR*, abs/2002.08791, 2020.