ANALYTICAL SOLUTIONS FOR A FAMILY OF SINGLE LAYER NEURAL NETWORK REGRESSION PROBLEMS

ABSTRACT

In this paper we show analytically that the optimal weights for a family of penalized single layer neural network regression problems wherein the response variable has all non-positive entries will always be $[0, 0, \ldots, 0, 0]^T$

PROBLEM SETUP

The loss function for a model which regresses a vector $\mathbf{y}$, on a data matrix $\mathbf{X}$, using a single layer neural network with RELU non-linearities is given by

$$f(\beta) = ||y - \max(0, \mathbf{X}\beta)||_2^2 + \lambda_1 ||\beta||_1 + \lambda_2 ||\beta||_2^2, \tag{1}$$

where $\lambda_1 \geq 0$ and $\lambda_2 \geq 0$. In this paper, we show analytically that when $\mathbf{y}$ has all non-positive entries (i.e., $\mathbf{y}[i] = -\alpha_i$ with $\alpha_i \geq 0$ for all i), and at least one of λ_1 or λ_2 is strictly greater than 0, then $\beta^* = [0, 0, \ldots, 0, 0]^T$ is the unique minimizer of (1)

PROOF

With the i^{th} row of $\mathbf{X}$ given by $\mathbf{X}_{i,:} = x_i^T$, and $\mathbf{y}[i] = -\alpha_i$ with $\alpha_i \geq 0$ as described above, define

$$h_i(\beta) = (\alpha_i + \max(0, x_i^T \beta))^2 \tag{2}$$

and

$$g(\beta) = \lambda_1 ||\beta||_1 + \lambda_2 ||\beta||_2^2. \tag{3}$$

With a minor re-arrangement of the terms, equation (1) can be rewritten as

$$f(\beta) = \sum_{i=1}^{i=N} h_i(\beta) + g(\beta) \tag{4}$$

Our proof proceeds in two parts. First, we show that $f(\beta)$ is a convex function of β, and therefore, every local optimal solution is also a global optimal solution. Next, we show that $\beta^* = [0, 0, \ldots, 0, 0]^T$ minimizes $f(\beta)$, and that for every $\beta' \neq \beta^*, f(\beta') > f(\beta^*)$. As a result, β^* is the unique minimizer of $f(\beta)$. We omit proofs for the well established facts that $g(\beta)$ is a convex function of β, and that $\beta^* = [0, 0, \ldots, 0, 0]^T$ is the unique minimizer of $g(\beta)$ whenever $\lambda_1 > 0$ and/or $\lambda_2 > 0$.

Proposition 1 $f(\beta)$ *is a convex function of* β

Proof Using example 3.5 (page 80) and section 3.2.1 (page 79) in Boyd et al. (2004), we have that $(\alpha_i + \max(0, x_i^T \beta))$ is a convex function of β. Furthermore, since $(\alpha_i + \max(0, x_i^T \beta)) \geq 0$ for all values of β, (2) implies $h_i(\beta)$ is a convex function of β for every i (bullet point 4 in example 3.13 (page 86) of Boyd et al. (2004)). Finally, since $g(\beta)$ is a convex function of β, (4) implies that $f(\beta)$ is a convex function of β. Therefore, any locally optimal point in $f(\beta)$ is also globally optimal ∎

Proposition 2 $\beta^* = [0, 0, \ldots, 0, 0]^T$ *is the unique global minimizer of* $f(\beta)$

Proof From (2), $h_i(\beta)$ attains its minimum value whenever $\max(0, x_i^T \beta) = 0$. Since β^* satisfies this condition, β^* minimizes $h_i(\beta)$ for every i. Furthermore, since β^* also minimizes $g(\beta)$, (4) implies $\beta^* = \arg\min_\beta f(\beta)$.

To prove the uniqueness of the solution, we note that when $\lambda_1 > 0$ and/or $\lambda_2 > 0$, β^* is the unique minimizer of $g(\beta)$. Therefore for any $\beta' \neq \beta^*$, we have $g(\beta) > g(\beta^*)$ and $h_i(\beta') \geq h_i(\beta^*)$ for every i. Combining the inequalities, we have $\sum_{i=1}^{i=N} h_i(\beta') + g(\beta') > \sum_{i=1}^{i=N} h_i(\beta^*) + g(\beta^*)$, i.e., $f(\beta') > f(\beta^*)$.

■

URM Statement

The authors acknowledge that at least one key author of this work meets the URM criteria of ICLR 2023 Tiny Papers Track.

References

Stephen Boyd, Stephen P Boyd, and Lieven Vandenberghe. *Convex optimization*. Cambridge university press, 2004.