FAST REGRESSION FOR STRUCTURED INPUTS

ABSTRACT

L_p Regression on Structured Inputs is an important problem in data analysis and machine learning where we find a vector $\mathbf{x} \in \mathbb{R}^d$ that minimizes $\|\mathbf{Ax} - \mathbf{b}\|_p$ for a *structured* matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ and response vector $\mathbf{b} \in \mathbb{R}^n$. Unfortunately, for many common classes of matrices, sampling-based algorithms for approximately solving L_p regression require runtime that is exponential in p, e.g., $d^{\mathcal{O}(p)}$, which is prohibitively expensive. We show that for a large class of structured inputs, such as combinations of low-rank matrices, sparse matrices, and Vandermonde matrices, L_p regression can be approximately solved using runtime that is polynomial in p. For example, we show that L_p regression on Vandermonde matrices can be approximately solved using time $\tilde{\mathcal{O}}(T(\mathbf{A}) \log n + (dp)^\omega \cdot \text{polylog}\, n)$, where $T(\mathbf{A})$ is the time to multiply $\mathbf{Ax}$ for an arbitrary vector $\mathbf{x} \in \mathbb{R}^d$, and ω is the exponent of matrix multiplication. The polynomial dependence on p also crucially allows our algorithms to extend naturally to sublinear time algorithms for L_∞ regression. Of independent interest, we develop a new algorithm for solving L_p regression for arbitrary matrices, which is significantly faster in practice for every $p \geq 4$.

1 INTRODUCTION

Given a matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ and a vector $\mathbf{b} \in \mathbb{R}^n$, the goal of linear regression is to find a vector $\mathbf{x} \in \mathbb{R}^d$ that best fits $\mathbf{A}$ to $\mathbf{b}$. However for $d < n$, the linear system is overconstrained and thus there may not exist a vector $\mathbf{x}$ such that $\mathbf{Ax} = \mathbf{b}$. Thus the L_p regression problem seeks to find a vector $\tilde{\mathbf{x}} \in \mathbb{R}^d$ such that

$$\|\mathbf{A}\tilde{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \varepsilon) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p,$$

for some approximation parameter $\varepsilon > 0$. Here for a vector $y \in \mathbb{R}^n$, $\|y\|_p^p = \sum_{i=1}^n |y_i|^p$. The L_p regression problem is a central question in statistical data analysis that has numerous applications in machine learning, data science, and applied mathematics (Friedman et al., 2001; Chatterjee & Hadi, 2006) and thus has received a considerable amount of attention. A direct way to solve the L_p regression problem is to use an iterative solver (Adil et al., 2019b;a) that makes $\text{poly}(n)$ calls to a linear system solver. However, the linear system solver uses time $\text{poly}(n, d)$, which can be prohibitively slow for $n \gg d$.

Therefore, a standard approach for L_p regression is to first sample a number of rows of $\mathbf{A}$ down to a smaller matrix $\mathbf{M}$ that generally has $\text{poly}\left(d, \frac{1}{\varepsilon}, \log n\right)$ rows by generating a sampling and rescaling matrix $\mathbf{S}$ such that $\mathbf{M} = \mathbf{SA}$ and $\|\mathbf{SAx} - \mathbf{Sb}\|_p$ is a "good" approximation to $\|\mathbf{Ax} - \mathbf{b}\|_p$ for all $\mathbf{x} \in \mathbb{R}^d$. Subsequently, an iterative solver can now be used on $\mathbf{M}$, which serves as a proxy for the original problem but has smaller input size, thus allowing for more efficient computation. Unfortunately, to handle worst-case inputs, $\mathbf{M}$ requires sampling $\mathcal{O}\left(d^{\max(1, p/2)}\right)$ rows, so that not only is the overall runtime exponential in p, but also the sampling is only helpful if $n \gg d^{p/2}$.

Fortunately, there are many applications in which the design matrix $\mathbf{A}$ is known to have additional structure that can be leveraged to design more efficient algorithms. For example, Vandermonde matrices are used in the polynomial regression problem, which has been studied for over 200 years (Gergonne, 1974) and recently motivated by applications to machine learning (Kalai et al., 2008), applied statistics (MacNeill, 1978), and computer graphics (Pratt, 1987) . The goal is to fit a continuous signal, which can be measured at time choices $t_1, \ldots, t_n$ to form a measurement vector $\mathbf{b} \in \mathbb{R}^n$, using the best degree d polynomial whose coefficients can be represented by the vector $\mathbf{x} \in \mathbb{R}^{d+1}$. Thus, entry $A_{i,j}$ of the feature matrix $\mathbf{A}$ is the value of the j-th polynomial basis at time t_i so that the

problem has been restated as the L_p regression problem with a Vandermonde matrix $\mathbf{A}$, where the entries of each row $\mathbf{a}_i$ form a sequence $1, t_i, (t_i)^2, \ldots, (t_i)^{d-1}$ for some parameter t_i. Regression problems with Vandermonde matrices also arise in the settings of Fourier fitting (Avron et al., 2019) and Toeplitz covariance estimation (Eldar et al., 2020). Shi & Woodruff (2019) show that for Vandermonde matrices $\mathbf{A}$, the time used to compute the smaller subsampled matrix $\mathbf{M}$ can be improved upon the naïve approach by using fast Vandermonde matrix-vector multiplication. However, the number of rows of $\mathbf{M}$ is still exponential in p so that unfortunately, the runtime of the iterative solver for L_p regression still uses time exponential in p, which suffices for small p but crucially does not obtain any improvements for the important problem of L_∞ regression.

1.1 OUR CONTRIBUTIONS

We first show that for L_p regression on Vandermonde matrices and its noisy perturbations, the size of the subsampled matrix $\mathbf{M}$ and therefore the runtime of the iterative solver only needs polynomial dependence in p.

Theorem 1.1. *Given $\varepsilon \in (0,1)$ and $p \geq 1$, a Vandermonde matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$, and $\mathbf{b} \in \mathbb{R}^d$, there exists an algorithm that creates a sampling matrix $\mathbf{S} \in \mathbb{R}^{m \times n}$ with $m = \mathcal{O}\left(\frac{p^2 d}{\varepsilon^3} \log^2 n\right)$ rows so that with high probability,*

$$(1-\varepsilon)\|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p \leq \|\mathbf{S}\mathbf{A}\mathbf{x} - \mathbf{S}\mathbf{b}\|_p \leq (1+\varepsilon)\|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p,$$

for all $\mathbf{x} \in \mathbb{R}^d$. The algorithm uses $\mathcal{O}\left(n \log^2 n\right) + p^3 d^{3/2} \operatorname{poly}\left(\frac{1}{\varepsilon}, \log n\right)$ total time and with high probability, returns a vector $\widehat{\mathbf{x}} \in \mathbb{R}^d$ such that

$$\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1+\varepsilon) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p.$$

The best known previous work has runtime either (1) $\mathcal{O}\left(n \log^2 n + d^q \cdot \operatorname{poly}(1/\varepsilon)\right)$ time (Shi & Woodruff, 2019), where $q = \omega \in [2,3)$ is the exponent for matrix multiplication for $p \in [1,4]$ and $q = p/2 + C$ for some fixed constant $C > 0$ for $p > 4$, which has an exponential dependence in p, or (2) using $\operatorname{poly}(n)$ solves on the $n \times d$ linear system Adil et al. (2019b;a). By contrast, our work only has a polynomial dependence on p.

In fact, we show that to obtain a $(1+\varepsilon)$-approximation to the fundamental problem of L_∞ regression, it suffices to consider L_p regression for $p = \mathcal{O}\left(\frac{\log n}{\varepsilon}\right)$ and since our results have polynomial dependence on p rather than exponential, we obtain *sublinear size* coresets for L_∞ regression:

Theorem 1.2. *Given $\varepsilon \in (0,1)$, a Vandermonde matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$, and $\mathbf{b} \in \mathbb{R}^d$, there exists an algorithm that uses $\mathcal{O}\left(n \log^2 n\right) + d^{3/2} \operatorname{poly}\left(\frac{1}{\varepsilon}, \log n\right)$ time that computes a sampling matrix $\mathbf{S} \in \mathbb{R}^{m \times n}$ with $m = d \operatorname{poly}\left(\frac{1}{\varepsilon}, \log n\right)$ such that with high probability,*

$$(1-\varepsilon)\|\mathbf{A}\mathbf{x} - \mathbf{b}\|_\infty \leq \|\mathbf{S}\mathbf{A}\mathbf{x} - \mathbf{S}\mathbf{b}\|_\infty \leq (1+\varepsilon)\|\mathbf{A}\mathbf{x} - \mathbf{b}\|_\infty$$

for all $\mathbf{x} \in \mathbb{R}^d$. In particular,

$$\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{S}\mathbf{A}\mathbf{x} - \mathbf{S}\mathbf{b}\|_\infty \leq (1+\varepsilon) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_\infty.$$

To the best of our knowledge, this is the first known dimensionality reduction for L_∞ regression with provable guarantees for any (nontrivial) input matrix.

We introduce the following definition to capture the "true dimension" of regression problems for structured input matrices.

Definition 1.3 (Rank of Regression Problem). *Given an integer $p \geq 1$ and a matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$, suppose there exists a matrix $\mathbf{M} \in \mathbb{R}^{n \times t}$ and a fixed function $f : \mathbb{R}^d \to \mathbb{R}^t$ so that for all $\mathbf{x} \in \mathbb{R}^d$,*

$$|\langle \mathbf{a}_i, \mathbf{x} \rangle|^p = |\langle \mathbf{m}_i, f(\mathbf{x}) \rangle|.$$

Then we call t the rank of the L_p regression problem.

Our result relies on the following key structural property that we prove about the p-fold tensor product of rows of the Vandermonde matrix.

Rows Sampled, L_p Regression	Rows Sampled, L_∞ Regression	Reference
$d^p \operatorname{poly}\left(\log n, \frac{1}{\varepsilon}\right)$	n	Avron et al. (2013)
$d^p \operatorname{poly}\left(\log n, \frac{1}{\varepsilon}\right)$	n	Shi & Woodruff (2019)
$dp^2 \operatorname{poly}\left(\log n, \frac{1}{\varepsilon}\right)$	$d \operatorname{poly}\left(\log n, \frac{1}{\varepsilon}\right)$	Our Results

Table 1: Sample complexity for regression on Vandermonde matrices

Lemma 1.4. *For integer $p \geq 1$, the rank of the L_p regression problem on a Vandermonde matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ is $\mathcal{O}\left(dp\right)$.*

Lemma 1.4 implies that the L_p loss function for a row of a Vandermonde matrix can potentially be expressed as a linear combination of $\mathcal{O}\left(dp\right)$ variables even though the entries of the measurement vector $\mathbf{b}$ can be arbitrary. By comparison, the L_p loss function for a row $\mathbf{a}_i$ of a general matrix $\mathbf{A}$ is $|\langle \mathbf{a}_i, \mathbf{x} \rangle - b_i|_p^p$ can only be expressed as a linear combination of $\mathcal{O}\left(pd^p\right)$ variables, corresponding to each of the d^k k-wise product of coordinates of $\mathbf{x}$, for each $k \in [p]$. As a corollary of Lemma 1.4, we obtain a small coreset for L_p regression (as well as L_∞ regression) on a Vandermonde matrix, which can thus be used as a preconditioner. We summarize these results in Table 1.

We generalize Lemma 1.4 to similar guarantees for L_p regression on a matrix $\mathbf{A}$ that is the sum of a low-rank matrix $\mathbf{L}$ and a sparse matrix $\mathbf{S}$ that has at most s non-zero entries per row.

Theorem 1.5. *Given $\varepsilon \in (0, 1)$ and $p \geq 1$, a matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ such that $\mathbf{A} = \mathbf{K} + \mathbf{S}$ for a rank k matrix $\mathbf{K}$ and a s-sparse matrix $\mathbf{S}$, and $\mathbf{b} \in \mathbb{R}^d$, there exists an algorithm that uses $\mathcal{O}\left(nk + ns\right) + \operatorname{poly}\left(d^s, k^p, s^p, \frac{1}{\varepsilon}\right)$ time, and with high probability, returns a vector $\widehat{\mathbf{x}} \in \mathbb{R}^d$ with*

$$\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \varepsilon) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p.$$

Similarly, we obtain efficient guarantees for L_p regression on a matrix $\mathbf{A}$ that is the sum of a Vandermonde matrix $\mathbf{V}$ and a sparse matrix $\mathbf{S}$ that has at most s non-zero entries per row.

Theorem 1.6. *Given $\varepsilon \in (0, 1)$ and $p \geq 1$, a matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ such that $\mathbf{A} = \mathbf{V} + \mathbf{S}$ for a Vandermonde matrix $\mathbf{V}$ and an s-sparse matrix $\mathbf{S}$, and $\mathbf{b} \in \mathbb{R}^d$, there exists an algorithm that uses $\mathcal{O}\left(T(\mathbf{A})\right) \log n + \operatorname{poly}\left(p^2 d, d^s, s^p, \log n, \frac{1}{\varepsilon}\right)$ time and with high probability, returns a vector $\widehat{\mathbf{x}} \in \mathbb{R}^d$ such that*

$$\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \varepsilon) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p.$$

Surprisingly, our methods even yield practical algorithms for general matrices *with absolutely no structure*. Although the rank of the L_p regression problem for general matrices is $\mathcal{O}\left(d^p\right)$, we still obtain the optimal sample complexity(see Li et al. (2021) for a lower bound) of roughly $\mathcal{O}\left(d^{p/2}\right)$. Furthermore, the advantage of our approach is that we now only need to perform L_q Lewis weight sampling for $q \in [1, 4]$. There are known efficient iterative methods for computing the L_q Lewis weights for $q \in [1, 4]$, e.g., see Section 2.1 or Section 3 in Cohen & Peng (2015). By contrast, previous methods relied on computing general L_p weights for $p > 4$, which required solving a large convex program, e.g., through semidefinite programming, see Section 4 in Cohen & Peng (2015).

Finally, we experimentally validate our theory on synthetic data. In particular, we consider matrices and response vectors that are motivated by existing lower bounds for subsampling and sketching methods. For Vandermonde regression, our experiments demonstrate that the number of rows needed is subexponential in p, reinforcing how structured matrices outperform the worst-case bound. For unstructured matrix L_p regression, we demonstrate that our L_q Lewis Weight subsampling scheme is effective and accurate. For instance, we can find a near-optimal solution vector by solving the regression problem on just 4% of the input matrix's rows.

1.2 OVERVIEW OF OUR TECHNIQUES

Our algorithmic contributions rely on two key observations, which we describe below. For the sake of presentation, we assume p is an integer in this overview, though we handle arbitrary real values of p in the subsequent algorithms and analyses.

Reduced rank of the L_p regression problem on structured inputs. The first main ingredient is the simple yet powerful observation that the rank of the L_p regression problem on structured inputs such as Vandermonde matrices $\mathbf{A} \in \mathbb{R}^{n \times d}$ does not need to be the $\mathcal{O}(d^p)$ rank of the L_p regression problem on general matrices. For a general matrix, $\langle \mathbf{a}_i, \mathbf{x} \rangle^p$ for an integer p can be rewritten as a linear combination $\sum \alpha_i y_i$ of $\mathcal{O}(d^p)$ terms, where each coefficient α_i is a p-wise product of entries of the row vector $\mathbf{a}_i \in \mathbb{R}^d$ and similarly each y_i is a p-wise product of entries of the vector $\mathbf{x} \in \mathbb{R}^d$. However, when $\mathbf{A}$ is a Vandermonde matrix, then the j-th entry of $\mathbf{a}_i$ is simply $A_{i,2}^{j-1}$. Thus each coefficient in $\langle \mathbf{a}_i, \mathbf{x} \rangle^p$ can be written as a linear combination of $1, A_{i,2}, (A_{i,2})^2, \ldots, (A_{i,2})^{p(d-1)}$ and so we can express $\langle \mathbf{a}_i, \mathbf{x} \rangle^p$ as a linear combination of $\mathcal{O}(dp)$ variables rather than $\mathcal{O}(d^p)$ variables.

We can similarly show that the rank of the L_p regression problem on rank k-matrix $\mathbf{A}$ is $\mathcal{O}(k^p)$ by writing each row $\mathbf{a}_i$ as a linear combination of basis vectors $\mathbf{v}_1, \ldots, \mathbf{v}_k$. Then, using the Hadamard Product-Kronecker Product mixed-product property, we can rewrite $\langle \mathbf{a}_i, \mathbf{x} \rangle^p$ as a linear combination of p-wise products of the variables $\langle \mathbf{v}_1, \mathbf{x} \rangle, \ldots, \langle \mathbf{v}_k, \mathbf{x} \rangle$, i.e., a linear combination of $\mathcal{O}(k^p)$ variables rather than $\mathcal{O}(d^p)$ variables. It follows that the rank of the L_p regression problem on a matrix $\mathbf{A}$ whose rows have at most s non-zero entries is $\mathcal{O}(d^s s^p)$ by noting that (1) there are $\binom{d}{s} = \mathcal{O}(d^s)$ sparsity patterns and that (2) for a fixed sparsity pattern, the rank of the induced matrix is at most s, which induces a linear combination of $\mathcal{O}(s^p)$ variables for the L_p regression problem. These decomposition techniques can also be generalized to show that the rank of the L_p regression problem is low on matrices $\mathbf{A}$ such that $\mathbf{A} = \mathbf{K} + \mathbf{V}$, $\mathbf{A} = \mathbf{K} + \mathbf{S}$ or $\mathbf{A} = \mathbf{V} + \mathbf{S}$, where $\mathbf{K}$ is a low-rank matrix, $\mathbf{V}$ is a Vandermonde matrix, and $\mathbf{S}$ is a sparse matrix.

However, we remark that although the observation that the rank of the L_p regression problem on structured inputs can be low, this itself does not yield an algorithm for L_p regression. This is because the loss function is $|\langle \mathbf{a}_i, \mathbf{x} \rangle - b_i|^p$ rather than $\langle \mathbf{a}_i, \mathbf{x} \rangle^p$ and there can be n possible different values of b_i across all $i \in [n]$.

Rounding and truncating the measurement vector: a novel algorithmic technique. Thus, the second main ingredient is manipulating the measurement vector $\mathbf{b} \in \mathbb{R}^n$ so that the loss function can utilize the low-rank property of the L_p regression problem on structured inputs. A natural approach would be to round the entries of $\mathbf{b}$, say to the nearest power of $(1 + \varepsilon)$. Unfortunately, such a rounding approach would roughly preserve $\|\mathbf{b}\|_p$ up to a multiplicative $(1 + \varepsilon)$ factor, but it would not preserve $\|\mathbf{Ax} - \mathbf{b}\|_p$. For example, suppose $\langle \mathbf{a}_i, \mathbf{x} \rangle = b_i = N$ for some arbitrarily large value N. Then $|\langle \mathbf{a}_i, \mathbf{x} \rangle - b_i|^p = 0$, but if b_i were rounded to $(1+\varepsilon)N$, we would have $|\langle \mathbf{a}_i, \mathbf{x} \rangle - b_i|^p = \varepsilon^p N^p$, which can be arbitrarily large.

The lesson from this counterexample is that when b_i is significantly larger than $\langle \mathbf{a}_i, \mathbf{x} \rangle - b_i$, any rounding technique can be arbitrarily bad because $\|\mathbf{b}\|_p$ can be significantly larger than $\text{OPT} := \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p$. On the other hand, if $\|\mathbf{b}\|_p$ is a constant factor approximation to OPT, then the previous counterexample cannot happen because either (1) b_i is large relative to $\|\mathbf{b}\|_p$ and $\langle \mathbf{a}_i, \mathbf{x} \rangle$ cannot be too close to b_i so that rounding b_i will not significantly affect the difference $\langle \mathbf{a}_i, \mathbf{x} \rangle - b_i$ or (2) b_i is small relative to $\|\mathbf{b}\|_p$ and so any rounding of b_i will not affect the contribution of the i-th row of $\mathbf{A}$ in the overall loss. Thus our task is reduced to manipulating the input so that $\|\mathbf{b}\|_p = \mathcal{O}(\text{OPT})$.

To that end, we note that if $\|\mathbf{A}\tilde{\mathbf{x}} - \mathbf{b}\|_p = \mathcal{O}(\text{OPT})$ for a vector $\tilde{\mathbf{x}} \in \mathbb{R}^d$, then by a triangle inequality argument, we have that the residual vector $\mathbf{b}' = \mathbf{b} - \mathbf{A}\tilde{\mathbf{x}} \in \mathbb{R}^d$ satisfies $\|\mathbf{b}'\|_p = \mathcal{O}(\text{OPT})$. Namely, we show that to find such a vector $\tilde{\mathbf{x}}$, it suffices by triangle inequality to find a subspace embedding for $\mathbf{A}$, i.e., a matrix $\mathbf{M} \in \mathbb{R}^{m \times d}$ with $m \ll n$ such that

$$(1 - C)\|\mathbf{Ax}\|_p \leq \|\mathbf{Mx}\|_p \leq (1 + C)\|\mathbf{Ax}\|_p$$

for some constant $C \in (0, 1)$. Typically, such a matrix $\mathbf{M}$ can be found by sampling the rows of $\mathbf{A}$ according to their L_p Lewis weights to generate a matrix with $m = \mathcal{O}\left(d^{\max(1,p/2)}\right)$ rows. However, using our observation that the rank of the L_p regression problem on structured inputs can be low, we can sample $\mathcal{O}(dp)$ rows of $\mathbf{A}$ with probabilities proportional to their L_q Lewis weights, where $q = \frac{p}{2^r}$ for an integer r such that $2^r \leq p < 2^{r+1}$. Crucially, we can find such a vector $\tilde{\mathbf{x}}$ *without reading the coordinates* of $\mathbf{b}$ since we only require to read the rows of $\mathbf{A}$ to perform Lewis weight sampling in this phase. Thus we can efficiently find a residual vector $\mathbf{b}'$ such that $\|\mathbf{b}'\|_p = \mathcal{O}(\text{OPT})$.

Partitioning the matrix into groups. We then round the entries of $\mathbf{b}'$ to obtain a vector $\mathbf{b}''$ with at most $\ell := \mathcal{O}\left(\frac{\log n}{\varepsilon}\right)$ unique values, truncating all entries that are less in magnitude than $\frac{1}{\text{poly}(n)}$ to zero instead. We now solve the L_p regression problem $\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}''\|_p$ by partitioning the rows of $\mathbf{A}$ into ℓ groups $G_1, \ldots, G_\ell$ based on their corresponding values of $\mathbf{b}''$. The main point is that all rows in each group G_k have the same entry t_k in $\mathbf{b}''$. Thus we can again observe that $|\langle \mathbf{a}_i, \mathbf{x} \rangle - t_k|^p$ can be written as a linear combination of $\mathcal{O}\left(p^2 d\right)$ variables and therefore use L_q Lewis weight sampling (for the same q) to reduce each group G_k down to $\mathcal{O}\left(\frac{p^2 d}{\varepsilon^2} \log d\right)$ rows.

Since there are $\ell = \mathcal{O}\left(\frac{\log n}{\varepsilon}\right)$ groups, then there are roughly $\mathcal{O}\left(p^2 d\right)$ total rows that have been sampled across all the groups. It follows from the decomposition of the L_p loss function across each group that the matrix $\mathbf{T}$ formed by these rows is a coreset for the L_p regression problem. Therefore, by solving the L_p regression problem on $\mathbf{T}$, which has significantly smaller dimension than the original input matrix $\mathbf{A}$, we obtain a vector $\widehat{\mathbf{x}}$ with the desired property that

$$\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \mathcal{O}(\varepsilon)) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p.$$

Practical L_p regression for arbitrary matrices. We now describe a practical procedure for L_p regression on general matrices that avoids the necessity for convex programming to approximate the L_p Lewis weights for $p > 4$. We first pick an integer r such that $\frac{p}{2^r} \in [2, 4)$. By the same structural argument as in Lemma 1.4, we can tensor product each row $\mathbf{a}_i$ of $\mathbf{A}$ with itself 2^r times, thus obtaining an extended matrix of size $n \times d'$, where $d' = d^{2^r}$, independent of the entries in $\mathbf{b}$. We then $L_{p/2^r}$ Lewis weight sample on the extended matrix, using the iterative method in Figure 1 since $\frac{p}{2^r} < 4$ rather than solving a convex program for L_p Lewis weight sampling for $p > 4$. Since $\frac{p}{2^r} \in [2, 4)$, it follows from Theorem 2.3 that Lewis weight sampling requires roughly $(d')^{p/2}$ rows. Thus we obtain a matrix with $\mathcal{O}\left((d^{2^r})^{p/2^{r+1}}\right) = \mathcal{O}\left(d^{p/2}\right)$ rows, from which we can compute a residual vector $\mathbf{b}'$ such that $\|\mathbf{b}'\|_p = \mathcal{O}(\text{OPT})$, where $\text{OPT} := \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p$.

We then round and truncate the entries of $\mathbf{b}'$ using the subroutine RoundTrunc to obtain a vector $\mathbf{b}''$, which allows us to partition the rows of $\mathbf{A}$ and the entries of $\mathbf{b}''$ into $\mathcal{O}\left(\frac{\log n}{\varepsilon}\right)$ groups. Due to the constant values of $\mathbf{b}''$ in each group, we can again $L_{p/2^r}$ Lewis weight sample on an extended matrix using an iterative method and finally solve the L_p regression problem on the subsequent rows that have been sampled. We give our algorithm in full in Algorithm 3.

2 L_p REGRESSION FOR VANDERMONDE MATRICES

2.1 PRELIMINARIES ON LEWIS WEIGHT SAMPLING

There are a number of known sampling distributions for dimensionality reduction for the L_p regression problem, such as the L_p leverage scores, e.g., (Cormode et al., 2018; Dasgupta et al., 2008) and the L_p sensitivities, e.g., (Clarkson et al., 2019; Braverman et al., 2020; Musco et al., 2021a); in this paper we focus on the L_p Lewis weights, e.g., (Cohen & Peng, 2015; Durfee et al., 2018; Chen & Derezinski, 2021; Parulekar et al., 2021).

Definition 2.1 (L_p Lewis Weights, Cohen & Peng (2015)). *Given a matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ and $p \geq 1$, the L_p Lewis weights $w_1(\mathbf{A}), \ldots, w_n(\mathbf{A})$ are the **unique** quantities that satisfy*

$$(w_i(\mathbf{A}))^{2/p} = \mathbf{a}_i^\top (\mathbf{A}^\top \mathbf{W}^{1-2/p} \mathbf{A})^{-1} \mathbf{a}_i$$

for all $i \in [n]$, where $\mathbf{W} \in \mathbb{R}^{n \times n}$ is the diagonal matrix with $W_{i,i} = w_i(\mathbf{A})$ for all $i \in [n]$.

Definition 2.2 (Lewis Weight Sampling). *For an input matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ and m samples, let the sampling matrix $\mathbf{S} \in \mathbb{R}^{m \times n}$ be generated by independently setting each row of $\mathbf{S}$ to be the i-th standard basis vector multiplied by $\left(\frac{d}{m \cdot w_i^p(\mathbf{A})}\right)^{1/p}$ with probability $\frac{w_i^p(\mathbf{A})}{d}$.*

Theorem 2.3 (L_p Subspace Embedding from Lewis Weight Sampling, (Cohen & Peng, 2015)). *For any $\mathbf{A} \in \mathbb{R}^{n \times d}$, let the matrix $\mathbf{S} \in \mathbb{R}^{m \times n}$ be generated from Lewis weight sampling with*

$m = \mathcal{O}\left(\frac{d^{\max(1,p/2)} \log(d/\delta) \log(1/\varepsilon)}{\varepsilon^\gamma}\right)$, *where* $\gamma = 2$ *for* $p \in [1, 2]$ *and* $\gamma = 5$ *for* $p > 2$. *Then with probability at least* $1 - \delta$, *we have that simultaneously for all* $\mathbf{x} \in \mathbb{R}^d$,

$$(1 - \varepsilon)\|\mathbf{A}\mathbf{x}\|_p \leq \|\mathbf{S}\mathbf{A}\mathbf{x}\|_p \leq (1 + \varepsilon)\|\mathbf{A}\mathbf{x}\|_p.$$

Since the Lewis weights are implicitly defined, it may not be clear how to compute them exactly. In fact, it suffices to compute constant factor approximations to the Lewis weights. Cohen & Peng (2015) show that for $p \leq 4$, there exists a simple iterative approach to compute a constant factor approximation to the L_p Lewis weights in input sparsity time, which we present in Figure 1.

$\mathbf{w} = \mathsf{LewisIterate}(\mathbf{A}, p, \beta, \mathbf{w})$

    (1) For $i = 1, \ldots, n$

        (a) Let τ_i be a constant factor approximation to $\mathbf{a}_i^\top (\mathbf{A}^\top \mathbf{W}^{1-2/p} \mathbf{A})^{-1} \mathbf{a}_i$

        (b) $w_i \leftarrow (\tau_i)^{p/2}$

        (c) Return $\mathbf{w}$

$\mathbf{w} = \mathsf{ApproxLewisWeights}(\mathbf{A}, p, \beta, T)$

    (1) Intialize $w_i = 1$ for all $i \in [n]$.

    (2) For $t = 1, \ldots, T$

        (a) $\mathbf{w} \leftarrow \mathsf{LewisIterate}(\mathbf{A}, p, \beta, \mathbf{w})$

    (3) Return $\mathbf{w}$

Fig. 1: Iterative algorithm for approximate L_p Lewis weights with $p < 4$.

At a high level, the correctness of Figure 1 follows from Banach's fixed point theorem and the fact that the subroutine LewisIterate is a contraction mapping, because $|2/p - 1| < 1$ for $p < 4$ (Cohen & Peng, 2015). However, for $p \geq 4$, Figure 1 no longer works. Instead, approximating L_p Lewis weights for $p > 4$ seems to require solving the convex program

$$\mathbf{Q} = \underset{\mathbf{M}}{\mathrm{argmax}}\, \det(\mathbf{M}), \qquad \text{subject to } \sum_i (\mathbf{a}_i^\top \mathbf{M} \mathbf{a})^{p/2} \leq d, \qquad \mathbf{M} \succeq 0,$$

and setting $w_i = (\mathbf{a}_i^\top \mathbf{Q} \mathbf{a})^{p/2}$. Unfortunately, this is often infeasible in practice, and we could not obtain empirical results using it. Therefore, a nice advantage of our algorithms, both for structured and unstructured matrices, is that we only use L_q Lewis weight sampling for $q \leq [1, 4)$, even if $p \geq 4$, whereas previous algorithms required using L_p Lewis weight sampling for $p > 4$.

2.2 ALGORITHM AND ANALYSIS

We first describe the general framework of our algorithm for efficient L_p regression, so that given $\mathbf{A} \in \mathbb{R}^{n \times d}$ and $\mathbf{b} \in \mathbb{R}^d$, the goal is to approximately compute $\mathrm{OPT} := \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p$. Recall that in order to apply our structural results, we first require a measurement vector $\mathbf{b}''$ with a small number of distinct entries. We obtain $\mathbf{b}''$ by first finding a constant factor approximation, i.e., $\tilde{\mathbf{x}} \in \mathbb{R}^d$ such that $\|\mathbf{A}\tilde{\mathbf{x}} - \mathbf{b}\|_p \leq \mathcal{O}(\mathrm{OPT})$. We can L_p Lewis weight sample the rows of $\mathbf{A}$ to do this, but the time to solve the subsequent polynomial regression problem would have exponential dependence on p, due to Theorem 2.3. Instead, we use our structural properties to implicitly create a matrix $\mathbf{M}$ from $\mathbf{A}$ with fewer than d^p columns and then $L_{p/2^r}$ Lewis weight sample the rows of $\mathbf{A}$, where r is the integer that satisfies $2^r \leq p < 2^{r+1}$, and then we solve the L_p regression problem on the sampled rows of $\mathbf{A}$ and entries of $\mathbf{b}$ to obtain $\tilde{\mathbf{x}}$. We set $\mathbf{b}'$ to be the residual vector $\mathbf{b} - \mathbf{A}\tilde{\mathbf{x}}$ so that $\|\mathbf{b}'\|_p = \mathcal{O}(\mathrm{OPT})$. We then use the procedure RoundTrunc, i.e., as in Algorithm 1, which sets the entries of $\mathbf{b}'$ that are the smallest in magnitude to zero, and rounds the remaining entries of $\mathbf{b}'$ to the nearest power of $(1 + \varepsilon)$.

Since $\|\mathbf{b}'\|_p = \mathcal{O}(\mathrm{OPT})$, it then follows by triangle inequality that it suffices to approximately solve the L_p regression problem on the vector $\mathbf{b}''$ instead. We partition the rows of $\mathbf{A}$ into groups

---

Algorithm 1 RoundTrunc: Round and truncate coordinates of input vector $\mathbf{b}$

Input: Vector $\mathbf{b} \in \mathbb{R}^n$, accuracy parameter $\varepsilon > 0$
Output: Vector $\mathbf{x} \in \mathbb{R}^n$ that is a rounded and truncated version of $\mathbf{b}$
1: $M \leftarrow \max_{i \in n} |b_i|$
2: **for** $i = 1$ to $i = n$ **do**
3:     $p_i \leftarrow \lfloor \log_{1+\varepsilon} |b_i| \rfloor$
4:     $x_i \leftarrow (1 + \varepsilon)^{p_i} \cdot \text{sign}(b_i)$
5:     **if** $|x_i| \leq \frac{M}{n^5}$ **then**
6:         $x_i \leftarrow 0$
7: **return** $\mathbf{x}$

$G_1, \ldots, G_\ell$ based on the values of $\mathbf{b}''$ and perform $L_{p/2^r}$ Lewis weight sampling again on an implicit matrix that we create from the rows of each group. Here we leverage our theory about the rank of the L_p regression problem for structured inputs. It then suffices to solve the L_p regression problem on the matrix formed by the sampled rows across all the groups.

Algorithm 2 Faster regression for Vandermonde matrices

Input: Vandermonde matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$, measurement vector $\mathbf{b}$, accuracy parameter $\varepsilon > 0$
Output: $\widehat{\mathbf{x}} \in \mathbb{R}$ with $\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \mathcal{O}(\varepsilon)) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p$
1: $r \leftarrow \lfloor \log p \rfloor$ $\triangleright 2^r \leq p < 2^{r+1}$.
2: Extend $\mathbf{A}$ to a Vandermonde matrix $\mathbf{M}$ with dimensions $n \times (2^r(d-1) + 1)$.
3: Use $L_{p/2^r}$-Lewis weight sampling on $\mathbf{M}$ to find a set S of $\mathcal{O}(d \log d)$ indices in $[n] = \{1, 2, \ldots, n\}$, and corresponding rescaling factors.
4: Let $\mathbf{A}'$ be the corresponding submatrix of $\mathbf{A}$ with indices in S, and scaled accordingly.
5: Compute $\tilde{\mathbf{x}} \leq 5 \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}'\mathbf{x} - \mathbf{b}\|_p$.
6: $\mathbf{b}' \leftarrow \mathbf{b} - \mathbf{A}\tilde{\mathbf{x}}$, $\mathbf{b}'' \leftarrow \text{RoundTrunc}(\mathbf{b}')$, $\ell \leftarrow O\left(\frac{\log n}{\varepsilon}\right)$
7: Partition the rows of $\mathbf{A}$ into groups $G_1, \ldots, G_\ell$, each containing all rows with the same value of $\mathbf{b}''$
8: Let $\mathbf{G}_k$ be the submatrix of G_k extended to $2^{2r}(d-1) + 1$ columns.
9: Let t_k be the coordinate of $\mathbf{b}''$ corresponding to G_k for each $k \in [\ell]$.
10: Use $L_{p/2^r}$-Lewis weight sampling on $[\mathbf{G}_k; t_k]$ to find a set S'_k of $O\left(\frac{1}{\varepsilon} d \log d\right)$ indices in $[n]$ and rescaling factors.
11: Let $\mathbf{T}_k$ be the corresponding rows with indices in S'_k and scaled accordingly.
12: $\mathbf{T} \leftarrow [\mathbf{T}_1; \ldots; \mathbf{T}_k]^\top$
13: Compute $\widehat{\mathbf{x}} \leq (1 + \varepsilon) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{T}\mathbf{x} - \mathbf{b}\|_p$.
14: **return** $\widehat{\mathbf{x}}$

We show in the supplementary material that Algorithm 2 satisfies the guarantees of Theorem 1.1. Moreover, because Theorem 1.1 has polynomial dependence on p rather than exponential dependence, we obtain the first known sublinear size coreset for the important problem of L_∞ regression. We use the following structural property.

Lemma 2.4. *Let* $\mathbf{x} \in \mathbb{R}^n$ *and* $p = \Omega\left(\frac{\log n}{\varepsilon}\right)$. *Then* $\|\mathbf{x}\|_\infty \leq \|\mathbf{x}\|_p \leq (1+\varepsilon)\|\mathbf{x}\|_\infty$.

Lemma 2.4 implies that to solve L_∞ regression, we can instead solve L_p regression for $p = \Omega\left(\frac{\log n}{\varepsilon}\right)$. Then Theorem 1.2 follows from the fact that even for $p = \Omega\left(\frac{\log n}{\varepsilon}\right)$, the matrix $\mathbf{T}$ in Algorithm 2 satisfies

$$(1 - \varepsilon)\|\mathbf{T}\mathbf{x} - \mathbf{b}''\|_p \leq \|\mathbf{A}\mathbf{x} - \mathbf{b}''\|_p \leq (1 + \varepsilon)\|\mathbf{T}\mathbf{x} - \mathbf{b}''\|_p$$

for all $\mathbf{x} \in \mathbb{R}^d$. Hence, we can solve the L_p regression problem on the smaller matrix $\mathbf{T}$ to solve the L_∞ regression on $\mathbf{A}$.

The results of Theorem 1.1 can be further extended to matrices with block Vandermonde structure.

Corollary 2.5. *Given* $\varepsilon \in (0, 1)$ *and* $p \geq 1$, $\mathbf{A} \in \mathbb{R}^{n \times dq}$, *and* $\mathbf{b} \in \mathbb{R}^d$, *suppose* $\mathbf{A} = [\mathbf{A}_1 | \ldots | \mathbf{A}_q]$ *for Vandermonde matrices* $\mathbf{A}_1, \ldots, \mathbf{A}_q \in \mathbb{R}^{n \times d}$. *Then there exists an algorithm that, with high*

probability, returns a vector $\widehat{\mathbf{x}} \in \mathbb{R}^d$ such that

$$\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \varepsilon) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p,$$

using $\mathcal{O}\left(T(\mathbf{A})(dp)^{q-1}\log n + \mathrm{poly}((dp)^q \log n, 1/\varepsilon)\right)$ time, where $T(\mathbf{A})$ is the runtime of multiplying the matrix $\mathbf{A}$ by an arbitrary vector. For $q = 2$, this can be further optimized to $\mathcal{O}\left(nd^{\omega_2/2-1} + \mathrm{poly}((dp)^2, \log n, 1/\varepsilon)\right)$ time, where $\mathcal{O}(n^{\omega_2})$ is the time to multiply an $n \times n$ matrix with an $n \times n^2$ matrix, so that $\omega_2 \in [3, 4]$.

We obtain similar algorithms for noisy low-rank matrices and noisy Vandermonde matrices, i.e., Theorem 1.5 and Theorem 1.6. We defer presentation and justification for these results to the supplementary material.

Algorithm 3 Faster L_p regression for general matrices

Input: Matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$, measurement vector $\mathbf{b}$, accuracy parameter $\varepsilon > 0$
Output: $\widehat{\mathbf{x}} \in \mathbb{R}$ with $\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \varepsilon) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p$, parameter $p \geq 4$
 1: $r \leftarrow \lfloor \log p \rfloor - 1$ ▷ $2^{r+1} \leq p < 2^{r+2}$.
 2: Extend $\mathbf{A}$ to a matrix $\mathbf{M}$ with dimension $n \times \mathcal{O}\left(d^{2^r}\right)$ so that each row in M_i is the 2^r-fold tensor product of $\mathbf{a}_i$ reshaped into a d^{2^r} length vector.
 3: Use $L_{p/2^r}$-Lewis weight sampling on $\mathbf{M}$ to find a set S of $\mathcal{O}\left(d^{p/2}\log d\right)$ indices in $[n]$ and rescaling factors.
 4: Let $\mathbf{A}'$ be the corresponding submatrix of $\mathbf{A}$ with indices in S and scaled accordingly.
 5: Compute $\tilde{\mathbf{x}} \leq 5\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}'\mathbf{x} - \mathbf{b}\|_p$.
 6: $\mathbf{b}' \leftarrow \mathbf{b} - \mathbf{A}\tilde{\mathbf{x}}$, $\mathbf{b}'' \leftarrow \mathsf{RoundTrunc}(\mathbf{b}')$, $\ell \leftarrow \mathcal{O}\left(\frac{\log n}{\varepsilon}\right)$
 7: Partition the rows of $\mathbf{A}$ into groups $G_1, \ldots, G_\ell$, each containing all rows with the same value of $\mathbf{b}''$
 8: Let $\mathbf{G}_k$ be the corresponding submatrix and t_k be the coordinate of $\mathbf{b}''$ corresponding to G_k for each $k \in [\ell]$.
 9: Use $L_{p/2^r}$-Lewis weight sampling on $[\mathbf{G}_k; t_k]$ to find a set S'_k of $\mathcal{O}\left(\frac{1}{\varepsilon^5}d^{p/2}\log d\right)$ indices in $[n]$ and rescaling factors.
10: Let $\mathbf{T}_k$ be the corresponding rows with indices in S'_k and scaled accordingly.
11: $\mathbf{T} \leftarrow [\mathbf{T}_1; \ldots; \mathbf{T}_k]^\top$
12: Compute $\widehat{\mathbf{x}} \leq (1 + \varepsilon)\min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{T}\mathbf{x} - \mathbf{b}\|_p$.
13: **return** $\widehat{\mathbf{x}}$

Finally, we describe in Algorithm 3 a practical approach for L_p regression for arbitrary matrices *without requiring any structural assumptions*.

Theorem 2.6. *Algorithm 3 outputs a vector $\widehat{\mathbf{x}}$ such that $\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1+\varepsilon)\min\|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p$. The runtime of Algorithm 3 is $T(n, d^{p/2}) + \mathrm{poly}\left(d^{p/2}, \log n, \frac{1}{\varepsilon}\right)$, where $T(n, d^p)$ is the time to multiply a matrix of size $n \times d^p$ by a vector of length d^p.*

Theorem 2.6 achieves the same optimal sample complexity as previous L_p Lewis weight algorithms of roughly $d^{p/2}$. However, the main advantage of Algorithm 3 is that it performs L_q Lewis weight sampling for some $q \in [2, 4)$, which is quite efficient because we can use an iterative method rather than solving a convex program.

3 EMPIRICAL VERIFICATION

We provide empirical evidence to validate our core statistical claim about Vandermonde regression: that the relative error ε achieved by $L_{p/2^r}$ Lewis Weight subsampling is polynomial in p, and not exponential in p. More precisely, we compute the gap between the error achieved from exact L_p Vandermonde regression and the error achieved from the subsampled regression:

$$\varepsilon_{empirical} = \frac{\|\mathbf{V}\widehat{\mathbf{x}} - \mathbf{b}\|_p - \|\mathbf{V}\mathbf{x}^* - \mathbf{b}\|_p}{\|\mathbf{V}\mathbf{x}^* - \mathbf{b}\|_p}$$

(a) Empirical Relative Error ($\varepsilon_{empirical}$) versus Lp parameter p. We ran the experiment 200 times and plot the median, 25^{th} quartile, and 75^{th} quartile for each value of p.

(b) Empirical Relative Error ($\varepsilon_{empirical}$) versus subsample complexity m. We ran the experiment 10 times and plot the median, 25^{th} quartile, and 75^{th} quartile for each value of p.

Our theory tells us that $(\varepsilon_{empirical})^3 \leq \tilde{\mathcal{O}}\left(\frac{dp^2}{m}\right)$, where m is the total number of subsampled rows. The prior work on unstructured matrices instead suggests $(\varepsilon_{empirical})^c \leq \tilde{\mathcal{O}}\left(\frac{d^{\mathcal{O}(p)}}{m}\right)$ (Shi & Woodruff, 2019). So, to visually distinguish these two settings, we look at the logarithm of both sides:

$$\log(\varepsilon_{empirical}) \leq \mathcal{O}\left(\ln(p) + \ln(\tfrac{d}{m})\right) \qquad \text{OR} \qquad \log(\varepsilon_{empirical}) \leq \mathcal{O}\left(p\ln(d) + \ln(\tfrac{1}{m})\right)$$

In particular, our work suggests a *logarithmic* dependence on p, while the prior work suggests a *linear* dependence on p.

To validate our theory, we plot $\log(\varepsilon_{empirical})$ versus p on synthetic data. Specifically, we generate $n = 25,000$ i.i.d. $N(0,1)$ time samples and let $\mathbf{b}$ take value 1 in its first 10 entries, and take value 0 otherwise. This sparse response vector mimics the worst-case response vector used in Lewis Weight subsampling lower bounds. Then, for $d = 20$ and $m = 1000$, we vary $p \in [2, 25]$ and compute both $\widehat{\mathbf{x}}$ and $\mathbf{x}^*$. Notably, in order to compute $\widehat{\mathbf{x}}$, we omit the rounding procedure in our code, since the rounding is designed for worst-case inputs. Instead, we simply compute $\widehat{\mathbf{x}}$ by solving $\min_{\mathbf{x}} \|\widehat{\mathbf{V}}\mathbf{x} - \widehat{\mathbf{b}}\|_p$ where $\widehat{\mathbf{V}}$ and $\widehat{\mathbf{b}}$ are computed by sampling and rescaling $\mathbf{V}$ and $\mathbf{b}$ with the $L_{p/2^r}$ Lewis Weights.

Figure 2a shows the result of the experiment, which was run in Julia 1.6.1, on Windows 10 with an Intel i7-7700K CPU and 16Gb RAM. The trendline clearly better fits a logarithmic model, as opposed to a linear model. This reinforces our analysis by showing that the dependence on p is notably sub-exponential, beating the known bounds for L_p subsampling on unstructured matrices.

We take a similar approach to verify that $L_{p/2^r}$ Lewis Weight sampling is correct for L_p regression on unstructured matrices. For this test we fix $n = 25,000$, $d = 10$, and $p = 6$, while varying $m \in [100, 3000]$. $\mathbf{A}$ is an i.i.d. $N(0,1)$ matrix. Due to numerical instability, we cannot apply Lewis Weight iteration to the expanded matrix $\mathbf{M}$ directly. Instead, we compute 10 steps of Lewis Weight Iteration on the random sketch $\mathbf{MG}$ where $\mathbf{G} \in \mathbb{R}^{(d^{\frac{p}{2}} \times 100)}$ has i.i.d. $N(0,1)$ entries. We then subsample $\mathbf{A}$ and $\mathbf{b}$ with respect to these Lewis Weights, and directly solve L_p regression on the subsampled matrices. Note that we again omit the rounding procedure on $\mathbf{b}$.

Figure 2b shows the result of this test, and we clearly see that the error shrinks quickly in m. For instance, to achieve 5% error, taking $m = 1000$, or just 4% of the rows of $\mathbf{A}$ suffices. Also note this approach is much more practical than the prior Lewis Weight approximation method for unstructured matrices when $p > 4$. That approach required solving a non-linearly constrained SDP $O(\log n)$ times (Cohen & Peng, 2015), while our method requires only a Gaussian sketch matrix and the standard Lewis Weight Iteration, which converges very quickly.

REFERENCES

Deeksha Adil, Rasmus Kyng, Richard Peng, and Sushant Sachdeva. Iterative refinement for l_p-norm regression. In *Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA*, pp. 1405–1424, 2019a. 1, 2, 15

Deeksha Adil, Richard Peng, and Sushant Sachdeva. Fast, provably convergent IRLS algorithm for p-norm linear regression. In *Advances in Neural Information Processing Systems 32, NeurIPS*, pp. 14166–14177, 2019b. 1, 2, 15

Haim Avron, Vikas Sindhwani, and David P. Woodruff. Sketching structured matrices for faster nonlinear regression. In *27th Annual Conference on Neural Information Processing Systems. Proceedings*, pp. 2994–3002, 2013. 3

Haim Avron, Michael Kapralov, Cameron Musco, Christopher Musco, Ameya Velingker, and Amir Zandieh. A universal sampling method for reconstructing signals with simple fourier transforms. In *Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing, STOC*, pp. 1051–1063, 2019. 2

Vladimir Braverman, Petros Drineas, Cameron Musco, Christopher Musco, Jalaj Upadhyay, David P. Woodruff, and Samson Zhou. Near optimal linear algebra in the online and sliding window models. In *61st IEEE Annual Symposium on Foundations of Computer Science, FOCS*, pp. 517–528, 2020. 5, 22

Samprit Chatterjee and Ali S Hadi. *Regression analysis by example*, volume 607. John Wiley and Sons, 2006. 1

Xue Chen and Michal Derezinski. Query complexity of least absolute deviation regression via robust uniform convergence. In *Conference on Learning Theory, COLT*, pp. 1144–1179, 2021. 5, 22

Rachit Chhaya, Jayesh Choudhari, Anirban Dasgupta, and Supratim Shit. Streaming coresets for symmetric tensor factorization. In *Proceedings of the 37th International Conference on Machine Learning, ICML*, pp. 1855–1865, 2020. 22

Kenneth L. Clarkson and David P. Woodruff. Low rank approximation and regression in input sparsity time. In *Symposium on Theory of Computing Conference, STOC*, pp. 81–90, 2013. 22

Kenneth L. Clarkson, Petros Drineas, Malik Magdon-Ismail, Michael W. Mahoney, Xiangrui Meng, and David P. Woodruff. The fast cauchy transform and faster robust linear regression. *SIAM J. Comput.*, 45(3):763–810, 2016. 22

Kenneth L. Clarkson, Ruosong Wang, and David P. Woodruff. Dimensionality reduction for tukey regression. In *Proceedings of the 36th International Conference on Machine Learning, ICML*, pp. 1262–1271, 2019. 5, 22

Michael B. Cohen and Richard Peng. l_p row sampling by lewis weights. In *Proceedings of the Forty-Seventh Annual ACM on Symposium on Theory of Computing, STOC*, pp. 183–192, 2015. 3, 5, 6, 9, 15, 22

Graham Cormode, Charlie Dickens, and David P. Woodruff. Leveraging well-conditioned bases: Streaming and distributed summaries in minkowski p-norms. In *Proceedings of the 35th International Conference on Machine Learning, ICML*, pp. 1048–1056, 2018. 5, 22

Anirban Dasgupta, Petros Drineas, Boulos Harb, Ravi Kumar, and Michael W. Mahoney. Sampling algorithms and coresets for l_p regression. In *Proceedings of the Nineteenth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA*, pp. 932–941, 2008. 5, 22

David Durfee, Kevin A. Lai, and Saurabh Sawlani. l_1 regression using lewis weights preconditioning and stochastic gradient descent. In *Conference On Learning Theory, COLT*, pp. 1626–1656, 2018. 5, 22

Yonina C. Eldar, Jerry Li, Cameron Musco, and Christopher Musco. Sample efficient toeplitz covariance estimation. In *Proceedings of the 2020 ACM-SIAM Symposium on Discrete Algorithms, SODA*, pp. 378–397, 2020. 2

Jerome Friedman, Trevor Hastie, Robert Tibshirani, et al. *The elements of statistical learning*. Springer series in statistics New York, 2001. 1

JD Gergonne. The application of the method of least squares to the interpolation of sequences. *Historia Mathematica*, 1(4):439–447, 1974. 1

I Gohberg and V Olshevsky. Complexity of multiplication with vectors for structured matrices. *Linear Algebra and Its Applications*, 202:163–192, 1994. 15

Jerrad Hampton and Alireza Doostan. Coherence motivated sampling and convergence analysis of least squares polynomial chaos regression. *Computer Methods in Applied Mechanics and Engineering*, 290:73–97, 2015. 20

Adam Tauman Kalai, Adam R Klivans, Yishay Mansour, and Rocco A Servedio. Agnostically learning halfspaces. *SIAM Journal on Computing*, 37(6):1777–1805, 2008. 1, 20

Daniel Kane, Sushrut Karmalkar, and Eric Price. Robust polynomial regression up to the information theoretic limit. In *58th IEEE Annual Symposium on Foundations of Computer Science, FOCS*, pp. 391–402, 2017. 20, 21

Kiran S. Kedlaya and Christopher Umans. Fast polynomial factorization and modular composition. *SIAM J. Comput.*, 40(6):1767–1802, 2011. 16

Yi Li, Ruosong Wang, and David P. Woodruff. Tight bounds for the subspace sketch problem with applications. *SIAM J. Comput.*, 50(4):1287–1335, 2021. 3, 22

Ian B MacNeill. Properties of sequences of partial sums of polynomial regression residuals with applications to tests for change of regression at unknown times. *The Annals of Statistics*, pp. 422–433, 1978. 1

Xiangrui Meng and Michael W. Mahoney. Low-distortion subspace embeddings in input-sparsity time and applications to robust linear regression. In *Symposium on Theory of Computing Conference, STOC*, pp. 91–100. ACM, 2013. 22

Raphael Meyer, Cameron Musco, Christopher Musco, David Woodruff, and Samson Zhou. Cheybshev sampling is universal for l_p polynomial regression, 2021. 20, 21, 22

Cameron Musco, Christopher Musco, and David P. Woodruff. Active sampling for linear regression beyond the l_2 norm, 2021a. (manuscript). 5, 22

Cameron Musco, Christopher Musco, David P. Woodruff, and Taisuke Yasuda. Active sampling for linear regression beyond the l_2 norm. *CoRR*, abs/2111.04888, 2021b. 20

Jelani Nelson and Huy L. Nguyen. OSNAP: faster numerical linear algebra algorithms via sparser subspace embeddings. In *54th Annual IEEE Symposium on Foundations of Computer Science, FOCS*, pp. 117–126, 2013. 22

Michael Nüsken and Martin Ziegler. Fast multipoint evaluation of bivariate polynomials. In *Algorithms - ESA, 12th Annual European Symposium, Proceedings*, pp. 544–555, 2004. 16

Aditya Parulekar, Advait Parulekar, and Eric Price. l_1 regression with lewis weights subsampling. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques, APPROX/RANDOM*, pp. 49:1–49:21, 2021. 5, 22

Vaughan Pratt. Direct least-squares fitting of algebraic surfaces. *ACM SIGGRAPH computer graphics*, 21(4):145–152, 1987. 1, 20

Christopher De Sa, Albert Gu, Rohan Puttagunta, Christopher Ré, and Atri Rudra. A two-pronged progress in structured dense matrix vector multiplication. In *Proceedings of the Twenty-Ninth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA*, pp. 1060–1079, 2018. 16

Tamás Sarlós. Improved approximation algorithms for large matrices via random projections. In *47th Annual IEEE Symposium on Foundations of Computer Science (FOCS), Proceedings*, pp. 143–152. IEEE Computer Society, 2006. 22

Xiaofei Shi and David P. Woodruff. Sublinear time numerical linear algebra for structured matrices. In *The Thirty-Third AAAI Conference on Artificial Intelligence*, pp. 4918–4925, 2019. 2, 3, 9, 13

Christian Sohler and David P. Woodruff. Subspace embeddings for the l_1-norm with applications. In *Proceedings of the 43rd ACM Symposium on Theory of Computing, STOC*, pp. 755–764, 2011. 22

Ruosong Wang and David P. Woodruff. Tight bounds for l_p oblivious subspace embeddings. In *Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA*, pp. 1825–1843, 2019. 22

David P. Woodruff and Qin Zhang. Subspace embeddings and l_p-regression using exponential random variables. In *COLT 2013 - The 26th Annual Conference on Learning Theory*, pp. 546–567, 2013. 22

BROADER IMPACT

Our work is focused on both theoretical and practical methods for improving the runtime efficiency of algorithms for L_p regression on structured input. As such, our work could have significant broader impacts by allowing users to more accurately or efficiently solve regression problems on large datasets. Although it may be possible that applications of our work to real data could potentially result in ethical considerations, we believe this is an indirect and unintended misuse of our work and we do not condone any such behavior. Our empirical evaluations use publicly available datasets to evaluate the performance of our algorithms. As a result, we believe no ethical considerations should be raised from our experiments.

A   MISSING PROOFS FROM SECTION 2

In this section, we give the missing proofs from Section 2.

A.1 L_p REGESSION ON VANDERMONDE MATRICES

We first prove a simple statement that shows a "good" solution to the L_p regression problem on a subspace embedding $\mathbf{SA}$ is also a "good" solution to the original input matrix $\mathbf{A}$.

Lemma A.1. *Let $\mathbf{S}$ be a sampling and rescaling matrix such that*

$$\frac{11}{12}\|\mathbf{SAx}\|_p \leq \|\mathbf{Ax}\|_p \leq \frac{13}{12}\|\mathbf{SAx}\|_p, \qquad \mathbb{E}\left[\|\mathbf{SAx}\|_p^p\right] = \|\mathbf{Ax}\|_p^p$$

for all $\mathbf{x} \in \mathbb{R}^d$. Let $OPT = \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{Ax} - \mathbf{b}\|_p$ and let $\mathbf{x} \in \mathbb{R}^d$ be any vector for which $\|\mathbf{SAx} - \mathbf{Sb}\|_p \leq 5OPT$. Then with probability at least 0.79,

$$\|\mathbf{Ax} - \mathbf{b}\|_p \leq 12OPT.$$

Proof. Let $\mathbf{x}^*$ be a minimizer of $\|\mathbf{Ax} - \mathbf{b}\|_p$ so that $OPT = \|\mathbf{Ax}^* - \mathbf{b}\|_p$. We will prove the claim by contrapositive, so we first suppose that $\|\mathbf{Ax} - \mathbf{b}\|_p \leq 12OPT$. Then by the triangle inequality,

$$\|\mathbf{SAx} - \mathbf{Sb}\|_p \geq \|\mathbf{SA}(\mathbf{x} - \mathbf{x}^*)\|_p - \|\mathbf{SA}^* - \mathbf{Sb}\|_p.$$

Since $\frac{11}{12}\|\mathbf{SAx}\|_p \leq \|\mathbf{Ax}\|_p \leq \frac{13}{12}\|\mathbf{SAx}\|_p$ for all $\mathbf{x} \in \mathbb{R}^d$, then

$$\|\mathbf{SAx} - \mathbf{Sb}\|_p \geq \frac{11}{12}\|\mathbf{A}(\mathbf{x} - \mathbf{x}^*)\|_p - \|\mathbf{SA}^* - \mathbf{Sb}\|_p.$$

By the triangle inequality,

$$\|\mathbf{SAx} - \mathbf{Sb}\|_p \geq \frac{11}{12}\left(\|\mathbf{Ax} - \mathbf{b}\|_p - \|\mathbf{Ax}^* - \mathbf{b}\|_p\right) - \|\mathbf{SA}^* - \mathbf{Sb}\|_p.$$

Note that by Jensen's inequality, $\mathbb{E}[\|\mathbf{SAx}^* - \mathbf{Sb}\|_p] \leq OPT$, so that by Markov's inequality,

$$\mathbf{Pr}\left[\|\mathbf{SAx}^* - \mathbf{Sb}\|_p \geq 5OPT\right] \leq \frac{1}{5}.$$

Thus with probability at least 0.8,

$$\|\mathbf{S}\mathbf{A}\mathbf{x} - \mathbf{S}\mathbf{b}\|_p \geq \frac{11}{12}\left(\|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p - \|\mathbf{A}\mathbf{x}^* - \mathbf{b}\|_p\right) - 5\text{OPT}.$$

Thus if $\|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p \leq 12\text{OPT}$, then

$$\|\mathbf{S}\mathbf{A}\mathbf{x} - \mathbf{S}\mathbf{b}\|_p \geq \frac{11}{12}\left(12\text{OPT} - \text{OPT}\right) - 5\text{OPT} > 5\text{OPT},$$

as desired. $\qquad\square$

Before justifying the correctness of Algorithm 2, we first recall the following algorithm for efficient L_p subspace embeddings.

Theorem A.2. *(Shi & Woodruff, 2019) Given $\varepsilon \in (0,1)$ and $p \in [1,2]$, a Vandermonde matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$, and $\mathbf{b} \in \mathbb{R}^d$, let $T(\mathbf{A})$ be the time it takes to perform matrix-vector multiplication, i.e., compute $\mathbf{A}\mathbf{v}$ for an arbitrary $\mathbf{v} \in \mathbb{R}^d$. There exists an algorithm that uses $\mathcal{O}\left(T(\mathbf{A})\log n + d^q \cdot \text{poly}(1/\varepsilon)\right)$ time, where $q = \omega$ for $p \in [1,4)$ and $q = p/2 + C$ for some fixed constant $C > 0$ for $p > 4$, and with high probability, returns $\tilde{\mathbf{x}} \in \mathbb{R}^d$ such that*

$$\|\mathbf{A}\tilde{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \varepsilon)\min_{\mathbf{x} \in \mathbb{R}^d}\|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p.$$

We now justify the correctness of Algorithm 2.

Lemma A.3. *Given $\varepsilon \in (0,1)$ and $p \geq 1$, a Vandermonde matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$, and $\mathbf{b} \in \mathbb{R}^d$, then with high probability, Algorithm 2 returns a vector $\widehat{\mathbf{x}} \in \mathbb{R}^d$ such that*

$$\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \mathcal{O}(\varepsilon))\min_{\mathbf{x} \in \mathbb{R}^d}\|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p.$$

Proof. Consider Algorithm 2 and let r be an integer so that $2^r \leq p < 2^{r+1}$. Note that

$$\sum_{i \in [n]}|\langle \mathbf{a}_i, \mathbf{x}\rangle|^p = \sum_{i \in [n]}|(\langle \mathbf{a}_i, \mathbf{x}\rangle)^{2^r}|^{p/2^r}.$$

Since $\mathbf{A}$ is Vandermonde, then

$$(\langle \mathbf{a}_i, \mathbf{x}\rangle)^{2^r} = \left(\sum_{j=1}^d a_{i,j}x_j\right)^{2^r} = \left(\sum_{j=1}^d a_{i,1}^j x_j\right)^{2^r} = \sum_{j=1}^{2^r(d-1)+1} a_{i,2}^{j-1}y_j,$$

where each y_j is a fixed function of the coordinates of $\mathbf{x}$. Notably, the fixed function *is the same across all $i \in [n]$*. Hence, the L_{2^r} subspace embedding problem on an input Vandermonde matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ can be reshaped as a constrained L_1 subspace embedding problem on an input Vandermonde matrix of size $n \times (2^r(d-1)+1)$. Thus, for L_p regression with $p \in [2^r, 2^{r+1})$, we have

$$\sum_{i \in [n]}|\langle \mathbf{a}_i, \mathbf{x}\rangle|^p = \sum_{i \in [n]}\left|\sum_{j=1}^{2^r(d-1)+1} a_{i,2}^{j-1}y_j\right|^{p/2^r},$$

which is a constrained $L_{p/2^r}$ regression problem on an input Vandermonde matrix $\mathbf{M}$ of size $n \times (2^r(d-1)+1)$.

By Theorem A.2, we can use $L_{p/2^r}$ Lewis weight sampling to find a matrix $\mathbf{M}'$ such that

$$\frac{1}{5}\|\mathbf{M}\mathbf{y}\|_{p/2^r}^{p/2^r} \leq \|\mathbf{M}'\mathbf{y}\|_{p/2^r}^{p/2^r} \leq 5\|\mathbf{M}\mathbf{y}\|_{p/2^r}^{p/2^r}$$

for all $\mathbf{y} \in \mathbb{R}^{2^r(d-1)+1}$ with high probability. Note that by the above argument, if we take the matrix $\mathbf{A}'$ corresponding to the scaled rows of $\mathbf{A}$ that are sampled by $\mathbf{M}'$, then we also have

$$\frac{11}{12}\|\mathbf{A}\mathbf{x}\|_p^p \leq \|\mathbf{A}'\mathbf{x}\|_p^p \leq \frac{13}{12}\|\mathbf{A}\mathbf{x}\|_p^p$$

and thus

$$\frac{11}{12}\|\mathbf{A}\mathbf{x}\|_p \leq \|\mathbf{A}'\mathbf{x}\|_p \leq \frac{13}{12}\|\mathbf{A}\mathbf{x}\|_p$$

for all $\mathbf{x} \in \mathbb{R}^{d+1}$ with high probability. Thus by Lemma A.1, we can find a vector $\tilde{\mathbf{x}}$ such that

$$\|\mathbf{A}\tilde{\mathbf{x}} - \mathbf{b}\|_p \leq 12 \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p.$$

We use the subroutine RoundTrunc to create the vector $\mathbf{b}''$ which is the vector with all entries of $\mathbf{b}'$ rounded to the nearest power of $(1 + \varepsilon)$, starting at the maximum entry of $\mathbf{b}'$ in absolute value, and stopping after we are $\frac{1}{\text{poly}(n)}$ times that, and replacing all remaining entries with 0. By the triangle inequality, we have

$$\|\mathbf{A}\mathbf{x} - \mathbf{b}''\|_p \leq \|\mathbf{A}\mathbf{x} - \mathbf{b}'\|_p + \|\mathbf{b}' - \mathbf{b}''\|_p \leq \|\mathbf{A}\mathbf{x} - \mathbf{b}'\|_p + 12\varepsilon\text{OPT} \leq (1 + 12\varepsilon)\|\mathbf{A}\mathbf{x} - \mathbf{b}''\|_p,$$

for any $\mathbf{x} \in \mathbb{R}^{d+1}$.

Note that $\mathbf{b}''$ has discretized the values of $\mathbf{b}'$ into $\ell = \mathcal{O}\left(\frac{\log n}{\varepsilon}\right)$ possible values. We partition the rows of $\mathbf{A}$ into ℓ groups $G_1, \ldots, G_\ell$, based on the corresponding values of $\mathbf{b}''$. Suppose that for a group G_k, the corresponding values of $\mathbf{b}''$ are all t_k. Then we have

$$\|\mathbf{A}\mathbf{x} - \mathbf{b}''\|_p^p = \sum_k \sum_{i \in G_k} |\langle \mathbf{a}_i, \mathbf{x}\rangle - t_k|^p.$$

Since $\mathbf{A}$ is Vandermonde, for each $i \in G_k$,

$$(\langle \mathbf{a}_i, \mathbf{x}\rangle - t_k)^{2^r} = \left(-t_k + \sum_{j=1}^d a_{i,j}x_j\right)^{2^r} = \left(-t_k + \sum_{j=1}^d a_{i,2}^{j-1}x_j\right)^{2^r} = \sum_{j=1}^{2^{2r}(d-1)+1} a_{i,2}^{j-1}t_{k,j}y_j,$$

for some fixed values $t_{k,1}, t_{k,2}, \ldots$ that can be computed from t_k, where again each y_j can be a different function of the coordinates of $\mathbf{x}$. In particular, there can be 2^r choices for the exponent of t_k. For a fixed choice of the exponent of t_k, the exponent of $a_{i,2}$ can range from 0 to $2^r(d-1)$.

Hence, the L_{2^r} regression problem on a submatrix of an input Vandermonde matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ *with the same fixed measurement values*, i.e., the corresponding coordinates of $\mathbf{b}''$ are all the same, can be reshaped to a constrained L_1 regression problem on an input Vandermonde matrix with $2^{2r}(d-1)+1$ columns times a $(2^{2r}(d-1)+1) \times (2^{2r}(d-1)+1)$ diagonal matrix.

Hence, by invoking Theorem A.2 to sample rows of $\mathbf{M}$ corresponding to their $L_{p/2^r}$ Lewis weights in the submatrix induced by the rows of G_k, we obtain a matrix $\mathbf{R}_k$ such that with high probability,

$$(1-\varepsilon)\|\mathbf{R}_k\mathbf{y} - t_k\|_{p/2^r}^{p/2^r} \leq \sum_{i \in G_k} \left(\sum_{j=1}^{2^r(d-1)+1} a_{i,2}^{j-1}t_{k,j}y_j\right)^{p/2^r} \leq (1+\varepsilon)\|\mathbf{R}_k\mathbf{y} - t_k\|_{p/2^r}^{p/2^r}.$$

Note that by the above argument, if we take the matrix $\mathbf{T}_k$ corresponding to the scaled rows of $\mathbf{A}$ that are sampled by $\mathbf{R}_k$, then we also have

$$(1-\varepsilon)\|\mathbf{T}_k\mathbf{x} - t_k\|_p^p \leq \sum_{i \in G_k} |\langle \mathbf{a}_i, \mathbf{x}\rangle - t_k|^p \leq (1+\varepsilon)\|\mathbf{T}_k\mathbf{x} - t_k\|_p^p,$$

with high probability. Summing over all k, we have

$$(1-\varepsilon)\sum_k \|\mathbf{T}_k\mathbf{x} - t_k\|_p^p \leq \sum_k \sum_{i \in G_k} |\langle \mathbf{a}_i, \mathbf{x}\rangle - t_k|^p \leq (1+\varepsilon)\sum_k \|\mathbf{T}_k\mathbf{x} - t_k\|_p^p,$$

with high probability.

Let $\mathbf{T} = \mathbf{T}_1 \circ \ldots \circ \mathbf{T}_\ell$, so that $\sum_k \|\mathbf{T}_k\mathbf{x} - t_k\|_p^p = \|\mathbf{T}\mathbf{x} - \mathbf{b}''\|_p^p$. Observe that $\|\mathbf{A}\mathbf{x} - \mathbf{b}''\|_p^p = \sum_k \sum_{i \in G_k} |\langle \mathbf{a}_i, \mathbf{x}\rangle - t_k|^p$. Hence,

$$(1-\varepsilon)\|\mathbf{T}\mathbf{x} - \mathbf{b}''\|_p^p \leq \|\mathbf{A}\mathbf{x} - \mathbf{b}''\|_p^p \leq (1+\varepsilon)\|\mathbf{T}\mathbf{x} - \mathbf{b}''\|_p^p$$

and thus

$$(1-\varepsilon)\|\mathbf{T}\mathbf{x} - \mathbf{b}''\|_p \leq \|\mathbf{A}\mathbf{x} - \mathbf{b}''\|_p \leq (1+\varepsilon)\|\mathbf{T}\mathbf{x} - \mathbf{b}''\|_p.$$

Thus we can compute a vector $\widehat{\mathbf{x}} \in \mathbb{R}^d$ such that

$$\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \mathcal{O}(\varepsilon)) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p.$$

$\square$

Before analyzing the time complexity of Algorithm 2, we first recall the following algorithms for Vandermonde matrix-vector multiplication and approximate L_p regression.

Theorem A.4 (Vandermonde Matrix-Vector Multiplication Runtime, e.g., Table 1 in Gohberg & Olshevsky (1994)). *The runtime of computing $\mathbf{A}\mathbf{x}$ for a Vandermonde matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ and a vector $\mathbf{x} \in \mathbb{R}^d$ for $d \leq n$ is $\mathcal{O}\left(n \log^2 n\right)$.*

Theorem A.5 (Approximate L_p Regression Runtime). *(Adil et al., 2019b) Given $\mathbf{A} \in \mathbb{R}^{n \times d}$ and $p \geq [2, \infty)$, there exists an algorithm that makes $\mathcal{O}\left(\sqrt{n} \log \frac{n}{\varepsilon}\right)$ calls to a linear system solver and computes a vector $\tilde{\mathbf{x}}$ such that*

$$\|\mathbf{A}\tilde{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \varepsilon) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p.$$

We now analyze the runtime of Algorithm 2.

Lemma A.6. *Algorithm 2 runs in $\mathcal{O}\left(n \log^2 n + \frac{p^3 d^{3/2}}{\varepsilon^{7/2}} \log^4 n \log \frac{n}{\varepsilon}\right)$ time.*

Proof. Observe that Algorithm 2 has three main bottlenecks for runtime. Since we only need to Lewis weight sample from the extended matrices, we do not need to explicitly form them, which would otherwise require $\Omega(ndp^2)$ time just to list to entries. Hence the first bottleneck is performing the Lewis weight sampling procedure on the extended matrices. The second bottleneck is solving the L_p regression problem on the final subsampled matrix. The only remaining procedure is rounding and truncating the coordinates of $\mathbf{b} \in \mathbb{R}^n$ to form a vector $\mathbf{b}'' \in \mathbb{R}^n$ using the procedure RoundTrunc and then forming the groups $G_1, \ldots, G_\ell$, which clearly takes $\mathcal{O}(n)$ arithmetic operations combined. We thus analyze each of the three main runtime bottlenecks.

First observe that the extended matrix $\mathbf{M}$ is a Vandermonde matrix with $\mathcal{O}\left(dp^2\right)$ columns. Cohen & Peng (2015) show that $\mathcal{O}(\log n)$ matrix-vector multiplication operations can be done to compute approximate Lewis weights for the purposes of L_p Lewis weight sampling. By Theorem A.4, each matrix-vector multiplication uses time $\mathcal{O}\left(n \log^2 n\right)$. Hence computing the extended matrix $\mathbf{M}$ uses $\mathcal{O}\left(n \log^2 n\right)$ time. Similarly, the extended matrix for each group G_k is the product of a Vandermonde matrix with $\mathcal{O}\left(dp^2\right)$ columns and a diagonal matrix. Thus by Theorem A.4, the extended matrices for all the groups G_k can be formed using $\mathcal{O}\left(n \log^2 n\right)$ time in total.

Since each Vandermonde matrix has $\mathcal{O}\left(dp^2\right)$ rows, then observe that each group G_k samples $\mathcal{O}\left(\frac{dp^2}{\varepsilon^2} \log d\right)$ rows and there are $\ell = \mathcal{O}\left(\frac{1}{\varepsilon} \log n\right)$ such groups $k \in [\ell]$. Thus the resulting subsampled matrix has $\mathcal{O}\left(\frac{dp^2}{\varepsilon^3} \log^2 n\right)$ rows for $d \leq n$. To approximately solve the L_p regression problem, Theorem A.5 notes that for $p \geq 2$ and a subsampled matrix of size $\mathcal{O}\left(\frac{dp^2}{\varepsilon^3} \log^2 d\right)$, we require only $\mathcal{O}\left(\frac{p\sqrt{d}}{\varepsilon^{3/2}} \log n \log \frac{d}{\varepsilon}\right)$ calls to a linear system solver. Moreover, on an iteration t of the L_p regression algorithm of Adil et al. (2019a) used in Theorem A.5, the linear system solves the equation $\mathbf{x}_t \leftarrow (\mathbf{A}^\top \mathbf{R}_T \mathbf{A})^{-1} \mathbf{A}^\top \mathbf{R}_t \mathbf{b}$ for a diagonal matrix $\mathbf{R}_T$. Thus by Theorem A.4, each linear system solve can be done in $\mathcal{O}\left(\frac{dp^2}{\varepsilon^3} \log^4 n\right)$ time for a Vandermonde matrix $\mathbf{A}$. Hence, the total time to approximately solve the L_p regression problem is $\mathcal{O}\left(\frac{p^3 d^{3/2}}{\varepsilon^{7/2}} \log^4 n \log \frac{n}{\varepsilon}\right)$. Therefore, the total runtime is $\mathcal{O}\left(n \log^2 n + \frac{p^3 d^{3/2}}{\varepsilon^{7/2}} \log^4 n \log \frac{n}{\varepsilon}\right)$. $\square$

Lemma 2.4. *Let $\mathbf{x} \in \mathbb{R}^n$ and $p = \Omega\left(\frac{\log n}{\varepsilon}\right)$. Then $\|\mathbf{x}\|_\infty \leq \|\mathbf{x}\|_p \leq (1 + \varepsilon)\|\mathbf{x}\|_\infty$.*

Proof. For any vector $\mathbf{x} \in \mathbb{R}^d$, we have

$$\|\mathbf{x}\|_\infty = \max_{i \in n} |x_i| \leq \left(\sum_{i \in} |x_i|^p\right)^{1/p} \leq n^{1/p} \cdot \max_{i \in n} |x_i|.$$

Since $(1+\varepsilon)^{3/eps} > e$ for all $\varepsilon > 0$, then $(1+\varepsilon)^p > n$ for $p = \Omega\left(\frac{\log n}{\varepsilon}\right)$. Therefore,

$$n^{1/p} \cdot \max_{i \in n} |x_i| \leq (1+\varepsilon) \max_{i \in n} |x_i| = (1+\varepsilon)\|\mathbf{x}\|_\infty.$$

$\square$

Corollary 2.5. *Given $\varepsilon \in (0,1)$ and $p \geq 1$, $\mathbf{A} \in \mathbb{R}^{n \times dq}$, and $\mathbf{b} \in \mathbb{R}^d$, suppose $\mathbf{A} = [\mathbf{A}_1| \ldots |\mathbf{A}_q]$ for Vandermonde matrices $\mathbf{A}_1, \ldots, \mathbf{A}_q \in \mathbb{R}^{n \times d}$. Then there exists an algorithm that, with high probability, returns a vector $\widehat{\mathbf{x}} \in \mathbb{R}^d$ such that*

$$\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1+\varepsilon) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p,$$

using $\mathcal{O}\left(T(\mathbf{A})(dp)^{q-1} \log n + \text{poly}((dp)^q \log n, 1/\varepsilon)\right)$ time, where $T(\mathbf{A})$ is the runtime of multiplying the matrix $\mathbf{A}$ by an arbitrary vector. For $q = 2$, this can be further optimized to $\mathcal{O}\left(nd^{\omega_2/2-1} + \text{poly}((dp)^2, \log n, 1/\varepsilon)\right)$ time, where $\mathcal{O}\left(n^{\omega_2}\right)$ is the time to multiply an $n \times n$ matrix with an $n \times n^2$ matrix, so that $\omega_2 \in [3, 4]$.

Proof. Recall that a key part in the proof of Theorem 1.1 was to first the L_{2^r} regression on a Vandermonde matrix with dimension $\mathbb{R}^{n \times d}$ to L_1 regression on a Vandermonde matrix with dimension $\mathbb{R}^{n \times (2^r(d-1)+1)}$, where $2^r \leq p < 2^{r+1}$. For a matrix $\mathbf{A}$ with block Vandermonde structure, we can similarly write

$$(\langle \mathbf{a}_i, \mathbf{x} \rangle)^{2^r} = \left(\sum_{j=1}^{dq} a_{i,j}x_j\right)^{2^r} = \left(\sum_{k=1}^q \sum_{j=1}^d a_{i,2+(k-1)d}^{j-1} x_{j+(k-1)d}\right)^{2^r}$$

$$= \sum_{j=1}^{(2^r(d-1))^q+1} \prod_{k \in [q], \sum p_{j,k}=2^r}^q a_{i,2+(k-1)d}^{p_{j,k}} y_j,$$

where again each y_j is a fixed function of the coordinates of $\mathbf{x}$. Thus we can reshape the L_{2^r} regression problem on a matrix $\mathbf{A}$ with dimension $\mathbb{R}^{n \times dq}$ with block Vandermonde structure to an L_1 regression problem on a matrix $\tilde{\mathbf{A}}$ with dimension $\mathbb{R}^{n \times (2^r(d-1))^q+1}$. Moreover, we can further reshape $\tilde{\mathbf{A}}$ into the concatenation of $(dp)^{q-1}$ Vandermonde matrices, where each Vandermonde matrix has columns that are geometrically growing in $a_{i,2}$ but are multiplied by all $(dp)^{q-1}$ products $\prod_{k=1}^{q-1} a_{i,2+kd}^{p_k}$, where $p_k \in [dp]$.

We can now use matrix-vector multiplication on each of the $(dp)^{q-1}$ Vandermonde matrices. Thus by Theorem A.2, we can L_1 Lewis weight sample from the rows of the reshaped $\tilde{\mathbf{A}}$, using $\mathcal{O}\left(T(\mathbf{A})(dp)^{q-1} \log n + (dp)^{\omega q}\right)$ time. We can similarly write $(\langle \mathbf{a}_i, \mathbf{x} - t_k \rangle)^{2^r}$ for each t_k among the discretized values of the updated $\mathbf{b}$ vector as a sum of $(2^r(d-1))^q + 1$ terms that are all products of powers of the bases $a_{i,2}, a_{i,2+d}, \ldots$ and a variables y_j, as in the proof of Theorem 1.1. Thus we can partition the L_{2^r} regression problem into $\ell = \mathcal{O}\left(\frac{\log n}{\varepsilon}\right)$ instances of a constrained L_1 regression problem on $(dp)^{q-1}$ Vandermonde matrices, each with at most $2^r(d-1))^q + 1$ columns. To approximately solve the L_p regression problem, we can thus sample rows by their $L_{p/2^r}$ Lewis weights, as in Theorem 1.1. Since there are up to $(dp)^{q-1}$ Vandermonde matrices, each with at most $2^r(d-1))^q + 1$ columns, then by Theorem A.2, the total time required is $\mathcal{O}\left(T(\mathbf{A})(dp)^{q-1} \log n + (dp)^{\omega q} \cdot \text{poly}(\log n, 1/\varepsilon)\right)$. $\square$

We remark that for the special case of $q = 2$, Sa et al. (2018) noted an efficient bivariate matrix multiplication algorithm of Nüsken & Ziegler (2004); Kedlaya & Umans (2011).

Theorem A.7. *(Nüsken & Ziegler, 2004; Kedlaya & Umans, 2011; Sa et al., 2018) Given a q-variate polynomial $f(X_1, \ldots, X_q)$ such that each variable has degree at most $d - 1$ and $N = d^q$ distinct points $x(i) = (x(i)_1, \ldots, x(i)_q)$ for $i \in [N]$, there exists an algorithm that uses $\mathcal{O}\left(d^{\omega_2(q-1)/2+1}\right)$ time to output the vector $(f(x(1)), \ldots, f(x(N)))$, where $\mathcal{O}\left(n^{\omega_2}\right)$ is the time to multiply an $n \times n$ matrix with an $n \times n^2$ matrix, so that $\omega_2 \in [3, 4]$.*

The case of a matrix-vector product for $q = 2$ corresponds to the evaluation of d^2 points in Theorem A.7. Thus we need to repeat the algorithm in Theorem A.7 a total of $\frac{n}{d^2}$ times to handle all n rows in the input matrix. Since each instance of the algorithm uses $\mathcal{O}\left(d^{w_2/2+1}\right)$ time, the total time for the matrix-vector product is $\mathcal{O}\left(nd^{\omega_2/2-1}\right)$, rather than the naïve $\mathcal{O}\left(nd\right)$ time (recall that $\omega_2 \in [3, 4]$).

Corollary A.8. *Given $\varepsilon \in (0, 1)$ and $p \geq 1$, $\mathbf{A} \in \mathbb{R}^{n \times 2d}$, and $\mathbf{b} \in \mathbb{R}^d$, suppose $\mathbf{A} = [\mathbf{A}_1 | \mathbf{A}_2]$ for Vandermonde matrices $\mathbf{A}_1, \mathbf{A}_2 \in \mathbb{R}^{n \times d}$. Then there exists an algorithm that with high probability returns a vector $\widehat{\mathbf{x}} \in \mathbb{R}^d$ such that*

$$\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \varepsilon) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p,$$

using $\mathcal{O}\left(nd^{\omega_2/2-1} + \text{poly}((dp)^2, \log n, 1/\varepsilon)\right)$, where $\mathcal{O}\left(n^{\omega_2}\right)$ time.

A.2 L_p REGESSION ON OTHER STRUCTURED INPUT

Algorithm 4 Faster regression for noisy low-rank matrices

Input: Sparse matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$, measurement vector $\mathbf{b}$, accuracy parameter $\varepsilon > 0$
Output: $\widehat{\mathbf{x}} \in \mathbb{R}$ with $\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \mathcal{O}(\varepsilon)) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p$
1: $r \leftarrow \lfloor \log p \rfloor$ $\triangleright 2^r \leq p < 2^{r+1}$.
2: Extend $\mathbf{A}$ to a matrix $\mathbf{M}$ with dimension $n \times \mathcal{O}\left(d^s(d + s)2^r\right)$ so that each entry $M_{i,j}$ is the coefficient of the j-th term in the tensor decomposition of $\mathbf{a}_i^{\otimes 2^r}$.
3: Use $L_{p/2^r}$-Lewis weight sampling on $\mathbf{M}$ to find a set S of $\mathcal{O}\left(d \log d\right)$ indices in $[n]$ and rescaling factors.
4: Let $\mathbf{A}'$ be the corresponding submatrix of $\mathbf{A}$ with indices in S and scaled accordingly.
5: Compute $\tilde{\mathbf{x}} \leq 5 \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}'\mathbf{x} - \mathbf{b}\|_p$.
6: $\mathbf{b}' \leftarrow \mathbf{b} - \mathbf{A}\tilde{\mathbf{x}}$, $\mathbf{b}'' \leftarrow \mathsf{RoundTrunc}(\mathbf{b}')$, $\ell \leftarrow \mathcal{O}\left(\frac{\log n}{\varepsilon}\right)$
7: Partition the rows of $\mathbf{A}$ into groups $G_1, \ldots, G_\ell$, each containing all rows with the same value of $\mathbf{b}''$
8: Let $\mathbf{G}_k$ be the corresponding submatrix and t_k be the coordinate of $\mathbf{b}''$ corresponding to G_k for each $k \in [\ell]$.
9: Use $L_{p/2^r}$-Lewis weight sampling on $[\mathbf{G}_k; t_k]$ to find a set S'_k of $O\left(\frac{1}{\varepsilon}d \log d\right)$ indices in $[n]$ and rescaling factors.
10: Let $\mathbf{T}_k$ be the corresponding rows with indices in S'_k and scaled accordingly.
11: $\mathbf{T} \leftarrow [\mathbf{T}_1; \ldots; \mathbf{T}_k]^\top$
12: Compute $\widehat{\mathbf{x}} \leq (1 + \varepsilon) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{T}\mathbf{x} - \mathbf{b}\|_p$.
13: **return** $\widehat{\mathbf{x}}$

Lemma A.9. *Given $\varepsilon \in (0, 1)$ and $p \geq 1$, a matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ such that $\mathbf{A} = \mathbf{K} + \mathbf{S}$ for a rank k matrix $\mathbf{K}$ and an s-sparse matrix $\mathbf{S}$, and $\mathbf{b} \in \mathbb{R}^d$, there exists an algorithm that with high probability, returns a vector $\widehat{\mathbf{x}} \in \mathbb{R}^d$ such that*

$$\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \varepsilon) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p.$$

Proof. The proof is similar to Lemma A.3. We once again let r be an integer so that $2^r \leq p < 2^{r+1}$ and observe that

$$\sum_{i \in [n]} |\langle \mathbf{a}_i, \mathbf{x} \rangle|^p = \sum_{i \in [n]} |(\langle \mathbf{a}_i, \mathbf{x} \rangle)^{2^r}|^{p/2^r} = \sum_{i \in [n]} |(\langle \mathbf{k}_i + \mathbf{s}_i, \mathbf{x} \rangle)^{2^r}|^{p/2^r}.$$

Since $\mathbf{K}$ is a low-rank matrix, then for all $\mathbf{k}_i$, we can write

$$\mathbf{k}_i = \sum_{j=1}^k \alpha_{i,j} \mathbf{v}_j,$$

for a fixed set of basis vectors $\mathbf{v}_1, \ldots, \mathbf{v}_k \in \mathbb{R}^d$. Hence we have

$$\sum_{i \in [n]} |\langle \mathbf{a}_i, \mathbf{x} \rangle|^p = \sum_{i \in [n]} \left| \left\langle \mathbf{s}_i + \sum_{j=1}^k \alpha_{i,j} \mathbf{v}_j, \mathbf{x} \right\rangle^{2^r} \right|^{p/2^r}.$$

Since $\mathbf{S}$ has sparsity s, then we can further write

$$\sum_{i\in[n]}|\langle\mathbf{a}_i,\mathbf{x}\rangle|^p = \sum_{i\in[n]}\left|\left\langle\sum_{j=1}^{s}\beta_{i,j}\mathbf{e}_{i_j}+\sum_{j=1}^{k}\alpha_{i,j}\mathbf{v}_j,\mathbf{x}\right\rangle^{2^r}\right|^{p/2^r},$$

where $\mathbf{e}_1,\ldots,\mathbf{e}_d$ denote the elementary vectors. By the Hadamard Product-Kronecker Product mixed-product property, we have

$$\sum_{i\in[n]}|\langle\mathbf{a}_i,\mathbf{x}\rangle|^p = \sum_{i\in[n]}\left|(\mathbf{y}_1\otimes\ldots\mathbf{y}_{2^r})\odot\mathbf{x}^{\otimes(2^r)}\right|^{p/2^r},$$

where each $\mathbf{y}_k\in\{\alpha_{i,1}\mathbf{v}_1,\ldots,\alpha_{i,k}\mathbf{v}_k,\beta_{i,1}\mathbf{e}_{i_1},\ldots,\beta_{i,s}\mathbf{e}_{i_s}\}$ for $i\in[2^r]$. Thus for a fixed set of elementary vectors $\mathbf{e}_{i_1},\ldots,\mathbf{e}_{i_s}$, there are $(d+s)^{2^r}$ possible values for the tensor product $(\mathbf{y}_1\otimes\ldots\mathbf{y}_{2^r})$. Since there are $\binom{d}{s}$ choices for the elementary vectors $\mathbf{e}_{i_1},\ldots,\mathbf{e}_{i_s}$, then there are at most $(Cd^s(d+s)^{2^r})$ possible values for the tensor product for an absolute constant $C>0$. Therefore, the L_p subspace embedding problem on $\mathbf{A}\in\mathbb{R}^{n\times d}$ can be reshaped as a constrained $L_{p/2^r}$ subspace embedding problem on an input matrix of size $n\times(Cd^s(d+s)^{2^r})$. Hence for L_p regression with $p\in[2^r,2^{r+1})$, we have

$$\sum_{i\in[n]}|\langle\mathbf{a}_i,\mathbf{x}\rangle|^p = \sum_{i\in[n]}\left|\sum_{j=1}^{2^r(d-1)+1}M_{i,j}y_j\right|^{p/2^r},$$

which is a constrained $L_{p/2^r}$ regression problem on a matrix $\mathbf{M}$ of size $n\times(Cd^s(d+s)^{2^r})$ whose entries can be determined from the decomposition of each row of $\mathbf{A}$.

Using $L_{p/2^r}$ Lewis weight sampling, Theorem A.2 implies that we can find a matrix $\mathbf{M}'$ such that

$$\frac{11}{12}\|\mathbf{M}\mathbf{y}\|_{p/2^r}^{p/2^r}\le\|\mathbf{M}'\mathbf{y}\|_{p/2^r}^{p/2^r}\le\frac{13}{12}\|\mathbf{M}\mathbf{y}\|_{p/2^r}^{p/2^r}$$

for all $\mathbf{y}\in\mathbb{R}^{(Cd^s(d+s)^{2^r})}$ with high probability. By Lemma A.1, we can thus compute a vector $\tilde{\mathbf{x}}$ such that

$$\|\mathbf{A}\tilde{\mathbf{x}}-\mathbf{b}\|_p\le 12\min_{\mathbf{x}\in\mathbb{R}^d}\|\mathbf{A}\mathbf{x}-\mathbf{b}\|_p.$$

We again set $\mathbf{b}'=\mathbf{b}-\mathbf{A}\tilde{\mathbf{x}}$ and define $\text{OPT}=\min_{\mathbf{x}\in\mathbb{R}^d}\|\mathbf{A}\mathbf{x}-\mathbf{b}\|_p$, so that

$$\|\mathbf{b}'\|_p=\|\mathbf{A}\tilde{\mathbf{x}}-\mathbf{b}\|_p\le 12\text{OPT}.$$

Let $\mathbf{b}'=\mathbf{b}-\mathbf{A}\tilde{\mathbf{x}}$ and $\text{OPT}=\min_{\mathbf{x}\in\mathbb{R}^d}\|\mathbf{A}\mathbf{x}-\mathbf{b}\|_p$, so that

$$\|\mathbf{b}'\|_p=\|\mathbf{A}\tilde{\mathbf{x}}-\mathbf{b}\|_p\le 12\text{OPT}.$$

Let $\mathbf{b}''$ be the vector with all entries of $\mathbf{b}'$ rounded to the nearest power of $(1+\varepsilon)$, starting at the maximum entry of $\mathbf{b}'$ in absolute value, and stopping after we are $\frac{1}{\text{poly}(n)}$ of that and replacing all remaining entries with 0. By the triangle inequality, we have

$$\|\mathbf{A}\mathbf{x}-\mathbf{b}''\|_p\le\|\mathbf{A}\mathbf{x}-\mathbf{b}'\|_p+\|\mathbf{b}'-\mathbf{b}''\|_p\le\|\mathbf{A}\mathbf{x}-\mathbf{b}'\|_p+12\varepsilon\text{OPT}\le(1+12\varepsilon)\|\mathbf{A}\mathbf{x}-\mathbf{b}''\|_p,$$

for any $\mathbf{x}\in\mathbb{R}^{d+1}$.

Since the coordinates of $\mathbf{b}''$ can have $\ell=O\left(\frac{\log n}{\varepsilon}\right)$ possible distinct values, we can partition the rows of $\mathbf{A}$ into ℓ groups, $G_1,\ldots,G_\ell$, based on the corresponding values of $\mathbf{b}''$. Let t_m be the corresponding value of $\mathbf{b}''$ for all rows in a group G_m, so that

$$\|\mathbf{A}\mathbf{x}-\mathbf{b}''\|_p^p=\sum_m\sum_{i\in G_k}|\langle\mathbf{a}_i,\mathbf{x}\rangle-t_m|^p.$$

By the above argument, we have for each $i\in G_k$,

$$(\langle\mathbf{a}_i,\mathbf{x}\rangle-t_m)^{2^r}=\left|-t_m+\left\langle\sum_{j=1}^{s}\beta_{i,j}\mathbf{e}_{i_j}+\sum_{j=1}^{k}\alpha_{i,j}\mathbf{v}_j,\mathbf{x}\right\rangle\right|^{2^r}=\sum_{j=1}^{2^rCd^s(d+s)^{2^r}}B_{i,j}t_{m,j}y_j,$$

where (1) $B_{i,j}$ are entries of a matrix $\mathbf{B}$ with $2^r C d^s (d+s)^{2^r}$ columns that can be computed from $\mathbf{A}$, (2) $t_{m,1}, t_{m,2}, \ldots$ are fixed values that can be computed from m_k, and (3) each y_j is a fixed function of the coordinates of $\mathbf{x}$. Notably, $\mathbf{B}$ is the matrix formed by the concatenation of the coefficients of the decomposition of the α-fold tensor product of the row into the α-fold tensor products of the low-rank and sparse basis elements, for each $\alpha = 0, \ldots, p$. By comparison, the matrix $\mathbf{M}$ previously defined in this proof is only the decomposition for $\alpha = p$. Hence, the L_{2^r} regression problem on a submatrix of $\mathbf{A} \in \mathbb{R}^{n \times d}$ the same coordinate of $\mathbf{b}''$, can be reshaped as a constrained L_1 regression problem on a matrix with $2^r C d^s (d+s)^{2^r}$ columns.

The remainder of the proof follows from the same grouping argument as Theorem 1.1. We apply Theorem A.2 by sampling rows of $\mathbf{B}$ corresponding to their $L_{p/2^r}$ Lewis weights in the submatrix induced by the rows of G_m, we obtain a matrix $\mathbf{R}_m$ such that with high probability,

$$(1-\varepsilon)\|\mathbf{R}_m \mathbf{y} - t_m\|_{p/2^r}^{p/2^r} \leq \sum_{i \in G_m} \left(\sum_{j=1}^{2^r C d^s (d+s)^{2^r}} B_{i,j} t_{m,j} y_j \right)^{p/2^r} \leq (1+\varepsilon)\|\mathbf{R}_m \mathbf{y} - t_m\|_{p/2^r}^{p/2^r}.$$

Conditioning on the above inequality holding, then by taking the matrix $\mathbf{T}_m$ corresponding to the scaled rows of $\mathbf{A}$ that are sampled by $\mathbf{R}_m$, it follows that

$$(1-\varepsilon)\|\mathbf{T}_m \mathbf{x} - t_m\|_p^p \leq \sum_{i \in G_m} |\langle \mathbf{a}_i, \mathbf{x} \rangle - t_m|^p \leq (1+\varepsilon)\|\mathbf{T}_m \mathbf{x} - t_m\|_p^p.$$

Therefore by summing over all $m \in [\ell]$, we have that with high probability,

$$(1-\varepsilon) \sum_m \|\mathbf{T}_m \mathbf{x} - t_m\|_p^p \leq \sum_m \sum_{i \in G_m} |\langle \mathbf{a}_i, \mathbf{x} \rangle - t_m|^p \leq (1+\varepsilon) \sum_m \|\mathbf{T}_m \mathbf{x} - t_m\|_p^p.$$

For $\mathbf{T} = \mathbf{T}_1 \circ \ldots \circ \mathbf{T}_\ell$, we have $\sum_m \|\mathbf{T}_m \mathbf{x} - t_m\|_p^p = \|\mathbf{T}\mathbf{x} - \mathbf{b}''\|_p^p$. Since $\|\mathbf{A}\mathbf{x} - \mathbf{b}''\|_p^p = \sum_m \sum_{i \in G_m} |\langle \mathbf{a}_i, \mathbf{x} \rangle - t_m|^p$, then

$$(1-\varepsilon)\|\mathbf{T}\mathbf{x} - \mathbf{b}''\|_p^p \leq \|\mathbf{A}\mathbf{x} - \mathbf{b}''\|_p^p \leq (1+\varepsilon)\|\mathbf{T}\mathbf{x} - \mathbf{b}''\|_p^p.$$

Therefore for $p \geq 1$,

$$(1-\varepsilon)\|\mathbf{T}\mathbf{x} - \mathbf{b}''\|_p \leq \|\mathbf{A}\mathbf{x} - \mathbf{b}''\|_p \leq (1+\varepsilon)\|\mathbf{T}\mathbf{x} - \mathbf{b}''\|_p.$$

Because $\ell = O\left(\frac{\log n}{\varepsilon}\right)$ and $\sum \mathcal{O}(T(G_k)) = \mathcal{O}(T(\mathbf{A}))$, we can compute a vector $\widehat{\mathbf{x}} \in \mathbb{R}^d$ such that

$$\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \mathcal{O}(\varepsilon)) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p.$$

$\square$

Lemma A.10. *The runtime of Algorithm 4 is* $\mathcal{O}\left(p n^p k + p n s^p\right) + \text{poly}\left(d^s, k^p, s^p, \frac{1}{\varepsilon}\right)$.

Proof. We analyze the runtime of Algorithm 4. First note that we can compute the extended matrix in time $\mathcal{O}\left(n((k+1)^p d^s s^p)\right)$ to perform the $L_{p/2^r}$ Lewis weight sampling, where we recall that r is the unique integer such that $2^r \leq p < 2^{r+1}$. To perform Lewis weight sampling on the extended matrix, we require matrix-vector multiplication, which requires time $\mathcal{O}(nk)$ for a low-rank matrix and time $\mathcal{O}(ns)$ for a matrix whose rows have at most s nonzero entries.

After the first iteration of Lewis weight sampling, we can round and truncate the coordinates of $\mathbf{b} \in \mathbb{R}^n$ to form a vector $\mathbf{b}'' \in \mathbb{R}^n$ using the procedure RoundTrunc and then forming the groups $G_1, \ldots, G_\ell$, which clearly takes $\mathcal{O}(n)$ arithmetic operations combined. Once the groups are formed, we can compute the extended matrix in time $\mathcal{O}\left(n((k+1)^p d^s s^p)\right)$ and perform $L_{p/2^r}$ Lewis weight sampling on each group, which takes $\mathcal{O}(nk + ns)$ total time across all groups. To approximately solve the resulting L_p regression problem formed by the subsampled rows, we require $\text{poly}\left(d^s, k^p, s^p, \frac{1}{\varepsilon}\right)$ time. Therefore, the total runtime is $\mathcal{O}(nk + ns) + \text{poly}\left(d^s, k^p, s^p, \frac{1}{\varepsilon}\right)$. $\square$

We obtain similar results for a noisy Vandermonde matrix.

Lemma A.11. *Given $\varepsilon \in (0,1)$ and $p \geq 1$, a matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ such that $\mathbf{A} = \mathbf{V} + \mathbf{S}$ for a Vandermonde matrix $\mathbf{V}$ and an s-sparse matrix $\mathbf{S}$, and $\mathbf{b} \in \mathbb{R}^d$, there exists an algorithm that with high probability, returns a vector $\widehat{\mathbf{x}} \in \mathbb{R}^d$ such that*

$$\|\mathbf{A}\widehat{\mathbf{x}} - \mathbf{b}\|_p \leq (1 + \varepsilon) \min_{\mathbf{x} \in \mathbb{R}^d} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p.$$

Lemma A.12. *The runtime of Algorithm 4 is $\mathcal{O}(nk + ns) + \text{poly}\left(d^s, pd, s^p, \frac{1}{\varepsilon}\right)$.*

Theorem 1.6 then follows from Lemma A.11 and Lemma A.12.

B   APPLICATIONS TO POLYNOMIAL REGRESSION

In the polynomial regression problem, the goal is to find a degree d polynomial $\hat{q}$ such that

$$\|\widehat{q}(t) - f(t)\|_p^p \leq (1 + \varepsilon) \cdot \min_{q:\deg(q) \leq d} \|q(t) - f(t)\|_p^p,$$

where $\varepsilon > 0$ is an accuracy parameter given as input and $\|\cdot\|_p^p$ is the L_p norm to the p^{th} power, $\|g\|_p^p = \int_{-1}^1 |g(t)|^p dt$. The polynomial regression problem is a fundamental problem in statistics, computational mathematics, machine learning, and more. The problem has been studied as early as the 19th century with the work of Legendre and Gauss on least squares polynomial regression and has applications in learning half-spaces Kalai et al. (2008), solving parametric PDEs Hampton & Doostan (2015), and surface reconstruction Pratt (1987).

Given the flexibility to choose query locations $x_1, \ldots, x_s$, we can consider the polynomial regression problem as an *active learning* or *experimental design* problem. Thus we would like to minimize the number of queries s, as a function of the approximation degree d, the norm p, and the accuracy parameter ε, to find $\hat{q}$. Observe that $d+1$ queries are obviously necessary, but also that $d+1$ queries suffice when f can be exactly fit by a degree d polynomial, by using direct interpolation. In general, however, in the case when $\min_{q:\deg(q) \leq d} \|q(t) - f(t)\|_p^p \neq 0$ we require $s > d + 1$ queries. Our Vandermonde L_p regression results can be used to give the first result showing that for all $p \geq 1$, $d \exp(O(p)) \cdot \text{poly}\left(\frac{1}{\varepsilon}\right)$ queries suffice to obtain a $(1+\varepsilon)$-approximation to the best polynomial fit.

We require the following structural theorem reducing the L_p polynomial regression problem to a problem of solving L_p regression on Vandermonde matrices:

Theorem B.1. *Kane et al. (2017); Meyer et al. (2021) Suppose $s_1, \ldots, s_{n_0}$ are drawn uniformly from $[-1,1]$. Let $\mathbf{A} \in \mathbb{R}^{n_0 \times (d+1)}$ be the associated Vandermonde matrix, so that $\mathbf{A}_{i,j} = s_i^{j-1}$. Let $n_0 = \exp(O(p))\tilde{O}\left(\frac{1}{\varepsilon^{2+2p}} d^5\right)$ and let $\mathbf{b} \in \mathbb{R}^{n_0}$ be the evaluations of f, so that $\mathbf{b}_i = f(s_i)$. Then with probability $\frac{11}{12}$, the sketched solution $\hat{\mathbf{x}} = \text{argmin}_{\mathbf{x}} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p$ satisfies*

$$\|\mathcal{P}\mathbf{x} - f\|_p^p \leq (1 + \varepsilon) \min_{\mathbf{x} \in \mathbb{R}^{d+1}} \|\mathcal{P}\mathbf{x} - f\|_p^p.$$

Theorem B.1 states that we can uniformly sample $\exp(O(p)) \cdot \text{poly}\left(d, \frac{1}{\varepsilon}\right)$ points from $[-1,1]$. We can then form an L_p regression problem by using the evaluation each of the polynomial bases at the sampled points to form the design matrix $\mathbf{A}$ and querying the underlying signal f at the sampled points to form the measurement vector $\mathbf{b}$. Theorem B.1 says that the optimal solution $\hat{\mathbf{x}} = \text{argmin}_{\mathbf{x}} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p$ is a $(1+\varepsilon)$-approximation to the best fit degree d polynomial. We can naïvely approximately solve the L_p regression on the Vandermonde matrix $\mathbf{A}$ and the measurement vector $\mathbf{b}$ by standard L_p regression techniques such as Lewis weight sampling, which would result in total query complexity $\exp(O(p))\tilde{O}\left(\frac{1}{\varepsilon^{2+2p}} d^5\right)$. However, we can instead note that Lemma 1.4 implies we can instead solve an L_q regression problem on a Vandermonde matrix with $O(dp)$ columns for $q \in [1,2]$. Crucially for $q \in [1,2]$, there exist active L_q regression algorithms that only require reading $\tilde{O}(d) \cdot \text{poly}\left(\frac{1}{\varepsilon}\right)$ entries of $\mathbf{b}$:

Theorem B.2. *Musco et al. (2021b) Given $p \in [1,2]$ and an input matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$, there exists an algorithm that reads $\tilde{O}(d) \cdot \text{poly}\left(\frac{1}{\varepsilon}\right)$ entries of $\mathbf{b}$ and with probability at least 0.99, outputs $\tilde{\mathbf{x}}$ such that*

$$\|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p \leq (1 + \varepsilon) \min_{\mathbf{x}} \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_p.$$

Hence by Theorem B.2, we can approximately solve the L_q regression on a Vandermonde matrix with dp rows by reading $\tilde{O}(dp) \cdot \text{poly}\left(\frac{1}{\varepsilon}\right)$ entries of $\mathbf{b}$. By Lemma 1.4, the approximate solution will also be a $(1+\varepsilon)$-approximation to the optimal L_p regression on the Vandermonde matrix $\mathbf{A}$. By Theorem B.1, the approximate solution will also form the coefficient vector of a polynomial that is a $(1+\varepsilon)$-approximation to the polynomial L_p regression problem. Therefore, we obtain the following guarantees for the polynomial regression problem:

Theorem B.3. *For any degree d and norm $p \geq 1$, there exists an algorithm that queries f at $s = dp \, \text{poly}\left(\log(dp), \frac{1}{\varepsilon}\right)$ points and outputs a degree d polynomial $\widehat{q}(t)$ such that*

$$\|\widehat{q}(t) - f(t)\|_p^p \leq (1+\varepsilon) \cdot \min_{q:\deg(q)\leq d} \|q(t) - f(t)\|_p^p,$$

with probability at least $\frac{2}{3}$.

The previous-best algorithm Kane et al. (2017); Meyer et al. (2021) for L_p polynomial regression sampled $O(d^5)$ points uniform from $[-1,1]$ and then used standard L_p regression algorithms that required reading the signal at all $O(d^5)$ sampled points, for a total query complexity of $O(d^5)$. By comparison, Theorem B.3 only has linear dependency in d due to the structural property of Lemma 1.4. Since $\Omega(d)$ queries are clearly necessary, our result settles the dependency of d in the query complexity for L_p polynomial regression for all d and all $p \geq 1$.

C ADDITIONAL EXPERIMENTS

In this section, we included additional experiments comparing our algorithms to uniform sampling. Uniform sampling is a common heuristic that is used effectively on "average-case" inputs but lacks provable guarantees for "worst-case" inputs. By contrast, our theoretical analysis provides provable guarantees against "worst-case" inputs, but it may be possible that on "average-case" inputs, it could potentially perform worse than uniform sampling. We show that this is not the case on a standard 5000×20 sized Vandermonde matrix with equally spaced points on the grid. Indeed, our results in Figure 3a show that our algorithm significantly better performance than uniform sampling across many choices of the norm parameter p, while our results in Figure 3b show that our algorithm significantly better performance than sample complexity m.

(a) Empirical Relative Error ($\varepsilon_{empirical}$) versus L_p parameter p.

(b) Empirical Relative Error ($\varepsilon_{empirical}$) versus subsample complexity m.

D ADDITIONAL RELATED WORK

As previously mentioned, subspace embeddings are common tools used to approximately solve L_p regression. Given an input matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$, a subspace embedding is a matrix $\mathbf{M} \in \mathbb{R}^{m \times d}$ with $m \ll n$ such that

$$(1-\varepsilon)\|\mathbf{A}\mathbf{x}\|_p \leq \|\mathbf{M}\mathbf{x}\|_p \leq (1+\varepsilon)\|\mathbf{A}\mathbf{x}\|_p,$$

for all $\mathbf{x} \in \mathbb{R}^d$. Thus given an instance of L_p regression, where the goal is to minimize $\|\mathbf{Ax} - \mathbf{b}\|_p$, we can set $\mathbf{B} = [\mathbf{A}; \mathbf{b}]$, compute a subspace embedding for $\mathbf{B}$, and then solve a constrained L_p regression problem on the smaller matrix $\mathbf{B}$.

A subspace embedding of a matrix $\mathbf{A} \in \mathbb{R}^{n \times d}$ can be formed by sampling rows of $\mathbf{A}$ with probabilities proportional to their L_p leverage scores, e.g., (Cormode et al., 2018; Dasgupta et al., 2008) their L_p sensitivities, e.g., (Clarkson et al., 2019; Braverman et al., 2020; Musco et al., 2021a); or their L_p Lewis weights, e.g., (Cohen & Peng, 2015; Durfee et al., 2018; Chhaya et al., 2020; Chen & Derezinski, 2021; Parulekar et al., 2021; Meyer et al., 2021). In any of these cases, the resulting matrix $\mathbf{M}$ will contain a subset of rows of $\mathbf{A}$ that are rescaled by a function of their sampling probability, as to give an unbiased estimate of the actual L_p mass.

In addition to sampling methods, sketching is a common approach for subspace embeddings. In these cases, the subspace embedding $\mathbf{M}$ is formed by setting $\mathbf{M} = \mathbf{RA}$ for some (often random) matrix $\mathbf{R} \in \mathbb{R}^{m \times n}$. The advantage of sketching over sampling is that sometimes $\mathbf{R}$ can be computed oblivious to the structure of $\mathbf{A}$, whereas the sampling probabilities for each of the above distributions (L_p leverage scores, L_p sensitivities, and L_p Lewis weights) are data dependent and thus require a pass over the matrix $\mathbf{A}$. The sketching matrix can be generated from a family of random matrices whose entries are Cauchy random variables for $p = 1$, e.g., Sohler & Woodruff (2011); Meng & Mahoney (2013); Clarkson et al. (2016), or sub-Gaussian random variables for $p = 2$, e.g., Sarlós (2006); Nelson & Nguyen (2013); Clarkson & Woodruff (2013). More generally, exponential random variables can be used for $p \geq 2$, though the number of rows m in the resulting sketch matrix now has a polynomial dependency in n Woodruff & Zhang (2013). A line of recent works has studied the tradeoffs between oblivious linear sketches, sampling-based algorithms, and other sketches for L_p subspace embeddings Wang & Woodruff (2019); Li et al. (2021). In this paper, we focus on sampling-based algorithms for L_p regression due to preservation of structure when the input matrix $\mathbf{A}$ itself has structure.