Evaluating Age-related Anatomical Consistency in Synthetic Brain MR Images against Real-world Alzheimer's Disease Data.

Hadya Yassin[1,2]   HADYA.YASSIN@HPI.DE
Jana Fehr[1,2,3]   JANA.FEHR@BIH-CHARITE.DE
Wei-Cheng Lai[1,2]   WEI-CHENG.LAI@HPI.DE
Alina Krichevsky[1,2]   ALINA.KRICHEVSKY@HPI.UNI-POTSDAM.DE
Alexander Rakowski[1,2]   ALEXANDER.RAKOWSKI@HPI.DE
Christoph Lippert[1,2]   CHRISTOPH.LIPPERT@HPI.DE

[1] *Digital Engineering Faculty, University of Potsdam, Germany*

[2] *Digital Health and Machine Learning, Hasso-Plattner-Institute, Potsdam, Germany*

[3] *QUEST Center for Responsible Research, Berlin Institute of Health (BIH), Charité Universitätsmedizin Berlin, Germany*

Editors: Under Review for MIDL 2024

Abstract

This study examines the realism of medical images created with deep generative models, specifically their replication of aging and Alzheimer's disease (AD) related anatomical changes. Previous research focused on developing generative methods with limited attention to image fidelity. We aim to assess the resemblance of brain MR images generated by a StyleGAN3 model with causal controls to neurodegenerative changes. For a benchmark, we conducted a visual Turing test (VTT) to see if radiologists could distinguish between synthetic and real images. Then, we employed a U-Net-based model to segment hallmarks relevant to normal aging and AD. Finally, we conducted statistical tests for our hypothesis that no significant differences existed between real and synthetic images. VTT results showed radiologists struggled to differentiate between image types, highlighting VTT's limitations due to subjectivity and time constraints. We found slight hippocampus distribution differences ($P = 5.7$e-2) and significant lateral ventricle discrepancies (Ps < 5.0e-2), indicating higher hippocampus realism and ventricle size inconsistencies. The model more effectively simulated changes in the hippocampus than in the lateral ventricles, where difficulties were encountered with certain subgroups. We conclude that the VTT alone is inadequate for a comprehensive quality evaluation, promoting a more objective approach for detailed anatomical consistency analysis. Future research could adapt our approach to evaluate other generated medical images intended for different downstream tasks.

Keywords: Deep Generative Models, Synthetic Medical Images, Deep Learning Segmentation, Visual Turing Test, Anatomical Consistency, Alzheimer's Disease, Magnetic Resonance Imaging.

1. Introduction

Deep learning algorithms, essential for automating medical image analysis, rely heavily on access to abundant, high-quality datasets for training (Sarker, 2021). Acquiring such data is hindered by high costs associated with equipment, expert annotation, subject availability,

and privacy concerns (Diaz et al., 2021; Kaissis et al., 2020). Consequently, these challenges can lead to data biases that significantly impair the algorithms' performance in real-world applications, as they may not accurately reflect the diversity of real patient populations or the complexity of medical conditions encountered in clinical settings (Nittas et al., 2023).

Despite significant progress in synthetic medical image generation through generative adversarial networks (GANs) (Dash et al., 2023) and Diffusion models (Yang et al., 2023), traditional evaluation metrics like the Fréchet inception distance (FID) and Structural Similarity Index Measure (SSIM) primarily assess overall image quality, overlooking anatomical accuracy critical for medical applications. Thus, accurately mirroring human physiology and conditions like Alzheimer's disease (AD) is crucial for synthetic images to bridge dataset gaps. This requires "Biological Plausibility" metrics, introduced by (Treder et al., 2022) and designed to provide an anatomical analysis specific to clinical needs. Our evaluation method focuses on crucial neurodegenerative features for normal aging, which are more prominent in AD, including ventricular enlargement and hippocampal shrinkage (Frisoni et al., 2010; Katabathula et al., 2021).

Visual Turing test (VTT) is a standard for assessing synthetic images' biological plausibility in the generative field, used by Khader et al. (2022) to evaluate diffusion model performance. Yet, Treder et al. (2022) highlight VTT's limitations: subjectivity, high costs, and difficulty detecting subtle anatomical changes. These issues suggest VTT's insufficiency in ensuring synthetic images' clinical applicability. Our study benchmarks against VTT, promoting more objective, robust, and clinically relevant evaluation methods to improve synthetic images' clinical utility.

Longitudinal GAN studies on AD explore biological plausibility. Xia et al. (2021) analyze relative changes (RC) in the region of interest (ROI) volumes between baseline and follow-up images, real and synthetic. Peng et al. (2021) examine the ROI Absolute Volume Difference (AVD) between synthetic follow-ups and real manual segmentation. Fu et al. (2023) calculate mean absolute error (MAE) for ROI volumes in longitudinal data, differing from Ravi et al. (2022); Ribeiro et al. (2023) that randomly match samples by age, sex, and clinical dementia rating (CDR). The pairwise approach in longitudinal studies, tracking changes in the same individual, offers more accuracy than random matching. Ribeiro et al. (2023) evaluates model-predicted rather than actual volumes segmented from synthetic images, which may misrepresent true volumes. These methods aim to compare anatomical consistencies between real and synthetic data but reduce the evaluation to a single average value (MAE, RC, AVD), favoring benchmark comparisons of generative models over anatomical consistency and clinical relevance. Wilms et al. (2022) investigate age-related brain volume changes across generative models without real data comparison, omitting key evaluations of synthetic-to-real anatomical consistency.

Additionally, the mentioned methods segment ROIs using statistical and algorithmic strategies, such as FSL (FMRIB Software Library) and FreeSurfer (Jenkinson et al., 2012; Fischl, 2012), or multi-atlas methods (Wang et al., 2014; Doshi et al., 2013). In contrast, our method utilizes advanced capabilities of deep learning. Litjens et al. (2017); Shen et al. (2017) demonstrate that deep learning surpasses traditional methods in accuracy, efficiency, and detecting complex patterns.

Deep learning segmentation enables accurate anatomical consistency comparisons between synthetic ROI sizes and real counterparts, capturing essential variability in aging

and AD. Such analysis is crucial for clinical applications requiring high accuracy and transparency. Our objective is to assess the null hypothesis (H0) that there is no significant distributional difference in brain ROI areas between real Alzheimer's disease neuroimaging initiative (ADNI) images and their synthetic counterparts. We utilize statistical tests to examine differences in magnitude, certainty, direction, and distribution (distr) shapes. Our work, while similar to Jung et al. (2023); Xia et al. (2021) that perform statistical analysis to identify distr differences, focuses on anatomical consistencies across CDR groups through multi-hypothesis testing. Unlike these studies, our investigation extends to age and sex covariates, employing a different approach to circumvent multi-hypothesis testing.

2. Methodology:

Biological plausibility overview in Figure 1. Input: AD-focused Synthetic images generated by a conditional GAN with causal control and a subset of the ADNI dataset as the real-world reference (unseen in training and validation of both generative and segmentation models to reduce bias). VTT was selected as a benchmark for its established role in assessing synthetic image realism. Our method quantifies ROI area via a segmentation model and performs statistical tests to evaluate anatomical consistency between image types across different covariates. Understanding our analysis requires familiarity with real-world and synthetic data characteristics, synthetic image generation, and the generative model's capabilities.

Figure 1: Overview of the analysis

2.1. Use Case and Datasets:

Real Data. We sourced real magnetic resonance (MR) brain scans from the UK Biobank (UKB) (Sudlow et al., 2015) and ADNI (Alzheimer's Disease Neuroimaging Initiative (ADNI), 2021) datasets. The UKB provided 42,427 high-resolution T1-weighted 3T MR brain scans of individuals aged 40-69. Unlike the UKB, which serves a general cohort, the ADNI provides a focused dataset on specific diseases. The ADNI database (adni.loni.usc.edu), established in 2003, explores the use of imaging, biological markers, and clinical assessments to track the progression of mild cognitive impairment (MCI) and early AD. ADNI consists of high-resolution T1-weighted 3T MR brain scans from 9,183 participants aged 55 to 90 years (mean age: 74), including 3,273 cognitive normal (CN) cases (CDR=0), 4,943 MCI cases (CDR=0.5), and 967 AD cases (CDR≥1).

Synthetic Data. We generated 600,000 2D mid-slice images using a conditional style-based generative adversarial network (StyleGAN3) model (Karras et al., 2021) with causal control (Pawlowski et al., 2020). While Kocaoglu et al. (2017) introduced causal inference to StyleGAN3, our model combines a conditional GAN structure (Mirza and Osindero, 2014; Miyato and Koyama, 2018) with a custom causal model. To achieve a high-quality synthesis,

we initially trained the model on the UKB dataset for its extensive data volume, followed by fine-tuning it on the ADNI dataset. For both datasets, our causal model consistently incorporated covariates like **age**, **sex**, and **left and right lateral ventricle volumes**. Given the disease-specific nature of the ADNI, we added AD-associated conditional labels like **CDR** and **hippocampus volumes** (Frisoni et al., 2010; Katabathula et al., 2021) during its fine-tuning.

During the inference step, the images are synthesized by the causal model and the Style-GAN3 model. The causal model is used to provide conditional labels as vectors, *i.e.*, **age**, **sex**, **CDR**, **lateral ventricle volumes**, **cerebral cortex volumes**, and **hippocampus volumes**. Taking advantage of the causal model allows us to control labels by changing ages. This results in different volumes since our assumption is that age causally affects brain volumes. These conditional labels are fitted into the StyleGAN3 model to synthesize images with specific attributes related to age, sex, and either healthy or AD diagnoses, as indicated by CDR values, for more details on the conditional causal model (Appendix A).

2.2. Benchmark: Visual Turing Test (VTT)

To assess synthetic image realism, we conducted a VTT with two radiologists reviewing 100 image pairs: 50 from UKB and 50 from ADNI with their synthetic counterparts. Each pair had one real and one synthetic image, matched by age, sex, and CDR for ADNI images only. The radiologists identified the synthetic image, rated their confidence, and explained their choices, enabling a comprehensive realism evaluation. The test, done with ImFusion Labels software (ImF), was untimed to allow breaks and resumptions.

2.3. Biological Plausibility Analysis

ROI Quantification via Segmentation. Our generative model produces 2D mid-slice brain MR images, while established deep-learning segmentation models like SynthSeg (Billot et al., 2023) are designed for 3D volumes. To bridge this gap, we developed a 2.5D segmentation model suitable for 2D and 3D data. Using SynthSeg, we generated ground truth masks for UKB and ADNI datasets. To prevent bias in our segmentation towards ADNI (real-world reference data), we trained and validated state-of-the-art (SOTA) architectures solely on UKB.

During inference, the best-performing model adapted UNet (A-UNet), was selected for its highest median dice similarity coefficient (DSC) from external ADNI testing. This model, an adaptation of the original UNet (Ronneberger et al., 2015), was optimized by increasing initial features from 32 to 64 and adding a layer of depth. Details on model implementation and comparisons with other architectures are in Appendix C. We quantified ROI areas by counting pixels in segmented masks, then normalized these areas against the intracranial area to account for brain size variations. Min-max scaling was applied to these normalized values to ensure consistency with observed area ranges.

Statistical Analysis. In the VTT, a two-tailed binomial test evaluated radiologists' accuracy against random guessing and Cohen's kappa for inter-rater agreement. To evaluate anatomical consistency between image types, we used statistical tests to explore the hypothesis that no significant differences exist in the ROI distributions between real and

synthetic images, setting the significance level at 0.05. The permutation (permutation test (permute)), chosen for its non-parametric nature that requires no distribution assumptions, assesses the statistical significance of mean differences through repeated recalculations, minimizing false significance risks. Subsequently, Cohen's d was used to measure effect size and direction. Additionally, we employed 95% confidence interval (CI), based on recommendations from Lee (2016), to asses effect size (ES) estimation precision and indicate the statistical significance of our findings. Finally, The Kolmogorov-Smirnov test (KS) test further compared distribution shapes, providing added insights into discrepancies between real and synthetic image distributions.

3. Experiments and Results

Experiment Setup. Our experiments evaluated the impact of aligning covariate distributions (age, CDR, and sex) on image analysis, maintaining a 5:1 synthetic-to-real ROI area ratio throughout. We conducted experiments on matched and mismatched covariate distribution, where we would expect higher deviation between real and synthetic ROI distributions when deviating from matching distributions. The experiments are: a) matched covariate distributions, b) introduced variances by adjusting the median age of the synthetic samples' distribution by ± 5 years, c) CDR, and d) sex adjustments in synthetic distribution are detailed in Appendix D.

VTT Results. Radiologist 1 (R1) achieved a 51% accuracy rate, similar to random guessing ($P = 9.2$e-1), while radiologist 1 (R2) scored 15%, a significant difference ($P = \mathbf{4.8e\text{-}13}$). Cohen's Kappa of 0.167 indicates low agreement between radiologists, suggesting variability in image perception and image fidelity conclusions. R1 often indicated 'moderate certainty' and cited 'contrast problems,' while R2 often reported 'very low certainty' and highlighted 'anatomical inaccuracies' as their certainty level and reasons behind their decisions. Interestingly, neither of them considered an 'inaccurate representation of pathology'. A follow-up revealed R2's bias stemming from MR characteristics familiarity, leading to misclassifications by associating smoothing effects with real images despite their presence in both types. Figure 3 a) displays a misclassification by R2, also incorrectly identified by R1 (see Figure 6 in appendix), both for different reasons (anatomical inaccuracies and noise pattern, respectively), despite both images being real. The sole instance where both radiologists strongly agreed and accurately identified a synthetic image, citing inaccurate anatomy due to unusually small ventricles, is shown in Figure 3 b). Further VTT results in Appendix B.

Biological Plausibity Results. Regression analysis in Figure 2 reveals that Lateral Ventricles (LV) areas increase and Hippocampus (HC) areas decrease with age, aligning with neurodegeneration patterns in both real and synthetic images. Furthermore, Table 1 demonstrates significant differences for LV in **matched distrs** ($P = \mathbf{2.0e\text{-}3}$, **negative** $CI -_l -_u$), indicating synthetic images overestimate LV enlargement with age compared to real images (Negative ES, see Figure 2). LV KS test results show significant shape differences in distributions across **both matched and unmatched conditions** (Figure 8). In contrast, hippocampus results show no significant difference in permute tests, though a near-significant p (5.7e-2) and positive ES (7.0e-2) suggest minor overestimation in synthetic image ROI size reduction with age at matched distrs. Detailed analysis in Figure 2 indicates

smaller differences in LV within 75-80 and 80-85 age bins, with nearly overlapping boxes. For HC, 60-65 and 85-90 age bins display lower synthetic medians, while 80-85 shows higher synthetic medians.

Table 1: Statistical Comparison between Real vs. Synthetic ROI Areas Across Distribution. **Match:** same covariate distr for Image Types (Median age of 74); **Mismatch:** Synthetic distr age median shifts by $\pm$ **3 or 5 years**). $CI -_l -_u$: CI bounds. **Significant differences** highlighted.

ROI	Cond.	Permute	ES	CI_l	CI_u	KS
LV	-5	6.0e-2	7.1e-2	-5.8e-3	1.5e-1	**1.3e-2**
LV	-3	8.2e-1	9.5e-3	-6.7e-2	8.6e-2	**1.4e-2**
LV	Match	**2.0e-3**	-1.3e-1	**-2.0e-1**	**-5.0e-2**	**2.0e-8**
LV	+3	**0.0e-0**	-2.3e-1	**-3.1e-1**	**-1.6e-1**	**2.1e-14**
LV	+5	**5.0e-2**	-3.1e-1	**-3.9e-1**	**-2.3e-1**	**4.4e-16**
H	-5	**4.0e-3**	-1.1e-1	**-1.9e-1**	**-3.6e-2**	**1.1e-2**
H	-3	3.2e-1	-4.0e-2	-1.2e-01	3.7e-2	4.5e-1
H	Match	5.7e-2	7.0e-2	-6.7e-3	1.5e-1	8.7e-2
H	+3	**4.0e-3**	1.1e-1	**3.8e-2**	**1.9e-1**	**2.4e-3**
H	+5	**5.0e-2**	2.1e-1	**1.4e-1**	**2.9e-1**	**9.4e-8**

Mismatched Conditions with age adjustments (+3, +5) showed significant differences in LV size, with larger negative ES than the matched distr. Conversely, deviations of (-3, -5) lacked significant findings and had smaller ES, particularly at -3, confirming the generative model's overestimation of LV sizes in matched distrs (As seen in Figure 2). For HC, the -3 mismatch, like matched distr, showed no significant differences but had a lower ES, suggesting minor overestimation in matched conditions due to its higher ES and a closer P value to significance. Finally, all other conditions showed significance.

Outliers and the whisker ranges of box plots for both ROIs in Figure 2 show variability within and between image types, with synthetic images exhibiting a broader area size range than real ones. Closer analysis shows more outliers in real images for lateral ventricles and in synthetic images for the hippocampus. Notably, 0.82% of synthetic ventricles' areas were below the minimum observed areas in real data. These sizes are clinically significant, indicating implausibly small lateral ventricles for adults (Agreed on by R1 & R2). Figure 3 b) illustrates such a case, clearly showing the differences in the unusual ventricle sizes of synthetic vs. real images for the same covariates, highlighting the need for rigorous evaluation of synthetic image generation for clinical use. Despite the segmentation model's training on real data, the model accurately segmented the unusually small ventricles (unseen in training), demonstrating robustness.

4. Discussion

The VTT reveals the challenge radiologists face in distinguishing synthetic from real brain MR images, attributed to the synthetic images' compelling realism and resulting in varied accuracy values. R2's frequent misclassifications, driven by cognitive biases, hint that performance could potentially improve through retraining with varied image types. Nonetheless, the smoothness effect is not only present in synthetic images but also in real ADNI and UKB images. Additionally, R2 consistently reported low confidence, often citing 'anatomical inaccuracies' in real images identified as synthetic, emphasizing the complexity of identifying subtle changes in ROIs linked to normal aging or AD progression. Although both

Figure 2: Age-wise comparison of observed LV and HC areas in real versus synthetic MR brain images across matched matched distrs shows aging effect on brain ROI sizes.

radiologists agreed and correctly identified the unusually small ventricles, manually inspecting 600,000 synthetic images to find similar conditions is infeasible. In contrast, our method allows for efficient identification of such cases. Based on these challenges, VTT alone is insufficient as a definitive measure of image realism, necessitating a more objective approach.

Our study advances this objective by rigorously evaluating the generative model's ability to replicate age-related anatomical changes, revealing its strengths and pinpointing limitations. Unlike the VTT, our method provides precise insights into anatomical accuracy, especially highlighting how the model overestimates ventricular enlargement while performing better in simulating hippocampal changes. This nuanced understanding highlights the importance of employing focused methods to evaluate both the realism and anatomical precision of synthetic medical imagery. Additionally, Observing a wide range of area sizes in synthetic images and outliers, particularly implausibly small ventricles, illustrates the model's current limitations in capturing the full spectrum of individual variability in real-world data.

The presence of biases in synthetic data, especially those that do not accurately reflect real-world conditions, poses a significant challenge in research on aging and neurodegenerative diseases. When synthetic images inaccurately represent brain ventricles as larger than expected, this introduces an unrealistic bias that can mislead studies, potentially leading to incorrect conclusions about ventricular changes. This issue is compounded in areas like AD research, where patient data may be scarce, yet it's vital that the data used reflect genuine conditions to maintain the integrity of findings. The risk extends to diagnostic accuracy, where reliance on biased synthetic images could lead to misdiagnoses or the formulation of ineffective treatment plans. Similarly, training AI systems with these inaccurately biased images could lead to errors in clinical applications, particularly those dependent on precise identification of age-related changes. To mitigate these challenges, it's crucial to refine

generative models to more faithfully represent the complexities of aging. By tackling these biases head-on, we can enhance the realism and reliability of synthetic images, thereby supporting more accurate clinical research, diagnostics, and the effective use of AI in healthcare settings.

a) Synthetic (left) vs. Real (right) (Male, 66, CN).

b) Synthetic (left) vs. Real (middle & right) (Female, 81, MCI).

Figure 3: a) R2's misclassification and incorrectly noting 'anatomical inaccuracies' (Red brush) in a real image, b) Unusually small LV sizes vs. real counterparts with same covariates. Colored LVs are predicted mask from our segmentation model

These discrepancies may stem from training with the extensive UKB dataset and fine-tuning on the smaller ADNI dataset. A potential direction is integrating UKB's demographic diversity with ADNI's AD-specific features, potentially improving the synthetic representation of underrepresented conditions. Additionally, advanced techniques could balance the influence of each data source, ensuring both cohorts' unique characteristics are captured in generated images. Moreover, while causal models account for ROI volumes, the generative model's 2D training might restrict performance because of reduced contextual and spatial information. This limitation highlights the necessity for additional research as we evolve our 2D proof of concept to a more clinically relevant 3D model in future work.

5. Conclusion and Future Work

In conclusion, our study highlights the difficulties in differentiating synthetic from real brain MR images, emphasizing the limitations of the VTT and advocating for more objective evaluation methods. Our analysis reveals the generative model's strength in replicating age-related anatomical changes and simulating neurodegenerative features, alongside a tendency to introduce unrealistic biases, such as overestimated ventricular enlargement or implausibly small ventricles. These insights underline the necessity for robust evaluation methods for synthetic medical images, aiming to enhance image generation and facilitate their successful application in clinical settings.

While our evaluation primarily focuses on neurodegeneration, its relevance extends across various medical domains. For example, adapting our segmentation model to tumor analysis in different organs necessitates retraining for tumor segmentation. Our statistical pipeline would then be used to compare tumor shapes and sizes with those of actual tumors at various stages. In brain MRI studies, examining mid-line shifts and the progression effects of tumors on adjacent ROIs in both real and synthetic images offers a unique opportunity to discern consistencies and discrepancies between image types. Developing these evaluation methods, though challenging, is essential for the effective integration of synthetic images into clinical practice and enhancing the transparency of deep learning in such a critical field.

Acknowledgments

Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U19 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.

This research has been conducted using the UK Biobank Resource under Application ID Number 77717.

We would like to thank ImFusion GmbH for providing the software used to conduct the visual Turing test free of charge.

References

ImFusion - ImFusion labels. https://www.imfusion.com/products/imfusion-labels. Accessed: 2023-09-06.

Alzheimer's Disease Neuroimaging Initiative (ADNI). Adni dataset [data set]. https://adni.loni.usc.edu, 2021. Data spans ADNI1, ADNI-GO, ADNI2, and ADNI3 phases, reflecting comprehensive Alzheimer's disease progression research. Coordinated by the Alzheimer's Therapeutic Research Institute, University of Southern California, with extensive funding from NIH (Grant U19 AG024904) and DOD (award W81XWH-12-2-0012), among others.

Benjamin Billot, Douglas N. Greve, Oula Puonti, Axel Thielscher, Koen Van Leemput, Bruce Fischl, Adrian V. Dalca, and Juan Eugenio Iglesias. Synthseg: Segmentation of brain MRI scans of any contrast and resolution without retraining. *Medical Image Analysis*, 86:102789, 2023. ISSN 1361-8415. doi: 10.1016/j.media.2023.102789.

Ankan Dash, Junyi Ye, and Guiling Wang. A review of generative adversarial networks (gans) and its applications in a wide variety of disciplines: From medical to remote sensing. *IEEE Access*, 2023.

Oliver Diaz, Kaisar Kushibar, Richard Osuala, Akis Linardos, Lidia Garrucho, Laura Igual, Petia Radeva, Fred Prior, Polyxeni Gkontra, and Karim Lekadir. Data preparation for artificial intelligence in medical imaging: A comprehensive guide to open-access platforms and tools. *Physica medica*, 83:25–37, 2021.

Jimit Doshi, Guray Erus, Yangming Ou, Bilwaj Gaonkar, and Christos Davatzikos. Multi-atlas skull-stripping. *Academic radiology*, 20(12):1566–1576, 2013.

Bruce Fischl. Freesurfer. *Neuroimage*, 62(2):774–781, 2012.

GB Frisoni, NC Fox, and R Clifford Jr. Jack cr jr, scheltens p, thompson pm. the clinical use of structural mri in ad. *Nat Rev Neurol*, 6(2):67–77, 2010.

Jingru Fu, Antonios Tzortzakakis, José Barroso, Eric Westman, Daniel Ferreira, Rodrigo Moreno, and Alzheimer's Disease Neuroimaging Initiative. Fast three-dimensional image generation for healthy brain aging using diffeomorphic registration. Technical report, Wiley Online Library, 2023.

Mark Jenkinson, Christian F Beckmann, and TE Behrens. Woolrich mw, and smith sm. *Fsl. Neuroimage*, 62(2):782–790, 2012.

Euijin Jung, Miguel Luna, and Sang Hyun Park. Conditional gan with 3d discriminator for mri generation of alzheimer's disease progression. *Pattern Recognition*, 133:109061, 2023.

Georgios A Kaissis, Marcus R Makowski, Daniel Rückert, and Rickmer F Braren. Secure, privacy-preserving and federated machine learning in medical imaging. *Nature Machine Intelligence*, 2(6):305–311, 2020.

Tero Karras, Miika Aittala, Samuli Laine, Erik Härkönen, Janne Hellsten, Jaakko Lehti-nen, and Timo Aila. Alias-free generative adversarial networks. *Advances in Neural Information Processing Systems*, 34:852–863, 2021.

Sreevani Katabathula, Qinyong Wang, and Rong Xu. Predict alzheimer's disease using hippocampus mri data: a lightweight 3d deep convolutional network model with visual and global shape representations. *Alzheimer's Research & Therapy*, 13(1):1–9, 2021.

Firas Khader, Gustav Mueller-Franzes, Soroosh Tayebi Arasteh, Tianyu Han, Christoph Haarburger, Maximilian Schulze-Hagen, Philipp Schad, Sandy Engelhardt, Bettina Baessler, Sebastian Foersch, et al. Medical diffusion: Denoising diffusion probabilistic models for 3d medical image generation. *arXiv preprint arXiv:2211.03364*, 2022.

Murat Kocaoglu, Christopher Snyder, Alexandros G Dimakis, and Sriram Vishwanath. Causalgan: Learning causal implicit generative models with adversarial training. *arXiv preprint arXiv:1709.02023*, 2017.

Dong Kyu Lee. Alternatives to p value: confidence interval and effect size. *Korean journal of anesthesiology*, 69(6):555–562, 2016.

Geert Litjens, Thijs Kooi, Babak Ehteshami Bejnordi, Arnaud Arindra Adiyoso Setio, Francesco Ciompi, Mohsen Ghafoorian, Jeroen Awm Van Der Laak, Bram Van Ginneken, and Clara I Sánchez. A survey on deep learning in medical image analysis. *Medical image analysis*, 42:60–88, 2017.

Jonathan Long, Evan Shelhamer, and Trevor Darrell. Fully convolutional networks for semantic segmentation. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pages 3431–3440, 2015.

Mehdi Mirza and Simon Osindero. Conditional generative adversarial nets. *arXiv preprint arXiv:1411.1784*, 2014.

Takeru Miyato and Masanori Koyama. cgans with projection discriminator. *arXiv preprint arXiv:1802.05637*, 2018.

Vasileios Nittas, Paola Daniore, Constantin Landers, Felix Gille, Julia Amann, Shannon Hubbs, Milo Alan Puhan, Effy Vayena, and Alessandro Blasimme. Beyond high hopes: A scoping review of the 2019–2021 scientific discourse on machine learning in medical imaging. *PLOS Digital Health*, 2(1):e0000189, 2023.

Nick Pawlowski, Daniel Coelho de Castro, and Ben Glocker. Deep structural causal models for tractable counterfactual inference. *Advances in Neural Information Processing Systems*, 33:857–869, 2020.

Liying Peng, Yen-wei Chen, Zhanhao Mo, Roza M Vlasova, Sun Hyung Kim, Alan C Evans, Annette M Estes, Robert C McKinstry, Kelly N Botteron, Guido Gerig, et al. Longitudinal prediction of infant mr images with multi-contrast perceptual adversarial learning. *Frontiers in neuroscience*, 15:653213, 2021.

Daniele Ravi, Stefano B Blumberg, Silvia Ingala, Frederik Barkhof, Daniel C Alexander, Neil P Oxtoby, Alzheimer's Disease Neuroimaging Initiative, et al. Degenerative adversarial neuroimage nets for brain scan simulations: Application in ageing and dementia. *Medical Image Analysis*, 75:102257, 2022.

Fabio De Sousa Ribeiro, Tian Xia, Miguel Monteiro, Nick Pawlowski, and Ben Glocker. High fidelity image counterfactuals with probabilistic causal models. 2023.

Olaf Ronneberger, Philipp Fischer, and Thomas Brox. U-net: Convolutional networks for biomedical image segmentation. In *Medical Image Computing and Computer-Assisted Intervention–MICCAI 2015: 18th International Conference, Munich, Germany, October 5-9, 2015, Proceedings, Part III 18*, pages 234–241. Springer, 2015.

Iqbal H Sarker. Deep learning: a comprehensive overview on techniques, taxonomy, applications and research directions. *SN Computer Science*, 2(6):420, 2021.

Dinggang Shen, Guorong Wu, and Heung-Il Suk. Deep learning in medical image analysis. *Annual review of biomedical engineering*, 19:221–248, 2017.

Cathie Sudlow, John Gallacher, Naomi Allen, Valerie Beral, Paul Burton, John Danesh, Paul Downey, Paul Elliott, Jane Green, Martin Landray, et al. Uk biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS medicine*, 12(3):e1001779, 2015.

Matthias S Treder, Ryan Codrai, and Kamen A Tsvetanov. Quality assessment of anatomical mri images from generative adversarial networks: Human assessment and image quality metrics. *Journal of Neuroscience Methods*, 374:109579, 2022.

Jiahui Wang, Clement Vachet, Ashley Rumple, Sylvain Gouttard, Clémentine Ouziel, Emilie Perrot, Guangwei Du, Xuemei Huang, Guido Gerig, and Martin Styner. Multi-atlas segmentation of subcortical brain structures via the autoseg software pipeline. *Frontiers in neuroinformatics*, 8:7, 2014.

Matthias Wilms, Jordan J Bannister, Pauline Mouches, M Ethan MacDonald, Deepthi Rajashekar, Sönke Langner, and Nils D Forkert. Invertible modeling of bidirectional relationships in neuroimaging with normalizing flows: application to brain aging. *IEEE Transactions on Medical Imaging*, 41(9):2331–2347, 2022.

Tian Xia, Agisilaos Chartsias, Chengjia Wang, Sotirios A Tsaftaris, Alzheimer's Disease Neuroimaging Initiative, et al. Learning to synthesise the ageing brain without longitudinal data. *Medical Image Analysis*, 73:102169, 2021.

Ling Yang, Zhilong Zhang, Yang Song, Shenda Hong, Runsheng Xu, Yue Zhao, Wentao Zhang, Bin Cui, and Ming-Hsuan Yang. Diffusion models: A comprehensive survey of methods and applications. *ACM Computing Surveys*, 56(4):1–39, 2023.

Kelly H Zou, Simon K Warfield, Aditya Bharatha, Clare MC Tempany, Michael R Kaus, Steven J Haker, William M Wells III, Ferenc A Jolesz, and Ron Kikinis. Statistical validation of image segmentation quality based on a spatial overlap index1: scientific reports. *Academic radiology*, 11(2):178–189, 2004.

Funding

This research is supported through the German Federal Ministry of Education and Research (BMBF) within the project 'Syreal' (Grant No. 01/S21069A). The funders had no role in the study design, data analysis, interpretation, or report writing.

Data Availability

The ADNI dataset analyzed during the current study is available at the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (https://ida.loni.usc.edu) upon consenting to the data sharing agreement. Access to the UKB dataset used during the training of the segmentation models requires an application and is subject to approval. Details for accessing the UKB data can be found on their official website. The synthetic data used in the analysis is publically available at figshare. The Python code implementing the segmentation models and statistical analysis is accessible at github.com/hsyassin/BioPlausibleSynthImgEval.git.

Appendix A. Synthetic Data

Here we provide more details on the conditional causal model, which generates the latent variables for the generative model. In this work, the synthetic images are generated from a conditional StyleGAN3 model with a handcrafted causal model, which controls the latent variables, *e.g.* age, sex, and lateral ventricle volumes. Figure 4 depicts the causal modeling of the conditional variables. Age and sex are the confounders of the CDR and volumes. Furthermore, we use Maximum Likelihood Estimation to fit the training data into a handcrafted parametric model for all latent variables. Age is modeled by a beta distr, sex is a Bernoulli distr, while CDR is sampled as a multi-class softmax regression, conditioned on age and sex. There are three classes in CDR, "0" means a healthy subject, "0.5" means a cognitively impaired subject, and "1" means a dementia patient. Lastly, we use a Gaussian mixture regression model, conditioned on age, sex, and CDR, to model the volumes.

Therefore, with the trained causal model, we can randomly sample latent variables from the distr of the training set (Pawlowski et al., 2020). Furthermore, we can also sample specific variables, *e.g.* CDR and volumes, by conditioning on a specific age. After sampling the latent variables, we fit them into the conditional StyleGAN3 to generate images with specific characteristics.

Figure 4: DAG (Directed acyclic graph) of the causal model. A: Age, S: Sex, C: CDR, V: Ventricles (lateral ventricle volumes, cerebral cortex volumes, and hippocampus volumes.)

Appendix B. Visual Turing Test (VTT)

Figure 5 shows the accuracy of two radiologists during the VTT, alongside the reasons cited for identifying synthetic images. Upon further analysis of the results, insights into the certainty levels associated with the radiologists' decisions were gained. R1 often indicated 'moderate certainty' in their choices, whereas R2 was more inclined towards 'very low certainty.' This variation in certainty directly correlates with their accuracy rates. Nonetheless, the task's complexity—from evaluating a single mid-brain slice (256x256) instead of a full 3D volume, which is atypical in clinical practice—posed significant challenges. In a subsequent meeting, the radiologists concurred that a repeat of the test, even after practicing, would unlikely improve their accuracy, highlighting the fundamental challenge of the task.

Figure 5: Bar plots of Radiologists' Accuracy values in VTT and Reasons for Correctly Identifying Synthetic Images.

Figure 6: R1 misclassification of a real image on the left as synthetic (actually on the right), stating "Noise Patterns" as the reason, and selecting "Other" (Green label brush) with no additional comments. The covariates for both images are: Male, 66 years old, and CN.

Additionally, we provide a breakdown of the reasons each radiologist cited for their choices, particularly when they correctly identified a synthetic image (see Figure 5 b)). R1 frequently attributed their decisions to 'contrast problems,' whereas R2, biased towards associating smooth images with real ones despite this characteristic being present in both image types, leaned toward 'anatomical inaccuracies.' as their rationale. Interestingly, neither radiologist selected "Inaccurate representation of pathology" as the reason for their choice. Lastly, Figures 3 and 6 demonstrate an example where both R1 and R2 misclassified a real image as synthetic for two different reasons ('anatomical inaccuracies,' 'Noise Patterns,' respectively).

Appendix C. Segmentation Training and Evaluation

This appendix complements the main paper with detailed implementation, data-related specifics, and performance comparisons, crucial for reproducibility yet placed here to keep the main text focused.

C.1. Model Architecture

We aimed to precisely segment ROI from brain MR images using state-of-the-art deep learning models, laying a solid foundation for evaluating age-related biological changes. We employed two principal architectures and their variations: **UNet** (Ronneberger et al., 2015) (both original and adapted versions) and **FCN with ResNet backbones (FCN-Res)** (Long et al., 2015) (incorporating ResNet-50 and ResNet-101 variants). For the A-UNet, we increased the initial feature count to 64 from 32 and added an extra layer of depth, significantly boosting its segmentation efficiency for our dataset.

C.2. Model Implementations

For consistency, we used standard optimizers and loss functions in A-UNet (lr = 0.0001, Adam optimizer with betas (0.9, 0.999), epsilon of 1e-8, dice loss) and FCN-Res models (lr = 0.01, SGD with momentum of 0.9, weight decay of 1e-6, cross-entropy and Dice loss with auxiliary loss). We applied max normalization and augmentations (random rotations, flips). Customizing models for the high-resolution UKB dataset, we chose a batch size of 4 with gradient accumulation. Validation was conducted using a 5-fold cross-validation, dividing the UKB dataset into 60% training, 15% validation, and 25% testing. The training was performed exclusively on UKB to avoid bias towards real ADNI images selected as a reference for our evaluation approach.

We implemented our segmentation models on PyTorch Lightning and trained our models on 1 A100 GPU until the convergence of DSC score on a validation set. For consistency with synthetic images, we adapted 2D StylGAN3 pre-processing, registering MR scans and ground truth masks to the MNI152 atlas space, and resized images to 256X256, focusing on the central coronal slice for training and evaluation.

C.3. Models' Performance and Architectural Comparison

The DSC served as the primary metric to evaluate segmentation accuracy (Zou et al., 2004), reflecting the overlap between model predictions and SynthSeg ground truth masks. Initial optimization attempts started with models showing a DSC around 0.6, indicative of underfitting. Through iterative enhancements, the A-UNet model emerged as the top performer on internal testing on UKB and external testing on ADNI. The best-performing models were based on external testing, where A-UNet achieved the highest median DSC scores of 96.53% ± 0.16 for ventricles and 91.68% ± 0.28 for the hippocampus across 3 out of 5 folds (Table 2).

Comprehensive comparisons in Tables 2, 3 and 4 detail DSC scores across models and validation folds, illustrating A-UNet's superior performance and its selection for subsequent analyses.

Table 2: Comparison of model performance using **median DSC $\pm$ interquartile range of lateral ventricles and hippocampus ROIs** over 3 out of 5 folds cross-validation for the 2 best performing models, highlighting the highest obtained DSC values between both

Model	Lateral Ventricles (DSC)		Hippocampus (DSC)	
	Internal	External	Internal	External
A-UNet	**97.48% $\pm$ 0.04**	**96.53% $\pm$ 0.16**	**94.37% $\pm$ 0.04**	**91.68% $\pm$ 0.28**
FCN-Res101	94.20% $\pm$ 0.01	93.34% $\pm$ 0.42	91.62% $\pm$ 0.05	88.33% $\pm$ 0.27

Table 3: Comparison of model performance using **mean DSC of lateral ventricles and hippocampus ROIs** over the first fold in cross-validation for all models

Model	Lateral Ventricles		Hippocampus	
	Internal	External	Internal	External
A-UNet	**97.52%**	**96.68%**	**94.37%**	**91.61%**
UNet	96.16%	94.65%	91.60%	88.09%
FCN-Res101	94.21%	93.71%	91.45%	86.96%
FCN-Res50	94.20%	93.55%	91.13%	84.79%

Table 4: Comparison of model performance using **median DSC of lateral ventricles and hippocampus ROIs** over the first fold in cross-validation for models trained on Intracranial ROI

Model	Intracranial Areaa (DSC)	
	Internal	External
A-UNet	**99.37%**	**98.57%**
FCN-Res50	99.10%	98.08%

C.4. Outlier Analysis

In our final analysis phase, we investigated extreme outliers within the lateral ventricles and hippocampus regions to detect potential segmentation inaccuracies. Notably, errors were confined to a handful of cases in both real and synthetic images, mainly resulting from the lack of typical T1-weighted contrast inherent to each image type. Furthermore, we encountered two empty synthetic images. Nonetheless, excluding these outliers from our

dataset did not influence the overall significance of our results. It appears that the observed irregular contrast patterns can be traced back to anomalies in the real dataset, which were then mirrored in some synthetic images, potentially arising from errors in scanning protocols. These discrepancies introduce segmentation challenges by creating contrast variations distinct from those in the training set. For visual examples of these anomalies and the segmentation issues they induced, please see Figure 7.

Image	Lateral Ventricles	Hippocampus	Intracranial

Figure 7: The first row displays successful segmentation of three ROI masks in normal T1-weighted contrast used for training. In contrast, the second and third rows highlight segmentation challenges in contrasts different from training. The first column shows the image, followed by segmentation masks for lateral ventricles, hippocampus, and intracranial area. The first and second rows feature real images, while the third row shows a synthetic image.

Appendix D. Statistical Tests' Results

This section provided the detailed results of our statistical analyses depicted in Figures 8, 9 and 10, Tables 1, 5 and 6

Figure 8: Comparison of ROI areas in real vs. synthetic MR brain images across different age distributions: Matched distributions are centered, showcasing a median age of 74. On the sides, mismatched distributions where synthetic images have a higher or lower median age (± 3 and 5 years).

Table 5: **Statistical Comparison of ROI Areas in Real vs. Synthetic Images Across different CDR Distributions:** Analysis examines ROI areas under **Matched Conditions** (covariates align, MCI dominant) and Shifted Conditions (Synthetic images either are CN or AD dominant). **Significant differences** are highlighted.

ROI	Cond.	Permute	ES	CI_l	CI_u	KS
Lateral Ventricles	CN	2.00E-01	-4.80E-02	-1.30E-01	2.90E-02	**3.8e-05**
Lateral Ventricles	Match	**2.0e-03**	-1.30E-01	**-2.00E-01**	**-5.00E-02**	**2.0e-08**
Lateral Ventricles	AD	**0.0e+00**	-1.90E-01	**-2.70E-01**	**-1.10E-01**	**2.0e-12**
Hippocampus	CN	7.00E-01	1.50E-02	-6.20E-02	9.20E-02	2.20E-01
Hippocampus	Match	5.70E-02	7.00E-02	-6.70E-03	1.50E-01	8.70E-02
Hippocampus	AD	**0.0e+00**	2.10E-01	**1.30E-01**	**2.80E-01**	**7.5e-08**

Figure 9: Comparison of ROI areas in real vs. synthetic MR brain images across different age distributions: Matched distributions are centered, showcasing a median age of 74. On the sides, mismatched distributions where synthetic images have a higher or lower median age ($\pm$ 3 and 5 years).

Table 6: Statistical Comparison of ROI Areas in Real vs. Synthetic Images across different sex distributions: Analysis examines ROI areas under **Matched Conditions** (Approx. balanced Female to male ratio) and Shifted Conditions (Synthetic images either female or male dominant at 40% and 60%). **Significant differences** are highlighted.

ROI	Cond.	Permute	ES	CI_l	CI_u	KS
Lateral Ventricles	Females	2.10E-01	-5.00E-02	-1.30E-01	2.70E-02	**1.7e-04**
Lateral Ventricles	Match	**2.0e-03**	-1.30E-01	**-2.00E-01**	**-5.00E-02**	**2.0e-08**
Lateral Ventricles	Males	**1.2e-02**	-9.70E-02	**-1.70E-01**	**-2.00E-02**	**1.2e-07**
Hippocampus	Females	3.70E-01	-3.60E-02	-1.10E-01	4.10E-02	2.70E-01
Hippocampus	Match	5.70E-02	7.00E-02	-6.70E-03	1.50E-01	8.70E-02
Hippocampus	Males	2.10E-01	4.70E-02	-3.00E-02	1.20E-01	1.30E-01

D.1. Results

The observations from Table 5 and Table 6 provide pivotal insights into how synthetic and real data distrs compare, especially regarding the lateral ventricles. Notably, distrs predominantly featuring Cognitive Normal (CN) individuals more closely resemble real data than

Figure 10: Comparison of ROI areas in real vs. synthetic MR brain images across different age distributions: Matched distributions are centered, showcasing a median age of 74. On the sides, mismatched distributions where synthetic images have a higher or lower median age (± 3 and 5 years).

matched distr (MCI dominant). Similarly, distrs, where female subjects are predominant, show a greater resemblance to real data than distrs with a balanced sex ratio (matched distr). These results suggest that certain demographic characteristics significantly influence the model's ability to generate synthetic images that accurately reflect the diversity of real anatomical structures.

D.2. Discussion

The observed results underscore the significance of incorporating covariates in the training and generating synthetic medical images. Our approach, utilizing a causal model that accounts for these covariates, aims to address this need. The increased resemblance of CN-dominated distrs to real data suggests that our synthetic models may be more adept at capturing the anatomical nuances of cognitively normal subjects, possibly reflecting the varied anatomical characteristics associated with neurodegenerative conditions like MCI and AD. The closer alignment of female-dominated distrs with real data further prompts a review of the model's sensitivity to sex-specific anatomical distinctions, an aspect vital for applications demanding accurate representation of physiological variances.

D.3. Future Work

Future research should explore the underlying mechanisms contributing to these observed discrepancies in synthetic image generation. It is essential to investigate the model's training

data and algorithms for potential biases or limitations in capturing the full spectrum of human anatomical diversity. Additionally, expanding the model's training dataset to include a more diverse representation of ages, cognitive states, and sexes may help generate synthetic images that more accurately mirror the variability found in real-world data.