How Does Adaptive Optimization Impact Local Neural Network Geometry?

Abstract

Adaptive optimization methods are well known to achieve superior convergence relative to vanilla gradient methods. The traditional viewpoint in optimization, particularly in convex optimization, explains this improved performance by arguing that, unlike vanilla gradient schemes, adaptive algorithms mimic the behavior of a second-order method by adapting to the *global* geometry of the loss function. We argue that in the context of neural network optimization, this traditional viewpoint is insufficient. Instead, we advocate for a *local* trajectory analysis. For iterate trajectories produced by running a generic optimization algorithm OPT, we introduce $R_{\mathrm{med}}^{\mathrm{OPT}}$, a statistic that is analogous to the condition number of the loss Hessian evaluated at the iterates. Through extensive experiments, we show that adaptive methods such as Adam bias the trajectories towards regions where $R_{\mathrm{med}}^{\mathrm{Adam}}$ is small, where one might expect faster convergence. By contrast, vanilla gradient methods like SGD bias the trajectories towards regions where $R_{\mathrm{med}}^{\mathrm{SGD}}$ is comparatively large. We complement these empirical observations with a theoretical result that provably demonstrates this phenomenon in the simplified setting of a two-layer linear network. We view our findings as evidence for the need of a new explanation of the success of adaptive methods, one that is different than the conventional wisdom.

1 Introduction

The efficient minimization of a parameterized loss function is a core primitive in statistics, optimization and machine learning. Gradient descent (GD), which iteratively updates a parameter vector with a step along the gradient of the loss function evaluated at that vector, is a simple yet canonical algorithm which has been applied to efficiently solve such minimization problems with enormous success. However, in modern machine learning, and especially deep learning, one frequently encounters problems where the loss functions are high dimensional, non-convex and non-smooth. The optimization landscape of such problems is thus extremely challenging, and in these settings gradient descent often suffers from prohibitively high iteration complexity.

To deal with these difficulties and improve optimization efficiency, practitioners in recent years have developed many variants of GD. One prominent class of these GD variants is the family of *adaptive* algorithms (Duchi et al., 2011; Tieleman et al., 2012; Kingma & Ba, 2015). At a high level, adaptive methods scale the gradient with an adpatively selected preconditioning matrix, which is constructed via a moving average of past gradients. These methods are reminiscent of second order gradient descent, since they construct approximations to the Hessian of the loss functions, while remaining computationally feasible since they eschew full computation of the Hessian. A vast line of empirical work has demonstrated the superiority of adaptive methods over GD to optimize deep neural networks, especially on Natural Language Processing (NLP) tasks with transformers (Vaswani et al., 2017; Devlin et al., 2019).

From a theoretical perspective, adaptive methods are well understood in the traditional context of convex optimization. For instance, Duchi et al. (2011) show that when the loss function is convex, then the Adagrad algorithm yields regret guarantees that are provably as good as those obtained by using the best (diagonal) preconditioner in hindsight. The key mechanism that underlies this improved performance, is that the loss function has some global geometric property (such as sparsity or a coordinate wise bounded Lipschitz constant), and the algorithm adapts to this global geometry by adaptively selecting learning rates for features that are more informative.

However, in non-convex optimization, and deep learning in particular, it is highly unclear whether this simple characterization is sufficient to explain the superiority of adaptive methods over GD. Indeed, for large scale neural networks, global guarantees on the geometric properties of the loss are typically vacuous. For instance, for a 20-layer feedforward neural network, if we scale up the weights in each layer by a factor of 1.5, then the global Lipschitz constant of the network is scaled up by a factor of at least e^{10}. Hence it only makes sense to study convergence by looking at the local geometry of the loss along the trajectory of the optimization algorithm (Arora et al., 2018).

Figure 1: **(left)** Training losses of SGD+M starting from x_{SGD} and x_{Adam}. **(right)** The 10th largest value over median in the diagonal of loss Hessian (which can be viewed as a variant of $R_{\text{med}}^{\text{OPT}}(t)$ defined in eq. (1)) for Adam and SGD+M. Since the full Hessian is too big, here we selected several layers and randomly sampled 200 coordinates per layer to compute.

Figure 2: Training losses of Adam and SGD+M on the sentence classification task described in Section 4.1.

Moreover, the interaction between an optimization algorithm and neural network geometry is highly complex — recent work has shown that geometric characteristics of iterates encountered during optimization is highly dependent on the choice of optimization algorithm and associated hyperparameters (Lewkowycz et al., 2020; Cohen et al., 2021). For instance, Cohen et al. (2021) demonstrate that while training neural networks with GD, the maximum eigenvalue of the Hessian evaluated at the GD iterates first increases and then plateaus at a level 2/(step size). The viewpoint from convex optimization, where a loss function has some (potentially) non-uniform but fixed underlying geometry that we must adapt to, is thus insufficient for neural networks, since the choice of optimization algorithm can actually *interact* with and *influence* the observed geometry **significantly**.

To provide another example of this interactive phenomenon, we consider the following experiment. On the same network training loss function f, we run stochastic gradient descent with momentum (SGD+M) and Adam to obtain two different trajectories. We select an iterate x_{Adam} from the Adam trajectory and an iterate x_{SGD} from the SGD trajectory, such that $f(x_{\text{Adam}}) = f(x_{\text{SGD}})$. We then run SGD+M twice, once from x_{Adam} and once from x_{SGD}. If the underlying geometry of the loss function f was truly fixed, then we would not expect a significant difference in the performance of running SGD+M from either of the two iterates. However, as shown in Figure 1(left), running SGD+M from x_{Adam} achieves lower loss than that from x_{SGD}, suggesting that Adam may bias the trajectory towards a region which is more favorable for rapid training. This motivates the following question.

How does adaptive optimization impact the observed geometry of a neural network loss function, relative to SGD (with momentum)?

The remainder of this paper is dedicated to answering the above question. To this end, for each iterate in a trajectory produced by running an optimization algorithm OPT, where the Hessian of the tth iterate is given by $H^{(t)} \in \mathbb{R}^{d \times d}$, we define the second order statistic $R_{\text{med}}^{\text{OPT}}(t)$ in the following fashion. For the tth iterate in the trajectory, let $R_{\text{med}}^{\text{OPT}}(t)$ be the ratio of *maximum* of the absolute entries of the diagonal of $H^{(t)}$, to the *median* of the absolute entries of the diagonal of $H^{(t)}$. Concretely, we define

$$R_{\text{med}}^{\text{OPT}}(t) = \frac{\max\{|H_{ii}^{(t)}|\}_{i=1}^d}{\text{median } \{|H_{ii}^{(t)}|\}_{i=1}^d}. \tag{1}$$

This statistic thus measures the *uniformity* of the diagonal of Hessian, where a smaller value of $R_{\text{med}}^{\text{OPT}}(t)$ implies that the Hessian has a more uniform diagonal. It can also be viewed as a stable[1] variant of the

---

[1]Consider the case where one parameter has little impact on the loss, then the second derivative w.r.t. this parameter is almost zero, making $\frac{\max\{|H_{ii}^{(t)}|\}_{i=1}^d}{\min\{|H_{ii}^{(t)}|\}_{i=1}^d}$ infinity. So we consider *median* which is more stable.

condition number. Instead of eigenvalues, we choose diagonal entries because adaptive methods used in practice are *coordinate-wise*, which can be viewed as the diagonal scaling approaches.[2] Hence we believe the diagonal of Hessian is more relevant than the spectrum. As a supplementary result, in Appendix E, we demonstrate that the loss Hessian approaches diagonal during training for Adam and SGD+M. There has been prior theoretical work on overparameterized neural networks showing that a smaller condition number of Hessian, Neural Tangent Kernel (Jacot et al., 2018) etc. could yield to faster convergence rate for (S)GD (Liu et al., 2022). As for (diagonal) adaptive methods (*e.g.* Adagrad), they were original designed to adapt to the nonuniform diagonal geometry. Intuitively, a smaller $R_{\mathrm{med}}^{\mathrm{OPT}}(t)$, which implies more uniform diagonal geometry, could lead to faster convergence.

Armed with this statistic, we make the following contributions:

- On a wide variety of neural network transformer architectures and language modeling datasets, we conduct experiments to compare how $R_{\mathrm{med}}^{\mathrm{Adam}}(t)$ and $R_{\mathrm{med}}^{\mathrm{SGDM}}(t)$ evolve over time, when Adam and SGD+M are run from the same initialization and with their optimal (initial) learning rates respectively. In each case, we demonstrate that the Adam trajectory attains $R_{\mathrm{med}}^{\mathrm{Adam}}(t)$ values that are significantly smaller than the $R_{\mathrm{med}}^{\mathrm{SGDM}}(t)$ values found by SGD+M. We show a simple example of this phenomenon in Figure 1(right). This suggests that relative to SGD+M, Adam biases the optimization trajectory to a region where the Hessian diagonal is more uniform. We call this phenomenon *the uniformity of diagonal geometry* for adaptive methods. As an aside, we observe that larger improvements in optimization performance of Adam over SGD+M are correlated with larger gaps between $R_{\mathrm{med}}^{\mathrm{Adam}}(t)$ and $R_{\mathrm{med}}^{\mathrm{SGDM}}(t)$. This suggests that a region where the Hessian diagonal is more uniform is also a region that is more amenable to rapid optimization.

- We complement our empirical results with a theoretical analysis of this phenomenon in the simplified setting of large batch Adam and SGD+M, on a two-layer linear network with d-dimensional input and hidden layer, and one dimensional output. We show that for a wide range of t, $R_{\mathrm{med}}^{\mathrm{Adam}}(t) = 1 \pm o(1)$ but $R_{\mathrm{med}}^{\mathrm{SGDM}}(t) = \Omega(\log d)$. Our proof reveals that Adam induces the weight matrices to have low rank whose leading singular vectors have certain type of uniformity (see Section 6 for discussion), a fact that we also observe empirically in large scale neural networks, suggesting that this may be a mechanism by which adaptive methods bias trajectories to have uniformity of diagonal geometry.

2 RELATED WORK

Existing analyses of adaptive methods. The vast majority of prior theoretical work on adaptive methods has focused on the blackbox setting (Duchi et al., 2011; Kingma & Ba, 2015; Chen et al., 2020; Reddi et al., 2018; Ward et al., 2020; Défossez et al., 2020; Ene et al., 2021). These works make minimal assumptions about the structure of the loss function, beyond (possibly) some global properties such as convexity or smoothness. These global properties (governed by parameters such as the smoothness parameter) are assumed to hold over the entire domain. Hence this style of analysis is worst case, since the resulting convergence bounds depend on polynomially on these global parameters. However, as we show in Section 3.1, in neural networks these parameters are prohibitively large. This worst case analysis is hence unlikely to explain the success of adaptive methods on neural networks. By contrast, our focus is on analyzing the local trajectory that is induced by running the optimization method.

Existing analyses of (S)GD on neural networks. There is an extensive literature on the analysis of GD/SGD in the non-blackbox setting, *e.g.* overparameterized neural networks, (Du et al., 2018; Ji & Telgarsky, 2020; Allen-Zhu et al., 2019a;b; Arora et al., 2019a; Liu et al., 2022). However, it is unclear how to translate these analyses of GD/SGD, to an analysis that explains the gap between GD/SGD and adaptive methods.

Influence of algorithms on the loss geometry. In many simple convex settings, *e.g.* linear or logistic regression and the Neural Tangent Kernel (Jacot et al., 2018), the loss geometry is usually fixed and not influenced by learning algorithms. However, in neural networks the interaction between algorithms and loss landscapes is more complicated. Lewkowycz et al. (2020) find a so-called catapult

---

[2]Recall that the main theoretical bound in the original Adagrad paper (Duchi et al., 2011) is in terms of the diagonal scaling.

effect of initial learning rate on the training trajectory of SGD and related loss curvature. Cohen et al. (2021) demonstrate that while training neural networks with GD, the maximum eigenvalue of the Hessian evaluated at the GD iterates first increases and then plateaus at a level that is inversely proportional to the step size. However, Cohen et al. (2021) leave open the problem of whether similar interactive phenomena occur in algorithms that are not GD, including adaptive methods.

3 OVERVIEW OF RESULTS AND SETUP

3.1 ISSUES OF PRIOR ANALYSES ON ADAPTIVE METHODS

As is mentioned in Section 2, existing work on adaptive algorithms has mainly focused on black-box analysis assuming some global worst-case parameters. However, these global bounds can be extremely bad in complicated deep learning models, as is discussed in Section 1. To see this, we initialized a transformer model[3] with default initialization in Pytorch but chose a large gain[4], and computed the smoothness parameter (denoted as l) and the condition number (denoted as κ) of loss Hessian on one layer. We observed that setting the gain as a large constant (*e.g.* 800) results in extremely large l and κ ($l \geq 10^7$ and $\kappa \geq 10^{10}$), which makes the convergence rates in prior black-box analysis vacuous.

The failure of global worst-case analysis implies that we need to focus on the local trajectory of algorithms. However, it is unclear that when two optimization algorithms are used, they will have the same geometry in local trajectory. In particular, although in theory, adaptive algorithms can yield to a convergence rate with better dependency on certain local geometry of the function comparing to SGD (with momentum), it could still be the case that the local geometry along the trajectory of adaptive algorithm can be much worse than that of SGD (with momentum).

That motivates us to study the local geometry, especially that obtained by adaptive methods comparing to SGD (with momentum) in the paper. Motivated by the diagonal scaling of Adagrad and Adam for neural network training, we ask the follow **main question** in our paper:

How does the local diagonal geometry (diagonal of the loss Hessian) along the local trajectory of adaptive algorithms compare to that of SGD (with momentum)?

3.2 OVERVIEW OF THE EXPERIMENTS

As is discussed in Section 1, we consider $R_{\mathrm{med}}^{\mathrm{OPT}}(t)$ defined in eq. (1) as a measurement of the uniformity of the diagonal of the loss Hessian. We conduct experiments on different NLP tasks to examine $R_{\mathrm{med}}^{\mathrm{OPT}}(t)$, as in language models, adaptive methods have shown significantly faster convergence than SGD (with momentum). The details of these experiments will be shown in Section 4. To explore potential different patterns of different layers, we do the computation layer by layer. On a wide variety of transformer architectures and language modeling datasets from the same initialization, we observe that:

When we train the neural network using Adam, the uniformity of diagonal geometry, measured by $R_{\mathrm{med}}^{\mathrm{OPT}}(t)$ is smaller than that when we train using SGD+M from the same initialization, except for first several layers.

Table 1 shows a typical example of $R_{\mathrm{med}}^{\mathrm{Adam}}(t)$ compared to $R_{\mathrm{med}}^{\mathrm{SGDM}}(t)$ on a sentence classification task using BERT-small (Turc et al., 2019; Bhargava et al., 2021) (See Section 4.1 for details). We repeated the experiments for 12 times starting from the same initialization. Table 1 shows the averaged $R_{\mathrm{med}}^{\mathrm{Adam}}(t)$ and $R_{\mathrm{med}}^{\mathrm{SGDM}}(t)$ in some randomly selected layers (except for the first several). We also report the averaged $\frac{R_{\mathrm{med}}^{\mathrm{SGDM}}(t)}{R_{\mathrm{med}}^{\mathrm{Adam}}(t)}$ and their standard deviations in the brackets.[5] Figure 2 shows the corresponding training losses of one in these 12 experiments.

---

[3] https://pytorch.org/tutorials/beginner/transformer_tutorial.html

[4] This refers to the gain parameter in some commonly used initialization functions of Pytorch, *e.g.* torch.nn.init.xavier_uniform_().

[5] $R_{\mathrm{med}}^{\mathrm{SGDM}}(t)$ values in Table 1 for most layers are roughly 1.4 to 2 times $R_{\mathrm{med}}^{\mathrm{Adam}}(t)$ in corresponding layers. In practice, it can be considered significant because it might imply 1.4 to 2 times faster convergence.

Table 1: $R_{\text{med}}^{\text{Adam}}(t)$ and $R_{\text{med}}^{\text{SGDM}}(t)$ in some layers, on the sentence classification task (see Section 4.1).

Layer#	Iteration 0		Iteration 750			Iteration 1250		
	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$
9	15.7	15.7	12.76	9.65	1.45 (0.65)	11.43	14.24	0.94 (0.40)
12	22.63	22.63	13.17	7.41	1.92 (0.67)	10.62	9.67	1.33 (0.75)
15	9.35	9.35	80.57	53.52	1.65 (0.65)	100.65	61.80	2.01 (1.00)
17	82.37	82.37	405.02	223.56	1.91 (0.53)	423.28	337.32	1.43 (0.63)
18	31.32	31.32	17.07	13.24	1.43 (0.58)	18.15	15.63	1.21 (0.36)
22	47.13	47.13	233.72	72.67	3.54 (1.21)	158.38	93.13	2.28 (1.18)
24	31.17	31.17	17.52	17.34	1.13 (0.40)	13.51	14.23	1.05 (0.36)

To understand this phenomenon in a more principled point of view, we also provide a formal proof of the statement in a simplified setting: large batch Adam and SGD+M on a 2-layer linear network. Although simple, the choice of 2-layer linear network to understand learning dynamics is common in prior works (*e.g.* (Tian et al., 2021)). Section 3.3 below describes the theoretical setup.

3.3 Setup of the theoretical analysis

Notation Let $[d] = \{1, 2, ..., d\}$. We use $\|\cdot\|_2$ to denote the l_2 norm of a vector, and $\|\cdot\|_F$ to denote the Frobenius norm of a matrix. Let $\langle\cdot, \cdot\rangle$ be the Euclidean inner product between vectors or matrices. Let $\mathcal{N}(\mu, \sigma^2)$ be the one-dimensional Gaussian distribution with mean μ and variance σ^2. For a scalar (vector, matrix) A which evolves over time, we use $A^{(t)}$ to denote its value at time t.

Let there be m data points. The data matrix is $X \in \mathbb{R}^{d_x \times m}$ and the label matrix is $Y \in \mathbb{R}^{d_y \times m}$. We assume that the input dataset is whitened, i.e. $\Lambda_{xx} := \frac{1}{m}XX^T \in \mathbb{R}^{d_x \times d_x}$ is an identity matrix.

The parameters of a 2-layer linear network are given by $W := (W_2, W_1)$. Assume $W_i \in \mathbb{R}^{d_i \times d_{i-1}}$ for $i = 1, 2$. We have $d_2 = d_y, d_0 = d_x$. We consider the square loss $L(W) := \frac{1}{2m}\|W_2W_1X - Y\|_F^2$.

Denote $A := \frac{1}{m}YX^T \in \mathbb{R}^{d_y \times d_x}$. Arora et al. (2019b) show that with whitened dataset,

$$L(W) := \frac{1}{2m}\|W_2W_1X - Y\|_F^2 = \bar{L}(W) + c, \quad \bar{L}(W) := \frac{1}{2}\|W_2W_1 - A\|_F^2. \tag{2}$$

where c does not depend on W. We consider the following model with small Gaussian initialization.

Assumption 1 (Setup). *The input covariance $\Lambda_{xx} := \frac{1}{m}XX^T \in \mathbb{R}^{d_x \times d_x}$ is an identity matrix. The input and hidden layers are both of dimension d, i.e. $d_1 = d_0 = d$. Without loss of generality, we can assume that A is a row vector (i.e. $d_2 = 1$) whose coordinates are positive[6] and $\Theta(1)$ in terms of d.*

Assumption 2 (Gaussian Initialization). $\forall i, j : w_{2i}^{(0)} \sim \mathcal{N}(0, \frac{1}{d^{2\alpha}}), W_1^{(0)}[i, j] \sim \mathcal{N}(0, \frac{1}{d^{4\alpha}})$ *are independently initialized with sufficiently large $\alpha > 0$.*

Denote $\tilde{A}$ and $\tilde{\Lambda}_{xx}$ as the batch versions of A and Λ_{xx}. We make the following large-batch assumption. We emphasize that large batches are commonly used in NLP tasks (*e.g.* (Brown et al., 2020)).

Assumption 3 (Large Batch). *For the randomly selected batches, assume $\mathbb{E}[\tilde{A}] = A$, $\mathbb{E}[\tilde{\Lambda}_{xx}] = \Lambda_{xx}$. $\forall i, j \in [d] : \mathbb{E}\left[(\tilde{A}_i - A_i)^2\right] \leq \sigma^2$, $\mathbb{E}\left[(\tilde{\Lambda}_{xx}[i, j] - \Lambda_{xx}[i, j])^2\right] \leq \sigma^2$, and $\sigma^2 = \mathcal{O}(\frac{1}{poly(d)})$.*

Denote $\tilde{g}^{(t)}$ as the batch gradient at time t. The update rules of SGD+M and Adam are given by

$$\text{SGD+M:} \quad u^{(t+1)} = \beta u^{(t)} + \tilde{g}^{(t)}, \quad W^{(t+1)} = W^{(t)} - \eta u^{(t)},$$

$$\text{Adam:} \quad \eta_t = \eta \cdot \sqrt{1 - \beta_2^{t+1}}/(1 - \beta_1^{t+1}), \quad m^{(t+1)} = \beta_1 m^{(t)} + (1 - \beta_1)\tilde{g}^{(t)}, \tag{3}$$

$$v^{(t+1)} = \beta_2 v^{(t)} + (1 - \beta_2)\tilde{g}^{(t)} \odot \tilde{g}^{(t)}, \quad W^{(t+1)} = W^{(t)} - \eta_t \frac{m^{(t)}}{\sqrt{v^{(t)}} + \xi},$$

where η is the learning rate, β, β_1, β_2 are momentum parameters, and ξ is for numerical stability. All operations on vectors are element-wise. Here and throughout, the notation $f(x) = \mathcal{O}(g(x))$

---

[6]In Assumption 2 we assume Gaussian initialization. Due to the rotational invariance of Gaussian distribution, we can assume that all coordinates of A are positive without loss of generality.

(resp. $f(x) = \Omega(g(x))$, $f(x) = \Theta(g(x))$) means that there exist constants $C_1, C_2 > 0$ such that $f(x) \leq C_2 g(x)$ (resp. $f(x) \geq C_1 g(x)$, $C_1 g(x) \leq f(x) \leq C_2 g(x)$). We will also use the notation with $\sim$, i.e. $\tilde{\mathcal{O}}(\cdot), \tilde{\Omega}(\cdot), \tilde{\Theta}(\cdot)$ to hide factors that are logarithmic in d. In our theoretical analysis, "with high probability", or "w.h.p." for short, means that with probability at least $1 - \frac{1}{\text{poly}(d)}$.

4 THE UNIFORMITY OF DIAGONAL GEOMETRY

As is mentioned in Section 3.2, we computed $R_{\text{med}}^{\text{OPT}}(t)$ defined in eq. (1) on different language models. In this section, we present the results of SGD+M and Adam on different architectures and datasets. In Appendix A, we present the results of other adaptive algorithms.

During training we started from the same initial weights and used the same learning rate schedule (constant or decreasing) for SGD+M and Adam. We tuned and chose the best (initial) learning rate of SGD+M. The (initial) learning rate of Adam was set as a value under which Adam converged faster than SGD+M with its best learning rate. The concrete values will be stated in later parts of this section. We used large batch sizes to make the training procedure stable. When computing Hessian, we also used large batch sizes. Due to the extremely large dimension, we did the computation on some uniformly selected coordinates, more precisely, 200 coordinates per layer.

4.1 EXPERIMENTS ON REAL DATASETS

Sentence classification task on BERT-small We fine-tuned BERT-small (Turc et al., 2019; Bhargava et al., 2021) on the IMDB dataset (Maas et al., 2011): the task is to classify whether movie reviews are positive or negative.[7] The momentum parameter β in SGD was set as 0.9. The two momentum parameters (β_1, β_2) of Adam were set as (0.9, 0.999). We trained the model using linearly decreasing learning rates for 10 epochs (2500 iterations). The initial learning rates of SGD+M and Adam were 0.001 and 5e-5, respectively. As mentioned in Section 3.2, Figure 2 and Table 1 show the training losses and the comparison between $R_{\text{med}}^{\text{Adam}}(t)$ and $R_{\text{med}}^{\text{SGDM}}(t)$, respectively.

Translation task We trained a Seq2Seq network that uses Transformer to solve a machine translation task on Multi30k (Elliott et al., 2016)(CC BY-NC-SA 4.0): this task is to train a German to English translation model.[8] The momentum parameter β in SGD was set as 0.9. The two momentum parameters (β_1, β_2) of Adam were set as (0.9, 0.98). We trained the model using constant learning rates (0.03 for SGD+M and 1e-4 for Adam) for 60 epochs (1800 iterations). The experiments were repeated for 8 times starting from the same initialization. Figure 3(left) shows the training losses for one among them. Table 2a shows the averaged $R_{\text{med}}^{\text{Adam}}(t)$, $R_{\text{med}}^{\text{SGDM}}(t)$ and $\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$ (with standard deviation in the brackets) in some randomly selected layers.

Figure 3: Training losses of Adam and SGD+M for the translation task on (left) Multi30k (see Section 4.1) and (right) data with randomly generated targets (see Section 4.2).

Figure 4: Singular values and R_u of the weight matrix in the 27-th layer on the translation task (see Section 4.1).

4.2 EXPERIMENTS ON RANDOM DATASETS

We used the same model and momentum parameters as in the translation task described in Section 4.1 but generated random integers as targets. Similar to the setting on real targets, the model was trained

---

[7]https://huggingface.co/docs/transformers/v4.16.2/en/training
[8]https://pytorch.org/tutorials/beginner/translation_transformer.html

Table 2: $R_{\text{med}}^{\text{Adam}}(t)$ and $R_{\text{med}}^{\text{SGDM}}(t)$ in some layers for the translation task. (a) on Multi30k and (b) on data with randomly generated targets.

(a)

Layer#	Epoch 0		Epoch 30			Epoch 55		
	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$
3	4.27	4.27	5.14	2.41	2.16 (0.75)	3.14	2	1.58 (0.41)
5	7.09	7.09	36.11	18.33	2.00 (0.42)	52.12	16.59	3.16 (0.64)
7	5.79	5.79	5.91	3.87	1.55 (0.32)	7.52	3.08	2.45 (0.56)
9	18.11	18.11	28.93	20.74	1.43 (0.28)	36.67	18	2.05 (0.18)
12	11.1	11.1	6.64	7.25	0.95 (0.21)	9.27	5.06	1.88 (0.54)
15	83.15	83.15	52.41	7.5	7.15 (1.63)	46.27	5.69	8.6 (3.06)
18	14.99	14.99	4.19	4.22	1.17 (0.45)	3.09	2.72	1.2 (0.46)
21	93.5	93.5	30.29	5.36	5.72 (1.05)	19.27	4.8	4.09 (0.86)
24	36.63	36.63	6.14	4.66	1.35 (0.31)	5.02	3.2	1.6 (0.36)
28	18.47	18.47	3.07	1.95	1.58 (0.16)	2.9	1.59	1.83 (0.14)

(b)

Layer#	Epoch 0		Epoch 30			Epoch 55		
	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$
3	4.82	4.82	3.98	1.8	2.23 (0.36)	3.79	1.61	2.36 (0.32)
5	8.04	8.04	46.06	45.84	1.01 (0.17)	47.83	34.18	1.41 (0.30)
7	5.69	5.69	44.77	3.92	11.79 (2.37)	46.5	2.74	17.4 (2.99)
9	11.89	11.89	317.34	55.61	5.81 (0.70)	351.85	46.54	7.61 (0.87)
12	19.73	19.73	133.39	3.91	34.17 (4.51)	145.09	2.97	49.49 (13.40)
15	32.12	32.12	462.74	51.53	9.03 (0.91)	492.73	50.57	9.84 (1.03)
18	19.79	19.79	74.6	6.59	11.8 (3.33)	79.02	3.58	22.75 (6.01)
21	26.94	26.94	767.31	48.89	16.4 (3.38)	797.49	36.88	21.98 (3.40)
24	34.72	34.72	467.75	9.15	52.57 (11.16)	602.03	3.51	172.65 (18.85)
28	13.13	13.13	19.8	2.22	8.99 (1.74)	19	1.63	11.7 (1.48)

using constant learning rates (0.015 for SGD+M and 5e-5 for Adam) for 60 epochs (1800 iterations), and we repeated the experiments for 8 times starting from the same initialization. Figure 3(right) shows the training losses for one among them. Table 2b shows the averaged $R_{\text{med}}^{\text{Adam}}(t)$, $R_{\text{med}}^{\text{SGDM}}(t)$ and $\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$ (with standard deviation in the brackets) of the same 10 layers as in Table 2a.[9]

4.3 SUMMARIZATION OF THE EMPIRICAL RESULTS AND DISCUSSION

Overall, through extensive experiments on language models, we demonstrate that **starting from the same initialization, the $R_{\text{med}}^{\text{OPT}}(t)$ values found by Adam are smaller than those found by SGD+M, except for the first several layers.** This suggests that Adam is biased towards a region with more uniform diagonal Hessian than SGD+M.

Positive correlation between uniformity of diagonal Hessian and fast convergence. We observe that on random dataset, SGD+M plateaus after about 400 steps and thus converges much slower when compared to Adam than on real dataset (see Figure 3). On the other hand, the gaps of $R_{\text{med}}^{\text{SGDM}}(t)$ and $R_{\text{med}}^{\text{Adam}}(t)$ are more significant on random data than on real data (see Table 2b) as well. In Appendix A.4, we conduct another experiment where we switch from SGD to Adam in the middle and compare it with the model trained by Adam from the beginning. The observation is that both the loss gap and the gap of $R_{\text{med}}^{\text{OPT}}(t)$ are gradually closed after switching (see Figure 7 and Table 8). Hence we find a positive correlation between fast convergence and the uniformity of diagonal of loss Hessian, suggesting that a region with more uniform diagonal of Hessian is also a region that is more amenable to fast optimization. In Appendix A we **study other adaptive algorithms (Adagrad, RMSprop and AMSGrad) and get similar observation**: all these adaptive methods converge faster than SGD or SGD+M and also bias the trajectory to regions with smaller $R_{\text{med}}^{\text{OPT}}(t)$, suggesting that the uniformity of diagonal Hessian might be a universal mechanism (partially) explaining the faster optimization of adaptive algorithms than SGD (with momentum).

More discussions on the trajectory difference. Considering the fact that our comparison between $R_{\text{med}}^{\text{Adam}}(t)$ and $R_{\text{med}}^{\text{SGDM}}(t)$ is conditioned on the same iteration when SGD+M has larger training loss than Adam, there is a potential alternative explanation of the Hessian diagonal uniformity. That is, the global minimum has uniform Hessian, and Adam simply converges faster to it than SGD+M. To rule out this possibility, in Appendix A.3 we add a comparison of our measurements $R_{\text{med}}^{\text{Adam}}(t)$ and $R_{\text{med}}^{\text{SGDM}}(t')$, where t, t' are picked such that tth Adam iterate and t'th SGD+M iterate have the *same training loss*. The results (in Table 7) show that $R_{\text{med}}^{\text{Adam}}(t) < R_{\text{med}}^{\text{SGDM}}(t')$ for most layers, thus demonstrating that the trajectories of Adam and SGD+M are truly different and that the difference is because Adam biases the local geometry (as opposed to faster convergence).

Adding regularization. People in practice usually add weight decay (equivalent to l_2 regularization) to encourage better generalization ability. In Appendix A.7 we compare SGD+M and Adam when both using small weight decay values (0.001). The results in Figure 13a and Table 9 suggest that in this case, the positive correlation between $R_{\text{med}}^{\text{OPT}}(t)$ and convergence speed still holds: Adam converges faster than SGD+M and in most of the layers except for the first several, $R_{\text{med}}^{\text{Adam}}(t)$ values

---

[9]To prevent $R_{\text{med}}^{\text{OPT}}(t)$ from getting too large due to tiny median, we added an additional term $0.001 \max\{|H_{ii}^{(t)}|\}_{i=1}^d$ to the denominator of eq. (1) when computing.

are smaller than $R_{\text{med}}^{\text{SGDM}}(t)$. This reveals the robustness of our observation under weak regularization. However, under large weight decay parameters, we observed cases where Adam still converged faster but $R_{\text{med}}^{\text{Adam}}(t)$ values were larger rather than smaller. In the case of strong regularization, the adaptivity of Adam requires further exploration and we hope to find new mechanisms in the future.

Image tasks. Although in this paper we focus on language models where Adam shows significant fast convergence, we also add supplementary results in Appendix A.8 on image tasks where SGD+M performs better. On a residual network trained on CIFAR-10, we observed that Adam did not converge faster than SGD+M (see Figure 13b) and in the meantime, $R_{\text{med}}^{\text{Adam}}(t)$ values were no longer smaller than $R_{\text{med}}^{\text{SGDM}}(t)$ during training (see Table 10). This reveals the connection between the local diagonal geometry and the convergence speed from another perspective. That is, when the diagonal of Hessian of Adam is not more uniform than SGD+M, its convergence speed is not better, either.

5 THEORETICAL ANALYSIS

In Section 4, we empirically demonstrate the uniformity of diagonal geometry. In this section, we theoretically analyze this property for large batch Adam and SGD+M on a two-layer linear network with 1-dimensional output.

Since the weights and Hessians in different layers may have different magnitudes, we compute the $R_{\text{med}}^{\text{OPT}}(t)$ layer by layer. We denote $R_{\text{med},k}^{\text{SGDM}}(t)$ (resp. $R_{\text{med},k}^{\text{Adam}}(t)$) as the $R_{\text{med}}^{\text{OPT}}(t)$ found by SGD+M (resp. Adam) w.r.t. W_k at time t where $k = 1, 2$.

Theorem 1. *Under Assumption 1, 2 and 3, consider the weights $\left\{ W_{SGD}^{(t)} \right\}_{t \geq 0}$ (resp. $\left\{ W_{Adam}^{(t)} \right\}_{t \geq 0}$) obtained by SGD+M (resp. Adam) defined in* (3).

1. For any $p > 0$, pick $0 < \epsilon < \frac{1}{d^p}$, $\eta \leq \mathcal{O}\left(\frac{\epsilon}{d^{7\alpha/4+4}} \right)$ and $\alpha \geq 4(p+2)$. Suppose $\sigma \leq \frac{\eta^{3/2}}{d^{\alpha/2+1}}$, then there exists $T_{SGD,1}, T_{SGD,2}$ such that w.h.p., $\bar{L}\left(W_{SGD}^{(T_{SGD,1})} \right) = \Theta(d)$, $\bar{L}\left(W_{SGD}^{(T_{SGD,2})} \right) \leq \tilde{\mathcal{O}}\left(\frac{1}{d^p} \right)$, and

$$\forall t \in [T_{SGD,1}, T_{SGD,2}]: \quad R_{med,k}^{SGDM}(t) = \Omega(\log d), \quad k = 1, 2.$$

2. For any $p > 0$, pick $\eta \leq \mathcal{O}\left(\frac{1}{d^{3\alpha}} \right)$, $\xi \leq \sqrt{\frac{\eta}{d^{3\alpha-1}}}$, $\alpha \geq \frac{p+4}{3}$ and $\beta_2 = \beta_1^2$. Suppose $\sigma \leq \frac{\eta^{3/2}\xi^2}{d^{13/4}}$, Then $\exists T_{Adam,1}, T_{Adam,2}$ such that w.h.p., $\bar{L}\left(W_{Adam}^{(T_{Adam,1})} \right) = \Theta(d)$, $\bar{L}\left(W_{Adam}^{(T_{Adam,2})} \right) \leq \tilde{\mathcal{O}}\left(\frac{1}{d^p} \right)$, and

$$\forall t \in [T_{Adam,1}, T_{Adam,2}]: \quad R_{med,k}^{Adam}(t) = 1 \pm \tilde{\mathcal{O}}\left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}} \right), \quad k = 1, 2.$$

An immediate corollary of this theorem below gives the difference between iterates of Adam and SGD+M that have the same loss.

Corollary 1. *Under the setup in Theorem 1, w.h.p., for any $t \in [T_{SGD,1}, T_{SGD,2}]$ and $t' \in [T_{Adam,1}, T_{Adam,2}]$ such that $\bar{L}\left(W_{SGD}^{(t)} \right) = \bar{L}\left(W_{Adam}^{(t')} \right) \in \left[\tilde{\Omega}\left(\frac{1}{d^p} \right), \Theta(d) \right]$, we have*

$$R_{med,k}^{SGDM}(t) = \Omega(\log d), \quad R_{med,k}^{Adam}(t') = 1 \pm \tilde{\mathcal{O}}\left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}} \right), \quad k = 1, 2.$$

Theorem 1 and Corollary 1 tell us that during a long training period when the loss decreases from $\Theta(d)$ to $\tilde{\mathcal{O}}\left(\frac{1}{d^p} \right)$, the diagonal of loss Hessian for Adam keeps nice uniformity in the sense that for each layer, its diagonal elements have roughly the same value, i.e. $R_{\text{med},k}^{\text{Adam}}(t) = 1 \pm o(1), k = 1, 2$. On the other hand, the diagonal of loss Hessian for SGD+M is less uniform. Appendix B gives a proof sketch of Theorem 1. The detailed proof can be found in Appendix C and D.

6 THE LOW RANK STRUCTURE OF WEIGHT MATRICES AND UNIFORMITY OF LEADING SINGULAR VECTORS

The proof sketch in Appendix B highlights one crucial intuition of Theorem 1: After $T_{\text{SGD},1}$ (resp. $T_{\text{Adam},1}$) steps, W_1 of SGD+M (resp. Adam) becomes an approximately rank-1 matrix. Consider

the left singular vector $\boldsymbol{u} := [u_1, u_2, ..., u_d]^T$ which corresponds to the leading singular value σ_1. We can show that the distribution of $u_1^2, u_2^2, ..., u_d^2$ for Adam is more uniform than that of SGD+M. This property, we call *the uniformity of the leading singular vector*, is related to the uniformity of the diagonal of loss Hessian, see Appendix F for more details.

Similar low rank bias after training has been studied in prior works (*e.g.* (Gunasekar et al., 2017; Li et al., 2018; Chou et al., 2020)). For more complicated models, we want to check whether the weight matrices also have low rank structures and if so, whether we can still observe *the uniformity of leading singular vectors*. More formally, consider the weight matrix in some layer $W \in \mathbb{R}^{m \times n}$, we want to check

(A) Whether $W \in \mathbb{R}^{m \times n}$ is approximately a rank k matrix with $k \ll \min\{m, n\}$, and

(B) If (A) is true, then consider the top k singular values $\sigma_1, ..., \sigma_k$ and corresponding left singular vectors $\boldsymbol{u}_1, \boldsymbol{u}_2, ... \boldsymbol{u}_k$. Define a new vector $\tilde{\boldsymbol{u}} := \sum_{i=1}^{k} \sigma_i^2 \boldsymbol{u}_i \odot \boldsymbol{u}_i := [\tilde{u}_1, \tilde{u}_2, ..., \tilde{u}_d]^T$ and compute $R_u := \frac{\max_i \tilde{u}_i}{\text{median } \tilde{u}_i}$, which is a generalized version of $\frac{\max_i u_i^2}{\text{median } u_i^2}$ in the rank 1 case. We want to see whether R_u obtained by Adam is smaller than that of SGD+M.

After reviewing the weight matrices we got in different settings, we observed that (A) and (B) hold for many layers in those models. For example, on the translation task mentioned in Section 4.1, we found 12 layers which had approximately low rank structures and for 10 of them, R_u values (defined in (B)) obtained by Adam were smaller than those found by SGD+M. Figure 4 shows the result on one typical layer. Results of more layers can be found in Appendix A.5.

Remarks   1. The definition of R_u is based on the connection between diagonal of loss Hessian and weight matrices. Appendix F shows that for a 2-layer linear network, $R_{\text{med},2}^{\text{OPT}}(t) = \frac{\max_i \|W_1^{(t)}[i,:]\|_2^2}{\text{median} \|W_1^{(t)}[i,:]\|_2^2}$. When $W_1 \in \mathbb{R}^{m \times n}$ is approximately rank k, i.e. $W_1 \approx \sum_{i=1}^{k} \sigma_i \boldsymbol{u}_i \boldsymbol{v}_i^T$, denote $\boldsymbol{u}_i = [u_{i1}, u_{i2}, ..., u_{im}]^T$ and $\boldsymbol{v}_i = [v_{i1}, v_{i2}, ..., v_{in}]^T$, we have that for the j-th row, $\|W_1[j,:]\|_2^2 \approx \left\|\sum_{i=1}^{k} \sigma_i u_{ij} \boldsymbol{v}_i^T\right\|_2^2 = \sum_{i=1}^{k} \sigma_i^2 u_{ij}^2$. By defining $\tilde{\boldsymbol{u}} = [\tilde{u}_1, \tilde{u}_2, ..., \tilde{u}_d]^T := \sum_{i=1}^{k} \sigma_i^2 \boldsymbol{u}_i \odot \boldsymbol{u}_i$, we have that $\|W_1[j,:]\|_2^2 \approx \tilde{u}_j$. Although in multi-layer nonlinear neural networks, the connection between diagonal of loss Hessian and the weight matrices is more complicated and $R_{\text{med},2}^{\text{OPT}}(t)$ may depend on the product of many weight matrices rather than one single matrix, we still believe that this definition of R_u is a reasonable ratio to consider.

2. We may also want to consider the right singular vectors $\boldsymbol{v}_1, \boldsymbol{v}_2, ... \boldsymbol{v}_k$ and corresponding $\tilde{\boldsymbol{v}} = [\tilde{v}_1, \tilde{v}_2, ..., \tilde{v}_d]^T := \sum_{i=1}^{k} \sigma_i^2 \boldsymbol{v}_i \odot \boldsymbol{v}_i$ and compute $R_v := \frac{\max_i \tilde{v}_i}{\text{median } \tilde{v}_i}$ for Adam and SGD+M. However, on this translation task, among the 12 layers which were approximately low rank, for only 6 of them, R_v of Adam were smaller, i.e. we did not observe uniformity of the leading right singular vector for Adam. Results of R_v can be found in Appendix A.5. One possible reason is that for a weight matrix, its right singular vectors are closer to the input data than left singular vectors and more easily influenced by the data, therefore may not show uniformity.

7   Conclusion and future work

We demonstrate that adaptive optimization methods bias the training trajectory towards a region where the diagonal of loss Hessian is more uniform, through extensive experiments on language models and theoretical analysis in a simplified setting of two-layer linear networks. Although our findings may not directly lead to an improved algorithm for practical use, they provide a new way of thinking when designing new algorithms: in contrast with the traditional view which tries to design a method that performs better *in* the bad loss geometry, our findings suggest that we can design algorithms which *implicitly avoid* regions with bad geometry. There are a lot of future directions along this line. For example, our theoretical results on the two-layer linear networks may be able to generalize to multi-layer networks. In fact, people conjecture that the key-value-query structure in language models can be approximated by a three-layer linear network. Hence the generalization to multi-layer networks might provide more connection to real deep models and could be an interesting and challenging future direction. Moreover, it is also possible to relax our large-batch assumption (Assumption 3) and prove similar results in the general stochastic setting.

REPRODUCIBILITY STATEMENT

The training details (e.g. hyperparameters) of experiments are specified in Section 4. The source code is provided in the supplemental material. For the theoretical results, Section 3.3 states the full set of assumptions. Section C and Section D in the appendix provide complete proofs.

REFERENCES

Zeyuan Allen-Zhu, Yuanzhi Li, and Yingyu Liang. Learning and generalization in overparameterized neural networks, going beyond two layers. In *Advances in Neural Information Processing Systems*, volume 32. Curran Associates, Inc., 2019a.

Zeyuan Allen-Zhu, Yuanzhi Li, and Zhao Song. A convergence theory for deep learning via over-parameterization. In *International Conference on Machine Learning*, pp. 242–252. PMLR, 2019b.

Sanjeev Arora, Nadav Cohen, and Elad Hazan. On the optimization of deep networks: Implicit acceleration by overparameterization. In *35th International Conference on Machine Learning, ICML 2018*, pp. 372–389. International Machine Learning Society (IMLS), 2018.

Sanjeev Arora, Simon Du, Wei Hu, Zhiyuan Li, and Ruosong Wang. Fine-grained analysis of optimization and generalization for overparameterized two-layer neural networks. In *International Conference on Machine Learning*, pp. 322–332. PMLR, 2019a.

Sanjeev Arora, Noah Golowich, Nadav Cohen, and Wei Hu. A convergence analysis of gradient descent for deep linear neural networks. In *7th International Conference on Learning Representations, ICLR 2019*, 2019b.

Prajjwal Bhargava, Aleksandr Drozd, and Anna Rogers. Generalization in nli: Ways (not) to go beyond simple heuristics, 2021.

Tom Brown, Benjamin Mann, Nick Ryder, Melanie Subbiah, Jared D Kaplan, Prafulla Dhariwal, Arvind Neelakantan, Pranav Shyam, Girish Sastry, Amanda Askell, Sandhini Agarwal, Ariel Herbert-Voss, Gretchen Krueger, Tom Henighan, Rewon Child, Aditya Ramesh, Daniel Ziegler, Jeffrey Wu, Clemens Winter, Chris Hesse, Mark Chen, Eric Sigler, Mateusz Litwin, Scott Gray, Benjamin Chess, Jack Clark, Christopher Berner, Sam McCandlish, Alec Radford, Ilya Sutskever, and Dario Amodei. Language models are few-shot learners. In *Advances in Neural Information Processing Systems*, volume 33, pp. 1877–1901. Curran Associates, Inc., 2020.

Jinghui Chen, Dongruo Zhou, Yiqi Tang, Ziyan Yang, Yuan Cao, and Quanquan Gu. Closing the generalization gap of adaptive gradient methods in training deep neural networks. In Christian Bessiere (ed.), *Proceedings of the Twenty-Ninth International Joint Conference on Artificial Intelligence, IJCAI 2020*, pp. 3267–3275. ijcai.org, 2020. doi: 10.24963/ijcai.2020/452. URL https://doi.org/10.24963/ijcai.2020/452.

Hung-Hsu Chou, Carsten Gieshoff, Johannes Maly, and Holger Rauhut. Gradient descent for deep matrix factorization: Dynamics and implicit bias towards low rank. *arXiv preprint arXiv:2011.13772*, 2020.

Jeremy M. Cohen, Simran Kaur, Yuanzhi Li, J. Zico Kolter, and Ameet Talwalkar. Gradient descent on neural networks typically occurs at the edge of stability. In *9th International Conference on Learning Representations, ICLR 2021, Virtual Event, Austria, May 3-7, 2021*. OpenReview.net, 2021.

Alexandre Défossez, Léon Bottou, Francis Bach, and Nicolas Usunier. A simple convergence proof of adam and adagrad. *arXiv preprint arXiv:2003.02395*, 2020.

Jacob Devlin, Ming-Wei Chang, Kenton Lee, and Kristina Toutanova. BERT: pre-training of deep bidirectional transformers for language understanding. In *Proceedings of the 2019 Conference of the North American Chapter of the Association for Computational Linguistics: Human Language Technologies, NAACL-HLT 2019, Minneapolis, MN, USA, June 2-7, 2019, Volume 1 (Long and Short Papers)*, pp. 4171–4186. Association for Computational Linguistics, 2019. doi: 10.18653/v1/n19-1423. URL https://doi.org/10.18653/v1/n19-1423.

Simon S Du, Xiyu Zhai, Barnabas Poczos, and Aarti Singh. Gradient descent provably optimizes over-parameterized neural networks. In *International Conference on Learning Representations*, 2018.

John Duchi, Elad Hazan, and Yoram Singer. Adaptive subgradient methods for online learning and stochastic optimization. *Journal of machine learning research*, 12(7), 2011.

Desmond Elliott, Stella Frank, Khalil Sima'an, and Lucia Specia. Multi30k: Multilingual english-german image descriptions. In *Proceedings of the 5th Workshop on Vision and Language*, pp. 70–74. Association for Computational Linguistics, 2016. doi: 10.18653/v1/W16-3210. URL `http://www.aclweb.org/anthology/W16-3210`.

Alina Ene, Huy L. Nguyen, and Adrian Vladu. Adaptive gradient methods for constrained convex optimization and variational inequalities. In *Thirty-Fifth AAAI Conference on Artificial Intelligence, AAAI 2021, Thirty-Third Conference on Innovative Applications of Artificial Intelligence, IAAI 2021, The Eleventh Symposium on Educational Advances in Artificial Intelligence, EAAI 2021, Virtual Event, February 2-9, 2021*, pp. 7314–7321. AAAI Press, 2021. URL `https://ojs.aaai.org/index.php/AAAI/article/view/16898`.

Suriya Gunasekar, Blake E Woodworth, Srinadh Bhojanapalli, Behnam Neyshabur, and Nati Srebro. Implicit regularization in matrix factorization. *Advances in Neural Information Processing Systems*, 30, 2017.

Arthur Jacot, Franck Gabriel, and Clément Hongler. Neural tangent kernel: Convergence and generalization in neural networks. *Advances in neural information processing systems*, 31, 2018.

Ziwei Ji and Matus Telgarsky. Polylogarithmic width suffices for gradient descent to achieve arbitrarily small test error with shallow relu networks. In *8th International Conference on Learning Representations, ICLR 2020, Addis Ababa, Ethiopia, April 26-30, 2020*. OpenReview.net, 2020. URL `https://openreview.net/forum?id=HygegyrYwH`.

Kenji Kawaguchi. Deep learning without poor local minima. In D. Lee, M. Sugiyama, U. Luxburg, I. Guyon, and R. Garnett (eds.), *Advances in Neural Information Processing Systems*, volume 29. Curran Associates, Inc., 2016. URL `https://proceedings.neurips.cc/paper/2016/file/f2fc990265c712c49d51a18a32b39f0c-Paper.pdf`.

Diederik P. Kingma and Jimmy Ba. Adam: A method for stochastic optimization. In *3rd International Conference on Learning Representations, ICLR 2015, San Diego, CA, USA, May 7-9, 2015, Conference Track Proceedings*, 2015. URL `http://arxiv.org/abs/1412.6980`.

Matthieu Lerasle. Lecture notes: Selected topics on robust statistical learning theory. *arXiv preprint arXiv:1908.10761*, 2019.

Aitor Lewkowycz, Yasaman Bahri, Ethan Dyer, Jascha Sohl-Dickstein, and Guy Gur-Ari. The large learning rate phase of deep learning: the catapult mechanism. *arXiv preprint arXiv:2003.02218*, 2020.

Yuanzhi Li, Tengyu Ma, and Hongyang Zhang. Algorithmic regularization in over-parameterized matrix sensing and neural networks with quadratic activations. In Sébastien Bubeck, Vianney Perchet, and Philippe Rigollet (eds.), *Proceedings of the 31st Conference On Learning Theory*, volume 75 of *Proceedings of Machine Learning Research*, pp. 2–47. PMLR, 06–09 Jul 2018.

Chaoyue Liu, Libin Zhu, and Mikhail Belkin. Loss landscapes and optimization in over-parameterized non-linear systems and neural networks. *Applied and Computational Harmonic Analysis*, 2022.

Andrew L. Maas, Raymond E. Daly, Peter T. Pham, Dan Huang, Andrew Y. Ng, and Christopher Potts. Learning word vectors for sentiment analysis. In *Proceedings of the 49th Annual Meeting of the Association for Computational Linguistics: Human Language Technologies*, pp. 142–150, Portland, Oregon, USA, June 2011. Association for Computational Linguistics. URL `http://www.aclweb.org/anthology/P11-1015`.

Stephen Merity, Caiming Xiong, James Bradbury, and Richard Socher. Pointer sentinel mixture models. In *5th International Conference on Learning Representations, ICLR 2017, Toulon, France, April 24-26, 2017, Conference Track Proceedings*. OpenReview.net, 2017. URL `https://openreview.net/forum?id=Byj72udxe`.

Sashank J. Reddi, Satyen Kale, and Sanjiv Kumar. On the convergence of adam and beyond. In *6th International Conference on Learning Representations, ICLR 2018, Vancouver, BC, Canada, April 30 - May 3, 2018, Conference Track Proceedings*. OpenReview.net, 2018. URL `https://openreview.net/forum?id=ryQu7f-RZ`.

Yuandong Tian, Xinlei Chen, and Surya Ganguli. Understanding self-supervised learning dynamics without contrastive pairs. In *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pp. 10268–10278. PMLR, 18–24 Jul 2021.

Tijmen Tieleman, Geoffrey Hinton, et al. Lecture 6.5-rmsprop: Divide the gradient by a running average of its recent magnitude. *COURSERA: Neural networks for machine learning*, 4(2):26–31, 2012.

Iulia Turc, Ming-Wei Chang, Kenton Lee, and Kristina Toutanova. Well-read students learn better: On the importance of pre-training compact models. *arXiv preprint arXiv:1908.08962v2*, 2019.

Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N Gomez, Łukasz Kaiser, and Illia Polosukhin. Attention is all you need. *Advances in neural information processing systems*, 30, 2017.

Rachel Ward, Xiaoxia Wu, and Léon Bottou. Adagrad stepsizes: Sharp convergence over nonconvex landscapes. *Journal of Machine Learning Research*, 21(219):1–30, 2020.

A   More experiments of the uniformity of diagonal geometry

A.1   Vanilla SGD vs. Adagrad

In this section, we present the $R_{\text{med}}^{\text{OPT}}(t)$ values defined in eq. (1) obtained by vanilla SGD and Adagrad on a language modeling task[10]. The task is to assign a probability for the likelihood of a given word (or a sequence of words) to follow a sequence of words. We trained a transformer model to solve this problem on both Wikitext-2 (Merity et al., 2017)(CC BY-SA 3.0) and random dataset (generating random integers as targets). This model has roughly 8 layers (not counting normalization and dropout layers)

The setup is the same as in Section 3.2. We used the same learning rate schedule (constant or decreasing) for SGD and Adagrad. We tuned and chose the best (initial) learning rate of SGD. The (initial) learning rate of Adagrad was set as a value under which Adagrad converged faster than SGD with its best (initial) learning rate. We used large batch sizes to make the training procedure more stable. When computing Hessian, we also used large batch sizes. Due to the extremely large dimension, we did the computation on some uniformly selected coordinates, more precisely, 200 coordinates per layer.

We tried different initialization (normal and uniform) by using different gains of the Pytorch initialization schedule.

A.1.1   Experiments on real dataset

Figure 5a shows the training losses on real dataset (wikitext-2). Table 3 (resp. Table 4) shows the $R_{\text{med}}^{\text{OPT}}(t)$ for Adagrad and SGD under uniform (resp. normal) initialization with different gains.

| (a) Wikitext-2 | (b) Random dataset |

Figure 5: Training losses of Adagrad and SGD on wikitext-2 (left) and random data (right)

Table 3: $R_{\text{med}}^{\text{OPT}}(t)$ of Adagrad and SGD under uniform initialization with different gains

(a) Gain = 2

Layer#	Epoch 1		Epoch 20		Epoch 40	
	SGD	Adagrad	SGD	Adagrad	SGD	Adagrad
1	6.07	6.77	5.91	9.77	5.16	10.37
2	4.60	6.26	3.43	1.66	3.44	1.88
3	5.15	6.84	4.35	4.34	4.84	3.60
4	9.47	10.78	9.76	3.54	8.67	3.14
5	12.54	13.96	10.31	6.59	9.79	6.98
6	4.92	5.25	7.21	2.33	7.94	2.28
7	5.73	5.45	40.56	4.57	21.24	4.76
8	9.39	8.87	37.95	4.50	46.03	3.19

(b) Gain = 0.5

Layer#	Epoch 1		Epoch 20		Epoch 40	
	SGD	Adagrad	SGD	Adagrad	SGD	Adagrad
1	69.36	78.60	15.26	7.74	18.22	7.23
2	24.12	24.36	4.05	2.30	3.70	2.04
3	2.83	2.85	3.78	4.98	3.56	4.40
4	5.25	4.74	3.83	5.68	3.11	4.81
5	66.49	67.83	88.75	19.31	63.01	15.64
6	6.54	6.91	3.57	2.08	3.50	1.97
7	3.22	3.73	13.03	3.97	9.55	4.07
8	6.12	5.99	6.73	7.82	5.43	6.98

---

[10]https://pytorch.org/tutorials/beginner/transformer_tutorial.html

Table 4: $R_{\mathrm{med}}^{\mathrm{OPT}}(t)$ of Adagrad and SGD under normal initialization with different gains

(a) Gain = 1

Layer#	Epoch 1		Epoch 20		Epoch 40	
	SGD	Adagrad	SGD	Adagrad	SGD	Adagrad
1	6.76	6.06	8.27	12.28	9.69	11.17
2	9.51	6.61	3.19	1.87	3.21	1.73
3	7.38	7.35	8.61	3.38	9.25	3.94
4	18.02	15.63	6.45	4.86	7.49	4.44
5	12.70	9.35	11.69	11.23	15.07	12.18
6	12.76	11.86	3.84	2.32	3.20	2.09
7	11.79	8.58	17.95	4.32	14.99	4.50
8	17.09	12.73	26.70	5.16	26.91	6.73

(b) Gain = 0.5

Layer#	Epoch 1		Epoch 20		Epoch 40	
	SGD	Adagrad	SGD	Adagrad	SGD	Adagrad
1	9.12	14.46	10.90	8.00	10.19	8.55
2	10.70	15.42	8.52	2.12	8.88	2.04
3	5.73	5.94	10.16	2.80	6.05	2.99
4	16.62	12.94	8.90	3.91	8.12	4.14
5	15.98	16.98	42.57	10.76	18.45	10.16
6	4.84	6.46	7.92	2.66	5.30	2.46
7	6.52	6.55	107.51	3.14	136.38	2.73
8	8.39	8.20	337.34	5.18	315.21	4.48

A.1.2 EXPERIMENTS ON RANDOM DATASET

Figure 5b shows the training losses on random dataset and Table 5 shows the $R_{\mathrm{med}}^{\mathrm{OPT}}(t)$ in different layers.

Table 5: $R_{\mathrm{med}}^{\mathrm{OPT}}(t)$ of Adagrad and SGD for random data

Layer#	Epoch 1		Epoch 20		Epoch 40	
	SGD	Adagrad	SGD	Adagrad	SGD	Adagrad
1	10.88	10.98	9.99	18.66	9.67	22.37
2	9.47	12.15	14.98	4.43	13.01	3.99
3	7.45	8.52	459.71	6.09	451.16	5.11
4	9.84	10.42	135.37	7.22	126.91	6.04
5	7.09	7.88	103.60	353.89	184.61	190.17
6	7.68	8.58	18.38	4.08	18.69	2.73
7	7.81	5.40	294.68	62.72	229.25	29.76
8	13.51	9.16	329.12	20.59	203.70	9.57

A.2 RMSPROP AND AMSGRAD

In this section, we present the results of RMSprop and AMSGrad and compare them with SGD+M. The experiments were conducted on the translation task described in Section 4.1. The learning rates we used were 2.5e-5 for RMSprop, 0.0005 for AMSGrad and 0.03 for SGD+M. Both RMSprop and SGD+M used momentum parameter 0.9. The two momentum parameters (β_1, β_2) of AMSGrad were $(0.9, 0.98)$. Figure 6 shows the training losses and Table 6 shows the corresponding $R_{\mathrm{med}}^{\mathrm{OPT}}(t)$.

Figure 6: Training losses of RMSprop, AMSGrad and SGD+M on the translation task described in Section 4.1.

Table 6: $R_{\text{med}}^{\text{RMSprop}}(t)$, $R_{\text{med}}^{\text{AMSGrad}}(t)$ and $R_{\text{med}}^{\text{SGDM}}(t)$ in some layers, on the translation task described in Section 4.1

Layer#	Epoch 10			Epoch 20			Epoch 40		
	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{RMSprop}}(t)$	$R_{\text{med}}^{\text{AMSGrad}}(t)$	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{RMSprop}}(t)$	$R_{\text{med}}^{\text{AMSGrad}}(t)$	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{RMSprop}}(t)$	$R_{\text{med}}^{\text{AMSGrad}}(t)$
3	3.97	2.69	2.56	2.33	1.89	1.68	2.83	1.62	1.56
5	26.17	21.19	11.36	37.11	17.83	10.85	51.94	10.22	12.31
7	4.10	6.98	6.12	3.94	4.95	2.92	7.58	2.29	2.58
9	29.41	35.72	25.86	37.81	19.89	16.90	30.68	16.24	9.97
12	4.93	6.20	12.67	4.63	6.61	4.64	6.44	5.13	4.06
15	85.06	33.63	19.51	140.99	12.22	6.72	44.07	6.98	5.37
18	8.71	2.99	9.48	3.86	2.44	4.16	3.51	2.10	2.35
21	95.34	11.68	6.62	47.20	6.37	4.74	22.20	4.58	3.58
24	8.70	5.67	6.95	8.13	3.59	5.13	6.46	2.30	2.83
28	4.44	2.42	2.64	4.67	1.85	1.81	2.63	1.46	2.13

A.3 COMPARISON CONDITIONED ON THE SAME LOSS

In this section, we compare $R_{\text{med}}^{\text{SGDM}}(t)$ and $R_{\text{med}}^{\text{Adam}}(t)$ conditioned on the same training loss. More precisely, we make comparison of $R_{\text{med}}^{\text{Adam}}(t)$ and $R_{\text{med}}^{\text{SGDM}}(t')$, where t, t' are picked such that tth Adam iterate and t'th SGD+M iterate have the same training loss. The details of the tasks are described in in Section 4.1. Table 7 shows the results of $R_{\text{med}}^{\text{Adam}}(t)$ and $R_{\text{med}}^{\text{SGDM}}(t')$ in some layers.

Table 7: $R_{\text{med}}^{\text{Adam}}(t)$ and $R_{\text{med}}^{\text{SGDM}}(t')$ in some layers. Dataset and task: (a) sentence classification task on BERT-small, (b) translation task on Multi30k.

(a)

Layer#	Loss 0.251		Loss 0.170		Loss 0.133	
	$R_{\text{med}}^{\text{SGDM}}(t')$	$R_{\text{med}}^{\text{Adam}}(t)$	$R_{\text{med}}^{\text{SGDM}}(t')$	$R_{\text{med}}^{\text{Adam}}(t)$	$R_{\text{med}}^{\text{SGDM}}(t')$	$R_{\text{med}}^{\text{Adam}}(t)$
9	16.77	13.69	14.14	12.71	15.17	9.86
12	16.68	8.29	9.98	8.31	8.90	5.42
15	18.64	7.79	51.39	46.43	80.82	40.97
17	208.29	381.05	464.37	315.58	498.26	313.99
18	14.43	23.56	19.17	19.26	15.76	12.99
22	257.32	88.47	188.55	110.87	197.79	139.48
24	34.22	16.34	16.42	18.08	14.04	15.97

(b)

Layer#	Loss 3.72		Loss 2.78		Loss 1.90	
	$R_{\text{med}}^{\text{SGDM}}(t')$	$R_{\text{med}}^{\text{Adam}}(t)$	$R_{\text{med}}^{\text{SGDM}}(t')$	$R_{\text{med}}^{\text{Adam}}(t)$	$R_{\text{med}}^{\text{SGDM}}(t')$	$R_{\text{med}}^{\text{Adam}}(t)$
3	4.01	4.45	5.80	3.02	2.44	2.28
5	31.19	27.50	44.29	21.46	57.83	19.52
7	5.80	4.38	7.51	3.71	5.25	2.87
9	21.23	53.65	28.99	20.92	44.26	28.13
13	53.18	17.77	51.17	20.64	35.80	35.49
15	82.30	186.41	34.17	13.76	33.87	5.31
21	100.43	23.66	23.45	5.12	12.96	5.35
26	7.45	3.48	4.69	3.10	3.33	2.83
30	19.14	9.54	10.46	5.48	9.56	5.33

Figure 7: Training losses of SGD, Adam after SGD and Adam for the translation task

Table 8: $R_{\text{med}}^{\text{OPT}}(t)$ of SGD, Adam after SGD and Adam in some layers after roughly 2160 iterations

Layer#	SGD	Adam	Adam after SGD	Adam
13	294.76	150.02	332.96	150.02
14	14.34	5.84	5.33	5.84
15	36.38	16.66	11.86	16.66
16	6.47	7.05	3.76	7.05
17	17.17	6.05	4.76	6.05
26	5.68	3.53	2.30	3.53
27	14.33	15.93	21.76	15.93
28	9.10	1.71	1.71	1.71
29	8.22	3.04	2.82	3.04
30	11.39	5.12	5.29	5.12

A.4 EXPERIMENTS OF SWITCHING FROM SGD TO ADAM

In this section we describe another learning schedule: the "Adam after SGD" schedule, where we switch from SGD to Adam in the middle to see whether the loss and $R_{\text{med}}^{\text{OPT}}(t)$ can catch up with the model trained by Adam from the very beginning. Again, we used the same model as in the translation

task in Section 4.1. In this section, we did not add momentum term to SGD in order to get a larger gap between SGD and Adam than the case using momentum. We want to see whether this larger gap can be closed after switching to Adam in the middle.

As is shown in Figure 7 and Table 8, both the loss gap and the gap of $R_{\text{med}}^{\text{OPT}}(t)$ were closed after a period of training after switching algorithms, which provides evidence of the connection between convergence speed and uniformity of diagonal of loss Hessian.

A.5 THE LOW RANK STRUCTURE

In this section, we present more results for the experiments in Section 6.

We examined the weights of the model trained for the translation task in Section 4.1. Among roughly 30 layers, we observed that for 12 layers, at least the weight matrices obtained by Adam after training have approximately low rank structures.

Figure 8 shows the examples of layers with or without the low rank structure.

Figure 8: Examples of layers with approximately low rank structure (right) and without low rank structure (left)

We then studied the uniformity of leading singular vectors of these 12 layers, i.e. computed R_u and R_v defined in (B) and the second remark of Section 6. The observation is that for 10 out of these 12 layers, R_u values of Adam were smaller those of SGD, which implies the uniformity of leading left singular vectors of Adam. However, we did not observe significant uniformity for Adam in terms of leading right singular vectors (R_v). The second remark of Section 6 discusses possible reasons.

Figure 9 shows how R_u and R_v changed over time in some layers.

Figure 9: R_u and R_v for Adam and SGD with momentum in some layers

A.6 How the (adaptive) gradient aligns with diagonal of loss Hessian

In this section we present the uniformity of diagonal geometry of adaptive methods from another perspective. Denote H_{ii} as the (i, i)-th element of the loss Hessian H and g_i as the i-th element of the gradient. It is conjectured that when $|H_{ii}|$ is large, the corresponding $|g_i|$ is usually large as well. For adaptive methods, we can regard the update per step as the learning rate times the "adaptive gradient". Let's use $g_{\text{adapt},i}$ to represent the i-th component of the adaptive gradient. Through experiments on language models, we found that $|g_{\text{adapt},i}|$ for different i are quite uniform and do not align with $|H_{ii}|$ as the true gradient $|g_i|$ does.

In the experiments, we first sorted $|H_{ii}|$ in the ascent order: $|H_{i_1,i_1}| \leq |H_{i_2,i_2}| \leq ... \leq |H_{i_d,i_d}|$ (suppose $H \in \mathbb{R}^{d \times d}$), and then plotted the corresponding $|g_{i_k}|$ and $|g_{\text{adapt},i_k}|$ for $k \in [d]$.

A.6.1 SGD vs. Adagrad

Here we compare SGD and Adagrad on the language modeling task on wikitext-2 described in Section A.1. We observed that the figures of all layers are quite similar so we select one layer as an example, as is shown in Figure 10.

Figure 10: How the true gradient ($\{|g_{i_k}|\}_{k=1}^d$) and "adaptive gradient" ($\{|g_{\text{adapt},i_k}|\}_{k=1}^d$) align with diagonal of Hessian ($\{|H_{i_k,i_k}|\}_{k=1}^d$). Here coordinates are sorted such that $|H_{i_1,i_1}| \leq |H_{i_2,i_2}| \leq ... \leq |H_{i_d,i_d}|$ (suppose $H \in \mathbb{R}^{d \times d}$). Experiments were conducted on the model described in Section A.1. This figure shows the results on the 12-th layer.

A.6.2 SGD with momentum vs. Adam

Here we compare Adam and SGD+M on the tasks described in Section 4.1. Again, we select one layer as an example for each task. Figure 11 shows the results on the sentence classification task and Figure 12 shows the results on the translation task.

A.7 Adding regularization and other tricks

In this section, we add weight decay to both Adam and SGD+M on the translation task described in Section 4. The momentum parameter β in SGD was set as 0.9. The two momentum parameters (β_1, β_2) of Adam were set as $(0.9, 0.98)$. For both algorithms, we set the weight decay parameter as 0.001. We trained the model using constant learning rates for 60 epochs (1800 iterations). We tuned and chose the best learning rate 0.03 for SGD+M. The learning rate of Adam was set as 0.0001, under which Adam converged faster than SGD+M with its best learning rate 0.03. Figure 13a shows the

Figure 11: How the true gradient ($\{|g_{i_k}|\}_{k=1}^d$) and "adaptive gradient" ($\{|g_{\text{adapt},i_k}|\}_{k=1}^d$) align with diagonal of Hessian ($\{|H_{i_k,i_k}|\}_{k=1}^d$). Here coordinates are sorted such that $|H_{i_1,i_1}| \leq |H_{i_2,i_2}| \leq ... \leq |H_{i_d,i_d}|$ (suppose $H \in \mathbb{R}^{d\times d}$). Experiments were conducted on the sentence classification task described in Section 4.1. This figure shows the results on the 12-th layer.

Figure 12: How the true gradient ($\{|g_{i_k}|\}_{k=1}^d$) and "adaptive gradient" ($\{|g_{\text{adapt},i_k}|\}_{k=1}^d$) align with diagonal of Hessian ($\{|H_{i_k,i_k}|\}_{k=1}^d$). Here coordinates are sorted such that $|H_{i_1,i_1}| \leq |H_{i_2,i_2}| \leq ... \leq |H_{i_d,i_d}|$ (suppose $H \in \mathbb{R}^{d\times d}$). Experiments were conducted on the translation task described in Section 4.1. This figure shows the results on the 5-th layer.

training losses and Table 9 shows the values of $R_{\text{med}}^{\text{Adam}}(t)$, $R_{\text{med}}^{\text{SGDM}}(t)$ and $\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$ in some randomly selected layers.

A.8 RESULTS ON IMAGE TASKS

We trained a ResNet[11] on CIFAR-10 dataset and compared the convergence speed and $R_{\text{med}}^{\text{OPT}}(t)$ of SGD+M and Adam. The momentum parameter β in SGD was set as 0.9. The two momentum parameters (β_1, β_2) of Adam were set as (0.9, 0.98). The model was trained using constant learning

---

[11]We borrowed the implementation here https://pytorch-tutorial.readthedocs.io/en/latest/tutorial/chapter03_intermediate/3_2_2_cnn_resnet_cifar10/ and replace the "layers" array [2,2,2] with [1,1,1].

(a)   (b)

Figure 13: (a) Training losses of Adam and SGD+M for the translation task, both with weight decay. (b) Training losses of Adam and SGD+M for a ResNet trained on CIFAR-10.

Table 9: $R_{\text{med}}^{\text{Adam}}(t)$ and $R_{\text{med}}^{\text{SGDM}}(t)$ (both using weight decay) in some layers for the translation task.

Layer#	Epoch 0		Epoch 30			Epoch 55		
	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$
3	73.09	73.09	17.65	13.11	1.35	13.38	6.28	2.13
5	469.88	469.88	293.48	310.85	0.94	601.68	588.12	1.02
7	80.78	80.78	8.22	39.65	0.21	13.65	4.85	2.81
9	494.27	494.27	150.14	123.79	1.21	301.89	119.53	2.53
15	632.10	632.10	277.18	175.34	1.58	334.48	282.88	1.18
18	55.08	55.08	6.56	4.45	1.47	23.88	4.52	5.29
21	549.62	549.62	257.89	44.78	5.76	515.99	53.79	9.59
24	107.51	107.51	8.54	3.64	2.34	53.79	3.32	16.20
28	13.77	13.77	4.74	2.37	2.00	15.60	2.15	7.24
30	491.62	491.62	6.91	2.66	2.60	9.60	2.02	4.77

rates for 41 epochs (2050 iterations). We tuned and chose the best learning rates for both algorithms: 0.5 for SGD+M and 0.005 for Adam. Figure 13b shows the training losses and Table 10 shows the values of $R_{\text{med}}^{\text{Adam}}(t)$, $R_{\text{med}}^{\text{SGDM}}(t)$ and $\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$.

Table 10: $R_{\text{med}}^{\text{Adam}}(t)$ and $R_{\text{med}}^{\text{SGDM}}(t)$ for ResNet on CIFAR-10.

Layer#	Epoch 10			Epoch 20			Epoch 40		
	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$	$R_{\text{med}}^{\text{SGDM}}(t)$	$R_{\text{med}}^{\text{Adam}}(t)$	$\frac{R_{\text{med}}^{\text{SGDM}}(t)}{R_{\text{med}}^{\text{Adam}}(t)}$
1	6.88	25.34	0.27	3.74	39.35	0.09	4.39	15.80	0.28
2	110.19	35.93	3.07	32.97	36.27	0.91	60.69	28.06	2.16
3	40.89	16.92	2.42	13.98	15.92	0.88	11.70	37.01	0.32
4	28.56	23.66	1.21	11.48	13.04	0.88	7.99	14.51	0.55
5	13.47	23.78	0.57	8.64	12.07	0.72	6.52	14.23	0.46
6	18.72	12.49	1.50	12.19	8.80	1.38	8.96	21.69	0.41
7	18.85	39.25	0.48	9.00	12.81	0.70	13.87	11.42	1.22
8	13.79	19.91	0.69	8.87	11.72	0.76	7.48	9.34	0.80
9	12.50	14.85	0.84	9.62	8.06	1.19	11.35	8.08	1.41
10	14.89	14.53	1.02	8.15	5.80	1.41	6.21	8.89	0.70

B   PROOF SKETCH OF THEOREM 1

Now we give a proof sketch of Theorem 1, which contains three major steps. The detailed proof can be found in Appendix F, C and D.

First we relate the diagonal of Hessian to weight matrices W_1, W_2. Under Assumption 1, denote $W_1[i,:]$ as the i-th row of W_1 and $W_2 := [w_{2i}, w_{22}, ..., w_{2d}]$. Since the input dataset is whitened, we can show that

$$R_{\text{med},1}^{\text{OPT}}(t) = \frac{\max_i \left(w_{2i}^{(t)}\right)^2}{\text{median}\left(w_{2i}^{(t)}\right)^2}, \quad R_{\text{med},2}^{\text{OPT}}(t) = \frac{\max_i \left\|W_1^{(t)}[i,:]\right\|_2^2}{\text{median}\left\|W_1^{(t)}[i,:]\right\|_2^2}.$$

Next, due to the one-dimensional output, we can prove that W_1 converges to an approximately rank-1 matrix. More precisely, we have

$$W_1^{(t)} = \boldsymbol{u}^{(t)}\boldsymbol{v}^{(t)T} + R_1^{(t)},$$
$$W_2^{(t)} = c^{(t)}\boldsymbol{u}^{(t)T} + R_2^{(t)T}.$$

where $c^{(t)}$ is a scalar, $\boldsymbol{u}^{(t)}, \boldsymbol{v}^{(t)}, R_2^{(t)} \in \mathbb{R}^d$ and $R_1^{(t)} \in \mathbb{R}^{d \times d}$. Denote the i-th coordinate of $\boldsymbol{u}^{(t)}, \boldsymbol{v}^{(t)}, R_2^{(t)}$ as $u_i^{(t)}, v_i^{(t)}, R_{2i}^{(t)}$, respectively. Denote the (i,j)-th element of $R_1^{(t)}$ as $R_1^{(t)}[i,j]$. We have that $\forall i,j \in [d] : \left|R_{2i}^{(t)}\right| \ll c^{(t)}\left|u_i^{(t)}\right|$ and $\left|R_1^{(t)}[i,j]\right| \ll \left|u_i^{(t)}v_i^{(t)}\right|$.

Using the rank 1 structure, we can further simplify $R_{\text{med},1}^{\text{OPT}}(t)$ and $R_{\text{med},2}^{\text{OPT}}(t)$ by

$$R_{\text{med},k}^{\text{OPT}}(t) \approx \frac{\max_i \left(u_i^{(t)}\right)^2}{\text{median}\left(u_i^{(t)}\right)^2}, k = 1, 2. \tag{4}$$

The final step is the detailed analysis of $\boldsymbol{u}^{(t)}$.

For SGD+M, we can prove that $\boldsymbol{u}^{(t)} \approx C(t)[X_1, X_2, ..., X_d]^T$ where $C(t) \in \mathbb{R}$ and $X_i, i \in [d]$ are i.i.d. Gaussian variables. Then we have with high probability, $\frac{\max_i \left(u_i^{(t)}\right)^2}{\text{median}\left(u_i^{(t)}\right)^2} = \Omega(\log d)$. For Adam, we can prove that $\forall i \in [d] : u_i^{(t)} \in \{\pm 1\}$, which gives us $\frac{\max_i \left(u_i^{(t)}\right)^2}{\text{median}\left(u_i^{(t)}\right)^2} = 1$. Substituting into eq. (4) completes the proof.

C   ANALYSIS OF SGD+M

Note that $A = \frac{1}{m}YX^T$, $\Lambda_{xx} := \frac{1}{m}XX^T$. Denote $g_k^{(t)} := \nabla_{W_k}L(W^{(t)}), k = 1, 2$. We have that

$$g_1^{(t)} = W_2^{(t)T}\left(W_2^{(t)}W_1^{(t)} - A\right), \quad g_2^{(t)} = \left(W_2^{(t)}W_1^{(t)} - A\right)W_1^{(t)T}.$$

Let $\tilde{A}^{(t)}, \tilde{\Lambda}_{xx}^{(t)}$ and $\tilde{g}_k^{(t)}, k = 1, 2$ be the corresponding batch versions at time t. By eq. (2), the update rules of W_1 and W_2 for SGD+M are given by:

$$W_1^{(t+1)} = W_1^{(t)} - \eta\sum_{\tau=0}^t \beta^{t-\tau}W_2^{(\tau)T}\left(W_2^{(\tau)}W_1^{(\tau)} - A\right) - \eta\sum_{\tau=0}^t \beta^{t-\tau}Dg_1^{(\tau)},$$

$$W_2^{(t+1)} = W_2^{(t)} - \eta\sum_{\tau=0}^t \beta^{t-\tau}\left(W_2^{(\tau)}W_1^{(\tau)} - A\right)W_1^{(\tau)T} - \eta\sum_{\tau=0}^t \beta^{t-\tau}Dg_2^{(\tau)},$$

where

$$Dg_1^{(t)} := \tilde{g}_1^{(t)} - g_1^{(t)} = W_2^{(t)T}\left(W_2^{(t)}W_1^{(t)}\left(\tilde{\Lambda}_{xx}^{(t)} - \Lambda_{xx}\right) - \left(\tilde{A}^{(t)} - A\right)\right),$$

$$Dg_2^{(t)} := \tilde{g}_2^{(t)} - g_2^{(t)} = \left(W_2^{(t)}W_1^{(t)}\left(\tilde{\Lambda}_{xx}^{(t)} - \Lambda_{xx}\right) - \left(\tilde{A}^{(t)} - A\right)\right)W_1^{(t)T}.$$

Based on the magnitude of W_2 and W_1, we can intuitively divide the training procedure into 2 phases.

1. **First phase**: the first several iterations when W_1 and W_2 are "small" so that $W_2 W_1 - A \approx -A$.

2. **Second phase**: later iterations when $W_2 W_1$ cannot be ignored.

More formally, the boundary between the first and second phase is defined below.

Definition 1 (End of the first phase). *The end of the first phase (denoted as T_1) is defined as*
$$T_1 := \inf \left\{ t \geq 0 : \exists i, j \in [d] : \left| w_{2i}^{(t)} \right| \geq \frac{1}{d^{\frac{\alpha}{2}}} \text{ or } \left| W_1^{(t)}[i,j] \right| \geq \frac{1}{d^{\frac{\alpha}{2}}} \right\}.$$

Denote the j-th coordinate of $W_2 W_1$ and A as $(W_2 W_1)_j$ and A_j, respectively. By Assumption 2 and the assumption that $\forall j \in [d] : A_j > 0, A_j = \Theta(1)$, at the beginning, w.h.p., $\forall j \in [d] : (W_2 W_1)_j - A_j < 0$. During the training, each $(W_2 W_1)_j$ increases and approaches A_j. We hope that by choosing a small learning rate, when $(W_2 W_1)_j$ overshoots for some coordinate j, i.e. $(W_2 W_1)_j > A_j$, it will be close to convergence. To analyze this overshooting issue more carefully, let's first define the following "almost overshooting time".

Definition 2 (Almost overshooting time). *For $\epsilon > 0$, denote $\epsilon_0 := \frac{1}{d^{\frac{1}{4}\alpha - 1}} + \epsilon \log \sqrt{\frac{d}{\epsilon}}$. Define*
$$T_2 := \inf \left\{ t \geq 0 : \exists j \in [d] : \left(W_2^{(t)} W_1^{(t)} \right)_j - A_j \geq -\sqrt{\epsilon_0} \right\}.$$

Definition 3 (Convergence time). *Denote $E^{(t)} := W_2^{(t)} W_1^{(t)} - A$. For $\epsilon > 0$, we define the "convergence time" $T_3 := \inf \left\{ t \geq 0 : \left\| E^{(t)} \right\|_2^2 \leq \epsilon \right\}$.*

We can first show that after the first phase, i.e. when $t = T_1$, W_1 will become an approximately rank-1 matrix, as described in the following lemma.

Lemma 1. *Under Assumption 1, 2 and 3, suppose $\sigma \leq \frac{\eta^{3/2}}{d^{\alpha/2+1}}$. By picking $\eta \leq \mathcal{O}\left(\frac{1}{d^\alpha}\right)$, we have that when $t = T_1$, $\bar{L}\left(W^{(T_1)}\right) = \Theta(d)$, and that*
$$W_1^{(T_1)} = R_1^{(T_1)} + \boldsymbol{u}^{(T_1)} \boldsymbol{v}^{(T_1)T},$$
$$W_2^{(T_1)} = R_2^{(T_1)T} + c^{(T_1)} \boldsymbol{u}^{(T_1)T},$$

where $c^{(T_1)} \in \mathbb{R}$, $\boldsymbol{u}^{(T_1)}, \boldsymbol{v}^{(T_1)}, R_2^{(T_1)} \in \mathbb{R}^d$ and $R_1^{(T_1)} \in \mathbb{R}^{d \times d}$. Denote the i-th coordinate of $\boldsymbol{u}^{(T_1)}, \boldsymbol{v}^{(T_1)}, R_2^{(T_1)}$ as $u_i^{(T_1)}, v_i^{(T_1)}, R_{2i}^{(T_1)}$, respectively, and the (i,j)-th element of $R_1^{(T_1)}$ as $R_1^{(T_1)}[i,j]$. Then w.h.p.,
$$\forall 1 \leq i, j \leq d : \quad \frac{\left| R_1^{(T_1)}[i,j] \right|}{\left| u_i^{(T_1)} v_j^{(T_1)} \right|} \leq \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha - 1}}\right), \quad \frac{\left| R_{2i}^{(T_1)} \right|}{\left| c^{(T_1)} u_i^{(T_1)} \right|} \leq \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha - 1}}\right).$$

The following lemma tells us that this approximate rank-1 structure is preserved when $T_1 \leq t \leq \min\{T_2, T_3\}$.

Lemma 2. *Under Assumption 1, 2 and 3, suppose $\sigma \leq \frac{\eta^{3/2}}{d^{\alpha/2+1}}$. By picking $\eta \leq \mathcal{O}\left(\frac{\epsilon}{d^{\frac{7\alpha}{4}+4}}\right)$, for $T_1 \leq t \leq \min\{T_2, T_3\}$, we have that*
$$W_1^{(t)} = \boldsymbol{u}^{(T_1)} \boldsymbol{v}^{(t)T} + R_1^{(t)},$$
$$W_2^{(t)} = c^{(t)} \boldsymbol{u}^{(T_1)T} + R_2^{(t)T}.$$

We have that
$$\forall 1 \leq i, j \leq d : \quad \frac{\left| R_1^{(t)}[i,j] \right|}{\left| u_i^{(T_1)} v_j^{(t)} \right|} \leq \tilde{\mathcal{O}}(\epsilon_0), \quad \frac{\left| R_{2i}^{(t)} \right|}{\left| c^{(t)} u_i^{(T_1)} \right|} \leq \tilde{\mathcal{O}}(\epsilon_0),$$

where ϵ_0 is defined in Definition 2. Moreover, when $t = \min\{T_2, T_3\}$, $\bar{L}\left(W^{(t)}\right) = \mathcal{O}(\epsilon_0 d)$.

The following lemma gives us a more detailed description of $\boldsymbol{u}^{(T_1)}$.

Lemma 3. *The $\boldsymbol{u}^{(T_1)}$ in Lemma 1 and 2 can be written as $\boldsymbol{u}^{(T_1)} = X + Y$ where $X_i, i \in [d]$ are i.i.d Gaussian random variables and that w.h.p. $\forall i \in [d] : \frac{|Y_i|}{|X_i|} \leq \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha - \frac{1}{2}}}\right)$.*

Now we are ready to prove the SGD+M part of Theorem 1.

C.1 PROOF OF THE SGD+M PART OF THEOREM 1

Define $T_{\text{SGD},1} = T_1, T_{\text{SGD},2} = \min\{T_2, T_3\}$. By picking $\eta \leq \mathcal{O}\left(\frac{\epsilon}{d^{\frac{7\alpha}{4}+4}}\right)$, we can apply Lemma 1 and 2 to conclude that $\bar{L}\left(W^{(T_{\text{SGD},1})}\right) = \Theta(d)$ and $\bar{L}\left(W^{(T_{\text{SGD},2})}\right) = \mathcal{O}(\epsilon_0 d)$. For any $p > 0$, by picking $0 < \epsilon < \frac{1}{d^p}$ and $\alpha \geq 4(p+2)$, we have $\bar{L}\left(W^{(T_{\text{SGD},2})}\right) = \mathcal{O}(\epsilon_0 d) \leq \tilde{\mathcal{O}}\left(\frac{1}{d^p}\right)$.

Moreover, when $t \in [T_{\text{SGD},1}, T_{\text{SGD},2}]$, the conditions in Lemma 30 are satisfied with $\delta = \tilde{\mathcal{O}}(\epsilon_0)$. Then we can apply Lemma 30 and get that

$$R_{\text{med},1}^{\text{SGDM}}(t), R_{\text{med},2}^{\text{SGDM}}(t) \geq \left(\frac{1 - \tilde{\mathcal{O}}(\epsilon_0)}{1 + \tilde{\mathcal{O}}(\epsilon_0)}\right)^2 \cdot \frac{\max_i \left(u_i^{(T_1)}\right)^2}{\text{median} \left(u_i^{(T_1)}\right)^2}.$$

By Lemma 3, $\boldsymbol{u}^{(T_1)} = X + Y$ where w.h.p. $\forall i \in [d] : \frac{|Y_i|}{|X_i|} \leq \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha - \frac{1}{2}}}\right)$.

This fact yields

$$\forall i \in [d] : \frac{\max_i \left(u_i^{(T_1)}\right)^2}{\text{median} \left(u_i^{(T_1)}\right)^2} \geq \left(\frac{1 - \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha - \frac{1}{2}}}\right)}{1 + \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha - \frac{1}{2}}}\right)}\right)^2 \frac{\max_i X_i^2}{\text{median} X_i^2}.$$

Here $X_i, i \in [d]$ are i.i.d Gaussian random variables by Lemma 3. To prove the concentration of median X_i^2, we borrow the Proposition 12 in Chapter 2.3 of (Lerasle, 2019). By setting $K = N = d$ in this proposition, we have

$$\mathbb{P}\left(\left|\text{median } X_i^2 - \mathbb{E}[X_1^2]\right| > 2\sqrt{\text{Var}(X_1^2)}\right) \leq e^{-\frac{d}{8}}$$

Denote σ^2 as the variance of $X_i, i \in [d]$. Then $\mathbb{E}[X_i^2] = \sigma^2$ and $\text{Var}(X_i^2) = 2\sigma^4$. Hence

$$\mathbb{P}\left(\left|\text{median } X_i^2 - \sigma^2\right| > 2\sqrt{2}\sigma^2\right) \leq e^{-\frac{d}{8}}$$

That means with high probability, median $X_i^2 \leq C\sigma^2$ for some $C > 0$. By Lemma 34 in Appendix G, we know that w.h.p.

$$\max_{1 \leq i \leq d} X_i^2 = \sigma^2 \Omega(\log d),$$

which gives us w.h.p.

$$\frac{\max_{1 \leq i \leq d} X_i^2}{\text{median } X_i^2} = \Omega(\log d).$$

Hence we have proved that $R_{\text{med},1}^{\text{SGDM}}(t), R_{\text{med},2}^{\text{SGDM}}(t) \geq \Omega(\log d)$.

C.2 PROOF OF LEMMA 1

In the first phase, $W_2 W_1$ is "small", and we write the update equations in the following way

$$
\begin{aligned}
W_1^{(t+1)} &= W_1^{(t)} - \eta \sum_{\tau=0}^t \beta^{t-\tau} W_2^{(\tau)T} \left(W_2^{(\tau)} W_1^{(\tau)} - A \right) - \eta \sum_{\tau=0}^t \beta^{t-\tau} Dg_1^{(\tau)} \\
&= W_1^{(t)} + \eta \sum_{\tau=0}^t \beta^{t-\tau} W_2^{(\tau)T} A - \eta \sum_{\tau=0}^t \beta^{t-\tau} W_2^{(\tau)T} W_2^{(\tau)} W_1^{(\tau)} - \eta \sum_{\tau=0}^t \beta^{t-\tau} Dg_1^{(\tau)} \\
&= W_1^{(t)} + \eta W_2^{(t)T} A \sum_{\tau=0}^t \beta^{t-\tau} + \eta \sum_{\tau=0}^t \beta^{t-\tau} \left(W_2^{(\tau)T} - W_2^{(t)T} \right) A \\
&\quad - \eta \sum_{\tau=0}^t \beta^{t-\tau} W_2^{(\tau)T} W_2^{(\tau)} W_1^{(\tau)} - \eta \sum_{\tau=0}^t \beta^{t-\tau} Dg_1^{(\tau)} \\
&= W_1^{(t)} + \frac{\eta}{1-\beta} W_2^{(t)T} A + \frac{\eta}{1-\beta} r_1^{(t)},
\end{aligned}
\tag{5}
$$

where

$$
\begin{aligned}
r_1^{(t)} &= -\beta^{t+1} W_2^{(t)T} A + (1-\beta) \sum_{\tau=0}^t \beta^{t-\tau} \left(W_2^{(\tau)T} - W_2^{(t)T} \right) A \\
&\quad - (1-\beta) \sum_{\tau=0}^t \beta^{t-\tau} W_2^{(\tau)T} W_2^{(\tau)} W_1^{(\tau)} - (1-\beta) \sum_{\tau=0}^t \beta^{t-\tau} Dg_1^{(\tau)}.
\end{aligned}
$$

Similarly, we have

$$
\begin{aligned}
W_2^{(t+1)} &= W_2^{(t)} - \eta \sum_{\tau=0}^t \beta^{t-\tau} \left(W_2^{(\tau)} W_1^{(\tau)} - A \right) W_1^{(\tau)T} \\
&= W_2^{(t)} + \frac{\eta}{1-\beta} A W_1^{(t)T} + \frac{\eta}{1-\beta} r_2^{(t)},
\end{aligned}
\tag{6}
$$

where

$$
\begin{aligned}
r_2^{(t)} &= -\beta^{t+1} A W_1^{(t)T} + (1-\beta) \sum_{\tau=0}^t \beta^{t-\tau} A \left(W_1^{(\tau)T} - W_1^{(t)T} \right) \\
&\quad - (1-\beta) \sum_{\tau=0}^t \beta^{t-\tau} W_2^{(\tau)} W_1^{(\tau)} W_1^{(\tau)T} - (1-\beta) \sum_{\tau=0}^t \beta^{t-\tau} Dg_2^{(\tau)}.
\end{aligned}
$$

The following lemma gives us an explicit formula of $W_2^{(t)}$.

Lemma 4. *Let $\lambda_1 < \lambda_2$ be the two roots of the quadratic equation $x^2 - 2x + 1 - \frac{\eta^2}{(1-\beta)^2} \|A\|_2^2 = 0$. Pick $\eta < \frac{1-\beta}{\|A\|_2}$, then we have that*

$$
W_2^{(t)} = C_1 \lambda_1^t + \left(C_2 + r_5^{(t)} \right) \lambda_2^t,
$$

where $C_1 = -\frac{W_2^{(1)} - \lambda_2 W_2^{(0)}}{\lambda_2 - \lambda_1}$, $C_2 = \frac{W_2^{(1)} - \lambda_1 W_2^{(0)}}{\lambda_2 - \lambda_1}$. $r_5^{(t)}$ will be specified in the proof.

We can prove that in the first phase, $r_5^{(t)}$ is "small". More specifically, denote its i-th coordinate as $r_{5i}^{(t)}$, and the i-th coordinate of C_2 as C_{2i}. Then the following lemmas tell us that $\forall i \in [d], \left| r_{5i}^{(t)} \right| \leq \mathcal{O}\left(\frac{1}{d^{p(\alpha)}} \right)$, where w.h.p. $\mathcal{O}\left(\frac{1}{d^{p(\alpha)}} \right) \ll \min_{i \in [d]} |C_{2i}|$.

We first have the following bounds of $\left| r_{1i}^{(t)} \right|, \left| r_{2i}^{(t)} \right|$ and $\left| r_{5i}^{(t)} \right|$ for $i \in [d]$.

Lemma 5. *Under Assumption 1, 2 and 3, suppose $\sigma \leq \frac{\eta^{3/2}}{d^{\alpha/2+1}}$. For $t \leq T_1$, pick $\eta \leq \mathcal{O}\left(\frac{1}{d^{\alpha}} \right)$. If $t \geq \Omega(\log d)$ in the first phase, we have that $\forall i \in [d] : \left| r_1^{(t)}[i,j] \right| \leq \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3}{2}\alpha - 1}} \right), \left| r_{2i}^{(t)} \right| \leq \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3}{2}\alpha - 2}} \right).$*

Lemma 6. *Under conditions of Lemma 5, for* $t \le T_1$*, we have that* $\forall i \in [d] : \left| r_{5i}^{(t)} \right| \le \tilde{\mathcal{O}} \left(\frac{1}{d^{\frac{3}{2}\alpha - 1}} \right)$.

Next we prove upper and lower bounds of $|C_{1i}|$ and $|C_{2i}|$ for $i \in [d]$.

Lemma 7. *Under Assumption 1, 2 and 3, suppose* $\sigma \le \frac{\eta^{3/2}}{d^{\alpha/2+1}}$*. Pick* $\eta < \frac{1-\beta}{\|A\|_2}$*, we have that*
i) w.h.p., $\forall i \in [d] : |C_{1i}| \le \tilde{\mathcal{O}} \left(\frac{1}{d^{\alpha}} \right), |C_{2i}| \le \tilde{\mathcal{O}} \left(\frac{1}{d^{\alpha}} \right)$*;*
ii) C_2 *can be written as* $C_2 := \frac{1}{2} (C_3 + C_4)$ *where* $C_{3i}, i \in [d]$ *are i.i.d Gaussian random variables and that w.h.p.* $\forall i \in [d] : \frac{|C_{4i}|}{|C_{3i}|} \le \tilde{\mathcal{O}} \left(\frac{1}{d^{\frac{1}{4}\alpha - \frac{1}{2}}} \right)$*;*
iii) w.h.p., $\forall i \in [d], |C_{1i}| \ge \tilde{\Omega} \left(\frac{1}{d^{\frac{5}{4}\alpha}} \right), |C_{2i}| \ge \tilde{\Omega} \left(\frac{1}{d^{\frac{5}{4}\alpha}} \right)$.

Now we are ready to prove Lemma 1.

Lemma 4 tells us that

$$W_2^{(t)} = C_1 \lambda_1^t + \left(C_2 + r_5^{(t)} \right) \lambda_2^t,$$

where $\lambda_1 = 1 - \frac{\eta}{1-\beta} \|A\|_2$ and $\lambda_2 = 1 + \frac{\eta}{1-\beta} \|A\|_2$.

Under the conditions of Theorem 1 and pick $\eta \le \mathcal{O} \left(\frac{1}{d^{\alpha}} \right)$, by Lemma 6 and 7, we know that w.h.p. $\forall t \le T_1, \forall 1 \le i \le d$,

$$\left| r_{5i}^{(t)} \right| \le \tilde{\mathcal{O}} \left(\frac{1}{d^{\frac{3}{2}\alpha - 1}} \right), \quad |C_{2i}| \ge \tilde{\Omega} \left(\frac{1}{d^{\frac{5}{4}\alpha}} \right), \quad |C_{2i}| \le \tilde{\mathcal{O}} \left(\frac{1}{d^{\alpha}} \right), \quad \frac{\left| r_{5i}^{(t)} \right|}{|C_{2i}|} \le \tilde{\mathcal{O}} \left(\frac{1}{d^{\frac{1}{4}\alpha - 1}} \right). \tag{7}$$

Now let's only focus on the update of $W_2^{(t)}$ (ignoring $W_1^{(t)}$ for a while) and consider how large t should be so that $\left| w_{2i}^{(t)} \right|$ reaches $\frac{1}{d^{\alpha/2}}$ for some coordinate i. Define $t_1 = \inf \left\{ t \ge 0 : \exists i \in [d] : \left| w_{2i}^{(t)} \right| \ge \frac{1}{d^{\frac{\alpha}{2}}} \right\}$. We will verify later that for $t \le t_1, \forall i, j \in [d] : \left| W_1^{(t)}[i,j] \right| = o \left(\frac{1}{d^{\alpha/2}} \right)$, which means that t_1 is essentially T_1 in Definition 1.

Since $\left| w_{2i}^{(t_1)} \right|$ is dominated by $|C_{2i}| \lambda_2^{t_1}$, we have $|C_{2i_0}| \lambda_2^{t_1} = \Theta \left(\frac{1}{d^{\alpha/2}} \right)$ for some $i_0 \in [d]$, which means w.h.p.

$$\frac{\Theta \left(\frac{1}{d^{\frac{\alpha}{2}}} \right)}{\tilde{\mathcal{O}} \left(\frac{1}{d^{\alpha}} \right)} \le \lambda_2^{t_1} \le \frac{\Theta \left(\frac{1}{d^{\frac{\alpha}{2}}} \right)}{\tilde{\Omega} \left(\frac{1}{d^{\frac{5}{4}\alpha}} \right)}, \quad \Rightarrow \quad \tilde{\Omega} \left(d^{\frac{\alpha}{2}} \right) \le \lambda_2^{t_1} = \left(1 + \frac{\eta}{1-\beta} \|A\|_2 \right)^{t_1} \le \tilde{\mathcal{O}} \left(d^{\frac{3}{4}\alpha} \right),$$

$$\Rightarrow \quad t_1 = \Theta \left(\frac{\log d}{\eta \|A\|_2} \right). \tag{8}$$

Since we pick $\eta \le \mathcal{O} \left(\frac{1}{d^{\alpha}} \right)$, we have that $t_1 \ge \Omega(\log d)$ and hence satisfies the condition in Lemma 5. We also have that

$$\lambda_1^{t_1} = \left(1 - \frac{\eta}{1-\beta} \|A\|_2 \right)^{t_1} \in \left[\tilde{\Omega} \left(d^{-\frac{3}{4}\alpha} \right), \tilde{\mathcal{O}} \left(d^{-\frac{\alpha}{2}} \right) \right], \quad \Rightarrow \quad \frac{\lambda_1^{t_1}}{\lambda_2^{t_1}} \le \tilde{\mathcal{O}} \left(d^{-\alpha} \right). \tag{9}$$

Now we prove that when $t = t_1$, W_1 will be approximately a rank-1 matrix.

Note that

$$W_1^{(t)} = W_1^{(t-1)} + \frac{\eta}{1-\beta} W_2^{(t-1)T} A + \frac{\eta}{1-\beta} r_1^{(t-1)}$$

$$= W_1^{(0)} + \frac{\eta}{1-\beta} \sum_{\tau=0}^{t-1} W_2^{(\tau)T} A + \frac{\eta}{1-\beta} \sum_{\tau=0}^{t-1} r_1^{(\tau)}$$

$$= W_1^{(0)} + \frac{\eta}{1-\beta} \left(C_1 \sum_{\tau=0}^{t-1} \lambda_1^\tau + C_2 \sum_{\tau=0}^{t-1} \lambda_2^\tau + \sum_{\tau=0}^{t-1} \lambda_2^\tau r_5^{(\tau)} \right)^T A + \frac{\eta}{1-\beta} \sum_{\tau=0}^{t-1} r_1^{(\tau)}$$

$$= W_1^{(0)} + \frac{\eta}{1-\beta} \sum_{\tau=0}^{t-1} r_1^{(\tau)} + \frac{\eta}{1-\beta} \left(C_1 \sum_{\tau=0}^{t-1} \lambda_1^\tau + \sum_{\tau=0}^{t-1} \lambda_2^\tau r_5^{(\tau)} \right)^T A + \frac{\eta}{1-\beta} \sum_{\tau=0}^{t-1} \lambda_2^\tau C_2^T A$$

$$:= R_1^{(t)} + \boldsymbol{u}^{(t)} \boldsymbol{v}^{(t)T},$$

where $R_1^{(t)} = W_1^{(0)} + \frac{\eta}{1-\beta} \sum_{\tau=0}^{t-1} r_1^{(\tau)} + \left(C_1 \sum_{\tau=0}^{t-1} \lambda_1^\tau + \sum_{\tau=0}^{t-1} \lambda_2^\tau r_5^{(\tau)} \right)^T \boldsymbol{v}^{(t)T}$, $\boldsymbol{v}^{(t)T} = \frac{\eta}{1-\beta} A$
and

$$\boldsymbol{u}^{(t)} = \sum_{\tau=0}^{t-1} \lambda_2^\tau C_2^T. \tag{10}$$

Moreover, we have that

$$W_2^{(t)} = C_1 \lambda_1^t + \left(C_2 + r_5^{(t)} \right) \lambda_2^t := R_2^{(t)} + c^{(t)} \boldsymbol{u}^{(t)T},$$

$$\text{where } R_2^{(t)} = C_1 \lambda_1^t + r_5^{(t)} \lambda_2^t, \quad c^{(t)} = \frac{\lambda_2^t}{\sum_{\tau=0}^{t-1} \lambda_2^\tau}.$$

For $t \le t_1$, by eq. (7), we get that w.h.p.,

$$\forall 1 \le i \le d: \quad \frac{\left| \sum_{\tau=0}^{t-1} \lambda_2^\tau r_{5i}^{(\tau)} v_j^{(t)} \right|}{\left| u_i^{(t)} v_j^{(t)} \right|} \le \frac{\mathcal{O}\left(\frac{1}{d^{\frac{3}{2}\alpha-1}} \right) \sum_{\tau=0}^{t} \lambda_2^\tau}{\tilde{\Omega}\left(\frac{1}{d^{\frac{5}{4}\alpha}} \right) \sum_{\tau=0}^{t} \lambda_2^\tau} \le \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha-1}} \right),$$

$$\frac{\left| \lambda_2^t r_{5i}^{(t)} \right|}{\left| c^{(t)} u_i^{(t)} \right|} = \frac{\left| r_{5i}^{(t)} \right|}{\left| C_{2i} \right|} \le \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha-1}} \right).$$

For $t \le t_1$, by Lemma 5, $\forall 1 \le i, j \le d: \left| r_1^{(t)}[i,j] \right| \le \mathcal{O}\left(\frac{1}{d^{\frac{3}{2}\alpha-1}} \right)$. Then we have that w.h.p.

$$\frac{\left| \frac{\eta}{1-\beta} \sum_{\tau=0}^{t-1} r_1^{(\tau)}[i,j] \right|}{\left| u_i^{(t)} v_j^{(t)} \right|} = \frac{\left| \sum_{\tau=0}^{t-1} r_1^{(\tau)}[i,j] \right|}{\left| \sum_{\tau=0}^{t-1} \lambda_2^\tau C_{2i} A_j \right|} \le \frac{\left| \sum_{\tau=0}^{t-1} r_1^{(\tau)}[i,j] \right|}{\left| \sum_{\tau=0}^{t-1} C_{2i} A_j \right|} \le \frac{\sum_{\tau=0}^{t-1} \mathcal{O}\left(\frac{1}{d^{\frac{3}{2}\alpha-1}} \right)}{\sum_{\tau=0}^{t-1} \tilde{\Omega}\left(\frac{1}{d^{\frac{5}{4}\alpha}} \right)}$$

$$= \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha-1}} \right).$$

Here we used $\forall i \in [d] : A_i = \Theta(1)$ by Assumption 1.

When $t = t_1$, by Lemma 7, eq. (8) and eq. (9), we have w.h.p. $\forall 1 \le i \le d$,

$$\frac{\left| C_{1i} \sum_{\tau=0}^{t_1-1} \lambda_1^\tau v_j^{(t_1)} \right|}{\left| u_i^{(t_1)} v_j^{(t_1)} \right|} = \frac{\left| C_{1i} \sum_{\tau=0}^{t_1-1} \lambda_1^\tau \right|}{\left| C_{2i} \sum_{\tau=0}^{t_1-1} \lambda_2^\tau \right|} \le \frac{\tilde{\mathcal{O}}\left(\frac{1}{d^\alpha} \right)}{\tilde{\Omega}\left(\frac{1}{d^{\frac{5}{4}\alpha}} \right)} \cdot \frac{\sum_{\tau=0}^{t_1} \lambda_1^\tau}{\sum_{\tau=0}^{t_1} \lambda_2^\tau} \le \tilde{\mathcal{O}}\left(d^{\frac{1}{4}\alpha} \right) \frac{1-\lambda_1^{t_1}}{\lambda_2^{t_1}-1} \le \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha}} \right),$$

$$\frac{\left| C_{1i} \lambda_1^{t_1} \right|}{\left| c^{(t_1)} u_i^{(t_1)} \right|} = \frac{\left| C_{1i} \lambda_1^{t_1} \right|}{\left| C_{2i} \lambda_2^{t_1} \right|} \le \frac{\tilde{\mathcal{O}}\left(\frac{1}{d^\alpha} \right)}{\tilde{\Omega}\left(\frac{1}{d^{\frac{5}{4}\alpha}} \right)} \cdot \frac{\lambda_1^{t_1}}{\lambda_2^{t_1}} \le \tilde{\mathcal{O}}\left(d^{\frac{1}{4}\alpha} \right) \tilde{\mathcal{O}}\left(d^{-\alpha} \right) \le \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3}{4}\alpha}} \right).$$

Finally, we have that w.h.p. $\forall 1 \le i, j \le d$,

$$
\begin{aligned}
\left|u_i^{(t_1)} v_j^{(t_1)}\right| &= \frac{\eta}{1-\beta} \sum_{\tau=0}^{t_1-1} \lambda_2^\tau |C_{2i} A_j| = \frac{\eta}{1-\beta} \cdot \frac{\lambda_2^{t_1}-1}{\lambda_2-1} |C_{2i} A_j| \stackrel{(i)}{=} \frac{\lambda_2^{t_1}-1}{\|A\|_2} |C_{2i} A_j| \\
&\ge \frac{\tilde{\Omega}\left(d^{\alpha/2}\right)}{\Theta\left(\sqrt{d}\right)} \tilde{\Omega}\left(\frac{1}{d^{\frac{5}{4}\alpha}}\right) = \tilde{\Omega}\left(\frac{1}{d^{\frac{3}{4}\alpha+\frac{1}{2}}}\right).
\end{aligned}
$$

Here in (i) we used $\lambda_2 = 1 + \frac{\eta}{1-\beta}\|A\|_2$. Using the Gaussian tail bound and union bound, we have w.h.p. $\forall 1 \le i, j \le d : \left|W_1^{(0)}[i,j]\right| = \tilde{\mathcal{O}}\left(\frac{1}{d^{2\alpha}}\right)$. Combining the above bounds yields that when $t = t_1$, w.h.p. $\forall 1 \le i, j \le d$,

$$
\begin{aligned}
\frac{\left|R_1^{(t_1)}[i,j]\right|}{\left|u_i^{(t_1)} v_j^{(t_1)}\right|} &\le \frac{\left|W_1^{(0)}[i,j]\right|}{\left|u_i^{(t_1)} v_j^{(t_1)}\right|} + \frac{\left|\frac{\eta}{1-\beta}\sum_{\tau=0}^{t_1} r_1^{(\tau)}[i,j]\right|}{\left|u_i^{(t_1)} v_j^{(t_1)}\right|} + \frac{\left|C_{1i}\sum_{\tau=0}^{t_1-1} \lambda_1^\tau v_j^{(t_1)}\right|}{\left|u_i^{(t_1)} v_j^{(t_1)}\right|} + \frac{\left|\sum_{\tau=0}^{t_1-1} \lambda_2^\tau r_5^{(\tau)} v_j^{(t_1)}\right|}{\left|u_i^{(t_1)} v_j^{(t_1)}\right|} \\
&= \frac{\tilde{\mathcal{O}}\left(\frac{1}{d^{2\alpha}}\right)}{\tilde{\Omega}\left(\frac{1}{d^{\frac{3}{4}\alpha+\frac{1}{2}}}\right)} + \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha-1}}\right) + \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha}}\right) + \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha-1}}\right) \\
&= \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha-1}}\right),
\end{aligned}
$$

$$
\frac{\left|R_{2i}^{(t_1)}\right|}{\left|c^{(t_1)} u_i^{(t_1)}\right|} \le \frac{\left|C_{1i} \lambda_1^{t_1}\right|}{\left|c^{(t_1)} u_i^{(t_1)}\right|} + \frac{\left|\lambda_2^{t_1} r_{5i}^{(t_1)}\right|}{\left|c^{(t_1)} u_i^{(t_1)}\right|} \le \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3}{4}\alpha}}\right) + \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha-1}}\right) = \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha-1}}\right).
$$

The definition of t_1 only considers the time $\left|w_{2i}^{(t)}\right|$ reaches $\frac{1}{d^{\alpha/2}}$. Now we prove that for $t \le t_1$, $\forall i, j \in [d] : \left|W_1^{(t)}[i,j]\right| = o\left(\frac{1}{d^{\alpha/2}}\right)$, and therefore $t_1 = T_1$.

To prove this, first note that for $t \le t_1$, we have $\forall j \in [d]$,

$$
\frac{\left|v_j^{(t)}\right|}{\left|c^{(t)}\right|} = \frac{\eta A_j}{1-\beta} \cdot \frac{\sum_{\tau=0}^{t-1} \lambda_2^\tau}{\lambda_2^t} = \frac{\eta A_j}{1-\beta} \frac{\lambda_2^t-1}{\lambda_2^t(\lambda_2-1)} = \frac{A_j(\lambda_2^t-1)}{\lambda_2^t \|A\|_2} \le \frac{A_j}{\|A\|_2} = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right).
$$

which yields that $\left|u_i^{(t)} v_j^{(t)}\right| \le \Theta\left(\frac{1}{\sqrt{d}}\right) \left|c^{(t)} u_i^{(t)}\right| = \mathcal{O}\left(\frac{1}{d^{\frac{\alpha+1}{2}}}\right)$.

Next by some small changes of the calculation above, we can get that for $t \le t_1$, $\left|R_1^{(t)}[i,j]\right| \le \mathcal{O}\left(\frac{1}{d^{\frac{\alpha+1}{2}}}\right)$. Then we have that w.h.p. for $t \le t_1$,

$$
\left|W_1^{(t)}[i,j]\right| \le \left|R_1^{(t)}[i,j]\right| + \left|u_i^{(t)} v_j^{(t)}\right| = \mathcal{O}\left(\frac{1}{d^{\frac{\alpha+1}{2}}}\right) = o\left(\frac{1}{d^{\alpha/2}}\right).
$$

Finally, we consider the loss. Since $\forall j \in [d] : \left(W_2^{(T_1)} W_1^{(T_1)}\right)_j - A_j = -\Theta(1)$, we know that $\bar{L}\left(W^{(T_1)}\right) = \Theta(d)$.

C.3 PROOF OF LEMMA 3

Eq. (10) and the fact that $t_1 = T_1$ (see the definition of t_1 in Section C.2) tell us that $\boldsymbol{u}^{(T_1)} = \sum_{\tau=0}^{T_1-1} \lambda_2^\tau C_2^T$.

Lemma 7 tells us that C_2 can be written as $C_2 := \frac{1}{2}(C_3 + C_4)$ where $C_{3i}, i \in [d]$ are i.i.d Gaussian random variables and that w.h.p. $\forall i \in [d] : \frac{|C_{4i}|}{|C_{3i}|} \le \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha-\frac{1}{2}}}\right)$.

Combining these two facts together finishes the proof.

C.4 PROOF OF LEMMA 4

Replacing t by $t-1$ in eq. (6), we get

$$W_2^{(t)} = W_2^{(t-1)} + \frac{\eta}{1-\beta} A W_1^{(t-1)T} + \frac{\eta}{1-\beta} r_2^{(t-1)}. \tag{11}$$

Eq. (6)-(11) and substituting eq. (5) yield

$$W_2^{(t+1)} - W_2^{(t)} = W_2^{(t)} - W_2^{(t-1)} + \frac{\eta^2}{(1-\beta)^2} \|A\|_2^2 W_2^{(t-1)} + \frac{\eta^2}{(1-\beta)^2} A r_1^{(t-1)T}$$
$$+ \frac{\eta}{1-\beta} \left(r_2^{(t)} - r_2^{(t-1)} \right),$$
$$\Rightarrow \quad W_2^{(t+1)} = 2W_2^{(t)} - \left(1 - \frac{\eta^2}{(1-\beta)^2} \|A\|_2^2 \right) W_2^{(t-1)} + r_3^{(t)},$$

where $r_3(t) := \frac{\eta^2}{(1-\beta)^2} A r_1^{(t-1)T} + \frac{\eta}{1-\beta} \left(r_2^{(t)} - r_2^{(t-1)} \right)$.

For the equation $x^2 - 2x + 1 - \frac{\eta^2}{(1-\beta)^2} \|A\|_2^2 = 0$, the roots are $\lambda_1 = 1 - \frac{\eta}{1-\beta} \|A\|_2$ and $\lambda_2 = 1 + \frac{\eta}{1-\beta} \|A\|_2$. We have that

$$W_2^{(t+1)} - \lambda_2 W_2^{(t)} = \lambda_1 \left(W_2^{(t)} - \lambda_2 W_2^{(t-1)} \right) + r_3^{(t)}$$
$$\Rightarrow \quad W_2^{(t)} - \lambda_2 W_2^{(t-1)} = \lambda_1^{t-1} \left(W_2^{(1)} - \lambda_2 W_2^{(0)} \right) + \sum_{\tau=1}^{t-1} \lambda_1^{t-1-\tau} r_3^{(\tau)}$$
$$:= \lambda_1^{t-1} \left(W_2^{(1)} - \lambda_2 W_2^{(0)} \right) + r_4^{(t)}.$$

We further have

$$W_2^{(t)} = \lambda_2^t W_2^{(0)} + \sum_{\tau=0}^{t-1} \lambda_2^{t-1-\tau} \lambda_1^\tau \left(W_2^{(1)} - \lambda_2 W_2^{(0)} \right) + \sum_{\tau=1}^{t} \lambda_2^{t-\tau} r_4^{(\tau)}$$
$$= \lambda_2^t W_2^{(0)} + \frac{\lambda_2^t - \lambda_1^t}{\lambda_2 - \lambda_1} \left(W_2^{(1)} - \lambda_2 W_2^{(0)} \right) + \sum_{\tau=1}^{t} \lambda_2^{t-\tau} r_4^{(\tau)}$$
$$= C_1 \lambda_1^t + C_2 \lambda_2^t + \sum_{\tau=1}^{t} \lambda_2^{t-\tau} r_4^{(\tau)}$$
$$= C_1 \lambda_1^t + \left(C_2 + r_5^{(t)} \right) \lambda_2^t,$$

where $r_5^{(t)} = \sum_{\tau=1}^{t} \lambda_2^{-\tau} r_4^{(\tau)}$, $C_1 = -\frac{W_2^{(1)} - \lambda_2 W_2^{(0)}}{\lambda_2 - \lambda_1}$ and $C_2 = \frac{W_2^{(1)} - \lambda_1 W_2^{(0)}}{\lambda_2 - \lambda_1}$.

C.5 PROOF OF LEMMA 5

Write $r_1^{(t)} = q_{11}^{(t)} + q_{12}^{(t)} + q_{13}^{(t)} + q_{14}^{(t)}$ where we have $q_{11}^{(t)} = -\beta^{t+1} W_2^{(t)T} A$, $q_{12}^{(t)} = (1-\beta) \sum_{\tau=0}^{t} \beta^{t-\tau} \left(W_2^{(\tau)T} - W_2^{(t)T} \right) A$, $q_{13}^{(t)} = -(1-\beta) \sum_{\tau=0}^{t} \beta^{t-\tau} W_2^{(\tau)T} W_2^{(\tau)} W_1^{(\tau)}$ and $q_{14}^{(t)} = -(1-\beta) \sum_{\tau=0}^{t} \beta^{t-\tau} D g_1^{(\tau)}$. And $r_2^{(t)} = q_{21}^{(t)} + q_{22}^{(t)} + q_{23}^{(t)} + q_{24}^{(t)}$, where $q_{21}^{(t)} = -\beta^{t+1} A W_1^{(t)T}$, $q_{22}^{(t)} = (1-\beta) \sum_{\tau=0}^{t} \beta^{t-\tau} A \left(W_1^{(\tau)T} - W_1^{(t)T} \right)$, $q_{23}^{(t)} = -(1-\beta) \sum_{\tau=0}^{t} \beta^{t-\tau} W_2^{(\tau)} W_1^{(\tau)} W_1^{(\tau)T}$ and $q_{24}^{(t)} = -(1-\beta) \sum_{\tau=0}^{t} \beta^{t-\tau} D g_2^{(\tau)}$.

We first bound $\left| q_{11}^{(t)}[i,j] \right|$ and $\left| q_{21,i}^{(t)} \right|$. For $t \leq T_1$, we have for all $i,j \in [d]$, $\left| w_{2i}^{(t)} A_j \right| \leq \mathcal{O} \left(\frac{1}{d^{\alpha/2}} \right)$ and $\left| \sum_{j=1}^{d} A_j W_1^{(t)}[i,j] \right| \leq \mathcal{O} \left(\frac{1}{d^{\alpha/2-1}} \right)$. Then if $\beta^{t+1} \leq \mathcal{O} \left(\frac{1}{d^\alpha} \right)$, i.e. $t = \Omega \left(\log d \right)$, we will have $\left| q_{11}^{(t)}[i,j] \right| = \beta^{t+1} \left| w_{2i}^{(t)} A_j \right| \leq \mathcal{O} \left(\frac{1}{d^{3\alpha/2}} \right)$ and $\left| q_{21,i}^{(t)} \right| = \beta^{t+1} \left| \sum_{j=1}^{d} A_j W_1^{(t)}[i,j] \right| \leq \mathcal{O} \left(\frac{1}{d^{3\alpha/2-1}} \right)$.

Next we bound $\left|q_{12}^{(t)}[i,j]\right|$ and $\left|q_{22,i}^{(t)}\right|$. Denote $E^{(t)} = W_2^{(t)}W_1^{(t)} - A$. Since $E^{(t)}$ and $W_2^{(t)}W_1^{(t)}$ are row vectors, we use $E_i^{(t)}$ and $\left(W_2^{(t)}W_1^{(t)}\right)_i$ to represent their i-th coordinates, respectively. For any $\tau \leq T_1$, we have that

$$\forall i \in [d]: \quad \left|\left(W_2^{(\tau)}W_1^{(\tau)}\right)_i\right| = \left|\sum_{j=1}^d w_{2j}^{(\tau)}W_1^{(\tau)}[j,i]\right| \leq \sum_{j=1}^d \left|w_{2j}^{(\tau)}\right|\left|W_1^{(\tau)}[j,i]\right| \leq \sum_{i=1}^d \frac{1}{d^\alpha} = \frac{1}{d^{\alpha-1}},$$

and thus $\forall i \in [d]: \left|E_i^{(\tau)}\right| = \mathcal{O}(1)$. Then we have for all $i,j \in [d]$,

$$\left|W_1^{(\tau+1)}[i,j] - W_1^{(\tau)}[i,j]\right| \leq \eta \sum_{k=0}^\tau \beta^{\tau-k}\left|w_{2i}^{(k)}E_j^{(k)}\right| \leq \eta \sum_{k=0}^\tau \beta^{\tau-k}\mathcal{O}\left(\frac{1}{d^{\alpha/2}}\right) = \eta\mathcal{O}\left(\frac{1}{d^{\alpha/2}}\right),$$

$$\left|w_{2i}^{(\tau+1)} - w_{2i}^{(\tau)}\right| \leq \eta \sum_{k=0}^\tau \beta^{\tau-k}\sum_{j=1}^d\left|E_j^{(k)}W_1^{(k)}[i,j]\right| \leq \eta \sum_{k=0}^\tau \beta^{\tau-k}\mathcal{O}\left(\frac{1}{d^{\alpha/2-1}}\right) = \eta\mathcal{O}\left(\frac{1}{d^{\alpha/2-1}}\right).$$

That gives us $\forall i,j \in [d]$,

$$\left|q_{12}^{(t)}[i,j]\right| \leq (1-\beta)\sum_{\tau=0}^t \beta^{t-\tau}\left|\left(w_{2i}^{(\tau)} - w_{2i}^{(t)}\right)A_j\right|$$

$$\leq \eta(1-\beta)\sum_{\tau=0}^t \beta^{t-\tau}(t-\tau)\mathcal{O}\left(\frac{1}{d^{\alpha/2-1}}\right) = \eta\mathcal{O}\left(\frac{1}{d^{\alpha/2-1}}\right),$$

$$\left|q_{22,i}^{(t)}\right| \leq (1-\beta)\sum_{\tau=0}^t \beta^{t-\tau}\sum_{j=1}^d\left|A_j\left(W_1^{(\tau)}[i,j] - W_1^{(t)}[i,j]\right)\right|$$

$$\leq \eta(1-\beta)\sum_{\tau=0}^t \beta^{t-\tau}(t-\tau)\mathcal{O}\left(\frac{1}{d^{\alpha/2-1}}\right) = \eta\mathcal{O}\left(\frac{1}{d^{\alpha/2-1}}\right).$$

Then we bound $\left|q_{13}^{(t)}[i,j]\right|$ and $\left|q_{23,i}^{(t)}\right|$. We have for $\forall i,j \in [d]$,

$$\left|q_{13}^{(t)}[i,j]\right| \leq (1-\beta)\sum_{\tau=0}^t \beta^{t-\tau}\left|w_{2i}^{(\tau)}\left(W_2^{(\tau)}W_1^{(\tau)}\right)_j\right| \leq (1-\beta)\sum_{\tau=0}^t \beta^{t-\tau}\frac{1}{d^{\frac{\alpha}{2}}}\cdot\frac{1}{d^{\alpha-1}} = \mathcal{O}\left(\frac{1}{d^{\frac{3}{2}\alpha-1}}\right),$$

$$\left|q_{23,i}^{(t)}\right| \leq (1-\beta)\sum_{\tau=0}^t \beta^{t-\tau}\sum_{j=1}^d\left|\left(W_2^{(t)}W_1^{(t)}\right)_j W_1^{(t)}[i,j]\right|$$

$$\leq (1-\beta)\sum_{\tau=0}^t \beta^{t-\tau}\sum_{i=1}^d \frac{1}{d^{\alpha-1+\frac{\alpha}{2}}} = \mathcal{O}\left(\frac{1}{d^{\frac{3}{2}\alpha-2}}\right).$$

Finally we use Lemma 31 to bound $\left|q_{14}^{(t)}[i,j]\right|$ and $\left|q_{24,i}^{(t)}\right|$. For $t \leq T_1$, the $M_1^{(t)}, M_2^{(t)}$ in Lemma 31 are upper bounded by $\frac{1}{d^{\frac{\alpha}{2}}}$. In the theorem we consider the training period before $T_{\text{SGD},2}$ so the time T in Lemma 31 is set as $T_{\text{SGD},2}$. In the following sections, we will prove that $T_{\text{SGD},2} \leq \mathcal{O}\left(\frac{d^\alpha \log(\sqrt{d/\epsilon})}{\eta}\right)$. Then by Lemma 31, we have with probability at least $1 - \frac{1}{d}$, for $\forall t \leq T_1$ and

$\forall i, j \in [d]$,

$$\left| Dg_1^{(t)}[i,j] \right| = \left| \tilde{g}_1^{(t)}[i,j] - g_1^{(t)}[i,j] \right| \le \mathcal{O}\left(\frac{1}{d^{\frac{3\alpha}{2}-3}} \sigma \sqrt{\frac{d^{\alpha+1}}{\eta}} \log \frac{d}{\epsilon} \right) + \mathcal{O}\left(\frac{1}{d^{\frac{\alpha}{2}}} \sigma \sqrt{\frac{d^{\alpha+2}}{\eta}} \log \frac{d}{\epsilon} \right)$$

$$\le \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{\alpha}{2}}} \sigma \sqrt{\frac{d^{\alpha+2}}{\eta}} \right)$$

$$\left| Dg_{2i}^{(t)} \right| = \left| \tilde{g}_{2i}^{(t)} - g_{2i}^{(t)} \right| \le \mathcal{O}\left(\frac{1}{d^{\frac{3\alpha}{2}-4}} \sigma \sqrt{\frac{d^{\alpha+1}}{\eta}} \log \frac{d}{\epsilon} \right) + \mathcal{O}\left(\frac{1}{d^{\frac{\alpha}{2}-1}} \sigma \sqrt{\frac{d^{\alpha+2}}{\eta}} \log \frac{d}{\epsilon} \right)$$

$$\le \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{\alpha}{2}-1}} \sigma \sqrt{\frac{d^{\alpha+2}}{\eta}} \right)$$

By picking $\sigma \le \frac{\eta^{3/2}}{d^{\alpha/2+1}}$, we have w.h.p. for $\forall t \le T_1$ and $\forall i, j \in [d]$, $\left| Dg_1^{(t)}[i,j] \right| \le \eta \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{\alpha}{2}}} \right)$ and $\left| Dg_{2i}^{(t)} \right| \le \eta \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{\alpha}{2}-1}} \right)$, which yields

$$\left| q_{14}^{(t)}[i,j] \right| \le (1-\beta) \sum_{\tau=0}^{t} \beta^{t-\tau} \left| Dg_1^{(\tau)}[i,j] \right| \le (1-\beta) \sum_{\tau=0}^{t} \beta^{t-\tau} \eta \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{\alpha}{2}}} \right) = \eta \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{\alpha}{2}}} \right)$$

$$\left| q_{24,i}^{(t)} \right| \le (1-\beta) \sum_{\tau=0}^{t} \beta^{t-\tau} \left| Dg_{2i}^{(\tau)} \right| \le (1-\beta) \sum_{\tau=0}^{t} \beta^{t-\tau} \eta \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{\alpha}{2}-1}} \right) = \eta \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{\alpha}{2}-1}} \right)$$

Combining the above bounds together and substituting $\eta \le \mathcal{O}\left(\frac{1}{d^\alpha} \right)$ complete the proof.

C.6 PROOF OF LEMMA 6

Based on the bound in Lemma 5, we have

$$\left| r_{3i}^{(t)} \right| = \left| \frac{\eta^2}{(1-\beta)^2} \sum_{j=1}^{d} A_j r_1^{(t-1)}[i,j] + \frac{\eta}{1-\beta} \left(r_{2i}^{(t)} - r_{2i}^{(t-1)} \right) \right|$$

$$\le \frac{\eta^2}{(1-\beta)^2} \sum_{j=1}^{d} \left| A_j r_1^{(t-1)}[i,j] \right| + \frac{\eta}{1-\beta} \left| r_{2i}^{(t)} \right| + \frac{\eta}{1-\beta} \left| r_{2i}^{(t-1)} \right|$$

$$\le \eta^2 \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3}{2}\alpha-2}} \right) + 2\eta \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3}{2}\alpha-2}} \right)$$

$$= \eta \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3}{2}\alpha-2}} \right).$$

Since $\lambda_1 = 1 - \frac{\eta}{1-\beta}\|A\|_2, \lambda_2 = 1 + \frac{\eta}{1-\beta}\|A\|_2$, and note that $\|A\|_2 = \Theta\left(\sqrt{d} \right)$, we have that

$$\left| r_{4i}^{(t)} \right| = \left| \sum_{\tau=1}^{t-1} \lambda_1^{t-1-\tau} r_{3i}^{(\tau)} \right| \le \eta \sum_{\tau=1}^{t-1} \lambda_1^{t-1-\tau} \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3}{2}\alpha-2}} \right) \le \frac{\eta}{1-\lambda_1} \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3}{2}\alpha-2}} \right) = \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3}{2}(\alpha-1)}} \right),$$

$$\left| r_{5i}^{(t)} \right| = \left| \sum_{\tau=1}^{t} \lambda_2^{-\tau} r_{4i}^{(\tau)} \right| \le \eta \sum_{\tau=1}^{t} \lambda_2^{-\tau} \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3}{2}(\alpha-1)}} \right) \le \frac{\eta}{\lambda_2-1} \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3}{2}(\alpha-1)}} \right) = \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3}{2}\alpha-1}} \right).$$

C.7 PROOF OF LEMMA 7

For the equation $x^2 - 2x + 1 - \frac{\eta^2}{(1-\beta)^2}\|A\|_2^2 = 0$, the roots are $\lambda_1 = 1 - \frac{\eta}{1-\beta}\|A\|_2$ and $\lambda_2 = 1 + \frac{\eta}{1-\beta}\|A\|_2$, which gives us

$$
\begin{aligned}
C_2 &= \frac{W_2^{(1)} - \lambda_1 W_2^{(0)}}{\lambda_2 - \lambda_1} \\
&= \frac{W_2^{(0)} + \eta A W_1^{(0)T} + \eta \tilde{r}_2^{(0)} - W_2^{(0)} + \frac{\eta}{1-\beta}\|A\|_2 W_2^{(0)}}{\frac{2\eta}{1-\beta}\|A\|_2} \\
&= \frac{1}{2}W_2^{(0)} + \frac{1-\beta}{2\|A\|_2} A W_1^{(0)T} + \frac{1-\beta}{2\|A\|_2}\tilde{r}_2^{(0)},
\end{aligned}
\tag{12}
$$

where $\tilde{r}_2^{(0)} = -W_2^{(0)} W_1^{(0)} W_1^{(0)T} - Dg_2^{(0)}$. Note that this is slightly different from the definition of $r_2^{(0)}$ in eq. (6). Now let's bound the i-th coordinate of $\tilde{r}_2^{(0)}$.

In Section C.5 we have shown that w.h.p. for $\forall t \leq T_1$ and $\forall i, j \in [d]$, $\left|Dg_{2i}^{(t)}\right| \leq \eta\tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{\alpha}{2}-1}}\right) = \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3\alpha}{2}-1}}\right)$, which also applies to $t = 0$. Using the Gaussian tail bound and union bound, w.p. at least $1 - \delta$, for ever $1 \leq i, j \leq d$, we have that

$$
\left|w_{2i}^{(0)}\right| \leq \sqrt{\frac{2}{d^{2\alpha}}\log\frac{2d}{\delta}}, \quad \left|W_1^{(0)}[i,j]\right| \leq \sqrt{\frac{2}{d^{4\alpha}}\log\frac{2d^2}{\delta}}.
$$

Then we have that w.p. at least $1 - \delta$, $\forall 1 \leq i, j \leq d$:,

$$
\begin{aligned}
\left|\left(W_2^{(0)} W_1^{(0)}\right)_i\right| &= \left|\sum_{j=1}^d w_{2j}^{(0)} W_1^{(0)}[j,i]\right| \leq \sum_{j=1}^d \left|w_{2j}^{(0)}\right|\left|W_1^{(0)}[j,i]\right| \\
&\leq \sum_{i=1}^d \sqrt{\frac{2}{d^{2\alpha}}\log\frac{2d}{\delta}}\sqrt{\frac{2}{d^{4\alpha}}\log\frac{2d^2}{\delta}} \leq \frac{2}{d^{3\alpha-1}}\log\frac{2d^2}{\delta},
\end{aligned}
$$

$$
\begin{aligned}
\Rightarrow \quad \left|\tilde{r}_{2i}^{(0)}\right| &\leq \sum_{j=1}^d \left|\left(W_2^{(0)}W_1^{(0)}\right)_j\right|\left|W_1^{(0)}[i,j]\right| + \left|Dg_{2i}^{(0)}\right| \\
&\leq \sum_{i=1}^d \frac{2}{d^{3\alpha-1}}\log\frac{2d^2}{\delta}\sqrt{\frac{2}{d^{4\alpha}}\log\frac{2d^2}{\delta}} + \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3\alpha}{2}-1}}\right) = \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3\alpha}{2}-1}}\right).
\end{aligned}
\tag{13}
$$

Next, we bound the i-th coordinate of $W_2^{(0)} + \frac{1-\beta}{\|A\|_2} A W_1^{(0)T}$, i.e. $w_{2i}^{(0)} + \frac{1-\beta}{\|A\|_2} A\left(W_1^{(0)}[i,:]\right)^T$.

By independence under Assumption 2, we have that

$$
\begin{aligned}
\text{Var}\left(w_{2i}^{(0)} + \frac{1-\beta}{\|A\|_2} A\left(W_1^{(0)}[i,:]\right)^T\right) &= \text{Var}\left(w_{2i}^{(0)}\right) + \frac{(1-\beta)^2}{\|A\|_2^2}\sum_{j=1}^d A_j^2 \text{Var}\left(W_1^{(0)}[i,j]\right) \\
&= \frac{1}{d^{2\alpha}} + \frac{(1-\beta)^2}{\|A\|_2^2}\sum_{i=1}^d A_j^2 \frac{1}{d^{4\alpha}} = \mathcal{O}\left(\frac{1}{d^{2\alpha}}\right).
\end{aligned}
$$

Using the Gaussian tail bound and union bound, w.p. at least $1 - \delta$, for ever $1 \leq i \leq d$, we have that

$$
\left|w_{2i}^{(0)} + \frac{1-\beta}{\|A\|_2} A\left(W_1^{(0)}[i,:]\right)^T\right| \leq \mathcal{O}\left(\sqrt{\frac{1}{d^{2\alpha}}\log\frac{d}{\delta}}\right) = \tilde{\mathcal{O}}\left(\frac{1}{d^\alpha}\right).
$$

Since for $X \sim \mathcal{N}(0, \sigma^2)$, we have that $P(|X| \leq t) \leq \frac{2t}{\sqrt{2\pi}\sigma}$, then for a fixed i,

$$
P\left(\left|w_{2i}^{(0)} + \frac{1-\beta}{\|A\|_2} A\left(W_1^{(0)}[i,:]\right)^T\right| \leq \frac{1}{d^{\frac{5}{4}\alpha}}\right) \leq \mathcal{O}\left(\frac{2/d^{\frac{5}{4}\alpha}}{\sqrt{2\pi}\cdot\sqrt{1/d^{2\alpha}}}\right) = \Theta\left(\frac{1}{d^{\frac{\alpha}{4}}}\right).
$$

Then by union bound, we have that w.p. at least $1 - \frac{1}{d^{\frac{\alpha}{4}-1}}$, for every $1 \le i \le d$,

$$\left| w_{2i}^{(0)} + \frac{1-\beta}{\|A\|_2} A \left(W_1^{(0)}[i,:] \right)^T \right| \ge \Theta \left(\frac{1}{d^{\frac{5}{4}\alpha}} \right).$$

Now define $C_3 := W_2^{(0)} + \frac{1-\beta}{\|A\|_2} A W_1^{(0)T}$ and $C_4 := \frac{1-\beta}{2\|A\|_2} \tilde{r}_{2i}^{(0)}$. We get that $C_{3i}, i \in [d]$ are i.i.d Gaussian random variables and that $C_2 = \frac{1}{2}(C_3 + C_4)$, where w.h.p. for all $i \in [d]$,

$$|C_{3i}| \le \tilde{\mathcal{O}} \left(\frac{1}{d^\alpha} \right), \quad |C_{3i}| \ge \Theta \left(\frac{1}{d^{\frac{5}{4}\alpha}} \right), \quad |C_{4i}| \overset{(i)}{\le} \tilde{\mathcal{O}} \left(\frac{1}{d^{\frac{3\alpha}{2}-\frac{1}{2}}} \right), \tag{14}$$

where (i) follows from eq. (13) and the fact that $\|A\|_2 = \sqrt{d}$. Then we get that w.h.p.

$$\forall i \in [d]: \frac{|C_{4i}|}{|C_{3i}|} \le \frac{\tilde{\mathcal{O}} \left(\frac{1}{d^{\frac{3\alpha}{2}-\frac{1}{2}}} \right)}{\Omega \left(\frac{1}{d^{\frac{5}{4}\alpha}} \right)} = \tilde{\mathcal{O}} \left(\frac{1}{d^{\frac{1}{4}\alpha-\frac{1}{2}}} \right).$$

Substituting eq. (14) into eq. (12), we get that w.h.p.,

$$|C_{2i}| = \Theta \left(\left| w_{2i}^{(0)} + \frac{1-\beta}{\|A\|_2} A \left(W_1^{(0)}[i,:] \right)^T \right| \right) \in \left[\tilde{\Omega} \left(\frac{1}{d^{\frac{5}{4}\alpha}} \right), \tilde{\mathcal{O}} \left(\frac{1}{d^\alpha} \right) \right].$$

Similarly, note that

$$\begin{aligned}
C_1 &= -\frac{W_2^{(1)} - \lambda_2 W_2^{(0)}}{\lambda_2 - \lambda_1} \\
&= -\frac{W_2^{(0)} + \eta A W_1^{(0)T} + \eta \tilde{r}_2^{(0)} - W_2^{(0)} - \frac{\eta}{1-\beta}\|A\|_2 W_2^{(0)}}{\frac{2\eta}{1-\beta}\|A\|_2} \\
&= \frac{1}{2} W_2^{(0)} - \frac{1-\beta}{2\|A\|_2} A W_1^{(0)T} - \frac{1-\beta}{2\|A\|_2} \tilde{r}_2^{(0)},
\end{aligned}$$

we can use the same techniques to get that i) w.p. at least $1 - \delta, \forall i \in [d] : |C_{1i}| \le \tilde{\mathcal{O}} \left(\frac{1}{d^\alpha} \right)$, ii) w.p. at least $1 - \delta - \frac{1}{d^{\frac{\alpha}{4}-1}}, \forall i \in [d], |C_{1i}| \ge \tilde{\Omega} \left(\frac{1}{d^{\frac{5}{4}\alpha}} \right)$.

C.8 Proof of Lemma 2

The proof in Section C.2 tells us that at the end of the first phase (when $t = T_1$),

$$\begin{aligned}
W_1^{(T_1)} &= \boldsymbol{u}^{(T_1)} \boldsymbol{v}^{(T_1)T} + R_1^{(T_1)}, \\
W_2^{(T_1)} &= c^{(T_1)} \boldsymbol{u}^{(T_1)T} + R_2^{(T_1)T}, \\
\text{where } \boldsymbol{v}^{(T_1)T} &= \frac{\eta A}{1-\beta}, \quad c^{(T_1)} = \frac{\lambda_2^{T_1}}{\sum_{\tau=0}^{T_1-1} \lambda_2^\tau}.
\end{aligned} \tag{15}$$

Denote the i-th coordinate of $\boldsymbol{u}^{(t)}, \boldsymbol{v}^{(t)}, R_2^{(t)}$ as $u_i^{(t)}, v_i^{(t)}, R_{2i}^{(t)}$, respectively. Denote the (i,j)-th element of $R_1^{(t)}$ as $R_1^{(t)}[i,j]$. For $t \ge T_1$, we prove by induction that,

$$\begin{aligned}
W_1^{(t)} &= \boldsymbol{u}^{(T_1)} \boldsymbol{v}^{(t)T} + R_1^{(t)}, \\
W_2^{(t)} &= c^{(t)} \boldsymbol{u}^{(T_1)T} + R_2^{(t)T},
\end{aligned} \tag{16}$$

where

$$\begin{aligned}
\boldsymbol{v}^{(t+1)T} &= \boldsymbol{v}^{(t)T} - \eta_t c^{(t)} E^{(t)}, \\
R_1^{(t+1)} &= R_1^{(t)} - \eta_t R_2^{(t)} E^{(t)} + r_1^{(t)}, \\
c^{(t+1)} &= c^{(t)} - \eta_t E^{(t)} \boldsymbol{v}^{(t)}, \\
R_2^{(t+1)T} &= R_2^{(t)T} - \eta_t E^{(t)} R_1^{(t)T} + r_2^{(t)},
\end{aligned}$$

with $E^{(t)} := W_2^{(t)} W_1^{(t)} - A$, $\eta_t = \eta \sum_{\tau=0}^{t} \beta^{t-\tau}$, $r_1^{(t)} :=$ $\eta \sum_{\tau=0}^{t} \beta^{t-\tau} \left(W_2^{(t)T} E^{(t)} - W_2^{(\tau)T} E^{(\tau)} \right) - \eta \sum_{\tau=0}^{t} \beta^{t-\tau} Dg_1^{(\tau)}$ and $r_2^{(t)} =$ $\eta \sum_{\tau=0}^{t} \beta^{t-\tau} \left(E^{(t)} W_1^{(t)T} - E^{(\tau)} W_1^{(\tau)T} \right) - \eta \sum_{\tau=0}^{t} \beta^{t-\tau} Dg_2^{(\tau)}$. Note that the $r_1^{(t)}$ and $r_2^{(t)}$ here are different from those defined in Section C.2, but we abuse the notation and still use $r_1^{(t)}$ and $r_2^{(t)}$ to represent the error terms.

The base case is already given by eq. (15).

Suppose our lemma holds for t, then for $t+1$, using the same techniques as in eq. (5) and eq. (6), we have that

$$W_1^{(t+1)} = W_1^{(t)} - \eta \sum_{\tau=0}^{t} \beta^{t-\tau} W_2^{(\tau)T} E^{(\tau)} - \eta \sum_{\tau=0}^{t} \beta^{t-\tau} Dg_1^{(\tau)}$$
$$= W_1^{(t)} - \eta_t W_2^{(t)T} E^{(t)} + r_1^{(t)},$$

$$W_2^{(t+1)} = W_2^{(t)} - \eta \sum_{\tau=0}^{t} \beta^{t-\tau} E^{(\tau)} W_1^{(\tau)T} - \eta \sum_{\tau=0}^{t} \beta^{t-\tau} Dg_2^{(\tau)}$$
$$= W_2^{(t)} - \eta_t E^{(t)} W_1^{(t)T} + r_2^{(t)},$$

Plugging in the inductive hypothesis yields

$$W_1^{(t+1)} = W_1^{(t)} - \eta_t W_2^{(t)T} E^{(t)} + r_1^{(t)}$$
$$= \boldsymbol{u}^{(T_1)} \boldsymbol{v}^{(t)T} + R_1^{(t)} - \eta_t \left(c^{(t)} \boldsymbol{u}^{(T_1)} + R_2^{(t)} \right) E^{(t)} + r_1^{(t)}$$
$$= \boldsymbol{u}^{(T_1)} \left(\boldsymbol{v}^{(t)T} - \eta_t c^{(t)} E^{(t)} \right) + R_1^{(t)} - \eta_t R_2^{(t)} E^{(t)} + r_1^{(t)},$$

$$W_2^{(t+1)} = W_2^{(t)} - \eta_t E^{(t)} W_1^{(t)T} + r_2^{(t)}$$
$$= c^{(t)} \boldsymbol{u}^{(T_1)T} + R_2^{(t)T} - \eta_t E^{(t)} \left(\boldsymbol{v}^{(t)} \boldsymbol{u}^{(T_1)T} + R_1^{(t)T} \right) + r_2^{(t)}$$
$$= \left(c^{(t)} - \eta_t E^{(t)} \boldsymbol{v}^{(t)} \right) \boldsymbol{u}^{(T_1)T} + R_2^{(t)T} - \eta_t E^{(t)} R_1^{(t)T} + r_2^{(t)}.$$

It implies that our lemma holds for $t+1$, which completes the proof.

Now we analyze the error terms $\left| R_1^{(t)}[i,j] \right|$ and $\left| R_{2i}^{(t)} \right|$. Eq. (15) tells us that $c^{(T_1)}$ and $\forall i \in [d], v_i^{(T_1)}$ are all positive. We first prove by induction that for all $T_1 \le t \le T_2$, $c^{(t)} > 0, \forall i \in [d], v_i^{(t)} > 0$.

The above discussion already proves the base case. Suppose at time t, we have $c^{(t)} > 0, \forall i \in [d], v_i^{(t)} > 0$. Note that when $T_1 \le t < T_2, \forall i \in [d] : E_i^{(t)} \le 0$, then for $t+1$,

$$v_i^{(t+1)} = v_i^{(t)} - \eta_t c^{(t)} E_i^{(t)} > 0,$$

$$c^{(t+1)} = c^{(t)} - \eta_t \sum_{i=1}^{d} E_i^{(t)} v_i^{(t)} > 0.$$

Therefore by induction, we have proved that for all $T_1 \le t \le T_2$, $c^{(t)} > 0, \forall i \in [d], v_i^{(t)} > 0$.

Now we prove that for all $T_1 \le t \le T_2$,

$$\forall 1 \le i, j \le d: \quad 0 \le \frac{\left| R_1^{(t)}[i,j] \right|}{\left| u_i^{(T_1)} \right| \left| v_j^{(t)} \right|} \le \delta_i + \sum_{\tau=T_1}^{t-1} \epsilon_i^{(\tau)}, \quad 0 \le \frac{\left| R_{2i}^{(t)} \right|}{c^{(t)} \left| u_i^{(T_1)} \right|} \le \delta_i + \sum_{\tau=T_1}^{t-1} \epsilon_i^{(\tau)}, \quad (17)$$

where

$$\delta_i := \max \left\{ \max_j \frac{\left| R_1^{(T_1)}[i,j] \right|}{\left| u_i^{(T_1)} \right| \left| v_j^{(T_1)} \right|}, \frac{\left| R_{2i}^{(T_1)} \right|}{c^{(T_1)} \left| u_i^{(T_1)} \right|} \right\}, \quad \epsilon_i^{(t)} := \max \left\{ \max_j \frac{\left| r_1^{(t)}[i,j] \right|}{\left| u_i^{(T_1)} \right| \left| v_j^{(t)} \right|}, \frac{\left| r_{2i}^{(t)} \right|}{c^{(t)} \left| u_i^{(T_1)} \right|} \right\}.$$

The left hand sides of the inequalities are trivial since we have proved that $c^{(t)} > 0, \forall i \in [d], v_i^{(t)} > 0$ for all $T_1 \leq t \leq T_2$. Now we prove the right hand sides by induction.

The base case is already verified by the definition of δ_i. Suppose eq.(17) holds for $T_1 \leq t < T_2$. Then for $t + 1$, using $\forall i \in [d] : E_i^{(t)} \leq 0$ and $v^{(t+1)} \geq v^{(t)}, c^{(t+1)} \geq c^{(t)}$, we can get that $\forall 1 \leq i, j \leq d$

$$\frac{\left|R_1^{(t+1)}[i,j]\right|}{\left|u_i^{(T_1)}\right| v_j^{(t+1)}} = \frac{\left|R_1^{(t+1)}[i,j]\right| \frac{1}{\left|u_i^{(T_1)}\right|}}{v_j^{(t+1)}} \leq \frac{\left|R_1^{(t)}[i,j]\right| \frac{1}{\left|u_i^{(T_1)}\right|} + \eta_t \left|R_{2i}^{(t)}\right| \frac{1}{\left|u_i^{(T_1)}\right|} \left(-E_j^{(t)}\right)}{v_j^{(t)} + \eta_t c^{(t)} \left(-E_j^{(t)}\right)} + \frac{\left|r_1^{(t)}[i,j]\right|}{\left|u_i^{(T_1)}\right| v_j^{(t)}}$$

$$\leq \frac{\left(\delta_i + \sum_{\tau=T_1}^{t-1} \epsilon_i^{(\tau)}\right) v_j^{(t)} + \eta_t \left(\delta_i + \sum_{\tau=T_1}^{t-1} \epsilon_i^{(\tau)}\right) c^{(t)} \left(-E_j^{(t)}\right)}{v_j^{(t)} + \eta_t c^{(t)} \left(-E_j^{(t)}\right)} + \epsilon_i^{(t)} = \delta_i + \sum_{\tau=T_1}^{t} \epsilon_i^{(\tau)}.$$

Similarly, we have that $\forall 1 \leq i \leq d$

$$\frac{\left|R_{2i}^{(t+1)}\right|}{c^{(t+1)} \left|u_i^{(T_1)}\right|} = \frac{\left|R_{2i}^{(t+1)}\right| \frac{1}{\left|u_i^{(T_1)}\right|}}{c^{(t+1)}} \leq \frac{\left|R_{2i}^{(t)}\right| \frac{1}{\left|u_i^{(T_1)}\right|} + \eta_t \sum_{j=1}^{d} \left(-E_j^{(t)}\right) \left|R_1^{(t)}[i,j]\right| \frac{1}{\left|u_i^{(T_1)}\right|}}{c^{(t)} + \eta_t \sum_{j=1}^{d} \left(-E_j^{(t)}\right) v_j^{(t)}} + \frac{\left|r_{2i}^{(t)}\right|}{\left|u_i^{(T_1)}\right| c^{(t)}}$$

$$\leq \frac{\left(\delta_i + \sum_{\tau=T_1}^{t-1} \epsilon_i^{(\tau)}\right) c^{(t)} + \eta_t \left(\delta_i + \sum_{\tau=T_1}^{t-1} \epsilon_i^{(\tau)}\right) \sum_{j=1}^{d} \left(-E_j^{(t)}\right) v_j^{(t)}}{c^{(t)} + \eta_t \sum_{j=1}^{d} \left(-E_j^{(t)}\right) v_j^{(t)}} + \epsilon_i^{(t)}$$

$$= \delta_i + \sum_{\tau=T_1}^{t} \epsilon_i^{(\tau)}.$$

Therefore by induction, eq. (17) holds for all t in the second phase.

So far we have proved the rank 1 structure stated in Lemma 2. The remaining part of the proof is given by the following lemma, whose proof is deferred to Section C.9.

Lemma 8. *Under Assumption 1, 2 and 3, suppose $\sigma \leq \frac{\eta^{3/2}}{d^{\alpha/2+1}}$. By picking $\eta \leq \mathcal{O}\left(\frac{\epsilon}{d^{\frac{7\alpha}{4}+4}}\right)$, we have that for $T_1 \leq t \leq \min\{T_2, T_3\}$,*

$$\forall 1 \leq i, j \leq d: \quad 0 \leq \frac{\left|R_1^{(t)}[i,j]\right|}{\left|u_i^{(T_1)}\right| v_j^{(t)}} \leq \tilde{\mathcal{O}}(\epsilon_0), \quad 0 \leq \frac{\left|R_{2i}^{(t)}\right|}{c^{(t)} \left|u_i^{(T_1)}\right|} \leq \tilde{\mathcal{O}}(\epsilon_0). \tag{18}$$

Moreover, when $t = \min\{T_2, T_3\}$, we have $\left\|E^{(t)}\right\|_2^2 = \mathcal{O}(\epsilon_0 d)$.

C.9 Proof of Lemma 8

We first have the following lemma which describes the structure of $v^{(t)}$ for $t \geq T_1$.

Lemma 9. *Under Assumption 1, 2 and 3, for $t \geq T_1$, we can write $v^{(t)T}$ as $v^{(t)T} = a^{(t)} A + R_v^{(t)T}$, with $a^{(T_1)} = \frac{\eta}{1-\beta}, R_v^{(T_1)T} = [0, 0, ..., 0]$, and*

$$a^{(t+1)} = \left(1 - \eta_t c^{(t)} d^{(t)}\right) a^{(t)} + \eta_t c^{(t)},$$

$$R_v^{(t+1)} = \left(1 - \eta_t c^{(t)} d^{(t)}\right) R_v^{(t)} - \eta_t c^{(t)} R_3^{(t)},$$

where $d^{(t)} := c^{(t)} \left\|u^{(T_1)}\right\|^2 + R_2^{(t)T} u^{(T_1)}, \quad R_3^{(t)T} := c^{(t)} u^{(T_1)T} R_1^{(t)} + R_2^{(t)T} R_1^{(t)}.$

Moreover, we have that

$$W_2^{(t)} W_1^{(t)} = d^{(t)} v^{(t)T} + R_3^{(t)T} = d^{(t)} a^{(t)} A + d^{(t)} R_v^{(t)T} + R_3^{(t)T}. \tag{19}$$

We prove Lemma 8 by induction. Denote the i-th coordinate of $R_3^{(t)}$ and $R_v^{(t)}$ as $R_{3i}^{(t)}$ and $R_{vi}^{(t)}$, respectively. The following lemmas constitute the inductive part.

Lemma 10. *Under Assumption 1, 2 and 3, suppose $\sigma \leq \frac{\eta^{3/2}}{d^{\alpha/2+1}}$ and pick $\eta \leq \mathcal{O}\left(\frac{1}{d^\alpha}\right)$. Consider some t such that $T_1 \leq t < \min\{T_2, T_3\}$. Suppose for all $T_1 \leq \tau \leq t$, we have $\forall i,j \in [d] : \left|w_{2i}^{(\tau)}\right| \leq \mathcal{O}\left(d^{1/4}\right), \left|W_1^{(\tau)}[i,j]\right| \leq \mathcal{O}\left(\frac{1}{d^{1/4}}\right)$, then we have that $\forall i,j \in [d] : \left|r_1^{(t)}[i,j]\right| = \tilde{\mathcal{O}}\left(\eta^2 d^{11/4}\right), \left|r_{2i}^{(t)}\right| = \tilde{\mathcal{O}}\left(\eta^2 d^{13/4}\right)$. Moreover, we can get that $\forall i \in [d] : \epsilon_i^{(t)} = \tilde{\mathcal{O}}\left(\eta^2 d^{\frac{3}{4}\alpha + \frac{13}{4}}\right)$, where $\epsilon_i^{(t)}$ is defined in eq. (17).*

Lemma 11. *Under the conditions of Lemma 10 and pick $\eta \leq \mathcal{O}\left(\frac{\epsilon}{d^{\frac{7\alpha}{4}+4}}\right)$, we have that at time $t+1$,*

$$\forall 1 \leq i,j \leq d : \quad 0 \leq \frac{\left|R_1^{(t+1)}[i,j]\right|}{\left|u_i^{(T_1)}\right|\left|v_j^{(t+1)}\right|} \leq \tilde{\mathcal{O}}(\epsilon_0), \quad 0 \leq \frac{\left|R_{2i}^{(t+1)}\right|}{c^{(t+1)}\left|u_i^{(T_1)}\right|} \leq \tilde{\mathcal{O}}(\epsilon_0).$$

where ϵ_0 is defined in Definition 2.

Lemma 12. *Under the conditions of Lemma 10 and pick $\eta \leq \mathcal{O}\left(\frac{\epsilon}{d^{\frac{7\alpha}{4}+4}}\right)$, we have that at time $t+1$,*

$$0 \leq \frac{\left|R_2^{(t+1)T}\boldsymbol{u}^{(T_1)}\right|}{c^{(t+1)}\left\|\boldsymbol{u}^{(T_1)}\right\|^2} \leq \tilde{\mathcal{O}}(\epsilon_0), \quad \forall j \in [d] : \quad 0 \leq \frac{\left|R_{3j}^{(t+1)}\right|}{c^{(t+1)}\left\|\boldsymbol{u}^{(T_1)}\right\|^2 v_j^{(t)}} \leq \tilde{\mathcal{O}}(\epsilon_0).$$

Moreover,

$$\forall j \in [d] : \frac{\left|R_{3j}^{(t+1)}\right|}{A_j} \leq \tilde{\mathcal{O}}(\epsilon_0). \tag{20}$$

Lemma 13. *Under the conditions of Lemma 10 and pick $\eta \leq \mathcal{O}\left(\frac{\epsilon}{d^{\frac{7\alpha}{4}+4}}\right)$, if we further suppose that $\forall j \in [d] : \frac{v_j^{(t)}}{c^{(t)}} = \Theta\left(\frac{1}{\sqrt{d}}\right), \frac{\left|R_{3j}^{(t)}\right|}{A_j}$ and $\frac{\left|R_{vj}^{(t)}\right|}{a^{(t)} A_j}$ are of order $\tilde{\mathcal{O}}(\epsilon_0)$, then we have that at time $t+1$,*

(A) $\forall i,j \in [d] : \frac{E_j^{(t)}}{E_i^{(t)}} = \Theta(1)$,

(B) $\forall j \in [d] : \frac{v_j^{(t+1)}}{c^{(t+1)}} = \Theta\left(\frac{1}{\sqrt{d}}\right)$,

(C) $\forall i,j \in [d] : \left|w_{2i}^{(t+1)}\right| \leq \mathcal{O}\left(d^{1/4}\right), \left|W_1^{(t+1)}[i,j]\right| \leq \mathcal{O}\left(\frac{1}{d^{1/4}}\right)$,

(D) $\forall j \in [d], \frac{\left|R_{3j}^{(t+1)}\right|}{A_j}$ and $\frac{\left|R_{vj}^{(t+1)}\right|}{a^{(t+1)} A_j}$ are of order $\tilde{\mathcal{O}}(\epsilon_0)$.

By combining Lemma 10, 11 and 13, we can prove by induction that for all $T_1 \leq t \leq \min\{T_2, T_3\}$, eq. (18) holds (which follows from Lemma 11), and

$$\forall i,j \in [d] : \frac{E_j^{(t)}}{E_i^{(t)}} = \Theta(1), \tag{21}$$

which follows from the part (A) of Lemma 13. Now the only thing to verify is the base case, i.e. when $t = T_1$. More specifically, we want to prove that 1) $\forall i,j \in [d] : \left|w_{2i}^{(T_1)}\right| \leq \mathcal{O}\left(d^{1/4}\right), \left|W_1^{(T_1)}[i,j]\right| \leq \mathcal{O}\left(\frac{1}{d^{1/4}}\right)$ and that 2) $\forall j \in [d] : \frac{v_j^{(T_1)}}{c^{(T_1)}} = \Theta\left(\frac{1}{\sqrt{d}}\right)$, and that 3) $\frac{\left|R_{3j}^{(T_1)}\right|}{A_j}$ and $\frac{\left|R_{vj}^{(T_1)}\right|}{a^{(T_1)} A_j}$ are of order $\tilde{\mathcal{O}}(\epsilon_0)$. All of them can be verified by the proof in Section C.2 and the definition of $R_v^{(t)}, R_3^{(t)}$.

So far we have proved eq. (18) in Lemma 8. Now let's prove when $t = \min\{T_2, T_3\}$, we have that $\left\|E^{(t)}\right\|_2^2 = \mathcal{O}(\epsilon_0 d)$.

If $\min\{T_2, T_3\} = T_3$, by Definition 3, we have $\left\|E^{(t)}\right\|_2^2 \leq \epsilon$. If $\min\{T_2, T_3\} = T_2$, by Definition 2, there exists $j \in [d]$ such that $E_j^{(t)} = -\Theta\left(\sqrt{\epsilon_0}\right)$. Combining with eq. (21) gives us $\forall i \in [d] :$

$E_i^{(t)} = -\Theta\left(\sqrt{\epsilon_0}\right)$. Combining these two cases, we get that when $t = \min\{T_2, T_3\}$, $\left\|E^{(t)}\right\|_2^2 \leq \max\{\epsilon, \Theta\left(\epsilon_0 d\right)\} = \mathcal{O}\left(\epsilon_0 d\right)$.

C.10 PROOF OF LEMMA 9

We prove this lemma by induction. The base case ($t = T_1$) of $\boldsymbol{v}^{(t)}$ is verified by eq. (15).

Suppose at time t, $\boldsymbol{v}^{(t)T} = a^{(t)}A + R_v^{(t)T}$, then by eq. 16, we have that

$$
\begin{aligned}
W_2^{(t)}W_1^{(t)} &= \left(c^{(t)}\boldsymbol{u}^{(T_1)T} + R_2^{(t)T}\right)\left(\boldsymbol{u}^{(T_1)}\boldsymbol{v}^{(t)T} + R_1^{(t)}\right) \\
&= \left(c^{(t)}\left\|\boldsymbol{u}^{(T_1)}\right\|^2 + R_2^{(t)T}\boldsymbol{u}^{(T_1)}\right)\boldsymbol{v}^{(t)} + c^{(t)}\boldsymbol{u}^{(T_1)T}R_1^{(t)} + R_2^{(t)T}R_1^{(t)} \\
&= d^{(t)}\boldsymbol{v}^{(t)T} + R_3^{(t)T} \\
&= d^{(t)}a^{(t)}A + d^{(t)}R_v^{(t)T} + R_3^{(t)T},
\end{aligned}
$$

where $d^{(t)} := c^{(t)}\left\|\boldsymbol{u}^{(T_1)}\right\|^2 + R_2^{(t)T}\boldsymbol{u}^{(T_1)}$, $\quad R_3^{(t)T} := c^{(t)}\boldsymbol{u}^{(T_1)T}R_1^{(t)} + R_2^{(t)T}R_1^{(t)}$. That gives us

$$
\begin{aligned}
\boldsymbol{v}^{(t+1)T} &= \boldsymbol{v}^{(t)T} - \eta_t c^{(t)}E^{(t)} \\
&= a^{(t)}A + R_v^{(t)T} - \eta_t c^{(t)}\left(d^{(t)}a^{(t)}A + d^{(t)}R_v^{(t)T} + R_3^{(t)T} - A\right) \\
&= \left(\left(1 - \eta_t c^{(t)}d^{(t)}\right)a^{(t)} + \eta_t c^{(t)}\right)A + \left(1 - \eta_t c^{(t)}d^{(t)}\right)R_v^{(t)T} - \eta_t c^{(t)}R_3^{(t)T} \\
&:= a^{(t+1)}A + R_v^{(t+1)T}.
\end{aligned}
$$

Therefore we have proved by induction that for t in the second phase, $\boldsymbol{v}^{(t)} = a^{(t)}A + R_v^{(t)T}$. The above steps also proved eq. (19).

C.11 PROOF OF LEMMA 10

Write $r_1^{(t)} = q_{11}^{(t)} + q_{12}^{(t)}$ where we have $q_{11}^{(t)} = \eta\sum_{\tau=0}^{t}\beta^{t-\tau}\left(W_2^{(t)T}E^{(t)} - W_2^{(\tau)T}E^{(\tau)}\right)$, $q_{12}^{(t)} = -\eta\sum_{\tau=0}^{t}\beta^{t-\tau}Dg_1^{(\tau)}$. Write $r_2^{(t)} = q_{21}^{(t)} + q_{22}^{(t)}$ where $q_{22}^{(t)} = -\eta\sum_{\tau=0}^{t}\beta^{t-\tau}Dg_2^{(\tau)}$, $q_{21}^{(t)} = \eta\sum_{\tau=0}^{t}\beta^{t-\tau}\left(E^{(t)}W_1^{(t)T} - E^{(\tau)}W_1^{(\tau)T}\right)$.

Let's first bound $\left|q_{11}^{(t)}[i,j]\right|$ and $\left|q_{21,i}^{(t)}\right|$. By definition of T_2, we know that for $T_1 \leq \tau \leq t, \forall i \in [d]$: $\left|E_i^{(\tau)}\right| = \mathcal{O}(1)$. Then we have for all $i, j \in [d]$,

$$
\left|W_1^{(\tau+1)}[i,j] - W_1^{(\tau)}[i,j]\right| \leq \eta\sum_{k=0}^{\tau}\beta^{\tau-k}\left|w_{2i}^{(k)}E_j^{(k)}\right| \leq \eta\sum_{k=0}^{\tau}\beta^{\tau-k}\mathcal{O}\left(d^{1/4}\right) = \eta\mathcal{O}\left(d^{1/4}\right),
$$

$$
\left|w_{2i}^{(\tau+1)} - w_{2i}^{(\tau)}\right| \leq \eta\sum_{k=0}^{\tau}\beta^{\tau-k}\sum_{j=1}^{d}\left|E_j^{(k)}W_1^{(k)}[j,i]\right| \leq \eta\sum_{k=0}^{\tau}\beta^{\tau-k}\mathcal{O}\left(d^{3/4}\right) = \eta\mathcal{O}\left(d^{3/4}\right).
$$

$$\tag{22}$$

Note that

$$
\begin{aligned}
\left|E_j^{(\tau+1)} - E_j^{(\tau)}\right| &= \sum_{i=1}^{d}\left(\left(w_{2i}^{(\tau+1)} - w_{2i}^{(\tau)}\right)W_1^{(\tau)}[i,j] + w_{2i}^{(\tau)}\left(W_1^{(\tau+1)}[i,j] - W_1^{(\tau)}[i,j]\right)\right) \\
&\quad + \sum_{i=1}^{d}\left(\left(w_{2i}^{(\tau+1)} - w_{2i}^{(\tau)}\right)\left(W_1^{(\tau+1)}[i,j] - W_1^{(\tau)}[i,j]\right)\right).
\end{aligned}
$$

We can further get that for $\forall j \in [d]$,

$$
\begin{aligned}
\left|E_j^{(\tau+1)} - E_j^{(\tau)}\right| &\leq \eta d\mathcal{O}\left(d^{3/4}\right)\mathcal{O}\left(d^{-1/4}\right) + \eta d\mathcal{O}\left(d^{1/4}\right)\mathcal{O}\left(d^{1/4}\right) + \eta^2 d\mathcal{O}\left(d^{3/4}\right)\mathcal{O}\left(d^{1/4}\right) \\
&= \mathcal{O}\left(\eta d^{3/2} + \eta^2 d^2\right) = \mathcal{O}\left(\eta d^{3/2}\right).
\end{aligned}
$$

Combining the above inequalities gives us $\forall i, j \in [d]$,

$$
\begin{aligned}
\left| q_{11}^{(t)}[i,j] \right| = \eta \left| \sum_{\tau=0}^{t} \beta^{t-\tau} \left(w_{2i}^{(t)} E_j^{(t)} - w_{2i}^{(\tau)} E_j^{(\tau)} \right) \right| \\
\leq \eta \sum_{\tau=0}^{t} \beta^{t-\tau} \left(\left| w_{2i}^{(t)} - w_{2i}^{(\tau)} \right| \left| E_j^{(t)} \right| + \left| w_{2i}^{(\tau)} \right| \left| E_j^{(t)} - E_j^{(\tau)} \right| \right) \\
\leq \eta^2 \sum_{\tau=0}^{t} \beta^{t-\tau} (t-\tau) \left(\mathcal{O}\left(d^{3/4} \right) \mathcal{O}(1) + \mathcal{O}\left(d^{1/4} \right) \mathcal{O}\left(d^{3/2} \right) \right) \\
= \mathcal{O}\left(\eta^2 d^{7/4} \right),
\end{aligned}
$$

$$
\begin{aligned}
\left| q_{12,i}^{(t)} \right| = \eta \left| \sum_{\tau=0}^{t} \beta^{t-\tau} \sum_{j=1}^{d} \left(E_j^{(t)} W_1^{(t)}[i,j] - E_j^{(\tau)} W_1^{(\tau)}[i,j] \right) \right| \\
\leq \eta \sum_{\tau=0}^{t} \beta^{t-\tau} \sum_{j=1}^{d} \left(\left| E_j^{(t)} \right| \left| W_1^{(t)}[i,j] - W_1^{(\tau)}[i,j] \right| + \left| E_j^{(t)} - E_j^{(\tau)} \right| \left| W_1^{(\tau)}[i,j] \right| \right) \\
\leq \eta^2 d \sum_{\tau=0}^{t} \beta^{t-\tau} (t-\tau) \left(\mathcal{O}(1) \mathcal{O}\left(d^{1/4} \right) + \mathcal{O}\left(d^{3/2} \right) \mathcal{O}\left(d^{-1/4} \right) \right) \\
= \mathcal{O}\left(\eta^2 d^{9/4} \right).
\end{aligned}
$$

Next let's bound $\left| q_{12}^{(t)}[i,j] \right|$ and $\left| q_{22,i}^{(t)} \right|$. By the assumption of this lemma and the analysis before T_1, we know that for all $\tau \leq t$, the $M_1^{(\tau)}, M_2^{(\tau)}$ in Lemma 31 are upper bounded by $\mathcal{O}\left(\frac{1}{d^{1/4}} \right)$ and $\mathcal{O}\left(d^{1/4} \right)$, respectively. In the theorem we consider the training period before $T_{\text{SGD},2}$ so the time T in Lemma 31 is set as $T_{\text{SGD},2}$. In the following sections, we will prove that $T_{\text{SGD},2} \leq \mathcal{O}\left(\frac{d^{\alpha} \log(\sqrt{d/\epsilon})}{\eta} \right)$. Then by Lemma 31, we have with probability at least $1 - \frac{1}{d}$, for $\forall \tau \leq t$ and $\forall i, j \in [d]$,

$$
\begin{aligned}
\left| Dg_1^{(\tau)}[i,j] \right| = \left| \tilde{g}_1^{(\tau)}[i,j] - g_1^{(\tau)}[i,j] \right| \leq \mathcal{O}\left(d^{\frac{13}{4}} \sigma \sqrt{\frac{d^{\alpha+1}}{\eta} \log \frac{d}{\epsilon}} \right) + \mathcal{O}\left(d^{\frac{1}{4}} \sigma \sqrt{\frac{d^{\alpha+2}}{\eta} \log \frac{d}{\epsilon}} \right) \\
\leq \tilde{\mathcal{O}}\left(d^{\frac{13}{4}} \sigma \sqrt{\frac{d^{\alpha+1}}{\eta}} \right)
\end{aligned}
$$

$$
\begin{aligned}
\left| Dg_{2i}^{(\tau)} \right| = \left| \tilde{g}_{2i}^{(\tau)} - g_{2i}^{(\tau)} \right| \leq \mathcal{O}\left(d^{\frac{15}{4}} \sigma \sqrt{\frac{d^{\alpha+1}}{\eta} \log \frac{d}{\epsilon}} \right) + \mathcal{O}\left(d^{\frac{3}{4}} \sigma \sqrt{\frac{d^{\alpha+2}}{\eta} \log \frac{d}{\epsilon}} \right) \\
= \tilde{\mathcal{O}}\left(d^{\frac{15}{4}} \sigma \sqrt{\frac{d^{\alpha+1}}{\eta}} \right)
\end{aligned}
$$

By picking $\sigma \leq \frac{\eta^{3/2}}{d^{\alpha/2+1}}$, we have $\left| Dg_1^{(\tau)}[i,j] \right| \leq \eta \tilde{\mathcal{O}}\left(d^{\frac{11}{4}} \right)$ and $\left| Dg_{2i}^{(\tau)} \right| \leq \eta \tilde{\mathcal{O}}\left(d^{\frac{13}{4}} \right)$, which yields

$$
\left| q_{12}^{(t)}[i,j] \right| \leq \eta \sum_{\tau=0}^{t} \beta^{t-\tau} \left| Dg_1^{(\tau)}[i,j] \right| \leq \tilde{\mathcal{O}}\left(\eta^2 d^{\frac{11}{4}} \right)
$$

$$
\left| q_{22,i}^{(t)} \right| \leq \eta \sum_{\tau=0}^{t} \beta^{t-\tau} \left| Dg_{2i}^{(\tau)} \right| \leq \tilde{\mathcal{O}}\left(\eta^2 d^{\frac{13}{4}} \right)
$$

Combining the above bounds, we get that $\forall i, j \in [d]$,

$$
\left| r_1^{(t)}[i,j] \right| \leq \tilde{\mathcal{O}}\left(\eta^2 d^{\frac{11}{4}} \right), \quad \left| r_{2i}^{(t)} \right| \leq \tilde{\mathcal{O}}\left(\eta^2 d^{\frac{13}{4}} \right).
$$

By the analysis in Section C.2, we know that at time T_1, for some $i_0 \in [d]$, $c^{(T_1)} \left| u_{i_0}^{(T_1)} \right| = \Theta \left(\frac{1}{d^{\frac{\alpha}{2}}} \right)$, and for $\forall i, j \in [d]$, we have $c^{(T_1)} \left| u_i^{(T_1)} \right| = \tilde{\Omega} \left(\frac{1}{d^{\frac{3\alpha}{4}}} \right)$ and $\left| u_i^{(T_1)} \right| v_j^{(T_1)} = \tilde{\Omega} \left(\frac{1}{d^{\frac{3\alpha}{4}+\frac{1}{2}}} \right)$, which gives us $\forall i, j \in [d]$,

$$\frac{\left| r_1^{(t)}[i,j] \right|}{\left| u_i^{(T_1)} \right| v_j^{(t)}} \leq \frac{\left| r_1^{(t)}[i,j] \right|}{\left| u_i^{(T_1)} \right| v_j^{(T_1)}} = \tilde{\mathcal{O}} \left(\eta^2 d^{\frac{3}{4}\alpha + \frac{13}{4}} \right), \quad \frac{\left| r_{2i}^{(t)} \right|}{c^{(t)} \left| u_i^{(T_1)} \right|} \leq \frac{\left| r_{2i}^{(t)} \right|}{c^{(T_1)} \left| u_i^{(T_1)} \right|} = \tilde{\mathcal{O}} \left(\eta^2 d^{\frac{3}{4}\alpha + \frac{13}{4}} \right).$$

Hence we get the bound $\forall i \in [d] : \epsilon_i^{(t)} \leq \tilde{\mathcal{O}} \left(\eta^2 d^{\frac{3}{4}\alpha + \frac{13}{4}} \right)$.

C.12 PROOF OF LEMMA 11

Let's first try to bound the length of $\min\{T_2, T_3\}$. More formally, we prove that under the conditions of Lemma 10 and pick $\eta \leq \mathcal{O} \left(\frac{\epsilon}{d^{\frac{7\alpha}{4}+4}} \right)$, we have that $\min\{T_2, T_3\} \leq \mathcal{O} \left(\frac{d^\alpha \log(\sqrt{d/\epsilon})}{\eta} \right)$.

Under the conditions of Lemma 10, we know that

$$\forall j \in [d] : \quad \left| \left(r_2^{(t)} W_1^{(t)} \right)_j \right| \leq \sum_{i=1}^d \left| r_{2i}^{(t)} W_1^{(t)}[i,j] \right| = \mathcal{O} \left(\eta^2 d^4 \right),$$

$$\left| \left(W_2^{(t)} r_1^{(t)} \right)_j \right| \leq \sum_{i=1}^d \left| w_{2i}^{(t)} r_1^{(t)}[i,j] \right| = \mathcal{O} \left(\eta^2 d^4 \right).$$

Combining with eq. (22), we get

$$E^{(t+1)}$$
$$=E^{(t)} + \left(W_2^{(t+1)} - W_2^{(t)} \right) W_1^{(t)} + W_2^{(t)} \left(W_1^{(t+1)} - W_1^{(t)} \right) + \left(W_2^{(t+1)} - W_2^{(t)} \right) \left(W_1^{(t+1)} - W_1^{(t)} \right)$$
$$=E^{(t)} - \eta_t E^{(t)} W_1^{(t)T} W_1^{(t)} + r_2^{(t)} W_1^{(t)} - \eta_t W_2^{(t)} W_2^{(t)T} E^{(t)} + W_2^{(t)} r_1^{(t)} + \mathcal{O} \left(\eta^2 d \right)$$
$$=E^{(t)} \left(I - \eta_t W_1^{(t)T} W_1^{(t)} - \eta_t \left\| W_2^{(t)} \right\|_2^2 I \right) + \mathcal{O} \left(\eta^2 d^4 \right) + \mathcal{O} \left(\eta^2 d^4 \right) + \mathcal{O} \left(\eta^2 d \right).$$

Then we have

$$\left\| E^{(t+1)} \right\|_2 \leq \left\| E^{(t)} \right\|_2 \left\| I - \eta_t W_1^{(t)T} W_1^{(t)} - \eta_t \left\| W_2^{(t)} \right\|_2^2 I \right\|_2 + \mathcal{O} \left(\eta^2 d^4 \right)$$

$$\leq \left(1 - \eta_t \left\| W_2^{(t)} \right\|_2^2 \right) \left\| E^{(t)} \right\|_2 + \mathcal{O} \left(\eta^2 d^4 \right).$$

When $T_1 \leq t < T_2$, we have proved that $c^{(t)}$ is increasing over time in Section C.8, which implies that $\left\| W_2^{(t)} \right\|_2^2 \geq C \left\| W_2^{(T_1)} \right\|_2^2$ since $c^{(t)} \boldsymbol{u}^{(T_1)T}$ is the leading term of $W_2^{(t)}$. Combining with $\eta_t \geq \eta$ gives us

$$\left\| E^{(t+1)} \right\|_2 \leq \left(1 - \eta C \left\| W_2^{(T_1)} \right\|_2^2 \right) \left\| E^{(t)} \right\|_2 + \mathcal{O} \left(\eta^2 d^4 \right),$$

$$\Rightarrow \quad \left\| E^{(t)} \right\|_2 \leq \frac{\mathcal{O} \left(\eta^2 d^4 \right)}{\eta C \left\| W_2^{(T_1)} \right\|_2^2} + \left(1 - \eta C \left\| W_2^{(T_1)} \right\|_2^2 \right)^{t-T_1} \left(\left\| E^{(T_1)} \right\|_2 - \frac{\mathcal{O} \left(\eta^2 d^4 \right)}{\eta C \left\| W_2^{(T_1)} \right\|_2^2} \right)$$

$$\overset{(i)}{\leq} \mathcal{O} \left(\frac{\eta d^4}{\left\| W_2^{(T_1)} \right\|_2^2} \right) + \exp \left(-\eta C \left\| W_2^{(T_1)} \right\|_2^2 (t - T_1) \right) \mathcal{O} \left(\sqrt{d} \right),$$

where (i) uses $\left\|E^{(T_1)}\right\|_2 = \mathcal{O}(\sqrt{d})$. By picking $\eta \leq \mathcal{O}\left(\frac{\epsilon}{d^{\frac{7\alpha}{4}+4}}\right)$ and noticing that $\left\|W_2^{(T_1)}\right\|_2^2 \geq \Omega\left(\frac{1}{d^\alpha}\right)$, we have $\frac{\eta d^4}{\left\|W_2^{(T_1)}\right\|_2^2} < \frac{\sqrt{\epsilon}}{2}$. Hence when $t - T_1 \geq \Theta\left(\frac{\log\left(\sqrt{d/\epsilon}\right)}{\eta\left\|W_2^{(T_1)}\right\|_2^2}\right)$, we have that $\left\|E^{(t)}\right\|_2 \leq \sqrt{\epsilon}$, i.e. $\left\|E^{(t)}\right\|_2^2 \leq \epsilon$.

That means after at most $\mathcal{O}\left(\frac{\log\left(\sqrt{d/\epsilon}\right)}{\eta\left\|W_2^{(T_1)}\right\|_2^2}\right)$ steps from T_1, either $t \geq T_2$, or we have $\left\|E^{(t)}\right\|_2^2 \leq \epsilon$. In other words, $\min\{T_2, T_3\} \leq T_1 + \mathcal{O}\left(\frac{\log\left(\sqrt{d/\epsilon}\right)}{\eta\left\|W_2^{(T_1)}\right\|_2^2}\right) \leq \mathcal{O}\left(\frac{d^\alpha \log\left(\sqrt{d/\epsilon}\right)}{\eta}\right)$.

Now we are ready to bound eq. 17.

Combining $\min\{T_2, T_3\} \leq \mathcal{O}\left(\frac{d^\alpha \log(\sqrt{d/\epsilon})}{\eta}\right)$ and Lemma 10 yields that for $t + 1 \leq \min\{T_2, T_3\}$,

$$\forall i \in [d] : \sum_{\tau=T_1}^{t+1} \epsilon_i^{(\tau)} \leq (t + 1 - T_1)\tilde{\mathcal{O}}\left(\eta^2 d^{\frac{3}{4}\alpha+\frac{13}{4}}\right) \leq \tilde{\mathcal{O}}\left(\eta d^{\frac{7}{4}\alpha+\frac{13}{4}} \log\sqrt{\frac{d}{\epsilon}}\right) = \tilde{\mathcal{O}}\left(\epsilon \log\sqrt{\frac{d}{\epsilon}}\right).$$

Lemma 1 tells us that $\delta_i = \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{1}{4}\alpha-1}}\right)$. Substituting these bounds into eq. (17) completes the proof.

C.13 PROOF OF LEMMA 12

The proof in Section C.8 tells us that for $T_1 \leq \tau \leq T_2$, $c^{(\tau)} > 0, \forall j \in [d] : v_j^{(\tau)} > 0$, which gives us $0 \leq \frac{\left|R_2^{(t+1)T}\boldsymbol{u}^{(T_1)}\right|}{c^{(t+1)}\left\|\boldsymbol{u}^{(T_1)}\right\|^2}$ and $0 \leq \frac{\left|R_{3j}^{(t+1)}\right|}{c^{(t+1)}\left\|\boldsymbol{u}^{(T_1)}\right\|^2 v_j^{(t+1)}}$. By Lemma 11, we have that

$$\forall 1 \leq i, j \leq d : \quad 0 \leq \frac{\left|R_1^{(t+1)}[i,j]\right|}{\left|u_i^{(T_1)}\right| v_j^{(t+1)}} \leq \tilde{\mathcal{O}}(\epsilon_0), \quad 0 \leq \frac{\left|R_{2i}^{(t+1)}\right|}{c^{(t+1)}\left|u_i^{(T_1)}\right|} \leq \tilde{\mathcal{O}}(\epsilon_0),$$

which gives us

$$\frac{\left|R_2^{(t+1)T}\boldsymbol{u}^{(T_1)}\right|}{c^{(t+1)}\left\|\boldsymbol{u}^{(T_1)}\right\|^2} \leq \frac{\sum_{i=1}^d \left|u_i^{(T_1)}\right|\left|R_{2i}^{(t+1)}\right|}{c^{(t+1)}\sum_{i=1}^d \left|u_i^{(T_1)}\right|^2} \leq \frac{\tilde{\mathcal{O}}(\epsilon_0)c^{(t+1)}\sum_{i=1}^d \left|u_i^{(T_1)}\right|^2}{c^{(t+1)}\sum_{i=1}^d \left|u_i^{(T_1)}\right|^2} = \tilde{\mathcal{O}}(\epsilon_0).$$

Lemma 9 tells us that

$$R_3^{(t+1)T} = c^{(t+1)}\boldsymbol{u}^{(T_1)T}R_1^{(t+1)} + R_2^{(t+1)T}R_1^{(t+1)}.$$

And we have that

$$\frac{\left|\left(c^{(t+1)}\boldsymbol{u}^{(T_1)T}R_1^{(t+1)}\right)_j\right|}{c^{(t+1)}\left\|\boldsymbol{u}^{(T_1)}\right\|^2 v_j^{(t+1)}} \leq \frac{c^{(t+1)}\sum_{i=1}^d \left|u_i^{(T_1)}\right|\left|R_1^{(t+1)}[i,j]\right|}{c^{(t+1)}\sum_{i=1}^d \left|u_i^{(T_1)}\right|^2 v_j^{(t+1)}} \leq \frac{\tilde{\mathcal{O}}(\epsilon_0)c^{(t+1)}\sum_{i=1}^d \left|u_i^{(T_1)}\right|^2 v_j^{(t+1)}}{c^{(t+1)}\sum_{i=1}^d \left|u_i^{(T_1)}\right|^2 v_j^{(t+1)}}$$

$$= \tilde{\mathcal{O}}(\epsilon_0),$$

$$\frac{\left|\left(R_2^{(t+1)T}R_1^{(t+1)}\right)_j\right|}{c^{(t+1)}\left\|\boldsymbol{u}^{(T_1)}\right\|^2 v_j^{(t+1)}} \leq \frac{\sum_{i=1}^d \left|R_{2i}^{(t+1)}\right|\left|R_1^{(t+1)}[i,j]\right|}{c^{(t+1)}\sum_{i=1}^d \left|u_i^{(T_1)}\right|^2 v_j^{(t+1)}} \leq \frac{\tilde{\mathcal{O}}\left(\epsilon_0^2\right)c^{(t+1)}\sum_{i=1}^d \left|u_i^{(T_1)}\right|^2 v_j^{(t+1)}}{c^{(t+1)}\sum_{i=1}^d \left|u_i^{(T_1)}\right|^2 v_j^{(t+1)}}$$

$$= \tilde{\mathcal{O}}\left(\epsilon_0^2\right).$$

Therefore

$$\frac{\left|R_{3j}^{(t+1)}\right|}{c^{(t+1)}\left\|\boldsymbol{u}^{(T_1)}\right\|^2 v_j^{(t+1)}} \leq \frac{\left|\left(c^{(t+1)}\boldsymbol{u}^{(T_1)T}R_1^{(t+1)}\right)_j\right|}{c^{(t+1)}\left\|\boldsymbol{u}^{(T_1)}\right\|^2 v_j^{(t+1)}} + \frac{\left|\left(R_2^{(t+1)T}R_1^{(t+1)}\right)_j\right|}{c^{(t+1)}\left\|\boldsymbol{u}^{(T_1)}\right\|^2 v_j^{(t+1)}} \leq \tilde{\mathcal{O}}(\epsilon_0).$$

By Lemma 9,

$$W_2^{(t+1)} W_1^{(t+1)} = c^{(t+1)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2 \boldsymbol{v}^{(t+1)T} + R_2^{(t+1)T} \boldsymbol{u}^{(T_1)} \boldsymbol{v}^{(t+1)T} + R_3^{(t+1)T}.$$

Then we have that

$$\forall j \in [d]: \quad \left(W_2^{(t+1)} W_1^{(t+1)} \right)_j = c^{(t+1)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2 v_j^{(t+1)} \left(1 + e_j^{(t+1)} \right), \quad \text{where } \left| e_j^{(t+1)} \right| \leq \tilde{\mathcal{O}}(\epsilon_0). \tag{23}$$

Since $t < T_2$, we have $\forall j \in [d]: \dfrac{\left(W_2^{(t+1)} W_1^{(t+1)} \right)_j}{A_j} = \mathcal{O}(1)$, which yields

$$0 \leq \frac{c^{(t+1)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2 v_j^{(t+1)}}{A_i} \leq \mathcal{O}(1), \tag{24}$$

which proves eq. (20), since $\forall j \in [d]: \quad 0 \leq \dfrac{\left| R_{3j}^{(t+1)} \right|}{c^{(t+1)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2 v_j^{(t+1)}} \leq \tilde{\mathcal{O}}(\epsilon_0).$

C.14 PROOF OF LEMMA 13

(A) Since $\dfrac{\left| R_{vj}^{(t)} \right|}{a^{(t)} A_j} \leq \tilde{\mathcal{O}}(\epsilon_0)$, we get that

$$v_j^{(t)} = a^{(t)} A_j + R_{vj}^{(t)} = a^{(t)} A_j \left(1 + e_{vj}^{(t)} \right), \quad \text{where } \left| e_{vj}^{(t)} \right| \leq \tilde{\mathcal{O}}(\epsilon_0).$$

Substituting into eq.(20) yields

$$\forall j \in [d]: \quad \left(W_2^{(t)} W_1^{(t)} \right)_j = a^{(t)} c^{(t)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2 A_j \left(1 + \tilde{e}_j^{(t)} \right), \quad \text{where } \left| \tilde{e}_j^{(t)} \right| \leq \tilde{\mathcal{O}}(\epsilon_0),$$

That gives us

$$\forall j \in [d]: E_j^{(t)} = A_j \left(a^{(t)} c^{(t)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2 - 1 + a^{(t)} c^{(t)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2 \tilde{e}_j^{(t)} \right).$$

Since $t < T_2$, we have $E_j^{(t)} < -\sqrt{\epsilon_0}$. Combining with $A_j = \Theta(1)$, gives us $a^{(t)} c^{(t)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2 - 1 = -\Omega\left(\sqrt{\epsilon_0} \right)$. Then we can rewrite $E_j^{(t)}$ as $\forall j \in [d]$,

$$E_j^{(t)} = A_j \left(a^{(t)} c^{(t)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2 - 1 \right) \left(1 + \frac{a^{(t)} c^{(t)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2}{a^{(t)} c^{(t)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2 - 1} \tilde{e}_j^{(t)} \right)$$

$$:= A_j \left(a^{(t)} c^{(t)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2 - 1 \right) \left(1 + e_{Ej}^{(t)} \right),$$

where $\left| e_{Ej}^{(t)} \right| = \tilde{\mathcal{O}}(\sqrt{\epsilon_0})$. Hence $\forall i, j \in [d]: \dfrac{E_j^{(t)}}{E_i^{(t)}} = \Theta(1)$.

(B) Note that we assume $\forall j \in [d]: \quad \dfrac{v_j^{(t)}}{c^{(t)}} = \Theta\left(\frac{1}{\sqrt{d}} \right)$, then we have for $j \in [d]$,

$$\frac{-E^{(t)} \boldsymbol{v}^{(t)}}{c^{(t)} \left(-E_j^{(t)} \right)} = \frac{\sum_{i=1}^d \left(-E_i^{(t)} \right) v_i^{(t)}}{c^{(t)} \left(-E_j^{(t)} \right)} = \sum_{i=1}^d \frac{E_i^{(t)}}{E_j^{(t)}} \cdot \frac{v_i^{(t)}}{c^{(t)}} = \sum_{i=1}^d \Theta\left(\frac{1}{\sqrt{d}} \right) = \Theta\left(\sqrt{d} \right),$$

$$\Rightarrow \quad \frac{c^{(t)} \left(-E_j^{(t)} \right)}{-E^{(t)} \boldsymbol{v}^{(t)}} = \Theta\left(\frac{1}{\sqrt{d}} \right).$$

Then for $t + 1$, we have that for $j \in [d]$,

$$\frac{v_j^{(t+1)}}{c^{(t+1)}} = \frac{v_j^{(t)} + \eta_t c^{(t)} \left(-E_j^{(t)} \right)}{c^{(t)} + \eta_t \left(-E^{(t)} \right) \boldsymbol{v}^{(t)}} = \Theta\left(\frac{1}{\sqrt{d}} \right).$$

(C) Combining eq. (24) and $\forall j \in [d] : A_j = \Theta(1)$, we know that

$$c^{(t+1)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2 v_j^{(t+1)} \leq \mathcal{O}(1),$$

which yields $\forall j \in [d]$,

$$\left\| \boldsymbol{u}^{(T_1)} \right\|^2 \left(v_j^{(t+1)} \right)^2 \leq \frac{v_j^{(t+1)}}{c^{(t+1)}} \mathcal{O}(1) = \mathcal{O}\left(\frac{1}{\sqrt{d}} \right), \quad \left(c^{(t+1)} \right)^2 \left\| \boldsymbol{u}^{(T_1)} \right\|^2 \leq \frac{c^{(t+1)}}{v_j^{(t+1)}} \mathcal{O}(1) = \mathcal{O}\left(\sqrt{d} \right).$$

(25)

Hence $\forall i, j \in [d]$,

$$c^{(t+1)} \left| u_i^{(T_1)} \right| = \mathcal{O}\left(d^{1/4} \right) \quad \Rightarrow \quad \left| w_{2i}^{(t+1)} \right| \leq c^{(t+1)} \left| u_i^{(T_1)} \right| + \left| R_{2i}^{(t+1)} \right| \overset{(i)}{=} \mathcal{O}\left(d^{1/4} \right),$$

$$\left| u_i^{(T_1)} \right| v_j^{(t+1)} = \mathcal{O}\left(\frac{1}{d^{1/4}} \right) \quad \Rightarrow \quad \left| W_1^{(t+1)}[i,j] \right| \leq \left| u_i^{(T_1)} \right| v_j^{(t+1)} + \left| R_1^{(t+1)}[i,j] \right| \overset{(ii)}{=} \mathcal{O}\left(d^{1/4} \right),$$

where (i) and (ii) use Corollary 11.

(D) The fact that $\forall j \in [d] : \frac{\left| R_{3j}^{(t+1)} \right|}{A_j} \leq \tilde{\mathcal{O}}(\epsilon_0)$ was already proved in Lemma 12 in eq.(20). To analyze $\frac{\left| R_{vi}^{(t+1)} \right|}{a^{(t+1)} A_i}$, we first prove that $1 - \eta_t c^{(t)} d^{(t)} > 0$.

Note that eq.(25) also holds for time t. Recall that $d^{(t)} = c^{(t)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2 + R_2^{(t)T} \boldsymbol{u}^{(T_1)}$ and Lemma 12 tells us that $0 \leq \frac{\left| R_2^{(t)T} \boldsymbol{u}^{(T_1)} \right|}{c^{(t)} \left\| \boldsymbol{u}^{(T_1)} \right\|^2} \leq \tilde{\mathcal{O}}(\epsilon_0)$, then we have

$$c^{(t)} d^{(t)} = \left(c^{(t)} \right)^2 \left\| \boldsymbol{u}^{(T_1)} \right\|^2 + c^{(t)} R_2^{(t)T} \boldsymbol{u}^{(T_1)} \leq \mathcal{O}\left(\sqrt{d} \right).$$

Under the conditions of Lemma 10, we pick $\eta = \mathcal{O}\left(\frac{1}{d^\alpha} \right)$, then we have that $1 - \eta_t c^{(t)} d^{(t)} \geq 1 - \eta c^{(t)} d^{(t)} > 0$.

The assumption $\forall j \in [d] : \frac{\left| R_{3j}^{(t)} \right|}{A_j} \leq \tilde{\mathcal{O}}(\epsilon_0)$ together with $c^{(t)} > 0$ gives us

$$\frac{\eta_t c^{(t)} \left| R_{3j}^{(t)} \right|}{\eta_t c^{(t)} A_j} \leq \tilde{\mathcal{O}}(\epsilon_0).$$

Combining with the assumption $\frac{\left| R_{vi}^{(t)} \right|}{a^{(t)} A_i} \leq \tilde{\mathcal{O}}(\epsilon_0)$ yields

$$\forall i \in [d] : \quad \frac{\left| R_{vi}^{(t+1)} \right|}{a^{(t+1)} A_i} \leq \frac{\left(1 - \eta_t c^{(t)} d^{(t)} \right) \left| R_v^{(t)} \right| + \eta_t c^{(t)} \left| R_{3i}^{(t)} \right|}{\left(1 - \eta_t c^{(t)} d^{(t)} \right) a^{(t)} A_i + \eta_t c^{(t)} A_i} \leq \tilde{\mathcal{O}}(\epsilon_0).$$

D   ANALYSIS OF ADAM

Note that $A = \frac{1}{m} Y X^T$, $\Lambda_{xx} := \frac{1}{m} X X^T$. Denote $g_k^{(t)} := \nabla_{W_k} L(W^{(t)}), k = 1, 2$. We have that

$$g_1^{(t)} = W_2^{(t)T} \left(W_2^{(t)} W_1^{(t)} - A \right), \quad g_2^{(t)} = \left(W_2^{(t)} W_1^{(t)} - A \right) W_1^{(t)T}.$$

Let $\tilde{A}^{(t)}$, $\tilde{\Lambda}_{xx}^{(t)}$ and $\tilde{g}_k^{(t)}, k = 1, 2$ be the corresponding batch versions at time t. Let $E^{(t)} := W_2^{(t)} W_1^{(t)} - A$, and denote $E_j^{(t)}$ as the j-th component of $E^{(t)}$, then by eq. (2), the update equations

of Adam are given by

$$\eta_t = \eta \cdot \frac{\sqrt{1 - \beta_2^{t+1}}}{1 - \beta_1^{t+1}}, \quad g_1^{(t)}[i,j] = w_{2i}^{(t)} E_j^{(t)}, \quad g_{2i}^{(t)} = \left\langle E^{(t)}, W_1^{(t)}[i,:] \right\rangle,$$

$$\begin{aligned}
W_1^{(t+1)}[i,j] &= W_1^{(t)}[i,j] - \eta_t \frac{m_1^{(t)}[i,j]}{\sqrt{v_1^{(t)}[i,j]}} \\
&= W_1^{(t)}[i,j] - \eta_t \frac{(1 - \beta_1) \sum_{\tau=0}^{t} \beta_1^{t-\tau} \tilde{g}_1^{(\tau)}[i,j]}{\sqrt{(1 - \beta_2) \sum_{\tau=0}^{t} \beta_2^{t-\tau} \left(\tilde{g}_1^{(\tau)}[i,j] \right)^2 + \xi}} \\
&= W_1^{(t)}[i,j] - \eta_t \frac{(1 - \beta_1) \sum_{\tau=0}^{t} \beta_1^{t-\tau} g_1^{(\tau)}[i,j] + r_{n1}^{(t)}[i,j]}{\sqrt{(1 - \beta_2) \sum_{\tau=0}^{t} \beta_2^{t-\tau} \left(g_1^{(\tau)}[i,j] \right)^2 + r_{d1}^{(t)}[i,j] + \xi}},
\end{aligned} \quad (26)$$

$$\begin{aligned}
w_{2i}^{(t+1)} &= w_{2i}^{(t)} - \eta_t \frac{m_{2i}^{(t)}}{\sqrt{v_{2i}^{(t)}}} = w_{2i}^{(t)} - \eta_t \frac{(1 - \beta_1) \sum_{\tau=0}^{t} \beta_1^{t-\tau} \tilde{g}_{2i}^{(\tau)}}{\sqrt{(1 - \beta_2) \sum_{\tau=0}^{t} \beta_2^{t-\tau} \left(\tilde{g}_{2i}^{(\tau)} \right)^2 + \xi}} \\
&= w_{2i}^{(t)} - \eta_t \frac{(1 - \beta_1) \sum_{\tau=0}^{t} \beta_1^{t-\tau} g_{2i}^{(\tau)} + r_{2n,i}^{(t)}}{\sqrt{(1 - \beta_2) \sum_{\tau=0}^{t} \beta_2^{t-\tau} \left(g_{2i}^{(\tau)} + Dg_{2i}^{(\tau)} \right)^2 + \xi}}.
\end{aligned}$$

where $Dg_1^{(t)} := \tilde{g}_1^{(t)} - g_1^{(t)}$ and $Dg_2^{(t)} := \tilde{g}_2^{(t)} - g_2^{(t)}$, and

$$\begin{aligned}
r_{1n}^{(t)}[i,j] &:= (1 - \beta_1) \sum_{\tau=0}^{t} \beta_1^{t-\tau} Dg_1^{(\tau)}[i,j], \\
r_{1d}^{(t)}[i,j] &= (1 - \beta_2) \sum_{\tau=0}^{t} \beta_2^{t-\tau} \left(2g_1^{(\tau)}[i,j] Dg_1^{(\tau)}[i,j] + \left(Dg_1^{(\tau)}[i,j] \right)^2 \right), \\
r_{2n,i}^{(t)} &:= (1 - \beta_1) \sum_{\tau=0}^{t} \beta_1^{t-\tau} Dg_{2i}^{(\tau)}, \\
r_{2d,i}^{(t)} &= (1 - \beta_2) \sum_{\tau=0}^{t} \beta_2^{t-\tau} \left(2g_{2i}^{(\tau)} Dg_{2i}^{(\tau)} + \left(Dg_{2i}^{(\tau)} \right)^2 \right)
\end{aligned} \quad (27)$$

Denote the i-th coordinate of $W_2 W_1$ and A as $(W_2 W_1)_i$ and A_i, respectively. By Assumption 2 and the assumption that $\forall i \in [d] : A_i > 0, A_i = \Omega(1)$, at the beginning, w.h.p., $\forall i \in [d] : (W_2 W_1)_i - A_i < 0$. Based on this, we divide the training procedure into two phases (note that these two phases are different from those of GD).

1. **First phase**: when the error $(W_2 W_1)_i - A_i$ is negative and its absolute value is big for all $i \in [d]$.
2. **Second phase**: when $(W_2 W_1)_i - A_i$ is close to zero for some coordinate $i \in [d]$.

More formally, we define the boundary between the two phases below.

Definition 4 (End of the first phase). *The end of the first phase (denoted as T_1) is defined as* $T_1 = \inf \left\{ t > 0 : \exists i \in [d] : E_i^{(t)} \geq -\sqrt{\eta d} \right\}$.

In the second phase, we define some time points.

Definition 5. *Define* $T_g := \inf \left\{ t > T_1 : \exists i \in [d] : \left| g_{2i}^{(t)} \right| \leq d\sqrt{\eta} \right\}$.

For $t < T_1$, we have $\forall i \in [d] : E_i^{(t)} < 0$ by Definition 4. For $t > T_1$, some $E_i^{(t)}$ may flip the sign and become positive. For certain coordinate i, we define the following "flip time".

Definition 6. *Define* $T_{f,i} := \inf \left\{ t > T_1 : E_i^{(t)} \geq -\sqrt{\eta d} \right\}$. *Define* $T_f := \max_i T_{f,i}$ *as the largest* "flip time" *over all* $i \in [d]$, *i.e. the* "flip time" *of the last* E_i *which flips the sign. Moreover, denote* $\tilde{T} := \min \{T_g, T_f\}$.

We can first show that after a few steps in the first phase, W_1 will become an approximately rank-1 matrix, as described in the following lemma.

Lemma 14. *Under Assumption 1, 2 and 3, suppose* $\sigma \leq \frac{\eta^{3/2} \xi^2}{d^{13/4}}$. *By picking* $\eta \leq \mathcal{O}\left(\frac{1}{d^{3\alpha}}\right), \xi \leq \sqrt{\frac{\eta}{d^{3\alpha-1}}}$, *and* $\beta_2 = \beta_1^2$, *there exists* $t_{inc} > 0$ *such that for* $t_{inc} \leq t < T_1$, *we have*

$$\forall i, j \in [d]: \quad w_{2i}^{(t)} = sign\left(w_{2i}^{(0)}\right) \eta \left(t - t_{inc}\right) + R_{2i}^{(t)},$$

$$W_1^{(t)}[i,j] = sign\left(w_{2i}^{(0)}\right) \eta \left(t - t_{inc}\right) + R_1^{(t)}[i,j],$$

where $\frac{\left|R_1^{(t)}[i,j]\right|}{\eta(t-t_{inc})} = \tilde{\mathcal{O}}\left(\sqrt{\eta} + \frac{1}{\eta(t-t_{inc})d^\alpha}\right)$, $\frac{\left|R_{2i}^{(t)}\right|}{\eta(t-t_{inc})} = \tilde{\mathcal{O}}\left(\sqrt{\eta} + \frac{1}{\eta(t-t_{inc})d^\alpha}\right)$.

Specially, when $t = T_1$, *we have that*

$$\forall i, j \in [d]: \quad w_{2i}^{(T_1)} = sign\left(w_{2i}^{(0)}\right) \eta \left(T_1 - t_{inc}\right) + R_{2i}^{(T_1)},$$

$$W_1^{(T_1)}[i,j] = sign\left(w_{2i}^{(0)}\right) \eta (T_1 - t_{inc}) + R_1^{(T_1)}[i,j],$$

where

$$\eta \left(T_1 - t_{inc}\right) = \Theta\left(\frac{1}{\sqrt{d}}\right), \quad \frac{\left|R_1^{(T_1)}[i,j]\right|}{\eta \left(T_1 - t_{inc}\right)} = \tilde{\mathcal{O}}\left(\sqrt{\eta} + \frac{1}{d^{\alpha-\frac{1}{2}}}\right), \quad \frac{\left|R_{2i}^{(T_1)}\right|}{\eta \left(T_1 - t_{inc}\right)} = \tilde{\mathcal{O}}\left(\sqrt{\eta} + \frac{1}{d^{\alpha-\frac{1}{2}}}\right).$$

The following lemma tells us that this approximate rank-1 structure is preserved when $T_1 \leq t \leq \tilde{T}$.

Lemma 15. *Under Assumption 1, 2 and 3, suppose* $\sigma \leq \frac{\eta^{3/2} \xi^2}{d^{13/4}}$. *By picking* $\eta \leq \mathcal{O}\left(\frac{1}{d^{3\alpha}}\right), \xi \leq \sqrt{\frac{\eta}{d^{3\alpha-1}}}$, *and* $\beta_2 = \beta_1^2$, *we have for* $T_1 \leq t < \tilde{T}$,

$$\forall i, j \in [d]: \quad w_{2i}^{(t)} = sign\left(w_{2i}^{(0)}\right) c^{(t)} + R_{2i}^{(t)},$$

$$W_1^{(t)}[i,j] = sign\left(w_{2i}^{(0)}\right) V_j^{(t)} + R_1^{(t)}[i,j],$$

$$\text{where} \quad \frac{\left|R_{2i}^{(t)}\right|}{\left|c^{(t)}\right|} = \mathcal{O}\left(\sqrt{\eta} + \frac{1}{d^{\alpha-1/2}}\right), \quad \frac{\left|R_1^{(t)}[i,j]\right|}{\left|V_j^{(t)}\right|} \leq \mathcal{O}\left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}}\right).$$

Moreover, when $t = \tilde{T}$, *we have* $L\left(W^{(t)}\right) \leq \mathcal{O}\left(\eta d^4\right)$.

Now we are ready to prove the Adam part of Theorem 1.

D.1 PROOF OF THE ADAM PART OF THEOREM 1

Define $T_{\text{Adam},1} = t_{inc} + \frac{1}{\eta d^{\frac{\alpha}{2}}}$. Note that this choice of $T_{\text{Adam},1}$ gives $\eta \left(T_{\text{Adam},1} - t_{inc}\right) = \frac{1}{d^{\frac{\alpha}{2}}}$. By picking $\eta \leq \mathcal{O}\left(\frac{1}{d^{3\alpha}}\right), \xi \leq \sqrt{\frac{\eta}{d^{3\alpha-1}}}$ and $\beta_2 = \beta_1^2$, we can apply Lemma 14 to get that $\forall i, j \in [d]:$ $\left|w_{2i}^{(T_{\text{Adam},1})}\right| = \Theta\left(\frac{1}{d^{\frac{\alpha}{2}}}\right), \left|W_1^{(T_{\text{Adam},1})}[i,j]\right| = \Theta\left(\frac{1}{d^{\frac{\alpha}{2}}}\right)$, and therefore $\forall i \in [d] : E_i^{(T_{\text{Adam},1})} = -\Theta(1)$ and $L\left(W^{(T_{\text{Adam},1})}\right) = \Theta(d)$. Define $T_{\text{Adam},2} = \tilde{T}$. By Lemma 15, we have $L\left(W^{(T_{\text{Adam},2})}\right) = \mathcal{O}\left(\eta d^4\right)$. For any $p > 0$, by picking $\alpha \geq \frac{p+4}{3}$, we have $L\left(W^{(T_{\text{Adam},2})}\right) = \mathcal{O}\left(\eta d^4\right) \leq \tilde{\mathcal{O}}\left(\frac{1}{d^p}\right)$.

Moreover, combining Lemma 14 and 15, we get that when $t \in [T_{\text{Adam},1}, T_{\text{Adam},2}]$, the conditions in Lemma 30 are satisfied with $\delta = \tilde{\mathcal{O}}\left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}}\right)$. The i-th component of the $\boldsymbol{u}$ vector (denoted as

u_i) is sign $\left(w_{2i}^{(0)} \right)$. That means $\forall i \in [d] : u_i^2 = 1$ and $\frac{\max_i (u_i)^2}{\text{median}(u_i)^2} = 1$. Then we can apply Lemma 30 and get that

$$R_{\text{med},1}^{\text{Adam}}(t), R_{\text{med},2}^{\text{Adam}}(t) \in \left[\left(\frac{1-\delta}{1+\delta} \right)^2 \frac{\max_i (u_i)^2}{\text{median}(u_i)^2}, \left(\frac{1+\delta}{1-\delta} \right)^2 \frac{\max_i (u_i)^2}{\text{median}(u_i)^2} \right]$$

$$= \left[\left(\frac{1-\delta}{1+\delta} \right)^2, \left(\frac{1+\delta}{1-\delta} \right)^2 \right],$$

$$\Rightarrow \quad R_{\text{med},1}^{\text{Adam}}(t), R_{\text{med},2}^{\text{Adam}}(t) = 1 \pm \mathcal{O}(\delta) = 1 \pm \tilde{\mathcal{O}} \left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2} - \frac{1}{4}}} \right).$$

D.2 PROOF OF LEMMA 14

Let's first denote $\Delta w_{2i}^{(t)} := w_{2i}^{(t+1)} - w_{2i}^{(t)}, \Delta W_1^{(t)}[i,j] := W_1^{(t+1)}[i,j] - W_1^{(t)}[i,j]$. Now we introduce two conditions.

Condition 1.

$$\forall \tau \in [H] : sign \left(g_1^{(t-\tau)}[i,j] \right) = s_1[i,j], \quad (1-\beta_1) \left| \sum_{\tau=0}^{H} \beta_1^{(\tau)} g_1^{(t-\tau)}[i,j] \right| \geq \Omega(\xi).$$

Condition 2.

$$\forall \tau \in [H] : sign \left(g_{2i}^{(t-\tau)} \right) = s_{2i}, \quad (1-\beta_1) \left| \sum_{\tau=0}^{H} \beta_1^{(\tau)} g_{2i}^{(t-\tau)} \right| \geq \Omega(\xi).$$

Next prove that, under Assumption 1 and 2, by picking $\eta \leq \mathcal{O} \left(\frac{1}{d^{3\alpha}} \right), \xi \leq \sqrt{\frac{\eta}{d^{3\alpha-1}}}$, and $\beta_2 = \beta_1^2$, there exists $t_{\text{inc}} > 0$ such that for $t_{\text{inc}} \leq t < T_1$, the weights can be approximated in the following way.

$$W_1^{(t+1)}[i,j] = W_1^{(t)}[i,j] - \eta \left(\text{sign} \left(g_1^{(t)}[i,j] \right) + e_1^{(t)}[i,j] \right),$$

$$w_{2i}^{(t+1)} = w_{2i}^{(t)} - \eta \left(\text{sign} \left(g_{2i}^{(t)} \right) + e_{2i}^{(t)} \right), \tag{28}$$

where $\left| e_1^{(t)}[i,j] \right| = \mathcal{O} \left(\sqrt{\eta} \right), \quad \left| e_{2i}^{(t)} \right| = \mathcal{O} \left(\sqrt{\eta} \right).$

Before we dive into the proof, let's introduce some useful lemmas.

The following lemma reflects our key idea: converting the exponential average in Adam to a finite-step average, and trying to bound the stochastic error terms in eq. (27).

Lemma 16. *Under Assumption 1, 2 and 3 and pick* $\beta_2 = \beta_1^2$. *Denote* $M_1^{(t)} := \max_{i,j \in [d], \tau \leq t} \left| W_1^{(\tau)}[i,j] \right|$, $M_2^{(t)} := \max_{i,j \in [d], \tau \leq t} \left| w_{2i}^{(\tau)} \right|$, $G_1^{(t)} := \max_{i,j \in [d], \tau \leq t} \left| g_1^{(\tau)}[i,j] \right|$ *and* $G_2^{(t)} := \max_{i,j \in [d], \tau \leq t} \left| g_{2i}^{(\tau)} \right|$. *We have that w.h.p., for all* $t \leq \mathcal{O} \left(\frac{1}{\sqrt{d\eta}} \right)$ *and* $\forall i, j \in [d]$,

$$\Delta W_1^{(t)}[i,j] = -\eta_t \frac{(1-\beta_1) \sum_{\tau=0}^{H} \beta_1^\tau g_1^{(t-\tau)}[i,j] + \epsilon_{1n}^{(t)}[i,j]}{\sqrt{(1-\beta_2) \sum_{\tau=0}^{H} \beta_2^\tau \left(g_1^{(t-\tau)}[i,j] \right)^2 + \epsilon_{1d}^{(t)}[i,j] + \xi}},$$

$$\Delta w_{2i}^{(t)} = -\eta_t \frac{(1-\beta_1) \sum_{\tau=0}^{H} \beta_1^\tau g_{2i}^{(t-\tau)} + \epsilon_{2n,i}^{(t)}}{\sqrt{(1-\beta_2) \sum_{\tau=0}^{H} \beta_2^\tau \left(g_{2i}^{(t-\tau)} \right)^2 + \epsilon_{2d,i}^{(t)} + \xi}},$$

where $H \geq \frac{1}{1-\beta_1} \log \frac{\max \left\{ G_1^{(t)}, \left(G_1^{(t)} \right)^2 \right\}}{\eta \xi^2}, H \geq \frac{1}{1-\beta_1} \log \frac{\max \left\{ G_2^{(t)}, \left(G_2^{(t)} \right)^2 \right\}}{\eta \xi^2}$ *and*

$$\left| \epsilon_{1n}^{(t)}[i,j] \right| \leq \mathcal{O}(\eta \xi^2) + \mathcal{O} \left(D_1^{(t)} \right), \quad \left| \epsilon_{1d}^{(t)}[i,j] \right| \leq \mathcal{O}(\eta \xi^2) + \mathcal{O} \left(D_1^{(t)} G_1^{(t)} + \left(D_1^{(t)} \right)^2 \right),$$

$$\left| \epsilon_{2n,i}^{(t)} \right| \leq \mathcal{O}(\eta \xi^2) + \mathcal{O} \left(D_2^{(t)} \right), \quad \left| \epsilon_{2d,i}^{(t)} \right| \leq \mathcal{O}(\eta \xi^2) + \mathcal{O} \left(D_2^{(t)} G_2^{(t)} + \left(D_2^{(t)} \right)^2 \right),$$

with

$$D_1^{(t)} \leq \mathcal{O}\left(d^3 M_1^{(t)} \left(M_2^{(t)}\right)^2 \sigma \sqrt{\frac{d^{1/2}}{\eta}}\right) + \mathcal{O}\left(M_2^{(t)} \sigma \sqrt{\frac{d^{3/2}}{\eta}}\right),$$

$$D_2^{(t)} \leq \mathcal{O}\left(d^4 \left(M_1^{(t)}\right)^2 M_2^{(t)} \sigma \sqrt{\frac{d^{1/2}}{\eta}}\right) + \mathcal{O}\left(d M_1^{(t)} \sigma \sqrt{\frac{d^{3/2}}{\eta}}\right).$$

Corollary 2. *Under the conditions of Lemma 16 and suppose* $\sigma \leq \frac{\eta^{3/2}\xi^2}{d^{13/4}}$. *Consider any* $t \leq \mathcal{O}\left(\frac{1}{\sqrt{d}\eta}\right)$. *If* $M_1^{(t)}, M_2^{(t)} \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), G_1^{(t)} \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), G_2^{(t)} \leq \mathcal{O}\left(\sqrt{d}\right)$, *then H in Lemma 16 can be picked as* $\frac{1}{1-\beta_1} \log \frac{d}{\eta\xi^2}$ *and we can get that* $\forall i, j \in [d], \left|\epsilon_{1n}^{(t)}[i,j]\right|, \left|\epsilon_{1d}^{(t)}[i,j]\right|, \left|\epsilon_{2n,i}^{(t)}\right|, \left|\epsilon_{2d,i}^{(t)}\right| \leq \mathcal{O}(\eta\xi^2)$.

The following lemma analyzes the magnitude of weights during a short period at the beginning.

Lemma 17. *Under Assumption 1, 2 and 3, suppose* $\sigma \leq \frac{\eta^{3/2}\xi^2}{d^{13/4}}$. *Pick* $\xi \leq \frac{1}{d^{\frac{3}{2}\alpha}}$, *then there exists some time point* $t_{inc} \in (0, T_1)$, *such that w.h.p., for* $t \leq t_{inc}$, *for every* $i, j \in [d]$,

$$\left|\Delta W_1^{(t)}[i,j]\right| \leq \tilde{\mathcal{O}}(\eta), \left|\Delta w_{2i}^{(t)}\right| \leq \tilde{\mathcal{O}}(\eta),$$

$$\left|W_1^{(t)}[i,j]\right| \leq \mathcal{O}\left(\frac{1}{d^{\frac{3}{2}\alpha+1}}\right), \Omega\left(\frac{1}{d^{\frac{3}{2}\alpha}}\right) \leq \left|w_{2i}^{(t)}\right| \leq \mathcal{O}\left(\frac{1}{d^{\alpha}}\right).$$

Specifically, when $t = t_{inc}$, *we have* $\text{sign}\left(w_{2i}^{(t_{inc})}\right) = \text{sign}\left(W_1^{(t_{inc})}[i,j]\right) = \text{sign}\left(w_{2i}^{(0)}\right)$, $\left|W_1^{(t_{inc})}[i,j]\right| = \Theta\left(\frac{1}{d^{\frac{3}{2}\alpha+1}}\right)$ *and* $\left|g_1^{(t_{inc})}[i,j]\right| \geq \Omega(\xi), \left|g_{2i}^{(t_{inc})}\right| \geq \Omega(\xi)$. *Moreover, Condition 1 and 2 are satisfied for* $t = t_{inc}$. *The* $s_1[i,j]$ *and* s_{2i} *in the conditions are both* $-\text{sign}\left(w_{2i}^{(0)}\right)$.

The following lemma gives us lower bounds of $\left|g_1^{(t)}[i,j]\right|$ and $\left|g_{2i}^{(t)}\right|$.

Lemma 18. *Under Assumption 1, 2 and 3, suppose* $\sigma \leq \frac{\eta^{3/2}\xi^2}{d^{13/4}}$. *Pick* $\xi \leq \sqrt{\frac{\eta}{d^{3\alpha-1}}}, \eta \leq \mathcal{O}\left(\frac{1}{d^{3\alpha}}\right)$. *Consider* t_{inc} *in Lemma 17. For any* $t \in [t_{inc}, T_1)$, *we have that* $\forall i, j \in [d] : \left|g_1^{(t)}[i,j]\right| \geq \Omega\left(\sqrt{\eta}\right), \left|g_{2i}^{(t)}\right| \geq \Omega\left(\sqrt{\eta}d\right)$.

Moreover, we have $\forall \tau \leq t, \forall i, j \in [d] : \left|W_1^{(\tau)}[i,j]\right| \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), \left|w_{2i}^{(\tau)}\right| \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), \left|g_1^{(\tau)}[i,j]\right| \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), \left|g_{2i}^{(\tau)}\right| \leq \mathcal{O}\left(\sqrt{d}\right)$.

The following lemma shows that when $t_{inc} \leq t < T_1$, we have $\forall i, j \in [d] : \left|g_{2i}^{(t)}\right| \gg \left|g_{2i}^{(t)} - g_{2i}^{(t-1)}\right|$ and that $\left|g_1^{(t)}[i,j]\right| \gg \left|g_1^{(t)}[i,j] - g_1^{(t-1)}[i,j]\right|$.

Lemma 19. *Under Assumption 1, 2 and 3, suppose* $\sigma \leq \frac{\eta^{3/2}\xi^2}{d^{13/4}}$. *Pick* $\xi \leq \sqrt{\frac{\eta}{d^{3\alpha-1}}}, \eta \leq \mathcal{O}\left(\frac{1}{d^{3\alpha}}\right)$. *For* t_{inc} *in Lemma 17, we have that for* $t_{inc} \leq t < T_1$ *and* $\tau \leq t, \forall i, j \in [d]$,

$$\frac{\left|g_1^{(t)}[i,j] - g_1^{(t-\tau)}[i,j]\right|}{\left|g_1^{(t)}[i,j]\right|} = \mathcal{O}\left(\sqrt{\eta}\tau\right), \quad \frac{\left|g_{2i}^{(t)} - g_{2i}^{(t-\tau)}\right|}{\left|g_{2i}^{(t)}\right|} = \mathcal{O}\left(\sqrt{\eta}\tau\right), \tag{29}$$

$$\frac{\left|\left(g_1^{(t)}[i,j]\right)^2 - \left(g_1^{(t-\tau)}[i,j]\right)^2\right|}{\left(g_1^{(t)}[i,j]\right)^2} = \mathcal{O}\left(\sqrt{\eta}\tau\right) + \mathcal{O}\left(\eta\tau^2\right), \quad \frac{\left|\left(g_{2i}^{(t)}\right)^2 - \left(g_{2i}^{(t-\tau)}\right)^2\right|}{\left(g_{2i}^{(t)}\right)^2} = \mathcal{O}\left(\sqrt{\eta}\tau\right) + \mathcal{O}\left(\eta\tau^2\right).$$

$$\tag{30}$$

Equipped with these lemmas, now let's prove eq. (28).

For any $t \in [t_{\mathrm{inc}}, T_1)$, by Lemma 18, we know that $M_1^{(t)}, M_2^{(t)} \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$, and that $G_1^{(t)} \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), G_2^{(t)} \leq \mathcal{O}\left(\sqrt{d}\right)$. At the end of the proof for this lemma, we will show that $T_1 = \Theta\left(\frac{1}{\sqrt{d}\eta}\right)$. Then we can pick $H := \frac{1}{1-\beta_1} \log \frac{d}{\eta\xi^2}$ and apply Lemma 16 and Corollary 2 to get that, w.h.p., for all $t \in [t_{\mathrm{inc}}, T_1)$ and $\forall i, j \in [d]$, eq. (26) can be written as

$$
\begin{aligned}
\Delta W_1^{(t)}[i,j] &= -\eta_t \frac{(1-\beta_1)\sum_{\tau=0}^{H} \beta_1^{\tau} g_1^{(t-\tau)}[i,j] + \epsilon_{1n}^{(t)}[i,j]}{\sqrt{(1-\beta_2)\sum_{\tau=0}^{H} \beta_2^{\tau} \left(g_1^{(t-\tau)}[i,j]\right)^2 + \epsilon_{1d}^{(t)}[i,j] + \xi}}, \\
\Delta w_{2i}^{(t)} &= -\eta_t \frac{(1-\beta_1)\sum_{\tau=0}^{H} \beta_1^{\tau} g_{2i}^{(t-\tau)} + \epsilon_{2n,i}^{(t)}}{\sqrt{(1-\beta_2)\sum_{\tau=0}^{H} \beta_2^{\tau} \left(g_{2i}^{(t-\tau)}\right)^2 + \epsilon_{2d,i}^{(t)} + \xi}},
\end{aligned}
\tag{31}
$$

where $\forall i, j \in [d]$, $\left|\epsilon_{1n}^{(t)}[i,j]\right|, \left|\epsilon_{1d}^{(t)}[i,j]\right|, \left|\epsilon_{2n,i}^{(t)}\right|, \left|\epsilon_{2d,i}^{(t)}\right| \leq \mathcal{O}(\eta\xi^2)$.

Let's first look at the update of $W_1^{(t)}[i,j]$. For t in the first phase, we write the RHS of eq. (31) as

$$
\begin{aligned}
&\frac{(1-\beta_1)\sum_{\tau=0}^{H} \beta_1^{\tau} g_1^{(t-\tau)}[i,j] + \epsilon_{1n}^{(t)}[i,j]}{\sqrt{(1-\beta_2)\sum_{\tau=0}^{H} \beta_2^{\tau} \left(g_1^{(t-\tau)}[i,j]\right)^2 + \epsilon_{1d}^{(t)}[i,j] + \xi}} \\
&= \frac{(1-\beta_1)g_1^{(t)}[i,j]\sum_{\tau=0}^{H} \beta_1^{\tau} + (1-\beta_1)\sum_{\tau=0}^{H} \beta_1^{\tau}\left(g_1^{(t-\tau)}[i,j] - g_1^{(t)}[i,j]\right) + \epsilon_{1n}^{(t)}[i,j]}{\sqrt{(1-\beta_2)\left(g_1^{(t)}[i,j]\right)^2 \sum_{\tau=0}^{H} \beta_2^{\tau} + (1-\beta_2)\sum_{\tau=0}^{H} \beta_2^{\tau}\left(\left(g_1^{(t-\tau)}[i,j]\right)^2 - \left(g_1^{(t)}[i,j]\right)^2\right) + \epsilon_{1d}^{(t)}[i,j] + \xi}} \\
&:= \frac{g_1^{(t)}[i,j](1-\beta_1^{H+1}) + e_{1n}^{(t)}[i,j] + \epsilon_{1n}^{(t)}[i,j]}{\sqrt{\left(g_1^{(t)}[i,j]\right)^2 (1-\beta_2^{H+1}) + e_{1d}^{(t)}[i,j] + \epsilon_{1d}^{(t)}[i,j] + \xi}},
\end{aligned}
$$

where

$$
\begin{aligned}
e_{1n}^{(t)}[i,j] &:= (1-\beta_1)\sum_{\tau=0}^{H} \beta_1^{\tau}\left(g_1^{(t-\tau)}[i,j] - g_1^{(t)}[i,j]\right), \\
e_{1d}^{(t)}[i,j] &:= (1-\beta_2)\sum_{\tau=0}^{H} \beta_2^{\tau}\left(\left(g_1^{(t-\tau)}[i,j]\right)^2 - \left(g_1^{(t)}[i,j]\right)^2\right).
\end{aligned}
$$

We have already shown that $\left|\epsilon_{1n}^{(t)}[i,j]\right|, \left|\epsilon_{1d}^{(t)}[i,j]\right| \leq \mathcal{O}(\eta\xi^2)$. By Lemma 19, we have that $\forall i, j \in [d]$,

$$
\begin{aligned}
\left|e_{1n}^{(t)}[i,j]\right| &\leq (1-\beta_1)\sum_{\tau=0}^{H} \beta_1^{\tau}\left|g_1^{(t-\tau)}[i,j] - g_1^{(t)}[i,j]\right| \\
&\leq \left|g_1^{(t)}[i,j]\right|\mathcal{O}\left(\sqrt{\eta}\right)(1-\beta_1)\sum_{\tau=0}^{H} \beta_1^{\tau}\tau = \left|g_1^{(t)}[i,j]\right|\mathcal{O}\left(\sqrt{\eta}\right).
\end{aligned}
$$

Similarly, we have $\forall i, j \in [d]$,

$$
\begin{aligned}
\left|e_{1d}^{(t)}[i,j]\right| &\leq \left(g_1^{(t)}[i,j]\right)^2 \mathcal{O}\left(\sqrt{\eta}\right)(1-\beta_2)\sum_{\tau=0}^{H} \beta_1^{\tau}\tau + \left(g_1^{(t)}[i,j]\right)^2 \mathcal{O}\left(\sqrt{\eta}\right)(1-\beta_2)\sum_{\tau=0}^{H} \beta_1^{\tau}\tau^2 \\
&= \left(g_1^{(t)}[i,j]\right)^2 \mathcal{O}\left(\sqrt{\eta}\right).
\end{aligned}
$$

By Lemma 18, we know that $\left|g_1^{(t)}[i,j]\right| = \Omega\left(\sqrt{\eta}\right)$. Then we have that

$$\forall i,j \in [d]: \quad \left|\epsilon_{1n}^{(t)}[i,j]\right| \leq \mathcal{O}(\eta \xi^2) \leq \mathcal{O}\left(\sqrt{\eta}\right)\left|g_1^{(t)}[i,j]\right|, \quad \left|\epsilon_{1d}^{(t)}[i,j]\right| \leq \mathcal{O}\left(\sqrt{\eta}\right)\xi^2.$$

Therefore by Lemma 33 in Appendix G, we have

$$\frac{g_1^{(t)}[i,j]\left(1-\beta_1^{H+1}\right) + e_{1n}^{(t)}[i,j] + \epsilon_{1n}^{(t)}[i,j]}{\sqrt{\left(g_1^{(t)}[i,j]\right)^2\left(1-\beta_2^{H+1}\right) + e_{1d}^{(t)}[i,j] + \epsilon_{1d}^{(t)}[i,j] + \xi}} = \frac{1-\beta_1^{H+1}}{\sqrt{1-\beta_2^{H+1}}}\left(\text{sign}\left(g_1^{(t)}[i,j]\right) + e_1^{(t)}[i,j]\right),$$

where $\left|e_1^{(t)}[i,j]\right| = \mathcal{O}\left(\sqrt{\eta}\right)$. Combining all of the above yields that

$$\begin{aligned}
W_1^{(t+1)}[i,j] &= W_1^{(t)}[i,j] - \eta_t \frac{(1-\beta_1)\sum_{\tau=0}^t \beta_1^\tau g_1^{(t-\tau)}[i,j]}{\sqrt{(1-\beta_2)\sum_{\tau=0}^t \beta_2^\tau \left(g_1^{(t-\tau)}[i,j]\right)^2 + \xi}} \\
&= W_1^{(t)}[i,j] - \eta_t \frac{1-\beta_1^{H+1}}{\sqrt{1-\beta_2^{H+1}}}\left(\text{sign}\left(g_1^{(t)}[i,j]\right) + e_1^{(t)}[i,j]\right) \\
&= W_1^{(t)}[i,j] - \eta\left(\text{sign}\left(g_1^{(t)}[i,j]\right) + e_1^{(t)}[i,j]\right),
\end{aligned}$$

where $\left|e_1^{(t)}[i,j]\right| = \mathcal{O}\left(\sqrt{\eta}\right)$. The proof for $w_{2i}^{(t)}$ is similar.

So far we have successfully proved eq. (28), which gives us

$$\begin{aligned}
\forall i,j \in [d]: \quad w_{2i}^{(t)} &= \text{sign}\left(w_{2i}^{(0)}\right)\eta\left(t - t_{\text{inc}}\right) + R_{2i}^{(t)}, \\
W_1^{(t)}[i,j] &= \text{sign}\left(w_{2i}^{(0)}\right)\eta\left(t - t_{\text{inc}}\right) + R_1^{(t)}[i,j],
\end{aligned}$$

where $\frac{\left|R_1^{(t)}[i,j]\right|}{\eta(t-t_{\text{inc}})} = \tilde{\mathcal{O}}\left(\sqrt{\eta} + \frac{\left|W_1^{(t_{\text{inc}})}[i,j]\right|}{\eta(t-t_{\text{inc}})}\right)$ and $\frac{\left|R_{2i}^{(t)}\right|}{\eta(t-t_{\text{inc}})} = \tilde{\mathcal{O}}\left(\sqrt{\eta} + \frac{\left|w_{2i}^{(t_{\text{inc}})}\right|}{\eta(t-t_{\text{inc}})}\right)$. Now it suffices to show that $\forall i,j \in [d]: \left|w_{2i}^{(t_{\text{inc}})}\right| \leq \tilde{\mathcal{O}}\left(\frac{1}{d^\alpha}\right), \left|W_1^{(t_{\text{inc}})}[i,j]\right| \leq \mathcal{O}\left(\frac{1}{d^\alpha}\right)$, which is implied by Lemma 17.

When $t = T_1$, we have $\forall j \in [d]: \sum_{i=1}^d w_{2i}^{(T_1)} W_1^{(T_1)}[i,j] = \Theta(1)$. Combining with the above results, we know that $d\eta^2(T_1 - t_{\text{inc}})^2 = \Theta(1)$, i.e. $\eta(T_1 - t_{\text{inc}}) = \Theta\left(\frac{1}{\sqrt{d}}\right)$. In Section D.5, we will prove $t_{\text{inc}} = \Theta\left(\frac{1}{\eta d^{\frac{3}{2}\alpha+1}}\right)$. Then we have $T_1 = \Theta\left(\frac{1}{\sqrt{d}\eta}\right)$.

D.3 PROOF OF LEMMA 16

For certain t and H, we write eq. (26) as

$$\begin{aligned}
\Delta W_1^{(t)}[i,j] &= -\eta_t \frac{(1-\beta_1)\sum_{\tau=0}^H \beta_1^\tau g_1^{(t-\tau)}[i,j] + \epsilon_{1n}^{(t)}[i,j]}{\sqrt{(1-\beta_2)\sum_{\tau=0}^H \beta_2^\tau \left(g_1^{(t-\tau)}[i,j]\right)^2 + \epsilon_{1d}^{(t)}[i,j] + \xi}}, \\
\Delta w_{2i}^{(t)} &= -\eta_t \frac{(1-\beta_1)\sum_{\tau=0}^H \beta_1^\tau g_{2i}^{(t-\tau)} + \epsilon_{2n,i}^{(t)}}{\sqrt{(1-\beta_2)\sum_{\tau=0}^H \beta_2^\tau \left(g_{2i}^{(t-\tau)}\right)^2 + \epsilon_{2d,i}^{(t)} + \xi}},
\end{aligned}$$

where

$$\epsilon_{1n}^{(t)}[i,j] := (1-\beta_1)\underbrace{\sum_{\tau=H+1}^{t}\beta_1^\tau g_1^{(t-\tau)}[i,j]+r_{1n}^{(t)}[i,j]}_{:=q_{1n}^{(t)}[i,j]}, \epsilon_{2n,i}^{(t)} := (1-\beta_1)\underbrace{\sum_{\tau=H+1}^{t}\beta_1^\tau g_{2i}^{(t-\tau)}+r_{2n,i}^{(t)}}_{:=q_{2n,i}^{(t)}}$$

$$\epsilon_{1d}^{(t)}[i,j] := (1-\beta_2)\underbrace{\sum_{\tau=H+1}^{t}\beta_2^\tau\left(g_1^{(t-\tau)}[i,j]\right)^2+r_{1d}^{(t)}[i,j]}_{:=q_{1d}^{(t)}[i,j]}, \epsilon_{2d,i}^{(t)} = (1-\beta_2)\underbrace{\sum_{\tau=H+1}^{t}\beta_2^\tau\left(g_{2i}^{(t-\tau)}\right)^2+r_{2d,i}^{(t)}}_{:=q_{2d,i}^{(t)}}$$

and $r_{1n}^{(t)}[i,j], r_{1d}^{(t)}[i,j], r_{2n,i}^{(t)}, r_{2d,i}^{(t)}$ are defined in eq. (27).

Since $\beta_2 = \beta_1^2 < \beta_1$, then if we pick $H \geq \frac{1}{1-\beta_1}\log\frac{\max\left\{G_1^{(t)},\left(G_1^{(t)}\right)^2\right\}}{\eta\xi^2}$, $H \geq \frac{1}{1-\beta_1}\log\frac{\max\left\{G_2^{(t)},\left(G_2^{(t)}\right)^2\right\}}{\eta\xi^2}$, then we can get that $H \geq \frac{1}{1-\beta_1}\log\frac{G_1^{(t)}}{\eta\xi^2}, H \geq \frac{1}{1-\beta_2}\log\frac{\left(G_1^{(t)}\right)^2}{\eta\xi^2}$, $H \geq \frac{1}{1-\beta_1}\log\frac{G_2^{(t)}}{\eta\xi^2}, H \geq \frac{1}{1-\beta_2}\log\frac{\left(G_2^{(t)}\right)^2}{\eta\xi^2}$. Hence we can apply Lemma 32 in Appendix G to get that $\left|q_{1n}^{(t)}[i,j]\right|, \left|q_{1d}^{(t)}[i,j]\right|, \left|q_{2n,i}^{(t)}\right|, \left|q_{2d,i}^{(t)}\right| \leq \eta\xi^2$.

Pick T in Lemma 31 as of order $\mathcal{O}\left(\frac{1}{\sqrt{d}\eta}\right)$. By Lemma 31, we have with probability at least $1-\frac{1}{d}$, for all $t \leq T, \forall \tau \leq t$ and $\forall i, j \in [d]$,

$$\left|Dg_1^{(\tau)}[i,j]\right| = \left|\tilde{g}_1^{(\tau)}[i,j] - g_1^{(\tau)}[i,j]\right| \leq \mathcal{O}\left(d^3 M_1^{(t)}\left(M_2^{(t)}\right)^2\sigma\sqrt{\frac{d^{1/2}}{\eta}}\right) + \mathcal{O}\left(M_2^{(t)}\sigma\sqrt{\frac{d^{3/2}}{\eta}}\right)$$

$$:= D_1^{(t)}$$

$$\left|Dg_{2i}^{(\tau)}\right| = \left|\tilde{g}_{2i}^{(\tau)} - g_{2i}^{(\tau)}\right| \leq \mathcal{O}\left(d^4\left(M_1^{(t)}\right)^2 M_2^{(t)}\sigma\sqrt{\frac{d^{1/2}}{\eta}}\right) + \mathcal{O}\left(dM_1^{(t)}\sigma\sqrt{\frac{d^{3/2}}{\eta}}\right)$$

$$:= D_2^{(t)}$$

Plugging into eq. (27) gives us

$$\left|r_{1n}^{(t)}[i,j]\right| \leq (1-\beta_1)\sum_{\tau=0}^{t}\beta_1^{t-\tau}\left|Dg_1^{(\tau)}[i,j]\right| \leq \mathcal{O}\left(D_1^{(t)}\right),$$

$$\left|r_{1d}^{(t)}[i,j]\right| \leq (1-\beta_2)\sum_{\tau=0}^{t}\beta_2^{t-\tau}\left|2g_1^{(\tau)}[i,j]Dg_1^{(\tau)}[i,j]\right| + \left|Dg_1^{(\tau)}[i,j]\right|^2 \leq \mathcal{O}\left(D_1^{(t)}G_1^{(t)} + \left(D_1^{(t)}\right)^2\right),$$

$$\left|r_{2n,i}^{(t)}\right| \leq (1-\beta_1)\sum_{\tau=0}^{t}\beta_1^{t-\tau}\left|Dg_{2i}^{(\tau)}\right| \leq \mathcal{O}\left(D_2^{(t)}\right),$$

$$\left|r_{2d,i}^{(t)}\right| \leq (1-\beta_2)\sum_{\tau=0}^{t}\beta_2^{t-\tau}\left|2g_{2i}^{(\tau)}Dg_{2i}^{(\tau)}\right| + \left|Dg_{2i}^{(\tau)}\right|^2 \leq \mathcal{O}\left(D_2^{(t)}G_2^{(t)} + \left(D_2^{(t)}\right)^2\right)$$

D.4 PROOF OF COROLLARY 2

Since $G_1^{(t)} \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), G_2^{(t)} \leq \mathcal{O}\left(\sqrt{d}\right)$, then $H := \frac{1}{1-\beta_1}\log\frac{d}{\eta\xi^2}$ is bigger than $\frac{1}{1-\beta_1}\log\frac{\max\left\{G_1^{(t)},\left(G_1^{(t)}\right)^2\right\}}{\eta\xi^2}$ and $\frac{1}{1-\beta_1}\log\frac{\max\left\{G_2^{(t)},\left(G_2^{(t)}\right)^2\right\}}{\eta\xi^2}$.

By $M_1^{(t)}, M_2^{(t)} \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), G_1^{(t)} \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), G_2^{(t)} \leq \mathcal{O}\left(\sqrt{d}\right)$ and the assumption $\sigma \leq \frac{\eta^{3/2}\xi^2}{d^{13/4}}$, we get that $D_1^{(t)}$ and $D_2^{(t)}$ are upper bounded by $D_1^{(t)} \leq \mathcal{O}\left(d^{7/4}\sigma\eta^{-1/2}\right)$ and $D_2^{(t)} \leq \mathcal{O}\left(d^{11/4}\sigma\eta^{-1/2}\right)$, which yields $\forall i, j \in [d], \left|\epsilon_{1n}^{(t)}[i,j]\right|, \left|\epsilon_{1d}^{(t)}[i,j]\right|, \left|\epsilon_{2n,i}^{(t)}\right|, \left|\epsilon_{2d,i}^{(t)}\right| \leq \mathcal{O}(\eta\xi^2)$.

D.5 PROOF OF LEMMA 17

The proof is based on the following two lemmas.

Lemma 20. *Under Assumption 1 and 2, we have that w.p. at least* $1 - \frac{1}{d^{\frac{\alpha}{2}-1}}$, *for every* $1 \leq i \leq d$, $\frac{\sqrt{\pi}}{d^{\frac{3}{2}\alpha}} \leq \left|w_{2i}^{(0)}\right| \leq \sqrt{\frac{2}{d^{2\alpha}} \log \frac{2d}{\delta}}$, *and that w.p. at least* $1 - \delta$ *for any given* $\delta > 0$, $\left|W_1^{(0)}[i,j]\right| \leq \sqrt{\frac{2}{d^{4\alpha}} \log \frac{2d^2}{\delta}}$.

Lemma 21. *Under Assumption 1, 2 and 3, suppose* $\sigma \leq \frac{\eta^{3/2}\xi^2}{d^{13/4}}$. *Pick* $\beta_2 = \beta_1^2, \xi \in (0, 1), \eta < \frac{1}{4}$. *Consider any time point* $t \leq \mathcal{O}\left(\frac{1}{\sqrt{d}\eta}\right)$. *If* $\forall \tau \leq t, \forall i, j \in [d] : \left|W_1^{(\tau)}[i,j]\right| \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), \left|w_{2i}^{(\tau)}\right| \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$ *and* $\left|g_1^{(\tau)}[i,j]\right| \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), \left|g_{2i}^{(\tau)}\right| \leq \mathcal{O}\left(\sqrt{d}\right)$, *we will have*

$$\left|\Delta W_1^{(t)}[i,j]\right| \leq \tilde{\mathcal{O}}(\eta) \quad \left|\Delta w_{2i}^{(t)}\right| \leq \tilde{\mathcal{O}}(\eta),$$

where the $\tilde{\mathcal{O}}$ *notation depends on* $H = \frac{1}{1-\beta_1} \log \frac{d}{\eta\xi^2}$.

Furthermore, if for certain $i, j \in [d]$, *Condition 1 (resp. Condition 2) is satisfied, we will have*

$$\text{sign}\left(\Delta W_1^{(t)}[i,j]\right) = -s_1[i,j], \left|\Delta W_1^{(t)}[i,j]\right| = \Theta(\eta)$$

$$\left(\text{resp. sign}\left(\Delta w_{2i}^{(t)}\right) = -s_{2i}, \left|\Delta w_{2i}^{(t)}\right| = \Theta(\eta)\right).$$

Now we prove Lemma 17. Define $t_d := \inf\left\{t : \exists i, j : \left|W_1^{(t)}[i,j]\right| > \frac{1}{d} \text{ or } \left|w_{2i}^{(t)}\right| > \frac{1}{d}\right\}$. Now we want to find a time point t_{inc} before t_d for the lemma to hold. During the period $t < t_d$, we have $\forall j \in [d], E_j = -\Theta(1)$ (which means $t_d < T_1$) and therefore for all $i, j \in [d], \left|g_1^{(t)}[i,j]\right| \leq \frac{1}{d}$ and $\left|g_{2i}^{(t)}\right| \leq 1$. Then we can use Lemma 21 to get that for $t \leq \min\left\{t_d, \frac{1}{\sqrt{d}\eta}\right\}$, we have $\left|\Delta W_1^{(t)}[i,j]\right| \leq \tilde{\mathcal{O}}(\eta), \left|\Delta w_{2i}^{(t)}\right| \leq \tilde{\mathcal{O}}(\eta)$. Hence $t_d \geq \tilde{\Omega}\left(\frac{1}{\eta d}\right)$.

Define $t_{\text{sign}} = \inf\left\{t : \exists i \in [d] : \left|w_{2i}^{(t)}\right| \leq \frac{1}{d^{\frac{3}{2}\alpha}}\right\}$. By Lemma 20, w.h.p. $\forall i \in [d] : \left|w_{2i}^{(0)}\right| \geq \frac{\sqrt{\pi}}{d^{\frac{3}{2}\alpha}}$, combining with $\left|\Delta w_{2i}^{(t)}\right| \leq \tilde{\mathcal{O}}(\eta)$ gives us that w.h.p., $t_{\text{sign}} \geq \frac{\sqrt{\pi}-1}{d^{\frac{3}{2}\alpha}} / \tilde{\mathcal{O}}(\eta) = \tilde{\Omega}\left(\frac{1}{\eta d^{\frac{3}{2}\alpha}}\right)$.

Now let's analyze the behavior of W_1 during the period $t < \min\left\{t_d, t_{\text{sign}}, \frac{1}{\sqrt{d}\eta}\right\}$. Consider any t s.t. $t < \min\left\{t_d, t_{\text{sign}}, \frac{1}{\sqrt{d}\eta}\right\}$ and any $i, j \in [d]$. By definition, $\text{sign}\left(w_{2i}^{(t)}\right) = \text{sign}\left(w_{2i}^{(0)}\right)$. Note that $E_j^{(t)} < 0$, then we have $\text{sign}\left(g_1^{(t)}[i,j]\right) = -\text{sign}\left(w_{2i}^{(0)}\right)$ and that $\left|g_1^{(t)}[i,j]\right| = \Omega\left(\frac{1}{d^{\frac{3}{2}\alpha}}\right) = \Omega(\xi)$. By our choice of ξ, we know that Condition 1 is satisfied with $s_1[i,j] = -\text{sign}\left(w_{2i}^{(0)}\right)$ (for all $t \leq \min\{t_d, t_{\text{sign}}\}$), which yields $\text{sign}\left(\Delta W_1^{(t)}[i,j]\right) = \text{sign}\left(w_{2i}^{(0)}\right)$ and $\left|\Delta W_1^{(t)}[i,j]\right| = \Theta(\eta)$.

Lemma 20 tells us that w.h.p., $\forall i, j \in [d] : \left|W_1^{(0)}[i,j]\right| = \tilde{\mathcal{O}}\left(\frac{1}{d^{2\alpha}}\right)$. For any i, j, if initially $\text{sign}\left(W_1^{(0)}[i,j]\right) = \text{sign}\left(w_{2i}^{(0)}\right)$, then for the following steps before $\min\left\{t_d, t_{\text{sign}}, \frac{1}{\sqrt{d}\eta}\right\}$, we will have $\text{sign}\left(W_1^{(t)}[i,j]\right) = \text{sign}\left(w_{2i}^{(0)}\right)$. If initially $\text{sign}\left(W_1^{(0)}[i,j]\right) \neq \text{sign}\left(w_{2i}^{(0)}\right)$, then after at most $t_0 = \tilde{\mathcal{O}}\left(\frac{1}{\eta d^{2\alpha}}\right)$ steps, $W_1[i,j]$ will flip the sign. Note that $t_0 = \tilde{\mathcal{O}}\left(\frac{1}{\eta d^{2\alpha}}\right)$ is smaller than $\min\left\{t_d, t_{\text{sign}}, \frac{1}{\sqrt{d}\eta}\right\}$.

Hence we have shown that at some time point t_0, we have $\forall i, j \in [d] : \text{sign}\left(W_1^{(t)}[i,j]\right) = \text{sign}\left(w_{2i}^{(t)}\right) = \text{sign}\left(w_{2i}^{(0)}\right)$. Now we analyze the period $t \geq t_0$.

When $t_0 < t \leq \min\left\{t_d, t_{\text{sign}}, \frac{1}{\sqrt{d}\eta}\right\}$, we still have $\text{sign}\left(\Delta W_1^{(t)}[i,j]\right) = \text{sign}\left(w_{2i}^{(0)}\right)$ and $\left|\Delta W_1^{(t)}[i,j]\right| = \Theta(\eta)$. Combining these two with the fact $\text{sign}\left(W_1^{(t_0)}[i,j]\right) = \text{sign}\left(w_{2i}^{(0)}\right)$, we know that for all $t \in \left[t_0, \min\left\{t_d, t_{\text{sign}}, \frac{1}{\sqrt{d}\eta}\right\}\right]$, $\text{sign}\left(W_1^{(t)}[i,j]\right) = \text{sign}\left(w_{2i}^{(0)}\right)$ and that $\forall i, j \in [d] : \left|W_1^{(t+1)}[i,j]\right| = \left|W_1^{(t)}[i,j]\right| + \Theta(\eta)$. Then at certain step t_{inc} which satisfies $t_{\text{inc}} = t_0 + \Theta\left(\frac{1}{\eta d^{\frac{3}{2}\alpha+1}}\right) < \min\left\{t_d, t_{\text{sign}}, \frac{1}{\sqrt{d}\eta}\right\}$, we will have $\forall t_{\text{inc}} - H \leq \tau \leq t_{\text{inc}}, \forall i, j \in [d] : \left|W_1^{(\tau)}[i,j]\right| = \Theta\left(\frac{1}{d^{\frac{3}{2}\alpha+1}}\right)$ and therefore $\left|g_{2i}^{(\tau)}\right| = \left|\sum_{j=1}^d W_1^{(\tau)}[i,j]E_j^{(\tau)}\right| = \sum_{j=1}^d \left|W_1^{(\tau)}[i,j]E_j^{(\tau)}\right| = \Theta\left(\frac{1}{d^{\frac{3}{2}\alpha}}\right) = \Omega(\xi)$. For $t \leq t_{\text{inc}}$, we have $\forall i, j \in [d] : \left|W_1^{(t)}[i,j]\right| = \mathcal{O}\left(\frac{1}{d^{\frac{3}{2}\alpha+1}}\right)$.

Since $t_{\text{inc}} < t_{\text{sign}}$, we have $\left|w_{2i}^{t_{\text{inc}}}\right| = \Omega\left(\frac{1}{d^{\frac{3}{2}\alpha}}\right)$. For $t \leq t_{\text{inc}}$, note that $\left|\Delta w_{2i}^{(t)}\right| \leq \tilde{\mathcal{O}}(\eta)$, $t_{\text{inc}} = t_0 + \Theta\left(\frac{1}{\eta d^{\frac{3}{2}\alpha+1}}\right) = \Theta\left(\frac{1}{\eta d^{\frac{3}{2}\alpha+1}}\right)$, combining with the upper bound in Lemma 20 yields

$$\left|w_{2i}^{(t)}\right| \leq \left|w_{2i}^{(0)}\right| + t_{\text{inc}}\tilde{\mathcal{O}}(\eta) \leq \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{3}{2}\alpha+1}}\right) \leq \tilde{\mathcal{O}}\left(\frac{1}{d^{\alpha}}\right).$$

Moreover, $\forall t_{\text{inc}} - H \leq \tau \leq t_{\text{inc}}, \forall i \in [d] : \text{sign}\left(g_{2i}^{(\tau)}\right) = -\text{sign}\left(w_{2i}^{(0)}\right)$. Then Condition 2 is satisfied for $t = t_{\text{inc}}$. In the analysis of $g_1^{(t)}[i,j]$, we have already shown that for all $t \leq \min\left\{t_d, t_{\text{sign}}, \frac{1}{\sqrt{d}\eta}\right\}$ (and thus for $t = t_{\text{inc}}$), Condition 1 is satisfied, which completes the proof.

D.6 PROOF OF LEMMA 20

Since for $X \sim \mathcal{N}\left(0, \sigma^2\right)$, we have that $P(|X| \leq t) \leq \frac{2t}{\sqrt{2\pi}\sigma}$, then for a fixed i,

$$P\left(\left|w_{2i}^{(0)}\right| \leq \frac{\sqrt{\pi}}{d^{\frac{3}{2}\alpha}}\right) \leq \frac{2\sqrt{\pi}/d^{\frac{3}{2}\alpha}}{\sqrt{2\pi} \cdot \sqrt{2/d^{2\alpha}}} = \frac{1}{d^{\frac{\alpha}{2}}}.$$

Then by union bound, we have that w.p. at least $1 - \frac{1}{d^{\frac{\alpha}{2}-1}}$, for every $1 \leq i \leq d$, $\left|w_{2i}^{(0)}\right| \geq \frac{\sqrt{\pi}}{d^{\frac{3}{2}\alpha}}$.

As for the upper bounds, using the Gaussian tail bound and union bound, we have w.p. at least $1 - \delta$,

$$\forall i, j \in [d] : \quad \left|w_{2i}^{(0)}\right| \leq \sqrt{\frac{2}{d^{2\alpha}} \log \frac{2d}{\delta}}, \quad \left|W_1^{(0)}[i,j]\right| \leq \sqrt{\frac{2}{d^{4\alpha}} \log \frac{2d^2}{\delta}}.$$

D.7 PROOF OF LEMMA 21

Now we analyze the magnitude order of $\Delta W_1^{(t)}[i,j]$. The analysis of $\Delta w_{2i}^{(t)}$ is similar.

For $t \leq \mathcal{O}\left(\frac{1}{\sqrt{d}\eta}\right)$. By assumption, $M_1^{(t)}, M_2^{(t)} \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), G_1^{(t)} \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), G_2^{(t)} \leq \mathcal{O}\left(\sqrt{d}\right)$, and $\sigma \leq \frac{\eta^{3/2}\xi^2}{d^{13/4}}$. Hence we can pick $H := \frac{1}{1-\beta_1} \log \frac{d}{\eta\xi^2}$ and apply Lemma 16 and Corollary 2 to get that,

w.h.p., for all $t \in [t_{\text{inc}}, T_1)$ and $\forall i, j \in [d]$, eq. (26) can be written as

$$
\begin{aligned}
\Delta W_1^{(t)}[i,j] &= -\eta_t \frac{(1-\beta_1)\sum_{\tau=0}^{H}\beta_1^\tau g_1^{(t-\tau)}[i,j] + \epsilon_{1n}^{(t)}[i,j]}{\sqrt{(1-\beta_2)\sum_{\tau=0}^{H}\beta_2^\tau\left(g_1^{(t-\tau)}[i,j]\right)^2 + \epsilon_{1d}^{(t)}[i,j] + \xi}}, \\
\Delta w_{2i}^{(t)} &= -\eta_t \frac{(1-\beta_1)\sum_{\tau=0}^{H}\beta_1^\tau g_{2i}^{(t-\tau)} + \epsilon_{2n,i}^{(t)}}{\sqrt{(1-\beta_2)\sum_{\tau=0}^{H}\beta_2^\tau\left(g_{2i}^{(t-\tau)}\right)^2 + \epsilon_{2d,i}^{(t)} + \xi}},
\end{aligned}
\tag{32}
$$

where $\forall i, j \in [d]$, $\left|\epsilon_{1n}^{(t)}[i,j]\right|, \left|\epsilon_{1d}^{(t)}[i,j]\right|, \left|\epsilon_{2n,i}^{(t)}\right|, \left|\epsilon_{2d,i}^{(t)}\right| \le \mathcal{O}(\eta\xi^2)$.

On one hand, using $\left|\epsilon_{1n}^{(t)}[i,j]\right|, \left|\epsilon_{1d}^{(t)}[i,j]\right| \le \eta\xi^2$ and $\beta_2 = \beta_1^2$, and $\sqrt{x+y} \ge \sqrt{x} - \sqrt{|y|}$ when $x \ge 0, x+y \ge 0$, we get from eq. (32) that

$$
\begin{aligned}
\left|\Delta W_1^{(t)}[i,j]\right| &\le \eta_t \frac{(1-\beta_1)\left|\sum_{\tau=0}^{H}\beta_1^\tau g_1^{(t-\tau)}[i,j]\right| + \eta\xi^2}{\sqrt{(1-\beta_2)\sum_{\tau=0}^{H}\left(\beta_1^\tau g_1^{(t-\tau)}[i,j]\right)^2 - \sqrt{\eta}\xi + \xi}} \\
&\overset{(i)}{\le} \eta_t \frac{(1-\beta_1)\sqrt{H+1}\sqrt{\sum_{\tau=0}^{H}\left(\beta_1^\tau g_1^{(t-\tau)}[i,j]\right)^2} + \eta\xi^2}{\sqrt{(1-\beta_2)\sum_{\tau=0}^{H}\left(\beta_1^\tau g_1^{(t-\tau)}[i,j]\right)^2 + \xi/2}} \le \mathcal{O}(\eta),
\end{aligned}
$$

where (i) uses Cauchy-Schwarz inequality.

On the other hand, when $\text{sign}\left(g_1^{(t-H)}[i,j]\right) = \text{sign}\left(g_1^{(t-H+1)}[i,j]\right) = \ldots = \text{sign}\left(g_1^{(t)}[i,j]\right) = s_1[i,j]$, we have

$$
\text{sign}\left(\sum_{\tau=0}^{H}\beta_1^\tau g_1^{(t-\tau)}[i,j]\right) = s_1[i,j], \quad \left|\sum_{\tau=0}^{H}\beta_1^\tau g_1^{(t-\tau)}[i,j]\right| \ge \sqrt{\sum_{\tau=0}^{H}\left(\beta_1^\tau g_1^{(t-\tau)}[i,j]\right)^2}.
$$

If we further have $(1-\beta_1)\left|\sum_{\tau=0}^{H}\beta_1^{(\tau)}g_1^{(t-\tau)}[i,j]\right| \ge \Omega(\xi)$, then combining with $\left|\epsilon_{1n}^{(t)}[i,j]\right| \le \eta\xi^2 < \xi$ we will get

$$
\text{sign}\left(\Delta W_1^{(t)}[i,j]\right) = -\text{sign}\left(\sum_{\tau=0}^{H}\beta_1^\tau g_1^{(t-\tau)}[i,j] + \epsilon_{1n}^{(t)}[i,j]\right) = -\text{sign}\left(\sum_{\tau=0}^{H}\beta_1^\tau g_1^{(t-\tau)}[i,j]\right) = -s_1[i,j].
$$

Using $\sqrt{x+y} \le \sqrt{|x|} + \sqrt{|y|}$, we obtain that

$$
\begin{aligned}
\left|\Delta W_1^{(t)}[i,j]\right| &\ge \eta_t \frac{(1-\beta_1)\left|\sum_{\tau=0}^{H}\beta_1^{(\tau)}g_1^{(t-\tau)}[i,j]\right| - \eta\xi^2}{\sqrt{(1-\beta_2)\sum_{\tau=0}^{H}\left(\beta_1^\tau g_1^{(t-\tau)}[i,j]\right)^2 + \sqrt{\eta}\xi + \xi}} \\
&\ge \eta_t \frac{\frac{1-\beta_1}{2}\left|\sum_{\tau=0}^{H}\beta_1^{(\tau)}g_1^{(t-\tau)}[i,j]\right|}{2\max\left\{\sqrt{(1-\beta_2)\sum_{\tau=0}^{H}\left(\beta_1^\tau g_1^{(t-\tau)}[i,j]\right)^2}, \frac{3}{2}\xi\right\}} = \Omega(\eta).
\end{aligned}
$$

D.8 PROOF OF LEMMA 18

The proof is based on the following lemma, which gives a coarse analysis on the magnitude of weights and their increments per step during the first phase.

Lemma 22. *Under Assumption 1, 2 and 3, suppose $\sigma \leq \frac{\eta^{3/2}\xi^2}{d^{13/4}}$. Pick $\xi \leq \min\left\{\sqrt{\frac{\eta}{d^{3\alpha-1}}}, \frac{1}{d^{\frac{3}{2}\alpha}}\right\}$, for t_{inc} in Lemma 17, we have that for all $t_{inc} \leq t < T_1$, $\forall i, j \in [d]$.*

$$sign\left(\Delta W_1^{(t)}[i,j]\right) = sign\left(\Delta w_{2i}^{(t)}\right) = sign\left(w_{2i}^{(0)}\right), \quad \left|\Delta W_1^{(t)}[i,j]\right| = \Theta(\eta), \left|\Delta w_{2i}^{(t)}\right| = \Theta(\eta),$$

$$sign\left(W_1^{(t)}[i,j]\right) = sign\left(w_{2i}^{(t)}\right) = sign\left(w_{2i}^{(0)}\right), \quad \left|W_1^{(t)}[i,j]\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), \left|w_{2i}^{(t)}\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right).$$

Specially, at the end of the first phase ($t = T_1$), we have $\forall i, j \in [d]$, $\left|w_{2i}^{(T_1)}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right), \left|W_1^{(T_1)}[i,j]\right| = \Theta\left(\frac{1}{\sqrt{d}}\right).$

Now we go back to the proof of Lemma 18. For $t_{\text{inc}} \leq t < T_1$, since $E_j^{(t)} = \left(W_2^{(t)} W_1^{(t)}\right)_j - A_j = \sum_{i=1}^d w_{2i}^{(t)} W_1^{(t)}[i,j] - A_j$, we have,

$$\begin{aligned}
\Delta E_j^{(t)} &:= E_j^{(t+1)} - E_j^{(t)} \\
&= \sum_{i=1}^d \left(w_{2i}^{(t+1)} W_1^{(t+1)}[i,j] - w_{2i}^{(t+1)} W_1^{(t)}[i,j] + w_{2i}^{(t+1)} W_1^{(t)}[i,j] - w_{2i}^{(t)} W_1^{(t)}[i,j]\right) \\
&= \sum_{i=1}^d \left(w_{2i}^{(t+1)} \Delta W_1^{(t)}[i,j] + \Delta w_{2i}^{(t)} W_1^{(t)}[i,j]\right).
\end{aligned} \tag{33}$$

Combining Lemma 22 and eq. (33) gives us $\forall j \in [d]$,

$$\Delta E_j^{(t)} > 0, \quad \left|\Delta E_j^{(t)}\right| = \sum_{i=1}^d \left|w_{2i}^{(t+1)} \Delta W_1^{(t)}[i,j]\right| + \left|\Delta w_{2i}^{(t)} W_1^{(t)}[i,j]\right| \leq \sum_{i=1}^d \mathcal{O}\left(\eta \frac{1}{\sqrt{d}}\right) = \mathcal{O}\left(\eta\sqrt{d}\right). \tag{34}$$

Let's first analyze $g_1^{(t)}[i,j]$. Note that

$$\begin{aligned}
\Delta g_1^{(t)}[i,j] &= w_{2i}^{(t+1)} E_j^{(t+1)} - w_{2i}^{(t+1)} E_j^{(t)} + w_{2i}^{(t+1)} E_j^{(t)} - w_{2i}^{(t)} E_j^{(t)} \\
&= w_{2i}^{(t+1)} \Delta E_j^{(t)} + \Delta w_{2i}^{(t)} E_j^{(t)},
\end{aligned} \tag{35}$$

where $sign\left(w_{2i}^{(t+1)} \Delta E_j^{(t)}\right) = sign\left(w_{2i}^{(0)}\right)$ while $sign\left(\Delta w_{2i}^{(t)} E_j^{(t)}\right) = -sign\left(w_{2i}^{(0)}\right)$.

Now we analyze the sign of $g_1^{(t)}[i,j]$ when $t_{\text{inc}} \leq t < T_1$. Using $\left|w_{2i}^{(t_{\text{inc}}+1)}\right| = \tilde{\mathcal{O}}\left(\frac{1}{d^\alpha}\right)$ and eq. (34), we get that $\left|w_{2i}^{(t_{\text{inc}}+1)} \Delta E_j^{(t_{\text{inc}})}\right| \leq \tilde{\mathcal{O}}\left(\frac{1}{d^\alpha} \cdot \sqrt{d}\eta\right)$. While on the other hand, $\left|\Delta w_{2i}^{(t_{\text{inc}})} E_j^{(t_{\text{inc}})}\right| = \Theta(\eta)$. That means $sign\left(\Delta g_1^{(t_{\text{inc}})}[i,j]\right) = -sign\left(w_{2i}^{(0)}\right)$. Note that $sign\left(g_1^{(t_{\text{inc}})}[i,j]\right) = -sign\left(w_{2i}^{(t_{\text{inc}})}\right) = -sign\left(w_{2i}^{(0)}\right)$, we know that $\left|g_1^{(t)}[i,j]\right|$ will increase when $t = t_{\text{inc}}$.

In the following steps, $\left|g_1^{(t)}[i,j]\right|$ will keep increasing as long as $\left|\Delta w_{2i}^{(t)} E_j^{(t)}\right| > \left|w_{2i}^{(t+1)} \Delta E_j^{(t)}\right|$. Since $\left|W_1^{(t)}[i,j]\right|, \left|w_{2i}^{(t)}\right|$ keep increasing while $\left|\Delta W_1^{(t)}[i,j]\right|, \left|\Delta w_{2i}^{(t)}\right|$ remain $\Theta(\eta)$, by eq. (34), we know that the trend of $\left|\Delta E_j^{(t)}\right|$ is to increase. On the other hand, $\left|E_j^{(t)}\right|$ keeps decreasing since $E_j^{(t)} < 0$ while $\Delta E_j^{(t)} > 0$. Then after some time point we will have $\left|\Delta w_{2i}^{(t)} E_j^{(t)}\right| < \left|w_{2i}^{(t+1)} \Delta E_j^{(t)}\right|$ and in the following steps $\left|g_1^{(t)}[i,j]\right|$ will keep decreasing. Specially, when $t = T_1 - 1$, we have $\left|E_j^{(t)}\right| = \Theta\left(\sqrt{\eta d}\right)$ and $\left|W_1^{(t)}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right), \left|w_{2i}^{(t+1)}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right)$ by Lemma 22, which gives us

$$\left|\Delta E_j^{(t)}\right| = \sum_{i=1}^d \left|w_{2i}^{(t+1)} \Delta W_1^{(t)}[i,j]\right| + \left|\Delta w_{2i}^{(t)} W_1^{(t)}[i,j]\right| \leq \sum_{i=1}^d \Theta\left(\eta \frac{1}{\sqrt{d}}\right) = \Theta\left(\eta\sqrt{d}\right).$$

Hence $\left|w_{2i}^{(t+1)}\Delta E_j^{(t)}\right| = \Theta(\eta) > \left|\Delta w_{2i}^{(t)}E_j^{(t)}\right| = \Theta\left(\eta\sqrt{\eta d}\right)$.

Therefore we have proved that when $t_{\text{inc}} \leq t < T_1$, $\left|g_1^{(t)}[i,j]\right|$ will first increase and then decrease. In order to prove $\left|g_1^{(t)}[i,j]\right| = \Omega\left(\sqrt{\eta}\right)$, it suffices to show that $\left|g_1^{(t_{\text{inc}})}[i,j]\right| = \Omega\left(\sqrt{\eta}\right)$ and $\left|g_1^{(T_1)}[i,j]\right| = \Omega\left(\sqrt{\eta}\right)$.

When $t = t_{\text{inc}}$,

$$\left|g_1^{(t_{\text{inc}})}[i,j]\right| = \left|w_{2i}^{(t_{\text{inc}})}\right| \cdot \left|E_j^{(t_{\text{inc}})}\right| = \Omega\left(\frac{1}{d^{\frac{3}{2}\alpha}}\right) \cdot \Theta(1) = \Omega\left(\sqrt{\eta}\right).$$

When $t = T_1$, we have

$$\left|g_1^{(T_1)}[i,j]\right| = \left|w_{2i}^{(T_1)}\right| \cdot \left|E_j^{(t_1)}\right| = \Theta\left(\frac{1}{\sqrt{d}} \cdot \sqrt{\eta d}\right) = \Theta\left(\sqrt{\eta}\right).$$

As for $g_{2i}^{(t)}$, since for $\forall i \in [d]$, $W_1^{(t)}[i,j]$ for different j have the same sign. Combining with $\forall j \in [d] : E_j^{(t)} < 0$ gives us

$$\left|g_{2i}^{(t)}\right| = \left|\sum_{j=1}^{d} E_j^{(t)} W_1^{(t)}[i,j]\right| = \sum_{j=1}^{d} \left|E_j^{(t)} W_1^{(t)}[i,j]\right|.$$

Then it suffices to show that for $t_{\text{inc}} \leq t < T_1$, $\left|E_j^{(t)} W_1^{(t)}[i,j]\right| = \Omega\left(\sqrt{\eta}\right)$, which can be proven using the same argument as above.

Finally, for $\forall \tau \leq t, \forall i, j \in [d]$, note that the upper bounds of $\left|W_1^{(\tau)}[i,j]\right|$ and $\left|w_{2i}^{(\tau)}\right|$ are already given in Lemma 22. As for $\left|g_1^{(\tau)}[i,j]\right|$ and $\left|g_{2i}^{(\tau)}\right|$, we have $\left|g_1^{(\tau)}[i,j]\right| = \left|w_{2i}^{(\tau)}E_j^{(\tau)}\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), \left|g_{2i}^{(\tau)}\right| \leq \sum_{j=1}^{d}\left|E_j^{(\tau)}W_1^{(\tau)}[i,j]\right| = \mathcal{O}\left(\sqrt{d}\right)$.

D.9 PROOF OF LEMMA 22

For any $i, j \in [d]$, and any t in the interval $[t_{\text{inc}}, T_1)$, we prove by induction that

(A) $\left|W_1^{(t)}[i,j]\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), \left|w_{2i}^{(t)}\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right).$

(B) $\forall \tau \in [t - H, t] : \text{sign}\left(W_1^{(\tau)}[i,j]\right) = \text{sign}\left(w_{2i}^{(\tau)}\right) = \text{sign}\left(w_{2i}^{(0)}\right).$

(C) $\left|g_1^{(t)}[i,j]\right| \geq \Omega(\xi), \left|g_{2i}^{(t)}\right| \geq \Omega(\xi).$

The base case $t = t_{\text{inc}}$ was already proven by Lemma 17.

For $t \in [t_{\text{inc}}, T_1)$, suppose (B) and (C) hold for time t and (A) holds for all $\tau \in [t_{\text{inc}}, t]$. From (A), we get that $\forall \tau \in [t_{\text{inc}}, t] : \left|g_1^{(\tau)}[i,j]\right| = \left|w_{2i}^{(\tau)}E_j^{(\tau)}\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), \left|g_{2i}^{(\tau)}\right| \leq \sum_{j=1}^{d}\left|E_j^{(\tau)}W_1^{(\tau)}[i,j]\right| = \mathcal{O}\left(\sqrt{d}\right)$. Since when $t < T_1, \forall j \in [d] : E_j^{(t)} < 0$, from (B) we know that $\forall \tau \in [t - H, t] : \text{sign}\left(g_1^{(\tau)}[i,j]\right) = \text{sign}\left(g_{2i}^{(\tau)}\right) = -\text{sign}\left(w_{2i}^{(0)}\right)$. Combining with (C) tells us that Condition 1 and 2 are satisfied.

In Section D.2 we have shown that $T_1 = \Theta\left(\frac{1}{\sqrt{d}\eta}\right)$. Then for $t \in [t_{\text{inc}}, T_1)$, we can use Lemma 21 to get that $\forall t_{\text{inc}} \leq \tau \leq t, \forall i, j \in [d]$,

$$\text{sign}\left(\Delta W_1^{(\tau)}[i,j]\right) = \text{sign}\left(\Delta w_{2i}^{(\tau)}\right) = \text{sign}\left(w_{2i}^{(0)}\right), \quad \left|\Delta W_1^{(\tau)}[i,j]\right| = \Theta(\eta), \left|\Delta w_{2i}^{(\tau)}\right| = \Theta(\eta).$$

Since when $t = t_{\text{inc}}$, $\text{sign}\left(W_1^{(t_{\text{inc}})}[i,j]\right) = \text{sign}\left(w_{2i}^{(t_{\text{inc}})}\right) = \text{sign}\left(w_{2i}^{(0)}\right)$. We get that for $t_{\text{inc}} \leq \tau \leq t$,

$$\forall i,j \in [d]: \quad \left|W_1^{(\tau+1)}[i,j]\right| = \left|W_1^{(\tau)}[i,j]\right| + \Theta(\eta), \quad \left|w_{2i}^{(\tau+1)}\right| = \left|w_{2i}^{(\tau)}\right| + \Theta(\eta).$$

Now for $t + 1$, we have

$$\forall i,j \in [d]: \quad \text{sign}\left(W_1^{(t+1)}[i,j]\right) = \text{sign}\left(w_{2i}^{(0)}\right), \quad \left|W_1^{(t+1)}[i,j]\right| = \left|W_1^{(t_{\text{inc}})}[i,j]\right| + (t+1-t_{\text{inc}})\,\Theta(\eta),$$

$$\text{sign}\left(w_{2i}^{(t+1)}\right) = \text{sign}\left(w_{2i}^{(0)}\right), \quad \left|w_{2i}^{(t+1)}\right| = \left|w_{2i}^{(t_{\text{inc}})}\right| + (t+1-t_{\text{inc}})\,\Theta(\eta).$$

That means $\forall \tau \in [t+1-H, t+1] : \text{sign}\left(W_1^{(\tau)}[i,j]\right) = \text{sign}\left(w_{2i}^{(\tau)}\right) = \text{sign}\left(w_{2i}^{(0)}\right)$. This proves (B) for time $t + 1$.

On the other hand, we get that $\left|W_1^{(t+1)}[i,j]\right| \geq \left|W_1^{(t_{\text{inc}})}[i,j]\right| = \Theta\left(\frac{1}{d^{\frac{3}{2}\alpha+1}}\right)$ and $\left|w_{2i}^{(t+1)}\right| \geq \left|w_{2i}^{(t_{\text{inc}})}\right| = \Omega\left(\frac{1}{d^{\frac{3}{2}\alpha}}\right)$. Since $t + 1 \leq T_1$ which means $\forall j \in [d] : \left|E_j^{(t+1)}\right| \geq \sqrt{\eta d}$. Then

$$\left|g_1^{(t+1)}[i,j]\right| = \left|w_{2i}^{(t+1)} E_j^{(t+1)}\right| \geq \Omega\left(\frac{1}{d^{\frac{3}{2}\alpha}}\right)\sqrt{\eta d} = \Omega(\xi),$$

$$\left|g_{2i}^{(t+1)}\right| = \left|\sum_{j=1}^d E_j^{(t+1)} W_1^{(t+1)}[i,j]\right| = \sum_{j=1}^d \left|E_j^{(t+1)} W_1^{(t+1)}[i,j]\right| \geq d\Theta\left(\frac{1}{d^{\frac{3}{2}\alpha+1}}\right)\sqrt{\eta d} = \Omega(\xi).$$

This proves (C) at time $t + 1$.

Since $t + 1 \leq T_1$ which means $\forall j \in [d] : \left(W_2^{(t+1)} W_1^{(t+1)}\right)_j \leq \mathcal{O}(1)$, we obtain that

$$\sum_{i=1}^d w_{2i}^{(t+1)} W_1^{(t+1)}[i,j] = \sum_{i=1}^d \left|w_{2i}^{(t+1)}\right|\left|W_1^{(t+1)}[i,j]\right|$$

$$= \sum_{i=1}^d \left(\left|w_{2i}^{(t_{\text{inc}})}\right| + (t+1-t_{\text{inc}})\Theta(\eta)\right)\left(\left|W_1^{(t_{\text{inc}})}[i,j]\right| + (t+1-t_{\text{inc}})\Theta(\eta)\right) \leq \mathcal{O}(1).$$

Note that $\left|W_1^{(t_{\text{inc}})}[i,j]\right|, \left|w_{2i}^{(t_{\text{inc}})}\right| < \frac{1}{d}$ (since $t_{\text{inc}} < t_d$), we get that $(t+1-t_{\text{inc}})\Theta(\eta) = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$, which gives us $\left|w_{2i}^{(t+1)}\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$ and $\left|W_1^{(t+1)}[i,j]\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$ and hence (A) holds at time $t + 1$.

Therefore by induction, we can prove that (A), (B), (C) hold for all $t_{\text{inc}} \leq t \leq T_1$. Then applying Lemma 21, we get that for all $t_{\text{inc}} \leq t < T_1, \forall i,j \in [d] : \left|\Delta W_1^{(t)}[i,j]\right| = \Theta(\eta), \left|\Delta w_{2i}^{(t)}\right| = \Theta(\eta)$.

Specially, at the end of the first phase, we have $(T_1 - t_{\text{inc}})\Theta(\eta) = \Theta\left(\frac{1}{\sqrt{d}}\right)$, which gives us $\left|w_{2i}^{(T_1)}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right)$ and $\left|W_1^{(T_1)}[i,j]\right| = \Theta\left(\frac{1}{\sqrt{d}}\right)$ for $\forall i,j \in [d]$.

D.10 PROOF OF LEMMA 19

Let's first prove eq. (29).

By Lemma 18, for $t_{\text{inc}} \leq t < T_1$, we have $\forall i,j \in [d], \left|g_1^{(t)}[i,j]\right| = \Omega\left(\sqrt{\eta}\right), \left|g_{2i}^{(t)}\right| = \Omega\left(\sqrt{\eta d}\right)$. Then it suffices to show that for $t_{\text{inc}} \leq t < T_1, \left|g_1^{(t)}[i,j] - g_1^{(t-\tau)}[i,j]\right| = \tau\mathcal{O}(\eta)$ and $\left|g_{2i}^{(t)} - g_{2i}^{(t-\tau)}\right| = \tau\mathcal{O}(\eta d)$. It suffices to show that when $t < T_1, \left|g_1^{(t+1)}[i,j] - g_1^{(t)}[i,j]\right| = \mathcal{O}(\eta)$ and $\left|g_{2i}^{(t+1)} - g_{2i}^{(t)}\right| = \mathcal{O}(\eta d)$.

By Lemma 17 and 22, we know that when $t < T_1, \forall i,j \in [d], \left|\Delta W_1^{(t)}[i,j]\right| \leq \tilde{\mathcal{O}}(\eta), \left|\Delta w_{2i}^{(t)}\right| \leq \tilde{\mathcal{O}}(\eta)$ and that $\left|W_1^{(t)}[i,j]\right| \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right), \left|w_{2i}^{(t)}\right| \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$. Then the bound $\left|\Delta E_j^{(t)}\right| \leq \mathcal{O}\left(\eta\sqrt{d}\right)$ in

eq. (34) hold for $t < T_1$ (not only $t_{\mathrm{inc}} \le t < T_1$). Substituting these bounds into eq. (35) gives us $\forall t < T_1$,

$$\left| g_1^{(t+1)}[i,j] - g_1^{(t)}[i,j] \right| \le \left| w_{2i}^{(t+1)} \right| \left| \Delta E_j^{(t)} \right| + \left| \Delta w_{2i}^{(t)} \right| \left| E_j^{(t)} \right|$$

$$= \mathcal{O}\left(\frac{1}{\sqrt{d}} \right) \mathcal{O}\left(\eta \sqrt{d} \right) + \Theta(\eta)\mathcal{O}(1) = \mathcal{O}(\eta).$$

Similarly, we have that $\left| g_{2i}^{(t+1)} - g_{2i}^{(t)} \right| = \mathcal{O}(\eta d)$, which proves eq. (29).

Note that for $a, b \in \mathbb{R}$:

$$\frac{\left| a^2 - b^2 \right|}{a^2} = \frac{\left| a^2 - (a-b-a)^2 \right|}{a^2} = \frac{\left| 2a(a-b) - (a-b)^2 \right|}{a^2} \le 2\frac{|a-b|}{|a|} + \left(\frac{|a-b|}{|a|} \right)^2.$$

Then eq. (30) immediately follows from eq. (29).

D.11 Proof of Lemma 15

We divide Lemma 15 into the following three lemmas. Combining them together immediately gives us the whole proof.

The first lemma below gives us the structure of W_2 in the second phase and that of W_1 under some conditions.

Lemma 23. *Under Assumption 1, 2 and 3, suppose $\sigma \le \frac{\eta^{3/2}\xi^2}{d^{13/4}}$. By picking $\eta \le \mathcal{O}\left(\frac{1}{d^{3\alpha}} \right), \xi \le \sqrt{\frac{\eta}{d^{3\alpha-1}}}$, and $\beta_2 = \beta_1^2$, we have for $T_1 \le t < \tilde{T}$,*

$$\forall i \in [d]: \quad w_{2i}^{(t+1)} = w_{2i}^{(t)} - \eta\left(sign\left(g_{2i}^{(t)} \right) + e_{2i}^{(t)} \right), \quad where \ \left| e_{2i}^{(t)} \right| = \mathcal{O}\left(\sqrt{\eta} \right),$$

which gives us

$$\forall i \in [d]: \quad w_{2i}^{(t)} = sign\left(w_{2i}^{(0)} \right) c^{(t)} + R_{2i}^{(t)}, \quad where \quad \frac{\left| R_{2i}^{(t)} \right|}{c^{(t)}} = \mathcal{O}\left(\sqrt{\eta} + \frac{1}{d^{\alpha-1/2}} \right).$$

As for W_1, if for certain $i.j \in [d]$ and certain $t \in [T_1, \tilde{T})$ we have $\left| g_1^{(t)}[i,j] \right| = \Omega\left(\sqrt{\eta} \right)$, then

$$W_1^{(t+1)}[i,j] = W_1^{(t)}[i,j] - \eta\left(sign\left(g_1^{(t)}[i,j] \right) + e_1^{(t)}[i,j] \right), \quad where \ \left| e_1^{(t)}[i,j] \right| = \mathcal{O}\left(\sqrt{\eta} \right).$$

The second lemma below also analyzes the structure of W_1 but removes the conditions in Lemma 23.

Lemma 24. *Under Assumption 1, 2 and 3, suppose $\sigma \le \frac{\eta^{3/2}\xi^2}{d^{13/4}}$. By picking $\eta \le \mathcal{O}\left(\frac{1}{d^{3\alpha}} \right), \xi \le \sqrt{\frac{\eta}{d^{3\alpha-1}}}$, and $\beta_2 = \beta_1^2$, we have for $T_1 \le t < \tilde{T}$, $\forall i, j \in [d]$, $\left| W_1^{(t)}[i,j] \right| = \Omega\left(\frac{1}{\sqrt{d}} \right)$ and for any $j \in [d]$,*

$$W_1^{(t)}[i,j] = sign\left(w_{2i}^{(0)} \right) V_j^{(t)} + R_1^{(t)}[i,j], \quad where \quad \frac{\left| R_1^{(t)}[i,j] \right|}{\left| V_j^{(t)} \right|} \le \mathcal{O}\left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}} \right).$$

The third lemma proves the convergence of Adam at time $\tilde{T}$.

Lemma 25. *Under Assumption 1, 2 and 3, suppose $\sigma \le \frac{\eta^{3/2}\xi^2}{d^{13/4}}$. By picking $\eta \le \mathcal{O}\left(\frac{1}{d^{3\alpha}} \right), \xi \le \sqrt{\frac{\eta}{d^{3\alpha-1}}}$, and $\beta_2 = \beta_1^2$, at time $\tilde{T}$, we have that $\forall j \in [d]: \left| E_j^{(\tilde{T})} \right| \le \mathcal{O}\left(d\sqrt{\eta d} \right)$, which implies $\left\| E^{(\tilde{T})} \right\|_2^2 \le \mathcal{O}\left(\eta d^4 \right)$.*

D.12   PROOF OF LEMMA 23

The proof is based on the following lemma, which gives a coarse analysis on the magnitude of weights and their increments per step during the second phase.

Lemma 26. *Under Assumption 1, 2 and 3, suppose* $\sigma \leq \frac{\eta^{3/2}\xi^2}{d^{13/4}}$. *By picking* $\eta \leq \mathcal{O}\left(\frac{1}{d^{3\alpha}}\right), \xi \leq \sqrt{\frac{\eta}{d^{3\alpha-1}}}$, *and* $\beta_2 = \beta_1^2$, *we have for* $T_1 \leq t < \tilde{T}$,

$$\forall i \in [d]: \quad \left|w_{2i}^{(t+1)}\right| > \left|w_{2i}^{(t)}\right|, \text{ and } \left|w_{2i}^{(t+1)}\right| - \left|w_{2i}^{(t)}\right| = \Theta(\eta).$$

Moreover, we have that $\forall i, j \in [d]: \quad \left|w_{2i}^{(t)}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right), \left|W_1^{(t)}[i,j]\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right).$

Equipped with Lemma 26, we are ready to prove Lemma 23. We will only prove the results of $w_{2i}^{(t)}$. The proof for $W_1^{(t)}[i,j]$ uses the same techniques.

Lemma 26 gives us upper bounds of $\left|w_{2i}^{(t)}\right|, \left|W_1^{(t)}[i,j]\right|$ for all $i, j \in [d]$. Combining with the upper bounds of $\left|w_{2i}^{(t+1)} - w_{2i}^{(t)}\right|$ and $\left|W_1^{(t+1)}[i,j] - W_1^{(t)}[i,j]\right|$, we know that eq.(33) still holds, which gives us $\forall j \in [d]: \left|E_j^{(t+1)} - E_j^{(t)}\right| = \mathcal{O}\left(\sqrt{d}\eta\right)$. Then we can use the same strategy in Lemma 19 to prove that $\left|g_{2i}^{(t+1)} - g_{2i}^{(t)}\right| = \mathcal{O}(\eta d)$.

By definition, for $t < \tilde{T}$, we know that $\left|g_{2i}^{(t)}\right| = \Omega\left(d\sqrt{\eta}\right)$. Combining with the bound $\left|g_{2i}^{(t+1)} - g_{2i}^{(t)}\right| = \mathcal{O}(\eta d)$, we know that the $g_{2i}^{(t)}$ parts in eq.(29) and eq.(30) still hold. Then we can use the same strategy in Section D.2 to prove that the $w_{2i}^{(t)}$ part of eq. (28) still holds, which gives us

$$\forall i \in [d]: \quad w_{2i}^{(t+1)} = w_{2i}^{(t)} - \eta\left(\text{sign}\left(g_{2i}^{(t)}\right) + e_{2i}^{(t)}\right), \quad \text{where } \left|e_{2i}^{(t)}\right| = \mathcal{O}\left(\sqrt{\eta}\right).$$

By Lemma 14, we have that at the end of the first phase ($t = T_1$),

$$\forall i \in [d]: \quad w_{2i}^{(T_1)} = \text{sign}\left(w_{2i}^{(0)}\right)c^{(T_1)} + R_{2i}^{(T_1)}, \quad \text{where} \quad \frac{\left|R_{2i}^{(T_1)}\right|}{c^{(T_1)}} = \mathcal{O}\left(\sqrt{\eta} + \frac{1}{d^{\alpha-1/2}}\right).$$

Combining with $\forall i \in [d], \forall t \leq T_i : \text{sign}\left(g_{2i}^{(t)}\right) = -\text{sign}\left(w_{2i}^{(0)}\right)$ yields that during the second phase, for $t \leq \tilde{T}$, we have

$$\forall i \in [d]: \quad w_{2i}^{(t)} = \text{sign}\left(w_{2i}^{(0)}\right)c^{(t)} + R_{2i}^{(t)}, \quad \text{where} \quad \frac{\left|R_{2i}^{(t)}\right|}{c^{(t)}} = \mathcal{O}\left(\sqrt{\eta} + \frac{1}{d^{\alpha-1/2}}\right).$$

D.13   PROOF OF LEMMA 26

We prove by induction that when $T_1 \leq t < \tilde{T}, \forall i, j \in [d] : \quad \left|w_{2i}^{(t)}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right), \left|W_1^{(t)}[i,j]\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right).$

The base case ($t = T_1$) was already proven by Lemma 22. Now suppose for some t such that $T_1 \leq t < \tilde{T}$, for all τ such that $T_1 \leq \tau \leq t$, we have $\forall i, j \in [d] : \quad \left|w_{2i}^{(\tau)}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right), \left|W_1^{(\tau)}[i,j]\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$. Using these bounds, we get that $\forall j \in [d] : \left|E_j^{(\tau)}\right| \leq \sum_{i=1}^{d}\left|w_{2i}^{(\tau)}W_1^{(\tau)}[i,j]\right| + |A_j| = \mathcal{O}(1)$, which then yields two upper bounds $\left|g_1^{(\tau)}[i,j]\right| = \left|w_{2i}^{(\tau)}E_j^{(\tau)}\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$ and $\left|g_{2i}^{(\tau)}\right| \leq \sum_{j=1}^{d}\left|E_j^{(\tau)}W_1^{(\tau)}[i,j]\right| = \mathcal{O}\left(\sqrt{d}\right)$.

By definition of T_g, we know that for all $T_1 \leq \tau \leq t, \forall i \in [d] : \left|g_{2i}^{(\tau)}\right| \geq d\sqrt{\eta} = \Omega(\xi)$ and that $\text{sign}\left(g_{2i}^{(\tau)}\right) = -\text{sign}\left(w_{2i}^{(0)}\right)$, which implies that Condition 2 is satisfied for $\forall i \in [d]$. At the end of

the proof of this lemma, we will show that $\tilde{T} = \Theta\left(\frac{1}{\sqrt{d\eta}}\right)$. Together with the upper bound of $\left|g_{2i}^{(\tau)}\right|$, we can apply Lemma 21 to get that for $T_1 \leq \tau \leq t$, $\left|w_{2i}^{(\tau+1)}\right| - \left|w_{2i}^{(\tau)}\right| = \Theta(\eta)$. Specially, when $\tau = t$, we get the lower bound $\left|w_{2i}^{(t+1)}\right| \geq \left|w_{2i}^{(t)}\right| = \Omega\left(\frac{1}{\sqrt{d}}\right)$.

On the other hand, by definition of T_f, there exists $j_0 \in [d]$ such that $E_{j_0}^{(\tau)} < -\sqrt{\eta d}$ for $T_1 \leq \tau \leq t$. Then we have that $\forall i \in [d] : \left|g_1^{(\tau)}[i,j_0]\right| = \left|w_{2i}^{(\tau)}\right|\left|E_{j_0}^{(\tau)}\right| = \Omega\left(\sqrt{\eta}\right) = \Omega(\xi)$ and that $\text{sign}\left(g_1^{(\tau)}[i,j_0]\right) = -\text{sign}\left(w_{2i}^{(0)}\right)$. That means Condition 1 is satisfied for $\forall i \in [d]$ and j_0. Again, together with the upper bound of $\left|g_1^{(\tau)}[i,j]\right|$, we can apply Lemma 21 to get that for $T_1 \leq \tau \leq t$, $\left|W_1^{(\tau+1)}[i,j_0]\right| - \left|W_1^{(\tau)}[i,j_0]\right| = \Theta(\eta)$.

Now we analyze the magnitude order of $\left|w_{2i}^{(t+1)}\right|, \left|W_1^{(t+1)}[i,j]\right|$. Let's first analyze $\left|w_{2i}^{(t+1)}\right|$.

By Lemma 14, when $t = T_1$,

$$\forall i,j \in [d] : \frac{\left|w_{2i}^{(T_1)}\right|}{\left|w_{2j}^{(T_1)}\right|} = 1 \pm \mathcal{O}\left(\sqrt{\eta} + \frac{1}{d^{\alpha-1/2}}\right), \quad \forall i \in [d] : \frac{\left|W_1^{(T_1)}[i,j_0]\right|}{\left|w_{2i}^{(T_1)}\right|} = 1 \pm \mathcal{O}\left(\sqrt{\eta} + \frac{1}{d^{\alpha-1/2}}\right).$$

Combining with the facts that for $T_1 \leq \tau \leq t$, $\left|W_1^{(\tau+1)}[i,j_0]\right| = \left|W_1^{(\tau)}[i,j_0]\right| + \Theta(\eta)$ and $\left|w_{2i}^{(\tau+1)}\right| = \left|w_{2i}^{(\tau)}\right| + \Theta(\eta)$ yields $\frac{\left|W_1^{(t+1)}[i,j_0]\right|}{\left|w_{2i}^{(t+1)}\right|} = \Theta(1)$. Since

$$(W_2 W_1)_{j_0}^{(t+1)} = \sum_{i=1}^{d} w_{2i}^{(t+1)} W_1^{(t+1)}[i,j_0] = \sum_{i=1}^{d} \left|w_{2i}^{(t+1)}\right|\left|W_1^{(t+1)}[i,j_0]\right| = \mathcal{O}(1),$$

which gives us that $\left|w_{2i}^{(t+1)}\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$. Since we have shown $\left|w_{2i}^{(t+1)}\right| \geq \Omega\left(\frac{1}{\sqrt{d}}\right)$, then we get $\left|w_{2i}^{(t+1)}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right)$.

Now we prove $\left|W_1^{(t+1)}[i,j]\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$. The upper bound of $\left|g_1^{(\tau)}[i,j]\right|$ allows us to use Lemma 21 to get that for $\tau \leq t$, $\forall i,j \in [d]$, $\left|W_1^{(\tau+1)}[i,j] - W_1^{(\tau)}[i,j]\right| = \mathcal{O}(\eta)$. Note that for $w_{2i}^{(\tau)}$, we have $\left|w_{2i}^{(\tau+1)}\right| - \left|w_{2i}^{(\tau)}\right| = \Theta(\eta)$, then $\forall i,j \in [d]$,

$$\frac{\left|W_1^{(t+1)}[i,j]\right|}{\left|w_{2i}^{(t+1)}\right|} = \frac{\left|W_1^{(T_1)}[i,j]\right| + \sum_{\tau=T_1}^{t}\left|W_1^{(\tau+1)}[i,j] - W_1^{(\tau)}[i,j]\right|}{\left|w_{2i}^{(T_1)}\right| + \sum_{\tau=T_1}^{t}\left|w_{2i}^{(\tau+1)}\right| - \left|w_{2i}^{(\tau)}\right|}$$

$$\leq \frac{\left|W_1^{(T_1)}[i,j]\right| + (t+1-T_1)\mathcal{O}(\eta)}{\left|w_{2i}^{(T_1)}\right| + (t+1-T_1)\Theta(\eta)} = \mathcal{O}(1),$$

where the last equality uses $\frac{\left|W_1^{(T_1)}[i,j]\right|}{\left|w_{2i}^{(T_1)}\right|} = 1 \pm \mathcal{O}\left(\sqrt{\eta} + \frac{1}{d^{\alpha-1/2}}\right)$. Since we already proved that $\left|w_{2i}^{(t+1)}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right)$, we get $\left|W_1^{(t+1)}\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$.

Therefore by induction, for all t in the interval $[T_1, \tilde{T})$, we have $\forall i,j \in [d] : \left|w_{2i}^{(t)}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right)$, $\left|W_1^{(t)}[i,j]\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$. From the proof we also get $\forall i \in [d] : \left|w_{2i}^{(t+1)}\right| > \left|w_{2i}^{(t)}\right|$, and $\left|w_{2i}^{(t+1)}\right| - \left|w_{2i}^{(t)}\right| = \Theta(\eta)$.

Now we verify that $\tilde{T} = \Theta\left(\frac{1}{\sqrt{d}\eta}\right)$. Combining $\forall i, j \in [d] : \left|w_{2i}^{(\tilde{T})}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right)$ and $\forall t \in [T_1, \tilde{T})$, $\left|w_{2i}^{(t+1)}\right| - \left|w_{2i}^{(t)}\right| = \Theta(\eta)$, we immediately get that $\tilde{T} - T_1 = \Theta\left(\frac{1}{\sqrt{d}\eta}\right)$. In Section D.2 we have shown that $T_1 = \Theta\left(\frac{1}{\sqrt{d}\eta}\right)$, then we get $\tilde{T} = \Theta\left(\frac{1}{\sqrt{d}\eta}\right)$.

D.14 PROOF OF LEMMA 24

We prove this lemma by induction. The base case ($t = T_1$) can be verified by Lemma 14. Now suppose for t in the interval $[T_1, \tilde{T})$, we have $\forall i, j \in [d]$, $\left|W_1^{(t)}[i, j]\right| = \Omega\left(\frac{1}{\sqrt{d}}\right)$.

For $t \in [T_1, \tilde{T})$, by Lemma 26 and 18, we know that for $\forall \tau \leq t$, we have $\forall i, j \in [d] : \left|w_{2i}^{(\tau)}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right)$, $\left|W_1^{(\tau)}[i, j]\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$ and that $\left|g_1^{(\tau)}[i, j]\right| \leq \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$, $\left|g_{2i}^{(\tau)}\right| \leq \mathcal{O}\left(\sqrt{d}\right)$. In the proof of Lemma 26 (Section D.13), we have shown that $\tilde{T}_1 = \Theta\left(\frac{1}{\sqrt{d}\eta}\right)$. Then we can pick $H := \frac{1}{1-\beta_1} \log \frac{d}{\eta\xi^2}$ and apply Lemma 16 and Corollary 2 to get that, w.h.p., for all $t \in [T_1, \tilde{T})$ and $\forall i, j \in [d]$, the update of W_1 can be written as

$$W_1^{(t+1)}[i, j] = W_1^{(t)}[i, j] - \eta_t \frac{(1 - \beta_1) \sum_{\tau=0}^{H} \beta_1^\tau g_1^{(t-\tau)}[i, j] + \epsilon_{1n}^{(t)}[i, j]}{\sqrt{(1 - \beta_2) \sum_{\tau=0}^{H} \beta_2^\tau \left(g_1^{(t-\tau)}[i, j]\right)^2 + \epsilon_{1d}^{(t)}[i, j] + \xi}},$$

where $\left|\epsilon_{1n}^{(t)}[i, j]\right|, \left|\epsilon_{1d}^{(t)}[i, j]\right| \leq \eta\xi^2$. By Lemma 23, we have that for $1 \leq i, j \leq d$,

$$g_1^{(t)}[i, j] = w_{2i}^{(t)} E_j^{(t)} = c^{(t)} \mathrm{sign}\left(w_{2i}^{(0)}\right) E_j^{(t)} + R_{g,1}^{(t)}[i, j], \quad \text{where} \quad \frac{\left|R_{g,1}^{(t)}[i, j]\right|}{c^{(t)} \left|E_j^{(t)}\right|} = \mathcal{O}\left(\sqrt{\eta} + \frac{1}{d^{\alpha-1/2}}\right),$$

$$\Rightarrow \quad \sum_{\tau=0}^{H} \beta_1^\tau g_1^{(t-\tau)}[i, j] = \mathrm{sign}\left(w_{2i}^{(0)}\right) \sum_{\tau=0}^{H} \beta_1^\tau c^{(t-\tau)} E_j^{(t-\tau)} + \sum_{\tau=0}^{H} \beta_1^\tau R_{g,1}^{(t-\tau)}[i, j]. \quad (36)$$

Using the fact that for $a, b \in \mathbb{R}, \frac{|a^2 - b^2|}{a^2} \leq 2\frac{|a-b|}{|a|} + \left(\frac{|a-b|}{|a|}\right)^2$, we get that

$$\left(g_1^{(t)}[i, j]\right)^2 = \left(c^{(t)} \mathrm{sign}\left(w_{2i}^{(0)}\right) E_j^{(t)} + R_{g,1}^{(t)}[i, j]\right)^2 = \left(c^{(t)} E_j^{(t)}\right)^2 + R_{\mathrm{gsqr},1}^{(t)}[i, j],$$

where $\frac{\left|R_{\mathrm{gsqr},1}^{(t)}[i, j]\right|}{\left(c^{(t)} E_j^{(t)}\right)^2} = \mathcal{O}\left(\sqrt{\eta} + \frac{1}{d^{\alpha-1/2}}\right)$. That yields

$$\sum_{\tau=0}^{H} \beta_2^\tau \left(g_1^{(t-\tau)}[i, j]\right)^2 = \sum_{\tau=0}^{H} \beta_2^\tau \left(c^{(t-\tau)} E_j^{(t-\tau)}\right)^2 + \sum_{\tau=0}^{H} \beta_2^\tau R_{\mathrm{gsqr},1}^{(t-\tau)}[i, j]. \quad (37)$$

Since $\left(c^{(t-\tau)} E_j^{(t-\tau)}\right)^2 > 0$, in eq. (37) we have that

$$\frac{\left|\sum_{\tau=0}^{H} \beta_2^\tau R_{\mathrm{gsqr},1}^{(t-\tau)}[i, j]\right|}{\left|\sum_{\tau=0}^{H} \beta_2^\tau \left(c^{(t-\tau)} E_j^{(t-\tau)}\right)^2\right|} = \mathcal{O}\left(\sqrt{\eta} + \frac{1}{d^{\alpha-1/2}}\right). \quad (38)$$

However in eq. (36), we cannot similarly prove that $\left|\sum_{\tau=0}^{H} \beta_1^\tau R_{g,1}^{(t-\tau)}[i, j]\right| \ll \left|\sum_{\tau=0}^{H} \beta_1^\tau c^{(t-\tau)} E_j^{(t-\tau)}\right|$ because $c^{(t-\tau)} E_j^{(t-\tau)}$ may not have the same sign for $\tau = 0, 1, ..., H$. To deal with eq.(36), we need to consider the two cases where $\left|\sum_{\tau=0}^{H} \beta_1^\tau R_{g,1}^{(t-\tau)}[i, j]\right| \ll \left|\sum_{\tau=0}^{H} \beta_1^\tau c^{(t-\tau)} E_j^{(t-\tau)}\right|$ or $\left|\sum_{\tau=0}^{H} \beta_1^\tau R_{g,1}^{(t-\tau)}[i, j]\right| \ll \left|\sum_{\tau=0}^{H} \beta_1^\tau c^{(t-\tau)} E_j^{(t-\tau)}\right|$.

Case 1. $\left|(1-\beta_1)\sum_{\tau=0}^{H}\beta_1^\tau R_{g,1}^{(t-\tau)}[i,j]+\epsilon_{1n}^{(t)}[i,j]\right|\leq\left(\eta^{\frac{1}{4}}+\frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}}\right)\left|(1-\beta_1)\sum_{\tau=0}^{H}\beta_1^\tau c^{(t-\tau)}E_j^{(t-\tau)}\right|.$

Note that from eq. (38) we have

$$\left|(1-\beta_1)\sum_{\tau=0}^{H}\beta_2^\tau R_{\text{gsqr},1}^{(t-\tau)}[i,j]\right|\leq\mathcal{O}\left(\sqrt{\eta}+\frac{1}{d^{\alpha-1/2}}\right)\left|(1-\beta_1)\sum_{\tau=0}^{H}\beta_2^\tau\left(c^{(t-\tau)}E_j^{(t-\tau)}\right)^2\right|.$$

Combining with $\left|\epsilon_{1d}^{(t)}[i,j]\right|\leq\eta\xi^2\leq\mathcal{O}\left(\eta^{\frac{1}{4}}+\frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}}\right)^2\xi^2$, we can apply Lemma 33 to get that

$$W_1^{(t+1)}[i,j]-W_1^{(t)}[i,j]$$

$$=-\eta_t\frac{(1-\beta_1)\text{sign}\left(w_{2i}^{(0)}\right)\sum_{\tau=0}^{H}\beta_1^\tau c^{(t-\tau)}E_j^{(t-\tau)}+(1-\beta_1)\sum_{\tau=0}^{H}\beta_1^\tau R_{g,1}^{(t-\tau)}[i,j]+\epsilon_{1n}^{(t)}[i,j]}{\sqrt{(1-\beta_2)\sum_{\tau=0}^{H}\beta_2^\tau\left(c^{(t-\tau)}E_j^{(t-\tau)}\right)^2+(1-\beta_2)\sum_{\tau=0}^{H}\beta_2^\tau R_{gsqr,1}^{(t-\tau)}[i,j]+\epsilon_{1d}^{(t)}[i,j]+\xi}}$$

$$=-\eta_t\frac{1-\beta_1}{\sqrt{1-\beta_2}}\cdot\frac{\text{sign}\left(w_{2i}^{(0)}\right)\sum_{\tau=0}^{H}\beta_1^\tau c^{(t-\tau)}E_j^{(t-\tau)}}{\sqrt{\sum_{\tau=0}^{H}\beta_2^\tau\left(c^{(t-\tau)}E_j^{(t-\tau)}\right)^2}+\xi}\left(1+\tilde{e}_1^{(t)}[i,j]\right)$$

$$:=-\text{sign}\left(w_{2i}^{(0)}\right)v_j^{(t)}\left(1+\tilde{e}_1^{(t)}[i,j]\right),$$

where $\left|\tilde{e}_1^{(t)}[i,j]\right|=\mathcal{O}\left(\eta^{\frac{1}{4}}+\frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}}\right)$. Since $\left|W_1^{(t+1)}[i,j]-W_1^{(t)}[i,j]\right|=\mathcal{O}(\eta)$, we get that $\left|v_j^{(t)}\right|=\mathcal{O}(\eta)$.

Case 2. $\left|(1-\beta_1)\sum_{\tau=0}^{H}\beta_1^\tau R_{g,1}^{(t-\tau)}[i,j]+\epsilon_{1n}^{(t)}[i,j]\right|>\left(\eta^{\frac{1}{4}}+\frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}}\right)\left|(1-\beta_1)\sum_{\tau=0}^{H}\beta_1^\tau c^{(t-\tau)}E_j^{(t-\tau)}\right|.$

Since $\frac{\left|R_{g,1}^{(t)}[i,j]\right|}{c^{(t)}\left|E_j^{(t)}\right|}=\mathcal{O}\left(\sqrt{\eta}+\frac{1}{d^{\alpha-1/2}}\right)$, we have that

$$\left|(1-\beta_1)\sum_{\tau=0}^{H}\beta_1^\tau R_{g,1}^{(t-\tau)}[i,j]\right|\leq\mathcal{O}\left(\sqrt{\eta}+\frac{1}{d^{\alpha-1/2}}\right)(1-\beta_1)\sum_{\tau=0}^{H}\beta_1^\tau\left|c^{(t-\tau)}E_j^{(t-\tau)}\right|$$

$$\overset{(i)}{\leq}\mathcal{O}\left(\sqrt{\eta}+\frac{1}{d^{\alpha-1/2}}\right)\sqrt{(H+1)(1-\beta_1)\sum_{\tau=0}^{H}\beta_2^\tau\left(c^{(t-\tau)}E_j^{(t-\tau)}\right)^2}$$

$$\overset{(ii)}{=}\mathcal{O}\left(\sqrt{\eta}+\frac{1}{d^{\alpha-1/2}}\right)\sqrt{(1-\beta_1)\sum_{\tau=0}^{H}\beta_2^\tau\left(g_1^{(t-\tau)}[i,j]\right)^2},$$

where (i) uses Cauchy-Schwarz inequality and $\beta_2=\beta_1^2$, (ii) uses eq. (37) and (38).

Combining with $\left|\epsilon_{1n}^{(t)}[i,j]\right|\leq\eta\xi^2\leq\mathcal{O}\left(\sqrt{\eta}+\frac{1}{d^{\alpha-1/2}}\right)\left(\xi-\sqrt{\left|\epsilon_{1d}^{(t)}[i,j]\right|}\right)$ gives us

$$\left|(1-\beta_1)\sum_{\tau=0}^{H}\beta_1^\tau c^{(t-\tau)}E_j^{(t-\tau)}\right|<\frac{\left|(1-\beta_1)\sum_{\tau=0}^{H}\beta_1^\tau R_{g,1}^{(t-\tau)}[i,j]+\epsilon_{1n}^{(t)}[i,j]\right|}{\eta^{\frac{1}{4}}+\frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}}}$$

$$\leq\frac{\mathcal{O}\left(\sqrt{\eta}+\frac{1}{d^{\alpha-1/2}}\right)}{\eta^{\frac{1}{4}}+\frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}}}\left(\sqrt{\sum_{\tau=0}^{H}\beta_2^\tau\left(g_1^{(t-\tau)}[i,j]\right)^2}-\sqrt{\left|\epsilon_{1d}^{(t)}[i,j]\right|}+\xi\right)$$

$$\leq\mathcal{O}\left(\eta^{\frac{1}{4}}+\frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}}\right)\left(\sqrt{\sum_{\tau=0}^{H}\beta_2^\tau\left(g_1^{(t-\tau)}[i,j]\right)^2+\epsilon_{1d}^{(t)}[i,j]}+\xi\right),$$

which implies

$$\left| W_1^{(t+1)}[i,j] - W_1^{(t)}[i,j] \right| \leq \eta \mathcal{O}\left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}} \right).$$

Consider certain $i.j \in [d]$ and the period from T_1 to t. Denote $\mathcal{T}$ as the set of time points when Case 1 is satisfied, and define $\Delta W_1^{(t)}[i,j] := W_1^{(t+1)}[i,j] - W_1^{(t)}[i,j]$. By Lemma 26, we know that $\eta(t - T_1) = \mathcal{O}\left(\frac{1}{\sqrt{d}} \right)$, which gives us

$$\sum_{\tau \notin \mathcal{T}} \Delta W_1^{(\tau)}[i,j] \leq (t - T_1)\eta \mathcal{O}\left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}} \right) = \mathcal{O}\left(\frac{\eta^{\frac{1}{4}}}{d^{\frac{1}{2}}} + \frac{1}{d^{\frac{\alpha}{2}+\frac{1}{4}}} \right).$$

By the first phase analysis, we have that

$$W_1^{(T_1)}[i,j] = \text{sign}\left(w_{2i}^{(0)} \right) V_j^{(T_1)} + R_1^{(T_1)}[i,j],$$

where $V_j^{(T_1)} = \mathcal{O}\left(\frac{1}{\sqrt{d}} \right), \left| R_1^{(T_1)}[i,j] \right| = \mathcal{O}\left(\sqrt{\frac{\eta}{d}} + \frac{1}{d^{\alpha}} \right)$. Combining with the analysis of Case 1, we have that

$$W_1^{(T_1)}[i,j] + \sum_{\tau \in \mathcal{T}} \Delta W_1^{(\tau)}[i,j] = \text{sign}\left(w_{2i}^{(0)} \right) \left(V_j^{(T_1)} - \sum_{\tau \in \mathcal{T}} v_j^{(\tau)} \right) + R_{\mathcal{T}}[i,j],$$

where $|R_{\mathcal{T}}[i,j]| \leq \mathcal{O}\left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}} \right) \left(\left| V_j^{(T_1)} \right| + \sum_{\tau \in \mathcal{T}} \left| v_j^{(\tau)} \right| \right)$.

Since for $\tau \in \mathcal{T}, \left| v_j^{(\tau)} \right| = \mathcal{O}(\eta), V_j^{(T_1)} = \mathcal{O}\left(\frac{1}{\sqrt{d}} \right)$ and $\eta(t - T_1) = \mathcal{O}\left(\frac{1}{\sqrt{d}} \right)$, we can bound $|R_{\mathcal{T}}[i,j]|$ by

$$|R_{\mathcal{T}}[i,j]| \leq \mathcal{O}\left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}} \right) \left(\mathcal{O}\left(\frac{1}{\sqrt{d}} \right) + (t - T_1)\mathcal{O}(\eta) \right) \leq \mathcal{O}\left(\frac{\eta^{\frac{1}{4}}}{d^{\frac{1}{2}}} + \frac{1}{d^{\frac{\alpha}{2}+\frac{1}{4}}} \right).$$

Combining the above results together yields

$$W_1^{(t)}[i,j] = W_1^{(T_1)} + \sum_{\tau=T_1}^{t-1} \Delta W_1^{(\tau)}[i,j] = W_1^{(T_1)} + \sum_{\tau \in \mathcal{T}} \Delta W_1^{(\tau)}[i,j] + \sum_{\tau \notin \mathcal{T}} \Delta W_1^{(\tau)}[i,j]$$

$$= \text{sign}\left(w_{2i}^{(0)} \right) \left(v_j^{(T_1)} - \sum_{\tau \in \mathcal{T}} v_j^{(\tau)} \right) + \mathcal{O}\left(\frac{\eta^{\frac{1}{4}}}{d^{\frac{1}{2}}} + \frac{1}{d^{\frac{\alpha}{2}+\frac{1}{4}}} \right)$$

$$:= \text{sign}\left(w_{2i}^{(0)} \right) V_j^{(t)} + R_1^{(t)}[i,j], \text{ where } \left| R_1^{(t)}[i,j] \right| \leq \mathcal{O}\left(\frac{\eta^{\frac{1}{4}}}{d^{\frac{1}{2}}} + \frac{1}{d^{\frac{\alpha}{2}+\frac{1}{4}}} \right).$$

By assumption $\left| W_1^{(t)}[i,j] \right| = \Omega\left(\frac{1}{\sqrt{d}} \right)$, we get that $\left| V_j^{(t)} \right| = \Omega\left(\frac{1}{\sqrt{d}} \right)$, which gives us $\frac{\left| R_1^{(t)}[i,j] \right|}{\left| V_j^{(t)} \right|} \leq \mathcal{O}\left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}} \right)$.

Therefore, we have that for any $j \in [d]$,

$$\forall i_1, i_2 \in [d] : \frac{\left| W_1^{(t)}[i_1,j] \right|}{\left| W_1^{(t)}[i_2,j] \right|} = \frac{\left| \text{sign}\left(w_{2i_1}^{(0)} \right) V_j^{(t)} \left(1 \pm \mathcal{O}\left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}} \right) \right) \right|}{\left| \text{sign}\left(w_{2i_2}^{(0)} \right) V_j^{(t)} \left(1 \pm \mathcal{O}\left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}} \right) \right) \right|} = 1 \pm \mathcal{O}\left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}} \right).$$

By Lemma 23, we know that $\left| w_{2i}^{(t)} \right|$ with different i are also roughly equal, i.e. $\frac{\left| w_{2i_1}^{(t)} \right|}{\left| w_{2i_2}^{(t)} \right|} = 1 \pm \mathcal{O}\left(\sqrt{\eta} + \frac{1}{d^{\alpha-1/2}} \right)$. Then we have

$$(W_2 W_1)_j^{(t)} = \sum_{i=1}^{d} w_{2i}^{(t)} W_1^{(t)}[i,j] = \sum_{i=1}^{d} \left| w_{2i}^{(t)} \right| \left| W_1^{(t)}[i,j] \right| = \Theta\left(d \left| w_{21}^{(t)} \right| \left| W_1^{(t)}[k,j] \right| \right)$$

$$= \Theta\left(\sqrt{d} \left| W_1^{(t)}[k,j] \right| \right).$$

where k can be any index in $\{1, 2, ..., d\}$ and the last equality uses $\forall i \in [d] : \left|w_{2i}^{(t)}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right)$. Once $\left|W_1^{(t)}[k, j]\right|$ is below some value of order $\Theta\left(\frac{1}{\sqrt{d}}\right)$ such that $(W_2 W_1)_j^{(t)} < A_j - \sqrt{\eta d}$, i.e. $E_j^{(t)} < -\sqrt{\eta d}$, then $\left|W_1^{(t)}[k, j]\right|$ will increase again. Hence $\left|W_1^{(t+1)}[k, j]\right| \geq \left|W_1^{(t)}[k, j]\right| = \Omega\left(\frac{1}{\sqrt{d}}\right)$. Since k is arbitrary, we have proved that at time $t + 1$, $\forall i, j \in [d] : \left|W_1^{(t+1)}[i, j]\right| \geq \Omega\left(\frac{1}{\sqrt{d}}\right)$.

Therefore by induction, we conclude that when $T_1 \leq t < \tilde{T}$, for $\forall i, j \in [d], \left|W_1^{(t)}[i, j]\right| = \Omega\left(\frac{1}{\sqrt{d}}\right)$. The remaining part of this lemma has also been proved by the analysis above.

D.15 PROOF OF LEMMA 25

Lemma 26 tells us that for any $i \in [d]$, $\left|w_{2i}^{(t)}\right|$ keeps increasing when $t < \tilde{T}$. However, the behavior of $W_1^{(t)}[i, j]$ is more complicated. The following lemma tells us that $\left|W_1^{(t)}[i, j]\right|$ will increase until $T_{f,j}$. After that $\left|W_1^{(t)}[i, j]\right|$ and $E_j^{(t)}$ may zigzag, but $E_j^{(t)}$ will not fluctuate dramatically and will be trapped in a small interval around zero.

Lemma 27. *Under Assumption 1, 2 and 3, suppose $\sigma \leq \frac{\eta^{3/2} \xi^2}{d^{13/4}}$. Pick $\eta \leq \mathcal{O}\left(\frac{1}{d^{3\alpha}}\right), \xi \leq \sqrt{\frac{\eta}{d^{3\alpha-1}}}$, and $\beta_2 = \beta_1^2$. Consider certain coordinate j. For $T_1 \leq t < \min\left\{\tilde{T}, T_{f,j}\right\}$, we have $\forall i \in [d] :$ $\left|W_1^{(t)}[i, j]\right|$ keeps increasing. If $T_{f,j} < \tilde{T}$, then for $T_{f,j} \leq t < \tilde{T}$, we will have $-\mathcal{O}\left(\sqrt{\eta d}\right) \leq E_j^{(t)} \leq \mathcal{O}\left(\sqrt{\eta d}\right)$.*

Now we start proving Lemma 25. At time $\tilde{T}$, denote $S := \left\{j : T_{f,j} < \tilde{T}\right\}$, i.e. the set of coordinates whose E_j have passed its "flip time". By Lemma 27, we know that $\forall j \in S, \left|E_j^{\tilde{T}}\right| \leq \mathcal{O}\left(\sqrt{\eta d}\right)$. If $S^c = \phi$, which means $\forall j \in [d] : \left|E_j^{(\tilde{T})}\right| \leq \mathcal{O}\left(\sqrt{\eta d}\right)$, then our lemma will immediately follow. If $S^c \neq \phi$, we have $\tilde{T} = \min\{T_g, T_f\} = T_g$ and that $\forall j \in S^c : E_j^{(\tilde{T})} < 0$. By the definition of T_g, we know that $\exists i_0 \in [d] : \left|g_{2i_0}^{(\tilde{T})}\right| \leq \mathcal{O}\left(d\sqrt{\eta}\right)$. Then

$$\left|\sum_{j \in S^c}^{d} E_j^{(\tilde{T})} W_1^{(\tilde{T})}[i_0, j]\right| = \left|\sum_{j=1}^{d} E_j^{(\tilde{T})} W_1^{(\tilde{T})}[i_0, j] - \sum_{j \in S} E_j^{(\tilde{T})} W_1^{(\tilde{T})}[i_0, j]\right|$$

$$\leq \left|g_{2i_0}^{(\tilde{T})}\right| + \left|\sum_{j \in S} E_j^{(\tilde{T})} W_1^{(\tilde{T})}[i_0, j]\right|$$

$$\leq \mathcal{O}\left(d\sqrt{\eta}\right) + d\mathcal{O}\left(\sqrt{\eta d}\right)\mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$$

$$= \mathcal{O}\left(d\sqrt{\eta}\right).$$

For $j \in S^c$, we have $T_{f,j} > \tilde{T}$ and then Lemma 27 tells us that $\left|W_1^{(t)}[i, j]\right|$ keeps increasing for $t < \tilde{T}$ and any $i \in [d]$. Then $\left|W_1^{(\tilde{T})}[i, j]\right| = \Omega\left(\frac{1}{\sqrt{d}}\right)$. Since in Lemma 27, we also proved $\forall i \in [d] : \text{sign}\left(W_1^{(\tilde{T})}[i, j]\right) = \text{sign}\left(w_{2i}^{(0)}\right)$, combining with $\forall j \in S^c : E_j^{(\tilde{T})} < 0$ gives us that $E_j^{(\tilde{T})} W_1^{(\tilde{T})}[i_0, j]$ for different j have the same sign. Therefore for any $j_0 \in S^c$,

$$\mathcal{O}\left(d\sqrt{\eta}\right) \geq \left|\sum_{j \in S^c}^{d} E_j^{(\tilde{T})} W_1^{(\tilde{T})}[i_0, j]\right| = \sum_{j \in S^c}^{d} \left|E_j^{(\tilde{T})} W_1^{(\tilde{T})}[i_0, j]\right| \geq \left|E_{j_0}^{(\tilde{T})} W_1^{(\tilde{T})}[i_0, j_0]\right| \geq \left|E_{j_0}^{(T)}\right| \Omega\left(\frac{1}{\sqrt{d}}\right)$$

$$\Rightarrow \left|E_{j_0}^{(T)}\right| \leq \mathcal{O}\left(d\sqrt{\eta d}\right).$$

Note that the above inequality holds for any $j_0 \in S^c$, which means $\forall j \in S^c : \left| E_j^{(\tilde{T})} \right| \leq \mathcal{O}\left(d\sqrt{\eta d} \right)$. Combining with the fact that $\forall j \in S : \left| E_j^{(\tilde{T})} \right| \leq \mathcal{O}\left(d\sqrt{\eta d} \right)$ completes the proof.

D.16   PROOF OF LEMMA 27

When $t < \min\left\{ \tilde{T}, T_{f,j} \right\}$, we have that $E_j^{(t)} < -\sqrt{\eta d}$. Therefore we can use the same argument as before to prove that $\left| W_1^{(t)}[i,j] \right|$ keeps increasing, and $\text{sign}\left(W_1^{(t)}[i,j] \right) = \text{sign}\left(w_{2i}^{(0)} \right)$ for all $i \in [d]$.

At time the "flip time" $t = T_{f,j}$, by definition, $E_j^{(t)} \geq -\sqrt{\eta d}$. After that $E_j^{(t)}$ may oscillate. Now we prove that once $E_j^{(t)} \geq \sqrt{\eta d}$ (or $E_j^{(t)} \leq -\sqrt{\eta d}$), after a short period $E_j^{(t)}$ will decrease (or increase) until $E_j^{(t)} \leq \mathcal{O}\left(\sqrt{\eta d} \right)$ (or $E_j^{(t)} \geq -\mathcal{O}\left(\sqrt{\eta d} \right)$). Moreover, during this period, $E_j^{(t)}$ won't change too much.

We first recall that when $t < \tilde{T}$, Lemma 26 gives us for all $i \in [d]$, $\left| w_{2i}^{(t)} \right| = \Theta\left(\frac{1}{\sqrt{d}} \right)$, $\left| W_1^{(t)}[i,j] \right| = \mathcal{O}\left(\frac{1}{\sqrt{d}} \right)$. Then eq.(33) we obtained in the first phase analysis still holds, which tells us that the change of $E_j^{(t)}$ per step satisfies $\left| E_j^{(t+1)} - E_j^{(t)} \right| = \mathcal{O}\left(\eta\sqrt{d} \right)$ for all $t < \tilde{T}$.

We divide the analysis into two cases, based on whether these $E_j^{(t)} \geq \sqrt{\eta d}$ or $E_j^{(t)} \leq -\sqrt{\eta d}$. By Lemma 24, we know that when $T_1 \leq t < \tilde{T}$, $\left| W_1^{(t)}[i,j] \right| = \Omega\left(\frac{1}{\sqrt{d}} \right)$, which means the sign of $W_1^{(t)}[i,j]$ remains unchanged during this period and $\text{sign}\left(W_1^{(t)}[i,j] \right) = \text{sign}\left(W_1^{(T_1)}[i,j] \right) = \text{sign}\left(w_{2i}^{(0)} \right)$ for all $i,j \in [d]$.

By the analysis of $w_{2i}^{(t)}$ in Lemma 23, we have for all $i \in [d]$, $w_{2i}^{(t+1)} = w_{2i}^{(t)} + \text{sign}\left(w_{2i}^{(0)} \right) \Delta_{2i}^{(t)}$, where $\Delta_{2i}^{(t)} = \eta\left(1 \pm \mathcal{O}\left(\sqrt{\eta} \right) \right)$.

Case 1.   Consider some time point t such that $E_j^{(t)} \leq -\sqrt{\eta d}$. Note that for all $i \in [d]$, $\left| g_1^{(t)}[i,j] \right| = \left| w_{2i}^{(t)} E_j^{(t)} \right| = \Omega\left(\sqrt{\eta} \right)$ and that $\text{sign}\left(g_1^{(t)}[i,j] \right) = -\text{sign}\left(w_{2i}^{(t)} \right) = -\text{sign}\left(w_{2i}^{(0)} \right)$. By Lemma 23, for all $i \in [d]$ we have $W_1^{(t+1)}[i,j] = W_1^{(t)}[i,j] + \text{sign}\left(w_{2i}^{(0)} \right) \Delta_1^{(t)}[i,j]$ with $\Delta_1^{(t)}[i,j] = \eta\left(1 \pm \mathcal{O}\left(\sqrt{\eta} \right) \right)$. That gives us

$$E_j^{(t+1)} = \sum_{i=1}^d w_{2i}^{(t+1)} W_1^{(t+1)}[i,j] = \sum_{i=1}^d \left(w_{2i}^{(t)} + \text{sign}\left(w_{2i}^{(0)} \right) \Delta_{2i}^{(t)} \right)\left(W_1^{(t)}[i,j] + \text{sign}\left(w_{2i}^{(0)} \right) \Delta_1^{(t)}[i,j] \right)$$

$$= \sum_{i=1}^d \left(w_{2i}^{(t)} W_1^{(t)}[i,j] + \text{sign}\left(w_{2i}^{(0)} \right)\left(w_{2i}^{(t)} \Delta_1^{(t)}[i,j] + \Delta_{2i}^{(t)} W_1^{(t)}[i,j] \right) + \Delta_{2i}^{(t)} \Delta_1^{(t)}[i,j] \right)$$

$$\overset{(i)}{=} E_j^{(t)} + \sum_{i=1}^d \left(\left| w_{2i}^{(t)} \right| \Delta_1^{(t)}[i,j] + \Delta_{2i}^{(t)} \left| W_1^{(t)}[i,j] \right| + \Delta_{2i}^{(t)} \Delta_1^{(t)}[i,j] \right),$$

$$\Rightarrow \quad E_j^{(t+1)} > E_j^{(t)},$$

where (i) is because $\text{sign}\left(w_{2i}^{(t)} \right) = \text{sign}\left(W_1^{(t)}[i,j] \right) = \text{sign}\left(w_{2i}^{(0)} \right)$. Therefore we have proved that $E_j^{(t)}$ will increase in the next step. After that for $\tau \geq t+1$, as long as $E_j^{(\tau)} \leq -\sqrt{\eta d}$, the above analysis will hold and $E_j^{(\tau)}$ will keep increasing (until $E_j^{(\tau)} > -\sqrt{\eta d}$).

Case 2. Consider some time point t such that $E_j^{(t)} \geq \sqrt{\eta d}$. We will prove that $E_j^{(t)}$ will decrease after a short period, and during this period, the change of it is at most $\mathcal{O}\left(\sqrt{\eta d}\right)$.

By similar arguments as in Case 1, we can get that $W_1^{(t+1)}[i,j] = W_1^{(t)}[i,j] - \text{sign}\left(w_{2i}^{(0)}\right)\Delta_1^{(t)}[i,j]$, where $\Delta_1^{(t)}[i,j] = \eta\left(1 \pm \mathcal{O}\left(\sqrt{\eta}\right)\right)$, Then

$$E_j^{(t+1)} = \sum_{i=1}^{d} w_{2i}^{(t+1)} W_1^{(t+1)}[i,j] = \sum_{i=1}^{d}\left(w_{2i}^{(t)} + \text{sign}\left(w_{2i}^{(0)}\right)\Delta_{2i}^{(t)}\right)\left(W_1^{(t)}[i,j] - \text{sign}\left(w_{2i}^{(0)}\right)\Delta_1^{(t)}[i,j]\right)$$

$$= \sum_{i=1}^{d}\left(w_{2i}^{(t)}W_1^{(t)}[i,j] - \text{sign}\left(w_{2i}^{(0)}\right)\left(w_{2i}^{(t)}\Delta_1^{(t)}[i,j] - \Delta_{2i}^{(t)}W_1^{(t)}[i,j]\right) - \Delta_{2i}^{(t)}\Delta_1^{(t)}[i,j]\right)$$

$$\overset{(i)}{=} E_j^{(t)} - \sum_{i=1}^{d}\left(\left|w_{2i}^{(t)}\right|\Delta_1^{(t)}[i,j] - \Delta_{2i}^{(t)}\left|W_1^{(t)}[i,j]\right| + \Delta_{2i}^{(t)}\Delta_1^{(t)}[i,j]\right),$$

where (i) is because $\text{sign}\left(w_{2i}^{(t)}\right) = \text{sign}\left(W_1^{(t)}[i,j]\right) = \text{sign}\left(w_{2i}^{(0)}\right)$. $E_j^{(t+1)}$ may not be smaller than $E_j^{(t)}$, but we will show that after at most t_s steps for some t_s, we will have $E_j^{(t+t_s+1)} < E_j^{(t+t_s)}$.

To see this, first note that by the bounds of $\Delta_1^{(t)}[i,j]$ and $\Delta_{2i}^{(t)}$. we get $\Delta_1^{(t)}[i,j] \geq \Delta_{2i}^{(t)}[i,j] - \eta\mathcal{O}\left(\sqrt{\eta}\right)$. Since $\left|w_{2i}^{(t)}\right|$ increases by $\Theta(\eta)$ per step, and $\left|W_1^{(t)}[i,j]\right|$ keeps decreasing, then after t_s steps for some t_s, we have $\forall i \in [d]: \left|w_{2i}^{(t+t_s)}\right| \geq \left|W_1^{(t+t_s)}[i,j]\right| + \sqrt{\eta}$.

Let's first deal with the case $t + t_s \leq \tilde{T}$. If $E_j^{(t+t_s)} < \sqrt{\eta d}$, then it's already what we want. Otherwise we will have $\Delta_1^{(t+t_s)}[i,j] = \eta\left(1 \pm \mathcal{O}\left(\sqrt{\eta}\right)\right)$. Hence

$$E_j^{(t+t_s)} - E_j^{(t+t_s+1)}$$

$$= \sum_{i=1}^{d}\left(\left|w_{2i}^{(t+t_s)}\right|\Delta_1^{(t+t_s)}[i,j] - \Delta_{2i}^{(t+t_s)}\left|W_1^{(t+t_s)}[i,j]\right| + \Delta_{2i}^{(t+t_s)}\Delta_1^{(t+t_s)}[i,j]\right)$$

$$\geq \sum_{i=1}^{d}\left(\left|W_1^{(t+t_s)}[i,j]\right|\left(\Delta_1^{(t+t_s)}[i,j] - \Delta_{2i}^{(t+t_s)}\right) + \sqrt{\eta}\Delta_1^{(t+t_s)}[i,j] + \Delta_{2i}^{(t+t_s)}\Delta_1^{(t+t_s)}[i,j]\right)$$

$$\geq \sum_{i=1}^{d}\left(-\eta\mathcal{O}\left(\sqrt{\eta}\right)\left|W_1^{(t+t_s)}[i,j]\right| + \eta\sqrt{\eta} + \Delta_{2i}^{(t+t_s)}\Delta_1^{(t+t_s)}[i,j]\right) > 0,$$

where the last inequality uses $\forall i, j \in [d]: \left|W_1^{(t+t_s)}[i,j]\right| = \mathcal{O}\left(\frac{1}{\sqrt{d}}\right)$. Therefore $E_j^{(t+t_s+1)} < E_j^{(t+t_s)}$. After that for $\tau \geq t + t_s + 1$, as long as $E_j^{(\tau)} \geq \sqrt{\eta d}$, the above analysis will hold and $E_j^{(\tau)}$ will keep decreasing (until $E_j^{(\tau)} < \sqrt{\eta d}$).

Now we prove that during these at most t_s steps, the change of E_j is $\mathcal{O}\left(\sqrt{\eta d}\right)$.

Since at each step the difference $|w_{2i}| - |W_1[i,j]|$ will be enlarged by $\Omega(\eta)$, then we know that $t_s = \sqrt{\eta}/\Omega(\eta) = \mathcal{O}\left(\frac{1}{\sqrt{\eta}}\right)$. Combining with the fact that for all $\tau \leq \tilde{T}$, $\left|E_j^{(\tau+1)} - E_j^{(\tau)}\right| = \mathcal{O}\left(\eta\sqrt{d}\right)$ gives us

$$E_j^{(t+t_s)} - E_j^{(t)} \leq \mathcal{O}\left(\eta t_s \sqrt{d}\right) = \mathcal{O}\left(\sqrt{\eta d}\right).$$

If $t + t_s > \tilde{T}$, we also have the same bound, because $E_j^{\tilde{T}} - E_j^{(t)} \leq \mathcal{O}\left(\eta(\tilde{T} - t)\sqrt{d}\right) \leq \mathcal{O}\left(\eta t_s \sqrt{d}\right) = \mathcal{O}\left(\sqrt{\eta d}\right)$.

Combining the above two cases, we find that if for some t, $E_j^{(t)} \geq \sqrt{\eta d}$, then after at most t_s steps E_j will decrease and keeps decreasing until $E_j < \sqrt{\eta d}$. During these steps, E_j can increase at most

$\mathcal{O}\left(\sqrt{\eta d}\right)$. If for some t, $E_j^{(t)} \leq -\sqrt{\eta d}$, then after one step it will increase and keeps increasing until $E_j > \sqrt{\eta d}$. That means once for some coordinate j, E_j overshoots, then it will zigzag in a small region around zero, which is $\left[-\mathcal{O}\left(\sqrt{\eta d}\right), \mathcal{O}\left(\sqrt{\eta d}\right)\right]$.

E   HESSIAN TENDS TO BECOME MORE AND MORE DIAGONAL DURING TRAINING

In this section, we empirically demonstrate that the trend of loss Hessian in practice is to become more and more diagonal during training. We also give a rigorous theoretical analysis on a two-layer network under Assumption 1 and 2.

E.1   EMPIRICAL RESULTS

Let's first define the diagonal domination of the i-th coordinate at time t.

$$r_{\mathrm{diag},i}^{\mathrm{OPT}}(t) := \frac{\sqrt{\sum_{j \neq i}\left(H^{(t)}[i,j]\right)^2}}{\left|H^{(t)}[i,i]\right|}.$$

To measure the diagonal domination of the whole Hessian, we need to consider the distribution of $r_{\mathrm{diag},i}^{\mathrm{OPT}}(t)$ for different i. Figure 14 shows the mean and median of $r_{\mathrm{diag},i}^{\mathrm{SGDM}}(t)$ and $r_{\mathrm{diag},i}^{\mathrm{Adam}}(t)$ on the sentence classification task using BERT-small. Here we chose 4 layers (Layer #6, 12, 17 and 22) and computed the Hessians across these 4 layers. Since the number of parameters is very large, we did the computation by random sampling. As we can see, for both $r_{\mathrm{diag},i}^{\mathrm{SGDM}}(t)$ and $r_{\mathrm{diag},i}^{\mathrm{Adam}}(t)$, the trend of their mean or median is to decrease over time, although there might be some oscillation.

(a) Mean   (b) Median

Figure 14: Mean and median of $r_{\mathrm{diag},i}^{\mathrm{SGDM}}(t)$ and $r_{\mathrm{diag},i}^{\mathrm{Adam}}(t)$ for the full hessian across the four layers (#6,12,17,22)

E.2   THEORETICAL ANALYSIS

To simplify the theoretical analysis, we consider the mean of $r_{\mathrm{diag},i}^{\mathrm{OPT}}(t)$ over all coordinate and define

$$R_{\mathrm{diag}}^{\mathrm{OPT}}(t) := \mathrm{mean}\left(r_{\mathrm{diag},i}^{\mathrm{OPT}}(t)\right). \tag{39}$$

We consider a 2-layer network under Assumption 1 and 2, and have two goals in our proof:

1. To show that $R_{\mathrm{diag}}^{\mathrm{OPT}}(t)$ after training is smaller than that before training ($t = 0$).

2. Note that in our setting (see in Assumption 1), the Hessian is a $(d^2 + d) \times (d^2 + d)$ matrix. For a completely "uniform" matrix with the same size, we have that $R_{\mathrm{diag}}^{\mathrm{OPT}}(t) = \Theta\left(\sqrt{d^2 + d}\right) = \Theta(d)$. Hence our second goal is to show that the $R_{\mathrm{diag}}^{\mathrm{OPT}}(t)$ after training is on lower order than $\Theta(d)$.

Theorem 2. *Consider the ratio $R_{diag}^{OPT}(t)$ defined in eq. (39). Under Assumption 1 and 2, we have that before training ($t = 0$), with high probability,*

$$R_{diag}^{OPT}(0) \geq \tilde{\Omega}\left(d^{4\alpha - \frac{3}{2}}\right). \tag{40}$$

For SGD+M defined in eq. (3). For any $p > 0$, by picking the same hyperparameters as in Theorem 1, for $T_{SGD,1}, T_{SGD,2}$ mentioned in Theorem 1, when $t \in [T_{SGD,1}, T_{SGD,2}]$, we have with constant probability,

$$R_{diag}^{SGDM}(t) \leq \tilde{\mathcal{O}}\left(\sqrt{d}\right) + q^{(t)}, \tag{41}$$

where the trend of $q^{(t)}$ is to decrease over time and $q^{(T_{SGD,2})} \leq \tilde{\mathcal{O}}\left(\frac{1}{d^{p/2-1}}\right) = \tilde{o}(d)$.

For Adam defined in eq. (3). For any $p > 0$, by picking the same hyperparameters as in Theorem 1, for $T_{Adam,1}, T_{Adam,2}$ mentioned in Theorem 1, when $t \in [T_{Adam,1}, T_{Adam,2}]$, we have with high probability,

$$R_{diag}^{Adam}(t) \leq \mathcal{O}\left(\sqrt{d}\right) + r^{(t)}, \tag{42}$$

where the trend of $r^{(t)}$ is to decrease over time and $r^{(T_{Adam,2})} \leq \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{p-1}{2}}}\right) = \tilde{o}\left(\sqrt{d}\right)$.

E.3 PROOF OF THEOREM 2

Lemma 4.3 of (Kawaguchi, 2016) gives us the following forms of Hessian.

For any $k \in \{1, 2, ..., H+1\}$, we know that

$$\nabla_{vec(W_k)}(\nabla_{vec(W_k)}L(W)) = ((W_{H+1}\ldots W_{k+1})^T(W_{H+1}\ldots W_{k+1}) \otimes (W_{k-1}\ldots W_1)(W_{k-1}\ldots W_1)^T,$$

and for $k \in \{2, 3, ..., H+1\}$,

$$\nabla_{vec(W_k)}(\nabla_{vec(W_1)}L(W))$$
$$=(C^T(W_{H+1}\ldots W_{k+1}) \otimes (W_{k-1}\ldots W_1)^T)$$
$$\quad + [(W_{k-1}\ldots W_2)^T \otimes I][I_{d_{k-1}} \otimes (r(W_{H+1}\ldots W_{k+1}))_{.,1}\cdots I_{d_{k-1}} \otimes (r(W_{H+1}\ldots W_{k+1}))_{.,d_k}],$$

where $r = (W_{H+1}\ldots W_1 - A)^T, C = W_{H+1}W_H\cdots W_2$.

For the 2-layer linear network, write the Hessian as

$$H := \left[\begin{array}{cc} H_{22} & H_{21}^T \\ H_{21} & H_{11} \end{array}\right],$$

then we have that

$$H_{11} = (W_2^T W_2) \otimes I_d \in \mathbb{R}^{d^2 \times d^2},$$
$$H_{22} = W_1 W_1^T \in \mathbb{R}^{d \times d},$$
$$H_{21} = W_2^T \otimes W_1^T + I_d \otimes (W_2 W_1 - A)^T \in \mathbb{R}^{d^2 \times d}.$$

Intuitively, before training the elements of W_1 and W_2 are very close to zero, and $W_2 W_1 - A \approx -A$. Since the elements of A are $\Theta(1)$, we know that the magnitudes of elements of H_{21} are much bigger than those of H_{11} and H_{22}.

After training, for both SGD+M and Adam, $W_2 W_1 - A \approx 0$. Then $H_{21} \approx (W_2)^T \otimes (W_1)^T$ and the magnitudes of its elements are no longer much larger than those of H_{11} and H_{22}. From the formula of H_{11}, we know that all the diagonal entries are nonzero, and among the $d^4 - d^2$ off-diagonal entries, there are only $d^3 - d^2$ nonzero entries, which helps us to bound $R_{diag}^{OPT}(t)$.

E.3.1 PROOF OF EQ. (40)

Let's first analyze the weights and Hessian before training ($t = 0$). For ease of notation, we omit the superscript (t).

For the i-th row where $1 \leq i \leq d$, i.e. the i-th row of the submatrix $[H_{22} \quad H_{21}^T]$, we have

$$\sum_{j \neq i} H^2[i, j] = \sum_{j \neq i} H_{22}^2[i, j] + \sum_{j=1}^{d^2} H_{21}^2[j, i]$$
$$\geq \sum_{j=(i-1)d}^{id} H_{21}^2[j, i] = \sum_{j=1}^{d} (w_{2i}W_1[i, j] + (W_2 W_1 - A)_j)^2 = \Theta(d).$$

On the other hand, for the diagonal elements, we have

$$|H[i,i]| = |H_{22}[i,i]| = \|W_1[i,:]\|_2^2 = \sum_{j=1}^{d} W_1^2[i,j] \leq \tilde{\mathcal{O}}\left(\frac{1}{d^{4\alpha-1}}\right).$$

Then we have that for $1 \leq i \leq d$,

$$\frac{\sqrt{\sum_{j\neq i} H^2[i,j]}}{|H[i,i]|} \geq \frac{\sqrt{\Omega(d)}}{\tilde{\mathcal{O}}\left(\frac{1}{d^{4\alpha-1}}\right)} = \tilde{\Omega}\left(d^{4\alpha-\frac{1}{2}}\right).$$

For the $(id+k)$-th row where $1 \leq i \leq d, 1 \leq k \leq d$, i.e. the $((i-1)d+k)$-th row of the submatrix $[H_{21} \quad H_{11}]$, we have

$$\sum_{j\neq id+k} H^2[i,j] = \sum_{j\neq(i-1)d+k} H_{11}^2[i,j] + \sum_{j=1}^{d} H_{21}^2[(i-1)d+k,j]$$

$$\geq H_{21}^2[(i-1)d+k,i] = (w_{2i}W_1[i,k] + (W_2W_1 - A)_k)^2 = \Theta(1).$$

On the other hand, for the diagonal elements, we have

$$|H[id+k,id+k]| = |H_{11}[(i-1)d+k,(i-1)d+k]| = w_{2i}^2 \leq \tilde{\mathcal{O}}\left(\frac{1}{d^{2\alpha}}\right).$$

Then we have that for $1 \leq i \leq d, 1 \leq k \leq d$,

$$\frac{\sqrt{\sum_{j\neq id+k} H^2[i,j]}}{|H[id+k,id+k]|} \geq \frac{\sqrt{\Omega(1)}}{\tilde{\mathcal{O}}\left(\frac{1}{d^{2\alpha}}\right)} = \tilde{\Omega}\left(d^{2\alpha}\right).$$

Taking the average, we obtain that before training, i.e. when $t = 0$,

$$R_{\text{diag}}^{(0)} \geq \frac{d\tilde{\Omega}\left(d^{4\alpha-\frac{1}{2}}\right) + d^2\tilde{\Omega}\left(d^{2\alpha}\right)}{d^2 + d} = \tilde{\Omega}\left(d^{4\alpha-\frac{3}{2}}\right).$$

E.3.2 PROOF OF EQ. (41)

The proof is based on the lemma below.

Lemma 28. *Suppose the weight matrices have the following structure:*

$$W_1 = \boldsymbol{u}\boldsymbol{v}^T + R_1,$$
$$W_2 = c\boldsymbol{u}^T + R_2^T,$$

where $\forall 1 \leq i,j \leq d: \quad \frac{|R_1[i,j]|}{|u_iv_j|} \leq \delta, \quad \frac{|R_{2i}|}{|cu_i|} \leq \delta, \quad \delta \in (0,1)..$

Then we have for $1 \leq i \leq d$,

$$\frac{\sqrt{\sum_{j\neq i} H^2[i,j]}}{|H[i,i]|} \leq \frac{1+\delta}{1-\delta}\left(1 + \frac{|c|}{\|\boldsymbol{v}\|_2}\right)\sqrt{\frac{\sum_{j=1}^{d} u_j^2}{u_i^2}} + \frac{\|E\|_2}{(1-\delta)^2 u_i^2 \|v\|_2^2},$$

and for $1 \leq i \leq d, 1 \leq k \leq d$,

$$\frac{\sqrt{\sum_{j\neq id+k} H^2[i,j]}}{|H[id+k,id+k]|} \leq \frac{1+\delta}{1-\delta}\left(1 + \frac{|v_k|}{|c|}\right)\sqrt{\frac{\sum_{j=1}^{d} u_j^2}{u_i^2}} + \frac{|E_k|}{(1-\delta)^2 c^2 u_i^2}.$$

Now we are ready to prove eq. (41).

By the analyses in Section C.1, we know that for $t \in [T_{\text{SGD},1}, T_{\text{SGD},2}]$, the weights obtained by GD with momentum satisfy

$$W_1^{(t)} = \boldsymbol{u}^{(T_1)}\boldsymbol{v}^{(t)T} + R_1^{(t)},$$
$$W_2^{(t)} = c^{(t)}\boldsymbol{u}^{(T_1)T} + R_2^{(t)T},$$

where $T_{\text{SGD},1} = T_1$ and

$$\forall 1 \leq i, j \leq d: \quad \frac{\left| R_1^{(t)}[i,j] \right|}{\left| u_i^{(T_1)} v_j^{(t)} \right|} \leq \tilde{\mathcal{O}}(\epsilon_0), \quad \frac{\left| R_{2i}^{(t)} \right|}{\left| c^{(t)} u_i^{(T_1)} \right|} \leq \tilde{\mathcal{O}}(\epsilon_0).$$

Here ϵ_0 is defined in Definition 2. Since $\boldsymbol{u}^{(T_1)}$ doesn't depend on $T_1 = T_{\text{SGD},1} < t \leq T_{\text{SGD},2}$, we write $\boldsymbol{u}^{(T_1)}$ as $\boldsymbol{u}$ for ease of notation.

Hence by Lemma 28, when $t \in [T_{\text{SGD},1}, T_{\text{SGD},2}]$, we have for $1 \leq i \leq d$,

$$\frac{\sqrt{\sum_{j \neq i} \left(H^{(t)}[i,j] \right)^2}}{\left| H^{(t)}[i,i] \right|} \leq \frac{1 + \tilde{\mathcal{O}}(\epsilon_0)}{1 - \tilde{\mathcal{O}}(\epsilon_0)} \left(1 + \frac{|c^{(t)}|}{\left\| \boldsymbol{v}^{(t)} \right\|_2} \right) \sqrt{\frac{\sum_{j=1}^d u_j^2}{u_i^2}} + \frac{\left\| E^{(t)} \right\|_2}{\left(1 - \tilde{\mathcal{O}}(\epsilon_0) \right)^2 u_i^2 \left\| \boldsymbol{v}^{(t)} \right\|_2^2}$$

$$= \mathcal{O} \left(1 + \frac{|c^{(t)}|}{\left\| \boldsymbol{v}^{(t)} \right\|_2} \right) \sqrt{\frac{\sum_{j=1}^d u_j^2}{u_i^2}} + \mathcal{O} \left(\frac{\left\| E^{(t)} \right\|_2}{u_i^2 \left\| \boldsymbol{v}^{(t)} \right\|_2^2} \right), \tag{43}$$

and for $1 \leq i \leq d, 1 \leq k \leq d$,

$$\frac{\sqrt{\sum_{j \neq id+k} \left(H^{(t)}[i,j] \right)^2}}{\left| H^{(t)}[id+k, id+k] \right|} \leq \frac{1 + \tilde{\mathcal{O}}(\epsilon_0)}{1 - \tilde{\mathcal{O}}(\epsilon_0)} \left(1 + \frac{|v_k^{(t)}|}{|c^{(t)}|} \right) \sqrt{\frac{\sum_{j=1}^d u_j^2}{u_i^2}} + \frac{\left| E_k^{(t)} \right|}{\left(1 - \tilde{\mathcal{O}}(\epsilon_0) \right)^2 \left(c^{(t)} \right)^2 u_i^2}$$

$$= \mathcal{O} \left(1 + \frac{|v_k^{(t)}|}{|c^{(t)}|} \right) \sqrt{\frac{\sum_{j=1}^d u_j^2}{u_i^2}} + \mathcal{O} \left(\frac{\left| E_k^{(t)} \right|}{\left(c^{(t)} \right)^2 u_i^2} \right). \tag{44}$$

By Lemma 3, we have $\boldsymbol{u} = X + Y$ where $X_i, i \in [d]$ are i.i.d Gaussian random variables and w.h.p.,

$$\forall i \in [d]: \frac{|Y_i|}{|X_i|} \leq \frac{\tilde{\mathcal{O}} \left(\frac{1}{d^{5\alpha - \frac{3}{2}}} \right)}{\Omega \left(\frac{1}{d^{\frac{5}{4}\alpha}} \right)} = \tilde{\mathcal{O}} \left(\frac{1}{d^{\frac{15}{4}\alpha - \frac{3}{2}}} \right) := \delta_{xy}, \tag{45}$$

which yields that

$$\forall i \in [d]: \frac{\sqrt{\sum_{j=1}^d u_j^2}}{|u_i|} \leq \left(\frac{1 + \delta_{xy}}{1 - \delta_{xy}} \right)^2 \frac{\sqrt{\sum_{j=1}^d X_j^2}}{|X_i|}, \quad \frac{1}{u_i^2} \leq \left(\frac{1}{1 - \delta_{xy}} \right)^2 \frac{1}{X_i^2}. \tag{46}$$

By the proof in Section C.8, we know that for $t \in [T_{\text{SGD},1}, T_{\text{SGD},2}], \forall i \in [d]: v_i^{(t)}, c^{(t)}$ are positive. The induction in Section C.9 further gives us that for $t \in [T_{\text{SGD},1}, T_{\text{SGD},2}]$, w.h.p. $\forall k \in [d]: \frac{v_k^{(t)}}{c^{(t)}} = \Theta \left(\frac{1}{\sqrt{d}} \right)$, which yields $\frac{c^{(t)}}{\left\| \boldsymbol{v}^{(t)} \right\|_2} = \Theta(1)$. Combining with eq. (46), we obtain

$$\left(1 + \frac{|c^{(t)}|}{\left\| \boldsymbol{v}^{(t)} \right\|_2} \right) \sqrt{\frac{\sum_{j=1}^d u_j^2}{u_i^2}} \leq \mathcal{O} \left(\frac{\sqrt{\sum_{j=1}^d X_j^2}}{|X_i|} \right),$$

$$\left(1 + \frac{|v_k^{(t)}|}{|c^{(t)}|} \right) \sqrt{\frac{\sum_{j=1}^d u_j^2}{u_i^2}} \leq \mathcal{O} \left(\frac{\sqrt{\sum_{j=1}^d X_j^2}}{|X_i|} \right). \tag{47}$$

By the proof in Section C.8, we know that for $t \in [T_{\text{SGD},1}, T_{\text{SGD},2}], \forall i \in [d]: v_i^{(t)}, c^{(t)}$ are positive and monotonically increasing. On the other hand, the proof in Section C.2 and C.9 tells us that w.h.p. $\left\| E^{(t)} \right\|_2$ (resp. $\forall k \in [d], \left| E_k^{(t)} \right|$) decreases from $\Theta(\sqrt{d})$ (resp. $\Theta(1)$) when $t = T_{\text{SGD},1}$ to $\mathcal{O}(\sqrt{\epsilon_0 d})$ (resp. $\mathcal{O}(\sqrt{\epsilon_0})$) when $t = T_{\text{SGD},2}$. Therefore, the trend of $\frac{\left\| E^{(t)} \right\|_2}{u_i^2 \left\| \boldsymbol{v}^{(t)} \right\|_2^2}$ and $\frac{\left| E_k^{(t)} \right|}{\left(c^{(t)} \right)^2 u_i^2}$ is to decrease over time, and when $t = T_{\text{SGD},2}$, we have w.h.p.

$$\forall k \in [d]: \left| E_k^{(t)} \right| = \mathcal{O} \left(\sqrt{\epsilon_0} \right), \quad \left\| E^{(t)} \right\|_2 = \mathcal{O} \left(\sqrt{\epsilon_0 d} \right). \tag{48}$$

Moreover, when $t = T_{\text{SGD},2}$, the inequality in eq. (25) becomes equality, i.e. $c^2\|\boldsymbol{u}\|_2^2 = \Theta\left(\sqrt{d}\right)$ and $\forall j \in [d] : \|\boldsymbol{u}\|_2^2 v_j^2 = \Theta\left(\frac{1}{\sqrt{d}}\right)$.

Using $\boldsymbol{u} = X + Y$ and eq. (45), we have

$$c^2\|X\|_2^2 = \Theta\left(\sqrt{d}\right), \quad \forall j \in [d] : \|X\|_2^2 v_j^2 \Theta\left(\frac{1}{\sqrt{d}}\right), \quad \Rightarrow \quad \|X\|_2^2\|\boldsymbol{v}\|_2^2 = \Theta\left(\sqrt{d}\right),$$

which together with the second inequality in eq. (46) yields

$$\frac{1}{u_i^2\|\boldsymbol{v}\|_2^2} \leq \left(\frac{1}{1-\delta_{xy}}\right)^2 \frac{1}{X_i^2\|\boldsymbol{v}\|_2^2} = \Theta\left(\frac{\sum_{j=1}^d X_j^2}{X_i^2\sqrt{d}}\right),$$

$$\frac{1}{c^2 u_i^2} \leq \left(\frac{1}{1-\delta_{xy}}\right)^2 \frac{1}{c^2 X_i^2} = \Theta\left(\frac{\sum_{j=1}^d X_j^2}{X_i^2\sqrt{d}}\right).$$

Combining with eq. (48), we get that

$$\frac{\left\|E^{(t)}\right\|_2}{u_i^2\left\|\boldsymbol{v}^{(t)}\right\|_2^2} \leq \mathcal{O}\left(\frac{\sum_{j=1}^d X_j^2}{X_i^2} \cdot \sqrt{\epsilon_0}\right), \quad \frac{\left|E_k^{(t)}\right|}{\left(c^{(t)}\right)^2 u_i^2} \leq \mathcal{O}\left(\frac{\sum_{j=1}^d X_j^2}{X_i^2} \cdot \sqrt{\frac{\epsilon_0}{d}}\right). \tag{49}$$

Substituting eq. (47) and (49) into eq. (43) and (44) gives us

$$\forall 1 \leq i \leq d : \frac{\sqrt{\sum_{j\neq i}\left(H^{(t)}[i,j]\right)^2}}{\left|H^{(t)}[i,i]\right|} \leq \mathcal{O}\left(\frac{\sqrt{\sum_{j=1}^d X_j^2}}{|X_i|}\right) + q_{1i}^{(t)},$$

where the trend of $q_{1i}^{(t)}$ is to decrease over time and $q_{1i}^{(T_{\text{SGD},2})} \leq \mathcal{O}\left(\frac{\sum_{j=1}^d X_j^2}{X_i^2} \cdot \sqrt{\epsilon_0}\right)$.

We also have

$$\forall 1 \leq i \leq d, 1 \leq k \leq d : \frac{\sqrt{\sum_{j\neq id+k}\left(H^{(t)}[i,j]\right)^2}}{\left|H^{(t)}[id+k,id+k]\right|} \leq \mathcal{O}\left(\frac{\sqrt{\sum_{j=1}^d X_j^2}}{|X_i|}\right) + q_{2i}^{(t)},$$

where the trend of $q_{2i}^{(t)}$ is to decrease over time and $q_{2i}^{(T_{\text{SGD},2})} \leq \mathcal{O}\left(\frac{\sum_{j=1}^d X_j^2}{X_i^2} \cdot \sqrt{\frac{\epsilon_0}{d}}\right)$.

Hence

$$R_{\text{diag}}^{\text{SGDM}}(t) = \mathcal{O}\left(\frac{1}{d}\sum_{i=1}^d \frac{\sqrt{\sum_{j=1}^d X_j^2}}{|X_i|}\right) + \frac{1}{d^2+d}\sum_{i=1}^d q_{1i}^{(t)} + \frac{d}{d^2+d}\sum_{i=1}^d q_{2i}^{(t)}$$

$$:= \mathcal{O}\left(\frac{1}{d}\sum_{i=1}^d \frac{\sqrt{\sum_{j=1}^d X_j^2}}{|X_i|}\right) + q^{(t)},$$

where the trend of $q^{(t)}$ is to decrease over time and

$$q^{(T_{\text{SGD},2})} \leq \frac{1}{d^2+d}\sum_{i=1}^d \mathcal{O}\left(\frac{\sum_{j=1}^d X_j^2}{X_i^2} \cdot \sqrt{\epsilon_0}\right) + \frac{d}{d^2+d}\sum_{i=1}^d \mathcal{O}\left(\frac{\sum_{j=1}^d X_j^2}{X_i^2} \cdot \sqrt{\frac{\epsilon_0}{d}}\right)$$

$$\leq \mathcal{O}\left(\frac{1}{d^2+d}\sum_{i=1}^d \frac{\sum_{j=1}^d X_j^2}{X_i^2} \cdot \sqrt{\epsilon_0 d}\right) = \mathcal{O}\left(\frac{1}{d}\sum_{i=1}^d \frac{\sum_{j=1}^d X_j^2}{X_i^2} \cdot \sqrt{\frac{\epsilon_0}{d}}\right).$$

Denote σ^2 as the variance of X_i for $i \in [d]$. By concentration of chi-squared distribution, we know that with probability at least $1 - \delta$ for $\delta > 0$,

$$\sum_{i=1}^d X_i^2 \leq \sigma^2 d + \sigma^2\mathcal{O}\left(\sqrt{d\log\frac{1}{\delta}}\right).$$

By Lemma 35 in Appendix G, we know that with constant probability $\frac{1}{d} \sum_{i=1}^{d} \frac{1}{|X_i|} = \mathcal{O}\left(\frac{1}{\sigma} \log d\right)$.

Then with constant probability, $\frac{1}{d} \sum_{i=1}^{d} \frac{1}{X_i^2} \leq \frac{1}{d} \left(\sum_{i=1}^{d} \frac{1}{|X_i|}\right)^2 = \mathcal{O}\left(\frac{d}{\sigma^2} \log^2 d\right)$. Hence

$$\frac{1}{d} \sum_{i=1}^{d} \frac{\sqrt{\sum_{j=1}^{d} X_j^2}}{|X_i|} = \tilde{\mathcal{O}}(\sqrt{d}), \quad \frac{1}{d} \sum_{i=1}^{d} \frac{\sum_{j=1}^{d} X_j^2}{X_i^2} = \tilde{\mathcal{O}}\left(d^2\right).$$

Therefore with constant probability,

$$R_{\text{diag}}^{\text{SGDM}}(t) = \tilde{\mathcal{O}}\left(\sqrt{d}\right) + q^{(t)},$$

where the trend of $q^{(t)}$ is to decrease over time and $q^{(T_{\text{SGD},2})} \leq \tilde{\mathcal{O}}\left(d\sqrt{\epsilon_0 d}\right)$. For any $p > 0$, by picking the same hyperparameters as in Theorem 1, we have $\epsilon_0 d \leq \tilde{\mathcal{O}}\left(\frac{1}{d^p}\right)$ and hence $q^{(T_{\text{SGD},2})} \leq \tilde{\mathcal{O}}\left(\frac{1}{d^{p/2-1}}\right) = \tilde{o}(d)$.

E.3.3 PROOF OF EQ. (42)

By the analyses in Section D.1, we know that for $t \in [T_{\text{Adam},1}, T_{\text{Adam},2}]$, the weights obtained by Adam satisfy

$$W_1^{(t)} = \boldsymbol{u}\boldsymbol{v}^{(t)T} + R_1^{(t)},$$
$$W_2^{(t)} = c^{(t)}\boldsymbol{u}^T + R_2^{(t)T},$$

where $\forall i \in [d] : u_i = \text{sign}(w_{2i}^{(0)}) \in \{\pm 1\}$ and

$$\forall 1 \leq i, j \leq d : \quad \frac{\left|R_1^{(t)}[i,j]\right|}{\left|u_i v_j^{(t)}\right|} \leq \delta := \tilde{\mathcal{O}}\left(\eta^{\frac{1}{4}} + \frac{1}{d^{\frac{\alpha}{2}-\frac{1}{4}}}\right), \quad \frac{\left|R_{2i}^{(t)}\right|}{\left|c^{(t)} u_i\right|} \leq \delta.$$

Hence by Lemma 28, when $t \in [T_{\text{Adam},1}, T_{\text{Adam},2}]$, we have for $1 \leq i \leq d$,

$$\begin{aligned}
\frac{\sqrt{\sum_{j\neq i}\left(H^{(t)}[i,j]\right)^2}}{\left|H^{(t)}[i,i]\right|} &\leq \frac{1+\delta}{1-\delta}\left(1 + \frac{\left|c^{(t)}\right|}{\left\|\boldsymbol{v}^{(t)}\right\|_2}\right)\sqrt{\frac{\sum_{j=1}^{d} u_j^2}{u_i^2}} + \frac{\left\|E^{(t)}\right\|_2}{(1-\delta)^2 u_i^2 \left\|\boldsymbol{v}^{(t)}\right\|_2^2} \\
&= \mathcal{O}\left(1 + \frac{\left|c^{(t)}\right|}{\left\|\boldsymbol{v}^{(t)}\right\|_2}\right)\sqrt{d} + \mathcal{O}\left(\frac{\left\|E^{(t)}\right\|_2}{u_i^2 \left\|\boldsymbol{v}^{(t)}\right\|_2^2}\right),
\end{aligned} \tag{50}$$

and for $1 \leq i \leq d, 1 \leq k \leq d$,

$$\begin{aligned}
\frac{\sqrt{\sum_{j\neq id+k}\left(H^{(t)}[i,j]\right)^2}}{\left|H^{(t)}[id+k,id+k]\right|} &\leq \frac{1+\delta}{1-\delta}\left(1 + \frac{\left|v_k^{(t)}\right|}{\left|c^{(t)}\right|}\right)\sqrt{\frac{\sum_{j=1}^{d} u_j^2}{u_i^2}} + \frac{\left|E_k^{(t)}\right|}{(1-\delta)^2 \left(c^{(t)}\right)^2 u_i^2} \\
&= \mathcal{O}\left(1 + \frac{\left|v_k^{(t)}\right|}{\left|c^{(t)}\right|}\right)\sqrt{d} + \mathcal{O}\left(\frac{\left|E_k^{(t)}\right|}{\left(c^{(t)}\right)^2 u_i^2}\right).
\end{aligned} \tag{51}$$

Recall the following facts of Adam.

(A) By Lemma 14, we know that for $t \in [T_{\text{Adam},1}, T_1]$ (where T_1 is defined in Definition 4), w.h.p. $\forall k \in [d] : v_k^{(t)} = c^{(t)} = \eta(t - t_{\text{inc}})$. Specially, when $t = T_{\text{Adam},1}, \forall k \in [d] :$ $v_k^{(t)} = c^{(t)} = \frac{1}{d^{\frac{\alpha}{2}}}$. Lemma 24 and 26 tell us that for $t \in [T_1, T_{\text{Adam},2}]$ w.h.p. $\forall i, j \in [d] :$ $|W_1[i,j]| = \Theta\left(\frac{1}{\sqrt{d}}\right), |w_{2i}| = \Theta\left(\frac{1}{\sqrt{d}}\right)$, which gives us $\forall k \in [d] : \left|v_k^{(t)}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right)$ and $\left|c^{(t)}\right| = \Theta\left(\frac{1}{\sqrt{d}}\right)$. That means when $t \in [T_{\text{Adam},1}, T_{\text{Adam},2}], \forall k \in [d] : \left|v_k^{(t)}\right|$ and $\left|c^{(t)}\right|$ increase from $\frac{1}{d^{\frac{\alpha}{2}}}$ to $\Theta(\frac{1}{\sqrt{d}})$ and $\frac{\left|v_k^{(t)}\right|}{\left|c^{(t)}\right|} = \Theta(1), \frac{\left|c^{(t)}\right|}{\left\|v^{(t)}\right\|_2} = \Theta\left(\frac{1}{\sqrt{d}}\right)$.

(B) Lemma 14 and 25 tell us that w.h.p. $\left\|E^{(t)}\right\|_2$ (resp. $\forall k \in [d], \left|E_k^{(t)}\right|$) decreases from $\Theta(d)$ (resp. $\Theta(1)$) when $t = T_{\text{Adam},1}$ to $\mathcal{O}\left(d^2\sqrt{\eta}\right)$ (resp. $\mathcal{O}\left(d\sqrt{\eta d}\right)$) when $t = T_{\text{Adam},2}$.

Combining (A) and (B), we get that the trend of $\dfrac{\left\|E^{(t)}\right\|_2}{u_i^2\left\|\boldsymbol{v}^{(t)}\right\|_2^2}$ and $\dfrac{\left|E_k^{(t)}\right|}{\left(c^{(t)}\right)^2 u_i^2}$ is to decrease over time, and when $t = T_{\text{Adam},2}$, we have w.h.p.

$$\frac{\left\|E^{(t)}\right\|_2}{u_i^2\left\|\boldsymbol{v}^{(t)}\right\|_2^2} \leq \mathcal{O}\left(d^2\sqrt{\eta}\right), \quad \frac{\left|E_k^{(t)}\right|}{\left(c^{(t)}\right)^2 u_i^2} \leq \mathcal{O}\left(d^2\sqrt{\eta d}\right). \tag{52}$$

Substituting (A) and eq. (52) into eq. (50) and (51) gives us w.h.p.,

$$\forall 1 \leq i \leq d : \frac{\sqrt{\sum_{j \neq i}\left(H^{(t)}[i,j]\right)^2}}{\left|H^{(t)}[i,i]\right|} \leq \mathcal{O}\left(\sqrt{d}\right) + r_{1i}^{(t)},$$

where the trend of $r_{1i}^{(t)}$ is to decrease over time and $r_{1i}^{(T_{\text{Adam},2})} \leq \mathcal{O}\left(d^2\sqrt{\eta}\right)$.

We also have

$$\forall 1 \leq i \leq d, 1 \leq k \leq d : \frac{\sqrt{\sum_{j \neq id+k}\left(H^{(t)}[i,j]\right)^2}}{\left|H^{(t)}[id+k, id+k]\right|} \leq \mathcal{O}\left(\sqrt{d}\right) + r_{2i}^{(t)},$$

where the trend of $r_{2i}^{(t)}$ is to decrease over time and $r_{2i}^{(T_{\text{Adam},2})} \leq \mathcal{O}\left(d^2\sqrt{\eta d}\right)$.

Hence $R_{\text{diag}}^{\text{SGDM}}(t) = \mathcal{O}\left(\sqrt{d}\right) + \frac{1}{d^2+d}\sum_{i=1}^d r_{1i}^{(t)} + \frac{d}{d^2+d}\sum_{i=1}^d r_{2i}^{(t)} := \mathcal{O}\left(\sqrt{d}\right) + r^{(t)}$ where the trend of $r^{(t)}$ is to decrease over time and

$$r^{(T_{\text{Adam},2})} \leq \frac{1}{d^2+d}\sum_{i=1}^d \mathcal{O}\left(d^2\sqrt{\eta}\right) + \frac{d}{d^2+d}\sum_{i=1}^d \mathcal{O}\left(d^2\sqrt{\eta d}\right) \leq \mathcal{O}\left(d^2\sqrt{\eta d}\right).$$

For any $p > 0$, by picking the same hyperparameters as in Theorem 1, we have $\eta d^4 \leq \tilde{\mathcal{O}}\left(\frac{1}{d^p}\right)$ and hence $r^{(T_{\text{Adam},2})} \leq \tilde{\mathcal{O}}\left(\frac{1}{d^{\frac{p-1}{2}}}\right) = \tilde{o}\left(\sqrt{d}\right)$.

E.4   PROOF OF LEMMA 28

By the assumed weight structure, we get that

$$\forall i \in [d] : (1-\delta)^2(cu_i)^2 \leq (w_{2i})^2 \leq (1+\delta)^2(cu_i)^2,$$
$$(1-\delta)^2(u_i)^2\|\boldsymbol{v}\|_2^2 \leq \|W_1[i,:]\|_2^2 \leq (1+\delta)^2(u_i)^2\|\boldsymbol{v}\|_2^2.$$

For the i-th row where $1 \leq i \leq d$, i.e. the i-th row of the submatrix $[H_{22} \quad H_{21}^T]$, by triangle inequality, we have

$$\sqrt{\sum_{j \neq i} H^2[i,j]} \leq \sqrt{\sum_{j \neq i} H_{22}^2[i,j]} + \sqrt{\sum_{j=1}^{d^2} H_{21}^2[j,i]}$$

$$\leq \sqrt{\sum_{j \neq i}\langle W_1[i,:], W_1[j,:]\rangle^2} + \sqrt{\sum_{j=1}^d w_{2j}^2 \sum_{k=1}^d W_1^2[i,k]} + \|E\|_2$$

$$\leq \|W_1[i,:]\|_2\left(\sqrt{\sum_{j \neq i}\|W_1[j,:]\|_2^2} + \sqrt{\sum_{j=1}^d w_{2j}^2}\right) + \|E\|_2.$$

Then we have that for $1 \leq i \leq d$,

$$
\begin{aligned}
\frac{\sqrt{\sum_{j \neq i} H^2[i,j]}}{|H[i,i]|} &\leq \frac{\|W_1[i,:]\|_2 \sqrt{\sum_{j \neq i} \|W_1[j,:]\|_2^2} + \sqrt{\sum_{j=1}^d w_{2j}^2}}{\|W_1[i,:]\|_2^2} + \frac{\|E\|_2}{\|W_1[i,:]\|_2^2} \\
&= \sqrt{\frac{\sum_{j \neq i} \|W_1[j,:]\|_2^2}{\|W_1[i,:]\|_2^2}} + \sqrt{\frac{\sum_{j=1}^d w_{2j}^2}{\|W_1[i,:]\|_2^2}} + \frac{\|E\|_2}{\|W_1[i,:]\|_2^2} \\
&\leq \sqrt{\frac{(1+\delta)^2}{(1-\delta)^2} \cdot \frac{\sum_{j \neq i} u_j^2 \|\boldsymbol{v}\|_2^2}{u_i^2 \|\boldsymbol{v}\|_2^2}} + \sqrt{\frac{(1+\delta)^2}{(1-\delta)^2} \cdot \frac{c^2 \sum_{j=1}^d u_j^2}{u_i^2 \|\boldsymbol{v}\|_2^2}} + \frac{\|E\|_2}{(1-\delta)^2 u_i^2 \|v\|_2^2} \\
&= \frac{1+\delta}{1-\delta}\left(1 + \frac{|c|}{\|\boldsymbol{v}\|_2}\right) \sqrt{\frac{\sum_{j=1}^d u_j^2}{u_i^2}} + \frac{\|E\|_2}{(1-\delta)^2 u_i^2 \|v\|_2^2}.
\end{aligned}
$$

For the $(id+k)$-th row where $1 \leq i \leq d, 1 \leq k \leq d$, i.e. the $((i-1)d+k)$-th row of the submatrix $[H_{21} \quad H_{11}]$, by triangle inequality again, we have

$$
\begin{aligned}
\sqrt{\sum_{j \neq id+k} H^2[i,j]} &\leq \sqrt{\sum_{j \neq (i-1)d+k} H_{11}^2[i,j]} + \sqrt{\sum_{j=1}^d H_{21}^2[(i-1)d+k,j]} \\
&\leq \sqrt{\sum_{j \neq i} w_{2i}^2 w_{2j}^2} + \sqrt{\sum_{j=1}^d w_{2i}^2 W_1^2[j,k]} + |E_k| \\
&= |w_{2i}|\left(\sqrt{\sum_{j \neq i} w_{2j}^2} + \sqrt{\sum_{j=1}^d W_1^2[j,k]}\right) + |E_k|.
\end{aligned}
$$

Then we have that for $1 \leq i \leq d, 1 \leq k \leq d$,

$$
\begin{aligned}
\frac{\sqrt{\sum_{j \neq id+k} H^2[i,j]}}{|H[id+k, id+k]|} &\leq \frac{|w_{2i}|\sqrt{\sum_{j \neq i} w_{2j}^2} + \sqrt{\sum_{j=1}^d W_1^2[j,k]}}{w_{2i}^2} + \frac{|E_k|}{w_{2i}^2} \\
&= \sqrt{\frac{\sum_{j \neq i} w_{2j}^2}{w_{2i}^2}} + \sqrt{\frac{\sum_{j=1}^d W_1^2[j,k]}{w_{2i}^2}} + \frac{|E_k|}{w_{2i}^2} \\
&\leq \sqrt{\frac{(1+\delta)^2}{(1-\delta)^2} \cdot \frac{\sum_{j \neq i} c^2 u_j^2}{c^2 u_i^2}} + \sqrt{\frac{(1+\delta)^2}{(1-\delta)^2} \cdot \frac{v_k^2 \sum_{j=1}^d u_j^2}{c^2 u_i^2}} + \frac{|E_k|}{(1-\delta)^2 c^2 u_i^2} \\
&= \frac{1+\delta}{1-\delta}\left(1 + \frac{|v_k|}{|c|}\right) \sqrt{\frac{\sum_{j=1}^d u_j^2}{u_i^2}} + \frac{|E_k|}{(1-\delta)^2 c^2 u_i^2}.
\end{aligned}
$$

F CONNECTION BETWEEN DIAGONAL OF LOSS HESSIAN AND WEIGHTS

The partial derivative at W_i of the cost function for each i is given by:

$$
\nabla_{W_i} L(W) = W_{i+1}^T \ldots W_{H+1}^T (W_{H+1} W_H \ldots W_1 - A) W_1^T \ldots W_{i-1}^T \tag{53}
$$

In our experiments, we were interested in the diagonal elements of the hessian. These are given by:

$$
\nabla_{(W_i)_{a,b}} (\nabla_{W_i} L(W))_{a,b} = \nabla_{(W_i)_{a,b}} (W_{i+1}^T \ldots W_{H+1}^T (W_{H+1} W_H \ldots W_1 - A) W_1^T \ldots W_{i-1}^T)_{a,b}
$$

for each possible i, a, b. For ease in notation, define for each i, the quantities $M_i := W_{i+1}^T \ldots W_{H+1}^T$ and $N_i := W_1^T \ldots W_{i-1}^T$. Then we have the following lemma.

Lemma 29. *The diagonal elements of the hessian of the cost function are given by:*

$$
\nabla_{(W_i)_{a,b}} (\nabla_{W_i} L(W))_{a,b} = (M_i M_i^T)_{a,a} (N_i^T N_i)_{b,b}
$$

for each possible i, a, b.

Proof. We have:

$$
\begin{aligned}
\nabla_{W_i} L(W) &= W_{i+1}^T \ldots W_{H+1}^T (W_{H+1} W_H \ldots W_1 - A) W_1^T \ldots W_{i-1}^T \\
&= M_i (W_{H+1} W_H \ldots W_1 - A) N_i \\
&= M_i W_{H+1} W_H \ldots W_1 N_i - M_i Y N_i.
\end{aligned}
$$

This implies that:

$$
\begin{aligned}
\nabla_{(W_i)_{a,b}} (\nabla_{W_i} L(W))_{a,b} &= \nabla_{(W_i)_{a,b}} \left(M_i W_{H+1} W_H \ldots W_1 N_i - M_i Y N_i \right)_{a,b} \\
&= \nabla_{(W_i)_{a,b}} \left(M_i W_{H+1} W_H \ldots W_1 N_i \right)_{a,b},
\end{aligned}
$$

where the last step follows since M_i and N_i are not functions of W_i.

Note that $M_i := W_{i+1}^T \ldots W_{H+1}^T$ and $N_i := W_1^T \ldots W_{i-1}^T$. Now define $C_i := M_i W_{H+1} W_H \ldots W_{i+1} = M_i M_i^T$ and $D_i := W_{i-1} \ldots W_2 W_1 N_i = N_i^T N_i$ so that:

$$
\nabla_{(W_i)_{a,b}} (\nabla_{W_i} L(W))_{a,b} = \nabla_{(W_i)_{a,b}} (C_i W_i D_i)_{a,b},
$$

where C_i and D_i are not functions of W_i. Now, Equation 74 in the Matrix Cookbook[12] shows us that for any matrices A and X we have:

$$
\nabla_{X_{mn}} (XA)_{ij} = \delta_{im} A_{nj}.
$$

Note that $W_i \in \mathbb{R}^{d_i \times d_{i-1}}$, then we can apply this to obtain that:

$$
\begin{aligned}
\nabla_{(W_i)_{a,b}} (\nabla_{W_i} L(W))_{a,b} &= \nabla_{(W_i)_{a,b}} (C_i W_i D_i)_{a,b} \\
&= \nabla_{(W_i)_{a,b}} \left[\sum_{k=1}^{d_i} (C_i)_{a,k} (W_i D_i)_{k,b} \right] \\
&= \sum_{k=1}^{d_i} (C_i)_{a,k} \nabla_{(W_i)_{a,b}} (W_i D_i)_{k,b} \\
&= \sum_{k=1}^{d_i} (C_i)_{a,k} \delta_{ak} (D_i)_{b,b} \\
&= (C_i)_{a,a} (D_i)_{b,b} \\
&= (M_i M_i^T)_{a,a} (N_i^T N_i)_{b,b}.
\end{aligned}
$$

This completes the proof. □

For ease of notation, let's now drop the superscript OPT and (t) and write $R_{\text{med},1}^{\text{OPT}}(t)$ as $R_{\text{med},1}$ and $R_{\text{med},2}^{\text{OPT}}(t)$ as $R_{\text{med},2}$. For a 2-layer linear network, $H = 1$. Consider the Hessian w.r.t W_1, we have $M_1 M_1^T = W_2^T W_2$ and $N_1^T N_1$ is an identity matrix. Under Assumption 1, we know that W_2 is a row vector, which can be denoted as $W_2 = [w_{21}, w_{22}, ..., w_{2d_1}]$. Then we have

$$
(M_1 M_1^T)_{a,a} = w_{2a}^2, (N_1^T N_1)_{b,b} = 1, \quad \Rightarrow \quad R_{\text{med},1} = \frac{\max_i (w_{2i})^2}{\text{median}(w_{2i})^2}.
$$

Similarly, consider the Hessian w.r.t. W_2, we have that $M_1 M_1^T$ is an identity matrix and $N_1^T N_1 = W_1 W_1^T$. Therefore,

$$
(M_1 M_1^T)_{a,a} = 1, (N_1^T N_1)_{b,b} = \|W_1[b,:]\|_2^2, \quad \Rightarrow \quad R_{\text{med},2} = \frac{\max_i \|W_1[i,:]\|_2^2}{\text{median}\|W_1[i,:]\|_2^2}.
$$

Hence we have related the uniformity of diagonal Hessian to that of weight matrices. In the detailed analysis, for both GD and Adam, we can prove that W_1 converges to an approximately rank 1 matrix. The following lemma allows us to use this rank 1 structure to compute $R_{\text{med},1}$ and $R_{\text{med},2}$.

---

[12]https://www.math.uwaterloo.ca/~hwolkowi/matrixcookbook.pdf

Lemma 30. *Suppose $W_1 \in \mathbb{R}^{d \times d}$ and $W_2 \in \mathbb{R}^{1 \times d}$ have the following structure:*

$$W_1 = \boldsymbol{u}\boldsymbol{v}^T + R_1,$$
$$W_2 = c\boldsymbol{u}^T + R_2,$$

where $\boldsymbol{u} \in \mathbb{R}^d, \boldsymbol{v} \in \mathbb{R}^d, R_1 \in \mathbb{R}^{d \times d}, R_2 \in \mathbb{R}^{1 \times d}$ and that

$$\forall 1 \le i, j \le d: \quad \frac{|R_1[i,j]|}{|u_i v_j|} \le \delta, \quad \frac{|R_{2i}|}{|cu_i|} \le \delta, \quad \delta \in (0,1).$$

Then we have

$$R_{med,1}, R_{med,2} \in \left[\frac{(1-\delta)^2}{(1+\delta)^2} \cdot \frac{\max_i u_i^2}{median\ u_i^2}, \frac{(1+\delta)^2}{(1-\delta)^2} \cdot \frac{\max_i u_i^2}{median\ u_i^2}\right].$$

Proof. Let's first consider $R_{\text{med},1}$. we have

$$\forall i \in [d]: (1-\delta)^2 (cu_i)^2 \le w_{2i}^2 \le (1+\delta)^2 (cu_i)^2$$
$$\Rightarrow \quad (1-\delta)^2 \max_i (cu_i)^2 \le \max_i w_{2i}^2 \le (1+\delta)^2 \max_i (cu_i)^2$$
$$(1-\delta)^2 \text{median}\ (cu_i)^2 \le \text{median}\ w_{2i}^2 \le (1+\delta)^2 \text{median}\ (cu_i)^2,$$

which yields

$$\frac{(1-\delta)^2}{(1+\delta)^2} \cdot \frac{\max_i u_i^2}{\text{median}\ u_i^2} \le R_{\text{med},1} = \frac{\max_i w_{2i}^2}{\text{median}\ w_{2i}^2} \le \frac{(1+\delta)^2}{(1-\delta)^2} \cdot \frac{\max_i u_i^2}{\text{median}\ u_i^2}.$$

Similarly, for $R_{\text{med},2}$. We have that

$$\forall i, j \in [d]: (1-\delta)^2 (u_i v_j)^2 \le W_1^2[i,j] \le (1+\delta)^2 (u_i v_j)^2$$
$$\Rightarrow \quad (1-\delta)^2 u_i^2 \|\boldsymbol{v}\|_2^2 \le \|W_1[i,:]\|_2^2 \le (1+\delta)^2 u_i^2 \|\boldsymbol{v}\|_2^2$$
$$\Rightarrow \quad (1-\delta)^2 \max_i u_i^2 \|\boldsymbol{v}\|_2^2 \le \max_i \|W_1[i,:]\|_2^2 \le (1+\delta)^2 \max_i u_i^2 \|\boldsymbol{v}\|_2^2$$
$$(1-\delta)^2 \text{median}\ u_i^2 \|\boldsymbol{v}\|_2^2 \le \text{median}\ \|W_1[i,:]\|_2^2 \le (1+\delta)^2 \text{median}\ u_i^2 \|\boldsymbol{v}\|_2^2,$$

which yields

$$\frac{(1-\delta)^2}{(1+\delta)^2} \cdot \frac{\max_i u_i^2 \|\boldsymbol{v}\|_2^2}{\text{median}\ u_i^2 \|\boldsymbol{v}\|_2^2} \le R_{\text{med},2} = \frac{\max_i \|W_1[i,:]\|_2^2}{\text{median}\|W_1[i,:]\|_2^2} \le \frac{(1+\delta)^2}{(1-\delta)^2} \cdot \frac{\max_i u_i^2 \|\boldsymbol{v}\|_2^2}{\text{median}\ u_i^2 \|\boldsymbol{v}\|_2^2}.$$

That means

$$\frac{(1-\delta)^2}{(1+\delta)^2} \cdot \frac{\max_i u_i^2}{\text{median}\ u_i^2} \le R_{\text{med},2} \le \frac{(1+\delta)^2}{(1-\delta)^2} \cdot \frac{\max_i u_i^2}{\text{median}\ u_i^2}.$$

$\square$

G AUXILIARY LEMMAS

Lemma 31. *Let $A = \frac{1}{m} Y X^T, \Lambda_{xx} := \frac{1}{m} X X^T, g_k^{(t)} = \nabla_{W_k} L(W^{(t)}), k = 1, 2$. Denote $\tilde{A}^{(t)}, \tilde{\Lambda}_{xx}^{(t)}$ and $\tilde{g}_k^{(t)}, k = 1, 2$ as the corresponding batch versions at time t. Let $M_1^{(t)} = \max_{i,j} \left|W_1^{(t)}[i,j]\right|$ and $M_2^{(t)} = \max_i \left|w_{2i}^{(t)}\right|$. Under Assumption 3, we have with probability at least $1 - \frac{1}{d}$, for $\forall t \le T$ and $\forall i, j \in [d]$,*

$$\left|\tilde{g}_1^{(t)}[i,j] - g_1^{(t)}[i,j]\right| \le d^3 M_1^{(t)} \left(M_2^{(t)}\right)^2 \sigma\sqrt{dT} + M_2^{(t)} \sigma\sqrt{d^2 T}$$

$$\left|g_{2i}^{(t)} - g_{2i}^{(t)}\right| \le d^4 \left(M_1^{(t)}\right)^2 M_2^{(t)} \sigma\sqrt{dT} + d M_1^{(t)} \sigma\sqrt{d^2 T}$$

Proof. By Assumption 3 and Chebyshev's inequality, we have for fixed $i, j \in [d]$ and $t \leq T$,

$$\mathbb{P}\left(\left|\tilde{A}_i^{(t)} - A_i\right| > \lambda\right) \leq \frac{\sigma^2}{\lambda^2}, \quad \mathbb{P}\left(\left|\tilde{\Lambda}_{xx}^{(t)}[i,j] - \Lambda_{xx}[i,j]\right| > \lambda\right) \leq \frac{\sigma^2}{\lambda^2}$$

Applying the union bound gives us

$$\mathbb{P}\left(\exists i \in [d], \exists t \leq T : \quad \left|\tilde{A}_i^{(t)} - A_i\right| > \lambda\right) \leq \frac{Td\sigma^2}{\lambda^2},$$

$$\mathbb{P}\left(\exists i, j \in [d], \exists t \leq T : \quad \left|\tilde{\Lambda}_{xx}^{(t)}[i,j] - \Lambda_{xx}[i,j]\right| > \lambda\right) \leq \frac{Td^2\sigma^2}{\lambda^2}$$

which gives us with probability at least $1 - \frac{1}{d}$, for $\forall t \leq T, \forall i, j \in [d]$,

$$\left|\tilde{A}_i^{(t)} - A_i\right| \leq \sigma\sqrt{d^2 T}, \quad \left|\tilde{\Lambda}_{xx}^{(t)}[i,j] - \Lambda_{xx}[i,j]\right| \leq \sigma d\sqrt{dT}$$

Now we are ready to bound $\tilde{g}_k^{(t)} - g_k^{(t)}$ for $k = 1, 2$ and $t \leq T$.

Note that for all $t \leq T$ and $\forall i \in [d]$,

$$\left|\left(W_2^{(t)} W_1^{(t)}\right)_i\right| = \left|\sum_{j=1}^{d} w_{2j}^{(t)} W_1^{(t)}[j,i]\right| \leq \sum_{j=1}^{d} \left|w_{2j}^{(t)}\right| \left|W_1^{(t)}[j,i]\right| \leq dM_1^{(t)} M_2^{(t)}$$

Then we have with probability at least $1 - \frac{1}{d}$, for all $t \leq T$ and $\forall i \in [d]$,

$$\left|\left(W_2^{(t)} W_1^{(t)} \left(\tilde{\Lambda}_{xx}^{(t)} - \Lambda_{xx}\right)\right)_i\right| \leq \sum_{j=1}^{d} \left|\left(W_2^{(t)} W_1^{(t)}\right)_j\right| \left|\tilde{\Lambda}_{xx}^{(t)}[j,i] - \Lambda_{xx}[j,i]\right| \leq d^3 M_1^{(t)} M_2^{(t)} \sigma\sqrt{dT}$$

Combining with $\tilde{g}_1^{(t)} - g_1^{(t)} = W_2^{(t)T}\left(W_2^{(t)} W_1^{(t)} \left(\tilde{\Lambda}_{xx}^{(t)} - \Lambda_{xx}\right) - \left(\tilde{A}^{(t)} - A\right)\right)$, we get that with probability at least $1 - \frac{1}{d}$, for all $t \leq T$ and $\forall i, j \in [d]$,

$$\left|\tilde{g}_1^{(t)}[i,j] - g_1^{(t)}[i,j]\right| \leq \left|w_{2i}^{(t)}\right| \left|\left(W_2^{(t)} W_1^{(t)} \left(\tilde{\Lambda}_{xx}^{(t)} - \Lambda_{xx}\right)\right)_j\right| + \left|w_{2i}^{(t)}\right| \left|\tilde{A}_j^{(t)} - A_j\right|$$

$$\leq d^3 M_1^{(t)} \left(M_2^{(t)}\right)^2 \sigma\sqrt{dT} + M_2^{(t)} \sigma\sqrt{d^2 T}$$

Similarly, note that $\tilde{g}_{2i}^{(t)} - g_{2i}^{(t)} = \left(W_2^{(t)} W_1^{(t)} \left(\tilde{\Lambda}_{xx}^{(t)} - \Lambda_{xx}\right) - \left(\tilde{A}^{(t)} - A\right)\right) W_1^{(t)T}$, we then have that with probability at least $1 - \frac{1}{d}$, for all $t \leq T$ and $\forall i, j \in [d]$,

$$\left|\tilde{g}_{2i}^{(t)} - g_{2i}^{(t)}\right| \leq \sum_{j=1}^{d} \left|\left(W_2^{(t)} W_1^{(t)} \left(\tilde{\Lambda}_{xx}^{(t)} - \Lambda_{xx}\right)\right)_j\right| \left|W_1^{(t)}[i,j]\right| + \sum_{j=1}^{d} \left|\tilde{A}_j^{(t)} - A_j\right| \left|W_1^{(t)}[i,j]\right|$$

$$\leq d^4 \left(M_1^{(t)}\right)^2 M_2^{(t)} \sigma\sqrt{dT} + dM_1^{(t)} \sigma\sqrt{d^2 T}$$

$\square$

Lemma 32. *Consider two sequences $\{a^{(t)}\}_{t \geq 0}, \{b^{(t)}\}_{t \geq 0}$, which satisfy*

$$a^{(t)} = (1 - \beta) \sum_{\tau=0}^{t} \beta^\tau b^{(t-\tau)}, \beta \in (0, 1).$$

Suppose $\forall \tau \leq t : \left|b^{(t)}\right| \leq B$, then for any $\epsilon > 0$, the following truncated version

$$\tilde{a}^{(t)} = (1 - \beta) \sum_{\tau=0}^{H} \beta^\tau b^{(t-\tau)}$$

with $H \geq \frac{1}{1-\beta} \log \frac{B}{\epsilon} = \tilde{\Omega}\left(\frac{1}{1-\beta}\right)$ satisfies

$$\left|a^{(t)} - \tilde{a}^{(t)}\right| \leq \epsilon.$$

Proof. We have that

$$\left| a^{(t)} - \tilde{a}^{(t)} \right| \leq \left| (1-\beta) \sum_{\tau=H+1}^{t} \beta^\tau b^{(t-\tau)} \right| \leq (1-\beta) \sum_{\tau=H+1}^{t} \beta^\tau B \leq B\beta^{H+1}.$$

To make it less than ϵ, it suffices to choose $H \geq \log(\frac{\epsilon}{B})/\log \beta$.

Since $\beta \in (0,1)$, we know that $\log \beta \leq \beta - 1 < 0$. We also have $\log \frac{\epsilon}{B} < 0$. Then it suffices to choose

$$H \geq \frac{\log(\epsilon/B)}{\beta - 1} \geq \frac{\log(\epsilon/B)}{\log \beta} \quad \Rightarrow \quad H \geq \frac{1}{1-\beta} \log \frac{B}{\epsilon} = \tilde{\Omega}\left(\frac{1}{1-\beta}\right).$$

$\square$

Lemma 33. *Suppose* $a, b, c, e_a, e_b, e_c \in \mathbb{R}, b > 0, c > 0$ *satisfy* $b + e_b + e_c > 0, |e_a| \leq \delta|a|, |e_b| \leq \delta b, |e_c| \leq \delta^2 c^2$ *with* $0 < \delta \ll 1$, *then we have*

$$\frac{a + e_a}{\sqrt{b + e_b + e_c} + c} = \frac{a}{\sqrt{b} + c}(1 + R), \quad \text{where } |R| = \mathcal{O}(\delta).$$

Proof. We have

$$\frac{a + e_a}{\sqrt{b + e_b + e_c} + c} = \frac{a}{\sqrt{b} + c} + \frac{a}{\sqrt{b + e_b + e_c} + c} - \frac{a}{\sqrt{b} + c} + \frac{e_a}{\sqrt{b + e_b + e_c} + c}$$

$$= \frac{a}{\sqrt{b} + c}\left(1 + \underbrace{\frac{\sqrt{b} + c}{\sqrt{b + e_b + e_c} + c} - 1}_{q_1} + \underbrace{\frac{e_a}{a} \cdot \frac{\sqrt{b} + c}{\sqrt{b + e_b + e_c} + c}}_{q_2}\right).$$

Define $R := q_1 + q_2$. The term $|q_1|$ can be bounded by

$$|q_1| = \frac{\left| \sqrt{b} - \sqrt{b + e_b + e_c} \right|}{\sqrt{b + e_b + e_c} + c}$$

$$= \frac{|e_b + e_c|}{(\sqrt{b + e_b + e_c} + c)\left(\sqrt{b} + \sqrt{b + e_b + e_c}\right)}$$

$$= \frac{|e_b|}{(\sqrt{b + e_b + e_c} + c)\left(\sqrt{b} + \sqrt{b + e_b + e_c}\right)} + \frac{|e_c|}{(\sqrt{b + e_b + e_c} + c)\left(\sqrt{b} + \sqrt{b + e_b + e_c}\right)}$$

$$\leq \frac{|e_b|}{(\sqrt{b + e_b + e_c} + c)\sqrt{b}} + \frac{\sqrt{|e_c|}}{c} \cdot \frac{\sqrt{|e_c|}}{\sqrt{b} + \sqrt{b + e_b + e_c}}$$

$$\leq \underbrace{\frac{|e_b|}{(\sqrt{b + e_b + e_c} + c)\sqrt{b}}}_{q_3} + \delta \underbrace{\frac{\sqrt{|e_c|}}{\sqrt{b} + \sqrt{b + e_b + e_c}}}_{q_4},$$

where $|q_3|$ can be bounded by

$$|q_3| \overset{(i)}{\leq} \frac{\delta b}{(\sqrt{b + e_b} - \sqrt{|e_c|} + c)\sqrt{b}} \leq \frac{\delta \sqrt{b}}{\sqrt{b(1-\delta)} + c(1-\delta)} \leq \frac{\delta \sqrt{b}}{\sqrt{b(1-\delta)}} = \mathcal{O}(\delta).$$

Here the denominator of (i) uses $b + e_b \geq b(1-\delta) > 0$ and $\sqrt{x+y} \geq \sqrt{x} - \sqrt{|y|}$ when $x \geq 0, x + y \geq 0$.

Now let's bound $|q_4|$. If $e_c > 0$, we have $e_c = |e_c|$ and $|q_4| \leq \frac{\sqrt{e_c}}{\sqrt{e_c}} = 1$ since $b + e_b \geq b(1-\delta) > 0$.

If $e_c \leq 0$, note that $b + e_b + e_c > 0$, we have $|e_c| < b + b_e \leq b(1 + \delta)$, which yields $|q_4| \leq \frac{\sqrt{|e_c|}}{\sqrt{b}} = \mathcal{O}(1)$. Combining the above bounds give us $|q_1| \leq |q_3| + \delta|q_4| = \mathcal{O}(\delta)$.

On the other hand, $|q_2|$ can be bounded by

$$|q_2| \leq \delta \frac{\sqrt{b}+c}{\sqrt{b+e_b}-\sqrt{|e_c|}+c} \leq \delta \frac{\sqrt{b}+c}{\sqrt{b(1-\delta)}+c(1-\delta)} = \mathcal{O}(\delta).$$

Then $|R| \leq |q_1| + |q_2| = \mathcal{O}(\delta)$ □

Lemma 34. *Suppose $X_1, X_2, ..., X_d$ are i.i.d Gaussian with mean 0 and variance σ^2, then for $0 < \delta < \frac{1}{e}$, we have with probability at least $1 - \delta$,*

$$\max_{1 \leq i \leq d} X_i^2 \geq \sigma^2 \left(C_1 \log d - C_2 \log \log \frac{1}{\delta} \right)$$

for some $C_1, C_2 > 0$.

Proof. It suffices to assume that $\sigma^2 = 1$ and prove that w.p. at least $1 - \delta$, $\max_{1 \leq i \leq d} X_i^2 \geq C_1 \log d - C_2 \log \log \frac{1}{\delta}$.

First, by the lower bound of Gaussian tail, there exists $\alpha, \beta > 0$ such that $\mathbb{P}(|X_i| > x) = 2\mathbb{P}(X_i > x) \geq \alpha e^{-\beta x^2}$ for $x \geq 0$. Then by i.i.d., we have

$$\mathbb{P}(\max_i |X_i| \leq x) = \mathbb{P}\left(\bigcap_{i=1}^{d} \{|X_i| \leq x\} \right)$$

$$= \prod_{i=1}^{d} \mathbb{P}(|X_i| \leq x) = (1 - \mathbb{P}(|X_i| > x))^d$$

$$\leq (1 - \alpha e^{-\beta x^2})^d$$

$$\leq \exp(-d\alpha e^{-\beta x^2}),$$

where the last inequality uses $x \in \mathbb{R}, 1 - x \leq e^{-x}$. Let $\exp(-d\alpha e^{-\beta x^2}) = \delta$, we get that w.p. at least $1 - \delta$,

$$\max_{1 \leq i \leq d} |X_i| \geq \sqrt{\frac{1}{\beta} \left(\log(\alpha d) - \log \log \frac{1}{\delta} \right)}.$$

Then we have w.p. at least $1 - \delta$,

$$\max_{1 \leq i \leq d} X_i^2 = \left(\max_{1 \leq i \leq d} |X_i| \right)^2 \geq \frac{1}{\beta} \left(\log(\alpha d) - \log \log \frac{1}{\delta} \right).$$

□

Lemma 35. *Suppose $X_1, X_2, ..., X_d$ are i.i.d Gaussian with mean 0 and variance σ^2, then we have with constant probability,*

$$\frac{1}{d} \sum_{i=1}^{d} \frac{1}{|X_i|} \leq \mathcal{O}\left(\frac{1}{\sigma} \log d \right).$$

Proof. It suffices to assume that $\sigma^2 = 1$ and prove that w.h.p., $\frac{1}{d} \sum_{i=1}^{d} \frac{1}{|X_i|} \leq \mathcal{O}(\log d)$.

Consider X_i for some fixed i. Since $X_i \sim \mathcal{N}(0, 1)$, we have $\mathbb{P}(|X_i| \leq t) \leq \frac{2t}{\sqrt{2\pi}}$. Then we know that with probability at least $1 - \Theta(d^{-1})$, $|X_i| \geq \frac{C}{d}$ for some $C > 0$. Then by union bound, with constant probability, $\forall i \in [d] : |X_i| \geq \frac{C}{d}$.

Now we split the interval $[\frac{C}{d}, 1]$ into several subintervals $\mathcal{I}_k = \{i : |X_i| \in [2^{-k-1}, 2^{-k}]\}$ for $k = 0, 1, ..., \lceil \log_2 \frac{d}{C} \rceil - 1$. Let $p_k = \mathbb{P}(|X_i| \in [2^{-k-1}, 2^{-k}])$, we know that $|\mathcal{I}_k| \sim \text{Binomial}(d, p_k)$ and $p_k \leq C_1 \cdot 2^{-k-1}$. Then by the concentration of binomial variables, we have w.p. at least $1 - d^{-p}$

for $p > 0$, $|\mathcal{I}_k| = \mathcal{O}\left(dp_k + \sqrt{dp_k \log d} + \log d\right) = \mathcal{O}\left(d \cdot 2^{-k-1} + \sqrt{d \cdot 2^{-k-1} \log d} + \log d\right)$.
Then we have

$$\sum_{i \in \mathcal{I}_k} \frac{1}{|X_i|} \leq |\mathcal{I}_k| 2^{k+1} = \mathcal{O}\left(d + \sqrt{d \cdot 2^{k+1} \log d} + 2^{k+1} \log d\right), \quad k = 0, 1, ..., \lceil \log_2 \frac{d}{C} \rceil - 1.$$

Therefore, with constant probability,

$$
\begin{aligned}
\sum_{i=1}^{d} \frac{1}{|X_i|} &= \sum_{k=0}^{\lceil \log_2 \frac{d}{C} \rceil - 1} \sum_{i \in \mathcal{I}_k} \frac{1}{|X_i|} + \sum_{|X_i| > 1} \frac{1}{|X_i|} \\
&\leq \sum_{k=0}^{\lceil \log_2 \frac{d}{C} \rceil - 1} \mathcal{O}\left(d + \sqrt{d \cdot 2^{k+1} \log d} + 2^{k+1} \log d\right) + d \\
&= \mathcal{O}\left(d \log_2 \frac{d}{C}\right) + \mathcal{O}\left(\sqrt{d \log d} \cdot (\sqrt{2})^{\lceil \log_2 \frac{d}{C} \rceil + C_2}\right) + 2^{\lceil \log_2 \frac{d}{C} \rceil + 1} \log d + d \\
&= \mathcal{O}\left(d \log d\right),
\end{aligned}
$$

which means with constant probability, $\frac{1}{d} \sum_{i=1}^{d} \frac{1}{|X_i|} = \mathcal{O}\left(\log d\right)$. $\qquad\square$