Differentiable Causal Discovery Under Latent Interventions

Abstract

Recent work has shown promising results in causal discovery by leveraging interventional data with gradient-based methods, even when the intervened variables are unknown. However, previous work assumes that the correspondence between samples and interventions is known, which is often unrealistic. We envision a scenario with an extensive dataset sampled from multiple intervention distributions and one observation distribution, but where we do not know which distribution originated each sample and how the intervention affected the system, *i.e.*, interventions are entirely latent. We propose a method based on neural networks and variational inference that addresses this scenario by framing it as learning a shared causal graph among a infinite mixture (under a Dirichlet process prior) of intervention structural causal models . Experiments with synthetic and real data show that our approach and its semi-supervised variant are able to discover causal relations in this challenging scenario.

Keywords: causal discovery, latent interventions, variational inference, Dirichlet process.

1. Introduction

Discovering causal relations among variables has countless applications in many scientific fields (Peters et al., 2017). However, causal graphs are hard to learn from data; only with strong assumptions can we ensure that we will learn the correct causal structures from observations alone (even with an unlimited supply of data). Recent work by Brouillard et al. (2020) has shown very promising results in causal discovery by leveraging interventional data with gradient-based methods, even when the intervened variables (targets) are not known. However, that work assumes that the *correspondence* between samples and interventions is known beforehand—often an unrealistic assumption.

Our work addresses this limitation/drawback by proposing a method for causal discovery under *fully latent* interventions, through a neural-based variational approach which infers the correspondences between samples and interventions from data. Our framework falls into the class of continuous constrained optimization methods for finding the DAG structure; other methods include constraint-based methods (Spirtes et al., 1993; Pearl, 2000; Spirtes et al., 2013). and score methods (Chickering and Heckerman, 1997; Bouckaert, 1993; Heckerman et al., 1994; Kuipers et al., 2021). We assume (as Brouillard et al. (2020) do) that the data is clustered into intervention groups, but we relax their assumption that the correspondence between intervention groups and samples is known, opening the door to more realistic scenarios. We model these *latent* interventions as being generated by a Dirichlet process prior and formulate the problem as one of end-to-end maximization of an evidence lower bound (ELBO). We summarize our contributions as follows:

- We formulate causal discovery under latent interventions as searching for the shared causal graph among an infinite mixture of intervened structural causal models.

- We propose a variational formulation with a Dirichlet process prior to model this infinite mixture. We use latent intervention embeddings with shared parameters to enable modeling an unlimited number of interventions.

- We develop a semi-supervised variant of our method; for when we know the correspondence for a subset of the samples.

Experiments on synthetic and real-world data show that our method is able to discover causal structures, outperforming several baselines and reaching similar performance to correspondence-aware methods.

2. Background

2.1. Structural causal models

We assume a structured causal model (SCM) $\mathcal{M} := (S, p(\mathcal{E}))$ over d *endogenous* variables $X = \{x_1, \ldots, x_d\}$, associated to d independent *exogenous* variables $\mathcal{E} = \{\varepsilon_1, \ldots, \varepsilon_d\}$, functions $\mathcal{F} = \{\zeta_1, \ldots, \zeta_d\}$, and a collection of d assignments,

$$x_j := \zeta_j(\mathbf{PA}_j^{\mathcal{G}}, \varepsilon_j), \qquad j = 1, \ldots, d, \tag{1}$$

where $\mathbf{PA}_j^{\mathcal{G}} \subseteq X \setminus \{x_j\}$ is the set of *parents* of x_j according to a *directed acyclic graph* (DAG) $\mathcal{G}$, *i.e.*, $x_i \in \mathbf{PA}_j^{\mathcal{G}}$ if and only if there is a directed edge $x_i \to x_j$ in $\mathcal{G}$. An SCM $\mathcal{M}$ defines a unique joint distribution for X, usually referred to as the *entailed* distribution $p_{\mathcal{M}}(X)$, which can be factorized as

$$p_{\mathcal{M}}(X) = \prod_{j=1}^{d} p_{\mathcal{M}}(x_j | \mathbf{PA}_j^{\mathcal{G}}), \tag{2}$$

where $p_{\mathcal{M}}(x_j | \mathbf{PA}_j^{\mathcal{G}})$ is the conditional distribution of x_j, given its parents.

2.2. Interventions

Given an SCM $\mathcal{M}$, we obtain an *interventioned* SCM $\tilde{\mathcal{M}}$ by replacing one (or more) of the original assignments. Let I be the set of variables targeted by the intervention; if $I = \emptyset$, $\tilde{\mathcal{M}} = \mathcal{M}$. For each variable $j \in I$, the intervention consists in one or multiple of the following actions: replacing the assignment function ζ_j by $\tilde{\zeta}_j$; replacing the parents $\mathbf{PA}_j^{\mathcal{G}}$ by a subset $\tilde{\mathbf{PA}}_j^{\mathcal{G}}$; changing the noise variable from ϵ_j to $\tilde{\epsilon}_j$. The SCM $\tilde{\mathcal{M}}$ generally has a different entailed distribution, called the *intervention distribution*:

$$p_{\tilde{\mathcal{M}}}(X) = \prod_{j \notin I} p_{\mathcal{M}}(x_j | \mathbf{PA}_j^{\mathcal{G}}) \prod_{j \in I} p_{\mathcal{M}; \mathrm{do}\left(x_j := \tilde{\zeta}_j(\tilde{\mathbf{PA}}_j^{\mathcal{G}}, \tilde{\varepsilon}_j)\right)}(x_j | \mathbf{PA}_j^{\mathcal{G}}). \tag{3}$$

If there are K possible interventions, we denote the corresponding sets of target variables as $I^{(k)}$, for $k = 1, \ldots, K$, and the corresponding SCMs by $\tilde{\mathcal{M}}^{(k)}$.

We divide the types of interventions into: *atomic*, if the target variable x_j is set to a constant value; *stochastic*, if x_j is set to a random variable $\tilde{\varepsilon}_j$; *imperfect* (or *soft*), if the intervention embedding and the set of parents are changed, as long as it does not become empty. We do not consider interventions that are able to add new elements to $\mathbf{PA}_j^{\mathcal{G}}$. This means that the intervention graph only differs from the observational by the removal of edges.

2.3. Faithfulness and Markov equivalence classes

Given a set $\mathcal{F}$ where each element ζ_i is *sufficiently* dependent on all of arguments $\mathbf{PA}_i^{\mathcal{G}}$, we obtain an SCM $\mathcal{M}$ whose computations strictly follow the structure of $\mathcal{G}$. In this scenario, $\mathcal{G}$ and $p_{\mathcal{M}}(X)$ are said to be mutually *faithful* since $\mathcal{G}$ encodes all and only the conditional independencies that hold in the entailed distribution. The set of faithful graphs that could entail a particular joint distribution is called the *Markov equivalence class* (MEC) (Verma and Pearl, 2013). If there is access to intervention data (in a set of interventions $\mathcal{I}$), it is possible to shrink the MEC to the so-called $\mathcal{I}$-MEC (Hauser and Bühlmann, 2012): the subset of graphs in the MEC that have the same conditional independencies after applying the interventions in $\mathcal{I}$.

2.4. Continuous constrained optimization for structure learning

Our work builds on a recent line of research that uses continuous constrained optimization to address causal discovery, initiated by Zheng et al. (2018) and extended by Brouillard et al. (2020) to cases where there is data from intervention distributions. In general, these methods adopt the *maximum a posteriori* (MAP) criterion (a.k.a. penalized maximum likelihood). Based on a generative/sampling model $p(\mathcal{D}|\mathcal{G}, \theta)$ for data $\mathcal{D}$, given the graph structure $\mathcal{G}$ and parameters θ, and on a *prior* $p(\mathcal{G})$ over graphs, they seek a graph that maximizes the score function

$$\mathcal{S}(\mathcal{G}) := \max_{\theta} \log p(\mathcal{D}|\mathcal{G}, \theta) + \log p(\mathcal{G}). \tag{4}$$

The prior $p(\mathcal{G})$ penalizes graph complexity to avoid over-fitting. A typical choice is $p(\mathcal{G}) \propto \exp(-\lambda|\mathcal{G}|)$, for $\lambda > 0$ and $|\mathcal{G}|$ is some graph complexity measure (*e.g.*, number of edges). With finite data, exact independence seldom occurs, thus graphs maximizing $\log p(\mathcal{D}|\mathcal{G}, \theta)$ alone would almost always be fully connected. If $\mathcal{D}$ is a collection of i.i.d. observations, then $p(\mathcal{D}|\mathcal{G}, \theta) = \prod_{i=1}^{n} p(x_i|\mathcal{G}, \theta)$.

Central to this class of methods is the weighted adjacency matrix $W^{\mathcal{G}} \in \mathbb{R}_{\geq 0}^{d \times d}$, where $W_{ij}^{\mathcal{G}} > 0$ is equivalent to $(i, j) \in \mathcal{G}$, which is treated as a parameter itself or as a function of the parameters. To ensure the estimated graph is a DAG, Zheng et al. (2018) proposed the constraint

$$\text{trace}\big(e^{W^{\mathcal{G}}}\big) - d = 0, \tag{5}$$

where $e^{W^{\mathcal{G}}}$ is the matrix exponential. Several other methods apply non-linear models such as neural networks (Lachapelle et al., 2019; Zheng et al., 2020) and define $W^{\mathcal{G}}$ differently.

The works from Ng et al. (2020); Kalainathan et al. (2020); Brouillard et al. (2020) treat the adjacency matrix as a random variable and relax the score from Eq. 4 in the following way:

$$\mathcal{S}^{\star}(\Lambda) := \max_{\theta} \mathbb{E}_{\mathcal{G} \sim \text{Bern}\big(\mathcal{G}; \sigma(\Lambda)\big)} \Big[\log p(\mathcal{D}|\mathcal{G}, \theta) + \log p(\mathcal{G}) \Big], \tag{6}$$

where $\sigma(\Lambda)$ is the sigmoid transformation applied element-wise to the parameter matrix $\Lambda \in \mathbb{R}^{d \times d}$, $\text{Bern}\big(\mathcal{G}; \sigma(\Lambda)\big)$ is a distribution over graphs, with mutually independent edges, with expected value $\sigma(\Lambda)$. This score tends asymptotically to $\mathcal{S}(\mathcal{G})$ as $\sigma(\Lambda)$ progressively concentrates its mass on a single DAG $\mathcal{G}$.

3. Differentiable causal discovery under latent interventions

In this section, we present a score for perfect or imperfect fully latent interventions, and show how this score can be approximately maximized by using an efficient variational optimization algorithm.

3.1. Mixture of intervention distributions

We assume that the dataset $\mathcal{D}$ is produced by a mixture of SCMs, each resulting from an intervention applied to a base SCM $\mathcal{M}$. More specifically, $\mathcal{D}$ is partitioned into $K + 1$ exchangeable groups, with group k containing i.i.d. samples from the intervention SCM $\tilde{\mathcal{M}}^{(k)}$ resulting from applying the k^{th} intervention to the base SCM $\mathcal{M}$; the index $k = 0$ indicates the absence of intervention, *i.e.*, $\tilde{\mathcal{M}}^{(0)} = \mathcal{M}$ (observational model). We denote by $\tilde{\mathcal{M}} = (\tilde{\mathcal{M}}^{(0)}, \ldots, \tilde{\mathcal{M}}^{(K)})$ the ensemble of SCMs.

The latent variables $z^{(i)} \in \{0, \ldots, K\}$ indicate which SCM generated each sample: $z^{(i)} = k$ if and only if $x^{(i)}$ is a sample of the SCM $\tilde{\mathcal{M}}^{(k)}$. Treating these correspondences $z^{(i)}$ as latent is a distinctive aspect of our work; while Brouillard et al. (2020) also assume unknown $\tilde{\mathcal{M}}$, they assume that $z^{(i)}$ is observed, not latent. We call the scenario where both $\tilde{\mathcal{M}}$ and $z^{(i)}$ are unknown as *fully latent interventions*.

Marginalizing with respect to the latent $z^{(i)}$ yields the mixture model

$$p(x^{(i)}|\tilde{\mathcal{M}}) = \sum_{k=0}^{K} p(z^{(i)} = k)\, p(x^{(i)}|z^{(i)} = k, \tilde{\mathcal{M}}) = \sum_{k=0}^{K} \tau_k\, p_{\tilde{\mathcal{M}}^{(k)}}(x^{(i)}), \tag{7}$$

where we define $\tau_k = p(z^{(i)} = k)$. Conditioning on $z^{(i)}$ and invoking Eq. 3 leads to

$$p(x^{(i)}|z^{(i)}, \tilde{\mathcal{M}}) = \sum_{k=0}^{K} \mathbb{I}(z^{(i)} = k) \prod_{j \notin I^{(k)}} p_{\mathcal{M}}(x_j^{(i)}|\mathbf{PA}_j^{\mathcal{G}}) \prod_{j \in I^{(k)}} p_{\mathcal{M};do(x_j := \tilde{\zeta}_j^k(\mathbf{PA}_j^{\mathcal{G}}, \tilde{\varepsilon}_j))}(x_j^{(i)}|\mathbf{PA}_j^{\mathcal{G}}).$$

We also consider that, like the group memberships, the set of targets $I^{(k)}$ of each intervention k is unknown (except for $I^{(0)} = \emptyset$).

3.2. Distribution over causal graphs

We represent the causal graph $\mathcal{G}$ via the adjacency matrix $A^{\mathcal{G}} \in \{0, 1\}^{d \times d}$. Following previous work, our prior models each entry $A_{ij}^{\mathcal{G}}$, corresponding to edge $x_i \to x_j$, as a Bernoulli variable independent of all the others,

$$p(\mathcal{G}) = \prod_{i,j=1}^{d} \sigma(\lambda_{ij})^{A_{ij}^{\mathcal{G}}} \big(1 - \sigma(\lambda_{ij})\big)^{1 - A_{ij}^{\mathcal{G}}}, \tag{8}$$

where $\sigma(u) = e^u/(1 + e^u)$ is the usual logistic transformation (sigmoid) and the λ_{ij} are hyperparameters. This prior over graphs is simplistic since it does not encode that $\mathcal{G}$ has to be a DAG. In this paper, we set $\lambda_{ij} = \lambda_{\mathcal{G}}$, for all i, j; however, in practice, a domain expert using the proposed method can embed prior knowledge in these hyper-parameters (our method can be straightforwardly adapted to that case).

The adoption of a probabilistic prior $p(\mathcal{G})$ should not be seen as expressing that it is in fact a random object, but rather as a subjective prior in the context of epistemic uncertainty about it.

3.3. Intervention embeddings and shared intervention space

We use density estimators, *e.g.*, neural networks and normalizing flows (Rezende and Mohamed, 2015), to model the conditional densities in both the observational and interventional distributions.

With this goal in mind, we use an appropriate encoding of the changes in the intervened assignments and the intervention targets. The set of targets in the k^{th} intervention $\tilde{\mathcal{M}}^{(k)}$ is indicated by a d-dimensional binary vector $r_k = [r_{k1}, \ldots, r_{kd}]$, where $r_{kj} = 1$ if and only if $j \in I^{(k)}$, and $r_{kj} = 0$ otherwise. Since $I^{(0)}$ has no targets (it corresponds to the observational SCM $\mathcal{M}$), we have $r_0 = [0, 0, \ldots, 0]$. To encode the type of intervention, we introduce the *intervention embedding vector* $u_k \in \mathbb{R}^h$, where h is an hyper-parameter. The vector u_k represents the changes in the affected assignments for intervention $\tilde{\mathcal{M}}^{(k)}$. We denote by $\mathcal{R} = [r_0, r_1, \ldots, r_K] \in \mathbb{R}^{K \times d}$ the matrix of intervention targets, and by $\mathcal{U} = [u_0, u_1, \ldots, u_K] \in \mathbb{R}^{K \times h}$ the matrix of intervention embeddings. We use $(\mathcal{R}, \mathcal{U})$ as the representation of the interventions $\tilde{\mathcal{M}}$.

Putting everything together, when given the graph $\mathcal{G}$, interventions $\tilde{\mathcal{M}}$, indicator z, and assuming the intervention is imperfect, the log-probability of single data-point x is given by

$$\log p(x|z, \tilde{\mathcal{M}}, \mathcal{G}; \theta) =$$
$$= \sum_{k=0}^{K} \mathbb{I}(z = k) \sum_{j=1}^{d} (1 - r_{kj}) \log \underbrace{g_j(x_j|A_j^{\mathcal{G}} \odot x, u_0; \theta_j)}_{=: \, p_{\mathcal{M}}(x_j|\mathbf{PA}_j^{\mathcal{G}})} + r_{kj} \log \underbrace{g_j(x_j|A_j^{\mathcal{G}} \odot x, u_k; \theta_j)}_{=: \, p_{\tilde{\mathcal{M}}^{(k)}}(x_j|\mathbf{PA}_j^{\mathcal{G}})}$$
$$= \sum_{j=1}^{d} \log g_j(x_j|A_j^{\mathcal{G}} \odot x, (e_z^\top \mathcal{R})_j \, (e_z^\top \mathcal{U} - u_0) + u_0; \theta_j), \tag{9}$$

where $e_z \in \mathbb{R}^K$ is a one-hot vector indicating z, and $A_j^{\mathcal{G}} \odot x$ is the Hadamard (elementwise) product between the j^{th} column of the adjacency matrix of $\mathcal{G}$ and x, which is equivalent to selecting the entries of x in $\mathbf{PA}_j^{\mathcal{G}}$. The parameters $\theta = (\theta_1, \ldots, \theta_d)$ parameterize the conditional densities $g_1, \ldots, g_d$. We will consider in the sequel several forms for these conditional densities, *e.g.*, using parametric families and normalizing flows. Crucially, each conditional density g_j is a distribution of x_j, and the parameters θ_j are *shared* between all of the interventions—only the intervention-specific intervention embedding vector u_k changes depending on the intervention. This enables dealing with an unlimited number of interventions, as we shall see.

3.4. Modeling the conditional densities

A simple nonlinear model for the conditional densities g_j can be constructed using neural networks. We use a neural network $\text{NN}([u_k, A_j^{\mathcal{G}} \odot x]; \theta_j) : \mathbb{R}^{(h+d)} \to \mathbb{R}^m$, a parametric non-linear mapping, parameterized by θ_j, that receives the concatenation of the parents of x_j and the intervention embedding u_k and outputs m parameters of some distribution $f(x_j; \text{NN}([u_k, A_j^{\mathcal{G}} \odot x]; \theta_j))$ of the variable x_j. There are many possible choices for the distribution f, depending on the problem at hand and on whether x_j is discrete or continuous: Poisson ($m = 1$), Bernoulli ($m = 1$), univariate Gaussian ($m = 2$), categorical, etc. In this paper, we focus on three density families in $\mathbb{R}$, which we experiment with in Section 4.

Linear Gaussian We use a neural network $\text{NN}(u_k; \theta_j)$ to output coefficients $\tilde{a}_j \in \mathbb{R}^d$, and $\tilde{\sigma}_j, \tilde{b}_j \in \mathbb{R}$. Then, we use these as parameters of a Gaussian distribution whose mean is an affine transformation of the values of the parents of x_j:

$$g_j(x_j|A_j^{\mathcal{G}} \odot x, u_k) = \mathcal{N}\big(\tilde{a}_j^\top (A_j^{\mathcal{G}} \odot x) + \tilde{b}_j, \tilde{\sigma}_j^2\big).$$

Non-Linear Gaussian   We use a neural network $\text{NN}([u_k, A_j^{\mathcal{G}} \odot x]; \theta_j)$ to output coefficients $\tilde{\mu}_j \in \mathbb{R}$ and $\tilde{\sigma}_j \in \mathbb{R}$, given the values of the parents of x_j as input. Then, we use these as parameters of a Gaussian distribution:

$$g_j(x_j | A_j^{\mathcal{G}} \odot x, u_k) = \mathcal{N}\big(\tilde{\mu}_j, \tilde{\sigma}_j^2\big).$$

Normalizing flows   To model non-linear non-Gaussian conditional densities, we employ normalizing flows. A normalizing flow (Rezende and Mohamed, 2015) is a transformation of a base probability density (in our case a Gaussian) through a sequence of invertible mappings $\tau(x_j; \tilde{W}_j) = \tau_l \circ \tau_{l-1} \cdots \circ \tau_1(x_j; \omega_1)$, where $\tilde{W}_j = \{\omega_1, \ldots, \omega_l\}$. We use a model introduced in Huang et al. (2018), called Deep Sigmoidal Flows (DSF), where each of the invertible mappings has the following form:

$$\tau_l(x) = \sigma^{-1}(w_l^{\top} \sigma(a_l x + b_l)), \qquad w_l \in \triangle_{F-1}, \ a_l \in \mathbb{R}_+^F, \ b_l \in \mathbb{R}^F.$$

where $\triangle_{F-1}$ is the probability simplex and F is an hyper-parameter. We use a neural network $\text{NN}([u_k, A_j^{\mathcal{G}} \odot x]; \theta_j)$ to output the parameters $\tilde{W}_j$. With the former, we obtain a controllable flow $\tau(x_j; \tilde{W}_j)$ that when given x_j, outputs the parameters of a Gaussian distribution $\tilde{\mu}, \tilde{\sigma}$. Altogether, the joint density has the following form:

$$g_j(x_j | A_j^{\mathcal{G}} \odot x, u_k) = \left| \det\left(\frac{\partial \tau(x_j; \tilde{W}_j)}{\partial x_j} \right) \right| \mathcal{N}\big(\tilde{\mu}, \tilde{\sigma}^2\big).$$

3.5. Modeling latent interventions with a Dirichlet process

To obtain a complete statistical description of the data-generating process, we still need to design a prior distribution for the latent interventions, apart from the sampling model g_j. Namely, we need a prior distribution for the correspondence z, the intervention embeddings $\mathcal{U}$, and the intervention targets $\mathcal{R}$. Furthermore, while experimentally, we can design scenarios where K, the number of latent interventions, is known, in general, given a data set, it will not be clear what the number of latent interventions is. Therefore, we will formulate the model to support a potentially non-specified number of latent interventions K.

We do this by using a *Dirichlet process prior* with a stick-breaking process representation (Ferguson, 1973; Sethuraman, 1991) as the prior distribution of the variables associated with the latent interventions. The generative story is as follows. We first draw the graph $\mathcal{G}$ from $p(\mathcal{G})$, as described in Eq. 8. Then, for $k = 0, 1, \ldots$, we sample the variables u_k and r_k, associated with each intervention $\tilde{\mathcal{M}}^{(k)}$, as well as the probability β_k of picking that intervention as a stick-breaking process, with scaling parameter $\alpha > 0$ (which controls the clustering effect of the Dirichlet process) and hyperparameter γ (which controls the sparsity of the intervention targets), as follows:

$$
\begin{aligned}
u_k &\sim \mathcal{N}(0, I_h) \\
r_{kj} &\sim \text{Bern}(\sigma(\gamma)), \quad j = 1, \ldots, d \\
\beta_k &= v_k \prod_{k'=0}^{k-1} (1 - v_{k'}), \quad \text{with } v_k \sim \text{Beta}(1, \alpha).
\end{aligned}
\tag{10}
$$

Figure 1: Graphical model representation of the Dirichlet Process Mixture model augmented for the causal discovery problem.

Then, to generate the data, we first sample the intervention index $z^{(i)}$, and then sample a point $x^{(i)}$ conditioning on the corresponding intervention $\tilde{\mathcal{M}}^{(z^{(i)})}$:

$$z^{(i)} \sim \text{Cat}(\beta_0, \beta_1, \ldots, \beta_k, \ldots)$$

$$x_j^{(i)} \sim \begin{cases} g_j(x_j | A_j^{\mathcal{G}} \odot x^{(i)}, u_0) & \text{if } r_{z^{(i)}j} = 0 \\ g_j(x_j | A_j^{\mathcal{G}} \odot x^{(i)}, u_{z^{(i)}}) & \text{if } r_{z^{(i)}j} = 1. \end{cases} \tag{11}$$

Figure 1 contains a graphical model representation of this joint distribution.

3.6. Variational approximation

To obtain the marginal likelihood $p(\mathcal{D}|\mathcal{G}; \theta)$, it is necessary to marginalize over the joint distribution of the model present in Section 3.5, *i.e.*:

$$p(\mathcal{D}|\mathcal{G}; \theta) = \prod_{i=1}^{N} p(x^{(i)}|\mathcal{G}; \theta) = \prod_{i=1}^{N} \mathbb{E}_{z^{(i)}, \tilde{\mathcal{M}} \sim p(z^{(i)}, \tilde{\mathcal{M}})} \left[p(x^{(i)}|z^{(i)}, \tilde{\mathcal{M}}, \mathcal{G}; \theta) \right]. \tag{12}$$

However, the exact maximization of this marginal log-likelihood (which involves a product of Gaussians, Beta distributions, and complex conditional distributions generated by neural networks) is intractable. Therefore, we resort to *approximate variational inference* (Blei et al., 2017). We design a family of tractable variational distributions $q_\phi(z^{(i)}, \tilde{\mathcal{M}})$ to approximate the true posterior $p(z^{(i)}, \tilde{\mathcal{M}}|x^{(i)}, \mathcal{G}; \theta)$. For the variables associated with latent interventions $\tilde{\mathcal{M}}$ we propose the fully factorized and finite variational family

$$q(\tilde{\mathcal{M}}) = \prod_{k=0}^{K} \left(\prod_{j=1}^{d} q_R(r_{kj}) \right) \left(\prod_{l=1}^{h} q_U(u_{kl}) \right) q_V(v_k), \tag{13}$$

where here the hyper-parameter K defines the truncation level of the variational approximation, and the distributions associated with v_k, u_{kl}, and r_{kj} take the following form:

$$q_V(v_k; \rho_k, w_k) = \text{Beta}(\rho_k w_k, (1 - \rho_k)w_k);$$
$$q_U(u_{kl}; \mu_{kl}, \sigma_{kl}) = \mathcal{N}(\mu_{kl}, \sigma_{kl}^2); \tag{14}$$
$$q_R(r_{kj}; \pi_{kj}) = \text{Bern}(\pi_{kj}).$$

where $\pi_{kj}, \mu_{kl}, \sigma_{kl}, \rho_k, w_k$ are free parameters to be optimized, for each $k \in \{0, \ldots, K\}$, $l \in \{1, \ldots, h\}$ and $j \in \{1, \ldots, d\}$. For the distribution of interventional assignments z, we propose the following variational posterior:

$$q_Z(z) \propto \exp\left(\frac{u_z^\top \text{NN}(x; \phi_Z)}{\sqrt{h}}\right), \qquad k = 1, \ldots, K, \tag{15}$$

where $\text{NN}(x; \phi_Z) : \mathbb{R}^d \to \mathbb{R}^h$ is a neural network. We provide details in Appendix A about the derivation and, in the case of the beta distribution, closed-form approximation of the Kullback–Leibler divergence between the proposals and the prior; and we describe the architecture of the neural networks used in the experiments in Appendix C. We use the shorthand ϕ to denote the vector of all variational parameters, which includes $\phi_Z, \mu_{kl}, \sigma_{kl}, \rho_k, w_k, \pi_{kj}$ for all $k \in \{0, \ldots K\}$, $l \in \{1, \ldots, h\}$, and $j \in \{1, \ldots, d\}$.

Given the ingredients above, we obtain the following lower bound for the marginal likelihood from Eq. 12:

$$\log p(\mathcal{D}|\mathcal{G}; \theta) \geq \sum_{i=1}^{N} \underbrace{\mathbb{E}_{z^{(i)}, \tilde{\mathcal{M}} \sim q(z^{(i)}, \tilde{\mathcal{M}}; \phi)}\left[\log p(x^{(i)}|z^{(i)}, \tilde{\mathcal{M}}, \mathcal{G}; \theta)\right] - D_{KL}\left[q(z^{(i)}, \tilde{\mathcal{M}})||p(z^{(i)}, \tilde{\mathcal{M}})\right]}_{\text{ELBO}_{q(z^{(i)}, \tilde{\mathcal{M}}; \phi)}(x^{(i)}, \mathcal{G}; \theta)}.$$

For different choices of ϕ, we get different lower bound approximations to the marginal likelihood. By maximizing the ELBO w.r.t. ϕ we minimize the approximation gap, which equals he KL divergence between the approximate and the true posterior.

3.7. A score for latent interventions

Using the log-likelihood proposed in Eq. 12, we write a new score function $\mathcal{S}(\mathcal{G})$ for our model with latent interventions, with an associated relaxation to support a weighted adjacency $\sigma(\Lambda)$, as shown in Eq. 6. As discussed in Section 3.6, in general we will not be able to exactly maximize this score. However, using the variational approximation above, and associated variational family $\mathcal{Q}$, we can approximate the score $\mathcal{S}(\mathcal{G})$ with a surrogate score $\mathcal{S}^{\mathcal{Q}}$ as follows:

$$\mathcal{S}(\Lambda) \geq \underbrace{\max_{\theta, \phi} \mathbb{E}_{\mathcal{G} \sim \text{Bern}\left(\mathcal{G}; \sigma(\Lambda)\right)}\left[\sum_{i=1}^{N} \text{ELBO}_{q(z^{(i)}, \tilde{\mathcal{M}}; \phi)}(x^{(i)}, \mathcal{G}; \theta) + \log p(\mathcal{G})\right]}_{\mathcal{S}^{\mathcal{Q}}(\Lambda)}. \tag{16}$$

The gap between the relaxed score $\mathcal{S}(\Lambda)$ and our surrogate $\mathcal{S}^{\mathcal{Q}}(\Lambda)$ tends asymptotically to the KL divergence between the best approximate posterior from $\mathcal{Q}$ and the true posterior, as $\sigma(\Lambda)$ progressively concentrates its mass on single DAG $\mathcal{G}$, more concretely:

$$\mathcal{S}(\Lambda) - \mathcal{S}^{\mathcal{Q}}(\Lambda) = \mathbb{E}_{\mathcal{G} \sim \text{Bern}\left(\mathcal{G}; \sigma(\Lambda)\right)}\left[D_{KL}\left[q(z^{(i)}, \tilde{\mathcal{M}}; \phi^*)||p(z^{(i)}, \tilde{\mathcal{M}}|x^{(i)}, \mathcal{G}; \theta^*)\right]\right], \tag{17}$$

where θ^* and ϕ^* are respectively the model and variational parameters that maximize Eq. 16.

3.8. Inference algorithm

The surrogate score, coupled with the acyclicity constraint from Zheng et al. (2018), enables us to formulate causal discovery under latent interventions as the following optimization problem:

$$\Lambda^* = \arg\max_{\Lambda} \mathcal{S}^{\mathcal{Q}}(\Lambda) \quad \text{s.t.} \quad \underbrace{\text{Tr}\left(e^{\sigma(\Lambda)}\right) - d = 0}_{h(\Lambda)}, \tag{18}$$

Following Zheng et al. (2018), we use the augmented Lagrangian procedure (Hestenes, 1969; Powell, 1969; Glowinski and Marroco, 1975) to transform the problem in Eq. 18 into a sequence of unconstrained optimization subproblems. When we estimate the gradients using the path-wise gradient estimator (Kingma and Welling, 2014; Rezende et al., 2014), each unconstrained optimization subproblem reduces to sampling the graph and the latent variables from the variational posteriors using the reparametrization trick, minimizing the following objective:

$$\mathcal{L}(\theta, \phi, \Lambda, z, \tilde{\mathcal{M}}, \mathcal{G}; x, \mu_t, \varphi_t) = -\log p(x|z, \tilde{\mathcal{M}}, \mathcal{G}; \theta) + \Omega(\phi) - \lambda_{\mathcal{G}}||\Lambda||_1 + \varphi_t h(\Lambda) + \frac{\mu_t}{2} h(\Lambda)^2,$$

where $\Omega(\phi)$ denotes the sum of the Kullback–Leibler divergences, and μ_t and φ_t are the parameters of the augmented Lagrangian at the t^{th} iteration. For estimating the gradients of Λ, $q(z)$ and $q(r_{kj})$ we used a Gumbel-softmax continuous relaxation (Jang et al., 2017; Maddison et al., 2017) which, for the causal graph's distribution, was combined with the straight-through gradient estimator (Bengio et al., 2013), to make sure the graph samples actually represented the hard dependencies of the SCM, instead of fractional ones. Having estimated the gradients, for each sample, we average them and feed them to the first order stochastic optimization algorithm Adam (Kingma and Ba, 2015). We implemented our method with the PyTorch framework (Li et al., 2020).

3.9. Semi-supervised extensions

The causal discovery algorithm can be extended to cases where we have information about the correspondences $z^{(i)}$ and/or the intervention targets $I^{(k)}$. To achieve this, we only have to ignore the corresponding variational posteriors and use the observed values $z^{(i)}$ and $r^{(i)}$ as constants. We designate the original version of our model the **latent** variant, the one with observed $z^{(i)}$ the **unknown** variant, and the one with observed $z^{(i)}$ and $r^{(i)}$ the **known** variant. It is straightforward to extend our method to a semi-supervised setting—a scenario where we only observe the correspondence z for a fraction of the samples. Under this scenario, we still use the samples from the variational posterior to predict the unobserved values of z and use the observed ones to improve the variational posterior. In essence, with the semi-supervised variant, we obtain a model that interpolates between the latent variant and the unknown variant. We follow the approach from Kingma et al. (2014) and extend the ELBO objective (in our case, the lower bound of the distribution $p(\mathcal{D}|\mathcal{G}; \theta)$). Let $\mathcal{O}$ be the set of the indices for which the intervention assignment z is known. We write the semi-supervised score $\mathcal{S}^{\text{SSL}}(\Lambda)$ as the surrogate score from Eq. 16, replacing the ELBO with:

$$\mathcal{L}^{\text{SSL}}(\mathcal{D}, \mathcal{G}) = \sum_{i=1}^{N} \text{ELBO}_{\theta, q(z^{(i)}, \tilde{\mathcal{M}})}(x^{(i)}, \mathcal{G}) + \kappa \, \mathbb{E}_{\tilde{\mathcal{M}} \sim q(\tilde{\mathcal{M}})} \Big[\sum_{i \in \mathcal{O}} \log q(z^{(i)}|x^{(i)}, \tilde{\mathcal{M}}; \phi_z) \Big],$$

where in the first term, when $i \in \mathcal{O}$, we do not sample from $q(z^{(i)}|x^{(i)}, \tilde{\mathcal{M}}; \phi_z)$, using the observed value instead. The variable $\kappa \in]0, 1[$ is an hyper-parameter that controls the relative importance of supervised component of the objective.

4. Experiments

We tested our method on synthetic and real data. The synthetic datasets allow us to do a systematic, controlled comparison of different methods in different scenarios (graph size and density, intervention and assignment types). In order to generate SCMs, perform interventions on them, and sample the corresponding entailed distributions, we created a Pytorch package for this purpose.[1]

Single-node interventions. We generated SCMs with $d = 10$ variables with a Erdős-Rényi scheme with expected number of edges per node $e \in \{1, 4\}$. We provide a detailed description of how we generate SCMs in Appendix B. We generated 10 SCMs for each combination of e, conditional density (linear Gaussian, non-linear Gaussian, and normalizing flow), and intervention type (stochastic and imperfect). In each of the SCMs, we performed 1 interventions for every variable. For each SCM within each configuration, we explored the following hyperparameter range: $\lambda_{\mathcal{G}} \in \{-.1, -.01, 0, .01, .1\}$, and $\gamma \in \{-.1, -.01\}$. For the remaining hyperparameters, we set $\alpha = 9$, $h = 248$ and the truncation level $K = 11$—the hyperparameter configuration that achieved the best log-likelihood on a validation set. From the generated SCM, we produced a dataset with $n = 10000$ samples, where each intervention has $\lfloor \frac{n}{d+1} \rfloor$ elements. This dataset was split into training (80%) and validation (20%). The models were trained for 500 epochs for the first iteration of the augmented Lagrangian and 50 epochs for the remaining ones, with a full batch($B = 8000$) until $h(\Lambda) < 10^{-8}$, and learning rate of $10^{-2.5}$.

The results from these experiments are shown in Table 1. The metric we use is Hamming distance (HD) between the adjacency matrices of the ground-truth graph and the estimated one. The columns from Table 1 correspond to the different variants we present in Section 3.9: **latent** refers to our model, **unknown** assumes that the correspondences are known (as Brouillard et al. (2020)), and **known** assumes that both correspondences and intervention targets are known. We additionally include a "naive" observational model (a baseline which ignores the existence of intervention data, *i.e.*, that assumes $K = 1$ as if all data was generated by the observational model). The experimental results preponderantly show that our method (the latent variant) consistently outperforms the observational baseline and is worst than the unknown variant, as expected, but only slightly. This indicates that taking account latent interventions, when these are present, improves the recovery of the causal graph.

Figure 2: Histogram of Hamming distance in the experiments with cause-effect pairs.

Single-node interventions on cause-effect pairs. We generated two-variable linear Gaussian SCMs (cause-effect pairs) with edge probability $e = \frac{2}{3}$ (equal probability for each of the three possible graphs). The SCMs were generated as described in Appendix B. We sampled 60 SCMs, and generate a dataset of $n = 999$ samples, where each intervention has 333 elements. The sampled graph $\mathcal{G}$, with variables A and B is one of 3 possible graphs, A causes B, B causes A, or A and B have no cause-effect relation. All of the possible graphs share

---

1. The code will be provided upon acceptance.

Model Type	e	latent	unknown	known	observational
Stochastic Interventions:					
Linear Gaussian		5.9 ± 6.2	3.4 ± 3.2	0.5 ± 1.3	10.3 ± 7.8
Non-Linear Gaussian	1	12.2 ± 3.9	10.3 ± 2.5	7.0 ± 3.6	13.7 ± 3.8
Non-Linear Non-Gaussian		8.7 ± 6.6	8.0 ± 2.7	6.6 ± 2.2	11.3 ± 5.0
Linear Gaussian		27.2 ± 6.2	24.1 ± 5.8	15.6 ± 6.0	39.6 ± 5.0
Non-Linear Gaussian	4	35.8 ± 3.8	30.3 ± 5.3	27.7 ± 4.3	37.5 ± 5.2
Non-Linear Non-Gaussian		36.1 ± 4.4	35.5 ± 8.1	31.5 ± 5.6	40.2 ± 6.9
Imperfect Interventions:					
Linear Gaussian		5.8 ± 4.2	6.2 ± 3.06	4.7 ± 3.6	10.4 ± 2.9
Non-Linear Gaussian	1	9.3 ± 2.4	8.9 ± 2.5	7.8 ± 3.9	10.5 ± 2.8
Non-Linear Non-Gaussian		8.8 ± 3.0	9.1 ± 3.5	7.9 ± 1.4	11.5 ± 5.4
Linear Gaussian		35.9 ± 8.3	29.7 ± 5.6	17.7 ± 7.9	39.1 ± 9.1
Non-Linear Gaussian	4	32.1 ± 6.0	32.6 ± 5.8	32.8 ± 5.4	39.8 ± 9.3
Non-Linear Non-Gaussian		30.4 ± 12.2	30.2 ± 11.2	25.8 ± 3.9	36.7 ± 9.8

Table 1: Hamming distances on synthetic 10 variable SCMs.

the MEC. We compare the variants presented in Section 3.9 in addition to the naive observational baseline. Figure 2 contains an histogram of Hamming distances for each of the variants. Edges in the anti-causal direction cost HD = 2, missing edge or wrong edge is HD = 1, and correct graph is HD = 0. The results show for this simple problem that cannot be identified using observational data alone, that our method correctly identifies a significant majority of the cause-effect pairs, even without information about the intervention assignments.

Real-world data set Finally, we tested our method on the flow cytometry data set of Sachs et al. (2005). Table 2 compares the estimated graph, under different conditional density assumptions but assuming imperfect interventions, to the consensus graph from Sachs et al. (2005). The results from the real-world data set show that our method outperforms several baselines, even with methods that use information regarding intervention correspondences and targets. The reasons that might justify relatively good results on this standard problem is that i) the causal sufficiency assumption may not hold, ii) the interventions may not be as specific as stated, and iii) the ground truth network is possibly not a DAG since feedback loops are common in cellular signaling networks as noted by Brouillard et al. (2020). These reasons can potentially be detrimental to the other methods and our method appears to be robust to them.

5. Related Work

The previous work that is closest to ours in that by Brouillard et al. (2020). As they do, we assume that the data is clustered in intervention groups, but we relax their assumption that the correspondence between intervention groups and samples is known, opening the door to more realistic scenarios.

The so-called "known" and "unknown" variants of our method share the assumptions made by Brouillard et al. (2020); however, contrary to them, the number of distinct neural networks in our model is not dependent on the number of interventions, which allows scaling well to many inter-

	SHD	tp	fn	fp	rev	F_1 score
IGSP (Wang et al., 2017)	18	4	6	5	7	0.42
GIES (Hauser and Bühlmann, 2012)	38	10	0	41	7	0.33
CAM (Bühlmann et al., 2014)	35	12	1	30	4	0.51
DCDI-G (Brouillard et al., 2020)	36	6	2	25	9	0.31
DCDI-DSF (Brouillard et al., 2020)	33	6	2	22	9	0.33
Linear Gaussian (imperfect) **(ours)**	33	7	11	22	3	0.30
Non-Linear Gaussian (imperfect) **(ours)**	19	7	11	8	0	0.42
Normalizing Flow (imperfect) **(ours)**	30	9	9	21	1	0.38

Table 2: Results for the flow cytometry dataset. The results for the baselines are reproduced from Brouillard et al. (2020).

ventions. We achieve this by encoding the change in assignments associated with each intervention using a specific latent variable u that we call intervention embedding and conditioning a shared model on it when computing the log-probability.

Our method can be seen as a *variational autoencoder* (VAE) with a Dirichlet process (DP) prior, with a stick-breaking process representatoion. The work by Nalisnick and Smyth (2017) introduces a VAE where the stick-breaking weights are the latent variables. Our model differs from theirs in that we use the entire DP (including the atoms). We only use the stick-breaking weights in the KL divergence of the correspondence variable z. This KL divergence has a closed-form, so we do not sample the stick-breaking weights as they do. van den Oord et al. (2017) proposes a VQ-VAE model with discrete latent variables, each represented as a latent embedding vector (an atom). Our approach is similar in that we use atoms (the intervention embeddings and intervention targets) to represent discrete latent variables. However, we do it in a statistically sounder way that more naturally fits our application.

6. Conclusion

We introduced an efficient variational optimization algorithm for causal structure learning under latent interventions. Our results are competitive with other state-of-the-art algorithms on the flow cytometry data set. Experiments with synthetic data show that our approach can recover causal relations even in our more challenging scenario, and it can consistently outperform our purely observational alternative.

A limitation of our method is the variational family we use. The proposal for variational posterior considers that the stick-breaking weights, the intervention embeddings, and intervention targets are independent of each other. In some cases, this can potentially create a surrogate score whose maximum structure is not an element of the $\mathcal{I}$-Markov equivalence class.

There are many avenues for future work. Our framework is particularly appealing for problems where performing interventions explicitly is expensive or unethical, but where interventions occur naturally in the data without being explicitly observed. Experimenting with more flexible variational families also seems appealing, albeit this may come at the cost of the closed-form expressions for the KL divergences.

References

Y. Bengio, N. Léonard, and A. Courville. Estimating or propagating gradients through stochastic neurons for conditional computation, 2013.

D. Blei, A. Kucukelbirb, and J. McAuliffe. Variational inference: A review for statisticians. *Journal of the American Statistical Association*, 112(518):859–877, 2017.

R. R. Bouckaert. Probabilistic network construction using the minimum description length principle. In Michael Clarke, Rudolf Kruse, and Serafín Moral, editors, *Symbolic and Quantitative Approaches to Reasoning and Uncertainty*, pages 41–48, Berlin, Heidelberg, 1993. Springer Berlin Heidelberg. ISBN 978-3-540-48130-0.

P. Brouillard, S. Lachapelle, A. Lacoste, S. L. Julien, and A. Drouin. Differentiable causal discovery from interventional data. *CoRR*, abs/2007.01754, 2020. URL https://arxiv.org/abs/2007.01754.

P. Bühlmann, J. Peters, and J. Ernest. CAM: Causal additive models, high-dimensional order search and penalized regression. *The Annals of Statistics*, 42(6):2526 – 2556, 2014. doi: 10.1214/14-AOS1260. URL https://doi.org/10.1214/14-AOS1260.

D. M. Chickering and D. Heckerman. Efficient approximations for the marginal likelihood of bayesian networks with hidden variables. In *Machine Learning*, pages 29–181, 1997.

T. S. Ferguson. A Bayesian Analysis of Some Nonparametric Problems. *The Annals of Statistics*, 1(2):209 – 230, 1973. doi: 10.1214/aos/1176342360. URL https://doi.org/10.1214/aos/1176342360.

R. Glowinski and A. Marroco. Sur l'approximation, par éléments finis d'ordre un, et la résolution, par pénalisation-dualité d'une classe de problèmes de dirichlet non linéaires. 1975.

A. Hauser and P. Bühlmann. Characterization and greedy learning of interventional markov equivalence classes of directed acyclic graphs. *J. Mach. Learn. Res.*, 13(1):2409–2464, August 2012. ISSN 1532-4435.

D. Heckerman, D. Geiger, and D. Chickering. Learning bayesian networks: The combination of knowledge and statistical data. In R. L.de Mantaras and D. Poole, editors, *Uncertainty Proceedings 1994*, pages 293–301. Morgan Kaufmann, San Francisco (CA), 1994. ISBN 978-1-55860-332-5. doi: https://doi.org/10.1016/B978-1-55860-332-5.50042-0. URL https://www.sciencedirect.com/science/article/pii/B9781558603325500420.

M. R. Hestenes. Multiplier and gradient methods. *Journal of Optimization Theory and Applications*, 4(5):303–320, 1969. doi: 10.1007/bf00927673. URL https://app.dimensions.ai/details/publication/pub.1018609722.

C. W. Huang, D. Krueger, A. Lacoste, and A. C. Courville. Neural autoregressive flows. *CoRR*, abs/1804.00779, 2018. URL http://arxiv.org/abs/1804.00779.

E. Jang, S. Gu, and B. Poole. Categorical reparameterization with gumbel-softmax, 2017.

D. Kalainathan, O. Goudet, I. Guyon, D. Lopez-Paz, and M. Sebag. Structural agnostic modeling: Adversarial learning of causal graphs, 2020.

D. P. Kingma and J. Ba. Adam: A method for stochastic optimization. *CoRR*, abs/1412.6980, 2015.

D. P Kingma and M. Welling. Auto-encoding variational bayes, 2014.

D. P. Kingma, D. J. Rezende, S. M., and M. Welling. Semi-supervised learning with deep generative models. *CoRR*, abs/1406.5298, 2014. URL http://arxiv.org/abs/1406.5298.

J. Kuipers, G. Moffa, and D. Heckerman. Addendum on the scoring of gaussian directed acyclic graphical models, 2021.

S. Lachapelle, P. Brouillard, T. Deleu, and S. L. Julien. Gradient-based neural DAG learning. *CoRR*, abs/1906.02226, 2019. URL http://arxiv.org/abs/1906.02226.

S. Li, Y. Zhao, R. Varma, O. Salpekar, P. Noordhuis, T. Li, A. Paszke, J. Smith, B. Vaughan, P. Damania, and S. Chintala. Pytorch distributed: Experiences on accelerating data parallel training. *CoRR*, abs/2006.15704, 2020. URL https://arxiv.org/abs/2006.15704.

C. J. Maddison, A. Mnih, and Y. W. Teh. The concrete distribution: A continuous relaxation of discrete random variables, 2017.

E. Nalisnick and P. Smyth. Stick-breaking variational autoencoders, 2017.

I. Ng, Z. Fang, S. Zhu, Z. Chen, and J. Wang. Masked gradient-based causal structure learning, 2020.

J. Pearl. Causality: Models, reasoning, and inference, second edition. *Causality*, 29, 01 2000. doi: 10.1017/CBO9780511803161.

J. Peters, D. Janzing, and B. Schölkopf. *Elements of Causal Inference - Foundations and Learning Algorithms*. Adaptive Computation and Machine Learning Series. The MIT Press, Cambridge, MA, USA, 2017.

M.J.D. Powell. A method for nonlinear constraints in minimization problems. *In: Fletcher, R., Ed., Optimization, Academic Press*, pages 283–298, 1969.

D. Rezende and S. Mohamed. Variational inference with normalizing flows. In *International Conference on Machine Learning (ICML)*, 2015.

D. J. Rezende, S. M., and D. Wierstra. Stochastic backpropagation and approximate inference in deep generative models, 2014.

K. Sachs, O. Perez, D. Pe'er, D. A. Lauffenburger, and G. P. Nolan. Causal protein-signaling networks derived from multiparameter single-cell data. *Science*, 308(5721):523–529, 2005. doi: 10.1126/science.1105809.

J. Sethuraman. A constructive definition of Dirichlet priors. 1991.

P. Spirtes, C. Glymour, and R. Scheines. *Causation, Prediction, and Search*, volume 81. 01 1993. ISBN 978-1-4612-7650-0. doi: 10.1007/978-1-4612-2748-9.

P. Spirtes, C. Meek, and T. S. Richardson. Causal inference in the presence of latent variables and selection bias. *CoRR*, abs/1302.4983, 2013. URL http://arxiv.org/abs/1302.4983.

A. van den Oord, O. Vinyals, and K. Kavukcuoglu. Neural discrete representation learning. *CoRR*, abs/1711.00937, 2017. URL http://arxiv.org/abs/1711.00937.

T. S. Verma and J. Pearl. On the equivalence of causal models. *CoRR*, abs/1304.1108, 2013. URL http://arxiv.org/abs/1304.1108.

Y. Wang, L. Solus, K. D. Yang, and C. Uhler. Permutation-based causal inference algorithms with interventions, 2017.

X. Zheng, B. Aragam, P. Ravikumar, and E. P. Xing. Dags with no tears: Continuous optimization for structure learning, 2018.

X. Zheng, C. Dan, B. Aragam, P. Ravikumar, and E. P. Xing. Learning sparse nonparametric dags, 2020.

Appendix A. Kullback–Leibler divergences

$$D_{KL}\big[q(\tilde{\mathcal{M}}, z^{(i)})||p(\tilde{\mathcal{M}}, z^{(i)})\big] =$$

$$= \mathbb{E}_{\tilde{\mathcal{M}}, z \sim q(\tilde{\mathcal{M}}, z)}\left[\log\frac{q(z|u_0, \ldots, u_T)\prod_{k=0}^{T}\left(\prod_{j=1}^{d}q_R(r_{kj})\right)\left(\prod_{l=1}^{h}q_U(u_{kl})\right)q_V(v_k)}{p(z|\beta_0, \ldots, \beta_T)\prod_{k=0}^{T}\left(\prod_{j=1}^{d}p(r_{kj}|\gamma)\right)\left(\prod_{l=1}^{h}p(u_{kl})\right)p(v_k|\alpha)}\right]$$

$$= \sum_{k=0}^{T}\sum_{l=1}^{h}D_{KL}\big(q(u_{kl})||\mathcal{N}(0,1)\big) + \sum_{k=0}^{T}\sum_{j=1}^{d}D_{KL}\big(q(r_{kj})||\mathrm{Bern}(\gamma)\big)$$

$$+ \sum_{k=0}^{T}D_{KL}\big(q(v_k)||p_\theta(v_k|\alpha)\big) + \mathbb{E}_{\tilde{\mathcal{M}}, z \sim q(\tilde{\mathcal{M}}, z)}\left[\log\frac{q(z|u_0, \ldots, u_T)}{p(z|\beta_0, \ldots, \beta_T)}\right]$$

$$(19)$$

Stick-breaking weights

$$D_{KL}\big(q(v_k)||p_\theta(v_k|\alpha)\big) = \log\left(\frac{\mathrm{B}\big(\rho_k w_k, (1-\rho_k)w_k\big)}{\mathrm{B}(1, \alpha)}\right) + (1-\rho_k w_k)\psi(1)+$$

$$\big(\alpha - (1-\rho_k)w_k\big)\psi(\alpha) + (-1 + w_k - \alpha)\psi(1+\alpha)$$

$$(20)$$

Intervention embeddings

$$D_{KL}\big(q(u_k)||\mathcal{N}(\mathbf{0}, I_h)\big) = \sum_{j=1}^{h}\frac{\sigma_{kj}^2 + \mu_{kj}^2 - 1}{2} - \log\sigma_{kj} \tag{21}$$

Intervention targets

$$D_{KL}\big(q(r_{kj})||\mathrm{Bernoulli}(\gamma)\big) = \pi_{kj}\Big(\mathrm{logit}(\pi_{kj}) - \gamma\Big) + \log\frac{1-\pi_{kj}}{1-\sigma(\gamma)} \tag{22}$$

Correspondence

$$\mathbb{E}_{\mathcal{V}^k, u_0, \ldots, u_K \sim q(\mathcal{V}^k)q(u_0)\ldots q(u_K)}\Big[D_{KL}\big(q(z)||p_\theta(z|\beta_0, \beta_1, \ldots, \beta_K)\big)\Big] =$$

$$= \mathbb{E}_{u_k \sim q(u_k)}\left[\sum_{k=0}^{K}q(z_k)\big(\log q(z_k) - \mathbb{E}_{\mathcal{V}^k \sim q(\mathcal{V}^k)}[\log\beta_k]\big)\right]$$

$$(23)$$

where

$$\mathbb{E}_{\mathcal{V}^k \sim q(\mathcal{V}^k)}[\log\beta_k] = \mathbb{E}_{v_k \sim q(v_k)}[\log v_k] + \sum_{k'=0}^{k-1}\mathbb{E}_{v_{k'} \sim q(v_{k'})}[\log 1 - v_{k'}]$$

$$= \psi(\rho_k w_k) + \sum_{k'=0}^{k-1}\psi\big((1-\rho_{k'})w_{k'}\big) - \sum_{k'=0}^{k}\psi(w_{k'}),$$

$$(24)$$

and ψ is the digamma function. We approximate using the Taylor series expansion, where we use the reparametrization trick on u_k to estimate the expectations.

Appendix B. Description of the synthetic datasets

To generate the SCMs, we first generated the causal graph. The graphs were generated going in order from node 1 to node N and adding edges from the already visited nodes to the current node i with probability p obtaining the adjacency matrix $A^{\mathcal{G}}$. In order to obtain graphs that weren't topologically ordered we applied a random permutation P to $A^{\mathcal{G}}$. Then, given the causal graph, we sampled the noise distributions and the assignments as described in Table 3. For every SCM, we generate d interventions, one for each variable. How we sample the new assignment for each intervened variables is described in Table 4. Before fitting the model, the data is always normalized. We subtract the mean and divide by the standard deviation.

For each setting, we sample $n/(d+1)$ examples. We used $n = 10000$ in the 10 variable data set. Before using our method, the data is always normalized. We subtract the mean and divide by the standard deviation. For stochastic interventions in variable j, we sampled a new Gaussian variable ε_j, as described in Table 4, and assigned it to x_j. Using imperfect interventions, we go to the last layer of the neural network (in the case of linear model its the weight vectors) representing the assignment, and sample new weights according to Table 4. Table 3 contains a description of how we sample assignments. Particularly, the architecture of the neural network, when it is non-linear SCM. Figure 3 contains scatter plots obtained after sampling the generated two variable SCMs.

Model Name	$p(\mathcal{E})$	weights	activation	layers	#hidden units
Linear Gaussian	$\prod_{j=1}^{d} \mathcal{N}(0, 0.015)$	$\mathcal{N}(0, 2.0)$	identity	1	-
Non-Linear Gaussian	$\prod_{j=1}^{d} \mathcal{N}(0, 0.015)$	$\mathcal{N}(0, 2.0)$	relu	2	5
Non-Linear Non-Gaussian	$\prod_{j=1}^{d} \mathcal{N}(0, 0.015)$	$\mathcal{N}(0, 2.0)$	relu	2	5

Table 3: Where $k \sim \mathcal{N}(0.1, 0.005)$ and $\ell \sim \mathcal{N}(0, 0.4)$.

Intervention type	new $p(\mathcal{E})$	new weights
Atomic	$(1 - 2b) \cdot u$	0
Stochastic	$\mathcal{N}((1 - 2b) \cdot u, 0.1)$	0
Imperfect	$\mathcal{N}((1 - 2b) \cdot u, 0.1)$	$\mathcal{N}(0, 2.0)$

Table 4: Where $u \sim \mathcal{U}[1.2, 2.2]$ and $b \sim \text{Bernoulli}(0.5)$

Figure 3: The underlying graph is has two variables and one edge. The cause is in the horizontal axis, the effect is in the vertical axis. The rows correspond to different models, particularly, Linear Gaussian, Non-Linear Gaussian and Non-Linear Non-Gaussian. The columns correspond to distinct types of interventions. Particularly, atomic, stochastic and imperfect. Red observational. Green intervention on the effect variable. blue intervention on the causal variables.

Appendix C. Neural network details

The neural network we used is a feed-forward (fully connected) network. Given an vector $x \in \mathbb{R}^{d_{in}}$, we parameterize with a vector of weights θ a non-linear mapping FFN : $\mathbb{R}^{d_{in}} \to \mathbb{R}^{d_{out}}$. This mapping is a composition of linear transformation interchanged with a non-linear element-wise operation. We use many tricks that have been shown to improve the results with neural network modules, particularly dropout, SiLU activation function, layer normalization and residual connections. We group these techniques together into a sub-layer that form the building block of our relatively more complex neural network models, as show in Figure 4.

Figure 4: Diagram of the NN-Block sub-layer.

To enable the use of skip connections, we have to make sure that the inputs and outputs of the sub-layer have the same dimension. For this reason before applying the stack of sub-layers we apply a linear transformation from $\mathbb{R}^{d_{in}}$ to $\mathbb{R}^{d_h}$ and after passing the stack from $\mathbb{R}^{d_h}$ to $\mathbb{R}^{d_{out}}$, as shown in the diagram in Figure 5.

Figure 5: Diagram of the Feed Forward Network(FFN) we use as functional approximator in our model.

Appendix D. Semi-supervised learning experiments

We test our semi-supervised method, on 10 variable linear SCMs generated according to Appendix B. We consider stochastic and imperfect interventions. We sampled a 10000 samples dataset from each SCM. For each dataset, we created multiple semi-supervised learning datasets by splitting the original 10000 samples into labelled (with intervention assignments) and unlabelled (without intervention assignments) according to some fraction f. The fraction f goes from 0 to 1, sampled every interval of 0.2. This means that, for each fraction f, we have 10 distinct semi-supervised datasets for both perfect and imperfect interventions. We picked the default hyper-parameters, particularly, $\lambda_{\mathcal{G}} = -0.1$, and $\gamma = -0.01$. The results are contained in Figure 6.

The results suggest, there is no significant improvement from using a the correspondences for a fraction of the samples under the tested conditions.

Figure 6: Semi supervised experiments on 10 variable Linear SCMs with an expected value of 1 edge per node. On the left, perfect interventions. On the right, imperfect interventions.