No Double Descent in PCA: Training and Pre-Training in High Dimensions

Abstract

With the recent body of work on overparameterized models the gap between theory and practice in contemporary machine learning is shrinking. While many of the present state-of-the-art models have an encoder-decoder architecture, there is little theoretical work for this model structure. To improve our understanding in this direction, we consider linear encoder-decoder models, specifically PCA with linear regression on data from a low-dimensional manifold. We present an analysis for fundamental guarantees of the risk and asymptotic results for isotropic data when the model is trained in a supervised manner. The results are also verified in simulations. Furthermore, we extend our analysis to the popular setting where parts of the model are pre-trained in an unsupervised manner by pre-training the PCA encoder with subsequent supervised training of the linear regression. We show that the overall risk depends on the estimates of the eigenvectors in the encoder and present a sample complexity requirement through a concentration bound. The results highlight that using more pre-training data decreases the overall risk only if it improves the eigenvector estimates. Therefore, we stress that the eigenvalue distribution determines whether more pre-training data is useful or not.

1 Introduction

Many recent success stories of deep learning employ an encoder-decoder structure, where parts of the model are pre-trained in an unsupervised or self-supervised way. Examples can be found in computer vision (Caron et al., 2020; Chen et al., 2020; Goyal et al., 2021), natural language processing (Vaswani et al., 2017; Devlin et al., 2019; Raffel et al., 2020) or multi-modal models (Ramesh et al., 2021; Alayrac et al., 2022). Understanding the properties of this model structure might shed light on how to reliably build large-scale models.

We add to the theoretical understanding of encoder-decoder based models by studying a model consisting of PCA and a linear regression head. We analyse this model for the supervised case and for the case where unsupervised pre-training is followed by supervised linear regression. Our model can be viewed as a simplified, linear example of a large pre-trained deep neural network in combination with linear probing (Devlin et al., 2019; Schneider et al., 2019). While linear models do not reveal the whole picture, they are studied as a tractable, first step towards deeper understanding. Indeed, research on linear models has previously provided important insights into relevant mechanisms (Saxe et al., 2014; Lampinen & Ganguli, 2019; Arora et al., 2019; Gidel et al., 2019; Pesme et al., 2021).

We utilize data generated from a low-dimensional manifold. This is motivated by the manifold hypothesis (Fefferman et al., 2016) which states that real-world high-dimensional data often have an underlying low-dimensional representation. Our PCA encoder can then exploit this data structure effectively. While we keep the low-dimensional data structure fixed, we vary the number of features w.r.t. the number of training data points which allows us to analyse what is often referred to as overparameterization, i.e. using more data features or model parameters than training samples (Belkin et al., 2019). We do not consider parameter count, since for our model the number of parameters, i.e. the linear regressors, stay fixed due to the PCA encoding. Instead we analyse high-dimensional settings. Studying overparameterization gives theoretical justification of the success of modern large-scale neural networks such as Szegedy et al. (2016); Dosovitskiy et al. (2021).

Theoretical grounding is exceeded by the empirical success of machine learning and specifically deep learning methods through new model structures (Krizhevsky et al., 2017; He et al., 2016; Vaswani et al., 2017) or training methods (Erhan et al., 2010; Ioffe & Szegedy, 2015; Ba et al., 2016). In recent years our theoretical understanding grew e.g. through the analysis of implicit regularization (Gunasekar et al., 2017; Chizat & Bach, 2020; Smith et al., 2021). But also experimental work contributed to our understanding (Keskar et al., 2017; Zhang et al., 2017). The goal of this paper is to extend our understanding of the successful encoder-decoder model structure through theoretical analysis of PCA-regression and by extensive numerical simulations. We generalize results for linear regression and combine classical analysis of overparamterization with pre-training of model components. Our contributions can be summarized as:

- In the **supervised case**, we provide **theoretical guarantees** for the risk and parameter norm of the PCA-regression model. For isotropic data we extend the results to the limit where the number of data points n and features m tend to infinity such that $m/n \to \gamma$.

- Through **simulations**, we confirm our theory for isotropic data, explain the model behavior on data from a low-dimensional manifold and establish connections to ridge regression.

- We extend our analysis to the popular scenario of unsupervised **pre-training of the encoder** and show that the correct estimation of feature covariance eigenvectors is crucial for low risk. These estimates are highly dependent on the data structure through the **eigenvalue decay rate**. We challenge the common wisdom that more pre-training data improves the overall risk and show that this is the case only if it improves the estimate of the eigenvectors in the encoder which is e.g. the case in data with rapidly decaying eigenvalues.

2   RELATED WORK

Overparameterization   The study of overparameterized models offers a natural route to gain theoretical understanding when it comes to the successes of large models with good generalization properties (Neyshabur et al., 2015; Zhang et al., 2017). The observation of the double descent Belkin et al. (2019) boosted research in this direction but generalization of large models was also studied before Bartlett & Mendelson (2002); Dziugaite & Roy (2017); Belkin et al. (2018); Advani et al. (2020). While double descent has been observed in deep and state-of-the-art models (d'Ascoli et al., 2020; Nakkiran et al., 2021), most of the theoretical studies focus on simple models: Examples are found for linear regression (Bartlett et al., 2020; Muthukumar et al., 2020; Hastie et al., 2022), ensembles (LeJeune et al., 2020), classification (Wang et al., 2021; Deng et al., 2022), random features (Belkin et al., 2019; Mei & Montanari, 2022) or small neural networks trained using gradient descent (Goldt et al., 2019; Advani et al., 2020). We add to the understanding of machine learning models by analysing the neglected class of encoder-decoder models with the PCA-regression model.

Analysis of pre-training   The introduction of pre-training of neural networks was a paradigm shift in deep learning research. The first attempts to understand the mechanisms relied on empirical work (Erhan et al., 2010; Raghu et al., 2019). Theoretical work explored the sample complexity when common representations are shared (Tripuraneni et al., 2020; Du et al., 2021) and the effect on out-of-distribution test samples (Kumar et al., 2022). For unsupervised pre-training there is extensive analysis of contrastive self-supervised methods (Wang & Isola, 2020; Von Kügelgen et al., 2021; Zimmermann et al., 2021). Encoder-decoder models are analysed in the form of unsupervised trained autoencoders for the training dynamics (Nguyen et al., 2019; 2021) and for overparameterization (Radhakrishnan et al., 2019; 2020; Buhai et al., 2020; Zhang et al., 2020), which identified patterns of training data memorization. In contrast, we analyse a pre-trained PCA encoder in the overparameterized setting and relate the risk to the covariance estimation quality in the encoder.

PCA—eigenvector estimation   Using PCA (Jolliffe, 1982) is common—discussions focus on the choice of principle components (Breiman & Freedman, 1983) or its use for high-dimensional data (Lee et al., 2012). PCA-regression is also investigated in Xu & Hsu (2019); Teresa et al. (2022). While the former assumes a known covariance matrix, the latter has no asymptotic results but compares experimentally with more projection methods. In the case with pre-training, we rely on sample complexity for the covariance estimation in the PCA for which we generalize the result of Loukas (2017). The data covariance structure turns out to be crucial as Wainwright (2019) points out.

3 PROBLEM FORMULATION

Data generator We generate a data set $\{\boldsymbol{x}_i, y_i\}_{i=1}^n$ according to a latent variable data generator

$$\boldsymbol{x}_i = \boldsymbol{D}\boldsymbol{z}_i + \boldsymbol{e}_i, \tag{1}$$

$$y_i = \boldsymbol{\theta}\boldsymbol{z}_i + \varepsilon_i, \tag{2}$$

by mapping the latent variable $\boldsymbol{z}_i \in \mathbb{R}^d$ with $\boldsymbol{D} \in \mathbb{R}^{m \times d}$ into the observed features $\boldsymbol{x}_i \in \mathbb{R}^m$ and with $\boldsymbol{\theta} \in \mathbb{R}^d$ into the observed outputs $y_i \in \mathbb{R}$. We create $\boldsymbol{D}$ randomly such that $\|\boldsymbol{D}\|_F^2 = dc^2$ with c as correction factor to control the signal-to-noise ratio (SNR), defined in (30). Similarly, to control the outcome-noise-ratio we create $\boldsymbol{\theta}$ such that $\mathbb{E}\left[\|\boldsymbol{\theta}^\top \boldsymbol{z}\|_2^2\right] = r_{\boldsymbol{\theta}}^2$. Feature noise $\boldsymbol{e}_i \sim \mathcal{N}(\boldsymbol{0}, \boldsymbol{I}_m)$ and output noise $\varepsilon_i \sim \mathcal{N}(0, \sigma_y^2)$ are added. The latent variables are generated such that the singular values of the features have an exponential decay controlled by the decay rate $\alpha \geq 0$ according to

$$\boldsymbol{z}_i \sim \mathcal{N}(0, \lambda_i^2 \boldsymbol{I}_d) \quad \text{with} \quad \lambda_i^2 = \exp(-i\alpha). \tag{3}$$

Our theoretical results do not specifically require an exponential decay of the eigenvalues or a specific rate. However, fast decaying eigenvalues occur in many real-world examples, see Appendix A. We distinguish between two data generators:

1. *Isotropic data.* This is a special case of (1), (2) with $d = m$, $\boldsymbol{D} = \boldsymbol{I}_m$, $\alpha = 0$ and $\boldsymbol{e} = \boldsymbol{0}$ to generate isotropic features. It allows us to rewrite the data generator as

$$y_i = \boldsymbol{\theta}\boldsymbol{x}_i + \varepsilon_i \quad \text{with} \quad \boldsymbol{x}_i \sim \mathcal{N}(0, \boldsymbol{I}_m). \tag{4}$$

2. *Latent variable data.* We distinguish between 1) $\alpha = 0$ leading to an isotropic but low-dimensional signal and 2) $\alpha > 0$ which has dominant, but rapidly decaying eigenvalues of the feature covariance matrix. The latter data generator is motivated since many real-world data sets have a low-dimensional signal manifold with rapidly decaying eigenvalues.

Note that while our latent variable data generator is similar to the latent space model from Hastie et al. (2022), we use our PCA-regression model instead of direct regression from features to outputs. A graphical model of our data generator is provided in Figure 1 and details are in Appendix B.

Figure 1: **Problem formulation.** *Left:* Data generator. *Right:* PCA and linear regression model.

Model We use a linear model which resembles an encoder-decoder based architecture. The input data $\boldsymbol{x}_i$ is encoded into a lower $\hat{d}$-dimensional space $\hat{\boldsymbol{z}}_i$ via the use of PCA, where $\hat{d}$ is chosen during model selection. When it comes to the decoder we employ a linear regression model with parameters $\hat{\boldsymbol{\theta}} \in \mathbb{R}^{\hat{d}}$. The resulting model is visualized in Figure 1 and we can formulate it as

$$\hat{\boldsymbol{z}}_i = \hat{\boldsymbol{V}}^\top \boldsymbol{x}_i, \tag{5}$$

$$\hat{y}_i = \hat{\boldsymbol{\theta}}^\top \hat{\boldsymbol{z}}_i. \tag{6}$$

Collecting the features as rows of the data matrix $\boldsymbol{X} = \begin{bmatrix} \boldsymbol{x}_1^\top & \dots & \boldsymbol{x}_n^\top \end{bmatrix}^\top$, we compute the principal components $\hat{\boldsymbol{V}} \in \mathbb{R}^{m \times \hat{d}}$ as the $\hat{d}$ first right singular vectors of the rank $\hat{d}$ reducing SVD $\boldsymbol{X} \approx \hat{\boldsymbol{U}}\hat{\boldsymbol{\Sigma}}\hat{\boldsymbol{V}}^\top$. Note that the estimated latent dimension $\hat{d}$ can be different from the true latent dimension d of the latent variable data generator. This construction is referred to as the *PCA-regression* model. In our numerical results we compare with a model which directly regresses the outcomes from the features, referred to as the *direct regression* model.

4 ANALYZING THE SUPERVISED CASE

Training the complete PCA-regression model in a supervised way represents a situation commonly encountered in high-dimensional real-world applications. Examples of using this model are in exploratory statistical research (Massy, 1965), econometrics (Geweke, 1996), genetics (Wang & Abbott, 2008), robotics (Vijayakumar & Schaal, 2000) and many more.

4.1 THEORETICAL ANALYSIS

For our analysis, we are interested in closed form solutions for the risk and parameter norm in order to obtain fundamental guarantees for the PCA-regression model. We decompose our solution into bias and variance terms similar to classic decompositions and interpret the results.

Bias-variance decomposition   We stack all outputs in the vector $\boldsymbol{y} \in \mathbb{R}^n$ and all estimated latent variables as rows in the matrix $\hat{\boldsymbol{Z}} \in \mathbb{R}^{n \times \hat{d}}$. The solution to the unregularized linear regression yields

$$\hat{\boldsymbol{\theta}} = (\hat{\boldsymbol{Z}}^\top \hat{\boldsymbol{Z}})^+ \hat{\boldsymbol{Z}}^\top \boldsymbol{y} = (\hat{\boldsymbol{V}}^\top \boldsymbol{X}^\top \boldsymbol{X} \hat{\boldsymbol{V}})^+ \hat{\boldsymbol{V}}^\top \boldsymbol{X}^\top \boldsymbol{y}, \tag{7}$$

where $(\cdot)^+$ denotes the Moore-Penrose pseudoinverse. We can rewrite our data generator directly from features to outputs as $\boldsymbol{y} = \boldsymbol{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$ with $\boldsymbol{\beta} \in \mathbb{R}^m$. Here, we transformed the noise to be sampled without loss of generality from $\epsilon_i \sim \mathcal{N}(0, \sigma_\epsilon^2)$. Following Appendix C.2 the solution becomes

$$\hat{\boldsymbol{\theta}} = (\hat{\boldsymbol{\Sigma}}^\top \hat{\boldsymbol{\Sigma}})^+ \hat{\boldsymbol{\Sigma}}^\top \hat{\boldsymbol{U}}^\top (\boldsymbol{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}) = \hat{\boldsymbol{V}}^\top \boldsymbol{\beta} + \hat{\boldsymbol{\Sigma}}^+ \hat{\boldsymbol{U}}^\top \boldsymbol{\epsilon}. \tag{8}$$

Lemma 1. *Let the sample covariance of centered features be $\hat{\boldsymbol{C}} = \frac{1}{n}\boldsymbol{X}^\top \boldsymbol{X}$ and the true covariance be $\boldsymbol{C}$. Define the orthogonal projectors $\boldsymbol{\Phi} = \hat{\boldsymbol{V}}\hat{\boldsymbol{V}}^\top$ and $\boldsymbol{\Pi} = \boldsymbol{I}_m - \boldsymbol{\Phi}$, where $\boldsymbol{\Phi}$ is the projection onto the column space of the $\hat{d}$ first right singular vectors of $\boldsymbol{X}$. Then, the risk of the PCA-regression model $R(\hat{\boldsymbol{\theta}}) = \mathbb{E}_{y_0}\left[(y_0 - \hat{y}_0)^2\right]$ and the parameter norm $\|\hat{\boldsymbol{\theta}}\|_2^2 = \hat{\boldsymbol{\theta}}^\top \hat{\boldsymbol{\theta}}$ are given by*

$$\mathbb{E}_{\boldsymbol{\epsilon}}\left[R(\hat{\boldsymbol{\theta}})\right] = \boldsymbol{\beta}^\top \boldsymbol{\Pi} \boldsymbol{C} \boldsymbol{\Pi} \boldsymbol{\beta} + \frac{\sigma_\epsilon^2}{n} \operatorname{Tr}(\hat{\boldsymbol{V}}^\top \boldsymbol{C} \hat{\boldsymbol{V}} \hat{\boldsymbol{V}}^\top \hat{\boldsymbol{C}}^+ \hat{\boldsymbol{V}}) + \sigma_\epsilon^2, \tag{9}$$

$$\mathbb{E}_{\boldsymbol{\epsilon}}\left[\|\hat{\boldsymbol{\theta}}\|_2^2\right] = \boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} + \frac{\sigma_\epsilon^2}{n} \operatorname{Tr}(\hat{\boldsymbol{V}}^\top \hat{\boldsymbol{C}}^+ \hat{\boldsymbol{V}}). \tag{10}$$

The proofs are in Appendices C.3, C.5. The variance (second) term is controlled by the estimated singular vectors $\hat{\boldsymbol{V}}$, which project the covariance matrices $\boldsymbol{C}, \hat{\boldsymbol{C}}$ down to a $\hat{d}$-dimensional subspace and therefore contain less noise. Hence, we expect the variance term to decrease constantly for larger γ and that the PCA-regression model therefore avoids the "interpolation peak" at $\gamma = 1$ which linear regression has. The results generalize Lemma 1 in Hastie et al. (2022) for the risk of direct regression models since we obtain the same form when choosing $\hat{d} = m$, i.e. no dimensionality reduction.

Asymptotics for isotropic features   Using results from random matrix theory, and Lemma 1 we derive asymptotics for the risk and parameter norm in the case of isotropic features $\boldsymbol{C} = \boldsymbol{I}_m$.

Theorem 1. *Assume isotropic features $\boldsymbol{C} = \boldsymbol{I}_m$ and choose constant $d, \hat{d}$. Then, as $m, n \to \infty$, such that $\frac{m}{n} \to \gamma$, the expected risk and parameter norm satisfy almost surely*

$$\mathbb{E}_{\boldsymbol{\epsilon}}\left[R(\hat{\boldsymbol{\theta}})\right] \to \sigma_\epsilon^2 \frac{m}{n} \int_{\bar{s}}^\infty \frac{1}{s} dF_\gamma(s) + \sigma_\epsilon^2 + \begin{cases} \boldsymbol{\beta}^\top \boldsymbol{\beta} \left(1 - \min(\hat{d}, m)/m\right) & for \quad \gamma < 1 \\ \boldsymbol{\beta}^\top \boldsymbol{\beta} \left(1 - \min(\hat{d}, n)/m\right) & for \quad \gamma > 1 \end{cases}, \tag{11}$$

$$\mathbb{E}_{\boldsymbol{\epsilon}}\left[\|\hat{\boldsymbol{\theta}}\|_2^2\right] \to \sigma_\epsilon^2 \frac{m}{n} \int_{\bar{s}}^\infty \frac{1}{s} dF_\gamma(s) + \begin{cases} \boldsymbol{\beta}^\top \boldsymbol{\beta} \min(\hat{d}, m)/m & for \quad \gamma < 1 \\ \boldsymbol{\beta}^\top \boldsymbol{\beta} \min(\hat{d}, n)/m & for \quad \gamma > 1 \end{cases}, \tag{12}$$

with F_γ the Marčenko-Pastur law (Marčenko & Pastur, 1967) and $\bar{s}$ the value in $\mathbb{R}$ that satisfies the integral equation $\frac{\hat{d}}{m} = \int_{\bar{s}}^\infty dF_\gamma$. The first term represents the variance and the last one the bias.

The proofs are in Appendices C.4, C.6. Again, we obtain the same risk when choosing $\hat{d} = m$ as for direct regression models on isotropic data, see Theorem 1 in Hastie et al. (2022). Contrary to direct regression, the PCA-model will always have a bias term since $\hat{d} < m, n$ in general.

4.2 NUMERICAL RESULTS

In this section we give numerical results for the different data generators and compare these results with those from the analysis above. We compare our PCA-regression model with 1) the learnt *direct regression* model and 2) a model that predicts always zero which we denote as *null risk*.

Isotropic features We generate $n = 400$ data points for training and testing according to our isotropic data generator (4) with $\sigma_\varepsilon^2 = 1$ and $r_\theta^2 = 1$. Each sample has $m = \gamma n$ features where we vary $\gamma \in [0.3, 20]$, i.e. from underparameterized ($\gamma < 1$) to overparameterized ($\gamma > 1$). We compute risk and parameter norm as defined in Lemma 1 and average over 200 realizations. The results are compared with analytical solutions from Theorem 1.

Figure 2: **Supervised results on isotropic data.** The solid lines are averaged simulation results; '×' mark analytical solutions from Theorem 1; '∘' is the null risk. *Left:* Risk. *Right:* Parameter norm.

Figure 2 depicts the results for different values of $\hat{d}$; we can make several observations: 1) The numerical results, i.e. the solid lines, and the analytical solutions, i.e. the '×' marks, align perfectly and therefore support our theoretical analysis. The expected decrease of the variance term and a nonzero bias term for all γ can be observed in Figure 10 where we show the bias-variance decomposition according to Theorem 1; 2) For sufficiently large $\hat{d}$ the results of the PCA model match the direct regression results in the limit of small and large γ. For isotropic data, every singular direction is equally important and the PCA requires sufficiently many components, i.e. large $\hat{d}$ to achieve reasonable results; 3) The PCA-regression model does not suffer from the singularity at $\gamma = 1$ as we predicted from Lemma 1. The PCA alleviates the bad conditioning of the matrix $\boldsymbol{X}^\top \boldsymbol{X}$ which has to be inverted for the least squares solution. Below we will see that ridge regression has a similar effect; and, 4) The parameter norm is constantly decreasing for larger γ. We observe this for all models, which implies that we obtain smooth solutions which are beneficial to avoid overfitting.

Latent variable data We use the latent variable data generator with $d = 20$, $r_\theta^2 = 1$, $\sigma_y^2 = 0$ and feature SNR $\rho_{\boldsymbol{x}} = 1$ to generate $n = 400$ training and testing data points and average over 200 realizations. The results for the risks are depicted in Figure 3 for eigenvalue decay of $\alpha = 0$ (left) and for $\alpha = 0.25$ (right). Corresponding plots for the parameter norm are in Figure 11.

Figure 3: **Supervised risk on latent variable data.** *Left:* Risk of models for data generated with feature covariance eigenvalue decay of $\alpha = 0$. *Right:* Equivalent results with $\alpha = 0.25$.

We observe for $\alpha = 0$ (left plot) if $\hat{d} \geq d$ then the PCA-regression model approaches the direct regression results for small and large γ. The plots for $\hat{d} = 20$ and $\hat{d} = 40$ overlay since both

are larger than d and capture all information. However, for $\hat{d} < d$ the solution obtained for the PCA-regression is suboptimal. Following Lemma 1, by choosing $\hat{d} < d$ we remove important eigendirections and therefore observe an increased risk. Similar conclusions can be drawn for the results for data with $\alpha > 0$ (right plot) but with less penalty on the risk for suboptimal choices of $\hat{d}$. Surprisingly in this case we observe that for a certain range of $\gamma > 1$, the optimal PCA-regression model is worse than the direct regression model. Further work is required to clarify this behavior.

Ridge regression So far, we focused on the unregularized linear regression problem. In this part we compare with the regularized ridge regression solution and λ as the ridge parameter

$$\hat{\boldsymbol{\theta}} = (\boldsymbol{X}^\top \boldsymbol{X} + \lambda \boldsymbol{I}_m)^+ \boldsymbol{X}^\top \boldsymbol{y}. \tag{13}$$

We use the same latent variable data as above without eigenvalue decay ($\alpha = 0$) and compare our PCA-regression model with $\hat{d} = d$ to solutions using different ridge parameters. The results are in Figure 4 and indicate a clear connection between both models for large values of λ. The observation is theoretically justified. Ridge regression lifts all eigenvalues $\boldsymbol{S}$ of the features by a value of λ

$$\boldsymbol{X}^\top \boldsymbol{X} + \lambda \boldsymbol{I}_m = \boldsymbol{V}^\top (\boldsymbol{S} + \lambda \boldsymbol{I}_d) \boldsymbol{V}. \tag{14}$$

Here, $\boldsymbol{V}$ are the true, non-truncated eigenvectors. In contrast, the PCA-regression model cuts the eigenvalues off at a threshold chosen by $\hat{d}$, which is clear in (7). The main difference is that with ridge regression, there is a smooth change of the risk controlled by the ridge parameter whereas with PCA-regression there is a hard cut-off.

Figure 4: **Comparison with ridge regression.** We highlight the similarity of the results obtained with large ridge parameter to our PCA-regression model. *Left:* Risk. *Right:* Parameter norm.

5   PRE-TRAINING THE PCA ENCODER

So far, we analysed the case when the complete model is trained supervised. Now we extend to the popular case of pre-training parts of the model in an unsupervised way. In this context we can view our model as a simple, linear version of large pre-trained neural networks with linear probing. Our analysis therefore yields insights to their understanding. The pre-training extension requires a generalization of our theory because we deal with different data sets of varying size.

5.1   GENERALISATION OF PROBLEM FORMULATION

First, we pre-train the PCA on a so called pre-training data set $\{\boldsymbol{x}_i\}_{i=1}^{n_p}$ without output values y_i. It can therefore only be used for unsupervised pre-training. Second, we train only the linear regression head on the PCA features with the training data set $\{\boldsymbol{x}_i, y_i\}_{i=1}^{n}$. Note that the number of samples n_p in the pre-training data set differs from the number of samples n in the training data set.

Data generator In this section, we focus on the latent variable data generator. We change our feature generation from (1) to simplify the theoretical analysis. We orthogonalise the signal $\boldsymbol{z}$ (generated by (3)) and the noise $\boldsymbol{e}$ by introducing $\boldsymbol{D}_\perp$ such that $\boldsymbol{D}^\top \boldsymbol{D}_\perp = \boldsymbol{0}$. We use the following feature generator for both, the training and pre-training data set features

$$\boldsymbol{x}_i = \boldsymbol{D}\boldsymbol{z}_i + \boldsymbol{D}_\perp \boldsymbol{e}_i. \tag{15}$$

Model The model is the same as in the supervised case. Since the PCA is performed on the pre-training data set, we rename the first $\hat{d}$ estimated eigenvectors of the feature covariance matrix from the pre-training data set to $\hat{H}$. We do so to distinguish it from the eigenvectors $\hat{V}$ estimated using the training data set as in the supervised case. Hence, in the pre-training case we obtain

$$\hat{z}_i = \hat{H}^\top x_i. \tag{16}$$

5.2 THEORETICAL ANALYSIS

As in section 4.1 we want to establish a connection to the complete training risk as fundamental model guarantee. Since we extend the setting to pre-train the encoder on a different data set, we have to deal with sample complexities for the estimation of eigenvectors in the PCA.

With the orthogonal feature generator (15), we recover the true latent variables from features

$$z_i = D^+(x_i - D_\perp e_i) = D^+ x_i. \tag{17}$$

Comparing it with the projection from the model in (16), we notice that the estimated latent space depends on how well $\hat{H}^\top$ estimates D^+. Hence, the risk analysis problem in the case with pre-training turns into a sample complexity problem of the eigenvectors. Note that $D^+ = D^\top/c^2$ with correction factor c for SNR control, see Appendix B.

Estimation of eigenvectors The sample complexity of eigenvectors is thoroughly studied by Loukas (2017). Here, we review some of their results and adapt them to our setting. The PCA loss of encoding x into the (estimated) latent space is given by

$$\mathcal{L}(D) = \mathbb{E}\left[\|x\|_2^2 - \|D^+ x\|_2^2\right] = \sum_{i=d+1}^{m} s_i, \tag{18}$$

$$\mathcal{L}(\hat{H}) = \mathbb{E}\left[\|x\|_2^2 - \|\hat{H}^\top x\|_2^2\right] = \sum_{i=1}^{m} s_i - \sum_{i=1}^{\hat{d}}\sum_{j=1}^{m} (\hat{h}_i^\top h_j)^2 s_j. \tag{19}$$

Here, s_i is the ith eigenvalue and h_i is the ith eigenvector of the true feature covariance matrix. The difference of the PCA losses, quantifies how well a sample x is projected with the estimated eigenvectors $\hat{H}$ into the latent space compared to a projection with the true eigenvectors $(D^+)^\top$.

Lemma 2. *Define the loss of projecting a sample x with D or $\hat{H}$ as in (18), (19). Then, we can write the loss difference as $\mathcal{L}(\hat{H}) - \mathcal{L}(D) = \mathbb{E}\left[\|z\|_2^2 - \|\hat{z}\|_2^2\right]$ and formulate it as*

$$\mathcal{L}(\hat{z}) - \mathcal{L}(z) = \sum_{i=1}^{\min(d,\hat{d})}\sum_{j=1}^{m}(\hat{h}_i^\top h_j)^2(s_i - s_j) + \underbrace{\sum_{i=\hat{d}}^{d} s_i}_{=0 \text{ for } \hat{d}\geq d} + \underbrace{\sum_{i=d}^{\hat{d}}\sum_{j=1}^{m}(\hat{h}_i^\top h_j)^2 s_j}_{=0 \text{ for } \hat{d}\leq d}. \tag{20}$$

The result indicates that if we have perfect encoding ($\hat{d} = d$), then only the first term remains. If also all eigenvalues are equal, then there is no loss difference and the estimation of the direction of eigenvectors $\hat{H}$ does not matter since we are dealing with the isotropic case. However, for more natural scenarios such as exponentially decaying eigenvalues, the eigenvalue difference is nonzero and correct estimation of the eigenvectors $\hat{H}$ is crucial for a small loss difference. If we are dealing with imperfect encoding, there is either an additional term due to misalignment of the estimated eigenvalues ($\hat{d} < d$) or due to encoding of noise ($\hat{d} > d$). The proof is in Appendix E.1.

Theorem 2. *Define a real $t > 0$, using Corollary 4.1 from Loukas (2017), and with $k_j^2 = s_j(s_j + \text{Tr}(C))$ from Corollary 4.3 in Loukas (2017), then we obtain the concentration inequality*

$$P\left(\mathcal{L}(\hat{H}) - \mathcal{L}(D) > t\right) \leq$$

$$\leq \frac{4}{t\,n_p}\left(\sum_{i=1}^{\min(d,\hat{d})}\sum_{j=i+1}^{m}\frac{k_j^2}{|s_i - s_j|} + \sum_{i=\hat{d}}^{d}\sum_{j=1}^{m}\frac{k_j^2 s_i}{(s_i - s_j)^2} + \sum_{i=d}^{\hat{d}}\sum_{j=1}^{m}\frac{k_j^2 s_j}{(s_i - s_j)^2}\right). \tag{21}$$

This theorem states, that in addition to the implications of Lemma 2, there are two main scenarios where we obtain a lower right hand side and therefore tighter bound. 1) When the feature covariance matrix has rapidly decaying eigenvalues, i.e. large $|s_i - s_j| \geq 0$, since $j > i$ or 2) when we have access to more pre-training samples n_p. The proof is in Appendix E.2.

Connection to the risk   We define the risk between features x and outcomes y in the same way as for the supervised case, see Lemma 1. The extension in this section is, that the model is trained on two different data sets. Therefore, we will discuss the effects of both training steps individually:

1. Step—PCA: We estimate the eigenvectors $\hat{H}$ from pre-training data and compare them to the true eigenvectors $(D^+)^\top$. Theorem 2 states that more pre-training samples yields a tighter concentration bound. Hence, the projection with $\hat{H}$ into the estimated latent space $\hat{z}$ will be close to the true latent space z with high probability for large pre-training data sets.

2. Step—Linear regression: Now two scenarios arise. 2.1) When the projection into $\hat{z}$ is almost perfect, we perform the linear regression effectively between z and y instead of between $\hat{z}$ and y. Therefore, the risk for the PCA-regression with pre-training is equal to the risk of linear regression from z to y for which Lemma 1 in Hastie et al. (2022) gives the result

$$\mathbb{E}_\epsilon \left[R(\hat{\theta}) \right] = \theta^\top \Psi \Lambda \Psi \theta + \sigma_\epsilon^2/n \operatorname{Tr}(\Lambda \hat{\Lambda}^+). \tag{22}$$

The feature covariance in this scenario is $\Lambda = \mathbb{E}\left[zz^\top \right]$ and we have the projection matrix $\Psi := I_d - \hat{\Lambda}^+\hat{\Lambda}$. 2.2) When the projection into $\hat{z}$ is imperfect due to a large difference between $\hat{H}$ and $(D^+)^\top$, then we perform the linear regression between $\hat{z}$ and y. Hence, we regress the outputs from a incorrect feature space, which increases the overall risk. Therefore, the estimation of the covariance matrix eigenvectors $\hat{H}$ is crucial, which can be improved by more pre-training data.

5.3   Numerical results

Here, we present numerical results when using pre-training. We denote the relation of pre-training samples to training samples as $\mu = \frac{n_p}{n}$ with $\mu \geq 1$ since we could use the training data set also for pre-training. We choose $d = 20$, $r_\theta^2 = 1$, $\sigma_y^2 = 0$, $\rho_x = 1$ and focus on $\hat{d} = d$ as the effects of misspecified models is elaborated in Section 4.2. We generate $n = 200$ training samples and $n_p = n\mu$ pre-training samples by varying $\mu \in [1, 10]$ and average the computed risk over 100 realizations.

Figure 5: **Pre-training risk on latent variable data.** On the x-axis we increase the number of features m and therefore the degree of overparameterization γ. On the y-axis we increase amount of pre-training n_p compared to training data n. *Left:* Risk for latent variable data generated with feature covariance eigenvalue decay of $\alpha = 0$. *Right:* Same setup but for $\alpha = 0.25$.

In Figure 5 the risk for data with two different eigenvalue decay rates are depicted. We make three main observations: 1) Horizontally for $\mu = \text{const.}$, in both plots the risk decreases similar to the supervised case in Figure 3 and therefore follows Lemma 1. 2) Vertically for $\gamma = \text{const.}$, in the right plot ($\alpha = 0.25$) the risk decreases as expected from Theorem 2 when using more pre-training samples. The effect is most significant for large overparameterization γ. 3) Vertically for $\gamma = \text{const.}$,

in the left plot ($\alpha = 0$), we notice that more pre-training data does *not* decrease the risk. Since we have perfect encoding $\hat{d} = d$ and two blocks of constant eigenvalues, the eigenvector estimation is by Lemma 2 almost perfect and barely improves with more pre-training, see Theorem 2. Therefore, using more pre-training data does also *not* improve the overall risk. The observation supports our finding that more pre-training data decreases the risk only if it improves the eigenvector estimation. Hence, the eigenvalue distribution is crucial for the necessity of pre-training.

Figure 6 shows horizontal slices of Figure 5 (right) and compares it with fully supervised models. We notice that all pre-trained models outperform the fully supervised models for $\gamma > 1$. Interestingly, in the results for $\mu = 1$ (blue solid line) we use the same amount of data to learn the PCA $n_p = n$ as in the fully supervised case (black dashed line). While for the pre-trained model we use a different data set of the same size to learn the regression, we use use the exact same data in the supervised case. We see that the loss with pre-training is lower, which indicates that using different data sets and therefore more diverse data is advantageous.

Figure 6: **Pre-training risk for different μ.** Comparing horizontal slices of Figure 5 (right, $\alpha = 0.25$) for pre-trained models with different amount of pre-training data μ to 1) a fully supervised direct regression model and 2) a fully supervised PCA-regression model, comparable to Figure 3.

6 CONCLUSION

Limitations Our proofs in the supervised case rely on random matrix theory for which we can find asymptotic results in the isotropic case. However, it is not trivial to find solutions for the general case, including our latent variable data generator, which requires more research. Similarly, an open question is how to obtain a closed form solution for the complete risk in the scenario with pre-training based on our sample complexity bounds. Furthermore, our empirical results are not validated against real-world data sets. Those experiments come with their own challenges which are outside the scope of this paper but that we believe can be important future work.

Supervised case Our theoretical analysis generalizes the results for linear regression (Hastie et al., 2022) which is a special case of PCA-regression without dimensionality reduction ($\hat{d} = m$). Without analytical solutions, Teresa et al. (2022) indicate similar observations in simulation and hence they independently support our theory. Selecting the correct latent dimension $\hat{d}$ for data from a low dimensional manifold is crucial for the risk as Lemma 2 suggests. This is in line with the discovery of latent factors in variational autoencoders from the disentanglement literature (Higgins et al., 2017; Kumar et al., 2018). While our results that PCA mitigates the "interpolation peak" due to its regularizing behavior may not surprise, they provide formal guarantees for the performance of a commonly used model on real-world data structures. Practitioners can now rely on these fundamental guarantees for model development, but more research is needed for general data structures.

Pre-training Our results from Figure 5 that a certain decay rate of the data covariance eigenvalues is necessary for pre-training to have its expected effect (more pre-training data is better) may be surprising at first. However, from Theorem 2 it becomes clear that more pre-training data only helps to improve the eigenvector estimation. If however, the eigenvectors are already estimated perfectly such as for two blocks of isotropic data (e.g. latent variable data with decay rate $\alpha = 0$), then using more pre-training data has no effect. Hence, we provide a fundamental insight into the mechanisms of pre-training which highlight that we have to be more aware of the data structure instead of following the general philosophy of adding more pre-training data. We do believe that our simple PCA-regression model is suitable for extensive studies of different phenomena of pre-training. Therefore, this study lays the groundwork for future research and opens up many questions.

REPRODUCIBILITY

Code for reproducibility is attached as Jupyter notebook in the supplementary material and will be published online upon acceptance; all simulation parameters are explained in detail in the paper and copied in the code. All of our numerical simulations are run on Intel Core i7-6850K CPUs @ 3.60GHz in a matter of minutes. The computationally most heavy experiment is for pre-training with large μ, see Figure 5 which takes for one run about 15 minutes for the fine-grained grid that we show in the paper. Averaging over multiple runs for more accurate results increase the computational cost linearly.

REFERENCES

Madhu S Advani, Andrew M Saxe, and Haim Sompolinsky. High-dimensional dynamics of generalization error in neural networks. *Neural Networks*, 132:428–446, 2020. (Cited on p. 2)

Jean-Baptiste Alayrac, Jeff Donahue, Pauline Luc, Antoine Miech, Iain Barr, Yana Hasson, Karel Lenc, Arthur Mensch, Katie Millican, Malcolm Reynolds, et al. Flamingo: a visual language model for few-shot learning. *arXiv preprint arXiv:2204.14198*, 2022. (Cited on p. 1)

Sanjeev Arora, Nadav Cohen, Noah Golowich, and Wei Hu. A convergence analysis of gradient descent for deep linear neural networks. In *International Conference on Learning Representations*, 2019. (Cited on p. 1)

Jimmy Lei Ba, Jamie Ryan Kiros, and Geoffrey E Hinton. Layer normalization. *arXiv preprint arXiv:1607.06450*, 2016. (Cited on p. 2)

Peter L Bartlett and Shahar Mendelson. Rademacher and Gaussian complexities: Risk bounds and structural results. *Journal of Machine Learning Research*, 3(Nov):463–482, 2002. (Cited on p. 2)

Peter L Bartlett, Philip M Long, Gábor Lugosi, and Alexander Tsigler. Benign overfitting in linear regression. *Proceedings of the National Academy of Sciences*, 117(48):30063–30070, 2020. (Cited on p. 2)

Mikhail Belkin, Daniel J Hsu, and Partha Mitra. Overfitting or perfect fitting? risk bounds for classification and regression rules that interpolate. *Advances in neural information processing systems*, 31, 2018. (Cited on p. 2)

Mikhail Belkin, Daniel J Hsu, Siyuan Ma, and Soumik Mandal. Reconciling modern machine-learning practice and the classical bias–variance trade-off. *Proceedings of the National Academy of Sciences*, 116(32):15849–15854, 2019. (Cited on p. 1, 2)

Leo Breiman and David Freedman. How many variables should be entered in a regression equation? *Journal of the American Statistical Association*, 78(381):131–136, 1983. (Cited on p. 2)

Rares-Darius Buhai, Yoni Halpern, Yoon Kim, Andrej Risteski, and David Sontag. Empirical study of the benefits of overparameterization in learning latent variable models. In *International Conference on Machine Learning*, pp. 1211–1219. PMLR, 2020. (Cited on p. 2)

Mathilde Caron, Ishan Misra, Julien Mairal, Priya Goyal, Piotr Bojanowski, and Armand Joulin. Unsupervised learning of visual features by contrasting cluster assignments. *Advances in Neural Information Processing Systems*, 33:9912–9924, 2020. (Cited on p. 1)

Ting Chen, Simon Kornblith, Mohammad Norouzi, and Geoffrey Hinton. A simple framework for contrastive learning of visual representations. In *International conference on machine learning*, pp. 1597–1607. PMLR, 2020. (Cited on p. 1)

Lenaic Chizat and Francis Bach. Implicit bias of gradient descent for wide two-layer neural networks trained with the logistic loss. In *Conference on Learning Theory*, pp. 1305–1338. PMLR, 2020. (Cited on p. 2)

Zeyu Deng, Abla Kammoun, and Christos Thrampoulidis. A model of double descent for high-dimensional binary linear classification. *Information and Inference: A Journal of the IMA*, 11(2):435–495, 2022. (Cited on p. 2)

Jacob Devlin, Ming-Wei Chang, Kenton Lee, and Kristina Toutanova. BERT: Pre-training of deep bidirectional transformers for language understanding. In *Proceedings of the 2019 Conference of the North American Chapter of the Association for Computational Linguistics: Human Language Technologies, Volume 1 (Long and Short Papers)*, pp. 4171–4186, Minneapolis, Minnesota, 2019. Association for Computational Linguistics. (Cited on p. 1)

Alexey Dosovitskiy, Lucas Beyer, Alexander Kolesnikov, Dirk Weissenborn, Xiaohua Zhai, Thomas Unterthiner, Mostafa Dehghani, Matthias Minderer, Georg Heigold, Sylvain Gelly, Jakob Uszkoreit, and Neil Houlsby. An image is worth 16x16 words: Transformers for image recognition at scale. In *International Conference on Learning Representations*, 2021. (Cited on p. 1)

Simon Shaolei Du, Wei Hu, Sham M. Kakade, Jason D. Lee, and Qi Lei. Few-shot learning via learning the representation, provably. In *International Conference on Learning Representations*, 2021. (Cited on p. 2)

Dheeru Dua and Casey Graff. UCI machine learning repository, 2017. URL http://archive.ics.uci.edu/ml. (Cited on p. 16)

Gintare Karolina Dziugaite and Daniel M. Roy. Computing nonvacuous generalization bounds for deep (stochastic) neural networks with many more parameters than training data. In *Proceedings of the Thirty-Third Conference on Uncertainty in Artificial Intelligence, UAI 2017, Sydney, Australia, August 11-15, 2017*. AUAI Press, 2017. (Cited on p. 2)

Stéphane d'Ascoli, Maria Refinetti, Giulio Biroli, and Florent Krzakala. Double trouble in double descent: Bias and variance (s) in the lazy regime. In *International Conference on Machine Learning*, pp. 2280–2290. PMLR, 2020. (Cited on p. 2)

Dumitru Erhan, Aaron Courville, Yoshua Bengio, and Pascal Vincent. Why does unsupervised pre-training help deep learning? In *Proceedings of the thirteenth international conference on artificial intelligence and statistics*, pp. 201–208. JMLR Workshop and Conference Proceedings, 2010. (Cited on p. 2)

Charles Fefferman, Sanjoy Mitter, and Hariharan Narayanan. Testing the manifold hypothesis. *Journal of the American Mathematical Society*, 29(4):983–1049, 2016. (Cited on p. 1)

John Geweke. Bayesian reduced rank regression in econometrics. *Journal of econometrics*, 75(1): 121–146, 1996. (Cited on p. 3)

Gauthier Gidel, Francis Bach, and Simon Lacoste-Julien. Implicit regularization of discrete gradient dynamics in linear neural networks. *Advances in Neural Information Processing Systems*, 32, 2019. (Cited on p. 1)

Sebastian Goldt, Madhu Advani, Andrew M Saxe, Florent Krzakala, and Lenka Zdeborová. Dynamics of stochastic gradient descent for two-layer neural networks in the teacher-student setup. *Advances in neural information processing systems*, 32, 2019. (Cited on p. 2)

Priya Goyal, Mathilde Caron, Benjamin Lefaudeux, Min Xu, Pengchao Wang, Vivek Mangalore Pai, Mannat Singh, Vitaliy Liptchinsky, Ishan Misra, Armand Joulin, and Piotr Bojanowski. Self-supervised pretraining of visual features in the wild. *arXiv preprint arXiv:2103.01988*, 2021. (Cited on p. 1)

Suriya Gunasekar, Blake E Woodworth, Srinadh Bhojanapalli, Behnam Neyshabur, and Nati Srebro. Implicit regularization in matrix factorization. *Advances in Neural Information Processing Systems*, 30, 2017. (Cited on p. 2)

Trevor Hastie, Andrea Montanari, Saharon Rosset, and Ryan J Tibshirani. Surprises in high-dimensional ridgeless least squares interpolation. *The Annals of Statistics*, 50(2):949–986, 2022. (Cited on p. 2, 3, 4, 8, 9, 23)

Kaiming He, Xiangyu Zhang, Shaoqing Ren, and Jian Sun. Deep residual learning for image recognition. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pp. 770–778, 2016. (Cited on p. 2)

Irina Higgins, Loic Matthey, Arka Pal, Christopher Burgess, Xavier Glorot, Matthew Botvinick, Shakir Mohamed, and Alexander Lerchner. beta-VAE: Learning basic visual concepts with a constrained variational framework. In *International Conference on Learning Representations*, 2017. (Cited on p. 9)

Sergey Ioffe and Christian Szegedy. Batch normalization: Accelerating deep network training by reducing internal covariate shift. In *International conference on machine learning*, pp. 448–456. PMLR, 2015. (Cited on p. 2)

Ian T Jolliffe. A note on the use of principal components in regression. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 31(3):300–303, 1982. (Cited on p. 2)

Nitish Shirish Keskar, Dheevatsa Mudigere, Jorge Nocedal, Mikhail Smelyanskiy, and Ping Tak Peter Tang. On large-batch training for deep learning: Generalization gap and sharp minima. In *International Conference on Learning Representations*, 2017. (Cited on p. 2)

Alex Krizhevsky, Ilya Sutskever, and Geoffrey E Hinton. Imagenet classification with deep convolutional neural networks. *Communications of the ACM*, 60(6):84–90, 2017. (Cited on p. 2)

Abhishek Kumar, Prasanna Sattigeri, and Avinash Balakrishnan. Variational inference of disentangled latent concepts from unlabeled observations. In *International Conference on Learning Representations*, 2018. (Cited on p. 9)

Ananya Kumar, Aditi Raghunathan, Robbie Matthew Jones, Tengyu Ma, and Percy Liang. Fine-tuning can distort pretrained features and underperform out-of-distribution. In *International Conference on Learning Representations*, 2022. (Cited on p. 2)

Andrew K. Lampinen and Surya Ganguli. An analytic theory of generalization dynamics and transfer learning in deep linear networks. In *International Conference on Learning Representations*, 2019. (Cited on p. 1)

Yann LeCun, Corinna Cortes, and CJ Burges. Mnist handwritten digit database. *ATT Labs [Online]. Available: http://yann.lecun.com/exdb/mnist*, 2, 2010. (Cited on p. 16)

Young Kyung Lee, Eun Ryung Lee, and Byeong U Park. Principal component analysis in very high-dimensional spaces. *Statistica Sinica*, pp. 933–956, 2012. (Cited on p. 2)

Daniel LeJeune, Hamid Javadi, and Richard Baraniuk. The implicit regularization of ordinary least squares ensembles. In *International Conference on Artificial Intelligence and Statistics*, pp. 3525–3535. PMLR, 2020. (Cited on p. 2)

Andreas Loukas. How close are the eigenvectors of the sample and actual covariance matrices? In *International Conference on Machine Learning*, pp. 2228–2237. PMLR, 2017. (Cited on p. 2, 7, 27)

V A Marčenko and L A Pastur. Distribution of eigenvalues for some sets of random matrices. *Mathematics of the USSR-Sbornik*, 1(4):457–483, 1967. (Cited on p. 4, 21)

William F Massy. Principal components regression in exploratory statistical research. *Journal of the American Statistical Association*, 60(309):234–256, 1965. (Cited on p. 3)

Song Mei and Andrea Montanari. The generalization error of random features regression: Precise asymptotics and the double descent curve. *Communications on Pure and Applied Mathematics*, 75(4):667–766, 2022. (Cited on p. 2)

Vidya Muthukumar, Kailas Vodrahalli, Vignesh Subramanian, and Anant Sahai. Harmless interpolation of noisy data in regression. *IEEE Journal on Selected Areas in Information Theory*, 1(1):67–83, 2020. (Cited on p. 2)

Preetum Nakkiran, Gal Kaplun, Yamini Bansal, Tristan Yang, Boaz Barak, and Ilya Sutskever. Deep double descent: Where bigger models and more data hurt. *Journal of Statistical Mechanics: Theory and Experiment*, 2021(12):124003, 2021. (Cited on p. 2)

Behnam Neyshabur, Ryota Tomioka, and Nathan Srebro. In search of the real inductive bias: On the role of implicit regularization in deep learning. In *ICLR (Workshop)*, 2015. URL http://arxiv.org/abs/1412.6614. (Cited on p. 2)

Thanh V Nguyen, Raymond KW Wong, and Chinmay Hegde. On the dynamics of gradient descent for autoencoders. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pp. 2858–2867. PMLR, 2019. (Cited on p. 2)

Thanh V. Nguyen, Raymond K. W. Wong, and Chinmay Hegde. Benefits of jointly training autoencoders: An improved neural tangent kernel analysis. *IEEE Transactions on Information Theory*, 67(7):4669–4692, 2021. (Cited on p. 2)

Scott Pesme, Loucas Pillaud-Vivien, and Nicolas Flammarion. Implicit bias of sgd for diagonal linear networks: a provable benefit of stochasticity. *Advances in Neural Information Processing Systems*, 34:29218–29230, 2021. (Cited on p. 1)

Adityanarayanan Radhakrishnan, Mikhail Belkin, and Caroline Uhler. Memorization in overparameterized autoencoders. In *ICML 2019 Workshop on Identifying and Understanding Deep Learning Phenomena*, 2019. (Cited on p. 2)

Adityanarayanan Radhakrishnan, Mikhail Belkin, and Caroline Uhler. Overparameterized neural networks implement associative memory. *Proceedings of the National Academy of Sciences*, 117 (44):27162–27170, 2020. (Cited on p. 2)

Colin Raffel, Noam Shazeer, Adam Roberts, Katherine Lee, Sharan Narang, Michael Matena, Yanqi Zhou, Wei Li, Peter J Liu, et al. Exploring the limits of transfer learning with a unified text-to-text transformer. *J. Mach. Learn. Res.*, 21(140):1–67, 2020. (Cited on p. 1)

Maithra Raghu, Chiyuan Zhang, Jon Kleinberg, and Samy Bengio. Transfusion: Understanding transfer learning for medical imaging. *Advances in neural information processing systems*, 32, 2019. (Cited on p. 2)

Aditya Ramesh, Mikhail Pavlov, Gabriel Goh, Scott Gray, Chelsea Voss, Alec Radford, Mark Chen, and Ilya Sutskever. Zero-shot text-to-image generation. In *International Conference on Machine Learning*, pp. 8821–8831. PMLR, 2021. (Cited on p. 1)

Andrew M Saxe, James L McClelland, and Surya Ganguli. Exact solutions to the nonlinear dynamics of learning in deep linear neural networks. In *International Conference on Learning Representations*, 2014. (Cited on p. 1)

Steffen Schneider, Alexei Baevski, Ronan Collobert, and Michael Auli. wav2vec: Unsupervised Pre-Training for Speech Recognition. In *Proc. Interspeech 2019*, pp. 3465–3469, 2019. (Cited on p. 1)

Samuel L Smith, Benoit Dherin, David Barrett, and Soham De. On the origin of implicit regularization in stochastic gradient descent. In *International Conference on Learning Representations*, 2021. (Cited on p. 2)

Christian Szegedy, Vincent Vanhoucke, Sergey Ioffe, Jon Shlens, and Zbigniew Wojna. Rethinking the inception architecture for computer vision. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pp. 2818–2826, 2016. (Cited on p. 1)

Ningyuan Teresa, David W Hogg, and Soledad Villar. Dimensionality reduction, regularization, and generalization in overparameterized regressions. *SIAM Journal on Mathematics of Data Science*, 4(1):126–152, 2022. (Cited on p. 2, 9)

Nilesh Tripuraneni, Michael Jordan, and Chi Jin. On the theory of transfer learning: The importance of task diversity. *Advances in Neural Information Processing Systems*, 33:7852–7862, 2020. (Cited on p. 2)

Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N Gomez, Łukasz Kaiser, and Illia Polosukhin. Attention is all you need. *Advances in neural information processing systems*, 30, 2017. (Cited on p. 1, 2)

Sethu Vijayakumar and Stefan Schaal. Locally weighted projection regression: An o(n) algorithm for incremental real time learning in high dimensional space. In *Proceedings of the seventeenth international conference on machine learning (ICML 2000)*, volume 1, pp. 288–293, 2000. (Cited on p. 3)

Julius Von Kügelgen, Yash Sharma, Luigi Gresele, Wieland Brendel, Bernhard Schölkopf, Michel Besserve, and Francesco Locatello. Self-supervised learning with data augmentations provably isolates content from style. *Advances in neural information processing systems*, 34:16451–16467, 2021. (Cited on p. 2)

Martin J Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge University Press, 2019. (Cited on p. 2)

Kai Wang and Diana Abbott. A principal components regression approach to multilocus genetic association studies. *Genet. Epidemiol.*, 32(2):108–118, 2008. (Cited on p. 3)

Ke Wang, Vidya Muthukumar, and Christos Thrampoulidis. Benign overfitting in multiclass classification: All roads lead to interpolation. *Advances in Neural Information Processing Systems*, 34: 24164–24179, 2021. (Cited on p. 2)

Tongzhou Wang and Phillip Isola. Understanding contrastive representation learning through alignment and uniformity on the hypersphere. In *International Conference on Machine Learning*, pp. 9929–9939. PMLR, 2020. (Cited on p. 2)

Ji Xu and Daniel J Hsu. On the number of variables to use in principal component regression. *Advances in neural information processing systems*, 32, 2019. (Cited on p. 2)

Chiyuan Zhang, Samy Bengio, Moritz Hardt, Benjamin Recht, and Oriol Vinyals. Understanding deep learning requires rethinking generalization. In *International Conference on Learning Representations*, 2017. (Cited on p. 2)

Chiyuan Zhang, Samy Bengio, Moritz Hardt, Michael C. Mozer, and Yoram Singer. Identity crisis: Memorization and generalization under extreme overparameterization. In *International Conference on Learning Representations*, 2020. (Cited on p. 2)

Roland S Zimmermann, Yash Sharma, Steffen Schneider, Matthias Bethge, and Wieland Brendel. Contrastive learning inverts the data generating process. In *International Conference on Machine Learning*, pp. 12979–12990. PMLR, 2021. (Cited on p. 2)

Appendix

CONTENTS

A   EIGENVALUE DISTRIBUTION OF REAL-WORLD DATA SETS

In Figure 7 we plot the eigenvalue distribution of four real-world data sets. Each of them has a low number of significant eigenvalues with a sharp exponential decay. For some data sets such as e.g. Steel Plates Fault there is even a low-dimensional data embedding up to about eigenvalue 12 visible. All data sets except MNIST were downloaded from the UCI Machine Learning Repository (Dua & Graff, 2017) through the OpenML interface.

Figure 7: **Eigenvalue distribution of real-world data sets.** *Top left:* Distribution for MNIST digit 0 of the test data set (LeCun et al., 2010). Notice that many of the 784 eigenvalues are almost zero. *Top right:* Complete features of the spambase data set from UCI (Dua & Graff, 2017). *Bottom left:* Breastcancer data set from UCI (Dua & Graff, 2017). *Bottom right:* Steel-plates-fault data set provided by Semeion, Research of Sciences of Communication, Via Sersale 117, 00128, Rome, Italy.

B   DETAILS ON THE DATA GENERATOR

In this appendix we concentrate without loss of generality on the latent variable data generator with orthogonal features introduced in (2) and (15). In matrix notation when collecting all features in rows we can write the data generator as

$$X = ZD^\top + ED_\perp^\top, \tag{23}$$
$$y = Z\theta^\top + \varepsilon. \tag{24}$$

Singular value and eigenvalue decomposition   Approximating the data matrix with an estimated singular values decomposition and reducing the rank to $\hat{d}$ yields

$$X = \hat{U}\hat{\Sigma}\hat{V}^\top, \tag{25}$$

with estimated singular values $\hat{\boldsymbol{\Sigma}} = \operatorname{diag}(\hat{\sigma}_1, \ldots, \hat{\sigma}_d)$. Similarly we can define the eigenvalue decomposition of the sample covariance matrix as

$$\hat{\boldsymbol{C}} = \frac{1}{n}\boldsymbol{X}^\top \boldsymbol{X} = \frac{1}{n}\hat{\boldsymbol{V}}\hat{\boldsymbol{\Sigma}}^\top \hat{\boldsymbol{\Sigma}}\hat{\boldsymbol{V}}^\top = \hat{\boldsymbol{V}}\hat{\boldsymbol{S}}\hat{\boldsymbol{V}}^\top, \tag{26}$$

with estimated eigenvalue matrix $\boldsymbol{S} = \operatorname{diag}(\hat{s}_1, \ldots, \hat{s}_d)$.

Covariance matrices   The feature covariance matrix can be written as

$$\boldsymbol{C} = \mathbb{E}\left[\boldsymbol{X}^\top \boldsymbol{X}\right] = [\boldsymbol{D} \quad \boldsymbol{D}_\perp]\,\mathbb{E}\left[\begin{bmatrix} \boldsymbol{Z}^\top \boldsymbol{Z} & \boldsymbol{Z}^\top \boldsymbol{E} \\ \boldsymbol{E}^\top \boldsymbol{Z} & \boldsymbol{E}^\top \boldsymbol{E} \end{bmatrix}\right]\begin{bmatrix} \boldsymbol{D}^\top \\ \boldsymbol{D}_\perp^\top \end{bmatrix} = \boldsymbol{V}\boldsymbol{S}\boldsymbol{V}^\top, \tag{27}$$

where $\boldsymbol{V} := [\boldsymbol{D} \quad \boldsymbol{D}_\perp]$ are the true eigenvectors—compare with the sample eigenvectors denoted by $\hat{\boldsymbol{V}}$. The eigenvalue matrix $\boldsymbol{S}$ can be written as

$$\boldsymbol{S} = \operatorname{diag}(s_1, \ldots, s_m) = \operatorname{diag}(\lambda_1, \ldots, \lambda_d, 1, \ldots, 1) = \begin{bmatrix} \boldsymbol{\Lambda} & \boldsymbol{0} \\ \boldsymbol{0} & \boldsymbol{I}_{m-d} \end{bmatrix}. \tag{28}$$

Signal-to-noise ratio control   For the orthogonal latent variable data generator we can compute the SNR $\rho_{\boldsymbol{x}}$ of the features as

$$\rho_{\boldsymbol{x}} = \frac{\mathbb{E}\left[\|\boldsymbol{D}\boldsymbol{z}\|_2^2\right]}{\mathbb{E}\left[\|\boldsymbol{E}\boldsymbol{D}_\perp\|_2^2\right]} = \frac{\operatorname{Tr}(\boldsymbol{D}\boldsymbol{\Lambda}\boldsymbol{D}^\top)}{\operatorname{Tr}(\boldsymbol{D}_\perp \boldsymbol{D}_\perp^\top)} = \frac{\operatorname{Tr}(c^2 \boldsymbol{\Lambda})}{m - d}, \tag{29}$$

since $\operatorname{Tr}(\boldsymbol{D}\boldsymbol{D}^\top) = \operatorname{Tr}(\boldsymbol{I}_{m-d})$ and $\boldsymbol{D}^\top \boldsymbol{D} = c^2 \boldsymbol{I}_d$. Here c is a correction factor which controls the SNR. We define is as

$$c = \sqrt{\frac{\rho_{\boldsymbol{x}}(m - d)}{d}}\sqrt{\frac{d}{\operatorname{Tr}(\boldsymbol{\Lambda})}}. \tag{30}$$

If non-orthogonal noise is used, then the first factor reduces to $\sqrt{\rho_{\boldsymbol{x}} m / d}$.

In the same way, we can compute the SNR ρ_y of the outputs as

$$\rho_y = \frac{\mathbb{E}\left[\|\boldsymbol{\theta}^\top \boldsymbol{z}\|_2^2\right]}{\mathbb{E}\left[\|\boldsymbol{\varepsilon}\|_2^2\right]} = \frac{\operatorname{Tr}(\boldsymbol{\theta}^\top \boldsymbol{\Lambda}\boldsymbol{\theta})}{\sigma_y^2} = \frac{r_{\boldsymbol{\theta}}^2}{\sigma_y^2}, \tag{31}$$

with $r_{\boldsymbol{\theta}}^2 = 1$ usually.

Implementation details for data generation   For our latent variable orthogonal feature generator, we generate the matrices $\boldsymbol{D}$ and $\boldsymbol{D}_\perp$ by first sampling an auxiliary random variable $\boldsymbol{A} \in \mathbb{R}^{m \times d}$ and then orthogonalizing it with a QR-decomposition

$$A_{i,j} \sim \mathcal{N}(0, 1), \tag{32}$$
$$\boldsymbol{Q}\boldsymbol{R} = \boldsymbol{A}, \tag{33}$$

where the columns of $\boldsymbol{Q}$ are orthogonal. Hence, we can define

$$\boldsymbol{D} = c\boldsymbol{Q}_{:d,:}, \tag{34}$$
$$\boldsymbol{D}_\perp = \boldsymbol{Q}_{d:,:}, \tag{35}$$

where the SNR-correction factor c is defined in (30).

In order to hold that $\mathbb{E}\left[\|\boldsymbol{\theta}^\top \boldsymbol{z}\|_2^2\right] = r_{\boldsymbol{\theta}}^2$, we generate $\boldsymbol{\theta}$ as

$$\boldsymbol{\theta} = \frac{\sqrt{r_{\boldsymbol{\theta}}^2 d}}{\sqrt{d\operatorname{Tr}(\boldsymbol{\Lambda})}} \begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix}_d. \tag{36}$$

C   PROOFS FOR THE SUPERVISED CASE

In this appendix we detail and proof the results shown in section 4.1. After stating some notation and definition in C.1, we first show the result for the linear regression solution in C.2. Subsequently, we will derive the result for the parameter norm in C.3 and its asymptotic in the isotropic case in C.4. Finally, the result for the risk is proven in C.5 and its asymptotic in the isotropic case is derived in C.6. For simplicity of notation, we will replace in the indices $\hat{d}$ by d–it is clear from context that we use $\hat{d}$ for the estimated latent space and d for the true one.

C.1   GENERAL

Note on notation in main text   While in the main paper for the derivation of the linear regression solution in (8) we denote for simplicity the estimated singular value matrix of the data $\boldsymbol{X}$ as $\hat{\boldsymbol{\Sigma}}$, here we are more precise. We distinguish between the case of $\gamma > 1$ with $m > n$ (left) and the case of $\gamma < 1$ with $m < n$ (right)

$$\hat{\boldsymbol{\Sigma}} = \left[\begin{array}{ccc|c} \hat{\sigma}_1 & & \mathbf{0} & \\ & \ddots & & \mathbf{0} \\ \mathbf{0} & & \hat{\sigma}_n & \end{array} \right] \in \mathbb{R}^{n \times m}, \quad \hat{\boldsymbol{\Sigma}} = \left[\begin{array}{ccc} \hat{\sigma}_1 & & \mathbf{0} \\ & \ddots & \\ \mathbf{0} & & \hat{\sigma}_m \\ \hline & \mathbf{0} & \end{array} \right] \in \mathbb{R}^{n \times m}. \tag{37}$$

When truncating with $d \ll \min(m,n)$ singular values, we obtain in both cases

$$\hat{\boldsymbol{\Sigma}}_d = \left[\begin{array}{ccc} \hat{\sigma}_1 & & \mathbf{0} \\ & \ddots & \\ \mathbf{0} & & \hat{\sigma}_d \\ \hline & \mathbf{0}_{n-d \times d} & \end{array} \right] \in \mathbb{R}^{n \times \hat{d}}. \tag{38}$$

In (8) we write $\hat{\boldsymbol{\Sigma}}$ instead of $\hat{\boldsymbol{\Sigma}}_d$ and $\hat{\boldsymbol{\Sigma}}^{-1}$ instead of $(\hat{\boldsymbol{\Sigma}}_d^\top \hat{\boldsymbol{\Sigma}}_d)^{-1} \hat{\boldsymbol{\Sigma}}_d^\top$ in order to not overload notation and simplify reading without harming the results. In the following we will use the notation with subscript in order to highlight the zero rows or columns. Further, below we simplify the square matrix with only the first $\hat{d}$ singular values on the diagonal as $\hat{\boldsymbol{\Sigma}}_{dd}$ to indicate its dimensions.

Sample covariance matrix   We can define the feature sample covariance matrix $\hat{\boldsymbol{C}} \in \mathbb{R}^{m \times m}$ and its Moore-Penrose pseudoinverse as

$$\hat{\boldsymbol{C}} = \frac{1}{n} \boldsymbol{X}^\top \boldsymbol{X}, \qquad \hat{\boldsymbol{C}}^+ = n(\boldsymbol{X}^\top \boldsymbol{X})^+. \tag{39}$$

Using the definition of the truncated SVD, we can rewrite the sample covariance as

$$\hat{\boldsymbol{C}} = \frac{1}{n} \hat{\boldsymbol{V}} \hat{\boldsymbol{\Sigma}}_d^\top \hat{\boldsymbol{U}}^\top \hat{\boldsymbol{U}} \hat{\boldsymbol{\Sigma}}_d \hat{\boldsymbol{V}}^\top = \frac{1}{n} \hat{\boldsymbol{V}} \hat{\boldsymbol{\Sigma}}_d^\top \hat{\boldsymbol{\Sigma}}_d \hat{\boldsymbol{V}}^\top = \frac{1}{n} \hat{\boldsymbol{V}} \hat{\boldsymbol{\Sigma}}_{dd}^2 \hat{\boldsymbol{V}}^\top = \hat{\boldsymbol{V}} \hat{\boldsymbol{S}} \hat{\boldsymbol{V}}^\top, \tag{40}$$

$$\hat{\boldsymbol{C}}^+ = n(\hat{\boldsymbol{V}} \hat{\boldsymbol{\Sigma}}_d^\top \hat{\boldsymbol{\Sigma}}_d \hat{\boldsymbol{V}}^\top)^+ = n\hat{\boldsymbol{V}} (\hat{\boldsymbol{\Sigma}}_d^\top)^+ \hat{\boldsymbol{\Sigma}}_d^+ \hat{\boldsymbol{V}}^\top = n\hat{\boldsymbol{V}} \hat{\boldsymbol{\Sigma}}_{dd}^{-2} \hat{\boldsymbol{V}}^\top = \hat{\boldsymbol{V}} \hat{\boldsymbol{S}}^{-1} \hat{\boldsymbol{V}}^\top. \tag{41}$$

The above formulation also implies the following which will be useful

$$\hat{\boldsymbol{\Sigma}}_{dd}^{-2} = \frac{1}{n} \hat{\boldsymbol{V}}^\top \hat{\boldsymbol{C}}^+ \hat{\boldsymbol{V}}. \tag{42}$$

C.2   LINEAR REGRESSION

We consider the unregularized linear regression solution between the latent variables $\hat{\boldsymbol{Z}}$ and the outcome $\boldsymbol{y}$:

$$\hat{\boldsymbol{\theta}} = (\hat{\boldsymbol{Z}}^\top \hat{\boldsymbol{Z}})^+ \hat{\boldsymbol{Z}}^\top \boldsymbol{y} \tag{43}$$

$$= (\hat{\boldsymbol{\Sigma}}_d^\top \hat{\boldsymbol{U}}^\top \hat{\boldsymbol{U}} \hat{\boldsymbol{\Sigma}}_d)^+ \hat{\boldsymbol{\Sigma}}_d^\top \hat{\boldsymbol{U}}^\top \boldsymbol{y} \tag{44}$$

with $\hat{\boldsymbol{U}}^\top \hat{\boldsymbol{U}} = \boldsymbol{I}$ and $\boldsymbol{y} = \boldsymbol{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$

$$\hat{\boldsymbol{\theta}} = (\hat{\boldsymbol{\Sigma}}_d^\top \hat{\boldsymbol{\Sigma}}_d)^+ \hat{\boldsymbol{\Sigma}}_d^\top \hat{\boldsymbol{U}}^\top (\boldsymbol{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}) \tag{45}$$

$$= (\hat{\boldsymbol{\Sigma}}_d^\top \hat{\boldsymbol{\Sigma}}_d)^+ \hat{\boldsymbol{\Sigma}}_d^\top \hat{\boldsymbol{\Sigma}} \hat{\boldsymbol{V}}^\top \boldsymbol{\beta} + (\hat{\boldsymbol{\Sigma}}_d^\top \hat{\boldsymbol{\Sigma}}_d)^+ \hat{\boldsymbol{\Sigma}}_d^\top \hat{\boldsymbol{U}}^\top \boldsymbol{\epsilon} \tag{46}$$

Where we used $\boldsymbol{X} = \hat{\boldsymbol{U}}\hat{\boldsymbol{\Sigma}}\hat{\boldsymbol{V}}^\top$. Now we combine the singular value matrices. We indicate dimensions of combined matrices. Note that $\hat{\boldsymbol{\Sigma}}_d$ and $\hat{\boldsymbol{\Sigma}}$ are of different sizes.

$$
\hat{\boldsymbol{\theta}} = \begin{bmatrix} \frac{1}{\hat{\sigma}_1^2} & & \boldsymbol{0} \\ & \ddots & \\ \boldsymbol{0} & & \frac{1}{\hat{\sigma}_d^2} \end{bmatrix}_{d \times d} \begin{bmatrix} \hat{\sigma}_1^2 & & \boldsymbol{0} & \\ & \ddots & & \boldsymbol{0} \\ \boldsymbol{0} & & \hat{\sigma}_d^2 & \end{bmatrix}_{d \times m} \hat{\boldsymbol{V}}^\top \boldsymbol{\beta} \;\; +
$$

$$
+ \begin{bmatrix} \frac{1}{\hat{\sigma}_1} & & \boldsymbol{0} & \\ & \ddots & & \boldsymbol{0} \\ \boldsymbol{0} & & \frac{1}{\hat{\sigma}_d} & \end{bmatrix}_{d \times n} \hat{\boldsymbol{U}}^\top \boldsymbol{\epsilon} \tag{47}
$$

$$
= [\boldsymbol{I}_d \quad \boldsymbol{0}] \hat{\boldsymbol{V}}^\top \boldsymbol{\beta} + [\hat{\boldsymbol{\Sigma}}_{dd}^{-1} \quad \boldsymbol{0}] \hat{\boldsymbol{U}}^\top \boldsymbol{\epsilon} \tag{48}
$$

Summarizing the matrices by truncating $\hat{\boldsymbol{V}}^\top$ and $\hat{\boldsymbol{U}}^\top$ yields the following solution for the regression parameter estimation

$$
\boxed{\hat{\boldsymbol{\theta}} = \hat{\boldsymbol{V}}_d^\top \boldsymbol{\beta} + \hat{\boldsymbol{\Sigma}}_{dd}^{-1} \hat{\boldsymbol{U}}_d^\top \boldsymbol{\epsilon}} \tag{49}
$$

C.3 PARAMETER NORM

Here, we prove the parameter norm part of Lemma 1.

Proof. In order to evaluate the parameter norm $\|\hat{\boldsymbol{\theta}}\|_2^2 = \hat{\boldsymbol{\theta}}^\top \hat{\boldsymbol{\theta}}$, we consider

$$
\hat{\boldsymbol{\theta}}^\top \hat{\boldsymbol{\theta}} = \boldsymbol{\beta}^\top \hat{\boldsymbol{V}} \hat{\boldsymbol{V}}^\top \boldsymbol{\beta} + \mathrm{Tr}(\boldsymbol{\epsilon}^\top \hat{\boldsymbol{U}} \hat{\boldsymbol{\Sigma}}_{dd}^{-1} \hat{\boldsymbol{\Sigma}}_{dd}^{-1} \hat{\boldsymbol{U}}^\top \boldsymbol{\epsilon}) + 2\boldsymbol{\beta}^\top \hat{\boldsymbol{V}} \hat{\boldsymbol{\Sigma}}_{dd}^{-1} \hat{\boldsymbol{U}}^\top \boldsymbol{\epsilon} \tag{50}
$$

where the second term is scalar and hence equal to its trace. Now, we can make use of the cyclic property of the trace. Furthermore, define $\boldsymbol{\Phi} := \hat{\boldsymbol{V}} \hat{\boldsymbol{V}}^\top$ as an orthogonal projector.

$$
\hat{\boldsymbol{\theta}}^\top \hat{\boldsymbol{\theta}} = \boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} + \mathrm{Tr}(\hat{\boldsymbol{U}}^\top \boldsymbol{\epsilon} \boldsymbol{\epsilon}^\top \hat{\boldsymbol{U}} \hat{\boldsymbol{\Sigma}}_{dd}^{-2}) + 2\boldsymbol{\beta}^\top \hat{\boldsymbol{V}} \hat{\boldsymbol{\Sigma}}_{dd}^{-1} \hat{\boldsymbol{U}}^\top \boldsymbol{\epsilon} \tag{51}
$$

Note that the properties of the orthogonal projector with $\boldsymbol{\Phi}^\top = \boldsymbol{\Phi}$ and $\boldsymbol{\Phi}\boldsymbol{\Phi} = \boldsymbol{\Phi}$ hold for our definition.

We take the expectation with respect to the noise

$$
\mathbb{E}_\epsilon \left[\hat{\boldsymbol{\theta}}^\top \hat{\boldsymbol{\theta}} \right] = \boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} + \mathrm{Tr}(\hat{\boldsymbol{U}}^\top \mathbb{E}_\epsilon \left[\boldsymbol{\epsilon} \boldsymbol{\epsilon}^\top \right] \hat{\boldsymbol{U}} \hat{\boldsymbol{\Sigma}}_{dd}^{-2}) \tag{52}
$$

$$
= \boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} + \sigma_\varepsilon^2 \mathrm{Tr}(\hat{\boldsymbol{U}}^\top \hat{\boldsymbol{U}} \hat{\boldsymbol{\Sigma}}_{dd}^{-2}) \tag{53}
$$

$$
= \boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} + \sigma_\varepsilon^2 \mathrm{Tr}(\hat{\boldsymbol{\Sigma}}_{dd}^{-2}) \tag{54}
$$

using (42) we can write

$$
\boxed{\mathbb{E}_\epsilon \left[\hat{\boldsymbol{\theta}}^\top \hat{\boldsymbol{\theta}} \right] = \boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} + \frac{\sigma_\epsilon^2}{n} \mathrm{Tr}(\hat{\boldsymbol{V}}^\top \hat{\boldsymbol{C}}^+ \hat{\boldsymbol{V}})} \tag{55}
$$

The second term uses the sample covariance matrix $\hat{\boldsymbol{C}}$ projected down on the $\hat{d}$ dimensional eigenvector space using $\hat{\boldsymbol{V}}$. $\qquad \square$

C.4 LIMITING PARAMETER NORM FOR ISOTROPIC FEATURES

Here, we prove the parameter norm part of Theorem 1.

Proof. We can analyze the two terms in (55) independently in the limit of $m, n \to \infty$ such that $\frac{m}{n} \to \gamma \in (0, \infty)$ almost surely. Furthermore we assume isotropic features $\mathrm{Cov}(\boldsymbol{x}_i) = \boldsymbol{C} = \boldsymbol{I}_m$.

First term We can write with the definition of our orthogonal projector.

$$\boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} = \boldsymbol{\beta}^\top \hat{\boldsymbol{V}} \hat{\boldsymbol{V}}^\top \boldsymbol{\beta} \tag{56}$$

We can write $\hat{\boldsymbol{V}}^\top$ with the SVD definition as $\hat{\boldsymbol{V}}^\top = \hat{\boldsymbol{\Sigma}}_d^+ \hat{\boldsymbol{U}}^\top \boldsymbol{X}$ which yields

$$\boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} = \boldsymbol{\beta}^\top \boldsymbol{X}^\top \hat{\boldsymbol{U}} \hat{\boldsymbol{\Sigma}}_d^{+\top} \hat{\boldsymbol{\Sigma}}_d^+ \hat{\boldsymbol{U}}^\top \boldsymbol{X} \boldsymbol{\beta} \tag{57}$$

For the special case of i.i.d. matrix entries $\boldsymbol{x}_i \sim \mathcal{N}(0,1)$ we have by rotational invariance that the distribution of $\boldsymbol{X}$ and $\boldsymbol{X}\boldsymbol{P}$ are equal for any orthogonal $\boldsymbol{P} \in \mathbb{R}^{m \times m}$

$$\boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} = \boldsymbol{\beta}^\top \boldsymbol{P}^\top \boldsymbol{X}^\top \hat{\boldsymbol{U}} \hat{\boldsymbol{\Sigma}}_d^{+\top} \hat{\boldsymbol{\Sigma}}_d^+ \hat{\boldsymbol{U}}^\top \boldsymbol{X} \boldsymbol{P} \boldsymbol{\beta} \tag{58}$$

Choose $\boldsymbol{P}$ such that $\boldsymbol{P}\boldsymbol{\beta} = \boldsymbol{\beta} \boldsymbol{e}_i$ with $\boldsymbol{e}_i$ as the ith normal vector and then average over $i = 1, \ldots, m$

$$\boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} = \boldsymbol{\beta}^\top \boldsymbol{\beta} \operatorname{Tr}(\boldsymbol{X}^\top \hat{\boldsymbol{U}} \hat{\boldsymbol{\Sigma}}_d^{+\top} \hat{\boldsymbol{\Sigma}}_d^+ \hat{\boldsymbol{U}}^\top \boldsymbol{X})/m \tag{59}$$

Now use again the definition of $\boldsymbol{X} = \hat{\boldsymbol{U}} \hat{\boldsymbol{\Sigma}} \hat{\boldsymbol{V}}^\top$ yields

$$\boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} = \boldsymbol{\beta}^\top \boldsymbol{\beta} \operatorname{Tr}(\hat{\boldsymbol{V}} \hat{\boldsymbol{\Sigma}}^\top \hat{\boldsymbol{U}}^\top \hat{\boldsymbol{U}} \hat{\boldsymbol{\Sigma}}_d^{+\top} \hat{\boldsymbol{\Sigma}}_d^+ \hat{\boldsymbol{U}}^\top \hat{\boldsymbol{U}} \hat{\boldsymbol{\Sigma}} \hat{\boldsymbol{V}}^\top)/m \tag{60}$$

$$= \boldsymbol{\beta}^\top \boldsymbol{\beta} \operatorname{Tr}(\hat{\boldsymbol{V}} \hat{\boldsymbol{\Sigma}}^\top \hat{\boldsymbol{\Sigma}}_d^{+\top} \hat{\boldsymbol{\Sigma}}_d^+ \hat{\boldsymbol{\Sigma}} \hat{\boldsymbol{V}}^\top)/m \tag{61}$$

Using the same arguments as for the linear regression parameter solution by combining the singular value matrices, we obtain

$$\boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} = \boldsymbol{\beta}^\top \boldsymbol{\beta} \operatorname{Tr}(\hat{\boldsymbol{V}} \hat{\boldsymbol{V}}^\top)/m \tag{62}$$

Here we again identify our orthogonal projector $\boldsymbol{\Phi}$

$$= \boldsymbol{\beta}^\top \boldsymbol{\beta} \operatorname{Tr}(\boldsymbol{\Phi})/m \tag{63}$$

Since $\boldsymbol{\Phi}$ is symmetric positive definite and since its components $\hat{\boldsymbol{V}}$ are orthogonal, all eigenvalues of $\boldsymbol{\Phi}$ are equal to one, yielding

$$\boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} = \boldsymbol{\beta}^\top \boldsymbol{\beta} \operatorname{rank}(\boldsymbol{\Phi})/m \tag{64}$$

For $m/n \to \gamma$ we have to distinguish between $\gamma < 1$ and $\gamma > 1$. Therefore, we obtain the final version for the first term of the limiting parameter norm:

$$\boxed{\boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} = \begin{cases} \boldsymbol{\beta}^\top \boldsymbol{\beta} \min(\hat{d}, m)/m & \text{for} \quad \gamma < 1 \\ \boldsymbol{\beta}^\top \boldsymbol{\beta} \min(\hat{d}, n)/m & \text{for} \quad \gamma > 1 \end{cases}} \tag{65}$$

Checking the results with considering all principal components, i.e. choosing $\hat{d} = m$ (with $m > n$ for $\gamma > 1$), we obtain

$$\boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} = \begin{cases} \boldsymbol{\beta}^\top \boldsymbol{\beta} & \text{for} \quad \gamma < 1 \\ \boldsymbol{\beta}^\top \boldsymbol{\beta} \frac{1}{\gamma} & \text{for} \quad \gamma > 1 \end{cases}$$

which is the same results as for the case of direct regression between $\boldsymbol{X}$ and $\boldsymbol{y}$.

Second term For the second term of the parameter norm we can write the trace as the sum over the eigenvalues s_i of $\hat{\boldsymbol{C}}$ but limited to the first $\hat{d}$ eigenvalues due to the projection using $\hat{\boldsymbol{V}}$

$$\frac{\sigma_\epsilon^2}{n} \operatorname{Tr}(\hat{\boldsymbol{V}}^\top \hat{\boldsymbol{C}}^+ \hat{\boldsymbol{V}}) = \sigma_\epsilon^2 \frac{1}{n} \sum_{i=1}^{\hat{d}} \frac{1}{s_i} \tag{66}$$

$$= \sigma_\epsilon^2 \frac{m}{n} \int_{s_f}^\infty \frac{1}{s} dF_{\hat{C}}(s) \tag{67}$$

where the summation is rewritten as integral over the spectral measure $F_{\hat{C}}$ of $\hat{\boldsymbol{C}}$ as and s_f is the $\hat{d}$ largest eigenvalue of $\hat{\boldsymbol{C}}$. We know that in the limit $m, n \to \infty$ the spectral measure will almost surely converge to the Marčenko-Pastur distribution F_γ which describes the distribution of the eigenvalues of $\hat{\boldsymbol{C}}$

$$\boxed{\frac{\sigma_\epsilon^2}{n} \operatorname{Tr}(\hat{\boldsymbol{V}}^\top \hat{\boldsymbol{C}}^+ \hat{\boldsymbol{V}}) \to \sigma_\epsilon^2 \frac{m}{n} \int_{s_f}^\infty \frac{1}{s} dF_\gamma(s)} \tag{68}$$

There are now two steps to solve this integral. First, we need to find out the lower integral bound s_f and second, solve the integral itself. For $s_f = -\infty$, one can use the closed form solution of the Stieltjes transformation $f(z)$ of the Marčenko-Pastur distribution and evaluate it at $z = 0$. However, there is no known closed form solution for general s_f. We therefore solve this part numerically.

Step 1 obtain the lower bound s_f: We can view the spectral measure as $F_{\hat{\Sigma}}$ as a series of m impulses at s_i with magnitude $1/m$ because the sum is normalized to 1. Since we only consider the $\hat{d}$ largest eigenvalues, we know that their sum is $\hat{d}/m$, see Figure 8a. This sum is the same as the integral from s_f over the Marčenko-Pastur distribution, see Figure 8b. Therefore we can find the lower integral bound s_f by solving

$$\frac{\hat{d}}{m} = \int_{s_f}^{\infty} dF_\gamma(s) \tag{69}$$

$$= \int_{s_f}^{s_+} \frac{1}{2\pi} \frac{\sqrt{(s_+ - s)(s - s_-)}}{\gamma s} ds \tag{70}$$

for s_f numerically with $s_\pm = (1 \pm \sqrt{\gamma})^2$ where $s_\pm$ is the lowest/highest eigenvalue. Note that $s \in [s_-, s_+]$.

Step 2 solve integral of interest: Now we can solve the integral in (68) numerically from s_f to the upper bound s_+. Therefore, we obtain a solution for the second term, which is not based on data but the properties of our data matrix, especially γ and $\hat{d}$. This concludes the full proof for the asymptotics of the parameter norm. $\qquad\square$

(a)   (b)

Figure 8: **Visualization of steps for variance term derivation**. *(a)* spectral measure impulses and and lower integral bound of integral s_f. *(b)* Marčenko-Pastur distribution (Marčenko & Pastur, 1967) for $\gamma = 0.3$ with specific lower integration bound. The area under the distribution from that threshold is equal to $\hat{d}/m$.

C.5 RISK

Here, we prove the risk part of Lemma 1.

Proof. We define the risk as the expectation over the mean squared error, and then use $y_0 = \boldsymbol{\beta}^\top \boldsymbol{x}_0 + \epsilon$, $\hat{y}_0 = \hat{\boldsymbol{\theta}}^\top \hat{\boldsymbol{z}}$ and $\hat{\boldsymbol{z}} = \hat{\boldsymbol{V}}^\top \boldsymbol{x}_0$ to obtain

$$R(\hat{\boldsymbol{\theta}}) = \mathbb{E}_{\boldsymbol{x}_0} \left[(y_0 - \hat{y}_0)^2 \right] \tag{71}$$

$$= \mathbb{E}_{\boldsymbol{x}_0} \left[(\boldsymbol{\beta}^\top \boldsymbol{x}_0 + \epsilon - \hat{y}_0)^2 \right] \tag{72}$$

$$= \mathbb{E}_{\boldsymbol{x}_0} \left[(\boldsymbol{\beta}^\top \boldsymbol{x}_0 + \epsilon - \hat{\boldsymbol{\theta}}^\top \hat{\boldsymbol{z}})^2 \right] \tag{73}$$

$$= \mathbb{E}_{\boldsymbol{x}_0} \left[(\boldsymbol{\beta}^\top \boldsymbol{x}_0 + \epsilon - \hat{\boldsymbol{\theta}}^\top \hat{\boldsymbol{V}}^\top \boldsymbol{x}_0)^2 \right] \tag{74}$$

$$= \mathbb{E}_{\boldsymbol{x}_0} \left[((\boldsymbol{\beta} - \hat{\boldsymbol{V}}\hat{\boldsymbol{\theta}})^\top \boldsymbol{x}_0 + \epsilon)^2 \right] \tag{75}$$

$$= (\boldsymbol{\beta} - \hat{\boldsymbol{V}}\hat{\boldsymbol{\theta}})^\top \boldsymbol{C} (\boldsymbol{\beta} - \hat{\boldsymbol{V}}\hat{\boldsymbol{\theta}}) + \epsilon\epsilon^\top \tag{76}$$

For simplicity we first rephrase the term in the bracket using the solution of the regression parameter estimation in (49). We re-use our orthogonal projector $\boldsymbol{\Phi} = \hat{\boldsymbol{V}}\hat{\boldsymbol{V}}^\top$ and define another orthogonal projector with $\boldsymbol{\Pi} = \boldsymbol{I}_m - \boldsymbol{\Phi}$ to obtain

$$\boldsymbol{\beta} - \hat{\boldsymbol{V}}\hat{\boldsymbol{\theta}} = \boldsymbol{\beta} - \hat{\boldsymbol{V}}(\hat{\boldsymbol{V}}^\top\boldsymbol{\beta} + \hat{\boldsymbol{\Sigma}}_{dd}^{-1}\hat{\boldsymbol{U}}^\top\epsilon) \tag{77}$$

$$= \boldsymbol{\beta} - \hat{\boldsymbol{V}}\hat{\boldsymbol{V}}^\top\boldsymbol{\beta} - \hat{\boldsymbol{V}}\hat{\boldsymbol{\Sigma}}_{dd}^{-1}\hat{\boldsymbol{U}}^\top\epsilon \tag{78}$$

$$= (\boldsymbol{I}_m - \boldsymbol{\Phi})\boldsymbol{\beta} - \hat{\boldsymbol{V}}\hat{\boldsymbol{\Sigma}}_{dd}^{-1}\hat{\boldsymbol{U}}^\top\epsilon \tag{79}$$

$$= \boldsymbol{\Pi}\boldsymbol{\beta} - \hat{\boldsymbol{V}}\hat{\boldsymbol{\Sigma}}_{dd}^{-1}\hat{\boldsymbol{U}}^\top\epsilon \tag{80}$$

Now we use this expression to take the expectation of the risk with respect to the noise. This yields

$$\mathbb{E}_\epsilon\left[R(\hat{\boldsymbol{\theta}})\right] = \boldsymbol{\beta}^\top\boldsymbol{\Pi}\boldsymbol{C}\boldsymbol{\Pi}\boldsymbol{\beta} + \mathbb{E}_\epsilon\left[\mathrm{Tr}(\epsilon^\top\hat{\boldsymbol{U}}\hat{\boldsymbol{\Sigma}}_{dd}^{-1}\hat{\boldsymbol{V}}^\top\boldsymbol{C}\hat{\boldsymbol{V}}\hat{\boldsymbol{\Sigma}}_{dd}^{-1}\hat{\boldsymbol{U}}^\top\epsilon)\right] + \mathbb{E}_\epsilon\left[\epsilon\epsilon^\top\right] \tag{81}$$

Here we made use of the Trace since the expression is scalar. Hence, we can use the cyclic property of the trace and pull the expectation inside

$$= \boldsymbol{\beta}^\top\boldsymbol{\Pi}\boldsymbol{C}\boldsymbol{\Pi}\boldsymbol{\beta} + \mathrm{Tr}(\hat{\boldsymbol{V}}^\top\boldsymbol{C}\hat{\boldsymbol{V}}\hat{\boldsymbol{\Sigma}}_{dd}^{-1}\hat{\boldsymbol{U}}^\top\mathbb{E}_\epsilon\left[\epsilon\epsilon^\top\right]\hat{\boldsymbol{U}}\hat{\boldsymbol{\Sigma}}_{dd}^{-1}) + \mathbb{E}_\epsilon\left[\epsilon\epsilon^\top\right] \tag{82}$$

with $\mathbb{E}_\epsilon\left[\epsilon\epsilon^\top\right] = \sigma_\epsilon^2$ and $\hat{\boldsymbol{U}}^\top\hat{\boldsymbol{U}} = \boldsymbol{I}$

$$= \boldsymbol{\beta}^\top\boldsymbol{\Pi}\boldsymbol{C}\boldsymbol{\Pi}\boldsymbol{\beta} + \sigma_\epsilon^2\,\mathrm{Tr}(\hat{\boldsymbol{V}}^\top\boldsymbol{C}\hat{\boldsymbol{V}}\hat{\boldsymbol{\Sigma}}_{dd}^{-2}) + \sigma_\epsilon^2 \tag{83}$$

using (42) for $\hat{\boldsymbol{\Sigma}}_{dd}^{-2}$

$$\boxed{= \boldsymbol{\beta}^\top\boldsymbol{\Pi}\boldsymbol{C}\boldsymbol{\Pi}\boldsymbol{\beta} + \frac{\sigma_\epsilon^2}{n}\,\mathrm{Tr}(\hat{\boldsymbol{V}}^\top\boldsymbol{C}\hat{\boldsymbol{V}}\hat{\boldsymbol{V}}^\top\hat{\boldsymbol{C}}^+\hat{\boldsymbol{V}}) + \sigma_\epsilon^2} \tag{84}$$

Again, similarly to the parameter norm, the second term here uses the covariance matrices projected onto the $\hat{d}$ dimensional eigenvector space. $\qquad\square$

C.6   Limiting risk for isotropic features

Here, we prove the risk part of Theorem 1.

Proof. Since we use isotropic features, we have $\boldsymbol{C} = \boldsymbol{I}_m$. Similar to the limiting parameter norm we split the analysis for the two first parts of (84).

First term: limiting bias   Using isotopic features and the definition of the orthogonal projector, we have

$$\boldsymbol{\beta}^\top\boldsymbol{\Pi}\boldsymbol{C}\boldsymbol{\Pi}\boldsymbol{\beta} = \boldsymbol{\beta}^\top\boldsymbol{\Pi}\boldsymbol{\beta} \tag{85}$$

$$= \boldsymbol{\beta}^\top(\boldsymbol{I}_m - \hat{\boldsymbol{V}}\hat{\boldsymbol{V}}^\top)\boldsymbol{\beta} \tag{86}$$

now we can use the same arguments as for the first term in the parameter norm. Namely, rewrite $\hat{\boldsymbol{V}}^\top = \hat{\boldsymbol{\Sigma}}_d^+\hat{\boldsymbol{U}}^\top\boldsymbol{X}$ in terms of $\boldsymbol{X}$, assume $\boldsymbol{x}_i \sim \mathcal{N}(0,1)$ and by rotation invariance the distribution of $\boldsymbol{X}$ and $\boldsymbol{X}\boldsymbol{P}$ are equal, where $\boldsymbol{P}$ is any orthogonal matrix. Then we choose $\boldsymbol{P}\boldsymbol{\beta} = \beta\boldsymbol{e}_i$ and average over all $i = 1, \dots, m$.

$$= \boldsymbol{\beta}^\top\left(\boldsymbol{I}_m - \boldsymbol{P}^\top\boldsymbol{X}^\top\hat{\boldsymbol{U}}\hat{\boldsymbol{\Sigma}}_d^{+\top}\hat{\boldsymbol{\Sigma}}_d^+\hat{\boldsymbol{U}}^\top\boldsymbol{X}\boldsymbol{P}\right)\boldsymbol{\beta} \tag{87}$$

$$= \boldsymbol{\beta}^\top\boldsymbol{\beta}\left(1 - \mathrm{Tr}(\boldsymbol{X}^\top\hat{\boldsymbol{U}}\hat{\boldsymbol{\Sigma}}_d^{+\top}\hat{\boldsymbol{\Sigma}}_d^+\hat{\boldsymbol{U}}^\top\boldsymbol{X})/m\right) \tag{88}$$

$$= \boldsymbol{\beta}^\top\boldsymbol{\beta}\left(1 - \mathrm{Tr}(\hat{\boldsymbol{V}}\hat{\boldsymbol{V}}^\top)/m\right) \tag{89}$$

$$= \boldsymbol{\beta}^\top\boldsymbol{\beta}\left(1 - \mathrm{Tr}(\boldsymbol{\Phi})/m\right) \tag{90}$$

$$= \boldsymbol{\beta}^\top\boldsymbol{\beta}\left(1 - \mathrm{rank}(\boldsymbol{\Phi})/m\right) \tag{91}$$

in the limit of $m, n \to \infty$ we have $\frac{m}{n} \to \gamma$ almost surely. We therefore obtain the final solution for the limiting bias as

$$
\boxed{\boldsymbol{\beta}^\top \boldsymbol{\Pi} \boldsymbol{C} \boldsymbol{\Pi} \boldsymbol{\beta} = \begin{cases} \boldsymbol{\beta}^\top \boldsymbol{\beta} \left(1 - \min(\hat{d}, m)/m \right) & \text{for} \quad \gamma < 1 \\ \boldsymbol{\beta}^\top \boldsymbol{\beta} \left(1 - \min(\hat{d}, n)/m \right) & \text{for} \quad \gamma > 1 \end{cases}}
\tag{92}
$$

Again, checking the results with considering all principal components, i.e. choosing $\hat{d} = m$ (with $m > n$ for $\gamma > 1$), we obtain

$$
\boldsymbol{\beta}^\top \boldsymbol{\Phi} \boldsymbol{\beta} = \begin{cases} 0 & \text{for} \quad \gamma < 1 \\ \boldsymbol{\beta}^\top \boldsymbol{\beta} \left(1 - \frac{1}{\gamma} \right) & \text{for} \quad \gamma > 1 \end{cases}
$$

which is the same results as for the case of direct regression between $\boldsymbol{X}$ and $\boldsymbol{y}$.

Second term: limiting variance   Using isotropic features we have

$$
\frac{\sigma_\epsilon^2}{n} \operatorname{Tr}(\hat{\boldsymbol{V}}^\top \boldsymbol{C} \hat{\boldsymbol{V}} \hat{\boldsymbol{V}}^\top \hat{\boldsymbol{C}}^+ \hat{\boldsymbol{V}}) = \frac{\sigma_\epsilon^2}{n} \operatorname{Tr}(\hat{\boldsymbol{V}}^\top \hat{\boldsymbol{C}}^+ \hat{\boldsymbol{V}})
\tag{93}
$$

this is the same form as the second term for the parameter norm and therefore yields the same numeric solution by solving

$$
\boxed{\frac{\sigma_\epsilon^2}{n} \operatorname{Tr}(\hat{\boldsymbol{V}}^\top \boldsymbol{C} \hat{\boldsymbol{V}} \hat{\boldsymbol{V}}^\top \hat{\boldsymbol{C}}^+ \hat{\boldsymbol{V}}) = \sigma_\epsilon^2 \frac{m}{n} \int_{s_f}^\infty \frac{1}{s} dF_\gamma(s)}
\tag{94}
$$

$\square$

D   ADDITIONAL NUMERICAL RESULTS FOR SUPERVISED CASE

Isotropic data   In Figure 9 we extend Figure 2. We additionally show the results for the PCA-regression model with $\hat{d} = m$, which corresponds to a PCA without compression and therefore a direct regression between input $\boldsymbol{x}$ and output y. We compare the analytical results from Theorem 1 of the PCA-model to 1) the direction regression model from simulations (solid line) and 2) the analytical solution for direct regression of isotropic data from Hastie et al. (2022). The authors give the solution for the risk in their Lemma 1 and we derive the parameter norm in the same way:

$$
\mathbb{E}_\epsilon \left[R(\hat{\boldsymbol{\theta}}) \right] \to \sigma_\epsilon^2 + \begin{cases} 0 + \sigma_\epsilon^2 \frac{\gamma}{1-\gamma} & \text{for} \quad \gamma < 1 \\ \boldsymbol{\beta}^\top \boldsymbol{\beta} \left(1 - \frac{1}{\gamma} \right) + \sigma_\epsilon^2 \frac{1}{\gamma-1} & \text{for} \quad \gamma > 1 \end{cases},
\tag{95}
$$

$$
\mathbb{E}_\epsilon \left[\|\hat{\boldsymbol{\theta}}\|_2^2 \right] \to \begin{cases} \boldsymbol{\beta}^\top \boldsymbol{\beta} + \sigma_\epsilon^2 \frac{\gamma}{1-\gamma} & \text{for} \quad \gamma < 1 \\ \boldsymbol{\beta}^\top \boldsymbol{\beta} \frac{1}{\gamma} + \sigma_\epsilon^2 \frac{1}{\gamma-1} & \text{for} \quad \gamma > 1 \end{cases},
\tag{96}
$$

We see in Figure 9 that the theory form Hastie et al. (2022) for direct regression, the numerical results for direct regression and our PCA-regression results without compression ($\hat{d} = m$) match and therefore further support our theory.

Complementary to Figure 2, we can decompose the risk and parameter norm according to Theorem 1 in a bias and variance term. The results for this decomposition are shown in Figure 10. We can see that the bias term is nonzero for all γ and increases for larger γ. Further, we observe a decrease of the variance term. In contrast, in the direct regression model, the variance term reaches a peak at $\gamma = 1$ and therefore forms the classical bias-variance decomposition trade-off for $\gamma < 1$.

Latent variable data   Complementary to the results of the risk for $\alpha = 0$ and $\alpha = 0.25$ in Figure 3, we show the results for the parameter norm in Figure 11. We observe that similarly to the isotropic case, the parameter norm decreases monotonically for larger γ. This indicates simpler solution for larger γ also in the latent variable data case.

In Figure 12 we show the results for risk and parameter norm for different values of σ_y^2. In the main text we only present results for $\sigma_y^2 = 0$ but the results here show that our conclusion hold also for additive output noise. Of course, the associated risk increases with the noise level but interestingly, the found solution as the same parameter norm.

Figure 9: **Supervised results on isotropic data.** We compare analytical solutions from Theorem 1 for the PCA-regression without compression ($\hat{d} = m$) with 1) analytical solution for direct regression and 2) simulations for direct regression

Figure 10: **Bias-variance decomposition of supervised results on isotropic data.** Same as Figure 2 but decomposed in the bias and variance terms of Theorem 1. In all plots the direct regression results are given as comparison. *Left:* Risk. *Right:* Parameter norm. *Top:* Bias terms. *Bottom:* Variance terms.

E PROOFS FOR THE CASE WITH PRE-TRAINING

This appendix derives and proofs the results of section 5.2. First, we derive the results for the estimation of the eigenvectors with the PCA loss in E.1. Then, we show derive the concentration inequality for the PCA loss in E.2.

Figure 11: **Supervised parameter norm on latent variable data**. *Left:* Parameter norm of models for data generated with $\alpha = 0$. *Right:* Equivalent results with $\alpha = 0.25$. This figure complements Figure 3

Figure 12: **Supervised results for different σ_y^2.** We have the same experimental setup as in Figure 3 with $\alpha = 0$ but vary the amount of output noise. Full lines show the PCA-model with $\hat{d} = d$; dashed lines show the direct regression model for the specific noise level.

E.1 ESTIMATION OF EIGENVECTORS

Here, we prove Lemma 2 for the loss difference for the projection into the true or estimated latent space from (20). We will look at both losses induced by the two projections separately,

Proof. First, define the loss for the projection into the true latent space as in (18)

$$\mathcal{L}(\boldsymbol{D}) = \mathbb{E}\left[\|\boldsymbol{x}\|_2^2 - \|\boldsymbol{D}^+\boldsymbol{x}\|_2^2\right]. \tag{97}$$

We can write the second term as the following where we use the cyclic property of the trace on scalars and apply the expectation on $\boldsymbol{x}\boldsymbol{x}^\top$

$$\mathbb{E}\left[\|\boldsymbol{D}^+\boldsymbol{x}\|_2^2\right] = \mathbb{E}\left[\boldsymbol{x}^\top(\boldsymbol{D}^+)^\top\boldsymbol{D}^+\boldsymbol{x}\right] \tag{98}$$

$$= \mathbb{E}\left[\mathrm{Tr}(\boldsymbol{x}^\top(\boldsymbol{D}^+)^\top\boldsymbol{D}^+\boldsymbol{x})\right] \tag{99}$$

$$= \mathrm{Tr}(\boldsymbol{D}^+\boldsymbol{V}\boldsymbol{S}\boldsymbol{V}^\top(\boldsymbol{D}^+)^\top) \tag{100}$$

Since $\boldsymbol{V} = [\boldsymbol{D} \quad \boldsymbol{D}_\perp]$ defined in (27), we obtain

$$\mathbb{E}\left[\|\boldsymbol{D}^+\boldsymbol{x}\|_2^2\right] = \mathrm{Tr}\left([\boldsymbol{I}_d \quad \boldsymbol{0}_{d\times m-m}]\,\boldsymbol{S}\begin{bmatrix}\boldsymbol{I}_d \\ \boldsymbol{0}_{m-m\times d}\end{bmatrix}\right) \tag{101}$$

$$= \mathrm{Tr}(\boldsymbol{\Lambda}) = \sum_{i=1}^{d} s_i \tag{102}$$

Hence, the loss $\mathcal{L}(\boldsymbol{D})$ becomes

$$\mathcal{L}(\boldsymbol{D}) = \sum_{i=1}^{m} s_i - \sum_{i=1}^{d} s_i = \sum_{i=d+1}^{m} s_i = \sum_{i=d+1}^{m} s_i \tag{103}$$

Second, define the loss for the projection into the estimated latent space as in (19)

$$\mathcal{L}(\hat{\boldsymbol{H}}) = \mathbb{E}\left[\|\boldsymbol{x}\|_2^2 - \|\hat{\boldsymbol{H}}^\top \boldsymbol{x}\|_2^2\right]. \tag{104}$$

Again, we can write the second term as the following by using the same arguments as for the first term

$$\mathbb{E}\left[\|\hat{\boldsymbol{H}}\boldsymbol{x}\|_2^2\right] = \mathrm{Tr}(\hat{\boldsymbol{H}}^\top \boldsymbol{V}\boldsymbol{S}\boldsymbol{V}^\top \hat{\boldsymbol{H}}) = \sum_{i=1}^{\hat{d}} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 s_j. \tag{105}$$

The last equality holds by switching to vector notation where factors can be combined to squared terms. Hence, the loss $\mathcal{L}(\hat{\boldsymbol{H}})$ becomes

$$\mathcal{L}(\hat{\boldsymbol{H}}) = \sum_{i=1}^{m} s_i - \sum_{i=1}^{\hat{d}} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 s_j. \tag{106}$$

Combining both solutions, the loss difference yields

$$\mathcal{L}(\hat{\boldsymbol{H}}) - \mathcal{L}(\boldsymbol{D}) = \sum_{i=d+1}^{m} s_i - \sum_{i=1}^{m} s_i + \sum_{i=1}^{\hat{d}} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 s_j \tag{107}$$

$$= \sum_{i=1}^{d} s_i + \sum_{i=1}^{\hat{d}} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 s_j \tag{108}$$

We can multiply the first term by $1 = \|\boldsymbol{v}_j\|_2^2 = \boldsymbol{v}_j^\top \boldsymbol{I}\boldsymbol{v}_j = \boldsymbol{v}_j^\top \hat{\boldsymbol{h}}_i \hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j = \|\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j\|_2^2$,

$$\mathcal{L}(\hat{\boldsymbol{H}}) - \mathcal{L}(\boldsymbol{D}) = \sum_{i=1}^{d} (\hat{\boldsymbol{h}}_i \boldsymbol{v}_j)^2 s_i + \sum_{i=1}^{\hat{d}} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 s_j \tag{109}$$

Now we can combine terms but have to be careful due to different end indices of the sum

$$\mathcal{L}(\hat{\boldsymbol{H}}) - \mathcal{L}(\boldsymbol{D}) = \begin{cases} \sum_{i=1}^{\hat{d}} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 (s_i - s_j) + \sum_{i=\hat{d}}^{d} s_i & \text{for} \quad \hat{d} < d \\ \sum_{i=1}^{\hat{d}} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 (s_i - s_j) & \text{for} \quad \hat{d} = d \\ \sum_{i=1}^{d} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 (s_i - s_j) + \sum_{i=d}^{\hat{d}} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 s_j & \text{for} \quad \hat{d} > d \end{cases} \tag{110}$$

which we can combine into

$$\mathcal{L}(\hat{\boldsymbol{H}}) - \mathcal{L}(\boldsymbol{D}) = \sum_{i=1}^{\min(d,\hat{d})} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 (s_i - s_j) + \underbrace{\sum_{i=\hat{d}}^{d} s_i}_{=0 \text{ for } \hat{d} \ge d} + \underbrace{\sum_{i=d}^{\hat{d}} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 s_j}_{=0 \text{ for } \hat{d} \le d}. \tag{111}$$

which concludes the proof. $\qquad\square$

E.2 CONCENTRATION INEQUALITY

Here, we prove the concentration inequality presented in Theorem 2.

Proof. Write the loss difference (20) with the same argument as in its derivation by including the factor $\|\boldsymbol{v}_j\|_2^2 = 1$ to the second summation yields

$$\mathcal{L}(\hat{\boldsymbol{H}}) - \mathcal{L}(\boldsymbol{D}) = \sum_{i=1}^{\min(d,\hat{d})} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 (s_i - s_j) + \sum_{i=\hat{d}}^{d} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 s_i + \sum_{i=d}^{\hat{d}} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 s_j \tag{112}$$

Notice that $s_i - s_j \geq 0$ for $j > i$ and $s_i - s_j \leq 0$ for $j < i$. Therefore, we can upper bound it by removing the negative indices from the first summation

$$\mathcal{L}(\hat{\boldsymbol{H}}) - \mathcal{L}(\boldsymbol{D}) \leq \sum_{i=1}^{\min(d,\hat{d})} \sum_{j>i}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 |s_i - s_j| + \sum_{i=\hat{d}}^{d} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 s_i + \sum_{i=d}^{\hat{d}} \sum_{j=1}^{m} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 s_j \tag{113}$$

We simplify by denoting the three terms as

$$\mathcal{L}(\hat{\boldsymbol{H}}) - \mathcal{L}(\boldsymbol{D}) = a + b + c \tag{114}$$

Now we define the probability that this upper bound on the loss difference is larger than a chosen real t. We can upper bound this expression by applying the union bound

$$P(a + b + c > t) \leq P(a > t) + P(b > t) + P(c > t) \tag{115}$$

Recall Corollary 4.1 from Loukas (2017): We have that for any weights w_{ij} and real $t > 0$ that

$$P\left(\sum_{i \neq j} w_{ij} (\hat{\boldsymbol{h}}_i^\top \boldsymbol{v}_j)^2 > t\right) \leq \sum_{i \neq j} \frac{4 w_{ij} k_j^2}{n_p t (s_i - s_j)^2} \tag{116}$$

where $k_j^2 = \mathbb{E}\left[\|\boldsymbol{x}\boldsymbol{x}^\top \boldsymbol{v}_j\|_2^2\right] - s_j^2$ and $w_{ij} \neq 0$ when $s_i \neq s_j$ and $sgn(s_i - s_j) 2 s_i > sgn(s_i - s_j)(s_i + s_j)$.

In accordance with this Corollary, we define

$$w_{ij} = \begin{cases} |s_i - s_j| & \text{for} \quad i \leq \min(d, \hat{d}), j > i \\ s_i & \text{for} \quad \hat{d} \leq i \leq d, \forall j \\ s_j & \text{for} \quad d \leq i \leq \hat{d}, \forall j \\ 0 & \text{otherwise} \end{cases} \tag{117}$$

Hence, we obtain

$$P\left(\mathcal{L}(\hat{\boldsymbol{H}}) - \mathcal{L}(\boldsymbol{D}) > t\right) \leq$$
$$\leq \frac{4}{t\, n_p} \left(\sum_{i=1}^{\min(d,\hat{d})} \sum_{j=i+1}^{m} \frac{k_j^2}{|s_i - s_j|} + \sum_{i=\hat{d}}^{d} \sum_{j=1}^{m} \frac{k_j^2 s_i}{(s_i - s_j)^2} + \sum_{i=d}^{\hat{d}} \sum_{j=1}^{m} \frac{k_j^2 s_j}{(s_i - s_j)^2} \right), \tag{118}$$

with $k_j^2 = s_j(s_j + \text{Tr}(\boldsymbol{C}))$ from Corollary 4.3 in Loukas (2017). This Corollary holds for our latent variable data generator. This concludes the proof. □