Outlier-Robust Sparse Mean Estimation for Heavy-Tailed Distributions

Ilias Diakonikolas
University of Wisconsin-Madison
`ilias@cs.wisc.edu`

Daniel M. Kane
University of California, San Diego
`dakane@cs.ucsd.edu`

Jasper C.H. Lee
University of Wisconsin-Madison
`jasper.lee@wisc.edu`

Ankit Pensia
University of Wisconsin-Madison
`ankitp@cs.wisc.edu`

Abstract

We study the fundamental task of outlier-robust mean estimation for heavy-tailed distributions in the presence of sparsity. Specifically, given a small number of corrupted samples from a high-dimensional heavy-tailed distribution whose mean μ is guaranteed to be sparse, the goal is to efficiently compute a hypothesis that accurately approximates μ with high probability. Prior work had obtained efficient algorithms for robust sparse mean estimation of light-tailed distributions. In this work, we give the first sample-efficient and polynomial-time robust sparse mean estimator for heavy-tailed distributions under mild moment assumptions. Our algorithm achieves the optimal asymptotic error using a number of samples scaling logarithmically with the ambient dimension. Importantly, the sample complexity of our method is optimal as a function of the failure probability τ, having an *additive* $\log(1/\tau)$ dependence. Our algorithm leverages the stability-based approach from the algorithmic robust statistics literature, with crucial (and necessary) adaptations required in our setting. Our analysis may be of independent interest, involving the delicate design of a (non-spectral) decomposition for positive semi-definite matrices satisfying certain sparsity properties.

1   Introduction

One of the most fundamental problem setups in statistics is as follows: given n i.i.d. samples drawn from an unknown distribution P chosen arbitrarily from some known distribution family $\mathcal{P}$, infer some particular property of P from the data. This generic model captures a range of statistical problems of interest, for example, parameter estimation (such as the mean and (co)variance of P), as well as hypothesis testing. While long lines of work have given us a deep understanding on the statistical and computational possibilities and limits on these problems, these results are not always applicable in real-world settings due to (i) modelling issues, that the underlying distribution P might not actually be in the known family $\mathcal{P}$ but only being close to it, and (ii) the fact that the n samples supplied might be corrupted, for example by nefarious actors in high-stakes applications [ABH$^+$72].

The field of *robust statistics* aims to design estimators and testers that can tolerate up to a *constant* fraction of corrupted samples, independent of the potentially high dimensionality of the data [Tuk60, HR09]. Classical works in the field have identified and resolved the statistical limits of problems in this setup, both in terms of constructing estimators and proving impossibility results [Yat85, DL88, DG92, HR09]. However, the proposed estimators were not computationally efficient, often requiring exponential time to compute either in the number of samples or the number of dimensions [HR09].

36th Conference on Neural Information Processing Systems (NeurIPS 2022).

A recent line of work, originating in the computer science community, has developed the subfield of *algorithmic* robust statistics, aiming to design estimators that not only attain tight statistical guarantees, but are also computable in polynomial time. This line of research has provided computationally and statistically efficient estimators in a variety of problem settings (e.g., mean estimation, covariance estimation, and linear regression) under different assumptions (e.g., the distribution might be assumed to be (sub-)Gaussian, or can be heavy tailed); see [DK19] for a recent survey of results.

The focus of this paper is the robust mean estimation problem under sparsity constraints on the mean vector. Sparsity is an important structural constraint that is both relevant in practice, especially in the face of increasing dimensionality of modern data, and extensively studied for statistical estimation (see, e.g., the books [HTW15, EK12, van16]). In the specific context of robust sparse mean estimation, prior works have studied the case where the underlying distribution has light-tails, e.g., sub-exponential tails [BDLS17, DKK⁺19b, CDG⁺, DKK⁺22]. In particular, the case of a spherical Gaussian distribution is now rather well-understood both in terms of the optimal information-theoretic estimation error, as well as the conjectured *computational-statistical tradeoff* — namely, that there is a gap between the statistical performance of computationally efficient and inefficient estimators [DKS17, BB20]. In this work, we initiate the investigation of robust sparse mean estimation for *heavy-tailed* distributions, under only mild moment assumptions. Our main result is the first computationally efficient robust mean estimator in the heavy-tailed setting which leverages sparsity to reduce sample complexity from depending polynomially on the dimensionality to a logarithmic dependence. Importantly, our algorithm also achieves the optimal dependence on the failure probability τ as it tends to 0; see the next two subsections for further discussion.

1.1 Problem Setup

We first define the input contamination model before formally stating the statistical problem.

Definition 1.1 (Strong Contamination Model). *Given a corruption parameter $\epsilon \in (0, 1/2)$ and a distribution P on uncorrupted samples, an algorithm takes samples from P with ϵ-contamination as follows: (i) The algorithm specifies the number n of samples it requires. (ii) n i.i.d. samples from P are drawn but not yet shown to the algorithm. (iii) An arbitrarily powerful adversary then inspects the entirety of the n i.i.d. samples, before deciding to replace any subset of $\lceil \epsilon n \rceil$ samples with arbitrarily corrupted points, and returning the modified set of n samples to the algorithm.*

Define the $\ell_{2,k}$-norm of a vector v, denoted by $\|v\|_{2,k}$, as the ℓ_2-norm of the largest k entries of a vector v in magnitude. The goal is to estimate the mean vector in this sparse norm.

Problem 1. *Fix a corruption parameter $\epsilon \in (0, 1/2)$, error parameter $\delta > 0$, failure probability $\tau \in (0, 1)$, and distribution family $\mathcal{D}$ over $\mathbb{R}^d$. Suppose we have access to ϵ-contaminated samples drawn from an unknown distribution $P \in \mathcal{D}$ with mean μ. The problem is to compute an estimate $\hat{\mu}$ such that $\|\hat{\mu} - \mu\|_{2,k}$ is upper bounded by error δ with probability at least $1 - \tau$ over n samples. The goal is then to give an estimator with the minimal sample complexity $n(k, \epsilon, \delta, \tau)$.*

The above problem is slightly more general than sparse mean estimation in the following sense. To estimate a k-sparse mean vector μ to error δ, it suffices (see [CDG⁺, Lemma 3.2]) to: 1) get an estimate $\tilde{\mu}$ with $\|\tilde{\mu} - \mu\|_{2,k} \leq \delta/3$, and 2) round $\tilde{\mu}$ to the k entries with the largest magnitude, and zero out all the other entries. The result of our paper solves robust mean estimation in the $\ell_{2,k}$ norm.

A key aspect of robust statistics is that, depending on the distribution family $\mathcal{D}$ we consider, the above problem is generally not solvable for all error parameters $\delta > 0$. This work focuses on sparse mean estimation for *heavy-tailed* distributions, where a commonly used model for heavy-tailedness is imposing only the mild assumption that the covariance is bounded by I, without any further tail assumptions (see Section 1.4 for more discussion). Even when $d = 1$ and even when there are infinitely many samples [DK19], it is known that in the heavy-tailed setting, the minimum δ achievable is in the order of $\sqrt{\epsilon}$. This immediately implies the same lower bound of $\Omega(\sqrt{\epsilon})$ for the minimum achievable δ in Problem 1.

Before discussing the algorithmic results in this paper, we first state known information-theoretic bounds on the sample complexity that applies to all estimators, efficient or not, for Problem 1 on distributions with covariance bounded by I, and for $\delta = \Theta(\sqrt{\epsilon})$.

Fact 1.2 (Information-theoretic sample complexity: computationally-inefficient). *In Problem 1, for the distribution family $\mathcal{D}_2$ which is the set of distributions with covariance $\Sigma \preceq I$, and for $\delta = \Theta(\sqrt{\epsilon})$,*

we have that $n(k, \epsilon, \delta, \tau) \asymp (k \log(d/k) + \log(1/\tau))/\epsilon$. That is, all algorithms need these many samples and there exists a (computationally-inefficient) estimator with this sample complexity. The upper bound is from [Dep20a, PBR20], and the lower bound follows from [LM19b] even in the absence of outliers (see also Footnote 2 in [DL22a]) and even when we restrict to the distribution family $\mathcal{D}_{Gaussian}$ which is the set of the Gaussian distributions with identity covariance.

An interesting aspect of robust sparse mean estimation is that there is a conjectured statistical-computational tradeoff, namely that efficient algorithms require a qualitatively larger sample complexity than inefficient algorithms. There is evidence (in the form of SQ lower bounds and reduction-based hardness) that all efficient algorithms have a quadratically-worse dependence on k, that is, even for constant ϵ, δ, τ, and $\mathcal{D}_{Gaussian}$ being identity-covariance Gaussians in Problem 1, the sample complexity of all efficient algorithms is at least $\tilde{\Omega}(k^2)$, as opposed to $\tilde{O}(k)$ in Fact 1.2. See [DKS17, BB20] for a detailed discussion.

Both the information-theoretic bound and the conjectured computational lower bound serve as benchmarks for our algorithm to match.

1.2 Our Result

We now state the results of this paper, before discussing in the next section our algorithmic approach and also the assumptions required in the results.

Our main result is a computationally efficiently robust mean estimator in the $\ell_{2,k}$ norm, with performance matching the conjectured computational-statistical tradeoff, under the standard heavy-tailed assumption that the covariance $\Sigma \preceq I$ and the additional mild assumption that the 4^{th} moment is bounded in all axis directions. In the following (and the rest of the paper), we use the notation $a \gg b$ for $a, b \in \mathbb{R}$ to mean there exists a sufficiently large constant c with $a \geq cb$.

Theorem 1.3 (Our result: computationally efficient). *Let $\epsilon \in (0, \epsilon_0)$ for small constant $\epsilon_0 > 0$. Let P be a multivariate distribution over $\mathbb{R}^d$, where the mean and covariance of P are μ and Σ respectively. Suppose $\Sigma \preceq I$ and further suppose that for all $j \in [d]$, $\mathbb{E}[(X_j - \mu_j)^4] = O(1)$. Then, there is an algorithm such that, on input (i) the corruption parameter ϵ, (ii) the failure probability τ, (iii) the sparsity parameter k, and (iv) T, an ϵ-corrupted set of $n \gg (k^2 \log d + \log(1/\tau))/\epsilon)$ i.i.d. samples from P, it outputs $\hat{\mu}$ satisfying $\|\hat{\mu} - \mu\|_{2,k} = O(\sqrt{\epsilon})$ with probability $1 - \tau$ in $\text{poly}(n, d)$ time.*

Phrased in a slightly different language, when our estimator is given a sufficiently large number n of ϵ-corrupted samples, it outputs an estimate $\hat{\mu}$ satisfying $\|\hat{\mu} - \mu\|_{2,k} = O\left(\sqrt{\frac{k^2 \log d}{n}} + \sqrt{\epsilon} + \sqrt{\frac{\log(1/\tau)}{n}}\right)$ with probability $1 - \tau$. We note also the guarantees of our algorithm remain the same under a weaker assumption on Σ: we need only $\|\Sigma\|_{\mathcal{X}_k} \leq 1$ ($\|\cdot\|_{\mathcal{X}_k}$ defined in Definition 1.4) instead of spectral norm. Informally, the $\mathcal{X}_k$ norm of a square matrix A is a convex relaxation of finding the maximum of $v^\top A v$ over k-sparse vectors v. See Theorem D.12 for the stronger version of the main result, which assumes only that $\|\Sigma\|_{\mathcal{X}_k} \leq 1$.

As outlined above, the dependence of our sample complexity result on k is tight with respect to the conjectured lower bound for efficient algorithms, and its dependence on τ and ϵ are also tight with respect to the information-theoretic lower bounds, even in the Gaussian case. In terms of the smallest achievable asymptotic error (even given infinitely many samples), we show in Lemma E.1 that, even after adding the mild axis-wise 4^{th} moment assumption in Theorem 1.3, the asymptotic error remains lower bounded by $\Omega(\sqrt{\epsilon})$ when k is sufficiently large. The restriction on k is very mild, and covers most parameter regimes of interest.

We also note that the sample complexity has a dependence on the failure probability that is $\log 1/\tau$, and importantly, this is an additive term in the complexity instead of multiplicative. This additive dependence was non-trivial to achieve even in the optimal rates for heavy-tailed mean estimation in the non-robust (and non-sparse) setting. See the [LM19a] survey for a more detailed discussion.

1.3 Our Approach

The algorithm we propose fits into the stability-based filtering approach that was first proposed by [DKK+16] (also see the recent survey [DK19]). Using a filtering algorithm is a by-now-standard technique in algorithmic robust statistics, and the approach can be summarized as follows: 1) with high

probability over the sampling of the n uncorrupted samples, there exists a large subset of uncorrupted samples (say, a $1 - O(\epsilon)$ fraction) satisfying a "stability" condition with respect to the mean of the uncorrupted distribution, and 2) a filtering algorithm taking as input a *corrupted* version of the stable samples will remove some of the samples, such that the sample mean of the remaining points is guaranteed to be close to the true mean, which can then be returned as the final estimate. The notion of "stability" depends crucially at the task at hand, and is defined below for the sparse estimation problem.

Stability-Based Algorithm for Sparsity   Informally speaking, we say a set S is stable when the mean and the covariance of S do not deviate too much when we remove a few elements from S. For the context of sparse estimation, we would like to measure the deviation only along the k-sparse directions. However, it is NP-hard to calculate the maximum of $v^\top A v$ over k-sparse unit vectors for an arbitrary matrix A (this is known as the sparse PCA problem [TP14]). Following [BDLS17], our definition of stability would involve a convex relaxation of the above optimization problem, using the following definition of the set $\mathcal{X}_k$ and the matrix norm $\| \cdot \|_{\mathcal{X}_k}$.

Definition 1.4 (The set $\mathcal{X}_k$ and the norm $\| \cdot \|_{\mathcal{X}_k}$). *The set $\mathcal{X}_k$ is defined as the set of positive semi-definite matrices that have trace 1 and ℓ_1-norm at most k when flattened as a vector. The matrix norm $\|A\|_{\mathcal{X}_k}$ is then defined as $\sup_{M \in \mathcal{X}_k} |A \bullet M|$, where $A \bullet M$ denotes the trace product $\mathrm{tr}(A^\top M)$.*

Note that for any square matrix A, $\|A\|_{\mathcal{X}_k}$ is always upper bounded by its spectral norm. Furthermore, observe that for any square matrix A, the maximum of $v^\top A v$ over k-sparse unit vectors is upper bounded by $\|A\|_{\mathcal{X}_k}$, and the latter can be calculated efficiently using a convex program. We are now ready to define the stability condition.

Definition 1.5 (Stability Condition). *For $0 < \epsilon < 0.5$ and $\epsilon \le \delta$, a set S is (ϵ, δ, k)-stable with respect to $\mu \in \mathbb{R}^d$ and $\sigma \in \mathbb{R}_+$ if it satisfies the following condition: for all subsets $S' \subset S$ with $|S'| \ge (1 - \epsilon)|S|$, the following holds: (i) $\|\mu_{S'} - \mu\|_{2,k} \le \sigma\delta$, and (ii) $\|\overline{\Sigma}_{S'} - \sigma^2 I\|_{\mathcal{X}_k} \le \sigma^2\delta^2/\epsilon$, where $\mu_{S'} = (1/|S'|) \sum_{x \in S'} x$ is the sample mean of S' and $\overline{\Sigma}_{S'} = (1/|S'|) \sum_{x \in S'} (x - \mu)(x - \mu)^\top$ is the second moment of S'.*

[Definition 1.5] is intended for distributions with covariance matrices at most σ^2 times identity. We will omit μ and σ above when they are clear from context. Focusing on the Gaussian distribution with identity covariance, [BDLS17] gave a computationally-efficient algorithm with $k^2 \log d$ samples.[1]

By using the standard median-of-means pre-processing described in [Section 2], we can reduce the problem to the case when the corruption parameter ϵ is constant, say 0.01, and aim to achieve only a constant estimator error in the $\ell_{2,k}$ norm. For this regime, we state the guarantees of robust sparse mean estimation algorithm of [BDLS17] (developed for the Gaussian setting) as follows[2]:

Fact 1.6. *Let S be a set in $\mathbb{R}^d$ such that there exists a set $S' \subseteq S$ such that (i) $|S'| \ge 0.99|S|$ and (ii) S' is an $(0.01, O(1), k)$-stable with respect to (unknown) μ and (unknown) σ. There is a $\mathrm{poly}(|S|, d)$-time algorithm that takes as input T, an 0.01-corruption of S, and returns a mean estimate $\hat{\mu}$ such that $\|\hat{\mu} - \mu\|_{2,k} \le O(\sigma)$.*

Given the prior algorithmic result, the remaining challenge is to show that, even in the setting of heavy-tailed data, a large subset of the uncorrupted samples satisfies the "stability" condition with high probability. In the dense setting, [DKP20, HLZ20] showed that $O(d)$ samples suffice for stability, which is too large for our setting.

Truncation is Necessary for Stability   Recall that our goal is to show that, if we sample $k^2 \log d$ samples from a heavy-tailed distribution, then it contains a large stable subset. For the light-tailed data (Gaussian), this was shown in [BDLS17]. However, this statement is not true for general heavy-tailed distributions. Consider the standard setting for modelling heavy-tailed data, that the covariance Σ of the uncorrupted distribution is upper bounded by the identity. For simplicity, also assume that the sparsity parameter k, corruption parameter ϵ and failure probability τ are all constants. Thus, our goal is to show that, with high probability, there is a large stable subset among $\log d$ samples. Yet, as we show in [Example 1] in [Section 3], there exists a distribution where deterministically for *any* set of even $o(d)$ many uncorrupted samples, *no* large subset can be stable. This distribution is the one

---

[1]The additional factor of k in their sample complexity (cf. [Fact 1.2]) is because the convex relaxation involving $\mathcal{X}_k$ norm can be loose. However, [DKS17, BB20] suggest that k^2 samples are needed for efficient algorithms.

[2]See also [ZJS21] for a related algorithm.

returning a vector of length $\sqrt{d}$ from a randomly chosen axis direction, which has unit covariance. Essentially, the long length of $\sqrt{d}$ along directions as sparse as the axis directions causes stability to fail to hold.

In order to circumvent this obstacle, we propose to "truncate" all the samples in ℓ_∞ norm, before using a stability-based filtering robust mean estimation algorithm: compute an initial mean estimate, then clip each sample coordinate-wise to within a radius of $\Theta(\sqrt{k})$ of the initial mean estimate. This radius is chosen carefully to ensure that the mean of the original distribution and the clipped distribution is close in $\ell_{2,k}$-norm. Ensuring that the clipped distribution also has small variance turns out to be non-trivial as we detail below.

Necessity of Bounded Higher Moments   After truncation, no point is too far from the true mean, but truncation can potentially also *rotate* a point about the true mean, in the sense that for a sample, the direction of its difference from the true mean may change after such truncation. In general, this rotation effect can cause much of the mass of the distribution to rotate and concentrate towards certain directions, and significantly *increase* the variance in those directions. (See Appendix A.4 for more details.) In this work, we identify the mild condition that the 4[th] moment is bounded along each axis direction by some constant, on top of our assumption that $\Sigma \preceq I$, to be sufficient to show that truncation can only increase variance in directions that are non-sparse, in the sense that the resulting covariance will still have bounded $\mathcal{X}_k$ norm (see Lemma 3.1). Thus, under these mild conditions, we can safely truncate our samples (which is necessary for stability to hold as outlined above), and modify our goal to show this truncated distribution contains a large stable set with high probability.

Stability of Truncated Samples with High Probability   Even after truncation and after imposing an axis-wise 4[th] moment bound, it remains challenging to show that, with high probability, there is a large subset of samples that are stable with respect to the true mean.

As we see in Section 5, the analysis reduces to showing that with high probability over the uncorrupted samples, for every matrix $M \in \mathcal{X}_k$, there exists a large subset of samples S whose empirical covariance $\overline{\Sigma}_S$ has a small inner product with M, namely that $M \bullet \overline{\Sigma}_S$ is bounded. In the non-sparse setting, the strategy used in [DL22b] and [DKP20] is to first show a high probability event for all $M = vv^\top$ for unit vectors v, and then to show that the event for all $M = vv^\top$ deterministically implies that the event holds also for all $M \succeq 0$ with $\mathrm{tr}(M) = 1$. This strategy is important because although the cover of PSD matrices would roughly be exponential in d^2, the cover of $vv^\top$ is only exponential in d. Thus the first step holds with roughly d samples, and the second step crucially uses the spectral decomposition (SVD) of positive semi-definite (PSD) matrices. On the other hand, in our sparse setting, if we applied the usual SVD to the PSD matrices $M \in \mathcal{X}_k$, the resulting decomposition will generally not yield sparse components, and thus not allowing us to leverage sparsity. Instead, inspired by some matrix norm results derived by Li [Li18], we carefully design a (non-spectral) decomposition that does yield k^2-sparse components and can be covered with $k^2 \log d$ samples, as well as a more delicate argument to complete the second step, namely that the event holding for all components M in the decomposition implies the event holding for all $M \in \mathcal{X}_k$. The intricacies of these arguments also allow us to get a sample complexity that ultimately yields an additive (instead of multiplicative) dependence on $\log 1/\tau$, which as described in the previous section is a crucial feature of our result, and in line with the non-robust non-sparse sub-Gaussian mean estimation setting.

1.4   Related Work

Algorithmic Robust Statistics   The goal of algorithmic robust statistics is to obtain dimension-independent asymptotic error even in the presence of constant fraction of outliers in high dimensions in a computationally efficient way. Since the dissemination of [DKK+16, LRV16], which focused on high-dimensional robust mean estimation, the body of work in the field has grown rapidly. For example, prior work has obtained dimension-independent guarantees for various problems such as linear regression [KKM18, DKS19] and convex optimization [PSBR20, DKK+19a]. See the recent survey [DK19] for a more detailed description. Most relevant to us are the works for robust mean estimation that leverage the sparsity constraints and obtain improved sample complexity. The algorithms developed in [BDLS17, DKK+19b, CDG+, DKK+22] obtain optimal asymptotic error for light-tailed distributions such as Gaussians. However, these algorithms crucially rely on the light-tails and, as outlined in Section 1.3, provably do not work for heavy-tailed distributions.

Heavy-Tailed Estimation   The recent decades also saw a growing interest in studying statistics in heavy-tailed settings. Even for the basic question of univariate mean estimation without sample corruption, the statistical limits are only recently resolved by a line of work started by Catoni [Cat12] and ending with Lee and Valiant [LV21] (see also [Min22] for an alternative estimator). There is also much research effort on resolving the statistical limits of various high-dimensional estimation tasks for heavy-tailed distributions, for example, mean estimation in ℓ_2 norm [LM19c] and in other norms [LM19b, DL22a], covariance estimation [MZ20], and stochastic convex optimization [BM22]. In absence of contamination, the goal is to obtain sample complexity as if the distribution were Gaussian. Roughly speaking, this corresponds to an additive dependence on the logarithm of failure probability in various estimation tasks (as we achieve also in this work). We refer the reader to the recent survey for more details [LM19a]. This line of work focuses on the statistical limits, and the estimators developed are generally computationally inefficient.

A closely-related body of research aims to obtain *efficient* algorithms for heavy-tailed distributions with optimal statistical performance, ideally matching the above guarantees. These works include high-dimensional (dense) mean estimation [Hop20, CFB19, DL22b, LLVZ20, DKP20, HLZ20, CTBJ22, LV22], linear regression [CHK$^+$20, PJL20, Dep20b], and covariance estimation [CHK$^+$20]. We note that many of these works are inspired by the algorithmic robust statistics literature, and can also tolerate a constant fraction of contaminated data.

To the best of our knowledge, none of these works studies sparse estimation under heavy-tailed distributions (even in absence of outliers), and our work is the first result with sample complexity that is additive in the logarithm of the failure probability.

2   Preliminaries

Notations   Here we define the notations we use in the rest of the paper. For a (multi-)set $S \subset \mathbb{R}^d$, we denote $\mu_S = (1/|S|) \sum_{x \in S} x$ and $\Sigma_S = (1/|S|)(\sum_{x \in S}(x - \mu_S)(x - \mu_S)^\top)$. When the vector μ notation is clear from context, we use $\overline{\Sigma}_S$ to denote $(1/|S|) \sum_{x \in S}(x - \mu)(x - \mu)^\top$.

Let $\mathcal{U}_k$ denote the set of k-sparse unit vectors in $\mathbb{R}^d$. For two vectors x and y, $\langle x, y \rangle$ denotes the dot product $x^\top y$. For a vector $x \in \mathbb{R}^d$, we use $\|x\|_{2,k} := \sup_{v \in \mathcal{U}_k} \langle x, v \rangle$ and $\|x\|_\infty$ to denote $\max_j |x_j|$. For a matrix M, we use $\|M\|_1$ to denote $\sum_{i,j} |M_{i,j}|$ and $\|M\|_0$ to denote the number of non-zero entries of M. For two matrices A and B, we use $A \bullet B$ to denote the trace inner product $\mathrm{tr}(A^\top B)$. Define $\mathcal{X}_k := \{M : M \succeq 0, \mathrm{tr}(M) = 1, \|M\|_1 \leq k\}$. For a matrix A, we define $\|A\|_{\mathcal{X}_k} := \sup_{M \in \mathcal{X}_k} |A \bullet M|$. For an $n \in \mathbb{N}$, we use $[n]$ to denote the set $\{1, \ldots, n\}$.

Coordinate-wise Median-of-Means   We use the coordinate-wise median-of-means algorithm to robustly obtain a preliminary mean estimate, with guarantees captured by the following fact.

Fact 2.1. *The coordinate-wise median-of-means algorithm satisfies the following guarantee: given the corruption parameter ϵ, failure probability τ, and a set T of n many ϵ-corrupted samples from a distribution D with mean μ and axis-wise variance $\mathbb{E}_{X \sim D}[(X_j - \mu_j)^2] \leq \sigma^2$ for all $j \in [d]$, then with probability at least $1 - \tau$ over the sample set T, the output of the algorithm $\hat{\mu}$ is such that $\|\hat{\mu} - \mu\|_\infty \leq \sigma O(\sqrt{\epsilon} + \sqrt{(\log(d/\tau))/n})$.*

Median-of-Means Pre-Processing   Another standard technique we use in this paper is the median-of-means pre-processing, which is a distinct technique from the coordinate-wise median-of-means algorithm mentioned right above. Recall that in Theorem 1.3, the asymptotic error term is $\sqrt{\epsilon}$, which tends to 0 as the corruption parameter $\epsilon \to 0$. The following pre-processing step allows us to reduce the problem from the $\epsilon \to 0$ case to a constant ϵ case: Split the samples randomly into g equally-sized groups of size $m = n/g$ where $g = \Theta(\epsilon n)$, and replace each group by the sample mean of the group. The effects of this pre-processing is captured by the following Fact 2.2, which we prove for completeness.

Fact 2.2 (Median-of-Means Pre-Processing)**.** *Suppose there is an efficient algorithm such that, on input $\sigma \in \mathbb{R}_+$ and a 0.1-corrupted set of $n \gg k^2 \log d + \log(1/\tau)$ samples from a distribution D with mean μ and covariance Σ with $\|\Sigma\|_{\mathcal{X}_k} \leq \sigma^2$ and $\mathbb{E}_{X \sim D}[(X_j - \mu_j)^4] = O(\sigma^4)$ for each coordinate $j \in [d]$, returns $\hat{\mu}$ such that $\|\hat{\mu} - \mu\|_{2,k} \leq O(\sigma)$ with probability at least $1 - \tau$.*

Then, there is an efficient algorithm such that, on input $\epsilon \in (0, 0.1)$ and an ϵ-corrupted set of $n \gg (k^2 \log d + \log(1/\tau))/\epsilon$ samples from a distribution with mean μ and covariance Σ, satisfying $\|\Sigma\|_{\mathcal{X}_k} \leq 1$ satisfying $\mathbb{E}_{X \sim D}[(X_j - \mu_j)^4] = O(1)$ for every coordinate $j \in [d]$, returns a mean estimate $\hat{\mu}$ such that $\|\hat{\mu} - \mu\|_{2,k} \leq O(\sqrt{\epsilon})$ with probability at least $1 - \tau$.

3 Truncation Pre-Processing

The general approach for using a stability-based filtering algorithm, for algorithmic robust statistics, is to show that given sufficiently many samples, there exists a large (say $1 - O(\epsilon)$ fraction) subset of the samples that are stable with respect to the true mean μ. However, the following simple example shows that it is not possible for i.i.d. samples from a heavy-tailed distribution to satisfy the sparse stability in $\text{poly}(\log d)$ samples.

Example 1. *For any number of moments $t \geq 2$, there is a distribution X satisfying the following conditions: (i) The mean of X is 0, and for every unit vector v, the t^{th} moment in direction v is upper bounded by 1, that is, $\mathbb{E}[|\langle v, x \rangle|^t] \leq 1$ for $t \geq 2$, (ii) If S is an arbitrary set of $n \leq o(d^{2/t})$ points from the support of X, then the set S cannot be $(\epsilon, O(\sqrt{\epsilon}), k)$-stable, for any $\epsilon > 0$, with respect to the mean of the distribution. As a corollary, no subset of S can be stable either.*

We show Example 1 in Appendix B.

Thus, we need to modify the algorithm if we want to do robust sparse mean estimation using $\text{poly}(k, \log d)$ samples. Our approach is to perform an initial truncation of the samples, before using a stability-based robust mean estimator. A balance needs to be struck, in order to truncate sufficiently for stability to hold (with high probability over the samples), but also to truncate mildly enough such that the mean (and covariance) of the truncated distribution does not shift too much.

For a scalar $a \in \mathbb{R}_+$ and a vector $b \in \mathbb{R}^d$, let $h_{a,b} : \mathbb{R}^d \to \mathbb{R}^d$ be the following thresholding function:

$$\forall i \in [d], \quad h_{a,b}(x)_i = \begin{cases} x_i, & \text{if } |x_i - b_i| \leq a \\ b_i + a & \text{if } x_i - b_i \geq a \\ b_i - a & \text{if } x_i - b_i \leq -a \end{cases} . \tag{1}$$

Note that $h_{a,b}(x)$ projects the point x to the ℓ_∞ ball of radius a around b.

As explained in the Introduction, truncation in general rotates a point about the true mean, and thus can in fact cause the covariance of the distribution to grow in certain directions. The following lemma captures the fact that, if we make the further mild assumption that the distribution has bounded 4^{th} moment along all the axis directions, then we will at least be able to preserve the $\mathcal{X}_k$ norm of the covariance matrix. The proof of Lemma 3.1 is also in Appendix B.

Lemma 3.1 (Truncation in ℓ_∞). *Let P be a distribution over $\mathbb{R}^d$ with mean μ_P and covariance Σ_P, with $\|\Sigma\|_{\mathcal{X}_k} \leq \sigma^2$ for some $\sigma^2 > 0$. Let $X \sim P$ and assume that for all $j \in [d]$, $\mathbb{E}[(X - \mu_P)_j^4] \leq \sigma^4 \nu^4$ for some $\nu \geq 1$. Let $b \in \mathbb{R}^d$ be such that $\|b - \mu\|_\infty \leq a/2$ and $a := 2\sigma\sqrt{k/\epsilon}$ for some $\epsilon \in (0, 1)$. Define Q to be the distribution of $Y := h_{a,b}(X)$. Let the mean and covariance of Q be μ_Q and Σ_Q respectively. Then the following hold:*

(1) $\|\mu_P - \mu_Q\|_\infty \leq \sigma\sqrt{\epsilon/k}$

(2) $\|\mu_P - \mu_Q\|_{2,k} \leq \sigma\sqrt{\epsilon}$

(3) $\|\Sigma_P - \Sigma_Q\|_{\mathcal{X}_k} \leq 3\sigma^2 \epsilon \nu^4$

(4) For all $i \in [d]$, $\mathbb{E}[(Y - \mu_Q)_i^4] \leq 8\nu^4 \sigma^4$

(5) $\|Y - \mu_Q\|_\infty \leq 2a = 4\sigma\sqrt{k/\epsilon}$ *almost surely.*

In Lemma 3.1 above, b represents the initial mean estimate, and $\tilde{\mu}$ will be obtained by Fact 2.1.

4 Algorithm and Analysis

The high-level algorithm we propose is stated as follows.

Algorithm 1 Robust Sparse Mean Estimation with High Probability

---

1. Input: An ϵ-corrupted sample set $S \subseteq \mathbb{R}^d$ of size n

2. Median-of-Means pre-processing: Group points into g groups, each of size $m = n/g$, where $g = \Theta(\epsilon n)$, and take the sample mean of a group to be a new point

3. Compute initial coordinate-wise median-of-means estimate $\tilde{\mu}$

4. Truncate all points to within $B_\infty(\tilde{\mu}, 2\sqrt{k})$, namely, given a point x, we replace it with the point $h_{2\sqrt{k},\tilde{\mu}}(x)$, where $h_{a,b}$ is defined in Equation (1).

5. Run the stability-based robust *sparse* mean estimator from Fact 1.6 on the samples after the processing of Step 4.

We note that this algorithm is shift and scale invariant, based on the same invariance of the median-of-means pre-processing as well as the invariance of the robust sparse mean estimator from Fact 1.6.

We will now prove that Algorithm 1 satisfies the guarantees of Theorem 1.3 and its stronger version, Theorem D.12. Our analysis uses Theorem D.11, which states that, with high probability, there exists a set consisting of most of the truncated samples that is stable with respect to some point close to the true mean in ℓ_∞ norm. In Section 5, we outline the central part (Theorem 5.1) of the proof of this theorem. We have not optimized constants in our analysis.

Proof of Theorems 1.3 and D.12. By Fact 2.2, it suffices to show that, for every $\sigma > 0$, Steps 3–5 in Algorithm 1 yields an $O(\sigma)$ estimation error in $\ell_{2,k}$ norm when given corrupted samples from a distribution D with covariance bounded by $\sigma^2 I$ and axis-wise 4th moment bounded by $O(\sigma^4)$.

Theorem D.11 states that, with probability at least $1 - \tau$, the samples after the processing of Step 4 are such that there exists a 95% of the samples that form a $(0.1, O(1), k)$-stable subset with respect to some vector μ' and σ with $\|\mu' - \mu\|_\infty \leq O(\sigma/\sqrt{k})$. Fact 1.6 then guarantees that, on input such a set of samples, the algorithm we invoke Step 5 of Algorithm 1 will return a mean estimate $\hat{\mu}$ such that $\|\hat{\mu} - \mu'\|_{2,k} \leq O(\sigma)$. Further, since $\|\mu' - \mu\|_\infty \leq O(\sigma/\sqrt{k})$, we have that $\|\mu' - \mu\|_{2,k} \leq O(\sigma)$, and therefore we can conclude with the triangle inequality that the mean estimate $\hat{\mu}$ satisfies $\|\hat{\mu} - \mu\|_{2,k} \leq O(\sigma)$. $\qquad\square$

5 Stability After Removing Points: Additive dependence on $\log(1/\tau)$

In this section, we sketch the main stability result in this paper. Recall, via the median-of-means pre-processing, that we only need to consider the constant contamination case ($\epsilon = \Theta(1)$). Thus, the goal is to show (Theorem D.11) that with high probability, after truncation according to the coordinate-wise median-of-means preliminary estimate, there exists a large subset of uncontaminated samples that is $(\Theta(1), O(\sigma), k)$-stable with respect to the *true mean* of the distribution where $\sigma = \|\Sigma\|_{\mathcal{X}_k}$.

Theorem 5.1 below, an informal version of Theorem C.1 in Appendix C, captures the core of the argument. The key difference between Theorems 5.1 and D.11 is that the former is a stability result concerning uncontaminated samples truncated according to some *fixed* vector close to the true mean. On the other hand, the final stability result we require concerns samples truncated according to the coordinate-wise median-of-means estimate, which itself depends on the samples and is not fixed. Appendix D shows the delicate argument going from Theorem 5.1 to Theorem D.11.

Theorem 5.1 (Informal version of Theorem C.1). *Let S be a set of n i.i.d. data points from a distribution P over $\mathbb{R}^d$. Let the mean of P be μ, and covariance Σ such that $\|\Sigma\|_{\mathcal{X}_k} \leq \sigma^2$, and for all $j \in [d]$, $\mathbb{E}[X_j^4] = O(\sigma^4)$. Suppose P is supported over the set $\{x : \|x - \mu\|_\infty = O(\sigma\sqrt{k})\}$. If $n = \Omega(k^2 \log d + \log(1/\tau))$, then, with probability $1 - \tau$, there exists a set $S' \subset S$ such that:*

1. *$|S'| \geq 0.98n$*

2. *S' is $(0.01, \delta, k)$-stable with respect to μ and σ where $\delta = O(1)$.*

For the rest of the section, we sketch the proof of Theorem 5.1.

Proof sketch for Theorem 5.1. In the following proof sketch, we hide most multiplicative constants. The proof of Theorem C.1 in Appendix C will make explicit the calculations for completeness. For the rest of the sketch, we will also assume $\mu = 0$ without loss of generality.

Instead of directly showing the existence of subset $S' \subseteq S$ (with high probability over the samples S) that is stable, we will be working with weights/distributions over the set S of samples. Given a weight vector w and $S = \{x_1, \ldots, x_n\} \subset \mathbb{R}^d$, denote $\overline{\Sigma}_w$ by $\sum_i w_i x_i x_i^\top$. Then, Proposition A.1 in Appendix A lets us show the following simpler condition instead: let $\Delta_{n,\epsilon}$ be the set of weights/distributions w such that $w_i \leq 1/(1 - \epsilon)$, then there exists a weighting $w \in \Delta_{n,0.01}$ such that $\|\overline{\Sigma}_w\|_{\mathcal{X}_k} \leq O(\sigma^2)$. That is, for the following proof, we just need to prove that $\min_{w \in \Delta_{n,0.01}} \|\overline{\Sigma}_w\|_{\mathcal{X}_k} = O(\sigma^2)$.

We proceed as follows:

$$\min_{w \in \Delta_{n,0.01}} \|\overline{\Sigma}_w\|_{\mathcal{X}_k} = \min_{w \in \Delta_{n,0.01}} \max_{M \in \mathcal{X}_k} M \bullet \overline{\Sigma}_w = \max_{M \in \mathcal{X}_k} \min_{w_M \in \Delta_{n,0.01}} M \bullet \overline{\Sigma}_w$$

where the last equality is a straightforward application of the minimax theorem for a minimax optimization problem with independent convex domains and a bilinear objective. It thus suffices to show the following: with probability $1 - \tau$,

$$\forall M \in \mathcal{X}_k : \; |\{x \in S : x_i^\top M x_i \gg \sigma^2\}| \leq 0.01|S| \tag{2}$$

and we can construct the weight w_M as uniform distribution over the elements outside the above set.

Define the following sets of sparse matrices, the first of which also appears in [Li18]:

$$\mathcal{A}_k := \left\{ A \in \mathbb{R}^{d \times d} : \|A\|_0 \leq k^2, \|A\|_F \leq 1 \right\},$$
$$\mathcal{A}_{k,P} := \left\{ A \in \mathcal{A}_k : \mathbb{P}_{x \sim P} \left(x^\top A x \gg \sigma^2 \right) \leq \text{small constant} \right\}. \tag{3}$$

If $n \gg k^2 \log d + \log(1/\tau)$, then a standard covering/VC-dimension bound (see Lemma B.2 for details) implies that the following event holds with probability $1 - \tau$:

$$\forall A \in \mathcal{A}_{k,P} : \; |\{x \in S : x^\top A x \gg \sigma^2\}| \leq 0.002 \cdot |S|. \tag{4}$$

We will now show that the event in Equation (4) implies that the event in Equation (2) holds. Note that the constants hidden in the $\gg$ notation in Equations Equation (2) and Equation (4) are different, which is the slackness that makes the implication possible. As mentioned, we will make these constants explicit in the formal proof in Appendix C.

Suppose, for the sake of contradiction, that the event in Equation (2) does not hold. Then there exists an $M \in \mathcal{X}_k$ such that $|\{x \in S : x_i^\top M x_i \gg \sigma^2\}| > 0.01|S|$. We will show the existence of a matrix Q violating event Equation (4), via the probabilistic method, to reach the desired contradiction.

To achieve this, we define a *decomposition* of $M \in \mathcal{X}_k$ into components in $\mathcal{A}_{k,P}$, in the sense of a distribution over matrices Q such that 1) $\mathbb{E}[Q] = M$ and 2) a small constant multiple of Q belongs to $\mathcal{A}_{k,P}$ with high constant probability. We will additionally show that this decomposition Q roughly preserves quadratic forms, in that if M violates Equation (2) then Q violates Equation (4) with non-trivial probability.

Fixing an M that violates Equation (2), consider the random matrix Q where each entry $Q_{i,j}$ is independently set to be $M_{i,j}/p_{i,j}$ with probability $p_{i,j}$ and 0 otherwise, where $p_{i,j} := \min(1, k|M_{i,j}|/c)$ for a large constant $c > 0$.

We will show that the following events hold simultaneously with non-zero probability, leading to a contradiction to event Equation (4): (i) $Q/c' \in \mathcal{A}_{k,P}$ for a large constant c' and (ii) $|\{x \in S : x_i^\top (Q/c') x_i \gg \sigma^2\}| > 0.002 \cdot |S|$. Using different techniques, we will show that the first condition holds with probability at least $1 - 2 \times 10^{-6}$ and the second condition holds with probability at least 4×10^{-6}, thus implying that the events hold simultaneously with non-zero probability.

Condition (i), that $Q/c' \in \mathcal{A}_{k,P}$ with high constant probability. Showing that Q/c' belongs to $\mathcal{A}_k$ is straightforward: by the construction of Q, it has small expected sparsity as well as small expected Frobenius norm. An application of Markov's inequality shows that $Q/c' \in \mathcal{A}_k$ with high constant probability (Lemma C.2).

The trickier part is to show that Q/c' is also in $\mathcal{A}_{k,P}$, namely that $\mathbb{P}_{x\sim P}(x^\top (Q/c')x \gg \sigma^2)$ is upper bounded by the small constant. We consider the distribution of $x^\top Qx$ over the probability of independently drawing $x \sim P$ and a random Q, and show that $x^\top Qx$ is small with high probability over this joint distribution (Lemma C.4), which requires using the axis-wise 4^{th} moment bounds on P as well as the fact that $M \in \mathcal{X}_k$. Lemma C.3 implies that with high probability, we will draw a Q satisfying $\mathbb{P}_{x\sim P}(x^\top Qx \gg \sigma^2)$ being bounded by a small constant.

Condition (ii), that if M violates Equation (2) then Q violates Equation (4) with non-trivial probability   We consider the random variable $Z = \sum_i \mathbb{1}_{x_i^\top Qx_i \gg \sigma^2}$, over the randomness of Q. The overall strategy is to lower bound $\mathbb{E}[Z] \geq cn$ for a small $c > 0$ and use the Paley-Zygmund inequality to show that with at least small constant probability, $Z \geq cn/2$, where we use Z always is less than n. The expectation of Z is just the average over all samples the probability that $x_i^\top Qz_i \gg \sigma^2$. Since the fraction of x_is with $x_i^\top Mx_i \gg \sigma^2$ is at least a constant, to lower bound $\mathbb{E}[Z]$, it suffices for us to lower bound this probability for each such x_i. Fix any such x for the remainder of the argument.

Lower bounding the probability that $x^\top Qx \gg \sigma^2$ is non-trivial, since, the support of the quadratic form can scale with k and given a lower bound on its expectation, it could be the case that it is 0 with high probability and has a huge value with small probability.

We will argue that this does not happen by considering two cases. First, and the simpler setting, is when $x^\top Qx$ has small variance. In this case, we can use Chebyshev's inequality to lower bound the probability that $x^\top Qx \gg \sigma^2$. Essentially, this is the case when $x^\top Qx$ is "concentrated".

The second, and the more involved setting, is when $x^\top Qx$ has large variance. Here, we would like to argue that $x^\top Qx$ is sufficiently "anti-concentrated" like a Gaussian, in the sense that there is a constant probability mass above its mean. Noting that $x^\top Qx = \sum_{i\in[d],j\in[d]}(x)_i(x)_j Q_{i,j}$ and the entries of Q are independent, this is a sum of independent terms, and we can reasonably expect Gaussian-like behavior. Concretely, we will show this using the Berry-Esseen theorem, which requires us to upper bound the third moment of $x^\top Qx$. This upper bound crucially uses the facts that 1) the samples here are truncated, which implies that each $(x)_i(x)_j$ has bounded magnitude and 2) each $Q_{i,j}$ is a (scaled) Bernoulli which has exact and simple expressions for its second and third moments.

Summarizing, we argued that conditions (i) and (ii) hold simultaneously with non-zero probability, thus showing the existence of a Q violating Equation (4), reaching a contradiction assuming the existence of some M violating Equation (2). Therefore, Equation (4) implies Equation (2), which completes the argument for Theorem 5.1. $\qquad\square$

Acknowledgements

We thank the anonymous reviewers for helpful comments on the paper. Ilias Diakonikolas was supported by NSF Medium Award CCF-2107079, NSF Award CCF-1652862 (CAREER), a Sloan Research Fellowship, and a DARPA Learning with Less Labels (LwLL) grant. Daniel Kane was supported by NSF Medium Award CCF-2107547, NSF Award CCF-1553288 (CAREER), and a grant from CasperLabs. Jasper C.H. Lee was supported in part by the generous funding of a Croucher Fellowship for Postdoctoral Research, NSF award DMS-2023239, NSF Medium Award CCF-2107079 and NSF AiTF Award CCF-2006206. Ankit Pensia was supported by NSF grants NSF Award CCF-1652862 (CAREER), DMS-1749857, and CCF-1841190.

Part of this work was done while Ilias Diakonikolas, Jasper C.H. Lee and Ankit Pensia were visitors at the Simons Institute for the Theory of Computing.

References

[ABH+72] D. F. Andrews, P. J. Bickel, F. R. Hampel, P. J. Huber, W. H. Rogers, and J. W. Tukey. *Robust Estimates of Location: Survey and Advances*. Princeton University Press, 1972.

[AV19] M. E. Ahsen and M. Vidyasagar. An approach to one-bit compressed sensing based on probably approximately correct learning theory. *Journal of Machine Learning Research*, 20(11):1–23, 2019.

[BB20]   M. Brennan and G. Bresler. Reducibility and statistical-computational gaps from secret leakage. In *Proc. 33rd Annual Conference on Learning Theory (COLT)*, 2020.

[BDLS17] S. Balakrishnan, S. S. Du, J. Li, and A. Singh. Computationally efficient robust sparse estimation in high dimensions. In *Proc. 30th Annual Conference on Learning Theory*, 2017.

[BM22]   D. Bartl and S. Mendelson. On Monte-Carlo methods in convex stochastic optimization. *The Annals of Applied Probability*, 32(4):3146 – 3198, 2022.

[Cat12]   O. Catoni. Challenging the empirical mean and empirical variance: A deviation study. *Annales de l'Institut Henri Poincaré, Probabilités et Statistiques*, 48(4):1148–1185, 2012.

[CDG$^+$] Y. Cheng, I. Diakonikolas, R. Ge, S. Gupta, D. M. Kane, and M. Soltanolkotabi. Outlier-robust sparse estimation via non-convex optimization. In *To appear in NeurIPS 2022*.

[CFB19]   Y. Cherapanamjeri, N. Flammarion, and P. L. Bartlett. Fast mean estimation with sub-gaussian rates. In *Proc. 32nd Annual Conference on Learning Theory (COLT)*, 2019.

[CHK$^+$20] Y. Cherapanamjeri, S. B. Hopkins, T. Kathuria, P. Raghavendra, and N. Tripuraneni. Algorithms for heavy-tailed statistics: Regression, covariance estimation, and beyond. In *Proc. 52nd Annual ACM Symposium on Theory of Computing (STOC)*, 2020.

[CTBJ22] Y. Cherapanamjeri, N. Tripuraneni, P. L. Bartlett, and M. I. Jordan. Optimal Mean Estimation without a Variance. In *Proc. 35th Annual Conference on Learning Theory (COLT)*, 2022.

[Dep20a] J. Depersin. Robust subgaussian estimation with VC-dimension. *CoRR*, abs/2004.11734, 2020.

[Dep20b] J. Depersin. A spectral algorithm for robust regression with subgaussian rates. *CoRR*, abs/2007.06072, 2020.

[DG92]   D. L. Donoho and M. Gasko. Breakdown properties of location estimates based on halfspace depth and projected outlyingness. *Ann. Statist.*, 20(4):1803–1827, 12 1992.

[DK19]   I. Diakonikolas and D. M. Kane. Recent advances in algorithmic high-dimensional robust statistics. *arXiv preprint arXiv:1911.05911*, 2019.

[DKK$^+$16] I. Diakonikolas, G. Kamath, D. M. Kane, J. Li, A. Moitra, and A. Stewart. Robust estimators in high dimensions without the computational intractability. In *Proc. 57th IEEE Symposium on Foundations of Computer Science (FOCS)*, 2016.

[DKK$^+$19a] I. Diakonikolas, G. Kamath, D. M. Kane, J. Li, J. Steinhardt, and A. Stewart. Sever: A robust meta-algorithm for stochastic optimization. In *Proc. 36th International Conference on Machine Learning (ICML)*, 2019.

[DKK$^+$19b] I. Diakonikolas, S. Karmalkar, D. Kane, E. Price, and A. Stewart. Outlier-robust high-dimensional sparse estimation via iterative filtering. In *Advances in Neural Information Processing Systems, (NeurIPS)*, volume 32, 2019.

[DKK$^+$22] I. Diakonikolas, D. M. Kane, S. Karmalkar, A. Pensia, and T. Pittas. Robust sparse mean estimation via sum of squares. In *Proc. 35th Annual Conference on Learning Theory (COLT)*, 2022.

[DKP20]   I. Diakonikolas, D. M. Kane, and A. Pensia. Outlier Robust Mean Estimation with Subgaussian Rates via Stability. In *Advances in Neural Information Processing Systems, (NeurIPS)*, volume 33, 2020.

[DKS17]   I. Diakonikolas, D. M. Kane, and A. Stewart. Statistical query lower bounds for robust estimation of high-dimensional Gaussians and Gaussian mixtures. In *Proc. 58th IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 73–84, 2017.

[DKS19] I. Diakonikolas, W. Kong, and A. Stewart. Efficient algorithms and lower bounds for robust linear regression. In *Proc. 30th Annual Symposium on Discrete Algorithms (SODA)*, 2019.

[DL88] D. L. Donoho and R. C. Liu. The "Automatic" Robustness of Minimum Distance Functionals. *The Annals of Statistics*, 16(2):552–586, 1988.

[DL22a] J. Depersin and G. Lecué. Optimal robust mean and location estimation via convex programs with respect to any pseudo-norms. *Probab. Theory Relat. Fields*, 183:997–1025, 2022.

[DL22b] J. Depersin and G. Lecué. Robust sub-Gaussian estimation of a mean vector in nearly linear time. *The Annals of Statistics*, 50(1):511–536, February 2022.

[EK12] Y. C. Eldar and G. Kutyniok. *Compressed sensing: theory and applications*. Cambridge university press, 2012.

[GKT51] D. Gale, H. W. Kuhn, and A. W. Tucker. Linear programming and the theory of games. *Activity analysis of production and allocation*, 13:317–335, 1951.

[HLZ20] S. B. Hopkins, J. Li, and F. Zhang. Robust and Heavy-Tailed Mean Estimation Made Simple, via Regret Minimization. In *Advances in Neural Information Processing Systems, (NeurIPS)*, volume 33, 2020.

[Hop20] S. B. Hopkins. Mean estimation with sub-Gaussian rates in polynomial time. *Annals of Statistics*, 48(2):1193–1213, 2020.

[HR09] P. J. Huber and E. M. Ronchetti. *Robust statistics*. Wiley New York, 2009.

[HTW15] T. Hastie, R. Tibshirani, and M. Wainwright. *Statistical Learning with Sparsity: The Lasso and Generalizations*. Chapman & Hall/CRC, 2015.

[KKM18] A. Klivans, P. Kothari, and R. Meka. Efficient algorithms for outlier-robust regression. In *Proc. 31st Annual Conference on Learning Theory (COLT)*, 2018.

[Li18] J. Li. *Principled Approaches to Robust Machine Learning and Beyond*. PhD thesis, Massachusetts Institute of Technology, 2018.

[LLVZ20] Z. Lei, K. Luh, P. Venkat, and F. Zhang. A fast spectral algorithm for mean estimation with sub-gaussian rates. In *Proc. 33rd Annual Conference on Learning Theory (COLT)*, 2020.

[LM19a] G. Lugosi and S. Mendelson. Mean Estimation and Regression Under Heavy-Tailed Distributions: A Survey. *Foundations of Computational Mathematics*, 19(5):1145–1190, 2019.

[LM19b] G. Lugosi and S. Mendelson. Near-optimal mean estimators with respect to general norms. *Probability Theory and Related Fields*, 175(3-4):957–973, 2019.

[LM19c] G. Lugosi and S. Mendelson. Sub-Gaussian estimators of the mean of a random vector. *The Annals of Statistics*, 47(2):783–794, 2019.

[LRV16] K. A. Lai, A. B. Rao, and S. Vempala. Agnostic estimation of mean and covariance. In *Proc. 57th IEEE Symposium on Foundations of Computer Science (FOCS)*, 2016.

[LV21] J. C. H. Lee and P. Valiant. Optimal Sub-Gaussian Mean Estimation in $\mathbb{R}$. In *Proc. 62nd IEEE Symposium on Foundations of Computer Science (FOCS)*, 2021.

[LV22] J. C. H. Lee and P. Valiant. Optimal sub-gaussian mean estimation in very high dimensions. In *Proc. 13th Innovations in Theoretical Computer Science Conference (ITCS)*, 2022.

[Min22] S. Minsker. U-statistics of growing order and sub-gaussian mean estimators with sharp constants. *CoRR*, abs/2202.11842, 2022.

[MZ20]   S. Mendelson and N. Zhivotovskiy. Robust covariance estimation under $L_4 - L_2$ norm equivalence. *Annals of Statistics*, 48(3):1648–1664, 2020.

[PBR20]   A. Prasad, S. Balakrishnan, and P. Ravikumar. A robust univariate mean estimator is all you need. In *Proc. 23rd International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2020.

[PJL20]   A. Pensia, V. Jog, and P. Loh. Robust regression with covariate filtering: Heavy tails and adversarial contamination. *CoRR*, abs/2009.12976, 2020.

[PSBR20]   A. Prasad, A. S. Suggala, S. Balakrishnan, and P. Ravikumar. Robust estimation via robust gradient estimation. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 82(3):601–627, July 2020.

[TP14]   A. M. Tillmann and M. E. Pfetsch. The Computational Complexity of the Restricted Isometry Property, the Nullspace Property, and Related Concepts in Compressed Sensing. *IEEE Transactions on Information Theory*, 60(2):1248–1259, 2014.

[Tuk60]   J. W. Tukey. A survey of sampling from contaminated distributions. *Contributions to probability and statistics*, 2:448–485, 1960.

[van16]   S. van de Geer. *Estimation and Testing Under Sparsity*. Number 2159 in École d'Été de Probabilités de Saint-Flour. Springer, 2016.

[Yat85]   Y. G. Yatracos. Rates of convergence of minimum distance estimators and Kolmogorov's entropy. *Annals of Statistics*, 13:768–774, 1985.

[ZJS21]   B. Zhu, J. Jiao, and J. Steinhardt. Robust estimation via generalized quasi-gradients. *Information and Inference: A Journal of the IMA*, 11(2):581–636, 08 2021.

Supplementary Material

Additional Notation. For a vector $x \in \mathbb{R}^d$ and $H \subset [d]$, we denote v_H to denote the vector that is equal to v on $i \in H$, and zero otherwise. For a real-valued random variable X and $m \in \mathbb{N}$, we use $\|X\|_{L_m}$ to denote $(\mathbb{E}\,|X|^m)^{1/m}$. For a set $S \subseteq \mathbb{R}^d$ and a function f, we also define the set function notation $f(S)$ as $\{f(x)\,|\,x \in S\}$.

A   Miscellaneous Lemmas and Facts

A.1   Finding a stable subset from a stable weighted subset

For a set S on n points, we define $\Delta_{n,\epsilon}$ as the set of weights $w \in \mathbb{R}^n$ such that $w_i \in [0, 1/((1-\epsilon)n]$ for all $i \in [n]$ and $\sum_i w_i = 1$. For a fixed vector $\mu \in \mathbb{R}^d$ that will be clear from context, a set of n points $S = \{x_1, \ldots, x_n\}$, and weights $w \in \Delta_{n,\epsilon}$ over S, we use $\overline{\Sigma}_w$ to denote $\sum_i w_i(x_i - \mu)(x_i - \mu)^\top$.

The goal of this section is to show Proposition A.1, which states that if we have a weight w over S such that $\overline{\Sigma}_w$ (with respect to some vector μ) has bounded $\mathcal{X}_k$ norm proportional to σ^2 for some $\sigma > 0$, then there must exists some large subset $S' \subseteq S$ that is stable with respect to μ and σ.

Proposition A.1. *Let S be a set of n points in $\mathbb{R}^d$. Let $\Delta_{n,\epsilon}$ be the set of weights defined above, and define the notation $\overline{\Sigma}_w = \sum_{x_i \in S} w_i(x_i - \mu)(x_i - \mu)^\top$ for some given vector $\mu \in \mathbb{R}^d$. Suppose that there exists a $w \in \Delta_{n,\epsilon}$ such that $\|\overline{\Sigma}_w\|_{\mathcal{X}_k} \leq B\sigma^2$ for some vector μ. Then there exists a subset $S' \subseteq S$ such that $(i)|S'| \geq (1 - 2\epsilon)n$ and (ii) S' is (ϵ, δ, k)-stable with respect to μ and σ, where $\delta = O(\sqrt{B} + 1)$.*

Observe that $\|\overline{\Sigma}_w\|_{\mathcal{X}_k} \leq B\sigma^2$ implies $\|\overline{\Sigma}_w - \sigma^2 I\|_{\mathcal{X}_k} \leq (B + 1)\sigma^2$ by the triangle inequality. In order to show Proposition A.1, we show Lemma A.2, which is a weakening of Proposition A.1 where we additionally assume that $\mu_w = \sum_i w_i x_i$ is close to μ, where μ is the vector we use to define $\overline{\Sigma}_w$ as well as the vector that we want to find a large sample subset S' to be stable with respect to. To use Lemma A.2, we additionally show Proposition A.4, which states that $\|\overline{\Sigma}_w\|_{\mathcal{X}_k} \leq B\sigma^2$ is enough to imply that μ_w is close to μ. We combine Lemma A.2 and Proposition A.4 to prove Proposition A.1 at the end of Appendix A.1.

Lemma A.2. *Suppose, for some $\epsilon \leq \frac{1}{3}$ and for some $\delta \geq \sqrt{\epsilon}$, there exist a $w \in \Delta_{n,\epsilon}$ over a set of n samples $S = \{x_1, \ldots, x_n\}$, a $\mu \in \mathbb{R}^d$ and a $\sigma > 0$ such that*

- $\|\mu_w - \mu\|_{2,k} \leq \delta\sigma$,

- $\|\sum_{i \in [n]} w_i(x_i - \mu)(x_i - \mu)^\top - \sigma^2 I\|_{\mathcal{X}_k} \leq \sigma^2 \frac{\delta^2}{\epsilon}$.

Then, there exists a subset $S' \subseteq S$ of samples such that

- $|S'| \geq (1 - 2\epsilon)|S|$,

- S' *is (ϵ, δ', k)-stable with respect to μ and σ, where $\delta' = O(\delta + \sqrt{\epsilon})$.*

Proof. Without loss of generality, we will only handle the $\sigma = 1$ case to simplify notation.

The main step is to show the existence of a large subset S' whose mean is within $10\delta + 10\sqrt{\epsilon}$ of μ and whose variance is at most $9(1 + \delta^2/\epsilon)$. In fact, we can simply choose S' to be the subset whose weights w_i are the largest.

Without loss of generality, assume $\mu = 0$ and that ϵn is an integer. We also order the samples in decreasing order of weight in w, namely, $1/((1 - \epsilon)n) \geq w_1 \geq w_2 \geq \ldots \geq w_n$.

First, we will lower bound each w_i. We have that for each $k \in [n]$,

$$1 = \sum_i w_i \leq \frac{k}{(1 - \epsilon)n} + (n - k)w_k,$$

which upon rearranging implies that

$$w_k \geq \frac{(1 - \epsilon)n - k}{(1 - \epsilon)n(n - k)}.$$

In particular, for $k = (1 - 2\epsilon)n$, we have

$$w_{(1-2\epsilon)n} \geq \frac{1}{2(1-\epsilon)n}.$$

Letting S' to be the $(1 - 2\epsilon)n$ points with largest weight, we have that for all $i \in S'$, $w_i \geq \frac{1}{2(1-\epsilon)n}$. We will use this to now bound the $\mathcal{X}_k$ norm of $\Sigma_{S'} = \frac{1}{|S'|} \sum_{i \in S'} x_i x_i^\top$. Consider an arbitrary $M \in \mathcal{X}_k$, we have

$$\sum_{i \in S'} \frac{1}{|S'|} \langle x_i x_i^\top, M \rangle = \sum_{i \in S'} \frac{1}{(1-2\epsilon)n} \langle x_i x_i^\top, M \rangle$$

$$\leq \sum_{i \in S'} \frac{2(1-\epsilon)}{1-2\epsilon} w_i \langle x_i x_i^\top, M \rangle$$

$$\leq \sum_{i \in S} \frac{2(1-\epsilon)}{1-2\epsilon} w_i \langle x_i x_i^\top, M \rangle$$

$$\leq 9 \left(1 + \frac{\delta^2}{\epsilon} \right).$$

Since $\delta \geq \sqrt{\epsilon}$, this in turn implies the (rather loose in constants) inequality that $\|\Sigma_{S'} - I\|_{\mathcal{X}_k} \leq 20(\delta^2/\epsilon)$.

Next, we show that the mean $\mu_{S'}$ of S' is $10\delta + 10\sqrt{\epsilon}$-close to $\mu = 0$. This will essentially follow from 1) the uniform distribution $U_{S'}$ over S' is close in total variation distance to w and 2) the contribution of the tail to the mean of a bounded-covariance distribution is small.

For 1), using the notation that U_S is the uniform distribution over S (analogous to the S' notation just before), it is immediate that by the triangle inequality,

$$d_{\mathrm{TV}}(w, U_{S'}) \leq d_{\mathrm{TV}}(w, U_S) + d_{\mathrm{TV}}(U_S, U_{S'}) \leq \epsilon + 2\epsilon = 3\epsilon.$$

A standard consequence is that there exists distributions $p^{(1)}$, $p^{(2)}$ and $p^{(3)}$ such that

$$w = (1 - 3\epsilon)p^{(1)} + 3\epsilon p^{(2)} \quad \text{and} \quad U_{S'} = (1 - 3\epsilon)p^{(1)} + 3\epsilon p^{(3)}.$$

Intuitively, treating $p^{(2)}$ and $p^{(3)}$ as the "tails", we will bound their contributions to the mean under the boundedness of the covariance of w and $U_{S'}$.

Take any k-sparse unit vector direction $v \in \mathcal{U}_k$, we can bound the following variances in the direction of v:

$$3\epsilon \sum_i p_i^{(2)} \langle x_i, v \rangle^2 \leq \sum_i w_i \langle x_i, v \rangle^2 \leq 1 + \frac{\delta^2}{\epsilon},$$

$$3\epsilon \sum_i p_i^{(3)} \langle x_i, v \rangle^2 \leq \sum_i U_{S',i} \langle x_i, v \rangle^2 \leq 9 \left(1 + \frac{\delta^2}{\epsilon} \right),$$

where we used the fact that $vv^\top$ is in $\mathcal{X}_k$ for a k-sparse unit vector v.

By Jensen's inequality, we can then conclude that

$$\left| 3\epsilon \sum_i p_i^{(2)} \langle x_i, v \rangle \right| \leq \sqrt{3\epsilon} \sqrt{3\epsilon \sum_i p_i^{(2)} \langle x_i, v \rangle^2} \leq \sqrt{3\epsilon} \sqrt{1 + \frac{\delta^2}{\epsilon}} \leq \sqrt{3}(\sqrt{\epsilon} + \delta),$$

$$\left| 3\epsilon \sum_i p_i^{(3)} \langle x_i, v \rangle \right| \leq \sqrt{3\epsilon} \sqrt{3\epsilon \sum_i p_i^{(3)} \langle x_i, v \rangle^2} \leq 3\sqrt{3\epsilon} \sqrt{1 + \frac{\delta^2}{\epsilon}} \leq 3\sqrt{3}(\sqrt{\epsilon} + \delta).$$

Finally, since $U_{S'} = w - 3\epsilon p^{(2)} + 3\epsilon p^{(3)}$, by the triangle inequality, we have

$$
\begin{aligned}
|\langle \mu_{S'} - \mu, v \rangle| &= \left| \sum_i U_{S',i} \langle x_i, v \rangle \right| \\
&\leq \left| \sum_i w_i \langle x_i, v \rangle \right| + \left| 3\epsilon \sum_i p_i^{(2)} \langle x_i, v \rangle \right| + \left| 3\epsilon \sum_i p_i^{(3)} \langle x_i, v \rangle \right| \\
&\leq \delta + \sqrt{3}(\sqrt{\epsilon} + \delta) + 3\sqrt{3}(\sqrt{\epsilon} + \delta) \\
&\leq 10\delta + 10\sqrt{\epsilon},
\end{aligned}
$$

where the second inequality uses the above bounds as well as the assumption that $\|\mu_w - \mu\|_{2,k} \leq \delta$. Now that we have shown that $\mu_{S'}$ is close to μ in $2, k$ norm and $\Sigma_{S'}$ is small in the $\mathcal{X}_k$ norm, we will use the following lemma (Lemma A.3) to show that the set S' is $(\epsilon, O(\delta + \sqrt{\epsilon})$-stable with respect to μ. $\qquad \square$

Lemma A.3 (Bounded Mean and Covariance implies $O(\sqrt{\epsilon})$ stability). *Let $\mu \in \mathbb{R}^d$ and let S' be a set of samples such that $\|\mu_{S'} - \mu\|_{2,k} \leq \delta$ and $\left\| \frac{1}{|S'|} \sum_{x \in S'} (x - \mu)(x - \mu)^\top - I \right\|_{\mathcal{X}_k} \leq \frac{\delta^2}{\epsilon}$ for some $0 \leq \epsilon \leq \delta$ and $\epsilon \leq 0.5$. Then S' is (ϵ, δ', k)-stable with respect to μ where $\delta' = O(\delta + \sqrt{\epsilon})$ and $\delta' \geq \sqrt{\epsilon}$.*

Proof. Consider an arbitrary large subset $S'' \subseteq S'$ where $|S''| \geq (1 - \epsilon)|S'|$. Without loss of generality, take $\mu = 0$. Then, for an arbitrary $M \in \mathcal{X}_k$,

$$
\langle \overline{\Sigma}_{S''} - I, M \rangle = \frac{1}{S''} \sum_{i \in S''} \langle x_i x_i^\top, M \rangle - 1,
$$

which is trivially at least $-1 \geq -(\delta'^2)/\epsilon$ for $\delta' \geq \sqrt{\epsilon}$. As for the upper bound, we have

$$
\begin{aligned}
\langle \overline{\Sigma} - I, M \rangle &= \frac{1}{S''} \sum_{i \in S''} \langle x_i x_i^\top, M \rangle - 1 \\
&\leq \left(\frac{1}{S''} \sum_{i \in S'} \langle x_i x_i^\top, M \rangle \right) - 1 \\
&\leq \frac{1}{1 - \epsilon} \left(1 + \frac{\delta^2}{\epsilon} \right) - 1 \\
&= \frac{\frac{\delta^2}{\epsilon} + \epsilon}{1 - \epsilon} \\
&\leq \frac{2}{\epsilon}(\delta^2 + \epsilon^2) \\
&\leq \frac{\delta'^2}{\epsilon},
\end{aligned}
$$

for some $\delta' = \Theta(\delta + \sqrt{\epsilon})$.

We now bound the error in the mean of S'' in $2, k$ norm. First, observe that, for an arbitrary k-sparse unit vector v,

$$\left| \frac{1}{|S'|} \sum_{i \in S' \setminus S''} \langle x_i, v \rangle \right| = \left| \frac{1}{|S'|} \sum_{i \in S'} \mathbb{1}[x_i \in S' \setminus S''] \langle x_i, v \rangle \right|$$

$$\leq \frac{1}{|S'|} \sum_{i \in S'} |\mathbb{1}[x_i \in S' \setminus S''] \langle x_i, v \rangle|$$

$$\leq \sqrt{\epsilon} \sqrt{\frac{1}{|S'|} \sum_{i \in S'} \langle x_i, v \rangle^2}$$

$$\leq \sqrt{\epsilon} \sqrt{1 + \frac{\delta^2}{\epsilon}}$$

$$= \sqrt{\epsilon + \delta^2} ,$$

where the second inequality is an application of Hölder's inequality, and the third inequality uses the fact that for a unit k-sparse vector v, $vv^\top$ is in $\mathcal{X}_k$.

Thus, again for an arbitrary k-sparse unit vector v,

$$|\langle \mu_{S''}, v \rangle| = \left| \frac{1}{|S''|} \sum_{i \in S''} \langle x_i, v \rangle \right|$$

$$\leq \frac{1}{1 - \epsilon} \left| \frac{1}{|S'|} \sum_{i \in S''} \langle x_i, v \rangle \right|$$

$$\leq 2 \left(\left| \frac{1}{|S'|} \sum_{i \in S'} \langle x_i, v \rangle \right| + \left| \frac{1}{|S'|} \sum_{i \in S' \setminus S''} \langle x_i, v \rangle \right| \right)$$

$$\leq 2(\delta + \sqrt{\epsilon + \delta^2}) = O(\delta + \sqrt{\epsilon}) = \delta'.$$

$\square$

Proposition A.4 (Bounded Covariance and Stability). *Let $\mu \in \mathbb{R}^d$ and let S be a set of n samples. Let $w \in \Delta_{n,\epsilon}$ over the set of samples S such that $\|\sum_i w_i(x_i - \mu)(x_i - \mu)^\top\|_{\mathcal{X}_k} \leq r$ for some $r > 0$. Then $\|\mu_w - \mu\|_{2,k} \leq \sqrt{r}$.*

Proof. For every k-sparse unit vector v, $vv^\top$ is in $\mathcal{X}_k$, and thus for every sparse unit vector v, we have that $\sum_i w_i \langle x_i - \mu, v \rangle^2 \leq r$. Applying Cauchy-Schwarz inequality, we get that for any sparse unit vector v, it follows that $\sum_i w_i \langle x_i - \mu, v \rangle \leq \sqrt{\sum_i w_i \langle x_i - \mu, v \rangle^2} \leq \sqrt{r}$. $\square$

With Proposition A.4 and Lemma A.2, we can prove Proposition A.1.

Proof of Proposition A.1. Without loss of generality, we will assume that $\sigma = 1$. By Proposition A.4, we have that $\|\mu_w - \mu\|_{2,k} \leq \sqrt{B}$. We thus have a weighting $w \in \Delta_{n,\epsilon}$, where $\|\mu_w - \mu\|_{2,k} \leq \delta_0$ and $\|\overline{\Sigma}_w - I\|_{\mathcal{X}_k} \leq \delta_0^2/\epsilon$ for $\delta_0 = \sqrt{B} + 1$, where we use triangle inequality on the $\|\cdot\|_{\mathcal{X}_k}$ norm. By Lemma A.2, we know that there exists a set S' such that $|S'| \geq (1 - 2\epsilon)n$ and S' is (ϵ, δ, k)-stable with respect to μ and σ, where $\delta = O(\delta_0 + \sqrt{\epsilon}) = O(\sqrt{\epsilon} + \sqrt{B} + 1) = O(\sqrt{B} + 1)$. $\square$

A.2 Median of Means

Fact 2.2 (Median-of-Means Pre-Processing). *Suppose there is an efficient algorithm such that, on input $\sigma \in \mathbb{R}_+$ and a 0.1-corrupted set of $n \gg k^2 \log d + \log(1/\tau)$ samples from a distribution D with mean μ and covariance Σ with $\|\Sigma\|_{\mathcal{X}_k} \leq \sigma^2$ and $\mathbb{E}_{X \sim D}[(X_j - \mu_j)^4] = O(\sigma^4)$ for each coordinate $j \in [d]$, returns $\hat{\mu}$ such that $\|\hat{\mu} - \mu\|_{2,k} \leq O(\sigma)$ with probability at least $1 - \tau$.*

Then, there is an efficient algorithm such that, on input $\epsilon \in (0, 0.1)$ and an ϵ-corrupted set of $n \gg (k^2 \log d + \log(1/\tau))/\epsilon$ samples from a distribution with mean μ and covariance Σ, satisfying

$\|\Sigma\|_{\mathcal{X}_k} \leq 1$ *satisfying* $\mathbb{E}_{X \sim D}[(X_j - \mu_j)^4] = O(1)$ *for every coordinate* $j \in [d]$, *returns a mean estimate* $\hat{\mu}$ *such that* $\|\hat{\mu} - \mu\|_{2,k} \leq O(\sqrt{\epsilon})$ *with probability at least* $1 - \tau$.

Proof. The new algorithm simply performs median-of-means preprocessing as defined in Section 2 before the fact statement, yielding g new samples that are fed into the algorithm that works with constant corruption. The uncorrupted new samples, namely the ones that are the sample mean of groups containing no originally corrupted samples, are distributed i.i.d. according to the distribution D' which has mean μ, and covariance $\Sigma' = (g/n)\Sigma$, with axis-wise fourth moment $\mathbb{E}_{Y \sim D'}[(Y_j - \mu_j)^4]$ being bounded by $C(g^2/n^2)\,\mathbb{E}_{X \sim D}[(X_j - \mu_j)^4]$ for every $j \in [d]$ for some constant $C > 0$, obtained by the following fact:

Fact A.5. *(Marcinkiewicz-Zygmund inequality) Recall the notation* $\|X\|_{L_s}$ *for a centered random variable* X, *defined as* $\mathbb{E}[|X|^s]^{1/s}$. *Let* $W_1, \ldots, W_m, W$ *be identical and independent centered random variables on* $\mathbb{R}$ *with a finite* $\|W\|_{L_s}$ *norm for* $s \geq 2$. *Then,*

$$\left\| \frac{1}{m} \sum_{i=1}^{m} W_i \right\|_{L_s} \leq \frac{3\sqrt{s}}{\sqrt{m}} \|W\|_{L_s}.$$

First note that we give g samples to the original algorithm, and $g = \Omega(\epsilon n) = \Omega(k^2 \log d + \log(1/\tau))$ by definition. Next, we need to check that the *normalized* axis-wise 4th moment of D' is $O(1)$ times the (bound on the) $\mathcal{X}_k$-norm of the covariance matrix, that is, for all $j \in [d]$, it holds that $(\mathbb{E}_{X \sim D'}[(X_j - \mu_j)^4])^{1/4} \leq O(\sigma^4)$ and $\|\Sigma'\|_{\mathcal{X}_k} = O(\sigma^2)$. By the calculations at the end of the previous paragraph and the assumptions in the statement, we note that this is true for $\sigma = O(\sqrt{g/n})$.

Lastly, we check that, by the scale-invariance of the original algorithm that works with constant corruption, the estimation error of the final algorithm is upper bounded by $O(\sigma\|\Sigma\|_{\mathcal{X}_k}) = O(\sqrt{(g/n)\|\Sigma\|_{\mathcal{X}_k}}) = O(\sqrt{g/n}) = O(\sqrt{\epsilon})$ as desired. $\qquad\square$

A.3 $\mathcal{X}_k$-Norm

Lemma A.6. *Let* $A \in \mathbb{R}^{d \times d}$ *be a symmetric matrix such that* $|A_{i,i}| \leq \eta_1$ *for each* $i \in [d]$, *and* $|A_{i,j}| \leq \eta_2$ *for each* $i \neq j \in [d] \times [d]$. *Then* $\|A\|_{\mathcal{X}_k} \leq \eta_1 + k\eta_2$.

Proof. Let $A = B + C$, where B is a diagonal matrix and C is diagonal-free. Then we have the following using triangle inequality: $\|A\|_{\mathcal{X}_k} \leq \|B\|_{\mathcal{X}_k} + \|C\|_{\mathcal{X}_k}$. Thus it suffices to bound each of these terms by 1.

$$\|B\|_{\mathcal{X}_k} \leq \sup_{M: \sum_{i=1}^{d} |M_{i,i}| \leq 1} \langle B, M \rangle = \|B\|_\infty \leq \eta_1,$$

where we use that B is a diagonal matrix with entry at most η_1.

$$\|C\|_{\mathcal{X}_k} \leq \sup_{M: \|M\|_1 \leq k} \langle C, M \rangle = \sup_{M: \|M\|_1 \leq k} \|C\|_\infty \|M\|_1 \leq k\eta_2.$$

$\qquad\square$

A.4 Truncation

We show how truncation can increase the spectral norm of covariance from 1 to $\omega(1)$.

Consider the distribution which, with probability $1/(2k)$, returns a vector where each coordinate is independent $-\sqrt{k}$ with probability $2/3$ and $2\sqrt{k}$ with probability $1/3$. Otherwise, with probability $1 - 1/(2\sqrt{k})$, the distribution returns the origin. The mean of the distribution is the origin, and the covariance is I.

Now consider the truncation $h_{0,\sqrt{k}}$, which truncates at distance $\sqrt{k}$ from the origin. Let Y be the resulting random variable. The mean of Y, μ', is thus equal to $(1/2k)(-\sqrt{k}/3, \ldots, -\sqrt{k}/3) = -1/(6\sqrt{k})\mathbf{v}$, where $\mathbf{v}$ is the all ones vector. The norm of μ' is $\Theta(\sqrt{d/k})$. Since the distribution returns the origin with constant probability (asymptotically tending to 1), the variance of Y along the direction of μ', which is $\mathbf{v}/\sqrt{d}$, is at least $\Omega(d/k) = \omega(1)$.

B   Concentration and Truncation

Example 1. *For any number of moments $t \geq 2$, there is a distribution X satisfying the following conditions: (i) The mean of X is 0, and for every unit vector v, the t^{th} moment in direction v is upper bounded by 1, that is, $\mathbb{E}[|\langle v, x \rangle|^t] \leq 1$ for $t \geq 2$, (ii) If S is an arbitrary set of $n \leq o(d^{2/t})$ points from the support of X, then the set S cannot be $(\epsilon, O(\sqrt{\epsilon}), k)$-stable, for any $\epsilon > 0$, with respect to the mean of the distribution. As a corollary, no subset of S can be stable either.*

Proof. For $j \in [d]$, let e_j be the vector that is 1 on the j-th coordinate and 0 otherwise. For a fixed r, consider the distribution P, supported uniformly on the $2d$ points $S = \{\pm r e_1, \pm r e_2, \ldots, \pm r e_d\}$.

It follows that P is a zero mean distribution. The covariance of the distribution P is $\sum_j (1/d) r^2 e_i e_i^\top = (r^2/d) I$.

Furthermore, for any unit vector v and $t \geq 2$, we have that the t-th moment in the direction v is bounded as follows:

$$\mathbb{E}[|v \cdot X|^t] = \sum_{j=1}^{d} \frac{1}{d} |v_j|^t r^t = \frac{r^t}{d} \|v\|_t^t \leq \frac{r^t}{d} \|v\|_2^t \leq \frac{r^t}{d},$$

where we use that $t \geq 2$ and $\|v\|_t \leq \|v\|_2$ for any vector v. Thus, we choose $r = d^{1/t}$ for the distribution.

Now we show the second claim, that *any* set of at most $\Omega(d^{2/t})$ samples from this distribution cannot be stable.

Let S be any (multi-)set of n points from the support of X. Let $x_1 \in S$. Since x_1 is 1-sparse and has ℓ_2 norm r, we have that $x_1 x_1^\top / r^2$ belongs to $\mathcal{X}_k$. Thus we have the following:

$$\left\| \frac{1}{n} \sum_{i \in S'} x_i x_i^\top \right\|_{\mathcal{X}_k} \geq \left\langle \frac{1}{n} \sum_{i \in S'} x_i x_i^\top, \frac{1}{r^2} x_1 x_1^\top \right\rangle \geq \frac{\|x_1\|^4}{r^2 n} = \frac{r^2}{n}$$

Thus, for r^2/n to be upper bounded by a constant, n has to be $\Omega(d^{2/t})$. $\qquad\square$

Lemma 3.1 (Truncation in ℓ_∞). *Let P be a distribution over $\mathbb{R}^d$ with mean μ_P and covariance Σ_P, with $\|\Sigma\|_{\mathcal{X}_k} \leq \sigma^2$ for some $\sigma^2 > 0$. Let $X \sim P$ and assume that for all $j \in [d]$, $\mathbb{E}[(X - \mu_P)_j^4] \leq \sigma^4 \nu^4$ for some $\nu \geq 1$. Let $b \in \mathbb{R}^d$ be such that $\|b - \mu\|_\infty \leq a/2$ and $a := 2\sigma\sqrt{k/\epsilon}$ for some $\epsilon \in (0, 1)$. Define Q to be the distribution of $Y := h_{a,b}(X)$. Let the mean and covariance of Q be μ_Q and Σ_Q respectively. Then the following hold:*

(1) $\|\mu_P - \mu_Q\|_\infty \leq \sigma\sqrt{\epsilon/k}$

(2) $\|\mu_P - \mu_Q\|_{2,k} \leq \sigma\sqrt{\epsilon}$

(3) $\|\Sigma_P - \Sigma_Q\|_{\mathcal{X}_k} \leq 3\sigma^2 \epsilon \nu^4$

(4) For all $i \in [d]$, $\mathbb{E}[(Y - \mu_Q)_i^4] \leq 8\nu^4 \sigma^4$

(5) $\|Y - \mu_Q\|_\infty \leq 2a = 4\sigma\sqrt{k/\epsilon}$ almost surely.

Proof. Let $Y := h_{a,b}(X)$ and denote $\mu := \mu_P$. Fix a $i \in [d]$. Since $|\mu_i - b_i| \leq a/2$ and we threshold at the radius a, we have the following:

$$|Y_i - \mu_i| \leq |X_i - \mu_i|, \quad \text{and} \quad |X_i - Y_i| \leq |X_i - \mu_i|. \tag{5}$$

Let $\mathcal{E}_i$ be the event that $Y_i \neq X_i$. We get the following by Markov's inequality and moment bounds:

$$\mathcal{P}(\mathcal{E}_i) = \mathbb{P}(|X_i - b_i| > a) \leq \mathbb{P}(|X_i - \mu_i| \geq a/2) \leq \min\left(4\frac{\sigma^2}{a^2}, 16\frac{\sigma^4 \nu^4}{a^4}\right) = \min\left(\frac{\epsilon}{k}, \frac{\epsilon^2 \nu^4}{k^2}\right). \tag{6}$$

1. We can verify the following relation using Equation (5):
$$|Y_i - X_i| \le \mathbb{1}_{\mathcal{E}_i} \cdot (|Y_i - X_i|) \le \mathbb{1}_{\mathcal{E}_i} \cdot (|X_i - \mu_i|). \tag{7}$$
Applying Cauchy-Scharz on the above inequality gives the desired conclusion:
$$|\mathbb{E}[Y_i] - \mu_i| = |\mathbb{E}[Y_i - X_i]| \le \mathbb{E}\left[\mathbb{1}_{\mathcal{E}} \cdot (|X_i - \mu_i|)\right] \le \sqrt{\mathbb{P}(\mathcal{E})}\sqrt{\mathbb{E}\left[|X_i - \mu_i|^2\right]} \le \sigma\sqrt{\frac{\epsilon}{k}},$$
where we use that variance of X_i is at most σ^2 and use Equation (6).

2. This follows directly from above.

3. By Lemma A.6, it suffices to show that $\|\Sigma_Q - \Sigma_P\|_\infty \le 3\sigma^2\epsilon\nu^4/k$. Using triangle inequality, we obtain the following:
$$\begin{aligned}
\|\Sigma_P - \Sigma_Q\|_\infty &= \left\| \mathbb{E}[(X - \mu_P)(X - \mu_P)^\top] - \mathbb{E}[(Y - \mu_P)(Y - \mu_P)^\top] \right. \\
&\qquad + \left. (\mu_Q - \mu_P)(\mu_Q - \mu_P)^\top \right\|_\infty \\
&\le \left\| \mathbb{E}[(X - \mu_P)(X - \mu_P)^\top] - \mathbb{E}[(Y - \mu_P)(Y - \mu_P)^\top] \right\|_\infty \\
&\qquad + \left\| (\mu_Q - \mu_P)(\mu_Q - \mu_P)^\top \right\|_\infty.
\end{aligned}$$
By the first part above, we have that $\|(\mu_Q - \mu_P)(\mu_Q - \mu_P)^\top\|_\infty \le \sigma^2\epsilon/k \le \sigma^2\nu^4\epsilon/k$, where we use that $\nu \ge 1$. We will thus focus on the first term. Without loss of generality, we will assume that $\mu_P = 0$ for the remainder of this proof. Thus for any $i, j \in [d]$, we thus need to upper bound $\mathbb{E}[|X_iX_j - Y_iY_j|]$.
$$\begin{aligned}
\mathbb{E}[|X_iX_j - Y_iY_j|] &\le \mathbb{E}[|X_i\|X_j - Y_j|] + \mathbb{E}[|Y_j\|X_i - Y_i|] \\
&\le \mathbb{E}[|X_i\|X_j| \cdot \mathbb{1}_{\mathcal{E}_j}] + \mathbb{E}[|X_i\|X_j| \cdot \mathbb{1}_{\mathcal{E}_i}] \qquad \text{(Using Equation (7))} \\
&\le \sqrt{\mathbb{E}[|X_iX_j|^2]}\left(\sqrt{\mathbb{P}(\mathcal{E}_i)} + \sqrt{\mathbb{P}(\mathcal{E}_j)}\right) \\
&\le (\mathbb{E}[X_i^4])^{1/4}(\mathbb{E}[X_j^4])^{1/4}\left(\sqrt{\mathbb{P}(\mathcal{E}_i)} + \sqrt{\mathbb{P}(\mathcal{E}_j)}\right) \\
&= \sigma^2\nu^2\left(2\frac{\epsilon\nu^2}{k}\right) \\
&= \frac{2\sigma^2\nu^4\epsilon}{k}.
\end{aligned}$$
Combining the above with Lemma A.6, we get that the $\|\Sigma_P - \Sigma_Q\|_{\mathcal{X}_k} \le 3\sigma^2\epsilon\nu^4$.

4. Fix an $i \in [d]$. We use the triangle inequality and Equation (7) to get the following:
$$\mathbb{E}[(Y - \mu_Q)_i^4] \le 4(\mathbb{E}[(Y - \mu_P)_i^4]) + 4\|\mu_P - \mu_Q\|_\infty^4 \le 4\sigma^4\nu^4 + 4\sigma^4\epsilon^2/k^2 \le 8\sigma^4\nu^4,$$
where the last inequality uses that $\nu \ge 1$ and $\epsilon \le 1$.

5. This follows by definition of the random variable Y, the function $h_{a,b}$, and the parameter a.

$\square$

Fact B.1 (VC inequality). *Let $\mathcal{F}$ be a family of boolean functions over $\mathcal{X}$ with VC dimension r and let $S = \{x_1, \ldots, x_n\}$ be a set of n i.i.d. data points from a distribution P over $\mathcal{X}$. If $n \gg c(r + \log(1/\tau))/\gamma^2$, then with probability $1 - \tau$, for all $f \in \mathcal{F}$, we have that*
$$\left|\sum_{i=1}^n \frac{f(x_i)}{n} - \mathbb{E}_P[f(x)]\right| \le \gamma.$$

Lemma B.2 (Uniform concentration over $\mathcal{A}_{k,P}$). *Let S be a set of n i.i.d. data points from a distribution P, and let $\mathcal{A}_{k,P}$ be as defined in Equation (9). There exists a constant $c > 0$ such that if $n \ge c(k^2\log d + \log(1/\tau))/(q^2)$, then Equation (10) holds with probability at least $1 - \tau$ over the set S of n i.i.d. points from distribution P.*

Proof. Let Q be the distribution of $y := xx^\top$. Let $\mathcal{F} := \{\mathbb{1}_{y \cdot A > s_1} : A \in \mathcal{A}_k\}$. Suppose for now that VC dimension of $\mathcal{F}$ is less than $Ck^2 \log(d)$. Then the standard VC inequality (Fact B.1) implies that if $n \geq c(k^2 \log d + \log(1/\tau))/(q^2)$, then Equation (10) holds because under $y \sim Q$, $\mathbb{P}(y \cdot A > s_1) \leq q$ for all $A \in \mathcal{A}_{k,P}$. Thus it remains to show an upper bound on the VC dimension of $\mathcal{F}$. Since $\mathcal{F}$ corresponds to a family of linear functions that are k^2-sparse in d^2 dimensional space, [AV19, Theorem 6] implies that the VC dimesion is at most $4k^2 \log(3d)$. This completes the proof. $\qquad\square$

C Stability with High Probability

Theorem C.1. *Let S be a set of n i.i.d. data points from a distribution P over $\mathbb{R}^d$. Let the mean of P be μ, and covariance Σ such that $\|\Sigma\|_{\mathcal{X}_k} \leq \sigma^2$, and for all $j \in [d]$, $\mathbb{E}[(X_j - \mu)^4] \leq \nu^4$. Suppose P is supported over the set $\{x : \|x - \mu\|_\infty \leq \sigma \times r \times \sqrt{k}\}$. If $n = \Omega(k^2 \log d + \log(1/\tau))$, then, with probability $1 - \tau$, there exists a set $S' \subset S$ such that:*

1. *$|S'| \geq 0.98n$*

2. *S' is $(0.01, \delta, k)$-stable with respect to μ and σ where $\delta = O(\max(1, r^2, \nu^2/\sigma^2))$.*

Proof. In the following proof, we will use notations q, s_1, s_2, s_3, V_Z and B, all of which are either constants or functions of σ, r and ν in the theorem statement. The functions are explicitly chosen in Appendix C.1.

We will assume $\mu = 0$ without loss of generality. Instead of directly showing the existence of subset $S' \subseteq S$ (with high probability over the samples S) that is stable, Proposition A.1 in Appendix A lets us show the following simpler condition: let $\Delta_{n,\epsilon}$ be the set of weights/distributions w such that $w_i \leq 1/(1 - \epsilon)$, then there exists a weighting $w \in \Delta_{n,0.01}$ such that $\|\Sigma_w\|_{\mathcal{X}_k} \leq B$ for the function B chosen in Appendix C.1, which satisfies $B = O(\sigma^2 \max(1, r^2, \nu^2/\sigma^2))$. That is, for the following proof, we just need to prove that $\min_{w \in \Delta_{n,0.01}} \|\Sigma_w\|_{\mathcal{X}_k} \leq B$.

We proceed as follows:

$$\min_{w \in \Delta_{n,0.01}} \|\Sigma_w\|_{\mathcal{X}_k} = \min_{w \in \Delta_{n,0.01}} \max_{M \in \mathcal{X}_k} \langle M, \Sigma_w \rangle = \max_{M \in \mathcal{X}_k} \min_{w_M \in \Delta_{n,0.01}} \langle M, \Sigma_w \rangle$$

where the last equality is a straightforward application of the minimax theorem for a minimax optimization problem with independent convex domains and a bilinear objective. It thus suffices to show the following: with probability $1 - \tau$,

$$\forall M \in \mathcal{X}_k : \ |\{x \in S : x_i^\top M x_i > B\}| \leq 0.01|S| \tag{8}$$

from which we can construct the weighting w_M as uniform distribution over the elements outside the above set.

Define the following sets of sparse matrices:

$$\mathcal{A}_k := \left\{ A \in \mathbb{R}^{d \times d} : \|A\|_0 \leq k^2, \|A\|_F \leq 1 \right\},$$
$$\mathcal{A}_{k,P} := \left\{ A \in \mathcal{A}_k : \mathbb{P}\left\{ x^\top A x \geq s_1 \right\} \leq q \right\}. \tag{9}$$

where q and s_1 are chosen in Appendix C.1. If $n \gtrsim (k^2 \log d + \log(1/\tau))/(q^2)$, then a standard covering/VC-dimension bound (see Lemma B.2 for details) implies that the following event holds with probability $1 - \tau$:

$$\forall A \in \mathcal{A}_{k,P} : \ |\{x \in S : x_i^\top A x_i > s_1\}| \leq 2 \times q \cdot |S|. \tag{10}$$

Our choice of q is a constant (cf. Appendix C.1) and thus the required sample complexity for Equation (10) to hold is $\Omega(k^2 \log d + \log(1/\tau))$. We will now show that the event in Equation (10) implies that the event in Equation (8) holds.

Suppose, for the sake of contradiction, that the event in Equation (8) does not hold. Then there exists a $M \in \mathcal{X}_k$ such that $|\{x \in S : x_i^\top M x_i > B\}| > 0.01|S|$. We will show the existence of a matrix Q violating event Equation (10), via the probabilistic method, to reach the desired contradiction.

Fixing an M that violates Equation (8), consider the random matrix Q where each entry $Q_{i,j}$ is sampled independently from the following distribution, defined using the constant s_2 chosen in Appendix C.1:

$$Q_{i,j} := \begin{cases} M_{i,j}, & \text{with prob. 1 if } |M_{i,j}| \geq s_2/k, \\ \frac{s_2}{k}\operatorname{sign}(M_{i,j}), & \text{with prob. } |kM_{i,j}|/s_2 \text{ if } |M_{i,j}| \leq s_2/k, \\ 0, & \text{with remaining prob. if } |M_{i,j}| \leq s_2/k \end{cases} \tag{11}$$

Defining $p_{i,j}$ to be $\min(1, k|M_{i,j}|/s_2)$, then $Q_{i,j}$ is equivalently $M_{i,j}/p_{i,j}$ with probability $p_{i,j}$ and 0 otherwise.

We will show that the following events hold simultaneously with non-zero probability, leading to a contradiction to event Equation (10):

1. $Q \in s_3 \mathcal{A}_{k,P}$

2. $|\{x \in S : x_i^\top Q x_i > s_3 \times s_1\}| > 2 \times q \cdot |S|$

where s_3 is also a constant, larger than 2, and explicitly chosen in Appendix C.1. Using different techniques, we will show that the first condition holds with probability at least $1 - 2 \times 10^{-6}$ and the second condition holds with probability at least 4×10^{-6}, thus implying that the events hold simultaneously with non-zero probability.

Condition 1 We begin with the following lemma showing that Q lies in $s_3 \mathcal{A}_k$ with high probability.

Lemma C.2 (Q lies in $s_3 \mathcal{A}_k$ with high probability). *Let Q be generated as described in Equation (11), for an $M \in \mathcal{X}_k$. Then with probability except $(1/s_2) + (s_2/s_3^2)$, we have that that $Q \in s_3 A_k$.*

Proof. The expected sparsity of Q is at most $\sum_{i,j} \frac{k}{s_2}|M_{i,j}| \leq \frac{k^2}{s_2}$ since $|M|_1 \leq k$. Thus, by Markov's inequality, except with $1/s_2$ probability, Q and hence Q/s_3 is k^2-sparse. We also have to show that with probability at least $1 - 10^8$, $\|Q\|_F \leq s_3$.

$$\mathbb{E}\|Q\|_F^2 \leq \sum_{i,j} \left(\frac{s_2}{k}\right)^2 \left(\frac{k|M_{i,j}|}{s_2}\right) = \frac{s_2|M_{i,j}|}{k} = s_2. \tag{12}$$

Again, by Markov's inequality, we get that with probability except s_2/s_3^2, the Frobenius norm of Q is at most s_3. The lemma statement follows from the union bound. $\square$

Our choice of constants in Appendix C.1 would ensure that the failure probability in Lemma C.2 is at most 10^{-6}.

$$\frac{1}{s_2} + \frac{s_2}{s_3^2} \leq 10^{-6}. \tag{13}$$

It remains to show that Q belongs to $s_3 \mathcal{A}_{k,p}$ with high (constant) probability, i.e., with probability 10^{-6} over sampling of Q, we have that $\mathbb{P}_{x \sim P}(x^\top Q x > s_3 \times s_1 | Q) \leq q$. Let $R := x^\top Q x$, where both x and Q are sampled independently from P and Equation (11) respectively.

To show this, we use the following the lemma for a sufficient condition involving sampling both x and Q.

Lemma C.3. *Consider a probability space over the randomness of independent variables X and Y. Suppose the event E (over pairs (X, Y)) happens with probability at least $1 - \alpha\beta$ for some $\alpha, \beta \in [0, 1]$. Then, it must be the case that, with probability at least $1 - \alpha$ over the sampling of X, the conditional probability of E given X is at least $1 - \beta$.*

Proof. For the sake of contradiction, suppose the lemma conclusion is false. Then

$$\mathbb{P}_{X,Y}(E) = \int_Y \mathbb{P}(E|X) \, d\mathbb{P}(X) < (1 - \alpha) + \alpha(1 - \beta) = 1 - \alpha\beta,$$

which contradicts the premise. $\square$

It thus suffices to show that with probability $1 - 10^{-6} \times q$ over both x and Q, $R \leq s_3 \times s_1$.

Lemma C.4. *Let $R = x^\top Q x$, where Q is independently drawn from the distribution in [Equation](11) and x is drawn independently from P. Under the assumptions of [Theorem C.1](),*

$$\mathbb{P}\{R > s_3 \times s_1\} \leq \frac{\sigma^2}{s_1} + \frac{4}{s_3} + \frac{s_2 \times \nu^4}{s_3 \times s_1^2}, \tag{14}$$

Proof. We consider three exhaustive events, over x and Q, of $\mathcal{E} := \{R > s_3 \times s_1\}$, and bound the probability of each of them :

1. $\mathcal{E}_1 := \{(x, Q) : \mathbb{E}[R|x] > s_1\}$. Since $\mathbb{E}[R|x] = x^\top M x$, the event corresponds to $\{x : x^\top M x > s_1\}$. We have that $\mathbb{E}[x^\top M x] = \langle \Sigma, M \rangle \leq \|\Sigma\|_{\mathcal{X}_k} = \sigma^2$. By Markov's inequality, $\mathbb{P}(\mathcal{E} \cap \mathcal{E}_1) \leq \mathbb{P}(\mathcal{E}_1) \leq \sigma^2/(s_1)$.

2. $\mathcal{E}_2 := \{(x, Q) : x \in \mathcal{F}\}$, where $\mathcal{F}$ is the following event over x: $\mathcal{F} = \{x : \mathbb{E}[R|x] \leq s_1, \mathrm{Var}(R|x) \leq s_3 \times s_1^2\}$. Observe that conditioned on $x \in \mathcal{F}$, we have that $R|x$ is a random variable with mean at most s_1 and variance at most $s_3 \times s_1^2$. Thus for each such $x \in \mathcal{F}$, the conditional probability that $R > s_3 \times s_1$ is at most $s_3 s_1^2/((s_3 - 1)^2 s_1^2)$ by Chebyshev's inequality. We thus get that $\mathbb{P}(\mathcal{E}_2 \cap \mathcal{E}) \leq \mathbb{P}(\mathcal{E}|\mathcal{E}_2) = \mathbb{P}(\mathcal{E}|x \in \mathcal{F}) \leq 4/(s_3)$, where we use that $s_3 \geq 2$.

3. $\mathcal{E}_3 := \{(x, Q) : \mathrm{Var}(R|x) \geq s_3 \times s_1^2\}$. We will upper bound $\mathbb{P}(\mathcal{E}_3)$. We first calculate the $\mathrm{Var}(R|x)$ using the independence of entries of Q as follows:

$$\mathrm{Var}(R|x) = \sum_{i,j} x_i^2 x_j^2 \, \mathrm{Var}(Q_{i,j}) = \sum_{i,j:|M_{i,j}| \leq s_2/k} x_i^2 x_j^2 |M_{i,j}| \left(\frac{s_2}{k} - |M_i, j|\right).$$

To show that $\mathrm{Var}(R|x)$ is small with high probability, we will upper bound $\mathbb{E}[\mathrm{Var}(R|x)]$.

$$
\begin{aligned}
\mathbb{E}[\mathrm{Var}(R|x)] &= \sum_{i,j:|M_{i,j}| \leq s_2/k} |M_{i,j}| \left(\frac{s_2}{k} - |M_i, j|\right) \mathbb{E}[x_i^2 x_j^2] \\
&\leq \sum_{i,j} \frac{s_2}{k} |M_{i,j}| \, \mathbb{E}[x_i^2 x_j^2] \\
&\leq \frac{s_2 \times \|M\|_1 \times \nu^4}{k} \qquad \text{(using } \mathbb{E}[x_i^2 x_j^2] \leq \sqrt{\mathbb{E}[x_i^4] \, \mathbb{E}[x_j^4]} = \nu^4\text{)} \\
&\leq s_2 \times \nu^4. \qquad\qquad\qquad \text{(using } \|M\|_1 \leq k\text{)}
\end{aligned}
$$

Thus Markov's inequality implies that $\mathbb{P}(\mathcal{E} \cap \mathcal{E}_3) \leq \mathbb{P}(\mathcal{E}_3) \leq (s_2 \times \nu^4)/(s_3 \times s_1^2)$.

Taking the union bound, we get the desired result. $\qquad\qquad\qquad\qquad\qquad\qquad\square$

As reasoned above, we want the failure probability in [Equation (14)]() to be less than $10^{-6} \times q$. That is,

$$\frac{\sigma^2}{s_1} + \frac{4}{s_3} + \frac{s_2 \times \nu^4}{s_3 \times s_1^2} \leq 10^{-6} \times q. \tag{15}$$

In [Appendix C.1](), we choose s_1, s_2, s_3 and q such that the bound holds. This, by the reasoning after [Lemma C.3](), guarantees that Q satisfies the extra condition for $s_3 \mathcal{A}_{k,p}$ (on top of being in $s_3 \mathcal{A}_k$) with probability at least $1 - 10^{-6}$.

Taking a union bound, with failure probabilities 10^{-6} (for Q being in $s_3 \in \mathcal{A}_k$, [Lemma C.2]()) and 10^{-6} for satisfying the additional criterion for being in $s_3 \mathcal{A}_{k,p}$, we conclude that Condition 1 happens with probability $1 - 2 \cdot 10^{-6}$.

Condition 2 The second condition makes crucial use of the Berry-Esseen theorem, which we restate as follows.

Fact C.5 (Berry-Esseen Theorem for sums of independent variables). *Consider a random variable* $\xi = \sum_i \xi_i$, *where the variables* ξ_i *are independent (but not necessarily identical) and each of them has finite third moment. Denote* μ_i *as* $\mathbb{E}[\xi_i]$, σ_i^2 *as* $\mathrm{Var}(\xi_i)$ *and* ρ_i *as the third central absolute moment, namely* $\rho_i = \mathbb{E}[|\xi_i - \mu_i|^3]$. *Then,*

$$d_{\mathrm{K}}(\xi, \mathcal{N}(\sum_i \mu_i, \sigma_i^2)) \le 0.57 \frac{\sum_i \rho_i}{(\sum_i \sigma_i^2)^{1.5}} = 0.57 \frac{\sum_i \rho_i}{(\mathrm{Var}(\xi))^{1.5}}$$

where d_{K} *is the Kolmogorov distance between two distributions (namely, the* ℓ_∞ *distance between the cumulative density functions).*

Define the random variable Z to be

$$Z = \sum_i \mathbb{1}_{(x_i x_i^\top) \bullet Q > s_3 \times s_1}. \tag{16}$$

The second condition is equivalent to saying that $Z > 2 \times q \times |S|$, which we show to happen with probability at least 4×10^{-6}.

The strategy is to lower bound $\mathbb{E}[Z]$, and then use Paley-Zygmund to show that Z is large with constant probability. To lower bound the expectation, for any i such that $(x_i x_i^\top) \bullet M > B$, we want to lower bound $\mathbb{P}_Q((x_i x_i^\top) \bullet Q > s_3 \times s_1)$, using either Chebyshev's inequality or the Berry-Esseen theorem. First, note that for these i, $\mathbb{E}[(x_i x_i^\top) \bullet Q] = (x_i x_i^\top) \bullet M > B$ by our assumption. If $\mathrm{Var}[(x_i x_i^\top) \bullet Q] \le V_Z$, where V_Z is a fixed function of r and σ chosen in Appendix C.1, then by Chebyshev's inequality, we have

$$\mathbb{P}\left((x_i x_i^\top) \bullet Q \ge s_3 \times s_1\right) \ge \mathbb{P}\left((x_i x_i^\top) \bullet Q \ge B - 10 \times \sqrt{V_Z}\right) \ge 0.99. \tag{17}$$

where the first inequality is true by our choice of s_1, s_3, V_Z and B in Appendix C.1. Otherwise, we have the case where $\mathrm{Var}[(x_i x_i^\top) \bullet Q] > V_Z$. In this case, we treat $(x_i x_i^\top) \bullet Q$ as a sum of independent variables

$$(x_i x_i^\top) \bullet Q = \sum_{s,t} (x_i)_s (x_i)_t Q_{s,t}$$

and use the Berry-Esseen theorem, which requires bounding the sum of the third central absolute moment of the summands. Let $\rho_{s,t}$ be the third central absolute moment of $(x_i)_s (x_i)_t Q_{s,t}$. For any (s,t) such that $0 < |M_{s,t}| \le s_2/k$, we can calculate its third moment as follows:

$$
\begin{aligned}
\rho_{s,t} &= \mathbb{E}[|(x_i)_s (x_i)_t Q_{s,t} - \mathbb{E}[(x_i)_s (x_i)_t Q_{s,t}]|^3] \\
&= |(x_i)_s|^3 |(x_i)_t|^3 \frac{|M_{s,t}|^3}{p_{s,t}^3} \mathbb{E}[|\mathrm{Ber}(p_{s,t}) - p_{s,t}|^3] \\
&= |(x_i)_s|^3 |(x_i)_t|^3 \frac{|M_{s,t}|^3}{p_{s,t}^3} p_{s,t}(1 - p_{s,t})(1 - 2p_{s,t} + 2p_{s,t}^2) \\
&\le |(x_i)_s|^3 |(x_i)_t|^3 \frac{|M_{s,t}|^3}{p_{s,t}^3} p_{s,t}(1 - p_{s,t}) && \text{for all } p_{s,t} \in [0,1] \\
&\le (\sigma^2 \times r^2 \times s_2)(x_i)_s^2 (x_i)_t^2 \frac{M_{s,t}^2}{p_{s,t}^2} p_{s,t}(1 - p_{s,t}) \\
&&& \text{since } |x_i|_\infty \le \sigma \times r \times \sqrt{k} \text{ and } |M_{s,t}|/p_{s,t} = s_2/k
\end{aligned}
$$

The same inequality holds trivially for (s,t) where $|M_{s,t}| \ge s_2/k$ or $M_{s,t} = 0$ since $\rho_{s,t} = 0$ in both of these edge cases. Thus, the sum of the third central absolute moment of the summands we need for Berry-Esseen is

$$
\begin{aligned}
\sum_{s,t} \rho_{s,t} &\le (\sigma^2 \times r^2 \times s_2) \sum_{s,t} (x_i)_s^2 (x_i)_t^2 \frac{M_{s,t}^2}{p_{s,t}^2} p_{s,t}(1 - p_{s,t}) \\
&= (\sigma^2 \times r^2 \times s_2) \mathrm{Var}\left((x_i x_i^\top) \bullet Q\right)
\end{aligned}
$$

where the last equality is a simple calculation to calculate the term-by-term variance for $(x_i x_i^\top) \bullet Q$. Thus, Fact C.5 implies that the Kolmogorov distance between the distribution of $(x_i x_i^\top) \bullet Q$ and the Gaussian with the same mean and variance is at most

$$0.57 \frac{\sum_{s,t} \rho_{s,t}}{(\mathrm{Var}((x_i x_i^\top) \bullet Q))^{1.5}} \le 0.57(\sigma^2 \times r^2 \times s_2) \frac{\mathrm{Var}\left((x_i x_i^\top) \bullet Q\right)}{\mathrm{Var}^{1.5}\left((x_i x_i^\top) \bullet Q\right)} \le \frac{0.57(\sigma^2 \times r^2 \times s_2)}{\sqrt{V_Z}}$$

where the inequality comes from the assumption that the variance is at least V_Z. Therefore, $(x_i x_i^\top) \bullet Q$ has at least probability $0.5 - \frac{0.57(\sigma^2 \times r^2 \times s_2)}{\sqrt{V_Z}}$ of exceeding its expectation. By our choice of quantities in Appendix C.1, this probability is at least 0.4. Furthermore, $\mathbb{E}((x_i x_i^\top) \bullet Q) = x_i x_i^\top \bullet M$ is bigger than B and in turn bigger than $s_3 \times s_1$ (by our choice for these quantities). Thus, with probability at least 0.4, $(x_i x_i^\top) \bullet Q$ exceeds $s_3 \times s_1$.

Combined with the guarantee that $\mathbb{P}_Q((x_i x_i^\top) \bullet Q > s_3 \times s_1) > 0.99$ in the case where $\mathrm{Var}((x_i x_i^\top) \bullet Q) \le V_Z$ (cf. Equation (17)), we have shown that in all cases, $\mathbb{P}_Q((x_i x_i^\top) \bullet Q > s_3 \times s_1) > 0.4$ whenever $x_i x_i^\top \bullet M > B$.

Thus, we have shown that $\mathbb{E}[Z] = \sum_i \mathbb{P}_Q((x_i x_i^\top) \bullet Q > s_3 \times s_1) > 0.4 \times 0.01n = 0.004n$, since at least 0.01 fraction of points satisfy $x_i x_i^\top \bullet M > B$ and thus also satisfy $\mathbb{P}_Q((x_i x_i^\top) \bullet Q > s_3 \times s_1) > 0.4$. Note also that $Z \in [0, n]$ always, meaning that $\mathbb{E}[Z^2] \le n^2$. Since $0.004 \ge 4 \times q$ by our choice of q, it then follows from the Paley-Zygmund inequality that

$$\mathbb{P}(Z > 2 \times q \times |S|) \ge 0.25 \frac{(\mathbb{E}[Z])^2}{n^2} > \frac{0.25 \times 0.004^2 n^2}{n^2} = 4 \cdot 10^{-6}$$

showing the second claim above, and completing the proof of this lemma.

$\square$

C.1 Choice of Numerical Constants

This section shows how to pick the numerical constants q, s_1, s_2, s_3, V_Z and B. In the proof of Theorem C.1, these constants need to satisfy the following constraints:

1. $s_3 \ge 2$
2. q is at least a small constant since the sample complexity is inversely proportional to $1/q^2$.
3. See (13)

$$\frac{1}{s_2} + \frac{s_2}{s_3^2} \le 10^{-6}$$

4. See (15)

$$\frac{\sigma^2}{s_1} + \frac{4}{s_3} + \frac{s_2 \times \nu^4}{s_3 \times s_1^2} \le 10^{-6} \times q$$

5. See (17) $B \ge s_3 \times s_1 + 10\sqrt{V_Z}$
6. See (18) $\frac{0.57(\sigma^2 \times r^2 \times s_2)}{\sqrt{V_Z}} \le 0.1$
7. Paley-Zygmund $0.004 \ge 4 \times q$

Therefore, we pick the constants as follows:

1. ν, σ and r are numbers we get from the ℓ_∞ truncation, nothing to choose here.
2. $q = 0.001$
3. $s_2 = 10^7$
4. $s_3 = 10^{10}$
5. Solve for s_1 in terms of above in Constraint 4. Suffices to take $s_1 = \max(\sigma^2, \nu^2) \times 10^{10}$
6. Solve for $\sqrt{V_Z}$ using Constraint 6. Suffices to take $V_Z = 10^{16} \sigma^4 r^4$.
7. Solve for B using Constraint 5. Suffices to take $B = \max(\sigma^2, \sigma^2 r^2, \nu^2) \times 10^{20}$.

D Smoothness of Stability

The goal of this appendix is to prove Theorem D.11, the stability result we use in the proof of Theorem 1.3.

In Section 5, we sketched the proof of Theorem 5.1, which we formalized as Theorem C.1 in Appendix C. The key difference between Theorems C.1 and D.11 is that the former is a stability result concerning uncontaminated samples truncated according to some *fixed* vector close to the true mean. On the other hand, Theorem D.11 concerns samples truncated according to the coordinate-wise median-of-means estimate, which itself depends on the samples and is not fixed. Thus, much of this appendix is dedicated to showing the "Lipschitzness" of the stability of samples, as we truncate using different preliminary mean estimates.

We start with showing Lemma D.1, which states that with high probability, there exists a large subset of samples where in each dimension, at most a negligible fraction of the points have large magnitude. Then, Lemma D.9 shows that we can take the intersection between this large subset and the large subset of samples that are stable. Finally, Lemma D.10 and Theorem D.11 show that this subset is stable as long as the truncation is centered at any point close to the true mean, thus yielding the final stability result we desire.

For a vector $X_i \in \mathbb{R}^d$, we will use $X_{i,j}$ to refer to the j-th coordinate of X_i.

Lemma D.1. *Let P be a distribution over $\mathbb{R}^d$. For $X \sim P$, suppose for all $j \in [d]$, $\mathbb{E}[X_j^4] \leq \nu^4$. Then let S be a set of n i.i.d. points from P. Then there exists positive constants c_1 such that the following holds with probability $1 - \tau$ if $n \geq c_1(k^{1.5} + \log(1/\tau))$: there is a set $S' \subset S$ such that the following hold simultaneously:*

1. *$|S'| \geq 0.99|S|$*

2. *For each j in $[d]$, the number of points in S' with j-th coordinate larger in magnitude than $2\nu\sqrt{k}$ is at most $n/k^{1.5}$. Equivalently,*

$$\forall j \in [d]: \quad \sum_{i \in S} \mathbb{1}_{|X_{i,j}| \geq 2\nu\sqrt{k}} \leq \frac{n}{k^{1.5}} \tag{18}$$

Before providing the proof of Lemma D.1, we highlight why the result is not obvious. The first approach that one may try is to show that the original set S directly satisfies the claim, i.e., (with high probability) in each coordinate, the fraction of points with large magnitude in that coordinate is at most $k^{-1.5}$. At the population level, this is indeed true by the fourth moment assumption, i.e., for any fixed $i \in [n]$ and $j \in [d]$, the probability that $|X_{i,j}|$ is large is at most $O(1/k^2)$. However, for this to hold with probability $1 - \tau$, one requires roughly $k^{1.5}\log(1/\tau)$ samples even in 1 dimension[3], which would give a multiplicative dependence on $\log(1/\tau)$ instead of additive dependence.

The second approach that one may try would be the following: define S' to be the set of all "good" samples, where we say a sample is "good" if all of its coordinates are smaller than $c\nu\sqrt{k}$. For any fixed coordinate $j \in [d]$, the probability that the j-th coordinate may be larger than $c\nu\sqrt{k}$ may be as large as $1/k^2$. Thus the probability that a particular sample is bad may be arbitrarily close to 1 — for example, when coordinates are independent — and the resulting set S' will be too small with high probability.

Proof. We will assume that $k \geq C$ for a large enough constant. If k is smaller than the constant, then the result follows by applying Bernstein inequality and taking $S' = S$.

---

[3]The upper bound follows from a Chernoff bound, and the lower bound follows from the fact that Chernoff bounds are essentially tight for Bernoulli coins.

Let $S = \{Y_1, \ldots, Y_n\}$. For $i \in [n]$ and $j \in [d]$, we use $Z_{i,j}$ to denote $\mathbb{I}_{|Y_{i,j}| \geq c_2 \nu \sqrt{k}}$. For simplicity, we set $\alpha = k^{-1.5}/3$. Our goal is to show that the following integer program is feasible:

$$
\begin{aligned}
\text{variables} \quad & p_1, \ldots, p_n \\
\text{subject to} \quad & \forall j \in [d] : \sum_{i=1}^{n} p_i Z_{i,j} \leq 3\alpha n \\
& \sum_{i=1}^{n} p_i \geq 0.99n \\
& \forall i \in [n] : p_i \in \{0, 1\}.
\end{aligned}
\tag{F1}
$$

As argued above in the prose after the statement, one needs to argue about all the samples, and their coordinates, simultaneously to prove the statement. Since directly handling the feasibility program (F1) seems difficult, our argument will go in the following steps: (i) first consider the LP relxation of (F1), (ii) using duality theory, the LP relaxation is feasible iff the dual LP is infeasible, (iii) Simplify the dual LP and show that, with high probability, the resulting program is infeasible.

We begin by considering the LP relaxation.

$$
\begin{aligned}
\text{variables} \quad & p_1, \ldots, p_n \\
\text{subject to} \quad & \forall j \in [d] : \sum_{i=1}^{n} p_i Z_{i,j} \leq \alpha n \\
& \sum_{i=1}^{n} p_i \geq 0.999n \\
& \forall i \in [n] : p_i \in [0, 1].
\end{aligned}
\tag{F2}
$$

We first show that if the following LP relaxation, (F2), is feasible, then (F1) is also feasible.

Claim D.2 (Feasibility of (F2) implies feasibility of (F1)). *Suppose $n > 10^6$ and $\alpha \geq (4 \log n)/n$. If (F2) is feasible, then (F1) is also feasible.*

Proof. Let $p_1, \ldots, p_n$ be the feasible solution to (F2). Consider the following random assignment, for $i \in [n]$, $P_i \sim \text{Ber}(p_i)$ independently. We will show that, with non-zero probability, P_i's satisfy (F1). We will use the following inequality:

Fact D.3 (Chernoff Inequality). *Let $a_1, \ldots, a_n$ such that $a_i \in \{0, 1\}$. Let $W_1, \ldots, W_n$ be independent Bernoulli random variables and consider the random variable $Z = \sum_{i=1}^{n} W_i$. Then, with probability $1 - \tau$, $Z \leq 2(\mathbb{E} Z + \log(1/\tau))$.*

By Fact D.3, we get that the each of the inequalities in (F1) holds with probability $1 - 1/(2n)$ as long as $n\alpha \geq 2 \log(2n)$ and $n > 1000 \log(2n)$. The latter holds when $n \geq 10^6$. $\qquad\square$

Since $n > 10^6$ in our setting (as k is large and choosing c_1 to be large enough) and $\alpha = 1/(3k^{1.5})$, we have that $\alpha \geq 4(\log n)/n$ is equivalent to $n \geq 12k^{1.5} \log n$, which is satisfied when $n \geq 100k^{1.5} \log k$. The latter holds when $n \geq ck^{1.5} \log d$ for a large enough constant c. Thus in the remainder of this section, we will show that, with high probability, this LP program is indeed feasible. We begin by considering the following dual program:

$$
\begin{aligned}
\text{variables} \quad & w_1, \ldots, w_d, y_1, \ldots, y_n, x \\
\text{subject to} \quad & \sum_{i=1}^{n} y_i + \alpha n \sum_{j=1}^{d} w_j < 0.999nx \\
& \forall i \in [n] : y_i + \sum_{j=1}^{d} Z_{i,j} w_j \geq x \\
& z \geq 0, \quad \forall i \in [n] : y_i \geq 0, \quad \forall j \in [d] : w_j \geq 0.
\end{aligned}
\tag{F3}
$$

Suppose for the sake of contradiction that (F2) is infeasible. By Farkas' Lemma [GKT51], it means that the (dual) program in (F3) is feasible. Formally, we have the following claim:

Claim D.4 (LP Duality for (F2))**.** *(F3) is infeasible if and only if (F2) is feasible.*

Claim D.4 follows from Farkas' Lemma. We will argue that (F3) is infeasible by showing that the following program, which is feasible whenever (F3) is feasible, is infeasible.

$$
\begin{aligned}
\text{variables} \quad & w_1, \ldots, w_d, A \\
\text{subject to} \quad & \forall i \in A : \sum_{j=1}^{d} Z_{i,j} w_j \geq \alpha \Big(\sum_{j=1}^{d} w_j \Big) \\
& \forall j \in [d] : w_j \geq 0, \\
& A \subset [n], |A| \geq 10^{-3} n
\end{aligned} \tag{F4}
$$

(F4) states that for at least 10^{-3} fraction of i's in n, the following inequality holds: $\sum_{j=1}^{d} Z_{i,j} w_j \geq \alpha \|w\|_1$. The following claim relates the two programs above.

Claim D.5. *If (F3) is feasible, then (F4) is feasible.*

Proof. Let $y_1, \ldots, Y_n, w_1, \ldots, w_d, x$ be any feasible solution to (F3). Then the first constraint in (F3) that the average of y_i's is less than $0.999x - \alpha(\sum_{j=1}^{d} w_j)$. By Markov's inequality, the fraction of the y_i's such $y_i \geq (x - \alpha(\sum_{j=1}^{d} w_j))$ is at most $\frac{0.999x - \alpha(\sum_{j=1}^{d} w_j)}{(x - \alpha(\sum_{j=1}^{d} w_j))} \leq 0.999$. Thus the fraction of y_i's such that $y_i < (x - \alpha(\sum_{j=1}^{d} w_j))$ is at least 0.001.

Let $A \subset [n]$ be the set of such indices. For any $i \in A$, the second constraint in (F3) implies that $\sum_{j=1}^{d} Z_{i,j} w_j \geq x - y_i \geq \alpha(\sum_{j=1}^{d} w_j)$. This implies that (F4) is feasible. $\square$

In order to argue that (F4) is infeasible, we first consider a particular w. Using calculations provided below, it can be seen that the probability that a particular w satisfies (F4) is exponentially small in n. However, a direct approach at covering w seems difficult since w is a dense vector in $\mathbb{R}^d$ and $n = o(d)$. Using a randomized rounding mechanism, we show that it suffices to consider only sparse w as follows:

$$
\begin{aligned}
\text{variables} \quad & w_1, \ldots, w_d, A \\
\text{subject to} \quad & \forall i \in A : \sum_{j=1}^{d} Z_{i,j} w_j \geq 1 \\
& \forall j \in [d] : w_j \in \{0, 1\}, \\
& \sum_{j=1}^{d} w_j \leq \frac{2 \times 10^7}{\alpha} \\
& A \subset [n], |A| \geq 10^{-4} n
\end{aligned} \tag{F5}
$$

The following claim shows that if (F4) is feasible then (F5) is also feasible.

Claim D.6. *If (F4) is feasible, then (F5) is also feasible.*

Proof. Let $w_1, \ldots, w_d$ and A be the feasible solution to (F5). Set $q_j = \min(1, w_j/(\alpha\|w\|_1))$ for $j \in [d]$. Consider the following random assignment: set $W_j \sim \text{Ber}(q_j)$ independently for $j \in [d]$. We will show that with non-zero probability W_j's satisfy (F5). Consider the following events:

$$
\mathcal{E}_1 := \left\{ \sum_{j=1}^{d} W_j \leq 2 \times 10^{-7}(1/\alpha) \right\}, \quad \text{and} \quad \mathcal{E}_2 := \left\{ |\{i : \sum_{j=1}^{d} Z_{i,j} W_j \geq 1\}| \geq 10^{-4} n \right\} \tag{19}
$$

We will show that $\mathbb{P}\{cE_1\} \geq 1 - 5 \times 10^{-8}$ and $\mathcal{P}\{cE_2\} \geq 10^{-7}$. By a union bound, we will have that $\mathcal{E}_1 \cap \mathcal{E}_2$ has non-zero probability and thus (F5) is feasible.

Let $F := \{j \in [d] : W_j = 1\}$ be the set of coordinates where W_j is non-zero. Then $\mathbb{E}[|F|] = \mathbb{E}[\sum_{j=1}^d W_j] = \sum_{j=1}^d q_j \leq 1/\alpha$. Thus with probability at least $1 - 5 \times 10^{-8}$, we have that the number of non-zero W_j's is at most $\frac{2 \times 10^7}{\alpha}$. Equivalently, $\mathbb{P}\{\mathcal{E}_1\} \geq 1 - 5 \times 10^{-8}$.

We now focus on the second event $\mathcal{E}_2$. Let $S_1, \ldots, S_n$ be the subsets of $[d]$ such that $S_i = \{j \in [d] : Z_{i,j} = 1\}$, i.e., for each sample i, S_i is the set of indices where the coordinates are large. Consider the random variables $R_1, \ldots, R_n$, where for $i \in [n]$, $R_i := \sum_{j=1}^d Z_{i,j} W_j = \sum_{j \in S_i} Z_{i,j} W_j$. (F5) requires that for at least 10^{-4} fraction of i's, $R_i \geq 1$. Since $Z_{i,j}$'s are binary and fixed, we have that R_i is distributed as Binomial random variable and is thus anti-concentrated.

Fact D.7 (Anti-concentration of Binomial). *Let $X \sim Binomial(n, p)$ for some $n \in \mathbb{N}$ and $p \in [0, 1]$. Suppose $\mathbb{E}[X] \geq 1$. Then $\mathbb{P}\{X \geq 1\} \geq (1 - 1/e)$.*

Proof. Using the fact that $1 + x \leq e^x$ for all $x \in \mathbb{R}$, we get the following:
$$\mathbb{P}\{X \geq 1\} = 1 - \mathbb{P}\{X = 0\} = 1 - (1 - p)^n \geq 1 - (e^{-p})^n = 1 - e^{-np} \geq (1 - 1/e). \qquad \square$$

Consider a fixed $i \in A$. Then either there exists a $j \in S_i$ such that $q_j = 1$, or for all $j \in S_i$, $q_j < 1$. In the former case, we have that R_i is at least one since $W_j = 1$.

In the latter setting, we have that $q_j = w_j/(\alpha\|w\|_1)$ for all $j \in S_i$, and thus $\mathbb{E}[R_i] = \sum_{j \in S_i} Z_{i,j} q_j = \sum_{j=1}^d Z_{i,j} w_j/(\alpha\|w\|_1) \geq 1$. Applying Fact D.7 to any such $i \in A$, we get that the probability of R_i being positive is at least $1 - 1/e$. Let A' be set of i's such that $R_i \geq 1$, i.e., $A' = \{i : R_i \geq 1\}$. Thus combining the two cases above, we have the following:
$$\forall i \in A : \mathbb{P}\{i \in A'\} \geq 0.5. \tag{20}$$
Thus $\mathbb{E}[|A'|] \geq 0.5|A| \geq 5 \times 10^{-4}$. Since $|A'|$ lies in in $[0, n]$, applying Paley-Zygmund inequality to the random variable $|A'|$, we get the following:
$$\mathbb{P}\{|A'| \geq 10^{-4} n\} \geq \mathbb{P}\{|A'| \geq 0.2 \mathbb{E}[|A'|]\} \geq 0.64 \frac{(\mathbb{E}[|A'|])^2}{n^2} \geq 0.64 \times 25 \times 10^{-8} > 10^{-7}. \tag{21}$$
Equivalently, $\mathbb{P}\{\mathcal{E}_2\} \geq 10^{-7}$. This completes the proof. $\qquad \square$

Thus it suffices to show that, with high probability, (F5) is infeasible.

Lemma D.8 (Infeasibility of (F5)). *Under the setting of Lemma D.1 and when $k > 10^{26}$, there exists a constant $c_1 > 0$ such that if $n \geq c_1(k^{1.5} \log d + \log(1/\tau))$, then with probability $1 - \tau$, (F5) is infeasible.*

Proof. First consider any fixed $w = (w_1, \ldots, w_d)$ such that $w_i \in \{0, 1\}$ and $\sum_{j=1}^d w_j \leq 2 \times 10^7 \cdot (1/\alpha)$.

Consider the integer-valued random variables $R_1, \ldots, R_n$ such that $R_i = \sum_{j=1}^d Z_{i,j} w_j$, and observe that R_i's are i.i.d. random variables (since X_i's are i.i.d. random variables). Thus (F5) requires that at least $10^{-4}\%$ of R_i's are non-zero.

By the fourth moment bound on each coordinate, we have that $\mathbb{E}[Z_{i,j}] = \mathbb{P}\{X_{i,j} \geq 2\nu\sqrt{k}\} \leq 1/k^2$ for each i and j. Thus the expectation of each R_i is at most $\sum_{j=1}^d w_j \mathbb{E}[Z_{i,j}] \leq \sum_{j=1}^d w_j(1/k^2) \leq (2 \times 10^7)/(k^2\alpha) = (2 \times 10^7)/(k^2\alpha) = (6 \times 10^7)/\sqrt{k}$, which is less than 10^{-5} for k large enough. By Markov's inequality, the probability that $\mathbb{P}\{R_1 \geq 1\} \leq 10^{-5}$.

Thus by Chernoff bound (since R_i's are independent), with probability at least $1 - \exp(-c'n)$, the fraction of R_i's that are non-zero is at most 5×10^{-5}. Hence, with the same probability, this particular choice of w does not satisfy (F5). Since there are at most $d^{(2 \times 10^7)/\alpha}$ such choices of w, applying a union bound, we get that (F5) is infeasible with probability at least $1 - \exp((2 \times 10^7)/\alpha) \cdot \log d - c'n)$. The failure probability is at most τ when $n \gtrsim \log(1/\tau) + k^{1.5} \log d$. This concludes the proof. $\quad \square$

Since we assumed k is large enough, Lemma D.8 is applicable. Lemma D.8 implies that, with high probability, the program (F5) is infeasible. Hence, with the same high probability, the programs (F4) and (F3) are also infeasible, and the programs (F1) and (F2) are feasible. This completes the proof. $\qquad \square$

D.1 Lipschitz Argument

Let $h_{a,b}$ be as defined in Equation (1).

Lemma D.9 (Lipschitzness of Truncation Under Coordinatewise Regularity). *Let μ, μ' be vectors in $\mathbb{R}^d$ and let $a \in \mathbb{R}_+$ be greater than 2. Let $S = \{x_1, \ldots, x_n\} \subseteq \mathbb{R}^d$ be the set of n points. Suppose there exist a set $S_1 \subset [n]$ satisfying the following:*

$$|S_1| \geq 0.98n \ \text{ and } \ \left\| \frac{1}{|S_1|} \sum_{i \in S_1} (h_{a,\mu}(x_i) - \mu')(h_{a,\mu}(x_i) - \mu')^\top \right\|_{\mathcal{X}_k} \leq r. \tag{22}$$

Suppose also that, for some $\alpha \in (0,1)$, there exist a set $S_2 \subset [n]$ satisfying the following:

$$|S_2| \geq 0.99n \ \text{ and } \ \forall j \in [d]: \sum_{i \in S_2} \mathbb{I}_{|x_{i,j} - \mu_j| \geq a/2} \leq \alpha n. \tag{23}$$

Then, we have the following: there exists a set $S_3 \subset [n]$ such that for all $b \in \mathbb{R}^d$ satisfying $\|b - \mu\|_\infty \leq a/2$ and $\|b - \tilde{\mu}\|_\infty \leq a$, we have that

$$|S_3| \geq 0.97n \ \text{ and } \ \left\| \frac{1}{|S_3|} \sum_{i \in S_3} (h_{a,b}(x_i) - \mu')(h_{a,b}(x_i) - \mu')^\top \right\|_{\mathcal{X}_k} \leq 1.1r + 5a\alpha k\|b - \mu\|_\infty. \tag{24}$$

Proof. We will take $S_3 = S_1 \cap S_2$, which directly implies that $|S_3| \geq 0.97n$. For any $M \in Chi_k$, since $xx^\top \bullet M \geq 0$, we have the following:

$$\left\langle M, \frac{1}{|S_3|} \sum_{i \in S_3} (h_{a,\mu}(x_i) - \mu')(h_{a,\mu}(x_i) - \mu')^\top \right\rangle \leq \left\langle M, \frac{1}{|S_3|} \sum_{i \in S_1} (h_{a,\mu}(x_i) - \mu')(h_{a,\mu}(x_i) - \mu')^\top \right\rangle$$

$$\leq \frac{1}{0.97}r.$$

Let $F(b)$ be the following matrix:

$$F(b) = \frac{1}{|S_3|} \sum_{i \in S_3} (h_{a,b}(x_i) - \mu')(h_{a,b}(x_i) - \mu')^\top$$

We will establish that $\|F(b) - F(\mu)\|_{\chi_k} \leq 5a\alpha k \|b - \mu\|_\infty$, which establishes the lemma statement by the triangle inequality. In order to do that, we will show that $\|F(b) - F(\mu)\|_\infty \leq 5a\alpha\|b - \mu\|_\infty$ and then use Lemma A.6.

Consider an arbitrary (j, ℓ)-entry of these matrices. By abusing notation, when x and y are scalar, we use $h_{a,y}(x)$ to be the function from $\mathbb{R} \to \mathbb{R}$ defined analogously to Equation (1). Let $g(\cdot, \cdot)$ be the following function that is equal to the (j, ℓ) entry of the matrix $F(b)$, which is explicitly

$$g(b_j, b_\ell) = \frac{1}{|S_3|} \sum_{i \in S_3} (h_{a,b_j}(x_j) - \mu'_j)(h_{a,b_\ell}(x_\ell) - \mu'_\ell).$$

We will show that $g(\cdot, \cdot)$ is locally Lipschitz in its arguments. Consider a particular $i \in S_3$ and define the following:

$$g_i(b_j, b_\ell) = (h_{a,b_j}(x_{i,j}) - \mu'_j)(h_{a,b_\ell}(x_{i,\ell}) - \mu'_\ell)$$

Then, we can bound the difference for each sample by

$$|g_i(b_j, b_\ell) - g_i(\mu_j, \mu_\ell)|$$
$$= |(h_{a,b_j}(x_{i,j}) - \mu'_j)(h_{a,b_\ell}(x_{i,\ell}) - \mu'_\ell) - (h_{a,\mu_j}(x_{i,j}) - \mu'_j)(h_{a,\mu_\ell}(x_{i,\ell}) - \mu'_\ell)|$$
$$\leq |(h_{a,b_j}(x_{i,j}) - h_{a,\mu_j}(x_{i,j}))(h_{a,b_\ell}(x_{i,\ell}) - \mu'_\ell)| + |(h_{a,\mu_j}(x_{i,j}) - \mu'_j)(h_{a,b_\ell}(x_{i,\ell}) - h_{a,\mu_\ell}(x_{i,\ell}))|$$
$$\leq (a + \|b - \mu'\|_\infty) \cdot \|b - \mu\|_\infty \left(\mathbb{1}_{|x_{i,j} - \mu_j| \geq a - \|b - \mu\|_\infty} + \mathbb{1}_{|x_{i,\ell} - \mu_\ell| \geq \|b - \mu\|_\infty} \right)$$
$$\leq (a + \|b - \mu'\|_\infty) \cdot \|b - \mu\|_\infty \left(\mathbb{1}_{|x_{i,j} - \mu_j| \geq a/2} + \mathbb{1}_{|x_{i,\ell} - \mu_\ell| \geq a/2} \right)$$
$$\leq 2a \cdot \|b - \mu\|_\infty \left(\mathbb{1}_{|x_{i,j} - \mu_j| \geq a/2} + \mathbb{1}_{|x_{i,\ell} - \mu_\ell| \geq a/2} \right),$$

where we use that $|h_{a,y}(x) - h_{a,z}(x)| \leq |y - z|$, $|h_{a,y}(x) - z| \leq |a + y - z|$, and $|h_{a,y}(x) - h_{a,z}(x)|$ is non-zero only if $|x - y| \geq a - |y - z|$.

Combined with assumption Equation (23), this implies that

$$|g(b_j, b_\ell) - g(\mu_j, \mu_\ell)| \leq 2a \cdot \|b - \mu\|_\infty \cdot \frac{2\alpha}{0.97} \leq 5a\alpha\|b - \mu\|_\infty.$$

By Lemma A.6, we have the following:

$$\|F(b) - F(\mu)\|_{\mathcal{X}_k} \leq k\|F(b) - F(\mu)\|_\infty \leq 5a\alpha k \cdot \|b - \mu\|_\infty.$$

$\square$

D.2 Main Theorem

Lemma D.10. *Let S be a set of n i.i.d. data points from a distribution P over $\mathbb{R}^d$. Let the mean of P be μ, and covariance Σ such that $\|\Sigma\|_{\mathcal{X}_k} \leq \sigma^2$, and for all $i \in [d]$, $\mathbb{E}[X_i^4] \leq O(\sigma^4)$. Suppose $n = \Omega(k^2 \log d + \log(1/\tau))$. Let $a = \sigma\sqrt{k}$. With probability $1 - \tau$ over S, there exists a subset $S' \subset S$ with $|S'| \geq 0.95n$ such that for any b satisfying $\|b - \mu\|_\infty = O(\sigma)$, we have $h_{a,b}(S')$ is $(0.01, O(1), k)$-stable with respect to some μ' and σ with $\|\mu' - \mu\|_\infty \leq O(\sigma/\sqrt{k})$.*

Proof. Let P' be distribution of $h_{a,\mu}(P)$ and let μ' and Σ' be the mean and covariance of P'. This will be the μ' in the lemma statement. By Lemma 3.1, we get that (i) $\|\mu' - \mu\|_\infty \leq \sigma/\sqrt{k}$, , (ii) $\|\Sigma - \Sigma'\|_{\mathcal{X}_k} \leq O(\sigma^2)$, and (iii) P' is supported on the set $\{x : \|x - \mu\|_\infty \leq a\}$.

Applying Theorem C.1 to P' states that with probability at least $1 - \tau$, there exists a subset $S_1 \subset S$ with $|S_1| \geq 0.98n$ such that

$$\left\| \frac{1}{|S_1|} \sum_{i \in S_1} (h_{a,\mu}(x_i) - \mu')(h_{a,\mu}(x_i) - \mu')^\top \right\|_{\mathcal{X}_k} \leq O(\sigma^2). \tag{25}$$

By applying Lemma D.1 to $P - \mu$, with probability at least $1 - \tau$, there exists a subset $S_2 \subset S$ with $|S_2| \geq 0.99n$ such that

$$\forall j \in [d] : \sum_{i \in S_2} \mathbb{I}_{|x_{i,j} - \mu_j| \geq a/2} \leq O(k^{-1.5})n. \tag{26}$$

We can then apply Lemma D.9 to show that, conditioned on the above events, there exists a subset $S_3 \subset S$ with $|S_3| \geq 0.97n$ such that for all b such that $\|b - \mu\|_\infty \leq O(\sigma)$ and $\|b - \mu'\|_\infty \leq O(\sigma)$ (the latter holds by the triangle inequality for all b such that $\|b - \mu\|_\infty \leq O(\sigma)$), we have that

$$\left\| \frac{1}{|S_3|} \sum_{i \in S_3} (h_{a,b}(x_i) - \mu')(h_{a,b}(x_i) - \mu')^\top \right\|_{\mathcal{X}_k} \leq O(\sigma^2) + O(ak^{-1.5}k\|b - \mu\|_\infty) \leq O(\sigma^2). \tag{27}$$

By Proposition A.1, this implies S_3 contains a set S' satisfying the following: (i) $|S'| \geq 0.95n$ is $(0.1, O(1), k)$-stable with respect to μ' and σ. Thus, we choose S' in the lemma statement to be this set.

Taking a union bound, all the above events fail with probability at most $O(\tau)$. Reparameterizing yields the lemma statement.

$\square$

Theorem D.11. *Let S be a set of n i.i.d. data points from a distribution P over $\mathbb{R}^d$, and let T be a 0.01-corruption of S. Let $\tilde{\mu}$ be the coordinate-wise median-of-means estimate computed from set T. Let the mean of P be μ, and covariance Σ such that $\|\Sigma\|_{\mathcal{X}_k} \leq \sigma^2$, and for all $i \in [d]$, $\mathbb{E}[X_i^4] \leq O(\sigma^4)$. Suppose $n = \Omega(k^2 \log d + \log(1/\tau))$. Let $a = \sigma\sqrt{k}$. With probability $1 - \tau$ over S, for all T we have that there exists a subset $S' \subset T$ with $|S'| \geq 0.95n$ such that $h_{a,\tilde{\mu}}(S')$ is $(0.01, O(1), k)$-stable with respect to some μ' and σ with $\|\mu' - \mu\|_\infty \leq O(\sigma/\sqrt{k})$.*

Proof. By Fact 2.1, we know that with probability at least $1 - \tau$, we have $\|\tilde{\mu} - \mu\|_\infty \leq O(\sigma)O(1 + (\log(d/\tau))/n) = O(\sigma)$ by the assumption that n is sufficiently large.

Thus, we use $\tilde{\mu}$ as "b" in Lemma D.10 to yield the stability guarantee in the theorem statement.

The total failure probability is at most 2τ, and reparameterizing yields the theorem statement. $\square$

In Section 4, we use Theorem D.11 as the main technical ingredient to prove Theorem D.12, which in turn implies Theorem 1.3.

Theorem D.12 (Stronger version of our result). *Let $\epsilon \in (0, \epsilon_0)$ for small constant $\epsilon_0 > 0$. Let P be a multivariate distribution over $\mathbb{R}^d$, where the mean and covariance of P are μ and Σ respectively. Suppose $\|\Sigma\|_{\mathcal{X}_k} \leq 1$ and further suppose that for all $j \in [d]$, $\mathbb{E}[(X_j - \mu_j)^4] = O(1)$. Then, there is an algorithm such that, on input (i) the corruption parameter ϵ, (ii) the failure probability τ, (iii) the sparsity parameter k, and (iv) T, an ϵ-corrupted set of $n \gg (k^2 \log d + \log(1/\tau))/\epsilon)$ i.i.d. samples from P, it outputs $\widehat{\mu}$ satisfying $\|\widehat{\mu} - \mu\|_{2,k} = O(\sqrt{\epsilon})$ with probability $1 - \tau$ in $\mathrm{poly}(n, d)$ time.*

E   Information-Theoretic Error

Let $\mathcal{D}_k$ be the family of all distributions over $\mathbb{R}^d$ that satisfy the following:

1. For every $D \in \mathcal{D}_k$, the mean of D is k-sparse,

2. For every D in $\mathcal{D}_k$ the covariance of $\mathcal{D}$ is upper bounded by I in spectral norm, and

3. For every $D \in \mathcal{D}_k$ we have that $\mathbb{E}[(X_i - \mathbb{E}[X_i])^4] = O(1)$, where $X = (X_1, \ldots, X_d) \sim D$.

Lemma E.1. *Let $k \geq 1/\sqrt{\epsilon}$. Then there exist two distributions in D_1 and D_2 in $\mathcal{D}_k$ such that the following hold: (i) $d_{TV}(D_1, D_2) = \epsilon$, and (ii) The means of D_1 and D_2 are separated by $\Omega(\sqrt{\epsilon})$ in $\ell_{2,k}$-norm.*

Before giving the proof of Lemma E.1, we remark that the assumption of $k \geq 1/\sqrt{\epsilon}$ is mild. First, the assumption is independent of the ambient dimensionality d—the most challenging parameter regime in algorithmic robust statistics is when we fix a small ϵ and then take the dimensionality d to ∞. Second, the typical interesting sparsity regime is when k is super-constant but grows very slowly in d, say, logarithmically. The assumption that $k \geq 1/\sqrt{\epsilon}$ applies readily to the above regime.

Proof. Let D_1 be the distribution that places all of its mass at origin, i.e., $(0, \ldots, 0)$. Let D_2 be the distribution that places $(1 - \epsilon)$ probability mass at origin and places ϵ probability mass at y, where the first k-coordinates of y are α and the remaining $d - k$ coordinates are 0.

It is easy to see that the total variation distance between D_1 and D_2 is ϵ, and that $D_1 \in \mathcal{D}_k$. We will now show that $D_2 \in \mathcal{D}_k$ for a suitable value of α.

1. First the mean of D_2 is ϵy, which is k-sparse by construction.

2. We have that the covariance of D_2 is $\epsilon yy^\top - \epsilon^2 yy^\top = \epsilon(1 - \epsilon)yy^\top \preceq \epsilon yy^\top$, which is upper bounded by 1 in spectral norm if $\|y\|_2 \leq 1/\sqrt{\epsilon}$. Since $\|y\|_2 = \sqrt{k}\alpha$, we want that $\alpha \leq 1/\sqrt{k\epsilon}$.

3. Finally, let $X \sim D_2$. For every $i > k$, we have that $\mathbb{E}[(X_i - \mathbb{E}[X_i])^4] = 0$. For $i \in [k]$, $\mathbb{E}[(X_i - \mathbb{E}[X_i])^4] = \mathbb{E}[(X_i - \epsilon\alpha)^4] \leq 8(\mathbb{E}[X_i^4 + \epsilon^4\alpha^4]) = 8(\epsilon\alpha^4 + \epsilon^4\alpha^4) \leq 16\epsilon\alpha^4$, which is less than 16, if $\alpha \leq \epsilon^{-1/4}$.

Thus the above construction goes through as long as $\alpha \leq \min(1/\sqrt{k\epsilon}, \epsilon^{-1/4})$. When $k \geq 1/\sqrt{\epsilon}$, it suffices that $\alpha = 1/\sqrt{k\epsilon}$. Finally, we note that the difference in means of D_1 and D_2 is $\epsilon\|y\|_2 = \epsilon\sqrt{k}\alpha = \sqrt{\epsilon}$ for the chosen value of α. $\square$